

Bounds for the minimum diameter of integral point sets

SASCHA KURZ REINHARD LAUE

*University of Bayreuth
Department of Mathematics
D-95440 Bayreuth
Germany*

{sascha.kurz, reinhard.laue}@uni-bayreuth.de

Abstract

Geometrical objects with integral sides have attracted mathematicians for ages. For example, the problem to prove or to disprove the existence of a perfect box, that is, a rectangular parallelepiped with all edges, face diagonals and space diagonals of integer lengths, remains open. More generally an integral point set \mathcal{P} is a set of n points in the m -dimensional Euclidean space \mathbb{E}^m with pairwise integral distances where the largest occurring distance is called its diameter. From the combinatorial point of view there is a natural interest in the determination of the smallest possible diameter $d(m, n)$ for given parameters m and n . We give some new upper bounds for the minimum diameter $d(m, n)$ and some exact values.

1 Introduction

Geometrical objects with integral sides have long attracted mathematicians. One of the earliest results is due to the Pythagoreans and characterizes the smallest rectangle with integral sides and diagonals, more precisely, the integral rectangle with the smallest possible diameter where diameter denotes the largest occurring distance of the points. This is a rectangle with edge lengths 3 and 4 so that the diagonal has length 5 by Pythagoras' Theorem. In this context, a famous old open problem is to show the existence of a perfect box, a rectangular parallelepiped with all edges, face diagonals and space diagonals of integer lengths [2, 10]. Because this problem seems to be too hard for our current state of mathematics, the authors of [15] considered combinatorial boxes, i.e., convex bodies with six quadrilaterals as faces, and gave 20 examples of integral combinatorial boxes, one of which is proven to be minimal with regard to the diameter in [6]. In [1], it is shown that there exist infinitely many integral combinatorial boxes.

Generally, an integral point set \mathcal{P} is a set of n points in the m -dimensional Euclidean space \mathbb{E}^m with pairwise integral distances, where not all n points are contained in a hyperplane. From the combinatorial point of view, there is a natural interest in the minimum possible diameter $d(m, n)$ for given parameters m and n .

In the following, we will focus on bounds and exact numbers for $d(m, n)$. For a more general overview and applications on integral point sets and similar structures, we refer to [3]. Clearly, the condition $n \geq m + 1$ is necessary for an m -dimensional point set. Due to general constructions, see e.g. [4], the condition is also sufficient for the existence of an m -dimensional integral point set consisting of n points.

Theorem 1.1 *For $n \geq m + 1$ we have*

$$(a) \quad d(m, n) \leq \begin{cases} 2^{n-m+1} - 2 & \text{for } n - m \equiv 0 \pmod{2}, \\ 3(2^{n-m} - 1) & \text{for } n - m \equiv 1 \pmod{2}, \end{cases} \quad [4]$$

$$(b) \quad d(m, n) \leq (n - m)^{c \log \log(n-m)} \text{ for a sufficiently large constant } c, \quad [5]$$

$$(c) \quad \sqrt{\frac{3}{2m}} n^{1/m} < d(m, n), \quad [8]$$

$$(d) \quad \frac{1}{\sqrt[14]{14}} n^{1/2} < d(3, n) \text{ for } n \geq 5, \quad [8]$$

$$(e) \quad cn \leq d(2, n) \text{ for a sufficiently small constant } c, \quad [17]$$

$$(f) \quad d(n, n + 1) = 1,$$

$$(g) \quad 3 \leq d(m, n) \leq 4 \text{ for } m + 2 \leq n \leq 2m \text{ and } d(m, 2m) = 4, \quad [7, 16]$$

$$(h) \quad d(m, 2m + 1) \leq 8, \quad [16]$$

$$(i) \quad d(m, 2m + 2) \leq 13, \quad [16]$$

$$(j) \quad d(m, 3m) \leq 109, \quad [9]$$

$$(k) \quad \text{and } d(m, n - 1) \leq d(m, n).$$

We conjecture that $d(m - 1, n) \geq d(m, n)$. Each of the known bounds is increasing in n for fixed m and decreasing in m for fixed n . Several functional relations f between m and n exist for which $d(m, f(m))$ can be bounded from above by a constant. Examples are the inequalities of Theorem 1.1 (g, h, i, j) and of Theorem 1.2 (a) below.

Aside from general bounds, some exact values of $d(m, n)$ have been determined (the bold printed value $d(3, 9) = 16$ was incorrectly stated as $d(3, 9) = 17$ in the literature, see [3, 16]):

$$(d(2, n))_{n=3, \dots, 89} = 1, 4, 7, 8, 17, 21, 29, 40, 51, 63, 74, 91, 104, 121, 134, 153, 164, 196, 212, 228, 244, 272, 288, 319, 332, 364, 396, 437, 464, 494, 524, 553, 578, 608, 642, 667, 692, 754, 816, 897, 959, 1026, 1066, 1139, 1190, 1248, 1306, 1363, 1410, 1460, 1514, 1564, 1614, 1675, 1727, 1770, 1817, 1887, 1906, 2060, 2140, 2169, 2231, 2299, 2432, 2494, 2556, 2624, 2692, 2827, 2895, 2993, 3098, 3196, 3294, 3465, 3575, 3658, 3749, 3885, 3922, 4223, 4380, 4437, 4559, 4693, 4883; \quad [3, 13, 14]$$

$$(d(3, n))_{n=4, \dots, 23} = 1, 3, 4, 8, 13, \mathbf{16}, 17, 17, 17, 56, 65, 77, 86, 99, 112, 133, 154, 195, 212, 228; \quad [3, 12, 13, 16]$$

- $d(3, 5) = d(6, 8) = d(8, 10) = 3;$ [3]
- $d(m, m + 2) = 3$ for $8 \leq m \leq 23;$ [13]
- $d(m, n) = 4$ for $3 \leq m \leq 12$ and $m + 3 \leq n \leq 2m;$ [13]
- $d(m, n) = 4$ for $13 \leq m \leq 23$ and $2m - 9 \leq n \leq 2m.$ [13]

Our main results are

Theorem 1.2

- (a) $d(m, m^2 + m) \leq 17,$
- (b) $d(m, n - 2 + m) \leq d(2, n)$ for $9 \leq n \leq 122.$

And the exact values:

$d(2, n)_{n=90, \dots, 122} = 5018, 5109, 5264, 5332, 5480, 5603, 5738, 5938, 5995, 6052,$
 $6324, 6432, 6630, 6738, 6939, 7061, 7245, 7384, 7568, 7752, 7935, 8119, 8321,$
 $8406, 8648, 8729, 8927, 9052, 9211, 9423, 9534, 9794, 9905$

$d(3, 24) = 244,$

and the following two constructions:

Theorem 1.3 *If \mathcal{P} is a plane integral point set with diameter $diam(\mathcal{P})$ consisting of n points, where $n - 1$ points are situated on a line \overline{AB} , then $d(m, n - 2 + m) \leq diam(\mathcal{P}).$*

Theorem 1.4 *If \mathcal{P} is a planar integral point set consisting of n points, where $n - 1$ points are situated on a line \overline{AB} , the n -th point has distance h to the line \overline{AB} , and \mathcal{P}' is an $(m - 1)$ -dimensional point set consisting of n' points on an $(m - 1)$ -dimensional sphere of radius h , then we have for $m \geq 2$ that*

$$d(m, n + n' - 1) \leq \max(diam(\mathcal{P}), diam(\mathcal{P}')).$$

Aside from these results, we have:

Conjecture

- (a) $d(m, n) > (n - m)^{c \log \log(n - m)}$ for each fixed m and suitable large n and $c,$
- (b) $d(m, m + 2) = 3$ for $m \geq 8,$
- (c) $d(m - 1, n) \geq d(m, n),$
- (d) the bound of Theorem 1.3 is sharp for $m = 2, n \geq 9; m = 3, n \geq 22,$ and $m \geq 4, n \geq m^2 + m + 1,$ respectively,
- (e) $d(m, n - 2 + m) \leq d(2, n)$ for $m \geq 2.$

2 Proofs

The exact values of $d(m, n)$ were obtained by exhaustive enumeration via the methods described in [12, 13, 14]. For future improvements due to faster computers, we refer the reader to [11]. By looking at the plane integral point sets with diameter at most 10000, it turns out that those with minimum diameter and $9 \leq n \leq 122$ points




Figure 1: Plane integral point set \mathcal{P} with $n - 1$ points on a line.

have a very simple structure [13, 14]. They consist of $n - 1$ points situated on a line l plus one point P apart from l ; see Figure 1. An easy method is given in [13, 14] to construct such integral point sets with diameters at most $n^{c \log \log n}$ for a suitably large constant c , by choosing integers h^2 with many divisors. If we replace the point P by an $(m - 2)$ -dimensional regular simplex \mathcal{S} with edge length 1, we obtain an m -dimensional integral point set with the same diameter, which proves Theorem 1.3 and Theorem 1.2(b).

If we assume that we have a plane integral point set \mathcal{P} consisting of a line l with $n - 2$ points and a parallel line with two points P_1 and P_2 (see Figure 2), we can




Figure 2: Plane point set with points on two parallel lines.

slightly modify the construction of Theorem 1.3 and blow up P_1 and P_2 to regular $(m - 2)$ -dimensional simplices \mathcal{S}_1 and \mathcal{S}_2 of side length v . An example is given in Figure 3. Because the distance of two points $p_1 \in \mathcal{S}_1$ and $p_2 \in \mathcal{S}_2$ is either f or $w := \sqrt{f^2 + v^2}$ we have to choose a suitable v so that w is an integer.

Theorem 2.1 *If \mathcal{P} is a plane integral point set consisting of $n - 2$ points on a line and two points P_1 and P_2 on a parallel line with distance $\overline{P_1 P_2} = f$, and there exist positive integers u, w with $f^2 + v^2 = w^2$ then*

$$d(m, n - 2 + 2(m - 2)) \leq \max(w, \text{diam}(\mathcal{P})).$$




Figure 3: 3-dimensional integral point sets consisting of 8 points with minimum diameter.

Theorem 2.1 is tight in the cases $m = 2, n = 4, 7, 8$ and $m = 3, n = 8$, and also gives $d(m, 2m + 2) \leq 13$ (cf. Figure 3).

Besides blowing up points to regular simplices, another technique to construct integral point sets of arbitrary dimension is to truncate simplices. By truncating regular m -dimensional simplices of side length a at all vertices of a regular m -dimensional simplex of side length $b + 2a$, we get a point set \mathcal{P} with $m^2 + m$ points. For $m = 2$, we can easily determine the set of distances of \mathcal{P} to be $\{a, b, a + b, \sqrt{a^2 + ab + b^2}\}$, so $\text{diam}(\mathcal{P}) = a + b$. The smallest integral example is depicted in Figure 4 (here the two missing lines have edge length 7). It is indeed the smallest integral point set with $m = 2$ and $n = 6$.




Figure 4: Smallest integral hexagon.

For $m \geq 3$, the occurring distances of \mathcal{P} are given by

$$\mathcal{D} = \{a, b, a + b, \sqrt{a^2 + ab + b^2}, \sqrt{a^2 + 2ab + 2b^2}\},$$

so $\text{diam}(\mathcal{P}) = \sqrt{a^2 + 2ab + 2b^2}$. The smallest integral solution is given by $a = 7$ and $b = 8$ which lead to the m -dimensional integral point set with diameter 17 consisting of $m^2 + m$ points and proves Theorem 1.2(a). We have depicted this integral point set for $m = 2$ and $m = 3$ in Figure 5.

$\{7, 8\}$ and $\{2\,021\,231, 8\,109\,409\}$ are the only coprime pairs of integers with $a, b \leq 10\,000\,000$, where all values of \mathcal{D} are integers. It is not known whether infinitely many such parameter sets exist. We remark that a generalization of this approach to the other platonic solids does not lead to integral point sets by our methods.




Figure 5: 3-dimensional integral point sets from a truncated tetrahedron.

There is another important construction of integral point sets for. In Figure 6, we have depicted a plane integral point set consisting of 12 points of which 11 are col-




Figure 6: 2-dimensional integral point set with $n = 12$ and diameter 77.

linear. If we rotate the point P_i around the line \overline{AB} , it forms a circle with radius $8\sqrt{15}$. If we scale the plane integral point set of Figure 7 by a factor of 15, we can arrange 4 points on this circle, so that we get a 3-dimensional integral point set consisting of 15 points with diameter $d(3, 15) = 77$.




Figure 7: Smallest plane integral point set with $n = 4$ and no three points on a line.

Generally, for $m \geq 2$ we may consider an $(m-1)$ -dimensional sphere S that intersects P_i , has its center on the line \overline{AB} , and spans a hyperplane that is orthogonal to \overline{AB} . (If $m = 2$, then S consists of the point P_i and its reflection in \overline{AB} .) If it is possible to place n' points on this sphere with pairwise integral distances, then these points together with the points of the line \overline{AB} form an m -dimensional integral point set consisting of $n + n' - 1$ points. This gives the proof of Theorem 1.4. It is tight for $m = 3$, $13 \leq n \leq 21$. Nevertheless, we conjecture that Theorem 1.3 yields better bounds for $m = 3$ and $n \geq 22$.

Acknowledgement

The authors thank the anonymous referee for his/her helpful comments and suggestions on the paper.

References

- [1] R. C. Alperin, A quartic surface of integer hexahedra, *Rocky Mt. J. Math.* 31(1) (2001), 37–43.
- [2] R. K. Guy, *Unsolved problems in number theory. 2nd ed.* Unsolved Problems in Intuitive Mathematics. 1. New York, NY: Springer-Verlag. xvi, 285 pp., 1994.
- [3] H. Harborth, Integral distances in point sets, in *Butzer, P. L. (ed.) et al., Karl der Grosse und sein Nachwirken. 1200 Jahre Kultur und Wissenschaft in Europa. Band 2: Mathematisches Wissen. Turnhout: Brepols*, (1998), 213–224.
- [4] H. Harborth and A. Kemnitz, Diameters of integral point sets, in *Intuitive geometry, Pap. Int. Conf., Siófok/Hung. 1985, Colloq. Math. Soc. János Bolyai 48*, vol. 48, 255–266. North-Holland Publishing Company, 1987.
- [5] H. Harborth, A. Kemnitz and M. Möller, An upper bound for the minimum diameter of integral point sets, *Discrete Comput. Geom.*, 9(4) (1993), 427–432.
- [6] H. Harborth and M. Möller, Smallest integral combinatorial box, in G. E. Bergum et al., editors, *Applications of Fibonacci numbers*, vol. 7, 153–159. Kluwer Academic Publishers, Dordrecht, 1996.
- [7] H. Harborth and L. Piepmeyer, Points sets with small integral distances, in P. Gritzmann and B. Sturmfels, eds., *Applied geometry and discrete mathematics, Festschr. 65th Birthday Victor Klee*, vol. 4 of *DIMACS, Ser. Discret. Math. Theor. Comput. Sci.*, 319–324. American Mathematical Society, 1991.
- [8] H.-J. Kanold, Über Punktmengen im k-dimensionalen euklidischen Raum, *Abh. Braunschw. Wiss. Ges.*, 32 (1981), 55–65.
- [9] A. Kemnitz, Punktmengen mit ganzzahligen Abständen, Habilitationsschrift, TU Braunschweig, 1988.
- [10] V. Klee and S. Wagon, *Old and new unsolved problems in plane geometry and number theory*. The Dolciani Mathematical Expositions 11, Washington, DC: Mathematical Association of America, xv, 333 pp., 1991.
- [11] S. Kurz, <http://www.wm.uni-bayreuth.de/index.php?id=241>.
- [12] S. Kurz, On the characteristic of integral point sets in \mathbb{E}^m , *Australas. J. Combin.* 36 (2006), 241–248.

- [13] S. Kurz, *Konstruktion und Eigenschaften ganzzahliger Punktmengen*, PhD Thesis, Bayreuth. Math. Schr. 76. Universität Bayreuth, 2006.
- [14] S. Kurz and A. Wassermann, On the minimum diameter of plane integral point sets, (preprint).
- [15] B. E. Peterson and J. H. Jordan, Integer hexahedra equivalent to perfect boxes, *Amer. Math. Monthly* 102(1) (1995), 41–45.
- [16] L. Piepmeyer, Räumliche ganzzahlige Punktmengen, Master's Thesis, TU Braunschweig, 1988.
- [17] J. Solymosi, Note on integral distances, *Discrete Comput. Geom.* 30(2) (2003), 337–342.

(Received 16 Oct 2006; revised 20 Apr 2007)