

HOW RANDOM ARE A LEARNER'S MISTAKES ?

JOEL RATSABY
RATSABY@ARIEL.AC.IL

Department of Electrical and Electronics Engineering
Ariel University Center
Ariel 40700, ISRAEL

ABSTRACT. Given a random binary sequence $X^{(n)}$ of random variables, X_t , $t = 1, 2, \dots, n$, for instance, one that is generated by a Markov source (teacher) of order k^* (each state represented by k^* bits). Assume that the probability of the event $X_t = 1$ is constant and denote it by β . Consider a learner which is based on a parametric model, for instance a Markov model of order k , who trains on a sequence $x^{(m)}$ which is randomly drawn by the teacher. Test the learner's performance by giving it a sequence $x^{(n)}$ (generated by the teacher) and check its predictions on every bit of $x^{(n)}$. An error occurs at time t if the learner's prediction Y_t differs from the true bit value X_t . Denote by $\xi^{(n)}$ the sequence of errors where the error bit ξ_t at time t equals 1 or 0 according to whether the event of an error occurs or not, respectively. Consider the subsequence $\xi^{(\nu)}$ of $\xi^{(n)}$ which corresponds to the errors of predicting a 0, i.e., $\xi^{(\nu)}$ consists of the bits of $\xi^{(n)}$ only at times t such that $Y_t = 0$. In this paper we compute an estimate on the deviation of the frequency of 1s of $\xi^{(\nu)}$ from β . The result shows that the level of randomness of $\xi^{(\nu)}$ decreases relative to an increase in the complexity of the learner.

1. INTRODUCTION

Let $X^{(n)} = X_1, \dots, X_n$ be a sequence of binary random variables drawn according to some unknown joint probability distribution $\mathbb{P}(X^{(n)})$. Consider the problem of learning to predict the next bit in a binary sequence drawn according to \mathbb{P} . For training, the learner is given a finite sequence $x^{(m)}$ of bits $x_t \in \{0, 1\}$, $1 \leq t \leq m$, drawn according to \mathbb{P} and estimates a model \mathcal{M} that can be used to predict the next bit of a partially observed sequence. After training, the learner is tested on another sequence $x^{(n)}$ drawn according to the same unknown distribution \mathbb{P} . Using \mathcal{M} he produces the bit y_t as a prediction for x_t , $1 \leq t \leq n$. Denote by $\xi^{(n)}$ the corresponding binary sequence of mistakes where $\xi_t = 1$ if $y_t \neq x_t$ and is 0 otherwise. We pose the following main question: how random is $\xi^{(n)}$?

It is clear that the sequence of mistakes should be random since the test sequence $x^{(n)}$ is random. It may also be that because the learner is using a model of a finite structure (or a finite description-length) then it may somehow introduce dependencies and cause $\xi^{(n)}$ to be less random than $x^{(n)}$. And yet by another intuition, perhaps the fact that the learner is of a finite complexity limits its ability to 'deform' (or distort) randomness of $x^{(n)}$? These are all valid initial guesses that relate to this main question. We note that our basis for saying that \mathcal{M} has a finite structure stems from it being an element of some regular hypothesis class, for instance, having a finite VC-dimension as is often the case in a learning setting (see for instance structural risk minimization of [18]). In the current paper, we are not interested in the learner's performance (as modeled for instance by Valiant's

PAC framework [17, 15]) but instead we take a data-centric view and ask how much influence does the learner has on the stochastic properties of the errors. We view the learner as an entity that 'interferes' with the randomness that is inherent in the sequence to be predicted and through his predictions creates a sequence of mistakes that has a different stochastic character.

To the best of the our knowledge, this main question has not been raised nor studied in information or learning theory. Our aim in this paper is to make a first attempt at answering it. Our approach will be a practical one, where we build on a specific learning setting and use it for our analysis. In this setting we consider a teacher that uses a probability distribution \mathbb{P} based on a Markov model with a certain complexity. The learner has access to a hypothesis class of Boolean decision rules that are also based on Markov models. Hence, learning amounts to the estimation of parameters of a finite-order Markov model (this has been studied extensively, see for instance [7, 13]). As this is only a first attempt, it is obvious that many different settings can be analyzed, in particular, more general ones. For instance, considering a learner that in addition to parametric estimation, does statistical-model-selection [4].

The remainder of the paper is organized as follows: in section 2 we give a brief introduction to the notion of randomness, in section 3 we define the problem and state our result (theorem), and in section 4 we prove the theorem.

2. RANDOMNESS OF A FINITE SEQUENCE

The notion of randomness of finite objects (binary sequences) aims to explain the intuitive idea that a sequence, whether finite or infinite, should be measured as being more unpredictable if it possess fewer regularities (patterns). There is no formal definition of randomness but there are three main properties that a random binary string of length n must intuitively satisfy [16]. The first property is the so-called *stochasticity* or frequency stability of the sequence which means that any binary word of length $k \leq n$ must have the same frequency limit (equal to 2^{-k}). This is basically the notion of normality that Borel introduced and is related to the degree of unpredictability of the sequence. The second property is *chaoticity* or disorderliness of the sequence. A sequence is less chaotic (less complex) if it has a short description, i.e., if the minimal length of a program that generates the sequence is short. The third property is *typicalness*. A random sequence is a typical representative of the class Ω of all binary sequences. It has no specific features distinguishing it from the rest of the population. An infinite binary sequence is typical if each small subset E of Ω does not contain it (the correct definition of a 'small' set was given by Martin Lőf [12]).

As mentioned in section 1, our interest in this paper is essentially to ask what 'interference' does a learner have on the randomness of a test sequence. It appears essential that we look not only on the randomness of the object itself (the test sequence $x^{(n)}$) but also at the interfering entity—the learner, specifically, its algorithmic component that is used for prediction. Related to this, there is an area of research that studies algorithmic randomness which is the relationship between complexity and stochasticity of finite and infinite binary sequences [3]. Algorithmic randomness was first considered by von Mises in 1919 who defined an infinite binary sequence α of zeros and ones as random if it is unbiased, i.e. if the frequency of zeros goes to $1/2$, and every subsequence of α that we can extract using an admissible selection rule (see definition below) is also not biased. Kolmogorov and Loveland [11, 10] proposed a more permissive definition of an admissible selection rule as any (partial) computable process which, having read any n bits of an infinite binary sequence α , picks a bit that has not been read yet, decides whether it should be

selected or not, and then reads its value. When subsequences selected by such a selection rule pass the unbiasedness test they are called Kolmogorov-Loveland stochastic (KL-stochastic for short). Martin Löff [12] introduced a notion of randomness which is now considered by many as the most satisfactory notion of algorithmic randomness. His definition says precisely which infinite binary sequences are random and which are not. The definition is probabilistically convincing in that it requires each random sequence to pass every algorithmically implementable statistical test of randomness.

Let us briefly define what is meant by a selection rule. As mentioned above, this is a principal concept used as part of tests of randomness of sequences. Let $\{0, 1\}^*$ be the space of all finite binary sequences and denote by $\{0, 1\}^n$ the set of all finite binary sequences of length n . An admissible *selection rule* R is defined ([8, 19]) based on three partial recursive functions f, g and h on $\{0, 1\}^*$. Let $x^{(n)} = x_1, \dots, x_n$. The process of selection is recursive. It begins with an empty sequence \emptyset . The function f is responsible for selecting possible candidate bits of $x^{(n)}$ as elements of the subsequence to be formed. The function g examines the value of these bits and decides whether to include them in the subsequence. Thus f does so according to the following definition: $f(\emptyset) = i_1$, and if at the current time k a subsequence has already been selected which consists of elements x_{i_1}, \dots, x_{i_k} then f computes the index of the next element to be examined according to element $f(x_{i_1}, \dots, x_{i_k}) = i$ where $i \notin \{i_1, \dots, i_k\}$, i.e., the next element to be examined must not be one which has already been selected (notice that maybe $i < i_j$, $1 \leq j \leq k$, i.e., the selection rule can go backwards on x). Next, the two-valued function g selects this element x_i to be the next element of the constructed subsequence of x if and only if $g(x_{i_1}, \dots, x_{i_k}) = 1$. The role of the two-valued function h is to decide when this process must be terminated. This subsequence selection process terminates if $h(x_{i_1}, \dots, x_{i_k}) = 1$ or $f(x_{i_1}, \dots, x_{i_k}) > n$. Let $R(x^{(n)})$ denote the selected subsequence. By $K(R|n)$ we mean the length of the shortest program computing the values of f, g and h given n .

From the previous discussion, we know that there are two principal measures related to the information content in a finite sequence $x^{(n)}$, stochasticity (unpredictability) and chaoticity (complexity). An infinitely long binary sequence is regarded random if it satisfies the principle of stability of the frequency of 1s for any of its subsequences that are obtained by an admissible selection rule [8]. Kolmogorov showed that the stochasticity of a finite binary sequence x may be precisely expressed by the deviation of the frequency of ones from some $0 < p < 1$, for any subsequence of $x^{(n)}$ selected by an admissible selection rule R of finite complexity $K(R|n)$ where for an object x given another object y he defined in [9] the complexity of x as

$$K(x|y) = \min\{l(\pi) : \phi(\pi, y) = x\} \quad (2.1)$$

where $l(\pi)$ is the length of the sequence π , ϕ is a universal partial recursive function which acts as a description method, i.e., when provided with input (π, y) it gives a specification for x (for more on that see the nice paper by [19]). The chaoticity of $x^{(n)}$ is large if its complexity is close to its length n . The classical work of [8, 1, 2, 19] relates chaoticity to stochasticity. In [1, 2] it is shown that chaoticity implies stochasticity. For a binary sequence s , let us denote by $\|s\|$ the number of 1s in s , then this can be seen from the following relationship (with $p = 1/2$):

$$\left| \frac{\|R(x^{(n)})\|}{l(R(x^{(n)}))} - \frac{1}{2} \right| \leq c \sqrt{\frac{n - K(x^{(n)}|n) + K(R|n) + 2 \log K(R|n)}{l(R(x^{(n)}))}} \quad (2.2)$$

where $l(R(x^{(n)}))$ is the length of the subsequence selected by R and $c > 0$ is some absolute constant. From this it is apparent that as the chaoticity of $x^{(n)}$ grows the stochasticity of the selected subsequence $R(x^{(n)})$ grows (the bias from $1/2$ decreases). Also, the information content of the selection rule, namely $K(R|n)$, has a direct effect on this relationship: the lower $K(R|n)$ the stronger the stability (smaller deviation of the frequency of 1s from $1/2$). In [5] the other direction which shows that stochasticity implies chaoticity is proved.

So referring back to the initial guesses we made in section 1 concerning our expectation about the randomness of the error sequence $\xi^{(n)}$, we now have a better clue and expect that the more algorithmically complex a learner's prediction rule is the more that it can distort (introduce bias into) the randomness of the test sequence $x^{(n)}$. As will be shown, rather than resorting to algorithmic randomness theory (which requires dealing with the non-practical and hard to analyze notion of Kolmogorov complexity) a direct combinatorial will do.

3. PROBLEM DEFINITION

Let us denote by $\{0, 1\}^*$ the space of all finite binary sequences. The learning problem consists of predicting the next bit value in a sequence $X^{(n)} = X_1, X_2, \dots, X_n$ of binary random variables drawn randomly according to a probability distribution \mathbb{P} which is defined based on a Markov chain with a finite number of states s . For convenience, we let the state space be the set of natural numbers between 0 and $2^k - 1$ and represent each state $s \in \mathbb{S}_k \equiv \{0, 1, \dots, 2^k - 1\}$ by its unique binary vector $b = [b(1), b(2), \dots, b(k)] \in \{0, 1\}^k$. We alternatively refer to states either by their decimal number s or their binary vector b .

Associated with these states is the transition matrix T where the i^{th} row represents the conditional probability distribution given state i . Consider drawing a random sequence $X^{(n)}$ using the chain by repeatedly making a transition from the current state S_t at time t to the next state S_{t+1} as dictated by T_k . Suppose that $S_t = i$ and $S_{t+1} = j$ then the teacher emits for X_{t+1} the bit value that is appended to b_i in order to obtain b_{t+1} , i.e., the value X_{t+1} satisfies

$$b_j = [b_i(2), b_i(3), \dots, b_i(k), X_{t+1}]$$

where b_j and b_i are the binary vectors corresponding to the states j and i , respectively. Clearly, the structure of the Markov model allows only two outgoing transitions from any given state since X_{t+1} is binary; we call them a *type-1* and *type-0* transitions. Let us denote by \mathcal{M}_k a Markov model (chain) based on transition matrix T_k . We use k^* to denote the order of the teacher's Markov chain (on which \mathbb{P} is based). For any binary sequence $x^{(k+n)}$ of length at least $k > 0$ if we let $b_k \equiv [b_k(1), \dots, b_k(k)] = [x_1, \dots, x_k]$ and define recursively the value of

$$b_t = [b_{t-1}(2), \dots, b_{t-1}(k), x_t] \tag{3.1}$$

for all times $t > k$, where $x_t \in \{0, 1\}$ then the probability of $x^{(k+n)}$ with respect to the teacher's model is defined by

$$\begin{aligned} \mathbb{P}(x^{(k+n)}) &= \mathbb{P}(S_1 = b_k) \mathbb{P}(S_2 = b_{1+k} | S_1 = b_k) \\ &\quad \cdots \mathbb{P}(S_{n+1} = b_{n+k} | S_n = b_{n-1+k}). \end{aligned} \tag{3.2}$$

Henceforth, all random binary sequences are assumed to be drawn according to this probability distribution \mathbb{P} which is based on model \mathcal{M}_{k^*} . Neither the value k^* nor the parameters of \mathcal{M}_{k^*} are known to the learner. From basic theory on finite Markov chains, since the matrix T_{k^*} is stochastic (i.e., the sum of the elements in

any row equals 1) then \mathcal{M}_{k^*} has a stationary probability distribution, which we denote by \mathbb{P}^* . Let us denote by

$$\beta = \mathbb{P}^*(X_t = 1) \quad (3.3)$$

at any time t .

As a learner, we consider an algorithm that assumes a Markov model \mathcal{M}_k of dimension $k \geq 1$. The learner estimates the probability parameters

$$p_{ij} \equiv \mathbb{P}(S_{t+1} = j | S_t = i), \quad i, j \in \mathbb{S}_k$$

by

$$\hat{p}_{ij} = \frac{\#\{S_{t+1} = j, S_t = i\}}{\#\{S_t = i\}}$$

where $\#\{S_t = i\}$ denotes the number of times state i appeared in the training sequence $x^{(k+m)}$ which drawn randomly by the teacher according to \mathbb{P} and so \hat{p}_{ij} are the frequency of transitions. The first k bits of $x^{(k+m)}$ indicate the initial state of the learner's model as it reads the training sequence, m is the number of state transitions taken by the teacher's model to generate the sequence. Note that p_{ij} are unknowns since they represent the probability of transition from state i to state j in the learner's model \mathcal{M}_k given a random sequence generated according to the teacher probability distribution \mathbb{P} (which is based on the unknown model \mathcal{M}_{k^*}).

After training, the learner is tested on a random test sequence $X^{(k+n)}$ obtained from the teacher based on \mathcal{M}_{k^*} . The learner is repeatedly asked to predict the next bit for each of the last n random bits X_{k+1}, \dots, X_{k+n} , where as above, the first k bits of $X^{(k+n)}$ indicate the starting state of the learner's model as it reads the test sequence. The learner computes the posterior probability $P(X_{t+1} = 1 | S_t)$, based on the learnt model \mathcal{M}_k , which is the probability that the next bit $X_{t+1} = 1$ given that the current state is S_t (at any given time t the current state consists of the last k bits seen in the test sequence up to t). The learner's decision (prediction) is based on the maximum *a posteriori* probability which is defined as follows: suppose that the current state is i then the decision is

$$d(i) \equiv \begin{cases} 1 & \text{if } \hat{p}(1|i) > 1 - \hat{p}(1|i) \\ 0 & \text{otherwise,} \end{cases} \quad (3.4)$$

$i \in \mathbb{S}_k$, where $\hat{p}(1|i)$ is defined as \hat{p}_{ij} for the state j whose $b_j = [b_i(2), \dots, b_i(k), 1]$ (a type-1 transition from state i) and the corresponding true probability (measured according to \mathbb{P}) is denoted by $p(1|i) = p_{ij}$. Note that (3.4) may be expressed alternatively as

$$d(i) = \begin{cases} 1 & \text{if } \hat{p}(1|i) > \frac{1}{2} \\ 0 & \text{otherwise.} \end{cases} \quad (3.5)$$

Denote by $m_i \geq 0$ the number of times state i was entered as the teacher scans the training sequence $x^{(k+m)}$ from $t = k + 1$ up to $t = k + m$ (as mentioned above, the initial state at time $t = k + 1$ is the state whose $b = [x_1, x_2, \dots, x_k]$). We will sometimes refer to m_i as the i^{th} subsample size. Note that $m_i, i \in \mathbb{S}_k$, are dependent random variables since the Markov chain may visit each state a random number of times and they all must satisfy $\sum_{i=0}^{2^k-1} m_i = m$. We claim that the $\hat{p}(1|i), i \in \mathbb{S}_k$, are independent random variables when conditioned on the vector $[m_0, \dots, m_{2^k-1}]$ (which we henceforth denote by \underline{m}). In order to see this, consider a training sequence $x^{(m+k)}$ generated by the teacher according to (3.2). Let us denote the corresponding sequence of states by $\sigma^{(m)} = (\sigma_1, \dots, \sigma_m)$ with $\sigma_i \in \mathbb{S}_k$.

Then by (3.2) we have

$$\begin{aligned} \mathbb{P}\left(x^{(m+k)}\right) &= \mathbb{P}\left(\sigma^{(m)}\right) \\ &= \mathbb{P}\left(S_1 = \sigma_1\right) \mathbb{P}\left(S_2 = \sigma_2 \mid S_1 = \sigma_1\right) \mathbb{P}\left(S_3 = \sigma_3 \mid S_2 = \sigma_2\right) \\ &\quad \cdots \mathbb{P}\left(S_m = \sigma_m \mid S_{m-1} = \sigma_{m-1}\right) \end{aligned}$$

For any $i \in \mathbb{S}_k$ denote by $N_\sigma(1|i)$ the number of type-1 transitions from state i in the sequence σ . Without loss of generality, let us assume that always initially the state is $i = 0$ so that we can write for the first factor $\mathbb{P}\left(S_1 = \sigma_1 \mid 0\right)$. Since all state transitions are either type-0 or type-1 transition then we have

$$\mathbb{P}\left(x^{(m+k)} \mid \underline{m}\right) = \prod_{i \in \mathbb{S}_k} (p(1|i))^{N_\sigma(1|i)} (1 - p(1|i))^{m_i - N_\sigma(1|i)} \quad (3.6)$$

where $p(1|i)$ was defined above. Let α be a non-negative integer constant and define the vector function $N(i) \equiv [N(1|i), \alpha - N(1|i)]$. Associate a conditional probability function for the random variable $N(i)$ as

$$\mathbb{P}\left(N(i) = [\ell, \alpha - \ell] \mid \alpha\right) = (p(1|i))^\ell (1 - p(1|i))^{\alpha - \ell}.$$

Then (3.6) may be written as

$$\mathbb{P}\left(x^{(m+k)} \mid \underline{m}\right) = \prod_{i \in \mathbb{S}_k} \mathbb{P}\left(N(i) = [N_\sigma(1|i), m_i - N_\sigma(1|i)] \mid m_i\right). \quad (3.7)$$

Note, conditioned on m_i , the event " $N(i) = [N_\sigma(1|i), m_i - N_\sigma(1|i)]$ " is equivalent to the event " $\hat{p}(1|i) = \frac{N_\sigma(1|i)}{m_i}$ ". Hence alternatively (3.7) can be expressed as

$$\mathbb{P}\left(x^{(m+k)} \mid \underline{m}\right) = \prod_{i \in \mathbb{S}_k} \mathbb{P}\left(\hat{p}(1|i) = \frac{N_\sigma(1|i)}{m_i} \mid \underline{m}\right). \quad (3.8)$$

The right side of (3.8) is a product of conditional probability functions of the random variables $\hat{p}(1|i)$. So conditioned on \underline{m} , the event that a sequence $x^{(m+k)}$ is generated by the teacher is equivalent to the event that a sequence of state transitions has frequencies $\hat{p}(1|i)$ that independently take the particular values $N_\sigma(1|i)/m_i$ according to $x^{(m+k)}$.

Let us state the main result of this paper.

Theorem 1. *Let $0 < \delta < 1$ and k, ℓ, m, n be positive integers. Let \mathbb{P} be an unknown probability distribution based on a Markov chain and denote by \mathbb{P}^* the corresponding stationary probability distribution. Suppose that a teacher generates a binary sequence $X^{(n)} = X_1, X_2, \dots, X_n$ by repeatedly drawing X_t according \mathbb{P} . Denote by $\beta = \mathbb{P}^*(X_t = 1)$ and let $x^{(k+m)}$ be a given randomly drawn training sequence.*

Suppose that a learner uses $x^{(k+m)}$ to estimate the values of the parameters of a Markov model \mathcal{M}_k with 2^k states and denote by ρ the average of the probabilities (according to \mathbb{P}) that the frequency (with respect to $x^{(k+m)}$) of emitting a 1 (when transitioning from a current state to the next state in \mathcal{M}_k) is larger than $1/2$. Suppose that the learner is tested incrementally on a randomly drawn sequence $x^{(k+n)}$ generated according to \mathbb{P} . The learner predicts an output bit y_t for every input bit x_t in $x^{(k+n)}$ using \mathcal{M}_k .

Denote by $\xi^{(k+n)}$ the sequence of mistakes where $\xi_t = 1$ if $y_t \neq x_t$, and $\xi_t = 0$ otherwise, $k + 1 \leq t \leq k + n$. Suppose that the subsequence $\xi^{(\nu)}$ of mistakes

corresponding to 0-valued predictions is of length $\nu \geq \ell$. Denote by

$$\epsilon(\ell, k, \rho, \delta) \equiv \sqrt{\max \left\{ \frac{1}{\ell} \left(\ln \left(\frac{4}{\delta} \right) + 3\rho 2^{k-1} \left(\ln \left(\frac{1}{\rho} \right) + 1 \right) + k \right), \frac{1}{2^{k-2}\rho} \ln \left(\frac{4}{\delta} \right) \right\}}.$$

Then for any $0 < \delta < 1$, with confidence at least $1 - \delta$ the deviation between β and the frequency of 1s of $\xi^{(\nu)}$ is bounded as

$$\left| \frac{1}{\nu} \sum_{j=1}^{\nu} \xi_j - \beta \right| \leq \epsilon(\ell, k, \rho, \delta)$$

where it is assumed that $(1 + \epsilon(\ell, k, \rho, \delta))\rho < 1/2$ and $\epsilon(\ell, k, \rho, \delta) < 1/2$.

4. PROOF OF THEOREM 1

Referring to (3.5), as the decision rule of the learner let us denote by the binary vector

$$d = [d(0), \dots, d(2^k - 1)]. \quad (4.1)$$

The $d(i)$ depend on the random variables $\hat{p}(1|i)$ hence d fully describes the learner's prediction rule after having learnt the model \mathcal{M}_k based on the random training sequence $x^{(k+m)}$. That is, d describes every possible prediction that can be made for all possible situations (present states).

We have from (3.5) that

$$\mathbb{P}(d(i) = 1) = \mathbb{P} \left(\hat{p}(1|i) > \frac{1}{2} \right) \equiv \rho_i \quad (4.2)$$

from which it is clear that the elements of d are nonidentically distributed Bernoulli random variables. Note that conditioned on \underline{m} they are independent (see above argument concerning the conditional independence of the random variables $\hat{p}(1|i)$).

For a binary vector d denote by $\|d\|$ the l_1 -norm (or Hamming weight) of d . Denoting by

$$\rho \equiv \left(\frac{1}{2^k} \sum_{i=0}^{2^k-1} \rho_i \right) \quad (4.3)$$

then the expected number of 1s in d is

$$\mathbb{E}[\|d\|] = \mathbb{E} \left[\sum_{i=0}^{2^k-1} d(i) \right] = 2^k \rho.$$

Let us define the following set,

$$A_\epsilon^{(k)} = \left\{ d \in \{0, 1\}^{2^k} : 1 - \epsilon \leq \frac{\|d\|}{2^k \rho} \leq 1 + \epsilon \right\}. \quad (4.4)$$

The probability of not falling in $A_\epsilon^{(k)}$ is

$$\mathbb{P} \left(d \notin A_\epsilon^{(k)} \right) = \sum_{\underline{m}} \mathbb{P} \left(d \notin A_\epsilon^{(k)} \mid \underline{m} \right) \mathbb{P}(\underline{m}) \quad (4.5)$$

where the sum ranges over all non-negative sub-sample size vectors $[m_0, \dots, m_{2^k-1}]$ that satisfy $\sum_i m_i = m$. We now bound the first factor inside the sum by a quantity which only depends on m (not on the specific vector \underline{m}). We have,

$$\begin{aligned} \mathbb{P}\left(d \notin A_\epsilon^{(k)} \mid \underline{m}\right) &= \mathbb{P}\left(\{d : \|d\| > (1 + \epsilon)2^k \rho\} \cup \{d : \|d\| < (1 - \epsilon)2^k \rho\} \mid \underline{m}\right) \\ &\leq \mathbb{P}\left(\|d\| > (1 + \epsilon)2^k \rho \mid \underline{m}\right) + \mathbb{P}\left(\|d\| < (1 - \epsilon)2^k \rho \mid \underline{m}\right). \end{aligned} \quad (4.6)$$

As stated above, conditioned on \underline{m} the $d(i)$ are independent thus $\|d\|$ is a sum of independent non-identically distributed Bernoulli random variables (also known as Poisson trials). We will use the following lemma.

Lemma 2. *Let X_1, \dots, X_n be independent Bernoulli random variables $P(X_i = 1) = p_i$ where $0 \leq p_i \leq 1$ and denote by $p = \frac{1}{n} \sum_{i=1}^n p_i$. Then for any $0 < \epsilon \leq 1$ we have*

$$P\left(\frac{1}{n} \sum_{i=1}^n X_i < (1 - \epsilon)p\right) < e^{-np\epsilon^2/2}$$

and

$$P\left(\frac{1}{n} \sum_{i=1}^n X_i > (1 + \epsilon)p\right) \leq e^{-np\epsilon^2/4}.$$

The slight asymmetry in the bounds can be seen from the proof of the lemma which is based on applying Chernoff bound on the tail probability of the sum of Poisson trials (see Theorem 4.1 and 4.3, in [14]).

By the above lemma and from (4.4), (4.5), (4.6) it follows that the probability that a random d does not fall in $A_\epsilon^{(k)}$ is bounded as

$$\mathbb{P}\left(d \notin A_\epsilon^{(k)}\right) \leq 2e^{-2^k \rho \epsilon^2/4}. \quad (4.7)$$

Next, we estimate the cardinality of the set $A_\epsilon^{(k)}$. From (4.4) we have,

$$\left|A_\epsilon^{(k)}\right| \leq \sum_{i=\lfloor (1-\epsilon)2^k \rho \rfloor}^{\lceil (1+\epsilon)2^k \rho \rceil} \binom{2^k}{i}.$$

Denote by $B(k, n) = \binom{n}{k}$, then it is easy to verify (see for instance [6]) that the ratio

$$\phi(k) = \frac{B(k, n)}{B(k-1, n)} = \frac{n-k+1}{k}$$

decreases monotonically as k increases from 1 to n . For $k > n/2$ this ratio is smaller than 1 hence it follows that for

$$\frac{\binom{n}{k+v}}{\binom{n}{k}} = \phi(k+1)\phi(k+2)\cdots\phi(k+v) \leq \left(\frac{n-k}{k+1}\right)^v.$$

It follows that for any $c > n/2$ if we denote by $\alpha_{c+1} \equiv \phi(c+1) = \frac{n-c}{c+1}$ then the following upper bound holds,

$$\sum_{k=c}^{c+v} \binom{n}{k} \leq \binom{n}{c} (1 + \alpha_{c+1} + \alpha_{c+1}^2 + \cdots + \alpha_{c+1}^v) \leq \binom{n}{c} \frac{1}{1 - \alpha_{c+1}}.$$

Similarly, the inverse $\phi^{-1}(k)$ increases monotonically as k increases and for $k < n/2$ is smaller than 1 hence it follows that for any $c < n/2$ we have

$$\sum_{k=c-v}^c \binom{n}{k} \leq \binom{n}{c} (1 + \alpha_c^{-1} + \alpha_c^{-2} + \cdots + \alpha_c^{-v}) \leq \binom{n}{c} \frac{1}{1 - \alpha_c^{-1}}.$$

Therefore, as a bound on the cardinality of $A_\epsilon^{(k)}$ we have

$$\left| A_\epsilon^{(k)} \right| \leq \begin{cases} \binom{2^k}{\lfloor (1-\epsilon)2^k \rho \rfloor} \frac{1}{1-\alpha_+} & \text{if } \epsilon < 1 - \frac{1}{2\rho} \\ \binom{2^k}{\lceil (1+\epsilon)2^k \rho \rceil} \frac{1}{1-\alpha_-^{-1}} & \text{if } \epsilon < \frac{1}{2\rho} - 1 \end{cases} \quad (4.8)$$

where

$$\alpha_+ = \frac{2^k - \lfloor (1-\epsilon)2^k \rho \rfloor}{\lfloor (1-\epsilon)2^k \rho \rfloor + 1}$$

and

$$\alpha_-^{-1} = \frac{\lceil (1+\epsilon)2^k \rho \rceil}{2^k - \lceil (1+\epsilon)2^k \rho \rceil + 1}.$$

By the assumption of the theorem, ρ_i (defined in (4.2)) have an average value (4.3) that satisfies $(1+\epsilon)\rho < \frac{1}{2}$ so the bottom bound in (4.8) applies. Hence the bound simplifies to

$$\left| A_\epsilon^{(k)} \right| \leq \binom{2^k}{(1+\epsilon)2^k \rho} \frac{1}{1-\alpha_-^{-1}} \quad (4.9)$$

where we henceforth drop the $\lceil \cdot \rceil$ from the lower entry and leave it implicit.

We now continue the analysis in order to obtain a bound on the possible deviation in randomness of the learner's mistake sequence. Let us denote by $R_d : \{0, 1\}^* \rightarrow \{0, 1\}$ the learner's decision rule which is defined based on the model \mathcal{M}_k learnt by the learner where d is defined in (4.1). When given a finite random binary sequence $X^{(t)}$, R_d produces a binary prediction at time $t+1$, referred to as an *output* bit, which equals

$$\begin{aligned} Y_{t+1} &= R_d(X^{(t)}) \\ &= d(S_t) \end{aligned}$$

where S_t is the state of the learner at time t and $d(S_t)$ is as defined in (3.4). Let us denote by $\xi^{(n)}$ the sequence of errors where the error bit ξ_t at time t equals 1 or 0 according to whether the event of an error in prediction occurs or not, respectively, that is, for a given input sequence $x^{(n)}$ and a prediction sequence $y^{(n)}$ we define

$$\xi_t = \begin{cases} 1 & \text{if } y_t \neq x_t \\ 0 & \text{otherwise.} \end{cases}$$

Consider the subsequence $\xi^{(\nu)}$ of $\xi^{(n)}$ corresponding to the errors associated with the prediction of a 0, i.e., $\xi^{(\nu)}$ consists of the bits of $\xi^{(n)}$ at times t when the prediction bit $y_t = 0$. Clearly, $\xi^{(\nu)}$ is a subsequence of the input $x^{(n)}$ since when the prediction is 0 the error bit equals the input bit. The length ν of this subsequence is a random variable since it depends on the learnt model \mathcal{M}_k . Since $\xi^{(\nu)}$ is a subsequence of the error resulting from prediction by R_d and is also a subsequence of the input $x^{(k+n)}$ we associate a selection rule Γ_d (see section 2) with the decision rule R_d and say that Γ_d *selects* $\xi^{(\nu)}$ from $x^{(n)}$.

Let $E_{d,\epsilon}^{(\ell)}$ denote the event that based on a given *fixed* rule Γ_d the selected subsequence $\xi^{(\nu)}$ from a random input sequence $x^{(n)}$ is of length at least ℓ and its frequency of 1s deviates from the expected value by at least ϵ , formally,

$$E_{d,\epsilon}^{(\ell)} = \left\{ x^{(n)} : \xi^{(\nu)} = \Gamma_d(x^{(n)}), \nu \geq \ell, \left| \frac{\|\xi^{(\nu)}\|}{\nu} - \beta \right| > \epsilon \right\},$$

where $\|\xi^{(\nu)}\|$ denotes the number of 1s in the binary sequence $\xi^{(\nu)}$ of length ν .

We use the following lemma which states a rate on the strong law of large numbers for Bernoulli trials.

Lemma 3. [10] *Let X_1, \dots, X_n be n i.i.d. Bernoulli random variables with $\mathbb{E}X_1 = p$. Then*

$$P\left(\sup_{k \geq n} \left| \frac{1}{k} \sum_{i=1}^k X_i - p \right| > \epsilon\right) \leq 2e^{-2n\epsilon^2(1-\epsilon)}.$$

From Lemma 3 it follows that for any fixed $d \in \{0, 1\}^{2^k}$ we have

$$\begin{aligned} \mathbb{P}\left(E_{d,\epsilon}^{(\ell)}\right) &= \sum_{\nu \geq \ell} \mathbb{P}\left(\left|\frac{\|\xi^{(\nu)}\|}{\nu} - \beta\right| > \epsilon \middle| \nu\right) \mathbb{P}(\nu) \\ &\leq \sum_{\nu \geq \ell} \mathbb{P}\left(\sup_{\nu \geq \ell} \left|\frac{\|\xi^{(\nu)}\|}{\nu} - \beta\right| > \epsilon\right) \mathbb{P}(\nu) \\ &\leq 2e^{-2\ell\epsilon^2(1-\epsilon)} \sum_{\nu \geq \ell} \mathbb{P}(\nu) \\ &\leq 2e^{-2\ell\epsilon^2(1-\epsilon)}. \end{aligned} \tag{4.10}$$

Denote by \hat{d} the binary vector associated with the learnt model \mathcal{M}_k (which is based on a random training sequence $x^{(k+m)}$). We are interested in the probability of the event $E_{\hat{d},\epsilon}^{(\ell)}$ that after learning, given a random test sequence $x^{(n)}$ for prediction, the learner based on the selection rule $\Gamma_{\hat{d}}$ selects a subsequence $\xi^{(\nu)}$ from $x^{(n)}$ of length at least ℓ which is biased away from β by an amount greater than ϵ .

Denoting by $A_\epsilon^{(k)c}$ the complement of the set $A_\epsilon^{(k)}$ then we have

$$\begin{aligned} \mathbb{P}\left(E_{\hat{d},\epsilon}^{(\ell)}\right) &= \mathbb{P}\left(E_{\hat{d},\epsilon}^{(\ell)} \middle| \hat{d} \in A_\epsilon^{(k)}\right) \mathbb{P}\left(A_\epsilon^{(k)}\right) + \mathbb{P}\left(E_{\hat{d},\epsilon}^{(\ell)} \middle| \hat{d} \notin A_\epsilon^{(k)}\right) \mathbb{P}\left(A_\epsilon^{(k)c}\right) \\ &= \mathbb{P}\left(\bigcup_{d \in A_\epsilon^{(k)}} E_{d,\epsilon}^{(\ell)}\right) \mathbb{P}\left(A_\epsilon^{(k)}\right) + \mathbb{P}\left(E_{\hat{d},\epsilon}^{(\ell)} \middle| \hat{d} \notin A_\epsilon^{(k)}\right) \mathbb{P}\left(A_\epsilon^{(k)c}\right) \\ &\leq \mathbb{P}\left(\bigcup_{d \in A_\epsilon^{(k)}} E_{d,\epsilon}^{(\ell)}\right) + \mathbb{P}\left(A_\epsilon^{(k)c}\right) \end{aligned} \tag{4.11}$$

$$\leq 2 \left|A_\epsilon^{(k)}\right| e^{-2\ell\epsilon^2(1-\epsilon)} + 2e^{-2^k \rho \epsilon^2/4} \tag{4.12}$$

where the last inequality follows from (4.7) and (4.10). To have the right side be no larger than $\delta > 0$ it suffices to ensure that ϵ satisfies as a first condition

$$\epsilon \leq 2\sqrt{\frac{1}{2^k \rho} \ln \frac{4}{\delta}}$$

and, assuming that

$$\epsilon \leq \frac{1}{2}$$

which implies that $\frac{1}{2}\epsilon^2 \leq \epsilon^2(1-\epsilon)$, a second condition on ϵ is

$$\epsilon \leq \sqrt{\frac{1}{\ell} \left(\ln \left|A_\epsilon^{(k)}\right| + \ln \left(\frac{4}{\delta}\right) \right)}. \tag{4.13}$$

Now, from (4.9),

$$\left|A_\epsilon^{(k)}\right| \leq \binom{2^k}{(1+\epsilon)2^k \rho} \frac{1}{1-\alpha_-^{-1}}.$$

Hence we may bound

$$\ln |A_\epsilon^{(k)}| \leq \ln \left(\frac{2^k}{(1+\epsilon)2^k\rho} \right) + \ln \frac{1}{1-\alpha_-^{-1}}. \quad (4.14)$$

Now, the following bound on the combination number is easy to verify,

$$\binom{n}{k} \leq \frac{n^k}{k!}$$

from which we obtain

$$\begin{aligned} \ln \binom{n}{k} &\leq k \ln n - \ln \prod_{j=1}^k j \\ &= k \ln n - \sum_{j=1}^k \ln j \\ &\leq k \left(\ln \left(\frac{n}{k} \right) + 1 \right) \end{aligned}$$

where we used $\sum_{j=1}^k \ln j \geq \int_1^k \ln x dx$. Hence the first term of (4.14) is bounded as

$$\ln \left(\frac{2^k}{(1+\epsilon)2^k\rho} \right) \leq (1+\epsilon)\rho 2^k \left(\ln \left(\frac{1}{(1+\epsilon)\rho} \right) + 1 \right) \leq 3\rho 2^{k-1} \left(\ln \left(\frac{1}{\rho} \right) + 1 \right).$$

Now,

$$\alpha_-^{-1} = \frac{\lceil (1+\epsilon)2^k\rho \rceil}{2^k - \lceil (1+\epsilon)2^k\rho \rceil + 1} \leq \frac{2^{k-1}}{2^k - 2^{k-1} + 1} = \frac{2^{k-1}}{2^{k-1} + 1}$$

hence

$$\frac{1}{1-\alpha_-^{-1}} \leq 2^{k-1} + 1 \leq 2^k.$$

It follows that the second term of (4.14) is bounded from above by k . Therefore (4.14) is bounded from above by

$$3\rho 2^{k-1} \left(\ln \left(\frac{1}{\rho} \right) + 1 \right) + k.$$

Hence, for any $0 < \delta \leq 1$, with confidence at least $1 - \delta$ the deviation between the frequency of 1s and β of the subsequence $\xi^{(\nu)}$ selected by the rule Γ_j based on the learnt model \mathcal{M}_k is bounded as

$$\left| \frac{1}{\nu} \sum_{j=1}^{\nu} \xi_j - \beta \right| \leq \sqrt{\max \left\{ \frac{1}{\ell} \left(\ln \left(\frac{4}{\delta} \right) + 3\rho 2^{k-1} \left(\ln \left(\frac{1}{\rho} \right) + 1 \right) + k \right), \frac{1}{2^{k-2}\rho} \ln \left(\frac{4}{\delta} \right) \right\}}.$$

This concludes the proof of Theorem 1.

5. CONCLUSIONS

Concerning Theorem 1, it is clear that the more complex the learner (larger k) the higher the bound $\epsilon(\ell, k, \rho, \delta)$ which implies that the mistake sequence $\xi^{(\nu)}$ can deviate considerably in randomness from a typical random subsequence of the input test sequence $x^{(k+n)}$. Note that the only explicit dependence of the bound on the teacher's probability distribution \mathbb{P} comes through ρ (and not k^*).

However, implicitly, it does depend on k^* since when the learner model has order $k \ll k^*$ then the 'memory' of the teacher is much larger than the learner's window size (recall that k represents the number of bits per state). The learner scans the training sequence using a small window (compared to the teacher's memory) and estimates the state-transition probabilities p_{ij} . Thus $\hat{p}(1|i)$ will be on average

close to $1/2$ and so will the ρ_i and hence ρ . If k is close to k^* then the learner's window is close to the teacher's memory length which means that (in general) the \hat{p}_{ij} may deviate considerably from $1/2$. In this case the ρ_i can be close to zero hence make ρ small which will increase $\epsilon(\ell, k, \rho, \delta)$.

Acknowledgements The author thanks Dr. Vitaly Maiorov from the department of Mathematics of the Technion for reviewing an earlier draft of the paper.

REFERENCES

- [1] A. E. Asarin. Some properties of Kolmogorov δ random finite sequences. *SIAM Theory of Probability and its Applications*, 32:507–508, 1987.
- [2] A. E. Asarin. On some properties of finite objects random in an algorithmic sense. *Soviet Mathematics Doklady*, 36(1):109–112, 1988.
- [3] L. Bienvenu. Kolmogorov-loveland stochasticity and kolmogorov complexity. In *24th Annual Symposium on Theoretical Aspects of Computer Science (STACS 2007)*, volume LNCS 4393, pages 260–271, 2007.
- [4] L. Devroye, L. Györfi, and G. Lugosi. *A Probabilistic Theory of Pattern Recognition*. Springer Verlag, 1996.
- [5] B. Durand and N. Vereshchagin. Kolmogorov-Loveland stochasticity for finite strings. *Information Processing Letters*, 91(6):263–269, 2004.
- [6] W. Feller. *An Introduction to Probability Theory and Its Applications*, volume 1. Wiley, New York, second edition, 1957.
- [7] J.G. Kemeny and J.L. Snell. *Finite Markov Chains*. Springer, New York., 1976.
- [8] A. N. Kolmogorov. On tables of random numbers. *Sankhyaa, The Indian Journal of Statistics*, A(25):369–376, 1963.
- [9] A. N. Kolmogorov. Three approaches to the quantitative definition of information. *Problems of Information Transmission*, 1:1–17, 1965.
- [10] A. N. Kolmogorov. On tables of random numbers. *Theoretical Computer Science*, 207(2):387–395, 1998.
- [11] D. W. Loveland. A new interpretation of the von Mises' concept of random sequence. *Zeitschrift für mathematische Logik und Grundlagen der Mathematik*, 12:279–294, 1966.
- [12] P. Martin-Löf. The definition of random sequences. *Information and Control*, 9:602–619, 1966.
- [13] J. Medhi. *Stochastic processes*. New Age Int'l, 1994.
- [14] P. Raghavan Motwani, R. *Randomized Algorithms*. Cambridge University Press, 1995.
- [15] J. Ratsaby. PAC learning. In M. Y. Kao, editor, *Encyclopedia of Algorithms*, volume 15, pages 622–624. Springer, 2008.
- [16] V.A. Uspenskii, A. L. Semenov, and A. Kh. Shen. Can and individual sequence of zeros and ones be random ? *Russian Mathematical Surveys*, 45(1):121–189, 1990.
- [17] L. G. Valiant. A theory of the learnable. *Comm. ACM*, 27(11):1134–1142, 1984.
- [18] V. N. Vapnik. *Statistical Learning Theory*. Wiley, 1998.
- [19] V. V. Vyugin. Algorithmic complexity and stochastic properties of finite binary sequences. *The Computer Journal*, 42:294–317, 1999.