Phenomenological Aspects of Hořava-Lifshitz Cosmology

Shinji Mukohyama^(a), Kazunori Nakayama^(b), Fuminobu Takahashi^(a), Shuichiro Yokoyama^(c)

^a Institute for the Physics and Mathematics of the Universe, University of Tokyo, Chiba 277-8568, Japan

^b Institute for Cosmic Ray Research,
 University of Tokyo, Kashiwa 277-8582, Japan,
 ^c Department of Physics and Astrophysics,
 Nagoya University, Aichi 464-8502, Japan
 (Dated: November 18, 2018)

Abstract

We show that, assuming the dispersion relation proposed recently by Hořava in the context of quantum gravity, radiation energy density exhibits a peculiar dependence on the scale factor; the radiation energy density decreases proportional to a^{-6} . This simple scaling can have an impact on cosmology. As an example, we show that the resultant baryon asymmetry as well as the stochastic gravity waves can be enhanced. We also discuss current observational constraint on the dispersion relation.

I. INTRODUCTION

Recently a new class of quantum gravity was proposed by Hořava [1], motivated by the solid state physics. This theory is fundamentally non-relativistic and, in the ultraviolet (UV), exhibits the Lifshitz scale invariance

$$t \to b^z t, \quad \vec{x} \to b\vec{x},$$
 (1)

with dynamical critical exponent z = 3. It is this anisotropic rescaling that makes Hořava's theory power-counting renormalizable. The relativistic scaling with z = 1 is restored in the infrared (IR) due to deformation by relevant operators, and the Lorentz symmetry emerges as an accidental symmetry.

The z = 3 scaling implies that, in the UV, a physical degree of freedom should have a dispersion relation of the form

$$\omega^2 \simeq \frac{k^6}{M^4},\tag{2}$$

where M is a characteristic mass scale. Since the Planck mass is an emergent quantity in Hořava's theory, M does not have to be the same order of magnitude as the Planck mass. Moreover, M can differ for different physical degrees of freedom: graviton, photon, quarks, leptons, scalar fields, etc. The fundamental symmetry of Hořava's theory is invariance under foliation-preserving diffeomorphism:

$$x^i \to \tilde{x}^i(x^j, t), \quad t \to \tilde{t}(t).$$
 (3)

Any value of M is consistent with this symmetry and, thus, there is no symmetry reason to expect any fundamental relations among M's for different species unless additional assumptions are made.

On the other hand, among relevant deformations to the dispersion relation (2), the coefficient of the k^2 term should flow to the squared speed of light at the IR fixed point. (See Eq. (4).) This applies to all species, since relativity would not emerge in the IR, otherwise.

Various aspects of Hořava-Lifshitz gravity were already explored in the literature. For example, cosmology based on Hořava-Lifshitz gravity was investigated in [2, 3, 4, 5, 6, 7, 8, 9, 10]. Some solutions in Hořava-Lifshitz gravity were presented in [11, 12, 13, 14]. Possible extensions of the theory were proposed [15, 16, 17, 18].

The dispersion relation (2) leads to interesting cosmological consequences, such as generation of scale-invariant cosmological perturbations [5] and time-delays in Gamma-ray bursts [10]. The purpose of this paper is to explore yet another cosmological consequence of the z = 3 dispersion relation. We shall point out that the radiation energy density in the UV epoch $(T \gg M)$ is proportional to a^{-6} and, thus, decays faster than in the IR epoch $(T \ll M)$ or in relativistic theories. This leads to intriguing cosmological consequences such as enhancement of baryon asymmetry and stochastic gravity waves. We shall also discuss current observational constrains on the dispersion relation.

This paper is organized as follows. In Sec. II cosmological impacts of dispersion relation (2) are explained with some explicit examples. In Sec. III observational constraints on the dispersion relation, in particular, the value of M is discussed. Sec. IV is devoted to summary.

II. COSMOLOGICAL IMPACT OF DISPERSION RELATION

We first derive the peculiar dependence of radiation energy density on the scale factor of the Universe. Suppose that the constituent particles satisfy the dispersion relation,

$$\omega^2 \simeq \frac{k^6}{M^4} + \kappa \frac{k^4}{M^2} + k^2, \tag{4}$$

where we set the speed of light c to be unity, for simplicity. If the typical momentum of the particle (~ temperature) is much lower than M, we recover the ordinary dispersion relation, $\omega^2 \simeq k^2$. On the other hand, if the momentum is much larger than M, the dispersion relation can be approximated to be

$$\omega^2 \simeq \frac{k^6}{M^4}.$$
 (5)

Assuming the relation (5), the energy density of the radiation, ρ_r , decreases as

$$\rho_r \sim \omega n \propto a^{-6},\tag{6}$$

where ω is a typical energy of the particles, n the number density, and a represents the scale factor. This corresponds to the fluid with the equation of state $p_r = \rho_r$ where p_r is the pressure of the radiation. Here we have used a fact that the physical momentum k redshifts as $k \propto a^{-1}$ due to the cosmic expansion, and the number density decreases as $n \propto a^{-3}$. The peculiar dependence (6) should be contrasted to the ordinary scaling, $\rho_r \propto a^{-4}$. Since the energy density of the radiation decreases more quickly than the ordinary case, the estimate on cosmological abundance of e.g. the baryon number, the stochastic gravity waves, etc. can be modified. Let us see this below.

A. Baryon asymmetry

As an example we take up non-thermal baryon production from an inflaton decay.^{#1} Suppose that, after inflation ends, the inflaton ϕ decays into radiation, which is heated up to a temperature higher than M. At the same time, the baryon asymmetry is assumed to be generated from the inflaton decay. This is indeed the case of non-thermal leptogenesis [19, 20, 21]. As the Universe expands, the temperature will decrease. We assume that the ordinary dispersion, $\omega = k$, is restored at a temperature below M. Then, the final baryon-to-entropy ratio can be estimated as

$$\frac{n_B}{s}\Big|_{T=M} = \left.\frac{\sqrt{\rho_r}}{s}\right|_{T=M} \left.\frac{n_B}{\sqrt{\rho_r}}\right|_{T=M} = \left.\frac{\sqrt{\rho_r}}{s}\right|_{T=M} \left(\frac{\rho_\phi}{\sqrt{\rho_r}}\frac{n_B}{\rho_\phi}\right)\Big|_{Reheating},\tag{7}$$

where ρ_{ϕ} denotes the inflaton energy density.^{#2} The final baryon asymmetry will be

$$\frac{n_B}{s}\Big|_{T=M} \sim \frac{\varepsilon}{M} \frac{\Gamma_{\phi} M_P}{m_{\phi}},\tag{8}$$

where $M_P \simeq 2.4 \times 10^{18} \,\text{GeV}$ is the reduced Planck mass, Γ_{ϕ} denotes the decay rate of the inflaton, and ε is the baryon asymmetry generated from the decay of one inflaton quanta. This result should be compared with the ordinary result,

$$\frac{n_B}{s} \sim \varepsilon \frac{\sqrt{\Gamma_{\phi} M_P}}{m_{\phi}}.$$
(9)

Thus, we conclude that the baryon asymmetry is enhanced by

$$\Delta \sim \frac{\sqrt{\Gamma_{\phi} M_P}}{M},\tag{10}$$

compared to the case of the ordinary dispersion relation. Such an enhancement may make an inefficient baryogenesis scenario viable, which would result in too small baryon asymmetry, otherwise.

B. Gravitational waves

Following the similar arguments, it is also possible to enhance abundance of the relic gravity waves. Although the amplitude of the gravitational waves remain constant when

^{#1} We call a field as inflaton, which dominates the energy density of the Universe and decays into radiation. It does not affect our discussion whether or not the field actually induces the inflationary expansion.

^{#2} Note that the entropy density is proportional to T^3 for T < M, but this could be modified for T > M. The calculation here does not depend on the definition of the entropy for T > M.

the corresponding mode lies outside the horizon, once it enters the horizon its amplitude decreases inversely proportional to the scale factor. Since the timing at which a mode enters the horizon depends on the equation of state of the Universe, the thermal history of the Universe is imprinted in the gravitational wave spectrum [22, 23, 24, 25, 26]. The present spectrum of the stochastic gravitational wave background in terms of the density parameter $\Omega_{\rm gw}(k)$ is given by

$$\Omega_{\rm gw}(k) \equiv \frac{1}{\rho_{c0}} \frac{d\rho_{\rm gw}}{d\ln k} = \frac{1}{12H_0^2} k^2 \Delta_h^{\rm (prim)}(k)^2 \left(\frac{a_{\rm in}(k)}{a_0}\right)^2,\tag{11}$$

where ρ_{c0} is the critical energy density at present, $d\rho_{gw}/d \ln k$ represents the energy density of the gravitational waves per logarithmic frequency, H_0 is the present Hubble parameter, $a_{in}(k)$ denotes the scale factor when the mode with comoving wave number k enters the horizon, and $\Delta_h^{(\text{prim})}(k)$ is the dimensionless power spectrum of the primordial gravitational waves, which is assumed to be scale invariant. From this equation we can see that the resulting spectrum scales as $\Omega_{gw}(k) \propto k^{(2-4p)/(1-p)}$ if the Universe expands as $a \propto t^p$ when the mode k enters the horizon. Therefore we get

$$\Omega_{\rm gw}(k) \propto \begin{cases} k^{-2} & \text{for } p = 2/3 \ (w = 0), \\ k^0 & \text{for } p = 1/2 \ (w = 1/3), \\ k^1 & \text{for } p = 1/3 \ (w = 1), \end{cases}$$
(12)

where w represents the equation of state of the Universe. The last line corresponds to the case where the radiation in the UV regime fills the Universe. Thus the mode with $k > k_M$, where k_M is the comoving Hubble scale at which the radiation transits from UV to IR regime, is enhanced by the factor $\Delta = k/k_M$, compared to the case of ordinary dispersion relation. The transition wavenumber k_M is estimated as

$$k_M \simeq 2.6 \times 10^3 \text{ Hz} \left(\frac{g_{*s}(T=M)}{106.75}\right)^{1/6} \left(\frac{M}{10^{11} \text{ GeV}}\right),$$
 (13)

where g_{*s} denotes the relativistic effective degrees of freedom. As we will see in the next section, M is observationally bounded below as $M > 10^{11}$ GeV and hence the frequency (13) lies outside the range covered by future space-based gravitational wave detectors, such as BBO and DECIGO [27]. However, it may be possible to relax this lower bound as $M > 10^7$ GeV as explained in the next section. In this case characteristic spectral shape may be observed by BBO/DECIGO.



FIG. 1: Present gravitational wave background spectrum. Solid (dashed) black line corresponds $M = 10^{11}(10^8)$ GeV. A mode enters the horizon in the matter dominated era, radiation dominated era with IR regime and radiation dominated era with UV regime as shown by MD, RD(IR) and RD(UV) on the top of the figure, respectively. Sensitivity of ultimate-DECIGO is also shown by blue dotted line.

Fig. 1 shows the gravitational wave background spectrum. Solid (dashed) black line corresponds $M = 10^{11}(10^8)$ GeV. A mode enters the horizon in the matter dominated era, radiation dominated era with IR regime and radiation dominated era with UV regime as shown by MD, RD(IR) and RD(UV) on the top of the figure, respectively. Sensitivity of ultimate-DECIGO is also shown by blue dotted line. Here the tensor-to-scalar ratio is assumed to be $0.1.^{\#3}$ It is seen that future gravitational wave detectors may have a chance to detect characteristic spectral shape of the gravitational waves if M happens to be close to the current observational bound.

We have considered two simple examples above, but the enhancement can occur in other situations. For instance, if the dark matter (e.g. gravitino) is produced from the inflaton

^{#3} Throughout this section we have assumed that the gravity waves satisfy the ordinary relativistic dispersion relation. This requires a scale $M_{\rm gw}$, above which the dispersion relation for the gravity waves is modified like (4), to be larger than the Hubble parameter during inflation. Since $M_{\rm gw}$ may be different from M, this is not a serious issue. If $M_{\rm gw}$ is smaller than the Hubble parameter during inflation, the amplitude of the gravity waves will be given by $\sim M_{\rm gw}/M_P$ instead of $H_{\rm inf}/M_P$, according to the discussion along with Ref. [5]. Even if $M_{\rm gw}$ equals to M, the enhancement of the gravity waves still exists, although the detection of the stochastic gravity waves by DECIGO becomes slightly difficult due to smaller tensor-to-scalar ratio.

decay, its abundance will be similarly enhanced. Or, if preheating occurs soon after the inflaton begins to oscillate, significant amount of gravitational waves may be produced with peak frequency corresponding to the comoving Hubble scale after inflation [28, 29, 30, 31]. These gravitational waves are also enhanced in a similar way.

III. OBSERVATIONAL CONSTRAINT ON THE DISPERSION RELATION

In the previous section we have assumed that the radiation, including the standard-model particles, are subject to the dispersion relation (5) at a temperature higher than M. There are observational constraints on the possible deviation from the ordinary dispersion relation, using the arrival timing of the high-energy gamma rays coming from far distant sources [32], such as gamma-ray bursts and active galactic nuclei. In fact, the dispersion relation (4) leads to the energy-dependent photon velocity,

$$v = \frac{d\omega}{dk} = c^2 \frac{k}{\omega} \left(1 + 2\kappa \frac{k^2}{M^2} + 3\frac{k^4}{M^4} \right),\tag{14}$$

where we have restored the speed of light c in the IR limit. In the IR limit $(k \ll M)$, we obtain

$$v \simeq c \left[1 + \frac{3}{2} \kappa \frac{k^2}{M^2} + \left(\frac{5}{2} - \kappa^2 \right) \frac{k^4}{M^4} \right].$$
 (15)

If κ is O(1), the second term gives dominant contribution to the variation of the propagation speed. According to the MAGIC collaboration [33], the lower bound on the scale M in this case reads^{#4}

$$M > 10^{11} \,\mathrm{GeV},$$
 (16)

for $\kappa = O(1)$. The essence of the Hořava-Lifshitz quantum gravity is the appearance of the first term in the r.h.s. of (4), and so, the precise value of κ may be different from order unity. In particular, if κ is negligibly small, it is k^4 term in Eq. (15) that is subject to observational constraints. In this case the constraint on M becomes relaxed as^{#5}

$$M > 10^7 \,\text{GeV}.$$
 (17)

^{#4} More or less similar bound was also found by the H.E.S.S. [34] and Fermi [35] collaborations.

^{#5} Even tighter constraint may come from observations of ultra high energy cosmic rays with energy higher than 10^7 GeV, which may push the bound as $M > 10^{11}$ GeV.

Note that, since the coefficient of k^4 -term in Eq. (15) is positive, it is not possible to attribute the observed delay in the arrival time of gamma-rays to the energy dependence in the speed of light.^{#6}

In order to see the enhancement of the relic gravity waves by the future experiments, M must be smaller than 10⁸ GeV or so, as we have seen in the previous section. This can be made compatible with the current observational constraint by assuming $\kappa \ll 1$. On the other hand, the enhancement of the baryon asymmetry can be viable for a broader parameter space.

IV. CONCLUSIONS

We have shown that the non-trivial dispersion relation suggested by the Hořava-Lifshitz quantum gravity theory leads to a peculiar dependence of the radiation energy density on the scale factor of the Universe. Such a dependence leads to an enhancement of the baryon asymmetry, gravity waves, dark matter, and so on. We have demonstrated this enhancement in simple examples.

These phenomenological aspects are not limited to the Hořava-Lifshitz theory of gravity, but applied to any quantum gravity model in which the dispersion relation is modified at high energy.

Acknowledgments

Three of the authors (S.M., F.T. and S.Y.) thank K. Kohri for discussion. The authors thank the Yukawa Institute for Theoretical Physics at Kyoto University, where this work was initiated during the YITP-W-09-01 on "Non-linear cosmological perturbations". This work is supported by World Premier International Research Center Initiative (WPI Initiative), MEXT, Japan. S.M. is supported in part by MEXT through a Grant-in-Aid for Young Scientists (B) No. 17740134, and by JSPS through a Grant-in-Aid for Creative Scientific Research No. 19GS0219. K.N. would like to thank the Japan Society for the Promotion of Science for financial support. S.Y. is supported in part by Grant-in-Aid for Scientific Re-

 $^{^{\#6}}$ We thank Q. G. Huang for pointing out this issue.

search on Priority Areas No. 467 "Probing the Dark Energy through an Extremely Wide and Deep Survey with Subaru Telescope". He also acknowledges the support from the Grand-in-Aid for the Global COE Program "Quest for Fundamental Principles in the Universe: from Particles to the Solar System and the Cosmos" from the Ministry of Education, Culture, Sports, Science and Technology (MEXT) of Japan.

- [1] P. Horava, "Quantum Gravity at a Lifshitz Point," Phys. Rev. D 79, 084008 (2009)
 [arXiv:0901.3775 [hep-th]].
- [2] T. Takahashi and J. Soda, "Chiral Primordial Gravitational Waves from a Lifshitz Point," arXiv:0904.0554 [hep-th].
- [3] G. Calcagni, "Cosmology of the Lifshitz universe," arXiv:0904.0829 [hep-th].
- [4] E. Kiritsis and G. Kofinas, "Horava-Lifshitz Cosmology," arXiv:0904.1334 [hep-th].
- [5] S. Mukohyama, "Scale-invariant cosmological perturbations from Horava-Lifshitz gravity without inflation," arXiv:0904.2190 [hep-th].
- [6] R. Brandenberger, "Matter Bounce in Horava-Lifshitz Cosmology," arXiv:0904.2835 [hep-th].
- [7] H. Nikolic, "Horava-Lifshitz gravity, absolute time, and objective particles in curved space," arXiv:0904.3412 [hep-th].
- [8] Y. S. Piao, "Primordial Perturbation in Horava-Lifshitz Cosmology," arXiv:0904.4117 [hep-th].
- X. Gao, "Cosmological Perturbations and Non-Gaussianities in Hořava-Lifshitz Gravity," arXiv:0904.4187 [hep-th].
- [10] B. Chen and Q. G. Huang, "Field Theory at a Lifshitz Point," arXiv:0904.4565 [hep-th].
- [11] H. Lu, J. Mei and C. N. Pope, "Solutions to Horava Gravity," arXiv:0904.1595 [hep-th].
- [12] H. Nastase, "On IR solutions in Horava gravity theories," arXiv:0904.3604 [hep-th].
- [13] R. G. Cai, L. M. Cao and N. Ohta, "Topological Black Holes in Horava-Lifshitz Gravity," arXiv:0904.3670 [hep-th].
- [14] E. O. Colgain and H. Yavartanoo, "Dyonic solution of Horava-Lifshitz Gravity," arXiv:0904.4357 [hep-th].
- [15] J. Kluson, "Branes at Quantum Criticality," arXiv:0904.1343 [hep-th].
- [16] K. I. Izawa, "A Note on Quasi-Riemannian Gravity with Higher Derivatives," arXiv:0904.3593

[hep-th].

- [17] R. G. Cai, Y. Liu and Y. W. Sun, "On the z=4 Horava-Lifshitz Gravity," arXiv:0904.4104 [hep-th].
- [18] T. Sotiriou, M. Visser and S. Weinfurtner, "Phenomenologically viable Lorentz-violating quantum gravity," arXiv:0904.4464 [hep-th].
- [19] M. Fukugita and T. Yanagida, Phys. Lett. B 174, 45 (1986).
- [20] H. Murayama and T. Yanagida, Phys. Lett. B **322**, 349 (1994) [arXiv:hep-ph/9310297];
 K. Hamaguchi, H. Murayama and T. Yanagida, Phys. Rev. D **65**, 043512 (2002) [arXiv:hep-ph/0109030].
- [21] T. Asaka, K. Hamaguchi, M. Kawasaki and T. Yanagida, Phys. Lett. B 464, 12 (1999)
 [arXiv:hep-ph/9906366]; Phys. Rev. D 61, 083512 (2000) [arXiv:hep-ph/9907559]; M. Endo,
 F. Takahashi and T. T. Yanagida, Phys. Rev. D 74, 123523 (2006) [arXiv:hep-ph/0611055].
- [22] N. Seto and J. Yokoyama, J. Phys. Soc. Jap. 72, 3082 (2003) [arXiv:gr-qc/0305096].
- [23] H. Tashiro, T. Chiba and M. Sasaki, Class. Quant. Grav. 21, 1761 (2004) [arXiv:gr-qc/0307068].
- [24] L. A. Boyle and P. J. Steinhardt, Phys. Rev. D 77, 063504 (2008) [arXiv:astro-ph/0512014];
 L. A. Boyle and A. Buonanno, Phys. Rev. D 78, 043531 (2008) [arXiv:0708.2279 [astro-ph]].
- [25] K. Nakayama, S. Saito, Y. Suwa and J. Yokoyama, Phys. Rev. D 77, 124001 (2008)
 [arXiv:0802.2452 [hep-ph]]; JCAP 0806, 020 (2008) [arXiv:0804.1827 [astro-ph]].
- [26] S. Kuroyanagi, T. Chiba and N. Sugiyama, arXiv:0804.3249 [astro-ph].
- [27] N. Seto, S. Kawamura and T. Nakamura, Phys. Rev. Lett. 87, 221103 (2001).
- [28] S. Y. Khlebnikov and I. I. Tkachev, Phys. Rev. D 56, 653 (1997) [arXiv:hep-ph/9701423].
- [29] R. Easther and E. A. Lim, JCAP 0604, 010 (2006) [arXiv:astro-ph/0601617]; R. Easther,
 J. T. . Giblin and E. A. Lim, Phys. Rev. Lett. 99, 221301 (2007) [arXiv:astro-ph/0612294].
- [30] J. Garcia-Bellido and D. G. Figueroa, Phys. Rev. Lett. 98, 061302 (2007); J. Garcia-Bellido,
 D. G. Figueroa and A. Sastre, Phys. Rev. D 77, 043517 (2008).
- [31] J. F. Dufaux, A. Bergman, G. N. Felder, L. Kofman and J. P. Uzan, Phys. Rev. D 76, 123517
 (2007) [arXiv:0707.0875 [astro-ph]].
- [32] G. Amelino-Camelia, J. R. Ellis, N. E. Mavromatos, D. V. Nanopoulos and S. Sarkar, Nature 393, 763 (1998) [arXiv:astro-ph/9712103].
- [33] J. Albert et al. [MAGIC Collaboration and Other Contributors Collaboration], Phys. Lett. B

668, 253 (2008) [arXiv:0708.2889 [astro-ph]].

- [34] J. Bolmont, R. Buehler, A. Jacholkowska, S. J. Wagner and f. t. H. Collaboration, arXiv:0904.3184 [gr-qc].
- [35] A. A. Abdo et al. [Fermi LAT and Fermi GBM Collaborations], Science 323, 1688 (2009).