

The Next-to-Minimal Supersymmetric Standard Model

Ulrich Ellwanger^a, Cyril Hugonie^b and Ana M. Teixeira^{a,c} ¹

^a *LPT, UMR 8627, CNRS, Université de Paris-Sud, 91405 Orsay, France*

^b *LPTA, UMR 5207, CNRS/IN2P3,*

Université de Montpellier II, 34095 Montpellier, France

^c *CFTP, Departamento de Física, Instituto Superior Técnico, 1049-001 Lisboa, Portugal*

Abstract

We review the theoretical and phenomenological aspects of the Next-to-Minimal Supersymmetric Standard Model: the Higgs sector including radiative corrections and the 2-loop β -functions for all parameters of the general NMSSM; the tadpole and domain wall problems, baryogenesis; NMSSM phenomenology at colliders, B physics and dark matter; specific scenarios as the constrained NMSSM, Gauge Mediated Supersymmetry Breaking, $U(1)'$ -extensions, CP and R-parity violation.

¹Address after October 2009: LPC Clermont-Ferrand, UMR 6533, CNRS/IN2P3, Université Blaise Pascal, 63177 Aubière, France

Contents

1	Introduction	3
2	Lagrangian of the general NMSSM	6
2.1	Tree level potential and mass matrices	6
2.2	Limiting cases: the effective MSSM and approximate global symmetries . . .	13
2.2.1	The effective MSSM	13
2.2.2	The Peccei-Quinn symmetry limit	14
2.2.3	The R-symmetry limit	15
3	Radiative corrections	16
3.1	Renormalisation group equations	16
3.2	Radiative corrections to the Higgs masses, and the upper bound on the lightest CP-even Higgs mass	18
3.3	Radiative corrections to coupling constants	22
4	The tadpole and domain wall problems, baryogenesis, hybrid inflation	25
4.1	The tadpole problem	25
4.2	The domain wall problem	26
4.3	Electroweak baryogenesis	27
4.4	Hybrid inflation	28
5	NMSSM phenomenology at colliders	29
5.1	The Higgs sector	29
5.1.1	LEP and e^+e^- colliders	29
5.1.2	Tevatron and LHC	32
5.2	The neutralino sector	34
5.2.1	Neutralino pair production at e^+e^- colliders	35
5.2.2	Decays into and of neutralinos	36
5.2.3	Displaced vertices	36
6	b physics and the anomalous magnetic moment of the muon	38
6.1	B physics	38
6.2	Υ and η_b physics	42
6.3	The anomalous magnetic moment of the muon	44
6.4	Combined constraints on a light CP-odd Higgs	45
7	Specific scenarios	47
7.1	The constrained NMSSM	47
7.1.1	Constraints from the absence of charge and colour breaking minima	48
7.1.2	The constrained NMSSM with dark matter constraints	49
7.1.3	The constrained NMSSM with relaxed universality conditions	52
7.2	The NMSSM and Gauge Mediated Supersymmetry Breaking	52
7.3	The nMSSM	54

7.4	The fine tuning problem in the NMSSM	55
8	Variants of the NMSSM	59
8.1	CP violation	59
8.1.1	Spontaneous CP violation	60
8.1.2	Explicit CP violation	62
8.2	R-parity violation and neutrino masses in the NMSSM	65
8.2.1	Massive neutrinos in the NMSSM	66
8.3	$U(1)'$ -extensions of the NMSSM	69
9	Dark Matter in the NMSSM	75
9.1	Neutralino relic density	75
9.1.1	Annihilation	76
9.1.2	NMSSM-specific annihilation processes	78
9.2	Dark matter detection in the NMSSM	79
9.2.1	Direct dark matter detection	79
9.2.2	Indirect dark matter detection	81
9.3	Specific scenarios	82
10	Conclusions and outlook	88
Appendix A. Mixing matrices and tree level Higgs couplings		89
A.1	Mixing matrices	89
A.2	Higgs couplings	90
Appendix B. Renormalisation group equations		98
B.1	Gauge and Yukawa couplings	98
B.2	Gaugino masses	99
B.3	Trilinear couplings	99
B.4	Squark and slepton masses	101
B.5	Higgs masses	103
B.6	Additional parameters of the general NMSSM	104
Appendix C. Radiative corrections to the Higgs masses		106
C.1	Yukawa and gauge couplings at the scale M_{SUSY}	106
C.2	Higgs wave function renormalisation constants	108
C.3	Corrections to the Higgs mass matrices	109
C.4	Pole masses	116
Appendix D. Public computer tools		118
References		119

1 Introduction

Supersymmetric extensions of the Standard Model (SM) are motivated by a solution of the hierarchy problem [1–5], an automatic unification of the running gauge couplings at a Grand Unified (GUT) scale M_{GUT} [6–9], and the possibility to explain the dark matter relic density in terms of a stable neutral particle [10, 11].

It is well known that a supersymmetric extension of the Higgs sector of the SM [12, 13] requires the introduction of two Higgs $SU(2)$ -doublets H_u and H_d , where vacuum expectation values (vevs) of H_u and H_d generate masses for up-type quarks and down-type quarks and charged leptons, respectively. The model with this minimal field content in the Higgs sector is denoted as the Minimal Supersymmetric Standard Model (MSSM) (for reviews see, e. g., [14–16]). The Lagrangian of the MSSM must contain a supersymmetric (SUSY) mass term μ for H_u and H_d , which has to be of the order of the SUSY breaking scale M_{SUSY} for phenomenological reasons (see below). This spoils a potentially attractive property of supersymmetric extensions of the SM: the electroweak scale generated by the Higgs vevs could depend only on M_{SUSY} , which would be the only scale asking for an explanation to why it is far below M_{GUT} or the Planck scale M_{Planck} . The question how a supersymmetric mass parameter μ can assume a value of the order of M_{SUSY} is denoted as the “ μ -problem” [17] of the MSSM.

A simple and elegant way to solve this problem consists in generating an effective (supersymmetric) mass term μ in a way similar to the generation of quark and lepton masses in the SM: the mass term μ is replaced by a Yukawa coupling of H_u and H_d to a scalar field, and the vev of the scalar field – induced by the soft SUSY breaking terms – is of the desired order. Since the μ parameter carries no $SU(3) \times SU(2) \times U(1)_Y$ quantum numbers, the field to be introduced has to be a singlet S (the complex scalar component of a chiral superfield \widehat{S}), and the resulting model is the Next-to-Minimal Supersymmetric Standard Model (NMSSM), sometimes also denoted as the (M+1)SSM.

In fact, already the first attempts to construct supersymmetric extensions of the SM employed such a singlet field [12, 13, 18]. A singlet was also present in most of the first globally supersymmetric GUT models [5, 19–22]. Then one realised that spontaneous supersymmetry breaking in the framework of supergravity (SUGRA) leads in a simple way to the desired soft SUSY breaking terms in the Lagrangian; see [14] for an early review. Within SUGRA, a μ term of the order of M_{SUSY} can actually be generated if one assumes the presence of a particular Higgs-dependent structure in the Kähler potential [23]. Still, the first locally supersymmetric extensions of the SM [24–26] as well as most GUT models within SUGRA [27–34] used a singlet field in the Higgs sector leading to variants of the NMSSM at the weak or SUSY breaking scale $\lesssim 1$ TeV. (See also SUGRA models motivated by string theory [35–44].)

Expanding around the vacuum with non-vanishing vevs of the neutral CP-even components of H_u , H_d and S , one finds that the scalar components of \widehat{S} mix with the neutral scalar components of \widehat{H}_u and \widehat{H}_d leading, in the absence of complex parameters (corresponding to the absence of explicit CP violation), to three CP-even and two CP-odd neutral scalars (see [45–47] for some reviews). Likewise, the fermionic superpartner of \widehat{S} mixes with the neutral fermionic superpartners of \widehat{H}_u , \widehat{H}_d (and the neutral electroweak gauginos) leading

to five neutralinos. As a consequence, both the Higgs and the neutralino sectors of the NMSSM can get considerably modified compared to the MSSM.

In the Higgs sector, important alterations with respect to the MSSM are a possibly larger mass of the Higgs scalar with SM-like couplings to gauge bosons, and additional possibly light states with reduced couplings to gauge bosons. Notably a light CP-odd scalar with vanishing couplings to two gauge bosons like all CP-odd scalars (but with possibly even enhanced couplings to quarks and leptons) can appear in the Higgs spectrum, allowing for new Higgs-to-Higgs decays. Under these circumstances, the detection of Higgs bosons at colliders can become considerably more complicated. At present it is not even guaranteed that a single Higgs scalar can be observed at the LHC within the NMSSM, see Section 5. In addition, a light CP-odd scalar can affect “low energy” observables in B physics, Υ physics and the anomalous magnetic moment of the muon.

The modifications within the neutralino sector are particularly relevant if the additional singlet-like neutralino is the lightest one and, simultaneously, the lightest supersymmetric particle (LSP). This would have an important impact on all decay chains of supersymmetric particles (sparticles), and hence on their signatures at colliders. For instance, the next-to-lightest supersymmetric particle (NLSP) can have a long life time leading to displaced vertices. Also, the LSP relic density has to be reconsidered in this case.

Hence, apart from the theoretical motivations for the NMSSM, its phenomenological consequences must be worked out in order not to miss (or misinterpret) both Higgs and sparticles signals – or the absence thereof – at past, present and future colliders.

In the present paper we review theoretical and phenomenological aspects of the NMSSM: Higgs masses and couplings including radiative corrections, issues related to the NMSSM Higgs potential as the nature of the electroweak phase transition, Higgs and sparticle spectra within specific assumptions on the origin of supersymmetry breaking as minimal SUGRA and gauge mediated supersymmetry breaking (GMSB), NMSSM specific Higgs and sparticle signatures at colliders, possible impact on low energy observables, and the dark matter relic density and its detection. Possible variants of the simplest NMSSM as explicit CP violation in the Higgs sector, R-parity violation (in connection with neutrino masses) and extra $U(1)'$ gauge symmetries are sketched briefly. Relevant formulae like Feynman rules, renormalisation group (RG) equations and details of the radiative corrections to the Higgs masses are given in the Appendices.

Clearly, we cannot present all details of all results that have been obtained within the NMSSM up to now. However, in all Sections we attempt to reference to the complete available literature, where the various subjects are discussed.

Let us conclude the Introduction by recalling the arguments for a μ parameter of the order of M_{SUSY} , whose necessity constitutes the main motivation for the NMSSM: both complex Higgs scalars H_u and H_d of the MSSM have to be components of chiral superfields which contain, in addition, fermionic $SU(2)$ -doublets ψ_u and ψ_d . The Lagrangian of the MSSM can contain supersymmetric mass terms for these fields, i.e. identical positive masses squared μ^2 for $|H_u|^2$ and $|H_d|^2$, and a Dirac mass μ for ψ_u and ψ_d . In the presence of a SUSY mass term $\sim \mu$ in the Lagrangian, a soft SUSY breaking mass term $B\mu H_u H_d$ can also appear, where the soft SUSY breaking parameter B has the dimension of a mass.

For various reasons the mass parameter μ *cannot* vanish. First, a Dirac mass μ for ψ_u

and ψ_d is required for phenomenological reasons: both fermionic $SU(2)$ -doublets ψ_u and ψ_d contain electrically charged components. Together with the fermionic superpartners of the W^\pm bosons, they constitute the so-called chargino sector (two charged Dirac fermions) of SUSY extensions of the SM. Due to the fruitless searches for a chargino at LEP, the lighter chargino has to have a mass above ~ 103 GeV [48]. Analysing the chargino mass matrix, one finds that this lower limit implies that the Dirac mass μ for ψ_u and ψ_d – for arbitrary values of the other parameters – has to satisfy the constraint $|\mu| \gtrsim 100$ GeV.

Second, an analysis of the Higgs potential shows that a non-vanishing term $B\mu H_u H_d$ is a necessary condition for that *both* neutral components of H_u and H_d are non-vanishing at the minimum. This, in turn, is required in order to generate masses for up-type quarks, down-type quarks and leptons by the Higgs mechanism. Moreover, the numerical value of the product $B\mu$ should be roughly of the order of the electroweak scale (M_Z^2).

Third, $\mu = 0$ would generate a Peccei-Quinn symmetry in the Higgs sector, and hence an unacceptable massless axion [17].

However, $|\mu|$ must not be too large: the Higgs potential must be unstable at its origin $H_u = H_d = 0$ in order to generate the electroweak symmetry breaking. Whereas the soft SUSY breaking mass terms for H_u and H_d of the order of the SUSY breaking scale M_{SUSY} can generate such a desired instability, the μ -induced masses squared for H_u and H_d are always positive, and must *not* dominate the negative soft SUSY breaking mass terms. Consequently the μ parameter must obey $|\mu| \lesssim M_{\text{SUSY}}$. Hence, both “natural” values $\mu = 0$ and very large μ ($\sim M_{\text{GUT}}$ or $\sim M_{\text{Planck}}$) are ruled out, and the need for an explanation of $\mu \approx M_{\text{SUSY}}$ is the μ -problem.

Within the NMSSM, where μ is generated by the vev $\langle S \rangle$ of a singlet S , $\langle S \rangle$ has to be of the order of M_{SUSY} ; this is easy to obtain with the help of soft SUSY breaking negative masses squared (or trilinear couplings) of the order of M_{SUSY} for S . Then, M_{SUSY} is the only scale in the theory. In this sense, the NMSSM is the simplest supersymmetric extension of the SM in which the weak scale is generated by the supersymmetry breaking scale M_{SUSY} only.

2 Lagrangian of the general NMSSM

In this Section, we present the scalar potential (at tree level) and the mass matrices in the general NMSSM. Their diagonalization and conventions for the mixing matrices are deferred to the Appendix A, where we also list the Feynman rules for the physical eigenstates.

2.1 Tree level potential and mass matrices

As in any softly broken supersymmetric theory, the Lagrangian of the NMSSM is specified by the supersymmetric gauge interactions, the superpotential and the soft supersymmetry breaking gaugino masses, scalar masses and trilinear couplings.

To begin with, we consider the general NMSSM defined as the MSSM with an additional gauge singlet chiral superfield \widehat{S} , including the most general renormalisable couplings in the superpotential and the corresponding soft SUSY breaking terms in $\mathcal{L}_{\text{soft}}$. (Here we limit ourselves, however, to the R-parity and CP conserving case. Chiral superfields are denoted by hatted capital letters; unhatted capital letters indicate their complex scalar components.) In the general NMSSM, the terms in the superpotential W_{Higgs} depending exclusively on the Higgs superfields \widehat{H}_u , \widehat{H}_d and \widehat{S} are (here we follow the SLHA2 conventions in [49] where, however, \widehat{H}_u is denoted as \widehat{H}_2 , and \widehat{H}_d as \widehat{H}_1):

$$W_{\text{Higgs}} = (\mu + \lambda \widehat{S}) \widehat{H}_u \cdot \widehat{H}_d + \xi_F \widehat{S} + \frac{1}{2} \mu' \widehat{S}^2 + \frac{\kappa}{3} \widehat{S}^3 \quad (2.1)$$

where the terms $\sim \lambda$, κ are dimensionless Yukawa couplings, the terms $\sim \mu$, μ' are supersymmetric mass terms, and ξ_F of dimension $mass^2$ parametrizes a (supersymmetric) tadpole term. To (2.1) we have to add the Yukawa couplings of the quark and lepton superfields:

$$W_{\text{Yukawa}} = h_u \widehat{Q} \cdot \widehat{H}_u \widehat{U}_R^c + h_d \widehat{H}_d \cdot \widehat{Q} \widehat{D}_R^c + h_e \widehat{H}_d \cdot \widehat{L} \widehat{E}_R^c \quad (2.2)$$

where the Yukawa couplings h_u , h_d , h_e and the superfields \widehat{Q} , \widehat{U}_R^c , \widehat{D}_R^c , \widehat{L} and \widehat{E}_R^c should be understood as matrices and vectors in family space, respectively.

In (2.1) and (2.2), the $SU(2)$ doublets are

$$\widehat{Q} = \begin{pmatrix} \widehat{U}_L \\ \widehat{D}_L \end{pmatrix}, \quad \widehat{L} = \begin{pmatrix} \widehat{\nu}_L \\ \widehat{E}_L \end{pmatrix}, \quad \widehat{H}_u = \begin{pmatrix} \widehat{H}_u^+ \\ \widehat{H}_u^0 \end{pmatrix}, \quad \widehat{H}_d = \begin{pmatrix} \widehat{H}_d^0 \\ \widehat{H}_d^- \end{pmatrix}, \quad (2.3)$$

and the products of two $SU(2)$ doublets are defined as, e. g.,

$$\widehat{H}_u \cdot \widehat{H}_d = \widehat{H}_u^+ \widehat{H}_d^- - \widehat{H}_u^0 \widehat{H}_d^0. \quad (2.4)$$

The corresponding soft SUSY breaking masses and couplings are, again in the SLHA2 conventions [49],

$$\begin{aligned} -\mathcal{L}_{\text{soft}} = & m_{H_u}^2 |H_u|^2 + m_{H_d}^2 |H_d|^2 + m_S^2 |S|^2 + m_Q^2 |Q|^2 + m_U^2 |U_R|^2 \\ & + m_D^2 |D_R|^2 + m_L^2 |L|^2 + m_E^2 |E_R|^2 \\ & + (h_u A_u Q \cdot H_u U_R^c - h_d A_d Q \cdot H_d D_R^c - h_e A_e L \cdot H_d E_R^c \\ & + \lambda A_\lambda H_u \cdot H_d S + \frac{1}{3} \kappa A_\kappa S^3 + m_3^2 H_u \cdot H_d + \frac{1}{2} m_S^2 S^2 + \xi_S S + \text{h.c.}) . \quad (2.5) \end{aligned}$$

(Sometimes the definitions $m_3^2 = B\mu$, $m_S^2 = B'\mu'$ are used.)

Clearly, the dimensionful supersymmetric parameters μ , μ' and ξ_F in the superpotential (2.1) (and the associated soft SUSY breaking parameters m_3^2 , m_S^2 and ξ_S in (2.5)) have to be of the order of the weak or SUSY breaking scale, in contradiction to one of the theoretical motivations for the NMSSM mentioned in the Introduction. Although some of these terms are non-vanishing in various scenarios, one considers mostly the simpler NMSSM with a scale invariant superpotential where $\mu = \mu' = \xi_F = 0$,

$$W_{\text{sc.inv.}} = \lambda \widehat{S} \widehat{H}_u \cdot \widehat{H}_d + \frac{\kappa}{3} \widehat{S}^3 \quad (2.6)$$

and vanishing parameters m_3^2 , m_S^2 and ξ_S in (2.5). Then, a vev s of \widehat{S} of the order of the weak or SUSY breaking scale generates an effective μ -term with

$$\mu_{\text{eff}} = \lambda s, \quad (2.7)$$

which solves the μ -problem of the MSSM.

As any supersymmetric theory with a scale invariant (cubic) superpotential, the complete Lagrangian – including the soft SUSY breaking terms – specified by (2.6) possesses an accidental \mathbb{Z}_3 -symmetry corresponding to a multiplication of all components of all chiral superfields by a phase $e^{2\pi i/3}$. In the following we denote the version with the scale invariant superpotential (2.6) as the “ \mathbb{Z}_3 -invariant NMSSM”. Any of the dimensionful terms in the general superpotential (2.1) breaks the \mathbb{Z}_3 -symmetry explicitly. Subsequently, the version corresponding to the general superpotential (2.1) will be denoted as the “general NMSSM”. In the literature, “NMSSM” stands mostly for the \mathbb{Z}_3 -invariant NMSSM. In the following we will retain this convention.

The Higgs sector of the \mathbb{Z}_3 -invariant NMSSM is specified by the seven parameters

$$\lambda, \kappa, m_{H_u}^2, m_{H_d}^2, m_S^2, A_\lambda \text{ and } A_\kappa. \quad (2.8)$$

Expressions for the Higgs mass matrices in the \mathbb{Z}_3 -invariant NMSSM can be found, e. g., in [45, 46, 50–52]; in the following we discuss, for completeness, the general NMSSM from which the \mathbb{Z}_3 -invariant NMSSM can always be obtained by setting $m_3^2 = m_S^2 = \xi_S = \mu = \mu' = \xi_F = 0$.

From the SUSY gauge interactions, the F - and the soft SUSY breaking terms one obtains the Higgs potential:

$$\begin{aligned} V_{\text{Higgs}} = & \left| \lambda (H_u^+ H_d^- - H_u^0 H_d^0) + \kappa S^2 + \mu' S + \xi_F \right|^2 \\ & + (m_{H_u}^2 + |\mu + \lambda S|^2) \left(|H_u^0|^2 + |H_u^+|^2 \right) + (m_{H_d}^2 + |\mu + \lambda S|^2) \left(|H_d^0|^2 + |H_d^-|^2 \right) \\ & + \frac{g_1^2 + g_2^2}{8} \left(|H_u^0|^2 + |H_u^+|^2 - |H_d^0|^2 - |H_d^-|^2 \right)^2 + \frac{g_2^2}{2} |H_u^+ H_d^{0*} + H_u^0 H_d^{-*}|^2 \\ & + m_S^2 |S|^2 + (\lambda A_\lambda (H_u^+ H_d^- - H_u^0 H_d^0) S + \frac{1}{3} \kappa A_\kappa S^3 + m_3^2 (H_u^+ H_d^- - H_u^0 H_d^0) \\ & + \frac{1}{2} m_S^2 S^2 + \xi_S S + \text{h.c.}) \end{aligned} \quad (2.9)$$

where g_1 and g_2 denote the $U(1)_Y$ and $SU(2)$ gauge couplings, respectively.

The neutral physical Higgs fields (with index R for the CP-even, index I for the CP-odd states) are obtained by expanding the full scalar potential (2.9) around the real neutral vevs v_u , v_d and s as

$$H_u^0 = v_u + \frac{H_{uR} + iH_{uI}}{\sqrt{2}}, \quad H_d^0 = v_d + \frac{H_{dR} + iH_{dI}}{\sqrt{2}}, \quad S = s + \frac{S_R + iS_I}{\sqrt{2}}; \quad (2.10)$$

where the vevs have to be obtained from the minima of

$$\begin{aligned} V_{\text{Higgs}} = & \left(-\lambda v_u v_d + \kappa s^2 + \mu' s + \xi_F\right)^2 + \frac{g_1^2 + g_2^2}{8} (v_u^2 - v_d^2)^2 \\ & + (m_{H_u}^2 + (\mu + \lambda s)^2) v_u^2 + (m_{H_d}^2 + (\mu + \lambda s)^2) v_d^2 \\ & + m_S^2 s^2 - 2\lambda A_\lambda v_u v_d s + \frac{2}{3} \kappa A_\kappa s^3 - 2m_3^2 v_u v_d + m_S'^2 s^2 + 2\xi_S s, \end{aligned} \quad (2.11)$$

The signs of some parameters in a Lagrangian have no physical meaning, since they can be changed by field redefinitions $\phi \rightarrow -\phi$. Analysing all possible field redefinitions in the Lagrangian above, one finds that one can choose positive Yukawa couplings λ , h_t , h_b , h_τ (the latter corresponding to the (3,3) components in family space of h_u , h_d and h_e in (2.2)) and positive vevs v_u and v_d , whereas κ , s and all dimensionful parameters can have both signs. In addition, in the general NMSSM one of the dimensionful S -dependent parameters can be removed by a constant shift of the real component of S . In particular, one can put the MSSM-like μ -term in the superpotential (2.1) to zero by a redefinition $s \rightarrow s - \mu/\lambda$ (assuming $\lambda \neq 0$) and corresponding redefinitions of the other dimensionful parameters; in the following we will assume, for simplicity, that this convention is used (which does *not* imply, in general, that $m_3^2 = 0$!).

We define, as usual,

$$\tan \beta = \frac{v_u}{v_d} \quad (2.12)$$

(positive by assumption), and we have

$$M_Z^2 = g^2 v^2 \quad \text{where } g^2 \equiv \frac{g_1^2 + g_2^2}{2}, \quad v^2 = v_u^2 + v_d^2 \simeq (174 \text{ GeV})^2. \quad (2.13)$$

Furthermore it is convenient to define, together with μ_{eff} as in (2.7),

$$B_{\text{eff}} = A_\lambda + \kappa s, \quad \widehat{m}_3^2 = m_3^2 + \lambda(\mu' s + \xi_F) \quad (2.14)$$

where we have used the convention $\mu = 0$. B_{eff} plays the rôle of the MSSM-like B -parameter, and \widehat{m}_3^2 vanishes in the \mathbb{Z}_3 -invariant NMSSM.

It is possible to use the three minimisation equations of the potential (2.11) with respect to v_u , v_d and s in order to replace the three parameters $m_{H_u}^2$, $m_{H_d}^2$ and m_S^2 by v_u , v_d and s . The minimisation equations are given by

$$\begin{aligned} v_u \left(m_{H_u}^2 + \mu_{\text{eff}}^2 + \lambda^2 v_d^2 + \frac{g_1^2 + g_2^2}{4} (v_u^2 - v_d^2) \right) - v_d (\mu_{\text{eff}} B_{\text{eff}} + \widehat{m}_3^2) &= 0, \\ v_d \left(m_{H_d}^2 + \mu_{\text{eff}}^2 + \lambda^2 v_u^2 + \frac{g_1^2 + g_2^2}{4} (v_d^2 - v_u^2) \right) - v_u (\mu_{\text{eff}} B_{\text{eff}} + \widehat{m}_3^2) &= 0, \\ s \left(m_S^2 + m_S'^2 + \mu'^2 + 2\kappa \xi_F + \kappa A_\kappa s + 2\kappa^2 s^2 + 3\kappa s \mu' + \lambda^2 (v_u^2 + v_d^2) - 2\lambda \kappa v_u v_d \right) \\ &+ \xi_S + \xi_F \mu' - \lambda v_u v_d (A_\lambda + \mu') = 0. \end{aligned} \quad (2.15)$$

From the first two of these equations one can derive

$$\frac{v_u v_d}{v^2} \equiv \frac{1}{2} \sin 2\beta = \frac{\mu_{\text{eff}} B_{\text{eff}} + \widehat{m}_3^2}{m_{H_u}^2 + m_{H_d}^2 + 2\mu_{\text{eff}}^2 + \lambda^2 v^2} ; \quad (2.16)$$

hence, in order to have both v_u and v_d different from zero ($\tan \beta \neq 0, \infty$), we need $\mu_{\text{eff}} B_{\text{eff}} + \widehat{m}_3^2 \neq 0$ in the general NMSSM, and $\mu_{\text{eff}} B_{\text{eff}} \neq 0$ in the \mathbb{Z}_3 -invariant NMSSM.

Given M_Z , one can choose as six independent parameters in the Higgs sector of the \mathbb{Z}_3 -invariant NMSSM

$$\lambda, \kappa, A_\lambda, A_\kappa, \tan \beta, \mu_{\text{eff}}, \quad (2.17)$$

to which one has to add in the general NMSSM (in the convention $\mu = 0$) the five parameters

$$m_3^2, \mu', m_S^2, \xi_F \text{ and } \xi_S . \quad (2.18)$$

Let us consider the conditions arising from a phenomenologically acceptable minimum of the potential: both v_u and v_d must not vanish, and – in the absence of a μ -term – s must be large enough to generate a sufficiently large effective μ -term $\mu_{\text{eff}} = \lambda s \gtrsim 100$ GeV (see the Introduction). In the \mathbb{Z}_3 -invariant NMSSM, the dominant terms for large s in the potential (2.11) are

$$V_{\text{Higgs}}(s) \sim m_S^2 s^2 + \frac{2}{3} \kappa A_\kappa s^3 + \kappa^2 s^4 . \quad (2.19)$$

One easily finds that $A_\kappa^2 \gtrsim 8 m_S^2$ is a condition for $s \neq 0$ [53, 54], and

$$A_\kappa^2 \gtrsim 9 m_S^2 \quad (2.20)$$

a condition for an absolute minimum with

$$s \simeq \frac{1}{4\kappa} \left(-A_\kappa - \sqrt{A_\kappa^2 - 8m_S^2} \right) . \quad (2.21)$$

We note that in the case of a scale invariant superpotential (2.6) and without *any* soft terms for the singlet – as it can happen in GMSB models [55] – a vev $s \neq 0$ is still triggered by the s -dependent terms in the potential (2.11) neglected in (2.19). However, the resulting value of s is too small in order to give $\mu_{\text{eff}} \gtrsim 100$ GeV [55].

Depending on the parameters, the Higgs potential of the \mathbb{Z}_3 -invariant NMSSM can possess several local minima (see, e. g., [56]). Notably, one should verify whether vacua where one of the vevs v_u , v_d or s vanishes (typically preferred during the cosmological evolution) are not deeper. A general analysis (taking the radiative corrections to the effective potential into account) is quite involved, but typically one obtains upper bounds on κ at least of the type $\kappa^2 < \lambda^2$ [54].

The tree level Higgs mass matrices are obtained by expanding the full scalar potential (2.9) around the real neutral vevs v_u , v_d and s as in (2.10). Then, the elements of the 3×3 CP-even mass matrix \mathcal{M}_S^2 read in the basis (H_{dR}, H_{uR}, S_R) after the elimination of

$m_{H_d}^2$, $m_{H_u}^2$ and m_S^2 (still in the general NMSSM, but assuming $\mu = 0$)

$$\begin{aligned}
\mathcal{M}_{S,11}^2 &= g^2 v_d^2 + (\mu_{\text{eff}} B_{\text{eff}} + \widehat{m}_3^2) \tan \beta , \\
\mathcal{M}_{S,22}^2 &= g^2 v_u^2 + (\mu_{\text{eff}} B_{\text{eff}} + \widehat{m}_3^2) / \tan \beta , \\
\mathcal{M}_{S,33}^2 &= \lambda(A_\lambda + \mu') \frac{v_u v_d}{s} + \kappa s (A_\kappa + 4\kappa s + 3\mu') - (\xi_S + \xi_F \mu') / s , \\
\mathcal{M}_{S,12}^2 &= (2\lambda^2 - g^2) v_u v_d - \mu_{\text{eff}} B_{\text{eff}} - \widehat{m}_3^2 , \\
\mathcal{M}_{S,13}^2 &= \lambda(2\mu_{\text{eff}} v_d - (B_{\text{eff}} + \kappa s + \mu') v_u) , \\
\mathcal{M}_{S,23}^2 &= \lambda(2\mu_{\text{eff}} v_u - (B_{\text{eff}} + \kappa s + \mu') v_d) .
\end{aligned} \tag{2.22}$$

Rotating the upper left 2×2 submatrix by an angle β , one finds that one of its diagonal elements reads

$$M_Z^2 \left(\cos^2 2\beta + \frac{\lambda^2}{g^2} \sin^2 2\beta \right) \tag{2.23}$$

which constitutes an upper bound on the lightest eigenvalue of \mathcal{M}_S^2 . The additional positive contribution $\sim \lambda^2 \sin^2 2\beta$ (as compared to the MSSM) in the NMSSM is highly welcome in view of the present lower LEP bound of ~ 114 GeV on the mass of a Higgs scalar with SM-like couplings to gauge bosons [57]. However, this additional contribution is relevant only for not too large $\tan \beta$; in fact, the expression inside the parenthesis in (2.23) is larger than one only for $\lambda^2 > g^2$, in which case it is maximal for small $\tan \beta$. Moreover, the actual lightest eigenvalue of \mathcal{M}_S^2 is smaller than the value given in (2.23) in general, see the discussion in Section 3.2.

In the general NMSSM, the elements of the 3×3 CP-odd mass matrix \mathcal{M}'_P read in the basis (H_{dI}, H_{uI}, S_I)

$$\begin{aligned}
\mathcal{M}'_{P,11} &= (\mu_{\text{eff}} B_{\text{eff}} + \widehat{m}_3^2) \tan \beta , \\
\mathcal{M}'_{P,22} &= (\mu_{\text{eff}} B_{\text{eff}} + \widehat{m}_3^2) / \tan \beta , \\
\mathcal{M}'_{P,33} &= \lambda(B_{\text{eff}} + 3\kappa s + \mu') \frac{v_u v_d}{s} - 3\kappa A_\kappa s - 2m_S'^2 - \kappa \mu' s - \xi_F \left(4\kappa + \frac{\mu'}{s} \right) - \frac{\xi_S}{s} , \\
\mathcal{M}'_{P,12} &= \mu_{\text{eff}} B_{\text{eff}} + \widehat{m}_3^2 , \\
\mathcal{M}'_{P,13} &= \lambda v_u (A_\lambda - 2\kappa s - \mu') , \\
\mathcal{M}'_{P,23} &= \lambda v_d (A_\lambda - 2\kappa s - \mu') .
\end{aligned} \tag{2.24}$$

\mathcal{M}'_P contains always a massless Goldstone mode G . Next we rotate this mass matrix into the basis (A, G, S_I) , where $A = \cos \beta H_{uI} + \sin \beta H_{dI}$:

$$\begin{pmatrix} H_{dI} \\ H_{uI} \\ S_I \end{pmatrix} = \begin{pmatrix} \sin \beta & -\cos \beta & 0 \\ \cos \beta & \sin \beta & 0 \\ 0 & 0 & 1 \end{pmatrix} \begin{pmatrix} A \\ G \\ S_I \end{pmatrix} . \tag{2.25}$$

Dropping the Goldstone mode, the remaining 2×2 mass matrix \mathcal{M}_P^2 in the basis (A, S_I)

has the elements

$$\begin{aligned}
\mathcal{M}_{P,11}^2 &= \frac{2(\mu_{\text{eff}} B_{\text{eff}} + \widehat{m}_3^2)}{\sin 2\beta}, \\
\mathcal{M}_{P,22}^2 &= \lambda(B_{\text{eff}} + 3\kappa s + \mu') \frac{v_u v_d}{s} - 3\kappa A_\kappa s - 2m_S'^2 - \kappa\mu' s - \xi_F \left(4\kappa + \frac{\mu'}{s}\right) - \frac{\xi_S}{s}, \\
\mathcal{M}_{P,12}^2 &= \lambda(A_\lambda - 2\kappa s - \mu') v.
\end{aligned} \tag{2.26}$$

In the \mathbb{Z}_3 -invariant NMSSM, \mathcal{M}_P^2 simplifies to

$$\begin{aligned}
\mathcal{M}_{P,11}^2 &= \frac{2\mu_{\text{eff}} B_{\text{eff}}}{\sin 2\beta}, \\
\mathcal{M}_{P,22}^2 &= \lambda(B_{\text{eff}} + 3\kappa s) \frac{v_u v_d}{s} - 3\kappa A_\kappa s, \\
\mathcal{M}_{P,12}^2 &= \lambda(A_\lambda - 2\kappa s) v
\end{aligned} \tag{2.27}$$

with B_{eff} as given in (2.14). The matrix element $\mathcal{M}_{P,11}^2$ corresponds to the mass squared M_A^2 of the (only physical) CP-odd scalar A of the MSSM.

Finally the charged Higgs mass matrix in the basis $(H_u^+, H_d^{-*} = H_d^+)$ is given by

$$\mathcal{M}'_{\pm} = \left(\mu_{\text{eff}} B_{\text{eff}} + \widehat{m}_3^2 + v_u v_d \left(\frac{g_2^2}{2} - \lambda^2 \right) \right) \begin{pmatrix} \cot \beta & 1 \\ 1 & \tan \beta \end{pmatrix}. \tag{2.28}$$

It contains one massless Goldstone mode, and one eigenstate with mass

$$\mathcal{M}_{\pm}^2 = \frac{2(\mu_{\text{eff}} B_{\text{eff}} + \widehat{m}_3^2)}{\sin 2\beta} + v^2 \left(\frac{g_2^2}{2} - \lambda^2 \right). \tag{2.29}$$

Due to the term $\sim \lambda^2$, the charged Higgs mass in the NMSSM can be somewhat smaller than in the MSSM (for a given value of $M_A^2 \equiv \mathcal{M}_{P,11}^2$). In contrast to the MSSM it is not even guaranteed within the NMSSM that $U(1)_{\text{em}}$ remains unbroken: considering again the \mathbb{Z}_3 -invariant NMSSM where $\widehat{m}_3^2 = 0$, the expression for the charged Higgs mass squared becomes negative for $s = \mu_{\text{eff}} = 0$, $\lambda^2 > g_2^2/2$, indicating a possible minimum in field space where the charged Higgs has a vev. Although radiative corrections have to be added and the depth of this minimum has to be compared to the physical one with $s \neq 0$, λ is bounded from above by the absence of a charged Higgs vev.

The diagonalization of all scalar mass matrices is carried out in Appendix A (together with the Feynman rules); next we consider the fermionic sector.

First, we have to consider the soft SUSY breaking gaugino mass terms, which do not differ from the MSSM. Denoting the $U(1)_Y$ gaugino by λ_1 , the $SU(2)$ gauginos by λ_2^i ($i = 1, 2, 3$) and the $SU(3)$ gauginos by λ_3^a ($a = 1, \dots, 8$), the soft SUSY breaking gaugino mass terms in the Lagrangian read

$$\mathcal{L} = \frac{1}{2} M_1 \lambda_1 \lambda_1 + \frac{1}{2} M_2 \lambda_2^i \lambda_2^i + \frac{1}{2} M_3 \lambda_3^a \lambda_3^a. \tag{2.30}$$

In the neutralino sector, λ_1 and λ_2^3 mix with the neutral higgsinos $\psi_d^0, \psi_u^0, \psi_S$ and generate a symmetric 5×5 mass matrix \mathcal{M}_0 . In the basis $\psi^0 = (-i\lambda_1, -i\lambda_2^3, \psi_d^0, \psi_u^0, \psi_S)$, the resulting mass terms in the Lagrangian read

$$\mathcal{L} = -\frac{1}{2}(\psi^0)^T \mathcal{M}_0 (\psi^0) + \text{h.c.} \quad (2.31)$$

where

$$\mathcal{M}_0 = \begin{pmatrix} M_1 & 0 & -\frac{g_1 v_d}{\sqrt{2}} & \frac{g_1 v_u}{\sqrt{2}} & 0 \\ & M_2 & \frac{g_2 v_d}{\sqrt{2}} & -\frac{g_2 v_u}{\sqrt{2}} & 0 \\ & & 0 & -\mu_{\text{eff}} & -\lambda v_u \\ & & & 0 & -\lambda v_d \\ & & & & 2\kappa s + \mu' \end{pmatrix} \quad (2.32)$$

and the term μ' in the (5,5) element appears in the general NMSSM only.

As in the MSSM, the charged $SU(2)$ gauginos are $\lambda^- = \frac{1}{\sqrt{2}}(\lambda_2^1 + i\lambda_2^2)$, and $\lambda^+ = \frac{1}{\sqrt{2}}(\lambda_2^1 - i\lambda_2^2)$, which mix with the charged higgsinos ψ_u^+ and ψ_d^- . Defining

$$\psi^+ = \begin{pmatrix} -i\lambda^+ \\ \psi_u^+ \end{pmatrix}, \quad \psi^- = \begin{pmatrix} -i\lambda^- \\ \psi_d^- \end{pmatrix}, \quad (2.33)$$

the corresponding mass terms in the Lagrangian can be written as

$$\mathcal{L} = -\frac{1}{2}(\psi^+, \psi^-) \begin{pmatrix} 0 & X^T \\ X & 0 \end{pmatrix} \begin{pmatrix} \psi^+ \\ \psi^- \end{pmatrix} + \text{h.c.} \quad (2.34)$$

with

$$X = \begin{pmatrix} M_2 & g_2 v_u \\ g_2 v_d & \mu_{\text{eff}} \end{pmatrix}. \quad (2.35)$$

Again, the diagonalization of the neutralino and chargino mass matrices will be described in Appendix A.

Finally we give the top and bottom squark and τ slepton mass-squared matrices. $\tilde{t}_L, \tilde{t}_R, \tilde{b}_L$ and \tilde{b}_R denote the scalar components of the third generation quark superfields $\widehat{U}_{L_3}, \widehat{U}_{R_3}, \widehat{D}_{L_3}$ and \widehat{D}_{R_3} ; $\tilde{\nu}_{\tau L}$ and $\tilde{\tau}_L$ the scalar components of the third generation lepton superfields $\widehat{\nu}_{L_3}$ and \widehat{E}_{L_3} , and $\tilde{\tau}_R$ the scalar component of the third generation lepton superfield \widehat{E}_{R_3} . (The superfields were given in (2.2) and (2.3).) $m_T^2 \equiv m_{U_3}^2, m_B^2 \equiv m_{D_3}^2, m_{Q_3}^2, m_{E_3}^2$ and $m_{L_3}^2$ are the soft SUSY breaking masses squared for the third generation (see $\mathcal{L}_{\text{soft}}$ in (2.5)), assumed to be diagonal in family space.

The top squark mass matrix reads in the basis $(\tilde{t}_R, \tilde{t}_L)$:

$$\begin{pmatrix} m_T^2 + h_t^2 v_u^2 - (v_u^2 - v_d^2) \frac{g_1^2}{3} & h_t (A_t v_u - \mu_{\text{eff}} v_d) \\ h_t (A_t v_u - \mu_{\text{eff}} v_d) & m_{Q_3}^2 + h_t^2 v_u^2 + (v_u^2 - v_d^2) \left(\frac{g_1^2}{12} - \frac{g_2^2}{4} \right) \end{pmatrix}. \quad (2.36)$$

The bottom squark mass matrix reads in the basis $(\tilde{b}_R, \tilde{b}_L)$:

$$\begin{pmatrix} m_B^2 + h_b^2 v_d^2 + (v_u^2 - v_d^2) \frac{g_1^2}{6} & h_b (A_b v_d - \mu_{\text{eff}} v_u) \\ h_b (A_b v_d - \mu_{\text{eff}} v_u) & m_{Q_3}^2 + h_b^2 v_d^2 + (v_u^2 - v_d^2) \left(\frac{g_1^2}{12} + \frac{g_2^2}{4} \right) \end{pmatrix}. \quad (2.37)$$

The tau slepton mass matrix reads in the basis $(\tilde{\tau}_R, \tilde{\tau}_L)$:

$$\begin{pmatrix} m_{E_3}^2 + h_\tau^2 v_d^2 + (v_u^2 - v_d^2) \frac{g_1^2}{2} & h_\tau (A_\tau v_d - \mu_{\text{eff}} v_u) \\ h_\tau (A_\tau v_d - \mu_{\text{eff}} v_u) & m_{L_3}^2 + h_\tau^2 v_d^2 - (v_u^2 - v_d^2) \left(\frac{g_1^2 - g_2^2}{4} \right) \end{pmatrix}. \quad (2.38)$$

The tau sneutrino ($\tilde{\nu}_{\tau_L}$) mass squared is:

$$m_{L_3}^2 - (v_u^2 - v_d^2) \left(\frac{g_1^2 + g_2^2}{4} \right). \quad (2.39)$$

Herewith we conclude the presentation of the tree level Lagrangian; clearly one has to add radiative corrections to the Lagrangian and all the resulting mass matrices. For the couplings/Feynman rules we refer to Appendix A.

2.2 Limiting cases: the effective MSSM and approximate global symmetries

2.2.1 The effective MSSM

As it becomes clear from the superpotential (2.1) of the general NMSSM, all couplings between the components of the singlet superfield \hat{S} and the MSSM Higgs superfields \hat{H}_u and \hat{H}_d vanish for $\lambda \rightarrow 0$. In order to generate a reasonably large value for μ_{eff} (in the absence of an MSSM-like μ -term in the superpotential), one should keep $\lambda s \gtrsim 100$ GeV in this limit. From (2.21) one finds that the vev s scales as $1/\kappa$, hence a reasonable decoupling limit is

$$\lambda \sim \kappa \rightarrow 0, \quad s \sim 1/\kappa \rightarrow \infty, \quad (2.40)$$

while keeping all dimensionful parameters fixed.

The corresponding parameters of the effective MSSM can be easily deduced from (2.7), (2.14) as well as the 1-2 components of the CP-even and CP-odd Higgs mass matrices (2.22) and (2.24) (and the charged Higgs and neutralino mass matrices):

$$\begin{aligned} \mu_{\text{eff}} &= \lambda s, \\ m_{3\text{eff}}^2 &= \hat{m}_3^2 + \mu_{\text{eff}} (A_\lambda + \kappa s). \end{aligned} \quad (2.41)$$

In the \mathbb{Z}_3 -invariant NMSSM \hat{m}_3^2 vanishes, and one has

$$\begin{aligned} m_{3\text{eff}}^2 &= \mu_{\text{eff}} B_{\text{eff}}, \\ B_{\text{eff}} &= A_\lambda + \kappa s. \end{aligned} \quad (2.42)$$

In the limit (2.40) all couplings between the MSSM sector (including quarks, leptons etc.) and the CP-even, CP-odd and fermionic singlet states vanish; it seems *à priori* impossible to distinguish the NMSSM from the MSSM, since the singlet-like states would never be produced. However, in this limit the singlino-like neutralino χ_S^0 with its mass given by $(\mathcal{M}_0)_{55} = 2\kappa s$ (in the \mathbb{Z}_3 -invariant NMSSM) can easily be the LSP. Then, assuming R-parity conservation and a small non-vanishing value of λ , all sparticle decay cascades will

first proceed as in the MSSM into the MSSM-like NLSP (which could be a charged slepton!) which, at the end, will decay into $\chi_S^0 + \text{SM-particles}$. The final decay of the MSSM-like NLSP involves necessarily a coupling $\sim \lambda$, implying a possibly very long lifetime of the NLSP leading to displaced vertices [58]. Then, the difference between the NMSSM and the MSSM can be spectacular even in the decoupling limit (2.40).

2.2.2 The Peccei-Quinn symmetry limit

If the term $\lambda \widehat{S} \widehat{H}_u \cdot \widehat{H}_d$ would be the only \widehat{S} dependent term in the superpotential of the \mathbb{Z}_3 -invariant NMSSM, the Lagrangian would be invariant under a Peccei-Quinn-like symmetry

$$H_u \rightarrow H_u e^{i\varphi_{PQ}}, \quad H_d \rightarrow H_d e^{i\varphi_{PQ}}, \quad S \rightarrow S e^{-2i\varphi_{PQ}} \quad (2.43)$$

which allows to solve the strong CP-problem [59, 60]. Since this global symmetry is spontaneously broken by the vevs v_u, v_d and s , a massless Nambu-Goldstone boson (the Peccei-Quinn axion) would appear in the CP-odd scalar sector, as can be verified by computing the determinant of \mathcal{M}_P^2 in the basis (A, S_I) in (2.27) for $\kappa = 0$.

The decomposition of the Peccei-Quinn axion A_{PQ} in terms of the weak eigenstates H_{uI}, H_{dI} and S_I is given by

$$\begin{aligned} A_{PQ} &= \frac{1}{N} (v \sin 2\beta A - 2s S_I), \quad \text{where} \\ N &= \sqrt{v^2 \sin^2 2\beta + 4s^2}, \quad A = \cos \beta H_{uI} + \sin \beta H_{dI}. \end{aligned} \quad (2.44)$$

Hence, in most of the parameter space where $s \gg v \sin 2\beta$, A_{PQ} is dominantly (but never purely) singlet-like.

Apart from the strong cosmological constraints on a Peccei-Quinn axion, it is not straightforward to stabilise the potential for the vev s of the NMSSM in the Peccei-Quinn limit; obviously the approximate expressions in (2.19) and (2.21) are no longer appropriate for $\kappa = 0$. In [61], couplings to additional singlets have been introduced in order to stabilise the potential for s . In [62–64], the limit $\lambda \ll 1$ has been considered, the potential including dominant radiative corrections in [65], and the Peccei-Quinn limit for large λ in [66]. In any case, the stability of the scalar potential imposes strong constraints on the parameters of the NMSSM in the strict Peccei-Quinn limit $\kappa = 0$ [51, 62–66].

If one prefers to avoid the cosmological constraints on a very light Peccei-Quinn axion, one can consider the situation where the Peccei-Quinn symmetry is explicitly broken by small additional terms in the superpotential of the NMSSM such as a small non-vanishing value for κ in the \mathbb{Z}_3 -invariant NMSSM [51] as it can be obtained in constructions of the NMSSM from the heterotic string [67]. Then the axion acquires a mass at tree level (without QCD contributions); if it is still light, it can lead to distinctive signatures at colliders due to the possible decay of the SM-like CP-even Higgs scalar into two pseudo-Goldstone bosons [68]. (For a scenario with an approximate Peccei-Quinn symmetry and large λ , see [66].)

In any case, the couplings of A_{PQ} are important for its phenomenological signatures: the couplings of A_{PQ} to gauge bosons, quarks and leptons are induced by its doublet

components indicated in (2.44), which are small for $s \gg v \sin 2\beta$. As a result, its couplings g_{Add} to down-type quarks and leptons are not enhanced for large $\tan\beta$, but given by

$$g_{Add} \sim \frac{v}{s} g_{Hdd}^{\text{SM}} \quad (2.45)$$

where g_{Hdd}^{SM} is the coupling of the SM-like CP-even Higgs scalar to down-type quarks and leptons. The couplings of A_{PQ} to all CP-even scalars H_i vanish in the strict Peccei-Quinn limit, but become non-zero as soon as the Peccei-Quinn symmetry is explicitly broken. Then, while small, these couplings can still induce a branching fraction for $H_i \rightarrow 2 A_{\text{PQ}}$ which is larger than the branching fraction for $H_i \rightarrow b\bar{b}$ [68].

2.2.3 The R-symmetry limit

For $A_\lambda, A_\kappa \rightarrow 0$, the Higgs sector of the \mathbb{Z}_3 -invariant NMSSM (specified by the superpotential (2.6) and the soft terms in (2.8)) is invariant under an R-symmetry under which the scalar fields transform as

$$H_u \rightarrow H_u e^{i\varphi_R}, \quad H_d \rightarrow H_d e^{i\varphi_R}, \quad S \rightarrow S e^{i\varphi_R}. \quad (2.46)$$

Again, this global symmetry is spontaneously broken by the vevs v_u, v_d and s , and a massless Nambu-Goldstone boson (now an R-axion) would appear in the CP-odd scalar sector, as can be verified by computing the determinant of \mathcal{M}_P^2 in (2.25) for $B_{\text{eff}} = \kappa s$, $A_\lambda = A_\kappa = 0$. The decomposition of the R-axion A_R in terms of the weak eigenstates H_{uI}, H_{dI} and S_I is given by

$$A_R = \frac{1}{N} (v \sin 2\beta A + s S_I), \quad \text{where } N = \sqrt{v^2 \sin^2 2\beta + s^2} \quad (2.47)$$

and A as in (2.44). Again, in most of the parameter space where $s \gg v \sin 2\beta$, A_R is dominantly (but never purely) singlet-like. However, in the full Lagrangian this R-symmetry is explicitly broken by the gaugino mass terms, which induce non-vanishing values for A_λ (and subsequently for A_κ) through radiative corrections, as can be seen from the β -functions for A_λ and A_κ in Appendix B.3. Even if the statement $A_\lambda = A_\kappa = 0$ is scale dependent (and hence unnatural), one can still consider the situation where both trilinear couplings are relatively small [69–71] implying a light “pseudo” R-Goldstone boson given approximately by A_R as in (2.47).

The couplings of A_R to down-type quarks and leptons (in the R-symmetry limit and for large $\tan\beta$) differ by a factor 2 from (2.45) and are given by

$$g_{Add} \sim \frac{2v}{s} g_{Hdd}^{\text{SM}}, \quad (2.48)$$

while the couplings to CP-even scalars H_i depend on the values of A_λ and A_κ . Again, this scenario can allow for possible (even dominant) $H_i \rightarrow 2 A_R$ decays with important consequences for Higgs searches at colliders [69–71].

Hence, light pseudoscalars can easily appear in the NMSSM in the form of (pseudo-) Nambu-Goldstone bosons; however, light pseudoscalars in the NMSSM can also result from accidental relations among the parameters in which case they would not suffer from suppressed couplings to down-type quarks and leptons. Phenomenological consequences of light pseudoscalars will be discussed in Sections 5 and 6.

3 Radiative corrections

3.1 Renormalisation group equations

Supersymmetry allows to formulate models at a very high scale, such as the GUT or Planck scale, while avoiding quadratically divergent quantum corrections involving an ultraviolet cutoff $\Lambda \sim M_{\text{GUT}}$ or $\Lambda \sim M_{\text{Planck}}$. However, assuming $M_{\text{SUSY}} \sim M_{\text{weak}}$, the quantum corrections still generate large logarithms $\sim \ln(\Lambda/M_{\text{SUSY}})$.

Fortunately, these can be summed up by the introduction of scale dependent parameters in the Lagrangian, where the scale dependence is described by the renormalisation group equations (RGEs) or β -functions. If the parameters are assumed to be given at a large scale $\Lambda \sim M_{\text{GUT}}$ or $\Lambda \sim M_{\text{Planck}}$, the large logarithms are accounted for by the integration of the RGEs down to a low scale $M_{\text{SUSY}} \sim M_{\text{weak}}$.

In the case of the three gauge couplings of the SM, this procedure allows for a successful unification at $M_{\text{GUT}} \sim 2 \times 10^{16}$ GeV in the MSSM [6–9]. It should be underlined that gauge coupling unification remains valid in the NMSSM: the additional gauge singlet field has no effect on the one-loop β -functions of the gauge couplings, and the additional Yukawa couplings λ and κ only appear in the two-loop terms of the gauge β -functions (see (B.1) in Appendix B). The resulting effect on the numerical values of the gauge couplings at the GUT scale is negligibly small (of the order of the unknown threshold effects at the GUT scale).

However, this is true only if the running Yukawa couplings λ , κ , h_t and h_b remain sufficiently small so that perturbation theory remains valid; typically one requires that Yukawa couplings remain below 1 or $\sqrt{4\pi}$. Once the running Yukawa couplings have assumed values of $\mathcal{O}(1)$ at a given scale, naïve extrapolations of the perturbative RGEs result in singularities (so-called Landau singularities) for the running couplings very close to that scale; as a result, upper bounds of 1 or $\sqrt{4\pi}$ have nearly the same consequences in practice. Since the running Yukawa couplings increase towards large scales (unless the one-loop contributions from gauge couplings are dominant, see the β -functions in Appendix B), avoiding Landau singularities implies upper bounds on the couplings at the scale $M_{\text{SUSY}} \sim M_{\text{weak}}$.

The β -functions for the Yukawa couplings have been derived for the NMSSM to one-loop order in [53], and to two-loop order in [72, 73], see Appendix B.1. (Studies of analytic solutions, quasi-fixed points and RG invariants have been performed in [74–78]; the impact of the NMSSM specific Yukawa couplings on h_b - h_τ unification was studied in [79].) Upper bounds on the Yukawa couplings at the scale M_{weak} are given in [46, 80–87] to one-loop order, and in [72] to two-loop order, with the following results: The top quark Yukawa coupling h_t at the scale m_t depends not only on the top quark mass m_t , but increases with decreasing $\tan\beta$ as given in (A.11). Hence, for small $\tan\beta$, h_t can become too large ($\gtrsim \sqrt{4\pi}$) at the GUT scale. Depending on the precise value of m_t and on the threshold corrections between the scales m_t and M_{SUSY} , one obtains a lower bound on $\tan\beta$ close to unity (typically between 1.5 and 2) both in the MSSM and the NMSSM. A similar reasoning based on h_b leads to an upper bound on $\tan\beta$, $\tan\beta \lesssim 80$.

The NMSSM specific Yukawa coupling λ plays an important rôle for the upper bound on the mass of the lightest CP-even Higgs scalar, see (2.23). The validity of perturbation

theory up to the GUT scale implies $\lambda < 0.7-0.8$ at the weak scale, depending on the precise value of m_t , on the threshold corrections, on κ and notably on $\tan\beta$: a large value of h_t (i. e. small values of $\tan\beta$) amplifies the increase of λ towards large scales, thus implying a decreasing upper limit on λ at low scales. Larger values of κ have the same effect on λ .

These features are visible in Fig. 1, where we plot λ_{\max} as function of $\tan\beta$ and κ (using NMSSMTools [52,88], see Appendix D). The black (dark) and red (grey) bands correspond to different choices of the Higgs and sparticle spectrum: the black bands correspond to a light spectrum with sfermion masses of 200 GeV, $\mu_{\text{eff}} = 100$ GeV, a bino mass M_1 of 50 GeV (with M_2, M_3 from (3.1) below) and Higgs masses below ~ 200 GeV. The red bands correspond to a heavy spectrum with sfermion masses, μ_{eff} and heavy Higgs bosons of 1 TeV, and $M_1 = 200$ GeV. (The squark/slepton trilinear couplings are chosen to vanish in both cases for simplicity.) Inside the black and red bands, the top quark mass is 171.2 ± 2.1 GeV [89], a larger top mass leading to a slightly lower upper bound λ_{\max} . However, the variation of λ_{\max} with the threshold corrections of the running gauge and Yukawa couplings induced by the unknown sparticle and (heavy) Higgs spectrum is numerically more important.




Figure 1: Left panel: upper bound on λ (λ_{\max}) as a function of $\tan\beta$ for fixed $\kappa = 0.01$. Right panel: λ_{\max} as a function of κ for fixed $\tan\beta = 10$. Black (lower) bands: light spectrum, red (upper) bands: heavy spectrum. Inside the bands the top quark mass is 171.2 ± 2.1 GeV.

On the left-hand side in Fig. 1, κ is fixed to 0.01; one sees both the decrease of the upper bound on λ for $\tan\beta \sim 1$ and very large $\tan\beta$ induced by very large values of h_t , h_b , respectively. On the right-hand side, $\tan\beta = 10$ and one sees the decrease of the upper bound on λ with increasing κ .

Motivated by simple models for supersymmetry breaking, the soft SUSY breaking parameters are often assumed to satisfy particularly simple relations at the GUT or the Planck

scale such as universal gaugino masses $M_{1/2}$, universal trilinear couplings A_0 and universal scalar masses m_0 (where, sometimes, the singlet mass and trilinear coupling are allowed to play a special rôle, see Section 7.1.3). Then, the RG evolution of these parameters from the high to the weak scale plays an important rôle.

The one-loop β -functions for all soft SUSY breaking parameters of the MSSM were computed in [90, 91], for the NMSSM in [53, 72], and for a general supersymmetric model in [92, 93]. The general two-loop β -functions for the soft terms can be found in [94, 95], from which those of the (general) NMSSM can be deduced. They are given in Appendix B.

Universal gaugino masses $M_1 = M_2 = M_3 = M_{1/2}$ at the GUT scale imply specific relations among these parameters (defined in (2.30)) at a low scale. At 1 TeV one finds (for squark masses $\lesssim 1$ TeV)

$$M_3 : M_2 : M_1 \sim 5.5 : 1.9 : 1 \quad (3.1)$$

with somewhat larger ratios (depending on threshold effects) below 1 TeV. These values hold for the MSSM as well as for the NMSSM.

In the case of universal scalar masses m_0 at the GUT scale (with $m_0^2 \geq 0$), the RG evolution down to the weak scale is essential for the Higgs masses: at the weak scale, at least one Higgs mass squared has to be negative in order to trigger electroweak symmetry breaking. Fortunately $m_{H_u}^2 < 0$ at the weak scale is induced nearly automatically by the large top Yukawa coupling [19], given the large top quark mass.

In models with GMSB, the soft SUSY breaking terms are induced radiatively at a messenger scale M_{mess} , which is typically far above the weak scale. In principle, all possible terms in the Lagrangian of the general NMSSM can be generated (see [96] and Section 7.2). Then, all these parameters have to be evolved by the RGEs from the messenger scale down to the weak scale. For completeness, we give the 2-loop β -functions for all parameters of the general NMSSM in Appendix B.

3.2 Radiative corrections to the Higgs masses, and the upper bound on the lightest CP-even Higgs mass

The Higgs mass matrices in Section 2, and notably the upper bound (2.23) on the mass of the lightest CP-even Higgs scalar, have been derived from the tree level potential (2.9), where all quartic terms are determined by supersymmetry through the superpotential and (electroweak) supersymmetric gauge interactions. This hypothesis would be justified only if the scale of supersymmetry breaking M_{SUSY} would be smaller than the Higgs vevs (or M_Z), which is obviously not the case.

However, for $M_{\text{SUSY}} > M_Z$ the deviation of the Higgs potential from (2.9) is calculable – as a function of the mass splittings among the superpartners – in the form of quantum corrections involving scales (momenta) Q^2 with $M_Z^2 \lesssim Q^2 \lesssim M_{\text{SUSY}}^2$. The dominant contributions to V_{Higgs} originate from top quark/squark loops, since these particles have the largest couplings to Higgs fields (to H_u), and lead to an increase of the upper bound (2.23) on the mass of the lightest CP-even Higgs scalar. This phenomenon had first been discussed in the MSSM to one-loop and subsequently to two-loop order in [97–103] where, at tree level, (2.23) with $\lambda = 0$ would imply a mass of the lightest CP-even Higgs scalar below M_Z .

An approximate formula for the mass M_{SM} of the SM-like Higgs scalar in the NMSSM in the limit $\kappa s \gg |A_\kappa|, |A_\lambda|$ (corresponding to a heavy singlet-like scalar), including the dominant top/stop radiative corrections, is given by

$$M_{\text{SM}}^2 \simeq M_Z^2 \cos^2 2\beta + \lambda^2 v^2 \sin^2 2\beta - \frac{\lambda^2}{\kappa^2} v^2 (\lambda - \kappa \sin 2\beta)^2 + \frac{3m_t^4}{4\pi^2 v^2} \left(\ln \left(\frac{m_T^2}{m_t^2} \right) + \frac{A_t^2}{m_T^2} \left(1 - \frac{A_t^2}{12m_T^2} \right) \right) \quad (3.2)$$

where v is defined in (2.12), the soft SUSY breaking stop masses squared in (2.36) are assumed to satisfy $m_T^2 \sim m_{Q_3}^2 \gg m_t^2$, A_t is the stop trilinear coupling assumed to satisfy $|A_t| \gg m_t, \mu_{\text{eff}}$; the terms $\sim \lambda^2$ are specific to the NMSSM, and the last term in the first line originates from the mixing with the singlet-like scalar. In the MSSM, where $\lambda = 0$, the LEP bound on M_{SM} implies that $\tan \beta$ has to be large such that $\cos 2\beta \sim 1$, m_T above ~ 300 GeV for maximal mixing ($A_t^2 \sim 6m_T^2$, maximising the second line in (3.2)), or $\gtrsim 1$ TeV otherwise.

In order to maximise M_{SM} in the NMSSM, λ should be as large as possible, and $\tan \beta$ should be small in order to avoid a suppression from $\sin^2 2\beta$. (As discussed before, λ is bounded from above by $\lambda \lesssim 0.7 - 0.8$ if one requires the absence of a Landau singularity below the GUT scale.) However, the negative contribution from the mixing with the singlet-like scalar should vanish; without neglecting A_λ , the relevant mixing term is proportional to $(\lambda - \sin 2\beta(\kappa + A_\lambda/(2s)))^2$ [104]. If this expression is not small, a larger value of λ can even generate a decrease of the mass of the Higgs scalar with SM-like couplings to the Z boson in the NMSSM.

The resulting upper bound on the lightest CP-even Higgs mass in the NMSSM has been studied in the leading log approximation in [82–84, 103, 105–109]. Full one-loop calculations of the corresponding upper bound involving top/bottom quark/squark loops have been carried out in [72, 85, 86, 110–117]. (Analyses at large values of $\tan \beta$ have been performed in [118–120], and upper bounds for more general supersymmetric Higgs sectors have been considered in [121–123].)

At present, additional known radiative corrections to the Higgs mass matrices in the NMSSM include MSSM-like electroweak together with the NMSSM-specific Higgs one-loop contributions [124, 125] and dominant two-loop terms [73, 87, 125–127]. In order to discuss these in detail, it is convenient to separate the quantum corrections involving scales Q^2 with $Q^2 \gtrsim M_{\text{SUSY}}^2$ from those with scales $Q^2 \lesssim M_{\text{SUSY}}^2$.

The result of the quantum corrections with $Q^2 \gtrsim M_{\text{SUSY}}^2$ is still a supersymmetric effective Lagrangian (including soft SUSY breaking terms), where all running parameters (couplings and masses) are defined, within a given subtraction scheme, at the scale $Q^2 \sim M_{\text{SUSY}}^2$. (If desired, the parameters at the scale Q^2 can be obtained in terms of parameters at a higher scale with the help of the RGEs.) Subsequently, the quantum corrections with $Q^2 \lesssim M_{\text{SUSY}}^2$ (i. e. with an ultraviolet cutoff M_{SUSY}^2) have to be evaluated, generating a non-supersymmetric effective action including an effective Higgs potential, effective couplings of fermions and wave function normalisation constants. From the effective potential and couplings one can derive the so-called running masses, which still differ somewhat from the physical pole masses (the poles of the propagators).

The terms in the effective action can be computed in a systematic expansion in powers of coupling constants and large logarithms where, to start with, the couplings are defined at the scale $Q^2 \sim M_{\text{SUSY}}^2$. Hence, some additional effort is necessary in order to relate the gauge couplings to those measured at the scale M_Z^2 , the Higgs vevs to M_Z , and the Yukawa couplings to the quark and lepton pole masses: the gauge and Yukawa couplings at the scale $Q^2 \sim M_{\text{SUSY}}^2$ have to be obtained from the measured couplings at lower scales by the integration of RGEs where, however, all possible threshold effects have to be taken into account.

The electroweak and NMSSM specific Higgs one-loop contributions, and the two-loop contributions $\propto h_t^2 \alpha_s$, have recently been computed in [125] without an expansion in large logarithms. However, subsequently we confine ourselves to electroweak and NMSSM-specific Higgs one-loop contributions involving large logarithms, and to two-loop contributions proportional to two powers of large logarithms. In Appendix C we summarise the corresponding formulae, which allow to determine the lightest CP-even Higgs mass in the NMSSM with the following accuracy (considering the tree level mass squared following from the first line in (3.2) to be of order g^2 or λ^2 , where g stands for g_1 or g_2):

- (i) One-loop contributions from top/bottom quarks/squarks are fully included, which generate contributions of order h_t^4 , h_b^4 and $g^2 h_t^2$ etc., possibly multiplied by a large logarithm $\ln(M_{\text{SUSY}}^2/m_t^2)$. (Here and below we identify M_{SUSY}^2 with an average value of the soft SUSY breaking squark mass terms, $M_{\text{SUSY}}^2 = m_{Q_3} m_T$. It is possible to allow for different definitions for M_{SUSY}^2 [52] in cases where other soft terms are larger than m_{Q_3}, m_T , which we do not consider here for simplicity.)
- (ii) Additional one-loop corrections considered are of the orders g^4 , $g^2 \lambda^2$, $g^2 \kappa^2$, λ^4 and κ^4 , if multiplied by a large logarithm. Here, loops from squarks and sleptons, charginos and neutralinos, Higgs and gauge bosons provide relevant contributions.
- (iii) At two-loop order, only dominant logarithms $\sim \ln^2(M_{\text{SUSY}}^2/m_t^2)$ multiplied by h_t^6 or $\alpha_s h_t^4$ (and h_t replaced by h_b , possibly relevant at large $\tan\beta$) are considered, including some subdominant effects due to the evaluation of h_t and α_s at the scale M_{SUSY} .

Taking these loop corrections into account, and requiring perturbative running Yukawa couplings h_t , λ and κ below the GUT scale, the upper bound on the lightest CP-even Higgs mass has been studied in [104] as a function of $\tan\beta$ and for different values of m_t in the NMSSM, and compared to the MSSM with the result shown in Fig. 2. (In Fig. 2, the upper bound is denoted as m_{max} .) The squark mass terms (and hence M_{SUSY}) have been chosen as 1 TeV; the upper bound would still increase slowly (logarithmically) with M_{SUSY} . In order to maximise the one-loop top/bottom (s)quark contributions to the lightest CP-even Higgs mass for these squark masses, the trilinear soft couplings are chosen as $A_t = A_b = 2.5$ TeV. The threshold effects depend somewhat on the gaugino masses, which are $M_1 = 150$ GeV, $M_2 = 300$ GeV and $M_3 = 1$ TeV (roughly in agreement with (3.1)); the remaining parameters λ , κ , A_λ , A_κ and μ_{eff} of the \mathbb{Z}_3 -invariant NMSSM have been chosen such that the upper bound is maximised, without encountering Landau singularities (which requires κ as small as possible) nor violating other constraints such as an unstable potential, which forbids $\kappa \rightarrow 0$.

The lower dashed lines in Fig. 2 refer to the MSSM, where the mass of the CP-odd scalar M_A – which can be chosen as the other independent parameter in the Higgs sector




Figure 2: Upper bound on the lightest Higgs mass in the NMSSM as a function of $\tan\beta$ for $m_t = 178$ GeV (M_A arbitrary: thick full line, $M_A = 1$ TeV: thick dotted line) and $m_t = 171.4$ GeV (thin full line: M_A arbitrary, thick dotted line: $M_A = 1$ TeV) and in the MSSM (with $M_A = 1$ TeV) for $m_t = 178$ GeV (thick dashed line) and $m_t = 171.4$ GeV (thin dashed line). Squark and gluino masses are 1 TeV and $A_t = A_b = 2.5$ TeV. (From [104].)

apart from $\tan\beta$ – is set to $M_A = 1$ TeV. The other parameters (and approximations) are the same as described above. In the MSSM, the increase of the upper bound with $\tan\beta$ originates from the tree level term (the first term $\sim \cos^2 2\beta$ in (3.2)), according to which m_{\max} is maximised for large $\tan\beta$. Due to the one-loop top (s)quark contributions, the upper bound m_{\max} increases with m_t . Numerically, a variation Δm_t of m_t implies nearly the same variation Δm_{\max} for large $\tan\beta$.

In the NMSSM, the second term $\sim \sin^2 2\beta$ in the tree level expression (3.2) dominates the first one for sufficiently large λ , and accordingly m_{\max} is maximal for low values of $\tan\beta$. On the other hand, the absence of a Landau singularity for λ below the GUT scale implies a decrease of the maximally allowed value of λ at M_{SUSY} with increasing h_t , i.e. with increasing m_t and decreasing $\tan\beta$, see Fig. 1. (At large $\tan\beta$, arbitrary variations of the NMSSM parameters λ , κ , A_λ , A_κ and μ_{eff} can imply a mass M_A of the MSSM-like CP-odd scalar far above 1 TeV. For comparison with the MSSM, m_{\max} in the NMSSM with $M_A \leq 1$ TeV is depicted as dotted lines in Fig. 2.)

Numerically, for $m_t = 171.4$ GeV, m_{\max} in the NMSSM (thin full line in Fig. 2) is 140 GeV, and assumed for $\tan\beta = 2$, $\lambda = 0.70$ and $\kappa = 0.05$, whereas for $m_t = 178$ GeV

(thick full line in Fig. 2), m_{\max} increases just to 141.5 GeV for $\tan\beta = 2.2$, $\lambda = 0.68$ and $\kappa = 0.07$.

It must be emphasised, however, that the non-observation of a Higgs boson in standard search channels with a mass below $\sim 140 - 145$ GeV (allowing for larger M_{SUSY} and remaining theoretical uncertainties) would *not* rule out the NMSSM: here the lightest CP-even Higgs scalar could have small couplings to all quarks, leptons and gauge bosons if it is dominantly singlet-like, in which case the mass of the Next-to-lightest Higgs scalar with SM-like couplings to gauge bosons is bounded from above, and can be ~ 20 GeV larger [127], see also Section 5.1. Also, the lightest CP-even Higgs scalar could have unconventional decay channels which can make its discovery at colliders quite difficult. In such a case one could possibly detect one of the heavier CP-even Higgs scalars of the NMSSM, whose masses and couplings can vary over a wide range. (Radiative corrections to their masses are also given in the Appendix C.)

Finally, m_{\max} in the NMSSM obviously increases if one allows for larger values of λ [128,129]. Since this would imply a Landau singularity below M_{GUT} for the particle content of the NMSSM, one can consider modifications of the theory at higher scales. For instance, one can allow for new strong gauge interactions at intermediate scales [130–132], large extra dimensions implying a lower effective ultraviolet cutoff [133], or replace the singlet S with its coupling λ by an unparticle operator [134] (in which case the Higgs content would not be NMSSM-like). Modifications of the β -functions due to additional $SU(5)$ multiplets with masses in the TeV range have been considered in [66]: additional $SU(5)$ multiplets lead to larger gauge couplings at larger scales which, in turn, affect the β -function for λ such that larger values of λ at the weak scale are allowed. In [135] an extra $SU(2)$ gauge symmetry has been introduced, acting on the third generation and the Higgs doublets only and allowing for $\lambda \sim 1$ at low energy, still consistent with perturbativity and GUT-scale gauge coupling unification. This, together with small values for $\tan\beta$ (possibly below 1) leads to an upper bound on the lightest Higgs mass of $m_{\max} \lesssim 250$ GeV. The phenomenology of large- λ scenarios with λ up to 1.5 has been studied in [136], again with the result that $\tan\beta$ must be small for $\lambda \rightarrow 1.5$.

3.3 Radiative corrections to coupling constants

Radiative corrections to coupling constants can be ultraviolet divergent or (ultraviolet) finite. In SUSY extensions of the Standard Model, all ultraviolet divergent radiative corrections can be absorbed into redefinitions of the SUSY preserving or soft SUSY breaking masses and couplings, which are effectively described by the RG equations for these parameters (valid between M_{SUSY} and, e. g., M_{GUT}).

However, ultraviolet finite quantum corrections involving scales between M_{weak} and M_{SUSY} can induce new couplings (or modify SUSY relations between existent couplings) which *cannot* be described in terms of an effective supersymmetric theory plus soft terms. Amongst others, such ultraviolet finite quantum corrections to the Higgs mass terms are responsible for the increase of the upper limit on the lightest CP-even Higgs mass with the logarithm of the top squark masses, as discussed in the previous subsection and in Appendix C.

Subsequently we briefly discuss quantum corrections to h_b and comment on the coupling

$Zb\bar{b}$. Radiative corrections to the Higgs self-couplings are given at the end of Appendix A. We leave aside the threshold effects for the running gauge and Yukawa couplings at the various sparticle and Higgs mass scales; for a recent NMSSM specific analysis see [137].

Large $\tan\beta$ corrections to h_b

The superpotentials of the MSSM and the NMSSM contain couplings of H_u to up-type quarks, and of H_d to down-type quarks and leptons. It is well known [138–140] that squark-gluino- and squark-chargino loops can induce a coupling of H_u to down-type quarks and leptons giving rise to an effective Lagrangian [141]

$$\mathcal{L}_{\text{eff}} = h_b H_d^0 b \bar{b} + \Delta h_b H_u^0 b \bar{b} + \dots \quad (3.3)$$

As a result, the physical b -quark mass m_b is given by

$$m_b = h_b v_d + \Delta h_b v_u \equiv h_b v_d (1 + \Delta m_b) \quad (3.4)$$

with

$$\Delta m_b = \frac{\Delta h_b}{h_b} \tan\beta. \quad (3.5)$$

Notably for large $\tan\beta$, large values of μ (or μ_{eff} in the NMSSM), large gluino masses and/or large values of $|A_t|$, the quantity Δm_b can be sizable; explicit expressions for Δm_b in the MSSM can be found in [138–141]. These hold also in the NMSSM, provided μ is replaced by μ_{eff} . As a first consequence, the formula for the Yukawa coupling h_b as function of m_b in (A.11) has to be corrected and should read

$$h_b^{\text{corr}} = \frac{m_b}{v \cos\beta (1 + \Delta m_b)}. \quad (3.6)$$

The corresponding value for h_b has to be used as boundary condition at the weak scale for the integration of the RGEs up to the GUT scale. Moreover, the couplings of the Higgs bosons to b -quarks in (A.12) have to be corrected as follows:

$$\begin{aligned} S_i b_L b_R^c &: \frac{h_b^{\text{corr}}}{\sqrt{2}} (S_{i1} + \Delta m_b \tan\beta S_{i2}) \\ A_i b_L b_R^c &: i \frac{h_b^{\text{corr}}}{\sqrt{2}} P_{i1} \\ H^- t_L b_R^c &: -h_b^{\text{corr}} \sin\beta, \end{aligned} \quad (3.7)$$

where the couplings of the CP-odd and charged Higgs bosons follow from the $SU(2)$ -invariant completion of the effective Lagrangian (3.3).

In the NMSSM, squark-gluino- and squark-chargino-loops also induce direct couplings of the singlet S to quarks and leptons [142]. They give rise to additional terms $\sim \Delta^S h_b S b \bar{b}$ in the effective Lagrangian (3.3), which have been worked out in the limit of large $\tan\beta$ in [142], together with the additional corrections to the Higgs b -quark couplings (3.7). The terms $\Delta^S h_b S b \bar{b}$ are proportional to the singlet-components of the Higgs mass eigenstates and to $v_{u,d}/s$; hence they are small for $s \gg v_{u,d}$.

The coupling $Zb\bar{b}$

In view of the discrepancy between the Standard Model and the forward-backward asymmetry A_{FB}^b in b -quark production at SLC and LEP [143], radiative corrections to the coupling $Zb\bar{b}$ are of particular interest. However, given that the measurements of $R_b = \Gamma(Z \rightarrow b\bar{b})/\Gamma(Z \rightarrow \text{hadrons})$ agree quite well with the Standard Model, explanations of the discrepancies only in the asymmetries are not easy. A recent analysis of radiative corrections to the $Zb\bar{b}$ vertex in supersymmetric extensions of the SM including the NMSSM is given in [144], including possible NMSSM specific contributions to other electroweak precision observables. The net result of this study is, however, that the NMSSM cannot improve the agreement with the measurements, since NMSSM specific contributions have the opposite sign of the one required.

4 The tadpole and domain wall problems, baryogenesis, hybrid inflation

4.1 The tadpole problem

In order to avoid a fine tuning problem in the NMSSM, none of the dimensionful terms in the superpotential W_{Higgs} (2.1) or in $\mathcal{L}_{\text{soft}}$ in (2.5) should be much larger than the weak or SUSY breaking scale.

However, if a singlet superfield \widehat{S} couples in the most general way to heavy fields (as it is possible in GUTs or GMSB), radiative corrections can induce very large terms in the effective action which are linear in \widehat{S} in the superpotential (denoted by ξ_F in (2.1)) or linear in S in $\mathcal{L}_{\text{soft}}$ (denoted by ξ_S in (2.5)). These terms are called tadpole terms, and if they are too large, the corresponding model has a “tadpole problem”. This problem has been discussed in the context of globally supersymmetric theories with soft supersymmetry breaking terms – as induced by a hidden sector in supergravity – in [29, 145–149], using general power counting rules and/or by explicitly evaluating Feynman diagrams.

Once non-renormalisable supergravity interactions suppressed by powers of M_{Planck} are taken into account, power counting rules signal again a potential tadpole problem [150], as confirmed by explicit calculations in [151–153].

In the case of soft supersymmetry breaking terms $\sim M_{\text{SUSY}}$ induced by a hidden sector in supergravity, the orders of magnitude of the radiatively induced tadpole terms are

$$\begin{aligned}\xi_F &\sim \Lambda M_{\text{SUSY}} , \\ \xi_S &\sim \Lambda M_{\text{SUSY}}^2 ,\end{aligned}\tag{4.1}$$

where $\Lambda \sim M_{\text{GUT}}$ in the presence of general couplings of \widehat{S} to GUT fields, or $\Lambda \sim M_{\text{Planck}}$ in the presence of the most general non-renormalisable supergravity interactions of \widehat{S} . In both cases, the induced orders of magnitude of ξ_F and ξ_S are too large.

In the case of GMSB (see Section 7.2), the phenomenologically required soft supersymmetry breaking mass terms of sparticles are induced radiatively; the source of supersymmetry breaking are so-called messenger fields $\widehat{\varphi}$ with SM gauge quantum numbers and supersymmetric mass terms $\sim M_{\text{mess}}$, but whose real and imaginary scalar components have different masses, split by a scale \widehat{m} . This kind of supersymmetry breaking is also denoted as F -type splitting, since it can be represented in terms of a non-vanishing F component of a spurion superfield which couples to the messenger fields $\widehat{\varphi}$. Then, M_{SUSY} in (4.1) is given by $M_{\text{SUSY}} \sim \widehat{m}^2/M_{\text{mess}}$, and Λ by M_{mess} . Hence, if M_{mess} as well as the F -type splitting \widehat{m} are not much larger than the weak scale, the singlet tadpole problem is absent [33, 34, 154–156]; if these scales are larger than the weak scale, the tadpole diagrams can be suppressed by small Yukawa couplings, see [96] and Section 7.2.

In order to circumvent the tadpole problem in the NMSSM in the case of soft supersymmetry breaking terms induced by a hidden sector in supergravity, we note that the dangerous terms (4.1) with $\Lambda \sim M_{\text{GUT}}$, M_{Planck} can be generated only if \widehat{S} is a singlet with respect to *all* continuous and discrete symmetries of the full theory. As soon as the heavy/hidden sector is invariant under a discrete symmetry under which \widehat{S} transforms, the

terms (4.1) are absent if the discrete symmetry is unbroken, or Λ is of the order of the scale of the breakdown of the discrete symmetry (possibly multiplied by high powers of coupling constants and loop factors $1/16\pi^2$). Hence, if the NMSSM is embedded in a GUT or supergravity theory with an exact or approximate discrete symmetry such that $\Lambda \lesssim M_{\text{SUSY}}$, the tadpole problem is avoided.

4.2 The domain wall problem

Discrete symmetries can generate another problem, however [157]: once they are spontaneously broken after a symmetric phase in the hot early universe, domain walls are generated which can dominate the energy density of the universe, creating unacceptably large anisotropies of the cosmic microwave background radiation and spoiling successful nucleosynthesis [157].

In particular, the NMSSM with a scale invariant superpotential is invariant under a \mathbb{Z}_3 -symmetry (see the discussion following (2.7)). Then, after electroweak symmetry breaking, the universe would consist of different “bubbles” of the same vacuum energy, but in each of which the phases of $\langle H_u \rangle$, $\langle H_d \rangle$ and $\langle S \rangle$ would differ by a \mathbb{Z}_3 -transformation; these bubbles are separated by domain walls [158–162].

However, since the \mathbb{Z}_3 -symmetry is just an accidental symmetry of any scale invariant superpotential, it is not expected that it will be preserved by Planck scale suppressed (non-renormalisable) gravitational interactions. Whereas such a violation of the \mathbb{Z}_3 -symmetry is typically sufficient to avoid the domain wall problem (since the vacuum energy within different bubbles would be slightly different, leading to a collapse of the ones with higher energy) [158–161], it may again lead to dangerously large tadpole diagrams [150–153, 162]. It has been believed [162], that the conflict between the domain wall and tadpole problems cannot be solved.

Subsequently solutions of this problem have been proposed in [163–165]: it is possible to impose constraints on \mathbb{Z}_3 -symmetry breaking non-renormalisable interactions or (renormalisable) hidden sectors in the form of various additional symmetries, such that \mathbb{Z}_3 -symmetry breaking *renormalisable* terms – as the tadpole terms above – are generated radiatively, but with very small coefficients. These \mathbb{Z}_3 -symmetry breaking terms can still solve the domain wall problem of the otherwise \mathbb{Z}_3 -invariant NMSSM, without having a visible impact on its phenomenology.

In [166] a \mathbb{Z}_5 -R-symmetry of the non-renormalisable interactions has been considered, as a consequence of which a tadpole term $\sim \xi_S$ is generated only at the six-loop level. If the order of magnitude of ξ_S is $\xi_S \ll M_{\text{SUSY}}^3$, it constitutes one of the possible terms that can solve the domain wall problem of the otherwise \mathbb{Z}_3 -invariant NMSSM. However, one can also assume that $\xi_S \sim M_{\text{SUSY}}^3$ [166]; in this case the trilinear term $\sim \frac{\kappa}{3} \hat{S}^3$ in the superpotential of the NMSSM is not even phenomenologically required and can be omitted. The resulting model has been denoted as the New Minimal MSSM (or nMSSM [167], if a term $\xi_F \sim M_{\text{SUSY}}^2$ is added to the superpotential as well), which we will discuss in more detail in Section 7.3.

4.3 Electroweak baryogenesis

A strongly first order electroweak phase transition is a necessary (albeit not sufficient) condition for baryogenesis relying on anomalous baryon number violating Standard Model processes [168]: during a first order phase transition bubbles are formed, within which the Higgs fields assume non-vanishing values corresponding to the absolute minimum of the zero temperature effective potential. These bubbles are separated by expanding “walls” from the phase with vanishing Higgs vevs corresponding to the minimum of the effective potential at high temperature. If the dynamics of the processes across the expanding walls sufficiently violates CP and baryon number, the baryon asymmetry of the present universe could be explained [168].

A first order electroweak phase transition is difficult to achieve in the SM and in simple supersymmetric extensions as the MSSM, given the present lower bounds on Higgs and stop masses (which play an important rôle for the one-loop corrections to the effective potential); see, e.g., [169]. On the other hand, the more complicated tree level Higgs potential in various versions of the NMSSM can easily give rise to a sufficiently strong electroweak phase transition [170–179]. However, the trajectory in the space of fields S , H_u^0 and H_d^0 must be studied carefully in order to avoid a phase transition in two steps where first S alone assumes a vev (implying an insufficient violation of CP) and subsequently H_u^0 and H_d^0 assume their vevs as in the MSSM with fixed μ_{eff} . To this end, the vevs of S , H_u^0 and H_d^0 should be of the same order along the trajectory in field space [172, 178], without violating present bounds on the Higgs sector. This seems easier to achieve in the presence of additional terms in the superpotential and/or $\mathcal{L}_{\text{soft}}$ beyond those of the \mathbb{Z}_3 -invariant NMSSM [171, 173, 174]. Simultaneously, such additional terms explicitly break the \mathbb{Z}_3 -symmetry avoiding the domain wall problem. The simplest additional terms are actually the tadpole terms $\sim \xi_S$ and/or $\sim \xi_F$, which allow to omit the terms $\sim \kappa$ of the \mathbb{Z}_3 -invariant NMSSM; the resulting model is the nMSSM mentioned above and discussed in Section 7.3. Electroweak baryogenesis in the nMSSM has been studied in [176, 177, 179].

In addition to a first order phase transition responsible for the formation of bubble walls, the processes inside the expanding bubble walls have to violate CP in order to generate a baryon asymmetry. CP violation can originate from the chargino sector (through a non-vanishing phase of $M_2 \times \mu_{\text{eff}}$ in the NMSSM) or the Higgs sector (the prospects for CP violation in the Higgs sector of the NMSSM are reviewed in Section 8.1). In both cases, strong constraints from the upper bound on the (neutron) electric dipole moments have to be respected, which is possible in the nMSSM [176, 179].

The possibility of spontaneous CP violation inside the bubble walls without CP violation at zero temperature has been advocated in [180, 181]; then, bounds on the neutron electric dipole moment are trivially satisfied. A similar mechanism inside the domain walls separating phases related by the \mathbb{Z}_3 -symmetry of the \mathbb{Z}_3 -invariant NMSSM has been considered in [161]. Electroweak baryogenesis in a $U(1)'$ model with a secluded $U(1)'$ -breaking sector (see Section 8.2.1), where one can have a sufficiently strong first order electroweak phase transition, has been studied in [182, 183]. Once the criteria for successful baryogenesis are fulfilled in the NMSSM, the background density of gravitational waves produced during the electroweak phase transition can be within reach of the gravitational wave experiment LISA [184, 185].

4.4 Hybrid inflation

A variant of the NMSSM without domain wall problem, solving the strong CP problem via an invisible axion, and allowing for hybrid inflation has been proposed and discussed in [186–191]. Two singlets ϕ and N are introduced, and the relevant part of the superpotential is given by

$$W = \lambda\phi H_u H_d + \kappa\phi N^2. \quad (4.2)$$

The soft SUSY breaking mass squared of ϕ is assumed to be very small and negative, and both Yukawa couplings λ and κ are assumed to be $\lambda \sim \kappa \sim 10^{-10}$ which can be motivated by brane constructions and/or type I string theory. The model has a Peccei-Quinn symmetry (see Section 2.2.2) with Peccei-Quinn charges -2 for ϕ and +1 for N . During the inflationary epoch the vevs of N , H_u and H_d vanish, and the inflaton ϕ rolls slowly along an almost flat direction of the scalar potential. Only near the end of inflation N , H_u and H_d develop vevs. Domain walls, created by the spontaneous breaking of the Peccei-Quinn symmetry, are diluted by the inflation. The vev of ϕ after inflation is automatically of the correct order such that $\mu_{\text{eff}} = \lambda\phi$ is of the order of the SUSY breaking scale, and the constraints on the parameters associated with inflation (number of e -folds, curvature perturbations and the spectral index) can be satisfied.

5 NMSSM phenomenology at colliders

Due to the additional singlet superfield \widehat{S} , the phenomenology of the NMSSM can differ strongly from the MSSM in the Higgs and neutralino sectors. Subsequently we discuss these subjects separately.

5.1 The Higgs sector

The mass matrices of the three CP-even and the two (physical) CP-odd neutral Higgs sectors are given in Section 2.1 at tree level, and the dominant radiative corrections in Appendix C. In general, after the diagonalization of the mass matrices as in Appendix A, the $SU(2)$ doublets mix with the singlet states in both sectors. As a consequence, the reduced couplings

$$\xi_i = \sin \beta S_{i2} + \cos \beta S_{i1} \quad (5.1)$$

of the 3 CP-even mass eigenstates H_i to the electroweak gauge bosons (normalised with respect to the couplings of the SM Higgs scalar) can be very small; however, they always satisfy the sum rule

$$\sum_{i=1}^3 \xi_i^2 = 1. \quad (5.2)$$

Another useful sum rule involving the eigenvalues $M_{H_i}^2$ of the CP-even mass matrix \mathcal{M}_S^2 has been given in [167]:

$$\begin{aligned} \sum_{i=1}^3 \xi_i^2 M_{H_i}^2 &= \cos^2 \beta (\mathcal{M}_S^2)_{11} + 2 \cos \beta \sin \beta (\mathcal{M}_S^2)_{12} + \sin^2 \beta (\mathcal{M}_S^2)_{22} \\ &= M_Z^2 \left(\cos^2 2\beta + \frac{\lambda^2}{g^2} \sin^2 2\beta \right) + \text{rad. corr.}, \end{aligned} \quad (5.3)$$

where the last line is the upper bound m_{\max}^2 on the lightest CP-even Higgs mass in the NMSSM (corresponding to (2.23) at tree level) including the radiative corrections which lift it up to ~ 140 GeV, see Fig. 2. Using $M_{H_3} \geq M_{H_2}$ (by definition) and eliminating $\xi_2^2 + \xi_3^2$ with the help of (5.2), one can derive from (5.3) [192]

$$M_{H_2}^2 \leq \frac{1}{1 - \xi_1^2} (m_{\max}^2 - \xi_1^2 M_{H_1}^2), \quad (5.4)$$

which will be useful below.

The extended CP-even and CP-odd Higgs sectors allow for the possibility of additional Higgs-to-Higgs decays compared to the MSSM. Subsequently we will discuss separately the regions in the parameter space of the NMSSM where Higgs-to-Higgs are allowed/disallowed.

5.1.1 LEP and $e^+ e^-$ colliders

To start with, the constraints from LEP1 and LEP2 must be translated carefully into constraints on the parameter space of the NMSSM. Early studies [113, 127, 193–196] confined

themselves to the standard Higgs search channels

$$e^+ e^- \rightarrow H Z, \quad H \rightarrow b\bar{b}, \tau^+ \tau^-, \quad (5.5)$$

neglecting possible unconventional Higgs decay modes.

Since the lightest CP-even scalar H_1 can have a dominant singlet component, its reduced coupling ξ_1 to Z bosons can be smaller than in the MSSM with, possibly, $\xi_1 \rightarrow 0$. The LEP Working group for Higgs Boson Searches [57] has published upper bounds on ξ^2 as function of M_H (valid for any H_i) combining results from the ALEPH, DELPHI, L3 and OPAL collaborations, as shown in Fig. 3. Clearly, even very light CP-even scalars are not ruled out if ξ_1 is sufficiently small, and such scenarios can indeed be realised within the NMSSM [113, 114, 127, 175, 192–199].




Figure 3: Upper bounds on ξ^2 from LEP [57], where SM branching ratios $H \rightarrow b\bar{b}$ and $H \rightarrow \tau^+ \tau^-$ are assumed. Full line: observed limit; dashed line: expected limit; dark (green) shaded band: within 68% probability; light (yellow) band: within 95% probability.

In the limit $\xi_1 \rightarrow 0$ corresponding to a light singlet-like CP-even Higgs scalar, (5.4) shows that the upper bound on the Higgs mass discussed in Section 3.2 applies now to the Next-to-lightest CP-even Higgs scalar H_2 , which helps to establish no-lose theorems at colliders (stating that at least one Higgs boson is detectable). In the case $\xi_1 \neq 0$, but ξ_1 small, the lightest CP-even scalar H_1 would still be very difficult to detect at the Tevatron and at the LHC, and Higgs searches will only be sensitive to H_2 (and possibly H_3) in the CP-even sector. For $\xi_1 \neq 0$, but $\xi_1^2 \leq \xi_2^2$, the upper bound on the mass M_{H_2} of the

observable H_2 is somewhat alleviated [127, 192]. Using again (5.4) and the upper bound on ξ_1^2 as a function of M_{H_1} from Fig. 3 (assuming SM decay branching ratios for H_1), one finds that the maximal possible value of M_{H_2} is assumed for $\xi_1^2 \sim \xi_2^2 \sim 1/2$, in which case Fig. 3 gives $M_{H_1} \gtrsim 110$ GeV implying $M_{H_2} \lesssim 162$ GeV.

The relevance of Higgs-to-Higgs decays in the NMSSM for Higgs searches was first mentioned in [195], and underlined in [68, 69] in the framework of scenarios with an approximate Peccei-Quinn or R-symmetry (see Section 2.2) where a CP-odd scalar A_1 can be very light allowing for $H_i \rightarrow A_1 A_1$ decays. Then the branching ratios for $H_i \rightarrow b\bar{b}$ and $H_i \rightarrow \tau^+ \tau^-$ would be suppressed, and the upper bounds on ξ^2 in Fig. 3 should be re-interpreted [52, 200–202]. In particular, a CP-even scalar H with a SM coupling to the Z boson ($\xi = 1$) but with $M_H < 114$ GeV can be compatible with the constraints from LEP [70, 71, 200–207].

On the other hand, once Higgs-to-Higgs decays as $H_i \rightarrow A_1 A_1$ (possibly also $H_i \rightarrow H_1 H_1$) are kinematically allowed, many additional Higgs search channels studied at LEP can be relevant. The constraints on the corresponding cross sections, after combining results from the ALEPH, DELPHI, L3 and OPAL collaborations, have also been worked out by the LEP Working group for Higgs Boson Searches [57], and those particularly relevant for the NMSSM have been summarised recently in [208]:

1. Searches for $e^+ e^- \rightarrow H Z$ independent of the H decay mode, looking for a peak of the M_X recoil mass distribution in $e^+ e^- \rightarrow X Z$, by OPAL [209]; these give $M_H > 82$ GeV if H couples to Z with SM strength ($\xi \sim 1$).
2. Searches for $H \rightarrow \Phi \Phi$ and $\Phi \rightarrow b\bar{b}$ ($\Phi =$ CP-even or CP-odd Higgs) by OPAL [210] and DELPHI [211]; once combined, these give $M_H > 110$ GeV for $\xi \sim 1$ [57].
3. Searches for $H \rightarrow \Phi \Phi$ and $\Phi \rightarrow gg$, $\Phi \rightarrow c\bar{c}$, $\Phi \rightarrow \tau^+ \tau^-$ by OPAL [212] exclude $45 \text{ GeV} < M_H < 86 \text{ GeV}$ for $\xi \sim 1$. A new analysis of ALEPH data [213] implies stronger limits on $H \rightarrow \Phi \Phi \rightarrow 4\tau$ as $M_H \gtrsim 109$ GeV for $M_\Phi = 10$ GeV and $\xi^2 \times BR(H \rightarrow \Phi \Phi) \times BR(\Phi \rightarrow \tau\tau)^2 \sim 1$.

Also possibly relevant could be [57] $e^+ e^- \rightarrow H Z$ with $H \rightarrow jj$ (two jets), $H \rightarrow \gamma\gamma$ and H decaying invisibly (as, e.g., into neutralinos), and $e^+ e^- \rightarrow H A$ together with $H A \rightarrow 3A \rightarrow 6b$, all of which are considered in the public code NMHDECAY [52, 88], which is part of NMSSMTools (see Appendix D).

In Fig. 3 one can note a slight excess of events for $M_H \sim 95 - 100$ GeV of 2.3σ statistical significance [214]. In the NMSSM this excess could be explained by

- a SM-like CP-even scalar of corresponding mass and $\xi \sim 1$, but with reduced $BR(H \rightarrow b\bar{b}) \sim 0.1$ [70, 202, 205] due to a dominant decay $H \rightarrow A A$ with a branching ratio ~ 0.8 . This scenario could alleviate the little fine tuning problem related to the non-observation of a CP-even scalar at LEP (see Section 7.4), but requires $M_A < 10.5$ GeV due to the constraints (2) on $H \rightarrow \Phi \Phi$ and $\Phi \rightarrow b\bar{b}$;
- a CP-even scalar H_1 of corresponding mass with $\xi \lesssim 0.4$ [215] as it can appear, e.g., in the constrained NMSSM (see Section 7.1) [216, 217] or for large $\lambda \approx 0.7 - 0.8$ [218]. Then, another CP-even scalar H_2 with $\xi \gtrsim 0.9$ and a mass not far above 115 GeV is present in the spectrum.

In the first scenario, difficulties for Higgs searches at the Tevatron and at the LHC are foreseeable (see below), whereas probably only H_2 would be detectable at hadron colliders within the second scenario.

If H_1 is singlet-like ($\xi_1^2 \leq \xi_2^2$) and decays unconventionally, the upper bound on M_{H_2} deduced above ($M_{H_2} \lesssim 162$ GeV from $M_{H_1} \gtrsim 110$ GeV) is no longer valid. Using the upper bound on M_{H_1} from constraints on $H_1 \rightarrow A_1 A_1 \rightarrow 4b$, (5.4) gives $M_{H_2} \lesssim 167$ GeV, whereas the weaker upper bound on M_{H_1} independent of its decay mode (using only the Z recoil mass) allows for $M_{H_2} \lesssim 192$ GeV.

The prospects for NMSSM Higgs searches at $e^+ e^-$ colliders of higher energy, as an ILC, have been investigated in [177, 192, 194, 219–222]; for a review see [223]. Due to the sum rule (5.2), one can derive a theoretical lower limit on the production cross section σ_i for at least one CP-even Higgs boson, obtained as $\sigma_i > 0.04$ pb for $\sqrt{s} = 300$ GeV in [192]. (Production cross sections at $\sqrt{s} = 500, 1000$ and 2000 GeV were computed in [194].)

As at LEP, one can search for Higgs bosons at an ILC independently from their decay modes using the recoil spectrum (the invariant mass M_X) in $e^+ e^- \rightarrow Z X$, $Z \rightarrow e^+ e^-$ and $Z \rightarrow \mu^+ \mu^-$. This allows to establish a no-lose theorem for arbitrary singlet extensions of the MSSM (assuming perturbativity of all Yukawa couplings up to 10^{16} GeV, implying upper bounds on the Higgs masses) for $\sqrt{s} = 500$ GeV and an integrated luminosity $\mathcal{L} = 100$ fb $^{-1}$ [219]. Also the ZZH couplings can be determined by this method [223]. The clean environment of $e^+ e^-$ colliders would allow for the study of NMSSM specific Higgs-to-Higgs decays [220], and for the discovery of light Higgs states that would be difficult to observe at the LHC [221, 222].

5.1.2 Tevatron and LHC

Next we turn to hadron colliders, starting with regions in the NMSSM parameter space where Higgs-to-Higgs decays are kinematically forbidden or occur only with small branching ratios. Then, the search for Higgs bosons in the NMSSM can proceed as in the MSSM (see [224] and references therein); of course, Higgs boson production cross sections and decay branching fractions in the NMSSM have to be rescaled (for tree level couplings) or re-evaluated (for loop corrected or loop induced couplings), employing the tree level couplings given in Appendix A.2. Important search channels are

1. $gg \rightarrow H$ (gluon-gluon fusion) with $H \rightarrow WW^{(*)} \rightarrow l^+ l^- \nu \bar{\nu}$, $H \rightarrow ZZ^{(*)} \rightarrow 4$ leptons, $H \rightarrow \gamma\gamma$ (the latter also for A instead of H);
2. Vector boson fusion corresponding to $q\bar{q} \rightarrow q\bar{q}VV$ with $VV \rightarrow H$ at the Tevatron, and $qq \rightarrow qqVV$ with $VV \rightarrow H$ at the LHC (where $V = W^\pm, Z$), and $H \rightarrow \gamma\gamma$, $H \rightarrow \tau^+ \tau^-$, $H \rightarrow WW^{(*)}$, H decaying invisibly or $H \rightarrow b\bar{b}$ (the latter decay suffers from a too large background at the LHC);
3. $q\bar{q} \rightarrow W^* \rightarrow WH$ and $q\bar{q} \rightarrow Z^* \rightarrow ZH$ (Higgs-strahlung);
4. Associated production with heavy quark pairs $q\bar{q}/gg \rightarrow Q\bar{Q}H$, where $Q = t, b$.

In some regions of the parameter space of the NMSSM, more Higgs bosons than in the MSSM could be observed (e. g. in the two photon channel [225]), which would allow

to distinguish the two models. On the other hand, since all ξ_i can be simultaneously less than 1 (respecting the sum rule (5.2)), it was not clear at first instance whether any of the H_i could always be detected after combining the results for Higgs searches at LEP2 with the prospects for the LHC [195, 198], even if Higgs-to-Higgs decays are assumed to be kinematically forbidden. Later on, the analysis of Higgs searches at the LHC was refined and additional Higgs production channels (notably vector boson fusion) have been considered. On the theoretical side, radiative corrections to the Higgs masses have been improved by dominant two-loop corrections; these lower the CP-even Higgs masses, increasing the region in the NMSSM parameter space excluded by LEP2. As a result, a no-lose theorem for the detection of at least one CP-even Higgs scalar in the NMSSM at the LHC with an integrated luminosity of 600 fb^{-1} (two detectors with 300 fb^{-1} each) could be established [201, 220, 226], however, under the assumption that Higgs-to-Higgs decays are kinematically forbidden.

Once Higgs-to-Higgs decays are possible (or even dominant), Higgs searches at the Tevatron and the LHC can become considerably more complicated; see [66, 201, 203, 208, 227–229] for discussions of possible scenarios and proposals for search channels. In principle, CP-even scalars H_2 can decay into a pair of CP-even H_1 in LEP-allowed regions of the NMSSM parameter space [201, 217, 228] (see [129] for very large $\lambda \approx 2$ where $M_{H_1} \approx 200 - 300 \text{ GeV}$, $M_{H_2} \approx 350 - 700 \text{ GeV}$); but most of the recent studies concentrated on decays into light CP-odd scalars A motivated by an approximate Peccei-Quinn or R-symmetry, and/or the $H \rightarrow AA$ explanation of the light excess of events at LEP.

For $M_A > 2m_b$ (in which case LEP constraints require $M_H \gtrsim 110 \text{ GeV}$ for SM-like couplings of H), the cascade $H \rightarrow AA$ would end mostly in a $4b$ final state. This final state has a very large background at hadron colliders (which seemed to make it invisible at least at the Tevatron, unless the H production rate is enhanced relative to the SM [230]). Hence, a first study proposed to consider the subdominant $H \rightarrow AA \rightarrow b\bar{b}\tau^+\tau^-$ final state, with H produced via vector boson fusion at the LHC [220, 231]. However, here the visibility of the signal depends strongly on the poorly known background from $t\bar{t}$ production. Subsequently the Higgs-strahlung process has been added, where the lepton(s) from W, Z decays can help to trigger on the relevant events [232–234]. Then, both $4b$ and $2b2\tau$ final states can be relevant for both the Tevatron and the LHC [234] (the $2b2\tau$ final state still being more promising at the LHC).

For $M_A < 2m_b$, the cascade $H \rightarrow AA$ would end mostly in a 4τ final state; clearly, the visible τ -decay products would not generate narrow peaks in the invariant masses of M_A or M_H . At the Tevatron, the prospects to detect the Higgs scalar in this case look pretty dim [204] unless the H production cross section is enhanced; the 4γ decay mode would require enhanced branching ratios for $A \rightarrow \gamma\gamma$ [235]. At the LHC particular efforts would also be required. Proposals for signals and cuts appropriate for the $AA \rightarrow 4\tau \rightarrow 2\mu + 2\text{jets}$ final state have been made in [236]; with 100 fb^{-1} of integrated luminosity, the expected rates after cuts are $\sim 8 \times 10^3$ from H production via vector boson fusion, and $\sim 10^3$ from H production via Higgs-strahlung ($W^\pm \rightarrow H + W^\pm$) where one can trigger on a lepton from W^\pm decays. In [237] it has been proposed to consider diffractive Higgs production ($pp \rightarrow pp + H$) in order to be sensitive to $H \rightarrow 4\tau$, which requires to install additional forward detectors. Using a track-based analysis in which all events with more than 6 tracks in the central region are discarded, a viable signal seems possible after accumulating

300 fb⁻¹ of integrated luminosity. In [238, 239], the subdominant $H \rightarrow AA \rightarrow 2\tau 2\mu$ final state was discussed: in spite of the small branching fraction (with 2μ from direct A decays) it was argued that, for $M_H \sim 102$ GeV, the Tevatron can see a signal over background for an integrated luminosity $\mathcal{L} \sim 10$ fb⁻¹, and the LHC already for $\mathcal{L} \sim 1$ fb⁻¹, with H being produced via gluon-gluon fusion.

Light A production in association with charginos [240] (which requires $\lambda \gtrsim 1$ to be observable), and from neutralino decays [241] has also been considered. The LHC discovery potential for $\lambda = 2$ – where M_{H_1} can be as large as 200 to 300 GeV – has been analysed in [129]. Further details of current ATLAS and CMS studies of benchmark scenarios including the $H \rightarrow AA \rightarrow 4\tau$ final state can be found in [228].

The charged Higgs decays $H^\pm \rightarrow W^\pm + H$ or $H^\pm \rightarrow W^\pm + A$ play only a marginal rôle in the MSSM. In the NMSSM, larger regions in the parameter space exist (at low $\tan\beta$) where $M_{H^\pm} \lesssim m_t$, but where these processes are kinematically allowed and the corresponding branching fractions are important [242]. In the case of a light A with a non-negligible $SU(2)$ doublet component, the second process (and also $pp \rightarrow H^\pm + A \rightarrow W^\pm + AA$) can be observable [243].

Higgs searches at the Tevatron have started to test regions of the NMSSM parameter space. Recall that, in the NMSSM, a CP-even Higgs H_2 with nearly SM-like couplings can have a mass up to 160 GeV (if a light singlet-like H_1 exists, escaping LEP constraints), or even up to ~ 190 GeV, if the singlet-like H_1 decays unconventionally (see above). The present results of searches for a SM Higgs at the Tevatron (CDF and DØ) from various production and decay channels have recently been combined [244], and exclude a Higgs scalar with SM couplings in the mass range $160 < m_H < 170$ GeV. While such a Higgs scalar would be impossible in the MSSM, the corresponding mass range touches the parameter space of the NMSSM.

In [245] the DØ collaboration has looked for CP-even or CP-odd scalars Φ produced in association with b quarks and decaying into $\tau^+\tau^-$, assuming a branching ratio of a SM-like Higgs scalar. In the mass range $90 \text{ GeV} < M_\Phi < 150 \text{ GeV}$, very large values of $\tan\beta \gtrsim 40 - 80$ (which would imply an enhanced production cross section) could be excluded. This result is relevant for both the MSSM and the NMSSM, although one usually expects $\tan\beta \lesssim 60$.

The DØ collaboration has also searched for $H \rightarrow A_1 A_1$ decays in the range $M_{A_1} < 2m_\tau$ where the branching ratio $A_1 \rightarrow \mu^+\mu^-$ is not too small. The 4μ signature has been searched, and an upper limit of about 10 fb on $\sigma(p\bar{p} \rightarrow H X) \times BR(H \rightarrow AA) \times (BR(A \rightarrow \mu^+\mu^-))^2$ has been set [246]. Assuming $M_H = 120$ GeV and $BR(H \rightarrow AA) = 100\%$, the negative result implies an upper bound on the $BR(A \rightarrow \mu^+\mu^-) \lesssim 10\%$. The final state $\mu^+\mu^-\tau^+\tau^-$ (relevant for $M_{A_1} > 2m_\tau$) has also been investigated, but the limits are still a factor ~ 4 above the theoretical expectations. Further results from the Tevatron can be expected in the near future.

5.2 The neutralino sector

The neutralino mass matrix in the general NMSSM is given in (2.32) as function of the vevs v_u , v_d and s . In the \mathbb{Z}_3 -invariant NMSSM, the last term $2\mu'$ in the (5, 5) element is

absent. In [58,247–249] approximate formulae for the mass eigenvalues and mixing matrices have been derived. From $M_{\chi_1^0}^2 \leq M_1^2 + M_Z^2 \sin^2 \theta_W$, and assuming the relations (3.1) for the gaugino masses, an upper bound on the lightest neutralino mass as function of the gluino mass M_3 can be deduced [250]. In the nMSSM (where $\kappa = 0$, see Section 7.3), the bound $M_{\chi_1^0} \leq \lambda v$ (with v as in (2.13)) holds [251].

The full set of neutralino couplings in the \mathbb{Z}_3 -invariant NMSSM, necessary to study neutralino production processes at $e^+ e^-$ colliders, in sfermion (squark/slepton) decays, chargino/neutralino cascade decays and Higgs decays, is given in [50].

Clearly, in the decoupling limit $\lambda \sim \kappa \rightarrow 0$, $s \sim 1/\kappa \rightarrow \infty$ of the \mathbb{Z}_3 -invariant NMSSM (see Section 2.2.1), it becomes practically impossible to distinguish the neutralino sector of the NMSSM from the one of the MSSM: in this limit, the singlino-like neutralino χ_S^0 ceases to mix with the MSSM-like neutralinos, and all production cross sections as well as partial widths of decays into χ_S^0 tend to zero. An important exception to this rule is the scenario where χ_S^0 is the LSP (for κs small enough), and R-symmetry is conserved: then all sparticle decay chains will first proceed as in the MSSM into the NLSP (which can now be charged!), and only at the end of the decay chain the NLSP finally decays into χ_S^0 . This additional decay (leading typically to extra leptons in the final state) should allow to distinguish the NMSSM from the MSSM in the singlino-LSP regime. Due to a possibly tiny value of the NLSP– χ_S^0 coupling, the NLSP lifetime can be quite long leading to displaced vertices at colliders, see below. Subsequently we briefly discuss neutralino production processes at $e^+ e^-$ colliders, in sfermion and cascade decays.

5.2.1 Neutralino pair production at $e^+ e^-$ colliders

Neutralino pair production at $e^+ e^-$ colliders can proceed via a Z boson in the s channel and via the exchange of a selectron in the t/u channels. Formulae for the corresponding cross sections are given in [252] where, however, the “photino/zino”-basis in the gaugino sector is employed, differing from the one used here. In [249], the formulae for the cross sections are given in the “bino-wino”-basis employed here.

Present constraints on the neutralino sector originate from its contribution to the invisible Z width and from bounds on $e^+ e^- \rightarrow \chi_i^0 \chi_j^0$ ($i \neq j$) from LEP. Note that bounds on the mass of the lightest neutralino χ_1^0 relying on MSSM-like relations between neutralino and chargino masses cannot be applied to the NMSSM. For earlier discussions of LEP constraints on the neutralino sector of the NMSSM see [253, 254].

Assuming 3 massless neutrinos, additional contributions to the invisible Z width Γ_Z^{inv} should not exceed ~ 2 MeV (see the Section on $\tilde{\nu}$ mass limits in [89]). This limit has to be compared to the partial Z width into neutralinos (if $m_{\chi_i^0} < M_Z/2$), given by

$$\Gamma_{Z \rightarrow \chi_i^0 \chi_i^0} = \frac{M_Z^3 G_F}{12\sqrt{2}\pi} (N_{i3}^2 - N_{i4}^2)^2 \left(1 - \frac{4m_{\chi_i^0}^2}{M_Z^2}\right)^{3/2}, \quad (5.6)$$

where the mixing matrix elements N_{ij} are defined in (A.7). Clearly, the mixing matrix elements N_{i3} and N_{i4} (proportional to the higgsino components of the neutralinos) are tiny for singlino and/or gaugino like neutralinos, which are thus not much constrained.

Signals from $e^+e^- \rightarrow \chi_i^0 \chi_j^0$ with subsequent χ_i^0 decays ($i > 1$) have been searched for by DELPHI [255] and OPAL [256] at LEP, with \sqrt{s} up to 209 GeV. The non-observation of such signals imposes upper bounds on the production cross sections $\sigma(e^+e^- \rightarrow \chi_1^0 \chi_i^0) \lesssim 10^{-2}$ pb and $\sigma(e^+e^- \rightarrow \chi_i^0 \chi_j^0) \lesssim 10^{-1}$ pb ($i, j > 1$).

Phenomenological analyses of pair production of neutralinos in the NMSSM at e^+e^- colliders at higher energies have been performed in [249, 252, 257–260]. Since the information on the neutralino sector from the LHC will be quite limited, an ILC-like e^+e^- collider can be crucial to distinguish the NMSSM neutralino sector from the one of the MSSM [259], although it cannot be guaranteed that the difference is visible if one is close to the decoupling limit mentioned above. This question has also been addressed in the radiative production of the lightest neutralino pair, $e^+e^- \rightarrow \chi_1^0 \chi_1^0 \gamma$, at an ILC with $\sqrt{s} = 500$ GeV in [260].

5.2.2 Decays into and of neutralinos

As in the MSSM, in the NMSSM sfermions \tilde{f}_i (squarks and sleptons) can decay into neutralinos or charginos via $\tilde{f}_i \rightarrow f_i \chi_j^0$ [249] or $\tilde{f}_i \rightarrow f_j \chi^\pm$, where the latter decay does not exhibit any NMSSM specific features. However, due to the modified neutralino sector, the processes $\tilde{f}_i \rightarrow f_i \chi_j^0$ can be quite different in the NMSSM: even a mostly singlino-like neutralino can be produced in decays of squarks and sleptons of the third generation, if a non-negligible higgsino component induced by a not too small value of λ [261] is present. In addition, gaugino components of the neutralinos also allow for their production in sfermion decays of all generations [249].

Higgs bosons could also decay invisibly into neutralinos; these processes can still be interesting for Higgs searches via the recoil spectrum of Z decay products in $Z^* \rightarrow ZH$, see above. An exception are GMSB models (see Section 7.2), where the neutralinos can subsequently decay into a gravitino and a photon; the present constraints on and the future prospects of this Higgs decay mode have been studied in [262].

Once produced, neutralinos can decay in many different ways (except for the LSP):

- into a fermion + sfermion (if kinematically allowed);
- via two-body decays (if kinematically allowed) $\chi_i^0 \rightarrow \chi_j^0 Z$, $\chi_i^0 \rightarrow \chi^\pm W^\mp$, $\chi_i^0 \rightarrow \chi_j^0 \Phi$, where Φ is a CP-even or CP-odd Higgs, and via the radiative decay $\chi_i^0 \rightarrow \chi_j^0 \gamma$;
- via three body decays $\chi_i^0 \rightarrow \chi_j^0 l^+ l^-$, $\chi_i^0 \rightarrow \chi_j^0 q \bar{q}$ and $\chi_i^0 \rightarrow \chi_j^0 \nu \bar{\nu}$ (via Z , Higgs and sfermion exchange).

Formulae for the relevant partial widths can be found in [58, 249, 252]. Note that the decay $\chi_i^0 \rightarrow \chi_j^0 A_1$ can be relevant for the search for a light pseudoscalar Higgs [241]. If the $m_{\chi_i^0} - m_{\chi_j^0}$ mass difference is small (e. g. in the case $\chi_i^0 \equiv \chi_2^0 \sim$ bino, $\chi_j^0 \equiv \chi_1^0 \sim$ singlino), soft leptons from $\chi_2^0 \rightarrow \chi_1^0 l^+ l^-$ can be an important signal for the NMSSM, but their detection at the LHC may require low cuts on lepton transverse momenta [263].

5.2.3 Displaced vertices

As stated above, for small λ the couplings between the NLSP and a singlino-like LSP in the NMSSM can be so small that the large NLSP lifetime leads to macroscopically displaced

vertices. This phenomenon has been studied first in the case of a bino NLSP in the scale invariant NMSSM in [58], and applied to searches at LEP in [264]. Such additional cascades in sparticle decays (as compared to the MSSM), with possibly displaced vertices, can also occur in more general extensions of the MSSM by singlets [265–267] as, for instance, in models designed to explain the CDF multi-muon events [268] where the cascades originate from Higgs decays.

Considering the case of a bino-like NLSP χ_2^0 (with a mass $\sim M_1$) and a singlino-like LSP χ_1^0 , approximations for the relevant neutralino mixing parameters can be derived for small λ , and an approximate expression for the (typically dominant) bino to singlino decay width via right-handed slepton (\tilde{E}_R) exchange can be obtained [58]:

$$\Gamma(\chi_2^0 \xrightarrow{\tilde{E}_R} \chi_1^0 l^+ l^-) \simeq 2 \times 10^{-6} \lambda^2 M_1 \left(\frac{M_Z}{m_{\tilde{E}_R}} \right)^4 I(\eta, \omega) \quad (5.7)$$

where $\eta = m_{\chi_1^0}/m_{\chi_2^0}$, $\omega = m_{\chi_2^0}/m_{\tilde{E}_R}$ and the phase space integral $I(\eta, \omega)$ is of $\mathcal{O}(10^{-1})$ for masses compatible with universal soft terms at the GUT scale [58]. The important point is that the bino decay width is proportional to λ^2 , implying a long lifetime for $\lambda \rightarrow 0$.

In the fully constrained NMSSM with its nearly unique sparticle spectrum (once a dark matter relic density complying with WMAP constraints has been imposed, see Section 7.1), the LSP is always singlino-like [216, 217], and the NLSP is the lighter stau. The stau mass must only be a few GeV above the LSP mass in order to allow for a sufficient reduction of the relic density via coannihilation, see section 9. Then, again using approximations for the relevant neutralino mixing parameters as in [58], the expression for the stau decay width is approximately given as [217]

$$\Gamma(\tilde{\tau}_1 \rightarrow \chi_1^0 \tau) \approx \lambda^2 \frac{\sqrt{\Delta m^2 - m_\tau^2}}{4\pi m_{\tilde{\tau}_1}} (\alpha \Delta m - \beta m_\tau), \quad (5.8)$$

where the coefficients α and β still depend on $m_{\tilde{\tau}_1}$ and $m_{\chi_1^0}$ (of the order $0.01 \gtrsim \alpha \sim \beta \gtrsim 0.0001$, decreasing with $m_{\tilde{\tau}_1}$), and $\Delta m \equiv m_{\tilde{\tau}_1} - m_{\chi_1^0}$. The stau decay width can be small due to a small value of λ and a possible phase space suppression for small $\Delta m - m_\tau$. For $\Delta m \sim$ a few GeV and $\lambda \sim 10^{-4}$, stau lengths of flight of $\mathcal{O}(\text{mm})$ are possible [217].

In the near future it will be important to study the possible impact of NMSSM specific displaced vertices for sparticle searches at hadron colliders.

6 b physics and the anomalous magnetic moment of the muon

In this Section we briefly discuss NMSSM specific effects in B physics, Υ physics and for the anomalous magnetic moment of the muon. In most cases an extensive literature exists on contributions from MSSM-like supersymmetric extensions of the Standard Model, which are beyond the scope of the present review. In general, the contributions in the NMSSM can differ from those of the MSSM due to the different Higgs and neutralino sectors. Correspondingly, most of the results within the MSSM – like constraints on the parameters or a possible improved agreement between theory and experiment – are valid in the NMSSM as well.

A notable exception is the region in the NMSSM parameter space corresponding to a light CP-odd Higgs scalar A_1 . Here, important new contributions “beyond the MSSM” can arise, which lead to constraints on this part of parameter space and/or to possible new phenomena (in the case of Υ/η_b physics).

6.1 B physics

The following B physics processes have been studied in the NMSSM: $BR(\bar{B} \rightarrow X_s \gamma)$ in [269, 270], $BR(\bar{B}_s \rightarrow \mu^+ \mu^-)$ and $\Delta M_{s(d)}$ in [269–271], $\bar{B}^+ \rightarrow \tau^+ \nu_\tau$ in [270], $BR(\bar{B} \rightarrow X_s l^+ l^-)$ in [269, 272], and $BR(\bar{B} \rightarrow \gamma l^+ l^-)$ in [272].

In most cases, the NMSSM specific contributions to the B physics observables originate essentially from a possibly light CP-odd Higgs scalar A_1 , and the fact that a flavour violating vertex b - q - A_1 (with $q = s, d$) is generated, amongst others, by squark-chargino loops (see, e.g., [269, 273]). Here and below, minimal flavour violation is assumed, i.e. the only flavour violation originates from the Yukawa sector and is parametrized by the CKM matrix. For large $\tan \beta$ and $|A_t|$, the loop-generated flavour violating vertex b - q - A_1 is roughly proportional to $\tan^2 \beta |A_t|$; as a consequence, the NMSSM specific contributions induced by a light CP-odd Higgs scalar A_1 increase strongly with $\tan \beta$ (similar to most of the other MSSM-like contributions). Subsequently we briefly discuss the various B physics processes.

$BR(\bar{B} \rightarrow X_s \gamma)$

The branching ratio $BR(\bar{B} \rightarrow X_s \gamma)$ is one of the most intensively studied quantity in B physics. In the past, constraints from $b \rightarrow s \gamma$ have been particularly severe, since the experimental world average for $BR(\bar{B} \rightarrow X_s \gamma)$ was somewhat below the Next-to-leading order (NLO) SM prediction, while the beyond the SM (BSM) contribution involving a charged Higgs boson in the relevant diagram is positive.

This situation has changed during the last years: the present world average estimated by the Heavy Flavour Averaging Group [274] reads (for a lower cut on the photon energy $E_\gamma > E_0 = 1.6$ GeV)

$$BR(\bar{B} \rightarrow X_s \gamma)|_{\text{exp}} = (3.52 \pm 0.23 \pm 0.09) \times 10^{-4}. \quad (6.1)$$

The SM Next-to-Next-to-Leading Order $\mathcal{O}(\alpha_s^2)$ corrections to the total $BR(\bar{B} \rightarrow X_s \gamma)$ branching fraction have recently been combined [275, 276], giving

$$BR(\bar{B} \rightarrow X_s \gamma)|_{\text{SM}} = (3.15 \pm 0.23) \times 10^{-4}. \quad (6.2)$$

In [277] the cut $E_\gamma > 1.6$ GeV on the photon energy has been treated differently, leading to an even lower SM prediction:

$$BR(\bar{B} \rightarrow X_s \gamma)|_{\text{SM}} = (2.98 \pm 0.26) \times 10^{-4}. \quad (6.3)$$

This result can be interpreted as a (weak) hint for positive BSM contributions to $b \rightarrow s \gamma$; in any case constraints on the parameter space of supersymmetric models have become less stringent.

In supersymmetric extensions of the SM, positive contributions to $BR(\bar{B} \rightarrow X_s \gamma)$ arise from loops with charged Higgs bosons. Additional contributions involving stop quarks and charginos are roughly proportional to $\tan \beta$ and to the trilinear coupling A_t , and can have either sign. For an analysis of the corresponding constraints on the parameter space of the MSSM see, e.g., [278]; in the meantime, the full two-loop SUSY QCD corrections of the MSSM contributions have been computed in [279].

NMSSM specific contributions to $BR(\bar{B} \rightarrow X_s \gamma)$ from the extended Higgs and neutralino sectors arise only at the two-loop level. These were considered in [269] and, in more detail, in [270] with the result that the effects are negligibly small, at least for a fixed charged Higgs mass (which can be somewhat lower in the NMSSM due to the negative term $\sim \lambda^2$ in (2.29)). Even contributions from a light CP-odd Higgs A_1 are not very important here, since A_1 appears only in loops and not in a possibly resonant s -channel.

Also in the case of $\bar{B}^+ \rightarrow \tau^+ \nu_\tau$ the NMSSM specific effects are merely indirect: this process is dominated by W^+ exchange at tree level in the SM; in the MSSM and NMSSM, additional contributions come from charged Higgs exchange, which can be somewhat lighter in the NMSSM.

$$BR(\bar{B}_s \rightarrow \mu^+ \mu^-)$$

The most up-to-date experimental result on this process is an upper bound (at 95% confidence level) from CDF [280]:

$$BR(\bar{B}_s \rightarrow \mu^+ \mu^-)|_{\text{exp}} < 5.8 \times 10^{-8}. \quad (6.4)$$

This is still one order of magnitude beyond the SM estimate [281]

$$BR(\bar{B}_s \rightarrow \mu^+ \mu^-)|_{\text{SM}} = (3.8 \pm 0.1) \times 10^{-9}, \quad (6.5)$$

so that there is some room for potentially large new physics contributions.

The SM contributions originate from box and gauge-penguin diagrams, which are small with respect to the experimental bound. The corresponding MSSM contributions can be found in [273] and have been generalized to the NMSSM in [269–271]. Here, the flavour changing b - s - A_1 vertex $\sim \tan^2 \beta$ can lead to significant contributions arising from penguin diagrams, notably if the CP-odd scalar A_1 in the s -channel (decaying into $\mu^+ \mu^-$) is close

to its mass shell, i.e. if its mass is close to $M_{\bar{B}_s}$. (Since the $A_1\text{-}\mu^+\text{-}\mu^-$ vertex is proportional to $\tan\beta$, the A_1 contribution to $BR(\bar{B}_s \rightarrow \mu^+\mu^-)$ is proportional to $\tan^6\beta$.) In fact, CP-odd Higgs masses equal to $M_{\bar{B}_s}$ are always excluded by the experimental bound on this process, but the width of the forbidden region depends on $\tan\beta$ and $|A_t|$. An example for a forbidden region in the $(M_{A_1}, \tan\beta)$ -plane is shown in Fig. 4 below (taken from [270]); for a study at very large $\tan\beta \sim 50$, see [271].

$\Delta M_{s(d)}$

Information on the mass differences $\Delta M_{s,d} \equiv m_{\bar{B}_{s,d}} - m_{B_{s,d}}$ originates from measurements of B meson oscillations. The present result for ΔM_s obtained by the CDF collaboration [282] is

$$\Delta M_s^{\text{exp}} = 17.77 \pm 0.12 \text{ ps}^{-1}. \quad (6.6)$$

The SM prediction depends on the hadronic factor f_{B_s} and CKM matrix elements; for a comparison with BSM contributions, the CKM matrix elements should be determined by tree level measurements such that their values are not ‘‘polluted’’ by BSM effects [283]. Using $|V_{ts}^*V_{tb}| = (41.3 \pm 0.7) \times 10^{-3}$ and $f_{B_s}\sqrt{\hat{B}_{B_s}} = 0.281 \pm 0.021 \text{ GeV}$ given by the HPQCD collaboration [284], one obtains [270]

$$\Delta M_s^{\text{SM}} = 20.5 \pm 3.1 \text{ ps}^{-1}. \quad (6.7)$$

Hence, despite the large uncertainty, a negative contribution from new physics seems favoured.

The measurement of ΔM_d is also quite precise [285]:

$$\Delta M_d^{\text{exp}} = 0.507 \pm 0.004 \text{ ps}^{-1}. \quad (6.8)$$

However, again the SM prediction suffers from large uncertainties:

$$\Delta M_d^{\text{SM}} = 0.59 \pm 0.19 \text{ ps}^{-1}, \quad (6.9)$$

obtained with a tree level determination of $|V_{td}^*V_{tb}| = (8.6 \pm 1.4) \times 10^{-3}$ [283] and a hadronic factor $f_{B_d}\sqrt{\hat{B}_{B_d}}$ calculated from $f_{B_s}\sqrt{\hat{B}_{B_s}} / \left(f_{B_d}\sqrt{\hat{B}_{B_d}} \right) = 1.216 \pm 0.041$ [286].

Supersymmetric contributions arise from box diagrams at the one-loop level, but also from double-penguin diagrams involving two flavour changing vertices like $b\text{-}s(d)\text{-}A_1$ [269–271], which are particularly important if A_1 is exchanged in the s -channel. Now the A_1 -contribution is proportional to $\tan^4\beta$, and leads again to an exclusion of A_1 masses near $M_B \sim 5 \text{ GeV}$, as shown in Fig. 4. In the MSSM, a relation between $\Delta M_{s(d)}$ and $BR(\bar{B}_s \rightarrow \mu^+\mu^-)$ at large $\tan\beta$ can be deduced [287]; as pointed out in [269], this relation is spoiled by the A_1 -exchange diagrams in the s -channel in the NMSSM.

$BR(B \rightarrow X_s l^+ l^-)$

This branching ratio (for $l = e$ or μ) has been measured by Babar [288] and Belle [289]. In order to avoid contributions from c -quark resonances, the regions $1 \text{ GeV}^2 < M_{l^+l^-}^2 <$

6 GeV² (low) and 14.4 GeV² < M_{l+l-}² < 25 GeV² (high) are considered separately:

$$BR(\bar{B} \rightarrow X_s l^+ l^-)_{\text{low}} = \begin{cases} (1.493 \pm 0.504_{-0.321}^{+0.411}) \times 10^{-6} & \text{Belle} \\ (1.8 \pm 0.7 \pm 0.5) \times 10^{-6} & \text{BaBar} \\ (1.60 \pm 0.50) \times 10^{-6} & \text{average} \end{cases} \quad (6.10)$$

$$BR(\bar{B} \rightarrow X_s l^+ l^-)_{\text{high}} = \begin{cases} (0.418 \pm 0.117_{-0.068}^{+0.061}) \times 10^{-6} & \text{Belle} \\ (0.5 \pm 0.25_{-0.07}^{+0.08}) \times 10^{-6} & \text{BaBar} \\ (0.44 \pm 0.12) \times 10^{-6} & \text{average} \end{cases} \quad (6.11)$$

The SM analysis has become quite refined (for a review, see [290]):

$$\begin{cases} BR(\bar{B} \rightarrow X_s \mu^+ \mu^-)_{\text{low}}^{\text{SM}} & = (1.59 \pm 0.11) \times 10^{-6} \\ BR(\bar{B} \rightarrow X_s \mu^+ \mu^-)_{\text{high}}^{\text{SM}} & = 2.40 \times 10^{-7} (1_{-0.26}^{+0.29}) \end{cases} \quad (6.12)$$

$$\begin{cases} BR(\bar{B} \rightarrow X_s e^+ e^-)_{\text{low}}^{\text{SM}} & = (1.64 \pm 0.11) \times 10^{-6} \\ BR(\bar{B} \rightarrow X_s e^+ e^-)_{\text{high}}^{\text{SM}} & = 2.09 \times 10^{-7} (1_{-0.30}^{+0.32}) \end{cases} \quad (6.13)$$

These values are well within 2σ of the experimental measurements.

In the NMSSM, the process $B \rightarrow X_s l^+ l^-$ is also sensitive to a light CP-odd scalar [269, 272]: as in the case of the $BR(\bar{B}_s \rightarrow \mu^+ \mu^-)$, the flavour changing b - s - A_1 vertex proportional to $\tan^2 \beta$ (with A_1 decaying into the $l^+ l^-$ pair) can lead to significant contributions, and again exclude regions of light A_1 masses as shown in Fig. 4. The contributions to the total branching ratio ($BR(\bar{B} \rightarrow X_s \mu^+ \mu^-)^{\text{exp}} = (4.3 \pm 1.2) \times 10^{-6}$ according to [291]) were studied in [272], but only the dependency on $\tan \beta$ was shown explicitly. Other processes such as $BR(\bar{B} \rightarrow X_s \tau^+ \tau^-)$ and $BR(\bar{B}_s \rightarrow l^+ l^- \gamma)$ are interesting for Higgs phenomenology as well [272], but here experimental data is not yet available.

Combined constraints from B physics

The combined constraints from $BR(\bar{B}_s \rightarrow \mu^+ \mu^-)$, $\Delta M_{d,s}$ and $BR(\bar{B} \rightarrow X_s \gamma)$ on the mass of a light CP-odd scalar in the NMSSM have been studied in [270], focusing on the dependency on parameters like $\tan \beta$, M_{A_1} and A_t . (In addition, LEP constraints on the Higgs sector were taken into account.) In Fig. 4 we show the excluded domains for the mass of a light CP-odd Higgs scalar as a function of $\tan \beta$ for $A_t = -2500$ GeV where, in addition, constraints from $BR(\bar{B} \rightarrow X_s \mu^+ \mu^-)$ are indicated.

As mentioned before, constraints from $BR(\bar{B}_s \rightarrow \mu^+ \mu^-)$ and $\Delta M_{d,s}$ exclude domains for a mass of a light CP-odd Higgs scalar around $M_B \sim 5$ GeV for all $\tan \beta$. Apart from the visible strong increase of the excluded region with $\tan \beta$, the excluded region decreases for smaller values of $|A_t|$ due to the smaller flavour violating A_1 -quark couplings induced by stop-chargino loops.

Constraints from $BR(\bar{B} \rightarrow X_s \mu^+ \mu^-)$ from $1 \text{ GeV} < M_{\mu^+ \mu^-} < \sqrt{6} \text{ GeV}$ and $\sqrt{14.4} \text{ GeV} < M_{\mu^+ \mu^-} < m_b$ exclude always masses of a light CP-odd Higgs scalar inside these domains (where A_1 would be on-shell). For larger $\tan \beta$, larger regions of M_{A_1} are excluded by this observable, although constraints from $\bar{B}_s \rightarrow \mu^+ \mu^-$ are typically more significant.




Figure 4: Excluded regions in the $\tan\beta$ - M_{A_1} plane for $A_t = -2500$ GeV: the gridded region is excluded by LEP, the green (vertically shaded) region is excluded by $\Delta M_{d,s}$, the brown (diagonally shaded) region by $BR(\bar{B}_s \rightarrow \mu^+\mu^-)$, the region inside the dashed lines by $BR(\bar{B} \rightarrow X_s\mu^+\mu^-)$ with $1 \text{ GeV} < M_{\mu^+\mu^-} < \sqrt{6} \text{ GeV}$ or $\sqrt{14.4} \text{ GeV} < M_{\mu^+\mu^-} < m_b$, and the lower yellow corner at $\tan\beta \gtrsim 12$ and small M_{A_1} by $BR(\bar{B} \rightarrow X_s\gamma)$ (from [270]).

6.2 Υ and η_b physics

Radiative Υ decays

Light CP-odd scalars could also be produced in radiative Υ decays $\Upsilon(nS) \rightarrow \gamma A_1$ with $A_1 \rightarrow l^+l^-$, $l = \mu, \tau$ [269, 292–299]. The branching ratio depends essentially, apart from M_{A_1} , on the coupling of A_1 to b -quarks. It is useful to introduce a reduced coupling X_d , which denotes the coupling of A_1 to b -quarks relative to the corresponding coupling of the SM Higgs boson. Using the decomposition of the mass eigenstate A_1 according to (A.2) in Appendix A (where P'_{ij} are the elements of an orthogonal 2×2 matrix) and the β -dependence of h_b as in (A.11), one obtains

$$X_d = \tan\beta P'_{11} \quad (6.14)$$

where P'_{11} can be of $\mathcal{O}(1)$. (Sometimes P'_{11} is denoted as $\cos\theta_{A_1}$.) Hence, for $\tan\beta \gg 1$, X_d can also be much larger than 1.

The $BR(\Upsilon(1S) \rightarrow \gamma A_1)$ is given by the Wilczek formula [300]

$$\frac{BR(\Upsilon(1S) \rightarrow \gamma A_1)}{BR(\Upsilon(1S) \rightarrow \mu^+\mu^-)} = \frac{G_F m_b^2 X_d^2}{\sqrt{2}\pi\alpha} \left(1 - \frac{m_{A_1}^2}{m_{\Upsilon(1S)}^2}\right) \times F \quad (6.15)$$

where α denotes the fine structure constant, and F is a correction factor which includes

three kinds of corrections to the leading-order Wilczek formula (the relevant formulae are summarised in [301]): bound state, QCD and relativistic corrections.

The CLEO collaboration recently published their negative results on the search for a CP-odd scalar (with a width below 10 MeV) [302]. Using the Wilczek formula (6.15), these results allow to put upper limits on X_d [293, 298].

Two caveats must be mentioned, however: first, the various (e.g. relativistic) corrections parametrized by F in (6.15) become large and unreliable for $m_{A_1} \gtrsim 8.8$ GeV, where F vanishes if the presently known corrections are extrapolated naïvely. Second, the experimental search assumed a width of A_1 below ~ 10 MeV which could be violated for m_{A_1} close to the Υ mass and/or for very large X_d .

Nevertheless, CLEO results impose strong constraints on CP-odd scalars with masses below ~ 8.8 GeV, which are shown as function of m_{A_1} and X_d (together with other constraints) in Fig. 6 in Section 6.4.

Mixing between the light CP-odd Higgs and η_b states

If the mass of the CP-odd Higgs is above ~ 9 GeV (as favoured by CLEO constraints), but below the $B\bar{B}$ threshold of ~ 10.5 GeV, a mixing between CP-odd hadronic resonances $\eta_b(nS)$ and the A_1 can become relevant in case of a large coupling X_d [294, 299, 303, 304]. Such a mixing could have a direct impact on the masses of the η_b system.

The BABAR collaboration has recently determined the $\eta_b(1S)$ mass $m_{\eta_b(1S)}$ with an error of only a few MeV from radiative decays $\Upsilon \rightarrow \gamma \eta_b$ of excited Υ states and the observation of peaks in the photon energy spectrum. The result from $\Upsilon(3S)$ decays is $m_{\eta_b(1S)} = 9388.9^{+3.1}_{-2.3}$ (stat) ± 2.7 (syst) MeV [305], and the result from $\Upsilon(2S)$ decays is $m_{\eta_b(1S)} = 9392.9^{+4.6}_{-4.8}$ (stat) ± 1.9 (syst) MeV [306]. The average gives [306] $m_{\eta_b(1S)} = 9390.9 \pm 3.1$ MeV, implying a hyperfine splitting $E_{hfs}(1S) = m_{\Upsilon(1S)} - m_{\eta_b(1S)}$ of

$$E_{hfs}^{\text{exp}}(1S) = 69.9 \pm 3.1 \text{ MeV} . \quad (6.16)$$

This result can be compared to predictions from QCD. Recent results based on perturbative QCD are in good agreement with each other and give $E_{hfs}^{\text{QCD}}(1S) = 44 \pm 11$ MeV [307] and $E_{hfs}^{\text{QCD}}(1S) = 39 \pm 14$ MeV [308] (see [309] and refs. therein; in phenomenological models as quark models $E_{hfs}(1S)$ varies over a wider range [310]). The discrepancy between $E_{hfs}^{\text{exp}}(1S)$ and $E_{hfs}^{\text{QCD}}(1S)$ could easily be cured in the presence of a mixing of the observed η_b with a CP-odd scalar A_1 with a mass somewhat above ~ 9.4 GeV [298].

On the one hand, a CP-odd Higgs scalar A_1 with a mass very close to 9.389 GeV (before mixing) *and* a strong $\eta_b - A_1$ mixing is excluded, since then the mass of the physical eigenstate (after mixing, i.e. after the diagonalization of the 2×2 mass matrix) could *not* be given by 9.389 GeV. This implies an upper bound on X_d for M_{A_1} near 9.389 GeV [298], which is also shown in Fig. 6.

On the other hand, a CP-odd Higgs scalar A_1 with a mass below 10.5 GeV can mix with all $\eta_b(nS)$ states ($n = 1, 2, 3$), possibly generating both a distorted spectrum as well as unusually large branching ratios into $\tau^+ \tau^-$ in the $\eta_b(nS) - A_1$ system [299]. Assuming that the large observed hyperfine splitting $E_{hfs}^{\text{exp}}(1S)$ compared to $E_{hfs}^{\text{QCD}}(1S)$ is induced by a mixing with A_1 , X_d can be determined as a function of M_{A_1} (leading to an increasing X_d with M_{A_1}), and predictions for the masses of the $\eta_b(nS) - A_1$ system can be made (within

errors). Denoting the 4 eigenstates of the $\eta_b(nS) - A_1$ system by η_i , $i = 1 \dots 4$, these masses are shown in Fig. 5 as function of M_{A_1} [299]. (M_{A_1} is denoted by m_A in Fig. 5.) For clarity we have indicated the masses $m_{\eta_b^0(nS)}$ before mixing as horizontal dashed lines. For $M_{A_1} \lesssim 9.8$ GeV, the state η_2 has a large A_1 component, but for $M_{A_1} \gtrsim 9.8$ GeV its mass drops below M_{A_1} due to the strong mixing with $\eta_b(2S)$.




Figure 5: Masses of all eigenstates η_i of the $\eta_b(nS) - A_1$ system as function of M_{A_1} , once m_{η_1} is forced to coincide with the BABAR result for $m_{\eta_b(1S)}$ (denoted by m_{obs} ; from [299]).

6.3 The anomalous magnetic moment of the muon

The measurement of the anomalous magnetic moment of the muon $a_\mu = (g_\mu - 2)/2$ by the E821 experiment at BNL has reached a level of precision which is sensitive to supersymmetric contributions. The latest experimental value obtained is [311]:

$$a_\mu^{\text{exp}} = 11\,659\,208.0(5.4)(3.3) \times 10^{-10}. \quad (6.17)$$

However, the evaluation of a_μ in the SM suffers from a relatively large uncertainty of the hadronic contribution (see, e.g., [312–314] and refs. therein). If it is determined using data from $e^+e^- \rightarrow \text{hadrons}$, one obtains a $\sim 3\sigma$ deviation between the SM and the experimental value [312–314] as, for instance, in [315]:

$$a_\mu^{\text{exp}} - a_\mu^{\text{SM}}(e^+e^-) = (27.7 \pm 9.3) \times 10^{-10}, \quad (6.18)$$

where the leading uncertainty originates both from the experimental measurement and the estimate of the hadronic contribution from e^+e^- data. Hence, a positive contribution to a_μ seems desirable.

However, if the hadronic contribution to a_μ is estimated from hadronic τ decays [312–314, 316], the discrepancy is reduced. The disadvantage of this approach is that it relies on assumptions on the pion form factor, isospin violating effects and vector meson mixings. On the other hand, recent (preliminary) measurements using the radiative return of $e^+e^- \rightarrow \pi^+\pi^-$ seem in better agreement with the estimate from hadronic τ decays [317, 318], according to which the discrepancy is reduced to $\sim 1.9\sigma$.

The various contributions to a_μ in MSSM-like supersymmetric models are reviewed and summarised in [319]. In the MSSM, the (dominant) one-loop contributions to a_μ originate from chargino/sneutrino or neutralino/smuon loops, which increase linearly with $\tan\beta$ and are positive for a positive μ -term of the MSSM. This phenomenon persists in the NMSSM with μ replaced by μ_{eff} ; the effects of the extended neutralino sector in the NMSSM are small [320].

Studies of a_μ in the NMSSM including a possibly light CP-odd Higgs scalar were performed in [315, 321, 322]; details of the necessary generalizations of all MSSM-like one- and two-loop contributions (see [319] and refs. therein) are given in [315]. The relevance of light CP-odd Higgs scalars for a_μ was actually first discussed in the context of general Two-Higgs-doublet extensions of the SM [323–325]: it is remarkable that positive contributions to a_μ from two-loop diagrams involving a closed fermion loop can dominate the negative contributions from one-loop Higgs-diagrams. The sum of both contributions from a light A_1 to a_μ has a (positive) maximum for $M_{A_1} \sim 6$ GeV. Below $M_{A_1} \sim 3$ GeV, the negative one-loop contribution dominates, which increases the discrepancy with respect to the measurement. Since the contributions of A_1 to a_μ are proportional to X_d^2 (its reduced coupling to charged leptons), one obtains upper limits on X_d for $M_{A_1} \lesssim 3$ GeV from (6.18), even after allowing for 2σ deviations, which are shown in Fig. 6 below.

6.4 Combined constraints on a light CP-odd Higgs

It is interesting to compare the constraints on a light A_1 in the NMSSM from CLEO, B physics, a_μ and from the measured $\eta_b(1S)$ mass. Clearly, constraints from B physics and a_μ (notably at large $\tan\beta$) also depend on MSSM-like parameters as M_{H^\pm} , chargino and slepton masses, which we will not review here and for which we refer to the corresponding literature. To some extent, the MSSM-like parameters of the NMSSM also affect the constraints from B physics and a_μ on a light A_1 in the NMSSM. It is possible, however, to scan over these parameters and obtain regions in the (X_d, A_1) -plane, which are *always* excluded. The results are shown in Fig. 6 taken from [298], where more details can be found.

We learn from Fig. 6 that, for $M_{A_1} \lesssim 9$ GeV, CLEO constraints are generally stronger than those from $B_s \rightarrow \mu^+\mu^-$, ΔM_q ($q = d, s$) and a_μ , with the exception of a small window for M_{A_1} near $M_B \sim 5$ GeV. In the range $9 \text{ GeV} \lesssim M_{A_1} \lesssim 10$ GeV, constraints due to the measured $\eta_b(1S)$ mass by BABAR play a rôle, but we recall that m_{A_1} slightly above 9.4 GeV can even have a desirable effect in the form of a negative shift of the $\eta_b(1S)$ mass, which would be interpreted as an unusually large $\Upsilon(1S) - \eta_b(1S)$ hyperfine splitting. In addition, constraints from LEP2 on a light CP-odd scalar below the $B - \bar{B}$ threshold are quite weak (cf. Section 5), which makes this mass range particularly interesting.

Future searches for a light A_1 in this mass range in radiative Υ decays will be challenging due to large backgrounds and the softness of the emitted photon. This problem can be




Figure 6: Upper bounds on X_d versus the A_1 mass for all parameters scanned over (see [298] for details). Indicated are constraints from $B_s \rightarrow \mu^+ \mu^-$ and ΔM_q , $q = d, s$ as a green dashed line, constraints from a_μ as a blue dashed line, the latest bounds from CLEO on $BR(\Upsilon \rightarrow \gamma \tau^+ \tau^-)$ as a black line and constraints due to the measured $\eta_b(1S)$ mass by BABAR as a red line.

overcome if one studies a breakdown of lepton universality in the inclusive leptonic decays of the Υ system as advocated in [298, 326, 327]: since A_1 would decay dominantly into $\tau^+ \tau^-$, the branching fraction into 2τ of an $\Upsilon(nS)$ state, decaying partially via A_1 , would be enhanced.

For a very light A_1 ($M_{A_1} \lesssim 1$ GeV), the decay $A_1 \rightarrow \mu^+ \mu^-$ could be dominant. The CLEO bounds mentioned above are valid for A_1 masses down to ~ 250 MeV, but the recent BABAR constraints from the absence of dimuon signals in $\Upsilon(3S)$, $\Upsilon(2S) \rightarrow \gamma A$ can be more relevant. These read [328]

$$BR(\Upsilon(3S) \rightarrow \gamma A) \times BR(A \rightarrow \mu^+ \mu^-) \lesssim 5.2 \times 10^{-6} \quad \text{for} \quad 0.212 \leq M_A \leq 9.3 \text{ GeV} \quad (6.19)$$

and give [329]

$$X_d \lesssim 0.4 \quad \text{if} \quad BR(A \rightarrow \mu^+ \mu^-) \sim 1. \quad (6.20)$$

Very light CP-odd scalars can also show up in Ω , Σ , K , η and π decays, which have been studied in the NMSSM in [269, 295, 329–333]. For $M_{A_1} \lesssim 360$ MeV, the decay $K^+ \rightarrow \pi^+ A_1$ is possible; for $M_{K^+} - M_{\pi^+} \gtrsim M_{A_1} \gtrsim 2 M_\mu$, the result of the HyperCP collaboration [334]

$$BR(K^+ \rightarrow \pi^+ \mu^+ \mu^-) = 9.8 \pm 1.0 \pm 0.5 \times 10^{-8} \quad (6.21)$$

implies $X_d(1 - \tan^{-2} \beta) \lesssim 0.06$ [329] within this specific range for M_{A_1} .

7 Specific scenarios

7.1 The constrained NMSSM

Even if one confines oneself to the \mathbb{Z}_3 -invariant NMSSM with a scale invariant superpotential, the number of independent parameters in the Higgs sector at tree level (six according to (2.17)) is considerably larger than the two parameters of the Higgs sector of the MSSM (typically chosen as $\tan\beta$ and M_A). However, one can make the assumption that the soft SUSY breaking terms are universal at a large scale like the GUT scale (not very different from the Planck scale), as it would be the case in mSUGRA with spontaneous supersymmetry breaking in a hidden sector and the mediation of supersymmetry breaking to the observable sector via flavour blind gravitational interactions [14]. The corresponding version of the \mathbb{Z}_3 -invariant NMSSM is denoted as the constrained NMSSM or cNMSSM.

Here, as in the constrained MSSM (cMSSM), all soft scalar masses squared, trilinear soft couplings and gaugino masses are assumed to be given by common values m_0^2 , A_0 and $M_{1/2}$, respectively, at the GUT scale. In fact, the number of independent parameters in the cNMSSM is the same as in the cMSSM: the additional parameters μ and B of the cMSSM [16] are replaced by the Yukawa couplings λ and κ in the superpotential (2.6). Apart from the gauge and quark/lepton Yukawa couplings, the Lagrangian of the cNMSSM depends on just five parameters m_0^2 , A_0 , $M_{1/2}$, λ and κ , and the correct value of M_Z reduces the dimension of the parameter space from five to four.

It is possible that the gravitational couplings (in the Kähler potential in supergravity) of the singlet S in the NMSSM differ from those of all other fields; then the singlet-dependent soft SUSY breaking terms could differ from universality (at the GUT or Planck scale). Therefore it is interesting to analyse models with relaxed universality conditions of the soft terms, which are slightly more general than the fully constrained NMSSM.

Below we will discuss first the cNMSSM, where the singlet-dependent soft SUSY breaking terms m_S and A_κ are assumed to coincide with m_0 and A_0 , respectively, at the GUT scale. Subsequently we will turn to the cNMSSM with relaxed universality constraints, where the values of m_S and/or A_κ at the GUT scale are allowed to differ from m_0 and A_0 .

First analyses of the phenomenologically acceptable regions of the cNMSSM parameter space [45, 46, 53] were handicapped by the still unknown top quark mass, which plays an important rôle due to the impact of the top Yukawa coupling on the RG equations [53] and the radiative corrections to the Higgs masses [110]. Somewhat later – but still before all present LEP2 constraints were available – the phenomenologically acceptable regions in this parameter space were investigated in [54, 72, 113, 197, 335, 336] (see [337] for an analysis of non-universal soft terms from orbifold string theory, and [338] for constraints on soft terms in dilaton-dominated SUSY breaking scenarios).

LEP2 constraints on the parameters of the Higgs sector of the cNMSSM have been analysed in [104, 127]. Concerning these, it is important to recall that larger values of λ do *not* necessarily imply an increase of the mass of the SM-like CP-even Higgs scalar: as indicated in and discussed below (3.2), the mixing of the SM-like Higgs with a heavy singlet-like Higgs (also proportional to λ) leads to a decrease of the SM-like Higgs mass. In the constrained parameter space of the cNMSSM, the off-diagonal matrix elements cannot

be fine tuned to 0. As a consequence, LEP constraints on the Higgs sector lead to an *upper* bound on λ within the cNMSSM:

$$\lambda \lesssim 0.3 . \quad (7.1)$$

Then the lower bound on $|\mu_{\text{eff}}| \equiv |\lambda s|$ of $|\mu_{\text{eff}}| \gtrsim 100$ GeV from the non-observation of a chargino lighter than ~ 100 GeV at LEP implies

$$|s| \gtrsim 300 \text{ GeV} . \quad (7.2)$$

For $|s| \gg v_u, v_d$, the dominant s -dependent terms in the potential (2.11) are given by (2.19) leading to the condition (2.20) for an absolute minimum with $s \neq 0$ [53, 54]:

$$A_\kappa^2 \gtrsim 9 m_S^2 . \quad (7.3)$$

For small λ and hence small κ from vacuum stability [54], the parameters A_κ and m_S are hardly renormalised between the GUT and the electroweak scales in the cNMSSM, and the above condition translates to (assuming $m_0^2 \geq 0$)

$$m_0 \lesssim \frac{1}{3} |A_0| . \quad (7.4)$$

Next, we consider the CP-odd Higgs boson mass matrix (2.27) whose diagonal matrix element $\mathcal{M}_{P,22}^2 \sim -3\kappa A_\kappa s$ (for large s as relevant here) must be positive. For positive s and κ this implies negative trilinear couplings

$$A_\kappa \sim A_0 < 0 . \quad (7.5)$$

In the region of the parameter space of the cNMSSM where $\lambda, \kappa \ll 1$, its phenomenology would be very close to the one of the cMSSM with corresponding soft SUSY breaking terms, since the additional Higgs and neutralino states of the NMSSM decouple and would never be produced – unless the additional singlino-like neutralino is the LSP (see below). For larger $\lambda \sim 0.3$ and low $\tan \beta \sim 2$, the mass of the lightest CP-even Higgs scalar with substantial couplings to gauge bosons and fermions can be about 10 GeV larger than in the cMSSM. However, the lightest CP-even Higgs scalar can also be singlet-like [54, 72, 113, 127, 197, 336], and escape LEP constraints due to its reduced coupling to the Z boson in spite of its small mass. The search for such a Higgs scalar would prove to be quite difficult at the LHC.

7.1.1 Constraints from the absence of charge and colour breaking minima

Additional constraints on the parameter space follow from the absence of charge and/or colour breaking (CCB) minima of the potential, where squark and/or slepton vevs do not vanish. A dangerous direction in field space is along the D -flat direction $|E_{R_1}| = |L_1| = |H_d|$ (with sleptons E_{R_1} and L_1 of the first generation with the smallest Yukawa coupling h_e). A deeper minimum is avoided if

$$\frac{1}{3} A_e^2 < m_{E_1}^2 + m_{L_1}^2 + m_{H_d}^2 \quad (7.6)$$

at the corresponding scale A_e/h_e [53, 339, 340]. (In the MSSM, an additional term μ^2 has to be added to the right-hand side of (7.6) [26]; in the NMSSM, s and hence μ_{eff} can vanish

in the dangerous minimum of the potential.) In principle, another dangerous direction in field space corresponds to $|T_R| = |Q_3| = |H_u|$; however, the corresponding constraints are never more relevant than the ones following from (7.6) in the cNMSSM.

In terms of the universal soft SUSY breaking parameters A_0 , $M_{1/2}$ and m_0 , (7.6) at the scale A_e/h_e becomes [340] (using the corresponding RG equations)

$$(A_0 - 0.5 M_{1/2})^2 \lesssim 9 m_0^2 + 2.67 M_{1/2}^2 \quad (7.7)$$

implying an upper bound on $|A_0|$.

Equally delicate could be the so-called unbounded-from-below (UFB) directions in field space, which are both D -flat and F -flat. In [339, 340], it has been clarified that such dangerous directions in the field space of the MSSM are still present in the NMSSM, although the singlet vev s gives an additional positive contribution to the potential. Analytic approximations to the potential along such dangerous directions have been studied in [341], with the conclusion that the inequality

$$m_0 \gtrsim (0.3 - 1.0) M_{1/2} \quad (7.8)$$

(where 1.0 corresponds to low $\tan \beta$, 0.3 to large $\tan \beta$) is an approximate condition for the absence of deeper minima in these directions. However, since the decay rate of the standard vacuum is usually much larger than the age of the universe [341], (7.8) can be violated if we assume that the early cosmology (temperature-induced positive masses squared for the squarks and sleptons) places us into the local standard minimum of the scalar potential.

7.1.2 The constrained NMSSM with dark matter constraints

Next one can require that the LSP of the cNMSSM provides the correct dark matter relic density, see Section 9. Early studies of correspondingly allowed regions of the cNMSSM parameter space have been performed in [335, 342]; in [339], the bounds (7.7) and (7.8) from the absence of CCB and UFB minima were taken into account. In the meantime, constraints on Higgs and sparticle masses as well on the dark matter relic density [343, 344] have become tighter (and RGEs/radiative corrections are known to a higher accuracy), with the result that the regions in the parameter space of the cNMSSM considered in these early studies are no longer phenomenologically viable.

As discussed above, the UFB constraint (7.8) can in principle be violated, and in the cNMSSM this is necessary in view of the up-to-date experimental constraints [216, 217]. On the other hand, the constraint on m_0 (7.4) (from $s \neq 0$) has to be respected in the cNMSSM, and it turns out that only regions where $m_0 \ll M_{1/2}$ are phenomenologically viable.

In the cMSSM [16], small values of $m_0 \lesssim \frac{1}{5} M_{1/2}$ result in a (charged) stau LSP $\tilde{\tau}_1$, which is excluded. In the cNMSSM, the additional singlino-like neutralino χ_1^0 can still be lighter than the lightest stau and be the true LSP – this allows for very small (or vanishing) values of m_0 . However, the χ_1^0 annihilation rate in the early universe must be large enough in order to avoid an excess of singlino-like dark matter. It turns out that, within the restricted parameter space of the cNMSSM, no s -channel resonances (Higgs or Z) with a mass twice the χ_1^0 mass are present, which could enhance the $\chi_1^0 - \chi_1^0$ annihilation rate (see Section 9).




Figure 7: The Higgs masses in the cNMSSM as a function of $M_{1/2}$ (in GeV). $m_0 = 0$ in the left panel, while $m_0 \sim M_{1/2}/10$ in the right panel. From below, the displayed lines correspond to the states h_1^0 (blue/dotted), h_2^0 (pink/dashed), a_1^0 (full/black) and a_2^0 (full/red) which is degenerate with the h_3^0 and h^\pm states (from [217]).

The only possibility to reduce the χ_1^0 abundance is then via coannihilation with $\tilde{\tau}_1$ (now the NLSP), whose mass has to be just somewhat above the χ_1^0 mass (see Section 9). Then the χ_1^0 abundance is reduced by the process $\chi_1^0 + X \rightarrow \tilde{\tau}_1 + X'$, and the $\tilde{\tau}_1 - \tilde{\tau}_1$ annihilation cross section is generally large enough. It should be mentioned that the singlino-like neutralino χ_1^0 would be practically invisible in direct or indirect dark matter detection experiments in this scenario.

The condition that the mass of $\tilde{\tau}_1$ is close to (just somewhat above) the χ_1^0 mass leads to $m_0 \lesssim \frac{1}{10}M_{1/2}$ and $A_0 \sim -\frac{1}{4}M_{1/2}$ [216, 217]; with these restrictions on the parameters, the LEP constraints on the Higgs sector are even more severe and require

$$\lambda \lesssim 0.02. \quad (7.9)$$

(As discussed in [216, 217] and in Section 9, the successful coannihilation of χ_1^0 with $\tilde{\tau}_1$ implies a lower bound on λ of $\sim 10^{-5}$.)

With A_0 being determined in terms of $M_{1/2}$ and the strong upper bounds on m_0 and λ , the spectrum of the cNMSSM is nearly completely determined by $M_{1/2}$. $\tan\beta$, which is no longer a free parameter, comes out relatively large with $\tan\beta > 25$.

The SM-like Higgs mass is 115-120 GeV for $M_{1/2} \gtrsim 400$ GeV, increasing with $M_{1/2}$. Within this range of $M_{1/2}$, all sparticle masses satisfy lower bounds from direct searches and from precision observables as those from B physics. If one requires that the SUSY contribution to the anomalous magnetic moment of the muon explains the $\sim 3\sigma$ deviation between the SM and the experimental value (see Section 6.3), the value of $M_{1/2}$ should be below ~ 1 TeV [216, 217, 315].

In Fig. 7 we display the masses of the neutral CP-even, CP-odd, and charged Higgs bosons as a function of the parameter $M_{1/2}$. On the left-hand side, we take $m_0 = 0$, while on the right-hand side we take the maximally viable value (from the dark matter relic density) $m_0 \sim \frac{1}{10}M_{1/2}$.

The CP-even Higgs boson with the dominant singlet component is the only Higgs state

whose mass depends – apart from $M_{1/2}$ – on m_0 . For small $M_{1/2}$ it is lighter than the SM-like Higgs boson, escaping LEP constraints due to the very small coupling to the Z boson. For increasing values of $M_{1/2}$, its mass increases until it becomes nearly degenerate with the SM-like CP-even Higgs state. In this region of parameter space, the singlet-like and SM-like Higgs states strongly mix; for a mass of the singlet-like Higgs state of ~ 100 GeV, the excess of events at this mass observed at LEP (cf. Section 5.1.1) could be explained [217].

The neutralino and slepton mass spectra are shown in Fig. 8. As for the Higgs bosons, $m_0 = 0$ in the left-hand panel, and $m_0 \sim \frac{1}{10}M_{1/2}$ in the right-hand panel. The two nearly degenerate sets of lower lines in both panels correspond to the masses of the χ_1^0 singlino-like LSP (blue/dotted) and the lighter stau $\tilde{\tau}_1$ NLSP (red/full). The mass difference between these two states is smaller than ~ 8 GeV, as required in order to obtain a cosmological relic density for the singlino χ_1^0 compatible with WMAP. The pattern for the masses of the charginos and the heavier neutralinos (blue/dotted lines) follows the one of the MSSM, once the proper relabeling of the states is made. Since the effective higgsino mass parameter μ_{eff} is generally quite large, $\mu_{\text{eff}} \gtrsim M_2$, the heavier neutralino states χ_4^0 and χ_5^0 are higgsino-like with masses $\sim \mu_{\text{eff}}$. The states χ_2^0 and χ_3^0 are, respectively, bino and wino-like with masses $m_{\chi_3^0} \approx 2m_{\chi_2^0} \approx M_2$ (with $M_2 \approx 0.75 M_{1/2}$). The charginos χ_1^\pm and χ_2^\pm are nearly degenerate in mass with, respectively, the wino-like χ_3^0 and the higgsino-like $\chi_{4,5}^0$ states.

In any case, the “smoking gun” for the cNMSSM would be the $\tilde{\tau}_1$ NLSP: all sparticle branching ratios into the singlino-like LSP are tiny for λ satisfying (7.9), hence all sparticles would decay at first into the $\tilde{\tau}_1$ NLSP. Only then the $\tilde{\tau}_1$ will decay into the χ_1^0 LSP and a τ lepton, which will thus appear in *every* sparticle decay chain. For very small λ (still larger than 10^{-5} , however) or a very small $\tilde{\tau}_1 - \chi_1^0$ mass difference, the $\tilde{\tau}_1$ life time can be so large that its decay vertices are visibly displaced [216, 217] (see Section 5.2.3). This signal would definitively be spectacular and serve to distinguish the cNMSSM from other supersymmetric extensions of the Standard Model.




Figure 8: Neutralino (blue/dotted lines), selectron (black/full lines) and stau (red/full lines) masses in the cNMSSM as a function of $M_{1/2}$ (in GeV); on the left-hand side $m_0 = 0$, while on the right-hand side $m_0 \sim M_{1/2}/10$. In both panels the states are ordered in mass as $m_{\chi_1^0} \lesssim m_{\tilde{\tau}_1} < m_{\tilde{e}_R} < m_{\chi_2^0} < m_{\tilde{\tau}_2} \lesssim m_{\tilde{e}_L} < m_{\chi_3^0} < m_{\chi_{4,5}^0}$. The charginos χ_1^\pm and χ_2^\pm are degenerate in mass with, respectively, χ_3^0 and $\chi_{4,5}^0$ (from [217]).

7.1.3 The constrained NMSSM with relaxed universality conditions

The constrained NMSSM with relaxed universality conditions in the singlet sector, allowing for both $m_S^2 \neq m_0^2$ and $A_\kappa \neq A_0$ at the GUT scale, has been first analysed – without requiring a dark matter relic density in agreement with WMAP constraints – in [345]. Then the bounds (7.4) (from the bound (7.3) on m_S, A_κ) and (7.5) no longer apply; notably m_0 is no longer bounded from above. Also the upper limit (7.1) on λ is relaxed to $\lambda \lesssim 0.55$. The freedom in the choice of A_κ allows for a possibly quite light CP-odd Higgs scalar A_1 . This opens new phenomenologically viable regions in parameter space, where the lightest CP-even Higgs mass M_{H_1} is well below 114 GeV, but LEP constraints are satisfied since H_1 decays dominantly into a pair of CP-odd Higgs scalars (see Section 5.1). On the other hand, the value of A_κ at the GUT scale must be chosen inside a narrow window in order to obtain $0 < M_{A_1} < \frac{1}{2} M_{H_1}$ [228, 345].

Since the detection of Higgs-to-Higgs decays will be particularly challenging at the LHC, some particular points in the parameter space of the semi-constrained NMSSM have been proposed as benchmark points in [228]. These include scenarios corresponding to a light CP-odd Higgs scalar A_1 ; allowing, in addition, the Higgs masses m_{H_u} and m_{H_d} to differ from m_0 at the GUT scale, scenarios with light CP-even singlet-like Higgs scalars can be obtained as well. Both cases require particular search strategies at the LHC, which are briefly surveyed in [228] and discussed in Section 5.1.

A complete analysis of the parameter space of the semi-constrained NMSSM, including dark matter constraints combined with those from LEP and other colliders on Higgs and sparticle searches, B -physics and the anomalous magnetic moment of the muon, has been performed in [346, 347]. In contrast to the fully constrained NMSSM, where the small values for m_0 imply a $\tilde{\tau}_1$ lighter than the bino-like neutralino, the LSP can now either be bino-, higgsino- or singlino-like. This, and the possible presence of light Higgs states in s -channels, lead to many additional regions in parameter space where the WMAP constraints on the dark matter relic density can be satisfied. The corresponding processes, as well as the prospects for the direct detection of the neutralino LSP [347], will be discussed in more detail in Section 9.

The potential of the LHC to unambiguously identify the semi-constrained NMSSM by observing at least 3 neutral Higgs states is confined to the regime of large values of $\tan\beta \sim 50$ and $\lambda \sim 0.1$, where one can benefit both from the enhanced couplings of the heavy Higgs states to b -quarks and from substantial Higgs doublet/singlet mixings [347].

The semi-constrained NMSSM with $m_S \neq m_0$, but imposing $A_\kappa = A_0$ at the GUT scale, has been investigated in [348], with the result that the phenomenologically viable regions in parameter space are still considerably larger than in the fully constrained NMSSM.

7.2 The NMSSM and Gauge Mediated Supersymmetry Breaking

The μ -problem of the MSSM is particularly acute in the framework of GMSB (see [349] for a review). The essential ingredients of GMSB models are a hidden or sequestered sector, where supersymmetry is spontaneously broken in such a way that the component F_X of at least one superfield \hat{X} – often considered as a spurion without kinetic term – does not vanish. In addition, a messenger sector $\hat{\varphi}_i$ with vector-like Standard Model gauge quantum

numbers exists, which couples to \widehat{X} in the superpotential:

$$W = \widehat{X} \widehat{\varphi}_i \widehat{\varphi}_i + \dots \quad (7.10)$$

The messengers $\widehat{\varphi}_i$ have a supersymmetric mass M_{mess} (the messenger scale), which can be parametrized by a vev $\langle X \rangle = M_{\text{mess}}/2$ of the scalar component of \widehat{X} . The non-vanishing vev of F_X induces a SUSY breaking mass splitting \widehat{m} between the scalar/pseudoscalar components of $\widehat{\varphi}_i$. Typically, dynamical supersymmetry breaking is assumed as the origin of $F_X \neq 0$ [349], but Kähler potentials in supergravity of the No-Scale type, together with Giudice-Masiero-like terms for the messengers, can also lead to models with GMSB [156]. Since the messengers $\widehat{\varphi}_i$ carry Standard Model gauge quantum numbers – often in the form of a complete $SU(5)$ multiplet – they induce SUSY breaking gaugino masses of the order

$$M_i \sim \frac{\alpha_i}{4\pi} \frac{F_X}{M_{\text{mess}}} \sim \frac{\alpha_i}{4\pi} \frac{\widehat{m}^2}{M_{\text{mess}}} \quad (7.11)$$

at one loop, and SUSY breaking scalar masses squared of the order

$$m_i^2 \sim \left(\frac{\alpha_i}{4\pi} \frac{\widehat{m}^2}{M_{\text{mess}}} \right)^2 \quad (7.12)$$

at two loops. However, neither a μ -term nor a $B\mu$ -term are generated, which would lead to serious phenomenological problems as summarised in the Introduction. The simplest solution to this problem is the introduction of a singlet superfield \widehat{S} with the scale invariant superpotential (2.6). However, since \widehat{S} is a gauge singlet, the radiative corrections mentioned before will *not* generally generate a soft mass m_S^2 and/or a trilinear coupling A_κ of the order $M_{\text{SUSY}} \sim M_i$ [55, 350], whereas at least one of these parameters (with $m_S^2 < 0$) is required in order to trigger a sufficiently large vev of s , cf. (2.19) – (2.21).

Numerous proposals have been put forward in order to successfully solve the μ -problem in GMSB models within the NMSSM or slight modifications thereof [55, 62, 71, 96, 156, 349–367]:

(i) Several gauge singlets (or non-renormalisable self-interactions of \widehat{S}) could trigger a vev s which is large enough [350–353, 357, 359, 363, 364, 366, 367]. Since gauge symmetries do not constrain the possible couplings of singlets, the choice of the terms kept in the superpotential should preferably be justified by a discrete (R-)symmetry.

(ii) If the messenger scale M_{mess} is large enough, the soft mass m_S^2 can become negative at low scales – as desired – due to a positive β -function and the RG evolution; however, additional positive terms in the β -function of m_S^2 beyond the ones of the NMSSM (see (B.6)) are generally necessary to this end. Such terms appear once \widehat{S} couples to additional light matter \widehat{Q}_i with vector-like Standard Model gauge quantum numbers in the form $\widehat{S} \widehat{Q}_i \widehat{Q}_i$, provided the soft SUSY breaking scalar masses of Q_i are large enough [71, 350, 351, 355]. Alternatively, the positive terms in the β -function of m_S^2 within the NMSSM (B.6) can already be large enough, if one allows for more general soft SUSY breaking Higgs (and squark) masses. This requires more than one spurion \widehat{X} and $SU(5)$ -breaking spurion-messenger couplings [365, 367].

(iii) The singlet \widehat{S} can couple directly to the messengers $\widehat{\varphi}_i$. The simplest coupling of the form $\eta \widehat{S} \widehat{\varphi}_i \widehat{\varphi}_i$ generates tadpole terms ξ_F and ξ_S [62, 96, 156, 351, 353] (defined in (2.1) and (2.5)) of the order

$$\begin{aligned}\xi_F &\sim \frac{\eta}{16\pi^2} F_X \sim \eta M_{\text{SUSY}} M_{\text{mess}} , \\ \xi_S &\sim \frac{\eta}{16\pi^2} \frac{F_X^2}{M_{\text{mess}}} \sim \eta M_{\text{SUSY}}^2 M_{\text{mess}}\end{aligned}\tag{7.13}$$

(see (4.1) with $\Lambda \sim M_{\text{mess}}$). On the one hand, such terms would trigger a vev s even for negligibly small parameters m_S^2 and A_κ ; on the other hand, the values of ξ_F and ξ_S can easily be too large for a messenger scale $M_{\text{mess}} \gg M_{\text{SUSY}}$ – this is nothing but the singlet tadpole problem already discussed in Section 4. However, a small Yukawa coupling $\eta \sim 10^{-5}$ can render such a scenario phenomenologically acceptable [96].

(iv) The singlet \widehat{S} can couple directly to a generalized messenger sector invariant under a discrete symmetry such that tadpole diagrams are forbidden [356]. Then parameters A_κ and $m_S^2 < 0$ of the desired order are radiatively generated to one- and two-loop order, respectively, leading to a consistent phenomenology [96, 362, 363].

The phenomenological consequences of the diverse scenarios can be quite different. First, the squark and slepton spectrum depends on the quantum numbers of the messengers and is generally easy to distinguish from, e.g., mSUGRA. Second, additional light matter fields \widehat{Q}_i as in (ii) above would typically be observable at the LHC. Finally, an approximate $U(1)_R$ symmetry in the NMSSM Higgs sector [68–70, 205, 368] in the context of GMSB [71, 96] can imply light CP-odd Higgs scalars, into which the SM-like Higgs scalar can decay; as discussed in Section 5, this scenario would have an important impact on the search for Higgs bosons at the LHC.

Finally we should mention that the μ -problem is also present in models with supersymmetry breaking from extra dimensions, Anomaly Mediated Supersymmetry Breaking and models where the MSSM sector couples to a superconformal field theory; again, the NMSSM and its variants as described above have been proposed as a way out [369–375].

7.3 The nMSSM

As discussed in Section 4, a solution of the domain wall problem of the scale invariant NMSSM invariant under a discrete \mathbb{Z}_3 symmetry consists in assuming the presence of tadpole terms $\sim \xi_S$ and/or $\sim \xi_F$ in the soft SUSY breaking Higgs potential and/or the superpotential. In order to solve the domain wall problem, these parameters can be very small such that their impact on the phenomenology can be neglected. On the other hand it has been proposed in [166] that notably ξ_S could be of the order $\xi_S \sim M_{\text{SUSY}}^3$ and, in [167], $\xi_F \sim M_{\text{SUSY}}^2$. Then, the tadpole terms in the potential trigger a vev $s \sim M_{\text{SUSY}}$ even for $\kappa \rightarrow 0$, i.e. the terms $\frac{\kappa}{3} \widehat{S}^3$ in the superpotential and $\frac{1}{3} \kappa A_\kappa S^3$ in $\mathcal{L}_{\text{soft}}$ can be omitted. The resulting version of the NMSSM, where the singlet only appears in the term $\lambda \widehat{S} \widehat{H}_u \cdot \widehat{H}_d$ in the superpotential and in the tadpole terms $\sim \xi_S$ and/or $\sim \xi_F$, has been denoted as the Minimal Non-minimal or new Minimal Supersymmetric Standard Model.

The phenomenological consequences of this class of models consist in a possibly quite light charged Higgs boson (lighter than the SM-like Higgs boson) into which top quarks can decay [167, 376], and a neutralino LSP which has always a large singlino component [167, 377]. Even so, as shown in [176, 378–380], the dark matter relic density can well be of the required magnitude: the neutralino LSP can annihilate either through the Z resonance (due to its small, but non-vanishing higgsino component) or through a CP-odd Higgs resonance in the s -channel (see Section 9). Bounds from $Z \rightarrow \chi_1^0 \chi_1^0$ have been checked; interestingly, the SM-like Higgs scalar could dominantly decay invisibly into $\chi_1^0 \chi_1^0$ [176]. All additional phenomenological constraints on supersymmetric models can be satisfied in this quite “economic” class of models [379, 380], and the prospects of neutralino, chargino and Higgs discoveries at the LHC and the ILC in the nMSSM have been worked out in [379]. Finally we recall that the nMSSM also allows for successful electroweak baryogenesis [176, 177, 179] (see Section 4).

7.4 The fine tuning problem in the NMSSM

One of the most important motivations of supersymmetric extensions of the SM is the solution of the hierarchy problem: if one assumes that the Lagrangian of the SM is valid up to a very high scale as M_{GUT} , due to quadratically divergent radiative corrections the Higgs mass squared has to be tuned with a relative precision of $(M_{\text{weak}}/M_{\text{GUT}})^2 \sim 10^{-28}$ in order to obtain a weak scale (given by the Higgs vev) 14 orders of magnitude below the GUT scale.

In supersymmetric extensions of the SM, the weak scale is naturally given by the SUSY breaking scale M_{SUSY} (if the μ problem is solved). The details of this relation depend, however, on the supersymmetric model considered: on its field content, and on the values of the unknown parameters. It is well known that the lower LEP bounds on a Higgs scalar with a SM-like coupling to the Z boson (and a large branching ratio into $b\bar{b}$), as well as the present lower bounds on sparticle masses, induce a “little fine tuning problem” in the MSSM: its parameters have to be tuned with a relative precision of $\sim 10^{-2}$ in order to satisfy these constraints. It has been suggested that the NMSSM (or variants thereof) can alleviate this problem, which we will discuss below (see [381] for a recent review).

There are different ways to quantify the amount of fine tuning within a given model: an analytic approach consists in identifying the independent parameters p_a of the fundamental Lagrangian, in terms of which all particle masses including M_Z are determined. To each parameter one associates a fine tuning measure Δ_a defined by

$$\Delta_a = \left| \frac{d \ln M_Z}{d \ln p_a} \right| \quad (7.14)$$

(or M_Z replaced by M_Z^2), and the “amount of fine tuning” is given by the maximum of all Δ_a . For the SM with a fundamental Lagrangian defined at the GUT scale one would obtain $\Delta_{\text{max}} \sim 10^{14}$, but all values much larger than 1 are considered as unnatural. In practice, after radiative corrections, the dependence of M_Z on all p_a is so complicated that the various Δ_a can only be approximatively estimated which is, however, sufficient to identify a potential fine tuning problem.

Alternatively one can use codes which compute the particle spectrum numerically in terms of input parameters: then one can perform scans over a large number of input parameters, and study which proportion of the parameter space is still consistent with present experimental constraints. However, the result of such an analysis is related to the required amount of fine tuning only if the input parameters correspond to the independent parameters p_a of the fundamental Lagrangian. Then, although the result depends somewhat on the measure and on the initial range used to scan over the parameter space, its physical interpretation is clearer than the one of the formal fine tuning parameters Δ_a introduced above.

Let us now consider the origins of the “little fine tuning problem”, starting with the generation of the weak scale in terms of the soft SUSY breaking Higgs masses m_{H_u} , m_{H_d} and μ_{eff} (in the NMSSM): neglecting the radiative corrections (which play no important rôle here), the minimisation equations (2.15) of the Higgs potential imply

$$M_Z^2 \simeq -2\mu_{\text{eff}}^2 + \frac{2(m_{H_d}^2 - \tan^2 \beta m_{H_u}^2)}{\tan^2 \beta - 1}. \quad (7.15)$$

In the absence of fine tuning, all terms on the right-hand side of (7.15) should be of comparable magnitude, and no large cancellations should occur; hence both μ_{eff}^2 and $-m_{H_u}^2$ should not be much larger than $\mathcal{O}(M_Z^2)$. The bound $|\mu_{\text{eff}}| \gtrsim 100$ GeV from the lower bound on chargino masses already starts to generate a slight tension, but the main problem stems typically from the value of $m_{H_u}^2$ once the soft SUSY breaking terms are generated at a large scale as M_{GUT} :

Fortunately, given the large value of the top Yukawa coupling, the dominant positive terms proportional to the stop masses squared in the one-loop RGE for $m_{H_u}^2$ (cf. (B.6)) generate easily $m_{H_u}^2 < 0$ at the weak scale as desired [19]. However, one typically obtains $m_{H_u}^2 \sim -m_T^2$, which is often much larger (in absolute value) than M_Z^2 : from the RGEs for m_T^2 one finds that m_T is never much smaller than the gluino mass M_3 (at M_{SUSY}) which, in turn, is bounded from below by ~ 300 GeV from searches at the Tevatron (see [89] and refs. therein).

Clearly these arguments are the same for the MSSM and the NMSSM, and could be alleviated in the case of the generation of the soft SUSY breaking terms at a much lower scale than M_{GUT} , as possible in GMSB. (See [62, 350, 354, 355, 367, 382] for discussions of the fine tuning problem in GMSB and proposals for solutions within the NMSSM or variants thereof.)

At least within the MSSM, another strong argument for large values of m_T originates from the LEP bound on the SM-like Higgs scalar mass, whose theoretical expression was given approximately in (3.2) according to which it increases proportional to $\ln\left(\frac{m_T^2}{m_t^2}\right)$. In the MSSM, large values for m_T are unavoidable in order to satisfy the LEP bound. Albeit large stop masses are consistent with the non-observation of stops, they would generate a too large value for $-m_{H_u}^2$ as discussed above.

In the NMSSM, two different strategies can be pursued in order to alleviate the fine tuning problem without assuming large values of m_T : first, one can try to use the first positive term in the second line of (3.2) to push the Higgs mass above the LEP bound. Then λ should be as large as possible, and $\tan \beta$ must be small in order to avoid a suppression

by $\sin^2 2\beta$. As discussed in Section 3, within the NMSSM λ is bounded from above by $\lambda \lesssim 0.7 - 0.8$ if one requires the absence of a Landau singularity below the GUT scale; nevertheless, a reduction of the required amount of fine tuning by the LEP bound has been observed in [175]. Otherwise one can enlarge the particle content or reduce the range of validity of the NMSSM to scales far below M_{GUT} in order to justify larger values of λ [128–133, 136, 383, 384].

Instead of trying to push the Higgs mass above the LEP bound with the help of the NMSSM specific terms, it is conceivable that the LEP bound of 114 GeV on the mass of a SM-like Higgs scalar H does not apply within the NMSSM (see Section 5): if H decays dominantly into a pair of light pseudoscalars $H \rightarrow AA$, the branching fraction for $H \rightarrow b\bar{b}$ can be much smaller than 1, and the Higgs mass M_H can be well below 114 GeV even for SM-like couplings of H to the Z boson. If A decays into $b\bar{b}$, combined LEP constraints still require $M_H > 110$ GeV [57], hence A should be lighter than 10.5 GeV. Then, M_H can be as low as 86 GeV [212] which is easily compatible with (3.2) without large stop masses. A reduction of the required fine tuning in the region of the parameter space of the NMSSM corresponding to light pseudoscalars has been discussed first in [200, 202] and was elaborated further in [70, 205, 215, 293]. (However, the latest constraints from ALEPH [213] should be taken care of.)

At first sight, light pseudoscalars appear naturally in the NMSSM in the case of an approximate global Peccei-Quinn- or R-symmetry (in the Higgs sector, see Section 2.2). It is not always straightforward, however, to realise a consistent scenario in these limits. In the Peccei-Quinn limit ($\kappa \rightarrow 0$ in the scale invariant NMSSM), two possible minima exist generically for the vev of the singlet s : one is the “large s solution” where $s \sim 1/\kappa$ (see (2.21)) would be very large. Then, in order to avoid an excessively large $\mu_{\text{eff}} \equiv \lambda s$, λ must be of the order of κ which corresponds to the decoupling limit (2.40). Since the light pseudoscalar would be nearly a pure singlet, the coupling HAA and hence the desired branching ratio would disappear. In the other possible minimum for s where $s \simeq \lambda A_\lambda \sin 2\beta/\lambda^2 + m_S^2/v^2$ (from the third of eqs. (2.15)), it is difficult to have $|\mu_{\text{eff}}| \equiv |\lambda s|$ large enough (above 100 GeV) and to ensure that it is the global minimum of the potential [368].

In the R-symmetry limit both trilinear soft terms A_λ and A_κ are assumed to be small; however, since the R-symmetry is broken by the gaugino masses M_i , A_λ and A_κ receive radiative corrections proportional to M_i (cf. the RGEs in (B.3)). Hence it is unnatural to assume that A_λ and A_κ are smaller than these radiative corrections, which requires some tuning in the (5 – 10)% range [70]. Hence, at least within the minimal NMSSM without enlarged particle content up to the GUT scale, the present experimental constraints also require some adjustment of parameters.

Besides these qualitative arguments for the required amount of fine tuning, probability distributions or likelihood analyses in parameter space can be studied numerically using scans over input parameters and suitable codes, which compute spectra and couplings and allow to check experimental constraints. As stated above, the input parameters should preferably correspond to the parameters of the fundamental Lagrangian (at the GUT scale) up to an overall mass scale of the soft terms, which can be fixed by M_Z . Then, quantities like $\tan\beta$ are obtained as output from the minimisation of the Higgs potential. This approach had been pursued in the fully constrained NMSSM in [54, 113, 127] with the result that

$\sim 90\%$ of the points in the parameter space still allowed by LEP1 are now eliminated by fruitless searches for the Higgs boson, charginos and sleptons at LEP2, implying a necessary fine tuning of at least $\sim 10\%$. (Also, large values $\tan \beta \gtrsim 10$ require fine tuned parameters at the GUT scale in the cNMSSM.)

Apart from the investigation of the required fine tuning, such analyses serve to study the reach of future experiments in the parameter space within specific scenarios. In the semi-constrained NMSSM (where the soft singlet mass squared $m_{\tilde{S}}^2$ is not required to unify with m_0^2 , see Section 7.1.3) such studies have recently been performed in [385, 386], where also an LSP dark matter relic density compatible with WMAP constraints was imposed. In these approaches, the code NMSPEC [345] was used, where $\tan \beta$ is an input parameter and $m_{\tilde{S}}^2$ and κ are determined by the minimisation equations of the Higgs potential, which can hide the required amount of fine tuning, notably for large soft SUSY breaking terms. Large values of $\tan \beta$ and/or quite heavy sparticle masses appear with relatively high probabilities in [385, 386], which would imply that the LHC cannot test large parts of the corresponding parameter space (whereas the prospects for direct dark matter detection in the future look better). It can be expected that, once fine tuning criteria are taken into account, the prospects of sparticle detection at the LHC would look much brighter.

8 Variants of the NMSSM

8.1 CP violation

In the SM, CP is explicitly violated by charged current interactions: complex Yukawa couplings lead to one physical phase in the Cabibbo-Kobayashi-Maskawa (CKM) quark mixing matrix (δ_{CKM}), which can successfully explain the observed direct and indirect CP violation (CPV) in the neutral kaon sector (ε'/ε and ε_K , respectively) and in the B -meson sector [89]. On the other hand, the non-observation of electric dipole moments (EDMs) of the electron, neutron and atoms (like Hg) puts severe constraints on flavour *conserving* phases, such as the strong CP phase, forcing them to be extremely small.

Supersymmetric extensions of the SM, like the MSSM and the NMSSM, introduce several potential new sources of explicit CPV: all couplings and fermion masses appearing in the superpotential or in the soft SUSY breaking Lagrangian can be complex, leading to non-trivial CPV phases. SUSY CPV can manifest itself in several low energy observables: in addition to the new mixings in the Higgs sector and the implications for the Higgs spectrum and decays, one can expect contributions to the EDMs, lepton polarisation asymmetries in semi-leptonic decays, CPV in B -meson mixings and decays, among many others. The measured values of (or bounds on) these observables lead to severe constraints on the new phases, which is sometimes denoted as the “SUSY CP problem”.

In particular, flavour conserving SUSY phases can induce very large contributions to the EDMs, several orders above the current bounds, even if the SUSY spectrum is relatively heavy (~ 1 TeV). In the absence of some cancellation mechanism (as it would be natural in minimal flavour violation, such as in the case of scenarios based on mSUGRA or GMSB), complying with observation forces the new flavour conserving phases to be very small, typically in the range $10^{-3} - 10^{-1}$. This holds both in the MSSM and the NMSSM.

In principle, generalizations of the Higgs sector of the SM allow for spontaneous CPV at the electroweak scale, an attractive scenario at first sight. However, in all supersymmetric models (including the NMSSM) where H_u couples to up-type and H_d to down-type quarks only, possible phases of $H_{u,d}$ can be rotated away by redefinitions of quark fields, and the CKM matrix is real. (This remains true if explicit CPV occurs in the Higgs sector only.) As a consequence, SM contributions to the CPV observables are absent, and all the observed values must be generated from SUSY contributions, which is difficult (but not excluded *à priori*). Furthermore, the structure of the Higgs potential often forbids stable minima with spontaneous CPV as in the MSSM and the \mathbb{Z}_3 -invariant NMSSM (see below).

Independently from CPV in charged or neutral current interactions, spontaneous or explicit CPV in the Higgs sector can have important phenomenological implications: in the CP conserving case, scalar and pseudoscalar Higgs bosons do not mix, leading to physical states with well defined CP parity. CPV in the Higgs sector may lead to mixings between scalar and pseudoscalar states (as the physical states no longer are CP eigenstates): the full neutral Higgs mass matrix will no longer be block diagonal, but a 5×5 matrix which can be parametrized as

$$\mathcal{M}_{H^0}^2 = \begin{pmatrix} \mathcal{M}_S^2 & \mathcal{M}_{SP}^2 \\ (\mathcal{M}_{SP}^2)^T & \mathcal{M}_P^2 \end{pmatrix}, \quad (8.1)$$

where $\mathcal{M}_{SP}^2 \neq 0$ implies CPV. Actually, at tree level no CP violation exclusively within the Higgs doublet sector occurs, since it can be “rotated away” by field redefinitions; at tree level only NMSSM specific doublet-singlet couplings can violate CP.

In general, the CPV couplings of the Higgs scalars to SM gauge bosons and fermions, to superpartners, and to Higgs bosons themselves can significantly differ from the CP conserving case, which requires to re-interpret present bounds on the Higgs spectrum, and has consequences for Higgs and sparticle detection at colliders. Finally, given the possibility of having an enhanced first order electroweak phase transition in the context of the NMSSM (see Section 4.3), additional sources of CPV, beyond those of the SM, could play an important rôle in obtaining successful electroweak baryogenesis, which is difficult within the MSSM.

In what follows, we briefly review some relevant aspects of NMSSM-like models with spontaneous or explicit CPV, focusing on the Higgs sector and the relevant literature. (For a review of CPV in connection with non-standard Higgs sectors see [387].)

8.1.1 Spontaneous CP violation

Assuming a CP conserving Lagrangian before electroweak symmetry breaking implies that all bilinear, trilinear and quartic terms can be chosen real (after possible field redefinitions). The Higgs fields can develop complex vevs, so that (2.10) generalizes to

$$\langle H_u \rangle = v_u e^{i\varphi_u}, \quad \langle H_d \rangle = v_d e^{i\varphi_d}, \quad \langle S \rangle = s e^{i\varphi_s}. \quad (8.2)$$

The two physical phases (up to a $U(1)_Y$ gauge transformation),

$$\theta = \varphi_u + \varphi_d + \varphi_s, \quad \delta = 3\varphi_s, \quad (8.3)$$

open the possibility of spontaneous CPV (SCPV). CP can be spontaneously broken at the minimum of the tree level Higgs potential, or only once radiative corrections to the effective potential are included – the latter case is sometimes denoted as “radiative SCPV”.

Tree level SCPV in the \mathbb{Z}_3 -invariant NMSSM

As stated in a no-go theorem in [388], spontaneous CP violation is not possible in the \mathbb{Z}_3 -symmetric NMSSM with a superpotential as in (2.6) at tree level. Minimising the scalar potential

$$\begin{aligned} V_{\text{Higgs}} = & \frac{g_1^2 + g_2^2}{8} (v_u^2 - v_d^2)^2 + (m_{H_u}^2 + \lambda^2 s^2) v_u^2 + (m_{H_d}^2 + \lambda^2 s^2) v_d^2 + m_S^2 s^2 \\ & + \kappa^2 s^4 + \lambda^2 v_u^2 v_d^2 + 2\kappa\lambda v_u v_d s \cos(\theta - \delta) \\ & + 2\lambda A_\lambda v_u v_d s \cos\theta + \frac{2}{3}\kappa A_\kappa s^3 \cos\delta \end{aligned} \quad (8.4)$$

with respect to the absolute values of the vevs and their phases, one finds that CP violating extrema are always local maxima and never local minima of the scalar potential, implying negative masses squared for at least one Higgs boson.

Radiatively induced SCPV in the \mathbb{Z}_3 -invariant NMSSM

Radiatively induced SCPV is theoretically possible in the NMSSM [389–391], but these scenarios are severely constrained by the non-observation of Higgs pair production off Z bosons at LEP: analogous to the MSSM, a very light spin-0, non-Goldstone boson appears in the spectrum. The presence of such a light boson is a consequence of the Georgi-Pais theorem [392], and current experimental bounds seem to lead to the exclusion of this class of models.

SCPV in the general NMSSM

In the general NMSSM, where dimensionful terms are present in the superpotential (2.1), the no-go theorem of [388] no longer holds and spontaneous CP violation is indeed viable already at tree level [393,394]. Combining the minimisation conditions for spontaneous CP violation with the constraints from ε_K , it was noticed that the theoretical upper bound on the lightest Higgs boson mass becomes stronger, leading to $M_{H_1} \lesssim 100$ GeV (still in agreement with LEP bounds due to reduced couplings to SM gauge bosons) [393,394]. However, these models re-introduce a μ -term in the superpotential so that a solution to the μ -problem of the MSSM is not obtained.

SCPV in the NMSSM with a tadpole term

As discussed in Section 7.3, the nMSSM contains singlet tadpole terms $\sim \xi_S, \xi_F$ in the scalar potential and/or superpotential [165–167,376,377], but $\kappa = 0$. Then, as in the MSSM, neither spontaneous nor explicit CPV in the Higgs sector is possible at tree level. Keeping $\kappa \neq 0$, but adding just a soft SUSY breaking tadpole term $\sim \xi_S$ to the otherwise \mathbb{Z}_3 -invariant NMSSM, allows already to circumvent the no-go theorem in [388], and one can find true minima of the scalar potential associated with non-trivial spontaneous CP violating phases which are phenomenologically viable [395]. Again some Higgs states can be quite light, but not excluded by current bounds due to their reduced coupling to gauge bosons. The new Higgs mass matrices, together with the relevant radiative corrections, can be found in [395].

Problems with SCPV

However, although elegant and apparently simple to realise, scenarios of SCPV are seldom viable: recalling that the CKM matrix is real, there are no SM contributions to either flavour conserving (EDMs) or flavour violating CP violating observables. SUSY contributions, with δ and θ as the only sources of CPV, would be the only means to saturate the observed values of $\varepsilon_K, \varepsilon'/\varepsilon$, the CP asymmetry of the B_d meson decay, etc.. In particular, the dominant contributions to ε_K should arise from the chargino-mediated box diagrams [395,396]. In general, complying with observation requires the phases δ and θ to be quite large, as well as maximal left-right squark mixing [396,397]. Large values for flavour conserving CP violating phases nearly inevitably lead to sizable contributions to the EDMs of the electron, neutron and atoms. Complying with the EDM bounds, either via a cancellation mechanism, or a heavy SUSY spectrum (not always possible as excessively heavy squarks would preclude saturating the values of ε_K etc.) represents the most serious challenge to the survival of all models with SCPV in the Higgs sector.

Spontaneous CP violation at finite temperature

As discussed in Section 4.3, additional sources of CP violation in the Higgs sector would be desirable for baryogenesis. An interesting hypothesis is to assume that CP is indeed conserved at $T = 0$, and to have spontaneous CP violation at finite temperature in the \mathbb{Z}_3 -invariant NMSSM. This would allow to evade constraints from EDMs, and to avoid a light spectrum in the Higgs sector (arising in the case of radiative CPV, as discussed before). Finite temperature effects can indeed trigger spontaneous CP violation inside the walls of the propagating bubbles present after the phase transition [180,181]. An extremely tiny explicit phase is nevertheless required in the effective potential, in order to lift the degeneracy between vacua (each generating identical baryon asymmetries, but of opposite signs). Such a phase, typically $\mathcal{O}(10^{-6} - 10^{-5})$, would give rise to negligible contributions to the EDMs.

8.1.2 Explicit CP violation

The NMSSM is the simplest SUSY extension of the SM where one can have explicit CP violation in the Higgs sector at tree level [45,398]. In the MSSM the explicit CP violating phases in the Higgs sector (in μ and B) can be rotated away by a redefinition of the H_u and H_d fields; this is not possible in the NMSSM due to the presence of the additional singlet couplings. Since now the standard CKM mechanism is assumed to be the dominant source of CPV in the quark sector, we will not discuss the SUSY contributions to flavour-dependent CP observables. MSSM-like phenomena (e.g. CP violation in the B -mesons) will also not be discussed here, as we will focus on the implications of CP violation for the NMSSM Higgs sector.

Explicit CPV at tree level in the \mathbb{Z}_3 -invariant NMSSM

In the NMSSM with a scale invariant superpotential (2.6), the parameters λ , κ , A_λ and A_κ can be complex. The associated phases only appear in a subset of terms of V_{Higgs} (2.9),

$$V_{\text{Higgs}}^{\text{phase}} = -\lambda\kappa^* H_u^0 H_d^0 S^{*2} - \lambda A_\lambda H_u^0 H_d^0 S + \frac{1}{3}\kappa A_\kappa S^3 + \text{h.c.} \quad (8.5)$$

By a redefinition of H_u , H_d and S , only one physical phase remains [399,400] which can be taken to be the phase of $\lambda\kappa^*$:

$$\phi \equiv \arg(\lambda\kappa^*) \quad (8.6)$$

The stationary conditions for the phases induce necessarily complex vevs (θ , $\delta \neq 0$) provided $\phi \neq 0$, π [400].

Regarding the tree level Higgs mass matrix (8.1), the diagonal blocks \mathcal{M}_S^2 and \mathcal{M}_P^2 are those already given in (2.22, 2.24), once the appropriate redefinitions have been performed:

$$\lambda A_\lambda \rightarrow \lambda A_\lambda \cos \theta, \quad \kappa \lambda \rightarrow \kappa \lambda \cos(\phi + \theta - \delta), \quad \kappa A_\kappa \rightarrow \kappa A_\kappa \cos \delta \quad (8.7)$$

The entries of the new off-diagonal block \mathcal{M}_{SP}^2 in (8.1) are all proportional to $\sin(\phi + \theta - \delta)$, and can be found in [399–403]. However, it is important to stress that, at tree level, explicit CP violation only induces scalar-pseudoscalar mixings between $H_{u,d}$ and S ; the scalar and pseudoscalar components of the Higgs doublets $H_{u,d}$ do not mix.

Nevertheless the impact of explicit CP violation on low energy phenomenology is not negligible. Although CPV effects are small in limiting cases for the singlet vev ($s \ll v_u, v_d$ or $s \gg v_u, v_d$) or very large $\tan\beta$, large scalar-pseudoscalar mixings can occur in regions where $s \sim \mathcal{O}(v)$ and $\tan\beta \sim \mathcal{O}(1)$ [400]. As a consequence, the lightest Higgs mass can decrease by 10–30 GeV compared to the CP conserving case. However, avoiding an excessive contribution to the neutron EDM would require comparatively heavy gauginos and squarks with masses $\sim \mathcal{O}(\text{TeV})$.

Explicit CPV in the \mathbb{Z}_3 -invariant NMSSM with radiative corrections

In order to obtain scenarios that are still in agreement with the bounds on EDMs, while at the same time inducing a moderate amount of CP violation (as desirable for electroweak baryogenesis), one can consider NMSSM scenarios where explicit CP violation in the Higgs sector is induced only through radiative corrections to the Higgs masses and couplings [404]. Explicit CP violation in the Higgs sector of the NMSSM at the one-loop level, including radiative corrections from third generation squarks, has been studied in [402, 403, 405, 406]: due to the possibly complex soft SUSY breaking terms in the squark sector, new CPV terms will appear in the neutral and charged Higgs effective Lagrangian via these radiative corrections. (The RGEs for the parameters of the NMSSM with explicit CP violation are given in [109, 407].)

Considering only the effect of the (dominant) corrections from third generation quark-squark loops, the new phases in addition to the three tree level phases in (8.5) are ϕ_{A_t} , ϕ_{A_b} , arising from A_t , A_b , respectively [402]. The minimisation conditions (with respect to H_{iI} and S_I) reduce the five phases to three. (Note that the complex phases of the third generation quark Yukawa couplings can be reabsorbed by redefinitions of the quark fields.) An important consequence of taking these higher order effects into account is that scalar-pseudoscalar mixings between the two Higgs *doublets* (absent at tree-level) can now occur.

CP violating effects originating from the chargino sector can also play a rôle. Contributions from loops involving charged particles (W^\pm bosons, charged Higgs and charginos) were taken into account in [405]. It was found that the relative phase between the soft breaking $SU(2)$ gaugino mass M_2 and λ , ϕ_C , yields corrections to *every* element of the 5×5 neutral Higgs mass matrix proportional to $\sin\phi_C$. Depending on the other low energy NMSSM parameters, the chargino induced CP violating corrections can lead to contributions to the lightest Higgs boson mass in the range $-24 \text{ GeV} \lesssim \delta_{\text{CPV}}^{\text{x}\pm} M_{H_1} \lesssim 16 \text{ GeV}$ [405]. Including all radiative corrections, $M_{H_1, \text{max}}$ increases with $\tan\beta$ (in contrast to the case without CP violation, see Section 3.2), and the upper bound of $M_{H_1, \text{max}} \sim 150 \text{ GeV}$ is saturated for $\tan\beta \sim 30$ [408].

Explicit CPV can also have implications on the production and decay rates of Higgs bosons at colliders [403, 405]: at a linear e^+e^- collider ($\sqrt{s} = 500 \text{ GeV}$ and 1 TeV), the production cross sections for the neutral Higgs bosons in the NMSSM with explicit CPV can be much smaller than in the CP conserving case. Explicit CPV in the NMSSM can possibly be tested at the ILC with $\sqrt{s} = 500 \text{ GeV}$, where at least one of the five neutral Higgs bosons would be produced via Higgs-strahlung, WW - and ZZ -fusion with corresponding cross sections of $\sim 12, 15$ and 1.5 fb , respectively. The decays of the neutral Higgs bosons

can also differ from those of the NMSSM without CP violation in the form of modified branching ratios of decays into pairs of $s\bar{s}$ and $c\bar{c}$ quarks [408].

In the nMSSM (with tadpole terms and $\kappa = 0$, see Section 7.3), explicit (as well as spontaneous) CPV is impossible at tree level, but radiative corrections from sbottom and stop squarks can also trigger CP violation in the Higgs sector. The associated phenomenology (mass spectrum and collider prospects) is analysed in [409].

In all cases of explicit CP violation in the Higgs sector, EDMs are expected to receive potentially large contributions. However, the phase combination responsible for the EDMs can be distinct from the one responsible for Higgs mixing, which potentially offers additional sources of CP violation for electroweak baryogenesis. It has been suggested [401] that the phases can be arranged in such a way that the combination inducing the neutron EDM can be suppressed, while the ones affecting the Higgs sector are allowed to be sizable. However, this requires a substantial fine tuning between the phases of the Higgs vevs and the phases of parameters in the Lagrangian. In order to ensure that SUSY contributions to the EDMs are below the experimental bounds, the SUSY spectra (notably gaugino masses) should be of $\mathcal{O}(1 \text{ TeV})$ [399, 400, 410].

Modifying the NMSSM by promoting the phases of gaugino masses and the trilinear couplings A_i to fields allows to address the strong and the SUSY CP problems simultaneously [411]. Then, however, observable SUSY CPV requires non-minimal flavour structures beyond those associated with the Yukawa couplings.

CP violation in multi-singlet extensions of the NMSSM

In order to keep sizable CPV in the Higgs sector (contributing to baryogenesis at the electroweak scale) without excessively large induced EDMs, multi-singlet extensions of the NMSSM have been considered. In [404] the NMSSM is extended by two additional vev-less singlets (S' and S''), and explicit CPV in the form of complex couplings is allowed only between fields which have no tree level couplings to quarks or leptons. Nevertheless, through loop corrections involving the new singlets, small CP violating phases can appear for the EDMs (consistent with current bounds).

Spontaneous CP violation in the NMSSM with two singlets S and S' , taking into account radiative effects, has also been proposed [412]. Both Higgs doublets and singlets can develop complex vevs, and one now has three independent phase combinations ($\theta = \varphi_u + \varphi_d + \varphi_s + \varphi_{s'}$, $\delta = 3\varphi_s$ and $\delta' = 3\varphi_{s'}$). Contrary to the NMSSM (with either spontaneous or explicit CPV), the addition of the second singlet implies that scalar-pseudoscalar mixings between the Higgs doublets can occur already at tree level. Although the Higgs spectrum still exhibits light states, it is possible to find regions in parameter space where the lightest Higgs mass is as large as 80 GeV, barely viable in view of LEP bounds. A general analysis of CP violation at the one-loop level in the Higgs sector of the MSSM extended by an arbitrary number singlet fields has been performed in [413].

8.2 R-parity violation and neutrino masses in the NMSSM

The phenomenological implications of R-parity violation (RpV) in the NMSSM are extensive (for a review of RpV in the MSSM, see [414]). Here we will briefly summarise some of the most relevant consequences, as neutrino mass generation.

R-parity is a discrete symmetry corresponding to the remnant of a group of continuous $U(1)_R$ transformations, whose charges R_p are assigned as [415]

$$R_p = (-1)^R \quad \text{with } R = \begin{cases} +1 & \text{for SM particles including all Higgs bosons} \\ -1 & \text{for the superpartners} \end{cases} \quad (8.8)$$

Historically, R-parity was introduced in early supersymmetric extensions of the SM in order to maintain the baryon (B) and lepton (L) number conservation laws of the SM [13, 18, 415–417]: conserved R-parity naturally leads to conserved B and L , while RpV requires a violation of at least B or L . If R-parity is conserved, R-odd particles are necessarily pair produced, the lightest R-odd particle (the LSP) is stable and, if neutral and colourless, a good dark matter candidate.

In the MSSM, the most general renormalisable superpotential including RpV terms is

$$W_{\text{MSSM}} = W_{\text{RpC}} + \mu_i \widehat{H}_u \cdot \widehat{L}_i + \frac{1}{2} \lambda_{ijk} \widehat{L}_i \cdot \widehat{L}_j \widehat{E}_{Rk}^c + \lambda'_{ijk} \widehat{L}_i \cdot \widehat{Q}_j \widehat{D}_{Rj}^c + \frac{1}{2} \lambda''_{ijk} \widehat{D}_i \cdot \widehat{D}_j \widehat{U}_{Rk} \ , \quad (8.9)$$

where W_{RpC} denotes the R-parity conserving part, and $i, j, k = 1, 2, 3$ are generation indices. The terms $\sim \mu_i$, $\sim \lambda_{ijk}$ and $\sim \lambda'_{ijk}$ violate L , and the term $\sim \lambda''_{ijk}$ violates B . (RpV can also occur spontaneously through the vev of a neutral R-odd scalar, necessarily a sneutrino in the MSSM or NMSSM, which also violates L .) Note that the four superfields \widehat{H}_d and \widehat{L}_i have the same gauge quantum numbers, and in the presence of the bilinear terms μ_i and μ it is no longer possible to distinguish them unambiguously.

If B and L are violated simultaneously, this can lead to fast proton decay (and other nuclear conversions). Generally, RpV can induce single production of neutralinos and charginos, LSP decays into SM particles, flavour violating Z couplings to leptons, violation of charged current universality in the lepton and quark sectors, additional sources of mixing (and CP violation, if the couplings are complex) in neutral mesons, new rare decays as lepton flavour violation (LFV) in the charged lepton sector. The baryon and/or lepton number violation induced by RpV can be an ingredient useful for baryogenesis.

One of the most desirable effects of the L number violation stemming from both bilinear and trilinear R-parity violating couplings (if the vevs of the sneutrinos are non-vanishing) is the mixing between Higgs bosons H_d and sleptons L_i (in the scalar sector) and between leptons and neutralinos/charginos. The latter implies enlarged chargino and neutralino mass matrices (respectively 5×5 and 7×7 , in the MSSM). These new mixtures allow to obtain massive neutrinos without the inclusion of right-handed neutrinos, and open the possibility of explaining neutrino data (mass squared differences and mixing angles) with a minimal superfield content. Bilinear R-parity violating couplings can induce, at the tree level, a single massive neutrino via a “seesaw-like” mechanism, where much heavier gauginos and higgsinos play the rôle of right-handed neutrinos. In order to account for realistic neutrino masses and mixings, higher order contributions (1-loop), arising from trilinear λ_{ijk} , λ'_{ijk} and/or bilinear R-parity violating couplings, must be taken into account.

In addition to the MSSM RpV terms, the NMSSM superpotential can contain a trilinear coupling

$$W_{\text{NMSSM}}^{\text{RpV}} = \lambda_i \widehat{S} \widehat{H}_u \cdot \widehat{L}_i, \quad (8.10)$$

and a corresponding contribution $\sim A_{\lambda_i} S H_u \widetilde{L}_i$ to the soft SUSY breaking Lagrangian. As in the MSSM with R-parity violating μ_i terms, the couplings λ_i in (8.10) induce numerous mixings (see e.g. [418]): the charged Higgs bosons mix with charged sleptons, and charginos with charged sleptons, leading to enlarged mass matrices. The neutral CP-even Higgs bosons mix with (left-handed) sneutrinos, now leading to 6×6 mass matrices. In the NMSSM one finds 8 neutral fermions resulting from the mixing of (left-handed) neutrinos and neutralinos.

Note that in the presence of bilinear RpV soft breaking terms, electroweak symmetry breaking generally leads to non-vanishing sneutrino vevs, and these have to be taken into account in the conditions for vacuum stability (UFB directions and CCB minima) in this class of models [419]. The RG equations of the R-parity violating couplings in the NMSSM have been given (and analysed for fixed points) in [420, 421], and all possible baryon and lepton number violating operators are classified in [421].

A common phenomenological feature of all SUSY models violating R-parity is the occurrence of an unstable LSP. The lightest neutralino (or the lightest stau) can decay via its R-parity violating couplings into SM particles. An unstable LSP can still remain a viable dark matter candidate provided its lifetime is sufficiently long (of the order of the age of the universe). This possibility has been investigated recently in the NMSSM – including constraints from neutrino data – in [422] with the result that a gravitino LSP can be sufficiently long-lived. If the LSP is short-lived (hence irrelevant to the solution of the dark matter problem), it must decay sufficiently fast as not to affect the successful predictions of Big-Bang nucleosynthesis. In both cases severe constraints on the R-parity violating couplings can be derived [414].

At colliders, one can expect displaced vertices from LSP decays [414]. Its branching fractions (depending on its composition) can shed light on the R-parity violating couplings: if the LSP is a singlino dominated neutralino, it would be possible to distinguish the R-parity violating NMSSM from the R-parity violating MSSM at future colliders [423].

Let us review some phenomenological implications of RpV, including the possibility of adding right-handed neutrino superfields to the NMSSM, and mechanisms of neutrino mass generation with and without RpV.

8.2.1 Massive neutrinos in the NMSSM

Neutrino masses from R-parity violation

Dirac neutrino-higgsino masses, which do not require the introduction of right-handed neutrino fields, can be generated from the R-parity violating MSSM-like first term $\sim \mu_i$ in (8.9). However, the resulting neutrino mass matrix is only of rank 1 at tree level, i. e. only one neutrino is massive, which must be improved by loop effects. Furthermore the smallness of the neutrino masses must be understood, which requires some fine tuning of parameters.

Adding only the terms $\sim \mu_i$ in (8.9) to the NMSSM, the required amount of fine tuning can be reduced due to possible compensations between gaugino and singlino exchange diagrams (at tree level) [424]. Such a compensation is equally possible if only the NMSSM specific terms (8.10) are present [418,424]. Once loop corrections are included, a hierarchical spectrum and mixing angles in agreement with observation can be obtained (albeit still at the expense of some fine tuning) [418]. Natural values for the heaviest neutrino mass lead to bounds on $\lambda_i \sim \mathcal{O}(10^{-5} - 10^{-2})$ [418,424].

If both terms $\sim \mu_i$ in (8.9) *and* those of (8.10) are present, two massive neutrino states can be obtained at tree-level (in contrast to the MSSM) [424,425]. This offers the appealing possibility of reproducing neutrino data without having to resort to loop effects. However, again the observed mass hierarchy and mixing angles can be obtained only at the price of substantial fine tuning of the parameters: although neutrino data by itself would allow for ratios $|\mu_i/\mu_{\text{eff}}|$ as large as $\sim 10^{-1}$, potential contributions to LFV processes (such as $\mu \rightarrow e\gamma$ or $\mu \rightarrow eee$) constrain the values of $|\mu_i/\mu_{\text{eff}}| \lesssim 10^{-2}$.

In [424] it has been pointed out that the NMSSM singlet can generate thermal leptogenesis (which would later be converted into a baryon asymmetry of the universe) via decays involving the R-parity violating trilinear couplings. The neutrino refraction indices (relevant for the neutrino propagation in matter, as in supernovae) have been studied in the NMSSM, including radiative corrections, in [426].

Right-handed neutrino sector

If one adds three generations of right-handed neutrino superfields \widehat{N}_{Ri} (with, by convention, negative R-parity of its scalar components in contrast to the singlet superfield), the superpotential can include the following additional terms:

$$W_{\text{NMSSM}}^{N_R^c} = h_{\nu ij} \widehat{L}_i \cdot \widehat{H}_u \widehat{N}_{Rj}^c + \frac{1}{2} M_{ij}^{\text{Maj}} \widehat{N}_{Ri}^c \widehat{N}_{Rj}^c + \frac{\kappa_i}{3} \widehat{N}_{Ri}^{c3} + \frac{1}{2} \lambda_{ij}^{\text{Maj}} \widehat{N}_{Ri}^c \widehat{N}_{Rj}^c \widehat{S} \quad (8.11)$$

and corresponding contributions to the soft SUSY breaking Lagrangian. The first two R-parity conserving terms are identical to those present in the MSSM with right-handed neutrinos, the third term $\sim \kappa_i$ is R-parity violating, and only the last R-parity conserving term is specific for the NMSSM. The fermion and scalar components of \widehat{N}_{Ri} will mix in general with all neutral fermions and scalars.

Models with right-handed neutrino superfields \widehat{N}_R and a scale and R-invariant superpotential ($M^{\text{Maj}} = 0$ and $\kappa_i = 0$ in (8.11)) allow for the spontaneous generation of mass terms for \widehat{N}_R through the last term $\sim \lambda^{\text{Maj}}$ in (8.11): when the singlet gets a vev s , a Majorana mass $M_{\text{eff}}^{\text{Maj}} = \lambda^{\text{Maj}} s$ (preferably of $\mathcal{O}(\text{TeV})$) is dynamically generated [427]. These models have been shown to possess regions in parameter space where left- and right-handed sneutrinos acquire vevs, so that R-parity is spontaneously broken and effective μ_i^{eff} -terms in (8.9) are generated as well. In this case, the model contains two distinct sources for neutrino mass generation (the ordinary seesaw with right-handed neutrinos and contributions from the μ_i^{eff} -terms) and both solar and atmospheric neutrino data can be accommodated.

Solving the μ -problem with right-handed neutrino superfields and RpV

From the point of view of particle content, an economic and interesting possibility is to replace the singlet \widehat{S} by the right-handed neutrino superfields \widehat{N}_{Ri} (which is not

compatible with an R-invariant superpotential). This class of models is also denoted as the $\mu\nu$ SSM [428]. In addition to quark and lepton Yukawa couplings – including the first term in (8.11) – the superpotential of the $\mu\nu$ SSM contains the following terms:

$$W = \tilde{\lambda}_i \hat{N}_{Ri}^c \hat{H}_u \cdot \hat{H}_d + \frac{\kappa_i}{3} \hat{N}_{Ri}^{c3}. \quad (8.12)$$

Then the vevs $\tilde{\nu}_{Ri}$ of the scalar components of \hat{N}_{Ri} generate an effective μ -term ($\mu_{\text{eff}} = \sum_i \tilde{\lambda}_i \tilde{\nu}_{Ri}$), and the Higgs vevs as well as the vevs $\tilde{\nu}_{Ri}$ generate both Dirac and Majorana masses for the neutrinos. The observed neutrino masses originate from two different sources: from the Yukawa couplings $h_{\nu ij}$ in (8.11) roughly of $\mathcal{O}(10^{-6})$ (and a seesaw mechanism with right-handed neutrino Majorana masses of the order of the electroweak scale from the terms $\sim \kappa_i$), and an R-parity violating mixing of neutrinos with neutralinos. (In an earlier study of the effects of a coupling $\sim \hat{N}_{Ri}^c \hat{H}_u \hat{H}_d$ [429], an explicit Majorana mass term for the right-handed neutrinos of the order of the GUT scale was added, and cosmological constraints on the lifetime of the neutralino LSP – unstable due to the R-parity violating couplings – were derived.)

The phenomenology of this class of models has been extensively studied in [423, 428, 430–432]; in large regions in parameter space, viable solutions can be found (avoiding Landau poles, and in agreement with collider constraints on Higgs and sparticle masses, and current neutrino data). In spite of the additional Higgs-sneutrino mixings, the upper bound on the lightest Higgs mass turns out to be similar to the one obtained in the framework of the general NMSSM ($M_{H_1} \lesssim 140$ GeV) [430]. Regarding the neutrino masses, both normal and inverted hierarchies can be obtained, and two large and one small mixing angle can be generated [423] even with a simple flavour diagonal structure for the neutrino Yukawa couplings. In [432] it was shown that bimaximal and tri-bimaximal neutrino mixing could be accommodated with diagonal and degenerated Yukawas for μ - and τ -neutrinos, with degenerated vevs for the left-handed sneutrinos. Allowing for complex Higgs and sneutrino vevs, spontaneous CP violation can occur in the lepton sector, leading to non-vanishing phases in the lepton mixing matrix.

R-parity conserving NMSSM with right-handed neutrinos

“Standard” seesaw mechanisms (type I, II and III) can be also incorporated in the R-parity conserving NMSSM, in which case additional higher dimensional operators (of dimension 6 or 7) can arise from integrating out heavy states [433]. Even with a TeV-scale seesaw mechanism, these can have sizable couplings to SM leptons, and may be detected at the LHC. Within minimal left-right extensions of the NMSSM one can aim at a simultaneous explanation of the smallness of neutrino masses via the seesaw mechanism, and a solution of the tachyon slepton problem arising in Anomaly Mediated SUSY breaking models [434].

Within the R-parity conserving NMSSM, the vev of the singlet can be the only source for non-vanishing Majorana masses for the right-handed neutrinos [435, 436] via the last term in (8.11). Remarkably, a right-handed sneutrino is a viable dark matter candidate in such a model (see also Section 9.3).

8.3 $U(1)'$ -extensions of the NMSSM

Extensions of the SM gauge group by one (or several) $U(1)'$ gauge symmetries can arise naturally from GUTs (see below), string-inspired constructions [437–440], solutions of the μ problem in GMSB [357] etc.; for recent reviews and a more complete list of references see [183, 441–443].

The simplest class of $U(1)'$ extended SUSY models involves one additional SM singlet S , charged under the additional $U(1)'$, whose vev is responsible for the $U(1)'$ breaking. The $U(1)'$ charges of the matter and $SU(2)$ doublet Higgs supermultiplets are not unique, but typically bilinear Higgs couplings (as the μ -term) are forbidden whereas a $\lambda\widehat{S}\widehat{H}_u\widehat{H}_d$ term is allowed, in which case $U(1)'$ models are similar to the NMSSM (sometimes they are denoted as the UMSSM). The term $\sim\kappa\widehat{S}^3$ in the superpotential of the NMSSM (which stabilises the potential for $s\rightarrow\infty$) is no longer possible, but its rôle is now played by the $U(1)'$ D -terms. Note that the Peccei-Quinn symmetry, that appears in the NMSSM in the absence of the cubic singlet term, is embedded in the gauged $U(1)'$ so that the would-be Peccei-Quinn axion is eaten by the Z' . Also the NMSSM domain wall problem is avoided, since the discrete \mathbb{Z}_3 -symmetry is embedded in the continuous $U(1)'$ as well.

The singlet vev generates simultaneously an effective μ -term (which should not be too large in order to avoid fine tuning, see Section 6.4) and the mass of the new Z' boson. Limits on $M_{Z'}$ (from direct searches at the Tevatron) are model-dependent, especially on the Z' couplings to quarks and leptons (see [89, 443] and references therein), and are typically $M_{Z'}\gtrsim 600-900$ GeV for a Z' that decays into SM fermions.

Loops from matter, charged under both the SM and the $U(1)'$ gauge groups, generate a $Z-Z'$ mixing mass term parametrized by the angle $\alpha_{ZZ'}$, which is constrained by electroweak precision data to be smaller than $\mathcal{O}(10^{-3})$, the exact value depending again on the chosen $U(1)'$ charges. Both constraints from $M_{Z'}$ and $\alpha_{ZZ'}$ require a quite large singlet vev $s\gtrsim\mathcal{O}(1\text{ TeV})$, and hence a sufficiently negative (and large) value of m_S^2 (see, e. g., [444]).

All possible $U(1)'$ charge assignments allowing for a $\lambda\widehat{S}\widehat{H}_u\widehat{H}_d$ term imply at least mixed anomalies between the $U(1)'$ and the SM symmetry groups. Their cancellation requires the introduction of new exotic fermions (hence superfields) which are vector-like with respect to the SM, but chiral under $U(1)'$ (see, e. g., [440, 445–448]). Then, unification of the gauge couplings implies the introduction of additional exotics which are charged, but non-chiral under both the SM and $U(1)'$ gauge symmetries. These new states do not contribute to the anomalies, but do affect the RGEs for the gauge couplings.

In what follows we briefly review some aspects of NMSSM-like models with one SM singlet S . Then we refer to some features of more general $U(1)'$ extended models.

The most relevant modifications of the Higgs potential with respect to the \mathbb{Z}_3 -invariant NMSSM are the absence of all terms $\sim\kappa$, but the presence of a new D term in the potential of the form

$$V_D^{U(1)'} = \frac{1}{2}g_{Z'}^2 (Q_S|S|^2 + Q_{H_d}|H_d|^2 + Q_{H_u}|H_u|^2 + \text{exotics})^2, \quad (8.13)$$

where $g_{Z'}$ is the $U(1)'$ gauge coupling, and Q_i are the $U(1)'$ charges which obey $Q_S + Q_{H_d} + Q_{H_u} = 0$ in order to allow for the term $\lambda\widehat{S}\widehat{H}_u\widehat{H}_d$ in the superpotential.

After symmetry breaking, the Higgs spectrum consists of a pair of charged Higgs bosons, three neutral CP-even scalars and only one pseudoscalar. A detailed overview of the Higgs mass matrices for various models, including the dominant top-stop one-loop corrections, can be found in [441].

Compared to the NMSSM, theoretically and phenomenologically allowed Higgs mass ranges have to be reconsidered: due to the additional quartic Higgs self-couplings from the D terms in (8.13), Higgs masses are generally larger; in particular, LEP bounds on Higgs bosons with SM-like couplings to the Z boson are easier to satisfy. The theoretical upper bound $M_{H_1\max}$ on the mass of the lightest neutral Higgs boson increases to $M_{H_1\max} \sim 170$ GeV [449, 450]. Radiative corrections from top-stop loops have been included in [449, 451, 452], as well as constraints from gauge coupling unification in [453].

Recalling that constraints from $M_{Z'}$ and $\alpha_{ZZ'}$ favour large values of s , the Higgs spectrum is generally similar to the MSSM with a large μ -term [440]: H_1 is SM-like, the heavy pseudoscalar is approximately degenerate with H_2 and H^\pm , completing a full $SU(2)$ doublet. The mass of the heaviest singlet-like neutral Higgs is typically of the order of $M_{Z'}$. Then, the lower bound on the lightest CP even Higgs mass implied by LEP constraints turns out to be in general similar to the one of the MSSM ($M_{H_1} \gtrsim 90$ GeV).

However, in particular cases one can obtain different configurations for the Higgs spectrum as large doublet-singlet mixing, light singlet dominated states, etc. [443, 454, 455]. In [456] a scenario with two light CP-even Higgs bosons has been proposed in order to explain the (light) excesses of events in Higgs searches at LEP (see Section 5). Note that in the absence of a light pseudoscalar one cannot relax the lower LEP bound on a doublet-like Higgs allowing for its decays into a pair of light pseudoscalars as in the NMSSM, see Section 5.1.

The neutralino sector of the NMSSM is extended by the Z' gaugino \tilde{Z}' with a soft SUSY breaking mass $M_{\tilde{Z}'}$, and possible $\tilde{B} - \tilde{Z}'$ mixing terms [457]. A detailed discussion of the 6×6 neutralino mixing matrix can be found in [378, 458–460]. In the limit $g_{Z'} s \gg M_{\tilde{Z}'}$, the singlino and the Z' gaugino mix to form an approximate Dirac fermion with mass $\sim M_{Z'}$, which mixes very little with the other four neutralinos. Conversely, in the large $M_{\tilde{Z}'}$ regime one has a very heavy Majorana fermion \tilde{Z}' , and a much lighter singlino with a “seesaw” mass $\sim M_{Z'}^2/M_{\tilde{Z}'}$. In all cases the new states will likely affect sparticle decay chains.

CPV has also been considered in $U(1)'$ extended models [183, 408, 461, 462]. However, at the tree level the Higgs sector is CP conserving, and spontaneous CPV (after including radiative corrections) cannot be realised in the allowed parameter space [408]. Explicit CPV in the Higgs sector can be possible via radiative corrections, through the explicit CPV phases present in the stop mass matrix or the additional phases in the exotic quark sector of the model. These can significantly alter the mixing between the pseudoscalar and the heaviest scalar [462]. Due to the extended neutralino sector, phases in λ and in the slepton trilinear couplings can contribute to the electron EDM [461]. EDM bounds strongly constrain the effective phase in the chargino mass matrix, but all other SUSY phases remain largely unconstrained, thus alleviating the SUSY CP problem [462].

$U(1)'$ extended models have also important cosmological implications, since the extra states can modify the nature of the LSP. Early studies [463] suggested that Z' mediated neutralino annihilation could provide important (or even dominant) contributions to the

correct relic density. More recent analyses [183, 378, 464–467] (for a general discussion, see [468] and references therein) have shown that compatibility with the WMAP bounds on the relic density can be achieved in the case of a singlino-like LSP, with a small higgsino component, via annihilation into a Z' ; enhanced couplings to the Z also allow annihilation via a Z resonance [464]. The prospects for observation of the LSP in dark matter direct detection experiments have been considered in [467]. These models offer also non-neutralino dark matter candidates: scalar right-handed sneutrinos can be viable dark matter candidates due to the possibility of annihilation through the Z' [469]; other possibilities include exotic LSPs [470], see also Section 9.3.

Dirac neutrino masses in the experimentally preferred range can also be generated within $U(1)'$ -extended generalized SUSY models [471–474]. If the $U(1)'$ symmetry forbids the Yukawa couplings to right-handed neutrinos in the superpotential, effective Dirac masses can be generated from (hard) SUSY breaking Yukawa couplings or, at the loop level, from non-holomorphic trilinear couplings to the “wrong” Higgs boson, which are naturally small [473]. SUSY models of sterile neutrinos can be naturally realised in $U(1)'$ extended models [472]: sterile neutrino masses of $\mathcal{O}(1\text{eV})$ and mixings among the active and sterile neutrinos can then be obtained via the high-dimensional operators, generated by integrating out the heavy fields. In $U(1)'$ extended models where the vev s generates Dirac neutrino masses through a dimension-4 term in the superpotential, the decays of the LSP and NLSP (the two lightest sneutrinos) could explain the PAMELA anomaly [475].

We recall that, once R-parity is violated, proton decay can be excessively fast since both baryon number and lepton number are violated in general. In $U(1)'$ models with RpV, the proton can still be sufficiently stable in a natural way, since the $U(1)'$ symmetry can forbid the renormalisable (dimension 3 and 4) and the most dangerous dimension 5 operators [445]. In [448] models are discussed which can still contain either lepton number violating or baryon number violating renormalisable interactions, but never simultaneously both. In [476] models have been put forward where bilinear lepton number violating terms in the superpotential allow to accommodate the observed pattern of neutrino masses and mixings, but the most dangerous dimension 3 and 4 baryon number violating operators (and, possibly, all $\Delta B = 1$ operators) are absent.

Production (and decays) of the additional Z' gauge boson and exotic states at colliders are reviewed in [443, 477]. The exotic quarks and squarks could appear in decays of the Z' , or may manifest themselves indirectly, since the production of the heaviest Higgs boson (via gluon fusion) can be dominantly mediated by loops of these exotics [452]. The production of neutral Higgs bosons in this class of models has been discussed in [478].

At e^+e^- colliders of higher energy, the Higgs sectors of $U(1)'$ extended models and the NMSSM can be distinguished in the limit of large trilinear Higgs couplings [199]. At the ILC, measurements of the $H_i^0 Z Z$ couplings may help to distinguish scenarios with a light, leptophobic Z' from the NMSSM, since the couplings are expected to be substantially smaller in the former case [479]. In the case of TeV Z' bosons, searches for Z' decays into higgsinos can also test if a gauge symmetry is indeed responsible for the absence of a bare μ -term in the superpotential [480].

There are innumerable possibilities for the construction of more general $U(1)'$ -extensions

of the MSSM including, for instance, flavour symmetries [481]. A systematic analysis and classification can be found in [445]. In what follows we consider some particular cases, some of which are motivated by GUTs or string-inspired constructions.

$U(1)'$ and a secluded singlet sector

A possible way to avoid the tension arising from a small enough μ_{eff} and a sufficiently large $M_{Z'}$ is to consider the secluded $U(1)'$ -extended MSSM (sMSSM), where a separation between μ_{eff} and $M_{Z'}$ is achieved by the introduction of four SM singlets S , S_1 , S_2 and S_3 [454]. The ordinary sector contains the NMSSM (or USSM) singlet S , while the three additional singlets S_i belong to a secluded sector which couples to the ordinary sector only via $U(1)'$ gauge and possibly soft SUSY breaking terms. All singlets are charged under $U(1)'$ such that their vevs contribute to $M_{Z'}$, but only S generates a μ_{eff} of the order of the electroweak scale. The SM singlet-dependent part of the superpotential is given by

$$W_H^{U(1)} = \lambda \widehat{S} \widehat{H}_d \widehat{H}_u + \lambda_S \widehat{S}_1 \widehat{S}_2 \widehat{S}_3. \quad (8.14)$$

The soft SUSY breaking Lagrangian contains soft SUSY breaking masses and trilinear couplings for the new singlets ($m_{S_i}^2$, $m_{S_i S_j}^2$, A_{λ_S}). Unwanted global symmetries have to be broken explicitly to avoid the potential occurrence of two massless experimentally excluded Goldstone bosons in the spectrum. For appropriate parameters, $S_{1,2,3}$ acquire large vevs along nearly F -flat and D -flat directions of the scalar potential so that one obtains $M_{Z'} \gg \mu_{\text{eff}}$ in a natural way. λ can be larger than in the NMSSM (without leading to a Landau singularity), since the additional gauge coupling in its β -function slows down its increase towards high scales. As a consequence, the lightest CP-even Higgs mass could be as large as ~ 174 GeV already at tree level [454]. In general the Higgs and neutralino sectors of these models are quite complicated, and we refer to [455] for a detailed discussion. A singlino-higgsino LSP can account for an acceptable cold dark matter relic density [464]. Regions in parameter space exist, that can explain the experimental deviation of the muon anomalous magnetic moment from the SM and yield an acceptable cold dark matter relic density without conflict with collider experimental constraints [482].

In the sMSSM, explicit CP violation in the Higgs sector can be induced at the tree level by the non-zero phase of $m_{S_1 S_2}^2$ [483]. Large values of the latter phase can allow for a neutral Higgs boson lighter than ~ 90 GeV, compatible with LEP constraints due to suppressed couplings to the Z boson. Unlike the MSSM and NMSSM, SCPV can occur at the tree level. The possibility of electroweak baryogenesis in the sMSSM has been discussed in [182].

Additional $U(1)'$ in string-inspired E_6

As mentioned before, gauge extensions of the SM gauge group by one (or several) non-anomalous $U(1)'$ gauge symmetries can naturally arise from a (string-inspired) GUT gauge group as E_6 (see, e. g., [484–486]).

In an E_6 GUT, the matter sector includes three 27-plets containing the three ordinary families of quarks and leptons (including right-handed neutrinos), three families of candidate Higgs doublets (H_{di} , H_{ui}) and singlets (S_i), and three families of extra colour triplets

(D'_i, \bar{D}'_i) . Anomalies are cancelled generation by generation within each complete 27 representation. Gauge coupling unification requires the addition of a further pair of Higgs-like multiplets, H' and \bar{H}' , arising from incomplete $27' + \bar{27}'$ representations. The breaking of E_6 occurs via

$$E_6 \rightarrow SO(10) \times U(1)_\psi \rightarrow SU(5) \times U(1)_\chi \times U(1)_\psi. \quad (8.15)$$

The low energy gauge group is assumed to be $SU(3) \times SU(2) \times U(1)_Y \times U(1)'$, where $U(1)' = U(1)_\chi$ or a specific linear combination of $U(1)_\chi$ and $U(1)_\psi$ motivated by the breaking of E_6 by Wilson lines.

In [484, 485], patterns of $U(1)'$ and electroweak symmetry breaking were studied in this class of E_6 models. E_6 -type relations between the Yukawa couplings are not imposed, and only those conserving baryon and lepton number are kept. Then, the most general superpotential allowed by gauge invariance is of the form [485]

$$W_H^{E_6} = h_{ijk}^u \hat{U}_i^c \hat{Q}_j \hat{H}_{uk} + h_{ijk}^d \hat{D}_i^c \hat{Q}_j \hat{H}_{dk} + \lambda_{ijk} \hat{S}_i \hat{H}_{dj} \hat{H}_{uk} + h_{ijk}^D \hat{S}_i \hat{D}_j^c \hat{D}_k^c, \quad (8.16)$$

where i, j, k are family indices. When some of the Higgs fields acquire vevs, electroweak and $U(1)'$ symmetries are broken. It is always possible to work in a basis where only one family of Higgs doublets and singlets develops a vev: one can define $S = S_3$, $H_d = H_{d3}$ and $H_u = H_{u3}$, while the remaining families are treated as additional exotic doublets and singlets without vevs. In order to obtain an acceptable low energy phenomenology, universality at the GUT scale of the soft SUSY breaking masses has to be relaxed. (Universal boundary conditions for the soft breaking terms could be accommodated by identifying $U(1)'$ with a more general linear combination of $U(1)_\chi$ and $U(1)_\psi$ [485].)

Another E_6 inspired version exhibiting an NMSSM-like low energy spectrum and allowing for explicit CP violation in the Higgs sector has been considered in [487]. The electroweak gauge symmetry of the SM is enlarged to $SU(2) \times U(1) \times U(1)_1 \times U(1)_2$, where both extra $U(1)$ symmetries originate from E_6 . The Higgs sector contains two doublets and two singlets (only one being allowed to couple to the doublets due to the underlying E_6 symmetry). Due to top-stop mediated radiative corrections the model allows for explicit CP violation in the Higgs sector at the one-loop level, leading to scalar-pseudoscalar mixings among the five neutral Higgs bosons.

The Exceptional SSM

Specific realisations of E_6 -inspired $U(1)'$ extensions offer the possibility of incorporating a seesaw mechanism (to explain the smallness of light neutrino masses) in a natural way. To this end the extra $U(1)'$ surviving at low energy (denoted by $U(1)_N$) must correspond to a particular combination of $U(1)_\chi \times U(1)_\psi$ in (8.15) under which the right-handed neutrinos (also contained in the 27 representation) transform trivially:

$$U(1)_N \equiv U(1)' = U(1)_\chi \cos \theta_N + U(1)_\psi \sin \theta_N \quad \text{with} \quad \theta_N = \arctan \sqrt{15}. \quad (8.17)$$

This is the so-called Exceptional SSM (E_6 SSM, see [488–495] and references therein).

As discussed in [488], due to the vanishing charges under $U(1)_N$, right-handed neutrinos can acquire very heavy Majorana masses suitable for the “standard” seesaw mechanism. This also opens the possibility of baryogenesis from leptogenesis, and avoids constraints on

the mass of the Z' arising from nucleosynthesis. Although $B - L$ is automatically conserved in E_6 -inspired SUSY models, some of the new Yukawa interactions could violate baryon number resulting in rapid proton decay. Proton stability, successful leptogenesis, non-zero neutrino masses and the absence of exotic coloured dark matter relics can be guaranteed with the help of a new conserved R-parity-like quantum number and corresponding baryon and/or lepton number assignments for the exotic particles. Furthermore, it is mandatory to impose a hierarchical structure for the additional Yukawa couplings as well as a discrete \mathbb{Z}_2 symmetry as to avoid excessive flavour changing neutral currents.

Due to the modified (two-loop) beta functions in the E_6 SSM, unification of the gauge couplings can be achieved for values of $\alpha_s(M_Z)$ closer to the measured central value than in the MSSM [490].

The E_6 SSM Higgs sector includes three CP-even, one CP-odd and two charged states. In the MSSM limit $\lambda \ll 1$, the new states become very heavy and decouple, rendering the model essentially indistinguishable from the MSSM. The analysis of RG flow of the gauge and Yukawa couplings shows that the absence of a Landau singularity allows for larger λ and lower $\tan\beta$ at the electroweak scale than in the MSSM and NMSSM, which affects the Higgs, neutralino and chargino spectra. When $\lambda \gtrsim g_1$, the lightest Higgs scalar can be somewhat heavier than in the NMSSM; including two-loop corrections, the maximally allowed value is around 150 GeV. Vacuum stability constraints push the masses of the heaviest CP-even, CP-odd and charged states above 1 TeV, so that only the lightest scalar would be within reach of the LHC and an ILC [489].

The superpartners of the Z' and the singlet contribute to the neutralino spectrum, while the chargino sector remains unchanged. For $\lambda \gtrsim g_1$, the heaviest neutralino and chargino states are the neutral and charged $H_{u,d}$ higgsinos; the lightest chargino is wino-like, while the LSP is a bino-like neutralino [488]. The dark matter prospects for this class models have been addressed in [491] (see Section 9.3).

The many new additional exotics form vector-like multiplets which could possibly be produced and detected at the LHC; the new Z' could be visible if lighter than 4-5 TeV. Within the constrained E_6 SSM (assuming universal soft SUSY breaking parameters at high energy) a detailed study of the phenomenology and impact for LHC discovery has been carried out in [492, 493]. Scenarios for early LHC discovery were investigated in the low $(m_0, M_{1/2})$ regime, and a set of benchmark points was proposed.

E_6 models based on a Δ_{27} family symmetry broken close to the GUT scale allow in addition to address the flavour problem: quark and lepton masses and mixing angles can be accounted for, with tri-bimaximal neutrino mixing [494, 495].

9 Dark Matter in the NMSSM

As in the MSSM, the lightest neutralino in the NMSSM is a candidate for cold dark matter (DM) in the form of WIMPs (weakly interacting massive particles), whose relic density should be in agreement with the constraints from WMAP, $0.094 \lesssim \Omega h^2 \lesssim 0.136$ at the 2σ level [343, 344]. (For a review of DM in the MSSM see [496].) Due to the differences in the neutralino and Higgs sectors, the properties of the LSP in the NMSSM can be significantly distinct from those in the MSSM, both in its nature and/or in the processes relevant for the LSP relic density and its detection: the LSP can be dominantly singlino-like, and/or the additional Higgs bosons can contribute to the LSP annihilation and detection processes. A neutralino LSP in the form of a higgsino-singlino mixed state in the NMSSM, giving rise to the required dark matter density, was considered first in [335, 497–500], where often universality constraints on the soft SUSY breaking terms were imposed.

The large number of processes relevant for the computation of the LSP relic density (annihilation and coannihilation cross sections) and of LSP detection in the NMSSM will not be discussed here in detail, but all of them are included in the code MicrOMEGAS (see Appendix D), and described in [501–503].

Subsequently we discuss some aspects of the LSP relic density and its detection in the NMSSM which differ from the MSSM: in Section 9.1, we discuss the processes relevant for the dark matter relic density; in Section 9.2 we consider the detection of an NMSSM dark matter candidate; finally, we will address specific NMSSM scenarios for DM in Section 9.3.

9.1 Neutralino relic density

The basic mechanisms that lead to a reduction of the neutralino relic density are essentially the same as in the MSSM: neutralino annihilation via s - and/or t -channel exchanges into gauge boson, Higgs boson and fermion pairs, and coannihilation of the neutralino with a heavier state (typically the NLSP). The cross section for these mechanisms can be strongly enhanced by s -channel resonances, occurring when $m_{\text{LSP}} + m_{(\text{N})\text{LSP}}$ is close to the mass of the particle exchanged in the s -channel.

Distinctive NMSSM scenarios can be manifest in several ways, depending on the regions in the NMSSM parameter space: the singlino component of the LSP can be sizable (e.g. mixed higgsino-singlino LSP), and one can even have (nearly) pure singlino LSPs. Secondly, the extra scalar and pseudoscalar Higgs bosons can lead to rapid LSP annihilation through s -channel Higgs resonances. Moreover, in the case where the additional scalar/pseudoscalars are light, new final states can be kinematically allowed: for example, annihilation into $Z H_1$, $H_1 H_1$, $H_1 A_1$ and $A_1 A_1$ can significantly contribute to the reduction of the neutralino relic density, either via s -channel Z , H_1 , A_1 , or t -channel (heavier) neutralino exchange. Here the NMSSM specific couplings λ and κ can play an important rôle.

A detailed discussion the viability of DM candidates in the (unconstrained) NMSSM, using recent cosmological constraints and numerical tools, is given in [501]. In what follows, we briefly discuss some illustrative examples of how the new features of the NMSSM are manifest, separately considering annihilation and coannihilation.

9.1.1 Annihilation

Here we briefly consider neutralino annihilation processes that reflect the new features of the NMSSM. For a detailed discussion, albeit in the framework of the MSSM, we refer to [496]. Detailed formulae for (MSSM) annihilation cross sections can be found, for example, in [504].

- Annihilation via s -channel Higgs exchange (CP-even or CP-odd)

The annihilation cross section is proportional to (neglecting interference terms)

$$\sigma(\chi_1^0 \chi_1^0 \xrightarrow{H_a} X X') \propto \left| \sum_a \frac{(H_a \chi_1^0 \chi_1^0) (H_a X X')}{s - M_{H_a}^2 + i\Gamma_{H_a} M_{H_a}} \right|^2 f_s(s, m_{\chi_1^0}^2, m_X^2, m_{X'}^2), \quad (9.1)$$

where X, X' denote SM fermions, gauge or Higgs bosons, $H_a \chi_1^0 \chi_1^0$ the Higgs-neutralino-neutralino couplings and $H_a X X'$ the Higgs couplings to the final states (see Appendix A). Γ_{H_a} is the width of H_a , and we have introduced a generic function f_s that depends on the s -channel momentum and on the masses of the initial and final states [504]. A similar expression is obtained for the exchange of a pseudoscalar (with the appropriate replacements in the couplings, masses, and different f_s).

- Annihilation via s -channel Z exchange

In this case, one approximately has

$$\sigma(\chi_1^0 \chi_1^0 \xrightarrow{Z} X X') \propto \left| \frac{(Z \chi_1^0 \chi_1^0) (Z X X')}{s - M_Z^2 + i\Gamma_Z M_Z} \right|^2 g_s(s, m_{\chi_1^0}^2, m_X^2, m_{X'}^2, M_Z^2), \quad (9.2)$$

where the SM $Z X X'$ couplings are proportional to electroweak gauge couplings, and $(Z \chi_1^0 \chi_1^0) = g_2 / (2 \cos \theta_W) (N_{13}^2 - N_{14}^2)$, with N_{13} and N_{14} (according to (A.7)) denoting the higgsino components of the lightest neutralino χ_1^0 . For the function g_s see [504].

- Annihilation via t -channel neutralino exchange

As in the MSSM, neutralino annihilation can also take place via t -channel sfermion, chargino and neutralino exchange. In the latter case, annihilation into a pair of singlet-like Higgs bosons ($H_1 H_1$, $A_1 A_1$, or $H_1 A_1$) can play a significant rôle for singlet-like LSPs [501]. As an example, the cross section for $\chi_1^0 \chi_1^0 \rightarrow H_1 A_1$ via t -channel neutralino exchange reads

$$\sigma(\chi_1^0 \chi_1^0 \xrightarrow{\chi_i^0} H_1 A_1) \propto (H_1 \chi_1^0 \chi_i^0)^2 (A_1 \chi_1^0 \chi_i^0)^2 h_s(s, m_{\chi_1^0}^2, m_{H_1}^2, m_{A_1}^2); \quad (9.3)$$

the scalar/pseudoscalar Higgs-neutralino-neutralino couplings are given in (A.14) and h_s denotes an auxiliary function encoding the mass dependence [504]. In this case, h_s decreases with the neutralino mass as $m_{\chi_1^0}^{-4}$.

- Resonant s -channel annihilation

LSP freeze-out occurs at temperatures $T \sim m_{\text{LSP}}/20$, where the LSPs are non-relativistic and $s \sim 4 m_{\text{LSP}}^2$. If neutralino annihilation proceeds via the s -channel

exchange of a particle (Higgs or gauge boson) with mass $\approx 2m_{\chi_1^0}$, the pole in the annihilation cross section can lead to a rapid decrease in the neutralino relic density. This resonance phenomenon can allow for compatibility with WMAP bounds in cases where the relic density would be otherwise too large. It is important to stress that for the case of resonant annihilation, the relic density is very sensitive to the mass differences of the LSPs and the exchanged particles. Hence, regions in the NMSSM (or MSSM) parameter space where WMAP compatibility is achieved via resonant annihilation typically correspond to narrow bands (for the NMSSM see, e.g., [346, 347, 501]).

- Coannihilation LSP-NLSP

In the MSSM and in the NMSSM scenarios exist where no efficient annihilation mechanism is available. However, the correct relic density can still be obtained by coannihilation processes. If the NLSP (typically the second lightest neutralino χ_2^0 or a sfermion) is not much heavier than the LSP, so that it is still present in the thermal plasma at the time of LSP freeze-out, coannihilation processes can efficiently reduce the LSP relic density.

In the NMSSM, coannihilations with heavier neutralinos or charginos can occur via s -channel Z , W^\pm or Higgs exchange, for example $\chi_1^0 + \chi_2^0 \rightarrow H \rightarrow X X'$, or via t -channel χ_i^0, χ_j^\pm exchange [501]. For a sfermion NLSP (usually a light, mostly right-handed stau), s -channel and t -channel (the latter via neutralino or chargino exchange) coannihilation lead to fermion final states. Just like in the case of s -channel LSP-LSP annihilation, resonances can also increase the LSP-NLSP annihilation cross section: in this case the mass of the exchanged particle should be $\approx m_{\text{LSP}} + m_{\text{NLSP}}$.

- NLSP-NLSP annihilation

This process, usually also labeled coannihilation, occurs when the dominant annihilation process of R-odd particles is via NLSP annihilation: $\text{NLSP} + \text{NLSP} \rightarrow X$. Provided the NLSP-LSP mass difference is small, the correct LSP relic density is achieved by processes that maintain the LSP and the NLSP in thermal equilibrium (implying $n_{\text{LSP}} \sim n_{\text{NLSP}}$ for the corresponding abundances): $\text{LSP} + X \rightleftharpoons \text{NLSP} + X'$, where X, X' are light SM particles (quarks and leptons).

The reaction rate for the NLSP annihilation process $\text{NLSP} + \text{NLSP} \rightarrow X$ is given by $(n_{\text{NLSP}})^2 \sigma$, where σ is the thermally averaged cross section. On the other hand, the reaction rate for the processes that help to keep the LSP and the NLSP in thermal equilibrium ($\text{LSP} + X \rightarrow \text{NLSP} + X'$ and its inverse) depends on $n_{\text{LSP}} n_X \sigma'$. Even if the latter cross section σ' is possibly suppressed (as, for instance, for a singlino-like LSP in the NMSSM), the process $\text{LSP} + X \rightarrow \text{NLSP} + X'$ is typically faster than the annihilation process $\text{NLSP} + \text{NLSP} \rightarrow X$, since near the freeze-out temperature the abundance n_X of quarks and leptons is $\sim 10^9$ larger than the abundances of the LSP and NLSP (for $m_{\text{LSP}} \sim m_{\text{NLSP}}$) [505]. This allows to dilute the LSP density as fast as the NLSP density, and such “assisted coannihilation” can play a very important rôle in reconciling the relic density of an NMSSM LSP with WMAP measurements [346, 501].

In the MSSM, the LSP is usually a bino- or a higgsino-like neutralino (or a bino-higgsino mixed state). These scenarios can also be found in regions of the NMSSM parameter space where, apart from the enlarged Higgs sector and heavy neutralinos, the spectrum is MSSM-like (see [346, 501]). Before turning to NMSSM specific annihilation processes, we briefly discuss these MSSM-like scenarios.

The relic density of a nearly pure bino-like LSP is typically difficult to reconcile with WMAP measurements: the only available annihilation channel for a pure bino is via t -channel sfermion exchange (as all couplings to gauge bosons require a higgsino component), or via coannihilation with a sfermion (typically stau or stop) NLSP. The former requires light sfermions (as the cross section is $\propto (m_{\tilde{f}})^{-4}$), while the latter requires the sfermion to have a mass close to the one of the bino-like LSP.

If the LSP has a non-negligible higgsino component, it can efficiently annihilate into a pair of gauge bosons (WW , ZZ) and doublet-like Higgs bosons ($W^\mp H^\pm$, ZH , HA) via s -channel Z or Higgs exchange, as well as through t -channel neutralino and chargino exchange. For sufficiently heavy LSPs ($m_{\chi_1^0} > m_t$), annihilation into a pair of top quarks via Higgs exchange can significantly contribute. In the large $\tan\beta$ regime, where the Higgs couplings to $b\bar{b}$ and $\tau^+\tau^-$ are enhanced, $b\bar{b}$ and $\tau^+\tau^-$ final states are possible via s -channel Higgs exchange (not necessarily resonant).

A tiny higgsino fraction already opens the possibility of annihilation through resonant s -channel Higgs exchange. However, to overcome the smallness of the Higgs- $\chi_1^0\chi_1^0$ coupling in this case, a significant fine tuning is required so that the corresponding Higgs boson has the appropriate mass $\sim 2m_{\chi_1^0}$, which is not always possible in the MSSM.

Higgsino-like LSPs can also coannihilate with neutralinos and charginos. In the MSSM, it can happen that coannihilation is excessively efficient in this case, so that the relic density is too small. In the NMSSM, a non-vanishing singlino component can reduce the LSP couplings (and hence the coannihilation cross sections), leading to viable values for the relic density [501].

9.1.2 NMSSM-specific annihilation processes

In the NMSSM, the richer scalar/pseudoscalar Higgs sector allows to have the correct relic density in large regions of the parameter space, from low to high $\tan\beta$ [346, 501]. The possibility of light scalars and pseudoscalars consistent with LEP constraints also implies that new final states are kinematically open [346, 501]. For example, nearly pure binos can annihilate via H_1 -resonances into A_1A_1 .

A distinctive feature of the NMSSM is the possibility of a nearly pure singlino-like LSP. However, due to the small couplings to SM particles, singlino-like LSPs tend to have a too large relic density due to comparatively small annihilation cross sections. Still, there are several possible mechanisms that allow to render a singlino-like LSP compatible with WMAP bounds [346, 501]:

The couplings of a singlino-like LSP (with $N_{15}^2 \sim 1$) to CP-even Higgs bosons H_a are approximately given by $\sim -\sqrt{2}S_{a3}\kappa$ (see (A.14)), with S_{a3} replaced by P_{a3} for CP-odd Higgs bosons. Accordingly any CP-even (or CP-odd) Higgs state, whose singlet component S_{a3} (or P_{a3}) is sufficiently large, can give an important contribution to LSP annihilation via s -channel exchange, if κ is not too small.

If the singlino-like LSP has a small higgsino component, resonant s -channel annihilation via scalar/pseudoscalar Higgs or Z bosons allows for viable NMSSM scenarios if $m_{\chi_1^0} \lesssim 50$ GeV (occurring in regimes of $\kappa \ll \lambda$, and not very large μ_{eff}), or even $m_{\chi_1^0} \sim$ a few GeV provided that $2m_{\chi_1^0} \approx M_{H_1}, M_{A_1}$. Annihilation can then proceed into $\chi_1^0 \chi_1^0 \rightarrow b\bar{b}$ or, for the case of H_1 exchange and a light pseudoscalar, into $\chi_1^0 \chi_1^0 \rightarrow A_1 A_1$.

For a heavier mixed singlino/higgsino-like LSP ($m_{\chi_1^0} \gtrsim 100$ GeV) and large λ , the annihilation can occur via t -channel χ_1^0 exchange into pairs of mostly singlet-like H_1 and A_1 due to the enhanced $\chi_1^0 \chi_1^0 H_1(A_1)$ couplings $\sim \lambda$.

An LSP with a large singlino component can also coannihilate efficiently with heavier neutralinos (provided their mass difference is $\lesssim 10$ GeV), especially with a higgsino-like χ_2^0 , leading to $\chi_1^0 + \chi_2^0 \rightarrow t\bar{t}, b\bar{b}$ via heavy doublet-like Higgs exchange. Such coannihilation cross sections can be further enhanced by s -channel resonances: if the heavy states H_3 and A_2 belong to the heavy Higgs doublet and are nearly degenerate, one may even obtain a ‘‘double resonance’’ with two s -poles from both H_3 and A_2 exchange.

For a nearly pure singlino ($N_{15}^2 \gtrsim 99\%$) in regimes of very small λ and κ ($\kappa \lesssim \lambda \lesssim 0.01$), no efficient annihilation mechanisms are available, and even LSP-NLSP coannihilation (e.g. with heavier neutralinos/charginos, or sfermions) fails to diminish the LSP relic density sufficiently. In this case, the LSP density can be reduced by NLSP-NLSP annihilation (assisted coannihilation) in cases of, for instance, a bino-like NLSP or a sfermion NLSP. Assisted coannihilation is relevant for the fully constrained NMSSM [216, 217] as discussed in Section 7.1, in which case it requires nearly degenerate χ_1^0 and $\tilde{\tau}_1$ masses.

However, to this end the reaction rate for the process $\text{LSP} + X \rightarrow \text{NLSP} + X'$ ($\sim n_{\text{LSP}} n_X \sigma'$) has to be at least as large as the NLSP annihilation rate ($\sim n_{\text{NLSP}}^2 \sigma$); otherwise the LSP will no longer be in thermal equilibrium with the NLSP near the freeze-out temperature, but can be considered as decoupled. This condition allows to derive a rough lower bound on λ : the process $\text{LSP} + X \rightarrow \text{NLSP} + X'$ is made possible only through the non-singlet component of the LSP; for a dominantly singlet-like LSP, the analysis of the neutralino mass matrix shows that this non-singlet component is proportional to λ . Hence, the cross section σ' is proportional to λ^2 , and $\sigma'/\sigma \sim \lambda^2/g_2^2$. As mentioned above, the density n_X of light SM particles is about $n_X \sim 10^9 n_{\text{NLSP}}$ near the freeze-out temperature. Then, since $n_{\text{NLSP}} \sim n_{\text{LSP}}$, the reaction rate for $\text{LSP} + X \rightarrow \text{NLSP} + X'$ is larger than the NLSP annihilation rate only if $\lambda^2/g_2^2 \gtrsim 10^{-9}$ or $\lambda \gtrsim 10^{-5}$ [216, 217].

9.2 Dark matter detection in the NMSSM

In the NMSSM, the prospects for direct and indirect detection of neutralino DM can also be substantially different from what is expected in the MSSM (for a review of WIMP detection see [496]); subsequently we discuss direct and indirect detection separately.

9.2.1 Direct dark matter detection

Direct detection of neutralino dark matter (and generically WIMP dark matter) proceeds via the measurement of the recoil energy deposited by the scattering of WIMPs on nuclei in a detector. The energy transfer arising from the elastic scatterings typically lies around

$\mathcal{O}(10 \text{ keV})$, with predicted event rates below 0.1 events/kg/day. WIMP-nucleus interactions can be spin-independent (SI) or spin-dependent (SD). The SD contribution (via the axial-vector interaction) is only non-zero if the target nucleus has a net non-vanishing spin (unpaired nucleons), which is assumed to be carried by “odd-group” nucleons: protons or neutrons, whichever is most unpaired. The SI (scalar) contribution is proportional to the mass squared of the nucleus, and almost always dominates for nuclei with $A \gtrsim 30$, like those used in most modern detectors (e. g. Germanium and Xenon). The total elastic cross section is the sum of both contributions; below we will focus on the SI contribution. To obtain the direct detection rate, one needs to take into account several astroparticle physics ingredients such as the WIMP density and the WIMP velocity distribution near the earth.

With one remarkable exception, all dark matter direct detection experiments have only set upper limits on the WIMP-nucleon scattering cross section (σ_N , $N = n, p$). Xenon [506] has reported $\sigma_N^{\text{SI}} \lesssim 4.5 \times 10^{-8}$ pb for a WIMP mass of 30 GeV ($\sigma_N^{\text{SI}} \lesssim 8.8 \times 10^{-8}$ pb for $m_{\text{WIMP}} \sim 100$ GeV) at 90% CL, and CDMS [507] has set a limit $\sigma_N^{\text{SI}} \lesssim 4.6 \times 10^{-8}$ pb for a WIMP mass of 60 GeV at 90% CL. The best bounds on SD cross sections are from Xenon [508] ($\sigma_n^{\text{SD}} \lesssim 5 \times 10^{-3}$ pb) and KIMS [509] ($\sigma_p^{\text{SD}} \lesssim 0.18$ pb).

On the other hand, a positive result has been reported by the DAMA experiment (which looks for the so-called annual modulation signature): $\sigma_N \approx 0.2 - 1.0 \times 10^{-5}$ pb for WIMP masses between 30-100 GeV [510]. It has been noted in [321] that, in the NMSSM, very light bino- or singlino-like neutralinos (with $m_{\chi_1^0} \sim 6 - 9$ GeV) could allow to reconcile the DAMA results with the negative searches by other collaborations.

In the absence of CP violation, and in the zero momentum transfer limit, the elastic scattering of a neutralino LSP with a quark in a nucleon can be described by the effective low energy four-fermion Lagrangian (in the Dirac fermion notation):

$$\mathcal{L}_{\text{eff}} = \alpha_q^{\text{SI}} \bar{\chi}_1^0 \chi_1^0 \bar{q} q + \alpha_q^{\text{SD}} \bar{\chi}_1^0 \gamma_5 \gamma_\mu \chi_1^0 \bar{q} \gamma_5 \gamma^\mu q, \quad (9.4)$$

where $\alpha_q^{\text{SI (SD)}}$ denotes the spin-independent (spin-dependent) interaction with up- and down-type quarks (described by spinors $q = u, d$). The contribution of the dominant SI term to the χ_1^0 -nucleon cross section is given by (see, e. g., [503])

$$\sigma_N^{\text{SI}} = \frac{4m_N m_{\chi_1^0}}{\pi(m_N + m_{\chi_1^0})} f_N^2 \quad (9.5)$$

where $f_N/m_N \propto \sum_q \alpha_q^{\text{SI}}/m_q$, and each term in the sum over quarks must be weighted by the appropriate hadronic matrix elements.

The scalar neutralino-nucleon interaction α_q^{SI} arises from s -channel squark and t -channel Higgs (or Z) exchange at the tree level, while neutralino-gluon interactions contribute at the one-loop level (where all (heavy) quark loops should be taken into account). The squark exchange term is identical to the MSSM, the only difference in the NMSSM arising if the LSP has an important singlino component implying a significant reduction of $\alpha_q^{\text{SI}, \tilde{q}}$.

Significant differences in the NMSSM can arise from the t -channel Higgs contributions [347, 467, 511–515], which can dominate the SI interaction provided the LSP higgsino component is large enough. One has (see, e. g., [514])

$$\alpha_q^{\text{SI}, H} \propto -\frac{1}{M_{H_a}^2} (H_a \chi_1^0 \chi_1^0) (H_a q q) \quad (9.6)$$

where the quark Yukawa couplings ($H_a qq$) and the Higgs-neutralino-neutralino Yukawa couplings ($H_a \chi_1^0 \chi_1^0$) are given in Appendix A. The light Higgs doublet contributions are often dominant, although in the large $\tan\beta$ regime the contributions of the heavier doublet-like Higgs can also be important, as their couplings to neutralinos and quarks are enhanced [347].

For a singlino-like LSP, the cross section for the scattering off, e. g., a strange-quark is proportional to

$$\sigma_N^{\text{SI}} \propto (\alpha_s^{\text{SI},H})^2 \propto \frac{\kappa^2 h_s^2}{M_{H_a}^4} S_{a3}^2 S_{a1}^2. \quad (9.7)$$

Hence the exchange of a light H_1 can lead to large direct detection cross sections within reach of the present generation of detectors [514, 515]. (For a singlino-like LSP, the MSSM-like s -channel squark exchange diagram is in general too small to lead to cross sections close to the experimental limits. NMSSM specific contributions to the spin-dependent cross section can occur from a higgsino-like LSP, scattering off nuclei via Z boson exchange [347].)

As an example, we display in Fig. 9 (from [515]) the SI cross sections in a large region of the unconstrained NMSSM parameter space, which is compatible with collider, cosmological and B physics constraints. (A heavy sfermion sector ~ 1 TeV is assumed, except for slepton masses of ~ 150 GeV to comply with $(g-2)_\mu$). Large detection cross sections can originate from the presence of light singlet-like Higgs ($S_{13}^2 \gtrsim 0.9$) with a mass $M_{H_1} \sim 50$ GeV, see the right-hand panel of Fig. 9. Neutralinos within the reach of dark matter experiments are typically mixed singlino-higgsino states, with a mass between 50 – 130 GeV. The upper bound on the neutralino mass is due to the lightest stau becoming the LSP. If the slepton mass is increased, heavier neutralinos can be the LSP, but the resulting SUSY contribution to $(g-2)_\mu$ is generally too small.

9.2.2 Indirect dark matter detection

It is also possible to detect dark matter indirectly, looking for distinctive signals of WIMP annihilation products: gamma-ray fluxes, peculiar gamma-ray spectral features (like a sharp monochromatic peak at $E_\gamma \sim m_{\chi_1^0}$ arising from the loop induced annihilation $\chi_1^0 \chi_1^0 \rightarrow \gamma\gamma$), sizable amounts of antimatter from pair annihilation in the galactic halo, and also energetic neutrino fluxes from annihilations in the center of the sun or earth. For a review in the MSSM we again refer to [496].

Indirect detection of NMSSM neutralino dark matter has been considered in [516]. Since the indirect detection rates generically scale as inverse powers of $m_{\chi_1^0}$ (the pair annihilation rate being fixed by the requirement of a correct relic abundance), light neutralinos – as possible in the NMSSM – could give significantly enhanced rates. Additionally, new contributions (annihilations through a very light pseudoscalar) can enhance monochromatic $\chi_1^0 \chi_1^0 \rightarrow \gamma\gamma$ gamma-ray lines. The neutrino production rates from neutralino annihilation inside the earth and sun could also be distinct from MSSM expectations (see, e.g., [516] and references therein).

NMSSM scenarios exist where the annihilation of sub-GeV neutralinos could explain the 511 keV gamma-ray emission reported by INTEGRAL/SPI [517]. These gamma-rays can originate from low energy positrons possibly arising from the annihilation of an exceptionally light DM candidate. In the NMSSM, a very light neutralino ($m_{\chi_1^0} \sim 100$ MeV) combined with a light pseudoscalar, could potentially produce the 511 keV gamma-rays [321].




Figure 9: Scatter plots of the spin-independent neutralino-nucleon cross section in the general NMSSM for $\tan\beta = 5$, with the remaining parameters in the ranges $0.01 \leq \lambda, \kappa \leq 0.7$, $110 \text{ GeV} \lesssim M_2 \lesssim 430 \text{ GeV}$, $110 \text{ GeV} < \mu_{\text{eff}} < 300 \text{ GeV}$, $-800 \text{ GeV} \lesssim A_\lambda \lesssim 800 \text{ GeV}$, and $-300 \text{ GeV} \lesssim A_\kappa \lesssim 300 \text{ GeV}$. All the points fulfil all the experimental constraints and have a relic density in agreement with $0.1 \lesssim \Omega h^2 \lesssim 0.3$ (astrophysical bound, grey dots) or the WMAP constraint (black dots). On the left as a function of the neutralino mass (GeV), and on the right as a function of the lightest scalar Higgs mass (GeV). From [515].

However, efficient LSP annihilation via resonant s -channel pseudoscalar exchange would require $M_{A_1} \approx 2m_{\chi_1^0} \pm 10 \text{ MeV}$.

Recently, the PAMELA experiment [518, 519] reported a cosmic ray positron excess, with a positron over electron fraction that appears to rise at energies from 10 to 100 GeV. However, the same detector does not report any obvious antiproton excess in the same energy range. In order to interpret such a positron excess (and the absence of a \bar{p} excess) as dark matter annihilation products, it would be required that the DM particles dominantly annihilate into leptons (either due to dominant couplings to SM leptons, or annihilation into intermediate particles, sufficiently light so that they cannot decay into hadrons). Neutralino annihilation in the NMSSM (for $m_{\chi_1^0} \sim 160 \text{ GeV}$, and a very light pseudoscalar) [329, 520, 521] has been put forward to explain the PAMELA excess: relatively light neutralinos (typically bino-higgsino mixtures) can annihilate as $\chi_1^0 \chi_1^0 \rightarrow H_1 A_1$ via A_2 exchange, with A_1 so light that only decays into a pair of leptons (muons, or even electrons, for $M_{A_1} \lesssim 2m_\mu$) are kinematically allowed. In this case, the lightest scalar H_1 also dominantly decays into $A_1 A_1$, so that the final state of the LSP annihilation indeed consists of very energetic leptons. Extensions of the MSSM by a singlet and an additional heavy lepton [522] (exhibiting two DM components) have also been proposed in order to accommodate the PAMELA data.

9.3 Specific scenarios

The (semi)-constrained NMSSM

The relic density in the cNMSSM was first discussed in [335,339,342]. In the meantime constraints on Higgs and sparticle masses as well on the dark matter relic density [343,344] have become tighter (and RGEs/radiative corrections are known to a higher accuracy), with the result that some of the regions in the parameter space of the cNMSSM considered in these early studies are no longer phenomenologically viable. On the other hand, a singlino-like LSP was believed to be excluded [339,342], which turned out to be incorrect after the inclusion of additional (co-)annihilation channels.

In more recent studies, the parameter space of the semi-constrained NMSSM (relaxing the universality requirements for $m_{\tilde{S}}^2$ and A_κ at the GUT scale, see Section 7.1.3) has been thoroughly investigated. Regions allowed by constraints from collider and flavour physics and a WMAP compatible relic density have been identified in [346,347]. In the semi-constrained NMSSM, less options for LSP annihilation than in the general NMSSM are available:

- (i) MSSM-like mechanisms: annihilation of a bino- or bino-higgsino-like LSP through sfermion or Higgs boson exchange; coannihilation of a bino-like LSP with sfermions;
- (ii) annihilation near a pseudoscalar singlet-like Higgs resonance in the s -channel ($\lambda \sim 0.1$, any $\tan\beta$);
- (iii) (assisted) coannihilation of a singlino-like LSP with a higgsino-like NLSP ($\lambda \ll 1$, large m_0 and $\tan\beta$);
- (iv) (assisted) coannihilation of singlino-like LSPs with bino-like NLSPs, with the bino rapidly annihilating through a Higgs resonance ($\lambda \ll 1$, large $\tan\beta$);
- (v) (assisted) coannihilation of a singlino-like LSP with a stau or stop NLSP ($\lambda \ll 1$, small m_0).

In Fig. 10 (from [347]), we display some regions in the $(m_0, M_{1/2})$ parameter space, where a DM relic density within (or below) the WMAP bound is obtained through different annihilation mechanisms. (Also indicated are other relevant theoretical and phenomenological constraints.) The dominant annihilation mechanisms depend on $(m_0, M_{1/2})$, and on the values of the remaining parameters λ , A_0 , A_κ and $\tan\beta$ of the semi-constrained NMSSM. On the left, these are chosen as $\lambda = 0.1$, $A_0 = -900$ GeV, $A_\kappa = -60$ GeV and $\tan\beta = 5$, and the LSP is bino-like. Within the green horizontal band around $M_{1/2} \sim 350$ GeV, the correct relic density is achieved through a pseudoscalar resonance (ii). In the green region next to the blue region excluded by a stau LSP (at small m_0), the dominant mechanism is bino-stau coannihilation (i), whereas for small m_0 and $M_{1/2} \sim 250$ GeV the dominant mechanism is bino-stop coannihilation. On the right we show an example with small λ and a singlino-like LSP. In the small m_0 regime efficient singlino annihilation is achieved via coannihilation with a nearly degenerate lightest stau (v).

The prospects for direct DM detection have also been considered in [347]. Present bounds already constrain such regions at large $\tan\beta$ where the heavy Higgs doublet is relatively light and/or its couplings to an LSP with a non-vanishing higgsino component and quarks are enhanced. Large cross sections near or above present limits can also be obtained for $\lambda \sim 0.1$, where the LSP annihilates via a pseudoscalar singlet resonance (ii), and for bino-stau coannihilation regions (i). Hence, the semi-constrained NMSSM parameter space with a dominantly bino- or higgsino-like LSP has good prospects of being probed by the future direct detection experiments. On the other hand, for small $\tan\beta$ or




Figure 10: Constraints in the $(m_0, M_{1/2})$ plane in the semi-constrained NMSSM: regions excluded by theoretical constraints or by LEP/Tevatron searches on sparticles are indicated in black, by charged sfermion LSP in blue, from violation of LEP limits on Higgs searches in red, by constraints from B -physics in pink and disfavoured by $(g - 2)_\mu$ in violet. The regions allowed by the upper bound on the DM relic density are displayed in green. On the left: $\lambda = 0.1$, $A_0 = -900$ GeV, $A_\kappa = -60$ GeV and $\tan\beta = 5$. On the right: $\lambda = 0.01$, $A_0 = 300$ GeV, $A_\kappa = 50$ GeV and $\tan\beta = 10$. Here, in addition, the regions with a singlino LSP are shown in grey/cyan (with a relic density above/below the WMAP limit). From [347].

TeV-range heavy Higgs and squarks, large scale detectors will be required. The detection of a singlino LSP is beyond the reach of (even far future) large scale detectors.

The cNMSSM where $A_\kappa = A_0$ is imposed at the GUT scale, but m_S is allowed to differ from m_0 , has been investigated in [348] (see Section 7.1.3). It is still possible to have an LSP with a large bino or higgsino component which satisfies WMAP constraints via neutralino-stop (and stau) coannihilation or via Higgs resonances. For most points in this parameter space, direct detection of DM would appear possible for upgraded large scale detectors. For larger λ , a singlino-like LSP can annihilate via a CP-odd Higgs resonance in the s -channel.

In the fully constrained NMSSM [216, 217], discussed in Section 7.1, the LSP is always singlino-like, and $\lambda \lesssim 10^{-2}$. The only available mechanism to reduce the relic density is “assisted coannihilation” (ν) with the stau NLSP (provided $\lambda \gtrsim 10^{-5}$), which requires a near degeneracy of the χ_1^0 and $\tilde{\tau}_1$ masses. Since direct (or indirect) detection of a singlino LSP relies on its non-singlet components of $\mathcal{O}(\lambda)$, the prospects to detect the LSP in the cNMSSM are quite dim: WIMP-nucleon cross sections are extremely small, and indirect detection of the products of the annihilation process $\chi_1^0 \chi_1^0 \rightarrow A_1$ (with a pseudoscalar sufficiently light to be produced on-shell) also appears impossible, as the $A_1 \chi_1^0 \chi_1^0$ coupling is proportional to κ which is also tiny in the cNMSSM ($\kappa \lesssim 10^{-3}$). Hence, the fully constrained NMSSM can be excluded by the direct or indirect detection of a WIMP-like dark matter candidate.

The nMSSM

The neutralino relic density has also been studied in the framework of the nMSSM (see Section 7.3). In [176, 379] it has been shown that, for large λ , one can simultaneously achieve the correct Ωh^2 and a strong first order phase transition leading to successful EW baryogenesis. This occurs for a very light LSP with a mass $m_{\chi_1^0} \sim 30 - 40$ GeV, mostly singlino-like but with a sizable higgsino component. The dominant annihilation modes are via (possibly resonant) s -channel Z , H_1 and A_1 exchange [176, 378–380], the latter being favoured by the constraint from $(g - 2)_\mu$ [380].

Regarding DM detection [379], the dominant contributions to SI and SD interactions arise typically from Higgs and Z t -channel exchange, respectively. Due to the smallness of the $\chi_1^0 \chi_1^0 Z$ coupling, the SD cross section lies beyond the reach of next generation of DM experiments. On the other hand, the SI cross section could be within reach of the current and next generation of DM experiments for sizable values of λ (leading to an enhancement of the $\chi_1^0 \chi_1^0 H_a$ coupling) and a for comparatively light Higgs spectrum.

$U(1)'$ extensions

In $U(1)'$ extensions (see Section 8.2.1), the inclusion of an additional $U(1)'$ gaugino (bino') and the absence of the singlet self-coupling term in the superpotential can significantly modify the nature and properties of neutralino dark matter relative to the MSSM and NMSSM [378, 463–468]. Just like for the nMSSM, the absence of the S^3 term in the superpotential restricts the annihilation modes of the singlino. Compared to the NMSSM, new annihilation channels include t -channel χ_i^0 mediated annihilation into a pair of Z 's, and Z' (resonant) s -channel exchange. The analysis of [468] shows that, depending on the nature of the LSP (higgsino, mixed singlino-higgsino-bino', singlino) and on the remaining spectrum, the correct DM relic density can be obtained via s -channel resonances ($H_2/A/Z'/H_3$). Pure singlino neutralinos would not directly couple to the singlet Higgs boson, but they couple strongly to the Z' so that even very heavy LSPs can efficiently annihilate. The prospects for direct detection of neutralino DM in $U(1)'$ extensions have been addressed in [463, 468].

Dark matter in supersymmetric $U(1)'$ models with a secluded $U(1)'$ breaking sector (sMSSM) has also been studied [464]. For a light singlino-higgsino LSP with a mass $\lesssim 100$ GeV and an enhanced coupling to the Z , annihilation through the Z -resonance leads to a viable relic density over a large part of the parameter space without violating LEP constraints.

A comparative study of unconstrained versions of the MSSM, NMSSM, nMSSM, $U(1)'$ extended models (UMSSM) and secluded models (sMSSM) has been carried out in [378, 467], identifying allowed ranges for the mass of the LSP in each case as displayed in Fig. 11. Prospects for direct detection of neutralino DM were also compared in [467]: the cross sections for the $U(1)'$ extended models are similar to those of the NMSSM, and can be within reach of SuperCDMS and WARP. In view of the (comparatively) lighter LSPs in secluded models as well as in the nMSSM (see Fig. 11), spin-independent proton scattering cross sections could be detectable at CDMS.

In the exceptional SSM (E_6 SSM), which in addition to the Z' comprises three families




Figure 11: Neutralino relic density versus the lightest neutralino mass for the MSSM and NMSSM-like extensions, with $0.099 \lesssim \Omega h^2 \lesssim 0.123$ within the blue lines. Too large annihilation cross sections (leading to a too small Ωh^2) can originate from the Higgs boson pole in the s -channel in the MSSM, NMSSM and UMSSM for $m_{\chi_1^0} \sim M_{H_1}/2 \sim 60$ GeV, and from the Z boson pole at $m_{\chi_1^0} \sim M_Z/2$ in the n/sMSSM. (From [467].)

of Higgs doublets and singlets, it has been shown in [491] that the two families of “inert” doublet higgsinos and singlino (i.e. whose scalar partners are vev-less) lead to a decoupled neutralino sector with a naturally light LSP, that successfully accounts for Ωh^2 provided that $\tan \beta \lesssim 2$.

As referred to in Section 8.2.1, non-neutralino DM candidates have also been put forward in the framework of $U(1)'$ extensions: exotic LSPs [470], and right-handed sneutrinos [469, 475]. The correct relic density of right-handed sneutrinos can be obtained via t -channel Z' exchange in large regions of the parameter space, and provided the Z' mass is not excessively large, s -channel resonant Z' exchange also provides an efficient annihilation mechanism. The prospects for the detection of these DM candidates have been investigated [469]. $U(1)'$ extended models with sneutrinos as the LSP as well as the NLSP [475] have been proposed to explain the PAMELA observations, while accounting for a relic density in agreement with the WMAP bounds.

Other variants

In large- λ models [128] (see Section 3.2), the LSP is higgsino-like in most of the phenomenologically viable regions of parameter space. Then, the relic density is in general too small; only for $\tan\beta \lesssim 1.7$ regions with a substantial DM abundance can be found: annihilation via s -channel Z exchange can be reduced by increasing the singlino component of the LSP, or via a partial cancellation of the higgsino- Z coupling.

In the NMSSM including right-handed neutrino superfields with couplings to the singlet Higgs (thus providing a dynamically generated non-vanishing Majorana neutrino mass term), the properties of sneutrino dark matter have been discussed in [436]. For natural values of the input parameters, right-handed sneutrinos with a mass in the range of 5 – 200 GeV can reproduce the observed dark matter relic density without being excluded by direct dark matter searches. In this case, the predicted direct detection cross sections are generally within the reach of future experiments.

10 Conclusions and outlook

In this paper we reviewed the present status of knowledge of the theoretical and phenomenological aspects of the NMSSM, including physics at colliders, low energy precision observables and the properties of the dark matter candidate, focusing on the possible differences with respect to the MSSM.

We hope that the direct production of sparticles and/or Higgs bosons at the Tevatron/LHC will provide us information about possible supersymmetric extensions of the SM in the near future. Subsequently it will be essential to study the precise nature of the supersymmetric extension of the SM as thoroughly as possible. Depending on the experimental results, it may not be easy to distinguish the MSSM from non-minimal extensions as the NMSSM or variants thereof, but possibly the properties of the Higgs sector and/or the neutralino sector (in particular the nature of the LSP) will signal the presence of an additional singlet superfield. Clearly, these signals must be well understood in order to be able to interpret the data correctly. In view of the various scenarios that are possible within the NMSSM (as unconventional Higgs decays, and/or a neutralino with a large singlino component), additional efforts in the form of simulations of events corresponding to such scenarios are still required.

On the other hand, the properties of the Higgs and/or the neutralino sectors could correspond to those expected within the MSSM – which would *not* prove, however, that the MSSM is the correct supersymmetric extension of the SM at the TeV scale: at least in the decoupling limit (see Section 2.2.1) – and without a singlino-like LSP – the physics of the NMSSM becomes indistinguishable from the one of the MSSM. The question, for which values of parameters (like λ) the two models can be distinguished experimentally, is complicated due to the large number of unknown parameters, and the answer depends on the nature and the precision of future experimental data. Also here, additional studies are desirable (at least once data is available).

In any case it will be essential to combine the various pieces of information from low energy precision observables, physics at colliders and the properties of dark matter; ideally, these will allow us to pin down the fundamental theory at the TeV scale without too many ambiguities.

Acknowledgements

Some of the results presented here had been developed in cooperation with colleagues, whose collaboration we acknowledge gratefully: G. Belanger, D. G. Cerdeño, A. Djouadi, F. Domingo, J. F. Gunion, C.-C. Jean-Louis, D. López-Fogliani, S. Moretti, C. Muñoz, A. Pukhov and M. A. Sanchis-Lozano. We are grateful to P. Richardson, F. Roberto, C. Spitzer, F. Staub and L. Suter for corrections of some formulas.

Appendix A. Mixing matrices and tree level Higgs couplings

The Feynman rules of the NMSSM have been described first in [50] (including the quartic couplings not given here), and subsequently in [52].

A.1 Mixing matrices

First we define the mixing matrices which diagonalize the Higgs, neutralino and chargino mass matrices (after the addition of radiative corrections).

The CP-even 3×3 mass matrix \mathcal{M}_S^2 (2.22) in the basis $H_i^{\text{weak}} = (H_{dR}, H_{uR}, S_R)$ (in the SLHA2 conventions [49]) is diagonalized by an orthogonal 3×3 matrix S_{ij} rotating the basis H_i^{weak} ,

$$H_i^{\text{mass}} = S_{ij} H_j^{\text{weak}} , \quad (\text{A.1})$$

such that the mass eigenstates H_i^{mass} are ordered in increasing mass.

The CP-odd 2×2 mass matrix \mathcal{M}_P^2 (2.24) in the basis $(A = \cos \beta H_{uI} + \sin \beta H_{dI}, S_I)$ was obtained after the rotation by the angle β (2.25), which allowed to omit the Nambu-Goldstone boson. It can be diagonalized by an orthogonal 2×2 matrix P'_{ij} such that the physical CP-odd states A_i^{mass} (ordered in mass) are

$$\begin{aligned} A_1^{\text{mass}} &= P'_{11} A + P'_{12} S_I , \\ A_2^{\text{mass}} &= P'_{21} A + P'_{22} S_I . \end{aligned} \quad (\text{A.2})$$

In terms of the weak eigenstates $A_i^{\text{weak}} = (H_{dI}, H_{uI}, S_I)$, the mass eigenstates $(A_1^{\text{mass}}, A_2^{\text{mass}})$ are given by

$$A_i^{\text{mass}} = P_{ij} A_j^{\text{weak}} , \quad (\text{A.3})$$

where

$$P_{i1} = \sin \beta P'_{i1} , \quad P_{i2} = \cos \beta P'_{i1} , \quad P_{i3} = P'_{i2} . \quad (\text{A.4})$$

The inverse relation reads explicitly (omitting the Nambu-Goldstone boson)

$$\begin{aligned} H_{dI} &= P_{11} A_1^{\text{mass}} + P_{21} A_2^{\text{mass}} , \\ H_{uI} &= P_{12} A_1^{\text{mass}} + P_{22} A_2^{\text{mass}} , \\ S_I &= P_{13} A_1^{\text{mass}} + P_{23} A_2^{\text{mass}} . \end{aligned} \quad (\text{A.5})$$

Omitting again the Goldstone boson, the charged weak eigenstates $H_{u,d}^{\pm}$ contain a component of the physical charged Higgs boson H^{\pm} given by

$$H_u^{\pm} = \cos \beta H^{\pm} , \quad H_d^{\pm} = \sin \beta H^{\pm} . \quad (\text{A.6})$$

The (symmetric) 5×5 neutralino mass matrix \mathcal{M}_0 (2.32) in the basis $\psi^0 = (-i\lambda_1, -i\lambda_2^3, \psi_d^0, \psi_u^0, \psi_S)$ is diagonalized by an orthogonal real matrix N_{ij} , such that

the physical masses $m_{\chi_i^0}$ ordered in $|m_{\chi_i^0}|$ are real, but not necessarily positive. Denoting the 5 eigenstates by χ_i^0 , we have

$$\chi_i^0 = N_{ij} \psi_j^0. \quad (\text{A.7})$$

The diagonalization of the (not symmetric) 2×2 chargino mass matrix X (2.35) in the basis ψ^-, ψ^+ (see (2.33), (2.34)) requires different rotations of ψ^- and ψ^+ into the mass eigenstates χ^-, χ^+ as

$$\chi^- = U\psi^-, \quad \chi^+ = V\psi^+ \quad (\text{A.8})$$

with

$$U = \begin{pmatrix} \cos \theta_U & \sin \theta_U \\ -\sin \theta_U & \cos \theta_U \end{pmatrix}, \quad V = \begin{pmatrix} \cos \theta_V & \sin \theta_V \\ -\sin \theta_V & \cos \theta_V \end{pmatrix}. \quad (\text{A.9})$$

Finally the top squark mass matrix (2.36) is diagonalized in terms of the mass eigenstates \tilde{t}_1, \tilde{t}_2 (with $m_{\tilde{t}_1} < m_{\tilde{t}_2}$), the bottom squark mass matrix (2.37) in terms of the mass eigenstates \tilde{b}_1, \tilde{b}_2 , and the tau slepton mass matrix (2.38) in terms of the mass eigenstates $\tilde{\tau}_1, \tilde{\tau}_2$ by the rotations

$$\begin{aligned} \tilde{t}_1 &= \cos \theta_T \tilde{t}_L + \sin \theta_T \tilde{t}_R, & \tilde{t}_2 &= \cos \theta_T \tilde{t}_R - \sin \theta_T \tilde{t}_L, \\ \tilde{b}_1 &= \cos \theta_B \tilde{b}_L + \sin \theta_B \tilde{b}_R, & \tilde{b}_2 &= \cos \theta_B \tilde{b}_R - \sin \theta_B \tilde{b}_L, \\ \tilde{\tau}_1 &= \cos \theta_\tau \tilde{\tau}_L + \sin \theta_\tau \tilde{\tau}_R, & \tilde{\tau}_2 &= \cos \theta_\tau \tilde{\tau}_R - \sin \theta_\tau \tilde{\tau}_L. \end{aligned} \quad (\text{A.10})$$

Now we proceed to give the Higgs couplings (Feynman rules) in terms of physical mass eigenstates.

A.2 Higgs couplings

Higgs-quarks/leptons

We consider Higgs couplings to quarks and leptons of the third generation only, from which the remaining couplings can be deduced easily. First we recall the relations between the quark and lepton masses m_t, m_b, m_τ and the Yukawa couplings h_t, h_b and h_τ , where $v^2 = v_u^2 + v_d^2$ (related to M_Z by (2.13)):

$$h_t = \frac{m_t}{v \sin \beta}, \quad h_b = \frac{m_b}{v \cos \beta}, \quad h_\tau = \frac{m_\tau}{v \cos \beta}. \quad (\text{A.11})$$

(Corrections to h_b are discussed in Section 3.3.) Below, the left and right-handed top (bottom) quark and lepton bi-spinors are denoted by t_L, t_R (b_L, b_R) and $\nu_{\tau L}, \tau_L, \tau_R$, and the quark couplings are diagonal in the colour indices. (For simplicity, the index ^{mass} of the CP-even mass eigenstates H_i^{mass} and the CP-odd mass eigenstates A_i^{mass} will be omitted subsequently.) The denominators $\sqrt{2}$ in the couplings originate from the rescaling of the kinetic terms of real Higgs bosons with respect to complex Higgs bosons.

$$\begin{aligned}
H_i t_L t_R^c &: -\frac{h_t}{\sqrt{2}} S_{i2} \\
H_i b_L b_R^c &: \frac{h_b}{\sqrt{2}} S_{i1} \\
H_i \tau_L \tau_R^c &: \frac{h_\tau}{\sqrt{2}} S_{i1} \\
A_i t_L t_R^c &: -i \frac{h_t}{\sqrt{2}} P_{i2} \\
A_i b_L b_R^c &: i \frac{h_b}{\sqrt{2}} P_{i1} \\
A_i \tau_L \tau_R^c &: i \frac{h_\tau}{\sqrt{2}} P_{i1} \\
H^+ b_L t_R^c &: h_t \cos \beta \\
H^- t_L b_R^c &: -h_b \sin \beta \\
H^- \nu_{\tau_L} \tau_R^c &: -h_\tau \sin \beta
\end{aligned} \tag{A.12}$$

Higgs-gauge bosons

$$\begin{aligned}
H_i Z_\mu Z_\nu &: g_{\mu\nu} \frac{g_1^2 + g_2^2}{\sqrt{2}} (v_d S_{i1} + v_u S_{i2}) \\
H_i W_\mu^+ W_\nu^- &: g_{\mu\nu} \frac{g_2^2}{\sqrt{2}} (v_d S_{i1} + v_u S_{i2}) \\
H_i(p) H^+(p') W_\mu^- &: \frac{g_2}{2} (\cos \beta S_{i2} - \sin \beta S_{i1}) (p - p')_\mu \\
A_i(p) H^+(p') W_\mu^- &: i \frac{g_2}{2} (\cos \beta P_{i2} + \sin \beta P_{i1}) (p - p')_\mu \\
H_i(p) A_j(p') Z_\mu &: -i \sqrt{\frac{g_1^2 + g_2^2}{2}} (S_{i1} P_{j1} - S_{i2} P_{j2}) (p - p')_\mu \\
H^+(p) H^-(p') Z_\mu &: \frac{g_1^2 - g_2^2}{\sqrt{g_1^2 + g_2^2}} (p - p')_\mu
\end{aligned} \tag{A.13}$$

Higgs-neutralinos/charginos

Here, the couplings to two neutralinos involve the combinations $\Pi_{ij}^{ab} = N_{ia} N_{jb} + N_{ib} N_{ja}$ of the neutralino mixing matrices:

$$\begin{aligned}
H_a \chi_i^+ \chi_j^- &: \frac{\lambda}{\sqrt{2}} S_{a3} U_{i2} V_{j2} + \frac{g_2}{\sqrt{2}} (S_{a2} U_{i1} V_{j2} + S_{a1} U_{i2} V_{j1}) \\
A_a \chi_i^+ \chi_j^- &: i \left(\frac{\lambda}{\sqrt{2}} P_{a3} U_{i2} V_{j2} - \frac{g_2}{\sqrt{2}} (P_{a2} U_{i1} V_{j2} + P_{a1} U_{i2} V_{j1}) \right) \\
H^+ \chi_i^- \chi_j^0 &: \lambda \cos \beta U_{i2} N_{j5} - \frac{\sin \beta}{\sqrt{2}} U_{i2} (g_1 N_{j1} + g_2 N_{j2}) + g_2 \sin \beta U_{i1} N_{j3} \\
H^- \chi_i^+ \chi_j^0 &: \lambda \sin \beta V_{i2} N_{j5} + \frac{\cos \beta}{\sqrt{2}} V_{i2} (g_1 N_{j1} + g_2 N_{j2}) + g_2 \cos \beta V_{i1} N_{j4} \\
H_a \chi_i^0 \chi_j^0 &: \frac{\lambda}{\sqrt{2}} (S_{a1} \Pi_{ij}^{45} + S_{a2} \Pi_{ij}^{35} + S_{a3} \Pi_{ij}^{34}) - \sqrt{2} \kappa S_{a3} N_{i5} N_{j5} \\
&\quad + \frac{g_1}{2} (S_{a1} \Pi_{ij}^{13} - S_{a2} \Pi_{ij}^{14}) - \frac{g_2}{2} (S_{a1} \Pi_{ij}^{23} - S_{a2} \Pi_{ij}^{24}) \\
A_a \chi_i^0 \chi_j^0 &: i \left(\frac{\lambda}{\sqrt{2}} (P_{a1} \Pi_{ij}^{45} + P_{a2} \Pi_{ij}^{35} + P_{a3} \Pi_{ij}^{34}) - \sqrt{2} \kappa P_{a3} N_{i5} N_{j5} \right. \\
&\quad \left. - \frac{g_1}{2} (P_{a1} \Pi_{ij}^{13} - P_{a2} \Pi_{ij}^{14}) + \frac{g_2}{2} (P_{a1} \Pi_{ij}^{23} - P_{a2} \Pi_{ij}^{24}) \right) \tag{A.14}
\end{aligned}$$

Triple Higgs couplings

$$\begin{aligned}
H_a H_b H_c &: \frac{\lambda^2}{\sqrt{2}} (v_d (\Pi_{abc}^{122} + \Pi_{abc}^{133}) + v_u (\Pi_{abc}^{211} + \Pi_{abc}^{233}) + s (\Pi_{abc}^{311} + \Pi_{abc}^{322})) \\
&\quad - \frac{\lambda \kappa}{\sqrt{2}} (v_d \Pi_{abc}^{323} + v_u \Pi_{abc}^{313} + 2s \Pi_{abc}^{123}) + \sqrt{2} \kappa^2 s \Pi_{abc}^{333} \\
&\quad - \frac{\lambda A_\lambda}{\sqrt{2}} \Pi_{abc}^{123} + \frac{\kappa A_\kappa}{3\sqrt{2}} \Pi_{abc}^{333} \\
&\quad + \frac{g_1^2 + g_2^2}{4\sqrt{2}} (v_d (\Pi_{abc}^{111} - \Pi_{abc}^{122}) - v_u (\Pi_{abc}^{211} - \Pi_{abc}^{222})) \\
&\quad + \frac{\mu'}{\sqrt{2}} (\kappa \Pi_{abc}^{333} - \lambda \Pi_{abc}^{123}) \tag{A.15}
\end{aligned}$$

where

$$\begin{aligned}
\Pi_{abc}^{ijk} &= S_{ai} S_{bj} S_{ck} + S_{ai} S_{cj} S_{bk} + S_{bi} S_{aj} S_{ck} \\
&\quad + S_{bi} S_{cj} S_{ak} + S_{ci} S_{aj} S_{bk} + S_{ci} S_{bj} S_{ak} . \tag{A.16}
\end{aligned}$$

$$\begin{aligned}
H_a A_b A_c &: \frac{\lambda^2}{\sqrt{2}} (v_d(\Pi_{abc}^{122} + \Pi_{abc}^{133}) + v_u(\Pi_{abc}^{211} + \Pi_{abc}^{233}) + s(\Pi_{abc}^{311} + \Pi_{abc}^{322})) \\
&+ \frac{\lambda\kappa}{\sqrt{2}} (v_d(\Pi_{abc}^{233} - 2\Pi_{abc}^{323}) + v_u(\Pi_{abc}^{133} - 2\Pi_{abc}^{313})) \\
&+ 2s(\Pi_{abc}^{312} - \Pi_{abc}^{123} - \Pi_{abc}^{213}) + \sqrt{2}\kappa^2 s\Pi_{abc}^{333} \\
&+ \frac{\lambda A_\lambda}{\sqrt{2}} (\Pi_{abc}^{123} + \Pi_{abc}^{213} + \Pi_{abc}^{312}) - \frac{\kappa A_\kappa}{\sqrt{2}} \Pi_{abc}^{333} \\
&+ \frac{g_1^2 + g_2^2}{4\sqrt{2}} (v_d(\Pi_{abc}^{111} - \Pi_{abc}^{122}) - v_u(\Pi_{abc}^{211} - \Pi_{abc}^{222})) \\
&+ \frac{\mu'}{\sqrt{2}} (\lambda (\Pi_{abc}^{312} - \Pi_{abc}^{123} - \Pi_{abc}^{213}) + \kappa \Pi_{abc}^{333})
\end{aligned} \tag{A.17}$$

where

$$\Pi_{abc}^{ijk} = S_{ai}(P_{bj}P_{ck} + P_{cj}P_{bk}) . \tag{A.18}$$

$$\begin{aligned}
H_a H^+ H^- &: \frac{\lambda^2}{\sqrt{2}} (s(\Pi_a^{311} + \Pi_a^{322}) - v_d \Pi_a^{212} - v_u \Pi_a^{112}) \\
&+ \sqrt{2}\lambda\kappa s \Pi_a^{312} + \frac{\lambda A_\lambda}{\sqrt{2}} \Pi_a^{312} \\
&+ \frac{g_1^2}{4\sqrt{2}} (v_u(\Pi_a^{211} - \Pi_a^{222}) + v_d(\Pi_a^{122} - \Pi_a^{111})) \\
&+ \frac{g_2^2}{4\sqrt{2}} (v_d(\Pi_a^{111} + \Pi_a^{122} + 2\Pi_a^{212}) + v_u(\Pi_a^{211} + \Pi_a^{222} + 2\Pi_a^{112})) \\
&+ \frac{\lambda\mu'}{\sqrt{2}} \Pi_a^{312}
\end{aligned} \tag{A.19}$$

where

$$\Pi_a^{ijk} = 2S_{ai}C_jC_k \text{ with } C_1 = \cos\beta, C_2 = \sin\beta . \tag{A.20}$$

In contrast to the previous couplings, the triple Higgs couplings in the general NMSSM differ slightly (by the terms $\sim \mu'$) from the ones in the scale invariant NMSSM. The quartic Higgs (and Higgs-gauge) couplings, which are less relevant for the Higgs phenomenology (i.e. Higgs-to-Higgs decays) can be found in [50].

Higgs-squarks/sleptons

Here we assume diagonal Yukawa couplings in family space which we denote as h_{ua}, h_{da}, h_{ea} , $a = 1, 2, 3$, with $h_{u3} \equiv h_t$, $h_{d3} \equiv h_b$, $h_{e3} \equiv h_\tau$. The soft trilinear couplings are A_{ua} with $A_{u3} \equiv A_t$, A_{da} with $A_{d3} \equiv A_b$, and A_{ea} with $A_{e3} \equiv A_\tau$.

The squarks are denoted as $U_{La}, D_{La}, U_{Ra}, D_{Ra}$, the charged sleptons as E_{La}, E_{Ra} , and the sneutrinos as $\tilde{\nu}_{La}$.

In the case of the first two families ($a = 1, 2$), these states can be considered as mass eigenstates. For the third family (apart from the tau sneutrino), the couplings of the mass eigenstates defined in (A.10) depend on the corresponding squark and slepton mixing angles. We denote the couplings of the squark/slepton weak eigenstates (= mass eigenstates for $a = 1, 2$) by $g(\dots)$ for all three families; the couplings of the mass eigenstates of the third family are subsequently expressed in terms of $g(\dots)$ with $a = 3$.

Higgs-squarks:

$$\begin{aligned}
g(H_i, U_{La}, U_{La}) &= \sqrt{2} \left(h_{ua}^2 v_u S_{i2} + \left(\frac{g_1^2}{12} - \frac{g_2^2}{4} \right) (v_u S_{i2} - v_d S_{i1}) \right), \\
g(H_i, D_{La}, D_{La}) &= \sqrt{2} \left(h_{da}^2 v_d S_{i1} + \left(\frac{g_1^2}{12} + \frac{g_2^2}{4} \right) (v_u S_{i2} - v_d S_{i1}) \right), \\
g(H_i, U_{Ra}, U_{Ra}) &= \sqrt{2} \left(h_{ua}^2 v_u S_{i2} - \frac{g_1^2}{3} (v_u S_{i2} - v_d S_{i1}) \right), \\
g(H_i, D_{Ra}, D_{Ra}) &= \sqrt{2} \left(h_{da}^2 v_d S_{i1} + \frac{g_1^2}{6} (v_u S_{i2} - v_d S_{i1}) \right), \\
g(H_i, U_{La}, U_{Ra}) &= \frac{-h_{ua}}{\sqrt{2}} (A_{ua} S_{i2} - \mu_{\text{eff}} S_{i1} - \lambda v_d S_{i3}), \\
g(H_i, D_{La}, D_{Ra}) &= \frac{h_{da}}{\sqrt{2}} (A_{da} S_{i1} - \mu_{\text{eff}} S_{i2} - \lambda v_u S_{i3}), \\
g(A_i, U_{La}, U_{Ra}) &= \frac{-h_{ua}}{\sqrt{2}} (A_{ua} P_{i2} + \mu_{\text{eff}} P_{i1} + \lambda v_d P_{i3}), \\
g(A_i, D_{La}, D_{Ra}) &= \frac{h_{da}}{\sqrt{2}} (A_{da} P_{i1} + \mu_{\text{eff}} P_{i2} + \lambda v_u P_{i3}), \\
g(H^+, U_{La}, D_{La}) &= \frac{v_u v_d}{v} (g_2^2 - h_{ua}^2 - h_{da}^2), \\
g(H^+, U_{La}, D_{Ra}) &= -h_{da} (\cos \beta \mu + \sin \beta A_{da}), \\
g(H^+, U_{Ra}, D_{La}) &= -h_{ua} (\sin \beta \mu + \cos \beta A_{ua}), \\
g(H^+, U_{Ra}, D_{Ra}) &= -\frac{h_{ua} h_{da}}{v}.
\end{aligned} \tag{A.21}$$

Third family squarks, CP-even scalars S_i :

$$\begin{aligned}
H_i \tilde{t}_1 \tilde{t}_1 & : \cos^2 \theta_T g(H_i, U_{L3}, U_{L3}) + \sin^2 \theta_T g(H_i, U_{R3}, U_{R3}) \\
& \quad + 2 \cos \theta_T \sin \theta_T g(H_i, U_{L3}, U_{R3}) \\
H_i \tilde{t}_2 \tilde{t}_2 & : \sin^2 \theta_T g(H_i, U_{L3}, U_{L3}) + \cos^2 \theta_T g(H_i, U_{R3}, U_{R3}) \\
& \quad - 2 \cos \theta_T \sin \theta_T g(H_i, U_{L3}, U_{R3}) \\
H_i \tilde{t}_1 \tilde{t}_2 & : \cos \theta_T \sin \theta_T (g(H_i, U_{R3}, U_{R3}) - g(H_i, U_{L3}, U_{L3})) \\
& \quad + (\cos^2 \theta_T - \sin^2 \theta_T) g(H_i, U_{L3}, U_{R3}) \\
H_i \tilde{b}_1 \tilde{b}_1 & : \cos^2 \theta_B g(H_i, D_{L3}, D_{L3}) + \sin^2 \theta_B g(H_i, D_{R3}, D_{R3}) \\
& \quad + 2 \cos \theta_B \sin \theta_B g(H_i, D_{L3}, D_{R3}) \\
H_i \tilde{b}_2 \tilde{b}_2 & : \sin^2 \theta_B g(H_i, D_{L3}, D_{L3}) + \cos^2 \theta_B g(H_i, D_{R3}, D_{R3}) \\
& \quad - 2 \cos \theta_B \sin \theta_B g(H_i, D_{L3}, D_{R3}) \\
H_i \tilde{b}_1 \tilde{b}_2 & : \cos \theta_B \sin \theta_B (g(H_i, D_{R3}, D_{R3}) - g(H_i, D_{L3}, D_{L3})) \\
& \quad + (\cos^2 \theta_B - \sin^2 \theta_B) g(H_i, D_{L3}, D_{R3})
\end{aligned} \tag{A.22}$$

Third family squarks, CP-odd scalars A_i :

$$\begin{aligned}
A_i \tilde{t}_1 \tilde{t}_1 & : 0 \\
A_i \tilde{t}_2 \tilde{t}_2 & : 0 \\
A_i \tilde{t}_1 \tilde{t}_2 & : g(A_i, U_{L3}, U_{R3}) \\
A_i \tilde{b}_1 \tilde{b}_1 & : 0 \\
A_i \tilde{b}_2 \tilde{b}_2 & : 0 \\
A_i \tilde{b}_1 \tilde{b}_2 & : g(A_i, D_{L3}, D_{R3})
\end{aligned} \tag{A.23}$$

Third family squarks, charged scalar H^+ :

$$\begin{aligned}
H^+ \tilde{t}_1 \tilde{b}_1 & : \cos \theta_T \cos \theta_B g(H^+, U_{L3}, D_{L3}) + \sin \theta_T \sin \theta_B g(H^+, U_{R3}, D_{R3}) \\
& \quad + \cos \theta_T \sin \theta_B g(H^+, U_{L3}, D_{R3}) + \sin \theta_T \cos \theta_B g(H^+, U_{R3}, D_{L3}) , \\
H^+ \tilde{t}_1 \tilde{b}_2 & : -\cos \theta_T \sin \theta_B g(H^+, U_{L3}, D_{L3}) + \sin \theta_T \cos \theta_B g(H^+, U_{R3}, D_{R3}) \\
& \quad + \cos \theta_T \cos \theta_B g(H^+, U_{L3}, D_{R3}) - \sin \theta_T \sin \theta_B g(H^+, U_{R3}, D_{L3}) , \\
H^+ \tilde{t}_2 \tilde{b}_1 & : -\sin \theta_T \cos \theta_B g(H^+, U_{L3}, D_{L3}) + \cos \theta_T \sin \theta_B g(H^+, U_{R3}, D_{R3}) \\
& \quad - \sin \theta_T \sin \theta_B g(H^+, U_{L3}, D_{R3}) + \cos \theta_T \cos \theta_B g(H^+, U_{R3}, D_{L3}) , \\
H^+ \tilde{t}_2 \tilde{b}_2 & : \sin \theta_T \sin \theta_B g(H^+, U_{L3}, D_{L3}) + \cos \theta_T \cos \theta_B g(H^+, U_{R3}, D_{R3}) \\
& \quad - \sin \theta_T \cos \theta_B g(H^+, U_{L3}, D_{R3}) - \cos \theta_T \sin \theta_B g(H^+, U_{R3}, D_{L3}) .
\end{aligned} \tag{A.24}$$

Higgs-sleptons:

$$\begin{aligned}
g(H_i, E_{La}, E_{La}) &= \sqrt{2} \left(h_{ea}^2 v_u S_{i1} + \left(-\frac{g_1^2}{4} + \frac{g_2^2}{4} \right) (v_u S_{i2} - v_d S_{i1}) \right) , \\
g(H_i, E_{Ra}, E_{Ra}) &= \sqrt{2} \left(h_{ea}^2 v_d S_{i1} + \frac{g_1^2}{2} (v_u S_{i2} - v_d S_{i1}) \right) , \\
g(H_i, E_{La}, E_{Ra}) &= \frac{h_{ea}}{\sqrt{2}} (A_{ea} S_{i1} - \mu_{\text{eff}} S_{i2} - \lambda v_u S_{i3}) , \\
g(H_i, \tilde{\nu}_{La}, \tilde{\nu}_{La}) &= \sqrt{2} \left(-\frac{g_1^2}{4} - \frac{g_2^2}{4} \right) (v_u S_{i2} - v_d S_{i1}) , \\
g(A_i, E_{La}, E_{Ra}) &= \frac{h_{ea}}{\sqrt{2}} (A_{ea} P_{i1} + \mu_{\text{eff}} P_{i2} + \lambda v_u P_{i3}) , \\
g(H^+, E_{La}, \tilde{\nu}_{La}) &= g_2^2 \frac{v_u v_d}{v} . \tag{A.25}
\end{aligned}$$

Third family sleptons (staus), CP-even scalars H_i (the coupling $H_i \tilde{\nu}_{L3} \tilde{\nu}_{L3}$ does not differ from (A.25)):

$$\begin{aligned}
H_i \tilde{\tau}_1 \tilde{\tau}_1 &: \cos^2 \theta_L g(H_i, E_{L3}, E_{L3}) + \sin^2 \theta_L g(H_i, E_{R3}, E_{R3}) \\
&\quad + 2 \cos \theta_L \sin \theta_L g(H_i, E_{L3}, E_{R3}) \\
H_i \tilde{\tau}_2 \tilde{\tau}_2 &: \sin^2 \theta_L g(H_i, E_{L3}, E_{L3}) + \cos^2 \theta_L g(H_i, E_{R3}, E_{R3}) \\
&\quad - 2 \cos \theta_L \sin \theta_L g(H_i, E_{L3}, E_{R3}) \\
H_i \tilde{\tau}_1 \tilde{\tau}_2 &: \cos \theta_L \sin \theta_L (g(H_i, E_{R3}, E_{R3}) - g(H_i, E_{L3}, E_{L3})) \\
&\quad + (\cos^2 \theta_L - \sin^2 \theta_L) g(H_i, E_{L3}, E_{R3}) \tag{A.26}
\end{aligned}$$

Third family sleptons (staus), CP-odd scalars A_i :

$$\begin{aligned}
A_i \tilde{\tau}_1 \tilde{\tau}_1 &: 0 \\
A_i \tilde{\tau}_2 \tilde{\tau}_2 &: 0 \\
A_i \tilde{\tau}_1 \tilde{\tau}_2 &: g(A_i, E_{L3}, E_{R3}) \tag{A.27}
\end{aligned}$$

Third family sleptons (staus), charged scalar H^+ :

$$\begin{aligned}
H^+ \tilde{\tau}_1 \tilde{\nu}_{L3} &: \cos \theta_L \frac{v_u v_d}{v} (g_2^2 - h_{e3}^2) - \sin \theta_L h_{e3} (\cos \beta \mu + \sin \beta A_{e3}) \\
H^+ \tilde{\tau}_2 \tilde{\nu}_{L3} &: -\sin \theta_L \frac{v_u v_d}{v} (g_2^2 - h_{e3}^2) - \cos \theta_L h_{e3} (\cos \beta \mu + \sin \beta A_{e3}) \tag{A.28}
\end{aligned}$$

Yukawa induced radiative corrections to triple Higgs couplings

As in the MSSM [523–528], the dominant radiative corrections to triple Higgs couplings originate from top/bottom-quark loops, which are regularized in the ultraviolet by the corresponding contributions from squark loops. Here we give these radiative corrections in the NMSSM, which just require to generalize the Higgs mixing matrices of the MSSM to the NMSSM, see Appendix A.1.

However, we confine ourselves to the contributions involving a potentially large logarithm: we assume approximately degenerate squark masses given by M_{SUSY} . The infrared cutoff inside the dominant logarithm is given by the mass m_q of the quark inside the loop or the mass of the decaying Higgs boson through this vertex (whichever is larger); we will denote it by M_H here. Then the potentially large logarithm is

$$t \equiv \ln (M_{\text{SUSY}}^2 / \max(m_q^2, M_H^2)) . \quad (\text{A.29})$$

The dominant corrections to the triple CP-even Higgs interactions (A.15) are then given by

$$\Delta H_a H_b H_c : \frac{3t}{8\sqrt{2}\pi^2} (h_t^4 v_u \Pi_{abc}^{222} + h_b^4 v_d \Pi_{abc}^{111}) \quad (\text{A.30})$$

with Π_{abc}^{ijk} as in (A.16). The dominant corrections to the CP-even/CP-odd Higgs interactions (A.17) read the same,

$$\Delta H_a A_b A_c : \frac{3t}{8\sqrt{2}\pi^2} (h_t^4 v_u \Pi_{abc}^{222} + h_b^4 v_d \Pi_{abc}^{111}) \quad (\text{A.31})$$

where now Π_{abc}^{ijk} is given in (A.18). The dominant corrections to the CP-even/charged Higgs couplings (A.19) read

$$\begin{aligned} \Delta H_a H^+ H^- : \frac{3t}{8\sqrt{2}\pi^2} & \left[h_t^2 v_u (h_t^2 \Pi_a^{211} + h_b^2 (\Pi_a^{222} + \Pi_a^{112})) \right. \\ & \left. + h_b^2 v_d (h_b^2 \Pi_a^{122} + h_t^2 (\Pi_a^{212} + \Pi_a^{111})) \right] \end{aligned} \quad (\text{A.32})$$

with Π_a^{ijk} as in (A.20).

Appendix B. Renormalisation group equations

In this Appendix we give the renormalisation group equations in the $\overline{\text{DR}}$ scheme, valid above the SUSY breaking scale, for the parameters of the general NMSSM. The conventions are $t = \ln Q^2$, and the $U(1)_Y$ gauge coupling g_1^2 is defined in the Standard Model normalisation (related to the GUT or $SU(5)$ normalisation by $g_1^2 = \frac{3}{5} (g_1^{\text{GUT}})^2$).

B.1 Gauge and Yukawa couplings

The RGEs are known to two-loop order [529–531] (for the NMSSM see [72, 73]); it suffices, however, to include the Yukawa couplings of the third family only. (The CKM mixing matrix is not considered here.)

$$\begin{aligned}
16\pi^2 \frac{dg_1^2}{dt} &= 11g_1^4 + \frac{g_1^4}{16\pi^2} \left(\frac{199}{9}g_1^2 + 9g_2^2 + \frac{88}{3}g_3^2 - \frac{26}{3}h_t^2 - \frac{14}{3}h_b^2 - 6h_\tau^2 - 2\lambda^2 \right), \\
16\pi^2 \frac{dg_2^2}{dt} &= g_2^4 + \frac{g_2^4}{16\pi^2} \left(3g_1^2 + 25g_2^2 + 24g_3^2 - 6h_t^2 - 6h_b^2 - 2h_\tau^2 - 2\lambda^2 \right), \\
16\pi^2 \frac{dg_3^2}{dt} &= -3g_3^4 + \frac{g_3^4}{16\pi^2} \left(\frac{11}{3}g_1^2 + 9g_2^2 + 14g_3^2 - 4h_t^2 - 4h_b^2 \right), \\
16\pi^2 \frac{dh_t^2}{dt} &= h_t^2 \left(6h_t^2 + h_b^2 + \lambda^2 - \frac{13}{9}g_1^2 - 3g_2^2 - \frac{16}{3}g_3^2 \right) \\
&\quad + \frac{h_t^2}{16\pi^2} \left(-22h_t^4 - 5h_b^4 - 3\lambda^4 - 5h_t^2h_b^2 - 3h_t^2\lambda^2 - h_b^2h_\tau^2 - 4h_b^2\lambda^2 \right. \\
&\quad - h_\tau^2\lambda^2 - 2\lambda^2\kappa^2 + 2g_1^2h_t^2 + \frac{2}{3}g_1^2h_b^2 + 6g_2^2h_t^2 + 16g_3^2h_t^2 \\
&\quad \left. + \frac{2743}{162}g_1^4 + \frac{15}{2}g_2^4 - \frac{16}{9}g_3^4 + \frac{5}{3}g_1^2g_2^2 + \frac{136}{27}g_1^2g_3^2 + 8g_2^2g_3^2 \right), \\
16\pi^2 \frac{dh_b^2}{dt} &= h_b^2 \left(6h_b^2 + h_t^2 + h_\tau^2 + \lambda^2 - \frac{7}{9}g_1^2 - 3g_2^2 - \frac{16}{3}g_3^2 \right) \\
&\quad + \frac{h_b^2}{16\pi^2} \left(-22h_b^4 - 5h_t^4 - 3h_\tau^4 - 3\lambda^4 - 5h_b^2h_t^2 - 3h_b^2h_\tau^2 - 3h_b^2\lambda^2 \right. \\
&\quad - 4h_t^2\lambda^2 - 2\lambda^2\kappa^2 + \frac{2}{3}g_1^2h_b^2 + \frac{4}{3}g_1^2h_t^2 + 2g_1^2h_\tau^2 + 6g_2^2h_b^2 + 16g_3^2h_b^2 \\
&\quad \left. + \frac{1435}{162}g_1^4 + \frac{15}{2}g_2^4 - \frac{16}{9}g_3^4 + \frac{5}{3}g_1^2g_2^2 + \frac{40}{27}g_1^2g_3^2 + 8g_2^2g_3^2 \right), \\
16\pi^2 \frac{dh_\tau^2}{dt} &= h_\tau^2 \left(4h_\tau^2 + 3h_b^2 + \lambda^2 - 3g_1^2 - 3g_2^2 \right) \\
&\quad + \frac{h_\tau^2}{16\pi^2} \left(-10h_\tau^4 - 9h_b^4 - 3\lambda^4 - 9h_\tau^2h_b^2 - 3h_\tau^2\lambda^2 - 3h_t^2h_b^2 - 3h_t^2\lambda^2 \right. \\
&\quad \left. - 2\lambda^2\kappa^2 + 2g_1^2h_\tau^2 - \frac{2}{3}g_1^2h_b^2 + 6g_2^2h_\tau^2 + 16g_3^2h_b^2 + \frac{75}{2}g_1^4 + \frac{15}{2}g_2^4 + 3g_1^2g_2^2 \right),
\end{aligned}$$

$$\begin{aligned}
16\pi^2 \frac{d\lambda^2}{dt} &= \lambda^2 \left(3h_t^2 + 3h_b^2 + h_\tau^2 + 4\lambda^2 + 2\kappa^2 - g_1^2 - 3g_2^2 \right) \\
&+ \frac{\lambda^2}{16\pi^2} \left(-10\lambda^4 - 9h_t^4 - 9h_b^4 - 3h_\tau^4 - 8\kappa^4 - 9\lambda^2 h_t^2 - 9\lambda^2 h_b^2 \right. \\
&- 3\lambda^2 h_\tau^2 - 12\lambda^2 \kappa^2 - 6h_t^2 h_b^2 + 2g_1^2 \lambda^2 + \frac{4}{3}g_1^2 h_t^2 - \frac{2}{3}g_1^2 h_b^2 + 2g_1^2 h_\tau^2 \\
&+ \left. 6g_2^2 \lambda^2 + 16g_3^2 h_t^2 + 16g_3^2 h_b^2 + \frac{23}{2}g_1^4 + \frac{15}{2}g_2^4 + 3g_1^2 g_2^2 \right), \\
16\pi^2 \frac{d\kappa^2}{dt} &= \kappa^2 \left(6\lambda^2 + 6\kappa^2 \right) + \frac{\kappa^2}{16\pi^2} \left(-24\kappa^4 - 12\lambda^4 - 24\kappa^2 \lambda^2 \right. \\
&- \left. 18h_t^2 \lambda^2 - 18h_b^2 \lambda^2 - 6h_\tau^2 \lambda^2 + 6g_1^2 \lambda^2 + 18g_2^2 \lambda^2 \right). \tag{B.1}
\end{aligned}$$

B.2 Gaugino masses

The RGEs to two-loop order can be found in [532, 533].

$$\begin{aligned}
16\pi^2 \frac{dM_1}{dt} &= 11g_1^2 M_1 + \frac{g_1^2}{16\pi^2} \left(\frac{398}{9}g_1^2 M_1 + 9g_2^2 (M_1 + M_2) + \frac{88}{3}g_3^2 (M_1 + M_3) \right. \\
&+ \left. \frac{26}{3}h_t^2 (A_t - M_1) + \frac{14}{3}h_b^2 (A_b - M_1) + 6h_\tau^2 (A_\tau - M_1) + 2\lambda^2 (A_\lambda - M_1) \right), \\
16\pi^2 \frac{dM_2}{dt} &= g_2^2 M_2 + \frac{g_2^2}{16\pi^2} \left(3g_1^2 (M_1 + M_2) + 50g_2^2 M_2 + 24g_3^2 (M_2 + M_3) \right. \\
&+ \left. 6h_t^2 (A_t - M_2) + 6h_b^2 (A_b - M_2) + 2h_\tau^2 (A_\tau - M_2) + 2\lambda^2 (A_\lambda - M_2) \right), \\
16\pi^2 \frac{dM_3}{dt} &= -3g_3^2 M_3 + \frac{g_3^2}{16\pi^2} \left(\frac{11}{3}g_1^2 (M_1 + M_3) + 9g_2^2 (M_2 + M_3) + 28g_3^2 M_3 \right. \\
&+ \left. 4h_t^2 (A_t - M_3) + 4h_b^2 (A_b - M_3) \right). \tag{B.2}
\end{aligned}$$

B.3 Trilinear couplings

The two-loop β -functions are known for the parameters of a general softly broken SUSY theory [94]. We concentrate on the trilinear couplings involving the squarks and sleptons of the third family; those of the first two families do not play an important phenomenological rôle, except for the muon trilinear coupling A_μ ($\equiv A_{e2}$) which enters the muon anomalous magnetic momentum $(g-2)_\mu$.

$$\begin{aligned}
16\pi^2 \frac{dA_t}{dt} &= 6h_t^2 A_t + h_b^2 A_b + \lambda^2 A_\lambda + \frac{13}{9} g_1^2 M_1 + 3g_2^2 M_2 + \frac{16}{3} g_3^2 M_3 \\
&+ \frac{1}{16\pi^2} \left(-44h_t^4 A_t - 10h_b^4 A_b - 6\lambda^4 A_\lambda - 5h_t^2 h_b^2 (A_t + A_b) \right. \\
&- 3h_t^2 \lambda^2 (A_t + A_\lambda) - h_b^2 h_\tau^2 (A_b + A_\tau) - 4h_b^2 \lambda^2 (A_b + A_\lambda) \\
&- h_\tau^2 \lambda^2 (A_\tau + A_\lambda) - 2\lambda^2 \kappa^2 (A_\lambda + A_\kappa) + 2g_1^2 h_t^2 (A_t - M_1) \\
&+ \frac{2}{3} g_1^2 h_b^2 (A_b - M_1) + 6g_2^2 h_t^2 (A_t - M_2) + 16g_3^2 h_t^2 (A_t - M_3) \\
&- \frac{2743}{81} g_1^4 M_1 - 15g_2^4 M_2 + \frac{32}{9} g_3^4 M_3 - \frac{5}{3} g_1^2 g_2^2 (M_1 + M_2) \\
&\left. - \frac{136}{27} g_1^2 g_3^2 (M_1 + M_3) - 8g_2^2 g_3^2 (M_2 + M_3) \right), \\
16\pi^2 \frac{dA_b}{dt} &= 6h_b^2 A_b + h_t^2 A_t + h_\tau^2 A_\tau + \lambda^2 A_\lambda + \frac{7}{9} g_1^2 M_1 + 3g_2^2 M_2 + \frac{16}{3} g_3^2 M_3 \\
&+ \frac{1}{16\pi^2} \left(-44h_b^4 A_b - 10h_t^4 A_t - 6h_\tau^4 A_\tau - 6\lambda^4 A_\lambda - 5h_b^2 h_t^2 (A_b + A_t) \right. \\
&- 3h_b^2 h_\tau^2 (A_b + A_\tau) - 3h_b^2 \lambda^2 (A_b + A_\lambda) - 4h_t^2 \lambda^2 (A_t + A_\lambda) \\
&- 2\lambda^2 \kappa^2 (A_\lambda + A_\kappa) + \frac{2}{3} g_1^2 h_b^2 (A_b - M_1) + \frac{4}{3} g_1^2 h_t^2 (A_t - M_1) \\
&+ 2g_1^2 h_\tau^2 (A_\tau - M_1) + 6g_2^2 h_b^2 (A_b - M_2) + 16g_3^2 h_b^2 (A_b - M_3) \\
&- \frac{1435}{81} g_1^4 M_1 - 15g_2^4 M_2 + \frac{32}{9} g_3^4 M_3 - \frac{5}{3} g_1^2 g_2^2 (M_1 + M_2) \\
&\left. - \frac{40}{27} g_1^2 g_3^2 (M_1 + M_3) - 8g_2^2 g_3^2 (M_2 + M_3) \right), \\
16\pi^2 \frac{dA_\tau}{dt} &= 4h_\tau^2 A_\tau + 3h_b^2 A_b + \lambda^2 A_\lambda + 3g_1^2 M_1 + 3g_2^2 M_2 + \frac{1}{16\pi^2} \left(\right. \\
&- 20h_\tau^4 A_\tau - 18h_b^4 A_b - 6\lambda^4 A_\lambda - 9h_\tau^2 h_b^2 (A_\tau + A_b) - 3h_\tau^2 \lambda^2 (A_\tau + A_\lambda) \\
&- 3h_t^2 h_b^2 (A_t + A_b) - 3h_t^2 \lambda^2 (A_t + A_\lambda) - 2\lambda^2 \kappa^2 (A_\lambda + A_\kappa) \\
&+ 2g_1^2 h_\tau^2 (A_\tau - M_1) - \frac{2}{3} g_1^2 h_b^2 (A_b - M_1) + 6g_2^2 h_\tau^2 (A_\tau - M_2) \\
&\left. + 16g_3^2 h_b^2 (A_b - M_3) - 75g_1^4 M_1 - 15g_2^4 M_2 - 3g_1^2 g_2^2 (M_1 + M_2) \right), \\
16\pi^2 \frac{dA_\mu}{dt} &= 3h_b^2 A_b + h_\tau^2 A_\tau + \lambda^2 A_\lambda + 3g_1^2 M_1 + 3g_2^2 M_2 + \frac{1}{16\pi^2} \left(\right. \\
&- 18h_b^4 A_b - 6h_\tau^4 A_\tau - 6\lambda^4 A_\lambda - 3h_t^2 h_b^2 (A_t + A_b) - 3h_t^2 \lambda^2 (A_t + A_\lambda) \\
&- 2\lambda^2 \kappa^2 (A_\lambda + A_\kappa) - \frac{2}{3} g_1^2 h_b^2 (A_b - M_1) + 2g_1^2 h_\tau^2 (A_\tau - M_1) \\
&\left. + 16g_3^2 h_b^2 (A_b - M_3) - 75g_1^4 M_1 - 15g_2^4 M_2 - 3g_1^2 g_2^2 (M_1 + M_2) \right),
\end{aligned}$$

$$\begin{aligned}
16\pi^2 \frac{dA_\lambda}{dt} &= 4\lambda^2 A_\lambda + 3h_t^2 A_t + 3h_b^2 A_b + h_\tau^2 A_\tau + 2\kappa^2 A_\kappa + g_1^2 M_1 + 3g_2^2 M_2 \\
&+ \frac{1}{16\pi^2} \left(-20\lambda^4 A_\lambda - 18h_t^4 A_t - 18h_b^4 A_b - 6h_\tau^4 A_\tau - 16\kappa^4 A_\kappa \right. \\
&- 9\lambda^2 h_t^2 (A_\lambda + A_t) - 9\lambda^2 h_b^2 (A_\lambda + A_b) - 3\lambda^2 h_\tau^2 (A_\lambda + A_\tau) \\
&- 12\lambda^2 \kappa^2 (A_\lambda + A_\kappa) - 6h_t^2 h_b^2 (A_t + A_b) + 2g_1^2 \lambda^2 (A_\lambda - M_1) \\
&+ \frac{4}{3} g_1^2 h_t^2 (A_t - M_1) - \frac{2}{3} g_1^2 h_b^2 (A_b - M_1) + 2g_1^2 h_\tau^2 (A_\tau - M_1) \\
&+ 6g_2^2 \lambda^2 (A_\lambda - M_2) + 16g_3^2 h_t^2 (A_t - M_3) + 16g_3^2 h_b^2 (A_b - M_3) \\
&\left. - 23g_1^4 M_1 - 15g_2^4 M_2 - 3g_1^2 g_2^2 (M_1 + M_2) \right), \\
16\pi^2 \frac{dA_\kappa}{dt} &= 6\kappa^2 A_\kappa + 6\lambda^2 A_\lambda + \frac{1}{16\pi^2} \left(-48\kappa^4 A_\kappa - 24\lambda^4 A_\lambda \right. \\
&- 24\kappa^2 \lambda^2 (A_\kappa + A_\lambda) - 18h_t^2 \lambda^2 (A_t + A_\lambda) - 18h_b^2 \lambda^2 (A_b + A_\lambda) \\
&\left. - 6h_\tau^2 \lambda^2 (A_\tau + A_\lambda) + 6g_1^2 \lambda^2 (A_\lambda - M_1) + 18g_2^2 \lambda^2 (A_\lambda - M_2) \right). \quad (\text{B.3})
\end{aligned}$$

B.4 Squark and slepton masses

Let us define the following quantities:

$$\begin{aligned}
M_t^2 &= m_{Q_3}^2 + m_{U_3}^2 + m_{H_u}^2 + A_t^2, \\
M_b^2 &= m_{Q_3}^2 + m_{D_3}^2 + m_{H_d}^2 + A_b^2, \\
M_\tau^2 &= m_{L_3}^2 + m_{E_3}^2 + m_{H_d}^2 + A_\tau^2, \\
M_\lambda^2 &= m_{H_u}^2 + m_{H_d}^2 + m_S^2 + A_\lambda^2, \\
M_\kappa^2 &= 3m_S^2 + A_\kappa^2, \\
\xi &= \text{Tr}[\mathbf{m}_Q^2 - 2\mathbf{m}_U^2 + \mathbf{m}_D^2 - \mathbf{m}_L^2 + \mathbf{m}_E^2] + m_{H_u}^2 - m_{H_d}^2, \\
\xi' &= h_t^2 (-m_{Q_3}^2 + 4m_{U_3}^2 - 3m_{H_u}^2) + h_b^2 (-m_{Q_3}^2 - 2m_{D_3}^2 + 3m_{H_d}^2) \\
&+ h_\tau^2 (m_{L_3}^2 - 2m_{E_3}^2 + m_{H_d}^2) + \lambda^2 (m_{H_d}^2 - m_{H_u}^2) \\
&+ g_1^2 \left(\text{Tr} \left[\frac{1}{18} \mathbf{m}_Q^2 - \frac{16}{9} \mathbf{m}_U^2 + \frac{2}{9} \mathbf{m}_D^2 - \frac{1}{2} \mathbf{m}_L^2 + 2\mathbf{m}_E^2 \right] + \frac{1}{2} (m_{H_u}^2 - m_{H_d}^2) \right) \\
&+ \frac{3}{2} g_2^2 \left(\text{Tr} [\mathbf{m}_Q^2 - \mathbf{m}_L^2] + m_{H_u}^2 - m_{H_d}^2 \right) + \frac{8}{3} g_3^2 \text{Tr} [\mathbf{m}_Q^2 - 2\mathbf{m}_U^2 + \mathbf{m}_D^2], \\
\sigma_1 &= g_1^2 \left(\text{Tr} \left[\frac{1}{3} \mathbf{m}_Q^2 + \frac{8}{3} \mathbf{m}_U^2 + \frac{2}{3} \mathbf{m}_D^2 + \mathbf{m}_L^2 + 2\mathbf{m}_E^2 \right] + m_{H_u}^2 + m_{H_d}^2 \right), \\
\sigma_2 &= g_2^2 \left(\text{Tr} [3\mathbf{m}_Q^2 + \mathbf{m}_L^2] + m_{H_u}^2 + m_{H_d}^2 \right), \\
\sigma_3 &= g_3^2 \text{Tr} [2\mathbf{m}_Q^2 + \mathbf{m}_U^2 + \mathbf{m}_D^2]. \quad (\text{B.4})
\end{aligned}$$

where \mathbf{m} denote matrices in family space. The two-loop RGEs then read

$$\begin{aligned}
16\pi^2 \frac{dm_{Q_a}^2}{dt} &= \delta_{a3} h_t^2 M_t^2 + \delta_{a3} h_b^2 M_b^2 - \frac{1}{9} g_1^2 M_1^2 - 3g_2^2 M_2^2 - \frac{16}{3} g_3^2 M_3^2 + \frac{1}{6} g_1^2 \xi \\
&+ \frac{1}{16\pi^2} \left(-10\delta_{a3} h_t^4 (M_t^2 + A_t^2) - 10\delta_{a3} h_b^4 (M_b^2 + A_b^2) \right. \\
&- \delta_{a3} h_t^2 \lambda^2 (M_t^2 + M_\lambda^2 + 2A_t A_\lambda) - \delta_{a3} h_b^2 h_\tau^2 (M_b^2 + M_\tau^2 + 2A_b A_\tau) \\
&- \delta_{a3} h_b^2 \lambda^2 (M_b^2 + M_\lambda^2 + 2A_b A_\lambda) + \frac{4}{3} \delta_{a3} g_1^2 h_t^2 (M_t^2 - 2M_1(A_t - M_1)) \\
&+ \frac{2}{3} \delta_{a3} g_1^2 h_b^2 (M_b^2 - 2M_1(A_b - M_1)) + \frac{199}{54} g_1^4 M_1^2 + \frac{33}{2} g_2^4 M_2^2 - \frac{64}{3} g_3^4 M_3^2 \\
&+ \frac{1}{3} g_1^2 g_2^2 (M_1^2 + M_2^2 + M_1 M_2) + \frac{16}{27} g_1^2 g_3^2 (M_1^2 + M_3^2 + M_1 M_3) \\
&\left. + 16g_2^2 g_3^2 (M_2^2 + M_3^2 + M_2 M_3) + \frac{1}{3} g_1^2 \xi' + \frac{1}{18} g_1^2 \sigma_1 + \frac{3}{2} g_2^2 \sigma_2 + \frac{8}{3} g_3^2 \sigma_3 \right), \\
16\pi^2 \frac{dm_{U_a}^2}{dt} &= 2\delta_{a3} h_t^2 M_t^2 - \frac{16}{9} g_1^2 M_1^2 - \frac{16}{3} g_3^2 M_3^2 - \frac{2}{3} g_1^2 \xi \\
&+ \frac{1}{16\pi^2} \left(-16\delta_{a3} h_t^4 (M_t^2 + A_t^2) - 2\delta_{a3} h_t^2 h_b^2 (M_t^2 + M_b^2 + 2A_t A_b) \right. \\
&- 2\delta_{a3} h_t^2 \lambda^2 (M_t^2 + M_\lambda^2 + 2A_t A_\lambda) - \frac{2}{3} \delta_{a3} g_1^2 h_t^2 (M_t^2 - 2M_1(A_t - M_1)) \\
&+ 6\delta_{a3} g_2^2 h_t^2 (M_t^2 - 2M_2(A_t - M_2)) + \frac{1712}{27} g_1^4 M_1^2 - \frac{64}{3} g_3^4 M_3^2 \\
&\left. + \frac{256}{27} g_1^2 g_3^2 (M_1^2 + M_3^2 + M_1 M_3) - \frac{4}{3} g_1^2 \xi' + \frac{8}{9} g_1^2 \sigma_1 + \frac{8}{3} g_3^2 \sigma_3 \right), \\
16\pi^2 \frac{dm_{D_a}^2}{dt} &= 2\delta_{a3} h_b^2 M_b^2 - \frac{4}{9} g_1^2 M_1^2 - \frac{16}{3} g_3^2 M_3^2 + \frac{1}{3} g_1^2 \xi \\
&+ \frac{1}{16\pi^2} \left(-16\delta_{a3} h_b^4 (M_b^2 + A_b^2) - 2\delta_{a3} h_b^2 h_t^2 (M_b^2 + M_t^2 + 2A_b A_t) \right. \\
&- 2\delta_{a3} h_b^2 h_\tau^2 (M_b^2 + M_\tau^2 + 2A_b A_\tau) - 2\delta_{a3} h_b^2 \lambda^2 (M_b^2 + M_\lambda^2 + 2A_b A_\lambda) \\
&+ \frac{2}{3} \delta_{a3} g_1^2 h_b^2 (M_b^2 - 2M_1(A_b - M_1)) + 6\delta_{a3} g_2^2 h_b^2 (M_b^2 - 2M_2(A_b - M_2)) \\
&+ \frac{404}{27} g_1^4 M_1^2 - \frac{64}{3} g_3^4 M_3^2 + \frac{64}{27} g_1^2 g_3^2 (M_1^2 + M_3^2 + M_1 M_3) \\
&\left. + \frac{2}{3} g_1^2 \xi' + \frac{2}{9} g_1^2 \sigma_1 + \frac{8}{3} g_3^2 \sigma_3 \right), \\
16\pi^2 \frac{dm_{L_a}^2}{dt} &= \delta_{a3} h_\tau^2 M_\tau^2 - g_1^2 M_1^2 - 3g_2^2 M_2^2 - \frac{1}{2} g_1^2 \xi + \frac{1}{16\pi^2} \left(-6\delta_{a3} h_\tau^4 (M_\tau^2 + A_\tau^2) \right. \\
&- 3\delta_{a3} h_\tau^2 h_b^2 (M_\tau^2 + M_b^2 + 2A_\tau A_b) - \delta_{a3} h_\tau^2 \lambda^2 (M_\tau^2 + M_\lambda^2 + 2A_\tau A_\lambda) \\
&+ 2\delta_{a3} g_1^2 h_\tau^2 (M_\tau^2 - 2M_1(A_\tau - M_1)) + \frac{69}{2} g_1^4 M_1^2 + \frac{33}{2} g_2^4 M_2^2 \\
&\left. + 3g_1^2 g_2^2 (M_1^2 + M_2^2 + M_1 M_2) - g_1^2 \xi' + \frac{1}{2} g_1^2 \sigma_1 + \frac{3}{2} g_2^2 \sigma_2 \right),
\end{aligned}$$

$$\begin{aligned}
16\pi^2 \frac{dm_{E_a}^2}{dt} &= 2\delta_{a3}h_\tau^2 M_\tau^2 - 4g_1^2 M_1^2 + g_1^2 \xi + \frac{1}{16\pi^2} \left(-8\delta_{a3}h_\tau^4 (M_\tau^2 + A_\tau^2) \right. \\
&- 6\delta_{a3}h_\tau^2 h_b^2 (M_\tau^2 + M_b^2 + 2A_\tau A_b) - 2\delta_{a3}h_\tau^2 \lambda^2 (M_\tau^2 + M_\lambda^2 + 2A_\tau A_\lambda) \\
&- 2\delta_{a3}g_1^2 h_\tau^2 (M_\tau^2 - 2M_1(A_\tau - M_1)) + 6\delta_{a3}g_2^2 h_\tau^2 (M_\tau^2 - 2M_2(A_\tau - M_2)) \\
&\left. + 156g_1^4 M_1^2 + 2g_1^2 \xi' + 2g_1^2 \sigma_1 \right). \tag{B.5}
\end{aligned}$$

B.5 Higgs masses

Similarly, the two-loop RGEs for the Higgs soft masses are

$$\begin{aligned}
16\pi^2 \frac{dm_{H_u}^2}{dt} &= 3h_t^2 M_t^2 + \lambda^2 M_\lambda^2 - g_1^2 M_1^2 - 3g_2^2 M_2^2 + \frac{1}{2}g_1^2 \xi \\
&+ \frac{1}{16\pi^2} \left(-18h_t^4 (M_t^2 + A_t^2) - 6\lambda^4 (M_\lambda^2 + A_\lambda^2) \right. \\
&- 3h_t^2 h_b^2 (M_t^2 + M_b^2 + 2A_t A_b) - 3h_b^2 \lambda^2 (M_b^2 + M_\lambda^2 + 2A_b A_\lambda) \\
&- h_\tau^2 \lambda^2 (M_\tau^2 + M_\lambda^2 + 2A_\tau A_\lambda) - 2\lambda^2 \kappa^2 (M_\lambda^2 + M_\kappa^2 + 2A_\lambda A_\kappa) \\
&+ \frac{4}{3}g_1^2 h_t^2 (M_t^2 - 2M_1(A_t - M_1)) + 16g_3^2 h_t^2 (M_t^2 - 2M_3(A_t - M_3)) \\
&+ \frac{69}{2}g_1^4 M_1^2 + \frac{33}{2}g_2^4 M_2^2 + 3g_1^2 g_2^2 (M_1^2 + M_2^2 + M_1 M_2) \\
&\left. + g_1^2 \xi' + \frac{1}{2}g_1^2 \sigma_1 + \frac{3}{2}g_2^2 \sigma_2 \right), \\
16\pi^2 \frac{dm_{H_d}^2}{dt} &= 3h_b^2 M_b^2 + h_\tau^2 M_\tau^2 + \lambda^2 M_\lambda^2 - g_1^2 M_1^2 - 3g_2^2 M_2^2 - \frac{1}{2}g_1^2 \xi \\
&+ \frac{1}{16\pi^2} \left(-18h_b^4 (M_b^2 + A_b^2) - 6h_\tau^4 (M_\tau^2 + A_\tau^2) \right. \\
&- 6\lambda^4 (M_\lambda^2 + A_\lambda^2) - 3h_b^2 h_t^2 (M_b^2 + M_t^2 + 2A_b A_t) \\
&- 3h_t^2 \lambda^2 (M_t^2 + M_\lambda^2 + 2A_t A_\lambda) - 2\lambda^2 \kappa^2 (M_\lambda^2 + M_\kappa^2 + 2A_\lambda A_\kappa) \\
&- \frac{2}{3}g_1^2 h_b^2 (M_b^2 - 2M_1(A_b - M_1)) + 2g_1^2 h_\tau^2 (M_\tau^2 - 2M_1(A_\tau - M_1)) \\
&+ 16g_3^2 h_b^2 (M_b^2 - 2M_3(A_b - M_3)) + \frac{69}{2}g_1^4 M_1^2 + \frac{33}{2}g_2^4 M_2^2 \\
&\left. + 3g_1^2 g_2^2 (M_1^2 + M_2^2 + M_1 M_2) - g_1^2 \xi' + \frac{1}{2}g_1^2 \sigma_1 + \frac{3}{2}g_2^2 \sigma_2 \right), \\
16\pi^2 \frac{dm_S^2}{dt} &= 2\lambda^2 M_\lambda^2 + 2\kappa^2 M_\kappa^2 + \frac{1}{16\pi^2} \left(-8\lambda^4 (M_\lambda^2 + A_\lambda^2) - 16\kappa^4 (M_\kappa^2 + A_\kappa^2) \right. \\
&- 6\lambda^2 h_t^2 (M_\lambda^2 + M_t^2 + 2A_\lambda A_t) - 6\lambda^2 h_b^2 (M_\lambda^2 + M_b^2 + 2A_\lambda A_b) \\
&- 2\lambda^2 h_\tau^2 (M_\lambda^2 + M_\tau^2 + 2A_\lambda A_\tau) - 8\lambda^2 \kappa^2 (M_\lambda^2 + M_\kappa^2 + 2A_\lambda A_\kappa) \\
&\left. + 2g_1^2 \lambda^2 (M_\lambda^2 - 2M_1(A_\lambda - M_1)) + 6g_2^2 \lambda^2 (M_\lambda^2 - 2M_2(A_\lambda - M_2)) \right). \tag{B.6}
\end{aligned}$$

B.6 Additional parameters of the general NMSSM

The two-loop RGEs for the SUSY conserving μ and μ' terms are

$$\begin{aligned}
32\pi^2 \frac{d\mu}{dt} &= \mu \left(3h_t^2 + 3h_b^2 + h_\tau^2 + 2\lambda^2 - g_1^2 - 3g_2^2 \right) + \frac{\mu}{16\pi^2} \left(-9h_t^4 - 9h_b^4 \right. \\
&\quad - 3h_\tau^4 - 6\lambda^4 - 6h_t^2 h_b^2 - 3h_t^2 \lambda^2 - 3h_b^2 \lambda^2 - h_\tau^2 \lambda^2 - 4\lambda^2 \kappa^2 + \frac{4}{3} g_1^2 h_t^2 \\
&\quad \left. - \frac{2}{3} g_1^2 h_b^2 + 2g_1^2 h_\tau^2 + 16g_3^2 h_t^2 + 16g_3^2 h_b^2 + \frac{23}{2} g_1^4 + \frac{15}{2} g_2^4 + 3g_1^2 g_2^2 \right), \\
16\pi^2 \frac{d\mu'}{dt} &= \mu' \left(2\lambda^2 + 2\kappa^2 \right) + \frac{\mu'}{16\pi^2} \left(-4\lambda^4 - 8\kappa^4 - 6h_t^2 \lambda^2 \right. \\
&\quad \left. - 6h_b^2 \lambda^2 - 2h_\tau^2 \lambda^2 - 8\lambda^2 \kappa^2 + 2g_1^2 \lambda^2 + 6g_2^2 \lambda^2 \right). \tag{B.7}
\end{aligned}$$

For the corresponding soft SUSY breaking terms m_3^2 and $m_S'^2$ (see (2.9)) we have

$$\begin{aligned}
32\pi^2 \frac{dm_3^2}{dt} &= 3h_t^2(m_3^2 + 2\mu A_t) + 3h_b^2(m_3^2 + 2\mu A_b) + h_\tau^2(m_3^2 + 2\mu A_\tau) \\
&\quad + 2\lambda^2(3m_3^2 + 2\mu A_\lambda) + 2\lambda\kappa m_S'^2 - g_1^2(m_3^2 - 2\mu M_1) - 3g_2^2(m_3^2 - 2\mu M_2) \\
&\quad + \frac{1}{16\pi^2} \left(-9h_t^4(m_3^2 + 4\mu A_t) - 9h_b^4(m_3^2 + 4\mu A_b) - 3h_\tau^4(m_3^2 + 4\mu A_\tau) \right. \\
&\quad - 2\lambda^4(7m_3^2 + 16\mu A_\lambda) - 6h_t^2 h_b^2(m_3^2 + 2\mu(A_t + A_b)) \\
&\quad - 3h_t^2 \lambda^2(5m_3^2 + 2\mu(3A_t + A_\lambda)) - 3h_b^2 \lambda^2(5m_3^2 + 2\mu(3A_b + A_\lambda)) \\
&\quad - h_\tau^2 \lambda^2(5m_3^2 + 2\mu(3A_\tau + A_\lambda)) - 4\lambda^2 \kappa^2(m_3^2 + 2\mu(A_\lambda + A_\kappa)) \\
&\quad - 8\lambda^3 \kappa(m_S'^2 + \mu' A_\lambda) - 8\lambda\kappa^3(m_S'^2 + \mu' A_\kappa) \\
&\quad + \frac{4}{3} g_1^2 h_t^2(m_3^2 + 2\mu(A_t - M_1)) - \frac{2}{3} g_1^2 h_b^2(m_3^2 + 2\mu(A_b - M_1)) \\
&\quad + 2g_1^2 h_\tau^2(m_3^2 + 2\mu(A_\tau - M_1)) + 4g_1^2 \lambda^2(m_3^2 - \mu M_1) \\
&\quad + 12g_2^2 \lambda^2(m_3^2 - \mu M_2) + 16g_3^2 h_t^2(m_3^2 + 2\mu(A_t - M_3)) \\
&\quad + 16g_3^2 h_b^2(m_3^2 + 2\mu(A_b - M_3)) + \frac{23}{2} g_1^4(m_3^2 - 4\mu M_1) \\
&\quad \left. + \frac{15}{2} g_2^4(m_3^2 - 4\mu M_2) + 3g_1^2 g_2^2(m_3^2 - 2\mu(M_1 + M_2)) \right), \\
16\pi^2 \frac{dm_S'^2}{dt} &= 2\lambda^2(m_S'^2 + 2\mu' A_\lambda) + 4\kappa^2(m_S'^2 + \mu' A_\kappa) + 4\lambda\kappa m_3^2 \\
&\quad + \frac{1}{16\pi^2} \left(-4\lambda^4(m_S'^2 + 4\mu' A_\lambda) - 8\kappa^4(2m_S'^2 + 5\mu' A_\kappa) \right. \\
&\quad - 8\lambda^2 \kappa^2(2m_S'^2 + \mu'(3A_\lambda + 2A_\kappa)) - 6\lambda^2 h_t^2(m_S'^2 + 2\mu'(A_\lambda + A_t)) \\
&\quad - 6\lambda^2 h_b^2(m_S'^2 + 2\mu'(A_\lambda + A_b)) - 2\lambda^2 h_\tau^2(m_S'^2 + 2\mu'(A_\lambda + A_\tau)) \\
&\quad - 8\lambda^3 \kappa(m_3^2 + \mu A_\lambda) - 12\lambda\kappa h_t^2(m_3^2 + \mu A_t) - 12\lambda\kappa h_b^2(m_3^2 + \mu A_b) \\
&\quad - 4\lambda\kappa h_\tau^2(m_3^2 + \mu A_\tau) + 4\lambda\kappa g_1^2(m_3^2 - \mu M_1) + 12\lambda\kappa g_2^2(m_3^2 - \mu M_2) \\
&\quad \left. + 2\lambda^2 g_1^2(m_S'^2 + 2\mu'(A_\lambda - M_1)) + 6\lambda^2 g_2^2(m_S'^2 + 2\mu'(A_\lambda - M_2)) \right). \tag{B.8}
\end{aligned}$$

Finally for the singlet tadpole terms, the two-loop RGEs read

$$\begin{aligned}
16\pi^2 \frac{d\xi_F}{dt} &= \xi_F \left(\lambda^2 + \kappa^2 \right) + \frac{\xi_F}{16\pi^2} \left(-2\lambda^4 - 4\kappa^4 - 3h_t^2 \lambda^2 \right. \\
&\quad \left. - 3h_b^2 \lambda^2 - h_\tau^2 \lambda^2 - 4\lambda^2 \kappa^2 + g_1^2 \lambda^2 + 3g_2^2 \lambda^2 \right), \\
16\pi^2 \frac{d\xi_S}{dt} &= \lambda^2 (\xi_S + 2A_\lambda \xi_F) + \kappa^2 (\xi_S + 2A_\kappa \xi_F) \\
&\quad + 2\lambda (m_3^2 (A_\lambda + \mu') + \mu (m_{H_u}^2 + m_{H_d}^2)) + \kappa (m_S'^2 (A_\kappa + \mu') + 2\mu' m_S^2) \\
&\quad + \frac{1}{16\pi^2} \left(-2\lambda^4 (\xi_S + 4A_\lambda \xi_F) - 4\kappa^4 (\xi_S + 4A_\kappa \xi_F) \right. \\
&\quad - 3\lambda^2 h_t^2 (\xi_S + 2(A_\lambda + A_t) \xi_F) - 3\lambda^2 h_b^2 (\xi_S + 2(A_\lambda + A_b) \xi_F) \\
&\quad - \lambda^2 h_\tau^2 (\xi_S + 2(A_\lambda + A_\tau) \xi_F) - 4\lambda^2 \kappa^2 (\xi_S + 2(A_\lambda + A_\kappa) \xi_F) \\
&\quad - 6\lambda h_t^2 \left(m_3^2 (A_\lambda + A_t + \mu') + \mu (M_t^2 + A_t (A_\lambda + \mu') + m_{H_u}^2 + m_{H_d}^2) \right) \\
&\quad - 6\lambda h_b^2 \left(m_3^2 (A_\lambda + A_b + \mu') + \mu (M_b^2 + A_b (A_\lambda + \mu') + m_{H_u}^2 + m_{H_d}^2) \right) \\
&\quad - 2\lambda h_\tau^2 \left(m_3^2 (A_\lambda + A_\tau + \mu') + \mu (M_\tau^2 + A_\tau (A_\lambda + \mu') + m_{H_u}^2 + m_{H_d}^2) \right) \\
&\quad - 4\lambda^3 \left(m_3^2 (2A_\lambda + \mu') + \mu (M_\lambda^2 + A_\lambda (A_\lambda + \mu') + m_{H_u}^2 + m_{H_d}^2) \right) \\
&\quad - 4\lambda^2 \kappa \left(m_S'^2 (A_\lambda + A_\kappa + \mu') + \mu' (M_\lambda^2 + A_\lambda (A_\kappa + \mu') + 2m_S^2) \right) \\
&\quad - 4\kappa^3 \left(m_S'^2 (2A_\kappa + \mu') + \mu' (M_\kappa^2 + A_\kappa (A_\kappa + \mu') + 2m_S^2) \right) \\
&\quad + \lambda g_1^2 \left(3m_3^2 (A_\lambda + \mu' - M_1) + 2\mu (m_{H_u}^2 + m_{H_d}^2 - A_\lambda M_1 - \mu' M_1 + 2M_1^2) \right. \\
&\quad \left. + \lambda (2\xi_F (A_\lambda - M_1) + \xi_S) \right) \\
&\quad + 3\lambda g_2^2 \left(3m_3^2 (A_\lambda + \mu' - M_2) + 2\mu (m_{H_u}^2 + m_{H_d}^2 - A_\lambda M_2 - \mu' M_2 + 2M_2^2) \right. \\
&\quad \left. + \lambda (2\xi_F (A_\lambda - M_2) + \xi_S) \right) \Big). \tag{B.9}
\end{aligned}$$

Appendix C. Radiative corrections to the Higgs masses

As explained in Section 3.2, the following steps are required for a systematic calculation of the radiative corrections to the Higgs masses in the present approach: i) the gauge and Yukawa couplings at the scale M_{SUSY} must be determined; ii) the effective potential and the Higgs wave function normalisation constants must be computed with an ultraviolet cutoff M_{SUSY} , and iii) the pole masses have to be evaluated from the effective action. Here we give the necessary formulae for the calculation of the radiative corrections to the Higgs masses to the order described in Section 3.2.

C.1 Yukawa and gauge couplings at the scale M_{SUSY}

First the Yukawa couplings (we confine ourselves to the quarks of the third family) have to be deduced from the pole masses. The running top quark Yukawa coupling at the scale m_t is given in terms of the top quark pole mass as²

$$h_t(m_t) = \frac{m_t^{\text{pole}}}{v_u} \left(1 + \frac{4\alpha_s(m_t)}{3\pi} + \frac{11\alpha_s^2(m_t)}{\pi^2} \right)^{-1}. \quad (\text{C.1})$$

For the computation of $h_t(M_{\text{SUSY}})$, we consider only contributions involving potentially large logarithms (the leading logarithmic approximation or LLA) as $\ln(M_{\text{SUSY}}^2/m_t^2)$ and more, which depend on the masses of the various particles appearing in the loops. For instance, a complete $SU(2)$ multiplet of Higgs bosons can have (nearly degenerate) masses given by M_A (the mass of the MSSM-like CP-odd scalar, cf. Section 2), if $M_A^2 \equiv \mathcal{M}_{P,11}^2 \sim \mathcal{M}_{\pm}^2 \gg M_Z^2$. These states will only contribute to the radiative corrections to h_t from scales $Q^2 > M_A^2$. In addition, charginos and neutralinos can have masses below M_{SUSY} and induce threshold effects; within the LLA it is sufficient, however, to approximate their masses by the diagonal elements of the corresponding mass matrices (2.35) and (2.32). Then, pure gaugino and higgsino states appear simultaneously in the Higgs wave function renormalisation diagrams, and only the mass of the heavier state appears in the potentially large logarithm. (On the other hand, effects from squark/gluino loops can be neglected in the LLA, since the squark masses define the ultraviolet cutoff M_{SUSY} .) One obtains (with

²Here we use the $\overline{\text{MS}}$ relation between the pole mass and the running mass. In the $\overline{\text{DR}}$ scheme the second factor $4/3\pi$ would read $5/3\pi$, which would decrease $h_t(m_t)$, and hence the lightest Higgs mass, by $\sim 1\%$. Since we have not included subdominant (single) logarithms in the two-loop corrections to the effective potential we are not sensitive to the scheme in which the running top quark mass is defined, which leads to a theoretical error of $\gtrsim 1\%$ on the mass of the lightest Higgs.

$\nu' \equiv \kappa s + \mu'/2$, and the scales of couplings without arguments are arbitrary in the LLA)

$$\begin{aligned}
h_t(M_{\text{SUSY}}) &= h_t(m_t) \left(1 + \frac{7}{4\pi} \alpha_s(m_t) \ln(M_{\text{SUSY}}^2/m_t^2) \right)^{-4/7} \\
&\times \left(1 + \frac{1}{64\pi^2} \left[\left(9h_t^2 + h_b^2 - \frac{17}{6}g_1^2 - \frac{9}{2}g_2^2 \right) \ln(M_{\text{SUSY}}^2/m_t^2) \right. \right. \\
&+ (-9 \cos^2 \beta h_t^2 + (3 \cos^2 \beta - 1)h_b^2 + 2 \cos^2 \beta h_\tau^2) \ln(M_A^2/m_t^2) \\
&- 2\lambda^2 \ln(\max(\mu_{\text{eff}}^2, 4\nu'^2)/M_{\text{SUSY}}^2) - g_1^2 \ln(\max(\mu_{\text{eff}}^2, M_1^2)/M_{\text{SUSY}}^2) \\
&\left. \left. - 3g_2^2 \ln(\max(\mu_{\text{eff}}^2, M_2^2)/M_{\text{SUSY}}^2) \right] \right). \tag{C.2}
\end{aligned}$$

In the case of h_b , we assume a given value of the running b quark mass $m_b(M_Z)$ (typically ~ 2.9 GeV), from which $h_b(M_Z)$ can be obtained as in (A.11). Integrating the one-loop (QCD) RGEs, $h_b(m_t)$ is given by

$$h_b(m_t) = h_b(M_Z) \left(1 - \frac{23}{12\pi} \alpha_s(m_t) \ln \frac{m_t^2}{M_Z^2} \right)^{12/23}, \tag{C.3}$$

and subsequently $h_b(M_{\text{SUSY}})$ by

$$\begin{aligned}
h_b(M_{\text{SUSY}}) &= h_b(m_t) \left(1 + \frac{7}{4\pi} \alpha_s(m_t) \ln(M_{\text{SUSY}}^2/m_t^2) \right)^{-4/7} \\
&\times \left(1 + \frac{1}{64\pi^2} \left[\left(9h_b^2 + h_t^2 + 2h_\tau^2 - \frac{5}{6}g_1^2 - \frac{9}{2}g_2^2 \right) \ln(M_{\text{SUSY}}^2/m_t^2) \right. \right. \\
&+ (-9 \sin^2 \beta h_b^2 + (3 \sin^2 \beta - 1)h_t^2 - 2 \sin^2 \beta h_\tau^2) \ln(M_A^2/m_t^2) \\
&- 2\lambda^2 \ln(\max(\mu_{\text{eff}}^2, 4\nu'^2)/M_{\text{SUSY}}^2) - g_1^2 \ln(\max(\mu_{\text{eff}}^2, M_1^2)/M_{\text{SUSY}}^2) \\
&\left. \left. - 3g_2^2 \ln(\max(\mu_{\text{eff}}^2, M_2^2)/M_{\text{SUSY}}^2) \right] \right). \tag{C.4}
\end{aligned}$$

The electroweak gauge couplings are assumed as given at the scale M_Z e.g. in the on-shell scheme in terms of the Fermi coupling G_F , M_Z and M_W as $g_2^2 = 4\sqrt{2}G_F M_W^2$, $g_1^2 = 4\sqrt{2}G_F(M_Z^2 - M_W^2)$ (the renormalisation scheme is irrelevant in the LLA). For their computation at the scale M_{SUSY} , we include threshold effects of the potentially heavy $SU(2)$ Higgs multiplet with a mass M_A , higgsinos with mass μ_{eff} , $SU(2)$ gauginos with mass M_2 , and sleptons of a common mass $M_{\tilde{l}}$ (in the LLA). Squarks with masses $\sim M_{\text{SUSY}}$ do not induce threshold effects at scales below M_{SUSY} , and the top quark threshold is not considered since $m_t \sim M_Z$ inside logarithms in the LLA.

$$\begin{aligned}
g_1^2(M_{\text{SUSY}}) &= g_1^2(M_Z) \left(1 - \frac{g_1^2}{16\pi^2} \left[\frac{41}{6} \ln(M_{\text{SUSY}}^2/M_Z^2) + \frac{1}{6} \ln(M_{\text{SUSY}}^2/M_A^2) \right. \right. \\
&\left. \left. + \frac{2}{3} \ln(M_{\text{SUSY}}^2/\mu_{\text{eff}}^2) + \frac{3}{2} \ln(M_{\text{SUSY}}^2/M_{\tilde{l}}^2) \right] \right)^{-1}, \tag{C.5}
\end{aligned}$$

$$\begin{aligned}
g_2^2(M_{\text{SUSY}}) &= g_2^2(M_Z) \left(1 + \frac{g_2^2}{16\pi^2} \left[\frac{19}{6} \ln(M_{\text{SUSY}}^2/M_Z^2) - \frac{1}{6} \ln(M_{\text{SUSY}}^2/M_A^2) \right. \right. \\
&\left. \left. - \frac{2}{3} \ln(M_{\text{SUSY}}^2/\mu_{\text{eff}}^2) - \ln(M_{\text{SUSY}}^2/M_{\text{slept}}^2) - \ln(M_{\text{SUSY}}^2/M_2^2) \right] \right)^{-1}. \tag{C.6}
\end{aligned}$$

The strong gauge coupling at M_{SUSY} is computed somewhat more precisely; the one-loop RGE is solved exactly below and above the top quark threshold:

$$g_3^2(m_t) = \frac{g_3^2(M_Z)}{1 + \frac{23g_3^2(M_Z)}{48\pi^2} \ln(m_t^2/M_Z^2)}, \quad (\text{C.7})$$

$$g_3^2(M_{\text{SUSY}}) = g_3^2(m_t) \left(1 + \frac{g_3^2(m_t)}{16\pi^2} \left[7 \ln(M_{\text{SUSY}}^2/m_t^2) - 2 \ln(M_{\text{SUSY}}^2/M_3^2) \right] \right)^{-1}. \quad (\text{C.8})$$

C.2 Higgs wave function renormalisation constants

Within the present approximation it suffices to consider wave function renormalisation constants for the weak eigenstates H_u , H_d and S . In the case of contributions from quarks and leptons of the third family we use an infrared cutoff m_t^2 , which simplifies the remaining top- and bottom-quark induced corrections to the pole masses in section C.4 below. In the case of contributions from gauge bosons, charginos and neutralinos we confine ourselves to potentially large logarithms, i.e. we use $M_W \approx M_Z$ and the diagonal elements of the chargino/neutralino mass matrices inside the logarithms.

If $M_A^2 \gg m_t^2$, we include the corresponding threshold effects: the wave function renormalisation constants of heavy Higgs states on their mass shell $\sim M_A^2$ do not receive contributions from momenta $< M_A^2$ in the LLA, and subsequently this effect is translated into the wave function renormalisation constants for the weak eigenstates H_u , H_d via a rotation by the angle β . Then we obtain for Z_{H_u} , Z_{H_d} and Z_S in the Landau gauge (we recall $\nu' \equiv \kappa s + \mu'/2$):

$$\begin{aligned} Z_{H_u} &= 1 + \frac{1}{16\pi^2} \left(3h_t^2 \ln(M_{\text{SUSY}}^2/m_t^2) - \frac{3}{4}(g_1^2 + 3g_2^2) \ln(M_{\text{SUSY}}^2/M_Z^2) \right. \\ &\quad + \cos^2 \beta (3h_b^2 + h_\tau^2 - 3h_t^2) \ln(M_A^2/m_t^2) + \frac{g_1^2}{2} \ln(M_{\text{SUSY}}^2/\max(\mu_{\text{eff}}^2, M_1^2)) \\ &\quad \left. + \frac{3g_2^2}{2} \ln(M_{\text{SUSY}}^2/\max(\mu_{\text{eff}}^2, M_2^2)) + \lambda^2 \ln(M_{\text{SUSY}}^2/\max(\mu_{\text{eff}}^2, 4\nu'^2)) \right), \quad (\text{C.9}) \end{aligned}$$

$$\begin{aligned} Z_{H_d} &= 1 + \frac{1}{16\pi^2} \left((3h_b^2 + h_\tau^2) \ln(M_{\text{SUSY}}^2/m_t^2) - \frac{3}{4}(g_1^2 + 3g_2^2) \ln(M_{\text{SUSY}}^2/M_Z^2) \right. \\ &\quad + \sin^2 \beta (3h_t^2 - h_\tau^2 - 3h_b^2) \ln(M_A^2/m_t^2) + \frac{g_1^2}{2} \ln(M_{\text{SUSY}}^2/\max(\mu_{\text{eff}}^2, M_1^2)) \\ &\quad \left. + \frac{3g_2^2}{2} \ln(M_{\text{SUSY}}^2/\max(\mu_{\text{eff}}^2, M_2^2)) + \lambda^2 \ln(M_{\text{SUSY}}^2/\max(\mu_{\text{eff}}^2, 4\nu'^2)) \right), \quad (\text{C.10}) \end{aligned}$$

$$Z_S = 1 + \frac{1}{8\pi^2} \left(\lambda^2 \ln(M_{\text{SUSY}}^2/\mu_{\text{eff}}^2) + \kappa^2 \ln(M_{\text{SUSY}}^2/4\nu'^2) \right). \quad (\text{C.11})$$

These wave function renormalisation constants will multiply the corresponding kinetic terms in the effective action, which results from the addition of quantum effects with $Q^2 < M_{\text{SUSY}}^2$ to the “tree level” Lagrangian, wherein all couplings are defined at $Q^2 = M_{\text{SUSY}}^2$. In the next subsection, the contributions of quantum effects with $Q^2 < M_{\text{SUSY}}^2$ to the Higgs mass matrices are considered, in which – to start with – the Higgs fields are *not*

(yet) properly normalised. We will denote the Higgs vevs before the rescaling by square roots of the wave function renormalisation constants by $v_u(Q)$, $v_d(Q)$ and $s(Q)$, which are related to the vevs v_u , v_d and s of the properly normalised Higgs fields by

$$v_u(Q) = v_u/\sqrt{Z_{H_u}}, \quad v_d(Q) = v_d/\sqrt{Z_{H_d}}, \quad s(Q) = s/\sqrt{Z_S}. \quad (\text{C.12})$$

In the LLA, v_u and v_d are related to M_Z and the to quark and lepton masses as in (2.13) and (A.11).

C.3 Corrections to the Higgs mass matrices

According to the philosophy employed here, we denote subsequently by “tree level” mass matrices those of Section 2 where i) all couplings are defined at the scale $Q^2 = M_{\text{SUSY}}^2$, and ii) all Higgs vevs are given by $v_u(Q)$, $v_d(Q)$ and $s(Q)$ as defined in (C.12).

The one-loop corrections to the effective potential are given by the Coleman-Weinberg formula (with an ultraviolet cutoff M_{SUSY}^2 , and in the $\overline{\text{DR}}$ scheme in agreement with the two-loop RGEs for the soft SUSY breaking terms in Appendix B):

$$\Delta V_{\text{eff}} = \frac{1}{64\pi^2} \text{STr } M^4 \left[\ln (M^2/M_{\text{SUSY}}^2) - \frac{3}{2} \right] \quad (\text{C.13})$$

where the couplings and Higgs vevs in the mass matrices M of the particles in the loops are still those at the scale M_{SUSY}^2 . As a result of this procedure, the two-loop contributions to ΔV_{eff} and hence to the Higgs mass matrices – at least in the approximation considered here – become quite simple.

The radiative corrections to the effective potential affect also the three minimisation equations with respect to $v_u(Q)$, $v_d(Q)$ and $s(Q)$, which serve to replace the three parameters $m_{H_u}^2$, $m_{H_d}^2$ and m_S^2 (μ_{eff} , B_{eff} and \widehat{m}_3^2 were defined in (2.7) and (2.14)):

$$\begin{aligned} v_u(Q) & \left(m_{H_u}^2 + \mu_{\text{eff}}^2 + \lambda^2 v_d^2(Q) + \frac{g_1^2 + g_2^2}{4} (v_u^2(Q) - v_d^2(Q)) \right) \\ & - v_d(Q) (\lambda s(Q) B_{\text{eff}} + \widehat{m}_3^2) + \frac{1}{2} \Delta V_{\text{eff}, v_u(Q)} = 0, \\ v_d(Q) & \left(m_{H_d}^2 + \mu_{\text{eff}}^2 + \lambda^2 v_u^2(Q) + \frac{g_1^2 + g_2^2}{4} (v_d^2(Q) - v_u^2(Q)) \right) \\ & - v_u(Q) (\lambda s(Q) B_{\text{eff}} + \widehat{m}_3^2) + \frac{1}{2} \Delta V_{\text{eff}, v_d(Q)} = 0, \\ s(Q) & \left(m_S^2 + m_S'^2 + \mu'^2 + 2\kappa\xi_F + \kappa A_\kappa s(Q) + 2\kappa^2 s^2(Q) \right. \\ & \left. + \lambda^2 (v_u^2(Q) + v_d^2(Q)) - 2\lambda\kappa v_u(Q)v_d(Q) \right) \\ & + \xi_S + \xi_F \mu' - \lambda v_u(Q)v_d(Q)(A_\lambda + \mu') + \frac{1}{2} \Delta V_{\text{eff}, s(Q)} = 0. \end{aligned} \quad (\text{C.14})$$

The contributions of ΔV_{eff} to the Higgs mass matrices are obtained by appropriate second derivatives of ΔV_{eff} (with respect to $v_u(Q)$, $v_d(Q)$ and $s(Q)$ in the CP-even case), taking

care of the modified replaced mass terms:

$$\begin{aligned}
\Delta\mathcal{M}_{ij}^2 &= \frac{1}{2}\Delta V_{\text{eff},v_i v_j} \text{ for } i \neq j, \\
\Delta\mathcal{M}_{ii}^2 &= \frac{1}{2}\Delta V_{\text{eff},v_i v_i} - \frac{1}{2v_i}\Delta V_{\text{eff},v_i}, \\
v_i &\equiv v_u(Q), v_d(Q), s(Q).
\end{aligned} \tag{C.15}$$

Below we give the corrections to the Higgs mass matrices for the CP-even, CP-odd and charged Higgs masses, separately for each particle species in the loops.

CP-even scalars

Top and bottom (s)quarks

The top and bottom squark mass matrices have been given in (2.36) and (2.37) in Section 2. They are diagonalized as in (A.10) with the help of angles θ_T and θ_B such that the eigenstates have masses $M_{\tilde{t}_1}$ and $M_{\tilde{t}_2}$ with $M_{\tilde{t}_1} < M_{\tilde{t}_2}$ (and corresponding eigenstates $M_{\tilde{b}_1}$ and $M_{\tilde{b}_2}$). It is useful to define the following quantities in terms of the mass eigenvalues and mixing angles:

$$\begin{aligned}
f_t &= \frac{1}{M_{\tilde{t}_2}^2 - M_{\tilde{t}_1}^2} \left(M_{\tilde{t}_2}^2 \ln \left(\frac{M_{\tilde{t}_2}^2}{M_{\text{SUSY}}^2} \right) - M_{\tilde{t}_1}^2 \ln \left(\frac{M_{\tilde{t}_1}^2}{M_{\text{SUSY}}^2} \right) \right) - 1, \\
g_t &= \sin^2 2\theta_T \left(\frac{M_{\tilde{t}_2}^2 + M_{\tilde{t}_1}^2}{M_{\tilde{t}_2}^2 - M_{\tilde{t}_1}^2} \ln \left(\frac{M_{\tilde{t}_2}^2}{M_{\tilde{t}_1}^2} \right) - 2 \right), \\
e_t &= -m_t \sin 2\theta_T \ln \left(\frac{M_{\tilde{t}_2}^2}{M_{\tilde{t}_1}^2} \right), \\
f_b &= \frac{1}{M_{\tilde{b}_2}^2 - M_{\tilde{b}_1}^2} \left(M_{\tilde{b}_2}^2 \ln \left(\frac{M_{\tilde{b}_2}^2}{M_{\text{SUSY}}^2} \right) - M_{\tilde{b}_1}^2 \ln \left(\frac{M_{\tilde{b}_1}^2}{M_{\text{SUSY}}^2} \right) \right) - 1, \\
g_b &= \sin^2 2\theta_B \left(\frac{M_{\tilde{b}_2}^2 + M_{\tilde{b}_1}^2}{M_{\tilde{b}_2}^2 - M_{\tilde{b}_1}^2} \ln \left(\frac{M_{\tilde{b}_2}^2}{M_{\tilde{b}_1}^2} \right) - 2 \right), \\
e_b &= -m_b \sin 2\theta_B \ln \left(\frac{M_{\tilde{b}_2}^2}{M_{\tilde{b}_1}^2} \right).
\end{aligned} \tag{C.16}$$

Some of the radiative corrections due to top/bottom squark loops to CP-even, CP-odd and charged Higgs masses can be described by a shift of the trilinear soft SUSY breaking coupling A_λ (at the scale M_{SUSY}^2) everywhere in the tree level mass matrices:

$$A_\lambda \rightarrow A'_\lambda = A_\lambda + \frac{3h_t^2}{16\pi^2} A_t f_t + \frac{3h_b^2}{16\pi^2} A_b f_b. \tag{C.17}$$

The remaining corrections $\sim h_t^2 \equiv h_t^2(M_{\text{SUSY}}^2)$ and $\sim h_b^2 \equiv h_b^2(M_{\text{SUSY}}^2)$ to the CP-even mass matrix \mathcal{M}_S^2 (2.20) in the weak basis (H_{dR}, H_{uR}, S_R) are given by

$$\begin{aligned}
\Delta\mathcal{M}_{S,11}^2 &= \frac{3h_b^2}{32\pi^2} \left(-A_b^2 g_b + 4A_b e_b + 4m_b^2 \ln \left(\frac{M_{b_1}^2 M_{b_2}^2}{m_b^4} \right) \right) \\
&\quad - \frac{3h_t^2}{32\pi^2} (\lambda s(Q))^2 g_t , \\
\Delta\mathcal{M}_{S,22}^2 &= \frac{3h_t^2}{32\pi^2} \left(-A_t^2 g_t + 4A_t e_t + 4m_t^2 \ln \left(\frac{M_{t_1}^2 M_{t_2}^2}{m_t^4} \right) \right) \\
&\quad - \frac{3h_b^2}{32\pi^2} (\lambda s(Q))^2 g_b , \\
\Delta\mathcal{M}_{S,33}^2 &= -\frac{3h_t^2}{32\pi^2} \lambda^2 v_d^2(Q) g_t - \frac{3h_b^2}{32\pi^2} \lambda^2 v_u^2(Q) g_b , \\
\Delta\mathcal{M}_{S,12}^2 &= \lambda s(Q) \left(\frac{3h_t^2}{32\pi^2} (A_t g_t - 2 e_t) + \frac{3h_b^2}{32\pi^2} (A_b g_b - 2 e_b) \right) , \\
\Delta\mathcal{M}_{S,13}^2 &= \frac{3h_b^2}{32\pi^2} \lambda v_u(Q) (A_b g_b - 2 e_b) + \frac{3h_t^2}{32\pi^2} \lambda^2 s(Q) v_d(Q) (4 f_t - g_t) , \\
\Delta\mathcal{M}_{S,23}^2 &= \frac{3h_t^2}{32\pi^2} \lambda v_d(Q) (A_t g_t - 2 e_t) + \frac{3h_b^2}{32\pi^2} \lambda^2 s(Q) v_u(Q) (4 f_b - g_b) .
\end{aligned} \tag{C.18}$$

Additional contributions proportional to the electroweak gauge couplings g_1^2 and g_2^2 do not generate large logarithms and would be the same as in the MSSM (see [534] and refs. therein).

Dominant two loop corrections

As mentioned above, the two-loop corrections to ΔV_{eff} and hence to the Higgs mass matrices are relatively simple within the present approach and within the approximation where we neglect all terms without two powers of large logarithms; the ones given below can be obtained by integrating the relevant RGEs. The only complication arises, as above, from diagrams where Higgs bosons are connected to two top or bottom-quark lines which generate a (logarithmic) dependence on M_A . As infrared cutoff (in the case of diagrams with b quark lines) we use $m_t \approx M_Z$ (in the LLA, inside the logarithms), since the running Higgs masses at this scale are closer to the Higgs pole masses. Then one obtains

$$\begin{aligned}
\Delta\mathcal{M}_{S,11}^2 &= \frac{3h_b^4 v_d^2(Q)}{128\pi^4} \left(\ln^2 \left(\frac{M_{\text{SUSY}}^2}{m_t^2} \right) (16g_3^2 - \frac{2}{3}g_1^2 + 3\sin^2 \beta h_t^2 - 3\cos^2 \beta h_b^2) \right. \\
&\quad \left. + \left[\ln^2 \left(\frac{M_A^2}{m_t^2} \right) - \ln^2 \left(\frac{M_{\text{SUSY}}^2}{m_t^2} \right) \right] (3\sin^2 \beta h_b^2 + (3\sin^2 \beta + 1) h_t^2) \right) , \\
\Delta\mathcal{M}_{S,22}^2 &= \frac{3h_t^4 v_u^2(Q)}{128\pi^4} \left(\ln^2 \left(\frac{M_{\text{SUSY}}^2}{m_t^2} \right) (16g_3^2 + \frac{4}{3}g_1^2 - 3\sin^2 \beta h_t^2 + 3\cos^2 \beta h_b^2) \right. \\
&\quad \left. + \left[\ln^2 \left(\frac{M_A^2}{m_t^2} \right) - \ln^2 \left(\frac{M_{\text{SUSY}}^2}{m_t^2} \right) \right] (3\cos^2 \beta h_t^2 + (3\cos^2 \beta + 1) h_b^2) \right) . \tag{C.19}
\end{aligned}$$

(Contributions $\sim g_2^2$ cancel.)

Dominant slepton contributions

Contributions proportional to the electroweak gauge couplings g_1^2 and g_2^2 from squark loops do not give large logarithms, since we assume the squarks to have masses close to the ultraviolet cutoff M_{SUSY} . Sleptons could be considerably lighter, and here we assume a common slepton mass $M_{\tilde{l}}$. The corresponding contributions are best expressed in terms of the weak mixing angle θ_W and the auxiliary quantity

$$\Delta_{\tilde{l}} = -\frac{g_1^2 + g_2^2}{32\pi^2} M_Z^2 (9 \sin^4 \theta_W + 3 \cos^4 \theta_W) \ln \left(\frac{M_{\text{SUSY}}^2}{M_{\tilde{l}}^2} \right). \quad (\text{C.20})$$

Then we have

$$\begin{aligned} \Delta \mathcal{M}_{S,11}^2 &= \Delta_{\tilde{l}} \cos^2 \beta, \\ \Delta \mathcal{M}_{S,22}^2 &= \Delta_{\tilde{l}} \sin^2 \beta, \\ \Delta \mathcal{M}_{S,12}^2 &= -\Delta_{\tilde{l}} \sin \beta \cos \beta. \end{aligned} \quad (\text{C.21})$$

Chargino/neutralino contributions

Here we assume that the gaugino mass terms M_1 and M_2 are of similar order of magnitude, $M_1 \sim M_2 \equiv M_{1,2}$ (inside logarithms). We define the following potentially large logarithms:

$$\begin{aligned} L_\mu &= \ln(\mu_{\text{eff}}^2/M_{\text{SUSY}}^2), \\ L_\nu &= \ln(4\nu'^2/M_{\text{SUSY}}^2), \\ L_{M_2\mu} &= \ln(\max(M_{1,2}^2, \mu_{\text{eff}}^2)/M_{\text{SUSY}}^2), \\ L_{\mu\nu} &= \ln(\max(4\nu'^2, \mu_{\text{eff}}^2)/M_{\text{SUSY}}^2). \end{aligned} \quad (\text{C.22})$$

Again, some of the radiative corrections due to chargino/neutralino loops to all CP-even, CP-odd and charged Higgs masses can be described by an additional shift of the trilinear soft SUSY breaking coupling A_λ on top of the corrections in (C.17):

$$A'_\lambda \rightarrow A''_\lambda = A'_\lambda + \frac{1}{16\pi^2} (g_1^2 M_1 + 3g_2^2 M_2) L_{M_2\mu}. \quad (\text{C.23})$$

The remaining contributions are given by

$$\begin{aligned} \Delta \mathcal{M}_{S,11}^2 &= \frac{1}{16\pi^2} \left[M_Z^2 (g_1^2 + g_2^2) \cos^2 \beta (-10 + 16 \sin^2 \theta_W - 8 \sin^4 \theta_W) L_{M_2\mu} \right. \\ &\quad \left. - 4 \left(\lambda^2 \mu_{\text{eff}} \nu' \tan \beta + \frac{2M_Z^2 \lambda^4 \cos^2 \beta}{g_1^2 + g_2^2} \right) L_{\mu\nu} \right], \\ \Delta \mathcal{M}_{S,22}^2 &= \frac{1}{16\pi^2} \left[M_Z^2 (g_1^2 + g_2^2) \sin^2 \beta (-10 + 16 \sin^2 \theta_W - 8 \sin^4 \theta_W) L_{M_2\mu} \right. \\ &\quad \left. - 4 \left(\frac{\lambda^2 \mu_{\text{eff}} \nu'}{\tan \beta} + \frac{2M_Z^2 \lambda^4 \sin^2 \beta}{g_1^2 + g_2^2} \right) L_{\mu\nu} \right], \\ \Delta \mathcal{M}_{S,33}^2 &= \frac{1}{16\pi^2} \left[16\kappa^2 (-2\nu'^2 + \frac{1}{2}\mu'\nu' + \frac{1}{4}\mu'^2 + \frac{\mu'^3}{8\kappa_s(Q)}) L_\nu \right. \\ &\quad \left. - 8\lambda^2 \mu_{\text{eff}}^2 L_\mu + \frac{\lambda^3 \mu' M_Z^2}{\mu_{\text{eff}} (g_1^2 + g_2^2)} (8\kappa - 4\lambda \sin \beta \cos \beta) L_{\mu\nu} \right], \end{aligned}$$

$$\begin{aligned}
\Delta\mathcal{M}_{S,12}^2 &= \frac{1}{16\pi^2} \left[4 \left(\lambda^2 \mu_{\text{eff}} \nu' - \frac{2\lambda^4 M_Z^2 \sin \beta \cos \beta}{g_1^2 + g_2^2} \right) L_{\mu\nu} \right. \\
&\quad \left. - 2M_Z^2 (g_1^2 + g_2^2) \sin \beta \cos \beta L_{M_2\mu} \right], \\
\Delta\mathcal{M}_{S,13}^2 &= \frac{1}{16\pi^2} \frac{\sqrt{2}M_Z}{\sqrt{g_1^2 + g_2^2}} \left[\lambda (g_1^2 + g_2^2) \mu_{\text{eff}} \cos \beta (-6 + 4 \sin^2 \theta_W) L_{M_2\mu} \right. \\
&\quad \left. + 4\lambda^2 \left(\lambda \sin \beta (2\nu' - \frac{\mu'}{2}) - \cos \beta (\lambda \mu_{\text{eff}} + 4\kappa \nu') \right) L_{\mu\nu} \right], \\
\Delta\mathcal{M}_{S,23}^2 &= \frac{1}{16\pi^2} \frac{\sqrt{2}M_Z}{\sqrt{g_1^2 + g_2^2}} \left[\lambda (g_1^2 + g_2^2) \mu_{\text{eff}} \sin \beta (-6 + 4 \sin^2 \theta_W) L_{M_2\mu} \right. \\
&\quad \left. + 4\lambda^2 \left(\lambda \cos \beta (2\nu' - \frac{\mu'}{2}) - \sin \beta (\lambda \mu_{\text{eff}} + 4\kappa \nu') \right) L_{\mu\nu} \right]. \tag{C.24}
\end{aligned}$$

Higgs contributions

The contributions from Higgs loops are quite involved even in the LLA: the eigenvalues of the tree level mass matrices (2.22), (2.26) and (2.28) can be anywhere between M_Z^2 and M_{SUSY}^2 , and consequently Higgs loops can generate many different possibly large logarithms. Here we confine ourselves to the loop corrections to the diagonal element of the CP-even mass matrix *after* the rotation by the angle β , which corresponds at tree level to (2.23), the upper bound on the lightest eigenvalue of \mathcal{M}_S^2 . These Higgs induced loop corrections will be denoted as Δ_{Higgs} . Subsequently the rotation by the angle β can be reversed, and the contributions can be written in terms of corrections to the upper left 2×2 CP-even submatrix. (This is the procedure employed equally in [124]; the present expression for Δ_{Higgs} differs however from the one in [124], since the conventions for the tree level mass matrices are different.) Hence, first we define

$$\begin{aligned}
\Delta_{\text{Higgs}} &= \frac{M_Z^2}{8\pi^2 (g_1^2 + g_2^2)} \left[\left\{ \frac{(g_1^2 + g_2^2)^2}{4} (-4 + \sin^2 2\beta + 2 \sin^2 \theta_W (1 + \sin^2 2\beta) - 2 \sin^4 \theta_W) \right. \right. \\
&\quad \left. \left. + \frac{g_1^2 + g_2^2}{2} \lambda^2 (2 + \sin^2 2\beta (1 - 2 \sin^2 \theta_W)) - 4\lambda^4 - 2\lambda^2 \sin^2 2\beta (\lambda^2 + \kappa^2) \right\} \ln \left(\frac{M_{\text{SUSY}}^2}{M_Z^2} \right) \right. \\
&\quad \left. + \left\{ \frac{(g_1^2 + g_2^2)^2}{2} (\sin^4 \theta_W - \sin^2 \theta_W (1 + \sin^2 2\beta)) - \frac{11}{4} \sin^4 2\beta + 5 \sin^2 2\beta + \frac{3}{4} \right\} \right. \\
&\quad \left. + \frac{g_1^2 + g_2^2}{4} \lambda^2 (4 \sin^2 \theta_W \sin^2 2\beta + 11 \sin^4 2\beta - 15 \sin^2 2\beta - 2) \right. \\
&\quad \left. + \lambda^4 \left(-\frac{11}{4} \sin^4 2\beta + \frac{5}{2} \sin^2 2\beta + 1 \right) \right\} \ln \left(\mathcal{M}_{P,11}^2 / M_Z^2 \right) \\
&\quad + \left\{ \lambda^2 (\lambda - \kappa \sin 2\beta)^2 - \frac{\lambda^4}{\mathcal{M}_{S,33}^4} (2\mu_{\text{eff}} - \sin 2\beta (A_\lambda + 2\nu'))^4 \right. \\
&\quad \left. + \frac{3\lambda^2}{\mathcal{M}_{S,33}^2} \left(\frac{g_1^2 + g_2^2}{2} + \left(\lambda^2 - \frac{g_1^2 + g_2^2}{2} \right) \sin^2 2\beta \right) (2\mu_{\text{eff}} - \sin 2\beta (A_\lambda + 2\nu'))^2 \right\} \ln \left(\mathcal{M}_{S,33}^2 / M_Z^2 \right) \\
&\quad \left. + \left\{ \frac{(g_1^2 + g_2^2)^2}{16} (1 - \sin^4 2\beta) + \frac{(g_1^2 + g_2^2)^2}{4} \lambda^2 (\sin^4 2\beta + \sin^2 2\beta - 2) \right\} \right]
\end{aligned}$$

$$\begin{aligned}
& + \lambda^4 \left(-\frac{1}{4} \sin^4 2\beta - \frac{1}{2} \sin^2 2\beta + 1 \right) + \lambda^2 (\lambda + \kappa \sin 2\beta)^2 \left\} \ln \left((\mathcal{M}_{P,11}^2 + \mathcal{M}_{P,22}^2) / M_Z^2 \right) \right. \\
& - \left\{ \frac{\mathcal{M}_{P,22}^2}{2(\mathcal{M}_{P,11}^2 + \mathcal{M}_{P,22}^2)} \left(\frac{g_1^2 + g_2^2}{2} - \lambda^2 \right)^2 \sin^2 2\beta (1 - \sin^2 2\beta) \right. \\
& - \lambda \frac{\mathcal{M}_{P,11}^2 \mathcal{M}_{P,22}^2}{(\mathcal{M}_{P,11}^2 + \mathcal{M}_{P,22}^2)^2} (\lambda + \kappa \sin 2\beta) \left(\frac{g_1^2 + g_2^2}{2} (1 - \sin^2 2\beta) - \lambda^2 (2 - \sin^2 2\beta) \right) \\
& + \frac{1}{(\mathcal{M}_{P,11}^2 + \mathcal{M}_{P,22}^2)^2} \left(\frac{\mathcal{M}_{P,11}^2 + \mathcal{M}_{P,22}^2}{2} \left(\frac{g_1^2 + g_2^2}{2} (1 - \sin^2 2\beta) - \lambda^2 (2 - \sin^2 2\beta) \right) \right. \\
& \left. \left. - \lambda \mathcal{M}_{P,11}^2 (\lambda + \kappa \sin 2\beta) - \lambda^2 (A_\lambda - 2\nu')^2 \right)^2 \right\} \ln \left(\frac{(\mathcal{M}_{P,11}^2 + \mathcal{M}_{P,22}^2)^2}{\mathcal{M}_{P,11}^2 \mathcal{M}_{P,22}^2 - \mathcal{M}_{P,12}^4} \right) \left. \right]. \quad (\text{C.25})
\end{aligned}$$

Then we have

$$\begin{aligned}
\Delta \mathcal{M}_{S,11}^2 &= \Delta_{\text{Higgs}} \cos^2 \beta, \\
\Delta \mathcal{M}_{S,22}^2 &= \Delta_{\text{Higgs}} \sin^2 \beta, \\
\Delta \mathcal{M}_{S,12}^2 &= \Delta_{\text{Higgs}} \sin \beta \cos \beta. \quad (\text{C.26})
\end{aligned}$$

Gauge boson contributions

These are relatively simple in the Landau gauge; first we define

$$\Delta_{\text{Gauge}} = \frac{M_Z^2 (g_1^2 + g_2^2)}{32\pi^2} (-9 + 12 \sin^2 \theta_W - 6 \sin^4 \theta_W) \ln(M_{\text{SUSY}}^2 / M_Z^2); \quad (\text{C.27})$$

then we obtain

$$\begin{aligned}
\Delta \mathcal{M}_{S,11}^2 &= \Delta_{\text{Gauge}} \cos^2 \beta, \\
\Delta \mathcal{M}_{S,22}^2 &= \Delta_{\text{Gauge}} \sin^2 \beta, \\
\Delta \mathcal{M}_{S,12}^2 &= \Delta_{\text{Gauge}} \sin \beta \cos \beta. \quad (\text{C.28})
\end{aligned}$$

Rescaling

After adding up all previous corrections to $\Delta \mathcal{M}_S^2$ (and performing the shifts (C.17) and (C.23) of A_λ in the tree level expressions for \mathcal{M}_S^2) we obtain the Higgs mass matrix in terms of Higgs fields which are not yet properly normalised, since the kinetic terms in the effective action are multiplied by the wave function renormalisation constants. Hence, the Higgs mass matrix elements have to be rescaled by appropriate powers of wave function renormalisation constants Z_i , $i = H_u, H_d, S$:

$$\mathcal{M}'_{S,ij} = \mathcal{M}_{S,ij}^2 / \sqrt{Z_i Z_j}. \quad (\text{C.29})$$

Herewith also dominant double logarithms (two-loop terms corresponding to an RG improved one-loop calculation) are generated on top of the ones already present in \mathcal{M}_S^2 .

In the basis $H_i^{\text{weak}} = (H_{dR}, H_{uR}, S_R)$, $\mathcal{M}'_{S,ij}$ is diagonalized by an orthogonal 3×3 matrix S_{ij} in analogy to the tree level procedure (A.1). The eigenvalues of $\mathcal{M}'_{S,ij}$ (ordered in mass) will be denoted as $\mathcal{M}'_{S,i}$. This concludes the evaluation of the radiative corrections to the CP-even Higgs mass matrix; the pole masses will be computed in the subsection C.4.

CP-odd scalars

Here we confine ourselves to radiative corrections of the order $h_{t,b}^4$ and $g^2 h_{t,b}^2$ to the mass matrix, except for the chargino/neutralino contributions $\sim g^4$, which are included as well. These radiative corrections are quite simple: all effects of the order $h_{t,b}^4$ are included by the shift (C.17) of A_λ , all effects induced by chargino/neutralino loops by the additional shift (C.23) of A_λ , and all contributions of the order $g^2 h_{t,b}^2$ – as before – by the rescaling by the wave function renormalisation constants where it is sufficient to consider the terms $\sim h_{t,b}^2$ in Z_{H_u, H_d} (which we will not indicate explicitly).

The only complication originates from the fact that the rescaling by the wave function renormalisation constants is performed for the 3×3 mass matrix before the identification of the Goldstone mode, and has to be translated into the 2×2 mass matrix after dropping the Goldstone mode. The result for the *rescaled* CP-odd mass matrix elements $\mathcal{M}'_{P,ij}$ in the basis specified before (2.26) is

$$\begin{aligned}
\mathcal{M}'_{P,11} &= (\mu_{\text{eff}} B_{\text{eff}} + \widehat{m}_3^2) \left(\frac{v_u(Q)}{Z_{H_d} v_d(Q)} + \frac{v_d(Q)}{Z_{H_u} v_u(Q)} \right), \\
\mathcal{M}'_{P,22} &= \lambda(B_{\text{eff}} + 3\kappa s(Q)) \frac{v_u(Q)v_d(Q)}{s(Q)} - 3\kappa A_\kappa s(Q) - 2m_S'^2 - \kappa\mu' s(Q) \\
&\quad - \xi_F \left(4\kappa + \frac{\mu'}{s(Q)} \right) - \frac{\xi_S}{s(Q)}, \\
\mathcal{M}'_{P',12} &= \lambda(A_\lambda'' - 2\kappa s(Q) - \mu') \sqrt{v_u^2(Q)/Z_{H_d} + v_d^2(Q)/Z_{H_u}}, \tag{C.30}
\end{aligned}$$

where μ_{eff} , B_{eff} and \widehat{m}_3^2 are defined as in (2.7, 2.14), but in terms of parameters (and s) at the scale $Q = M_{\text{SUSY}}$ and with A_λ'' instead of A_λ .

The conventions for the diagonalization of the radiatively corrected CP-odd mass matrix are the same as those at tree level corresponding to (A.2 - A.5) above, leading to two eigenvalues $\mathcal{M}'_{P,i}^2$ (ordered in mass).

Charged scalar

The precision of the radiative corrections is the same as in the CP-odd case. However, both the corrections of the order $h_{t,b}^4$ and those induced by chargino/neutralino, gauge boson and slepton loops give rise to some additional terms on top of the shifts of A_λ . Altogether one obtains

$$\begin{aligned}
\mathcal{M}'_{\pm} &= \left(\mu_{\text{eff}} B_{\text{eff}} + \widehat{m}_3^2 + v_u(Q)v_d(Q) \left(\frac{g_2^2}{2} - \lambda^2 \right) \right) \left(\frac{v_u(Q)}{Z_{H_d} v_d(Q)} + \frac{v_d(Q)}{Z_{H_u} v_u(Q)} \right) \\
&\quad + \frac{v_u^2(Q) + v_d^2(Q)}{16\pi^2} \left(6h_t^2 h_b^2 \ln(M_{\text{SUSY}}^2/m_t^2) - \frac{3}{4}g_2^4 \ln(M_{\text{SUSY}}^2/M_t^2) \right. \\
&\quad \left. + \frac{7g_1^2 g_2^2 - g_2^4}{4} \ln(M_{\text{SUSY}}^2/M_Z^2) + 2(g_1^2 g_2^2 - g_2^4) L_{M_2\mu} \right), \tag{C.31}
\end{aligned}$$

where $L_{M_2\mu}$ was defined in (C.22).

C.4 Pole masses

In order to obtain the Higgs pole masses with the precision defined in Section 3.2 (exact to the order h_t^4 , h_b^4 , $g^2 h_t^2$, etc., but only in the LLA in the case of four powers of electroweak gauge couplings and Yukawa couplings λ , κ) it suffices to consider top- and bottom-quark corrections to the pole masses, which can be deduced from [534]. There, the corrections $\Delta\Pi(p^2, Q^2)$ to the inverse tree level propagators $\Pi(p^2) = p^2 - M_{\text{tree}}^2$ were computed. In order to make contact with our approach, we first have to write

$$\Delta\Pi(p^2, Q^2) = \Delta\widehat{\Pi}(p^2, Q^2) + \Delta\Pi(p^2 = 0, Q^2). \quad (\text{C.32})$$

The quantity $\Delta\Pi(p^2 = 0, Q^2)$ corresponds to (the negative of) our radiative corrections to the masses squared before the rescaling by the wave function renormalisation constants. The quantity $\Delta\widehat{\Pi}(p^2, Q^2)$ proportional to p^2 still contains potentially large logarithms. We have already included these large logarithms in our wave function renormalisation constants in (C.9) – (C.11), but there remains a missing part $\delta\Pi(p^2)$ which is defined as

$$\begin{aligned} \Pi(p^2) + \Delta\Pi(p^2, Q^2) &\equiv p^2 + \Delta\widehat{\Pi}(p^2, Q^2) - M_{\text{tree}}^2 + \Delta\Pi(p^2 = 0, Q^2) \\ &= Z(Q^2) p^2 - M^2 + \delta\Pi(p^2) \end{aligned} \quad (\text{C.33})$$

where $M^2 \equiv M_{\text{tree}}^2 - \Delta\Pi(p^2 = 0, Q^2)$ are the radiatively corrected masses squared before the rescaling by the wave function renormalisation constants. Hence we obtain

$$\delta\Pi(p^2) = p^2 + \Delta\widehat{\Pi}(p^2, Q^2) - Z(Q^2) p^2 \equiv \Delta\widehat{\Pi}(p^2, m_t^2) \quad (\text{C.34})$$

where we have used the fact that the relevant terms $\sim h_{t,b}^2$ in our wave function renormalisation constants $Z(Q^2)$ in (C.9) – (C.11) were computed with an infrared cutoff m_t^2 . Hence, no large logarithms appear in the remaining pole corrections $\delta\Pi(p^2)$, which justifies the present method. Finally, one obtains for the pole mass from the last expression in (C.33) within the present approximation

$$M_{\text{pole}}^2 = M^2/Z(Q^2) - \delta\Pi(M^2/Z(Q^2)) \quad (\text{C.35})$$

where the first term corresponds to the results in Section C.3 (where $M^2/Z(Q^2) \equiv M'^2$), and $\delta\Pi$ can be deduced from [534] according to the formulae above. In the case of mass matrices, M_{pole}^2 is diagonalized by the same rotation which diagonalizes M'^2 within the present approximation.

In the CP-even case, one obtains in terms of the eigenvalues $\mathcal{M}'^2_{S,i}$ and of the elements of the diagonalization matrix S_{ij}

$$\begin{aligned} \mathcal{M}_{S,i}^{\text{pole } 2} &= \mathcal{M}'^2_{S,i} - \frac{3h_t^2}{16\pi^2} S_{i2}^2 \left(\mathcal{M}'^2_{S,i} - 4m_t^2 \right) B(\mathcal{M}'^2_{S,i}, m_t^2) \\ &\quad - \frac{3h_b^2}{16\pi^2} S_{i1}^2 \left(\mathcal{M}'^2_{S,i} \ln(m_t^2/m_b^2) + \left(\mathcal{M}'^2_{S,i} - 4m_b^2 \right) B(\mathcal{M}'^2_{S,i}, m_b^2) \right) \end{aligned} \quad (\text{C.36})$$

where the logarithm $\ln(m_t^2/m_b^2)$ in the second line originates from the infrared cutoff m_t^2 in Z_{H_d} in (C.10), and leads to a finite pole mass correction for $m_b \rightarrow 0$. $B(M^2, m^2)$ is defined

by

$$\begin{aligned}
B(M^2, m^2) &= 2 - \sqrt{1 - \frac{4m^2}{M^2}} \ln \left(\frac{1 + \sqrt{1 - \frac{4m^2}{M^2}}}{1 - \sqrt{1 - \frac{4m^2}{M^2}}} \right) \text{ if } M^2 > 4m^2 ; \\
B(M^2, m^2) &= 2 - 2\sqrt{\frac{4m^2}{M^2} - 1} \arctan \left(\sqrt{\frac{M^2}{4m^2 - M^2}} \right) \text{ if } M^2 < 4m^2 . \quad (\text{C.37})
\end{aligned}$$

A similar expression holds for the CP-odd Higgs pole masses in terms of the eigenvalues $\mathcal{M}'_{P,i}$ and the elements of the diagonalization matrix P_{ij} defined in (A.3):

$$\begin{aligned}
\mathcal{M}_{P,i}^{\text{pole } 2} &= \mathcal{M}'_{P,i} - \frac{3h_t^2}{16\pi^2} P_{i2}^2 \mathcal{M}'_{P,i} B(\mathcal{M}'_{P,i}, m_t^2) \\
&\quad - \frac{3h_b^2}{16\pi^2} P_{i1}^2 \left(\mathcal{M}'_{P,i} \ln(m_t^2/m_b^2) + \mathcal{M}'_{P,i} B(\mathcal{M}'_{P,i}, m_b^2) \right) . \quad (\text{C.38})
\end{aligned}$$

For the charged Higgs pole mass one obtains (neglecting m_b^2 inside logarithms)

$$\begin{aligned}
\mathcal{M}_{\pm}^{\text{pole } 2} &= \mathcal{M}'_{\pm} + \frac{3}{16\pi^2} \left\{ (h_t^2 \cos^2 \beta + h_b^2 \sin^2 \beta) \left(\right. \right. \quad (\text{C.39}) \\
&\quad \mathcal{M}'_{\pm} \left[\left(1 - \frac{m_t^2}{\mathcal{M}'_{\pm}} \right) \ln \left| \frac{\mathcal{M}'_{\pm} - m_t^2}{m_t^2} \right| - 2 \right] \\
&\quad \left. \left. + (m_t^2 + m_b^2) \left[\left(1 - \frac{m_t^2}{\mathcal{M}'_{\pm}} \right) \ln \left| \frac{m_t^2}{\mathcal{M}'_{\pm} - m_t^2} \right| + 1 \right] \right) \right. \\
&\quad \left. + 4h_t m_t h_b m_b \sin \beta \cos \beta \left[\left(1 - \frac{m_t^2}{\mathcal{M}'_{\pm}} \right) \ln \left| \frac{m_t^2}{\mathcal{M}'_{\pm} - m_t^2} \right| + 1 \right] \right\} .
\end{aligned}$$

Appendix D. Public computer tools

NMSSMTools [52, 88, 345] allows to compute the Higgs and sparticle spectrum, mixing angles and Higgs branching fractions. NMSSMTools includes NMHDECAY (where the parameters can be specified at the weak/SUSY scale), NMSPEC (where the parameters can be specified at the GUT scale) and a version where the soft terms satisfy GMSB-like boundary conditions. A connection to MicrOMEGAS is provided.

Link: <http://www.th.u-psud.fr/NMHDECAY/nmssmtools.html>

CalcHEP allows to compute cross sections in the NMSSM (used, e. g., by MicrOMEGAS).

Link: <http://theory.sinp.msu.ru/~pukhov/calchep.html>

MicrOMEGAS [502, 503] allows to compute all relevant cross sections for LSP annihilation and coannihilation, integrates the density evolution equations, determines the LSP relic density and detection cross sections. It can be called from NMSSMTools.

Link: <http://wwwlapp.in2p3.fr/laph/micromegas>

Spheno [535] allows to compute the sparticle two-body decay amplitudes in the NMSSM; a computation of the Higgs and sparticle spectrum is in progress.

Link: <http://ific.uv.es/~porod/SPheno.html>

WHIZARD is a multi-purpose Monte Carlo event generator, into which the NMSSM has been implemented [536].

Link: <http://projects.hepforge.org/whizard>

SuperIso [537] allows to compute flavour physics observables in the NMSSM such as $B \rightarrow X_s \gamma$, the isospin asymmetry of $B \rightarrow K^* \gamma$, and the branching ratios $B_s \rightarrow \mu^+ \mu^-$, $B \rightarrow \tau \nu_\tau$, $B \rightarrow D \tau \nu_\tau$, $K \rightarrow \mu \nu_\mu$, $D_s \rightarrow \tau \nu_\tau$ and $D_s \rightarrow \mu \nu_\mu$.

Link: <http://superiso.in2p3.fr>

References

- [1] E. Witten, Nucl. Phys. B **188** (1981) 513.
- [2] S. Dimopoulos and H. Georgi, Nucl. Phys. B **193** (1981) 150.
- [3] E. Witten, Phys. Lett. B **105** (1981) 267.
- [4] R. K. Kaul and P. Majumdar, Nucl. Phys. B **199** (1982) 36.
- [5] N. Sakai, Z. Phys. C **11** (1981) 153.
- [6] J. R. Ellis, S. Kelley and D. V. Nanopoulos, Phys. Lett. B **260** (1991) 131.
- [7] C. Giunti, C. W. Kim and U. W. Lee, Mod. Phys. Lett. A **6** (1991) 1745.
- [8] U. Amaldi, W. de Boer and H. Furstenau, Phys. Lett. B **260** (1991) 447.
- [9] P. Langacker and M. x. Luo, Phys. Rev. D **44** (1991) 817.
- [10] H. Pagels and J. R. Primack, Phys. Rev. Lett. **48** (1982) 223.
- [11] H. Goldberg, Phys. Rev. Lett. **50** (1983) 1419.
- [12] P. Fayet, Nucl. Phys. B **90** (1975) 104.
- [13] P. Fayet and S. Ferrara, Phys. Rept. **32** (1977) 249.
- [14] H. P. Nilles, Phys. Rept. **110** (1984) 1.
- [15] S. P. Martin, arXiv:hep-ph/9709356.
- [16] D. J. H. Chung, L. L. Everett, G. L. Kane, S. F. King, J. D. Lykken and L. T. Wang, Phys. Rept. **407** (2005) 1 [arXiv:hep-ph/0312378].
- [17] J. E. Kim and H. P. Nilles, Phys. Lett. B **138** (1984) 150.
- [18] P. Fayet, Phys. Lett. B **69** (1977) 489.
- [19] L. E. Ibanez and G. G. Ross, Phys. Lett. B **110** (1982) 215.
- [20] D. V. Nanopoulos and K. Tamvakis, Phys. Lett. B **113** (1982) 151.
- [21] J. R. Ellis, L. E. Ibanez and G. G. Ross, Phys. Lett. B **113** (1982) 283.
- [22] J. R. Ellis, L. E. Ibanez and G. G. Ross, Nucl. Phys. B **221** (1983) 29.
- [23] G. F. Giudice and A. Masiero, Phys. Lett. B **206** (1988) 480.
- [24] R. Barbieri, S. Ferrara and C. A. Savoy, Phys. Lett. B **119** (1982) 343.
- [25] H. P. Nilles, M. Srednicki and D. Wyler, Phys. Lett. B **120** (1983) 346.
- [26] J. M. Frere, D. R. T. Jones and S. Raby, Nucl. Phys. B **222** (1983) 11.
- [27] A. H. Chamseddine, R. L. Arnowitt and P. Nath, Phys. Rev. Lett. **49** (1982) 970.

- [28] P. Nath, R. L. Arnowitt and A. H. Chamseddine, Phys. Lett. B **121** (1983) 33.
- [29] S. Ferrara, D. V. Nanopoulos and C. A. Savoy, Phys. Lett. B **123** (1983) 214.
- [30] P. Nath, R. L. Arnowitt and A. H. Chamseddine, Nucl. Phys. B **227** (1983) 121.
- [31] R. L. Arnowitt, A. H. Chamseddine and P. Nath, Phys. Rev. Lett. **50** (1983) 232.
- [32] A. I. Veselov, M. I. Vysotsky and K. A. Ter-Martirosian, Sov. Phys. JETP **63** (1986) 489 [Zh. Eksp. Teor. Fiz. **90** (1986) 838].
- [33] N. Dragon, U. Ellwanger and M. G. Schmidt, Phys. Lett. B **154** (1985) 373.
- [34] U. Ellwanger, N. Dragon and M. G. Schmidt, Z. Phys. C **29** (1985) 209.
- [35] E. Cohen, J. R. Ellis, K. Enqvist and D. V. Nanopoulos, Phys. Lett. B **165** (1985) 76.
- [36] J. P. Derendinger, L. E. Ibanez and H. P. Nilles, Nucl. Phys. B **267** (1986) 365.
- [37] P. Binetruy, S. Dawson, I. Hinchliffe and M. Sher, Nucl. Phys. B **273** (1986) 501.
- [38] J. R. Ellis, K. Enqvist, D. V. Nanopoulos and F. Zwirner, Nucl. Phys. B **276** (1986) 14.
- [39] L. E. Ibanez and J. Mas, Nucl. Phys. B **286** (1987) 107.
- [40] G. Costa, F. Feruglio, F. Gabbiani and F. Zwirner, Nucl. Phys. B **286** (1987) 325.
- [41] I. Antoniadis, J. R. Ellis, J. S. Hagelin and D. V. Nanopoulos, Phys. Lett. B **194** (1987) 231.
- [42] B. A. Campbell, J. R. Ellis, J. S. Hagelin, D. V. Nanopoulos and R. Ticciati, Phys. Lett. B **198** (1987) 200.
- [43] I. Antoniadis, J. R. Ellis, J. S. Hagelin and D. V. Nanopoulos, Phys. Lett. B **205** (1988) 459.
- [44] J. R. Ellis, J. S. Hagelin, S. Kelley and D. V. Nanopoulos, Nucl. Phys. B **311** (1988) 1.
- [45] J. R. Ellis, J. F. Gunion, H. E. Haber, L. Roszkowski and F. Zwirner, Phys. Rev. D **39** (1989) 844.
- [46] M. Drees, Int. J. Mod. Phys. A **4** (1989) 3635.
- [47] M. Maniatis, arXiv:0906.0777 [hep-ph].
- [48] LEPSUSYWG, ALEPH, DELPHI, L3 and OPAL experiments, note LEPSUSYWG/01-03.1 (<http://lepsusy.web.cern.ch/lepsusy/Welcome.html>).
- [49] B. Allanach *et al.*, Comput. Phys. Commun. **180** (2009) 8 [arXiv:0801.0045 [hep-ph]].
- [50] F. Franke and H. Fraas, Int. J. Mod. Phys. A **12** (1997) 479 [arXiv:hep-ph/9512366].
- [51] D. J. Miller, R. Nevzorov and P. M. Zerwas, Nucl. Phys. B **681** (2004) 3 [arXiv:hep-ph/0304049].

- [52] U. Ellwanger, J. F. Gunion and C. Hugonie, JHEP **0502** (2005) 066 [arXiv:hep-ph/0406215].
- [53] J. P. Derendinger and C. A. Savoy, Nucl. Phys. B **237** (1984) 307.
- [54] U. Ellwanger, M. Rausch de Traubenberg and C. A. Savoy, Nucl. Phys. B **492** (1997) 21 [arXiv:hep-ph/9611251].
- [55] M. Dine and A. E. Nelson, Phys. Rev. D **48** (1993) 1277 [arXiv:hep-ph/9303230].
- [56] M. Maniatis, A. von Manteuffel and O. Nachtmann, Eur. Phys. J. C **49** (2007) 1067 [arXiv:hep-ph/0608314].
- [57] S. Schael *et al.* [ALEPH and DELPHI and L3 and OPAL Collaborations and LEP Working Group for Higgs Boson Searches], Eur. Phys. J. C **47** (2006) 547 [arXiv:hep-ex/0602042].
- [58] U. Ellwanger and C. Hugonie, Eur. Phys. J. C **5** (1998) 723 [arXiv:hep-ph/9712300].
- [59] R. D. Peccei and H. R. Quinn, Phys. Rev. D **16** (1977) 1791.
- [60] R. D. Peccei and H. R. Quinn, Phys. Rev. Lett. **38** (1977) 1440.
- [61] E. J. Chun, Phys. Lett. B **348** (1995) 111 [arXiv:hep-ph/9411290].
- [62] P. Ciafaloni and A. Pomarol, Phys. Lett. B **404** (1997) 83 [arXiv:hep-ph/9702410].
- [63] L. J. Hall and T. Watari, Phys. Rev. D **70** (2004) 115001 [arXiv:hep-ph/0405109].
- [64] B. Feldstein, L. J. Hall and T. Watari, Phys. Lett. B **607** (2005) 155 [arXiv:hep-ph/0411013].
- [65] D. J. Miller and R. Nevzorov, arXiv:hep-ph/0309143.
- [66] R. Barbieri, L. J. Hall, A. Y. Papaioannou, D. Pappadopulo and V. S. Rychkov, JHEP **0803** (2008) 005 [arXiv:0712.2903 [hep-ph]].
- [67] O. Lebedev and S. Ramos-Sanchez, Phys. Lett. B **684** (2010) 48 [arXiv:0912.0477].
- [68] B. A. Dobrescu, G. L. Landsberg and K. T. Matchev, Phys. Rev. D **63** (2001) 075003 [arXiv:hep-ph/0005308].
- [69] B. A. Dobrescu and K. T. Matchev, JHEP **0009** (2000) 031 [arXiv:hep-ph/0008192].
- [70] R. Dermisek and J. F. Gunion, Phys. Rev. D **75** (2007) 075019 [arXiv:hep-ph/0611142].
- [71] D. E. Morrissey and A. Pierce, Phys. Rev. D **78** (2008) 075029 [arXiv:0807.2259 [hep-ph]].
- [72] S. F. King and P. L. White, Phys. Rev. D **52** (1995) 4183 [arXiv:hep-ph/9505326].
- [73] M. Masip, R. Munoz-Tapia and A. Pomarol, Phys. Rev. D **57** (1998) 5340 [arXiv:hep-ph/9801437].

- [74] Y. Mambrini, G. Moultaka and M. Rausch de Traubenberg, Nucl. Phys. B **609** (2001) 83 [arXiv:hep-ph/0101237].
- [75] R. B. Nevzorov and M. A. Trusov, Phys. Atom. Nucl. **64** (2001) 1299 [Yad. Fiz. **64** (2001) 1375] [arXiv:hep-ph/0110363].
- [76] R. B. Nevzorov and M. A. Trusov, Phys. Atom. Nucl. **64** (2001) 1513 [Yad. Fiz. **64** (2001) 1589] [arXiv:hep-ph/0112301].
- [77] Y. Mambrini and G. Moultaka, Phys. Rev. D **65** (2002) 115011 [arXiv:hep-ph/0112353].
- [78] D. A. Demir, JHEP **0511** (2005) 003 [arXiv:hep-ph/0408043].
- [79] B. C. Allanach and S. F. King, Phys. Lett. B **328** (1994) 360 [arXiv:hep-ph/9403212].
- [80] L. Durand and J. L. Lopez, Phys. Lett. B **217** (1989) 463.
- [81] L. Durand and J. L. Lopez, Phys. Rev. D **40** (1989) 207.
- [82] P. Binetruy and C. A. Savoy, Phys. Lett. B **277** (1992) 453.
- [83] J. R. Espinosa and M. Quiros, Phys. Lett. B **279** (1992) 92.
- [84] T. Moroi and Y. Okada, Phys. Lett. B **295** (1992) 73.
- [85] T. Elliott, S. F. King and P. L. White, Phys. Lett. B **314** (1993) 56 [arXiv:hep-ph/9305282].
- [86] T. Elliott, S. F. King and P. L. White, Phys. Rev. D **49** (1994) 2435 [arXiv:hep-ph/9308309].
- [87] G. K. Yeghian, arXiv:hep-ph/9904488.
- [88] U. Ellwanger and C. Hugonie, Comput. Phys. Commun. **175** (2006) 290 [arXiv:hep-ph/0508022].
- [89] C. Amsler *et al.* [Particle Data Group], Phys. Lett. B **667** (2008) 1.
- [90] K. Inoue, A. Kakuto, H. Komatsu and S. Takeshita, Prog. Theor. Phys. **68** (1982) 927 [Erratum-ibid. **70** (1983) 330].
- [91] K. Inoue, A. Kakuto, H. Komatsu and S. Takeshita, Prog. Theor. Phys. **71** (1984) 413.
- [92] B. Gato, J. Leon, J. Perez-Mercader and M. Quiros, Nucl. Phys. B **253** (1985) 285.
- [93] N. K. Falck, Z. Phys. C **30** (1986) 247.
- [94] S. P. Martin and M. T. Vaughn, Phys. Rev. D **50** (1994) 2282 [Erratum-ibid. D **78** (2008) 039903] [arXiv:hep-ph/9311340].
- [95] Y. Yamada, Phys. Rev. D **50** (1994) 3537 [arXiv:hep-ph/9401241].
- [96] U. Ellwanger, C. C. Jean-Louis and A. M. Teixeira, JHEP **0805** (2008) 044 [arXiv:0803.2962 [hep-ph]].

- [97] Y. Okada, M. Yamaguchi and T. Yanagida, *Prog. Theor. Phys.* **85** (1991) 1.
- [98] J. R. Ellis, G. Ridolfi and F. Zwirner, *Phys. Lett. B* **257** (1991) 83.
- [99] Y. Okada, M. Yamaguchi and T. Yanagida, *Phys. Lett. B* **262** (1991) 54.
- [100] H. E. Haber and R. Hempfling, *Phys. Rev. Lett.* **66** (1991) 1815.
- [101] J. R. Ellis, G. Ridolfi and F. Zwirner, *Phys. Lett. B* **262** (1991) 477.
- [102] A. Yamada, *Phys. Lett. B* **263** (1991) 233.
- [103] J. R. Espinosa and M. Quiros, *Phys. Lett. B* **266** (1991) 389.
- [104] U. Ellwanger and C. Hugonie, *Mod. Phys. Lett. A* **22** (2007) 1581 [arXiv:hep-ph/0612133].
- [105] U. Ellwanger and M. Rausch de Traubenberg, *Z. Phys. C* **53** (1992) 521.
- [106] U. Ellwanger and M. Lindner, *Phys. Lett. B* **301** (1993) 365 [arXiv:hep-ph/9211249].
- [107] T. Elliott, S. F. King and P. L. White, *Phys. Lett. B* **305** (1993) 71 [arXiv:hep-ph/9302202].
- [108] M. A. Diaz, T. A. ter Veldhuis and T. J. Weiler, *Phys. Rev. Lett.* **74** (1995) 2876 [arXiv:hep-ph/9408319].
- [109] G. M. Asatrian and G. K. Egikian, *Mod. Phys. Lett. A* **10** (1995) 2943 [arXiv:hep-ph/9508263].
- [110] U. Ellwanger, *Phys. Lett. B* **303** (1993) 271 [arXiv:hep-ph/9302224].
- [111] P. N. Pandita, *Z. Phys. C* **59** (1993) 575.
- [112] P. N. Pandita, *Phys. Lett. B* **318** (1993) 338.
- [113] U. Ellwanger, M. Rausch de Traubenberg and C. A. Savoy, *Z. Phys. C* **67** (1995) 665 [arXiv:hep-ph/9502206].
- [114] F. Franke and H. Fraas, *Phys. Lett. B* **353** (1995) 234 [arXiv:hep-ph/9504279].
- [115] S. W. Ham, S. K. Oh and B. R. Kim, *J. Phys. G* **22** (1996) 1575 [arXiv:hep-ph/9604243].
- [116] Y. Daikoku and D. Suematsu, *Prog. Theor. Phys.* **104** (2000) 827 [arXiv:hep-ph/0003206].
- [117] R. B. Nevzorov and M. A. Trusov, *J. Exp. Theor. Phys.* **91** (2000) 1079 [*Zh. Eksp. Teor. Fiz.* **91** (2000) 1251] [arXiv:hep-ph/0106351].
- [118] B. Ananthanarayan and P. N. Pandita, *Phys. Lett. B* **353** (1995) 70 [arXiv:hep-ph/9503323].
- [119] B. Ananthanarayan and P. N. Pandita, *Phys. Lett. B* **371** (1996) 245 [arXiv:hep-ph/9511415].

- [120] B. Ananthanarayan and P. N. Pandita, *Int. J. Mod. Phys. A* **12** (1997) 2321 [arXiv:hep-ph/9601372].
- [121] G. L. Kane, C. F. Kolda and J. D. Wells, *Phys. Rev. Lett.* **70** (1993) 2686 [arXiv:hep-ph/9210242].
- [122] J. R. Espinosa and M. Quiros, *Phys. Lett. B* **302** (1993) 51 [arXiv:hep-ph/9212305].
- [123] J. R. Espinosa and M. Quiros, *Phys. Rev. Lett.* **81** (1998) 516 [arXiv:hep-ph/9804235].
- [124] U. Ellwanger and C. Hugonie, *Phys. Lett. B* **623** (2005) 93 [arXiv:hep-ph/0504269].
- [125] G. Degrossi and P. Slavich, *Nucl. Phys. B* **825** (2010) 119 [arXiv:0907.4682 [hep-ph]].
- [126] M. A. Diaz, T. A. ter Veldhuis and T. J. Weiler, *Phys. Rev. D* **54** (1996) 5855 [arXiv:hep-ph/9512229].
- [127] U. Ellwanger and C. Hugonie, *Eur. Phys. J. C* **25** (2002) 297 [arXiv:hep-ph/9909260].
- [128] R. Barbieri, L. J. Hall, Y. Nomura and V. S. Rychkov, *Phys. Rev. D* **75** (2007) 035007 [arXiv:hep-ph/0607332].
- [129] L. Cavicchia, R. Franceschini and V. S. Rychkov, *Phys. Rev. D* **77** (2008) 055006 [arXiv:0710.5750 [hep-ph]].
- [130] R. Harnik, G. D. Kribs, D. T. Larson and H. Murayama, *Phys. Rev. D* **70** (2004) 015002 [arXiv:hep-ph/0311349].
- [131] S. Chang, C. Kilic and R. Mahbubani, *Phys. Rev. D* **71** (2005) 015003 [arXiv:hep-ph/0405267].
- [132] A. Delgado and T. M. P. Tait, *JHEP* **0507** (2005) 023 [arXiv:hep-ph/0504224].
- [133] A. Birkedal, Z. Chacko and Y. Nomura, *Phys. Rev. D* **71** (2005) 015006 [arXiv:hep-ph/0408329].
- [134] N. G. Deshpande, X. G. He and J. Jiang, *Phys. Lett. B* **656** (2007) 91 [arXiv:0707.2959 [hep-ph]].
- [135] P. Batra, A. Delgado, D. E. Kaplan and T. M. P. Tait, *JHEP* **0406** (2004) 032 [arXiv:hep-ph/0404251].
- [136] J. Cao and J. M. Yang, *Phys. Rev. D* **78** (2008) 115001 [arXiv:0810.0989 [hep-ph]].
- [137] W. Wang, Z. Xiong and J. M. Yang, *Phys. Lett. B* **680** (2009) 167 [arXiv:0901.3818 [hep-ph]].
- [138] L. J. Hall, R. Rattazzi and U. Sarid, *Phys. Rev. D* **50** (1994) 7048 [arXiv:hep-ph/9306309].
- [139] R. Hempfling, *Phys. Rev. D* **49** (1994) 6168.
- [140] M. S. Carena, M. Olechowski, S. Pokorski and C. E. M. Wagner, *Nucl. Phys. B* **426** (1994) 269 [arXiv:hep-ph/9402253].

- [141] M. S. Carena, S. Mrenna and C. E. M. Wagner, Phys. Rev. D **62** (2000) 055008 [arXiv:hep-ph/9907422].
- [142] R. N. Hodgkinson and A. Pilaftsis, Phys. Rev. D **76** (2007) 015007 [arXiv:hep-ph/0612188].
- [143] LEP and SLD Collaborations, Phys. Rept. **427** (2006) 257 [arXiv:hep-ex/0509008].
- [144] J. Cao and J. M. Yang, JHEP **0812** (2008) 006 [arXiv:0810.0751 [hep-ph]].
- [145] J. Polchinski and L. Susskind, Phys. Rev. D **26** (1982) 3661.
- [146] H. P. Nilles, M. Srednicki and D. Wyler, Phys. Lett. B **124** (1983) 337.
- [147] A. B. Lahanas, Phys. Lett. B **124** (1983) 341.
- [148] L. J. Hall, J. D. Lykken and S. Weinberg, Phys. Rev. D **27** (1983) 2359.
- [149] L. Alvarez-Gaume, J. Polchinski and M. B. Wise, Nucl. Phys. B **221** (1983) 495.
- [150] U. Ellwanger, Phys. Lett. B **133** (1983) 187.
- [151] J. Bagger and E. Poppitz, Phys. Rev. Lett. **71** (1993) 2380 [arXiv:hep-ph/9307317].
- [152] V. Jain, Phys. Lett. B **351** (1995) 481 [arXiv:hep-ph/9407382].
- [153] J. Bagger, E. Poppitz and L. Randall, Nucl. Phys. B **455** (1995) 59 [arXiv:hep-ph/9505244].
- [154] D. Nemeschansky, Nucl. Phys. B **234** (1984) 379.
- [155] U. Ellwanger, N. Dragon and M. G. Schmidt, Nucl. Phys. B **255** (1985) 549.
- [156] U. Ellwanger, Phys. Lett. B **349** (1995) 57 [arXiv:hep-ph/9501227].
- [157] A. Vilenkin, Phys. Rept. **121** (1985) 263.
- [158] J. R. Ellis, K. Enqvist, D. V. Nanopoulos, K. A. Olive, M. Quiros and F. Zwirner, Phys. Lett. B **176** (1986) 403.
- [159] G. B. Gelmini, M. Gleiser and E. W. Kolb, Phys. Rev. D **39** (1989) 1558.
- [160] B. Rai and G. Senjanovic, Phys. Rev. D **49** (1994) 2729 [arXiv:hep-ph/9301240].
- [161] S. A. Abel and P. L. White, Phys. Rev. D **52** (1995) 4371 [arXiv:hep-ph/9505241].
- [162] S. A. Abel, S. Sarkar and P. L. White, Nucl. Phys. B **454** (1995) 663 [arXiv:hep-ph/9506359].
- [163] S. A. Abel, Nucl. Phys. B **480** (1996) 55 [arXiv:hep-ph/9609323].
- [164] C. F. Kolda, S. Pokorski and N. Polonsky, Phys. Rev. Lett. **80** (1998) 5263 [arXiv:hep-ph/9803310].
- [165] C. Panagiotakopoulos and K. Tamvakis, Phys. Lett. B **446** (1999) 224 [arXiv:hep-ph/9809475].

- [166] C. Panagiotakopoulos and K. Tamvakis, Phys. Lett. B **469** (1999) 145 [arXiv:hep-ph/9908351].
- [167] C. Panagiotakopoulos and A. Pilaftsis, Phys. Rev. D **63** (2001) 055003 [arXiv:hep-ph/0008268].
- [168] V. A. Kuzmin, V. A. Rubakov and M. E. Shaposhnikov, Phys. Lett. B **155** (1985) 36.
- [169] M. S. Carena, M. Quiros, M. Seco and C. E. M. Wagner, Nucl. Phys. B **650** (2003) 24 [arXiv:hep-ph/0208043].
- [170] M. Pietroni, Nucl. Phys. B **402** (1993) 27 [arXiv:hep-ph/9207227].
- [171] A. T. Davies, C. D. Froggatt and R. G. Moorhouse, Phys. Lett. B **372** (1996) 88 [arXiv:hep-ph/9603388].
- [172] S. J. Huber and M. G. Schmidt, Eur. Phys. J. C **10** (1999) 473 [arXiv:hep-ph/9809506].
- [173] S. J. Huber, P. John, M. Laine and M. G. Schmidt, Phys. Lett. B **475** (2000) 104 [arXiv:hep-ph/9912278].
- [174] S. J. Huber and M. G. Schmidt, Nucl. Phys. B **606** (2001) 183 [arXiv:hep-ph/0003122].
- [175] M. Bastero-Gil, C. Hugonie, S. F. King, D. P. Roy and S. Vempati, Phys. Lett. B **489** (2000) 359 [arXiv:hep-ph/0006198].
- [176] A. Menon, D. E. Morrissey and C. E. M. Wagner, Phys. Rev. D **70** (2004) 035005 [arXiv:hep-ph/0404184].
- [177] S. W. Ham, S. K. OH, C. M. Kim, E. J. Yoo and D. Son, Phys. Rev. D **70** (2004) 075001 [arXiv:hep-ph/0406062].
- [178] K. Funakubo, S. Tao and F. Toyoda, Prog. Theor. Phys. **114** (2005) 369 [arXiv:hep-ph/0501052].
- [179] S. J. Huber, T. Konstandin, T. Prokopec and M. G. Schmidt, Nucl. Phys. B **757** (2006) 172 [arXiv:hep-ph/0606298].
- [180] D. Comelli, M. Pietroni and A. Riotto, Phys. Rev. D **50** (1994) 7703 [arXiv:hep-ph/9406368].
- [181] D. Comelli, M. Pietroni and A. Riotto, Phys. Lett. B **343** (1995) 207 [arXiv:hep-ph/9410225].
- [182] J. Kang, P. Langacker, T. j. Li and T. Liu, Phys. Rev. Lett. **94** (2005) 061801 [arXiv:hep-ph/0402086].
- [183] J. Kang, P. Langacker, T. Li and T. Liu, arXiv:0911.2939.
- [184] R. Apreda, M. Maggiore, A. Nicolis and A. Riotto, Class. Quant. Grav. **18** (2001) L155 [arXiv:hep-ph/0102140].

- [185] R. Apreda, M. Maggiore, A. Nicolis and A. Riotto, Nucl. Phys. B **631** (2002) 342 [arXiv:gr-qc/0107033].
- [186] M. Bastero-Gil and S. F. King, Phys. Lett. B **423** (1998) 27 [arXiv:hep-ph/9709502].
- [187] M. Bastero-Gil and S. F. King, Nucl. Phys. B **549** (1999) 391 [arXiv:hep-ph/9806477].
- [188] M. Bastero-Gil, V. Di Clemente and S. F. King, Phys. Rev. D **71** (2005) 103517 [arXiv:hep-ph/0408336].
- [189] O. J. Eyton-Williams and S. F. King, Phys. Lett. B **610** (2005) 87 [arXiv:hep-ph/0411170].
- [190] O. J. Eyton-Williams and S. F. King, JHEP **0506** (2005) 040 [arXiv:hep-ph/0502156].
- [191] S. Antusch, O. J. Eyton-Williams and S. F. King, JHEP **0508** (2005) 103 [arXiv:hep-ph/0505140].
- [192] J. i. Kamoshita, Y. Okada and M. Tanaka, Phys. Lett. B **328** (1994) 67 [arXiv:hep-ph/9402278].
- [193] S. F. King and P. L. White, Phys. Rev. D **53** (1996) 4049 [arXiv:hep-ph/9508346].
- [194] S. W. Ham, H. Genten, B. R. Kim and S. K. Oh, Phys. Lett. B **383** (1996) 179 [arXiv:hep-ph/9606361].
- [195] J. F. Gunion, H. E. Haber and T. Moroi, *In the Proceedings of 1996 DPF / DPB Summer Study on New Directions for High-Energy Physics (Snowmass 96), Snowmass, Colorado, 25 Jun - 12 Jul 1996, pp LTH095* [arXiv:hep-ph/9610337].
- [196] S. W. Ham, S. K. Oh and B. R. Kim, Phys. Lett. B **414** (1997) 305 [arXiv:hep-ph/9612294].
- [197] U. Ellwanger, M. Rausch de Traubenberg and C. A. Savoy, Phys. Lett. B **315** (1993) 331 [arXiv:hep-ph/9307322].
- [198] N. V. Krasnikov, Mod. Phys. Lett. A **13** (1998) 893 [arXiv:hep-ph/9709467].
- [199] D. A. Demir, arXiv:hep-ph/9902468.
- [200] R. Dermisek and J. F. Gunion, Phys. Rev. Lett. **95** (2005) 041801 [arXiv:hep-ph/0502105].
- [201] U. Ellwanger, J. F. Gunion and C. Hugonie, JHEP **0507** (2005) 041 [arXiv:hep-ph/0503203].
- [202] R. Dermisek and J. F. Gunion, Phys. Rev. D **73** (2006) 111701 [arXiv:hep-ph/0510322].
- [203] S. Chang, P. J. Fox and N. Weiner, JHEP **0608** (2006) 068 [arXiv:hep-ph/0511250].
- [204] P. W. Graham, A. Pierce and J. G. Wacker, arXiv:hep-ph/0605162.
- [205] R. Dermisek and J. F. Gunion, Phys. Rev. D **76** (2007) 095006 [arXiv:0705.4387 [hep-ph]].

- [206] R. Dermisek, arXiv:0806.0847 [hep-ph].
- [207] R. Dermisek and J. F. Gunion, Phys. Rev. D **79** (2009) 055014 [arXiv:0811.3537 [hep-ph]].
- [208] S. Chang, R. Dermisek, J. F. Gunion and N. Weiner, Ann. Rev. Nucl. Part. Sci. **58** (2008) 75 [arXiv:0801.4554 [hep-ph]].
- [209] G. Abbiendi *et al.* [OPAL Collaboration], Eur. Phys. J. C **27** (2003) 311 [arXiv:hep-ex/0206022].
- [210] G. Abbiendi *et al.* [OPAL Collaboration], Eur. Phys. J. C **37** (2004) 49 [arXiv:hep-ex/0406057].
- [211] J. Abdallah *et al.* [DELPHI Collaboration], Eur. Phys. J. C **38** (2004) 1 [arXiv:hep-ex/0410017].
- [212] G. Abbiendi *et al.* [OPAL Collaboration], Eur. Phys. J. C **27** (2003) 483 [arXiv:hep-ex/0209068].
- [213] S. Schael *et al.* [ALEPH Collaboration], JHEP **1005** (2010) 049 [arXiv:1003.0705].
- [214] R. Barate *et al.* [LEP Working Group for Higgs boson searches and ALEPH Collaboration and and], Phys. Lett. B **565** (2003) 61 [arXiv:hep-ex/0306033].
- [215] R. Dermisek and J. F. Gunion, Phys. Rev. D **77** (2008) 015013 [arXiv:0709.2269 [hep-ph]].
- [216] A. Djouadi, U. Ellwanger and A. M. Teixeira, Phys. Rev. Lett. **101** (2008) 101802 [arXiv:0803.0253 [hep-ph]].
- [217] A. Djouadi, U. Ellwanger and A. M. Teixeira, JHEP **0904** (2009) 031 [arXiv:0811.2699 [hep-ph]].
- [218] L. Cavicchia, Nucl. Phys. B **813** (2009) 123 [arXiv:0807.3921 [hep-ph]].
- [219] J. R. Espinosa and J. F. Gunion, Phys. Rev. Lett. **82** (1999) 1084 [arXiv:hep-ph/9807275].
- [220] U. Ellwanger, J. F. Gunion, C. Hugonie and S. Moretti, arXiv:hep-ph/0305109.
- [221] D. J. 2. Miller and S. Moretti, arXiv:hep-ph/0403137.
- [222] G. Weiglein *et al.* [LHC/LC Study Group], Phys. Rept. **426** (2006) 47 [arXiv:hep-ph/0410364].
- [223] J. F. Gunion, H. E. Haber and R. J. Van Kooten, arXiv:hep-ph/0301023.
- [224] A. Djouadi, Phys. Rept. **459** (2008) 1 [arXiv:hep-ph/0503173].
- [225] S. Moretti and S. Munir, Eur. Phys. J. C **47** (2006) 791 [arXiv:hep-ph/0603085].
- [226] U. Ellwanger, J. F. Gunion and C. Hugonie, arXiv:hep-ph/0111179.
- [227] U. Aglietti *et al.*, arXiv:hep-ph/0612172.

- [228] A. Djouadi *et al.*, JHEP **0807** (2008) 002 [arXiv:0801.4321 [hep-ph]].
- [229] I. Rottlander and M. Schumacher, “Investigation of the LHC discovery potential for higgs bosons in the NMSSM,” ATL-PHYS-CONF-2008-009.
- [230] T. Stelzer, S. Wiesenfeldt and S. Willenbrock, Phys. Rev. D **75** (2007) 077701 [arXiv:hep-ph/0611242].
- [231] U. Ellwanger, J. F. Gunion, C. Hugonie and S. Moretti, arXiv:hep-ph/0401228.
- [232] S. Moretti, S. Munir and P. Poulose, Phys. Lett. B **644** (2007) 241 [arXiv:hep-ph/0608233].
- [233] K. Cheung, J. Song and Q. S. Yan, Phys. Rev. Lett. **99** (2007) 031801 [arXiv:hep-ph/0703149].
- [234] M. Carena, T. Han, G. Y. Huang and C. E. M. Wagner, JHEP **0804** (2008) 092 [arXiv:0712.2466 [hep-ph]].
- [235] S. Chang, P. J. Fox and N. Weiner, Phys. Rev. Lett. **98** (2007) 111802 [arXiv:hep-ph/0608310].
- [236] A. Belyaev, S. Hesselbach, S. Lehti, S. Moretti, A. Nikitenko and C. H. Shepherd-Themistocleous, arXiv:0805.3505 [hep-ph].
- [237] J. R. Forshaw, J. F. Gunion, L. Hodgkinson, A. Papaefstathiou and A. D. Pilkington, JHEP **0804** (2008) 090 [arXiv:0712.3510 [hep-ph]].
- [238] M. Lisanti and J. G. Wacker, Phys. Rev. D **79** (2009) 115006 [arXiv:0903.1377 [hep-ph]].
- [239] R. Dermisek and J. F. Gunion, Phys. Rev. D **81** (2010) 055001 [arXiv:0911.2460 [hep-ph]].
- [240] A. Arhrib, K. Cheung, T. J. Hou and K. W. Song, JHEP **0703** (2007) 073 [arXiv:hep-ph/0606114].
- [241] K. Cheung and T. J. Hou, Phys. Lett. B **674** (2009) 54 [arXiv:0809.1122 [hep-ph]].
- [242] M. Drees, M. Guchait and D.P. Roy, Phys. Lett. B **471** (1999) 39 [arXiv:hep-ph/9909266].
- [243] A. G. Akeroyd, A. Arhrib and Q. S. Yan, Eur. Phys. J. C **55** (2008) 653 [arXiv:0712.3933 [hep-ph]].
- [244] [CDF Collaboration and D0 Collaboration], “Combined CDF and DZero Upper Limits on Standard Model Higgs-Boson Production with up to 4.2 fb⁻¹ of Data,” arXiv:0903.4001 [hep-ex].
- [245] V. M. Abazov *et al.* [D0 Collaboration], Phys. Rev. Lett. **102** (2009) 051804 [arXiv:0811.0024 [hep-ex]].
- [246] D0 collaboration, D0 Note 5891-CONF (2009); see also L. Scodellaro, CDF and D0 Collaborations, “Searches for BSM Higgs at the Tevatron,” arXiv:0905.2554 [hep-ex].

- [247] P. N. Pandita, Phys. Rev. D **50** (1994) 571.
- [248] P. N. Pandita, Z. Phys. C **63** (1994) 659.
- [249] S. Y. Choi, D. J. Miller and P. M. Zerwas, Nucl. Phys. B **711** (2005) 83 [arXiv:hep-ph/0407209].
- [250] P. N. Pandita, Phys. Rev. D **53** (1996) 566.
- [251] S. Hesselbach, D. J. Miller, G. Moortgat-Pick, R. Nevzorov and M. Trusov, Phys. Lett. B **662** (2008) 199 [arXiv:0712.2001 [hep-ph]].
- [252] F. Franke and H. Fraas, Z. Phys. C **72** (1996) 309 [arXiv:hep-ph/9511275].
- [253] M. Drees and X. Tata, Phys. Rev. D **43** (1991) 2971.
- [254] F. Franke, H. Fraas and A. Bartl, Phys. Lett. B **336** (1994) 415 [arXiv:hep-ph/9408217].
- [255] J. Abdallah *et al.* [DELPHI Collaboration], Eur. Phys. J. C **31** (2004) 421 [arXiv:hep-ex/0311019].
- [256] G. Abbiendi *et al.* [OPAL Collaboration], Eur. Phys. J. C **35** (2004) 1 [arXiv:hep-ex/0401026].
- [257] S. Y. Choi, J. Kalinowski, G. A. Moortgat-Pick and P. M. Zerwas, Eur. Phys. J. C **22** (2001) 563 [Addendum-ibid. C **23** (2002) 769] [arXiv:hep-ph/0108117].
- [258] F. Franke and S. Hesselbach, Phys. Lett. B **526** (2002) 370 [arXiv:hep-ph/0111285].
- [259] G. A. Moortgat-Pick, S. Hesselbach, F. Franke and H. Fraas, JHEP **0506** (2005) 048 [arXiv:hep-ph/0502036].
- [260] R. Basu, P. N. Pandita and C. Sharma, Phys. Rev. D **77** (2008) 115009 [arXiv:0711.2121 [hep-ph]].
- [261] S. Kraml and W. Porod, Phys. Lett. B **626** (2005) 175 [arXiv:hep-ph/0507055].
- [262] J. D. Mason, D. E. Morrissey and D. Poland, Phys. Rev. D **80** (2009) 115015 [arXiv:0909.3523].
- [263] S. Kraml, A. R. Raklev and M. J. White, Phys. Lett. B **672** (2009) 361 [arXiv:0811.0011 [hep-ph]].
- [264] U. Ellwanger and C. Hugonie, Eur. Phys. J. C **13** (2000) 681 [arXiv:hep-ph/9812427].
- [265] S. Hesselbach, F. Franke and H. Fraas, Phys. Lett. B **492** (2000) 140 [arXiv:hep-ph/0007310].
- [266] S. P. Martin, Phys. Rev. D **62** (2000) 095008 [arXiv:hep-ph/0005116].
- [267] V. Barger, P. Langacker and G. Shaughnessy, Phys. Lett. B **644** (2007) 361 [arXiv:hep-ph/0609068].

- [268] F. Domingo and U. Ellwanger, *Mod. Phys. Lett. A* **24**, 557 (2009) [arXiv:0812.1167 [hep-ph]].
- [269] G. Hiller, *Phys. Rev. D* **70** (2004) 034018 [arXiv:hep-ph/0404220].
- [270] F. Domingo and U. Ellwanger, *JHEP* **0712** (2007) 090 [arXiv:0710.3714 [hep-ph]].
- [271] R. N. Hodgkinson and A. Pilaftsis, *Phys. Rev. D* **78** (2008) 075004 [arXiv:0807.4167 [hep-ph]].
- [272] Z. Heng, R. J. Oakes, W. Wang, Z. Xiong and J. M. Yang, *Phys. Rev. D* **77** (2008) 095012 [arXiv:0801.1169 [hep-ph]].
- [273] A. J. Buras, P. H. Chankowski, J. Rosiek and L. Slawianowska, *Nucl. Phys. B* **659** (2003) 3 [arXiv:hep-ph/0210145].
- [274] E. Barberio *et al.* [Heavy Flavor Averaging Group], arXiv:0808.1297 [hep-ex].
- [275] M. Misiak *et al.*, *Phys. Rev. Lett.* **98** (2007) 022002 [arXiv:hep-ph/0609232].
- [276] M. Misiak and M. Steinhauser, *Nucl. Phys. B* **764** (2007) 62 [arXiv:hep-ph/0609241].
- [277] T. Becher and M. Neubert, *Phys. Rev. Lett.* **98** (2007) 022003 [arXiv:hep-ph/0610067].
- [278] M. S. Carena, A. Menon, R. Noriega-Papaqui, A. Szyrkman and C. E. M. Wagner, *Phys. Rev. D* **74** (2006) 015009 [arXiv:hep-ph/0603106].
- [279] G. Degrossi, P. Gambino and P. Slavich, *Phys. Lett. B* **635** (2006) 335 [arXiv:hep-ph/0601135].
- [280] CDF Collaboration, “Search for $B_s^0 \rightarrow \mu^+\mu^-$ and $B_d^0 \rightarrow \mu^+\mu^-$ Decays in 2 fb^{-1} of $p\bar{p}$ Collisions with CDF II”, CDF Public Note 8956 (2007).
- [281] G. Buchalla, A. J. Buras and M. E. Lautenbacher, *Rev. Mod. Phys.* **68** (1996) 1125 [arXiv:hep-ph/9512380].
- [282] A. Abulencia *et al.* [CDF Collaboration], *Phys. Rev. Lett.* **97** (2006) 242003 [arXiv:hep-ex/0609040].
- [283] P. Ball and R. Fleischer, *Eur. Phys. J. C* **48** (2006) 413 [arXiv:hep-ph/0604249].
- [284] E. Dalgic *et al.*, *Phys. Rev. D* **76** (2007) 011501 [arXiv:hep-lat/0610104].
- [285] E. Barberio *et al.* [Heavy Flavor Averaging Group], “Averages of b-hadron and c-hadron Properties at the End of 2007,” arXiv:0808.1297 [hep-ex].
- [286] M. Okamoto, “Full determination of the CKM matrix using recent results from lattice QCD,” *PoS LAT2005* (2006) 013 [arXiv:hep-lat/0510113].
- [287] A. J. Buras, P. H. Chankowski, J. Rosiek and L. Slawianowska, *Phys. Lett. B* **546** (2002) 96 [arXiv:hep-ph/0207241].
- [288] B. Aubert *et al.* [BABAR Collaboration], *Phys. Rev. Lett.* **93** (2004) 081802 [arXiv:hep-ex/0404006].

- [289] M. Iwasaki *et al.* [Belle Collaboration], Phys. Rev. D **72** (2005) 092005 [arXiv:hep-ex/0503044].
- [290] T. Hurth, G. Isidori, J. F. Kamenik and F. Mescia, Nucl. Phys. B **808** (2009) 326 [arXiv:0807.5039 [hep-ph]].
- [291] W. M. Yao *et al.* [Particle Data Group], “Review of particle physics,” J. Phys. G **33** (2006) 1.
- [292] B. McElrath, Phys. Rev. D **72** (2005) 103508 [arXiv:hep-ph/0506151].
- [293] R. Dermisek, J. F. Gunion and B. McElrath, Phys. Rev. D **76** (2007) 051105 [arXiv:hep-ph/0612031].
- [294] E. Fullana and M. A. Sanchis-Lozano, Phys. Lett. B **653** (2007) 67 [arXiv:hep-ph/0702190].
- [295] M. L. Mangano and P. Nason, Mod. Phys. Lett. A **22** (2007) 1373 [arXiv:0704.1719 [hep-ph]].
- [296] R. N. Hodgkinson, Phys. Lett. B **665** (2008) 219 [arXiv:0802.3197 [hep-ph]].
- [297] D. McKeen, Phys. Rev. D **79** (2009) 015007 [arXiv:0809.4787 [hep-ph]].
- [298] F. Domingo, U. Ellwanger, E. Fullana, C. Hugonie and M. A. Sanchis-Lozano, JHEP **0901** (2009) 061 [arXiv:0810.4736 [hep-ph]].
- [299] F. Domingo, U. Ellwanger and M. A. Sanchis-Lozano, Phys. Rev. Lett. **103** (2009) 111802 [arXiv:0907.0348 [hep-ph]].
- [300] F. Wilczek, Phys. Rev. Lett. **39** (1977) 1304.
- [301] J. F. Gunion, H. E. Haber, G. L. Kane and S. Dawson, *The Higgs Hunter’s Guide* (Perseus Publishing, Cambridge (US), MA, 1990).
- [302] W. Love *et al.* [CLEO Collaboration], Phys. Rev. Lett. **101** (2008) 151802 [arXiv:0807.1427 [hep-ex]].
- [303] M. Drees and K. i. Hikasa, Phys. Rev. D **41** (1990) 1547.
- [304] M. A. Sanchis-Lozano, Nucl. Phys. Proc. Suppl. **142** (2005) 163 [arXiv:hep-ph/0407320].
- [305] B. Aubert *et al.* [BABAR Collaboration], Phys. Rev. Lett. **101** (2008) 071801 [Erratum-ibid. **102** (2009) 029901] [arXiv:0807.1086 [hep-ex]].
- [306] B. Aubert *et al.* [BABAR Collaboration], Phys. Rev. Lett. **103** (2009) 161801 [arXiv:0903.1124 [hep-ex]].
- [307] S. Recksiegel and Y. Sumino, Phys. Lett. B **578** (2004) 369 [arXiv:hep-ph/0305178].
- [308] B. A. Kniehl, A. A. Penin, A. Pineda, V. A. Smirnov and M. Steinhauser, Phys. Rev. Lett. **92** (2004) 242001 [arXiv:hep-ph/0312086].
- [309] A. A. Penin, “The mass of η_b ,” arXiv:0905.4296 [hep-ph].

- [310] N. Brambilla *et al.* [Quarkonium Working Group], arXiv:hep-ph/0412158.
- [311] G. W. Bennett *et al.* [Muon G-2 Collaboration], Phys. Rev. D **73** (2006) 072003 [arXiv:hep-ex/0602035].
- [312] M. Davier, Nucl. Phys. Proc. Suppl. **169** (2007) 288 [arXiv:hep-ph/0701163].
- [313] J. P. Miller, E. de Rafael and B. L. Roberts, Rept. Prog. Phys. **70** (2007) 795 [arXiv:hep-ph/0703049].
- [314] F. Jegerlehner, Acta Phys. Polon. B **38** (2007) 3021 [arXiv:hep-ph/0703125].
- [315] F. Domingo and U. Ellwanger, JHEP **0807** (2008) 079 [arXiv:0806.0733 [hep-ph]].
- [316] Z. Zhang, “Muon g-2: a mini review,” arXiv:0801.4905 [hep-ph].
- [317] M. Davier *et al.*, Eur. Phys. J. C **66** (2010) 127 [arXiv:0906.5443 [hep-ph]].
- [318] M. Davier, A. Hoecker, B. Malaescu, C. Z. Yuan and Z. Zhang, arXiv:0908.4300 [hep-ph].
- [319] D. Stockinger, J. Phys. G **34** (2007) R45 [arXiv:hep-ph/0609168].
- [320] H. Konig, Z. Phys. C **52** (1991) 159.
- [321] J. F. Gunion, D. Hooper and B. McElrath, Phys. Rev. D **73** (2006) 015011 [arXiv:hep-ph/0509024].
- [322] J. F. Gunion, JHEP **0908** (2009) 032 [arXiv:0808.2509 [hep-ph]].
- [323] D. Chang, W. F. Chang, C. H. Chou and W. Y. Keung, Phys. Rev. D **63** (2001) 091301 [arXiv:hep-ph/0009292].
- [324] K. m. Cheung, C. H. Chou and O. C. W. Kong, Phys. Rev. D **64** (2001) 111301 [arXiv:hep-ph/0103183].
- [325] M. Krawczyk, Acta Phys. Polon. B **33** (2002) 2621 [arXiv:hep-ph/0208076].
- [326] M. A. Sanchis-Lozano, Mod. Phys. Lett. A **17** (2002) 2265 [arXiv:hep-ph/0206156].
- [327] M. A. Sanchis-Lozano, Int. J. Mod. Phys. A **19** (2004) 2183 [arXiv:hep-ph/0307313].
- [328] B. Aubert [The BABAR Collaboration], arXiv:0902.2176 [hep-ex] and B. Aubert *et al.* [BABAR Collaboration], Phys. Rev. Lett. **103** (2009) 081803 [arXiv:0905.4539 [hep-ex]].
- [329] Y. Bai, M. Carena and J. Lykken, Phys. Rev. D **80** (2009) 055004 [arXiv:0905.2964 [hep-ph]].
- [330] X. G. He, J. Tandean and G. Valencia, Phys. Rev. Lett. **98** (2007) 081802 [arXiv:hep-ph/0610362].
- [331] G. Xiangdong, C. S. Li, Z. Li and H. Zhang, Eur. Phys. J. C **55** (2008) 317 [arXiv:0712.0257 [hep-ph]].

- [332] X. G. He, J. Tandean and G. Valencia, JHEP **0806** (2008) 002 [arXiv:0803.4330 [hep-ph]].
- [333] Q. Chang and Y. D. Yang, Phys. Lett. B **676** (2009) 88 [arXiv:0808.2933 [hep-ph]].
- [334] H. K. Park *et al.* [HyperCP Collaboration], Phys. Rev. Lett. **88** (2002) 111801 [arXiv:hep-ex/0110033].
- [335] S. A. Abel, S. Sarkar and I. B. Whittingham, Nucl. Phys. B **392** (1993) 83 [arXiv:hep-ph/9209292].
- [336] T. Elliott, S. F. King and P. L. White, Phys. Lett. B **351** (1995) 213 [arXiv:hep-ph/9406303].
- [337] P. Brax, U. Ellwanger and C. A. Savoy, Phys. Lett. B **347** (1995) 269 [arXiv:hep-ph/9411397].
- [338] G. V. Kraniotis, Z. Phys. C **68** (1995) 491 [arXiv:hep-ph/9501393].
- [339] A. Stephan, Phys. Rev. D **58** (1998) 035011 [arXiv:hep-ph/9709262].
- [340] U. Ellwanger and C. Hugonie, Phys. Lett. B **457** (1999) 299 [arXiv:hep-ph/9902401].
- [341] S. A. Abel and C. A. Savoy, Nucl. Phys. B **532** (1998) [arXiv:hep-ph/9803218].
- [342] A. Stephan, Phys. Lett. B **411** (1997) 97 [arXiv:hep-ph/9704232].
- [343] D. N. Spergel *et al.* [WMAP Collaboration], Astrophys. J. Suppl. **170** (2007) 377 [arXiv:astro-ph/0603449].
- [344] M. Tegmark *et al.* [SDSS Collaboration], Phys. Rev. D **74** (2006) 123507 [arXiv:astro-ph/0608632].
- [345] U. Ellwanger and C. Hugonie, Comput. Phys. Commun. **177** (2007) 399 [arXiv:hep-ph/0612134].
- [346] C. Hugonie, G. Belanger and A. Pukhov, JCAP **0711** (2007) 009 [arXiv:0707.0628 [hep-ph]].
- [347] G. Belanger, C. Hugonie and A. Pukhov, JCAP **0901** (2009) 023 [arXiv:0811.3224 [hep-ph]].
- [348] C. Balazs and D. Carter, Phys. Rev. D **78** (2008) 055001 [arXiv:0808.0770 [hep-ph]].
- [349] G. F. Giudice and R. Rattazzi, Phys. Rept. **322** (1999) 419 [arXiv:hep-ph/9801271].
- [350] A. de Gouvea, A. Friedland and H. Murayama, Phys. Rev. D **57** (1998) 5676 [arXiv:hep-ph/9711264].
- [351] M. Dine, A. E. Nelson and Y. Shirman, Phys. Rev. D **51** (1995) 1362 [arXiv:hep-ph/9408384].
- [352] M. Dine, A. E. Nelson, Y. Nir and Y. Shirman, Phys. Rev. D **53** (1996) 2658 [arXiv:hep-ph/9507378].

- [353] G. R. Dvali, G. F. Giudice and A. Pomarol, Nucl. Phys. B **478** (1996) 31 [arXiv:hep-ph/9603238].
- [354] P. Ciafaloni and A. Strumia, Nucl. Phys. B **494** (1997) 41 [arXiv:hep-ph/9611204].
- [355] K. Agashe and M. Graesser, Nucl. Phys. B **507** (1997) 3 [arXiv:hep-ph/9704206].
- [356] G. F. Giudice and R. Rattazzi, Nucl. Phys. B **511** (1998) 25 [arXiv:hep-ph/9706540].
- [357] P. Langacker, N. Polonsky and J. Wang, Phys. Rev. D **60** (1999) 115005 [arXiv:hep-ph/9905252].
- [358] S. L. Dubovsky, D. S. Gorbunov and S. V. Troitsky, Phys. Usp. **42** (1999) 623 [Usp. Fiz. Nauk **169** (1999) 705] [arXiv:hep-ph/9905466].
- [359] T. Han, D. Marfatia and R. J. Zhang, Phys. Rev. D **61** (2000) 013007 [arXiv:hep-ph/9906508].
- [360] Z. Chacko and E. Ponton, Phys. Rev. D **66** (2002) 095004 [arXiv:hep-ph/0112190].
- [361] D. Suematsu, Eur. Phys. J. C **38** (2004) 359 [arXiv:hep-ph/0308007].
- [362] A. Delgado, G. F. Giudice and P. Slavich, Phys. Lett. B **653** (2007) 424 [arXiv:0706.3873 [hep-ph]].
- [363] G. F. Giudice, H. D. Kim and R. Rattazzi, Phys. Lett. B **660** (2008) 545 [arXiv:0711.4448 [hep-ph]].
- [364] M. Dine and J. D. Mason, Phys. Rev. D **78** (2008) 055013 [arXiv:0712.1355 [hep-ph]].
- [365] T. Liu and C. E. M. Wagner, JHEP **0806** (2008) 073 [arXiv:0803.2895 [hep-ph]].
- [366] Z. Komargodski and N. Seiberg, JHEP **0903** (2009) 072 [arXiv:0812.3900 [hep-ph]].
- [367] J. D. Mason, Phys. Rev. D **80** (2009) 015026 [arXiv:0904.4485 [hep-ph]].
- [368] P. C. Schuster and N. Toro, arXiv:hep-ph/0512189.
- [369] Z. Chacko, M. A. Luty, I. Maksymyk and E. Ponton, JHEP **0004** (2000) 001 [arXiv:hep-ph/9905390].
- [370] D. E. Kaplan, G. D. Kribs and M. Schmaltz, Phys. Rev. D **62** (2000) 035010 [arXiv:hep-ph/9911293].
- [371] N. Arkani-Hamed, L. J. Hall, Y. Nomura, D. Tucker-Smith and N. Weiner, Nucl. Phys. B **605** (2001) 81 [arXiv:hep-ph/0102090].
- [372] T. Kobayashi and H. Terao, Phys. Rev. D **64** (2001) 075003 [arXiv:hep-ph/0103028].
- [373] R. Kitano, G. D. Kribs and H. Murayama, Phys. Rev. D **70** (2004) 035001 [arXiv:hep-ph/0402215].
- [374] L. M. Carpenter, P. J. Fox and D. E. Kaplan, arXiv:hep-ph/0503093.

- [375] K. Choi, K. S. Jeong and K. i. Okumura, JHEP **0509** (2005) 039 [arXiv:hep-ph/0504037].
- [376] C. Panagiotakopoulos and A. Pilaftsis, Phys. Lett. B **505** (2001) 184 [arXiv:hep-ph/0101266].
- [377] A. Dedes, C. Hugonie, S. Moretti and K. Tamvakis, Phys. Rev. D **63** (2001) 055009 [arXiv:hep-ph/0009125].
- [378] V. Barger, P. Langacker and H. S. Lee, Phys. Lett. B **630** (2005) 85 [arXiv:hep-ph/0508027].
- [379] C. Balazs, M. S. Carena, A. Freitas and C. E. M. Wagner, JHEP **0706** (2007) 066 [arXiv:0705.0431 [hep-ph]].
- [380] J. Cao, H. E. Logan and J. M. Yang, Phys. Rev. D **79** (2009) 091701 [arXiv:0901.1437 [hep-ph]].
- [381] R. Dermisek, Mod. Phys. Lett. A **24** (2009) 1631 [arXiv:0907.0297 [hep-ph]].
- [382] T. Kobayashi, H. Terao and A. Tsuchiya, Phys. Rev. D **74** (2006) 015002 [arXiv:hep-ph/0604091].
- [383] Z. Chacko, Y. Nomura and D. Tucker-Smith, Nucl. Phys. B **725** (2005) 207 [arXiv:hep-ph/0504095].
- [384] Y. Nomura, D. Poland and B. Tweedie, Nucl. Phys. B **745** (2006) 29 [arXiv:hep-ph/0509243].
- [385] D. E. Lopez-Fogliani, L. Roszkowski, R. R. de Austri and T. A. Varley, Phys. Rev. D **80** (2009) 095013 [arXiv:0906.4911 [hep-ph]].
- [386] C. Balazs and D. Carter, AIP Conf. Proc. **1178** (2009) 23 [JHEP **1003** (2010) 016] [arXiv:0906.5012 [hep-ph]].
- [387] E. Accomando *et al.*, arXiv:hep-ph/0608079.
- [388] J. C. Romao, Phys. Lett. B **173** (1986) 309.
- [389] K. S. Babu and S. M. Barr, Phys. Rev. D **49** (1994) 2156 [arXiv:hep-ph/9308217].
- [390] N. Haba, M. Matsuda and M. Tanimoto, Phys. Rev. D **54** (1996) 6928 [arXiv:hep-ph/9512421].
- [391] S. W. Ham, S. K. Oh and H. S. Song, Phys. Rev. D **61** (2000) 055010 [arXiv:hep-ph/9910461].
- [392] H. Georgi and A. Pais, Phys. Rev. D **10** (1974) 1246.
- [393] G. C. Branco, F. Kruger, J. C. Romao and A. M. Teixeira, JHEP **0107** (2001) 027 [arXiv:hep-ph/0012318].
- [394] A. T. Davies, C. D. Froggatt and A. Usai, Phys. Lett. B **517** (2001) 375 [arXiv:hep-ph/0105266].

- [395] C. Hugonie, J. C. Romao and A. M. Teixeira, JHEP **0306** (2003) 020 [arXiv:hep-ph/0304116].
- [396] A. Pomarol, Phys. Rev. D **47** (1993) 273 [arXiv:hep-ph/9208205].
- [397] O. Lebedev, Int. J. Mod. Phys. A **15** (2000) 2987 [arXiv:hep-ph/9905216].
- [398] S. M. Barr, Phys. Rev. D **47** (1993) 2025.
- [399] M. Matsuda and M. Tanimoto, Phys. Rev. D **52** (1995) 3100 [arXiv:hep-ph/9504260].
- [400] N. Haba, Prog. Theor. Phys. **97** (1997) 301 [arXiv:hep-ph/9608357].
- [401] K. Funakubo and S. Tao, Prog. Theor. Phys. **113** (2005) 821 [arXiv:hep-ph/0409294].
- [402] S. W. Ham, J. Kim, S. K. Oh and D. Son, Phys. Rev. D **64** (2001) 035007 [arXiv:hep-ph/0104144].
- [403] S. W. Ham, S. H. Kim, S. K. OH and D. Son, Phys. Rev. D **76** (2007) 115013 [arXiv:0708.2755 [hep-ph]].
- [404] R. Garisto, Phys. Rev. D **49** (1994) 4820 [arXiv:hep-ph/9311249].
- [405] S. W. Ham, S. K. Oh and D. Son, Phys. Rev. D **65** (2002) 075004 [arXiv:hep-ph/0110052].
- [406] S. W. Ham, Y. S. Jeong and S. K. Oh, arXiv:hep-ph/0308264.
- [407] G. M. Asatrian and G. K. Egüian, arXiv:hep-ph/9605458.
- [408] S. W. Ham, E. J. Yoo and S. K. Oh, Phys. Rev. D **76** (2007) 015004 [arXiv:hep-ph/0703041].
- [409] S. W. Ham, J. O. Im and S. K. OH, Eur. Phys. J. C **58** (2008) 579 [arXiv:0805.1115 [hep-ph]].
- [410] M. Boz, Mod. Phys. Lett. A **21** (2006) 243 [arXiv:hep-ph/0511072].
- [411] D. A. Demir, Phys. Rev. D **62** (2000) 075003 [arXiv:hep-ph/9911435].
- [412] S. W. Ham, S. K. Oh and D. Son, Phys. Rev. D **66** (2002) 015008 [arXiv:hep-ph/0110183].
- [413] S. W. Ham, S. K. OH, E. J. Yoo, C. M. Kim and D. Son, arXiv:hep-ph/0406070.
- [414] R. Barbier *et al.*, Phys. Rept. **420** (2005) 1 [arXiv:hep-ph/0406039].
- [415] G. R. Farrar and P. Fayet, Phys. Lett. B **76** (1978) 575.
- [416] S. Weinberg, Phys. Rev. D **26** (1982) 287.
- [417] N. Sakai and T. Yanagida, Nucl. Phys. B **197** (1982) 533.
- [418] M. Chemtob and P. N. Pandita, Phys. Rev. D **73** (2006) 055012 [arXiv:hep-ph/0601159].

- [419] M. Chemtob and P. N. Pandita, Phys. Rev. D **76** (2007) 095019 [arXiv:0708.1284 [hep-ph]].
- [420] P. N. Pandita and P. F. Paulraj, Phys. Lett. B **462** (1999) 294 [arXiv:hep-ph/9907561].
- [421] P. N. Pandita, Phys. Rev. D **64** (2001) 056002 [arXiv:hep-ph/0103005].
- [422] C. C. Jean-Louis and G. Moreau, arXiv:0911.3640.
- [423] P. Ghosh and S. Roy, JHEP **0904** (2009) 069 [arXiv:0812.0084 [hep-ph]].
- [424] A. Abada and G. Moreau, JHEP **0608** (2006) 044 [arXiv:hep-ph/0604216].
- [425] A. Abada, G. Bhattacharyya and G. Moreau, Phys. Lett. B **642** (2006) 503 [arXiv:hep-ph/0606179].
- [426] J. Gava and C. C. Jean-Louis, Phys. Rev. D **81** (2010) 013003 [arXiv:0907.3947 [hep-ph]].
- [427] R. Kitano and K. y. Oda, Phys. Rev. D **61** (2000) 113001 [arXiv:hep-ph/9911327].
- [428] D. E. Lopez-Fogliani and C. Munoz, Phys. Rev. Lett. **97** (2006) 041801 [arXiv:hep-ph/0508297].
- [429] R. Allahverdi, B. A. Campbell and K. A. Olive, Phys. Lett. B **341** (1994) 166 [arXiv:hep-ph/9402294].
- [430] N. Escudero, D. E. Lopez-Fogliani, C. Munoz and R. R. de Austri, JHEP **0812** (2008) 099 [arXiv:0810.1507 [hep-ph]].
- [431] A. Bartl, M. Hirsch, A. Vicente, S. Liebler and W. Porod, JHEP **0905** (2009) 120 [arXiv:0903.3596 [hep-ph]].
- [432] J. Fidalgo, D. E. Lopez-Fogliani, C. Munoz and R. Ruiz de Austri, JHEP **0908** (2009) 105 [arXiv:0904.3112 [hep-ph]].
- [433] I. Gogoladze, N. Okada and Q. Shafi, Phys. Lett. B **672** (2009) 235 [arXiv:0809.0703 [hep-ph]].
- [434] R. N. Mohapatra, N. Setzer and S. Spinner, JHEP **0804** (2008) 091 [arXiv:0802.1208 [hep-ph]].
- [435] D. G. Cerdeno, C. Munoz and O. Seto, Phys. Rev. D **79** (2009) 023510 [arXiv:0807.3029 [hep-ph]].
- [436] D. G. Cerdeno and O. Seto, JCAP **0908** (2009) 032 [arXiv:0903.4677 [hep-ph]].
- [437] D. Suematsu and Y. Yamagishi, Int. J. Mod. Phys. A **10** (1995) 4521 [arXiv:hep-ph/9411239].
- [438] M. Cvetič and P. Langacker, Phys. Rev. D **54** (1996) 3570 [arXiv:hep-ph/9511378]. [arXiv:hep-ph/9506380].

- [439] K. S. Babu, C. F. Kolda and J. March-Russell, Phys. Rev. D **54** (1996) 4635 [arXiv:hep-ph/9603212].
- [440] M. Cvetič, D. A. Demir, J. R. Espinosa, L. L. Everett and P. Langacker, Phys. Rev. D **56** (1997) 2861 [Erratum-ibid. D **58** (1998) 119905] [arXiv:hep-ph/9703317].
- [441] V. Barger, P. Langacker, H. S. Lee and G. Shaughnessy, Phys. Rev. D **73** (2006) 115010 [arXiv:hep-ph/0603247].
- [442] V. Barger, P. Langacker and G. Shaughnessy, New J. Phys. **9** (2007) 333 [arXiv:hep-ph/0702001].
- [443] P. Langacker, Rev. Mod. Phys. **81** (2008) 1199 [arXiv:0801.1345 [hep-ph]].
- [444] D. Suematsu and G. Zoupanos, JHEP **0106** (2001) 038 [arXiv:hep-ph/0102096].
- [445] J. Erler, Nucl. Phys. B **586** (2000) 73 [arXiv:hep-ph/0006051].
- [446] D. A. Demir, G. L. Kane and T. T. Wang, Phys. Rev. D **72** (2005) 015012 [arXiv:hep-ph/0503290].
- [447] D. E. Morrissey and J. D. Wells, Phys. Rev. D **74** (2006) 015008 [arXiv:hep-ph/0512019].
- [448] H. S. Lee, K. T. Matchev and T. T. Wang, Phys. Rev. D **77** (2008) 015016 [arXiv:0709.0763 [hep-ph]].
- [449] D. A. Demir and N. K. Pak, Phys. Rev. D **57** (1998) 6609 [arXiv:hep-ph/9809357].
- [450] Y. Daikoku and D. Suematsu, Phys. Rev. D **62** (2000) 095006 [arXiv:hep-ph/0003205].
- [451] H. Amini, New J. Phys. **5** (2003) 49 [arXiv:hep-ph/0210086].
- [452] S. W. Ham and S. K. OH, J. Phys. G **37** (2010) 045003 [arXiv:0906.5526 [hep-ph]].
- [453] A. Maloney, A. Pierce and J. G. Wacker, JHEP **0606** (2006) 034 [arXiv:hep-ph/0409127].
- [454] J. Erler, P. Langacker and T. j. Li, Phys. Rev. D **66** (2002) 015002 [arXiv:hep-ph/0205001].
- [455] T. Han, P. Langacker and B. McElrath, Phys. Rev. D **70** (2004) 115006 [arXiv:hep-ph/0405244].
- [456] D. A. Demir, L. Solmaz and S. Solmaz, Phys. Rev. D **73** (2006) 016001 [arXiv:hep-ph/0512134].
- [457] D. Suematsu, Phys. Rev. D **59** (1999) 055017 [arXiv:hep-ph/9808409].
- [458] D. Suematsu, Phys. Rev. D **57** (1998) 1738 [arXiv:hep-ph/9708413].
- [459] S. Hesselbach, F. Franke and H. Fraas, Eur. Phys. J. C **23** (2002) 149 [arXiv:hep-ph/0107080].

- [460] S. Y. Choi, H. E. Haber, J. Kalinowski and P. M. Zerwas, Nucl. Phys. B **778** (2007) 85 [arXiv:hep-ph/0612218].
- [461] D. Suematsu, Mod. Phys. Lett. A **12** (1997) 1709 [arXiv:hep-ph/9705412].
- [462] D. A. Demir and L. L. Everett, Phys. Rev. D **69** (2004) 015008 [arXiv:hep-ph/0306240].
- [463] B. de Carlos and J. R. Espinosa, Phys. Lett. B **407** (1997) 12 [arXiv:hep-ph/9705315].
- [464] V. Barger, C. Kao, P. Langacker and H. S. Lee, Phys. Lett. B **600** (2004) 104 [arXiv:hep-ph/0408120].
- [465] D. Suematsu, Phys. Rev. D **73** (2006) 035010 [arXiv:hep-ph/0511299].
- [466] S. Nakamura and D. Suematsu, Phys. Rev. D **75** (2007) 055004 [arXiv:hep-ph/0609061].
- [467] V. Barger, P. Langacker, I. Lewis, M. McCaskey, G. Shaughnessy and B. Yencho, Phys. Rev. D **75** (2007) 115002 [arXiv:hep-ph/0702036].
- [468] J. Kalinowski, S. F. King and J. P. Roberts, JHEP **0901** (2009) 066 [arXiv:0811.2204 [hep-ph]].
- [469] H. S. Lee, K. T. Matchev and S. Nasri, Phys. Rev. D **76** (2007) 041302 [arXiv:hep-ph/0702223].
- [470] T. Hur, H. S. Lee and S. Nasri, Phys. Rev. D **77** (2008) 015008 [arXiv:0710.2653 [hep-ph]].
- [471] J. h. Kang, P. Langacker and T. j. Li, Phys. Rev. D **71** (2005) 015012 [arXiv:hep-ph/0411404].
- [472] J. Kang and T. Li, Phys. Rev. D **71** (2005) 095011 [arXiv:hep-ph/0501101].
- [473] D. A. Demir, L. L. Everett and P. Langacker, Phys. Rev. Lett. **100** (2008) 091804 [arXiv:0712.1341 [hep-ph]].
- [474] D. Suematsu, T. Toma and T. Yoshida, arXiv:0910.3086 [hep-ph].
- [475] D. A. Demir, L. L. Everett, M. Frank, L. Selbuz and I. Turan, Phys. Rev. D **81** (2010) 035019 [arXiv:0906.3540 [hep-ph]].
- [476] R. S. Hundi, S. Pakvasa and X. Tata, Phys. Rev. D **79** (2009) 095011 [arXiv:0903.1631 [hep-ph]].
- [477] V. Barger, P. Langacker and G. Shaughnessy, Phys. Rev. D **75** (2007) 055013 [arXiv:hep-ph/0611239].
- [478] S. W. Ham, T. Hur, P. Ko and S. K. Oh, J. Phys. G **35** (2008) 095007 [arXiv:0801.2361 [hep-ph]].
- [479] S. W. Ham, E. J. Yoo, S. K. Oh and D. Son, Phys. Rev. D **77** (2008) 114011 [arXiv:0801.4640 [hep-ph]].

- [480] T. Cohen and A. Pierce, Phys. Rev. D **78** (2008) 055012 [arXiv:0803.0765 [hep-ph]].
- [481] Y. Daikoku and H. Okada, arXiv:0910.3370 [hep-ph].
- [482] V. Barger, C. Kao, P. Langacker and H. S. Lee, Phys. Lett. B **614** (2005) 67 [arXiv:hep-ph/0412136].
- [483] C. W. Chiang and E. Senaha, JHEP **0806** (2008) 019 [arXiv:0804.1719 [hep-ph]].
- [484] J. L. Hewett and T. G. Rizzo, Phys. Rept. **183** (1989) 193.
- [485] P. Langacker and J. Wang, Phys. Rev. D **58** (1998) 115010 [arXiv:hep-ph/9804428].
- [486] J. Kang and P. Langacker, Phys. Rev. D **71** (2005) 035014 [arXiv:hep-ph/0412190].
- [487] S. W. Ham, J. O. Im, E. J. Yoo and S. K. Oh, JHEP **0812** (2008) 017 [arXiv:0810.4194 [hep-ph]].
- [488] S. F. King, S. Moretti and R. Nevzorov, Phys. Rev. D **73** (2006) 035009 [arXiv:hep-ph/0510419].
- [489] S. F. King, S. Moretti and R. Nevzorov, Phys. Lett. B **634** (2006) 278 [arXiv:hep-ph/0511256].
- [490] S. F. King, S. Moretti and R. Nevzorov, Phys. Lett. B **650** (2007) 57 [arXiv:hep-ph/0701064].
- [491] J. P. Hall and S. F. King, JHEP **0908** (2009) 088 [arXiv:0905.2696 [hep-ph]].
- [492] P. Athron, S. F. King, D. J. . Miller, S. Moretti, R. Nevzorov and R. Nevzorov, Phys. Lett. B **681** (2009) 448 [arXiv:0901.1192 [hep-ph]].
- [493] P. Athron, S. F. King, D. J. Miller, S. Moretti and R. Nevzorov, Phys. Rev. D **80** (2009) 035009 [arXiv:0904.2169 [hep-ph]].
- [494] R. Howl and S. F. King, JHEP **0805** (2008) 008 [arXiv:0802.1909 [hep-ph]].
- [495] R. Howl and S. F. King, Phys. Lett. B **687** (2010) 355 [arXiv:0908.2067 [hep-ph]].
- [496] G. Jungman, M. Kamionkowski and K. Griest, Phys. Rept. **267** (1996) 195
- [497] B. R. Greene and P. J. Miron, Phys. Lett. B **168** (1986) 226.
- [498] K. A. Olive and D. Thomas, Nucl. Phys. B **355** (1991) 192.
- [499] R. Flores, K. A. Olive and D. Thomas, Phys. Lett. B **245** (1990) 509.
- [500] P. Gondolo and G. Gelmini, Nucl. Phys. B **360** (1991) 145.
- [501] G. Belanger, F. Boudjema, C. Hugonie, A. Pukhov and A. Semenov, JCAP **0509** (2005) 001 [arXiv:hep-ph/0505142].
- [502] G. Belanger, F. Boudjema, A. Pukhov and A. Semenov, Comput. Phys. Commun. **176** (2007) 367 [arXiv:hep-ph/0607059].

- [503] G. Belanger, F. Boudjema, A. Pukhov and A. Semenov, *Comput. Phys. Commun.* **180** (2009) 747 [arXiv:0803.2360 [hep-ph]].
- [504] T. Nihei, L. Roszkowski and R. Ruiz de Austri, *JHEP* **0203** (2002) 031 [arXiv:hep-ph/0202009].
- [505] K. Griest and D. Seckel, *Phys. Rev. D* **43** (1991) 3191.
- [506] J. Angle *et al.* [XENON Collaboration], *Phys. Rev. Lett.* **100** (2008) 021303 [arXiv:0706.0039 [astro-ph]].
- [507] Z. Ahmed *et al.* [CDMS Collaboration], *Phys. Rev. Lett.* **102** (2009) 011301 [arXiv:0802.3530 [astro-ph]].
- [508] J. Angle *et al.*, *Phys. Rev. Lett.* **101** (2008) 091301 [arXiv:0805.2939 [astro-ph]].
- [509] H. S. Lee. *et al.* [KIMS Collaboration], *Phys. Rev. Lett.* **99** (2007) 091301 [arXiv:0704.0423 [astro-ph]].
- [510] R. Bernabei *et al.* [DAMA Collaboration], *Phys. Lett. B* **480** (2000) 23.
- [511] R. A. Flores, K. A. Olive and D. Thomas, *Phys. Lett. B* **263** (1991) 425.
- [512] V. A. Bednyakov and H. V. Klapdor-Kleingrothaus, *Phys. Rev. D* **59** (1999) 023514 [arXiv:hep-ph/9802344].
- [513] V. A. Bednyakov and H. V. Klapdor-Kleingrothaus, *Phys. Atom. Nucl.* **62** (1999) 966 [*Yad. Fiz.* **62** (1999) 1033].
- [514] D. G. Cerdeno, C. Hugonie, D. E. Lopez-Fogliani, C. Munoz and A. M. Teixeira, *JHEP* **0412** (2004) 048 [arXiv:hep-ph/0408102].
- [515] D. G. Cerdeno, E. Gabrielli, D. E. Lopez-Fogliani, C. Munoz and A. M. Teixeira, *JCAP* **0706** (2007) 008 [arXiv:hep-ph/0701271].
- [516] F. Ferrer, L. M. Krauss and S. Profumo, *Phys. Rev. D* **74** (2006) 115007 [arXiv:hep-ph/0609257].
- [517] P. Jean *et al.*, *Astron. Astrophys.* **407** (2003) L55 [arXiv:astro-ph/0309484].
- [518] O. Adriani *et al.*, *Phys. Rev. Lett.* **102** (2009) 051101 [arXiv:0810.4994 [astro-ph]].
- [519] O. Adriani *et al.* [PAMELA Collaboration], *Nature* **458** (2009) 607 [arXiv:0810.4995 [astro-ph]].
- [520] D. Hooper and T. M. P. Tait, *Phys. Rev. D* **80** (2009) 055028 [arXiv:0906.0362 [hep-ph]].
- [521] W. Wang, Z. Xiong, J. M. Yang and L. X. Yu, *JHEP* **0911** (2009) 053 [arXiv:0908.0486 [hep-ph]].
- [522] J. H. Huh, J. E. Kim and B. Kyae, *Phys. Rev. D* **79** (2009) 063529 [arXiv:0809.2601 [hep-ph]].
- [523] Z. Kunszt and F. Zwirner, *Nucl. Phys. B* **385** (1992) 3 [arXiv:hep-ph/9203223].

- [524] V. D. Barger, M. S. Berger, A. L. Stange and R. J. N. Phillips, Phys. Rev. D **45** (1992) 4128.
- [525] A. Brignole and F. Zwirner, Phys. Lett. B **299** (1993) 72 [arXiv:hep-ph/9210266].
- [526] S. Heinemeyer and W. Hollik, Nucl. Phys. B **474** (1996) 32 [arXiv:hep-ph/9602318].
- [527] W. Hollik and S. Penaranda, Eur. Phys. J. C **23** (2002) 163 [arXiv:hep-ph/0108245].
- [528] F. Boudjema and A. Semenov, Phys. Rev. D **66** (2002) 095007 [arXiv:hep-ph/0201219].
- [529] D. R. T. Jones, Nucl. Phys. B **87** (1975) 127.
- [530] D. R. T. Jones and L. Mezincescu, Phys. Lett. B **136** (1984) 242.
- [531] P. C. West, Phys. Lett. B **137** (1984) 371.
- [532] S. P. Martin and M. T. Vaughn, Phys. Lett. B **318** (1993) 331 [arXiv:hep-ph/9308222].
- [533] Y. Yamada, Phys. Rev. Lett. **72** (1994) 25 [arXiv:hep-ph/9308304].
- [534] D. M. Pierce, J. A. Bagger, K. T. Matchev and R. j. Zhang, Nucl. Phys. B **491** (1997) 3 [arXiv:hep-ph/9606211].
- [535] W. Porod, Comput. Phys. Commun. **153** (2003) 275 [arXiv:hep-ph/0301101].
- [536] J. Reuter and F. Braam, AIP Conf. Proc. **1200** (2010) 470 [arXiv:0909.3059 [hep-ph]].
- [537] F. Mahmoudi, Comput. Phys. Commun. **180** (2009) 1718.