# Phase structure of Topologically massive gauge theory with fermion

Yuichi Hoshino Kushiro National College of Technology, Otanoshike-nishi 2-32-1, Kushro 084-0916, Hokkaido, Japan

April 21, 2022

#### Abstract

Using Bloch-Nordsieck approximation fermion propagator in 3-dimensional gauge theory with topological mass is studied. Infrared divergence of Chern-Simon term is soft, which modifes anomalous dimension. In unquenched QCD with 2-component spinor anomalous dimension has fractional value, where order parmeter is divergent.

# 1 Introduction

The Lagrangeans of Topologically massive gauge theory with fermion are(1)

$$L = \frac{1}{4}F_{\mu\nu}F^{\mu\nu} + \frac{1}{4}\theta\epsilon^{\mu\nu\rho}F_{\mu\nu}A_{\rho} + \overline{\psi}(i\gamma\cdot(\partial - ieA) - m)\psi + \frac{1}{2d}(\partial\cdot A)^{2}, \quad (1)$$

$$L = \frac{1}{4g^2} tr(F_{\mu\nu}F^{\mu\nu}) - \frac{\theta}{4g^2} \epsilon^{\mu\nu\rho} tr(F_{\mu\nu}A_\rho - \frac{2}{3}A_\mu A_\nu A_\rho)$$
  
+  $\overline{\psi}(i\gamma \cdot (\partial - ieA) - m)\psi + \frac{1}{2d}(\partial \cdot A)^2$  (2)

,where  $\theta = (g^2/4\pi)n$ ,  $(n = 0, \pm 1, \pm 2...)$  with  $\gamma$  matrix for 4-comonent fermion (2).In comparison with massless  $QED_3$ ,  $QCD_3$  topological mass term seems to soften the infrared divergence of massive fermion near its on-shell.In Minkowski metric we have 5  $\gamma$  matrices  $\{\gamma_{\mu}, \gamma_{\nu}\} = 2g_{\mu\nu}$ ,  $(\mu = 0, 1, 2)$ 

$$\begin{split} \gamma_0 &= \left( \begin{array}{cc} \sigma_3 & 0 \\ 0 & -\sigma_3 \end{array} \right), \gamma_{1,2} = -i \left( \begin{array}{cc} \sigma_{1,2} & 0 \\ 0 & -\sigma_{1,2} \end{array} \right), \gamma_4 = \left( \begin{array}{cc} 0 & I \\ I & 0 \end{array} \right), \gamma_5 = \left( \begin{array}{cc} 0 & -iI \\ iI & 0 \end{array} \right), \\ \tau &\equiv \frac{-i}{2} [\gamma_4, \gamma_5] = \operatorname{diag}\left( I, -I \right), \tau_\pm = \frac{1 \pm \tau}{2}. \end{split} \tag{3}$$

There are two redundant matrices  $\gamma_4$  and  $\gamma_5$  which anticommutes with other three  $\gamma$  matrices. There exists two kind of chiral transformation  $\psi \to \exp(i\alpha\gamma_4)\psi, \psi \to \exp(i\alpha\gamma_4)\psi$ 

 $\exp(\alpha\gamma_5)\psi$ . The matrices  $\{\gamma_4,\gamma_5,I_4,\tau\}$  generate a U(2) chiral symmetry containing massless spinor fields. This U(2) symmetry is broken down to  $U(1)\times \underline{U}(1)$  by a spinor mass term  $m_e\overline{\psi}\psi$  and parity violating mass  $m_o\overline{\psi}\tau\psi$ , where  $\overline{\psi}\tau\psi$  is a spin density. Here after we take 4-component spinors to study chiral symmetry breaking by  $m^e\overline{\psi}\psi$  in pure QED<sub>3</sub>, where  $\overline{\psi}\tau\psi$  is invariant under chiral transformation. After that the effects of Chern-Simon term will be studied by 2-component spinors.

# 2 Fermion spectral function

The spectral function of 4-component fermion and photon propagator(3) are defined as

$$S_F(x) = S_F^0(x) \exp(F(x)), S_F^0(x) = -(i\gamma \cdot \partial + m) \frac{\exp(-m\sqrt{-x^2})}{4\pi\sqrt{x^2}}.$$
 (4)

$$D_F^0(k) = -i(\frac{g_{\mu\nu} - k_{\mu}k_{\nu}/k^2 - i\theta\epsilon_{\mu\nu\rho}k^{\rho}/k^2}{k^2 - \theta^2 + i\epsilon}) + id\frac{k_{\mu}k_{\nu}}{k^4},\tag{5}$$

where F is an  $O(e^2)$  matrix element  $|T_1|^2$  for the process electron (p+k)  $\rightarrow$ electron (p)+photon (k) as

$$T_1 = -ie \frac{\epsilon_{\mu}(k,\lambda)}{\gamma \cdot (p+k) - m} \gamma^{\mu} \exp(i(p+k) \cdot x) U(p,s), \tag{6}$$

$$\sum_{\lambda,S} T_1 \overline{T_1} = -\frac{\gamma \cdot p + m}{2m} e^2 \left[ \frac{m^2}{(p \cdot k)^2} + \frac{1}{p \cdot k} + \frac{(d-1)}{k^2} \right] - \frac{\gamma \cdot p}{m} \frac{e^2}{4\theta} \frac{m}{p \cdot k}, \quad (7)$$

$$F = \int \frac{d^3k}{(2\pi)^2} \exp(ikx)\theta(k_0)\delta(k^2 - \theta^2) \sum_{\lambda \in S} T_1 \overline{T_1}.$$
 (8)

Here we use the retarded propagator to derive the function F

$$D_{+}(x) = \int \frac{d^{3}k}{i(2\pi)^{2}} \exp(ik \cdot x)\theta(k_{0})\delta(k^{2} - \theta^{2}) = \frac{\exp(-\theta\sqrt{-x^{2}})}{8\pi i\sqrt{-x^{2}}}$$
(9)

The function F is evaluated by  $\alpha$  integration for pure QED<sub>3</sub>(4; 3).

$$F = ie^2 m^2 \int_0^\infty \alpha d\alpha D_F(x + \alpha p) - e^2 \int_0^\infty d\alpha D_F(x + \alpha p) - i(d - 1)e^2 \frac{\partial}{\partial \theta^2} D_F(x, \theta^2)$$
$$= \frac{e^2}{8\pi} \left[ \frac{\exp(-\theta|x|) - \theta|x|E_1(\theta|x|)}{\theta} - \frac{E_1(\theta|x|)}{m} + \frac{(d - 1)\exp(-\theta|x|)}{2\theta} \right]. \tag{10}$$

It is well known that function F has linear and logarithmic infrared divergence with respect to  $\theta$  where  $\theta$  is a bare photon mass. Here we notice the followings. (1)  $\exp(F)$  includes all infrared divergences. (2) quenhed propagator has linear and logarithmic infrared divergences. Linear divergence is absent in a special gauge. In unquenched case  $\theta$  dependence of  $\exp(F)$  is modified by dressed boson spectral function to  $\int ds \rho_{\gamma}(s) \exp(F(x, \sqrt{s}))$ , where  $\pi \rho_{\gamma}(s) = -\Im(s - \Pi(s))^{-1}$ .

## 3 Phase strucutures

For short and long distance we have the approximate form of the function F

$$F_S \sim A - \theta |x| + (D + C|x|) \ln(\theta |x|)) - \frac{(d + 1 - 2\gamma)e^2|x|}{16\pi}, (\theta |x| \ll 1), F_L \sim 0, (1 \ll \theta |x|)..$$

From the above formulae we have

$$\exp(F) = A(\theta|x|)^{D+C|x|}(\theta|x| \ll 1), A = \exp(\frac{e^2(1+d)}{16\pi\theta} + \frac{e^2\gamma}{8\pi m}),$$

$$C = \frac{e^2}{8\pi}, D = \frac{e^2}{8\pi m},$$
(12)

where m is the physical mass and  $\gamma$  is an Euler's constant. Here we see D acts to change the power of |x|. For  $D=1, S_F(0)$  is finite and we have  $\langle \overline{\psi}\psi \rangle \neq 0$  (4; 6). Thus if we require D=1, we obtain  $m=e^2/8\pi$ . In the same way we add the contribution of Chern-Simon term. For simplicity we consider the 2-component spinors in (7). In the condensed phase we have the modified anomalous dimension for  $\theta>0$ 

$$D = \frac{e^2}{8\pi m} + \frac{e^2}{32\pi\theta} = 1, m' = \frac{e^2}{8\pi} / (1 - \frac{e^2}{32\pi\theta}), \Delta m = \frac{e^2}{8\pi} \frac{e^2}{32\pi\theta} / (1 - \frac{e^2}{32\pi\theta}).$$
 (13)

In unquenched case there are parity even and odd spectral function of gauge boson by vacuum polarization(1).In this case we separate D into parity even and odd contribution  $D^e = e^2/8\pi m, D^O = e^2/32\pi\theta$ . For Topologically Massive QCD, $\theta$  is quantized with  $n(0,\pm 1,\pm 2,..)$ . Thus we have  $D=e^2/(8\pi m)+1/8n=1$  for quenched case, and  $D^e=e^2/8\pi m=1, D^O=1/(8n)$  for unquenched case, which leads to  $\langle \overline{\psi}\psi \rangle = \infty$  for any  $n \neq 0$ . In the 4-component spinor free fermion propagator is decomposed into chiral representation

$$S_F(p) = \frac{1}{m_{\epsilon}I + m_O \tau - \gamma \cdot p} = \frac{(\gamma \cdot p + m_+)\tau_+}{p^2 - m_+^2 + i\epsilon} + \frac{(\gamma \cdot p + m_-)\tau_-}{p^2 - m_-^2 + i\epsilon}.$$
 (14)

The difference in two spectral functions is a opposite sign of each Chern-Simon contribution for  $\tau_{\pm}$ .In Toplogically massive gauge theory dynamical mass is parity even and Chern-Simon term shifts mass of different chirality with opposite sign. However shifted mass may not strongly depend on  $\theta$  but  $\langle \overline{\psi}\psi \rangle$  is proportional to  $\theta(12)(5)$ . Our approximation is convenient for unquenched case by the use of gauge boson spectral function(6).

## References

- [1] S.Deser, R.Jackiw & Templeton, Annals of Physics 281, 409-449 (2000).
- [2] C.J.Burden, Nucler Physics B387(1992)419-446, Kei-ichi Kondo, Int. J. Mod. Phys. A11;777-822, 1996.

- [3] R.Jackiw, L.Soloviev, Phys. Rev. 173.5(1968)1485.
- $[4]\,$  Yuichi Hoshino,<br/>in CONTINUOUS ADVANCES IN QCD 2008;361-372.
- [5] Toyoki Matsuyama, Hideko Nagahiro, Mod. Phys. Lett. A15 (2000) 2373-2386.
- $[6] \ C.S. Fisher, Reinhart\ Alkofer, T. Darm, P. Maris, Phys. Rev. \textbf{D70}: 073007, 2004.$