

A Tight Quantitative Version of Arrow’s Impossibility Theorem

Nathan Keller*

Faculty of Mathematics and Computer Science
 Weizmann Institute of Science
 P.O. Box 26, Rehovot 76100, Israel
 nathan.keller@weizmann.ac.il

September 23, 2021

Abstract

The well-known Impossibility Theorem of Arrow asserts that any Generalized Social Welfare Function (GSWF) with at least three alternatives, which satisfies Independence of Irrelevant Alternatives (IIA) and Unanimity and is not a dictatorship, is necessarily non-transitive. In 2002, Kalai asked whether one can obtain the following quantitative version of the theorem: For any $\epsilon > 0$, there exists $\delta = \delta(\epsilon)$ such that if a GSWF on three alternatives satisfies the IIA condition and its probability of non-transitive outcome is at most δ , then the GSWF is at most ϵ -far from being a dictatorship or from breaching the Unanimity condition. In 2009, Mossel proved such quantitative version, with $\delta(\epsilon) = \exp(-C/\epsilon^{21})$, and generalized it to GSWFs with k alternatives, for all $k \geq 3$.

In this paper we show that the quantitative version holds with $\delta(\epsilon) = C \cdot \epsilon^3$, and that this result is tight up to logarithmic factors. Furthermore, our result (like Mossel’s) generalizes to GSWFs with k alternatives. Our proof is based on the works of Kalai and Mossel, but uses also an additional ingredient: a combination of the Bonami-Beckner hypercontractive inequality with a reverse hypercontractive inequality due to Borell, applied to find simultaneously upper bounds and lower bounds on the “noise correlation” between Boolean functions on the discrete cube.

1 Introduction

Consider an election procedure in which a society of n members selects a ranking amongst k alternatives. In the voting process, each member of the society gives a ranking of the alternatives (the ranking is a full linear ordering; that is, indifference between alternatives is not allowed). The set of the rankings given by the individual members is called a *profile*. Given the profile, the ranking of the society is determined according to some function, called a *generalized social welfare function* (GSWF).

The GSWF is a function $F : (S_k)^n \rightarrow \{0, 1\}^{\binom{k}{2}}$, where S_k is the set of linear orderings on k elements. In other words, given the profile consisting of linear orderings supplied by the voters, the function determines the preference of the society amongst each of the $\binom{k}{2}$ pairs of alternatives. If the output of F can be represented as a full linear ordering of the k alternatives, then F is called a *social welfare function* (SWF).

*The author was partially supported by the Adams Fellowship Program of the Israeli Academy of Sciences and Humanities and by the Koshland Center for Basic Research.

Throughout this paper we consider GSWFs satisfying the *Independence of Irrelevant Alternatives* (IIA) condition: For any pair of alternatives A and B , the preference of the entire society between A and B depends only on the preference of each individual voter between A and B . This natural condition on GSWFs can be traced back to Condorcet [6].

The Condorcet’s paradox demonstrates that if the number of alternatives is at least three and the GSWF is based on the majority rule amongst every pair of alternatives, then there exist profiles for which the voting procedure cannot yield a full order relation. That is, there exist alternatives A, B , and C , such that the majority of the society prefers A over B , the majority prefers B over C , and the majority prefers C over A . Such situation is called *non-transitive outcome* of the election.

In his well-known Impossibility theorem [1], Arrow showed that such paradox occurs for any “reasonable” GSWF satisfying the IIA condition:

Theorem 1.1 (Arrow). *Consider a generalized social welfare function F with at least three alternatives. If the following conditions are satisfied:*

- *The IIA condition,*
- *Unanimity — if all the members of the society prefer some alternative A over another alternative B , then A is preferred over B in the outcome of F ,*
- *F is not a dictatorship (that is, the preference of the society is not determined by a single member),*

then the probability of a non-transitive outcome is positive (i.e., there necessarily exists a profile for which the outcome is non-transitive).

Since the existence of profiles leading to a non-transitive outcome has significant implications on voting procedures, an extensive research has been conducted in order to evaluate the probability of non-transitive outcome for various GSWFs. Most of the results in this area are summarized in [9]. In addition to its significance in Social Choice theory, this area of research leads to interesting questions in probabilistic and extremal combinatorics (see [19]).

In 2002, Kalai [14] suggested an analytic approach to this study. He showed that for GSWFs on three alternatives satisfying the IIA condition, the probability of a non-transitive outcome with respect to a uniform distribution of the individual preferences can be computed by a formula related to the Fourier-Walsh expansion of the GSWF. Using this formula he presented a new proof of Arrow’s impossibility theorem under additional assumption of neutrality (i.e., invariance of the GSWF under permutation of the alternatives), and established upper bounds on the probability of non-transitive outcome for specific classes of GSWFs.

While providing an analytic proof to Arrow’s theorem does not seem such an important goal (as there are several simple proofs of it, see [10]), Kalai aimed at establishing a *quantitative* version of the theorem. Such version would show that for any $\epsilon > 0$, there exists $\delta = \delta(\epsilon)$ such that if a GSWF on three alternatives satisfies the IIA condition and its probability of non-transitive outcome is at most δ , then the GSWF is at most ϵ -far from being a dictatorship or from breaching the Unanimity condition. Kalai indeed proved such statement for neutral GSWFs on three alternatives, with $\delta(\epsilon) = C \cdot \epsilon$ for a universal constant C .

Kalai [15] asked whether his techniques can be extended to general GSWFs, and suggested to use the Bonami-Beckner hypercontractive inequality [4, 3] in order to get such an extension. However, Keller [17] showed by an example that a direct extension cannot hold – if there exists

$\delta(\epsilon)$ as above, then it cannot depend linearly on ϵ . Keller asked whether for general GSWFs on three alternatives, a quantitative version holds with $\delta(\epsilon) = C \cdot \epsilon^2$.

A few months ago Mossel [20] succeeded to prove a quantitative version of Arrow’s theorem for general GSWFs on three alternatives. Furthermore, he generalized his result to GSWFs on more than three alternatives, and to more general probability distributions on the individual preferences. Unlike Kalai’s techniques, Mossel’s proof is quite complex. While Kalai’s proof uses only simple analytic tools but no combinatorial tools, Mossel’s proof extends and exploits a combinatorial proof of Arrow’s theorem given by Barbera [2]. Furthermore, it uses “heavier” analytic tools, including a reverse hypercontractive inequality of Borell [5] and a non-linear invariance principle introduced by Mossel et al. [19]. The only drawback in Mossel’s result is the dependence of δ on ϵ : $\delta(\epsilon) = \exp(-C/\epsilon^{21})$ for a universal constant C , which seems far from being optimal. Mossel conjectured that the “correct” dependence of δ on ϵ should be polynomial.¹

In this paper we present a tight quantitative version of Arrow’s theorem for general GSWFs. We show that the dependence of δ on ϵ is indeed polynomial, and compute the exact dependence, up to logarithmic factors.

Before we present our results, we should specify the notion of “the distance of a GSWF on k alternatives satisfying the IIA condition from a dictatorship or from breaching the Unanimity condition”. We consider two different definitions of this notion. In both definitions, the underlying probability measure is the uniform measure on $(S_k)^n$ (the set of all possible profiles).

The first definition measures the distance of the GSWF under examination from the family of GSWFs on k alternatives which satisfy the IIA condition and whose output is *always transitive*. This family was partially characterized by Wilson [22], and fully characterized by Mossel [20]. It essentially consists of combinations of dictatorships with constant functions (see Section 2.3 for the exact characterization).

Definition 1.2. *Denote by $\mathcal{F}_k(n)$ the family of GSWFs on k alternatives which satisfy the IIA condition and whose output is always transitive. For a GSWF F on k alternatives that satisfies the IIA condition, let*

$$D_1(F) = \min_{G \in \mathcal{F}_k(n)} \Pr[F \neq G].$$

We note that this is the definition that was used in [20]. Our main result with respect to this definition is the following:

Theorem 1.3. *There exists an absolute constant C such that for any k and for any GSWF F on k alternatives that satisfies the IIA condition, if the probability of non-transitive outcome in F is at most*

$$\delta(\epsilon) = C \cdot (\epsilon/k^2)^3,$$

then $D_1(F) \leq \epsilon$.

For the second definition, we note that a GSWF F on k alternatives that satisfies the IIA condition actually consists of $\binom{k}{2}$ independent Boolean functions F_{ij} that represent the choice functions amongst the pairs of alternatives (i, j) (for $1 \leq i < j \leq k$). The second definition is given in terms of these functions.

¹We note that Mossel also obtained another variant of his theorem, in which the dependence of δ on ϵ is $\delta(\epsilon) = C\epsilon^3 n^{-3}$, where n is the number of voters, and C is a “decent” constant. As follows from our results presented below, this variant is essentially tight for very small values of ϵ (dependent on n). Moreover, for certain choices of parameters (specifically, “relatively small” n and ϵ very small as a function of n), this result gives a stronger bound than our result, due to the better value of the constant.

Definition 1.4. Denote by $\mathcal{G}_2(n)$ the set of constant functions and dictatorships on two alternatives. For a GSWF F on k alternatives that satisfies the IIA condition, let

$$D_2(F) = \min_{1 \leq i < j \leq k} \min_{G \in \mathcal{G}_2(n)} \Pr[F_{ij} \neq G],$$

where $\{F_{ij}\}_{1 \leq i < j \leq k}$ are as defined above.

Our main result with respect to this definition is the following:

Theorem 1.5. *There exists an absolute constant C such that for any k and for any GSWF F on k alternatives that satisfies the IIA condition, if the probability of non-transitive outcome in F is at most*

$$\delta(\epsilon) = C \cdot \epsilon^{\frac{9(\sqrt{\log_2(1/\epsilon)+1/3})^2}{8 \log_2(1/\epsilon)}}, \tag{1}$$

then $D_2(F) \leq \epsilon$.

Note that for small values of ϵ , the exponent of ϵ in (1) tends to $9/8$.

We show that the dependence of δ on ϵ in Theorems 1.3 and 1.5 is tight, up to logarithmic factors in ϵ . The examples showing the tightness are GSWFs on three alternatives, in which all the three choice functions F_{12} , F_{23} , and F_{13} are monotone threshold functions. In the example of Theorem 1.3, the expectations of the choice functions are $0, 1 - \epsilon, 1 - \epsilon$ (in particular, one of the functions is constant!), and in the example of Theorem 1.5, the expectations are $\epsilon, 1/2, 1 - \epsilon$.

As in the works of Kalai and Mossel, the techniques we use are mainly analytic. Our proof essentially consists of three steps:

1. We consider a GSWF F on three alternatives, and use a modification of Kalai’s formula to express the probability of non-transitive outcome as a linear combination of “noise correlations” between the Boolean functions F_{12} , F_{23} , and F_{13} (see Section 2.2 for the definition of noise correlation).
2. We show that if at least one of the functions F_{12} , F_{23} , and F_{13} is close enough to a constant function, then the Bonami-Beckner hypercontractive inequality [4, 3] and a reverse hypercontractive inequality due to Borell [5] can be applied to obtain simultaneously upper bounds and lower bounds on the noise correlations. Combination of these bounds yields a lower bound on the probability of non-transitive outcome in terms of $D_1(F)$ or $D_2(F)$.
3. To complete the proof, we use the techniques of Mossel to “cover” all the remaining cases (i.e., functions with $D_1(F)$ or $D_2(F)$ greater than a fixed constant, etc.)

We note that since in the case where $D_1(F)$ or $D_2(F)$ is greater than a fixed constant we use Mossel’s result as a black box, the value of the constant we obtain in the dependence of $\delta(\epsilon)$ on ϵ is extremely low, and seems to be very far from optimality. Extension of our techniques to cover all the cases would make the proof free of non-linear invariance arguments, and lead to a “decent” value of the constant. This is one of the main open problems left in our paper.

This paper is organized as follows: In Section 2 we present the tools used in the later sections. In Section 3 we prove our main lemma. We deduce Theorems 1.3 and 1.5 from the main lemma in Section 4. In Section 5 we discuss the tightness of our results. We conclude the paper with questions for further research in Section 6.

2 Preliminaries

In this section we present the tools used in the next sections. First we describe the Fourier-Walsh expansion of functions on the discrete cube. We continue with the noise operator and the hypercontractive inequalities of Bonami-Beckner and of Borell. Finally, we cite the statements from Mossel's proof of the quantitative Arrow theorem [20] that are used as a black box in our proof.

2.1 Fourier-Walsh Expansion of Functions on the Discrete Cube

Throughout the paper we consider the discrete cube $\Omega = \{0, 1\}^n$, endowed with the uniform measure μ . Elements of Ω are represented either by binary vectors of length n , or by subsets of $\{1, 2, \dots, n\}$. Denote the set of all real-valued functions on Ω by X . The inner product of functions $f, g \in X$ is defined as usual as

$$\langle f, g \rangle = \mathbb{E}_\mu[fg] = \frac{1}{2^n} \sum_{x \in \{0, 1\}^n} f(x)g(x).$$

The Rademacher functions $\{r_i\}_{i=1}^n$, defined as $r_i(x_1, \dots, x_n) = 2x_i - 1$, constitute an orthonormal system in X . Moreover, this system can be completed to an orthonormal basis in X by defining

$$r_S = \prod_{i \in S} r_i,$$

for all $S \subset \{1, \dots, n\}$. Every function $f \in X$ can be represented by its Fourier expansion with respect to the system $\{r_S\}_{S \subset \{1, \dots, n\}}$:

$$f = \sum_{S \subset \{1, \dots, n\}} \langle f, r_S \rangle r_S.$$

This representation is called the Fourier-Walsh expansion of f . The coefficients in this expansion are denoted by

$$\hat{f}(S) = \langle f, r_S \rangle.$$

The Fourier-Walsh expansion allows to adapt tools from classical harmonic analysis to the discrete setting, and to use them in the study of Boolean functions. Since the introduction of such analytic methods in the landmark paper of Kahn, Kalai, and Linial [13] in 1988, they were intensively studied, and led to applications in numerous fields, including combinatorics, theoretical computer science, social choice theory, mathematical physics, etc. (see, e.g., the survey [16]).

The most basic analytic tool we use is the Parseval identity, asserting that for all $f, g \in X$,

$$\langle f, g \rangle = \sum_{S \subset \{1, \dots, n\}} \hat{f}(S)\hat{g}(S),$$

and in particular, $\sum_{S \subset \{1, \dots, n\}} \hat{f}(S)^2 = \|f\|_2^2$, for any $f \in X$.

The next simple tool we use is the close relation between the Fourier-Walsh expansions of a function and of the respective dual function.

Definition 2.1. Let $f : \{0, 1\}^n \rightarrow \{0, 1\}$. The dual function of f , which we denote by $\bar{f} : \{0, 1\}^n \rightarrow \{0, 1\}$, is defined by

$$\bar{f}(x_1, x_2, \dots, x_n) = 1 - f(1 - x_1, 1 - x_2, \dots, 1 - x_n).$$

Claim 2.2. Consider the Fourier-Walsh expansions of a Boolean function f and its dual function \bar{f} . For any $S \subset \{1, \dots, n\}$ with $|S| \geq 1$,

$$\widehat{\bar{f}}(S) = (-1)^{|S|-1} \hat{f}(S). \quad (2)$$

The simple proof of the claim is omitted. We use also a variant of the dual function: $f'(x) = 1 - \bar{f}(x)$, defined as

$$f'(x_1, x_2, \dots, x_n) = f(1 - x_1, 1 - x_2, \dots, 1 - x_n).$$

Similarly to Claim 2.2, it is easy to see that for any $|S| \geq 1$,

$$\hat{f}'(S) = (-1)^{|S|} \hat{f}(S). \quad (3)$$

In Kalai's proof of the quantitative Arrow theorem for neutral GSWFs [14], only the most basic analytic tools (like the Parseval identity) were used. Following the proof of Mossel [20], we use also more advanced analytic tools, related to the noise operator presented below.

2.2 The Noise Operator and Hypercontractive Inequalities

The *noise operator*, defined in [3, 4], is a convolution operator that represents the application of the function on a slightly perturbed input.

Definition 2.3. For $x \in \{0, 1\}^n$, the ϵ -noise perturbation of x , denoted by $N_\epsilon(x)$, is a distribution obtained from x by independently keeping each coordinate of x unchanged with probability $1 - \epsilon$, and replacing it by a random value with probability ϵ .

Definition 2.4. Let $f : \{0, 1\}^n \rightarrow \{0, 1\}$. For $0 \leq \epsilon \leq 1$, the noise operator T_ϵ applied to f is defined by

$$T_\epsilon f(x) = \mathbb{E}_{y \sim N_{1-\epsilon}(x)} [f(y)].$$

It is easy to see that the noise operator has a convenient representation in terms of the Fourier-Walsh expansion:

Claim 2.5. Consider a function f on the discrete cube with a Fourier-Walsh expansion $f = \sum_S \hat{f}(S) r_S$. The Fourier-Walsh expansion of $T_\epsilon f$ is given by:

$$T_\epsilon f = \sum_S \epsilon^{|S|} \hat{f}(S) r_S. \quad (4)$$

Since $T_\epsilon f$ represents the application of f on a noisy variant of the input, it makes sense to define the ϵ -noise correlation of two functions f and g as $\langle T_\epsilon f, g \rangle$. Using the Parseval identity, we get an equivalent definition in terms of the Fourier-Walsh expansion (note that the definition is symmetric between f and g):

Definition 2.6. Given two functions $f, g : \{0, 1\}^n \rightarrow \{0, 1\}$, the ϵ -noise correlation of f, g is

$$\langle T_\epsilon f, g \rangle = \sum_{S \subset \{1, 2, \dots, n\}} \epsilon^{|S|} \hat{f}(S) \hat{g}(S). \quad (5)$$

In the proof of Lemma 3.2, we express the probability of non-transitive outcome in a GSWF F on three alternatives in terms of the noise correlations between the Boolean choice functions F_{12}, F_{23} , and F_{13} . Then we obtain upper and lower bounds on the noise correlations using the hypercontractive inequalities presented below.

The first hypercontractive inequality we use is the Bonami-Beckner inequality, discovered independently by Bonami [4] in 1970 and by Beckner [3] in 1975.

Theorem 2.7 (Bonami, Beckner). *Let $f : \{0, 1\}^n \rightarrow \mathbb{R}$, and let $q_1 \geq q_2 \geq 1$. Then*

$$\|T_\epsilon f\|_{q_1} \leq \|f\|_{q_2}, \quad \text{for all } 0 \leq \epsilon \leq \left(\frac{q_2 - 1}{q_1 - 1}\right)^{1/2}.$$

In particular,

$$\|T_\epsilon f\|_2 \leq \|f\|_{1+\epsilon^2}, \quad \text{for all } 0 \leq \epsilon \leq 1.$$

This inequality was first applied in a combinatorial context in [13], and since then it was used in numerous papers in the field. We combine the Bonami-Beckner inequality with the Cauchy-Schwarz inequality to obtain an *upper bound* on the ϵ -noise correlation of Boolean functions. The upper bound is presented here for $\epsilon = 1/3$ since this is the case we use in the proof of Lemma 3.2, but it can be immediately generalized to any $0 \leq \epsilon \leq 1$.

Proposition 2.8. *Let $f, g : \{0, 1\}^n \rightarrow \{0, 1\}$, and denote $\mathbb{E}[f] = p_1$, and $\mathbb{E}[g] = p_2$. Then:*

$$\sum_S \left(\frac{1}{3}\right)^{|S|} \hat{f}(S) \hat{g}(S) \leq \min(p_1^{0.9} p_2^{0.5}, p_1^{0.75} p_2^{0.75}). \quad (6)$$

Proof: By Claim 2.5, the Parseval identity, the Cauchy-Schwarz inequality and the Bonami-Beckner hypercontractive inequality, we get:

$$\sum_S \left(\frac{1}{3}\right)^{|S|} \hat{f}(S) \hat{g}(S) = \langle T_{1/3} f, g \rangle \leq \|T_{1/3} f\|_2 \|g\|_2 \leq \|f\|_{1+1/9} \|g\|_2 = p_1^{0.9} p_2^{0.5}.$$

Similarly,

$$\begin{aligned} \sum_S \left(\frac{1}{3}\right)^{|S|} \hat{f}(S) \hat{g}(S) &= \langle T_{(1/\sqrt{3})} f, T_{(1/\sqrt{3})} g \rangle \leq \|T_{(1/\sqrt{3})} f\|_2 \|T_{(1/\sqrt{3})} g\|_2 \\ &\leq \|f\|_{1+1/3} \|g\|_{1+1/3} = p_1^{0.75} p_2^{0.75}. \end{aligned}$$

This completes the proof. \square

The second hypercontractive inequality we use is a reverse hypercontractive inequality, due to Borell [5]. This inequality asserts that under some conditions, a variant of the Bonami-Beckner inequality holds in the inverse direction.

Theorem 2.9 (Borell). *Let $f : \{0, 1\}^n \rightarrow \mathbb{R}_+$, and let $q_1 \leq q_2 \leq 1$. Then*

$$\|T_\epsilon f\|_{q_1} \geq \|f\|_{q_2}, \quad \text{for all } 0 \leq \epsilon \leq \left(\frac{q_2 - 1}{q_1 - 1}\right)^{1/2}.$$

Although Borell’s result dates back to 1982, it wasn’t used in the research of Boolean functions until recent years. In the last few years, Borell’s inequality was used in several papers [7, 18, 20]; it seems to be a useful tool that has yet to be fully developed.

We use Borell’s inequality to obtain a *lower bound* on the ϵ -noise correlation of Boolean functions through the following corollary, presented in ([18], Corollary 3.5):

Theorem 2.10. *Let $f, g : \{0, 1\}^n \rightarrow \{0, 1\}$. If $\mathbb{E}[f] = p_1$, and $\mathbb{E}[g] = p_2 = p_1^\alpha$ for some $\alpha \geq 0$, then for all $0 < \epsilon < 1$,*

$$\sum_S \epsilon^{|S|} \hat{f}(S) \hat{g}(S) \geq p_1 \cdot p_1^{(\sqrt{\alpha} + \epsilon)^2 / (1 - \epsilon^2)} = p_1 \cdot p_2^{\frac{(\sqrt{\alpha} + \epsilon)^2}{(1 - \epsilon^2)\alpha}}. \quad (7)$$

In the proof of Lemma 3.2, we apply Theorem 2.10 with $\alpha \geq 1$, and write the lower bound in the form $p_1 \cdot p_2^\beta$. We use a simple observation regarding properties of $\beta = \frac{(\sqrt{\alpha} + \epsilon)^2}{(1 - \epsilon^2)\alpha}$:

Observation 2.11. • *As a function of $\alpha = \log_{p_1}(p_2)$, β is monotone decreasing.*

- *For all $\alpha \geq 1$, we have $\beta \leq \frac{1 + \epsilon}{1 - \epsilon}$. When $\alpha \rightarrow \infty$, we have $\beta \rightarrow \frac{1}{1 - \epsilon^2}$.*

Finally, we use the following notation:

Notation 2.12. *We denote by $RHC(p_1, p_2)$ the lower bound obtained in Theorem 2.10 for $\mathbb{E}[f] = p_1$, $\mathbb{E}[g] = p_2$, and $\epsilon = 1/3$. In particular, $RHC(p, p) = p^3$, and*

$$RHC(1/2, p) = \frac{1}{2} \cdot p^{\frac{9(\sqrt{\alpha} + 1/3)^2}{8\alpha}}, \quad (8)$$

where $\alpha = \log_2(1/p)$. For a small value of p , the exponent tends to $9/8$.

2.3 Mossel’s Quantitative Arrow Theorem

In the proofs of Theorems 1.3 and 1.5 we use (as a “black box”) three major components of Mossel’s proof of his quantitative version of Arrow’s theorem [20]. The first is a quantitative Arrow theorem for GSWFs on three alternatives:

Theorem 2.13 ([20], Theorem 8.1). *There exists an absolute constant C such that for any GSWF F on three alternatives that satisfies the IIA condition, if the probability of non-transitive outcome in F is at most*

$$\delta(\epsilon) = \exp(-C/\epsilon^{21}),$$

then $D_1(F) \leq \epsilon$.

We use this theorem only in the case where ϵ is bigger than some fixed constant. Thus, the “bad” dependence of δ on ϵ affects our final result only by a constant factor.

The second component is a generic reduction lemma that allows to leverage results from GSWFs on three alternatives to GSWFs on k alternatives, for all $k \geq 3$. The reduction can be formulated as follows:

Theorem 2.14 ([20], Theorem 9.1 and Remark 9.2). *Suppose that there exists $\delta_0(\epsilon)$ such that for any GSWF F on three alternatives that satisfies the IIA condition, if the probability of non-transitive outcome in F is at most $\delta_0(\epsilon)$, then $D_1(F) \leq \epsilon$. Then we have the following quantitative Arrow theorem for GSWFs on k alternatives:*

For any GSWF F on k alternatives that satisfies the IIA condition, if the probability of non-transitive outcome in F is at most

$$\delta(\epsilon) = \delta_0(\epsilon/k^2),$$

then $D_1(F) \leq \epsilon$.

The third component is a complete characterization of the set $\mathcal{F}_k(n)$ of GSWFs on k alternatives that satisfy the IIA condition and whose output is always transitive. Though we use in our proof only the characterization of $\mathcal{F}_3(n)$, the result is presented here for a general k for the sake of completeness.

Theorem 2.15 ([20], Theorem 1.2). *The class $\mathcal{F}_k(n)$ consists exactly of all GSWFs F satisfying the following: There exists a partition of the set of alternatives into disjoint sets A_1, A_2, \dots, A_r such that:*

- For any profile, F ranks all the alternatives in A_i above all the alternatives in A_j , for all $i < j$.
- For all s such that $|A_s| \geq 3$, the restriction of F to the alternatives in A_s is a dictatorship (i.e., is equal either to the preference order of some voter j or to the reverse of such order).
- For all s such that $|A_s| = 2$, the restriction of F to the alternatives in A_s is an arbitrary non-constant function of the individual preferences between the two alternatives in A_s .

3 Proof of the Main Lemma

In this section we prove our main lemma, asserting that if F is a GSWF on three alternatives that satisfies the IIA condition, and at least one of the Boolean choice functions F_{12}, F_{23} , and F_{31} is “close enough” to a constant function, then the probability of non-transitive outcome can be bounded from below in terms of $D_1(F)$ and $D_2(F)$. Throughout this section we use the following notations:

Notation 3.1. *The Boolean choice functions F_{12}, F_{23} , and F_{31} are denoted by f, g , and h , respectively. This means that the preferences of the voters between alternatives 1 and 2 are denoted by a vector $(x_1, x_2, \dots, x_n) \in \{0, 1\}^n$, where $x_k = 1$ if the k 'th voter prefers alternative 1 over alternative 2, and $x_k = 0$ otherwise. Then, $f(x_1, \dots, x_n) = 1$ if in the output of F , alternative 1 is preferred over alternative 2, and $f(x_1, \dots, x_n) = 0$ otherwise. The functions g and h are defined similarly with respect to the pairs of alternatives (2,3) and (3,1). The expectations of the choice functions are denoted by*

$$\mathbb{E}[f] = p_1, \quad \mathbb{E}[g] = p_2, \quad \mathbb{E}[h] = p_3.$$

For each i , we denote $\bar{p}_i = \min(p_i, 1 - p_i)$, and let

$$D'_2(F) = \min_{1 \leq i \leq 3} \bar{p}_i.$$

Note that $D'_2(F)$ measures the distance of the Boolean choice functions of F from the family of constant functions. Finally, the probability of non-transitive outcome is denoted by $P(F)$.

Now we can formulate our main lemma:

Lemma 3.2. *Let F be a GSWF on three alternatives satisfying the IIA condition. If $D'_2(F) \leq 2^{-500000}$, then*

$$P(F) \geq \frac{1}{10} \cdot \max \left(RHC(D_1(F)/2, D_1(F)/2), RHC(D'_2(F), 1/2) \right),$$

where $D_1(F)$ is as defined in the introduction and $RHC(\cdot, \cdot)$ is as defined in Notation 2.12.

Proof: Our starting point is Kalai's formula [14] for the probability of non-transitive outcome in a GSWF on three alternatives satisfying the IIA condition:

$$\begin{aligned} P(F) = & p_1 p_2 p_3 + (1 - p_1)(1 - p_2)(1 - p_3) + \\ & + \sum_{S \neq \emptyset} \left(-\frac{1}{3}\right)^{|S|} \hat{f}(S) \hat{g}(S) + \sum_{S \neq \emptyset} \left(-\frac{1}{3}\right)^{|S|} \hat{g}(S) \hat{h}(S) + \sum_{S \neq \emptyset} \left(-\frac{1}{3}\right)^{|S|} \hat{h}(S) \hat{f}(S). \end{aligned} \quad (9)$$

The proof is divided into several cases, and in each case we use a different modification of Formula (9). In the following, we assume w.l.o.g. that $\bar{p}_1 \leq \bar{p}_2 \leq \bar{p}_3$. Moreover, we assume w.l.o.g. that $p_1 \leq 1/2$, since otherwise we can replace f, g, h by the dual functions without changing the value of the right hand side of (9).

3.1 Case 1: $p_1, p_2 \leq 1/2$

First, we note that if $p_3 \leq 1/2$, then the assertion follows easily from Kalai's Formula (9). Indeed, since by assumption we have $p_1 < 2^{-500000}$, it follows that $(1 - p_1)(1 - p_2)(1 - p_3) \geq \frac{1}{4}(1 - 2^{-500000})$. On the other hand, by the Parseval identity and the Cauchy-Schwarz inequality, we have

$$\left| \sum_{S \neq \emptyset} \left(-\frac{1}{3}\right)^{|S|} \hat{f}(S) \hat{g}(S) \right| \leq \frac{1}{3} \left(\langle f, g \rangle - p_1 p_2 \right) \leq \frac{1}{3} \cdot \frac{1}{2} \cdot 2^{-250000},$$

and similarly,

$$\left| \sum_{S \neq \emptyset} \left(-\frac{1}{3}\right)^{|S|} \hat{g}(S) \hat{h}(S) \right| \leq \frac{1}{3} \cdot \frac{1}{2} \cdot \frac{1}{2}, \quad \text{and} \quad \left| \sum_{S \neq \emptyset} \left(-\frac{1}{3}\right)^{|S|} \hat{h}(S) \hat{f}(S) \right| \leq \frac{1}{3} \cdot \frac{1}{2} \cdot 2^{-250000}.$$

Thus, by Formula (9),

$$P(F) \geq \frac{1}{4}(1 - 2^{-500000}) - \frac{1}{12} - 2 \cdot \frac{1}{3} \cdot \frac{1}{2} \cdot 2^{-250000} > 1/10.$$

Therefore, we can assume that $p_3 \geq 1/2$. Note that in this case, we have

$$D_1(F) \leq 2(1 - p_3). \quad (10)$$

Indeed, define a GSWF G on three alternatives by the choice functions:

$$f' = G_{12} = 0 \text{ (constant)}, \quad g' = G_{23} = g, \quad h' = G_{31} = 1 \text{ (constant)}.$$

It is clear that $G \in \mathcal{F}_3(n)$, since G always ranks alternative 1 at the bottom and thus its output is always transitive, and

$$\Pr[F \neq G] \leq \Pr[f \neq f'] + \Pr[g \neq g'] + \Pr[h \neq h'] \leq p_1 + (1 - p_3) \leq 2(1 - p_3).$$

Therefore, $D_1(F) \leq D(F, G) \leq 2(1 - p_3)$. Also, by the definition,

$$D_2'(F) = p_1. \quad (11)$$

We modify Formula (9) using the following identities:

$$\sum_{S \neq \emptyset} \left(-\frac{1}{3}\right)^{|S|} \hat{f}(S) \hat{g}(S) = \sum_{S \neq \emptyset} \left(\frac{1}{3}\right)^{|S|} \hat{f}'(S) \hat{g}(S) = \langle T_{1/3} f', g \rangle - p_1 p_2. \quad (12)$$

$$\begin{aligned} \sum_{S \neq \emptyset} \left(-\frac{1}{3}\right)^{|S|} \hat{g}(S) \hat{h}(S) &= \sum_{S \neq \emptyset} \left(-\frac{1}{3}\right)^{|S|} \widehat{\hat{g}}(S) \widehat{\hat{h}}(S) = \sum_{S \neq \emptyset} \left(\frac{1}{3}\right)^{|S|} \widehat{\hat{g}}(S) \widehat{1-h}(S) \\ &= \langle T_{1/3} \bar{g}, 1-h \rangle - (1-p_2)(1-p_3). \end{aligned} \quad (13)$$

$$\begin{aligned} \sum_{S \neq \emptyset} \left(-\frac{1}{3}\right)^{|S|} \hat{h}(S) \hat{f}(S) &= \langle T_{1/3} f', h \rangle - p_1 p_3 = \langle T_{1/3} f', 1 \rangle - \langle T_{1/3} f', 1-h \rangle - p_1 p_3 \\ &= p_1 - p_1 p_3 - \langle T_{1/3} f', 1-h \rangle. \end{aligned} \quad (14)$$

All three identities follow immediately from basic properties of the dual function (e.g., Equations (2) and (3)) and the Parseval identity. Substituting Equations (12), (13), and (14) into Formula (9), we get:

$$\begin{aligned} P(F) &= p_1 p_2 p_3 + (1-p_1)(1-p_2)(1-p_3) + \\ &\quad + \langle T_{1/3} f', g \rangle - p_1 p_2 + \langle T_{1/3} \bar{g}, 1-h \rangle - (1-p_2)(1-p_3) + p_1 - p_1 p_3 - \langle T_{1/3} f', 1-h \rangle = \\ &= \langle T_{1/3} f', g \rangle + \langle T_{1/3} \bar{g}, 1-h \rangle - \langle T_{1/3} f', 1-h \rangle. \end{aligned} \quad (15)$$

Equation (15) expresses $P(F)$ as a linear combination of “noise correlations” between the functions f, g, h , which are obviously nonnegative. Thus, if we obtain a lower bound on the noise correlations that appear in Equation (15) with a ‘+’ sign, and an upper bound on the correlation that appears with a ‘-’ sign, we will get a lower bound on $P(f)$. We shall obtain these bounds using the Bonami-Beckner hypercontractive inequality and Borell’s reverse hypercontractive inequality. We subdivide our case into two sub-cases.

3.1.1 Case 1a: $1 - p_3 < 1/32$.

We bound $\langle T_{1/3} f', 1-h \rangle$ from above using the Bonami-Beckner hypercontractive inequality. By Proposition 2.8, we get:

$$\langle T_{1/3} f', 1-h \rangle \leq p_1^{0.75} (1-p_3)^{0.75} \leq (1-p_3)^{1.5}. \quad (16)$$

We bound $\langle T_{1/3} \bar{g}, 1-h \rangle$ from below using Borell’s reverse hypercontractive inequality. By Theorem 2.10, we have

$$\langle T_{1/3} \bar{g}, 1-h \rangle \geq RHC(1-p_2, 1-p_3).$$

In order to estimate $RHC(1-p_2, 1-p_3)$, we write it in the form $(1-p_2) \cdot (1-p_3)^{\beta(\alpha)}$, where $\alpha = \log_{1-p_2}(1-p_3)$. By Observation 2.11, $\beta(\alpha)$ is a monotone decreasing function of α . Since by assumption, $1-p_2 > 31/32$ and $1-p_3 < 1/32$, we have $\alpha \geq \log_{31/32}(1/32) = 109.16$.

Substituting the value $\alpha = 109.16$ into the definition of $\beta(\alpha)$ and using the monotonicity of $\beta(\alpha)$, we get $\beta \leq 1.198$, and thus,

$$\langle T_{1/3}\bar{g}, 1 - h \rangle \geq (1 - p_2)(1 - p_3)^{1.198} \geq \frac{31}{32}(1 - p_3)^{1.198}. \quad (17)$$

Combining Inequalities (16) and (17), we get

$$\begin{aligned} \langle T_{1/3}f', 1 - h \rangle &\leq (1 - p_3)^{1.5} = \left(\frac{32}{31}(1 - p_3)^{0.302}\right) \left(\frac{31}{32}(1 - p_3)^{1.198}\right) \\ &\leq \left(\frac{32}{31}\left(\frac{1}{32}\right)^{0.302}\right) \langle T_{1/3}\bar{g}, 1 - h \rangle \leq 0.37 \langle T_{1/3}\bar{g}, 1 - h \rangle. \end{aligned} \quad (18)$$

Finally, substituting into Equation (15) we get:

$$\begin{aligned} P(F) &= \langle T_{1/3}f', g \rangle + \langle T_{1/3}\bar{g}, 1 - h \rangle - \langle T_{1/3}f', 1 - h \rangle \\ &\geq \langle T_{1/3}\bar{g}, 1 - h \rangle - \langle T_{1/3}f', 1 - h \rangle \geq 0.63 \langle T_{1/3}\bar{g}, 1 - h \rangle \\ &\geq 0.63 RHC(1 - p_2, 1 - p_3) \geq 0.63 \max(RHC(1 - p_3, 1 - p_3), RHC(1/2, p_1)), \end{aligned} \quad (19)$$

where the last inequality holds since $1 - p_2 \geq 1/2 \geq 1 - p_3 \geq p_1$ and since $RHC(\cdot, \cdot)$ is clearly non-decreasing in its arguments. The assertion of the lemma follows now from Inequalities (10) and (11).

3.1.2 Case 1b: $1 - p_3 \geq 1/32$.

As in the previous case, we bound $\langle T_{1/3}f', 1 - h \rangle$ from above using the Bonami-Beckner hypercontractive inequality. By Proposition 2.8, we get:

$$\langle T_{1/3}f', 1 - h \rangle \leq p_1^{0.9}(1 - p_3)^{0.5} \leq (1 - p_3)^{4.1}, \quad (20)$$

where the last inequality follows since by assumption $p_1 \leq 2^{-500000}$, and in particular, $p_1 \leq (1 - p_3)^4$. In order to bound $\langle T_{1/3}\bar{g}, 1 - h \rangle$ from below we use the reverse hypercontractive inequality. By Theorem 2.10, we have

$$\langle T_{1/3}\bar{g}, 1 - h \rangle \geq RHC(1 - p_2, 1 - p_3).$$

As in the previous case, we write $RHC(1 - p_2, 1 - p_3)$ in the form $(1 - p_2) \cdot (1 - p_3)^{\beta(\alpha)}$. Since by Observation 2.11, for any $\alpha \geq 1$, we have $\beta(\alpha) \leq \frac{1+\epsilon}{1-\epsilon} = 2$, we get

$$\langle T_{1/3}\bar{g}, 1 - h \rangle \geq (1 - p_2)(1 - p_3)^2 \geq 0.5(1 - p_3)^2. \quad (21)$$

Combination of Equations (20) and (21) yields

$$\begin{aligned} \langle T_{1/3}f', 1 - h \rangle &\leq (1 - p_3)^{4.1} = \left(2(1 - p_3)^{2.1}\right) \left(0.5(1 - p_3)^2\right) \\ &\leq 2(1 - p_3)^{2.1} \langle T_{1/3}\bar{g}, 1 - h \rangle \leq 0.5 \langle T_{1/3}\bar{g}, 1 - h \rangle, \end{aligned}$$

where the last inequality follows since $1 - p_3 \leq 1/2$. Finally,

$$\begin{aligned} P(F) &= \langle T_{1/3}f', g \rangle + \langle T_{1/3}\bar{g}, 1 - h \rangle - \langle T_{1/3}f', 1 - h \rangle \\ &\geq \langle T_{1/3}\bar{g}, 1 - h \rangle - \langle T_{1/3}f', 1 - h \rangle \geq 0.5 \langle T_{1/3}\bar{g}, 1 - h \rangle \geq 0.5 RHC(1 - p_2, 1 - p_3) \\ &\geq 0.5 \max(RHC(1 - p_3, 1 - p_3), RHC(1/2, p_1)), \end{aligned} \quad (22)$$

as asserted. This completes the proof of Case 1.

3.2 Case 2: $p_1 \leq 1/2$ and $p_2 \geq 1/2$

In this case, we have

$$D_1(F) \leq 2(1 - p_2), \quad (23)$$

since defining a GSWF G' on three alternatives by the choice functions:

$$f'' = G'_{12} = 0 \text{ (constant)}, \quad g'' = G'_{23} = 1 \text{ (constant)}, \quad h'' = G'_{31} = h,$$

we get $G' \in \mathcal{F}_3(n)$, and $D(F, G') \leq 2(1 - p_2)$. Also, it is clear that like in Case 1,

$$D_2(F) \leq D(f, \text{const}) = p_1. \quad (24)$$

This time we use a slightly different modification of Kalai's formula. Specifically, we interchange the roles of g and h in Equations (12), (13), and (14), and get the following modification of Equation (15):

$$P(F) = \langle T_{1/3}f', h \rangle + \langle T_{1/3}(1 - g), \bar{h} \rangle - \langle T_{1/3}f', 1 - g \rangle. \quad (25)$$

We subdivide this case into several sub-cases.

3.2.1 Case 2a: $(1 - p_2) \leq p_1^{0.45412}$.

By Proposition 2.8, we get:

$$\langle T_{1/3}f', 1 - g \rangle \leq p_1^{0.9}(1 - p_2)^{0.5} \leq p_1^{0.9}p_1^{0.22706} = p_1^{1.12706}. \quad (26)$$

On the other hand, by Theorem 2.10, we have

$$\langle T_{1/3}f', h \rangle + \langle T_{1/3}(1 - g), \bar{h} \rangle \geq RHC(p_3, p_1) + RHC(1 - p_3, 1 - p_2).$$

Since $p_1 \leq 1 - p_2$ and either p_3 or $1 - p_3$ is not less than $1/2$, we get

$$\langle T_{1/3}f', h \rangle + \langle T_{1/3}(1 - g), \bar{h} \rangle \geq RHC(1/2, p_1) \geq 0.5p_1^{1.12606}, \quad (27)$$

where the last inequality follows from Observation 2.11 since $p_1 \leq 2^{-500000}$. Combining Inequalities (26) and (27) we get

$$\langle T_{1/3}f', 1 - g \rangle \leq p_1^{1.12706} = (2p_1^{0.001})(0.5p_1^{1.12606}) \leq 0.5(\langle T_{1/3}f', h \rangle + \langle T_{1/3}(1 - g), \bar{h} \rangle).$$

Finally,

$$\begin{aligned} P(F) &= \langle T_{1/3}f', h \rangle + \langle T_{1/3}(1 - g), \bar{h} \rangle - \langle T_{1/3}f', 1 - g \rangle \\ &\geq 0.5(\langle T_{1/3}f', h \rangle + \langle T_{1/3}(1 - g), \bar{h} \rangle) \geq 0.5\left(RHC(p_3, p_1) + RHC(1 - p_3, 1 - p_2)\right) \\ &\geq 0.5 \max\left(RHC(1 - p_2, 1 - p_2), RHC(1/2, p_1)\right), \end{aligned} \quad (28)$$

where the last inequality holds since $1 - p_3 \geq 1 - p_2$. The assertion of the lemma follows now from Inequalities (23) and (24).

3.2.2 Case 2b: $(1 - p_2) > p_1^{0.45412}$ and $\bar{p}_3 \geq p_1^{0.2002}$.

The upper bound in this case is the same as in Case 2a:

$$\langle T_{1/3}f', 1 - g \rangle \leq p_1^{0.9}(1 - p_2)^{0.5}. \quad (29)$$

For the lower bound, we use the reverse hypercontractive inequality for the term $\langle T_{1/3}(1 - g), \bar{h} \rangle$, and get

$$\langle T_{1/3}(1 - g), \bar{h} \rangle \geq RHC(1 - p_3, 1 - p_2) \geq (1 - p_3)(1 - p_2)^2 \geq p_1^{0.2002}(1 - p_2)^2, \quad (30)$$

where the second inequality follows from Observation 2.11, and the third inequality follows from the assumption $\bar{p}_3 \geq p_1^{0.2002}$. Combination of Inequality (29) with Inequality (30) yields:

$$\begin{aligned} \langle T_{1/3}f', 1 - g \rangle &\leq p_1^{0.9}p_2^{0.5} = \left(p_1^{0.6998}p_2^{-1.5}\right) \left(p_1^{0.2002}(1 - p_2)^2\right) \\ &\leq p_1^{0.01862} \langle T_{1/3}(1 - g), \bar{h} \rangle \leq 0.5 \langle T_{1/3}(1 - g), \bar{h} \rangle, \end{aligned}$$

where the second to last inequality follows from the assumption $1 - p_2 \geq p_1^{0.45412}$, and the last inequality follows since $p_1 \leq 2^{-500000}$. Finally, if $p_3 \geq 1/2$ then $\langle T_{1/3}f', h \rangle \geq RHC(1/2, p_1)$, and otherwise, $\langle T_{1/3}(1 - g), \bar{h} \rangle \geq RHC(1/2, p_1)$. In both cases,

$$\begin{aligned} P(F) &= \langle T_{1/3}f', h \rangle + (\langle T_{1/3}(1 - g), \bar{h} \rangle - \langle T_{1/3}f', 1 - g \rangle) \\ &\geq \langle T_{1/3}f', h \rangle + 0.5 \langle T_{1/3}(1 - g), \bar{h} \rangle \\ &\geq 0.5 \max \left(RHC(1 - p_2, 1 - p_2), RHC(1/2, p_1) \right), \end{aligned} \quad (31)$$

and the assertion follows.

3.2.3 Case 2c: $p_3 \leq p_1^{0.2002}$.

In this case we use another modification of Kalai's formula (9), resulting from the following modification of Equation (25):

$$\begin{aligned} P(F) &= \langle T_{1/3}f', h \rangle + \langle T_{1/3}(1 - g), \bar{h} \rangle - \langle T_{1/3}f', 1 - g \rangle \\ &= \langle T_{1/3}f', h \rangle + (\langle T_{1/3}(1 - g), 1 \rangle - \langle T_{1/3}(1 - g), 1 - \bar{h} \rangle) - \langle T_{1/3}f', 1 - g \rangle \\ &= (1 - p_2) + \langle T_{1/3}f', h \rangle - \langle T_{1/3}(1 - g), 1 - \bar{h} \rangle - \langle T_{1/3}f', 1 - g \rangle. \end{aligned} \quad (32)$$

By Proposition 2.8,

$$\langle T_{1/3}(1 - g), 1 - \bar{h} \rangle \leq (1 - p_2)^{0.9}p_3^{0.5} \leq (1 - p_2)^{0.9}p_1^{0.1001} \leq (1 - p_2)^{1.0001}.$$

Similarly,

$$\langle T_{1/3}f', 1 - g \rangle \leq p_1^{0.9}(1 - p_2)^{0.5} \leq (1 - p_2)^{1.4}.$$

Hence,

$$\langle T_{1/3}(1 - g), 1 - \bar{h} \rangle + \langle T_{1/3}f', 1 - g \rangle \leq (1 - p_2)^{1.0001} + (1 - p_2)^{1.4} \leq 2(1 - p_2)^{1.0001} \leq 0.5(1 - p_2),$$

where the last inequality follows since

$$(1 - p_2)^{0.0001} \leq p_3^{0.0001} \leq p_1^{0.2002 \cdot 0.0001} < 1/2.$$

Finally, by Equation (32),

$$\begin{aligned} P(F) &\geq (1 - p_2) - \langle T_{1/3}(1 - g), 1 - \bar{h} \rangle - \langle T_{1/3}f', 1 - g \rangle \\ &\geq 0.5(1 - p_2) \geq 0.5 \max \left(RHC(1 - p_2, 1 - p_2), RHC(1/2, p_1) \right), \end{aligned} \quad (33)$$

as asserted.

3.2.4 Case 2d: $1 - p_3 \leq p_1^{0.2002}$

We use yet another modification of Equation (25):

$$\begin{aligned}
P(F) &= \langle T_{1/3}f', h \rangle + \langle T_{1/3}(1-g), \bar{h} \rangle - \langle T_{1/3}f', 1-g \rangle \\
&= (\langle T_{1/3}f', 1 \rangle - \langle T_{1/3}f', 1-h \rangle) + \langle T_{1/3}(1-g), \bar{h} \rangle - \langle T_{1/3}f', 1-g \rangle \\
&= p_1 - \langle T_{1/3}f', 1-h \rangle + \langle T_{1/3}(1-g), \bar{h} \rangle - \langle T_{1/3}f', 1-g \rangle.
\end{aligned} \tag{34}$$

Similarly to Case 2c, we have

$$\langle T_{1/3}f', 1-h \rangle \leq p_1^{0.9}(1-p_3)^{0.5} \leq p_1^{1.0001},$$

and $\langle T_{1/3}f', 1-g \rangle \leq p_1^{1.4}$, and thus,

$$\langle T_{1/3}f', 1-h \rangle + \langle T_{1/3}f', 1-g \rangle \leq p_1^{1.0001} + p_1^{1.4} \leq 0.5p_1.$$

Therefore,

$$\begin{aligned}
P(F) &= p_1 - \langle T_{1/3}f', 1-h \rangle + \langle T_{1/3}(1-g), \bar{h} \rangle - \langle T_{1/3}f', 1-g \rangle \\
&\geq 0.5p_1 + \langle T_{1/3}(1-g), \bar{h} \rangle \geq 0.5 \max\left(RHC(1-p_2, 1-p_2), RHC(1/2, p_1)\right).
\end{aligned} \tag{35}$$

This completes the proof of Lemma 3.2. \square

4 Proof of Theorems 1.3 and 1.5

In this section we present the proofs of Theorems 1.3 and 1.5. The proofs are based on Lemma 3.2, but also rely heavily on several components of Mossel's proof of his quantitative version of Arrow's theorem cited in Section 2.3. The general structure of both proofs is as follows:

1. We consider first GSWFs on three alternatives, and examine several cases:
 - (a) If $D_1(F)$ (resp., $D_2(F)$) is greater than a fixed constant, we deduce the assertion from Theorem 2.13.
 - (b) If F is close to a GSWF that always ranks one of the candidates at the top/bottom (resp., if at least one of the Boolean choice functions of F is close to a constant function), we deduce the assertion from Lemma 3.2.
 - (c) If F (resp., one of the Boolean choice functions of F) is close to a dictatorship of the i 'th voter, we split F into six GSWFs $\{F^\sigma\}_{\sigma \in S_3}$ according to the preferences of the i 'th voter. We further subdivide this case into two cases:
 - If for all $\sigma \in S_3$, $D_1(F^\sigma)$ (resp., $D'_2(F^\sigma)$) is small, we get a contradiction (resp., show directly that $P(F)$ cannot be small).
 - If there exists $\sigma_0 \in S_3$ such that $D_1(F^{\sigma_0})$ (resp., $D'_2(F^{\sigma_0})$) is not small, we deduce the assertion by applying Lemma 3.2 to the GSWF F^{σ_0} .
2. We leverage the result to GSWFs on k alternatives, for all $k \geq 3$. In the proof of Theorem 1.3 this requires the reduction technique of Theorem 2.14, and in the proof of Theorem 1.5, the generalization is immediate.

Since the proofs differ in many of the details, we present them separately. Throughout this section, we use the notations defined at the beginning of Section 3.

4.1 Proof of Theorem 1.3

Theorem 4.1. *There exists an absolute constant C such that for any GSWF F on three alternatives that satisfies the IIA condition, if the probability of non-transitive outcome in F is at most*

$$\delta(\epsilon) = \min\left(C, \frac{1}{50000} \cdot \epsilon^3\right),$$

then $D_1(F) \leq \epsilon$.

Proof: It is clearly sufficient to prove that for any $\epsilon > 0$, if $D_1(F) = \epsilon$, then $P(F) \geq \min\left(C, \frac{1}{50000} \cdot \epsilon^3\right)$, for a universal constant C . We shall prove this for

$$C = \exp\left(-\frac{C'}{(2^{-500003})^{21}}\right), \quad (36)$$

where C' is the constant in Mossel's Theorem 2.13.

Let F be a GSWF on three alternatives satisfying the IIA conditions, and denote the choice functions of F by f, g , and h , as in the proof of Lemma 3.2. If $D_1(F) \geq 2^{-500003}$, then by Theorem 2.13, $P(F) \geq C$. Thus, we may assume that $D_1(F) < 2^{-500003}$.

Let $G \in \mathcal{F}_3(n)$ satisfy $\Pr[F \neq G] = D_1(F)$ (such element exists by the definition of the distance $D_1(F)$). Denote the Boolean choice functions of G by f', g' , and h' . By Theorem 2.15, G either always ranks one alternative at the top/bottom or is a dictatorship. If G always ranks one alternative at the top/bottom, then at least two of the functions f', g' , and h' are constant. Assume w.l.o.g. that f' and g' are constant. Since

$$D_1(F) = \Pr[F \neq G] \geq \max(\Pr[f \neq f'], \Pr[g \neq g'], \Pr[h \neq h']),$$

it follows that either $\mathbb{E}[f] \leq D_1(F)$ or $\mathbb{E}[f] \geq 1 - D_1(F)$, and similarly for g . This implies that $D'_2(F) \leq D_1(F) < 2^{-500003}$, and thus we can apply Lemma 3.2 to F and get

$$P(F) \geq \frac{1}{10} \cdot RHC(D_1(F)/2, D_1(F)/2) \geq \frac{1}{10} \cdot (D_1(F)/2)^3 > \frac{1}{50000} \cdot D_1(F)^3,$$

as asserted. Thus, we may assume that G is a dictatorship.

The following part of the proof is similar to the proof of Theorem 7.1 in [20]. W.l.o.g., we assume that the output of G is determined by the first voter. We “split” the choice functions according to the first voter. Let

$$f^0(x_2, x_3, \dots, x_n) = f(0, x_2, x_3, \dots, x_n), \quad f^1(x_2, x_3, \dots, x_n) = f(1, x_2, x_3, \dots, x_n),$$

and similarly for g and h . Furthermore, for any profile $(\sigma_1, \sigma_2, \dots, \sigma_n) \in S_3^n$, denote

$$F^{\sigma_1}(\sigma_2, \sigma_3, \dots, \sigma_n) = F(\sigma_1, \sigma_2, \sigma_3, \dots, \sigma_n),$$

and similarly for G . The Boolean choice functions of F^σ are f^{a_1}, g^{a_2} , and h^{a_3} , where $(a_1, a_2, a_3) \in \{0, 1\}^3$ represents the preference σ of the first voter (note that only six of the eight possible combinations of (a_1, a_2, a_3) represent elements of S_3). Denote by $\bar{f}^{a_1}, \bar{g}^{a_2}, \bar{h}^{a_3}$ the choice functions of G^σ . Since G is a dictatorship of the first voter, the functions $\bar{f}^{a_1}, \bar{g}^{a_2}$, and \bar{h}^{a_3} are constant. Clearly, we have

$$D_1(F) = \Pr[F \neq G] = \frac{1}{6} \sum_{\sigma \in S_3} \Pr[F^\sigma \neq G^\sigma], \quad (37)$$

and thus, for all $\sigma \in S_3$,

$$\Pr[F^\sigma \neq G^\sigma] \leq 6D_1(F).$$

Since

$$\Pr[F^\sigma \neq G^\sigma] \geq \max(\Pr[f^{a_1} \neq \bar{f}^{a_1}], \Pr[g^{a_2} \neq \bar{g}^{a_2}], \Pr[h^{a_3} \neq \bar{h}^{a_3}]),$$

and since G^σ is constant, this implies that

$$\mathbb{E}[f^{a_1}] \leq 6D_1(F) \quad \text{or} \quad \mathbb{E}[f^{a_1}] \geq 1 - 6D_1(F), \quad (38)$$

and similarly for g^{a_2} and h^{a_3} .

The rest of the proof is divided into two cases:

- **Case A:** For all $\sigma \in S_3$ we have $D_1(F^\sigma) \leq D_1(F)/4$.
- **Case B:** There exists $\sigma_0 \in S_3$ such that $D_1(F^{\sigma_0}) > D_1(F)/4$.

We first show that Case A leads to a contradiction by constructing a GSWF $G' \in \mathcal{F}_3(n)$ such that $\Pr[F \neq G'] < D_1(F)$. Then we show that in Case B, the assertion of the theorem follows by applying Lemma 3.2 to the function F^{σ_0} .

Case A: Consider a GSWF G' whose choice functions f'' , g'' , and h'' are defined as follows: For $a_1 \in \{0, 1\}$,

$$f''(a_1, x_2, \dots, x_n) = \begin{cases} 1 \text{ (constant)}, & \mathbb{E}[f^{a_1}] \geq 1 - D_1(F)/4, \\ 0 \text{ (constant)}, & \mathbb{E}[f^{a_1}] \leq D_1(F)/4, \\ f^{a_1}, & \text{otherwise,} \end{cases}$$

and similarly for g' and h' . We claim that the output of G' is always transitive, and thus $G' \in \mathcal{F}_3(n)$. Indeed, by assumption, for any $\sigma \in S_3$, there exists $\bar{G}^\sigma \in \mathcal{F}_3(n-1)$ such that $\Pr[F^\sigma \neq \bar{G}^\sigma] \leq D_1(F)/4$. The GSWF \bar{G}^σ cannot be a dictatorship since by Equation (38), the choice functions f^{a_1} , g^{a_2} , and h^{a_3} of F^σ satisfy

$$\mathbb{E}[f^{a_1}] \leq 6D_1(F) \quad \text{or} \quad \mathbb{E}[f^{a_1}] \geq 1 - 6D_1(F),$$

and thus, for any dictatorship H ,

$$\Pr[F^\sigma \neq H] \geq 1/2 - 6D_1(F) > 1/2 - 2^{500000}.$$

Therefore, \bar{G}^σ always ranks one alternative at the top/bottom. Denote the choice functions of \bar{G}^σ by \tilde{f} , \tilde{g} , and \tilde{h} , and assume w.l.o.g. that \bar{G}^σ always ranks alternative 1 at the top, and thus $\tilde{f} = 1$ and $\tilde{h} = 0$. Since

$$D_1(F)/4 \geq \Pr[F^\sigma \neq \bar{G}^\sigma] \geq \max(\Pr[f^{a_1} \neq \tilde{f}], \Pr[g^{a_2} \neq \tilde{g}], \Pr[h^{a_3} \neq \tilde{h}]),$$

it follows that

$$\mathbb{E}[f^{a_1}] \geq 1 - D_1(F)/4, \quad \text{and} \quad \mathbb{E}[h^{a_3}] \leq D_1(F)/4.$$

Hence, by the definition of G' , its choice functions satisfy $f'' = 1$ and $h'' = 0$, which means that G' always ranks alternative 1 at the top, and is thus always transitive.

Therefore, $G' \in \mathcal{F}_3(n)$, and on the other hand, we have

$$\Pr[F \neq G'] \leq \Pr[f \neq f''] + \Pr[g \neq g''] + \Pr[h \neq h''] \leq 3 \cdot D_1(F)/4 < D_1(F),$$

contradicting the definition of $D_1(F)$.

Case B: Let $\sigma_0 \in S_3$ be such that $D_1(F^{\sigma_0}) > D_1(F)/4$. By Equation (38), the choice functions $f^{a_1}, g^{a_2}, h^{a_3}$ of F^{σ_0} satisfy

$$\mathbb{E}[f^{a_1}] \leq 6D_1(F) \quad \text{or} \quad \mathbb{E}[f^{a_1}] \geq 1 - 6D_1(F),$$

and thus (in the notation of Lemma 3.2), $D'_2(F^{\sigma_0}) \leq 6D_1(F) < 2^{-500000}$. Hence, we can apply Lemma 3.2 to the GSWF G^{σ_0} , and get

$$P(F^{\sigma_0}) \geq \frac{1}{10} \cdot RHC(D_1(F^{\sigma_0})/2, D_1(F^{\sigma_0})/2) \geq \frac{1}{10}(D_1(F)/8)^3 = \frac{1}{5120} \cdot D_1(F)^3.$$

Finally,

$$P(F) = \frac{1}{6} \sum_{\sigma \in S_3} P(F^\sigma) \geq \frac{1}{6} \cdot P(F^{\sigma_0}) > \frac{1}{50000} D_1(F)^3.$$

This completes the proof of the theorem. \square

Theorem 1.3 follows immediately from Theorem 4.1 using Theorem 2.14 (the generic reduction lemma of Mossel).

4.2 Proof of Theorem 1.5

Theorem 4.2. *There exists an absolute constant C such that for any GSWF F on three alternatives that satisfies the IIA condition, if the probability of non-transitive outcome in F is at most*

$$\delta(\epsilon) = \min \left(C, \frac{1}{10000} \cdot \epsilon^{\frac{9(\sqrt{\log_2(1/\epsilon)}+1/3)^2}{8 \log_2(1/\epsilon)}} \right), \quad (39)$$

then $D_2(F) \leq \epsilon$.

Proof: By Equation (8), it is sufficient to prove that for any $\epsilon > 0$, if $D_2(F) = \epsilon$, then

$$P(F) \geq \min \left(C, \frac{1}{5000} \cdot RHC(1/2, \epsilon) \right),$$

for a universal constant C . We shall prove this for

$$C = \exp \left(-\frac{C'}{(2^{-500003})^{21}} \right), \quad (40)$$

where C' is the constant in Mossel's Theorem 2.13. Let F be a GSWF on three alternatives satisfying the IIA conditions, and denote the choice functions of F by f, g , and h , as in the proof of Lemma 3.2.

First we consider the case $D_2(F) \geq 2^{-500003}$. We show that in general, $D_1(F) \geq D_2(F)$, and thus in this case we have $D_1(F) \geq D_2(F) \geq 2^{-500003}$, which by Theorem 2.13 implies that $P(F) \geq C$. Let $G \in \mathcal{F}_3(n)$ satisfy $\Pr[F \neq G] = D_1(F)$, and denote the Boolean choice functions of G by f', g' , and h' . Clearly,

$$D_1(F) = \Pr[F \neq G] \geq \max(\Pr[f \neq f'], \Pr[g \neq g'], \Pr[h \neq h']). \quad (41)$$

By Theorem 2.15, G either always ranks one alternative at the top/bottom or is a dictatorship. In the first case, at least two of the functions $f', g',$ and h' are constant, and thus Equation (41) implies that at least two of the functions $f, g,$ and h are at most $D_1(F)$ -far from a constant function. In the latter case, the functions $f', g',$ and h' are dictatorships, and thus Equation (41) implies that $f, g,$ and h are at most $D_1(F)$ -far from a dictatorship. Hence, in both cases,

$$D_2(F) = \min_{1 \leq i < j \leq 3} \min_{G \in \mathcal{G}_2(n)} \Pr[F_{ij} \neq G] \leq D_1(F),$$

as asserted.

Now we consider the case $D_2(F) < 2^{-500003}$. Assume w.l.o.g. that the minimal distance $\min_{G \in \mathcal{G}_2(n)} \Pr[F_{ij} \neq G]$ is obtained by the choice function f , and let $\tilde{f} \in \mathcal{G}_2(n)$ satisfy $\Pr[f \neq \tilde{f}] = D_2(F)$. If \tilde{f} is a constant function, then in the notations of Lemma 3.2, this implies that $D'_2(F) = D_2(F) < 2^{-500003}$, and thus we can apply Lemma 3.2 to F and get

$$P(F) \geq \frac{1}{10} \cdot RHC(1/2, D'_2(F)) > \frac{1}{5000} \cdot RHC(1/2, D_2(F)),$$

as asserted. Thus, we may assume that \tilde{f} is a dictatorship.

Assume w.l.o.g. that \tilde{f} is a dictatorship of the first voter. Define the functions $F^\sigma, f^0, f^1, g^0, g^1, h^0,$ and h^1 as in the proof of Theorem 4.1, and let

$$\tilde{f}^0(x_2, x_3, \dots, x_n) = \tilde{f}(0, x_2, x_3, \dots, x_n), \quad \text{and} \quad \tilde{f}^1(x_2, x_3, \dots, x_n) = \tilde{f}(1, x_2, x_3, \dots, x_n).$$

Clearly, we have

$$D_2(F) = \Pr[f \neq \tilde{f}] = \frac{1}{2}(\Pr[f^0 \neq \tilde{f}^0] + \Pr[f^1 \neq \tilde{f}^1]), \quad (42)$$

and thus, for $a_1 \in \{0, 1\}$,

$$\Pr[f^{a_1} \neq \tilde{f}^{a_1}] \leq 2D_2(F).$$

Since \tilde{f}^0 and \tilde{f}^1 are constant functions, this implies that

$$\mathbb{E}[f^{a_1}] \leq 2D_2(F) \quad \text{or} \quad \mathbb{E}[f^{a_1}] \geq 1 - 2D_2(F). \quad (43)$$

The rest of the proof is divided into two cases:

- **Case A:** For all $\sigma \in S_3$ we have $D'_2(F^\sigma) \leq D_2(F)/4$.
- **Case B:** There exists $\sigma_0 \in S_3$ such that $D'_2(F^{\sigma_0}) > D_2(F)/4$.

Case A: In this case, for any $\sigma \in S_3$, at least one of the choice functions of F^σ is at most $D_2(F)/4$ -far from a constant function. Note that if f^0 is at most $D_2(F)/4$ -far from a constant function, then f^1 must be at least $7D_2(F)/4$ -far from a constant function, since otherwise, f is less than $D_2(F)$ -far either from a constant function or from a dictatorship, contradicting the definition of $D_2(F)$. The same holds also for the pairs (g^0, g^1) and (h^0, h^1) . Thus, the only two possibilities are that either the functions f^1, g^1, h^1 or the functions f^0, g^0, h^0 are simultaneously at most $D_2(F)/4$ -far from a constant function. (For example, if $f^1, g^1,$ and h^0 are at most $D_2(F)/4$ -far from a constant function, then $f^0, g^0,$ and h^1 are at least $7D_2(F)/4$ -far from a constant function, and thus, for the preference $\sigma = (0, 0, 1)$, we have $D'_2(F^\sigma) \geq 7D_2(F)/4$, a contradiction. The other possibilities are discarded in a similar way). Assume w.l.o.g. that $f^1, g^1,$ and h^1 are at most $D_2(F)/4$ -far from a constant function. Furthermore, since amongst

the expectations $\mathbb{E}[f^1], \mathbb{E}[g^1], \mathbb{E}[h^1]$, at least two are close to one or at least two are close to zero, we can assume w.l.o.g. that

$$\Pr[f^1 \neq 1] \leq D_2(F)/4, \quad \text{and} \quad \Pr[g^1 \neq 1] \leq D_2(F)/4.$$

Consider the GSWF F^{σ_0} for the preference $\sigma_0 = (1, 1, 0)$. Since h^0 is at least $7D_2(F)/4$ -far from the constant zero function, it follows that

$$\begin{aligned} P(F^\sigma) &\geq \Pr_{\text{profile} \in (S_3)^{n-1}} [(f^1, g^1, h^0)(\text{profile}) = (1, 1, 1)] \\ &\geq 7D_2(F)/4 - D_2(F)/4 - D_2(F)/4 = 5D_2(F)/4, \end{aligned}$$

and thus,

$$P(F) = \frac{1}{6} \sum_{\sigma \in S_3} P(F^\sigma) \geq \frac{1}{6} \cdot P(F^{\sigma_0}) \geq \frac{5}{24} \cdot D_2(F) > \frac{1}{5000} \cdot RHC(1/2, D_2(F)),$$

as asserted.

Case B: Let $\sigma_0 \in S_3$ be such that $D'_2(F^{\sigma_0}) > D_2(F)/4$. By Equation (43), the choice function f^{a_1} of F^{σ_0} satisfies

$$\mathbb{E}[f^{a_1}] \leq 2D_2(F) \quad \text{or} \quad \mathbb{E}[f^{a_1}] \geq 1 - 2D_2(F),$$

and thus, $D'_2(F^{\sigma_0}) \leq 2D_2(F) < 2^{-500000}$. Hence, we can apply Lemma 3.2 to the GSWF G^{σ_0} , and get

$$P(F^{\sigma_0}) \geq \frac{1}{10} \cdot RHC(1/2, D'_2(F^{\sigma_0})) \geq \frac{1}{10} \cdot RHC(1/2, D_2(F)/4) \geq \frac{1}{640} \cdot RHC(1/2, D_2(F)).$$

Finally,

$$P(F) = \frac{1}{6} \sum_{\sigma \in S_3} P(F^\sigma) \geq \frac{1}{6} \cdot P(F^{\sigma_0}) > \frac{1}{5000} \cdot RHC(1/2, D_2(F)).$$

This completes the proof of the theorem. \square

The generalization to k alternatives for all $k \geq 3$ follows immediately by applying Theorem 4.2 to any subset of three alternatives.

5 Tightness of Results

In this section we show that for GSWFs on three alternatives, the assertions of Theorems 1.3 and 1.5 are tight up to logarithmic factors. In all our examples below, the Boolean choice functions f, g, h of the GSWF F are monotone threshold functions, that is, functions of the form:

$$(f(x) = 1) \Leftrightarrow \left(\sum_{i=1}^n x_i \geq l \right), \quad (44)$$

for different values of l . We note that in ([19], Theorem 2.9), Mossel et al. showed that amongst *neutral* GSWFs on three alternatives, a GSWF based on the majority rule is the “most rational” in the asymptotic sense (i.e., has the least probability of non-transitive outcome as the number of

voters tends to infinity). To some extent, our examples generalize this result to general GSWFs on three alternatives. The examples show that GSWFs based on monotone threshold Boolean choice functions are “close to be the most rational” amongst GSWFs whose choice functions have the same expectations, in the sense that their probability of non-transitive outcome is logarithmic close to the lower bound. In fact, we conjecture that such GSWFs are indeed the *most* rational amongst GSWFs whose choice functions have the same expectations. However, such exact result is not known even for neutral GSWFs.

We use the following proposition of Mossel et al. [18], showing that Borell’s reverse Bonami-Beckner inequality is essentially tight for diametrically opposed Hamming balls. Since we use the proposition only for noise of rate $\epsilon = 1/3$, we state it in this particular case.

Theorem 5.1 ([18], Proposition 3.9). *Fix $s, t > 0$, and let $f_n, g_n : \{0, 1\}^n \rightarrow \{0, 1\}$ be defined by*

$$(f_n(x) = 1) \Leftrightarrow \left(\sum_{i=1}^n x_i \leq \frac{n}{2} - \frac{s}{2} \sqrt{n} \right), \quad \text{and} \quad (g_n(x) = 1) \Leftrightarrow \left(\sum_{i=1}^n x_i \geq \frac{n}{2} + \frac{t}{2} \sqrt{n} \right).$$

Then

$$\lim_{n \rightarrow \infty} \sum_{S \subset \{1, \dots, n\}} \left(\frac{1}{3} \right)^{|S|} \hat{f}_n(S) \hat{g}_n(S) \leq \frac{\sqrt{8/9}}{2\pi s(s/3 + t)} \exp \left(-\frac{1}{2} \frac{s^2 + 2st/3 + t^2}{8/9} \right). \quad (45)$$

In order to show the tightness of Theorem 1.3, we fix a constant $\epsilon > 0$ and define the choice functions according to Equation (44), choosing the values of l such that

$$\mathbb{E}[f] = 0, \quad \mathbb{E}[g] = 1 - \epsilon, \quad \mathbb{E}[h] = 1 - \epsilon.$$

It is clear that $D_1(F) = \epsilon$. By Equation (25),

$$P(F) = \langle T_{1/3}(1 - g), \bar{h} \rangle.$$

By our construction, the pair of functions $(1 - g, \bar{h})$ is of the form considered in Theorem 5.1, with $s = t \approx \sqrt{2 \log(1/\epsilon)}$, and thus by the theorem, for n sufficiently large,

$$P(F) = \langle T_{1/3}(1 - g), \bar{h} \rangle \leq \frac{\sqrt{8/9}}{2\pi s(s/3 + t)} \exp \left(-\frac{1}{2} \frac{s^2 + 2st/3 + t^2}{8/9} \right) \approx C\epsilon^3 \log(1/\epsilon).$$

The lower bound asserted by Theorem 1.3 is $P(F) \geq C' \cdot \epsilon^3$, and thus the example shows the tightness of the assertion up to logarithmic factors.

The tightness of Theorem 1.5 is shown similarly, with choice functions chosen such that

$$\mathbb{E}[f] = \epsilon, \quad \mathbb{E}[g] = 1 - \epsilon, \quad \mathbb{E}[h] = 1/2.$$

It is clear that $D_2(F) = \epsilon$, and by Equation (25),

$$P(F) \leq \langle T_{1/3}f', h \rangle + \langle T_{1/3}(1 - g), \bar{h} \rangle.$$

The pairs (f', h) and $(1 - g, \bar{h})$ are both of the form considered in Theorem 5.1, and application of the theorem to both of them yields tightness up to a logarithmic factor, like in the previous case.

Finally, we note that while the examples above deal with GSWFs whose choice functions have constant expectation, it also makes sense to consider choice functions whose expectation tends to zero, as n (the number of voters) tends to infinity. In particular, one may ask what is the *least possible* probability of non-transitive outcome, as function of n , for GSWFs with $D_1(F) > 0$ or $D_2(F) > 0$. It appears that the question is of interest mainly for $D_2(F)$, as for $D_1(F)$, one can easily check that the minimal possible probability of 6^{-n} is obtained by a GSWF whose choice functions are chosen according to Equation (44), such that

$$\mathbb{E}[f] = 0, \quad \mathbb{E}[g] = 1 - 2^{-n}, \quad \mathbb{E}[h] = 1 - 2^{-n}.$$

For $D_2(F)$, it was shown in [17] that for a GSWF whose choice functions are chosen according to Equation (44), such that

$$\mathbb{E}[f] = 2^{-n}, \quad \mathbb{E}[g] = 1 - 2^{-n}, \quad \mathbb{E}[h] = 1/2,$$

we have $P(F) \leq 0.471^n$. Furthermore, it was conjectured that this is the most rational GSWF on three alternatives that satisfies the assumptions of Arrow’s theorem (and in particular, the minimal possible probability 6^{-n} is not obtained). Our results show that this function is at least “close” to be the most rational, as by Theorem 1.5, for any GSWF F such that $D_2(F) > 0$, we have

$$P(F) \geq C \cdot RHC(1/2, 2^{-n}) \approx C \cdot 0.458^n.$$

6 Questions for Further Research

We conclude this paper with several open problems related to our results.

- Our main lemma (Lemma 3.2) gives an essentially tight lower bound on the probability of non-transitive outcome for GSWFs in which at least one of the Boolean choice functions is “close” to a constant function. In the case where the distance from constant functions is greater than a fixed constant, our technique fails, and we use Mossel’s theorem [20] instead. As a result, the constant multiplicative factor in the assertions of Theorems 1.3 and 1.5 is extremely small, and clearly non-optimal. It will be interesting to find a “direct” proof also for GSWFs whose Boolean choice functions are “far” from constant functions, thus removing the reliance of the proof on the non-linear invariance principle (used in Mossel’s argument) that seems “unnatural” in our context, and improving the constant factor.
- While the results of Kalai [14] and Mossel [20] hold also for more general distributions of the individual preferences called “even product distributions” or “symmetric distributions” (see [17, 20]), our proof does not extend directly to such distributions. The reason is that for highly biased distributions of the preferences, the lower bound obtained by Borell’s reverse Bonami-Beckner inequality is weaker, and cannot “beat” the upper bound obtained by the Bonami-Beckner inequality. Thus, obtaining a tight quantitative version of Arrow’s theorem for general even product distributions of the preferences is an interesting open problem.
- We believe that GSWFs whose Boolean choice functions are monotone threshold functions are the most rational amongst GSWFs whose choice functions have the same expectations, not only in the asymptotic sense, but also for any particular (large enough) n . However, this conjecture seems quite challenging, as it includes the Majority is Stablest conjecture (whose proof by Mossel et al. [19] holds only in the limit as $n \rightarrow \infty$).

- Another direction of research is using our techniques to obtain quantitative versions of other theorems in social choice theory. In [8], Friedgut et al. presented a quantitative version of the Gibbard-Satterthwaite theorem [11, 21] for neutral GSWFs on three alternatives. Recently, Isaksson et al. [12] generalized the result of [8] to neutral GSWFs on k alternatives, for all $k \geq 4$. One of the main ingredients in the proof of [8] is Kalai’s quantitative Arrow theorem for neutral GSWFs. It seems interesting to find out whether our quantitative version of Arrow’s theorem can lead to a quantitative Gibbard-Satterthwaite theorem for general GSWFs (without the neutrality assumption).
- Finally, our results (as well as the previous results of Kalai [14] and Mossel [20]) apply only to GSWFs that satisfy the IIA condition, since such GSWFs can be represented by their Boolean choice functions, which allows to use the tools of discrete harmonic analysis. It will be very interesting to find a quantitative version of Arrow’s theorem that will not assume the IIA condition, but rather will relate the probability of non-transitive outcome to the distance of the GSWF from satisfying IIA.

7 Acknowledgements

It is a pleasure to thank Gil Kalai and Elchanan Mossel for numerous fruitful discussions that motivated our work.

References

- [1] K.J. Arrow, A Difficulty in the Concept of Social Welfare, *Journal of Political Economy* **58**(4) (1950), pp. 328-346.
- [2] S. Barbera, Pivotal Voters: A New Proof of Arrow’s Theorem, *Economics Letters* **6** (1980), pp. 13–16.
- [3] W. Beckner, Inequalities in Fourier Analysis, *Annals of Math.* **102** (1975), pp. 159–182.
- [4] A. Bonamie, Etude des Coefficients Fourier des Fonctions de $L^p(G)$, *Ann. Inst. Fourier* **20** (1970), pp. 335–402.
- [5] C. Borell, Positivity Improving Operators and Hypercontractivity, *Math. Zeitschrift*, **180**(2) (1982), pp. 225–234.
- [6] M. de Condorcet, An Essay on the Application of Probability Theory to Plurality Decision Making, 1785.
- [7] U. Feige, G. Kindler, and R. O’Donnell, Understanding Parallel Repetition Requires Understanding Foams, *Electronic Colloquium on Computational Complexity*, Report No. 43 (2007).
- [8] E. Friedgut, G. Kalai, and N. Nisan, Elections Can be Manipulated Often, Proc. 49-th Ann. Symp. on Foundations of Comp. Sci. (FOCS), pp. 243–249, 2009.
- [9] W.V. Gehrlein, Condorcet’s Paradox and the Condorcet Efficiency of Voting Rules, *Math. Japon.* **45** (1997), pp. 173–199.

- [10] J. Geneakoplos, Three Brief Proofs of Arrow's Impossibility Theorem, Cowels Foundation Discussion Paper number 1123R, Yale University, 1997. Available online at: <http://ideas.uqam.ca/ideas/data/Papers/cwlcwldpp1123R.html>
- [11] A. Gibbard, Manipulation of Voting Schemes: a General Result, *Econometrica* **41(4)** (1973), pp. 587–601.
- [12] M. Isaksson, G. Kindler, and E. Mossel, The Geometry of Manipulation - a Quantitative Proof of the Gibbard Satterthwaite Theorem, available at arXiv:0911.0517v3, 2009.
- [13] J. Kahn, G. Kalai, and N. Linial, The Influence of Variables on Boolean Functions, Proc. 29-th Ann. Symp. on Foundations of Comp. Sci. (FOCS), pp. 68–80, Computer Society Press, 1988.
- [14] G. Kalai, A Fourier-theoretic Perspective on the Condorcet Paradox and Arrow's Theorem, *Adv. in Appl. Math.* **29** (2002), no. 3, pp. 412–426.
- [15] G. Kalai, private communication, 2007.
- [16] G. Kalai and M. Safra, Threshold Phenomena and Influence, in: *Computational Complexity and Statistical Physics*, (A.G. Percus, G. Istrate and C. Moore, eds.), Oxford University Press, New York, 2006, pp. 25-60.
- [17] N. Keller, On The Probability of a Rational Outcome for Generalized Social Welfare Functions on Three Alternatives, *J. of Combinatorial Theory, Ser. A* **117** (2010), pp. 389–410.
- [18] E. Mossel, R. O'Donnell, O. Regev, J.E. Steif, and B. Sudakov, Non-Interactive Correlation Distillation, Inhomogeneous Markov Chains, and the Reverse Bonami-Beckner Inequality, *Israel J. Math.* **154** (2006), pp. 299–336.
- [19] E. Mossel, R. O'Donnell, and K. Oleszkiewicz, Noise Stability of Functions with Low Influences: Invariance and Optimality, *Annals of Math.*, to appear.
- [20] E. Mossel, A Quantitative Arrow Theorem, available at arXiv:0903.2574v4., 2009.
- [21] M. A. Satterthwaite, Strategy-proofness and Arrow's Conditions: Existence and Correspondence Theorems for Voting Procedures and Social Welfare Functions, *J. of Economic Theory* **10** (1975), pp. 187–217.
- [22] R. Wilson, Social Choice Theory Without the Pareto Principle, *Journal of Economic Theory* **5(3)** (1972), pp. 478–486.