

## Mean Width of a Regular Simplex

STEVEN R. FINCH

March 12, 2016

**ABSTRACT.** The mean width is a measure on  $n$ -dimensional convex bodies. An integral formula for the mean width of a regular  $n$ -simplex appeared in the electrical engineering literature in 1997. As a consequence, expressions for the expected range of a sample of  $n + 1$  normally distributed variables, for  $n \leq 6$ , carry over to widths of regular  $n$ -simplices. As another consequence, precise asymptotics for the mean width become available as  $n \rightarrow \infty$ .

Let  $C$  be a convex body in  $\mathbb{R}^n$ . A **width** is the distance between a pair of parallel  $C$ -supporting planes (linear varieties of dimension  $n - 1$ ). Every unit vector  $u \in \mathbb{R}^n$  determines a unique such pair of planes orthogonal to  $u$  and hence a width  $w(u)$ . Let  $u$  be uniformly distributed on the unit sphere  $S^{n-1} \subset \mathbb{R}^n$ . Then  $w$  is a random variable and

$$\mathbb{E}(w_3) = \frac{3}{2\pi} \arccos\left(-\frac{1}{3}\right)$$

for  $C =$  the regular 3-simplex (tetrahedron) in  $\mathbb{R}^3$  with edges of unit length and

$$\mathbb{E}(w_4) = \frac{10}{3\pi^2} \left[ 3 \arccos\left(-\frac{1}{3}\right) - \pi \right]$$

for  $C =$  the regular 4-simplex in  $\mathbb{R}^4$  with edges of unit length [1, 2]. Our contribution is to extend the preceding **mean width** results to regular  $n$ -simplices in  $\mathbb{R}^n$  for  $n \leq 6$ . We similarly extend the following **mean square width** result:

$$\mathbb{E}(w_3^2) = \frac{1}{3} \left( 1 + \frac{3 + \sqrt{3}}{\pi} \right)$$

which, as far as is known, first appeared in [3].

The key observation underlying our work is due to Sun [4], which in turn draws upon material in [5, 6]. It does not seem to have been acknowledged in the mathematics literature. After most of this paper was written, we found [7], which assigns priority to Hadwiger [8] and to Ruben [9] for closely related ideas.

---

<sup>0</sup>Copyright © 2011, 2016 by Steven R. Finch. All rights reserved.

## 1. ORDER STATISTICS

Let  $X_1, X_2, \dots, X_n$  denote a random sample from a Normal  $(0, 1)$  distribution, that is, with density function  $f$  and cumulative distribution  $F$ :

$$f(x) = \frac{1}{\sqrt{2\pi}} \exp\left(-\frac{x^2}{2}\right), \quad F(x) = \int_{-\infty}^x f(\xi) d\xi = \frac{1}{2} \operatorname{erf}\left(\frac{x}{\sqrt{2}}\right) + \frac{1}{2}.$$

The first two moments of the **range**

$$r_n = \max\{X_1, X_2, \dots, X_n\} - \min\{X_1, X_2, \dots, X_n\}$$

are given by [10, 11]

$$\mu_n = \mathbb{E}(r_n) = \int_{-\infty}^{\infty} \{1 - F(x)^n - [1 - F(x)]^n\} dx,$$

$$\nu_n = \mathbb{E}(r_n^2) = 2 \int_{-\infty}^{\infty} \int_{-\infty}^y \{1 - F(y)^n - [1 - F(x)]^n + [F(y) - F(x)]^n\} dx dy.$$

For small  $n$ , exact expressions are possible [12, 13, 14]:

$$\begin{aligned} \mu_2 &= \frac{2}{\sqrt{\pi}} = 1.128\dots, & \nu_2 &= 2, \\ \mu_3 &= \frac{3}{\sqrt{\pi}} = 1.692\dots, & \nu_3 &= 2 \left(1 + \frac{3\sqrt{3}}{2\pi}\right) = 3.653\dots, \\ \mu_4 &= \frac{6}{\sqrt{\pi}} (1 - 2S_2) = 2.058\dots, & \nu_4 &= 2 \left(1 + \frac{3+\sqrt{3}}{\pi}\right) = 5.012\dots, \\ \mu_5 &= \frac{10}{\sqrt{\pi}} (1 - 3S_2) = 2.325\dots, & \nu_5 &= 2 \left(1 + \frac{5\sqrt{3}}{2\pi} + \frac{30}{\pi}S_{1/2} - \frac{5\sqrt{3}}{\pi}S_3\right) = 6.156\dots, \\ \mu_6 &= \frac{15}{\sqrt{\pi}} (1 - 4S_2 + 2T_2) = 2.534\dots, & \nu_6 &= 2 \left(1 + \frac{5(9+2\sqrt{3})}{2\pi} - \frac{90}{\pi}S_2 - \frac{15\sqrt{3}}{\pi}S_3\right) = 7.142\dots, \\ \mu_7 &= \frac{21}{\sqrt{\pi}} (1 - 5S_2 + 5T_2) = 2.704\dots, & \nu_7 &= 2 \left(1 + \frac{35\sqrt{3}}{4\pi} + \frac{210}{\pi}S_{1/2} - \frac{105}{\pi}S_2 - \frac{35\sqrt{3}}{\pi}S_3 \right. \\ & & & \left. + \frac{35\sqrt{3}}{2\pi}T_3 + \frac{210}{\pi}U - \frac{420}{\pi}V\right) = 8.007\dots \end{aligned}$$

where

$$S_k = \frac{\sqrt{k}}{\pi} \int_0^{\pi/4} \frac{dx}{\sqrt{k + \sec(x)^2}} = \frac{1}{2\pi} \operatorname{arcsec}(k + 1),$$

$$T_k = \frac{\sqrt{k}}{\pi^2} \int_0^{\pi/4} \int_0^{\pi/4} \frac{dx dy}{\sqrt{k + \sec(x)^2 + \sec(y)^2}} = \frac{1}{2\pi^2} \int_0^{\pi S_k} \operatorname{arcsec}\left(1 + \frac{k(k+1)}{k - \tan(z)^2}\right) dz,$$

$$U = \frac{1}{\pi^2} \int_0^1 \frac{\operatorname{arcsec}(2t^2 + 4)}{(2t^2 + 1)\sqrt{2t^2 + 3}} dt, \quad V = \frac{1}{\pi^2} \int_0^1 \frac{\operatorname{arcsec}(t^2 + 5)}{(t^2 + 2)\sqrt{t^2 + 4}} dt.$$

The preceding table complements an analogous table in [15] for first and second moments of  $\max\{X_1, X_2, \dots, X_n\}$ . Similar expressions for  $\mu_8 = 2.847\dots$  and  $\nu_8 = 8.778\dots$  remain to be found.

## 2. KEY OBSERVATION

Let us rescale length so that the circumradius of the  $n$ -simplex is 1. Adjusted width will be denoted by  $\tilde{w}_n$ . Using optimality properties of the  $n$ -simplex, Sun [4] deduced a formula for mean half width:

$$\begin{aligned} \frac{1}{2}\mathbb{E}(\tilde{w}_n) &= \frac{n+1}{2} \sqrt{\frac{(n+1)n}{2\pi}} \frac{\Gamma(\frac{n}{2})}{\Gamma(\frac{n+1}{2})} \int_{-\infty}^{\infty} F\left(\frac{x}{\sqrt{2}}\right)^{n-1} f(x) dx \\ &= \frac{(n+1)^{3/2}}{\sqrt{2n}} \frac{\Gamma(\frac{n}{2})}{\Gamma(\frac{n+1}{2})} \int_{-\infty}^{\infty} x F(x)^n f(x) dx \end{aligned}$$

(see Corollary 2 on p. 1581 and its proof on p. 1585; his  $M$  is the same as our  $n+1$ ). We recognize the latter integral as  $\mu_{n+1}/(2(n+1))$ ; hence

$$\mathbb{E}(\tilde{w}_n) = \sqrt{\frac{n+1}{2n}} \frac{\Gamma(\frac{n}{2})}{\Gamma(\frac{n+1}{2})} \mu_{n+1}$$

and therefore

$$\mathbb{E}(w_n) = \frac{1}{2} \frac{\Gamma(\frac{n}{2})}{\Gamma(\frac{n+1}{2})} \mu_{n+1}$$

because, in our original scaling, the circumradius is  $\sqrt{n/(2(n+1))}$ .

No similar integral expression for  $\mathbb{E}(\tilde{w}_n^2)$  appears in [4]. We circumvent this difficulty by noticing that the formula [16, 17]

$$\mathbb{E}\left(\sqrt{\sum_{k=1}^n X_k^2}\right) = \sqrt{2} \frac{\Gamma(\frac{n+1}{2})}{\Gamma(\frac{n}{2})}$$

bears some resemblance to the coefficient of  $\mu_{n+1}$  in our expression for  $\mathbb{E}(w_n)$ . The square version

$$\mathbb{E}\left(\sum_{k=1}^n X_k^2\right) = n$$

is trivial and leads us to conjecture that

$$\mathbb{E}(w_n^2) = \frac{1}{2n} \nu_{n+1}$$

by analogy. Numerical confirmation for  $n \leq 6$  is possible via the computer algebra technique described in [3].

In summary, we have mean width results

$$\mathbb{E}(w_2) = \frac{3}{\pi} = 0.954929658551372\dots,$$

$$\mathbb{E}(w_3) = \frac{3}{2}(1 - 2S_2) = 0.912260171954089\dots,$$

$$\mathbb{E}(w_4) = \frac{20}{3\pi}(1 - 3S_2) = 0.874843256085440\dots,$$

$$\mathbb{E}(w_5) = \frac{45}{16}(1 - 4S_2 + 2T_2) = 0.842274297659162\dots,$$

$$\mathbb{E}(w_6) = \frac{56}{5\pi}(1 - 5S_2 + 5T_2) = 0.813743951590337\dots,$$

and mean square width results

$$\mathbb{E}(w_2^2) = \frac{1}{2} \left( 1 + \frac{3\sqrt{3}}{2\pi} \right) = 0.913496671566344\dots,$$

$$\mathbb{E}(w_3^2) = \frac{1}{3} \left( 1 + \frac{3 + \sqrt{3}}{\pi} \right) = 0.835419517991054\dots,$$

$$\begin{aligned} \mathbb{E}(w_4^2) &= \frac{1}{4} \left( 1 + \frac{5\sqrt{3}}{2\pi} + \frac{30}{\pi} S_{1/2} - \frac{5\sqrt{3}}{\pi} S_3 \right) \\ &= 0.769572883591771\dots, \end{aligned}$$

$$\begin{aligned} \mathbb{E}(w_5^2) &= \frac{1}{5} \left( 1 + \frac{5(9 + 2\sqrt{3})}{2\pi} - \frac{90}{\pi} S_2 - \frac{15\sqrt{3}}{\pi} S_3 \right) \\ &= 0.714241915072694\dots, \end{aligned}$$

$$\begin{aligned} \mathbb{E}(w_6^2) &= \frac{1}{6} \left( 1 + \frac{35\sqrt{3}}{4\pi} + \frac{210}{\pi} S_{1/2} - \frac{105}{\pi} S_2 - \frac{35\sqrt{3}}{\pi} S_3 \right. \\ &\quad \left. + \frac{35\sqrt{3}}{2\pi} T_3 + \frac{210}{\pi} U - \frac{420}{\pi} V \right) \\ &= 0.667314714095430\dots \end{aligned}$$

### 3. ASYMPTOTICS

We turn now to the asymptotic distribution of  $r_n$  as  $n \rightarrow \infty$ . Define  $a_n$  to be the positive solution of the equation [12, 18]

$$2\pi a_n^2 \exp(a_n^2) = n^2,$$

that is,

$$a_n = \sqrt{W\left(\frac{n^2}{2\pi}\right)} \sim \sqrt{2\ln(n)} - \frac{1}{2} \frac{\ln(\ln(n)) + \ln(4\pi)}{\sqrt{2\ln(n)}}$$

in terms of the Lambert  $W$  function [19]. It can be proved that the required density is a convolution [20, 21]:

$$\begin{aligned} \lim_{n \rightarrow \infty} \frac{d}{dy} \mathbb{P}\left(\sqrt{2\ln(n)}(r_n - 2a_n) < y\right) &= \int_{-\infty}^{\infty} \exp(-x - e^{-x}) \exp(-(y-x) - e^{-(y-x)}) dx \\ &= 2e^{-y} K_0(2e^{-y/2}) \end{aligned}$$

where  $K_0$  is the modified Bessel function of the second kind [22]. A random variable  $Y$ , distributed as such, satisfies

$$\mathbb{E}(Y) = 2\gamma, \quad \mathbb{E}(Y^2) = \frac{\pi^2}{3} + 4\gamma^2$$

where  $\gamma$  is the Euler-Mascheroni constant [23]. This implies that

$$\mu_n \sim 2 \left( a_n + \frac{\gamma}{\sqrt{2\ln(n)}} \right) \sim 2\sqrt{2\ln(n)} - \frac{\ln(\ln(n)) + \ln(4\pi) - 2\gamma}{\sqrt{2\ln(n)}}$$

and hence

$$\begin{aligned} \mathbb{E}(w_n) &= \frac{1}{2} \frac{\Gamma\left(\frac{n}{2}\right)}{\Gamma\left(\frac{n+1}{2}\right)} \mu_{n+1} = \frac{1}{2} \frac{\Gamma\left(\frac{n}{2}\right)}{\Gamma\left(\frac{n+1}{2}\right)} \cdot \frac{\mu_{n+1}}{\mu_n} \cdot \mu_n \\ &\sim \frac{1}{\sqrt{2n}} \left(1 + \frac{1}{4n}\right) \cdot \left(1 + \frac{1}{2n\ln(n)}\right) \cdot 2 \left(a_n + \frac{\gamma}{\sqrt{2\ln(n)}}\right) \\ &\sim 2\sqrt{\frac{\ln(n)}{n}} - \frac{\ln(\ln(n)) + \ln(4\pi) - 2\gamma}{2\sqrt{n\ln(n)}}. \end{aligned}$$

More terms in the asymptotic expansion are possible.

If we rescale length so that the inradius of the  $n$ -simplex is 1 and denote adjusted width by  $\widehat{w}_n$ , then

$$\mathbb{E}(\widehat{w}_n) \sim \sqrt{2n} \cdot 2\sqrt{\frac{\ln(n)}{n}} \sim 2\sqrt{2n \ln(n)}$$

because, in our original scaling, the inradius is  $\sqrt{1/(2n(n+1))}$ . This first-order approximation is consistent with [2].

#### 4. REGULAR OCTAHEDRON

As an aside, we return to the setting of  $\mathbb{R}^3$  and review our computational methods for  $C =$  the regular octahedron with edges of unit length.

For simplicity, let  $\diamond$  be the octahedron with vertices

$$\begin{aligned} v_1 &= (1, 0, 0), & v_2 &= (-1, 0, 0), & v_3 &= (0, 1, 0), \\ v_4 &= (0, -1, 0), & v_5 &= (0, 0, 1), & v_6 &= (0, 0, -1). \end{aligned}$$

At the end, it will be necessary to normalize by  $\sqrt{2}$ , the edge-length of  $\diamond$ .

Also let  $\widetilde{\diamond}$  be the union of six overlapping balls of radius  $1/2$  centered at  $v_1/2, v_2/2, v_3/2, v_4/2, v_5/2, v_6/2$ . Clearly  $\diamond \subset \widetilde{\diamond}$  and  $\widetilde{\diamond}$  has centroid  $(0, 0, 0)$ . A diameter of  $\widetilde{\diamond}$  is the length of the intersection between  $\widetilde{\diamond}$  and a line passing through the origin.

Computing all widths of  $\diamond$  is equivalent to computing all diameters of  $\widetilde{\diamond}$ . The latter is achieved as follows. Fix a point  $(a, b, c)$  on the unit sphere. The line  $L$  passing through  $(0, 0, 0)$  and  $(a, b, c)$  has parametric representation

$$x = ta, \quad y = tb, \quad z = tc, \quad t \in \mathbb{R}$$

and hence  $y = (b/a)x, z = (c/a)x$  assuming  $a \neq 0$ . The nontrivial intersection between first sphere and  $L$  satisfies

$$\left(x - \frac{1}{2}\right)^2 + \left(\frac{b}{a}x\right)^2 + \left(\frac{c}{a}x\right)^2 = \frac{1}{4}$$

thus  $x_1 = a^2$  since  $a^2 + b^2 + c^2 = 1$ ; the nontrivial intersection between second sphere and  $L$  satisfies

$$\left(x + \frac{1}{2}\right)^2 + \left(\frac{b}{a}x\right)^2 + \left(\frac{c}{a}x\right)^2 = \frac{1}{4}$$

thus  $x_2 = -a^2$ . The nontrivial intersection between third/fourth sphere and  $L$  satisfies

$$x^2 + \left(\frac{b}{a}x \mp \frac{1}{2}\right)^2 + \left(\frac{c}{a}x\right)^2 = \frac{1}{4}$$

thus  $x_3 = ab, x_4 = -ab$ . The nontrivial intersection between fifth/sixth sphere and  $L$  satisfies

$$x^2 + \left(\frac{b}{a}x\right)^2 + \left(\frac{c}{a}x \mp \frac{1}{2}\right)^2 = \frac{1}{4}$$

thus  $x_5 = ac$ ,  $x_6 = -ac$ .

We now examine all pairwise distances, squared, between the six intersection points:

$$(x_i - x_j)^2 + \left(\frac{b}{a}x_i - \frac{b}{a}x_j\right)^2 + \left(\frac{c}{a}x_i - \frac{c}{a}x_j\right)^2 = \begin{cases} 4a^2 & \text{if } i = 1, j = 2 \\ 1 - 2ab - c^2 & \text{if } i = 1, j = 3 \text{ or } i = 2, j = 4 \\ 1 + 2ab - c^2 & \text{if } i = 1, j = 4 \text{ or } i = 2, j = 3 \\ 1 - 2ac - b^2 & \text{if } i = 1, j = 5 \text{ or } i = 2, j = 6 \\ 1 + 2ac - b^2 & \text{if } i = 1, j = 6 \text{ or } i = 2, j = 5 \\ 4b^2 & \text{if } i = 3, j = 4 \\ (b - c)^2 & \text{if } i = 3, j = 5 \text{ or } i = 4, j = 6 \\ (b + c)^2 & \text{if } i = 3, j = 6 \text{ or } i = 4, j = 5 \\ 4c^2 & \text{if } i = 5, j = 6 \end{cases}$$

and define

$$g(a, b) = \max \{4a^2, 1 - 2ab - c^2, 1 + 2ab - c^2, 1 - 2ac - b^2, 1 + 2ac - b^2, 4b^2, (b - c)^2, (b + c)^2, 4c^2\}.$$

The mean width for  $C$  is

$$\frac{1}{\sqrt{2}} \frac{1}{4\pi} \int_0^{2\pi} \int_0^\pi \sqrt{g(\cos \theta \sin \varphi, \sin \theta \sin \varphi, \cos \varphi)} \sin \varphi \, d\varphi \, d\theta = \frac{3}{\pi} \arccos \left( \frac{1}{3} \right)$$

and the mean square width is

$$\frac{1}{2} \frac{1}{4\pi} \int_0^{2\pi} \int_0^\pi g(\cos \theta \sin \varphi, \sin \theta \sin \varphi, \cos \varphi) \sin \varphi \, d\varphi \, d\theta = \frac{2}{3} \left( 1 + \frac{2\sqrt{3}}{\pi} \right).$$

Here are details on the final integral. A plot of the surface

$$(\theta, \varphi) \mapsto \sqrt{\frac{g(\cos \theta \sin \varphi, \sin \theta \sin \varphi, \cos \varphi)}{2}}$$

appears in Figure 1, where  $0 \leq \theta \leq 2\pi$  and  $0 \leq \varphi \leq \pi$ . Figure 2 contains the same surface, but viewed from above. Our focus will be on the part of the surface to the right of the bottom center, specifically  $0 \leq \theta \leq \pi/4$  and  $\pi/2 \leq \varphi \lesssim 9/4$ . The volume under this part is  $1/24^{\text{th}}$  of the volume under the full surface.

We need to find the precise upper bound on  $\varphi$  as a function of  $\theta$ . Recall the formula for  $g$  as a maximum over nine terms; let  $g_\ell$  denote the  $\ell^{\text{th}}$  term, where  $1 \leq \ell \leq 9$ . Then the upper bound on  $\varphi$  is found by solving the equation

$$g_1(\cos \theta \sin \varphi, \sin \theta \sin \varphi, \cos \varphi) = g_9(\cos \theta \sin \varphi, \sin \theta \sin \varphi, \cos \varphi)$$

for  $\varphi$ . We obtain  $\varphi(\theta) = 2 \arctan(h(\theta))$ , where

$$h(\theta) = \cos \theta + \sqrt{\frac{3 + \cos(2\theta)}{2}}$$

and, in particular,

$$\varphi(0) = 2 \arctan(1 + \sqrt{2}) \approx 2.3562,$$

$$\varphi(\pi/4) = 2 \arctan\left(\frac{(1 + \sqrt{3})/\sqrt{2}}{\sqrt{2}}\right) \approx 2.1862.$$

It follows that  $g = g_1$  for  $0 \leq \theta \leq \pi/4$  and  $\pi/2 \leq \varphi \leq 2 \arctan(h)$ . Now we have

$$\begin{aligned} & \frac{1}{2} \frac{1}{4\pi} \int g_1(\cos \theta \sin \varphi, \sin \theta \sin \varphi, \cos \varphi) \sin \varphi d\varphi \\ &= \frac{(1 + \cos(2\theta)) (\cos(3\varphi) - 9 \cos(\varphi))}{48\pi} \end{aligned}$$

and

$$\cos(3\varphi)|_{\pi/2}^{2 \arctan(h)} = \frac{(1 + 4h + h^2) (1 - 4h + h^2) (1 - h^2)}{(1 + h^2)^3},$$

$$\cos(\varphi)|_{\pi/2}^{2 \arctan(h)} = \frac{1 - h^2}{1 + h^2},$$

therefore

$$\begin{aligned} & \frac{1}{2} \frac{1}{4\pi} \int_{\pi/2}^{2 \arctan(h)} g_1(\cos \theta \sin \varphi, \sin \theta \sin \varphi, \cos \varphi) \sin \varphi d\varphi \\ &= \frac{(h^4 + 4h^2 + 1) (h^2 - 1) (1 + \cos(2\theta))}{6\pi(1 + h^2)^3}. \end{aligned}$$

Integrating this expression from 0 to  $\pi/4$  gives the desired formula for  $\mathbb{E}(w_{\text{octa}}^2)$ .

## 5. $n$ -CUBES

After having written the preceding, we discovered [7], which gives the mean width for a regular  $n$ -simplex in  $\mathbb{R}^n$  as

$$\mathbb{E}(w_n) = \frac{n(n+1)}{\sqrt{2}\pi} \frac{\Gamma(\frac{n}{2})}{\Gamma(\frac{n+1}{2})} \int_{-\infty}^{\infty} e^{-2x^2} \left(\frac{1 + \operatorname{erf}(x)}{2}\right)^{n-1} dx.$$



Consistency is readily established; nothing is said in [7] about the connection between  $\mathbb{E}(w_n)$  and order statistics from a normal distribution (more precisely, the expected range  $\mu_{n+1}$ ).

By contrast, the mean width for an  $n$ -cube with edges of unit length is elementary:

$$\mathbb{E}(w_{n\text{-cube}}) = \frac{n}{\sqrt{\pi}} \frac{\Gamma\left(\frac{n}{2}\right)}{\Gamma\left(\frac{n+1}{2}\right)}$$

and we conjecture that

$$\mathbb{E}(w_{n\text{-cube}}^2) = 1 + \frac{2(n-1)}{\pi}.$$

## 6. $n$ -CROSSPOLYTOPES

A regular  $n$ -crosspolytope with edges of unit length has mean width [7, 24]

$$\mathbb{E}(w_{n\text{-crosspolytope}}) = \frac{2\sqrt{2}n(n-1)}{\pi} \frac{\Gamma\left(\frac{n}{2}\right)}{\Gamma\left(\frac{n+1}{2}\right)} \int_0^\infty e^{-2x^2} \operatorname{erf}(x)^{n-2} dx;$$

the case  $n = 3$  corresponds to the octahedron discussed earlier. It is not surprising that a connection exists with order statistics from a half-normal (folded) distribution. We will examine this later, as well as relevant expressions from [25]. An appropriate mean square conjecture also needs to be formulated in this scenario.

## REFERENCES

- [1] K. Böröczky, About the mean width of simplices, *Period. Polytech. Mech. Engrg.* 36 (1992) 291–297; MR1269513 (95b:52024).
- [2] K. Böröczky and R. Schneider, Circumscribed simplices of minimal mean width, *Beiträge Algebra Geom.* 48 (2007) 217–224; MR2326411 (2008j:52005).
- [3] S. R. Finch, Width distributions for convex regular polyhedra, arXiv:1110.0671.
- [4] Y. Sun, Stochastic iterative algorithms for signal set design for Gaussian channels and optimality of the L2 signal set, *IEEE Trans. Inform. Theory* 43 (1997) 1574–1587; MR1476788 (99a:94064).
- [5] C. L. Weber, *Elements of Detection and Signal Design*, Springer-Verlag, 1987, pp. 149–214.
- [6] A. V. Balakrishnan, A contribution to the sphere-packing problem of communication theory, *J. Math. Anal. Appl.* 3 (1961) 485–506; MR0219340 (36 #2423).

- [7] M. Henk, J. Richter-Gebert and G. M. Ziegler, Basic properties of convex polytopes, *Handbook of Discrete and Computational Geometry*, CRC Press, 1997, 243–270; MR1730169.
- [8] H. Hadwiger, Gitterpunktanzahl im Simplex und Wills'sche Vermutung, *Math. Annalen* 239 (1979) 271–288; MR0522784 (80d:52015).
- [9] H. Ruben, On the geometrical moments of skew-regular simplices in hyperspherical space, with some applications in geometry and mathematical statistics, *Acta Math.* 103 (1960) 1–23; MR0121713 (22 #12447).
- [10] L. H. C. Tippett, On the extreme individuals and the range of samples taken from a normal population, *Biometrika* 17 (1925) 364–387.
- [11] K. V. Mardia, Tippett's formulas and other results on sample range and extremes, *Annals Inst. Statist. Math.* 17 (1965) 85–91; MR0178522 (31 #2779).
- [12] H. A. David, *Order Statistics*, 2<sup>nd</sup> ed., Wiley, 1981, pp. 38–43, 53, 258–269; MR0099101 (20 #5545).
- [13] H. Ruben, On the moments of the range and product moments of extreme order statistics in normal samples, *Biometrika* 43 (1956) 458–460; MR0082769 (18,607d).
- [14] Y. Watanabe, M. Isida, S. Taga, Y. Ichijo, T. Kawase, G. Niside, Y. Takeda, A. Horisuzi, and I. Kuriyama, Some contributions to order statistics, *J. Gakugei, Tokushima Univ.* 8 (1957) 41–90; MR0099101 (20 #5545).
- [15] S. R. Finch, Extreme value constants, *Mathematical Constants*, Cambridge Univ. Press, 2003, pp. 363–367; MR2003519 (2004i:00001).
- [16] I. Ben Yaacov, Continuous and random Vapnik-Chervonenkis classes, *Israel J. Math.* 173 (2009) 309–333; MR2570671 (2011j:03072); arXiv:0802.0068.
- [17] H. O. Lancaster, Chi distribution, *Encyclopedia of Statistical Sciences*, v. 1, ed. S. Kotz, N. L. Johnson and C. B. Read, Wiley, 1982, p. 439; MR0646617 (83j:62001a).
- [18] P. Hall, On the rate of convergence of normal extremes, *J. Appl. Probab.* 16 (1979) 433–439; MR0531778 (80d:60025).
- [19] R. M. Corless, G. H. Gonnet, D. E. G. Hare, D. J. Jeffrey and D. E. Knuth, On the Lambert  $W$  function, *Adv. Comput. Math.* 5 (1996) 329–359; MR1414285 (98j:33015).

- [20] E. J. Gumbel, The distribution of the range, *Annals of Math. Statistics* 18 (1947) 384–412; MR0022331 (9,195a).
- [21] D. R. Cox, A note on the asymptotic distribution of range, *Biometrika* 35 (1948) 310–315; MR0028562 (10,466b).
- [22] F. W. J. Olver, Bessel functions of integer order, *Handbook of Mathematical Functions*, ed. M. Abramowitz and I. A. Stegun, Dover, 1992, pp. 374–377; MR1225604 (94b:00012).
- [23] S. R. Finch, Euler-Mascheroni constant, *Mathematical Constants*, Cambridge Univ. Press, 2003, pp. 28–40; MR2003519 (2004i:00001).
- [24] U. Betke and M. Henk, Intrinsic volumes and lattice points of crosspolytopes, *Monatsh. Math.* 115 (1993) 27–33; MR1223242 (94g:52010).
- [25] Z. Govindarajulu, Exact lower moments of order statistics in samples from the chi-distribution (1 d.f.), *Annals Math. Statist.* 33 (1962) 1292–1305; MR0141179 (25 #4590).
- [26] S. R. Finch, Simulations in R involving colliding dice and mean widths, <http://www.people.fas.harvard.edu/~sfinch/csolve/rsimul.html>.

Steven R. Finch  
Dept. of Statistics  
Harvard University  
Cambridge, MA, USA  
[steven\\_finch@harvard.edu](mailto:steven_finch@harvard.edu)




Figure 1: Surface plot of  $\sqrt{g/2}$ , where  $0 \leq \theta \leq 2\pi$  and  $0 \leq \varphi \leq \pi$ .




Figure 2: Another view of  $\sqrt{g/2}$ , with contours of intersection.