CANONICAL-TYPE CONNECTION ON ALMOST CONTACT MANIFOLDS WITH B-METRIC

MANCHO MANEV AND MIROSLAVA IVANOVA

ABSTRACT. The canonical-type connection on the almost contact manifolds with B-metric is constructed. It is proved that its torsion is invariant with respect to a subgroup of the general conformal transformations of the almost contact B-metric structure. The basic classes of the considered manifolds are characterized in terms of the torsion of the canonical-type connection.

Introduction

In the differential geometry of the manifolds with additional structures, there are important the so-called natural connections, i.e. linear connections with torsion such that the additional structures are parallel with respect to them. There exists a significant interest to these natural connections which have some additional geometric or algebraic properties, for instance about their torsion.

On an almost Hermitian manifold (M,J,g) there exists a unique natural connection ∇^C with a torsion T such that $T(J\cdot,J\cdot)=-T(\cdot,\cdot)$. This connection is known as the canonical Hermitian connection or the Chern connection. An example of the natural Hermitian connection is the first canonical connection of Lichnerowicz ∇^L [15, 16]. According to [9], there exists a one-parameter family of canonical Hermitian connections $\nabla^t = t\nabla^C + (1-t)\nabla^L$. The connection ∇^t obtained for t=-1 is called the Bismut connection or the KT-connection, which is characterized with a totally skew-symmetric torsion. The latter connection with a closed torsion 3-form has applications in type II string theory and in 2-dimensional supersymmetric σ -models [8, 26, 14]. In [2] and [3] all almost contact metric, almost Hermitian and G_2 -structures admitting a connection with totally skew-symmetric torsion tensor are described.

Natural connections of canonical type are considered on the Riemannian almost product manifolds in [10, 11, 12] and on the almost complex manifolds with Norden metric in [6, 4, 24]. The connection in [4] is the so-called B-connection, which is studied on the class of the locally conformal Kählerian manifolds with Norden metric.

In the present paper¹ we consider natural connections (i.e. preserving the structure) of canonical type on the almost contact manifolds with B-metric. These manifolds are the odd-dimensional extension of the almost complex manifolds with Norden metric and the case with indefinite metrics corresponding to the almost contact metric manifolds.

The paper is organized as follows. In Sec. 1 we give some necessary facts about the considered manifolds. In Sec. 2 we define a natural connection of canonical type on an

1

²⁰¹⁰ Mathematics Subject Classification. 53C05, 53C15, 53C50.

Key words and phrases. almost contact manifold, B-metric, natural connection, canonical connection, conformal transformation, torsion tensor.

¹Partially supported by project NI11-FMI-004 of the Scientific Research Fund, Paisii Hilendarski University of Plovdiv, Bulgaria and the German Academic Exchange Service (DAAD)

almost contact manifold with B-metric. We determine the class of the considered manifolds where this connection and a known natural connection coincide. In Sec. 3 we consider the group G of the general conformal transformations of the almost contact B-metric structure. We determine the invariant class of the considered manifolds and a tensor invariant of the group G. In Sec. 4 we establish that the torsion of the canonical-type connection is invariant exactly in the subgroup G_0 of G. We characterize the basic classes of the considered manifolds by the torsion of the canonical-type connection. In Sec. 5 we determine the class of the almost contact B-metric manifolds of Sasakian type and supply a relevant example.

1. ALMOST CONTACT MANIFOLDS WITH B-METRIC

Let (M, φ, ξ, η) be an almost contact manifold, i.e. M is a (2n+1)-dimensional differentiable manifold with an almost contact structure (φ, ξ, η) consisting of an endomorphism φ of the tangent bundle, a vector field ξ and its dual 1-form η such that the following algebraic relations are satisfied:

(1)
$$\varphi \xi = 0, \quad \varphi^2 = -\operatorname{Id} + \eta \otimes \xi, \quad \eta \circ \varphi = 0, \quad \eta(\xi) = 1.$$

Further, let us endow the almost contact manifold (M, φ, ξ, η) with a pseudo-Riemannian metric g of signature (n, n+1) determined by

(2)
$$g(\varphi x, \varphi y) = -g(x, y) + \eta(x)\eta(y)$$

for arbitrary x, y of the algebra $\mathfrak{X}(M)$ on the smooth vector fields on M. Then $(M, \varphi, \xi, \eta, g)$ is called an almost contact manifold with B-metric or an *almost contact B-metric manifold*. Further, x, y, z will stand for arbitrary elements of $\mathfrak{X}(M)$.

The associated metric \tilde{g} of g on M is defined by $\tilde{g}(x,y) = g(x,\varphi y) + \eta(x)\eta(y)$. Both metrics g and \tilde{g} are necessarily of signature (n,n+1). The manifold $(M,\varphi,\xi,\eta,\tilde{g})$ is also an almost contact B-metric manifold.

Let us remark that the 2n-dimensional contact distribution $H=\ker(\eta)$ generated by the contact 1-form η can be considered as the horizontal distribution of the sub-Riemannian manifold M. Then H is endowed with an almost complex structure determined as $\varphi|_H$ – the restriction of φ on H, as well as a Norden metric $g|_H$, i.e. $g|_H(\varphi|_{H^*}, \varphi|_{H^*}) = -g|_H(\cdot,\cdot)$. Moreover, H can be considered as a n-dimensional complex Riemannian manifold with a complex Riemannian metric $g^{\mathbb{C}} = g|_H + i\tilde{g}|_H$ [5].

The structural group of the almost contact B-metric manifolds is $(GL(n,\mathbb{C}) \cap O(n,n))$ $\times I_1$, i.e. it consists of real square matrices of order 2n+1 of the following type

$$\begin{pmatrix} A & B & \vartheta^T \\ \hline -B & A & \vartheta^T \\ \hline \vartheta & \vartheta & 1 \end{pmatrix}, \qquad \begin{array}{c} A^TA - B^TB = I_n, \\ B^TA + A^TB = O_n, \end{array} \quad A, B \in GL(n; \mathbb{R}),$$

where ϑ and its transpose ϑ^T are the zero row *n*-vector and the zero column *n*-vector; I_n and O_n are the unit matrix and the zero matrix of size *n*, respectively.

1.1. **The structural tensor** F**.** The covariant derivatives of φ , ξ , η with respect to the Levi-Civita connection ∇ play a fundamental role in the differential geometry on the almost contact manifolds. The structural tensor F of type (0,3) on $(M, \varphi, \xi, \eta, g)$ is defined by

(3)
$$F(x,y,z) = g((\nabla_x \varphi)y,z).$$

It has the following properties:

(4)
$$F(x,y,z) = F(x,z,y) = F(x,\varphi y,\varphi z) + \eta(y)F(x,\xi,z) + \eta(z)F(x,y,\xi).$$

The relations of $\nabla \xi$ and $\nabla \eta$ with F are:

$$(\nabla_x \eta) y = g(\nabla_x \xi, y) = F(x, \varphi y, \xi).$$

The following 1-forms are associated with F:

(5)
$$\theta(z) = g^{ij}F(e_i, e_j, z), \quad \theta^*(z) = g^{ij}F(e_i, \varphi e_j, z), \quad \omega(z) = F(\xi, \xi, z),$$

where g^{ij} are the components of the inverse matrix of g with respect to a basis $\{e_i; \xi\}$ $(i=1,2,\ldots,2n)$ of the tangent space T_pM of M at an arbitrary point $p \in M$. Obviously, the equality $\omega(\xi) = 0$ and the following relation are always valid:

$$\theta^* \circ \varphi = -\theta \circ \varphi^2.$$

A classification of the almost contact B-metric manifolds with respect to F is given in [7]. This classification includes eleven basic classes $\mathcal{F}_1, \mathcal{F}_2, \ldots, \mathcal{F}_{11}$. Their intersection is the special class \mathcal{F}_0 determined by the condition F(x,y,z)=0. Hence \mathcal{F}_0 is the class of almost contact B-metric manifolds with ∇ -parallel structures, i.e. $\nabla \varphi = \nabla \xi = \nabla \eta = \nabla g = \nabla \tilde{g} = 0$.

Further we use the following characteristic conditions of the basic classes:

$$\mathcal{F}_{1}: \qquad F(x,y,z) = \frac{1}{2n} \left\{ g(x,\varphi y)\theta(\varphi z) + g(\varphi x,\varphi y)\theta(\varphi^{2}z) \right\}_{(y \leftrightarrow z)};$$

$$\mathcal{F}_{2}: \qquad F(\xi,y,z) = F(x,\xi,z) = 0, \qquad \underset{x,y,z}{\mathfrak{S}} F(x,y,\varphi z) = 0, \qquad \theta = 0;$$

$$\mathcal{F}_{3}: \qquad F(\xi,y,z) = F(x,\xi,z) = 0, \qquad \underset{x,y,z}{\mathfrak{S}} F(x,y,z) = 0;$$

$$\mathcal{F}_{4}: \qquad F(x,y,z) = -\frac{\theta(\xi)}{2n} \left\{ g(\varphi x,\varphi y)\eta(z) + g(\varphi x,\varphi z)\eta(y) \right\};$$

$$\mathcal{F}_{5}: \qquad F(x,y,z) = -\frac{\theta^{2}(\xi)}{2n} \left\{ g(x,\varphi y)\eta(z) + g(x,\varphi z)\eta(y) \right\};$$

$$\mathcal{F}_{6/7}: \qquad F(x,y,z) = F(x,y,\xi)\eta(z) + F(x,z,\xi)\eta(y),$$

$$\qquad F(x,y,\xi) = \pm F(y,x,\xi) = -F(\varphi x,\varphi y,\xi), \qquad \theta = \theta^{*} = 0;$$

$$\mathcal{F}_{8/9}: \qquad F(x,y,z) = F(x,y,\xi)\eta(z) + F(x,z,\xi)\eta(y),$$

$$\qquad F(x,y,\xi) = \pm F(y,x,\xi) = F(\varphi x,\varphi y,\xi);$$

$$\mathcal{F}_{10}: \qquad F(x,y,z) = F(\xi,\varphi y,\varphi z)\eta(x);$$

$$\mathcal{F}_{11}: \qquad F(x,y,z) = \eta(x) \left\{ \eta(y)\omega(z) + \eta(z)\omega(y) \right\},$$

where (for the sake of brevity) we use the following denotations: $\{A(x,y,z)\}_{(x\leftrightarrow y)}$ — instead of $\{A(x,y,z)+A(y,x,z)\}$ for any tensor A(x,y,z); $\mathfrak S$ — for the cyclic sum by three arguments; and the former and latter subscripts of $\mathfrak F_{i/j}$ correspond to upper and down signs plus or minus, respectively.

1.2. **The Nijenhuis tensor** N**.** An almost contact structure (φ, ξ, η) on M is called *normal* and respectively (M, φ, ξ, η) is a *normal almost contact manifold* if the corresponding almost complex structure J generated on $M' = M \times \mathbb{R}$ is integrable (i.e. M' is a complex manifold) [25]. The almost contact structure is normal if and only if the Nijenhuis tensor of (φ, ξ, η) is zero [1].

The Nijenhuis tensor N of the almost contact structure is defined by

$$N:=[\varphi,\varphi]+\mathrm{d}\eta\otimes\xi,$$

where $[\varphi, \varphi](x, y) = [\varphi x, \varphi y] + \varphi^2[x, y] - \varphi[\varphi x, y] - \varphi[x, \varphi y]$ and $d\eta$ is the exterior derivative of the 1-form η . Obviously, N is an antisymmetric tensor, i.e. N(x, y) = -N(y, x). Hence, using $[x, y] = \nabla_x y - \nabla_y x$ and $d\eta(x, y) = (\nabla_x \eta) y - (\nabla_y \eta) x$, the tensor N has the following form in terms of the covariant derivatives with respect to the Levi-Civita connection

 ∇ :

(8)
$$N(x,y) = (\nabla_{\varphi x} \varphi) y - \varphi (\nabla_x \varphi) y + (\nabla_x \eta) y \cdot \xi \\ - (\nabla_{\varphi y} \varphi) x + \varphi (\nabla_y \varphi) x - (\nabla_y \eta) x \cdot \xi.$$

The corresponding Nijenhuis tensor of type (0,3) on $(M, \varphi, \xi, \eta, g)$ is defined by N(x,y,z) = g(N(x,y),z). Then, from (8) and (3), we have

(9)
$$N(x,y,z) = \left\{ F(\boldsymbol{\varphi}x,y,z) - F(x,y,\boldsymbol{\varphi}z) + F(x,\boldsymbol{\varphi}y,\boldsymbol{\xi})\boldsymbol{\eta}(z) \right\}_{[x \leftrightarrow y]}.$$

where we use the denotation $\{A(x,y,z)\}_{[x\leftrightarrow y]}$ instead of $\{A(x,y,z)-A(y,x,z)\}$ for any tensor A(x,y,z).

Lemma 1. The Nijenhuis tensor on an arbitrary almost B-metric manifold has the following properties:

$$N(\varphi x, \varphi y, \varphi z) = -N(\varphi^2 x, \varphi^2 y, \varphi z) = N(\varphi x, \varphi^2 y, \varphi^2 z),$$

$$N(\varphi^2 x, \varphi y, \varphi z) = N(\varphi x, \varphi^2 y, \varphi z) = -N(\varphi x, \varphi y, \varphi^2 z).$$

Proof. Bearing in mind properties (4) of F and relation (9), the equalities in the first line of the statement follow. They imply the equalities in the last line by virtue of (1) and (2). \Box

Lemma 2. The class $U_0 = \mathcal{F}_1 \oplus \mathcal{F}_2 \oplus \mathcal{F}_4 \oplus \mathcal{F}_5 \oplus \mathcal{F}_6 \oplus \mathcal{F}_8 \oplus \mathcal{F}_9 \oplus \mathcal{F}_{10} \oplus \mathcal{F}_{11}$ of the almost contact B-metric manifolds is determined by the condition $N(\boldsymbol{\varphi}, \boldsymbol{\varphi}) = 0$.

Proof. The statement follows from the following form of the tensor N for each of the basic classes \mathcal{F}_i (i = 1, 2, ..., 11) of $M = (M, \varphi, \xi, \eta, g)$:

$$(10) \begin{array}{ll} N(x,y) = 0, & M \in \mathcal{F}_1 \oplus \mathcal{F}_2 \oplus \mathcal{F}_4 \oplus \mathcal{F}_5 \oplus \mathcal{F}_6; \\ N(x,y) = 2 \left\{ \left(\nabla_{\varphi x} \varphi \right) y - \varphi \left(\nabla_x \varphi \right) y \right\}, & M \in \mathcal{F}_3; \\ N(x,y) = 4 \left(\nabla_x \eta \right) y \cdot \xi, & M \in \mathcal{F}_7; \\ N(x,y) = 2 \left\{ \eta \left(x \right) \nabla_y \xi - \eta \left(y \right) \nabla_x \xi \right\}, & M \in \mathcal{F}_8 \oplus \mathcal{F}_9; \\ N(x,y) = -\eta \left(x \right) \varphi \left(\nabla_\xi \varphi \right) y + \eta \left(y \right) \varphi \left(\nabla_\xi \varphi \right) x, & M \in \mathcal{F}_{10}; \\ N(x,y) = \left\{ \eta \left(x \right) \omega (\varphi y) - \eta \left(y \right) \omega (\varphi x) \right\}, & M \in \mathcal{F}_{11}. \end{array}$$

The calculations are made using (9) and (7).

2. φ-Canonical Connection

Definition 1. A linear connection D is called a natural connection on the manifold $(M, \varphi, \xi, \eta, g)$ if the almost contact structure (φ, ξ, η) and the B-metric g are parallel with respect to D, i.e. $D\varphi = D\xi = D\eta = Dg = 0$.

As a corollary, the associated metric \tilde{g} is also parallel with respect to a natural connection D on $(M, \varphi, \xi, \eta, g)$.

According to [22], a necessary and sufficient condition a linear connection D to be natural on $(M, \varphi, \xi, \eta, g)$ is $D\varphi = Dg = 0$.

If *T* is the torsion of *D*, i.e. $T(x,y) = D_x y - D_y x - [x,y]$, then the corresponding tensor of type (0,3) is determined by T(x,y,z) = g(T(x,y),z).

Let us denote the difference between the natural connection D and the Levi-Civita connection ∇ of g by $Q(x,y) = D_x y - \nabla_x y$ and the corresponding tensor of type (0,3) — by Q(x,y,z) = g(Q(x,y),z).

It is easy to establish (see, e.g. [19]) that a linear connection *D* is a natural connection on an almost contact B-metric manifold if and only if

(11)
$$Q(x,y,\varphi z) - Q(x,\varphi y,z) = F(x,y,z), \qquad Q(x,y,z) = -Q(x,z,y).$$

Therefore, according T(x,y) = Q(x,y) - Q(y,x), we have the equality of Hayden's theorem [13]

$$Q(x,y,z) = \frac{1}{2} \left\{ T(x,y,z) - T(y,z,x) + T(z,x,y) \right\}.$$

Definition 2. A natural connection D is called a φ -canonical connection on the manifold $(M, \varphi, \xi, \eta, g)$ if the torsion tensor T of D satisfies the following identity:

(12)
$$\left\{ T(x, y, z) - T(x, \varphi y, \varphi z) - \eta(x) \left\{ T(\xi, y, z) - T(\xi, \varphi y, \varphi z) \right\} - \eta(y) \left\{ T(x, \xi, z) - T(x, z, \xi) - \eta(x) T(z, \xi, \xi) \right\} \right\}_{[y \leftrightarrow z]} = 0.$$

Let us remark that the restriction of the φ -canonical connection D of the manifold $(M, \varphi, \xi, \eta, g)$ on the contact distribution H is the unique canonical connection of the corresponding almost complex manifold with Norden metric, studied in [6].

In [21], it is introduced a natural connection on $(M, \varphi, \xi, \eta, g)$, defined by

(13)
$$\nabla_x^0 y = \nabla_x y + Q^0(x, y),$$

where $Q^0(x,y) = \frac{1}{2} \{ (\nabla_x \varphi) \varphi y + (\nabla_x \eta) y \cdot \xi \} - \eta(y) \nabla_x \xi$. Therefore, we have

(14)
$$Q^{0}(x,y,z) = \frac{1}{2} \left\{ F(x,\varphi y,z) + \eta(z)F(x,\varphi y,\xi) - 2\eta(y)F(x,\varphi z,\xi) \right\}.$$

The torsion of the φ B-connection has the following form

(15)
$$T^{0}(x,y,z) = \frac{1}{2} \left\{ F(x,\varphi y,z) + \eta(z) F(x,\varphi y,\xi) + 2\eta(x) F(y,\varphi z,\xi) \right\}_{[x \leftrightarrow y]}.$$

In [23], the connection determined by (13) is called a φ *B-connection*. It is studied for some classes of $(M, \varphi, \xi, \eta, g)$ in [21, 17, 18, 23]. The restriction of the φ B-connection on H coincides with the B-connection on an almost complex manifold with Norden metric, studied for the class of the locally conformal Kählerian manifolds with Norden metric in [4].

We construct a linear connection ∇' as follows:

(16)
$$g(\nabla'_x y, z) = g(\nabla_x y, z) + Q'(x, y, z),$$

where

(17)
$$Q'(x,y,z) = Q^{0}(x,y,z) - \frac{1}{8} \left\{ N(\varphi^{2}z, \varphi^{2}y, \varphi^{2}x) + 2N(\varphi z, \varphi y, \xi) \eta(x) \right\}.$$

By direct computations, we check that ∇' satisfies conditions (11) and therefore it is a natural connection on $(M, \varphi, \xi, \eta, g)$. Its torsion is

(18)
$$T'(x,y,z) = T^{0}(x,y,z) - \frac{1}{8} \left\{ N(\varphi^{2}z, \varphi^{2}y, \varphi^{2}x) + 2N(\varphi z, \varphi y, \xi) \eta(x) \right\}_{[x \leftrightarrow y]}.$$

We verify immediately that T' satisfies (12) and thus ∇' , determined by (16) and (17), is a φ -canonical connection on $(M, \varphi, \xi, \eta, g)$.

The explicit expression (16), supported by (14) and (9), of the φ -canonical connection by the tensor F implies that the φ -canonical connection is unique.

Immediately we get the following

Proposition 3. A necessary and sufficient condition the φ -canonical connection to coincide with the φ B-connection is $N(\varphi, \varphi) = 0$.

Thus, Proposition 3 and Lemma 2 imply

Corollary 4. The φ -canonical connection and the φ B-connection coincide on an almost contact B-metric manifold $(M, \varphi, \xi, \eta, g)$ if and only if $(M, \varphi, \xi, \eta, g)$ is in the class \mathcal{U}_0 .

In [22], it is given a classification of the linear connections on the almost contact B-metric manifolds with respect to their torsion tensors T in 11 classes \mathfrak{T}_{ij} . The characteristic conditions of these basic classes are the following:

$$\begin{split} &\mathfrak{I}_{11/12}: \quad T(\xi,y,z) = T(x,y,\xi) = 0, \quad T(x,y,z) = -T(\varphi x,\varphi y,z) = \mp T(x,\varphi y,\varphi z); \\ &\mathfrak{I}_{13}: \quad T(\xi,y,z) = T(x,y,\xi) = 0, \quad T(x,y,z) - T(\varphi x,\varphi y,z) = \mathop{\mathfrak{S}}_{x,y,z} T(x,y,z) = 0; \\ &\mathfrak{I}_{14}: \quad T(\xi,y,z) = T(x,y,\xi) = 0, \quad T(x,y,z) - T(\varphi x,\varphi y,z) = \mathop{\mathfrak{S}}_{x,y,z} T(\varphi x,y,z) = 0; \\ &\mathfrak{I}_{21/22}: \quad T(x,y,z) = \eta(z)T(\varphi^2 x,\varphi^2 y,\xi), \quad T(x,y,\xi) = \mp T(\varphi x,\varphi y,\xi); \\ &\mathfrak{I}_{31/32}: \quad T(x,y,z) = \eta(x)T(\xi,\varphi^2 y,\varphi^2 z) - \eta(y)T(\xi,\varphi^2 x,\varphi^2 z), \\ & \qquad T(\xi,y,z) = \pm T(\xi,z,y) = -T(\xi,\varphi y,\varphi z); \\ &\mathfrak{I}_{33/34}: \quad T(x,y,z) = \eta(x)T(\xi,\varphi^2 y,\varphi^2 z) - \eta(y)T(\xi,\varphi^2 x,\varphi^2 z), \\ & \qquad T(\xi,y,z) = \pm T(\xi,z,y) = T(\xi,\varphi y,\varphi z); \\ &\mathfrak{I}_{41}: \quad T(x,y,z) = \eta(z)\left\{\eta(y)\hat{t}(x) - \eta(x)\hat{t}(y)\right\}. \end{split}$$

3. GENERAL CONTACTLY CONFORMAL GROUP G

In this section we consider the group of transformations of the φ -canonical connection generated by the general contactly conformal transformations of the almost contact B-metric structure.

Let $(M, \varphi, \xi, \eta, g)$ be an almost contact B-metric manifold. The general contactly conformal transformations of the almost contact B-metric structure are defined by

(19)
$$\bar{\xi} = e^{-w}\xi$$
, $\bar{\eta} = e^{w}\eta$, $\bar{g}(x,y) = \alpha g(x,y) + \beta g(x,\varphi y) + (\gamma - \alpha)\eta(x)\eta(y)$,

where $\alpha = e^{2u}\cos 2v$, $\beta = e^{2u}\sin 2v$, $\gamma = e^{2w}$ for differentiable functions u, v, w on M [18]. These transformations form a group denoted by G.

If w = 0, we obtain the contactly conformal transformations of the B-metric, introduced in [20]. By v = w = 0, the transformations (19) are reduced to the usual conformal transformations of g.

Let us remark that G can be considered as a complex conformal gauge group, i.e. the composition of an almost contact group, preserving H and a complex conformal transformation of the complex Riemannian metric $\overline{g^{\mathbb{C}}} = e^{2(u+iv)}g^{\mathbb{C}}$ on H.

Let $(M, \varphi, \xi, \eta, g)$ and $(M, \varphi, \bar{\xi}, \bar{\eta}, \bar{g})$ be contactly conformally equivalent with respect to a transformation from G. The Levi-Civita connection of \bar{g} is denoted by $\bar{\nabla}$. Using the formula

$$2g(\nabla_x y, z) = xg(y, z) + yg(x, z) - zg(x, y) + g([x, y], z) + g([z, x], y) + g([z, y], x),$$

by straightforward computations we get the following relation between ∇ and $\bar{\nabla}$:

$$2\left(\alpha^{2}+\beta^{2}\right)g\left(\bar{\nabla}_{x}y-\nabla_{x}y,z\right)=$$

$$=\frac{1}{2}\left\{-\alpha\beta\left[2F(x,y,\varphi^{2}z)-F(\varphi^{2}z,x,y)\right]-\beta^{2}\left[2F(x,y,\varphi z)-F(\varphi z,x,y)\right]\right.$$

$$\left.+\frac{\beta}{\gamma}\left(\alpha^{2}+\beta^{2}\right)\left[2F(x,y,\xi)-F(\xi,x,y)\right]\eta(z)\right.$$

$$\left.+2\left(\frac{\alpha}{\gamma}-1\right)\left(\alpha^{2}+\beta^{2}\right)F(\varphi^{2}x,\varphi y,\xi)\eta(z)\right.$$

$$\left.+2\alpha(\gamma-\alpha)\left[F(x,\varphi z,\xi)+F(\varphi^{2}z,\varphi x,\xi)\right]\eta(y)\right.$$

$$\left.-2\beta(\gamma-\alpha)\left[F(x,\varphi^{2}z,\xi)-F(\varphi z,\varphi x,\xi)\right]\eta(y)\right.$$

$$\left.-2\left[\alpha\mathrm{d}\alpha(x)+\beta\mathrm{d}\beta(x)\right]g(\varphi y,\varphi z)+2\left[\alpha\mathrm{d}\beta(x)-\beta\mathrm{d}\alpha(x)\right]g(y,\varphi z)\right.$$

$$\left.-\left[\alpha\mathrm{d}\alpha(\varphi^{2}z)+\beta\mathrm{d}\alpha(\varphi z)\right]g(\varphi x,\varphi y)+\left[\alpha\mathrm{d}\beta(\varphi^{2}z)+\beta\mathrm{d}\beta(\varphi z)\right]g(x,\varphi y)\right.$$

$$\left.+\left[\alpha\mathrm{d}\gamma(\varphi^{2}z)+\beta\mathrm{d}\gamma(\varphi z)\right]\eta(x)\eta(y)\right.$$

$$\left.+\frac{1}{\gamma}(\alpha^{2}+\beta^{2})\left\{\mathrm{d}\alpha(\xi)g(\varphi x,\varphi y)-\mathrm{d}\beta(\xi)g(x,\varphi y)\right\}\eta(z)\right.$$

$$\left.+\frac{1}{\gamma}(\alpha^{2}+\beta^{2})\left\{2\mathrm{d}\gamma(x)\eta(y)-\mathrm{d}\gamma(\xi)\eta(x)\eta(y)\right\}\eta(z)\right\}_{(x\leftrightarrow y)}.$$

Using (3) and (20), we obtain the following formula for the transformation by G of the tensor F:

$$2\bar{F}(x,y,z) = 2\alpha F(x,y,z) + \left\{ \beta \left\{ F(\varphi y,z,x) - F(y,\varphi z,x) + F(x,\varphi y,\xi) \eta(z) \right\} \right.$$

$$\left. + (\gamma - \alpha) \left\{ \left[F(x,y,\xi) + F(\varphi y,\varphi x,\xi) \right] \eta(z) + \left[F(y,z,\xi) + F(\varphi z,\varphi y,\xi) \right] \eta(x) \right\} \right.$$

$$\left. - \left[\mathrm{d}\alpha(\varphi y) + \mathrm{d}\beta(y) \right] g(\varphi x,\varphi z) - \left[\mathrm{d}\alpha(y) - \mathrm{d}\beta(\varphi y) \right] g(x,\varphi z) \right.$$

$$\left. + \eta(x) \eta(y) \mathrm{d}\gamma(\varphi z) \right\}_{(y \leftrightarrow z)}.$$

Theorem 5. The tensor $N(\phi \cdot, \phi \cdot)$ is an invariant of the group G on any almost contact B-metric manifold.

Proof. From (9), (21), (4) and (19), it follows the formula for the transformation by G of the Nijenhuis tensor:

(22)
$$\bar{N}(\varphi x, \varphi y, z) = \alpha N(\varphi x, \varphi y, z) + \beta N(\varphi x, \varphi y, \varphi z) + (\gamma - \alpha) N(\varphi x, \varphi y, \xi) \eta(z).$$

Thus, bearing in mind (19), we obtain immediately $\bar{N}(\varphi x, \varphi y) = N(\varphi x, \varphi y)$.

According to Lemma 2, we establish immediately the following

Corollary 6. The class \mathcal{U}_0 is closed by the action of the group G.

Proposition 7. Let the almost contact B-metric manifolds $(M, \varphi, \xi, \eta, g)$ and $(M, \varphi, \bar{\xi}, \bar{\eta}, \bar{g})$ be contactly conformally equivalent with respect to a transformation from G. Then the

corresponding φ -canonical connections $\bar{\nabla}'$ and ∇' as well as their torsions \bar{T}' and T' are related as follows:

(23)
$$\bar{\nabla}'_{x}y = \nabla'_{x}y - du(x)\varphi^{2}y + dv(x)\varphi y + dw(x)\eta(y)\xi$$

$$+ \frac{1}{2} \left\{ \left[du(\varphi^{2}y) - dv(\varphi y) \right] \varphi^{2}x - \left[du(\varphi y) + dv(\varphi^{2}y) \right] \varphi x$$

$$- g(\varphi x, \varphi y) \left[\varphi^{2}p - \varphi q \right] + g(x, \varphi y) \left[\varphi p + \varphi^{2}q \right] \right\},$$

$$\bar{T}'(x, y) = T'(x, y) + \frac{1}{2} \left\{ 2dw(x)\eta(y)\xi$$

$$+ \left[du(\varphi^{2}x) + dv(\varphi x) - 2du(\xi)\eta(x) \right] \varphi^{2}y$$

$$+ \left[du(\varphi x) - dv(\varphi^{2}x) + 2dv(\xi)\eta(x) \right] \varphi y \right\}_{[x \leftrightarrow y]},$$

where p = gradu, q = gradv.

Proof. Taking into account (13), we have the following equality on $(M, \varphi, \xi, \eta, g)$:

(25)
$$g\left(\nabla_{x}^{0}y - \nabla_{x}y, z\right) = \frac{1}{2}\left\{F(x, \varphi y, z) + F(x, \varphi y, \xi)\eta(z) - 2F(x, \varphi z, \xi)\eta(y)\right\}.$$

Then we can rewrite the corresponding equality on the manifold $(M, \varphi, \bar{\xi}, \bar{\eta}, \bar{g})$, which is the image of $(M, \varphi, \xi, \eta, g)$ by a transformation from G:

(26)
$$\bar{g}\left(\bar{\nabla}_{x}^{0}y - \bar{\nabla}_{x}y, z\right) = \frac{1}{2}\left\{\bar{F}(x, \varphi y, z) + \bar{F}(x, \varphi y, \bar{\xi})\bar{\eta}(z) - 2\bar{F}(x, \varphi z, \bar{\xi})\bar{\eta}(y)\right\}.$$

By virtue of (25), (26), (21), (20), we get the following formula of the transformation by G of the φ B-connection:

$$g\left(\overline{\nabla}_{x}^{0}y - \nabla_{x}^{0}y, z\right) = \frac{1}{8}\sin 4\nu N(\varphi z, \varphi y, \varphi x) - \frac{1}{4}\sin^{2} 2\nu N(\varphi^{2}z, \varphi^{2}y, \varphi^{2}x)$$

$$-\frac{1}{4}e^{2(w-u)}\sin 2\nu N(\varphi^{2}z, \varphi y, \xi)\eta(x) - \frac{1}{4}\left(1 - e^{2(w-u)}\cos 2\nu\right)N(\varphi z, \varphi y, \xi)\eta(x)$$

$$-du(x)g(\varphi y, \varphi z) + dv(x)g(y, \varphi z) + dw(x)\eta(y)\eta(z)$$

$$+\frac{1}{2}\left[du(\varphi^{2}y) - dv(\varphi y)\right]g(\varphi x, \varphi z) - \frac{1}{2}\left[du(\varphi y) + dv(\varphi^{2}y)\right]g(x, \varphi z)$$

$$-\frac{1}{2}\left[du(\varphi^{2}z) - dv(\varphi z)\right]g(\varphi x, \varphi y) + \frac{1}{2}\left[du(\varphi z) + dv(\varphi^{2}z)\right]g(x, \varphi y).$$

Taking into account (17), (22), (19) and (27), we get (23). As a consequence of (23), the torsions T' and \bar{T}' of ∇' and $\bar{\nabla}'$, respectively, are related as in (24).

The torsion forms associated with T' of the φ -canonical connection are defined, in a similar way of (5), as follows:

(28)
$$t'(x) = g^{ij}T'(x, e_i, e_j), \quad t'^*(x) = g^{ij}T'(x, e_i, \varphi e_j), \quad \hat{t}'(x) = T'(x, \xi, \xi).$$

Obviously, $\hat{t}(\xi) = 0$ is always valid.

Using (28), (18), (15), (4) and Lemma 1, we obtain that the torsion forms of the φ -canonical connection are expressed by the associated forms with F:

(29)
$$t' = \frac{1}{2} \{ \theta^* + \theta^*(\xi) \eta \}, \quad t'^* = -\frac{1}{2} \{ \theta + \theta(\xi) \eta \}, \quad \hat{t}' = -\omega \circ \varphi.$$

The equality (6) and (29) imply the following relation:

$$(30) t'^* \circ \varphi = -t' \circ \varphi^2.$$

4. General Contactly Conformal Subgroup G_0

Let us consider the subgroup G_0 of G defined by the conditions

(31)
$$du \circ \varphi^2 + dv \circ \varphi = du \circ \varphi - dv \circ \varphi^2 y = du(\xi) = dv(\xi) = dw \circ \varphi = 0.$$

By direct computations, from (7), (19), (21) and (31), we prove that each of the basic classes \mathcal{F}_i ($i=1,2,\ldots,11$) of the almost contact B-metric manifolds is closed by the action of the group G_0 . Moreover, G_0 is the largest subgroup of G preserving the 1-forms θ , θ^* , ω and the special class \mathcal{F}_0 .

Theorem 8. The torsion of the φ -canonical connection is invariant with respect to the general contactly conformal transformations if and only if these transformations belong to the group G_0 .

Proof. Proposition 7 and (31) imply immediately

(32)
$$\bar{\nabla}'_x y = \nabla'_x y - \mathrm{d}u(x)\varphi^2 y + \mathrm{d}v(x)\varphi y + \mathrm{d}w(\xi)\eta(x)\eta(y)\xi$$

$$- \mathrm{d}u(y)\varphi^2 x + \mathrm{d}v(y)\varphi x + g(\varphi x, \varphi y)p - g(x, \varphi y)q.$$

The statement follows from (32), or alternatively from (24) and (31).

Bearing in mind the invariance of \mathcal{F}_i $(i=1,2,\ldots,11)$ and T' with respect to the transformations of G_0 , we establish that each of the eleven basic classes of the manifolds $(M, \varphi, \xi, \eta, g)$ is characterized by the torsion of the φ -canonical connection. Then we give this characterization in the following

Proposition 9. The basic classes of the almost contact B-metric manifolds are characterized by conditions for the torsion of the φ -canonical connection as follows:

```
\begin{array}{ll} \mathcal{F}_1: & T'(x,y) = \frac{1}{2n} \left\{ t'(\varphi^2 x) \varphi^2 y - t'(\varphi^2 y) \varphi^2 x + t'(\varphi x) \varphi y - t'(\varphi y) \varphi x \right\}; \\ \mathcal{F}_2: & T'(\xi,y) = 0, \ \eta \left( T'(x,y) \right) = 0, \ T'(x,y) = T'(\varphi x, \varphi y), \ t' = 0; \\ \mathcal{F}_3: & T'(\xi,y) = 0, \ \eta \left( T'(x,y) \right) = 0, \ T'(x,y) = \varphi T'(x,\varphi y); \\ \mathcal{F}_4: & T'(x,y) = \frac{1}{2n} t'^*(\xi) \left\{ \eta(y) \varphi x - \eta(x) \varphi y \right\}; \\ \mathcal{F}_5: & T'(x,y) = \frac{1}{2n} t'(\xi) \left\{ \eta(y) \varphi^2 x - \eta(x) \varphi^2 y \right\}; \\ \mathcal{F}_6: & T'(x,y) = \eta(x) T'(\xi,y) - \eta(y) T'(\xi,x), \\ & T'(\xi,y,z) = T'(\xi,z,y) = -T'(\xi,\varphi y,\varphi z); \\ \mathcal{F}_{7/8}: & T'(x,y) = \eta(x) T'(\xi,y) - \eta(y) T'(\xi,x) + \eta(T'(x,y)) \xi, \\ & T'(\xi,y,z) = -T'(\xi,z,y) = \mp T'(\xi,\varphi y,\varphi z) = \frac{1}{2} T'(y,z,\xi) = \mp \frac{1}{2} T'(\varphi y,\varphi z,\xi); \\ \mathcal{F}_{9/10}: & T'(x,y) = \eta(x) T'(\xi,y) - \eta(y) T'(\xi,x), \\ & T'(\xi,y,z) = \pm T'(\xi,z,y) = T'(\xi,\varphi y,\varphi z); \\ \mathcal{F}_{11}: & T'(x,y) = \left\{ \hat{t'}(x) \eta(y) - \hat{t'}(y) \eta(x) \right\} \xi. \end{array}
```

Proof. According to Proposition 3, Corollary 4, (15) and (18), we have the following form of the torsion of the φ -canonical connection when $(M, \varphi, \xi, \eta, g)$ belongs to the classes \mathcal{F}_i $(i \in \{1, 2, ..., 11\}; i \neq 3, 7)$:

$$T'(x,y) = T^{0}(x,y) = \frac{1}{2} \left\{ (\nabla_{x} \varphi) \varphi y + (\nabla_{x} \eta) y \cdot \xi + 2 \eta(x) \nabla_{y} \xi \right\}_{[x \leftrightarrow y]}.$$

For the classes \mathcal{F}_3 and \mathcal{F}_7 , we use (18) and equalities (10).

Then, using (4), (29), (30) and (7), we obtain the characteristics in the statement.

According to the classification of the torsion tensors in [22] and Proposition 9, we get the following

Proposition 10. Let T' be the torsion of the φ -canonical connection on an almost contact B-metric manifold $M = (M, \varphi, \xi, \eta, g)$. The correspondence between the classes \mathfrak{F}_i of M and the classes \mathfrak{T}_{ik} of T' is given as follows:

$$\begin{array}{ll} \textit{M} \in \mathcal{F}_1 \ \Leftrightarrow \ \textit{T}' \in \mathcal{T}_{13}, \ t' \neq 0; \\ \textit{M} \in \mathcal{F}_2 \ \Leftrightarrow \ \textit{T}' \in \mathcal{T}_{13}, \ t' = 0; \\ \textit{M} \in \mathcal{F}_3 \ \Leftrightarrow \ \textit{T}' \in \mathcal{T}_{12}; \\ \textit{M} \in \mathcal{F}_3 \ \Leftrightarrow \ \textit{T}' \in \mathcal{T}_{31}, \ t' = 0, \ t'^* \neq 0; \\ \textit{M} \in \mathcal{F}_4 \ \Leftrightarrow \ \textit{T}' \in \mathcal{T}_{31}, \ t' = 0, \ t'^* \neq 0; \\ \textit{M} \in \mathcal{F}_5 \ \Leftrightarrow \ \textit{T}' \in \mathcal{T}_{31}, \ t' = 0, \ t'^* = 0; \\ \textit{M} \in \mathcal{F}_6 \ \Leftrightarrow \ \textit{T}' \in \mathcal{T}_{31}, \ t' = 0, \ t'^* = 0; \\ \textit{M} \in \mathcal{F}_{11} \ \Leftrightarrow \ \textit{T}' \in \mathcal{T}_{41}. \\ \end{array}$$

5. THE RELATION BETWEEN THE ALMOST CONTACT B-METRIC MANIFOLDS AND THE KÄHLERIAN MANIFOLDS WITH NORDEN METRIC

Firstly, let us consider the manifold $M^\times = M \times \mathbb{R}$, where $(M, \varphi, \xi, \eta, g)$ is a (2n+1)-dimensional almost contact B-metric manifold. The almost complex structure on M^\times is defined (as in [1]) by $J(x, a\partial_t) = (\varphi x - a\xi, \eta(x)\partial_t)$, where $x \in \mathfrak{X}(M)$, t is the coordinate on \mathbb{R} , $\partial_t = \frac{\mathrm{d}}{\mathrm{d}t}$ and a is a differentiable function on M^\times . Let us consider the product metric $g^\times(x^\times,y^\times) = g(x,y) - ab$ for $x^\times = (x,a\partial_t)$, $y^\times = (y,b\partial_t)$ on M^\times . Since g is a B-metric, then g^\times is a Norden metric, i.e. $g^\times(Jx^\times,Jy^\times) = -g^\times(x^\times,y^\times)$. Thus (M^\times,J,g^\times) is an almost Norden manifold. We consider M as a hypersurface of M^\times . Then the Gauss-Weingarten equations are $\nabla_x^\times y = \nabla_x y - g(Ax,y)\partial_t$, $\nabla_x^\times \partial_t = -Ax$, where ∇^\times is the Levi-Civita connection for g^\times and A is its Weingarten map. Then we have

$$\begin{split} &(\nabla_x^\times J)\,y = (\nabla_x \varphi)\,y - \eta(y)Ax - g(Ax,y)\xi + \{(\nabla_x \eta)\,y - g(Ax,\varphi y)\}\,\partial_t, \quad \left(\nabla_{\partial_t}^\times J\right)\partial_t = A\xi\,, \\ &(\nabla_x^\times J)\,\partial_t = -\nabla_x \xi + \varphi Ax + 2\eta(Ax)\partial_t, \quad \left(\nabla_{\partial_t}^\times J\right)y = \varphi Ay - A\varphi y + \eta(Ay)\partial_t. \end{split}$$

If we set $(M^{\times},J,g^{\times})$ to be a Kählerian manifold with Norden metric, i.e. $\nabla^{\times}J=0$, we obtain

$$(\nabla_x \varphi) y = \eta(y) Ax + g(Ax, y) \xi,$$
 $(\nabla_x \eta) y = g(Ax, \varphi y),$ $\nabla_x \xi = \varphi Ax,$ $\varphi \circ A = A \circ \varphi,$ $\eta(Ax) = 0,$ $A\xi = 0.$

Therefore we have

$$F(x, y, z) = F(x, y, \xi) \eta(z) + F(x, z, \xi) \eta(y), \quad F(x, y, \xi) = F(y, x, \xi) = -F(\varphi x, \varphi y, \xi).$$

Thus, bearing in mind (7), the manifold $(M, \varphi, \xi, \eta, g)$ belongs to the class $\mathcal{F}_4 \oplus \mathcal{F}_5 \oplus \mathcal{F}_6$, which may be viewed as the class of the almost contact B-metric manifolds of Sasakian type as an analogy to the contact metric geometry [1].

Secondly, in [7] it is given an example of the considered manifolds as follows. Let the vector space $\mathbb{R}^{2n+2} = \left\{ (u^1,\dots,u^{n+1};v^1,\dots,v^{n+1}) \mid u^i,v^i \in \mathbb{R} \right\}$ be considered as a complex Riemannian manifold with the canonical complex structure J and the metric g defined by $g(x,x) = -\delta_{ij}\lambda^i\lambda^j + \delta_{ij}\mu^i\mu^j$ for $x = \lambda^i\frac{\partial}{\partial u^i} + \mu^i\frac{\partial}{\partial v^i}$. Identifying the point $p \in \mathbb{R}^{2n+2}$ with its position vector it is considered the time-like sphere S:g(n,n)=-1 of g in \mathbb{R}^{2n+2} , where n is the unit normal to the tangent space T_pS at $p \in S$. It is set $g(n,Jn)=\tan \psi$, $\psi \in \left(-\frac{\pi}{2},\frac{\pi}{2}\right)$. Then the almost contact structure is introduced by $\xi=\sin \psi.n+\cos \psi.Jn$, $\eta=g(\cdot,\xi)$, $\varphi=J-\eta\otimes J\xi$. It is shown that (S,φ,ξ,η,g) is an almost contact B-metric manifold in the class $\mathcal{F}_4\oplus\mathcal{F}_5$.

Since the φ -canonical connection coincides with the φ B-connection on any manifold in $\mathcal{F}_4 \oplus \mathcal{F}_5$, according to Corollary 4, then by virtue of (15) we get the torsion tensor and the

torsion forms of the φ -canonical connection as follows:

$$T'(x,y,z) = \left\{ \eta(x) \left\{ \cos \psi g(y,\varphi z) - \sin \psi g(\varphi y,\varphi z) \right\} \right\}_{[x \leftrightarrow y]},$$

$$t' = 2n \sin \psi \eta, \qquad t'^* = -2n \cos \psi \eta, \qquad \hat{t}' = 0.$$

These equalities are in accordance with Proposition 9. Moreover, it follows that the statement $T' \in \mathcal{T}_{31}$ is valid, which confirms Proposition 10.

REFERENCES

- [1] Blair, D. E.: Riemannian Geometry of Contact and Symplectic Manifolds. Progress in Mathematics 203, Birkhäuser, Boston (2002)
- [2] Friedrich, T., Ivanov, S.: Parallel spinors and connections with skew-symmetric torsion in string theory. Asian J. Math. 6, 303–336 (2002)
- [3] Friedrich, T., Ivanov, S.: Almost contact manifolds, connections with torsion, and parallel spinors. J. Reine Angew. Math. 559, 217–236 (2003)
- [4] Ganchev, G., Gribachev, K., Mihova, V.: B-connections and their conformal invariants on conformally Kaehler manifolds with B-metric. Publ. Inst. Math. (Beograd) (N.S.) 42(56), 107–121 (1987)
- [5] Ganchev, G., Ivanov, S.: Characteristic curvatures on complex Riemannian manifolds. Riv. Mat. Univ. Parma (5) 1, 155–162 (1992)
- [6] Ganchev, G., Mihova, V.: Canonical connection and the canonical conformal group on an almost complex manifold with B-metric. Ann. Univ. Sofia Fac. Math. Inform. 81, 195–206 (1987)
- [7] Ganchev, G., Mihova, V., Gribachev, K.: Almost contact manifolds with B-metric. Math. Balkanica (N.S.) 7 (3-4), 261–276 (1993)
- [8] Gates, S. J., Hull, C. M., Roček, M.: Twisted multiplets and new supersymmetric non-linear σ -models. Nucl. Phys. B 248, 157–186 (1984)
- [9] Gauduchon, P.: Hermitian connections and Dirac operators. Boll. Unione Mat. Ital. 11, 257–288 (1997)
- [10] Gribacheva, D.: Natural connections on Riemannian product manifolds. Compt. rend. Acad. bulg. Sci. 64 (6), 799–806 (2011)
- [11] Gribacheva, D.: Natural connections on conformal Riemannian P-manifolds. Compt. rend. Acad. bulg. Sci. 65 (5), 581–590 (2012)
- [12] Gribacheva, D., Mekerov, D.: Canonical connection on a class of Riemannian almost product manifolds. J. Geom. 102 (1-2), 53–71 (2011)
- [13] Hayden, H.: Subspaces of a space with torsion. Proc. London Math. Soc. 34, 27–50 (1934)
- [14] Ivanov, S., Papadopoulos, G.: Vanishing theorems and string backgrounds. Classical Quant. Grav. 18, 1089– 1110 (2001)
- [15] Lichnerowicz, A.: Un théorème sur les espaces homogènes complexes. Arch. Math. 5, 207-215 (1954)
- [16] Lichnerowicz, A.: Généralisation de la géométrie kählérienne globale. Coll. de Géom. Diff. Louvain, 99– 122 (1955)
- [17] Maney, M.: Properties of curvature tensors on almost contact manifolds with B-metric. Proc. of Jubilee Sci. Session of Vassil Levsky Higher Mil. School, Veliko Tarnovo, 27, 221–227 (1993)
- [18] Manev, M.: Contactly conformal transformations of general type of almost contact manifolds with B-metric. Applications. Math. Balkanica (N.S.) 11 (3-4), 347–357 (1997)
- [19] Maney, M.: A connection with totally skew-symmetric torsion on almost contact manifolds with B-metric. Int. J. Geom. Methods Mod. Phys. 9 (5), 1250044 (20 pages) (2012)
- [20] Manev, M., Gribachev, K.: Contactly conformal transformations of almost contact manifolds with B-metric. Serdica Math. J. 19, 287–299 (1993)
- [21] Maney, M., Gribachev, K.: Conformally invariant tensors on almost contact manifolds with B-metric. Serdica Math. J. 20, 133–147 (1994)
- [22] Manev, M., Ivanova, M.: A classification of the torsion tensors on almost contact manifolds with B-metric. arXiv:1105.5715
- [23] Manev, M., Ivanova, M.: A natural connection on some classes of almost contact manifolds with B-metric. Compt. rend. Acad. bulg. Sci. 65 (4), 429–436 (2012)
- [24] Mekerov, D.: Canonical connection on quasi-Kähler manifolds with Norden metric. J. Tech. Univ. Plovdiv Fundam. Sci. Appl. Ser. A Pure Appl. Math. 14, 73–86 (2009)
- [25] Sasaki, S., Hatakeyama, Y.: On differentiable manifolds with certain structures which are closely related to almost contact structures II, Tôhoku Math. J. 13, 281–294 (1961)
- [26] Strominger, A.: Superstrings with torsion. Nucl. Phys. B 274, 253-284 (1986)

University of Plovdiv, Faculty of Mathematics and Informatics, Department of Algebra and Geometry, 236 Bulgaria Blvd, Plovdiv 4003, Bulgaria, Tel.: +359-32-261744, Fax: +359-32-261794

 $\textit{E-mail address} : \verb|mmanev@uni-plovdiv.bg|, \verb|mirkaiv@uni-plovdiv.bg|$