Non Asymptotic Performance of Some Markov Chain Order Estimators

A.R. Baigorri^{*} Mathematics Department UnB

October 28, 2012

Abstract

In what follows we study non asymptotic behavior of different well known estimators AIC([26]), BIC([24]) and EDC([27, 16]) in contrast with the Markov chain order estimator, named as Global Depency Level-GDL([6]).

The estimator GDL, is based on a different principle which makes it behave in a quite different form. It is strongly consistent and more efficient than AIC(inconsistent), outperforming the well established and consistent BIC and EDC, mainly on relatively small samples.

The estimators mentioned above mainly consist in the evaluation of the Markov chain's sample by different multivariate *deterministic* functions. The log likelihood approach, as in (11),

$$\mathbb{L}_{\left[\left(n\,,\,a_{1}^{\kappa}\right)\left(i,k\right)\right]}\left(\mathbb{X}_{a_{1}^{\kappa}}(i,k)\right) = \mathcal{L}_{1}\left(\mathcal{L}_{2}-\pi(i)\,\,\mathbb{X}_{a_{1}^{\kappa}}(i,k)\,\log\mathbb{X}_{a_{1}^{\kappa}}(i,k)\right) \tag{1}$$

with deterministic function

$$\boldsymbol{L}_{[(n, a_1^{\kappa})]} = \mathcal{L}_1 \Big(\mathcal{L}_2 - \pi(i) \, x(i, j) \, \log x(i, j), \quad \mathcal{L}_1 = const., \mathcal{L}_2 = const.$$

16

1

2

3

4

5

6

7

8

9

10

11

12

13

14

15

or, the GDL approach, as in (13)),

18

$$\mathbb{G}_{[(n, a_1^{\kappa})(i,j)]}(..., \mathbb{X}_{a_1^{\kappa}}(s,t), ...) = \sum_{i=1}^m \sum_{j=1}^m G_{[(n, a_1^{\kappa})(i,j)]}(..., \mathbb{X}_{a_1^{\kappa}}(s,t), ...)$$

^{*}baig@unb.br

with deterministic function

$$\boldsymbol{G}_{[(n, a_{1}^{\kappa})(i, j)]}\big(..., x(s, t), ...\big) = \frac{\left(x(i, j) - \left[\sum_{t=1}^{m} x(i, t)\right]\left[\sum_{s=1}^{m} x(s, j)\right]\right)^{2}}{\left(\sum_{t=1}^{m} x(i, t)\right)\left(\sum_{s=1}^{m} x(s, j)\right)}.$$

shall be analized in *Section 3*, exhibiting different structural properties. It will become clear the intimate differences existing between the variance of both estimators, which induce quite dissimilar performance, mainly for samples of moderated sizes.

25 1 Introduction

19

20

21

22

23

24

A Markov Chain is a discrete stochastic process $\mathbb{X} = \{X_n\}_{n\geq 0}$ with state space E, cardinality $|E| < \infty$ for which there is a $k \geq 1$ such that for $(x_1, \dots, x_n) \in E^n, n \geq k$

²⁹
$$P(X_1 = x_1, ..., X_n = x_n) = P(X_1 = x_1, ..., X_k = x_k) \prod_{i=k+1}^n Q(x_i | x_{i-k}, ..., x_{i-1})$$

for suitable transition probabilities Q(.|.). The class of processes that holds the above condition for a given $k \ge 1$ will be denoted by \mathcal{M}_k , and \mathcal{M}_0 will denote the class of i.i.d. processes. The **order of a Markov Chain** in $\cup_{i=0}^{\infty} \mathcal{M}_i$ is the **smallest integer** κ such that $\mathbb{X} = \{X_n\}_{n>0} \in \mathcal{M}_{\kappa}$.

Along the last few decades there has been a great number of research on
the estimation of the order of a Markov Chains, starting with M.S. Bartlett
[8], P.G. Hoel [18], I.J. Good [17], T.W. Anderson & L.A. Goodman [4],
P. Billingsley [9], [10] among others, and more recently, H. Tong [26], G.
Schwarz [24], R.W. Katz [19], I. Csiszar and P. Shields [13], L.C. Zhao et all
[27] had contributed with new Markov chain order estimators.

Since 1973, H. Akaike [1] entropic information criterion, known as AIC, has 40 had a fundamental impact in statistical model evaluation problems. The 41 AIC has been applied by Tong, for example, to the problem of estimating the 42 order of autoregressive processes, autoregressive integrated moving average 43 processes, and Markov chains. The Akaike-Tong (AIC) estimator was derived 44 as an asymptotic approximate estimate of the Kullback-Leibler information 45 discrepancy and provides a useful tool for evaluating models estimated by 46 the maximum likelihood method. Later on, Katz derived the asymptotic 47

distribution of the estimator and showed its inconsistency, proving that there
is a positive probability of overestimating the true order no matter how large
the sample size. Nevertheless, AIC is the most used and succesfull Markov
chain order estimator used at the present time, mainly because it is more
efficient than BIC for small sample.

⁵³ 2 Essentials on Some Estimators

54 2.1 Maximum Likelihood Methods

The main consistent estimator alternative, the BIC estimator, does not perform too well for relatively small samples, as it was pointed out by Katz [19] and Csiszar & Shields [13]. It is natural to admit that the expansion of the Markov Chain complexity (size of the state space and order) has significant influence on the sample size required for the identification of the unknown order, even though, most of the time it is difficult to obtain sufficiently large samples.

Katz(1981) [19] obtained the asymptotic distribution of $\hat{\kappa}_{AIC}$ and proved its inconsistency showing the existence of a positive probability to overestimate the order. See also Shibata(1976) [25]. On the contrary Schwarz (1978) [24] and Zhao(2001) [27] proved strong consistency for the estimators $\hat{\kappa}_{BIC}$ and $\hat{\kappa}_{EDC}$, respectively.

⁶⁷ Clearly, for a given η , $AIC(\eta)$ [26], $BIC(\eta)$ [24] and $EDC(\eta)$ [27, 16] contain ⁶⁸ much of the information concerning the sample's relative dependency, never-⁶⁹ theless numerical simulations as well as theoretical considerations anticipates ⁷⁰ a great deal of variability for small samples.

⁷¹ Let $X_1^n = (X_1, ..., X_n)$ be a sample from a multiple stationary Markov chain ⁷² $\mathbb{X} = \{X_n\}_{n\geq 1}$ of unknown order κ .

Assume that X take values on a finite state space $E = \{1, 2, ..., m\}$ with transition probabilities given by

75
$$p(x_{\kappa+1}|x_1^{\kappa}) = P(X_{n+1} = x_{n+1}|X_{n-\kappa+1}^n = x_1^{\kappa}) > 0$$
(2)

76 where
$$x_1^{\kappa} = (x_1, ..., x_{\kappa}) = x_1^j x_{j+1}^{\kappa} \in E^{\kappa}$$

77 Define

$$N(x_1^l | \mathbf{X}_1^n) = \sum_{j=1}^{n-l+1} 1(X_j = x_1, ..., X_{j+l-1} = x_l)$$
(3)

⁷⁹ i.e. the number of ocurrences of x_1^l in \mathbf{X}_1^n . If l = 0 we take $N(.|\mathbf{X}_1^n) = n$. The sums are taken over positive terms $N(x_1^{l+1}|\mathbf{X}_1^n) > 0$, or else, we ⁸¹ convention 0/0 or $0.\infty$ as 0.

82

78

Definition 2.1. For $a_1^{\eta} = (a_1, ..., a_{\eta}) \in E^{\eta}$ and $j \in E$, let $\mathbb{X}_{a_1^{\eta}}$ be the empirical random variables, extracted from the Markov chain sample $\mathbf{X}_1^n = (X_1, ..., X_n)$

$$\mathbb{X}_{a_1^{\eta}} : \mathbf{X}_1^n \longrightarrow \left(\mathbb{X}_{a_1^{\eta}}(1), \dots, \mathbb{X}_{a_1^{\eta}}(j), \dots, \mathbb{X}_{a_1^{\eta}}(m) \right)$$

$$\mathbb{X}_{a_1^{\eta}}(j) = \left(\frac{N(a_1^{\eta} j \mid \mathbf{X}_1^n)}{N(a_1^{\eta} \mid \mathbf{X}_1^n)} \right), \quad 1 \le j \le m$$

$$(4)$$

88 and

89
$$\mathbb{X}_{a_{1}^{\eta}}(i,j) : \mathbf{X}_{1}^{n} \longrightarrow \left(\mathbb{X}_{a_{1}^{\eta}i}(1), ..., \mathbb{X}_{a_{1}^{\eta}i}(j), ..., \mathbb{X}_{a_{1}^{\eta}i}(m)\right)$$
90
$$\mathbb{X}_{a_{1}^{\eta}}(i,j) = \left(\frac{N(a_{1}^{\eta}ij \mid \mathbf{X}_{1}^{n})}{N(a_{1}^{\eta}i \mid \mathbf{X}_{1}^{n})}\right), \quad 1 \le i, j \le m. \quad \bullet$$
(5)

⁹¹ Let us define for the *order* the log likelihood function

$$\log \hat{L}(\eta) = \sum_{j}^{m} N(a_{1}^{\eta} \mid \boldsymbol{X}_{1}^{n}) \left(\sum_{a_{1}^{\eta}} \frac{N(a_{1}^{\eta} j \mid \boldsymbol{X}_{1}^{n})}{N(a_{1}^{\eta} \mid \boldsymbol{X}_{1}^{n})} \log \mathbb{X}_{a_{1}^{\eta}}(j) \right)$$

$$\log \hat{L}(\eta) = \sum_{a_1^{\eta}} N(a_1^{\eta} \mid \mathbf{X}_1^n) \left(\sum_{j=1}^{m} \frac{N(a_1^{\eta} j \mid \mathbf{X}_1^n)}{N(a_1^{\eta} \mid \mathbf{X}_1^n)} \log \mathbb{X}_{a_1^{\eta}}(j) \right)$$

94
$$\log \hat{L}(\eta) = \sum_{a_1^{\eta}} \left[N(a_1^{\eta} | \mathbb{X}_1^{\eta}) \left(\sum_j^m \mathbb{X}_{a_1^{\eta}}(j) \log \mathbb{X}_{a_1^{\eta}}(j) \right) \right].$$
(6)

The estimators based on likelihood estimators and penalty functions, for 95 Markov chains of order κ are defined, under the following hypothesis: 96

$$There \ exist \ a \ known \ B \ so \ that \ 0 \ \leq \kappa \ \leq B$$

as 98

$$\widehat{\kappa}_{AIC} = \operatorname{argmin}\{AIC(\eta) \; ; \; \eta = 0, 1, ..., B\}$$
(7)

$$\widehat{\kappa}_{BIC} = \operatorname{argmin}\{BIC(\eta) : \eta = 0, 1, ..., B\}$$

$$\widehat{\kappa}_{n-1} = \operatorname{argmin}\{EDC(n) : n = 0, 1, ..., B\}$$

$$(8)$$

$$\widehat{\kappa}_{EDC} = \operatorname{argmin}\{EDC(\eta); \eta = 0, 1, ..., B\}$$
(9)

where 102

103
$$AIC(\eta) = -2\log \hat{L}(\eta) + |E|^{\eta+1} 2(|E|-1),$$

104
$$BIC(\eta) = -2\log \hat{L}(\eta) + |E|^{\eta+1} 2(|E|-1)\left(\frac{\log(n)}{2}\right)$$

$$BIC(\eta) = -2\log L(\eta) + |E|^{\eta+1} 2(|E|-1)\left(\frac{-8(\gamma)}{2}\right)$$

$$EDC(\eta) = -2\log \hat{L}(\eta) + |E|^{\eta+1} 2(|E|-1)\left(\frac{\log\log(n)}{2(|E|-1)}\right)$$

$$AIC(\eta) \leq EDC(\eta) \leq BIC(\eta).$$

$$AIC(\eta) \leq EDC(\eta) \leq BIC(\eta)$$

Finally, let us $fix a_1^{\eta}$ and consider the function

$$\mathbb{L}_{\left[\left(n\,,\,a_{1}^{\eta}\right)\right]}\,:\,\left(0,1\right)^{m^{2}}\,\rightarrow\,\mathbb{R}^{+}$$

defined as: 107

¹⁰⁸
$$\mathbb{L}_{[(n,a_{1}^{\eta})]}\left(...,x(i,j),...\right) = N(a_{1}^{\eta} \mid \boldsymbol{X}_{1}^{n}) \left(\sum_{i=1}^{m} \mathbb{X}_{a_{1}^{\eta}}(i) \log \mathbb{X}_{a_{1}^{\eta}}(i) - \sum_{i=1}^{m} \left[\frac{N(a_{1}^{\eta}i \mid \boldsymbol{X}_{1}^{n})}{N(a_{1}^{\eta} \mid \boldsymbol{X}_{1}^{n})}\right] \sum_{j=1}^{m} \mathbb{X}_{a_{1}^{\eta}}(i,j) \log \mathbb{X}_{a_{1}^{\eta}}(i,j)\right). (10)$$

Later on in *Section 3.1*, we shall analyse the behavior and derivatives of 110 $L_{[(n, a_1^{\kappa})(i,k)]}$ which is just a generic representation of $\mathbb{L}_{[(n, a_1^{\kappa})(i,k)]}$. 111

$$L_{[(n, a_1^{\kappa})(i,k)]} : (0,1) \to \mathbb{R}^+$$

$$\boldsymbol{L}_{[(n,a_1^{\kappa})(i,k)]}\Big(x(i,k)\Big) = \mathcal{L}_1\Big(\mathcal{L}_2 - \pi(i) \ x(i,j) \ \log x(i,j)\Big)$$
(11)

113 such that

$$\mathbb{L}_{[(n,a_1^{\kappa})(i,k)]}\left(\mathbb{X}_{a_1^{\kappa}}(i,k)\right) \equiv \boldsymbol{L}_{[(n,a_1^{\kappa})(i,k)]}\left(x(i,j)\right)$$

where $\mathcal{L}_1 = N(a_1^{\kappa} | \mathbf{X}_1^n)$ and $\mathcal{L}_2 = \sum_{i=1}^m \mathbb{X}_{a_1^{\kappa}}(i) \log \mathbb{X}_{a_1^{\kappa}}(i)$ are assumed **constants** with respect to the the variables x(i, j), with $x(i, j) = \mathbb{X}_{a_1^{\kappa}}(i, j)$ as in (5), κ the Markov chain order and

$$\pi(i) = \left[\frac{N(a_1^{\kappa}i \mid \mathbf{X}_1^n)}{N(a_1^{\kappa} \mid \mathbf{X}_1^n)}\right], \quad 1 \le i, j \le m.$$

115 2.2 χ^2 -divergence estimator

¹¹⁶ We now briefly recall this new Markov chain order's estimator referring the ¹¹⁷ reader to ([6]) for related details.

Definition 2.2. Let $X_n = \{X_i\}_{i=1}^n$ be a sample of a Markov chain X of order $\kappa \geq 0, X_{a_1^{\eta}}(i,j)$ as in (2.1), $\eta \geq 0$ and $\Delta_2(X_{a_1^{\eta}}(i,j))$ the random variable defined as follows

121
$$\Delta_2(\mathbb{X}_{a_1^\eta}(i,j)) =$$

122

$$= N(a_1^{\eta} | \mathbb{X}_1^n) \sum_{i=1}^m \sum_{j=1}^m \left(\frac{\left(\mathbb{X}_{a_1^{\eta}}(i,j) - \left[\sum_{t=1}^m \mathbb{X}_{a_1^{\eta}}(i,t) \right] \left[\sum_{s=1}^m \mathbb{X}_{a_1^{\eta}}(s,j) \right] \right)^2}{(\sum_{t=1}^m \mathbb{X}_{a_1^{\eta}}(i,t))(\sum_{s=1}^m \mathbb{X}_{a_1^{\eta}}(s,j))} \right) =$$

123
$$= \mathcal{G}_{(n,a_1^{\eta})} \sum_{i=1}^{m} \sum_{j=1}^{m} \mathbb{G}_{[(n,a_1^{\eta})(i,j)]}.$$

Assume that V is a χ^2 random variable with $(m-1)^2$ degrees of freedom where \mathcal{P} is the continuous strictly decreasing function $\mathcal{P}: \mathbb{R}^+ \longrightarrow [0, 1]$

126
$$\mathcal{P}(x) = P(V \ge x), \ x \in \mathbb{R}^+.$$

112

¹²⁷ The Local Dependency Level $LDL_n(a_1^{\eta})$ and the Global Dependency Level ¹²⁸ $GDL_n(\eta)$, respectively, are defined as follows

129
$$LDL_n(a_1^{\eta}) = \frac{\Delta_2(\mathbb{X}_{a_1^{\eta}}(i,j))}{2\log(\log(n))},$$

130

140

$$GDL_n(\eta) = \mathcal{P}\left(\sum_{a_1^\eta \in E^\eta} \left(\frac{N(a_1^\eta \mid \mathbb{X}_1^n)}{n}\right) LDL_n(a_1^\eta)\right). \quad \blacklozenge$$

Finally, let us define the Markov chain order estimator based on the information contained in the vector GDL_n .

Definition 2.3. Given a fixed number $0 < B \in \mathbb{N}$, let us define the set $\mathcal{S} = \{0, 1\}^{B+1}$ and the application $T : \mathcal{S} \to \mathbb{N}$

135
$$T(s) = -1 \iff s_i = 1, \ i = 0, 1, .., B$$

136
$$T(s) = \max_{0 \le i \le B} \{i : s_i = 0, s_{i+1} = \mathcal{P}(\mathcal{L})\}, \ s = (s_0, s_1, ..., s_B). \blacklozenge$$

137 **Definition 2.4.** Let $X_1^n = \{X_i\}_{i=1}^n$ be a sample for the Markov chain \mathbb{X} of 138 order κ , $0 \leq \kappa < B \in \mathbb{N}$ and $\{GDL_n(i)\}_{i=1}^B$ as above. We define the order's 139 estimator $\kappa_{GDL}(\mathbb{X}_1^n)$ as

1

$$\widehat{\kappa}_{GDL}(\mathbb{X}_1^n) = T(\sigma_n) +$$

141 with $\sigma_n \in \mathcal{S}$ so that $\forall s \in \mathcal{S}$

142
$$\sum_{i=0}^{B} \left(GDL_n(i) - \sigma_n(i) \right)^2 \le \sum_{i=0}^{B} (GDL_n(i) - s(i))^2. \quad \blacklozenge$$

Observe that the **Local Dependency Level** $LDL_n(a_1^{\eta})$ entirely relies on the just defined χ^2 -square divergence estimator which itself is the summation of several univariate random variables $\mathbb{G}_{[(n, a_1^{\eta})(i,j)]}, 1 \leq i, j \leq m$

$$\mathbb{G}_{[(n, a_{1}^{\eta})(i, j)]} = \qquad (12)$$

$$= \sum_{i=1}^{m} \sum_{j=1}^{m} \frac{\left(\mathbb{X}_{a_{1}^{\eta}}(i, j) - \left[\sum_{t=1}^{m} \mathbb{X}_{a_{1}^{\eta}}(i, t) \right] \left[\sum_{s=1}^{m} \mathbb{X}_{a_{1}^{\eta}}(s, j) \right] \right)^{2}}{\left(\sum_{t=1}^{m} \mathbb{X}_{a_{1}^{\eta}}(i, t) \right) \left(\sum_{s=1}^{m} \mathbb{X}_{a_{1}^{\eta}}(s, j) \right)}.$$

Later on in **Section 3.2**, we shall analyse the behavior of the deterministic function $G_{[(n, a_1^{\kappa})(i,j)]}, 1 \leq i, j \leq m$ and their derivatives

$$G_{[(n, a_1^{\kappa})(i, j)]} : (0, 1)^{2m} \to \mathbb{R}^+, \quad 1 \le i, j \le m$$

150 with

146

147

$$\boldsymbol{G}_{[(n, a_{1}^{\kappa})(i, j)]}(\dots, x(s, t), \dots) = \frac{\left(x(i, j) - \left[\sum_{t=1}^{m} x(i, t)\right] \left[\sum_{s=1}^{m} x(s, j)\right]\right)^{2}}{\left(\sum_{t=1}^{m} x(i, t))(\sum_{s=1}^{m} x(s, j)\right)}$$
(13)

151

152 such that

153

$$\mathbb{G}_{[(n, a_{1}^{\kappa})(i, j)]}\left(..., \mathbb{X}_{a_{1}^{\kappa}}(s, t), ...\right) \equiv \boldsymbol{G}_{[(n, a_{1}^{\kappa})(i, j)]}\left(..., x(s, t), ...\right)$$

154 with $x(s,t) = \mathbb{X}_{a_1^{\kappa}}(i,j), \ 1 \le i,j \le m, \ \mathbb{X}_{a_1^{\kappa}}(i,j)$ as in (5).

3 Deterministic Accessory Functions

¹⁵⁶ 3.1 Functions Related with AIC-Estimator

Let us calculate the derivatives of the deterministic function $L_{(n, a_1^{\kappa})}$ as in(11), which for the sake of notational simplicity and for a fixed n and a_1^{κ} , we'll *temporarily rename* the function

$$\boldsymbol{L} = \boldsymbol{L}_{[(n, a_1^{\kappa})(i, k)]}, \quad \boldsymbol{L} : D_{\boldsymbol{L}} \subseteq (0, 1) \to \mathbb{R},$$

$$D_{L} = \{x(i,k) : x(i,k) \in (0,1)\},$$

$$\boxed{L(x) = x(i,k) \log(x(i,k)).}$$
(14)

160

161 First Order Derivatives :

$$\frac{\partial \boldsymbol{L}}{\partial x(i,k)} = 1 + \log(x(i,k)), \quad 1 \le i, k \le m-1.$$

163 Second Order Derivatives :

$$\frac{\partial^2 \boldsymbol{L}}{\partial x^2(i,k)} = \frac{1}{x(i,k)}, \ 1 \le i,k \le m-1,$$
(15)

165

166

164

$$\frac{\partial^2 \boldsymbol{L}}{\partial x(j,l)\partial x(i,k)} = 0, \ 1 \le i, k, l \le m-1,$$
(16)

167 respectively.

Later on we shall obtain the gradient vector and the hessian matrix

 $\nabla_{\boldsymbol{L}}\left(\Lambda_{a_{1}^{\kappa}}(o)\right) , \mathcal{H}_{\boldsymbol{L}}\left(\Lambda_{a_{1}^{\kappa}}(o)\right)$

of the function \boldsymbol{L} at a point

$$\Lambda_{a_1^{\kappa}}(o) = \left(\dots, E\big(\mathbb{X}_{a_1^{\kappa}}(i,k)\big),\dots\big).$$

¹⁶⁸ 3.2 Functions Related with GDL-Estimator

Herein, we shall consider the deterministic multivariate set of functions, asin (13),

$$\mathbf{G}_{[(n, a_1^{\kappa})(i, k)]}\Big(x(i, k), h(i), v(k)\Big) = \frac{\left(x(i, k) - h(i)v(k)\right)^2}{h(i)v(k)}, \ 1 \le i, k \le m.$$

which, after fixing (i, k), we **temporarily rename** it as follows:

$$\boldsymbol{G} \equiv \boldsymbol{G}_{[(n, a_1^{\kappa})(i, k)]}, \qquad \boldsymbol{G} : D_{\boldsymbol{G}} \subseteq (0, 1)^3 \to \mathbb{R},$$
$$D_{\boldsymbol{G}} = \{\boldsymbol{x} \in (0, 1)^3 : \boldsymbol{x} = (x, h, v)\},$$

 $\boldsymbol{G}(\boldsymbol{x}) = \frac{(x - hv)^2}{hv}.$

(17)

174

179

173

175 First Order Derivatives :

$$\frac{\partial \boldsymbol{G}}{\partial x} = \frac{2x}{hv} - 2, \quad \frac{\partial \boldsymbol{G}}{\partial h} = \frac{-x^2}{vh^2} + v, \quad \frac{\partial \boldsymbol{G}}{\partial v} = \frac{-x^2}{hv^2} + h.$$

177 Second Order Derivatives :

$$\frac{\partial^2 \mathbf{G}}{\partial x^2} = \frac{2}{hv}, \qquad \frac{\partial^2 \mathbf{G}}{\partial h^2} = \frac{-x^2}{h^3 v}, \qquad \frac{\partial^2 \mathbf{G}}{\partial v^2} = \frac{-x^2}{v^3 h}, \tag{18}$$

$$\frac{\partial^2 \mathbf{G}}{\partial h \partial x} = \frac{-2x}{h^2 v}, \qquad \frac{\partial^2 \mathbf{G}}{\partial v \partial x} = \frac{-2x}{hv^2}, \qquad \frac{\partial^2 \mathbf{G}}{\partial v \partial h} = \frac{x^2}{h^2 v^2} + 1.$$
(19)

Likewise, as in the previous subsection we get the gradient vector and the hessian matrix $\nabla_{\boldsymbol{G}} (\Gamma_{a_1^{\kappa}}(o))$, $\mathcal{H}_{\boldsymbol{G}} (\Gamma_{a_1^{\kappa}}(o))$ of the function $\boldsymbol{G}(x,h,v)$ at a

point

$$\Gamma_{a_1^{\kappa}}(o) = \left(E(\mathbb{X}_{a_1^{\kappa}}(i,k)), E(\mathbb{H}_{a_1^{\kappa}}(i)), E(\mathbb{V}_{a_1^{\kappa}}(k)) \right).$$

180 4 Multivariate Variances

Focusing on $G_{[(n, a_1^{\kappa})(i,k)]}$ for fixed (i, k), κ the order of the Markov chain and $G_{[(n, a_1^{\kappa})(i,k)]}$ as in (13), let us recall the empirical random variables, introduced in Definition 2.1

$$\mathbb{X}_{a_{1}^{\kappa}}(i,k) = \frac{N(i \, a_{1}^{\kappa} \, k \, | \, \mathbf{X}_{1}^{n})}{N(a_{1}^{\kappa} \, | \, \mathbf{X}_{1}^{n}))}, \quad \mathbb{H}_{a_{1}^{\kappa}}(i) = \sum_{t=1}^{m} \mathbb{X}_{a_{1}^{\kappa}}(i,t), \quad \mathbb{V}_{a_{1}^{\kappa}}(k) = \sum_{s=1}^{m} \mathbb{X}_{a_{1}^{\kappa}}(s,k).$$

 184 Observe that the Markov chain we are interested in, has order κ and it is 185 clear that

$$\mathbb{X}_{a_1^{\kappa}}(i,k)$$
 independent random variables, $1 \leq i, k \leq m$,

187

$$G_{[(n, a_1^{\kappa})(i,j)]}, \quad 1 \le i, k \le m-1,$$
(20)

¹⁸⁸ are independent, with

¹⁸⁹
$$\boldsymbol{G}_{[(n, a_{1}^{\kappa})(i, j)]}\left(\mathbb{X}_{a_{1}^{\kappa}}(i, k), \mathbb{H}_{a_{1}^{\kappa}}(i), \mathbb{V}_{a_{1}^{\kappa}}(k)\right) = \frac{\left(\mathbb{X}_{a_{1}^{\kappa}}(i, k) - \mathbb{H}_{a_{1}^{\kappa}}(i) \mathbb{V}_{a_{1}^{\kappa}}(k)\right)^{2}}{\mathbb{H}_{a_{1}^{\kappa}}(i) \mathbb{V}_{a_{1}^{\kappa}}(k)}$$

190 as well as for *adequatly* sample size n the random variables

¹⁹¹
$$E\left(\mathbb{H}_{a_1^{\kappa}}(i)\right) \cong \frac{1}{m}, \quad E\left(\mathbb{V}_{a_1^{\kappa}}(k)\right) \cong m E\left(\mathbb{X}_{a_1^{\kappa}}(s,k)\right), \quad 1 \le s \le m.$$

For the sake of notation's simplicity, we temporarily rename

$$\boldsymbol{G}_{\left[\left(n,a_{1}^{\kappa}\right)\left(i,j\right)\right]}\left(\boldsymbol{x}(i,k),\boldsymbol{h}(i),\boldsymbol{v}(k)\right)$$

as G(x, h, v) where its derivatives, as well as the variances, covariances and related information of $\{ \mathbb{X}_{a_1^{\kappa}}(i, k), \mathbb{H}_{a_1^{\kappa}}(i), \mathbb{V}_{a_1^{\kappa}}(k) \}$ shall be as follows:

$$E\left(\frac{\mathbb{X}_{a_{1}^{\kappa}}(i,k)}{\mathbb{H}_{a_{1}^{\kappa}}(i)\mathbb{V}_{a_{1}^{\kappa}}(k)}\right) \cong \frac{E\left(\mathbb{X}_{a_{1}^{\kappa}}(i,k)\right)}{E\left(\mathbb{H}_{a_{1}^{\kappa}}(i)\right)E\left(\mathbb{V}_{a_{1}^{\kappa}}(k)\right)} \cong \frac{E(\mathbb{X}_{a_{1}^{\kappa}}(i,k))}{\frac{1}{m}\left(mE(\mathbb{X}_{a_{1}^{\kappa}}(i,k))\right)} \cong 1,$$

$$\begin{array}{ccc}
\sum_{\alpha,\beta\in\{x,h,v\}} \frac{\partial \boldsymbol{G}}{\partial \alpha}(x,h,v) \frac{\partial \boldsymbol{G}}{\partial \beta}(x,h,v) \\ \cong \\
\begin{array}{c}
\sum_{\alpha,\beta\in\{x,h,v\}} \frac{\partial \boldsymbol{G}}{\partial \alpha}(x,h,v) \frac{\partial \boldsymbol{G}}{\partial \beta}(x,h,v) \\ \cong \\
\left(1 - \left(\frac{x}{hv}\right)^{2}\right) \left(4 + h^{2} + v^{2}\right) + \frac{2x}{h} \left(\left(\frac{x}{hv}\right) - \left(\frac{x}{hv}\right)^{2}\right) + \\
\end{array} \\
\begin{array}{c}
\sum_{\alpha,\beta\in\{x,h,v\}} \frac{\partial \boldsymbol{G}}{\partial \alpha}(x,h,v) \frac{\partial \boldsymbol{G}}{\partial \beta}(x,h,v) \\ \cong \\
\left(1 - \left(\frac{x}{hv}\right)^{2}\right) \left(4 + h^{2} + v^{2}\right) + \frac{2x}{h} \left(\left(\frac{x}{hv}\right) - \left(\frac{x}{hv}\right)^{2}\right) + \\
\left(\frac{x}{hv}\right) - \left(\frac{x}{hv}\right)^{2}\right) + x \left(\left(\frac{x}{hv}\right)^{3} - 2\left(\frac{x}{hv}\right)^{2} + \left(\frac{x}{hv}\right)\right) \\
\end{array} \\
\begin{array}{c}
\sum_{\alpha,\beta\in\{x,h,v\}} \frac{\partial \boldsymbol{G}}{\partial \alpha}(x,h,v) \frac{\partial \boldsymbol{G}}{\partial \beta}(x,h,v) \\ \cong \\
\left(1 - \left(\frac{x}{hv}\right)^{2}\right) \left(4 + h^{2} + v^{2}\right) + \frac{2x}{h} \left(\left(\frac{x}{hv}\right) - \left(\frac{x}{hv}\right)^{2}\right) + \\
\left(\frac{x}{hv}\right)^{2} \left(\left(\frac{x}{hv}\right) - \left(\frac{x}{hv}\right)^{2}\right) + x \left(\left(\frac{x}{hv}\right)^{3} - 2\left(\frac{x}{hv}\right)^{2} + \left(\frac{x}{hv}\right)\right) \\
\end{array} \\
\begin{array}{c}
\sum_{\alpha,\beta\in\{x,h,v\}} \frac{\partial \boldsymbol{G}}{\partial \alpha}(x,h,v) \frac{\partial \boldsymbol{G}}{\partial \beta}(x,h,v) \\
\cong \\
\sum_{\alpha,\beta\in\{x,h,v\}} \frac{\partial \boldsymbol{G}}{\partial \alpha}(x,h,v) \frac{\partial \boldsymbol{G}}{\partial \beta}(x,h,v) \\
\end{array} \\
\xrightarrow{\left(1 - \left(\frac{x}{hv}\right)^{2}\right) \left(4 + h^{2} + v^{2}\right) + \frac{2x}{h} \left(\left(\frac{x}{hv}\right) - \left(\frac{x}{hv}\right)^{2}\right) + \\
\left(\frac{x}{hv}\right)^{2} \left(\left(\frac{x}{hv}\right) - \left(\frac{x}{hv}\right)^{2}\right) + x \left(\left(\frac{x}{hv}\right)^{3} - 2\left(\frac{x}{hv}\right)^{2} + \left(\frac{x}{hv}\right)\right) \\
\end{array} \\
\xrightarrow{\left(1 - \frac{x}{v}\right)^{2}} \left(\frac{x}{hv}\right)^{2} \left(\frac{x}{hv}\right) + x \left(\frac{x}{hv}\right)^{2} \left(\frac{x}{hv}\right)^{2} + \left(\frac{x}{hv}\right)^{2}\right) \\
\xrightarrow{\left(1 - \frac{x}{v}\right)^{2}} \left(\frac{x}{hv}\right)^{2} \left(\frac{x}{hv}\right)^{2} + \frac{x}{hv}\left(\frac{x}{hv}\right)^{2} \left(\frac{x}{hv}\right)^{2} \left(\frac{x}{hv}\right)^{2} + \frac{x}{hv}\left(\frac{x}{hv}\right)^{2}\right) \\
\xrightarrow{\left(1 - \frac{x}{hv}\right)^{2}} \left(\frac{x}{hv}\right)^{2} \left($$

Likewise,

$$\sum_{\alpha,\beta,\gamma\in\{x,h,v\}} \frac{\partial \boldsymbol{G}}{\partial \alpha}(x,h,v) \frac{\partial^{2} \boldsymbol{G}}{\partial \beta \partial \gamma}(x,h,v) \approx 0.$$

$$Cov_{(\mathbb{X}_{a_{1}^{\kappa}}(i,k),\mathbb{X}_{a_{1}^{\kappa}}^{2}(i,k))} \approx Cov_{(\mathbb{X}_{a_{1}^{\kappa}}(i,k),\mathbb{H}_{a_{1}^{\kappa}}^{2}(i))} \approx Cov_{(\mathbb{X}_{a_{1}^{\kappa}}(i,k),\mathbb{V}_{a_{1}^{\kappa}}^{2}(k))} \approx \sigma_{\mathbb{X}_{a_{1}^{\kappa}}}^{3}(i,k)}$$

$$Cov_{(\mathbb{X}_{a_{1}^{\kappa}}(i,k),\mathbb{H}_{a_{1}^{\kappa}}^{2}(i))} \approx Cov_{(\mathbb{X}_{a_{1}^{\kappa}}(i,k),\mathbb{V}_{a_{1}^{\kappa}}^{2}(k))} \approx \sigma_{\mathbb{X}_{a_{1}^{\kappa}}}^{3}(i,k)}$$

Finally,
Finally,

$$\sum_{\alpha,\beta\in\{x,h,v\}} \frac{\partial^2 \mathbf{G}}{\partial \alpha^2}(x,h,v) \frac{\partial^2 \mathbf{G}}{\partial \beta^2}(x,h,v) \approx \left(\frac{\partial^2 \mathbf{G}}{\partial x^2}\right)^2 + \frac{\partial^2 \mathbf{G}}{\partial x^2} \frac{\partial^2 \mathbf{G}}{\partial v^2} + \frac{\partial^2 \mathbf{G}}{\partial x^2} \frac{\partial^2 \mathbf{G}}{\partial v \partial x} + \left(\frac{\partial^2 \mathbf{G}}{\partial v \partial x}\right)^2 = \frac{1}{h^2 v^2} \left(4 + \frac{x^4}{v^4} + \frac{4x^3}{v^3} - \frac{8x}{v}\right) \approx$$

$$\approx \frac{1}{x^2} \left(4 + \frac{1}{m^4} + \frac{4}{m^3} - \frac{8}{m}\right) \approx \frac{4}{x^2} \left(1 - \frac{2}{m}\right) + \frac{1}{m^3} + \frac{1}{m^4} \approx \left[\frac{4}{x^2}\right]$$

²⁰⁹
$$\operatorname{cov}_{(\mathbb{X}^{2_{\kappa}}_{a_{1}^{\kappa}}(i,k),\mathbb{X}^{2_{\kappa}}_{a_{1}^{\kappa}}(i,k))} \cong \operatorname{cov}_{(\mathbb{X}^{2_{\kappa}}_{a_{1}^{\kappa}}(i,k),\mathbb{H}^{2_{\kappa}}_{a_{1}^{\kappa}}(i))} \cong \operatorname{cov}_{(\mathbb{X}^{2_{\kappa}}_{a_{1}^{\kappa}}(i,k),\mathbb{V}^{2_{\kappa}}_{a_{1}^{\kappa}}(k))} \cong \sigma^{4}_{\mathbb{X}^{4_{\kappa}}_{a_{1}^{\kappa}}(i,k)}$$

$$\operatorname{cov}_{(\mathbb{H}^{2}_{a_{1}^{\kappa}}(i),\mathbb{H}^{2}_{a_{1}^{\kappa}}(i))} \cong \operatorname{cov}_{(\mathbb{H}^{2}_{a_{1}^{\kappa}}(i),\mathbb{V}^{2}_{a_{1}^{\kappa}}(k))} \cong \operatorname{cov}_{(\mathbb{V}^{2}_{a_{1}^{\kappa}}(k),\mathbb{V}^{2}_{a_{1}^{\kappa}}(k))} \cong \sigma^{4}_{\mathbb{X}_{a_{1}^{\kappa}}(i,k)}.$$

211 Let us denote by $\mathcal{B} \in \mathbb{R}^3$, the unit ball centered at the point

$$\Gamma_{a_1^{\kappa}}(o) = \left(E(\mathbb{X}_{a_1^{\kappa}}(i,k)), E(\mathbb{H}_{a_1^{\kappa}}(i)), E(\mathbb{V}_{a_1^{\kappa}}(k)) \right)$$
$$\omega = \left(\mathbb{X}_{a_1^{\kappa}}(i,k), \mathbb{H}_{a_1^{\kappa}}(i), \mathbb{V}_{a_1^{\kappa}}(k) \right), \ \Delta = \omega - \Gamma_{a_1^{\kappa}}(o).$$

²¹² Taylor ([5]) showed that there exist $0 < c_g, c_l < 1$ such that

$$\mathbf{G}_{[(n, a_{1}^{\kappa})(i, j)]}(\omega) = \mathbf{G}_{(n, a_{1}^{\kappa})}[i, j] \Big(\Gamma_{a_{1}^{\kappa}}(o) \Big) + \nabla_{\mathbf{G}_{(n, a_{1}^{\kappa})}[i, j]} \Big(\Gamma_{a_{1}^{\kappa}}(o) \Big) \cdot (\omega - \Gamma_{a_{1}^{\kappa}}(o)) + \frac{1}{2!} (\omega - \Gamma_{a_{1}^{\kappa}}(o)) \cdot \mathcal{H}_{\mathbf{G}_{(n, a_{1}^{\kappa})}[i, j]} \Big(\Gamma_{a_{1}^{\kappa}}(o) + c_{g} \Delta \Big) \cdot (\omega - \Gamma_{a_{1}^{\kappa}}(o))^{t}$$

where the variance of

$$\boldsymbol{G}_{[(n, a_1^{\kappa})(i, k)]}\Big(\mathbb{X}_{a_1^{\kappa}}(i, k), \mathbb{H}_{a_1^{\kappa}}(i), \mathbb{V}_{a_1^{\kappa}}(k)\Big)$$

216 is

with

$$\begin{aligned} & 217 \qquad \sigma^{2}_{\boldsymbol{G}_{[(n, a_{1}^{\kappa})(i, j)]}} = \\ & 218 \qquad = E \left[\boldsymbol{G}_{(n, a_{1}^{\kappa})}[i, j](\omega) - \boldsymbol{G}_{(n, a_{1}^{\kappa})}[i, j](\Gamma_{a_{1}^{\kappa}}(o)) \right]^{2} = \\ & 219 \qquad = E \left[\nabla_{\boldsymbol{G}_{(n, a_{1}^{\kappa})}[i, j]}(\Gamma_{a_{1}^{\kappa}}(o)) \cdot \Delta + \frac{1}{2!}\Delta \cdot \mathcal{H}_{\boldsymbol{G}_{(n, a_{1}^{\kappa})}[i, j]}(\Gamma_{a_{1}^{\kappa}}(o) + c_{g}(\omega - \Gamma_{a_{1}^{\kappa}}(o))) \cdot \Delta^{t} \right]^{2}. \end{aligned}$$

$$\sigma_{\mathbb{G}[(n, a_{1}^{\kappa})(i, k)]}^{2} \approx \frac{1}{[2!]^{2}} \Phi_{g}^{\prime\prime} \Big(\Gamma_{a_{1}^{\kappa}}(o) + c_{g} \left(\omega - \Gamma_{a_{1}^{\kappa}}(o) \right) \Big) \sigma_{\mathbb{X}_{a_{1}^{\kappa}}(i, k)}^{4}$$

221 where
$$\omega = \left(\mathbb{X}_{a_1^{\kappa}}(i,k), \mathbb{H}_{a_1^{\kappa}}(i), \mathbb{V}_{a_1^{\kappa}}(k) \right) \in \mathcal{B}$$
 and

$$P_{g}''(x(i,k),h(i),v(k)) = \left(\frac{1}{h^{2}(i)v^{2}(k)}\right) \left[4 + \frac{x^{4}(i,k)}{v^{4}(k)} + 4\frac{x^{3}(i,k)}{v^{3}(k)} - 8\frac{x(i,k)}{v(k)}\right]$$

$$\approx \frac{4}{x(i,k)}$$

with variance for $1 \le i, k \le m - 1$

$$\sigma_{\mathbb{G}[(n,a_1^{\kappa})(i,k)]}^2 \approx \frac{\sigma_{\mathbb{X}_{a_1^{\kappa}}(i,k)}^4}{\left[\Gamma_{a_1^{\kappa}}(o) + c_g\left(\omega - \Gamma_{a_1^{\kappa}}(o)\right)\right]^2}.$$

and, by (20), the total variance

227
$$\sigma_{\mathbb{G}_{[(n, a_1^{\eta})]}}^2 = \sum_{i=1}^{m-1} \sum_{k=1}^{m-1} \sigma_{\mathbb{G}[(n, a_1^{\kappa})(i, k)]}^2.$$

Exactly as before we can obtain the total variance of $\mathbb{L}_{(n,a_1^{\kappa})}(\mathbb{X}_{a_1^{\kappa}}(i,k))$ and defining

$$\sigma^2_{\mathbb{L}_{[(n, a_1^{\kappa})(i, k)]}} \cong [1 + \ln(x(i, k))]^2 \sigma^2_{\mathbb{X}_{a_1^{\kappa}}(i, k)} + \Phi_l''(x(i, k)) \sigma^4_{\mathbb{X}_{a_1^{\kappa}}(i, k)}, 1 \le i, k \le m$$

where $\omega = (x(i,k), h(i), v(k)) \in \mathcal{B}$,

$$\varPhi_l''(x(i,k)) = \left[\frac{1}{x^2(i,k)}\right]$$

with variance for $1 \le i, k \le m-1$

$$\sigma_{\mathbb{L}_{[(n,a_{1}^{\kappa})(i,k)]}}^{2} \approx \left[1 + \ln(x(i,k))\right]^{2} \sigma_{\mathbb{X}_{a_{1}^{\kappa}}(i,k)}^{2} + \frac{\sigma_{\mathbb{X}_{a_{1}^{\kappa}}(i,k)}^{4}}{\left[\left(E(\mathbb{X}_{a_{1}^{\kappa}}) + c_{l}\left[x(i,k) - E(\mathbb{X}_{a_{1}^{\kappa}})\right]\right)\right]^{2}},$$

 $_{239}$ and, by (20), the total variance

$$\sigma^2_{\mathbb{L}_{[(n, a_1^{\kappa})]}} \approx \sum_{k=1}^{m-1} \sum_{i=1}^{m-1} \sigma^2_{\mathbb{L}_{(n, a_1^{\kappa})}(i,k)}.$$

241 5 Conclusion

The purpose of this work was the comparative analysis of the non asymptotic behavior for the estimators AIC(7), BIC(8), EDC(9), versus the estimator defined in Definition 2.2 and named as *Global Depency Level-GDL*, for details see ([6]).

The *GDL* uses a function different to the log likelihoog function applied to the sample, which makes the estimator perform in a quite different form. It is strongly consistent and more efficient than *AIC*(inconsistent), outperforming the well established and consistent *BIC* and *EDC*, mainly on reasonable small samples.

The estimators just mentioned are based on the composition of the empirical random variables with two different *deterministic* functions. The log likelihood approach, as in (11), with

$$\boldsymbol{L}_{[(n,a_1^{\kappa})(i,k)]}\Big(x(i,k)\Big) = \mathcal{L}_1\Big(\mathcal{L}_2 - \pi(i)\,x(i,k)\,\log x(i,k)\Big)$$

²⁵⁵ or, the GDL approach, as in (12), with

240

254

$$\boldsymbol{G}_{[(n, a_{1}^{\kappa})(i,j)]}(..., x(i,j), ...) = \frac{\left(x(i,j) - \left[\sum_{t=1}^{m} x(i,t)\right] \left[\sum_{s=1}^{m} x(s,j)\right]\right)^{2}}{\left(\sum_{t=1}^{m} x(i,t)\right)\left(\sum_{s=1}^{m} x(s,j)\right)}.$$

Since the sample only depends on the Markov chain X_1^n and its size **n**, once the sample is chosen, the entirely responsibles for the estimator's variance are the following random variables:

$$\mathbb{L}_{[(n,a_1^{\kappa})(i,k)]} = \mathcal{L}_1 \Big(\mathcal{L}_2 - \pi(i) \ \mathbb{X}_{a_1^{\kappa}}(i,k) \log \mathbb{X}_{a_1^{\kappa}}(i,k) \Big|$$

261 and

256

$$\mathbb{G}_{[(n,a_{1}^{\kappa})(i,k)]} = \sum_{k=1}^{m} \sum_{i=1}^{m} \frac{\left(\mathbb{X}_{a_{1}^{\kappa}}(i,k) - \left[\sum_{t=1}^{m} \mathbb{X}_{a_{1}^{\kappa}}(i,t)\right] \left[\sum_{s=1}^{m} \mathbb{X}_{a_{1}^{\kappa}}(s,k)\right]\right)^{2}}{\left(\sum_{t=1}^{m} \mathbb{X}_{a_{1}^{\kappa}}(i,t)\right)\left(\sum_{s=1}^{m} \mathbb{X}_{a_{1}^{\kappa}}(s,k)\right)}$$

with variances for $1 \le i, k \le m-1$

$$\sigma_{\mathbb{L}_{[(n,a_{1}^{\kappa})(i,k)]}}^{2} \approx \left[1 + \ln(x(i,k))\right]^{2} \sigma_{\mathbb{X}_{a_{1}^{\kappa}}(i,k)}^{2} + \frac{\sigma_{\mathbb{X}_{a_{1}^{\kappa}}(i,k)}^{4}}{\left[\left(E(\mathbb{X}_{a_{1}^{\kappa}}) + c_{l}\left[x(i,k) - E(\mathbb{X}_{a_{1}^{\kappa}})\right]\right)\right]^{2}},$$

265 and

266

Γ.

$$\sigma_{\mathbb{G}[(n, a_1^{\kappa})(i, k)]}^2 \approx \frac{4 \sigma_{\mathbb{X}_{a_1^{\kappa}}(i, k)}^4}{\left[\Gamma_{a_1^{\kappa}}(o) + c_g \left(\omega - \Gamma_{a_1^{\kappa}}(o)\right)\right]^2}.$$

²⁶⁷ respectively.

Finally the reader should notice that the log likelihood based estimators are heavily affected by $\log(x(i,k))$ which in cases where the Markov chain intrisically presents empirical random variables $\mathbf{X}_{a_1^{\kappa}}(i,k)$ with small expectations, the fluctuating values of x(i,k) converging to $E(\mathbf{X}_{a_1^{\kappa}}(i,k)) \cong 0$ imposes the coefficients $[1 + \log(x(i,k))]^2$ and its variance $\sigma^2_{\mathbf{L}_{(n,a_1^{\kappa})}}$ a great deal of instability or variance. ²⁷⁴ The following Appendix presents a few examples exhibiting such anomaly.

275 6 Appendix

276 6.1 Numerical Evidence

In what follows we shall compare the non-asymptotic performance, mainly 277 for small samples, of some of the most used Markov chains order estimators. 278 It is quite intuitive that the random information regarding the order of a 279 Markov chain, is spread over an exponentially growing set of empirical dis-280 tributions Θ with $|\Theta| = m^{B+1}$, where **B** is the maximum integer η , as in 281 $\alpha = (i_1 i_2 \dots i_\eta)$. It seems reasonable to think that a small viable sample, 282 i.e. samples able to retrieve enough information to estimate the chain order, 283 should have size $n \approx O(m^{B+1})$. Keeping in mind that for the present nu-284 merical simulation, the maximum length to be used is B = 5, from now on 285 the sample sizes for |E| = 3 and |E| = 4 should be $n \approx 1.500$ and $n \approx 5.000$, 286 respectively. 287

The following numerical simulation, based on an algorithm due to Raftery [23], starts on with the generation of a Markov chain transition matrix, $\mathbf{Q} = (q_{i_1 i_2 \dots i_k; i_{k+1}})$ with entries

$$q_{i_1 i_2 \dots i_{\kappa}; i_{\kappa+1}} = \sum_{t=1}^{\kappa} \lambda_{i_t} R(i_{\kappa+1}, i_t), \ 1 \le i_t, i_{\kappa+1} \le m.$$
(21)

²⁹² where the matrix

291

2

293
$$R(i,j), \ 0 \le i,j \le m, \quad \sum_{i=1}^{m} R(i,j) = 1, \ 1 \le j \le m$$

²⁹⁴ and the positive numbers

95
$$\{\lambda_i\}_{i=1}^\kappa, \ \sum_{i=1}^\kappa \lambda_i = 1$$

²⁹⁶ are arbitrarily chosen in advance.

- ²⁹⁷ Once the matrix $\mathbf{Q} = (q_{i_1 i_2 \dots i_{\kappa}; i_{\kappa+1}})$ is obtained, two hundreds replications of
- ²⁹⁸ the Markov chain sample of size n, space state E and transition matrix \mathbf{Q}

are generated to compare $GDL(\eta)$ performance against the standards, well known and already established order estimators just mentioned above.

³⁰¹ Finally, after applying all estimators to each one of the replicated samples,

³⁰² the final results two hundreds replications are registered in the form of tables.

303 Case I: Markov Chain Examples with $\kappa = 0$, |E| = 3.

Firstly, we choose the matrix $\{Q_1, Q_2, Q_3\}$ to produce samples with sizes 500 $\leq n \leq 2.000$, originated from Markov chains of order $\kappa = 0$ with quite different probability distributions.

| | | 0.33 | 0.335 | 0.335 | | 0.05 | 0.475 | 0.475 |] | 0.05 | 0.05 | 0.90 | |
|-----|---------|------|-------|---------|-----------|------|-------|-------|-----------|------|------|------|---|
| 307 | $Q_1 =$ | 0.33 | 0.335 | 0.335 | $, Q_2 =$ | 0.05 | 0.475 | 0.475 | $, Q_3 =$ | 0.05 | 0.05 | 0.90 | . |
| | | 0.33 | 0.335 | 0.335 _ | | 0.05 | 0.475 | 0.475 | | 0.05 | 0.05 | 0.90 | |

| 3 | 0 | 8 |
|---|---|---|
| | | |

| | $ E = 3 \qquad \leftrightarrow \qquad \kappa = 0 \qquad \leftrightarrow \qquad \lambda_i = 1/3, i = 1,2,3.$ | | | | | | | | | | | | | |
|---|--|------|------|-----|-----------|------|-------|-------|-----------|------|------|-----|--|--|
| | | Q | 1 | | | 4 | Q_1 | | Q_1 | | | | | |
| | | n = | 500 | | n = 1.000 | | | | n = 1.500 | | | | | |
| k | Aic | Bic | Edc | Gdl | Aic | Bic | Edc | Gdl | Aic | Bic | Edc | Gdl | | |
| 0 | 75.5% | 100% | 100% | 99% | 80% | 100% | 100% | 99.5% | 71.5% | 100% | 100% | 99% | | |
| 1 | 24.5% | | | 1% | 18% | | | 0.5% | 22.5% | | | 1% | | |
| 2 | | | | | 2% | | | | 6% | | | | | |
| 3 | | | | | | | | | | | | | | |
| 4 | | | | | | | | | | | | | | |

| | $ E = 3 \qquad \leftrightarrow \qquad \kappa = 0 \qquad \leftrightarrow \qquad \lambda_i = 1/3, i = 1,2,3.$ | | | | | | | | | | | | |
|---|---|-------|------|-----|-----------|-------------|------|-----|---------|------|-------|-----|--|
| | | Q | 2 | | | Q_2 Q_2 | | | | | P_2 | | |
| | | n = 1 | .000 | | n = 1.500 | | | | n = 500 | | | | |
| k | Aic | Bic | Edc | Gdl | Aic | Bic | Edc | Gdl | Aic | Bic | Edc | Gdl | |
| 0 | 63.5% | 100% | 100% | 99% | 63% | 100% | 100% | 99% | 59% | 100% | 100% | 99% | |
| 1 | 29% | | | 1% | 34.5% | | | 1% | 37% | | | 1% | |
| 2 | 7.5% | | | | 2.5% | | | | 4% | | | | |
| 3 | | | | | | | | | | | | | |
| 4 | | | | | | | | | | | | | |

| | $ E =3$ \leftrightarrow $\kappa=0$ \leftrightarrow $\lambda_i=1/3, i=1,2,3.$ | | | | | | | | | | | | | |
|---|--|-------|-------|-----|-------|-----------|-------|-----|-------|-----------|------|-----|--|--|
| | | Ģ |)3 | | Q_3 | | | | Q_3 | | | | | |
| | | n = 1 | 1.000 | | | n = 1.500 | | | | n = 2.000 | | | | |
| k | Aic | Bic | Edc | Gdl | Aic | Bic | Edc | Gdl | Aic | Bic | Edc | Gdl | | |
| 0 | 43% | 100% | 100% | 98% | 47% | 100% | 99.5% | 96% | 46% | 100% | 100% | 97% | | |
| 1 | 53% | | | 2% | 51.5% | | 0.5% | 4% | 50.5% | | | 2% | | |
| 2 | 4% | | | | 1.5% | | | | 3.5% | | | 1% | | |
| 3 | | | | | | | | | | | | | | |
| 4 | | | | | | | | | | | | | | |

Notice that for a fixed sample size $n = \{500, 1.000, 1.500, 2.000\}$, the order estimator $\hat{\kappa}_{AIC}$ steadily overestimate the real order $\kappa = 0$ with the excessiveness depending on the probability distribution of the Markov chain. Differently, the order estimators $\hat{\kappa}_{BIC}$, $\hat{\kappa}_{EDC}$ and $\hat{\kappa}_{GDL}$ show consistent performance, mainly obtaining the right order, free from the influence of the sample size and the generating matrix. Regarding $\hat{\kappa}_{BIC}$ and $\hat{\kappa}_{EDC}$ improved effect, most likely depends on their correcting factor, $\frac{\log(n)}{2}$ and $\left(\frac{\log\log(n)}{2(|E|-1)}\right)$ which tend to decrease the estimated order.

For |E| = 4 the greater complexity of a Markov chain of order $\kappa = 3$ impose the use of larger sample size for estimators to acomplish some reliability. Finally, we choose the matrix $\{Q_6, Q_7\}$ to produce samples with size n =5.000, originated from Markov chains of order $\kappa \in \{2, 3, 0\}$ like in the previous cases.

324

$$Q_6 = \begin{bmatrix} 0.05 & 0.05 & 0.05 & 0.85 \\ 0.05 & 0.05 & 0.85 & 0.05 \\ 0.05 & 0.85 & 0.05 & 0.05 \\ 0.85 & 0.05 & 0.05 & 0.05 \end{bmatrix}, \qquad Q_7 = \begin{bmatrix} 0.05 & 0.05 & 0.05 & 0.85 \\ 0.05 & 0.05 & 0.05 & 0.85 \\ 0.05 & 0.05 & 0.05 & 0.85 \\ 0.05 & 0.05 & 0.05 & 0.85 \end{bmatrix}$$

| | $ E = 4 \leftrightarrow n = 5.000$ | | | | | | | | | | | | | |
|---|---------------------------------------|------------|---------------------------|------------|--------|---|--------------|-----|-----|------|------|------|--|--|
| | Q_6 | $Q_6 \neq$ | $\Rightarrow \lambda_i =$ | 1/3, i = 1 | 1,2,3. | $Q_7 \Leftrightarrow \lambda_i = 1/3, i = 1,2,3.$ | | | | | | | | |
| | | | $\kappa = 3$ | | | | $\kappa = 0$ | | | | | | | |
| k | Aic | Bic | Edc | Gdl | Aic | Bic | Edc | Gdl | Aic | Bic | Edc | Gdl | | |
| 0 | | | | | | | | | 85% | 100% | 100% | 100% | | |
| 1 | | | | | | | | | 15% | | | | | |
| 2 | 100% | 100% | 100% | 100% | | 99% | | 4% | | | | | | |
| 3 | | | | | 100% | 1% | 100% | 96% | | | | | | |
| 4 | | | | | | | | | | | | | | |
| 5 | | | | | | | | | | | | | | |
| 6 | | | | | | | | | | | | | | |

325

For the order for |E| = 4, $\kappa = 0$, apparently $\hat{\kappa}_{AIC}$ keeps overestimating the order in some degree, while $\hat{\kappa}_{BIC}$ as in example $\kappa = 3$ severely underestimate the order, presumably due to the excessive weight of the correcting factors $\frac{\log(n)}{2}$. On the contrary $\hat{\kappa}_{EDC}$ and $\hat{\kappa}_{GDL}$ behaves quite well in same setting.

330 References

- [1] H. Akaike, "A New Look at the Statistical Model Identification", *IEEE Trans. Autom. Cont., vol. 19*, 1998.
- [2] H. Akaike & G. Kitagawa, (Eds.), "The Practice of Time Series Analysis", Springer-Verlag, New York, 1974.
- [3] M. S. Ali & D. Silvey, "A general class of coefficients of divergence of one distribution from another", *Journal of the Royal Statistical Society*, *Ser. B, No. 28, pp. 131-140*, 1966.
- T. W. Anderson & Leo A. Goodman, "Statistical Inference about Markov Chains", Annals of Mathematical Statistics, vol. 28, 1957.
- [5] Tom M. Apostol, "Calculus", John Wiley & Sons, Vol. 1, Second Edition, 1967.
- ³⁴² [6] A.R. Baigorri, C.R. Gonçalves & P.A. Resende, "Markov Chain Order ³⁴³ Estimation and χ^2 -divergence measure", arXiv:0910.0264v4[math.ST], ³⁴⁴ 2012.

- [7] A. Banerjee, X. Guo & H. Wang, "The Optimallity of Conditional Expectation as a Bregman Predictor", *IEEE Transactions on Information Theory, vol. 51, No 7*, 2005.
- [8] M. S. Bartlett, "The Frequency Goodness of Fit Test for Probability Chains", Math. Proc. Camb. Phil. Soc., vol. 9, 1951.
- [9] P. Billingsley, "Statistical Methods in Markov Chains", Annals of Mathematical Statistics, Vol. 32, No 1, 1961.
- [10] P. Billingsley, "Statistical Inference for Markov Chains", University of Chicago Press, Chicago, 1961.
- [11] W. Cochran, "The χ^2 Test of Goodness of Fit", Annals of Mathematical Statistics, vol. 23, No 3, 1952.
- [12] I. Csiszar, "Information-type measures of difference of probability distributions and indirect observation", *Studia Sci. Math. Hungar.*, Vol. 2, pp. 229-318, 1967.
- [13] I. Csiszar & P. Shields, "The Consistency of the BIC Markov Order
 Estimator", *The Annals of statistics, Vol 28, No 6, pp. 1601-1619*, 2000.
- [14] I. Csiszar & P. Shields, "Information Theory and Statistics: A Tutorial",
 Foundations and Trends in Communications and Information Theory,
 Vol. 1, No 4, pp. 417-528, 2004.
- [15] J.L. Doob, "Stochastic Processes(Wiley Publication in Statistics)", John
 Wiley & Sons. Inc., 1966.
- [16], C.C.Y. Dorea, "Optimal Penalty Term for EDC Markov Chain Order
 Estimator", Annales de l'Institut de Statistique de l'Universite de Paris
 (l'ISUP), v. 52, p. 15-26, 2008.
- [17] I.J. Good, "The Likelihood Ratio Test for Markov Chains", *Biometrica*, vol. 42, (3/4), pp. 531-533, 1955.
- ³⁷¹ [18] P. G. Hoel, "A test for Markov Chains", *Biometrika*, vol. 41, 1954.
- [19] R. W. Katz, "On Some Criteria for Estimating the Order Of Markov Chains", *Technometrics, vol. 23(3)*, 1981.
- S.P. Meyn & R.L. Tweedie, "Markov Chain and Stochastic Stability",
 Springer-Verlag, London, 1993.

- I. Pardo, "Statistical Inference Based on Divergence Measures", Taylor
 & Francis Group, LLC, New York, 2006.
- R Development Core Team, "R: A language and environment for statistical computing", R Foundation for Statistical Computing, Vienna, Austria. ISBN 3-900051-07-0, URL http://www.R-project.org, 2008.
- [23] A. E. Raftery, "A model for High-Order Markov Chains", Journal of the Royal Statistical Society, Serie B, vol. 47, No. 3, 1985.
- ³⁸³ [24] G. Schwarz, "Estimating the Dimension of a Model", The Annals of
 ³⁸⁴ Statistics, vol. 6, No 2, 1978.
- R. Shibata, "Selection of the Order of an Autoregressive model by
 Akaike's Information Criterion.", *Biometrika, vol. 6*, 1976.
- [26] H. Tong, "Determination of the order of a Markov Chain by Akaikes
 Information Criterion", Journal of Applied Probability, vol. 12, pp. 488-497, 1975.
- [27] H. C. Zhao, C.C.Y. Dorea & C.R. Gonçalves, "On Determination of the
 Order of a Markov Chain", *Stat. Infer. for Stoc. Processes, vol.* 4, 2001.