

On the Existence of Telescopers for Mixed Hypergeometric Terms

Shaoshi Chen^a, Frédéric Chyzak^b
Ruyong Feng^c, Guofeng Fu^c, Ziming Li^c

^a*Department of Mathematics, North Carolina State University, 27695-8205 Raleigh, USA*

^b*INRIA, 91120 Palaiseau, France*

^c*Key Laboratory of Math.-Mech., Chinese Academy of Sciences, 100190 Beijing, China*

Abstract

We present a criterion for the existence of telescopers for mixed hypergeometric terms, which is based on multiplicative and additive decompositions. The criterion enables us to determine the termination of Zeilberger's algorithms for mixed hypergeometric inputs.

Key words: Creative telescoping, Zeilberger's algorithms, Existence criteria.

1. Introduction

Given a sum $U_n := \sum_{k=0}^{\infty} u_{n,k}$ to be computed, *creative telescoping* is a process that determines a recurrence in n satisfied by the univariate sequence $U = (U_n)$ from a system of recurrences in n and k satisfied by the bivariate summand $u = (u_{n,k})$. A natural counterpart exists for integration. Algorithmic research on this topic has been initiated by Zeilberger in the early 1980s, leading in the 1990s to algorithms for summands and integrands described by first-order equations, that is for hypergeometric and hyperexponential creative telescoping (Zeilberger, 1990a, 1991; Almkvist and Zeilberger, 1990).

The termination problem of Zeilberger's algorithms has been extensively studied in the last two decades (Wilf and Zeilberger, 1992b; Abramov and Le, 2002; Abramov, 2003;

Email addresses: schen@amss.ac.cn, frederic.chyzak@inria.fr (Shaoshi Chen^a, Frédéric Chyzak^b), ryfeng@amss.ac.cn, fuguofeng@mmrc.iss.ac.cn, zmli@mmrc.iss.ac.cn (Ruyong Feng^c, Guofeng Fu^c, Ziming Li^c).

¹ In this work, S. Chen was supported by the NSF grant CCF-1017217. F. Chyzak was supported in part by the Microsoft Research-INRIA Joint Centre. G. Fu, R. Feng and Z. Li were supported in part by two grants of NSFC No. 60821002/F02 and No. 10901156.

Chen et al., 2005) and can be related to existence problems for other operations, like the computation of diagonals (Lipshitz, 1988). The main output of creative telescoping is the recurrence on the sum U . It is called a *telescoper* of u . Zeilberger’s algorithms terminate if and only if telescopers exist, whence the interest to discuss their existence. Zeilberger (1990b) shows that holonomicity, a notion borrowed from the theory of D-modules, implies the existence of telescopers. In particular, Wilf and Zeilberger (1992a) present an elementary proof, based on the ideas of Fasenmyer (1947) and Verbaeten (1974), that telescopers always exist for proper hypergeometric terms. However, holonomicity is merely a sufficient condition, i.e., there are cases in which the input functions are not holonomic but Zeilberger’s algorithms still terminate, see Chyzak et al. (2009). Therefore, a challenging problem is to find a necessary and sufficient condition that enables us to determine the existence of telescopers.

In view of the theoretical difficulty, special attention has been focused on the subclass of hypergeometric terms, hyperexponential functions, and mixed hypergeometric terms (see the definition in Section 2.4). In the continuous case, the results by Bernšteĭn (1971), Kashiwara (1978), Lipshitz (1988) and Takayama (1992) show that every hyperexponential function has a telescoper. This implies that Zeilberger’s algorithms always succeed on hyperexponential inputs. However, the situation in other cases turns out to be more involved. In the discrete case, the first complete solution to the termination problem has been given by Le (2001) and Abramov and Le (2002), by deciding whether telescopers exist for a given bivariate rational sequence in the (q) -discrete variables y_1 and y_2 . According to their criterion, the rational sequence

$$f = \frac{1}{y_1^2 + y_2^2}$$

has no telescoper. The criterion has been extended to the general case of bivariate hypergeometric terms by Abramov (2002, 2003). He proved that a hypergeometric term can be written as a sum of a hypergeometric-summable term and a proper one if it has a telescoper, see (Abramov, 2003, Theorem 10). Similar results have been obtained in the q -discrete case by Chen et al. (2005).

Almkvist and Zeilberger (1990) presented a continuous-discrete analogue of creative telescoping. This analogue is useful in the study of orthogonal polynomials (Koepf, 1998, Chapters 10–13). In analogy with the discrete case, not all mixed hypergeometric terms have telescopers. Therefore, an Abramov-like criterion is also needed in the mixed case.

In order to unify the various cases of mixed rational terms, Chen and Singer (2012) recently presented a criterion that is based on residues analysis for the existence of telescopers for bivariate rational functions. In the present paper, we give a criterion, Theorem 5.12, on the existence of telescopers for mixed hypergeometric terms, including continuous-discrete, continuous- q -discrete and discrete- q -discrete terms. Beside the termination problem of creative telescoping, the criterion, together with the results in (Hardouin and Singer, 2008; Schneider, 2010), can be used to determine if indefinite sums and integrals satisfy (possibly nonlinear) differential equations (see Example 6.4).

The rest of this paper is organized as follows. An algebraic setting for mixed hypergeometric terms is described in Section 2, and the existence problem of telescopers is stated in Section 3. We define the notion of exact terms, and introduce the additive decompositions in Section 4. The criterion for the existence of telescopers for mixed hypergeometric terms is presented in Section 5. Based on the criterion, we develop an algorithm for deciding the existence of telescopers, and present a few examples in Section 6.

2. Preliminaries

Throughout the paper, we let k be an algebraically closed field of characteristic zero, and q be a nonzero element of k . Assume further that q is not a root of unity. Let $k(x, y)$ be the field of rational functions in x and y over k . For a nonzero element $f \in k(x, y)$, the denominator and numerator of f are denoted by $\text{den}(f)$ and $\text{num}(f)$, respectively. They are two coprime polynomials in $k[x, y]$. For a ring R , R^* stands for $R \setminus \{0\}$. The symbol \mathbb{N} stands for the set of nonnegative integers.

This section contains five subsections. In §2.1, we describe a field that will serve as ground field in our subsequent algebraic constructions, and we define a (noncommutative) ring of Ore polynomials. In §2.2, we define the notion of mixed hypergeometric terms, and describe a (commutative) ring extension of the ground field. The ring extension contains mixed hypergeometric terms that occur in the study of existence of telescopers. In §2.3, we recall from (Chen et al., 2011) the structure of compatible rational functions, which leads to a multiplicative decomposition of mixed hypergeometric terms given in §2.4. Finally, we define in §2.5 the notion of split rational functions and the notions of proper terms that are meaningful in presence of two types of operators.

2.1. Fields endowed with a pair of operators

Let $\delta_x = \partial/\partial x$ and $\delta_y = \partial/\partial y$ be the usual derivations with respect to x and y , respectively. For an element $f \in k(x, y)$, we define the shift operators σ_x and σ_y as

$$\sigma_x(f(x, y)) = f(x + 1, y) \quad \text{and} \quad \sigma_y(f(x, y)) = f(x, y + 1),$$

and q -shift operators τ_x and τ_y as

$$\tau_x(f(x, y)) = f(qx, y) \quad \text{and} \quad \tau_y(f(x, y)) = f(x, qy).$$

To describe the mixed cases concisely, we introduce the following notation.

Notation. Let Θ denote the set

$$\{\delta_x, \sigma_x, \tau_x\} \times \{\delta_y, \sigma_y, \tau_y\} \setminus \{(\delta_x, \delta_y), (\sigma_x, \sigma_y), (\tau_x, \tau_y)\}.$$

A pair $(\xi_x, \eta_y) \in \Theta$ is called a *mixed* pair of operators.

Note that, for every $(\xi_x, \eta_y) \in \Theta$, ξ_x and η_y act on variables of different types. It follows that $\xi_x \circ \eta_y(f) = \eta_y \circ \xi_x(f)$ for all $f \in k(x, y)$.

In the sequel, $k(x, y)$ is usually endowed with a mixed pair (ξ_x, η_y) of operators. The resulting structure is denoted as $(k(x, y), (\xi_x, \eta_y))$. There are six cases listed in Figure 1, in which D- Δ stands for the differential and difference case, D- Δ_q for the differential and q -difference case, etc. Of course, the last three cases can be identified with the first three when we swap x and y in the field $k(x, y)$.

Given a field $(k(x, y), (\xi_x, \eta_y))$, one can define a ring of Ore polynomials (Chyzak and Salvy, 1998), which we denote here by $k(x, y)\langle \partial_x, \partial_y \rangle$. According to the different choices of (ξ_x, η_y) , we illustrate the corresponding rings of Ore polynomials in Figure 2. In particular, the commutation rules for the explicit case $f = x$ are as follows:

$$D_x x = x D_x + 1, \quad S_x x = (x + 1) S_x, \quad T_x x = q x T_x, \quad D_y x = x D_y, \quad S_y x = x S_y, \quad T_y x = x T_y.$$

Case	(ξ_x, η_y)	Field endowed with (ξ_x, η_y)
D- Δ	(δ_x, σ_y)	$(k(x, y), (\delta_x, \sigma_y))$
D- Δ_q	(δ_x, τ_y)	$(k(x, y), (\delta_x, \tau_y))$
Δ - Δ_q	(σ_x, τ_y)	$(k(x, y), (\sigma_x, \tau_y))$
Δ -D	(σ_x, δ_y)	$(k(x, y), (\sigma_x, \delta_y))$
Δ_q -D	(τ_x, δ_y)	$(k(x, y), (\tau_x, \delta_y))$
Δ_q - Δ	(τ_x, σ_y)	$(k(x, y), (\tau_x, \sigma_y))$

Fig. 1. Fields endowed with a mixed pair of operators

Case	(ξ_x, η_y)	Ring $k(x, y)\langle \partial_x, \partial_y \rangle$	Commutation rules, where $f \in k(x, y)$
D- Δ	(δ_x, σ_y)	$k(x, y)\langle D_x, S_y \rangle$	$D_x f = f D_x + \delta_x(f)$ and $S_y f = \sigma_y(f) S_y$
D- Δ_q	(δ_x, τ_y)	$k(x, y)\langle D_x, T_y \rangle$	$D_x f = f D_x + \delta_x(f)$ and $T_y f = \tau_y(f) T_y$
Δ - Δ_q	(σ_x, τ_y)	$k(x, y)\langle S_x, T_y \rangle$	$S_x f = \sigma_x(f) S_x$ and $T_y f = \tau_y(f) T_y$
Δ -D	(σ_x, δ_y)	$k(x, y)\langle S_x, D_y \rangle$	$S_x f = \sigma_x(f) S_x$ and $D_y f = f D_y + \delta_y(f)$
Δ_q -D	(τ_x, δ_y)	$k(x, y)\langle T_x, D_y \rangle$	$T_x f = \tau_x(f) T_x$ and $D_y f = f D_y + \delta_y(f)$
Δ_q - Δ	(τ_x, σ_y)	$k(x, y)\langle T_x, S_y \rangle$	$T_x f = \tau_x(f) T_x$ and $S_y f = \sigma_y(f) S_y$

Fig. 2. Rings of Ore polynomials

2.2. First-order mixed systems

A first-order mixed linear-functional system is of the form

$$\begin{cases} \xi_x(z) = az, \\ \eta_y(z) = bz, \end{cases} \quad (1)$$

where $(\xi_x, \eta_y) \in \Theta$ and $a, b \in k(x, y)$. For brevity, we call (1) a *first-order mixed system* or a *mixed system* in the sequel.

Example 2.1. Let $(\xi_x, \eta_y) = (\delta_x, \sigma_y)$, $a = y/x$ and $b = -x$. The system (1) becomes

$$\begin{cases} \delta_x(z) = \frac{y}{x} z, \\ \sigma_y(z) = -x z. \end{cases}$$

It is straightforward to verify that the expression $(-x)^y$ solves this mixed system. Moreover, this system does not have any nonzero rational solution in $k(x, y)$: if it had, we could write such a solution in the form

$$z = \frac{P}{Q} = \frac{p_m y^m + \cdots + p_0}{q_n y^n + \cdots + q_0}, \quad \text{where } p_i \text{ and } q_j \text{ are in } k(x) \text{ with } p_m q_n \neq 0.$$

By the equality $\sigma_y(z) = -xz$, we have $\sigma_y(P)Q = -xP\sigma_y(Q)$. Equating the leading

coefficients with respect to y yields $p_m q_n = -x p_m q_n$, which further implies that $x = -1$. This is a contradiction with the assumption that x is transcendental over k .

The example above shows that solving generally requires to extend the field $k(x, y)$. This motivates us to consider ring extensions of $k(x, y)$ endowed with a mixed pair of operators.

Definition 2.2. For a pair $(\xi_x, \eta_y) \in \Theta$, we call a triple $(R, (\bar{\xi}_x, \bar{\eta}_y))$ a *ring extension* of $(k(x, y), (\xi_x, \eta_y))$ if the following conditions are satisfied.

- (i) R is a commutative ring containing $k(x, y)$.
- (ii) $\bar{\xi}_x : R \rightarrow R$ is an extension of ξ_x , and $\bar{\eta}_y : R \rightarrow R$ is an extension of η_y .
- (iii) $\bar{\xi}_x$ is a derivation on R if $\xi_x = \delta_x$, and it is a monomorphism if $\xi_x = \sigma_x$ or $\xi_x = \tau_x$.
- (iv) $\bar{\eta}_y$ is a derivation on R if $\eta_y = \delta_y$, and it is a monomorphism if $\eta_y = \sigma_y$ or $\eta_y = \tau_y$.
- (v) $\bar{\xi}_x$ and $\bar{\eta}_y$ commute.

Moreover, such a ring extension is said to be *simple* if there does not exist any ideal I of R such that $\bar{\xi}_x(I) \subset I$ and $\bar{\eta}_y(I) \subset I$ except for $I = R$ and $I = \{0\}$.

Without any possible ambiguity, we shall denote the operators $\bar{\xi}_x$ and $\bar{\eta}_y$ obtained as in the definition above by ξ_x and η_y , respectively. The reader may find more general ring extensions endowed with derivations, shift and q -shift operators in (Hardouin and Singer, 2008).

Let $L = \sum_{i,j} a_{i,j} \partial_x^i \partial_y^j$ be an Ore polynomial in $k(x, y)\langle \partial_x, \partial_y \rangle$, where $k(x, y)$ is endowed with a mixed pair (ξ_x, η_y) of operators. Let $(R, (\xi_x, \eta_y))$ be a ring extension of $(k(x, y), (\xi_x, \eta_y))$. We define the application of L to an element $r \in R$ to be

$$L(r) = \sum_{i,j} a_{i,j} \xi_x^i \circ \eta_y^j(r).$$

It is straightforward to verify that, for $L_1, L_2 \in k(x, y)\langle \partial_x, \partial_y \rangle$,

$$L_1(L_2(r)) = (L_1 L_2)(r) \quad \text{for all } r \in R.$$

We are about to define the constants of a given field $(k(x, y), (\xi_x, \eta_y))$, by describing them in a uniform way as the solutions of specific operators. By application of elements in $k(x, y)\langle \partial_x, \partial_y \rangle$, we have

$$\Delta_y := \partial_y - \partial_y(1) = \begin{cases} D_y & \text{if } \eta_y = \delta_y, \\ S_y - 1 & \text{if } \eta_y = \sigma_y, \\ T_y - 1 & \text{if } \eta_y = \tau_y. \end{cases} \quad (2)$$

The same holds when we replace y by x and η_y by ξ_x , and we define Δ_x similarly. An element $c \in R$ is then called a *constant* with respect to the pair (ξ_x, η_y) if

$$\Delta_x(c) = \Delta_y(c) = 0.$$

One can easily verify that $c \in k(x, y)$ is a constant with respect to (ξ_x, η_y) if and only if c is an element of k .

Given a system of the form (1), a basic question is whether there exists a ring extension $(R, (\xi_x, \eta_y))$ containing a nonzero solution of the system. This question is related to compatibility conditions of (1), which we discuss in the next section.

Case	(ξ_x, η_y)	Compatibility condition
D- Δ	(δ_x, σ_y)	$\frac{\delta_x(b)}{b} = \sigma_y(a) - a$ and $b \neq 0$
D- Δ_q	(δ_x, τ_y)	$\frac{\delta_x(b)}{b} = \tau_y(a) - a$ and $b \neq 0$
Δ - Δ_q	(σ_x, τ_y)	$\frac{\sigma_x(b)}{b} = \frac{\tau_y(a)}{a}$ and $ab \neq 0$
Δ -D	(σ_x, δ_y)	$\sigma_x(b) - b = \frac{\delta_y(a)}{a}$ and $a \neq 0$
Δ_q -D	(τ_x, δ_y)	$\tau_x(b) - b = \frac{\delta_y(a)}{a}$ and $a \neq 0$
Δ_q - Δ	(τ_x, σ_y)	$\frac{\tau_x(b)}{b} = \frac{\sigma_y(a)}{a}$ and $ab \neq 0$

Fig. 3. Compatibility conditions

Case	(ξ_x, η_y)	$a, b \in k(x, y)$ compatible w.r.t. (ξ_x, η_y)
D- Δ	(δ_x, σ_y)	There exist $f \in k(x, y)$, $\alpha, \beta \in k(x)$, and $\gamma \in k(y)$ such that $a = \frac{\delta_x(f)}{f} + y \frac{\delta_x(\beta)}{\beta} + \alpha \quad \text{and} \quad b = \frac{\sigma_y(f)}{f} \cdot \beta \cdot \gamma.$
D- Δ_q	(δ_x, τ_y)	There exist $f \in k(x, y)$, $\alpha \in k(x)$, and $\beta \in k(y)$ such that $a = \frac{\delta_x(f)}{f} + \alpha \quad \text{and} \quad b = \frac{\tau_y(f)}{f} \cdot \beta.$
Δ - Δ_q	(σ_x, τ_y)	There exist $f \in k(x, y)$, $\alpha \in k(x)$, and $\beta \in k(y)$ such that $a = \frac{\sigma_x(f)}{f} \cdot \alpha \quad \text{and} \quad b = \frac{\tau_y(f)}{f} \cdot \beta.$
Δ -D	(σ_x, δ_y)	There exist $f \in k(x, y)$, $\alpha, \beta \in k(y)$, and $\gamma \in k(x)$ such that $a = \frac{\sigma_x(f)}{f} \cdot \beta \cdot \gamma \quad \text{and} \quad b = \frac{\delta_y(f)}{f} + x \frac{\delta_y(\beta)}{\beta} + \alpha.$
Δ_q -D	(τ_x, δ_y)	There exist $f \in k(x, y)$, $\alpha \in k(y)$, and $\beta \in k(x)$ such that $a = \frac{\tau_x(f)}{f} \cdot \beta \quad \text{and} \quad b = \frac{\delta_y(f)}{f} + \alpha.$
Δ_q - Δ	(τ_x, σ_y)	There exist $f \in k(x, y)$, $\alpha \in k(y)$, and $\beta \in k(x)$ such that $a = \frac{\tau_x(f)}{f} \cdot \beta \quad \text{and} \quad b = \frac{\sigma_y(f)}{f} \cdot \alpha.$

Fig. 4. Structure of compatible rational functions

2.3. Compatible rational functions

Let $k(x, y)$ be endowed with a mixed pair (ξ_x, η_y) of operators. If the first-order mixed system (1) has a nonzero solution h in a ring extension $(R, (\xi_x, \eta_x))$, by the commutativity of ξ_x and η_y , we have $\eta_y \circ \xi_x(h) = \xi_x \circ \eta_y(h)$. It follows from (1) that

$$\eta_y(ah) = \xi_x(bh), \quad a \neq 0 \text{ if } \xi_x \neq \delta_x, \text{ and } b \neq 0 \text{ if } \eta_y \neq \delta_y.$$

The above three conditions lead to six pairs of compatibility conditions listed in Figure 3. These are necessary conditions for System (1) to possess a solution.

Definition 2.3. Let $a, b \in k(x, y) \times k(x, y)$ and $(\xi_x, \eta_y) \in \Theta$. We say that a and b are *compatible* with respect to (ξ_x, η_y) if the compatibility conditions corresponding to (ξ_x, η_y) in Figure 3 are satisfied.

Theorem 1 in (Chen et al., 2011) describes the special structure of compatible rational functions. Applying this theorem to the six cases in Figure 3, we obtain Figure 4, which describes the structure of compatible bivariate rational functions. In fact, the conclusions in Figure 4 can also be derived from Lemmas 3.1, 3.2 and 3.3 in (Chen et al., 2011), because we are only concerned with bivariate compatible rational functions.

A first-order mixed system of the form (1) is said to be *compatible* if its coefficients a and b are compatible with respect to (ξ_x, η_y) . A first-order mixed system must be compatible with respect to (ξ_x, η_y) if it has a nonzero solution in some ring extension of $(k(x, y), (\xi_x, \eta_y))$.

2.4. Mixed hypergeometric terms

Hypergeometric terms are a common abstraction of geometric terms and factorials. They play an important role in combinatorics. The continuous analogue of hypergeometric terms is hyperexponential functions: they generalize usual exponential functions and simple radicals. In this paper, we will consider a class of functions in x and y that are solutions of first-order mixed systems, and are therefore intermediate objects between hypergeometric terms and hyperexponential functions.

Given a compatible mixed system of the form (1), Theorem 2 in (Bronstein et al., 2005) implies that there exists a simple ring extension $(\mathcal{R}, (\xi_x, \eta_y))$ of $(k(x, y), (\xi_x, \eta_y))$ containing a nonzero solution of (1). Such a simple ring is called a *Picard-Vessiot extension* associated to (1). Moreover, the set of constants in \mathcal{R} is equal to k due to the assumption that k is algebraically closed. These facts allow us to define the notion of mixed hypergeometric terms in a ring setting as follows.

Definition 2.4. Let $k(x, y)$ be a field endowed with a mixed pair (ξ_x, η_y) of operators. Assume that $(\mathcal{R}, (\xi_x, \eta_y))$ is a simple ring extension of $(k(x, y), (\xi_x, \eta_y))$, and that the set of constants in \mathcal{R} is equal to k . A nonzero element h of \mathcal{R} is called a *mixed hypergeometric term* over $(k(x, y), (\xi_x, \eta_y))$ if there exist $a, b \in k(x, y)$ such that

$$\xi_x(h) = ah \quad \text{and} \quad \eta_y(h) = bh.$$

We call a the *certificate of h with respect to ξ_x* , and b the *certificate with respect to η_y* .

For brevity, a mixed hypergeometric term will be called a mixed term in the sequel.

Lemma 2.5. Let the ring extension $(\mathcal{R}, (\xi_x, \eta_y))$ be given as in Definition 2.4.

- (i) Every mixed term is invertible.
- (ii) If two mixed terms have the same certificates, then their ratio belongs to k .

Proof. Let h be a mixed term in \mathcal{R} , and I be the ideal generated by h in \mathcal{R} . Then $\xi_x(h)$ and $\eta_y(h)$ belong to I . It follows that $\xi_x(I) \subset I$ and $\eta_y(I) \subset I$. Since \mathcal{R} is simple and h is nonzero, $I = \mathcal{R}$, that is, $1 \in I$. Consequently, h is invertible. The first assertion holds.

Let h_1 and h_2 be two mixed terms in \mathcal{R} . If they have the same certificates, then h_1/h_2 is a constant by a straightforward calculation, that is to say, $h_1 = ch_2$ for some $c \in k$. The second assertion holds. \square

Viewing mixed terms in an abstract ring allows us to compute their sums, products and inverses legitimately. Moreover, we will never encounter any analytic considerations,

such as singularities and the regions of definition. This choice will not do any harm, as the problem we are dealing with is purely algebraic.

Two mixed terms h_1 and h_2 are said to be *similar* if the ratio h_1/h_2 is in $k(x, y)$. It is easy to verify that similarity is an equivalence relation.

When studying the existence of telescopers, we will encounter at most finitely many mixed terms that are dissimilar to each other. These terms can be regarded as elements in a simple ring, because a finite number of Picard–Vessiot extensions associated to compatible first-order mixed systems can be embedded into a simple ring (Li et al., 2006, §2.2). From now on, we assume that \mathcal{R} is given as in Definition 2.4. It will be sufficient to consider mixed terms in \mathcal{R} .

By the second assertion of Lemma 2.5, two mixed terms having the same certificates differ by a multiplicative constant. These constants are irrelevant to the main result of this paper. So we introduce a notation to suppress them.

Let h be a mixed term in \mathcal{R} with ξ_x -certificate a and η_y -certificate b . Then a and b are compatible with respect to (ξ_x, η_y) because of the commutativity of ξ_x and η_y . Set

$$\mathcal{H}(a, b) = \{ch \mid c \in k\}.$$

The set consists of zero and mixed terms in \mathcal{R} whose respective certificates are a and b . Clearly, $\mathcal{H}(a, b)$ is a one-dimensional linear subspace over k . In the sequel, whenever the notation $\mathcal{H}(a, b)$ is used, a and b are assumed to be compatible rational functions, and $\mathcal{H}(a, b) \subset \mathcal{R}$. In particular, for any rational function $f \in k(x, y)$, the set $f\mathcal{H}(a, b)$ is a subset of \mathcal{R} . Indeed, it is the one-dimensional linear subspace spanned by fh . Moreover, let h' be another mixed term in \mathcal{R} , and a' and b' be the corresponding certificates. We consider

$$\mathcal{H}(a, b)\mathcal{H}(a', b') = \{rr' \mid r \in \mathcal{H}(a, b), r' \in \mathcal{H}(a', b')\},$$

which is equal to the one-dimensional linear subspace spanned by hh' . By the definition of certificates, we also have

$$\xi_x(\mathcal{H}(a, b)) = a\mathcal{H}(a, b) \quad \text{and} \quad \eta_y(\mathcal{H}(a, b)) = b\mathcal{H}(a, b).$$

More rules for calculations with $\mathcal{H}(a, b)$ are given below.

Lemma 2.6. For a field $(k(x, y), (\xi_x, \eta_y))$, we let $\mathcal{H}(a, b)$ and $\mathcal{H}(a', b')$ be given above, and f be a nonzero rational function in $k(x, y)$.

(i) If $\xi_x = \delta_x$ and $\eta_y \in \{\sigma_y, \tau_y\}$, then

$$f\mathcal{H}(a, b) = \mathcal{H}\left(a + \frac{\xi_x(f)}{f}, b \frac{\eta_y(f)}{f}\right) \quad \text{and} \quad \mathcal{H}(a, b)\mathcal{H}(a', b') = \mathcal{H}(a + a', bb').$$

(ii) If both ξ_x and η_y are automorphisms, then

$$f\mathcal{H}(a, b) = \mathcal{H}\left(a \frac{\xi_x(f)}{f}, b \frac{\eta_y(f)}{f}\right) \quad \text{and} \quad \mathcal{H}(a, b)\mathcal{H}(a', b') = \mathcal{H}(aa', bb').$$

(iii) If $\xi_x \in \{\sigma_x, \tau_x\}$ and $\eta_y = \delta_y$, then

$$f\mathcal{H}(a, b) = \mathcal{H}\left(a \frac{\xi_x(f)}{f}, b + \frac{\eta_y(f)}{f}\right) \quad \text{and} \quad \mathcal{H}(a, b)\mathcal{H}(a', b') = \mathcal{H}(aa', b + b').$$

Proof. Let h be a mixed term in $\mathcal{H}(a, b)$. It is straightforward to verify that the ξ_x -certificate and η_y -certificate of fh are $a + \xi_x(f)/f$ and $b\eta_y(f)/f$, respectively, if ξ_x is δ_x and η_y is an automorphism in $\{\xi_x, \eta_y\}$. Let $h \in \mathcal{H}(a, b)$ and $h' \in \mathcal{H}(a', b')$ with $hh' \neq 0$. Then the ξ_x -certificate of hh' is $a + a'$, and its η_y -certificate is bb' . It follows that

$$\mathcal{H}(a, b)\mathcal{H}(a', b') = \mathcal{H}(a + a', bb').$$

The first assertion holds. The other two assertions can be proved in a similar way. \square

Lemma 2.7. Let h_1 and h_2 be two mixed terms over $(k(x, y), (\xi_x, \eta_y))$. If h_1 and h_2 are similar, then

- (i) $h_1 + h_2$ is either equal to zero or similar to h_1 ;
- (ii) for any $L \in k(x, y)\langle \partial_x, \partial_y \rangle$, $L(h_1)$ is either equal to zero or similar to h_1 .

Proof. Let $r \in k(x, y)$ be equal to h_1/h_2 . Then the first assertion follows from the equality $h_1 + h_2 = (1 + 1/r)h_1$. Since h_1 is a mixed term, its successive derivatives and (q -)shifts are all similar to h_1 . The second assertion holds. \square

Remark 2.8. Let h, h_1, h_2 be three mixed terms. If $h = h_1 + h_2$, then the three terms are similar. This is because $\xi_x(h) = \xi_x(h_1) + \xi_x(h_2)$ and $\eta_y(h) = \eta_y(h_1) + \eta_y(h_2)$.

The next example illustrates how a linear differential or recurrence operator applies to mixed terms.

Example 2.9. Let us consider how to apply D_x and S_x to $r\mathcal{H}(u, v)$ with $r, u, v \in k(x, y)$. First, $D_x(r\mathcal{H}(u, v)) = (\delta_x(r) + ru)\mathcal{H}(u, v)$. Putting $L_1 = D_x + u$, we rewrite the above relation as $D_x(r\mathcal{H}(u, v)) = L_1(r)\mathcal{H}(u, v)$. An easy induction shows that

$$D_x^i(r\mathcal{H}(u, v)) = L_i(r)\mathcal{H}(u, v),$$

where $L_i \in k(x, y)\langle D_x \rangle$ whose coefficients have a common denominator that divides some power of $\text{den}(u)$. Moreover, the denominator of $L_i(r)$ divides $(\text{den}(u)\text{den}(r))^{i+1}$.

Let $M_i = \left(\prod_{j=0}^{i-1} \sigma_x^j(u)\right) S_x^i$ for $i > 0$. Then

$$S_x^i(r\mathcal{H}(u, v)) = M_i(r)\mathcal{H}(u, v).$$

So the denominator of $M_i(r)$ divides the product $\text{den}\left(\prod_{j=0}^{i-1} \sigma_x^j(u)\right) \cdot \text{den}(\sigma_x^i(r))$.

According to the structure of compatible rational functions given in Figure 4, we obtain a multiplicative decomposition of mixed terms, which is described in Figure 5. In each case of the figure, a mixed term h is re-expressed by a *structural decomposition* as $h = fh'$ (see Definition 2.11 below). An interesting property of such a decomposition is that the certificates of h' are expressed by *univariate* functions α, β and γ , as given in Figure 5. This fact will be crucial to prove our existence criterion for telescopers.

Let us verify the conclusion in the differential-difference case in Figure 5. Assume that a and b are the ξ_x -certificate and η_y -certificate of h , respectively. According to the conclusion given in the differential-difference case in Figure 4, there exist $f \in k(x, y)$, $\alpha, \beta \in k(x)$, and $\gamma \in k(y)$ such that

$$a = \frac{\delta_x(f)}{f} + y \frac{\delta_x(\beta)}{\beta} + \alpha \quad \text{and} \quad b = \frac{\sigma_y(f)}{f} \cdot \beta \cdot \gamma.$$

Case	(ξ_x, η_y)	Mixed term h
D- Δ	(δ_x, σ_y)	$\exists f \in k(x, y), \alpha, \beta \in k(x), \gamma \in k(y), h \in f\mathcal{H}\left(y\frac{\delta_x(\beta)}{\beta} + \alpha, \beta\gamma\right)$
D- Δ_q	(δ_x, τ_y)	$\exists f \in k(x, y), \alpha \in k(x), \beta \in k(y), h \in f\mathcal{H}(\alpha, \beta)$
Δ - Δ_q	(σ_x, τ_y)	$\exists f \in k(x, y), \alpha \in k(x), \beta \in k(y), h \in f\mathcal{H}(\alpha, \beta)$
Δ -D	(σ_x, δ_y)	$\exists f \in k(x, y), \alpha, \beta \in k(y), \gamma \in k(x), h \in f\mathcal{H}\left(\beta\gamma, x\frac{\delta_y(\beta)}{\beta} + \alpha\right)$
Δ_q -D	(τ_x, δ_y)	$\exists f \in k(x, y), \alpha \in k(y), \beta \in k(x), h \in f\mathcal{H}(\beta, \alpha)$
Δ_q - Δ	(τ_x, σ_y)	$\exists f \in k(x, y), \alpha \in k(y), \beta \in k(x), h \in f\mathcal{H}(\beta, \alpha)$

Fig. 5. Decomposition of mixed terms

Thus, the conclusion in the first row of Figure 5 holds by the first assertion in Lemma 2.6. The rest of conclusions in Figure 5 can be verified likewise.

Remark 2.10. In the case when both ξ_x and η_y are shift operators, the Ore–Sato theorem says that every hypergeometric term can be decomposed into the product of a rational function and a factorial term, (see also Corollary 4 in (Abramov and Petkovšek, 2002b)). The criterion for the existence of telescopers given by Abramov (2003) is based on the Ore–Sato theorem.

Definition 2.11. Let h be a mixed term over $(k(x, y), (\xi_x, \eta_y))$, as given in Figure 5. We say that fh' is a *structural decomposition* of h if f is a rational function given in the row corresponding to (ξ_x, η_y) in Figure 5, and h' is the mixed term such that $h = fh'$.

2.5. Split polynomials

In the discrete case, integer-linear polynomials are used to describe the existence criterion for telescopers of hypergeometric terms. Split polynomials play a similar role as integer-linear polynomials in the mixed cases.

Definition 2.12. A polynomial $p \in k[x, y]$ is said to be *split* if it is of the form $p_1(x)p_2(y)$ with $p_1 \in k[x]$ and $p_2 \in k[y]$. More generally, a rational function $r \in k(x, y)$ is said to be *split* if it is of the form $r_1(x)r_2(y)$ with $r_1 \in k(x)$ and $r_2 \in k(y)$.

A rational function $f \in k(x, y)$ can always be decomposed as $f_1(x)f_2(y)f_3(x, y)$, where $f_1 \in k(x)$, $f_2 \in k(y)$ and neither $\text{num}(f_3)$ nor $\text{den}(f_3)$ has split factors except constants. We call f_1f_2 and f_3 the *split* and *non-split parts* of f , respectively. Both are defined up to a nonzero multiplicative constant.

Remark 2.13. For a polynomial $p \in k[x, y]$, one may decide whether it is split by comparing all monic normalized coefficients of p with respect to y : p is split if and only if all those are equal. In an implementation, one would abort early as soon as a mismatch is found.

The next definition is fundamental for our existence criterion.

Definition 2.14. Let h be a mixed term over $(k(x, y), (\xi_x, \eta_y))$. Assume that fh' is a structural decomposition of h . We say that h is *proper* if $\text{den}(f)$ is split.

The lemma below shows that the properness of a mixed term is independent of the choice of its structural decompositions.

Lemma 2.15. Let $h = f_1 h_1 = f_2 h_2$ be a mixed term over $(k(x, y), (\xi_x, \eta_y))$, where f_1, f_2 are in $k(x, y)$. Assume that $f_1 h_1$ is a structural decomposition of h . Moreover, assume that $h_1 \in \mathcal{H}(a_1, b_1)$ and $h_2 \in \mathcal{H}(a_2, b_2)$. Then

$$\text{den}(f_1) \text{ is split} \quad \text{if and only if} \quad \text{den}(f_2) \text{ is split,}$$

provided that one of the conditions in the following table holds:

$(\xi_x, \eta_y) = (\delta_x, \sigma_y)$	either $\text{den}(a_2)$ or b_2 is split;
$(\xi_x, \eta_y) = (\delta_x, \tau_y)$	either $\text{den}(a_2)$ or b_2 is split;
$(\xi_x, \eta_y) = (\sigma_x, \delta_y)$	either a_2 or b_2 is split;
$(\xi_x, \eta_y) = (\sigma_x, \tau_y)$	either a_2 or $\text{den}(b_2)$ is split;
$(\xi_x, \eta_y) = (\tau_x, \delta_y)$	either a_2 or $\text{den}(b_2)$ is split;
$(\xi_x, \eta_y) = (\tau_x, \sigma_y)$	either a_2 or b_2 is split.

Proof. Assume that $(\xi_x, \eta_y) = (\delta_x, \sigma_y)$. Then, by the conclusion in the first row of Figure 5, there exist $\alpha, \beta \in k(x)$ and $\gamma \in k(y)$ such that

$$a_1 = y \frac{\delta_x(\beta)}{\beta} + \alpha \quad \text{and} \quad b_1 = \beta \gamma.$$

To show the equivalence in the first assertion, it suffices to prove that f_1/f_2 is split.

Case 1. Assume that $\text{den}(a_2)$ is split. Since $f_1 h_1 = f_2 h_2$, we have

$$\frac{\delta_x(f_1)}{f_1} + y \frac{\delta_x(\beta)}{\beta} + \alpha = \frac{\delta_x(f_2)}{f_2} + a_2,$$

which implies that

$$\frac{\delta_x(f_1/f_2)}{f_1/f_2} = a_2 - y \frac{\delta_x(\beta)}{\beta} - \alpha \quad (3)$$

The denominator of the right-hand side of (3) is split, since $\text{den}(a_2)$ is split and α, β are in $k(x)$. Suppose that $p \in k[x, y]$ is a nontrivial and non-split irreducible factor of $\text{num}(f_1/f_2) \cdot \text{den}(f_1/f_2)$. Then p divides the denominator of the logarithmic derivative of f_1/f_2 , a contradiction to (3). Thus, f_1/f_2 is split.

Case 2. Assume that b_2 is split. Since $f_1 h_1 = f_2 h_2$, we have

$$\frac{\sigma_y(f_1)}{f_1} \beta \gamma = \frac{\sigma_y(f_2)}{f_2} b_2,$$

which implies that

$$\frac{\sigma_y(f_1/f_2)}{f_1/f_2} = \frac{b_2}{\beta \gamma}. \quad (4)$$

Note that the right-hand side of (4) is split. Suppose that $p \in k[x, y]$ is a nontrivial and non-split irreducible factor of the polynomial $\text{num}(f_1/f_2) \cdot \text{den}(f_1/f_2)$. Then there

exists $\ell \in \mathbb{N}$ such that

$$\sigma_y^\ell(p) \mid \text{num}(f_1/f_2) \cdot \text{den}(f_1/f_2) \quad \text{but} \quad \sigma_y^{\ell+1}(p) \nmid \text{num}(f_1/f_2) \cdot \text{den}(f_1/f_2).$$

This implies that $\sigma_y^{\ell+1}(p)$ is a factor of either the numerator or the denominator of the rational function $\sigma_y(f_1/f_2)/(f_1/f_2)$, a contradiction to (4). Again, f_1/f_2 is split.

A similar argument proves that the assertion holds in the cases that $(\xi_x, \eta_y) = (\delta_x, \tau_y)$ and $(\xi_x, \eta_y) = (\sigma_x, \tau_y)$. The assertion for other cases can be proved by interchanging the roles of x and y . \square

3. Telescopers for mixed hypergeometric terms

The method of creative telescoping was first formulated and popularized in a series of papers by Zeilberger and his collaborators in the early 1990's ([Almkvist and Zeilberger, 1990](#); [Zeilberger, 1990a,b, 1991](#); [Wilf and Zeilberger, 1992a](#)). To illustrate the idea of this method, we consider the problem of finding a linear recurrence equation for the integral (if there exists one):

$$H(x) := \int_0^{+\infty} h(x, y) dy,$$

where $h(x, y)$ is a mixed term over $(k(x, y), (\sigma_x, \delta_y))$. Suppose that all integrals occurring in the derivation below are well-defined. The key step of creative telescoping tries to find a nonzero linear recurrence operator $L(x, S_x)$ in $k(x)\langle S_x \rangle$ such that

$$L(x, S_x)(h) = D_y(g), \tag{5}$$

for some mixed term g over $k(x, y)$.

Applying the integral sign to both sides of (5) yields

$$L(x, S_x)(H(x)) = g(x, +\infty) - g(x, 0).$$

This further implies that $L(x, S_x)$ is indeed the recurrence relation satisfied by $H(x)$ under certain nice boundary condition, say $g(x, +\infty) = g(x, 0)$. For example, consider the integral

$$A(x) = \int_0^{+\infty} y^{x-1} \exp(-y) dy.$$

The differential variant [Almkvist and Zeilberger \(1990\)](#) of Zeilberger's algorithm brings us a pair (L, g) with

$$L = S_x - x \quad \text{and} \quad g = -y^x \exp(-y).$$

Note that $g(x, +\infty) = g(x, 0) = 0$, which implies that $L(A(x)) = A(x+1) - xA(x) = 0$. So we recognize that $A(x) = \Gamma(x)$ since the initial value $A(1)$ is equal to 1. For more interesting examples, see the appendix of ([Almkvist and Zeilberger, 1990](#)) or Koepf's book ([Koepf, 1998](#), Chapters 10–13).

Definition 3.1. Let h be a mixed term over $(k(x, y), (\xi_x, \eta_y))$. A nonzero linear operator $L(x, \partial_x) \in k(x)\langle \partial_x \rangle$ is called a *telescoper of type (∂_x, ∂_y)* for h if there exists another mixed term g such that

$$L(x, \partial_x)(h) = \Delta_y(g). \tag{6}$$

For a given mixed term, when does a telescoper of certain type exist? And how one can construct telescopers? These are two basic problems related to the method of creative telescoping. In the subsequent sections, we will answer the first one for the mixed cases. More precisely, we solve the following problem, which is equivalent to the termination problem of creative-telescoping algorithms for mixed inputs.

Existence Problem for Telescopers. For a mixed term h over $(k(x, y), (\xi_x, \eta_y))$, find a necessary and sufficient condition on the existence of telescopers of type (∂_x, ∂_y) for h .

4. Exact terms and additive decompositions

For a univariate hypergeometric term $H(y)$, the Gosper algorithm (Gosper, 1978) decides whether it is hypergeometric summable with respect to y , i.e., whether $H = (S_y - 1)(G)$ for some hypergeometric term G . Based on the Gosper algorithm, Zeilberger (1990a,b) developed his fast version of creative-telescoping algorithms for bivariate hypergeometric terms. Almkvist and Zeilberger (1990) presented a continuous analogue of the Gosper algorithm for deciding the hyperexponential integrability, which leads to a fast algorithm for hyperexponential telescoping. From the viewpoint of creative telescoping, the Gosper algorithm and its continuous analogue decide whether the identity operator, 1, is a telescoper for the inputs.

The following notion of *exact terms* is motivated in the differential case by the existence of an underlying exact form. This differential-form point of view was used in a recent work of one of the authors (Chen et al., 2012).

Definition 4.1. Let h be a mixed term over $(k(x, y), (\xi_x, \eta_y))$. We say that h is *exact with respect to ∂_y* if there exists a mixed term g such that $h = \Delta_y(g)$, where Δ_y is defined in (2).

Remark 4.2. In (Abramov and Petkovšek, 2002a; Geddes et al., 2004), an exact term is traditionally called *(q-)hypergeometric summable* term in the discrete case, and a *hyperexponential integrable* function in the continuous case, respectively. For each choice of ∂_x in $\{D_x, S_x, T_x\}$, it is clear that every exact term with respect to ∂_y has a telescoper of type (∂_x, ∂_y) : for instance 1 is such a telescoper.

The notion of spread polynomials is defined by Abramov (2003) for establishing his criterion on the existence of telescopers for hypergeometric terms. The following definition translates it into the continuous and q -discrete cases.

Definition 4.3. Let K be a field of characteristic zero and $\delta_z, \sigma_z, \tau_z$ be the usual derivation, shift and q -shift operators over $K[z]$, respectively. For a polynomial $a \in K[z]$, we say that:

- (i) a is δ_z -spread if every nontrivial irreducible factor of a has multiplicity > 1 ;
- (ii) a is σ_z -spread if, for every nontrivial irreducible factor b of a , $\sigma_z^i(b) \mid a$ for some nonzero integer i ;
- (iii) a is τ_z -spread if, for every nontrivial irreducible factor b of a with $z \nmid b$, $\tau_z^i(b) \mid a$ for some nonzero integer i .

(Note that Case (iii) of this definition makes no constraint on the multiplicity of z in b . This is because we shall only consider τ_z -spread *non-split* polynomials in what follows.)

The following proposition is a mixed analogue of Theorem 8 in (Abramov, 2003), which relates exact terms to spread polynomials.

Proposition 4.4. Let $h = fh'$ be a mixed term over $(k(x, y), (\xi_x, \eta_y))$, where $f \in k(x, y)$ and $h' \in \mathcal{H}(u, v)$. Then the following statements hold:

- (i) Case $\eta_y = \delta_y$: If $\text{den}(v)$ is split and h is exact with respect to D_y , then the non-split part of $\text{den}(f)$ is δ_y -spread.
- (ii) Case $\eta_y = \sigma_y$: If v is split and h is exact with respect to S_y , then the non-split part of $\text{den}(f)$ is σ_y -spread.
- (iii) Case $\eta_y = \tau_y$: If v is split and h is exact with respect to T_y , then the non-split part of $\text{den}(f)$ is τ_y -spread.

Proof. Let p be a non-split irreducible factor of $\text{den}(f)$ with $\deg_x p > 0$ and $\deg_y p > 0$.

To prove the first assertion, we assume that $\text{den}(v)$ is split and $h = D_y(g)$, where g is either equal to zero or similar to h . Then $g = rh'$ for some $r \in k(x, y)$. It follows from $h = D_y(g)$ that $fh' = D_y(rh')$, which implies that

$$f = \delta_y(r) + rv.$$

Since $\text{den}(v)$ is split, p is an irreducible factor of $\text{den}(r)$. So there exists an integer $i > 1$ such that $p^i \mid \text{den}(\delta_y(r))$ and $p^i \nmid \text{den}(r)$. Therefore, $p^i \nmid \text{den}(rv)$ and $p^i \mid \text{den}(f)$. The first assertion holds.

To prove the second assertion, we assume that v is split and $h = (S_y - 1)(g)$, where g is either zero or similar to h . Then $g = rh'$ for some $r \in k(x, y)$. From $h = (S_y - 1)(g)$, it follows that $fh' = (S_y - 1)(rh')$. Hence, we get

$$f = v\sigma_y(r) - r.$$

Since v is split, $p \mid \text{den}(r)$ or $p \mid \text{den}(\sigma_y(r))$. So, the set

$$L := \{\ell \in \mathbb{Z} \text{ such that } \sigma_y^\ell(p) \mid \text{den}(r)\}$$

is finite and nonempty (consider $0 \in L$ and $-1 \in L$, respectively). Therefore, there exist $i, j \in \mathbb{Z}$ with $i > j$ such that $i - 1 \in L$, $i \notin L$, $j - 1 \notin L$, $j \in L$, that is

$$\sigma_y^i(p) \mid \text{den}(\sigma_y(r)), \quad \sigma_y^i(p) \nmid \text{den}(r), \quad \sigma_y^j(p) \nmid \text{den}(\sigma_y(r)), \quad \text{and} \quad \sigma_y^j(p) \mid \text{den}(r).$$

It follows that from the above equation both $\sigma_y^i(p)$ and $\sigma_y^j(p)$ divide $\text{den}(f)$. If i is nonzero, then both p and $\sigma^i(p)$ divide $\text{den}(f)$, so the non-split part of $\text{den}(f)$ is σ -spread. Otherwise, j must be nonzero. Then the non-split part is again σ -spread since both $\sigma^j(p)$ and p divide $\text{den}(f)$. In any case, the second assertion holds.

The third assertion can be proved in the same vein as in the second case. We only need to note that an irreducible factor p of the non-split part of $\text{den}(f)$ has at least two terms, which implies that $\tau_y^i(p)$ and $\tau_y^j(p)$ are coprime if $i \neq j$. \square

The next notion to be introduced, related to exact terms, is that of additive decompositions.

An algorithm by Abramov and Petkovšek (2001, 2002a) decomposes a hypergeometric term $H(y)$ into the sum $\Delta_y(H_1) + H_2$, where H_2 is minimal in some sense. Such a decomposition is called an *additive decomposition* for H with respect to y . Abramov

and Petkovšek's algorithm generalizes the capability of the Gosper algorithm in the sense that H is hypergeometric summable if and only if H_2 is zero. In the continuous case, an algorithm to decompose a hyperexponential function $H(y)$ as $D_y(H_1) + H_2$, where H_1 and H_2 are either zero or hyperexponential, is part of the proof of Lemma 4.2 in [Davenport \(1986\)](#). This remained unknown to [Geddes et al. \(2004\)](#), who later described a similar additive decomposition as a continuous analogue of Abramov and Petkovšek's algorithm, but also proved that H_2 satisfies certain minimality requirement. On the other hand, a q -discrete analogue is presented in [\(Chen et al., 2005\)](#). When H is a rational function, additive decompositions are more classical; they were presented by [Ostrogradskii \(1845\)](#) and [Hermite \(1872\)](#) for the continuous case, and by [Abramov \(1975, 1995\)](#) for the discrete and q -discrete cases.

For a mixed term h over $(k(x, y), (\xi_x, \eta_y))$, we can perform three kinds of additive decompositions with respect to y according to the choice of η_y . We recall now the notions related to additive decompositions for later use. Additive decompositions will be defined in Definition 4.11 below, after three steps of preliminary material.

First, we borrow from [\(Abramov and Petkovšek, 2001, 2002a\)](#), [\(Geddes et al., 2004\)](#), and [\(Chen et al., 2005\)](#) different notions of reduced rational functions.

Definition 4.5. Let K be a field of characteristic zero and f be a rational function in $K(z)$. Denote by δ_z, σ_z , and τ_z the usual derivation, shift and q -shift operators with respect to z on $K(z)$, respectively, and set $a = \text{num}(f)$, $b = \text{den}(f)$. We say that

- (i) f is δ_z -reduced if $\text{gcd}(b, a - i\delta_z(b)) = 1$ for all $i \in \mathbb{Z}$;
- (ii) f is σ_z -reduced if $\text{gcd}(b, \sigma_z^i(a)) = 1$ for all $i \in \mathbb{Z}$; and
- (iii) f is τ_z -reduced if $\text{gcd}(b, \tau_z^i(a)) = 1$ for all $i \in \mathbb{Z}$.

The next lemma reveals a connection between reduced rational functions and split ones. Its proof is based on the structure of compatible rational functions given in Figure 4.

Lemma 4.6. Let $u, v \in k(x, y)$ be two compatible rational functions with respect to $(\xi_x, \eta_y) \in \Theta$.

- (1) Case $\eta_y = \delta_y$ and δ_y -reduced v . If $\xi_x \in \{\sigma_x, \tau_x\}$, then both u and $\text{den}(v)$ are split.
- (2) Case $\eta_y = \sigma_y$ and σ_y -reduced v . Then:
 - (2a) if $\xi_x = \delta_x$, then both $\text{den}(u)$ and v are split;
 - (2b) if $\xi_x = \tau_x$, then both u and v are split.
- (3) Case $\eta_y = \tau_y$ and τ_y -reduced v . Then:
 - (3a) if $\xi_x = \delta_x$, then both $\text{den}(u)$ and v are split;
 - (3b) if $\xi_x = \sigma_x$, then both u and v are split.

Proof. Assume that $\xi_x = \sigma_x$ and $\eta_y = \delta_y$. By the conclusion in the fourth row of Figure 4, there exist $f \in k(x, y)$, $\gamma \in k(x)$, $\alpha, \beta \in k(y)$ such that

$$u = \frac{\sigma_x(f)}{f} \beta \gamma \quad \text{and} \quad v = \frac{\delta_y(f)}{f} + x \frac{\delta_y(\beta)}{\beta} + \alpha. \quad (7)$$

Since v is δ_y -reduced, every irreducible factor of $\text{den}(f) \cdot \text{num}(f)$ is either in $k[x]$ or in $k[y]$ by Lemma 2 in [\(Geddes et al., 2004\)](#) and the second equality in (7). Hence, $\text{den}(v)$ is split. As $\text{den}(f) \cdot \text{num}(f)$ is split, it also follows from the first equality that u is split. The first assertion holds when $\xi_x = \sigma_x$. Using the conclusion in the fifth row of Figure 4 and a similar argument, one can show that the assertion holds when $\xi_x = \tau_x$.

To prove the second assertion, we first assume that $\xi_x = \delta_x$ and $\eta_y = \sigma_y$. By the conclusion in the first row of Figure 4, there exist $f \in k(x, y)$, $\alpha, \beta \in k(x)$, and $\gamma \in k(y)$ such that

$$u = \frac{\delta_x(f)}{f} + y \frac{\delta_x(\beta)}{\beta} + \alpha \quad \text{and} \quad v = \frac{\sigma_y(f)}{f} \beta \gamma. \quad (8)$$

Let p be a non-split nontrivial irreducible factor of $\text{den}(f) \cdot \text{num}(f)$. Then there exists a nonnegative integer i such that

$$\sigma_y^i(p) \mid \text{den}(f) \cdot \text{num}(f) \quad \text{and} \quad \sigma_y^{i+\ell}(p) \nmid \text{den}(f) \cdot \text{num}(f) \quad \text{for all } \ell > 0,$$

which, together with the second equality in (8), $\beta \in k(x)$ and $\gamma \in k(y)$, implies that

- $\sigma_y^{i+1}(p) \mid \text{num}(v)$ and $\sigma_y^j(p) \mid \text{den}(v)$ if

$$\sigma_y^i(p) \mid \text{num}(f), \quad \sigma_y^j(p) \mid \text{num}(f), \quad \text{but} \quad \sigma_y^{j-1}(p) \nmid \text{num}(f);$$

- $\sigma_y^{i+1}(p) \mid \text{den}(v)$ and $\sigma_y^j(p) \mid \text{num}(v)$ if

$$\sigma_y^i(p) \mid \text{den}(f), \quad \sigma_y^j(p) \mid \text{den}(f), \quad \text{but} \quad \sigma_y^{j-1}(p) \nmid \text{den}(f).$$

Either case leads to a contradiction to the assumption that v is σ_y -reduced. Hence, f is split and so is v . By the first equality in (8), $\text{den}(u)$ is split. This proves Assertion (2a).

Let $\xi_x = \tau_x$. By the conclusion in the sixth row of Figure 4, there exist $f \in k(x, y)$, $\alpha \in k(y)$ and $\beta \in k(x)$ such that

$$u = \frac{\tau_x(f)}{f} \beta \quad \text{and} \quad v = \frac{\sigma_y(f)}{f} \alpha. \quad (9)$$

The same argument used in Case (2a) implies that f is split, and so is v . By the first equality in (9), u is also split.

Finally, we let $\eta_y = \tau_y$. Then Assertions (3a) and (3b) hold by the conclusions in second and third rows of Figure 4 and similar arguments used in the proofs of Assertions (2a) and (2b), respectively. \square

Second, we recall the notions of squarefree, shift-free and q -shift-free polynomials.

Definition 4.7. Let K be a field of characteristic zero, and a be a nonzero polynomial in $K[z]$. Denote by δ_z, σ_z , and τ_z the usual derivation, shift and q -shift operators with respect to z on $K(z)$, respectively.

- (i) a is said to be *squarefree* if $\gcd(a, \delta_z(a)) = 1$.
- (ii) a is said to be *shift-free* if $\gcd(a, \sigma_z^i(a)) = 1$ for every nonzero integer i .
- (iii) Let $a = z^s \tilde{a}$ with $\tilde{a} \in K[z]$ and $z \nmid \tilde{a}$. Then a is said to be *q -shift-free* if

$$\gcd(\tilde{a}, \tau_z^i(\tilde{a})) = 1 \quad \text{for every nonzero integer } i.$$

Note that every non-split polynomial in $k[x, y]$ has at least two terms. For a polynomial in $K[z]$ having at least two terms, it is not δ_z -spread (resp. σ_z -spread, τ_z -spread) if it is squarefree (resp. shift-free, q -shift-free) with respect to z . However, the converses are false, as shown in the examples below.

Example 4.8. Let $p = z^2(z+1)$. Since the multiplicity of $z+1$ in p is 1, p is not δ_z -spread. However, p is not squarefree with respect to z , because $\gcd(p, \delta_z(p)) = z \notin K$.

Example 4.9. Let $p = z(z+1)(z+1/2)$. Since any nontrivial shift of $(z+1/2)$ does not divide p , p is not σ_z -spread. However, p is not shift-free with respect to z , because $\gcd(p, \sigma_z(p)) = z+1$.

Example 4.10. Let $p = (z+1)(z+q)(z+2)$ and $q \in K$ such that $q^i \neq 2$ for any $i \in \mathbb{Z}$. Since $\tau_z^i(z+2) \nmid p$ for any nonzero $i \in \mathbb{Z}$, p is not τ_z -spread. However, p is not q -shift-free with respect to z , because $\gcd(p, \tau_z(p)) = z+1$.

Finally, we define the three additive decompositions in the setting of mixed terms.

Definition 4.11. Let h be a mixed term over $(k(x, y), (\xi_x, \eta_y))$. Assume that

$$h = \Delta_y(h_1) + h_2 \tag{10}$$

where h_1 is a mixed term, and h_2 is equal to either zero or a mixed term of the form

$$h_2 \in r\mathcal{H}(u, v) \tag{11}$$

for some $r, u, v \in k(x, y)$ satisfying

- (a) $\text{den}(r)$ is squarefree with respect to y , and v is δ_y -reduced if $\eta_y = \delta_y$;
- (b) $\text{den}(r)$ is shift-free with respect to y , and v is σ_y -reduced if $\eta_y = \sigma_y$;
- (c) $\text{den}(r)$ is q -shift-free with respect to y , and v is τ_y -reduced if $\eta_y = \tau_y$.

We call (10) an *additive decomposition* of h with respect to ∂_y .

Additive decompositions with respect to D_y can be computed by the algorithms described in (Davenport, 1986) and (Geddes et al., 2004). Additive decompositions with respect to S_y and T_y can be computed by the algorithms in (Abramov and Petkovšek, 2001) and (Chen et al., 2005), respectively. In particular, additive decompositions always exist, although they are not unique. We remark that the additive decompositions given in Definition 4.11 are weaker than those in (Abramov and Petkovšek, 2001) and (Geddes et al., 2004). For example, h_2 is not necessarily equal to zero when h is an exact term.

5. A criterion on the existence of telescopers for mixed terms

The Fundamental Theorem in (Wilf and Zeilberger, 1992a) states that there exist telescopers for proper mixed terms. However, properness is just a sufficient condition. In this section, we present a necessary and sufficient condition on the existence of telescopers for mixed terms.

Definition 5.1. A mixed term h over $(k(x, y), (\xi_x, \eta_y))$ is said to be *telescoperable of type* (∂_x, ∂_y) if it has a telescoper of type (∂_x, ∂_y) .

In this section, we show that every proper mixed term is telescoperable in §5.1, and that every telescoperable term is a sum of an exact term and a proper one in §5.2.

5.1. Proper terms are telescoperable

Wilf and Zeilberger present an elementary proof of the existence of telescopers for proper hypergeometric terms (Wilf and Zeilberger, 1992a, Theorem 3.1), and indicate that their argument should be applicable to the mixed setting. For the sake of completeness, we elaborate a proof that every proper mixed term is telescoperable. Before presenting the proof, we need a few lemmas.

Lemma 5.2. Let h and H be two mixed terms over $(k(x, y), (\xi_x, \eta_y))$ with $\Delta_y(H) = 0$, where Δ_y is defined in (2). If h is telescopable of type (∂_x, ∂_y) , so is Hh .

Proof. Let $L \in k(x)\langle\partial_x\rangle$ be a telescoper of h such that $L(h) = \Delta_y(g)$ for some mixed term g . Assume that $H \in \mathcal{H}(a, b)$. Since $\Delta_y(H) = 0$, $\eta_y(H) = \eta_y(1)H$. Thus, $b = \eta_y(1)$. It follows that $b = 0$ if $\eta_y = \delta_y$, and that $b = 1$ if either $\eta_y = \sigma_y$ or $\eta_y = \tau_y$.

By the compatibility conditions on a and b given in Figure 3, we derive that a is nonzero and, depending on η_y , that one of the following formulas holds: (i) $\delta_y(a) = 0$; (ii) $\sigma_y(a) = a$; (iii) $\tau_y(a) = a$. At this point, we claim that $a \in k(x)$. This is clear in case (i). In case (ii) or (iii), write $a = u/v$ for coprime u and v , and v monic with respect to y . Then, either $u/v = \sigma_y(u)/\sigma_y(v)$ or $u/v = \tau_y(u)/\tau_y(v)$, where all fractions are reduced. As a consequence, either $u = \alpha\sigma_y(u)$ and $v = \beta\sigma_y(v)$, or $u = \alpha\tau_y(u)$ and $v = \beta\tau_y(v)$, in each case for α and β nonzero in $k(x)$. Case (ii) leads to $\alpha = \beta = 1$, and $a \in k(x)$. In case (iii), from q not being a root of unity follows that u and v are monomials in y . By coprimeness, at most one of them has nonzero degree in y . If $u \in k(x)$, then α/β is equal to q^j where $j = \deg_y v$, so that, as $\alpha/\beta = 1$, v is in $k(x)$ as well. A symmetric argument leads to the fact that both u and v are in $k(x)$, and so is a .

We continue by distinguishing two cases according to ξ_x :

Case 1. Assume that $\xi_x = \delta_x$. Then $\partial_x(Hh) = \partial_x(H)h + H\partial_x(h)$, so that $(\partial_x - a)(Hh) = H\partial_x(h)$. By an easy induction, this implies that

$$(\partial_x - a)^i(Hh) = H\partial_x^i(h) \quad \text{for all } i \in \mathbb{N}. \quad (12)$$

Let L' be the Ore polynomial obtained by replacing ∂_x^i in L with $(\partial_x - a)^i$. Then L' belongs to $k(x)\langle\partial_x\rangle$ because a does. By (12), $L'(Hh) = HL(h)$. Hence,

$$L'(Hh) = HL(h) = H\Delta_y(g) = \Delta_y(Hg),$$

in which the last equality follows from $\Delta_y(H) = 0$. The product Hh is telescopable.

Case 2. Assume that $\xi_x = \sigma_x$ or $\xi_x = \tau_x$. Then $\partial_x(Hh) = \partial_x(H)\partial_x(h)$. So

$$a^{-1}\partial_x(Hh) = H\partial_x(h),$$

which, together with an easy induction, implies that

$$\left(\prod_{j=0}^{i-1} \sigma_x^j(a)\right)^{-1} \partial_x^i(Hh) = H\partial_x^i(h) \quad \text{for all } i \in \mathbb{N}.$$

Let L'' be the Ore polynomial obtained by replacing ∂_x^i in L with $\left(\prod_{j=0}^{i-1} \sigma_x^j(a)\right)^{-1} \partial_x^i$. Then L'' belongs to $k(x)\langle\partial_x\rangle$ because a does. Consequently, L'' is a telescoper of Hh of type (∂_x, ∂_y) by the same argument as the one used in Case 1. \square

Lemma 5.3. Let h_1 and h_2 be two similar mixed terms over $(k(x, y), (\xi_x, \eta_y))$. Let \mathbb{V}_y be the $k(x)$ -vector space spanned by

$$\{\Delta_y^i(h_j) \mid i \in \mathbb{N}, j \in \{1, 2\}\},$$

where $\Delta_y = \Delta_y$. If both h_1 and h_2 have telescopers of type (∂_x, ∂_y) , so does every nonzero element in \mathbb{V}_y .

Proof. Assume that, for $j = 1, 2$, $L_j(h_j) = \Delta_y(g_j)$ for some $L_j \in k(x)\langle\partial_x\rangle$ and for some mixed term g_j . Then

$$L_j(\Delta_y^i(h_j)) = \Delta_y(\Delta_y^i(g_j)).$$

Hence, every nonzero element of $\{\Delta_y^i(h_j) \mid i \in \mathbb{N}, j \in \{1, 2\}\}$ is telescopable of (∂_x, ∂_y) . By Lemma 5.2, a mixed term of the form $f\Delta_y^i(h_j)$ for every $f \in k(x)^*$ is also telescopable of the same type. Assume that H_1 and H_2 are two telescopable mixed terms. Then a common left multiple of their telescopers is a telescoper of the sum $H_1 + H_2$. Hence, every nonzero element of \mathbb{V}_y is telescopable. \square

The “noncommutative trick” below was first used in (Wegschaider, 1997, Theorem 3.2) to transform a nonzero linear recurrence operator not involving y to a telescoper. Here, we generalize it to transform an operator in $k(x)\langle\partial_x, \partial_y\rangle$ to a telescoper of type (∂_x, ∂_y) . The result, Lemma 5.5, bases on formulas depending on the operator types, which we state first. However, only the cases when η_y is δ_y or σ_y are necessary in the proof of Lemma 5.6, so Lemmas 5.4 and 5.5 do not consider q -discrete operators.

Lemma 5.4. The following statements hold.

(1) *Case $\eta_y = \sigma_y$.* For any $m \in \mathbb{N}$, there exists $Q_m \in k[y]\langle S_y \rangle$ satisfying

$$\frac{y^m}{(-1)^m m!} (S_y - 1)^m = 1 + (S_y - 1)Q_m.$$

(2) *Case $\eta_y = \delta_y$.* For any $m \in \mathbb{N}$, there exists $Q_m \in k[y]\langle D_y \rangle$ satisfying

$$\frac{y^m}{(-1)^m m!} D_y^m = 1 + D_y Q_m.$$

Proof. The first assertion is just Theorem 3.2 in (Wegschaider, 1997). The second assertion follows from a straightforward continuous analogue of Wegschaider’s proof for the first assertion. \square

Lemma 5.5. Let h be a mixed term over $(k(x, y), (\xi_x, \eta_y))$, under the restriction $\eta_y \in \{\delta_y, \sigma_y\}$. If there exists a nonzero operator $A \in k(x)\langle\partial_x, \partial_y\rangle$ such that $A(h) = 0$, then h has a telescoper of type (∂_x, ∂_y) .

Proof. Assume that $A(h) = 0$ for some nonzero operator A in $k(x)\langle\partial_x, \partial_y\rangle$. By a repeated use of left-hand division, we can write

$$A = \Delta_y^m (L(x, \partial_x) + \Delta_y M), \tag{13}$$

where m is in \mathbb{N} , Δ_y is defined in (2), L is a nonzero operator in $k(x)\langle\partial_x\rangle$ and M is in $k(x)\langle\partial_x, \partial_y\rangle$. By Lemma 5.4, we have

$$\frac{y^m}{(-1)^m m!} \Delta_y^m = \Delta_y Q + 1$$

for some $Q \in k[y]\langle\partial_y\rangle$, which, together with (13), implies that

$$\frac{y^m}{(-1)^m m!} A = L + \Delta_y N \quad \text{for some } N \in k(x, y)\langle\partial_x, \partial_y\rangle.$$

It follows from $A(h) = 0$ that $L(h) = \Delta_y(-N(h))$. Hence, L is a telescoper for h of type (∂_x, ∂_y) . \square

The next lemma paves the way to apply the multiplicative decompositions in Figure 5. Its proof is reminiscent of the linear algebra argument given by [Lipshitz \(1988\)](#); however, it bases on linear algebra and filtrations over $k(x)$ instead of k .

Lemma 5.6. Let h be a mixed term over $(k(x, y), (\xi_x, \eta_y))$.

- (i) If $(\xi_x, \eta_y) = (\delta_x, \sigma_y)$ and $h \in \mathcal{H}\left(y \frac{\delta_x(\beta)}{\beta}, \beta\gamma\right)$ with $\beta \in k(x)$ and $\gamma \in k(y)$, then h has a telescoper of type (D_x, S_y) .
- (ii) If $(\xi_x, \eta_y) = (\sigma_x, \delta_y)$ and $h \in \mathcal{H}\left(\beta, x \frac{\delta_y(\beta)}{\beta} + \alpha\right)$ with $\alpha, \beta \in k(y)$, then h has a telescoper of type (S_x, D_y) .

Proof. To show the first assertion, we let

$$a = \text{num}(\beta), \quad b = \text{den}(\beta), \quad s = \text{num}(\gamma), \quad t = \text{den}(\gamma),$$

and

$$u = \text{num}\left(\frac{\delta_x(\beta)}{\beta}\right), \quad v = \text{den}\left(\frac{\delta_x(\beta)}{\beta}\right).$$

A straightforward calculation yields that the δ_x -certificate and σ_y -certificate of h are,

$$y \frac{\delta_x(\beta)}{\beta} = \frac{yu}{v} \quad \text{and} \quad \beta\gamma = \frac{as}{bt},$$

respectively. Note that $a, b, u, v \in k[x]$ and $s, t \in k[y]$.

Let \mathcal{F}_N be the linear subspace spanned by $\{D_x^i S_y^j \mid i + j \leq N\}$ over $k(x)$. Let μ be the maximum of the degrees in y of s and t , and let

$$\mathcal{W}_N = \text{span}_{k(x)} \left\{ \frac{y^\ell h}{t(y + N - 1) \cdots t(y)} \mid \ell \leq (\mu + 1)N \right\}.$$

An easy induction on i and j yields

$$D_x^i S_y^j(h) = \frac{w(x, y)h}{t(y + j - 1) \cdots t(y)}, \quad \text{where } w \in k(x)[y] \text{ and } \deg_y(w) \leq j\mu + i.$$

Hence, $D_x^i S_y^j(h)$ belongs to \mathcal{W}_N if $i + j \leq N$. Accordingly, there is a well-defined $k(x)$ -linear map ϕ_N from \mathcal{F}_N to \mathcal{W}_N that sends L to $L(h)$ for all $L \in \mathcal{F}_N$. Since the dimension of \mathcal{F}_N over $k(x)$ is $\binom{N+2}{2}$, while that of \mathcal{W}_N is $(\mu + 1)N + 1$, the kernel of ϕ_N is nontrivial when N is sufficiently large. Let A be a nonzero element of $\ker(\phi_N)$. Then $A(h) = 0$. The first assertion follows from Lemma 5.5.

To show the second assertion, we let $a = \text{num}(\beta)$, $b = \text{den}(\beta)$,

$$u = \text{num}\left(x \frac{\delta_y(\beta)}{\beta} + \alpha\right) \quad \text{and} \quad v = \text{den}\left(y \frac{\delta_y(\beta)}{\beta} + \alpha\right).$$

A straightforward calculation yields that the y -certificate of h is

$$x \frac{\delta_y(\beta)}{\beta} + \alpha = \frac{u(x, y)}{v(y)}.$$

Note that $a, b, v \in k[y]$, $u \in k[x, y]$.

Consider the linear space \mathcal{F}_N spanned by $\{S_x^i D_y^j \mid i + j \leq N\}$ over $k(x)$. Let μ be the maximum of the degrees in y of a, b, u and v , and let

$$\mathcal{W}_N = \text{span}_{k(x)} \left\{ \frac{y^\ell h}{(bv)^N} \mid \ell \leq 2\mu N \right\}.$$

An easy induction on i and j yields

$$S_x^i D_y^j(h) = \frac{w(x, y)}{b^i \gamma^j} h, \quad \text{where } w \in k(x)[y] \text{ with } \deg_y(w) \leq (i + j)\mu.$$

Hence, $S_x^i D_y^j(h)$ belongs to \mathcal{W}_N if $i + j \leq N$. Accordingly, there is well-defined a $k(x)$ -linear map ψ_N from \mathcal{F}_N to \mathcal{W}_N that sends L to $L(h)$ for all $L \in \mathcal{F}_N$. Since the dimension of \mathcal{F}_N over $k(x)$ is $\binom{N+2}{2}$, while that of \mathcal{W}_N is $2\mu N + 1$, the kernel of ψ_N is nontrivial when N is sufficiently large. Let A be a nonzero element in $\ker(\psi_N)$. Then $A(h) = 0$. The second assertion follows from Lemma 5.5. \square

We are ready to show the main conclusion of this subsection.

Theorem 5.7. Let h be a mixed term over $(k(x, y), (\xi_x, \eta_y))$. If h is proper, then it is telescopic of type (∂_x, ∂_y) .

Proof. In the mixed setting, there are six cases to be considered.

First, we assume that $\xi_x = \delta_x$ and $\eta_y = \sigma_y$, and prove that h has a telescoper of type (D_x, S_y) . By the conclusion in the first row of Figure 5 and Lemma 2.15, h has a structural decomposition

$$h \in \frac{A}{B \cdot C} \mathcal{H} \left(y \frac{\delta_x(\beta)}{\beta} + \alpha, \beta\gamma \right),$$

where $A \in k[x, y]$, $B \in k[x]$, $C \in k[y]$, $\alpha, \beta \in k(x)$ and $\gamma \in k(y)$. By Lemma 2.6 (i),

$$h = AUV, \quad \text{where } U \in \mathcal{H} \left(\alpha + \frac{\delta_x(1/B)}{1/B}, 1 \right), \text{ and } V \in \mathcal{H} \left(y \frac{\delta_x(\beta)}{\beta}, \beta \frac{\sigma_y(1/C)}{1/C} \gamma \right).$$

Write $A = \sum_{i=0}^m a_i y^i$ with $a_i \in k[x]$. Then we have $h = \sum_{i=0}^m G_i H_i$, where

$$G_i = a_i U \in \mathcal{H} \left(\alpha + \frac{\delta_x(a_i/B)}{a_i/B}, 1 \right) \text{ and } H_i = y^i V \in \mathcal{H} \left(y \frac{\delta_x(\beta)}{\beta}, \beta \frac{\sigma_y(y^i/C)}{y^i/C} \gamma \right).$$

Note that

$$\alpha + \frac{\delta_x(a_i/B)}{a_i/B} \in k(x) \quad \text{and} \quad \frac{\sigma_y(y^i/C)}{y^i/C} \gamma \in k(y).$$

By Lemma 5.6 (i), the term H_i has a telescoper of type (D_x, S_y) for all i with $0 \leq i \leq m$. Since $(S_y - 1)(G_i) = 0$, $G_i H_i$ has a telescoper of type (D_x, S_y) for all i with $0 \leq i \leq m$ by Lemma 5.2. So h has a telescoper of the same type by Lemma 5.3. This completes the proof for the first case.

Second, we assume that $\xi_x = \sigma_x$ and $\eta_y = \delta_y$, and show that h has a telescoper of type (S_x, D_y) . By the conclusion in the fourth row of Figure 5 and Lemma 2.15, h has a structural decomposition

$$h \in \frac{A}{B \cdot C} \mathcal{H} \left(\beta\gamma, \alpha + x \frac{\delta_y(\beta)}{\beta} \right),$$

where $A \in k[x, y]$, $B \in k[x]$, $C \in k[y]$, $\alpha, \beta \in k(y)$ and $\gamma \in k(x)$.

Write $A = \sum_{i=0}^m a_i y^i$ with $a_i \in k[x]$. A similar consideration as above leads to

$$h = \sum_{i=0}^m G_i H_i, \quad \text{where } G_i \in \mathcal{H} \left(\gamma \frac{\sigma_x(a_i/B)}{a_i/B}, 0 \right) \text{ and } H_i \in \mathcal{H} \left(\beta, \alpha + x \frac{\delta_y(\beta)}{\beta} \right).$$

Note that $\gamma\sigma_x(a_i/B)/(a_i/B) \in k(x)$ and $\alpha \in k(y)$. By Lemma 5.6 (ii), the term H_i has a telescoper of type (S_x, D_y) for all i with $0 \leq i \leq m$. Since $D_y(G_i) = 0$, G_iH_i has a telescoper of type (S_x, D_y) by Lemma 5.2, so does h by Lemma 5.3.

In the other four cases, (ξ_x, η_y) is equal to (δ_x, τ_y) , (σ_x, τ_y) , (τ_x, δ_y) and (τ_x, σ_y) , respectively. The presence of q -shift operators leads to simpler structural decompositions, which enable us prove the existence of telescopers in a fairly straightforward way.

By Lemma 2.15 and the conclusions in the second, third, fifth and sixth rows of Figure 5, h has a structural decomposition

$$h \in \frac{A}{B \cdot C} \mathcal{H}(\alpha, \beta),$$

where $A \in k[x, y]$, $B \in k[x]$, $C \in k[y]$, $\alpha \in k(x)$ and $\beta \in k(y)$.

Write $A = \sum_{i=0}^m a_i y^i$ with $a_i \in k[x]$. It follows from Lemma 2.6 that

$$h = \sum_{i=0}^m G_i H_i, \quad \text{where } G_i \in \frac{a_i}{B} \mathcal{H}(\alpha, 1) \text{ and } H_i \in \frac{y^i}{C} \mathcal{H}(1, \beta).$$

By Lemma 5.3, it suffices to show that, for all i with $0 \leq i \leq m$, $G_i H_i$ has a telescoper of type (∂_x, ∂_y) . Since $\Delta_y(G_i) = 0$, it suffices to show that H_i has a telescoper by Lemma 5.2. Since $\Delta_x(H_i) = 0$, Δ_x is a telescoper of H_i . \square

5.2. Characterization of telescoperable terms

We have seen that exact terms and proper ones are all telescoperable. We now show that a telescoperable term is the sum of an exact term and a proper one. To this end, we present a few useful lemmas.

Lemma 5.8. Let h, h_1 and h_2 be three mixed terms over $(k(x, y), (\xi_x, \eta_y))$. Assume that $h = \Delta_y(h_1) + h_2$. Then L is a telescoper of type (∂_x, ∂_y) for h if and only if it is a telescoper for h_2 .

Proof. If h has a telescoper L in $k(x)\langle\partial_x\rangle$, then $L(h) = \Delta_y(g)$ for some mixed term g . Since L and Δ_y commute with each other, $L(h_2) = \Delta_y(g - L(h_1))$. So L is a telescoper for h_2 . The converse can be proved by the same argument. \square

The following lemma is a q -analogue of Theorem 7 in (Abramov and Petkovšek, 2002b).

Lemma 5.9. Let p be an irreducible polynomial in $k[x, y]$. Then

- (i) If there exist $i, j \in \mathbb{Z}$ with $i \neq 0$ and $c \in k$ such that $\sigma_x^i \tau_y^j(p) = cp$, then $p \in k[y]$.
- (ii) If there exist $i, j \in \mathbb{Z}$ with $j \neq 0$ and $c \in k$ such that $\sigma_x^i \tau_y^j(p) = cp$, then $p = \lambda y$ for some $\lambda \in k$ or $p \in k[x]$.

Proof. For the first assertion, assume $\sigma_x^i \tau_y^j(p) = cp$ for $i, j \in \mathbb{Z}$ and $c \in k$, with $i \neq 0$. We consider two cases. First, if $j = 0$ or $\deg_y(p) = 0$, then $\sigma_x^i(p) = cp$, so that $c = 1$ by comparing the leading coefficients. Upon setting $K = k(y)$ and applying Lemma 2 of (Abramov and Petkovšek, 2002b), we conclude that $p \in K$, and then $p \in k[y]$. For the second case, we assume $d = \deg_y(p) > 0$ and $j \neq 0$. Write $p = p_d(x)y^d + \cdots + p_0(x)$, where $p_0, \dots, p_d \in k[x]$ and $p_d \neq 0$. Then

$$\sigma_x^i \tau_y^j(p) = p_d(x+i)q^{jd}y^d + \cdots + p_0(x+i).$$

From the equality $\sigma_x^i \tau_y^j(p) = cp$, we get

$$p_\ell(x+i)q^{j\ell} = cp_\ell(x) \quad \text{for all } \ell \text{ with } 0 \leq \ell \leq d.$$

Assume that ℓ is an integer in $\{0, \dots, d\}$ such that $p_\ell(x) \neq 0$. Then $c = q^{j\ell}$ by the above equation. This implies that $p_\ell(x+i) = p_\ell(x)$. By Lemma 2 in (Abramov and Petkovšek, 2002b), $p_\ell \in k$ for all ℓ with $p_\ell(x) \neq 0$. Thus, $p \in k[y]$ again, and the first assertion holds.

To prove the second assertion, we first show that the case $i \neq 0$ reduces to the case $i = 0$. Indeed, if $\sigma_x^i \tau_y^j(p) = cp$ for $i, j \in \mathbb{Z}$ and $c \in k$, and if $i, j \neq 0$, the first assertion of the lemma implies that $p \in k[y]$. As a consequence, $\sigma_x^0 \tau_y^j(p) = cp$ as well. So we now focus to the special case $i = 0$, that is $\tau_y^j(p) = cp$. Suppose that there are d_1 and d_2 in \mathbb{N} with $d_1 > d_2$ such that

$$p = p_{d_1}y^{d_1} + p_{d_2}y^{d_2} + \text{terms of lower degree in } y,$$

where $p_{d_1}, p_{d_2} \in k[x]^*$. Applying τ_y^j to p yields

$$\tau_y^j(p) = p_{d_1}q^{jd_1}y^{d_1} + p_{d_2}q^{jd_2}y^{d_2} + \text{lower terms in } y.$$

The equality $\tau_y^j(p) = cp$ implies that $c = q^{jd_1}$ and $c = q^{jd_2}$. Hence, $q^{j(d_1-d_2)} = 1$, a contradiction to the assumption that q is not a root of unity. So $p = \lambda y^s$ for some λ in $k[x]$ and s in \mathbb{N} . Since p is irreducible, s is equal to 0 or 1, and if $s = 1$, then $p \in k$, again because it is irreducible. Therefore, the second assertion holds. \square

The following lemma is an analogue for the difference- q -difference case of Theorem 9 in (Abramov, 2003).

Lemma 5.10. Let $f = a/b \in k(x, y)^*$ with $\gcd(a, b) = 1$, let b be non-split, and let L be in $k(x, y)\langle \partial_x \rangle$ whose coefficients are all split. Then

- (i) If b is shift-free with respect to σ_y , and $\partial_x = T_x$, then the non-split part of the denominator of $L(f)$ is not σ_y -spread.
- (ii) If b is q -shift-free with respect to τ_y , and $\partial_x = S_x$, then the non-split part of the denominator of $L(f)$ is not τ_y -spread.

Proof. Let $L = \sum_{i=0}^{\rho} u_i \partial_x^i$, where $u_0, u_1, \dots, u_\rho \in k(x, y)$ are split and u_ρ is nonzero.

First, assume that b is shift-free with respect to σ_y , and that $\partial_x = T_x$. The first assertion clearly holds if $\rho = 0$, because a nontrivial and shift-free polynomial with respect to σ_y is not σ_y -spread. So it suffices to consider the case in which $\rho > 0$.

Applying L to f , we have that

$$L(f) = \sum_{i=0}^{\rho} u_i \tau_x^i \left(\frac{a}{b} \right). \quad (14)$$

Denote by B the non-split part of the denominator of $L(f)$, which belongs to $k[x, y]$. If B is in k , then the first assertion evidently holds. Assume further that B has positive degrees in both x and y . By (14), b must have a nontrivial irreducible factor, which is non-split. Let p be such a factor. Furthermore, we may assume without loss of generality that $\tau_x^\mu(p) \nmid b$ for any positive integer μ . Then $\tau_x^\rho(p) \mid B$, because all the u_i 's are split.

Suppose that B is σ_y -spread. Then there exists $j_0 \in \mathbb{Z}^*$ such that $\sigma_y^{j_0} \tau_x^\rho(p) \mid B$. By (14), B divides $\prod_{i=0}^{\rho} \tau_x^i(b)$. By the above two divisibilities, there exists ℓ_0 in $\{0, \dots, \rho\}$ such that $\sigma_y^{j_0} \tau_x^\rho(p) \mid \tau_x^{\ell_0}(b)$. Note that $\ell_0 \neq \rho$, for otherwise, both $\sigma_y^{j_0}(p)$ and p would

divide b , a contradiction to the assumption that b is shift-free. Therefore, $\ell_0 < \rho$. Since $\sigma_y^{j_0} \tau_x^{\rho - \ell_0}(p) \mid b$, there exists a non-negative integer i_0 such that

$$\sigma_y^{j_0} \tau_x^{\rho - \ell_0 + i_0}(p) \mid b \quad \text{but} \quad \sigma_y^{j_0} \tau_x^{\rho - \ell_0 + i_0 + \mu}(p) \nmid b$$

for any positive integer μ . It follows from $\rho - \ell_0 > 0$ that $\rho - \ell_0 + i_0 > 0$.

Repeating the above process for the irreducible factor $\sigma_y^{j_0} \tau_x^{\rho - \ell_0 + i_0}(p)$, we can find that $j_1 \in \mathbb{Z}^*$, $i_1 \in \mathbb{N}$ and $\ell_1 \in \{0, \dots, \rho - 1\}$ such that $\rho - \ell_1 + i_1 > 0$ and

$$\sigma_y^{j_0 + j_1} \tau_x^{(\rho - \ell_0 + i_0) + (\rho - \ell_0 + i_1)}(p) \mid b \quad \text{but} \quad \sigma_y^{j_0 + j_1} \tau_x^{(\rho - \ell_0 + i_0) + (\rho - \ell_1 + i_1) + \mu}(p) \nmid b$$

for any positive integer μ . Continuing this process yields a sequences of irreducible factors of b . Since b has only finitely many irreducible factors, there exist $m, n \in \mathbb{N}$ with $n < m$ such that

$$\sigma_y^{j_0 + \dots + j_n} \tau_x^{(\rho - \ell_0 + i_0) + \dots + (\rho - \ell_n + i_n)}(p) = c \sigma_y^{j_0 + \dots + j_m} \tau_x^{(\rho - \ell_0 + i_0) + \dots + (\rho - \ell_m + i_m)}(p)$$

for some $c \in k$. This implies that

$$\sigma_y^{-j_{n+1} - \dots - j_m} \tau_x^{-(\rho - \ell_{n+1} + i_{n+1}) - \dots - (\rho - \ell_m + i_m)}(p) = cp.$$

Note that $(\rho - \ell_{n+1} + i_{n+1}) + \dots + (\rho - \ell_m + i_m) \neq 0$. By Lemma 5.9 (ii), p is either in $k[x]$ or is equal to λy for some $\lambda \in k$, a contradiction to the assumption that p is non-split.

The second assertion can be proved similarly according to Lemma 5.9 (i). \square

We are ready to prove that a telescopic term is the sum of an exact term and a proper one.

Theorem 5.11. Let h be a mixed term over $(k(x, y), (\xi_x, \eta_y))$. Let

$$h = \Delta_y(h_1) + h_2$$

be an additive decomposition with respect to ∂_y . If h has a telescoper of type (∂_x, ∂_y) , then h_2 is either zero or proper.

Proof. By the definition of additive decompositions, there exist $r, u, v \in k(x, y)$ such that $h_2 \in r\mathcal{H}(u, v)$, where r, u and v satisfy the properties described in Definition 4.11. Clearly, we may assume that h_2 is nonzero. Assume further that h has a telescoper L of type (∂_x, ∂_y) . Then L is also a telescoper of h_2 by Lemma 5.8. Our goal is to show that h_2 is proper.

First, we let $\eta_y = \delta_y$. By Property (a) in Definition 4.11, $\text{den}(r)$ is squarefree with respect to y , and v is δ_y -reduced. By Lemma 4.6 (i), both u and $\text{den}(v)$ are split. Therefore, h_2 is proper if and only if $\text{den}(r)$ is split by Lemma 2.15 (iv) and (v). So it suffices to show that $\text{den}(r)$ is split.

Case 1.1. Let $\xi_x = \sigma_x$. Then we may write

$$L = \sum_{i=0}^{\rho} e_i S_x^i \in k(x) \langle S_x \rangle, \quad \text{where } e_0, \dots, e_\rho \in k(x) \text{ and } e_\rho \neq 0.$$

As in Example 2.9, we have

$$L(h_2) = \underbrace{\left(\sum_{i=0}^{\rho} u_i \sigma_x^i(r) \right)}_f g,$$

where $g \in \mathcal{H}(u, v) \setminus \{0\}$, $u_i = e_i \prod_{j=0}^{i-1} \sigma_x^j(u)$, and u_ρ is nonzero.

Suppose that $\text{den}(r)$ is non-split. Then there exists a non-split irreducible polynomial $p \in k[x, y]$ such that $p \mid \text{den}(r)$ and $\sigma_x^i(p) \nmid \text{den}(r)$ for all $i > 0$. Moreover, p has multiplicity one in $\text{den}(r)$, because $\text{den}(r)$ is squarefree. Since all the u_i 's are split, $\sigma_x^\rho(p)$ is an irreducible factor of $\text{den}(f)$ with multiplicity one. In particular, f is nonzero. It follows that the non-split part of $\text{den}(f)$ is not δ_y -spread. Since L is a telescoper of h_2 whose type is (S_x, D_y) , fg is an exact term with respect to D_y . Note that $g \in \mathcal{H}(u, v)$ and $\text{den}(v)$ is split. We can apply Proposition 4.4 (i) to $L(h_2)$, which implies that the non-split part of $\text{den}(f)$ is δ_y -spread, a contradiction. Thus, $\text{den}(r)$ is split.

Case 1.2. Let $\xi_x = \tau_x$. The same argument used in Case 1.1 shows that h_2 is proper.

Second, we let $\eta_y = \sigma_y$. By Property (b) in Definition 4.11, $\text{den}(r)$ is shift-free with respect to σ_y , and v is σ_y -reduced. Both $\text{den}(u)$ and v are split if $\xi_x = \delta_x$ by Lemma 4.6 (2a). Both u and v are split if $\xi_x = \tau_x$ by Lemma 4.6 (2b). Therefore, h_2 is proper if and only if $\text{den}(r)$ is split by Lemma 2.15 (i) and (vi). So it suffices to show that $\text{den}(r)$ is split.

Case 2.1. Let $\xi_x = \delta_x$. Then we may write

$$L = \sum_{i=0}^{\rho} e_i D_x^i \in k(x)\langle D_x \rangle, \quad \text{where } e_0, \dots, e_\rho \in k(x) \text{ and } e_\rho \neq 0.$$

As in Example 2.9, we have

$$L(h_2) = \underbrace{\left(e_\rho \delta_x^\rho(r) + \sum_{i=0}^{\rho-1} u_i \delta_x^i(r) \right)}_f g,$$

where $g \in \mathcal{H}(u, v) \setminus \{0\}$, $u_i \in k(x, y)$, and $\text{den}(u_i)$ is split for all i with $0 \leq i \leq \rho - 1$.

Suppose that $\text{den}(r)$ is non-split. Then there exists a non-split irreducible polynomial $p \in k[x, y]$ such that $p \mid \text{den}(r)$. Assume that m is the multiplicity of p in $\text{den}(r)$. Then the multiplicity of p in $\text{den}(\delta^i(r))$ is equal to $m + i$. In particular, its multiplicity in $\text{den}(e_\rho \delta^\rho(r))$ is equal to $m + \rho$, which is also the multiplicity of p in $\text{den}(f)$, because all the $\text{den}(u_i)$ are split. It follows that p is an irreducible factor of $\text{den}(f)$. In particular, f is nonzero. Since L is a telescoper of h_2 whose type is (D_x, S_y) , fg is an exact term with respect to S_y . Note that $g \in \mathcal{H}(u, v)$ and that v is split. We can apply Proposition 4.4 (ii) to fg , which concludes that the non-split part of $\text{den}(f)$ is σ_y -spread. Thus, there exists a nonzero integer j such that $\sigma_y^j(p)$ is an irreducible factor of $\text{den}(f)$. From the definitions of f and splitness of the $\text{den}(u_i)$, it follows that $\sigma_y^j(p)$ is also an irreducible factor of $\text{den}(r)$, a contradiction to the fact that $\text{den}(r)$ is shift-free. Hence, $\text{den}(r)$ is split.

Case 2.2. Let $\xi_x = \tau_x$. Then we may write $L = \sum_{i=0}^{\rho} e_i T_x^i \in k(x)\langle T_x \rangle$

$$L = \sum_{i=0}^{\rho} e_i T_x^i \in k(x)\langle T_x \rangle, \quad \text{where } e_0, \dots, e_\rho \in k(x) \text{ and } e_\rho \neq 0.$$

As in Example 2.9, we have

$$L(h_2) = \underbrace{\left(\sum_{i=0}^{\rho} u_i T_x^i(r) \right)}_f g,$$

Algorithm IsTelescopable**Input:** a mixed term $h \in \mathcal{H}(a, b)$ over $(k(x, y), (\xi_x, \eta_y))$.**Output:** true, if h has a telescoper of type (ξ_x, η_y) ; false, otherwise.

- (1) Compute an additive decomposition of
- h
- with respect to
- η_y
- and get

$$h = \Delta_y(h_1) + rg,$$

where $g \in \mathcal{H}(u, v)$ is as given by (10–11) in Definition 4.11.

- (2) Compute the primitive part
- p
- of
- $\text{den}(r)$
- with respect to
- y
- .
-
- (3) If
- p
- is in
- $k[y]$
- , then return true, otherwise, return false.

Fig. 6. Algorithms for deciding the existence of telescopers.

where $g \in \mathcal{H}(u, v) \setminus \{0\}$ and $u_i \in k(x, y)$ is split for all i with $0 \leq i \leq \rho$.

Suppose that $\text{den}(r)$ is non-split. By Lemma 5.10 (i), $\text{den}(f)$ is not σ_y -spread. Since L is a telescoper of h_2 whose type is (T_x, S_y) , fg is an exact term with respect to S_y . Note that $g \in \mathcal{H}(u, v)$ and v is split. Applying Proposition 4.4 (ii) to fg , we see that $\text{den}(f)$ is σ_y -spread, a contradiction. Hence, $\text{den}(r)$ is split.

Third, we let $\eta_y = \tau_y$. By Property (c) of Definition 4.11, $\text{den}(r)$ is q -shift-free with respect to y , and v is τ_y -reduced. Both $\text{den}(u)$ and v are split if $\xi_x = \delta_x$ by Lemma 4.6 (3a). Both u and v are split if $\xi_x = \tau_x$ by Lemma 4.6 (3b). Therefore, h_2 is proper if and only if $\text{den}(r)$ is split by Lemma 2.15 (ii) and (iii). So it suffices to show that $\text{den}(r)$ is split.

Case 3.1. Let $\xi_x = \delta_x$. The proof is similar to that in Case 2.1, in which one applies the third assertion of Proposition 4.4 instead of the second one.

Case 3.2. Let $\xi_x = \sigma_x$. The proof is similar that in Case 2.2, in which one applies the second assertion of Lemma 5.10 instead of the first one, and the third assertion of Proposition 4.4 instead of the second one. \square

Combining Theorems 5.7 and 5.11, we obtain a criterion for the existence of telescopers of a mixed term, which is the main result of this article.

Theorem 5.12. Let h be a mixed term over $(k(x, y), (\xi_x, \eta_y))$. Assume that

$$h = \Delta_y(h_1) + h_2$$

is an additive decomposition of h . Then h has a telescoper of type (∂_x, ∂_y) if and only if h_2 is either zero or a proper mixed term.

6. Algorithms and examples

For a given mixed term, we can decide the existence of telescopers by Theorem 5.12. First, we use the algorithms in (Abramov and Petkovšek, 2002a), (Geddes et al., 2004), and (Chen et al., 2005) to perform the respective additive decompositions. Second, we test whether the denominator of the rational part in the non-exact component is split or not by Remark 2.13. The decision procedure is given in Figure 6.

Example 6.1. It is possible that a mixed term has a telescoper of type (S_x, D_y) but no telescoper of type (D_x, S_y) or (D_x, T_y) . Consider the rational function

$$h = \frac{1}{(x+y)^2}.$$

Applying Hermite reduction to h with respect to δ_y yields

$$h = D_y \left(\frac{-1}{x+y} \right),$$

which implies that 1 is a telescoper of type (S_x, D_y) for h . Note that $h = \Delta_y(0) + h$ is an additive decomposition when $\eta_y = \sigma_y$ or $\eta_y = \tau_y$, because $\text{den}(h) = (x+y)^2$ is both shift-free and q -shift-free with respect to y . But h is not proper, because $x+y$ is not split. Hence, h has no telescoper of type (D_x, S_y) and (D_x, T_y) . Similarly, consider the rational function

$$h = \frac{1}{(x+y)(x+y+1)}.$$

Since $h = (S_y - 1)(-1/(x+y))$, 1 is a telescoper of type (D_x, S_y) for h . However, h has no telescoper of type (S_x, D_y) or (T_x, D_y) , because $(x+y)(x+y+1)$ is squarefree with respect to δ_y and it is not split.

As we mentioned before, properness is only a sufficient condition for the existence of telescopers. The following two examples illustrate this fact.

Example 6.2. Consider the mixed term over $(k(x, y), (\sigma_x, \delta_y))$

$$h = \frac{-y + 2xy + 2x^2}{(x+y)^2 x} \cdot y^x \cdot e^{-y}. \quad (15)$$

In the structural decomposition (15) given by $\alpha = -1$, $\beta = y$, $\gamma = 1$, and $f = h/(y^x e^{-y})$, the denominator $(x+y)^2$ is not split. By case (iv) in Lemma 2.15, it follows that h is not proper. But h has a telescoper of type (S_x, D_y) since it can be decomposed into

$$h = D_y \left(\frac{1}{x+y} \cdot y^x \cdot e^{-y} \right) + y^{x-1} \cdot e^{-y},$$

where $y^{x-1} \cdot e^{-y}$ is proper, because the rational function in the corresponding structural decomposition is 1, and therefore its denominator is split.

Example 6.3. Consider the mixed term over $(k(x, y), (\tau_x, \delta_y))$

$$h = \frac{y^2 + xy - x}{(x+y)^2 x} \cdot e^{-y},$$

which is not proper, because, in the structural decomposition given by $\alpha = -1$, $\beta = 1$, $f = h/e^{-y}$, the denominator $(x+y)^2$ is not split. But it has a telescoper of type (T_x, D_y) since h can be decomposed into

$$h = D_y \left(\frac{1}{x+y} \cdot e^{-y} \right) + \frac{e^{-y}}{x},$$

where e^{-y}/x is proper, because the rational function in the corresponding structural decomposition is $1/x$, and therefore its denominator is split.

The last example presents another application of Theorem 5.12.

Example 6.4. Let

$$f = \frac{1}{y^2 - x}.$$

Note that the denominator of f is non-split and shift-free with respect to σ_y . By Theorem 5.12, there is no linear differential operator $L(x, D_x) \in k(x)\langle D_x \rangle$ and $g \in k(x, y)$ such that $L(x, D_x)(f) = \Delta_y(g)$, which, together with Proposition 3.1 in (Hardouin and Singer, 2008) and the descent argument similar to that given in the proof of Corollary 3.2 in (Hardouin and Singer, 2008) (or Section 1.2.1 of (Di Vizio and Hardouin, 2012)), implies that the sum

$$F(x, y) := \sum_{i=1}^{y-1} \frac{1}{i^2 - x} \quad (\text{satisfying } S_y(F) - F = f)$$

satisfies no polynomial differential equation $P(x, y, F, D_x(F), D_x^2(F), \dots) = 0$.

References

- Abramov, S. A., 1975. The rational component of the solution of a first order linear recurrence relation with rational right hand side. *Ž. Vyčisl. Mat. i Mat. Fiz.* 15 (4), 1035–1039, 1090.
- Abramov, S. A., 1995. Indefinite sums of rational functions. In: Proceedings of the 1995 International Symposium on Symbolic and Algebraic Computation. ISSAC '95. ACM, New York, NY, USA, pp. 303–308.
URL <http://doi.acm.org/10.1145/220346.220386>
- Abramov, S. A., 2002. Applicability of Zeilberger's algorithm to hypergeometric terms. In: ISSAC'02: Proceedings of the 2002 International Symposium on Symbolic and Algebraic Computation. ACM, New York, pp. 1–7.
- Abramov, S. A., 2003. When does Zeilberger's algorithm succeed? *Adv. in Appl. Math.* 30 (3), 424–441.
URL [http://dx.doi.org/10.1016/S0196-8858\(02\)00504-3](http://dx.doi.org/10.1016/S0196-8858(02)00504-3)
- Abramov, S. A., Le, H. Q., 2002. A criterion for the applicability of Zeilberger's algorithm to rational functions. *Discrete Math.* 259 (1-3), 1–17.
URL [http://dx.doi.org/10.1016/S0012-365X\(02\)00442-9](http://dx.doi.org/10.1016/S0012-365X(02)00442-9)
- Abramov, S. A., Petkovšek, M., 2001. Minimal decomposition of indefinite hypergeometric sums. In: ISSAC'01: Proceedings of the 2001 International Symposium on Symbolic and Algebraic Computation. ACM, New York, pp. 7–14 (electronic).
- Abramov, S. A., Petkovšek, M., 2002. Rational normal forms and minimal decompositions of hypergeometric terms. *J. Symbolic Comput.* 33 (5), 521–543.
URL <http://dx.doi.org/10.1006/jsco.2002.0522>
- Abramov, S. A., Petkovšek, M., 2002. On the structure of multivariate hypergeometric terms. *Adv. Appl. Math.* 29 (3), 386–411.
- Almkvist, G., Zeilberger, D., 1990. The method of differentiating under the integral sign. *J. Symbolic Comput.* 10, 571–591.
- Bernštejn, J., 1971. Modules over a ring of differential operators. An investigation of the fundamental solutions of equations with constant coefficients. *Funkcional. Anal. i Priložen.* 5 (2), 1–16.

- Bronstein, M., Li, Z., Wu, M., 2005. Picard–Vessiot extensions for linear functional systems. In: ISSAC '05: Proceedings of the 2005 International Symposium on Symbolic and Algebraic Computation. ACM, New York, USA, pp. 68–75.
- Chen, S., Feng, R., Fu, G., Li, Z., 2011. On the structure of compatible rational functions. In: Leykin, A. (Ed.), ISSAC '11: Proceedings of the 2011 International Symposium on Symbolic and Algebraic Computation. ACM, pp. 91–98.
URL <http://algo.inria.fr/chen/>
- Chen, S., Kauers, M., Singer, M. F., 2012. Telescopers for rational and algebraic functions via residues. In: van der Hoeven, J., van Hoeij, M. (Eds.), ISSAC'92. ACM Press, pp. 130–137, Proceedings of the 37th International Symposium on Symbolic and Algebraic Computation (Grenoble, France, July 22–27, 2012).
- Chen, S., Singer, M. F., 2012. Residues and telescopers for bivariate rational functions. *Adv. in Appl. Math.* 49 (2), 111–133.
URL <http://dx.doi.org/10.1016/j.aam.2012.04.003>
- Chen, W. Y. C., Hou, Q.-H., Mu, Y.-P., 2005. Applicability of the q -analogue of Zeilberger's algorithm. *J. Symbolic Comput.* 39 (2), 155–170.
URL <http://dx.doi.org/10.1016/j.jsc.2004.09.002>
- Chyzak, F., Kauers, M., Salvy, B., 2009. A non-holonomic systems approach to special function identities. In: ISSAC '09: Proceedings of the 2009 International Symposium on Symbolic and Algebraic Computation. ACM, New York, NY, USA, pp. 111–118.
- Chyzak, F., Salvy, B., 1998. Non-commutative elimination in Ore algebras proves multivariate identities. *J. Symbolic Comput.* 26 (2), 187–227.
URL <http://dx.doi.org/10.1006/jsc.1998.0207>
- Davenport, J. H., 1986. The Risch differential equation problem. *SIAM J. Comput.* 15 (4), 903–918.
URL <http://dx.doi.org/10.1137/0215063>
- Di Vizio, L., Hardouin, C., 2012. Descent for differential galois theory of difference equations: confluence and q -dependence. *Pac. J. Math.* 256 (1), 79–104.
- Fasenmyer, M. C., 1947. Some generalized hypergeometric polynomials. *Bull. Amer. Math. Soc.* 53, 806–812.
- Geddes, K. O., Le, H. Q., Li, Z., 2004. Differential rational normal forms and a reduction algorithm for hyperexponential functions. In: ISSAC'04: Proceedings of the 2004 International Symposium on Symbolic and Algebraic Computation. ACM, New York, USA, pp. 183–190.
- Gosper, Jr., R. W., 1978. Decision procedure for indefinite hypergeometric summation. *Proc. Nat. Acad. Sci. U.S.A.* 75 (1), 40–42.
- Hardouin, C., Singer, M. F., 2008. Differential Galois theory of linear difference equations. *Math. Ann.* 342 (2), 333–377.
URL <http://dx.doi.org/10.1007/s00208-008-0238-z>
- Hermite, C., 1872. Sur l'intégration des fractions rationnelles. *Ann. Sci. École Norm. Sup.* (2) 1, 215–218.
- Kashiwara, M., 1978. On the holonomic systems of linear differential equations. II. *Invent. Math.* 49 (2), 121–135.
URL <http://dx.doi.org/10.1007/BF01403082>
- Koepf, W., 1998. Hypergeometric Summation. *Advanced Lectures in Mathematics*. Friedr. Vieweg & Sohn, Braunschweig, an algorithmic approach to summation and special function identities.

- Le, H. Q., 2001. On the q -analogue of Zeilberger's algorithm to rational functions. *Program. Comput. Softw.* 27 (1), 35–42.
- Li, Z., Singer, M. F., Wu, M., Zheng, D., 2006. A recursive method for determining the one-dimensional submodules of Laurent-Ore modules. In: *ISSAC'06: Proceedings of the 2006 International Symposium on Symbolic and Algebraic Computation*. ACM, New York, pp. 220–227.
URL <http://dx.doi.org/10.1145/1145768.1145806>
- Lipshitz, L. M., 1988. The diagonal of a D-finite power series is D-finite. *J. Algebra* 113 (2), 373–378.
- Ostrogradskii, M. V., 1845. De l'intégration des fractions rationnelles. *Bull. de la classe physico-mathématique de l'Acad. Impériale des Sciences de Saint-Petersbourg* 4, 145–167, 286–300.
- Schneider, C., 2010. Parameterized telescoping proves algebraic independence of sums. *Ann. Comb.* 14 (4), 533–552.
URL <http://dx.doi.org/10.1007/s00026-011-0076-7>
- Takayama, N., 1992. An approach to the zero recognition problem by Buchberger algorithm. *J. Symbolic Comput.* 14 (2-3), 265–282.
- Verbaeten, P., 1974. The automatic construction of pure recurrence relations. *SIGSAM Bull.* 8 (3), 96–98.
- Wegschaider, K., May 1997. Computer Generated Proofs of Binomial Multi-Sum Identities. Master's thesis, RISC, J. Kepler University.
- Wilf, H. S., Zeilberger, D., 1992a. An algorithmic proof theory for hypergeometric (ordinary and “ q ”) multisum/integral identities. *Invent. Math.* 108 (3), 575–633.
URL <http://dx.doi.org/10.1007/BF02100618>
- Wilf, H. S., Zeilberger, D., 1992b. Rational function certification of multisum/integral/“ q ” identities. *Bull. Amer. Math. Soc. (N.S.)* 27 (1), 148–153.
URL <http://dx.doi.org/10.1090/S0273-0979-1992-00297-5>
- Zeilberger, D., 1990a. A fast algorithm for proving terminating hypergeometric identities. *Discrete Math.* 80 (2), 207–211.
URL [http://dx.doi.org/10.1016/0012-365X\(90\)90120-7](http://dx.doi.org/10.1016/0012-365X(90)90120-7)
- Zeilberger, D., 1990b. A holonomic systems approach to special functions identities. *J. Comput. Appl. Math.* 32, 321–368.
- Zeilberger, D., 1991. The method of creative telescoping. *J. Symbolic Comput.* 11 (3), 195–204.