

Ordered Probability Mass Function

Zhengjun Cao¹, Lihua Liu²

¹Department of Mathematics, Shanghai University, China. caozhj@shu.edu.cn

²Department of Mathematics, Shanghai Maritime University, Shanghai, China.

Abstract Suppose that in the four tests Alice's scores are 90, 95, 85, 90, and Bob's scores are 85, 95, 90, 90. How to evaluate their scores? In this paper, we introduce the concept of ordered probability mass function which can be used to find a probability mass function with smaller variance. More interestingly, we can use it to distinguish sequences of positive numbers statistically.

Keywords: probability mass function, expectation, variance.

1 Introduction

A random variable that can take on at most a countable number of possible values is said to be discrete. Suppose that X is a discrete random variable whose possible values are x_1, x_2, \dots, x_n , $x_i \geq 0, i = 1, \dots, n$. We call the probability mass function

$$p(x_i) = P(X = x_i) = 1/n, \quad i = 1, \dots, n$$

as average probability mass function (APM for short), the probability mass function

$$p(x_i) = P(X = x_i) = \frac{x_i}{\sum_{i=1}^n x_i}, \quad i = 1, \dots, n$$

as general probability mass function (GPM). In a typical statistical problem, we have a random variable X of interest, but its probability mass function is not known. The above two probability mass functions are widely used in daily life.

The quantity $E[X]$ defined by $\sum_{x:p(x)>0} xp(x)$ is called the expected value of X , which is also called the *expectation* of X . The variance $[1, 2]$, which is equal to the expected square of the difference between X and its expected value, is a measure of the spread of the possible values of X . Practically speaking, a probability mass function with all positive probabilities and smaller variance is more appreciated.

How to find a probability mass function with smaller variance? We have observed that there were no methods which can be used to find it. We have also observed that there were no methods which can distinguish sequences of positive numbers statistically, because the common sample mean and sample variance do not depend on the order relation of elements in a sample space. In this paper, we introduce the concept of ordered probability mass function which can be used to find a probability mass function with smaller variance. More interestingly, it can be used to distinguish sequences of positive numbers statistically. To the best of our knowledge, it is the first time to introduce a method to cope with this problem.

2 Ordered probability mass function

In order to introduce the new concept of ordered probability mass function, we prove the following lemma.

Lemma 1. *If $abc = 1$, then*

$$\frac{a}{1+a+ab} + \frac{b}{1+b+bc} + \frac{c}{1+c+ca} = 1 \quad (1)$$

Proof. Since $abc = 1$, we have

$$\begin{aligned} \frac{b}{1+b+bc} &= \frac{ab}{a+ab+abc} = \frac{ab}{1+a+ab}, \\ \frac{c}{1+c+ca} &= \frac{abc}{ab+abc+abca} = \frac{1}{1+a+ab}. \end{aligned}$$

Thus Eq.(1) holds. □

Likewise, if $abcd = 1$, then

$$\frac{a}{1+a+ab+abc} + \frac{b}{1+b+bc+bcd} + \frac{c}{1+c+cd+cda} + \frac{d}{1+d+da+dab} = 1 \quad (2)$$

More generally, if $x_1x_2 \cdots x_n = 1$, then

$$\sum_{i=1}^n \frac{x_i}{1+x_i+x_ix_{i+1}+\cdots+x_i \cdots x_n+x_i \cdots x_nx_1+\cdots+x_i \cdots x_nx_1 \cdots x_{i-2}} = 1 \quad (3)$$

Suppose that $\frac{x_i}{1+x_i+x_ix_{i+1}+\cdots+x_i \cdots x_n+x_i \cdots x_nx_1+\cdots+x_i \cdots x_nx_1 \cdots x_{i-2}} > 0, i = 1, \cdots, n$, then Eq.(3) provides us a new probability mass function.

Definition 1. Suppose that X is a discrete random variable whose possible values are $x_1, x_2, \cdots, x_n, x_i > 0, i = 1, \cdots, n$. We call the probability mass function

$$\begin{aligned} p(x_i) &= P(X = x_i) \\ &= \frac{x_i}{1+x_i+x_ix_{i+1}+\cdots+x_i \cdots x_n+x_i \cdots x_nx_1+\cdots+x_i \cdots x_nx_1 \cdots x_{i-2}}, \quad i = 1, \cdots, n \quad (4) \end{aligned}$$

as ordered probability mass function (OPM) provided that $x_1x_2 \cdots x_n = 1$.

By the definition, the OPM for the sequence $\frac{5}{6}, 1, 1, \frac{6}{5}$ is computed as follows:

$$\begin{aligned}
 p\left(\frac{5}{6}\right) &= \frac{\frac{5}{6}}{1 + \frac{5}{6} + \frac{5}{6} \times 1 + \frac{5}{6} \times 1 \times 1} = \frac{5}{21}, \\
 p(1) &= \frac{1}{1 + 1 + 1 \times 1 + 1 \times 1 \times \frac{6}{5}} = \frac{5}{21}, \quad (\text{for the first element "1" in the sequence}) \\
 p(1) &= \frac{1}{1 + 1 + 1 \times \frac{6}{5} + 1 \times \frac{6}{5} \times \frac{5}{6}} = \frac{5}{21}, \quad (\text{for the second element "1" in the sequence}) \\
 p\left(\frac{6}{5}\right) &= \frac{\frac{6}{5}}{1 + \frac{6}{5} + \frac{6}{5} \times \frac{5}{6} + \frac{6}{5} \times \frac{5}{6} \times 1} = \frac{6}{21}.
 \end{aligned}$$

It is easy to find that the APM and GPM do not depend on the order relation of elements in a sample space, but the OPM depends on it. Here is an illustration example.

Example 1. Find the APM, GPM, OPM, expectations and variances of the following sequences.

sequence	probability mass	Expectation	Variance
$\frac{6}{5}, 1, 1, \frac{5}{6}$	APM: $\frac{1}{4}, \frac{1}{4}, \frac{1}{4}, \frac{1}{4}$	1.00833	0.0168750
$\frac{6}{5}, 1, 1, \frac{5}{6}$	GPM: $\frac{36}{121}, \frac{30}{121}, \frac{30}{121}, \frac{25}{121}$	1.02507	0.0170116
$\frac{6}{5}, \frac{5}{6}, 1, 1$	OPM: $\frac{6}{21}, \frac{5}{21}, \frac{5}{21}, \frac{5}{21}$	1.01746	0.0177375
$\frac{6}{5}, 1, \frac{5}{6}, 1$	OPM: $\frac{6}{22}, \frac{6}{22}, \frac{5}{22}, \frac{5}{22}$	1.01667	0.0169444
$\frac{6}{5}, 1, 1, \frac{5}{6}$	OPM: $\frac{6}{23}, \frac{6}{23}, \frac{6}{23}, \frac{5}{23}$	1.01594	<u>0.0162193</u>

Note that the sequence $\frac{6}{5}, 1, 1, \frac{5}{6}$, with ordered probability mass $\frac{6}{23}, \frac{6}{23}, \frac{6}{23}, \frac{5}{23}$, has the smallest variance. From the practical point of view, the probability distribution

$$\left\{ \left(\frac{6}{5}; \frac{6}{23} \right), \left(1; \frac{6}{23} \right), \left(1; \frac{6}{23} \right), \left(\frac{5}{6}; \frac{5}{23} \right) \right\}$$

is more appreciated.

Example 2. Compute the APM, GPM, OPM, expectations and variances of the following sequences.

sequence	probability mass	Expectation	Variance
$\frac{6}{5}, \frac{7}{6}, \frac{6}{7}, \frac{5}{6}$	APM: $\frac{1}{4}, \frac{1}{4}, \frac{1}{4}, \frac{1}{4}$	1.01429	0.0287868
$\frac{6}{5}, \frac{7}{6}, \frac{6}{7}, \frac{5}{6}$	GPM: $\frac{252}{852}, \frac{245}{852}, \frac{180}{852}, \frac{175}{852}$	1.04267	<u>0.0280154</u>
$\frac{6}{5}, \frac{7}{6}, \frac{6}{7}, \frac{5}{6}$	OPM: $\frac{6}{24}, \frac{7}{24}, \frac{6}{24}, \frac{5}{24}$	1.02817	0.0281971
$\frac{6}{5}, \frac{7}{6}, \frac{5}{6}, \frac{6}{7}$	OPM: $\frac{36}{143}, \frac{42}{143}, \frac{35}{143}, \frac{30}{143}$	1.02854	0.0284942
$\frac{6}{5}, \frac{6}{7}, \frac{7}{6}, \frac{5}{6}$	OPM: $\frac{42}{155}, \frac{36}{155}, \frac{42}{155}, \frac{35}{155}$	1.02854	0.0285634
$\frac{6}{5}, \frac{6}{7}, \frac{5}{6}, \frac{7}{6}$	OPM: $\frac{42}{143}, \frac{36}{143}, \frac{30}{143}, \frac{35}{143}$	1.02860	0.0286940
$\frac{6}{5}, \frac{5}{6}, \frac{6}{7}, \frac{7}{6}$	OPM: $\frac{42}{142}, \frac{35}{142}, \frac{30}{142}, \frac{35}{142}$	1.02897	0.0289964
$\frac{6}{5}, \frac{5}{6}, \frac{7}{6}, \frac{6}{7}$	OPM: $\frac{36}{131}, \frac{30}{131}, \frac{35}{131}, \frac{30}{131}$	1.02861	0.0286305

Note that $\frac{6}{5}, \frac{7}{6}, \frac{6}{7}, \frac{5}{6}$ with the general probability mass $\frac{252}{852}, \frac{245}{852}, \frac{180}{852}, \frac{175}{852}$, has the smallest variance.

3 Application

As we mentioned before, a probability mass function with smaller variance is more appreciated in practice. Equipped with the tool of ordered probability function, we now can handle the following problem: In the four tests, Alice's scores are 90, 95, 85, 90, and Bob's scores are 85, 95, 90, 90. How to evaluate their scores?

To meet the requirement that $x_1 x_2 \cdots x_n = 1$ in the definition of OPM, we have first to standardize their scores. Set $v = \sqrt[4]{90 \times 95 \times 85 \times 90}$, $a_1 = \frac{90}{v}, a_2 = \frac{95}{v}, a_3 = \frac{85}{v}, a_4 = \frac{90}{v}$, $b_1 = \frac{85}{v}, b_2 = \frac{95}{v}, b_3 = \frac{90}{v}, b_4 = \frac{90}{v}$. Hence, $a_1 a_2 a_3 a_4 = 1, b_1 b_2 b_3 b_4 = 1$. We then compute the ordered probability mass functions of sequences a_1, a_2, a_3, a_4 , and b_1, b_2, b_3, b_4 .

	sequence	OPM	Expectation	Variance
Alice	a_1, a_2, a_3, a_4	0.246667, 0.260572, 0.246286, 0.246476	1.00157	0.00156616
Bob	b_1, b_2, b_3, b_4	0.239718, 0.253231, 0.253427, 0.253623	1.00152	0.00152324

Since Bob's ordered probability mass function has smaller variance 0.00152324, we claim that Bob wins the four tests.

4 Conclusion

The order relation of elements in a sample space is ignored because the common sample mean and variance do not depend on the relation. Thus the common statistical methods can not

evaluate the sequence 90, 95, 85, 90, and the sequence 85, 95, 90, 90. The proposed concept of ordered probability function can be used to distinguish sequences of positive numbers. We believe that the novel method developed in this paper will have more practical applications in the future.

References

- [1] Hogg R. V., Mckean J. W., Craig A. T.: Introduction to mathematical statistics, seventh edition, Pearson Education (2012)
- [2] Ross, S. M.: A first course in probability, eighth edition, Pearson Education (2010)