

Average case polyhedral complexity of the maximum stable set problem

Gábor Braun¹, Samuel Fiorini², and Sebastian Pokutta³

¹ISyE, Georgia Institute of Technology, Atlanta, GA, USA. *Email:* gabor.braun@isye.gatech.edu

²Department of Mathematics, Université libre de Bruxelles CP 216, Bd. du Triomphe, 1050 Brussels, Belgium. *Email:* sfiorini@ulb.ac.be

³ISyE, Georgia Institute of Technology, Atlanta, GA, USA. *Email:* sebastian.pokutta@isye.gatech.edu

October 31, 2014

Abstract

We study the minimum number of constraints needed to formulate random instances of the maximum stable set problem via linear programs (LPs), in two distinct models. In the uniform model, the constraints of the LP are not allowed to depend on the input graph, which should be encoded solely in the objective function. There we prove a $2^{\Omega(n/\log n)}$ lower bound with probability at least $1 - 2^{-2^n}$ for every LP that is exact for a randomly selected set of instances; each graph on at most n vertices being selected independently with probability $p \geq 2^{-(n/4)+n}$. In the non-uniform model, the constraints of the LP may depend on the input graph, but we allow weights on the vertices. The input graph is sampled according to the $G(n, p)$ model. There we obtain upper and lower bounds holding with high probability for various ranges of p . We obtain a super-polynomial lower bound all the way from $p = \Omega\left(\frac{\log^{6+\epsilon} n}{n}\right)$ to $p = o\left(\frac{1}{\log n}\right)$. Our upper bound is close to this as there is only an essentially quadratic gap in the exponent, which currently also exists in the worst-case model. Finally, we state a conjecture that would close this gap, both in the average-case and worst-case models.

1 Introduction

In the last four years, extended formulations considerably gained interest in various areas, including discrete mathematics, combinatorial optimization, and theoretical computer science. The key idea underlying extended formulations is that with the right choice of variables, various combinatorial optimization problems can be *efficiently* expressed via linear programs (LPs). This asks for studying the intrinsic difficulty of expressing optimization problems through a single LP, in terms of the minimum number of necessary *constraints*. In turn, this leads to a complexity measure that we call loosely here ‘polyhedral complexity’ (precise definitions are given later in Section 2).

On the one hand, there is an ever expanding collection of examples of small size extended formulations. For instance, Williams [2002] has expressed the minimum spanning tree problem on a planar graph with only a linear number of (variables and) constraints, while in the natural edge variables the LP has an exponential number of constraints. There exist numerous other examples, see e.g., the surveys by Conforti et al. [2010] and Kaibel [2011].

On the other hand, a recent series of breakthroughs in lower bounds renewed interest for extended formulations [Rothvoß, 2011, Fiorini et al., 2012, Braun et al., 2012, 2014, Braverman and Moitra,

2013, Braun and Pokutta, 2013, Chan et al., 2013, Rothvoß, 2014]. These breakthroughs make it now conceivable to quantify the polyhedral complexity of *any* given combinatorial optimization problem *unconditionally*, that is, independently of conjectures such as P vs. NP, and without extra assumption on the structure of the LP.

Although a polynomial upper bound on the polyhedral complexity yields a polynomial upper bound on the true algorithmic complexity of the problem e.g., through interior point methods—provided that the LP can be efficiently constructed and also that the size of the coefficients is kept under control (see Rothvoß [2011] for a discussion of this last issue)—it is becoming increasingly clear that the converse does not hold. Recently, Chan et al. [2013] proved that every LP for MAXCUT with approximation factor at most $2 - \varepsilon$ needs at least $n^{\Omega\left(\frac{\log n}{\log \log n}\right)}$ constraints, while the approximation factor of the celebrated SDP-based polynomial time algorithm of Goemans and Williamson [1995] is close to 1.13. Even more recently, Rothvoß [2014] solved another major open problem in the area by showing a $2^{\Omega(n)}$ lower bound on the size of any LP expressing the perfect matching problem. Braun and Pokutta [2015] show that the matching polytope does not admit any fully-polynomial size relaxation scheme (the polyhedral equivalent of an FPTAS).

In this paper, we consider the problem of determining the *average case* polyhedral complexity of the maximum stable set problem, in two different models: ‘uniform’ and ‘non-uniform’, see Section 1.2 below. Roughly, the uniform model asks for a single LP that works for a given set of input graphs. In the non-uniform model the LP can depend on the input graph G but should work for every choice of weights on the vertices of G (in particular, for all induced subgraphs of G).

We show that the polyhedral complexity of the maximum stable set problem remains high in each of these models, when the input graph is sampled according to natural distributions. Therefore, we conclude that *the (polyhedral) hardness of the maximum stable set problem is not concentrated on a small mass of graphs but is spread out through all graphs.*

1.1 Related work

Our work is most directly related to Fiorini et al. [2012] and Braun et al. [2012, 2014], where the framework for bounding the size of approximate linear programming formulations was laid out. We also borrow a few ideas from Chan et al. [2013] to set up our uniform model. We will also employ a robustness theorem from Braun and Pokutta [2013] for dropping constraints and feasible solutions.

1.2 Contribution

We present the first strong and unconditional results on the average case size of LP formulations for the maximum stable set problem. In particular, we establish that the maximum stable set problem in two natural average case models does not admit a polynomial size linear programming formulation, even in the unlikely case that $P = NP$.

Uniform model In the *uniform model* the polytope P containing the feasible solutions is *independent* of the instances. The instances will be solely encoded into the objective functions. This ensures that no complexity of the problem is leaked into an instance-specific formulation. A good example of a uniform model is the TSP polytope over K_n with which we can test for Hamiltonian cycles in any graph with at most n vertices by choosing an appropriate objective function.

In the case of the maximum stable set problem, we shall use the natural encoding via characteristic vectors of stable sets. We consider a random collection of input graphs G , where each graph G with $V(G) \subseteq [n]$ is included in the collection with probability $p \geq 2^{-\binom{n/4}{2}+n}$ and show that with probability at least $1 - 2^{-2^n}$, every LP for the maximum stable set problem on such collection of input graphs has at least $2^{\Omega(n/\log n)}$ inequality constraints.

Non-uniform model In the *non-uniform model* we consider the stable set polytope for a *specific but random graph*. The polyhedral description may depend heavily on the chosen graph. We sample a graph G in the Erdős–Rényi $G(n, p)$ model, i.e., G has n vertices, and every pair of vertices is independently connected by an edge with probability p . We then analyze the stable set polytope $\text{STAB}(G)$ of G . If p is small enough, so that the obtained graph is sufficiently sparse, it will contain an induced subgraph allowing a polyhedral reduction from the correlation polytope, and of sufficient size. Via this reduction we derive strong lower bounds on the size of any LP expressing $\text{STAB}(G)$ that hold with high probability. In particular, we obtain superpolynomial lower bounds for p ranging between $\Omega(\frac{\log^{6+\varepsilon} n}{n})$ and $o(\frac{1}{\log n})$. For example for $p = n^{-\varepsilon}$ and $\varepsilon < 1/4$, any LP has at least $2^{\Omega(\sqrt{n^\varepsilon \log n})}$ constraints w.h.p. (with high probability), and for $p = \Omega(\frac{\log^{6+\varepsilon} n}{n})$, any LP has at least $n^{\log(3/2) \log^{\varepsilon/5} n}$ constraints w.h.p. Figure 1 illustrates our lower bounds.




Figure 1: Comparing lower bounds on $\text{fc}(G(n, p))$ for various regimes. For p close to $1/\sqrt[3]{n}$ the blue and green lines provide roughly the same bounds. For p significantly above $1/\sqrt[3]{n}$ the magenta line outperforms the green line.

1.3 Outline

In Section 2 we recall basics on extended formulations. We introduce the model of general linear programming formulations in Section 2.1. We then establish bounds on the average case complexity for the uniform model of the maximum stable set problem in Section 3. In Section 4 we consider the non-uniform model and derive lower bounds as well as upper bounds. We conclude with a conjecture in Section 5.

2 Preliminaries

We use \log to denote the base-2 logarithm. Let G be a graph. We denote by $\alpha(G)$ the maximum size of a stable set in G . This is the *stability number* of G . If S is any set, we let $G[S]$ denote the induced subgraph of G on $V(G) \cap S$. For a positive integer n we let $[n] := \{1, 2, \dots, n\}$.

2.1 Linear programming formulations

The notion of linear programming formulation we introduce here generalizes that appearing in the work of Chan et al. [2013] on MAX CSPs and is related to the notion of faithful linear encodings, their corresponding pairs of polyhedra and extended formulations thereof, see Braun et al. [2012, 2014]. Our aim is to obtain lower bounds that are *independent* of the actual encoding of the optimization problem as a linear program (see Braun and Pokutta [2014] for more details).

We will first specify the notion of an optimization problem. For the sake of exposition, we restrict ourselves to maximization problems.

Definition 2.1 (Maximization problem). A *maximization problem* $\Pi = (\mathcal{S}, \mathcal{F}, *)$ consists of a finite set \mathcal{S} of *feasible solutions*, a finite set \mathcal{F} of *objective functions* where $f: \mathcal{S} \rightarrow \mathbb{R}$ for each $f \in \mathcal{F}$, and an *approximation guarantee* $f^* \in \mathbb{R}$ so that $f^* \geq \max_{s \in \mathcal{S}} f(s)$ for each objective function $f \in \mathcal{F}$.

By solving such a maximization problem $\Pi = (\mathcal{S}, \mathcal{F}, *)$ we mean determining an approximation $\hat{f} \in \mathbb{R}$ of the optimum value $\max_{s \in \mathcal{S}} f(s)$ satisfying

$$\max_{s \in \mathcal{S}} f(s) \leq \hat{f} \leq f^*,$$

for each $f \in \mathcal{F}$. Our next definition specifies what it means to formulate the maximization problem as a linear program.

Definition 2.2 (LP formulation of a maximization problem). A *linear programming formulation* of a maximization problem $\Pi = (\mathcal{S}, \mathcal{F}, *)$ is a linear system $A^{\leq} x \leq b^{\leq}$, $A^= x = b^=$ with $x \in \mathbb{R}^d$ together with *realizations* of:

(i) **feasible solutions** as points $x^s \in \mathbb{R}^d$ for each $s \in \mathcal{S}$, such that

$$A^{\leq} x^s \leq b^{\leq}, A^= x^s = b^=; \quad (1)$$

(ii) **objective functions** as linear functions on \mathbb{R}^d , i.e., for each $f \in \mathcal{F}$ there is $w^f \in \mathbb{R}^d$ with

$$\langle w^f, x^s \rangle = f(s) \quad \text{for all } s \in \mathcal{S}. \quad (2)$$

Conditions (1) and (2) imply that

$$\max_{s \in \mathcal{S}} f(s) \leq \max \left\{ \langle w^f, x \rangle \mid A^{\leq} x \leq b^{\leq}, A^= x = b^= \right\} \quad \text{for all } f \in \mathcal{F}.$$

We additionally require that

$$\max \left\{ \langle w^f, x \rangle \mid A^{\leq} x \leq b^{\leq}, A^= x = b^= \right\} \leq f^* \quad \text{for all } f \in \mathcal{F}, \quad (3)$$

so that the optimum value of the LP provides an approximation of the optimum value $\max_{s \in \mathcal{S}} f(s)$. The *size* of the formulation is number of inequalities in the LP, that is, the number of constraints in $A^{\leq} x \leq b^{\leq}$. We define the *formulation complexity* $\text{fc}(\Pi)$ of the problem Π as the minimum size of all its LP formulations.

We point out that the above definition encompasses the concept of extended formulation, in the following sense. If instead of the constraints $A^{\leq} x \leq b^{\leq}$, $A^= x = b^=$ we used an extended formulation $E^{\leq} x + F^{\leq} y \leq g^{\leq}$, $E^= x + F^= y = g^=$ to express the set over which we want to maximize $\langle w^f, x \rangle$, where $y \in \mathbb{R}^k$ is an extra (vector) variable, we could redefine the ambient space, the points x^s ($s \in \mathcal{S}$) and the coefficient vectors w^f ($f \in \mathcal{F}$) to eliminate the extra variable y . Indeed, we could work directly in \mathbb{R}^{d+k} instead of \mathbb{R}^d , consider the vectors $(w^f, 0)$ instead of w^f and the points (x^s, y^s) instead of x^s , where $y^s \in \mathbb{R}^k$ is chosen so that (x^s, y^s) satisfies the constraints of the extended formulation. (Such a point y^s exists because by Condition (1), x^s is feasible for the LP.)

In Braun et al. [2012, 2014], we start from a *specific* encoding of the problem by points x^s ($s \in \mathcal{S}$) and coefficient vectors w^f ($f \in \mathcal{F}$), then infer from this a pair (P, Q) of nested polyhedra where

$$\begin{aligned} P &:= \text{conv} \{ x^s \mid s \in \mathcal{S} \}, \\ Q &:= \left\{ x \in \mathbb{R}^d \mid \langle w^f, x \rangle \leq f^*, \forall f \in \mathcal{F} \right\}, \end{aligned}$$

and finally consider any extended formulation of the pair (P, Q) . The approach taken here is at the same time more direct and more general.

Now we describe two ways in which the maximum stable set problem can be seen as a maximization problem $\Pi = (\mathcal{S}, \mathcal{F}, *)$ that we use in the rest of the paper.

Definition 2.3 (The maximum stable set problem—uniform model). We start with a family of graphs \mathcal{G} with $V(G) \subseteq [n]$ for each $G \in \mathcal{G}$. The goal is to approximate the stability number of the graphs in \mathcal{G} within a given relative error guarantee $\rho \geq 0$. To each graph $G \in \mathcal{G}$ we correspond an objective function f_G in the set \mathcal{F} of objective functions. The set \mathcal{S} of feasible solutions is taken to be the set of *all* subsets of $[n]$. This is natural since, typically, \mathcal{G} contains many graphs with many different vertex sets and many $S \subseteq [n]$ occur as a stable set in some $G \in \mathcal{G}$. We require

- (a) $\alpha(G) = \max_{S \in \mathcal{S}} f_G(S)$, for all $G \in \mathcal{G}$;
- (b) $f_G(S) = |V(G) \cap S|$ whenever $|V(G) \cap S| \leq 1$, for all $G \in \mathcal{G}$ and $S \in \mathcal{S}$.

We choose as approximation guarantee $f_G^* := (1 + \rho)\alpha(G)$ for $G \in \mathcal{G}$. This defines a *class* of maximization problems that we denote by $\text{STAB}^u(\mathcal{G}, \rho)$: every set \mathcal{F} of objective functions subject to the above conditions gives a valid approximated computation of the stability number in a graph G chosen from \mathcal{G} . The later derived lower bounds apply to every problem Π from this class and we define $\text{fc}(\text{STAB}^u(\mathcal{G}, \rho)) := \min \{\text{fc}(\Pi) \mid \Pi \in \text{STAB}^u(\mathcal{G}, \rho)\}$. In the exact case, that is, when $\rho = 0$, we use $\text{STAB}^u(\mathcal{G})$ to mean $\text{STAB}^u(\mathcal{G}, 0)$.

As an example, a concrete maximization problem $\Pi \in \text{STAB}^u(\mathcal{G}, \rho)$ is obtained by defining the objective functions as

$$f_G(S) := |V(G) \cap S| - |E(G[S])|.$$

Here $f_G(S)$ is a conservative lower bound on the size of a stable set, which one would naturally obtain from $G[S]$ by deleting one endpoint from every edge in that induced subgraph. Moreover we have $f_G(S) = |S|$ for stable sets S of G , i.e., in this case our choice is exact. We leave it to the reader to check Condition (a) while Condition (b) is immediate.

Definition 2.4 (The maximum stable set problem—non-uniform model). To each *fixed* graph G on n vertices we associate a problem that corresponds to the exact computation of the stability number of some induced subgraph H of G . We will denote this problem by $\text{STAB}^{uu}(G)$ or (later) simply $\text{STAB}(G)$. More precisely, $\text{STAB}^{uu}(G)$ has feasible solutions that are sets $S \subseteq V(G)$ which are stable sets of G , and objective functions of the form $f_H(S) := |S \cap V(H)|$ where H is an induced subgraph of G . We let $f_H^* := \alpha(H)$ for the approximation guarantee.

2.2 Size lower bounds from nonnegative rank

The basis of all lower bounds for extended formulations is Yannakakis's celebrated Factorization Theorem (see Yannakakis [1991, 1988]), equating the minimal size of an extended formulation with the nonnegative rank of a slack matrix. We will now derive a factorization theorem in a similar spirit that characterizes formulation complexity.

A *rank- r nonnegative factorization* of $M \in \mathbb{R}_+^{m \times n}$ is a factorization of $M = TU$ where $T \in \mathbb{R}_+^{m \times r}$ and $U \in \mathbb{R}_+^{r \times n}$. This is equivalent to $M = \sum_{i \in [r]} u_i v_i^T$ for some (column vectors) $u_i \in \mathbb{R}_+^m$, $v_i \in \mathbb{R}_+^n$ with $i \in [r]$. The *nonnegative rank* of M , denoted by $\text{rank}_+ M$, is the minimum r such that there exists a rank- r nonnegative factorization of M .

Theorem 2.5 (Factorization theorem for formulation complexity). *Consider a maximization problem $\Pi = (\mathcal{S}, \mathcal{F}, *)$ as in Definition 2.1. The slack matrix of Π is the nonnegative $\mathcal{F} \times \mathcal{S}$ matrix M , with entries*

$$M(f, s) := f^* - f(s).$$

Then

$$\text{rank}_+ M - 1 \leq \text{fc}(P) \leq \text{rank}_+ M.$$

Proof. To prove the first inequality, let $A^{\leq} x \leq b^{\leq}$, $A^= x = b^=$ be an arbitrary size- r LP formulation of Π , with realizations w^f ($f \in \mathcal{F}$) of objective functions and x^s ($s \in \mathcal{S}$) of feasible solutions. We shall construct a size- $(r + 1)$ nonnegative factorization of M . As $\max_{x: Ax \leq b} \langle w^f, x \rangle \leq f^*$ by Condition (3), via Farkas's lemma, we have

$$f^* - \langle w^f, x \rangle = T(f, 0) + \sum_{j=1}^r T(f, j) (b_j^{\leq} - \langle A_j^{\leq}, x \rangle) + \sum_{j=1}^k \lambda(f, j) (b_j^= - \langle A_j^=, x \rangle)$$

for some nonnegative multipliers $T(f, j) \in \mathbb{R}_+$ with $0 \leq j \leq r$, and arbitrary multipliers $\lambda(f, j)$ for $0 \leq j \leq k$. By taking $x = x^s$, and by $A^= x^s = b^=$, we obtain

$$M(f, s) = \sum_{j=0}^r T(f, j) U(j, s), \quad \text{with } U(j, s) := \begin{cases} 1 & \text{for } j = 0, \\ b_j^{\leq} - \langle A_j^{\leq}, x^s \rangle & \text{for } j > 0. \end{cases}$$

That is, $M = TU$. By construction, T is nonnegative. By Condition (1) we also obtain that U is nonnegative. Therefore $M = TU$ is a rank- $(r + 1)$ nonnegative factorization of M .

For the second inequality, let $M = TU$ be a size- r nonnegative factorization. We shall construct an LP formulation of size r . Let T^f denote the f -row of T for $f \in \mathcal{F}$, and U_s denote the s -column of U for $s \in \mathcal{S}$. Then

$$T^f U_s = M(f, s) = f^* - f(s). \quad (4)$$

In the following we represent the vectors $y \in \mathbb{R}^{r+1}$ via $y = (x, \alpha)$ with $x \in \mathbb{R}^r$ and $\alpha \in \mathbb{R}$. We claim that the linear system

$$x \leq 0, \alpha = 1 \quad (5)$$

with representations

$$w^f := (T^f, f^*) \quad \forall f \in \mathcal{F} \quad \text{and} \quad x^s := (-U_s, 1) \quad \forall s \in \mathcal{S}$$

satisfies the requirements of Definition 2.2. Condition (2) clearly follows from (4):

$$\langle w^f, x^s \rangle = -T^f U_s + f^* = f(s).$$

Moreover, the x^s satisfy the linear program (5), because U is nonnegative, so that Condition (1) is fulfilled. Finally, Condition (3) also follows readily:

$$\max \left\{ \langle w^f, (x, \alpha) \rangle \mid x \leq 0, \alpha = 1 \right\} = \max \left\{ T^f x + f^* \mid x \leq 0 \right\} = f^*,$$

as the nonnegativity of T implies $T^f x \leq 0$; equality holds e.g., for $x = 0$. Thus we have constructed an LP formulation with r inequalities, as claimed. \square

Remark 2.6. *The slack matrix of the problem $\text{STAB}^{\text{nu}}(G)$ is actually the submatrix of the slack matrix of the stable set polytope of G corresponding to the rank inequalities $\sum_{v \in V(H)} x_v \leq \alpha(H)$ for H an induced subgraph of G (see, e.g., Fiorini et al. [2012] for definitions). Thus the abuse of notation carried out by using $\text{STAB}(G)$ to denote the problem $\text{STAB}^{\text{nu}}(G)$ is not too severe.*

We will use the following elementary properties of nonnegative rank.

Lemma 2.7. *Let $A \in \mathbb{R}_+^{k \times m}$, $M \in \mathbb{R}_+^{m \times n}$ and $B \in \mathbb{R}_+^{n \times l}$ be nonnegative matrices. Then $\text{rank}_+(AMB) \leq \text{rank}_+ M$. In particular, the following operations do not increase the nonnegative rank of a matrix:*

- (i) *deleting, duplicating, permuting rows or columns;*
- (ii) *adding nonnegative linear combination of rows as a new row;*
- (iii) *adding nonnegative linear combination of columns as a new column.*

By considering the nonnegative factorizations $M = I_m M$ and $M = M I_n$, we immediately obtain

$$\text{rank}_+ M \leq \min\{m, n\}. \quad (6)$$

2.3 Unique disjointness

As we will demonstrate, the polyhedral hardness of the maximum stable set problem arises from the unique disjointness (partial) matrix. Recall that the unique disjointness (UDISJ) matrix, which we denote by $\text{UDISJ}(n)$ below, has 2^n rows and 2^n columns indexed by all size- n 0/1-vectors a and b . Its entries are:

$$\text{UDISJ}(n)(a, b) = \begin{cases} 0 & \text{if } a^\top b = 1 \\ 1 & \text{if } a^\top b = 0. \end{cases} \quad (7)$$

Although $\text{UDISJ}(n)$ is only a partial matrix, i.e., not all of its entries are defined, we will refer to it as a matrix from here on. The fact that it is only partial does not matter for our purpose, as we only care for whether this (partial) matrix occurs as a submatrix of some appropriate slack matrices. The UDISJ matrix has been studied in many disciplines, arguably the most notable being communication complexity.

Theorem 2.8. $\text{rank}_+ \text{UDISJ}(n) \geq 2^{n \cdot \log(3/2)}$.

The factor $\log(3/2) \approx 0.585$ in the exponent is the current best one due to Kaibel and Weltge [2013]; for various approximate case versions see Braun et al. [2012, 2014], Braverman and Moitra [2013], Braun and Pokutta [2013]. The first exponential lower bound was established in Fiorini et al. [2012] by combining the seminal work of Razborov [1992] together with an observation in Wolf [2003]. This was at the core of the first results establishing high extension complexity for the correlation polytope, cut polytope, stable set polytope, and the TSP polytope in Fiorini et al. [2012].

Braun et al. [2012, 2014] prove that *any* $2^n \times 2^n$ matrix M with rows and columns indexed by vectors in $\{0, 1\}^n$ satisfying (7) has superpolynomial nonnegative rank, and that this remains true even if we shift the entries of the matrix M by some number $\rho = O(n^{1/2-\epsilon})$. This result was then extended to shifts $\rho = O(n^{1-\epsilon})$ in Braverman and Moitra [2013] which then immediately leads to a polyhedral inapproximability of CLIQUE (in the uniform model!) of $O(n^{1-\epsilon})$, matching Håstad's hardness result for approximating CLIQUE.

The ρ -shifted UDISJ matrix is any $2^n \times 2^n$ matrix indexed by pairs (a, b) where $a, b \in \{0, 1\}^n$ such that

$$(\text{UDISJ}(n) + \rho J)(a, b) = \begin{cases} \rho & \text{if } a^\top b = 1 \\ 1 + \rho & \text{if } a^\top b = 0, \end{cases}$$

where J is the all-one matrix of compatible size.

In Braun and Pokutta [2013] an information-theoretic approach for studying the nonnegative rank has been developed. This approach allows to lower bound the nonnegative rank of various

‘deformations’ of the UDISJ matrix. The following theorem from Braun and Pokutta [2013] will allow us to analyze a specific type of deformation that we will use in the following. Informally speaking, the theorem shows that the UDISJ matrix has high nonnegative matrix almost everywhere. Below, we use $\text{UDISJ}(n, k)$ to denote the UDISJ matrix $\text{UDISJ}(n)$ restricted to subsets of size k .

Theorem 2.9. *Let M be any submatrix of the ρ -shifted UDISJ matrix $\text{UDISJ}(n, k) + \rho J$ obtained by deleting at most an α -fraction of rows and at most a β -fraction of columns for some $0 \leq \alpha, \beta < 1$. Then for $0 < \varepsilon < 1$:*

$$\text{rank}_+ M \geq 2^{(1/8(\rho+1) - (\alpha+\beta)\mathbb{H}[1/4])n - O(n^{1-\varepsilon})} \quad \text{for } k = n/4 + O(n^{1-\varepsilon}).$$

We finish this section with an easy example of embedding $\text{UDISJ}(n)$ into a slack matrix.

Example 2.10. Let us consider the family \mathcal{G} of all (non-empty) complete graphs, that is, all graphs G with $V(G) \subseteq [n]$ and $\alpha(G) = 1$ (in case $V(G) = \emptyset$ we have $\alpha(G) = 0$). Then the slack matrix M of any $\Pi \in \text{STAB}^u(\mathcal{G})$ contains $\text{UDISJ}(n)$ as a submatrix (without the row of the empty set $a = \emptyset$). To verify this, note that by Condition (b) for every pair of subsets $a, b \subseteq [n]$ (with $a \neq \emptyset$):

$$M(K_n[a], b) = 1 - |a \cap b| = \begin{cases} 0 & \text{if } |a \cap b| = 1 \\ 1 & \text{if } |a \cap b| = 0, \end{cases}$$

where K_n denotes the complete graph on $[n]$.

3 Average case complexity in the uniform model

We will now establish our main result regarding the average case complexity of the uniform model. We obtain that for any random collection of graphs $\mathcal{G} = \mathcal{G}(n, p)$ where each graph G with $V(G) \subseteq [n]$ is picked independently with probability p , the formulation complexity of $\text{STAB}^u(\mathcal{G})$ is high. Loosely speaking, the size of any “simultaneous” LP formulation of the maximum stable set problem for all graphs in this random collection \mathcal{G} is high. In a way, this indicates that the instances of the stable set problem resulting in high polyhedral complexity are not localized in a set of small density.

Main Theorem 3.1 (Super-polynomial fc of $\text{STAB}^u(\mathcal{G})$ w.h.p.). *Let $n \geq 40$ and $p \in [0, 1]$ with $p \geq 2^{-(\binom{n}{2})+n}$. Pick a random family $\mathcal{G} = \mathcal{G}(n, p)$ of graphs by adding each graph G with $V(G) \subseteq [n]$ to the family with probability p , independent of the other G . Then*

$$\mathbb{P} \left[\text{fc}(\text{STAB}^u(\mathcal{G})) \geq 2^{\Omega(n/\log n)} \right] \geq 1 - 2^{-2^n}.$$

A crucial point of the proof is a concentration result on $\alpha(G)$. It is well-known that almost all graphs G on n vertices have stability number $\alpha(G) \sim 2 \log n$. However, the following rough estimate will be sufficient for our purpose, see e.g. [Diestel, 2005, Proposition 11.3.4, page 304] for a proof.

Lemma 3.2. *Let $n \geq 10$. The probability that a uniformly sampled random graph G with $V(G) = [n]$ has $\alpha(G) \geq 3 \log n$ is at most n^{-1} .*

We are ready to prove the main theorem of this section.

Proof of Main Theorem 3.1. Consider any problem $\Pi \in \text{STAB}^u(\mathcal{G})$. Thus, $\Pi = (\mathcal{S}, \mathcal{F}, *)$ is a maximization problem that corresponds to determining the stability number of the graphs of \mathcal{G} . Recall that \mathcal{S} is the set of all subsets of $S \subseteq [n]$ and that \mathcal{F} contains an objective function f_G for each graph $G \in \mathcal{G}$. These functions satisfy $\max_{S \in \mathcal{S}} f_G(S) = \alpha(G)$ (see Condition (a)). Also, recall that we require $f_G(S) = |V(G) \cap S|$ whenever $|V(G) \cap S| \leq 1$ (see Condition (b)). Finally, since $\rho = 0$, the approximation guarantee is $f_G^* = \alpha(G)$.

Consider the slack matrix M of Π . We want to show that the nonnegative rank of M is high by embedding a large portion of UDISJ (n) into it. By Theorem 2.5, this will imply that $\text{fc}(\Pi)$ and thus $\text{fc}(\text{STAB}^u(\mathcal{G}))$ is high, since $\Pi \in \text{STAB}^u(\mathcal{G})$ is arbitrary.

The main idea of the proof is that, with extremely large probability, among all sets $a \subseteq [n]$ of size $\lceil n/4 \rceil$, the collection \mathcal{G} contains many graphs G with $V(G) = a$. For each of these sets a , there will be at least one corresponding graph G_a with $\alpha(G_a) \leq 3 \log n$. Restricting to these graphs G_a , the resulting slack matrix contains a large part of the $O(\log n)$ -shifted UDISJ matrix as a submatrix (in fact, a large fraction of the rows, and all the columns, survive). We apply Theorem 2.9 to conclude.

Now let us turn to the detailed proof. Consider a set a with $a \subseteq [n]$ and size $k := \lceil n/4 \rceil$. We say that a graph G is *good* for a if $V(G) = a$ and $\alpha(G) \leq 3 \log n$. Set a is said to be *good* if some graph $G \in \mathcal{G}$ is good for a . Otherwise, a is called *bad*.

We claim that, with high probability, the total fraction of bad sets among all k -sets a is at most $\alpha := 1/(24 \log n)$. By Lemma 3.2, the total number of graphs G with $V(G) = a$ that are not good for a fixed k -set a is at most $k^{-1} 2^{\binom{k}{2}}$. Thus

$$\begin{aligned} \mathbb{P}[a \text{ is bad}] &= \mathbb{P}[\mathcal{G} \text{ contains no good graph for } a] \\ &\leq (1-p)^{(1-k^{-1})2^{\binom{k}{2}}} \\ &\leq e^{-p(1-k^{-1})2^{\binom{k}{2}}} \\ &\leq 2^{-\frac{9}{10}2^n \log e} \\ &\leq \alpha 2^{-2^n}. \end{aligned} \tag{8}$$

where the third inequality follows from $k \geq n/4 \geq 10$ and $p \geq 2^{-(\frac{n}{4})+n}$ and the last inequality follows from our choice of α . Let X denote the random variable with value the number of bad k -sets a . By Markov's inequality,

$$\mathbb{P}\left[X \geq \alpha \binom{n}{k}\right] \leq 2^{-2^n}.$$

For each good k -set a , pick a good graph $G_a \in \mathcal{G}$ arbitrarily, i.e., with $\alpha(G_a) \leq 3 \log n$. We need two auxiliary nonnegative vectors. Let $\mathbf{1}$ denote the row vector with all entries 1, and entries indexed by subsets $S \subseteq [n]$. Let u be the vector with entries indexed by the G_a for all good k -sets a , and with entries

$$u_{G_a} := 3 \log n - \alpha(G_a) \geq 0.$$

Because the slack matrix M of Π satisfies

$$M(G_a, S) = \begin{cases} \alpha(G_a) - 1 & \text{if } |V(G) \cap S| = 1, \\ \alpha(G_a) & \text{if } |V(G) \cap S| = 0, \end{cases}$$

we obtain after applying the rank-1 shift $u\mathbf{1}$

$$(M + u\mathbf{1})(G_a, S) = \begin{cases} 3 \log n - 1 & \text{if } |V(G) \cap S| = 1, \\ 3 \log n & \text{if } |V(G) \cap S| = 0. \end{cases}$$

By the above, with probability at least $1 - 2^{-2^n}$, the fraction of bad k -sets $a \subseteq [n]$ among all k -sets is at most α , and hence the matrix $(M + u\mathbf{1})$ contains a $(3 \log n - 1)$ -shift of $\text{UDISJ}(n, k)$, with at most an α -fraction of the rows thrown away. From Theorem 2.9 (with $\beta = 0$), the nonnegative rank of M is at least

$$\text{rank}_+ M \geq \text{rank}_+(M + u\mathbf{1}) - 1 \geq 2^{(1/8(3 \log n + 1) - \alpha H[1/4]) \cdot n - O(n^{1-\varepsilon})} - 1 = 2^{\Omega(n/\log n)}.$$

□

Without much additional work, we can obtain a similar lower bound on the average case formulation complexity also in the approximate case, that is, when $\rho > 0$.

Corollary 3.3 (Super-polynomial xc of $\text{STAB}^u(\mathcal{G}, \rho)$ w.h.p.). *As in Main Theorem 3.1, let $\mathcal{G} = \mathcal{G}(n, \rho)$ be a random family of graphs such that each graph G with $V(G) \subseteq [n]$ is contained in \mathcal{G} with probability $p \geq 2^{-(\binom{n/4}{2} + n)}$ independent of the other graphs. Then for all $0 < \varepsilon < 1/2$ and $\rho \leq \frac{n^{1-\varepsilon}}{\log n}$, we have that $\text{STAB}^u(\mathcal{G}, \rho)$ has formulation complexity $2^{\Omega(n^\varepsilon)}$, with probability at least $1 - 2^{-2^n}$.*

Proof. The proof is identical to Theorem 3.1 subject to minor changes. First, now we use a different factor $\alpha := 1/[24(1 + \rho) \log n]$. The computation in Equation (8) still remains valid, as the different choice of α affects only the last inequality, which remains true, because $\rho \leq n^{1-\varepsilon}/\log n$. Therefore again with probability $1 - 2^{-2^n}$, with the exception of at most an $1/[24(1 + \rho) \log n]$ -fraction, all graphs $G \in \mathcal{G}$ are good, i.e., have clique number at most $\alpha(G) \leq 3 \log n$.

The second difference is that due to dilation, the slack entries of M are a bit different:

$$M(G, S) = \begin{cases} (1 + \rho)\alpha(G) - 1 & \text{if } |V(G) \cap S| = 1, \\ (1 + \rho)\alpha(G) & \text{if } |V(G) \cap S| = 0. \end{cases}$$

This provides an embedded copy of a $(3(1 + \rho) \log n - 1)$ -shift of $\text{UDISJ}(n, k)$ in $M + (1 + \rho)u\mathbf{1}$ with at most an α -fraction of rows missing. Hence Theorem 2.9 applies again, but now we replace the ε there with a ε' lying strictly between ε and $1/2$, but not depending on n . (E.g., $\varepsilon' := (\varepsilon + 1/2)/2$ is a good choice.) This will make the error term $O(n^{1-\varepsilon'}) = o(n^{1-\varepsilon})$ in the exponent negligible. We obtain the lower bound: $2^{(1/8(3(1+\rho) \log n + 1) - \alpha H[1/4]) \cdot n - O(n^{1-\varepsilon'})} - 1 = 2^{\Omega(n^{1-\varepsilon})}$ on formulation complexity, as claimed. □

Observe that the relative approximation guarantee ρ in Corollary 3.3 can be larger than $3 \log n$. The reason why this is possible, contradicting initial intuition, is that the hardness arises from having many different graphs and hence many objective functions to consider simultaneously and the encoding is highly non-monotone. Roughly speaking, graphs with different vertex sets are independent of each other, even if one is an induced subgraph of the other.

4 Average case complexity in the non-uniform model

We now turn our attention to the non-uniform problem $\text{STAB}^{uu}(G)$, see Definition 2.4. Thus G is a fixed graph with vertex set $[n]$, the feasible solutions are the stable sets of G and the objective functions correspond to induced subgraphs of G . For simplicity of notation, we index the objective functions with the supporting vertex set $a \subseteq [n]$ instead of the induced subgraph $G[a]$. Thus $f_a(S) = |S \cap a|$ for every stable set S of G and $a \subseteq [n]$, and $f_a^* = \alpha(G[a])$.

Notice that an LP formulation for the problem $\text{STAB}^{uu}(G)$ is provided by the linear description of the stable set polytope of G , or any extended formulation of the stable polytope of G . In

this sense, $\text{STAB}^{\text{nu}}(G)$ generalizes the stable set polytope. For the sake of brevity, we denote the problem $\text{STAB}^{\text{nu}}(G)$ simply by $\text{STAB}(G)$.

We lower bound the formulation complexity of $\text{STAB}(G(n, p))$ for the *random* Erdős–Rényi graph $G(n, p)$. Our strategy is to embed certain subdivisions of the complete graph K_t as induced subgraphs of G , with t as large as possible, using the probabilistic method.

Our construction is parametrized by an *even* integer $\ell \geq 0$. For a graph T , we let T^\sim denote the subdivision of T obtained by replacing each edge ij of T with a path P_{ij} with $2\ell + 3$ edges between i and j . We denote u_{ij} and v_{ij} the middle vertices of P_{ij} , see Figure 2. In total, T^\sim has $v := |V(T)| + (2\ell + 2)|E(T)|$ vertices and $e := (2\ell + 3)|E(T)|$ edges.




Figure 2: Path P_{ij} replacing edge ij of T in T^\sim . There are $\ell + 1$ edges between i and u_{ij} , as between v_{ij} and j . In the figure, $\ell = 0$.

Our next lemma proves that increasing the parameter ℓ decreases the average degree of induced subgraphs of the gadget graph T^\sim , which makes it easier to embed T^\sim in $G(n, p)$ for lower values of p .

Lemma 4.1. *For any graph T , the average degree of any induced subgraph of T^\sim is at most $2 + 1/(\ell + 1)$. For $\ell = 0$, the average degree is at most 3.*

Proof. Consider an induced subgraph of T^\sim with maximum average degree. Clearly, we may assume that T contains at least one edge, so the average degree of the induced subgraph is at least $3/2$. This implies that it does not contain any vertex of degree at most 1, because the deletion of any such vertex would increase the average degree. It follows that the induced subgraph is of the form H^\sim where H is a subgraph of T . The average degree of H^\sim can be expressed in terms of that of H as:

$$\bar{d}(H^\sim) = \frac{2|E(H^\sim)|}{|V(H^\sim)|} = \frac{2(2\ell + 3)|E(H)|}{|V(H)| + (2\ell + 2)|E(H)|} = \frac{(2\ell + 3)\bar{d}(H)}{1 + (\ell + 1)\bar{d}(H)}.$$

From the last expression we see that the average degree of H^\sim is an increasing function of the average degree of H that tends to $(2\ell + 3)/(\ell + 1) = 2 + 1/(\ell + 1)$ in the limit. \square

Lemma 4.2. *If graph G contains K_t^\sim as an induced subgraph, then*

$$\text{fc}(\text{STAB}(G)) \geq \text{rank}_+ \text{UDISJ}(t) - 2 \geq 2^{t \log(3/2)} - 2.$$

Proof. Let T be any graph with T^\sim being an induced subgraph of G . Later we will specialize to $T = K_t$. We choose representatives of subsets of $V(T)$ as stable sets and induced subgraphs of T^\sim .

For every $b \subseteq V(T)$, we choose an extension to a stable set $S(b)$ of T^\sim by adding as much internal vertices of each path P_{ij} as possible, see Figure 3. Let $ij \in E(T)$. On the part of P_{ij} from i to u_{ij} , as well on the part from j to v_{ij} , we alternate between vertices belonging to T^\sim and not belonging to T^\sim , with one exception: if both $i, j \in b$ then we drop either u_{ij} or v_{ij} in $S(b)$.

If we start from a maximum stable set b of T , we see that $S(b)$ has $|b| + (\ell + 1)|E(T)|$ vertices. Thus $\alpha(T^\sim) \geq \alpha(T) + (\ell + 1)|E(T)|$. This inequality is tight because no stable set S of T^\sim can have more vertices than $S(b)$, where $b := S \cap V(T)$. That is, $|S| \leq |b| + (\ell + 1)|E(T)| - |E(T[b])| \leq \alpha(T) + (\ell + 1)|E(T)|$. For any graph T , we get:

$$\alpha(T^\sim) = \alpha(T) + (\ell + 1)|E(T)|. \tag{9}$$




Figure 3: Possible stable sets $S(b)$ extending a given $b \subseteq V(T)$. Black vertices are those which are part of $S(b)$.

For every subset $a \subseteq V(T)$, consider the induced subgraph $T[a]^\sim$ of T^\sim . By (9), we have $\alpha(T[a]^\sim) = \alpha(T[a]) + (\ell + 1)|E(T[a])|$. We also consider the induced subgraph $T[a]^\circ$ on all the u_{ij} and v_{ij} in $T[a]^\sim$. This is a matching, so obviously $\alpha(T[a]^\circ) = |E(T[a])|$.

By construction, for all sets $a, b \subseteq V(T)$ we have:

$$\begin{aligned} |V(T[a]^\sim) \cap S(b)| &= |S \cap a| + (\ell + 1) |E(T[a])| - |E(T[a \cap b])|, \\ |V(T[a]^\circ) \cap S(b)| &= |E(T[a])| - |E(T[a \cap b])|. \end{aligned}$$

From the slack matrix M of $\text{STAB}(G)$, we construct a matrix N with rows and columns indexed by all subsets a, b of $V(T)$ with $a \neq \emptyset$, with entries

$$\begin{aligned} N(a, b) &:= M(T[a]^\sim, S(b)) + M(T[a]^\circ, S(b)) \\ &= \alpha(T[a]^\sim) - |V(T[a]^\sim) \cap S(b)| + \alpha(T[a]^\circ) - |V(T[a]^\circ) \cap S(b)| \\ &= \alpha(T[a]) + 2|E(T[a \cap b])| - |a \cap b|. \end{aligned}$$

Specializing to $T = K_t$, we obtain for $a \neq \emptyset$:

$$N(a, b) = 1 + 2 \binom{|a \cap b|}{2} - |a \cap b| = (1 - |a \cap b|)^2 = \begin{cases} 1 & \text{if } |a \cap b| = 0, \\ 0 & \text{if } |a \cap b| = 1. \end{cases}$$

Thus, N contains $\text{UDISJ}(t)$ as a submatrix without the row of the empty set. Now Theorem 2.5 followed by Lemma 2.7 implies

$$\text{fc}(\text{STAB}(G)) \geq \text{rank}_+ M - 1 \geq \text{rank}_+ N - 1 \geq \text{rank}_+ \text{UDISJ}(t) - 2 \geq 2^{t \log(3/2)} - 2.$$

□

4.1 Existence of gadgets in random graphs

In this section, we estimate the probability that a random Erdős–Rényi graph $G = G(n, p)$ contains an induced copy of a graph H . Recall that in the $G(n, p)$ model, each of the $\binom{n}{2}$ pairs of vertices is connected by an edge with probability p , independently from the other edges. The next lemma is key for proving lower bounds on the formulation complexity of $\text{STAB}(G(n, p))$ via embedding $H = T^\sim$ as an induced subgraph. The lemma is formulated in a general for future applications to many types of subgraphs H .

Lemma 4.3. Let H be a graph with v vertices and with all induced subgraphs having average degree at most d . Let $0 < p \leq 1/2$ and

$$g = g(n, p, v) := \frac{v^2 p^{-\frac{d}{2}} (1-p)^{-\frac{v}{2}}}{n-v}.$$

The probability of $G(n, p)$ not containing an induced copy of H satisfies

$$\mathbb{P} \left[H \not\subseteq^{\text{ind}} G(n, p) \right] \leq c_0 g^2 \approx 1.23 g^2,$$

where $c_0 := \exp(2W(1/\sqrt{2}))/2$ and W is the Lambert W -function, the inverse of $x \rightarrow x \exp x$.

Proof. The proof is via the second-moment method.

Let S be any graph isomorphic to H with $V(S) \subseteq V(G)$. Let X_S be the indicator random variable of S being an induced subgraph of G . Obviously, the total number X of induced subgraphs of G isomorphic to H satisfies $X = \sum_S X_S$. We estimate the expectation and variance of X . Let e denote the number of edges of H , and let $\text{Aut}(H)$ denote the automorphism group of H . The expectation is clearly

$$\mathbb{E}[X] = \sum_S \mathbb{E}[X_S] = \binom{n}{v} \frac{v!}{|\text{Aut}(H)|} p^e (1-p)^{\binom{v}{2}-e}.$$

The variance needs more preparations. Let now S and T be two graphs isomorphic to H with $V(S), V(T) \subseteq V(G)$. Using that X_S and X_T are independent and thus $\text{Cov}[X_S, X_T] = 0$ when $|V(S) \cap V(T)| \leq 1$ we get

$$\begin{aligned} \text{Var}[X] &= \sum_{S, T} \text{Cov}[X_S, X_T] \leq \sum_{|V(S) \cap V(T)| \geq 2} \mathbb{E}[X_S X_T] \\ &= \sum_{|V(S) \cap V(T)| \geq 2} \mathbb{E}[X_S] \mathbb{E}[X_T | X_S = 1] = \mathbb{E}[X] \sum_{T: |V(S) \cap V(T)| \geq 2} \mathbb{E}[X_T | X_S = 1]. \end{aligned}$$

Note that in the last sum S is fixed, and by symmetry, the sum is independent of the actual value of S . That is why we could factor it out. We obtain via Chebyshev's inequality,

$$\mathbb{P} \left[H \not\subseteq^{\text{ind}} G(n, p) \right] = \mathbb{P}[X = 0] \leq \frac{\text{Var}[X]}{\mathbb{E}[X]^2} \leq \frac{\sum_{T: |V(S) \cap V(T)| \geq 2} \mathbb{E}[X_T | X_S = 1]}{\mathbb{E}[X]}.$$

We shall estimate $\mathbb{E}[X_T | X_S = 1]$, which is the probability that H is induced in G provided S is induced in G , as a function of $k := |V(S) \cap V(T)|$. We assume that S and T coincide on $V(S) \cap V(T)$, and therefore have at most $dk/2$ edges in common, as their intersection is isomorphic to an induced subgraph of H , and therefore have average degree at most d by assumption. Hence as $p \leq 1/2$

$$\mathbb{E}[X_T | X_S = 1] = \mathbb{P} \left[T \subseteq^{\text{ind}} G \mid S \subseteq^{\text{ind}} G \right] \leq p^{e - \frac{d}{2}k} (1-p)^{\binom{v}{2} - e - \binom{k}{2} + \frac{d}{2}k}.$$

This is clearly also true if S and T do not coincide on $V(S) \cap V(T)$, as then the probability is 0. Now

we can continue our estimation by summing up for all possible T with $k \geq 2$:

$$\begin{aligned} \frac{\sum_T \mathbb{E}[X_T | X_S = 1]}{\mathbb{E}[X]} &\leq \frac{\sum_{k=2}^v \binom{v}{k} \binom{n-v}{v-k} \frac{v!}{|\text{Aut } H|} p^{e - \frac{d}{2}k} (1-p)^{\binom{v}{2} - e - \binom{k}{2} + \frac{d}{2}k}}{\binom{n}{v} \frac{v!}{|\text{Aut } H|} p^e (1-p)^{\binom{v}{2} - e}} \\ &= \sum_{k=2}^v \frac{\binom{v}{k} \binom{n-v}{v-k}}{\binom{n}{v}} p^{-\frac{d}{2}k} \left(\underbrace{(1-p)^{\frac{d+1-k}{2}}}_{\leq (1-p)^{-\frac{d}{2}}} \right)^k \leq \sum_{k=2}^v \frac{v^k}{2(k-2)!} \left(\frac{v}{n-v} \right)^k \left(p^{-\frac{d}{2}} (1-p)^{-\frac{d}{2}} \right)^k \\ &= \frac{1}{2} g^2 \sum_{k=2}^v \frac{1}{(k-2)!} g^{k-2} \leq \frac{1}{2} g^2 \exp(g), \end{aligned}$$

as

$$\frac{\binom{v}{k} \binom{n-v}{v-k}}{\binom{n}{v}} \leq \frac{\binom{v}{k} \frac{(n-v)^{v-k}}{(v-k)!}}{\frac{(n-v)^v}{v!}} = \binom{v}{k}^2 \frac{k!}{(n-v)^k} \leq \frac{1}{k!} \left(\frac{v}{n-v} \right)^k.$$

The lemma follows: the probability of H not being an induced subgraph is at most $e^g g^2/2$. This upper bound is 1 exactly if $g = 2W(1/\sqrt{2})$. For $g \leq 2W(1/\sqrt{2})$, we obtain the upper bound in the lemma. For $g \geq 2W(1/\sqrt{2})$, the upper bound in the lemma is at least 1, so the statement is obvious. \square

4.2 High formulation complexity with high probability

In order to obtain lower bounds on the formulation complexity of the maximum stable set problem of $G = G(n, p)$, via Lemmas 4.3 and 4.2, taking H to be K_t^\sim . We obtain the following result:

Main Theorem 4.4 (Super-polynomial xc of $\text{STAB}(G(n, p))$ w.h.p.). *With high probability, the maximum stable set problem of the random graph $G(n, p)$ has at least the following formulation complexity, depending on the size of p :*

(i) For $p = \omega(1/\sqrt[4]{n})$ and fixed $0 < c < 2/\sqrt{3} \approx 1.1547$, we have

$$\mathbb{P} \left[\text{fc}(\text{STAB}(G(n, p))) \geq 2 \sqrt{c \frac{\ln(np^4)}{p} \log(3/2)} \right] = 1 - o(1). \quad (10)$$

(ii) For $c > 0$ and $c/\sqrt[3]{n} \leq p = o(1)$ we have

$$\mathbb{P} \left[\text{fc}(\text{STAB}(G(n, p))) \geq 2 \sqrt{\frac{\log(3/2)}{p \ln(1/p)}} \right] = 1 - O(1/c^6). \quad (11)$$

(iii) Moreover, for any fixed $c > 0$ for all $1/n < p \leq c/\sqrt[3]{n}$ and $0 < \delta < 1$

$$\mathbb{P} \left[\text{fc}(\text{STAB}(G(n, p))) \geq 2^\delta \sqrt{\frac{\sqrt{pn}}{\ln(1/p)} \log(3/2)} \right] \geq 1 - O(\delta^8). \quad (12)$$

As an illustration of Main theorem 4.4, we include concrete lower bounds in special cases of interest.

Corollary 4.5. For every fixed $0 < \varepsilon < 1$, we have

$$\mathbb{P} \left[\text{fc}(\text{STAB}(G(n, n^{-\varepsilon}))) \geq 2^{\sqrt{(1-4\varepsilon)n^\varepsilon \ln n} \log(3/2)} \right] = 1 - o(1) \quad \text{for } \varepsilon < 1/4, \quad (13)$$

$$\mathbb{P} \left[\text{fc}(\text{STAB}(G(n, n^{-\varepsilon}))) \geq 2^{\frac{n^\varepsilon/2}{\sqrt{\varepsilon \ln n}} \log(3/2)} \right] = 1 - o(1) \quad \text{for } \varepsilon < 1/3, \quad (14)$$

$$\mathbb{P} \left[\text{fc}(\text{STAB}(G(n, n^{-\varepsilon}))) \geq 2^{\frac{n^{(1-\varepsilon)/4}}{\ln n} \log(3/2)} \right] = 1 - o(1) \quad \text{for } \varepsilon \geq 1/3. \quad (15)$$

Below the $p = n^{-\varepsilon}$ range, we obtain

$$\mathbb{P} \left[\text{fc}(\text{STAB}(G(n, (\ln^{6+\varepsilon} n)/n))) \geq 2^{\ln^{1+\varepsilon/5} n \cdot \log(3/2)} \right] = 1 - o(1), \quad (16)$$

and (at the other end of the range) for fixed $\delta > 0$,

$$\mathbb{P} \left[\text{fc}(\text{STAB}(G(n, \delta \ln^{-1} n))) \geq n^{\delta^{-1/2} \log(3/2)} \right] = 1 - o(1). \quad (17)$$

Proof. Equations (13) and (17) are special cases of (10). For Equation (13), we choose $p = n^{-\varepsilon}$ and $c = 1$. For Equation (10), we choose $p = \delta \ln^{-1} n$ and $c = 1.1$, a bit larger than 1, then the square root in (10) becomes

$$\sqrt{c \frac{\ln n p^4}{p}} = \sqrt{c \frac{\ln n + 4 \ln(\delta) + 4 \ln \ln^{-1} n}{\delta} \ln n} = c \delta^{-1/2} (1 + o(1)) \ln n > \delta^{-1/2} \ln n,$$

proving the equation.

Equation (14) follows from Equation (11) via $p = n^{-\varepsilon}$. Equations (15) and (16) are special cases of Equation (12). Equation (15) is the case $p = n^{-1+\varepsilon}$ and $\delta = \sqrt{\varepsilon} \ln^{-1} n$. For Equation (16), we choose $p = (\ln^{6+\varepsilon} n)/n$ and $\delta = \ln^{-\varepsilon/20} n$, then the interesting part of the exponent is

$$\delta \sqrt{\frac{\sqrt{pn}}{\ln(1/p)}} = \ln^{-\varepsilon/20} n \sqrt{\frac{\sqrt{\ln^{6+\varepsilon} n}}{\ln n - \ln \ln^{6+\varepsilon} n}} > \ln^{-\varepsilon/20} n \sqrt{\frac{\sqrt{\ln^{6+\varepsilon} n}}{\ln n}} = \ln^{1+\varepsilon/5} n$$

proving the claim. \square

Now we are going to prove the main theorem of Section 4.2.

Proof of Main Theorem 4.4. We apply Lemma 4.3 to the graph $H := K_t^\sim$ together with Lemma 4.2 to obtain:

$$\begin{aligned} \mathbb{P} \left[\text{fc}(\text{STAB}(G(n, p))) \geq 2^{t \log(3/2)} \right] &\geq \mathbb{P} \left[K_t^\sim \stackrel{\text{ind}}{\subseteq} G(n, p) \right] \\ &\geq 1 - c_0 \frac{v^4 p^{-d} (1-p)^{-v}}{(n-v)^2} \\ &\geq 1 - c_0 (1 + o(1)) \frac{v^4 p^{-d} e^{pv}}{n^2} \quad \text{if } v = o(n). \end{aligned}$$

Here v is the number of vertices of H , and every induced subgraph of H should have average degree at most d . We shall estimate the last fraction $v^4 p^{-d} e^{pv}/n^2$, using the d provided by Lemma 4.1. Below we will tacitly assume $t = \omega(1)$, which is w.l.o.g because $\text{fc}(\text{STAB}(G(n, p))) \geq n$ always.

Now we shall substitute various values for p, t, d, ℓ to obtain the equations of the theorem. We will verify $v = o(n)$ and $v^4 p^{-d} e^{pv} / n^2 = o(1)$ to obtain an $1 - o(1)$ lower bound from the last inequality.

For establishing (10), we choose

$$\ell := 0 \qquad t := \left\lceil c \sqrt{\frac{\ln(np^4)}{p}} \right\rceil \qquad d := 4.$$

Note that for $p \geq 1/\sqrt[4]{n}$,

$$v = t + 3 \binom{t}{2} = \left(\frac{3}{2} + o(1)\right) t^2 = \left(\frac{3}{2} + o(1)\right) c^2 \frac{\ln(np^4)}{p} \leq \left(\frac{3}{2} + o(1)\right) c^2 \sqrt[4]{n} \ln n = o(n),$$

and hence

$$\begin{aligned} \frac{v^4 p^{-d} e^{pv}}{n^2} &= \left(\frac{3}{2} + o(1)\right)^4 (pt^2)^4 e^{(3/2+o(1))pt^2 - 2\ln(np^4)} \\ &\leq \left(\frac{3}{2} + o(1)\right)^4 c^8 (\ln(np^4))^4 \exp\left\{\left[\left(\frac{3}{2} + o(1)\right) c^2 - 2\right] \ln(np^4)\right\} = o(1), \end{aligned}$$

as $np^4 = \omega(1)$ by assumption. This finishes the proof of (10).

We turn to (11) and (12). We will choose a positive ℓ to approximately minimize the fraction in terms of the other parameters. To ease computation, let

$$\gamma := \frac{2\ell + 3}{2} > 1.$$

Then the parameters v and d look like

$$\begin{aligned} d &= 2 + \frac{4}{2\ell + 3} = 2 + \frac{2}{\gamma}, \\ v &= t + (2\ell + 3) \binom{t}{2} = \gamma t^2 + (1 - \gamma)t < \gamma t^2. \end{aligned}$$

Hence

$$\frac{v^4 p^{-d} e^{pv}}{n^2} < \frac{\gamma^4 t^8 e^{p\gamma t^2 + 2(\ln(1/p))/\gamma}}{p^2 n^2}.$$

The γ minimizing the expression is

$$\frac{\sqrt{4 + 2pt^2 \ln(1/p)} - 2}{pt^2} = \frac{2 \ln(1/p)}{\sqrt{4 + 2pt^2 \ln(1/p)} + 2},$$

but we use an approximation as ℓ needs to be an even integer. Therefore we choose

$$\ell = 2 \left\lceil \frac{\ln(1/p)}{\sqrt{4 + 2pt^2 \ln(1/p)} + 2} - \frac{3}{4} \right\rceil.$$

We will verify later that actually $\ell = \omega(1)$. Hence

$$\gamma = (1 + o(1)) \frac{2 \ln(1/p)}{\sqrt{4 + 2pt^2 \ln(1/p)} + 2} = (1 + o(1)) \frac{\sqrt{4 + 2pt^2 \ln(1/p)} - 2}{pt^2},$$

and

$$\begin{aligned} \frac{v^4 p^{-d} e^{pv}}{n^2} &< (1 + o(1)) \left(\frac{2pt^2 \ln(1/p)}{\sqrt{np^3}} \right)^4 \frac{e^{(2+o(1))\sqrt{4+2pt^2 \ln(1/p)}}}{\left(\sqrt{4+2pt^2 \ln(1/p)} + 2 \right)^4} \\ &= (1 + o(1)) \frac{e^{(2+o(1))\sqrt{4+2pt^2 \ln(1/p)}} \left(\sqrt{4+2pt^2 \ln(1/p)} - 2 \right)^4}{(np^3)^2}. \end{aligned} \quad (18)$$

Now we shall substitute various values for p and t to obtain the equations of the theorem.

We will need to verify $\ell = \omega(1)$ and $v = o(n)$ for every choice.

For Equation (12), i.e., in the case $1/n < p \leq c/\sqrt[3]{n}$, we neglect the exponential term in (18) for the choice of t :

$$t = \left\lceil \delta \sqrt{\frac{\sqrt{pn}}{\ln(1/p)}} \right\rceil.$$

Here $0 < \delta < 1$ is an additional parameter. Rearranging gives us

$$2pt^2 \ln(1/p) = (1 + o(1))\delta^2 \sqrt{np^3} \leq (1 + o(1))\delta^2 c^{3/2} \leq (1 + o(1))c^{3/2},$$

so in particular,

$$\begin{aligned} \ell &\geq 2 \left\lceil \frac{\ln(\sqrt[3]{n}/c)}{\sqrt{4 + (1 + o(1))c^{3/2}} + 2} - \frac{3}{4} \right\rceil = \omega(1) \\ v &< \gamma t^2 = O(1/p) = O(\sqrt[3]{n}) = o(n). \end{aligned}$$

Finally,

$$\begin{aligned} \frac{v^4 p^{-d} e^{pv}}{n^2} &< (1 + o(1)) \frac{e^{(2+o(1))\sqrt{4+(1+o(1))c^{3/2}}} \left(\sqrt{4 + (1 + o(1))\delta^2 \sqrt{np^3}} - 2 \right)^4}{(np^3)^2} \\ &\leq (1 + o(1)) e^{(2+o(1))\sqrt{4+(1+o(1))c^{3/2}}} ((1/4 + o(1))\delta^2)^4 = O(\delta^8), \end{aligned}$$

as claimed.

For Equation (11), i.e., when $c/\sqrt[3]{n} \leq p = o(1)$, we choose

$$t = \left\lceil \frac{1}{\sqrt{p \ln(1/p)}} \right\rceil.$$

This provides the estimate

$$2pt^2 \ln(1/p) = 2 + o(1),$$

hence $\ell = \Theta(\ln(1/p)) = \omega(1)$, and $v < \gamma t^2 = O(1/p) = O(\sqrt[3]{n}) = o(n)$. Finally,

$$\begin{aligned} \frac{v^4 p^{-d} e^{pv}}{n^2} &= (1 + o(1)) \frac{e^{(2+o(1))\sqrt{4+(2+o(1))}} \left(\sqrt{4 + (2 + o(1))} - 2 \right)^4}{(np^3)^2} \\ &= O\left(\frac{1}{(np^3)^2}\right) = O(1/c^6), \end{aligned}$$

as $np^3 \geq c^3$. □

Main Theorem 4.4 gives super-polynomial lower bounds all the way from $p = \Omega(\frac{\log^{6+\varepsilon} n}{n})$ to $p = O(\frac{1}{\log n})$. The key for being able to cover the whole regime is to have the gadgets depend on the parameter choice. Notice that for $p < 1/n$ a random graph almost surely will have all its components of size $O(\log n)$, making the stable set problem easy to solve, so that we essentially leave only a small polylog gap.

4.3 Upper bound on formulation complexity with high probability

We now complement Main Theorem 4.4 with an upper bound, which is close to the lower bound, up to an essentially quadratic gap in the exponent.

Theorem 4.6 (Upper bound on the xc of $\text{STAB}(G(n, p))$ w.h.p.). *For $0 < p \leq 1/2$,*

$$\mathbb{P} \left[\text{fc}(\text{STAB}(G)) \geq 2^{\Omega(\frac{\ln^2 n}{p})} \right] \leq n^{-\Omega(\frac{\ln n}{p})}.$$

In particular, for $p = n^{-\varepsilon}$, we obtain $\mathbb{P} \left[\text{fc}(\text{STAB}(G)) \geq 2^{\Omega(n^\varepsilon \ln^2 n)} \right] = o(1)$ and similarly for $p = \delta \ln^{-1} n$, we get $\mathbb{P} \left[\text{fc}(\text{STAB}(G)) \geq n^{\Omega(\frac{\ln^3 n}{\delta})} \right] = o(1)$.

The upper bound stated in Theorem 4.6 is actually an upper bound on the number of stable sets in G , i.e., follows from (6).

Proof of Theorem 4.6. By standard arguments (see, e.g., [Diestel, 2005, Chapter 11, page 300]), for $G = G(n, p)$ we have

$$\mathbb{P} [\alpha(G) \geq r] \leq \left(n e^{-p(r-1)/2} \right)^r$$

and thus for $r = 4\frac{\ln n}{p}$ we get

$$\mathbb{P} \left[\alpha(G) \geq 4\frac{\ln n}{p} \right] \leq \left(\frac{n}{\sqrt{e}} \right)^{-4\frac{\ln n}{p}}.$$

Therefore, with very high probability, we have $\alpha(G) \leq 4\frac{\ln n}{p}$. Using the inequality $\sum_{i=0}^k \binom{n}{i} \leq (n+1)^k$, we get

$$\#(\text{stable sets in } G) \leq (n+1)^{\alpha(G)} = 2^{\log(n+1)\alpha(G)} = 2^{(\frac{1}{\ln 2} + o(1)) \ln(n)\alpha(G)}.$$

The result then follows directly from (6). □

5 Concluding remarks

We conclude with the following conjecture whose validity, we believe, is necessary to strengthen the result, close the remaining gap, as well as establishing truly exponential lower bounds on the extension complexity of further combinatorial problems.

Conjecture 5.1 (Sparse Graph Conjecture). *There exists an infinite family $(T_k)_{k \in \mathbb{N}}$ of template graphs such that, denoting by t_k the number of vertices of T_k : (i) $\text{fc}(\text{STAB}(T_k)) = 2^{\Omega(t_k)}$; (ii) T_k has bounded average degree; (iii) $t_k \leq t_{k+1}$ but at the same time $t_{k+1} = O(t_k)$.*

The existence of such a family would have various consequences.

Exact case. Assuming the Sparse Graph Conjecture we would obtain that the extension complexity of polytopes (see, e.g., Fiorini et al. [2012] for definitions) for important combinatorial problems considered in Fiorini et al. [2012], Avis and Tiwary [2013], Pokutta and Van Vyve [2013] including (among others) the stable set polytope, knapsack polytope, and the 3SAT polytope would have truly exponential extension complexity, that is $2^{\Omega(n)}$ extension complexity, where n is the *dimension* of the polytope.

The recent groundbreaking result of Rothvoß [2014] gives $2^{\Omega(n)}$ bounds for the extension complexity of the matching polytope and TSP polytope. These bounds are also tight up to constants, but this time the upper bound does not come from the number of vertices but rather from the number of facets and dynamic programming algorithms, respectively. Notice that the dimension of both polytopes is $d = \Theta(n^2)$, thus the bounds are in fact $2^{\Omega(\sqrt{d})}$.

Average case. As observed above, there is a quadratic gap in the best current lower and upper bounds on the worst-case extension complexity of the stable set polytope: $2^{\Omega(\sqrt{n})}$ versus 2^n respectively. This is reflected in the results we obtain here. Assuming the Sparse Graph Conjecture we could reduce the gap between upper and lower bounds to a logarithmic factor. Moreover, our results could be strengthened to establish super-polynomial lower bounds on the average-case formulation complexity of $\text{STAB}(G(n, p))$ up to constant probability p .

Acknowledgements

Research reported in this paper was partially supported by NSF grant CMMI-1300144.

References

- D. Avis and H. R. Tiwary. On the extension complexity of combinatorial polytopes. *ArXiv e-prints*, Feb. 2013.
- G. Braun and S. Pokutta. Common information and unique disjointness. In *IEEE 54th Annual Symp. on Foundations of Computer Science (FOCS 2013)*, pages 688–697, 2013. doi: 10.1109/FOCS.2013.79. <http://eccc.hpi-web.de/report/2013/056/>.
- G. Braun and S. Pokutta. Analyzing the size of LPs and SDPs without knowing extended formulations. *manuscript*, 2014.
- G. Braun and S. Pokutta. The matching polytope does not admit fully-polynomial size relaxation schemes. *To appear in Processings of SODA / preprint available at* <http://arxiv.org/abs/1403.6710>, 2015.
- G. Braun, S. Fiorini, S. Pokutta, and D. Steurer. Approximation Limits of Linear Programs (Beyond Hierarchies). In *53rd IEEE Symp. on Foundations of Computer Science (FOCS 2012)*, pages 480–489, 2012. ISBN 978-1-4673-4383-1. doi: 10.1109/FOCS.2012.10.
- G. Braun, S. Fiorini, S. Pokutta, and D. Steurer. Approximation Limits of Linear Programs (Beyond Hierarchies). In *To appear in Mathematics of Operations Research*, 2014. ISBN 978-1-4673-4383-1. doi: 10.1109/FOCS.2012.10.
- M. Braverman and A. Moitra. An information complexity approach to extended formulations. In *Proceedings of the 45th annual ACM symposium on Theory of computing*, pages 161–170, 2013.
- S. Chan, J. Lee, P. Raghavendra, and D. Steurer. Approximate constraint satisfaction requires large LP relaxations. In *IEEE 54th Annual Symp. on Foundations of Computer Science (FOCS 2013)*, pages 350–359, 2013. doi: 10.1109/FOCS.2013.45.

- M. Conforti, G. Cornuéjols, and G. Zambelli. Extended formulations in combinatorial optimization. *4OR*, 8:1–48, 2010. doi: 10.1007/s10288-010-0122-z.
- R. Diestel. *Graph Theory*. Springer-Verlag Heidelberg, New York, 2005.
- S. Fiorini, S. Massar, S. Pokutta, H. R. Tiwary, and R. de Wolf. Linear vs. Semidefinite Extended Formulations: Exponential Separation and Strong Lower Bounds. *Proc. STOC 2012*, 2012.
- M. X. Goemans and D. P. Williamson. Improved approximation algorithms for maximum cut and satisfiability problems using semidefinite programming. *J. Assoc. Comput. Mach.*, 42:1115–1145, 1995. doi: 10.1145/227683.227684.
- V. Kaibel. Extended formulations in combinatorial optimization. *Optima*, 85:2–7, 2011.
- V. Kaibel and S. Weltge. A Short Proof that the Extension Complexity of the Correlation Polytope Grows Exponentially. *ArXiv e-prints*, July 2013.
- S. Pokutta and M. Van Vyve. A note on the extension complexity of the knapsack polytope. *Operations Research Letters*, 41:347–350, 2013.
- A. A. Razborov. On the distributional complexity of disjointness. *Theoret. Comput. Sci.*, 106(2): 385–390, 1992.
- T. Rothvoß. Some 0/1 polytopes need exponential size extended formulations, 2011. arXiv:1105.0036.
- T. Rothvoß. The matching polytope has exponential extension complexity. *Proceedings of STOC*, 2014.
- J. Williams. A linear-size zero-one programming model for the minimum spanning tree problem in planar graphs. *Networks*, 39:53–60, 2002.
- R. d. Wolf. Nondeterministic quantum query and communication complexities. *SIAM Journal on Computing*, 32(3):681–699, 2003.
- M. Yannakakis. Expressing combinatorial optimization problems by linear programs (extended abstract). In *Proc. STOC 1988*, pages 223–228, 1988.
- M. Yannakakis. Expressing combinatorial optimization problems by linear programs. *J. Comput. System Sci.*, 43(3):441–466, 1991. doi: 10.1016/0022-0000(91)90024-Y.