

Improving the Minimax Rate of Active Learning

Lijun Zhang Mehrdad Mahdavi Rong Jin
Department of Computer Science and Engineering
Michigan State University
{zhanglij,mahdavim,rongjin}@msu.edu

Abstract

Active learning refers to the learning protocol where the learner is allowed to choose the data from which it learns. Previous studies have shown that, compared with passive learning, active learning is able to reduce the label complexity exponentially if the data are linearly separable or satisfy the Tsybakov noise condition with parameter $\kappa = 1$. In this paper, we propose a novel active learning algorithm based on a convex surrogate loss, with the goal to broaden the cases for which active learning achieves an exponential improvement in label complexity. We make use of a convex loss not only because it reduces the computational cost, but more importantly because it leads to a tight bound for the empirical process (i.e., the difference between the empirical estimation and the expectation) when the current solution is close to the optimal one. Under the assumption that the norm of the optimal classifier to the convex risk is available, our analysis shows that the introduction of the convex surrogate loss yields an exponential reduction in the label complexity even when the parameter κ of the Tsybakov noise is larger than 1. To the best of our knowledge, this is the first work that improves the minimax rate of active learning by utilizing some priori knowledge.

1 Introduction

The goal of active learning is to learn a classifier with a small binary excess risk using a small number of labeled examples [13]. In [10], the authors show that the minimax convergence rate for any active learning algorithm is bounded by $n^{-\frac{\kappa}{2\kappa-2}}$, where n is the number of labeled instances and $\kappa \geq 1$ is used in Tsybakov noise condition [32] to characterize the behavior of $\Pr(Y = 1|X = \mathbf{x})$ in the neighborhood of the decision boundary. This result implies that unless $\kappa = 1$, no active learning algorithm is able to achieve an exponential reduction in the label complexity for general data distributions. In this work, we develop theory and algorithm for active learning that aims to override this impossibility. We show that if $R = \|\mathbf{w}_*\|$ is known apriori, where \mathbf{w}_* minimizes both the binary risk and the convex risk, it is possible to achieve an exponential reduction in label complexity for certain family of distributions even when $\kappa > 1$. We emphasize that our result does not contradict with the minimax rate of active learning proved in [10] because we assume the prior knowledge of $R = \|\mathbf{w}_*\|$ is available to the learner.

Our work is closely related to the previous study of surrogate losses in passive and active learning [26], in which the authors showed that under appropriate conditions, the exponential reduction in label complexity can still be achieved when the binary loss function is replaced with a convex loss in order to improve the computational efficiency in active learning. In this study, we make one step further. We show that besides the computational efficiency, the introduction of

convex loss function could also benefit the convergence rates of active learning if the length of the optimal linear classifier for the convex risk is known a priori. The key idea is to explore the Lipschitz smoothness of the convex loss function and the technique of local Rademacher complexity [7, 27], i.e., the concentration bound gets tighter as the solution is approaching the optimal one. It is the improved concentration bound, due to the use of a convex loss function, that leads to the surprising result that under the favored condition, it is possible to achieve an exponential reduction in label complexity even when $\kappa > 1$.

2 Related Work

According to [21], most active learning algorithms can be classified into two categories: greedy algorithms and mellow algorithms. The greedy active learning algorithms are designed to select the most informative instances for labeling that will result in an approximately even split of the hypothesis space. Instead of trying to find the most informative instances for labeling, the mellow algorithms for active learning, sometimes referred to as selective sampling, solicit labels for instances as long as they satisfy a given criterion which is usually adjusted over iterations. In this work, we will focus on selective sampling for active learning.

Many studies show that active learning can result in exponential reduction of label complexity compared to passive learning when data are linearly separable [12, 15, 18]. More recent studies focus on agnostic active learning where data cannot be perfectly classified by a linear classifier [3–5, 9, 24–26, 28]. In particular, the studies in [2, 6] show that the exponential reduction in label complexity is still possible for the Tsybakov noise condition with parameter $\kappa = 1$, the malicious noise model, and the adversarial noise model. The minimax rate for active learning under the Tsybakov noise condition is proved in [10], which indicates it is impossible to reduce the sample complexity exponentially when $\kappa > 1$. In this study, we try to override the minimax rate for the Tsybakov noise condition by considering the scenario when $R = \|\mathbf{w}_*\|$ is known a priori, where \mathbf{w}_* minimizes both the binary risk and the convex risk.

Various algorithms have been developed for active learning. Several active learning algorithms (e.g. [3, 24]) require maintaining the subset of hypotheses that yield small binary excess risk, which may result in a high computational cost. This limitation was addressed by either making specific assumptions about data distribution [4, 6] or by maintaining only two hypotheses and soliciting the class label for an instance \mathbf{x} when the two hypotheses differ significantly [9, 14]. Several studies extend active learning theory to the online learning setup [11, 15, 16].

An important quantity in the analysis of active learning is disagreement coefficient [23, 24]. It was shown in [28] that the disagreement coefficient is closely related to the capacity function [1], and under the Massart low noise condition, the capacity function can be bounded by a constant. It was further shown in [19], the disagreement coefficient can be bounded by a constant for any smooth function. In [31], the capacity function was used to provide lower bounds for both passive and active learning.

3 Assumptions

We first introduce the notations used in this paper, and discuss the assumptions we made.

3.1 Notations

Let $\mathcal{X} \subseteq \mathbb{R}^d$ be the domain for the input patterns for classification, and $\mathcal{Y} = \{-1, +1\}$ be the binary class assignment. Let \mathcal{P}_{XY} be the joint distribution for input pattern $X \in \mathcal{X}$ and output binary class assignment $Y \in \mathcal{Y}$, and \mathcal{P}_X be the marginal distribution for \mathcal{X} . Let $\phi(z)$ be a convex loss function that is L -Lipschitz continuous. For a prediction function $f(\mathbf{x}) : \mathbb{R}^d \rightarrow \mathbb{R}$, we define the convex risk $\ell_\phi(\cdot)$ and binary risk $\ell_b(\cdot)$ as

$$\begin{aligned}\ell_\phi(f) &= \mathbb{E}_{(\mathbf{x}, y) \sim \mathcal{P}_{XY}}[\phi(yf(\mathbf{x}))], \\ \ell_b(f) &= \mathbb{E}_{(\mathbf{x}, y) \sim \mathcal{P}_{XY}}[\mathbb{1}(yf(\mathbf{x}) \leq 0)],\end{aligned}$$

where $\mathbb{1}(z)$ is an indicator function that outputs 1 when the predicate z is true and zero otherwise.

In this study, we focus on linear classifier like most studies of active learning. To keep the notation simple, we refer to a weight vector and the linear classifier with that weight vector interchangeably. Let \mathbf{w}_* be the optimal linear classifier that minimizes the convex risk $\ell_\phi(\cdot)$, i.e.,

$$\mathbf{w}_* = \operatorname{argmin}_{\mathbf{w} \in \mathbb{R}^d} \ell_\phi(\mathbf{w}).$$

Throughout the paper, we will use $\bar{\mathbf{w}}$ to represent the unit vector that is along the direction of \mathbf{w} , and use $\|\mathbf{w}\|$ to represent the ℓ_2 -norm of vector \mathbf{w} .

3.2 Assumptions

In this following, we discuss the assumptions made about the optimal solution \mathbf{w}_* and the binary risk $\ell_b(\mathbf{w})$ that are crucial to our analysis. They are:

- **Assumption (I):** $R = \|\mathbf{w}_*\|$ is bounded and known apriori to the learner.
- **Assumption (II):** \mathbf{w}_* minimizes both the convex risk $\ell_\phi(\cdot)$ and the binary risk $\ell_b(\cdot)$ over all the measurable functions.
- **Assumption (III):** There exists constants $\ell_-, \ell_+ > 0$ and $\gamma_- \geq \gamma_+ > 0$ such that for any $\mathbf{w} \in \mathbb{R}^d$

$$\ell_- \|\bar{\mathbf{w}} - \bar{\mathbf{w}}_*\|^{\gamma_-} \leq \ell_b(\mathbf{w}) - \ell_b(\mathbf{w}_*) \leq \ell_+ \|\bar{\mathbf{w}} - \bar{\mathbf{w}}_*\|^{\gamma_+}$$

To quantify the noise level, we choose the Tsybakov low noise condition [32], which combined with Assumption (II) leads to the following condition.

$$\Pr_{X \sim \mathcal{P}_X} \left\{ \operatorname{sign}(\mathbf{w}^\top X) \neq \operatorname{sign}(\mathbf{w}_*^\top X) \right\} \leq \mu (\ell_b(\mathbf{w}) - \ell_b(\mathbf{w}_*))^{1/\kappa}.$$

for some constants $\mu > 0$ and $\kappa \geq 1$.

Assumption (I) assumes the prior knowledge of $R = \|\mathbf{w}_*\|$. The knowledge of R may be obtained based on the assumption of the application domain. For instance, in the case of transfer learning [30], when instances in the target domain are subjected to a unknown unitary transformation of instances from a source domain, the length of the linear classifier will be preserved and transferred from the source domain to the target domain, leading to the knowledge of R for the target domain. This type of transfer learning problem appears in computer vision [22], where images of one domain are acquired by applying certain invariant transform to the images from another domain.

The combination of Assumptions (I) and (II) allows us to nicely connect the minimization of the binary loss with the minimization of a convex loss. It is this connection that makes it possible to improve the convergence rate of active learning. Assumption (III) is a key technical assumption to our analysis. It is introduced to ensure that as the binary risk is reduced, the estimated solution is getting closer to the optimal solution, which makes it possible to explore the local Rademacher complexity of minimizing the convex loss for faster convergence rate of active learning. Below we will justify Assumptions (II) and (III).

3.3 Justification for Assumption (II)

Define

$$\eta(\mathbf{x}) = \Pr(Y = 1|X = \mathbf{x}).$$

The optimal prediction function $\tau(\cdot)$ that minimizes $\ell_\phi(\cdot)$ over all measurable function is given by

$$\tau(\mathbf{x}) = \operatorname{argmin}_{z \in \mathbb{R}} \eta(\mathbf{x})\phi(z) + (1 - \eta(\mathbf{x}))\phi(-z).$$

The first part of Assumption (II) (i.e. \mathbf{w}_* minimizes $\ell_\phi(\cdot)$) assumes that $\tau(\cdot)$ is a linear function, which is used in several studies of the convex surrogate loss [26]. This assumption allows us to bound the binary excess risk in terms of convex excess risk [8].

Remark There are some special cases of $\eta(\cdot)$ and $\phi(\cdot)$ such that $\tau(\cdot)$ is certainly a linear function [8].

- Exponential loss $\phi(\alpha) = e^{-\alpha}$, and a logistic model $\eta(\mathbf{x}) = 1/[1 + \exp(-\mathbf{w}^\top \mathbf{x})]$. We have

$$\tau(\mathbf{x}) = \frac{1}{2} \log \left(\frac{\eta(\mathbf{x})}{1 - \eta(\mathbf{x})} \right) = \frac{1}{2} \mathbf{w}^\top \mathbf{x}.$$

- Truncated quadratic loss $\phi(\alpha) = [\max(0, 1 - \alpha)]^2$, and an affine model $\eta(\mathbf{x}) = \mathbf{w}^\top \mathbf{x} + 1/2$. We have

$$\tau(\mathbf{x}) = 2\eta(\mathbf{x}) - 1 = \mathbf{w}^\top \mathbf{x}.$$

The second part of this assumption (i.e. \mathbf{w}_* also minimizes the binary loss $\ell_b(\cdot)$) is a direct consequence of the first one if the convex loss is *classification-calibrated* [8].

Definition 1. A convex loss $\phi(\cdot)$ is *classification-calibrated* if, for any $\eta \neq 1/2$,

$$H^-(\eta) > H(\eta),$$

where

$$H^-(\eta) = \inf_{\alpha: \alpha(2\eta-1) \leq 0} (\eta\phi(\alpha) + (1 - \eta)\phi(-\alpha)), \text{ and } H(\eta) = \inf_{\alpha \in \mathbb{R}} (\eta\phi(\alpha) + (1 - \eta)\phi(-\alpha)).$$

We finally note that $\tau(\mathbf{x})$ only depends on conditional distribution $\eta(\mathbf{x})$ and the convex loss $\phi(\cdot)$, and is independent from the marginal distribution \mathcal{P}_X . Thus, Assumption (II) actually holds for any distribution of X , as long as $\eta(\mathbf{x})$ and $\phi(\cdot)$ remain the same.

3.4 Justification for Assumption (III)

This assumption will be used in our analysis to bound the distance between $\bar{\mathbf{w}}$ and $\bar{\mathbf{w}}_*$ using the difference in their binary risk. We first examine the lower bound in Assumption (III), i.e., $\ell_b(\mathbf{w}) - \ell_b(\mathbf{w}_*) \geq \ell_- \|\bar{\mathbf{w}} - \bar{\mathbf{w}}_*\|^{\gamma_-}$, and then discuss the upper bound $\ell_b(\mathbf{w}) - \ell_b(\mathbf{w}_*) \leq \ell_+ \|\bar{\mathbf{w}} - \bar{\mathbf{w}}_*\|^{\gamma_+}$.

Lower bound in Assumption (III) The following lemma bounds the values for ℓ_- and γ_- when X follows an isotropic log-concave distribution.

Lemma 1. *Assume \mathcal{P}_X is an isotropic log-concave in \mathbb{R}^d and assume that the Tsybakov condition holds with constants $\mu > 0$ and $\kappa \geq 1$. We have $\gamma_- = \kappa$ and $\ell_- \geq \frac{c\mu}{\mu^\kappa}$ for Assumption (III), where c is an universal constant defined in [6, lemma 3].*

Proof. We need the following lemma regarding the isotropic log-concave distribution [6].

Lemma 2. *([6, Lemma 3]) Assume \mathcal{P}_X is an isotropic log-concave in \mathbb{R}^d . Then, there exists constant $c > 0$ such that for any two unit vectors \mathbf{u} and \mathbf{v} we have*

$$c\theta(\mathbf{u}, \mathbf{v}) \leq \Pr_{X \sim \mathcal{P}_X} \left\{ \text{sign}(\mathbf{u}^\top X) \neq \text{sign}(\mathbf{v}^\top X) \right\},$$

where $\theta(\mathbf{u}, \mathbf{v})$ is the angle between \mathbf{u} and \mathbf{v} .

Using the fact $\sin(x) \leq x$ and the above lemma, we have

$$\begin{aligned} \|\bar{\mathbf{w}} - \bar{\mathbf{w}}_*\| &= \sqrt{2 - 2 \cos(\theta(\mathbf{w}, \mathbf{w}_*))} = 2 \sin\left(\frac{1}{2}\theta(\mathbf{w}, \mathbf{w}_*)\right) \\ &\leq \theta(\mathbf{w}, \mathbf{w}_*) \leq \frac{1}{c} \Pr_{X \sim \mathcal{P}_X} \left\{ \text{sign}(\mathbf{w}^\top X) \neq \text{sign}(\mathbf{w}_*^\top X) \right\}. \end{aligned}$$

From the Tsybakov low noise condition, we have

$$\|\bar{\mathbf{w}} - \bar{\mathbf{w}}_*\| \leq \frac{\mu}{c} (\ell_b(\mathbf{w}) - \ell_b(\mathbf{w}_*))^{1/\kappa}.$$

□

Upper bound in Assumption (III) The following lemma justifies the upper bound when \mathcal{P}_X is orthogonally invariant ¹.

Lemma 3. *Suppose \mathcal{P}_X is orthogonally invariant. We have $\gamma_+ = 1$ and $\ell_+ = 1/2$ for Assumption (III).*

Proof. First, we have

$$\ell_b(\mathbf{w}) - \ell_b(\mathbf{w}_*) \leq \Pr_{X \sim \mathcal{P}_X} \left\{ \text{sign}(\mathbf{w}^\top X) \neq \text{sign}(\mathbf{w}_*^\top X) \right\}.$$

Since the sign function is invariant respect to scaling, we have

$$\ell_b(\mathbf{w}) - \ell_b(\mathbf{w}_*) \leq \Pr_{X \sim \mathcal{P}_X} \left\{ \text{sign}(\mathbf{w}^\top X / \|X\|_2) \neq \text{sign}(\mathbf{w}_*^\top X / \|X\|_2) \right\}.$$

¹In the literature, orthogonally invariant is also referred to as isotropic [17], which is different from the definition of isotropic in [6].

Define $Y = X/\|X\|_2$. Because the distribution of X is orthogonally invariant, it is well-known that Y follows the uniform distribution on the n -dimensional sphere, which is denoted by σ_n [17]. Following [20, Lemma 3.2], we have

$$\ell_b(\mathbf{w}) - \ell_b(\mathbf{w}_*) \leq \Pr_{Y \sim \sigma_n} \left\{ \text{sign}(\mathbf{w}^\top Y) \neq \text{sign}(\mathbf{w}_*^\top Y) \right\} = \frac{1}{\pi} \theta(\mathbf{w}, \mathbf{w}_*). \quad (1)$$

From Jordan's inequality, we know $\sin(x) \geq 2x/\pi$ for $x \in [0, \pi/2]$. Thus, we have

$$\|\bar{\mathbf{w}} - \bar{\mathbf{w}}_*\| = 2 \sin \left(\frac{1}{2} \theta(\mathbf{w}, \mathbf{w}_*) \right) \geq \frac{2}{\pi} \theta(\mathbf{w}, \mathbf{w}_*) \stackrel{(1)}{\geq} 2\ell_b(\mathbf{w}) - \ell_b(\mathbf{w}_*).$$

□

A more general discussion for the upper bound in Assumption (III) can be found in the appendix.

4 Algorithm

Our algorithm works as follows. We divide the learning into m epochs. At the k th epoch, we have a hypothesis space $\Omega_k = \{\mathbf{w} \in \mathbb{R}^d : \|\mathbf{w}\| = 1, \|\mathbf{w} - \mathbf{w}_k\| \leq r_k = 2^{-k+2}\}$, where \mathbf{w}_k is a unit vector that is computed from the previous epoch, and r_k specifies the size of the domain. We sequentially scan through the pool of training examples, and request the class label for a training instance only when it interacts with the domain Ω_k . More specifically, we will request the class label for instance \mathbf{x} if

$$\text{sign}(\mathbf{w}^\top \mathbf{x}) \neq \text{sign}(\mathbf{w}_k^\top \mathbf{x}), \exists \mathbf{w} \in \Omega_k. \quad (2)$$

The following theorem simplifies this condition significantly.

Lemma 4. *The condition in (2) is equivalent to*

$$|\bar{\mathbf{x}}^\top \mathbf{w}_k| \leq \begin{cases} r_k \sqrt{1 - r_k^2/4}, & \text{if } r_k \leq 1; \\ 1, & \text{if } r_k = 2. \end{cases} \quad (3a)$$

$$(3b)$$

Here, $\bar{\mathbf{x}}$ is the unit vector along the direction of \mathbf{x} .

We denote by $\mathcal{D}_k = \{(\mathbf{x}_k^t, y_k^t), t = 1, \dots, n_k\}$ the collection of labeled training examples received at epoch k , where n_k is the number of labeled instances at epoch k . Using the training examples in \mathcal{D}_k , we compute a new classifier \mathbf{w}_{k+1} as the solution to

$$\min_{\mathbf{w} \in \Omega_k} \sum_{t=1}^{n_k} \mathbb{1} \left(y_k^t \neq \text{sign}(\mathbf{w}^\top \mathbf{x}_k^t) \right). \quad (4)$$

The new hypothesis space, denoted by Ω_{k+1} , is then updated as

$$\Omega_{k+1} = \left\{ \mathbf{w} \in \mathbb{R}^d : \|\mathbf{w}\| = 1, \|\mathbf{w} - \mathbf{w}_{k+1}\| \leq r_{k+1} = r_k/2 \right\},$$

where the size of the hypothesis space is reduced by half through each epoch. We note that the idea of reducing the hypothesis space by half for each epoch has been used in the margin-based active learning algorithm [4, 6].

Algorithm 1 Active Learning with Faster Convergence Rate

- 1: Set \mathbf{w}_1 as a random unit vector and $r_1 = 2$
 - 2: **for** $k = 1, 2, \dots, m$ **do**
 - 3: Label n_k training instances that satisfy (3a) or (3b).
 - 4: Learn a new classifier $\tilde{\mathbf{w}}_{k+1}$ solving either the non-convex optimization problem in (4) or the convex optimization in (5)
 - 5: Set $\mathbf{w}_{k+1} = \tilde{\mathbf{w}}_{k+1} / \|\tilde{\mathbf{w}}_{k+1}\|$ and $r_{k+1} = r_k/2$.
 - 6: **end for**
- Return** \mathbf{x}_f
-

One problem with the updating procedure given in (4) is that it requires solving a non-convex optimization problem, which could be computationally expensive when the number of training examples is large. To address this limitation, we propose to obtain $\tilde{\mathbf{w}}_{k+1}$ by solving the following convex optimization problem

$$\min_{\mathbf{w} \in \Delta_k} \sum_{t=1}^{n_k} \phi(y_k^t \mathbf{w}^\top \mathbf{x}_k^t), \quad (5)$$

where $\Delta_k = \{\mathbf{w} \in \mathbb{R}^d : \|\mathbf{w} - R\mathbf{w}_k\| \leq Rr_k\}$. Here $R = \|\mathbf{w}_*\|$ comes from the prior knowledge of \mathbf{w}_* due to Assumption (I). The final \mathbf{w}_{k+1} is obtained by normalizing $\tilde{\mathbf{w}}_{k+1}$ to a unit vector. Algorithm 1 summarizes the key steps of both approaches.

5 Analysis

We will first introduce the basic concepts that are commonly used in the analysis of active learning. We will then analyze the label complexity for solving the non-convex optimization problem in (4). The key result of this work is presented in Section 5.3, where we show that the exponential reduction can be achieved even when $\kappa > 1$ if we solve the convex optimization problem in (5) in Algorithm 1. Due to space limitations, most of the technical proofs are provided in the Appendix.

5.1 Basics

Similar to most active learning theories, we assume bounded disagreement coefficient [24]. We define the region of disagreement as, for any subset of hypothesis V ,

$$\text{DIS}(V) = \left\{ \mathbf{x} \in \mathcal{X} : \exists \mathbf{w}_1, \mathbf{w}_2 \in V \text{ s. t. } \text{sign}(\mathbf{w}_1^\top \mathbf{x}) \neq \text{sign}(\mathbf{w}_2^\top \mathbf{x}) \right\}.$$

For $r \in [0, 1]$, let

$$B(\mathbf{w}, r) = \left\{ \mathbf{w}' \in \mathcal{B} : \Pr \left\{ \text{sign}(\mathbf{w}^\top X) \neq \text{sign}([\mathbf{w}']^\top X) \right\} \leq r \right\},$$

where

$$\mathcal{B} = \{\mathbf{w} \in \mathbb{R}^d : \|\mathbf{w}\| = 1\}.$$

The disagreement coefficient of \mathbf{w} with respect to $B(\mathbf{w}, r)$ is then defined as

$$\theta_{\mathbf{w}}(\varepsilon) = \sup_{r \geq \varepsilon} \frac{\Pr(\text{DIS}(B(\mathbf{w}, r)))}{r}.$$

Define

$$\theta(\varepsilon) = \theta_{\mathbf{w}_*}(\varepsilon).$$

Note that we keep the dependence of ε in the definition of disagreement coefficient since it may include factor $\log(1/\varepsilon)$ as indicated in [6]. The disagreement coefficient allows us to connect the largest disagreement between two classifiers in a given hypothesis space with the percentage of the examples that are classified differently by at least two classifiers in the hypothesis space.

Since our work tries to bound the binary excess risk with a convex excess risk, we need the ψ -transform [8] stated below,

$$\psi(z) = \inf_{\alpha z \leq 0} \left(\frac{1+z}{2} \phi(\alpha) + \frac{1-z}{2} \phi(-\alpha) \right) - \inf_{\alpha \in \mathbb{R}} \left(\frac{1+z}{2} \phi(\alpha) + \frac{1-z}{2} \phi(-\alpha) \right).$$

Here are two examples of ψ -transform from [8]: (i) for exponential loss $\phi(\alpha) = e^{-\alpha}$, $\psi(z) = 1 - \sqrt{1-z^2} \geq z^2/2$, and (ii) for truncated quadratic loss $\phi(\alpha) = [\max(0, 1-\alpha)]^2$, $\phi(z) = z^2$.

The following theorem from [8, Theorem 1] shows that the binary excess risk can be bounded by the convex excess risk using ψ -transform.

Theorem 1. *For any non-negative loss function ϕ , any measurable function $f : \mathcal{X} \mapsto \mathbb{R}$, and any probability distribution on $\mathcal{X} \times \{-1, +1\}$, we have*

$$\psi(R(f) - R^*) \leq R_\phi(f) - R_\phi^*$$

where $R(f) = \mathbb{E}_{(X,Y) \sim \mathcal{P}_{XY}} [\mathbb{1}(yf(X) \leq 0)]$, $R^* = \min_f R(f)$, $R_\phi(f) = \mathbb{E}_{(X,Y) \sim \mathcal{P}_{XY}} [\phi(Yf(X))]$, and $R_\phi^* = \min_f R_\phi(f)$. Here the minimization is taken over all measurable functions.

5.2 Label Complexity for Non-Convex Optimization

Our analysis is based on induction. The key to our analysis is to show that $\bar{\mathbf{w}}_* \in \Omega_{k+1}$, if (i) $\bar{\mathbf{w}}_* \in \Omega_k$ and (ii) n_k , the number of labeling queries issued at epoch k , is sufficiently large.

Theorem 2. *Suppose Assumption (III) holds, and $\bar{\mathbf{w}}_* \in \Omega_k$. Then, with a probability $1 - \delta/m$, we have $\bar{\mathbf{w}}_* \in \Omega_{k+1}$, if*

$$n_k = 2c^2 \theta^2(\varepsilon) \left[\log \frac{4m}{\delta} + 2(d+1) \left(\log 8 + 2 \log \frac{c\theta(\varepsilon)}{r_k^{\gamma_- - \gamma_+ / \kappa}} \right) \right] r_k^{2(\frac{\gamma_+}{\kappa} - \gamma_-)}, \quad (6)$$

where $c = \mu \ell_+^{1/\kappa} 2^{\gamma_0} / \ell_-$ and $\gamma_0 = 2 + \gamma_- + \gamma_+ / \kappa$.

Theorem 3. *Suppose Assumption (III) holds. Let \mathbf{w}_{m+1} be the solution output from the proposed algorithm after m iterations, where $m = \lceil \log_2(2/\varepsilon) \rceil$. Then, with a probability $1 - \delta$, we have*

$$\|\mathbf{w}_{m+1} - \bar{\mathbf{w}}_*\| \leq \varepsilon,$$

and the total number of labeled instances is bounded by

$$n \leq \begin{cases} n_0 \frac{2^{2\alpha}}{2^{2\alpha} - 1} \left(\frac{4}{\varepsilon}\right)^{2\alpha}, & \alpha > 0 \\ n_0 \log_2 \frac{4}{\varepsilon}, & \alpha \leq 0 \end{cases}$$

where

$$\alpha = \gamma_- - \frac{\gamma_+}{\kappa},$$

$$n_0 = 2^{1-4\alpha} c^2 \theta^2(\varepsilon) \left[\log \frac{4m}{\delta} + 2(d+1) (\log 8 + 2 \log c\theta(\varepsilon) + 2m\alpha \log 2) \right].$$

Proof. From Theorem 2, with a probability $1 - \delta$, we have

$$\|\mathbf{w}_{m+1} - \bar{\mathbf{w}}_*\| \leq r_{m+1} = r_1 2^{-m} = 2^{-m+1} \leq \varepsilon.$$

and the number of labeled instances is bounded by

$$\begin{aligned} n &= \sum_{k=1}^m n_k \\ &= \sum_{k=1}^m 2^{1-4\alpha} c^2 \theta^2(\varepsilon) \left[\log \frac{4m}{\delta} + 2(d+1) (\log 8 + 2 \log c\theta(\varepsilon) + 2(k-2)\alpha \log 2) \right] 2^{2\alpha k} \\ &\leq n_0 \sum_{k=1}^m 2^{2\alpha k}. \end{aligned}$$

In the case that $\alpha \leq 0$, we have

$$n \leq n_0 m \leq n_0 \log_2 \frac{4}{\varepsilon}.$$

Otherwise, we have

$$n \leq n_0 \frac{2^{2\alpha}(2^{2\alpha m} - 1)}{2^{2\alpha} - 1} \leq n_0 \frac{2^{2\alpha}}{2^{2\alpha} - 1} 2^{2\alpha m} \leq n_0 \frac{2^{2\alpha}}{2^{2\alpha} - 1} \left(\frac{4}{\varepsilon}\right)^{2\alpha}.$$

□

Remark When the distribution \mathcal{P}_X is isotropic log-concave and orthogonally invariant, we have $\gamma_- = \kappa$ and $\gamma_+ = 1$, as discussed in Lemmas 1 and 3. In the case when $\kappa = 1$, $\alpha = 0$ and Algorithm 1 achieves exponential reduction in label complexity, consistent with the previous studies. In the next subsection, we show that it is possible to achieve exponential reduction in label complexity even $\kappa > 1$ provided Assumptions (I)-(III) hold.

5.3 Label Complexity for Convex Optimization

For the simplicity of the presentation, we first assume a bounded ℓ_2 -norm for the input X , i.e., $\|X\| \leq 1$, and discuss a relaxed condition in the end of this section.

5.3.1 A Special Case with Bounded ℓ_2 -norm

First, we give a concentration result bounding the empirical process.

Theorem 4. Assume $\phi(\cdot)$ to be Lipschitz continuous with constant L , and $\|X\| \leq 1$. Let $(\mathbf{x}_i, \mathbf{y}_i)_{i=1}^n$ be a set of i.i.d. samples drawn from an unknown distribution \mathcal{P}_{XY} . Then with probability at least $1 - \delta$, for every \mathbf{w} with $\|\mathbf{w} - \mathbf{w}_*\| \leq r$, we have

$$\ell_\phi(\mathbf{w}) - \ell_\phi(\mathbf{w}_*) - \frac{1}{n} \sum_{i=1}^n \left(\phi\left(y_i \mathbf{w}^\top \mathbf{x}_i\right) - \phi\left(y_i \mathbf{w}_*^\top \mathbf{x}_i\right) \right) \leq \frac{Lr}{\sqrt{n}} \left(4 + \sqrt{2 \log \frac{m}{\delta}} \right). \quad (7)$$

We observe the upper bound in (7) depends on the radius r of the solution space, and thus the upper bound becomes tighter as the solution is approaching to \mathbf{w}_* , a key idea used in the local Rademacher complexity [7, 27]. Notice that in the case of binary loss, we do not have this nice property.

Similar to Theorem 2, we have the following theorem bounding the number of label requests in each iterations.

Theorem 5. Suppose Assumptions (I)-(III) hold, and $\phi(\cdot)$ is L -Lipschitz continuous. Assume $\psi(z) \geq az^\gamma$ for any $z \in (0, 1)$ and $\bar{\mathbf{w}}_* \in \Omega_k$. Then, with a probability $1 - \delta/m$, we have $\bar{\mathbf{w}}_* \in \Omega_{k+1}$, if

$$n_k = \left(\frac{\mu \ell_+^{1/\kappa} 2^{\gamma_- + \gamma_+/\kappa} \theta(\varepsilon)}{\ell_-} \right)^{2\gamma} \left(\frac{2LR}{a} \left(4 + \sqrt{2 \log \frac{m}{\delta}} \right) \right)^2 r_k^{2(1 + \frac{\gamma_+}{\kappa} - \gamma\gamma_-)}. \quad (8)$$

Note that the assumption $\psi(z) \geq az^\gamma$ is met by the two examples of ψ -transform given in Section 3.4.

Following the same analysis as Theorem 3, we have the following results for the convex case.

Theorem 6. Suppose Assumption (I)-(IV) hold. Assume $\psi(z) \geq az^\gamma$ for any $z \in (0, 1)$. Let \mathbf{w}_{m+1} be the solution output from the proposed algorithm after m iterations, where $m = \lceil \log_2(2/\varepsilon) \rceil$. Then, with a probability $1 - \delta$, we have

$$\|\mathbf{w}_{m+1} - \bar{\mathbf{w}}_*\| \leq \varepsilon,$$

and the total number of labeled instances is bounded by

$$n \leq \begin{cases} n_0 \frac{2^{2\alpha}}{2^{2\alpha} - 1} \left(\frac{4}{\varepsilon}\right)^{2\alpha}, & \alpha > 0 \\ n_0 \log_2 \frac{4}{\varepsilon}, & \alpha \leq 0 \end{cases}$$

where

$$\alpha = \gamma\gamma_- - \frac{\gamma\gamma_+}{\kappa} - 1, \\ n_0 = 2^{-4\alpha} \left(\frac{\mu \ell_+^{1/\kappa} 2^{\gamma_- + \gamma_+/\kappa} \theta(\varepsilon)}{\ell_-} \right)^{2\gamma} \left(\frac{2LR}{a} \left(4 + \sqrt{2 \log \frac{m}{\delta}} \right) \right)^2.$$

Remark As indicated by Theorem 6, an exponential reduction in label complexity can be achieved if $\alpha \leq 0$. More specifically, the number of labeled instances requested by Algorithm 1 is

$$O\left((\theta(\varepsilon))^{2\gamma} \log \frac{1}{\varepsilon} \right).$$

Since $\alpha \leq 0$ implies

$$\kappa \leq \frac{\gamma\gamma_+}{\gamma\gamma_- - 1},$$

we have an exponential reduction in label complexity even when $\kappa > 1$ provided the above inequality holds. To be more concrete, consider the case when \mathcal{P}_X is orthogonally invariant, we have $\gamma_- = \kappa$ and $\gamma_+ = 1$, as discussed in Lemmas 1 and 3, and therefore

$$\kappa \leq \kappa_0 := \frac{1 + \sqrt{1 + 4\gamma^2}}{2\gamma}$$

will ensure $\alpha \leq 0$ and consequentially an exponential reduction in label complexity. We emphasize that our result does not contradict with the minimax rate of active learning proved in [10] because we assume the prior knowledge of $R = \|\mathbf{w}_*\|$ is available to the learner.

5.3.2 A General Case

In the following, we study a more general assumption that the ℓ_2 -norm of X is a sub-exponential or a sub-gaussian random variable, i.e.,

$$\| \|X\| \|_{\psi_1} \leq 1 \text{ or } \| \|X\| \|_{\psi_2} \leq 1.$$

For $\alpha > 0$, the ψ_α -norm of a random variable η , which is a special Orlicz norm [29, Section A.1], is defined as follows

$$\|\eta\|_{\psi_\alpha} = \inf \left\{ C > 0 : \mathbb{E} \left[\exp \left(\left(\frac{|\eta|}{C} \right)^\alpha \right) \right] \leq 2 \right\}.$$

The analysis for this case is almost the same as the previous one, except that we need a generalized version of Theorem 4.

Theorem 7. *Assume $\phi(\cdot)$ is Lipschitz continuous with constant L , and the marginal distribution P_X ensures $\| \|X\| \|_{\psi_1} \leq 1$ or $\| \|X\| \|_{\psi_2} \leq 1$. Let $(\mathbf{x}_i, \mathbf{y}_i)_{i=1}^n$ be a set of i.i.d. samples drawn from an unknown distribution P_{XY} . Then with probability at least $1 - \delta$, for every \mathbf{w} with $\|\mathbf{w} - \mathbf{w}_*\| \leq r$, we have*

$$\ell_\phi(\mathbf{w}) - \ell_\phi(\mathbf{w}_*) - \frac{1}{n} \sum_{i=1}^n \left(\phi(y_i \mathbf{w}^\top \mathbf{x}_i) - \phi(y_i \mathbf{w}_*^\top \mathbf{x}_i) \right) \leq \frac{CLr}{\sqrt{n}} \left(1 + \sqrt{\log \frac{m}{\delta}} \right). \quad (9)$$

for some constant C , provided that

$$n \geq 1 + 2 \log \frac{m}{\delta} \cdot \log \left(\frac{2}{e} \log \frac{m}{\delta} \right). \quad (10)$$

Comparing (7) and (9), we can see the bounds in Theorems 4 and 7 differ only by a constant factor, provided the condition in (10) holds. Thus, we just need to make the following modifications to Theorems 5 and 6: (i) replacing the factor $4 + \sqrt{2 \log \frac{m}{\delta}}$ with $C(1 + \sqrt{\log \frac{m}{\delta}})$, and (ii) adding constraints to ensure (10) is true. Notice that the condition in (10) only requires the number of labeled instances in each iteration is on the order of $\Omega(\log \log \frac{1}{\epsilon})$, and thus the total number of labeled instances is on the order of $\Omega(\log \frac{1}{\epsilon} \cdot \log \log \frac{1}{\epsilon})$. So, it does not affect the order of the sample complexity shown in Theorem 7.

6 Conclusion

In this paper, we study the active learning problem with a convex surrogate loss. Our results show that with some additional assumptions, the convex surrogate loss not only improves the computational efficiency of active learning, but also reduces the sample complexity. In particular, our analysis reveals that it is possible to achieve an exponential reduction in the label complexity even when the noisy level is high.

References

- [1] Kenneth S. Alexander. Rates of growth and sample moduli for weighted empirical processes indexed by sets. *Probability Theory and Related Fields*, 75(3):379–423, 1987.
- [2] Pranjal Awasthi, Maria Florina Balcan, and Philip M. Long. The power of localization for efficiently learning linear separators with noise. *ArXiv e-prints*, arXiv:1307.8371, 2013.
- [3] Maria-Florina Balcan, Alina Beygelzimer, and John Langford. Agnostic active learning. In *Proceedings of the 23rd international conference on Machine learning*, pages 65–72, 2006.
- [4] Maria-Florina Balcan, Andrei Broder, and Tong Zhang. Margin based active learning. In *Proceedings of the 20th Annual Conference on Learning Theory*, pages 35–50, 2007.
- [5] Maria-Florina Balcan, Steve Hanneke, and Jennifer Wortman Vaughan. The true sample complexity of active learning. *Machine Learning*, 80(2-3):111–139, 2010.
- [6] Maria Florina Balcan and Philip M. Long. Active and passive learning of linear separators under log-concave distributions. In *Proceedings of the 26th Annual Conference on Learning Theory*, pages 288–316, 2013.
- [7] Peter L. Bartlett, Olivier Bousquet, and Shahar Mendelson. Local rademacher complexities. *The Annals of Statistics*, 33(4):1497–1537, 2005.
- [8] Peter L. Bartlett, Michael I. Jordan, and Jon D. McAuliffe. Convexity, classification, and risk bounds. *Journal of the American Statistical Association*, 101(473):138–156, 2006.
- [9] Alina Beygelzimer, Daniel Hsu, John Langford, and Tong Zhang. Agnostic active learning without constraints. In *Advances in Neural Information Processing Systems 23*, pages 199–207, 2010.
- [10] Rui M. Castro and Robert D. Nowak. Minimax bounds for active learning. In *Proceedings of the 20th annual conference on Learning theory*, pages 5–19, 2007.
- [11] Giovanni Cavallanti, Nicolò Cesa-Bianchi, and Claudio Gentile. Learning noisy linear classifiers via adaptive and selective sampling. *Machine Learning*, 83(1):71–102, 2011.
- [12] Sanjoy Dasgupta. Coarse sample complexity bounds for active learning. In *Advances in Neural Information Processing Systems 18*, pages 235–242, 2006.
- [13] Sanjoy Dasgupta. Active learning theory. *Encyclopedia of Machine Learning*, pages 14–19, 2011.

- [14] Sanjoy Dasgupta, Daniel Hsu, and Claire Monteleoni. A general agnostic active learning algorithm. In *Advances in Neural Information Processing Systems 20*, pages 353–360, 2008.
- [15] Sanjoy Dasgupta, Adam Tauman Kalai, and Claire Monteleoni. Analysis of perceptron-based active learning. *Journal of Machine Learning Research*, 10:281–299, 2009.
- [16] Ofer Dekel, Claudio Gentile, and Karthik Sridharan. Selective sampling and active learning from single and multiple teachers. *Journal of Machine Learning Research*, 13:2655–2697, 2012.
- [17] Morris L. Eaton. On the projections of isotropic distributions. *The Annals of Statistics*, 9(2):391–400, 1981.
- [18] Yoav Freund, H. Sebastian Seung, Eli Shamir, and Naftali Tishby. Selective sampling using the query by committee algorithm. *Machine Learning*, 28(2-3):133–168, 1997.
- [19] Eric J. Friedman. Active learning for smooth problems. In *Proceedings of the 22nd Annual Conference on Learning Theory*, 2009.
- [20] Michel X. Goemans and David P. Williamson. Improved approximation algorithms for maximum cut and satisfiability problems using semidefinite programming. *Journal of the ACM*, 42(6):1115–1145, 1995.
- [21] Alon Gonen, Sivan Sabato, and Shai Shalev-Shwartz. Efficient active learning of halfspaces: An aggressive approach. *Journal of Machine Learning Research*, 14:2583–2615, 2013.
- [22] Thore Graepel and Ralf Herbrich. Invariant pattern recognition by semidefinite programming machines. In *Advances in Neural Information Processing Systems 16*, pages 33–40, 2004.
- [23] Steve Hanneke. A bound on the label complexity of agnostic active learning. In *Proceedings of the 24th International Conference on Machine Learning*, pages 353–360, 2007.
- [24] Steve Hanneke. Rates of convergence in active learning. *The Annals of Statistics*, 39(1):333–361, 2011.
- [25] Steve Hanneke and Liu Yang. Negative results for active learning with convex losses. *Proceedings of the 13th International Conference on Artificial Intelligence and Statistics*, pages 321–325, 2010.
- [26] Steve Hanneke and Liu Yang. Surrogate losses in passive and active learning. *ArXiv e-prints*, arXiv:1207.3772, 2012.
- [27] Vladimir Koltchinskii. Local rademacher complexities and oracle inequalities in risk minimization. *The Annals of Statistics*, 34(6):2593–2656, 2006.
- [28] Vladimir Koltchinskii. Rademacher complexities and bounding the excess risk in active learning. *Journal of Machine Learning Research*, 11:2457–2485, 2010.
- [29] Vladimir Koltchinskii. *Oracle Inequalities in Empirical Risk Minimization and Sparse Recovery Problems*. Springer, 2011.
- [30] Sinno Jialin Pan and Qiang Yang. A survey on transfer learning. *IEEE Transactions on Knowledge and Data Engineering*, 22(10):1345–1359, 2010.

- [31] Maxim Raginsky and Alexander Rakhlin. Lower bounds for passive and active learning. In *Advances in Neural Information Processing Systems 24*, pages 1026–1034, 2011.
- [32] Alexandre B. Tsybakov. Optimal aggregation of classifiers in statistical learning. *The Annals of Statistics*, 32:135–166, 2004.
- [33] Aad W. van der Vaart and Jon A. Wellner. *Weak Convergence and Empirical Processes*. Springer Series in Statistics. Springer, 1996.
- [34] Vladimir N. Vapnik. *Statistical Learning Theory*. Wiley-Interscience, 1998.
- [35] Roman Vershynin. Introduction to the non-asymptotic analysis of random matrices. In *Compressed Sensing, Theory and Applications*, chapter 5, pages 210–268. Cambridge University Press, 2012.

A More Discussion about the Upper Bound in Assumption (III)

We can have a more general result by exploiting the relationship between the binary excess risk $\ell_b(\mathbf{w}) - \ell_b(\mathbf{w}_*)$ and the convex excess risk $\ell_\phi(\mathbf{w}) - \ell_\phi(\mathbf{w}_*)$. Using Theorem 1, we have the following result for ℓ_+ and γ_+ .

Lemma 5. *Assume (i) ϕ is non-negative, (ii) $\ell_\phi(\mathbf{w})$ is L_ϕ -strongly smooth, that is,*

$$\ell_\phi(\mathbf{w}') \leq \ell_\phi(\mathbf{w}) + \langle \nabla \ell_\phi(\mathbf{w}), \mathbf{w}' - \mathbf{w} \rangle + \frac{L_\phi}{2} \|\mathbf{w}' - \mathbf{w}\|^2,$$

and (iii) $\psi(z) \geq az^\gamma$ for any $z \in [0, 1]$. We have

$$\ell_+ = \left(\frac{L_\phi R^2}{2a} \right)^{1/\gamma}, \text{ and } \gamma_+ = \frac{2}{\gamma},$$

where $R = \|\mathbf{w}_*\|$.

Proof. Combining Theorem 1 and Assumption (II), we have

$$\psi(\ell_b(\mathbf{w}) - \ell_b(\mathbf{w}_*)) \leq \ell_\phi(\mathbf{w}) - \ell_\phi(\mathbf{w}_*) \leq \frac{L_\phi}{2} \|\mathbf{w} - \mathbf{w}_*\|^2.$$

Since we can arbitrary scale \mathbf{w} without changing its binary risk $\ell_b(\mathbf{w})$, we have

$$\psi(\ell_b(\mathbf{w}) - \ell_b(\mathbf{w}_*)) = \psi(\ell_b(R\bar{\mathbf{w}}) - \ell_b(\mathbf{w}_*)) \leq \frac{L_\phi R^2}{2} \|\bar{\mathbf{w}} - \bar{\mathbf{w}}_*\|^2.$$

From the assumption $\psi(z) \geq az^\gamma$, we have

$$a(\ell_b(\mathbf{w}) - \ell_b(\mathbf{w}_*))^\gamma \leq \frac{L_\phi R^2}{2} \|\bar{\mathbf{w}} - \bar{\mathbf{w}}_*\|^2,$$

which completes the proof. □

An example of convex loss that satisfies the conditions in Lemma 5 is the truncated quadratic loss $\phi(\alpha) = [\max(0, 1 - \alpha)]^2$, which is 1-strongly smooth with $\psi(z) = z^2$.

B Proof of Lemma 4

We first consider the case $r_k \leq 1$ such that $2 \arcsin(r_k/2) < \frac{\pi}{2}$. Using a simple geometry argument, it is easy to show that (2) is equivalent to

$$\text{sign}(\mathbf{w}^\top \mathbf{x}) \neq \text{sign}(\mathbf{w}_k^\top \mathbf{x}), \exists \mathbf{w} \in \left\{ \mathbf{w} \in \mathbb{R}^d : \theta(\mathbf{w}, \mathbf{w}_k) \leq 2 \arcsin(r_k/2) \right\},$$

which is equivalent to

$$\frac{\pi}{2} - 2 \arcsin(r_k/2) \leq \theta(\mathbf{w}_k, \mathbf{x}) \leq \frac{\pi}{2} + 2 \arcsin(r_k/2),$$

leading to the following condition

$$|\cos(\theta(\mathbf{w}_k, \mathbf{x}))| \leq \sin(2 \arcsin(r_k/2)).$$

Using the fact

$$\sin(\theta) = 2 \sin(\theta/2) \cos(\theta/2) = 2 \sin(\theta/2) \sqrt{1 - \sin^2(\theta/2)},$$

we have

$$|\bar{\mathbf{x}}^\top \mathbf{w}_k| = |\cos(\theta(\mathbf{w}_k, \mathbf{x}))| \leq r_k \sqrt{1 - r_k^2/4}.$$

If $r_k = 2$, it is obvious that both (2) and (3b) become vacuous.

C Proof of Theorem 2

Let Ω_k be the subset of hypothesis obtained in epoch k with center \mathbf{w}_k and radius r_k , i.e.,

$$\Omega_k = \left\{ \mathbf{w} \in \mathbb{R}^d : \|\mathbf{w}\| = 1, \|\mathbf{w} - \mathbf{w}_k\| \leq r_k \right\}.$$

By the induction assumption, we have $\bar{\mathbf{w}}_* \in \Omega_k$, and thus for any $\mathbf{w} \in \Omega_k$

$$\|\mathbf{w} - \bar{\mathbf{w}}_*\| \leq \|\mathbf{w} - \mathbf{w}_k\| + \|\bar{\mathbf{w}}_* - \mathbf{w}_k\| \leq 2r_k.$$

Using the upper bound in Assumption (III), for any $\mathbf{w} \in \Omega_k$, we have

$$\ell_b(\mathbf{w}) - \ell_b(\bar{\mathbf{w}}_*) \leq \ell_+ \|\bar{\mathbf{w}} - \bar{\mathbf{w}}_*\|^{\gamma_+} \leq \ell_+ 2^{\gamma_+} r_k^{\gamma_+}.$$

According to the Tsybakov's low noise condition [32], for any $\mathbf{w} \in \Omega_k$, we have

$$\Pr_{X \sim \mathcal{P}_X} \left\{ \text{sign}(\mathbf{w}^\top X) \neq \text{sign}(\bar{\mathbf{w}}_*^\top X) \right\} \leq \mu (\ell_b(\mathbf{w}) - \ell_b(\bar{\mathbf{w}}_*))^{1/\kappa} \leq \mu \ell_+^{1/\kappa} 2^{\gamma_+/\kappa} r_k^{\gamma_+/\kappa}.$$

As a result, we have

$$\Omega_k \subseteq B(\mathbf{w}, \mu \ell_+^{1/\kappa} 2^{\gamma_+/\kappa} r_k^{\gamma_+/\kappa}) \Rightarrow \text{DIS}(\Omega_k) \subseteq \text{DIS} \left(B \left(\mathbf{w}, \mu \ell_+^{1/\kappa} 2^{\gamma_+/\kappa} r_k^{\gamma_+/\kappa} \right) \right).$$

From the definition of disagreement coefficient, we have

$$\Pr(\text{DIS}(\Omega_k)) \leq \Pr \left(\text{DIS} \left(B \left(\mathbf{w}, \mu \ell_+^{1/\kappa} 2^{\gamma_+/\kappa} r_k^{\gamma_+/\kappa} \right) \right) \right) \leq \mu \ell_+^{1/\kappa} 2^{\gamma_+/\kappa} r_k^{\gamma_+/\kappa} \theta(\varepsilon). \quad (11)$$

Define $\mathcal{C}_k = \mathcal{X} \setminus \text{DIS}(\Omega_k)$. Notice that $\text{DIS}(\Omega_k)$ is the subset of instances for which at least two classifiers from Ω_k will result in different predictions. Since $\mathbf{w}_* \in \Omega_k$, we have for any given classifier $\mathbf{w} \in \Omega_k$, $\text{sign}(\mathbf{w}^\top \mathbf{x}) = \text{sign}(\mathbf{w}_*^\top \mathbf{x})$ for all $\mathbf{x} \in \mathcal{C}_k$. Hence, we have, for any $\mathbf{w} \in \Omega_k$,

$$\ell_b(\mathbf{w}) - \ell_b(\mathbf{w}_*) = (\ell_b(\mathbf{w}|\text{DIS}(\Omega_k)) - \ell_b(\mathbf{w}_*|\text{DIS}(\Omega_k))) \Pr(\text{DIS}(\Omega_k)), \quad (12)$$

where $\ell_b(\mathbf{w}|T)$ is defined as the binary risk for instances sampled from the set T . It is easy to verify that the n_k training instances labeled at epoch k are i.i.d. samples from the space $\text{DIS}(\Omega_k)$. To bound the generalization error, we use the following theorem from [34].

Theorem 8. *For any distribution \mathcal{D} over $\mathcal{X} \times \{\pm 1\}$, with a probability at least $1 - \delta$ over the m i.i.d. samples $(X_i, Y_i)_{i=1}^m$ from \mathcal{D} , for every $h \in \mathcal{H}$, we have*

$$|er_Z(h) - er_D(h)| \leq G(m, \delta),$$

where

$$er_D(h) = \mathbb{E}_{(X,Y) \sim D} [\mathbb{1}(Yh(X) \leq 0)], \quad er_Z(h) = \frac{1}{m} \sum_{i=1}^m \mathbb{1}(Y_i h(X) \leq 0),$$

and

$$G(m, \delta) = \frac{1}{m} + \sqrt{\frac{\log(4/\delta) + d \log(2em/d)}{m}}.$$

Here d is the VC dimension of the function space \mathcal{H} .

Since the VC dimension of the linear classifier in \mathbb{R}^d is $d+1$, we have, with a probability $1 - \delta/m$,

$$\ell_b(\mathbf{w}|\text{DIS}(\Omega_k)) - \ell_b(\mathbf{w}_*|\text{DIS}(\Omega_k)) \leq \frac{2}{n_k} + 2\sqrt{\frac{\log(4m/\delta) + (d+1) \log(2en_k/(d+1))}{n_k}}, \quad (13)$$

and therefore

$$\ell_b(\mathbf{w}_{k+1}) - \ell_b(\mathbf{w}_*) \stackrel{(11), (12), (13)}{\leq} \eta_k r_k^{\gamma_+/\kappa},$$

where

$$\eta_k := \mu \ell_+^{1/\kappa} 2^{1+\gamma_+/\kappa} \theta(\varepsilon) \left(\frac{1}{n_k} + \sqrt{\frac{\log(4m/\delta) + (d+1) \log(2en_k/(d+1))}{n_k}} \right).$$

Using the lower bound in Assumption (III), we have

$$\ell_b(\mathbf{w}_{k+1}) - \ell_b(\mathbf{w}_*) \geq \ell_- \|\mathbf{w}_{k+1} - \bar{\mathbf{w}}_*\|^{\gamma_-}.$$

As a result, with a probability $1 - \delta/m$, we have

$$\|\mathbf{w}_{k+1} - \bar{\mathbf{w}}_*\| \leq \frac{\eta_k^{1/\gamma_-}}{\ell_-^{1/\gamma_-}} r_k^{\gamma_+ / (\gamma_- \kappa)}. \quad (14)$$

To ensure the R.H.S. of (14) is smaller than $r_{k+1} = r_k/2$, we need

$$\eta_k \leq \frac{\ell_-}{2^{\gamma_-}} r_k^{\gamma_- - \gamma_+ / \kappa},$$

which is

$$\frac{1}{n_k} + \sqrt{\frac{\log(4m/\delta) + (d+1)\log(2en_k/(d+1))}{n_k}} \leq \frac{\ell_- r_k^{\gamma_- - \gamma_+/\kappa}}{\mu \ell_+^{1/\kappa} 2^{1+\gamma_- + \gamma_+/\kappa} \theta(\varepsilon)}. \quad (15)$$

Since $(d+1)\log(2en_k/(d+1)) \geq 1$, we must have

$$\frac{1}{n_k} \leq \sqrt{\frac{\log(4m/\delta) + (d+1)\log(2en_k/(d+1))}{n_k}}.$$

Thus, to satisfy the condition in (15), it is sufficient to ensure

$$n_k \geq \left(\frac{\mu \ell_+^{1/\kappa} 2^{2+\gamma_- + \gamma_+/\kappa} \theta(\varepsilon)}{\ell_- r_k^{\gamma_- - \gamma_+/\kappa}} \right)^2 \left(\log \frac{4m}{\delta} + (d+1) \log \frac{2en_k}{d+1} \right). \quad (16)$$

A sufficient condition to (16) is

$$n_k \geq 2 \left(\frac{c\theta(\varepsilon)}{r_k^{\gamma_- - \gamma_+/\kappa}} \right)^2 \log \frac{4m}{\delta}, \quad (17)$$

and

$$n_k \geq 2 \left(\frac{c\theta(\varepsilon)}{r_k^{\gamma_- - \gamma_+/\kappa}} \right)^2 (d+1) \log \frac{2en_k}{d+1}, \quad (18)$$

where

$$c = \frac{\mu \ell_+^{1/\kappa} 2^{2+\gamma_- + \gamma_+/\kappa}}{\ell_-}.$$

To address the inequality in (18), define

$$a = 2 \left(\frac{c\theta(\varepsilon)}{r_k^{\gamma_- - \gamma_+/\kappa}} \right)^2.$$

Notice that

$$\log \frac{2en_k}{d+1} = \log \frac{n_k}{2a(d+1)} + \log 4ae \leq \frac{n_k}{2a(d+1)} - 1 + \log 4ae = \frac{n_k}{2a(d+1)} + \log 4a,$$

where we use the inequality $1 + \log x \leq x$. Thus, a sufficient condition to (18) is

$$n_k \geq a(d+1) \left(\frac{n_k}{2a(d+1)} + \log 4a \right),$$

which implies

$$n_k \geq 2a(d+1) \log 4a = 4 \left(\frac{c\theta(\varepsilon)}{r_k^{\gamma_- - \gamma_+/\kappa}} \right)^2 (d+1) \left(\log 8 + 2 \log \frac{c\theta(\varepsilon)}{r_k^{\gamma_- - \gamma_+/\kappa}} \right). \quad (19)$$

Then, it is clear that (6) is a sufficient condition to ensure both (17) and (19).

D Proof of Theorem 4

Define

$$P_n(\mathbf{w}) = \frac{1}{n} \sum_{i=1}^n \phi(y_i \mathbf{w}^\top \mathbf{x}_i) - \phi(y_i \mathbf{w}_*^\top \mathbf{x}_i), \quad P(\mathbf{w}) = \mathbb{E} \left[\phi(y \mathbf{w}^\top \mathbf{x}) - \phi(y \mathbf{w}_*^\top \mathbf{x}) \right],$$

and

$$\|P_n(\mathbf{w}) - P(\mathbf{w})\|_r = \sup \{ |P_n(\mathbf{w}) - P(\mathbf{w})| : \|\mathbf{w} - \mathbf{w}_*\| \leq r \}.$$

Since

$$|\phi(y \mathbf{w}^\top \mathbf{x}) - \phi(y \mathbf{w}_*^\top \mathbf{x})| \leq L \|\mathbf{w} - \mathbf{w}_*\| \|\mathbf{x}\| \leq Lr,$$

following [29, Theorem 2.5], we have, with a probability $1 - \delta/m$,

$$\|P_n(\mathbf{w}) - P(\mathbf{w})\|_r \leq Lr \sqrt{\frac{2 \log m / \delta}{n}} + \mathbb{E} [\|P_n(\mathbf{w}) - P(\mathbf{w})\|_r]. \quad (20)$$

Let $\{\varepsilon_i\}$ be Rademacher random variables, that is, ε_i takes the values 1 and -1 with probability $1/2$ each. Using symmetrization inequality of Rademacher complexity [29], we have

$$\mathbb{E} [\|P_n(\mathbf{w}) - P(\mathbf{w})\|_r] \leq 2\mathbb{E} \left[\frac{1}{n} \sup_{\|\mathbf{w} - \mathbf{w}_*\| \leq r} \left| \sum_{i=1}^n \varepsilon_i \left(\phi(y_i \mathbf{w}^\top \mathbf{x}_i) - \phi(y_i \mathbf{w}_*^\top \mathbf{x}_i) \right) \right| \right].$$

Since $\phi(\cdot)$ is L -Lipschitz continuous, one can use the contraction inequality [29] to get

$$\begin{aligned} \mathbb{E} [\|P_n(\mathbf{w}) - P(\mathbf{w})\|_r] &\leq \frac{4L}{n} \mathbb{E} \left[\sup_{\|\mathbf{w} - \mathbf{w}_*\| \leq r} \left| \sum_{i=1}^n \varepsilon_i (\mathbf{w} - \mathbf{w}_*)^\top \mathbf{x}_i \right| \right] \\ &\leq \frac{4Lr}{n} \mathbb{E} \left[\left\| \sum_{i=1}^n \varepsilon_i \mathbf{x}_i \right\| \right] \leq \frac{4Lr}{n} \sqrt{\mathbb{E} \left[\left\| \sum_{i=1}^n \varepsilon_i \mathbf{x}_i \right\|^2 \right]} \\ &= \frac{4Lr}{n} \sqrt{\mathbb{E} \left[\sum_{i=1}^n \|\mathbf{x}_i\|^2 + \sum_{i \neq j} \varepsilon_i \varepsilon_j \mathbf{x}_i^\top \mathbf{x}_j \right]} \leq \frac{4Lr}{n} \sqrt{n} = \frac{4Lr}{\sqrt{n}}, \end{aligned} \quad (21)$$

where in the last inequality we use the fact that ε_i 's are independent from \mathbf{x}_i 's such that

$$\mathbb{E} \varepsilon_i \varepsilon_j \mathbf{x}_i^\top \mathbf{x}_j = \mathbb{E} \left[\mathbf{x}_i^\top \mathbf{x}_j \mathbb{E}[\varepsilon_i \varepsilon_j] \right] = 0, \quad \text{when } i \neq j.$$

We complete the proof by combining (20) and (21).

E Proof of Theorem 5

Based on our induction assumption, we have $\|\tilde{\mathbf{w}}_* - \mathbf{w}_k\| \leq r_k$, implying $\|\mathbf{w}_* - R\mathbf{w}_k\| \leq Rr_k$. Since $\tilde{\mathbf{w}}_{k+1}$ is the optimal solution to (5) and $\mathbf{w}_* \in \Delta_k$, we have

$$\sum_{t=1}^{n_k} \phi \left(y_k^t \tilde{\mathbf{w}}_{k+1}^\top \mathbf{x}_k^t \right) - \sum_{t=1}^{n_k} \phi \left(y_k^t \mathbf{w}_*^\top \mathbf{x}_k^t \right) \leq 0. \quad (22)$$

Notice that for any $\mathbf{w} \in \Delta_k$, we have $\|\mathbf{w} - \mathbf{w}_*\| \leq 2Rr_k$. Following Theorem 4 and (22), we have, with a probability $1 - \delta/m$

$$\ell_\phi(\tilde{\mathbf{w}}_{k+1}|\text{DIS}(\Omega_k)) - \ell_\phi(\mathbf{w}_*|\text{DIS}(\Omega_k)) \leq \frac{2LRr_k}{\sqrt{n_k}} \left(4 + \sqrt{2 \log \frac{m}{\delta}} \right) \quad (23)$$

where $\ell_\phi(\cdot|T)$ is defined as the convex risk for instances sampled from the set T . From the discussion in the end of Section 3.3, we know that \mathbf{w}_* minimizes both the $\ell_b(\cdot|\text{DIS}(\Omega_k))$ and $\ell_\phi(\cdot|\text{DIS}(\Omega_k))$ over all measurement functions. Thus, we can apply Theorem 1 to bound the excess binary risk as follows

$$\begin{aligned} & \psi(\ell_b(\tilde{\mathbf{w}}_{k+1}|\text{DIS}(\Omega_k)) - \ell_b(\mathbf{w}_*|\text{DIS}(\Omega_k))) \\ & \leq \ell_\phi(\tilde{\mathbf{w}}_{k+1}|\text{DIS}(\Omega_k)) - \ell_\phi(\mathbf{w}_*|\text{DIS}(\Omega_k)) \\ & \stackrel{(23)}{\leq} \frac{2LRr_k}{\sqrt{n_k}} \left(4 + \sqrt{2 \log \frac{m}{\delta}} \right). \end{aligned}$$

Using the assumption that $\psi(z) \geq az^\gamma$, we have

$$\ell_b(\tilde{\mathbf{w}}_{k+1}|\text{DIS}(\Omega_k)) - \ell_b(\mathbf{w}_*|\text{DIS}(\Omega_k)) \leq \left(\frac{2LR}{a\sqrt{n_k}} \left(4 + \sqrt{2 \log \frac{m}{\delta}} \right) \right)^{1/\gamma} r_k^{1/\gamma}. \quad (24)$$

Following the same analysis as that for Theorem 3, we have, with a probability $1 - \delta$

$$\ell_b(\mathbf{w}_{k+1}) - \ell_b(\mathbf{w}_*) \stackrel{(11), (12), (24)}{\leq} \nu_k r_k^{\frac{1}{\gamma} + \frac{\gamma_+}{\kappa}},$$

where

$$\nu_k = \mu \ell_+^{1/\kappa} 2^{\gamma_+/\kappa} \theta(\varepsilon) \left(\frac{2LR}{a\sqrt{n_k}} \left(4 + \sqrt{2 \log \frac{m}{\delta}} \right) \right)^{1/\gamma}$$

Using the lower bound in Assumption (III), we have

$$\|\mathbf{w}_{k+1} - \bar{\mathbf{w}}_*\| \leq \frac{\nu_k^{1/\gamma_-}}{\ell_-^{1/\gamma_-}} r_k^{\frac{1}{\gamma_-} \left(\frac{1}{\gamma} + \frac{\gamma_+}{\kappa} \right)}. \quad (25)$$

To ensure the R.H.S. of (25) is smaller than $r_{k+1} = r_k/2$, we need

$$\nu_k \leq \frac{\ell_-}{2^{\gamma_-}} r_k^{\gamma_- - \frac{1}{\gamma} - \frac{\gamma_+}{\kappa}}.$$

which requires n_k to satisfy the condition in (8).

F Proof of Theorem 7

Similar to the proof of Theorem 4, we define

$$P_n(\mathbf{w}) = \frac{1}{n} \sum_{i=1}^n \phi(y_i \mathbf{w}^\top \mathbf{x}_i) - \phi(y_i \mathbf{w}_*^\top \mathbf{x}_i), \quad P(\mathbf{w}) = \mathbb{E} \left[\phi(y \mathbf{w}^\top \mathbf{x}) - \phi(y \mathbf{w}_*^\top \mathbf{x}) \right],$$

and

$$\|P_n(\mathbf{w}) - P(\mathbf{w})\|_r = \sup \{|P_n(\mathbf{w}) - P(\mathbf{w})| : \|\mathbf{w} - \mathbf{w}_*\| \leq r\}.$$

The difference is that we need to use the Adamczak bound [29, Section 2.3] to deal with the challenge that the function value may be unbounded. Based on the Adamczak bound, we have, with a probability $1 - \delta/m$,

$$\|P_n(\mathbf{w}) - P(\mathbf{w})\|_r \leq C_1 \left[\mathbb{E} [\|P_n(\mathbf{w}) - P(\mathbf{w})\|_r] + \sigma_r \sqrt{\frac{\log(m/\delta)}{n}} + U_r \frac{\log(m/\delta)}{n} \right], \quad (26)$$

where

$$\begin{aligned} \sigma_r^2 &\leq \sup_{\mathbf{w}: \|\mathbf{w} - \mathbf{w}_*\| \leq r} \mathbb{E} \left[\left(\phi(y\mathbf{w}^\top \mathbf{x}) - \phi(y\mathbf{w}_*^\top \mathbf{x}) \right)^2 \right], \\ U_r &= \left\| \max_{1 \leq i \leq n} \sup_{\mathbf{w}: \|\mathbf{w} - \mathbf{w}_*\| \leq r} \left| \phi(y_i \mathbf{w}^\top \mathbf{x}_i) - \phi(y_i \mathbf{w}_*^\top \mathbf{x}_i) \right| \right\|_{\psi_1}, \end{aligned}$$

and C_1 is some constant. In the following, we consider how to bound the three terms on the R.H.S. of (26).

Bounding $\mathbb{E} [\|P_n(\mathbf{w}) - P(\mathbf{w})\|_r]$ Following the same analysis in the proof of Theorem 4, we arrive at

$$\mathbb{E} [\|P_n(\mathbf{w}) - P(\mathbf{w})\|_r] \leq \frac{4Lr}{n} \sqrt{\mathbb{E} \left[\sum_{i=1}^n \|\mathbf{x}_i\|^2 \right]}. \quad (27)$$

Following the equivalence of sub-exponential (sub-gaussian) properties [35], we have

$$\mathbb{E} [\|\mathbf{x}_i\|^2] \leq \begin{cases} \left(C_2 \|\|\mathbf{x}_i\|\|_{\psi_1} 2 \right)^2 \leq 4C_2^2, & \text{if } \|\|X\|\|_{\psi_1} \leq 1 \\ \left(C_3 \|\|\mathbf{x}_i\|\|_{\psi_2} \sqrt{2} \right)^2 \leq 2C_3^2, & \text{if } \|\|X\|\|_{\psi_2} \leq 1 \end{cases} \quad (28)$$

where C_2 and C_3 are some constants. Thus, in both cases, we have $\mathbb{E} [\|\mathbf{x}_i\|^2] \leq C_4$ for some constant C_4 . Combining (27) and (28), we have

$$\mathbb{E} [\|P_n(\mathbf{w}) - P(\mathbf{w})\|_r] \leq \frac{C_5 L r}{\sqrt{n}}, \quad (29)$$

for some constant C_5 .

Bounding σ_r^2 Since $\phi(\cdot)$ is Lipschitz continuous with constant L , we have

$$\begin{aligned} \sigma_r^2 &\leq \sup_{\mathbf{w}: \|\mathbf{w} - \mathbf{w}_*\| \leq r} \mathbb{E} \left[\left(Ly(\mathbf{w} - \mathbf{w}_*)^\top \mathbf{x} \right)^2 \right] \\ &\leq L^2 \sup_{\mathbf{w}: \|\mathbf{w} - \mathbf{w}_*\| \leq r} \mathbb{E} [\|\mathbf{w} - \mathbf{w}_*\|^2 \|\mathbf{x}\|^2] \leq r^2 L^2 \mathbb{E} [\|\mathbf{x}\|^2] \stackrel{(28)}{\leq} C_6 r^2 L^2, \end{aligned} \quad (30)$$

for some constant C_6 .

Bounding U_r Based on the bound for the Orlicz norm of a finite maximum [33, Lemma 2.2.2], we have

$$\begin{aligned} U_r &\leq C_7 \sqrt{\log(n+1)} \max_{1 \leq i \leq n} \left\| \sup_{\mathbf{w}: \|\mathbf{w} - \mathbf{w}_*\| \leq r} \left| \phi(y_i \mathbf{w}^\top \mathbf{x}_i) - \phi(y_i \mathbf{w}_*^\top \mathbf{x}_i) \right| \right\|_{\psi_1} \\ &= C_7 \sqrt{\log(n+1)} \left\| \sup_{\mathbf{w}: \|\mathbf{w} - \mathbf{w}_*\| \leq r} \left| \phi(y_1 \mathbf{w}^\top \mathbf{x}_1) - \phi(y_1 \mathbf{w}_*^\top \mathbf{x}_1) \right| \right\|_{\psi_1}, \end{aligned}$$

where C_7 is some constant, and the last equality comes from the fact $\{(\mathbf{x}_i, y_i)\}$ follow the same distribution. Based on the Lipschitz continuity of the loss function $\psi(\cdot)$, we further have

$$U_r \leq C_7 \sqrt{\log(n+1)} \left\| \sup_{\mathbf{w}: \|\mathbf{w} - \mathbf{w}_*\| \leq r} \left| L(\mathbf{w} - \mathbf{w}_*)^\top \mathbf{x}_1 \right| \right\|_{\psi_1} \leq C_7 L r \sqrt{\log(n+1)} \|\mathbf{x}_1\|_{\psi_1}. \quad (31)$$

Thus, in the case that $\|X\|_{\psi_1} \leq 1$, we have

$$U_r \leq C_7 L r \sqrt{\log(n+1)}.$$

In the case that $\|X\|_{\psi_2} \leq 1$, we use the following relation between ψ_p - and ψ_q -norms [33, Page 95]

$$\|\eta\|_{\psi_p} \leq \|\eta\|_{\psi_q} (\log 2)^{1/q-1/p}, \quad p \leq q. \quad (32)$$

Then, we have

$$U_r \stackrel{(31),(34)}{\leq} C_7 (\log 2)^{-1/2} L r \sqrt{\log(n+1)} \|\mathbf{x}_1\|_{\psi_2} \leq C_7 (\log 2)^{-1/2} L r \sqrt{\log(n+1)}.$$

As a result, in both cases, we have

$$U_r \leq C_8 L r \sqrt{\log(n+1)}, \quad (33)$$

for some constant C_8 .

Substituting (29), (30), and (34) into (26), we have, with a probability $1 - \delta/m$,

$$\|P_n(\mathbf{w}) - P(\mathbf{w})\|_r \leq C_9 L r \left[\frac{1}{\sqrt{n}} + \sqrt{\frac{\log(m/\delta)}{n}} \left(1 + \sqrt{\log(n+1) \frac{\log(m/\delta)}{n}} \right) \right], \quad (34)$$

for some constant C_9 . From the condition in (10), we have

$$\begin{aligned} n &\geq \frac{n+1}{2} + \log \frac{m}{\delta} \cdot \log \left(\frac{2}{e} \log \frac{m}{\delta} \right) \\ &= \left(\frac{n+1}{2 \log(m/\delta)} + \log \left(\frac{2}{e} \log \frac{m}{\delta} \right) \right) \log \frac{m}{\delta} \\ &\geq \left(1 + \log \left(\frac{n+1}{2 \log(m/\delta)} \right) + \log \left(\frac{2}{e} \log \frac{m}{\delta} \right) \right) \log \frac{m}{\delta} \\ &= \log(n+1) \log \frac{m}{\delta}, \end{aligned} \quad (35)$$

where in the second inequality we use the fact that $1 + \log x \leq x$. Substituting (35) into (34), we have, with a probability $1 - \delta/m$,

$$\|P_n(\mathbf{w}) - P(\mathbf{w})\|_r \leq CLr \left(\frac{1}{\sqrt{n}} + \sqrt{\frac{\log(m/\delta)}{n}} \right),$$

for some constant C .