## **BSM** Primary Effects

Rick S. Gupta,<sup>1</sup> Alex Pomarol,<sup>2</sup> and Francesco Riva<sup>3</sup>

<sup>1</sup>IFAE, Universitat Autònoma de Barcelona, 08193 Bellaterra, Barcelona

<sup>2</sup>Dept. de Física, Universitat Autònoma de Barcelona, 08193 Bellaterra, Barcelona

<sup>3</sup>Institut de Théorie des Phénomènes Physiques, EPFL,1015 Lausanne, Switzerland

Using the predictive power of the effective field theory approach, we present a physical parametrization of the leading effects beyond the SM (BSM), that give us at present the best way to constrain heavy new-physics at low-energies. We show that other BSM effects are not independent from these ones, and we provide the explicit correlations. This information is useful to know where to primarily look for new physics in future experiments, and to know how this new physics is related to previous measurements, most importantly in electroweak-symmetry breaking processes or Higgs physics.

The absence at the LHC of new physics beyond the SM (BSM) suggests that the characteristic scale of new-physics  $\Lambda$  must be heavier than the electroweak (EW) scale. Assuming this, one can obtain an SM effective-theory as an expansion in SM fields and derivatives over  $\Lambda$ :  $\mathcal{L}_{\text{eff}} = \mathcal{L}_4 + \mathcal{L}_6 + \cdots$ , where  $\mathcal{L}_4$  is made of dimension-4 operators and defines what we call the SM Lagrangian, while  $\mathcal{L}_6$ , that contains dimension-6 operators, gives the leading BSM effects.<sup>1</sup> While the predictions from  $\mathcal{L}_4$  have been fully addressed and tested experimentally, it becomes now crucial to understand what are the complete predictions from  $\mathcal{L}_6$ .

This is the main goal of this article. Our approach however will differ from the usual one, in which one starts with the set of independent operators in  $\mathcal{L}_6$ [1, 2], and relates their Wilson coefficients to observables [3, 4]. We will follow here a bottom-up approach, our starting point being all possible new interactions among SM fields that can be induced by BSM physics. Not all these interactions can be obtained from  $\mathcal{L}_6$  and, of the possible ones, not all of them are independent. Our purpose is to find the set of independent new-interactions that can arise from  $\mathcal{L}_6$  and are, at present, the experimentally best tested ones. This set of physical quantities, that give us the best way to constrain new physics, will be called BSM primary effects. For electroweak processes these are, in some sense, a generalization of the well-known Sand T parameters [5].

Our main result will be to show which new-physics effects are not independent and are instead correlated with the BSM primaries (see also Ref. [6]). This information can be useful to understand where to prioritize new physics searches. Furthermore, in the future, if a deviation from the SM is measured, the correlations that we present will tell us where other BSM effects have to be found.

We start with the CP-conserving BSM-effects. It is useful to parametrize these effects as  $\hat{h}$ -dependent interactions, where  $\hat{h}$  is the neutral component of the Higgs-field:

$$\hat{h} \equiv v + h(x) \,, \tag{1}$$

where  $v \simeq 246$  GeV is the Higgs vacuum expectation value (VEV) and h the Higgs excitation. We first consider those effects which affect interactions involving only h, that arise from dimension-6 operators generated by multiplying operators in  $\mathcal{L}_4$  by  $|H|^2/\Lambda^2$  (H being the Higgs doublet). In the unitary gauge, that will be used henceforth, these effects can be captured by promoting the SM parameters, that we take to be  $e, s_{\theta_W}, g_s, Y_f, \lambda_h$ , and the Higgs kineticterm  $Z_h$ , <sup>2</sup> to  $\hat{h}$ -dependent functions:

$$e(\hat{h}), \ s_{\theta_W}(\hat{h}), \ g_s(\hat{h}), \ Y_f(\hat{h}), \ \lambda_h(\hat{h}), \ Z_h(\hat{h}).$$
 (2)

~

These functions can be expanded in powers of  $\hat{h}^2/\Lambda^2$ , e.g.  $e(\hat{h}) = e + \delta e \hat{h}^2/v^2 + \cdots$ , where (here and in what follows) we absorb powers of  $v^2/\Lambda^2$  in the expansion coefficients. In the vacuum  $\hat{h} = v$ , Eq. (2) only implies a redefinition of the SM parameters with no impact on physical processes, *i.e.*, these effects

 $<sup>^{1}</sup>$  Assuming lepton and baryon number conservation.

 $<sup>^2</sup>$  The effect of  $|H|^2/\Lambda^2$  multiplying the SM fermion kinetic-terms is redundant as can be eliminated by a redefinition of the SM fermions.

can only be probed in Higgs physics. To understand the effects from  $e(\hat{h}), s_{\theta_W}(\hat{h}), g_s(\hat{h})$ , it is convenient to write the SM gauge-interactions in a non-canonical way, since this is suitable to accommodate space-time dependent couplings while keeping gauge-invariance manifest. For the EW sector this is given by

$$\mathcal{L}_{\rm EW} = -\frac{1}{4e^2(\hat{h})} \left( A_{\mu\nu} + s_{\theta_W}^2(\hat{h}) Z_{\mu\nu} \right)^2 - \frac{c_{\theta_W}^2(\hat{h})}{4g^2(\hat{h})} Z_{\mu\nu}^2 - \frac{1}{2g^2(\hat{h})} W_{\mu\nu}^+ W^{-\mu\nu} + \frac{\hat{h}^2}{4} \left[ W_{\mu}^+ W^{-\mu} + \frac{1}{2} Z^{\mu} Z_{\mu} \right] + A_{\mu} J_{em}^{\mu} + Z_{\mu} J_3^{\mu} + W_{\mu}^+ J_{+}^{\mu} + W_{\mu}^- J_{-}^{\mu}, \qquad (3)$$

where  $g(\hat{h}) \equiv e(\hat{h})/s_{\theta_W}(\hat{h})$ , and we have defined  $J_{\pm}^{\mu} \equiv (J_1^{\mu} \pm iJ_2^{\mu})/\sqrt{2}, J_{em}^{\mu} \equiv J_3^{\mu} + J_Y^{\mu}, Z_{\mu\nu} \equiv \hat{Z}_{\mu\nu} - iW_{[\mu}^{+}W_{\nu]}^{-}, A_{\mu\nu} \equiv \hat{A}_{\mu\nu}, W_{\mu\nu}^{\pm} \equiv \hat{W}_{\mu\nu}^{\pm} \pm iW_{[\mu}^{\pm}(A+Z)_{\nu]}$ with  $\hat{V}_{\mu\nu} \equiv \partial_{\mu}V_{\nu} - \partial_{\nu}V_{\mu}$ , and  $J_a^{\mu}$  (a=1,2,3) and  $J_Y^{\mu}$ being respectively the  $SU(2)_L$  and  $U(1)_Y$  currents. In the vacuum  $\hat{h} = v$ , Eq. (3) gives the SM EWinteractions after substituting  $A_{\mu} \to A_{\mu} - s_{\theta_W}^2(v)Z_{\mu}$ and canonically normalizing the gauge-boson fields. The two independent deviations w.r.t. the SM, parametrized by  $e(\hat{h})$  and  $s_{\theta_W}(\hat{h})$ , can be projected orthogonally into two different physical processes, that we choose to be  $h \to \gamma\gamma$  and  $h \to Z\gamma$  for the accuracy to which they are experimentally constrained. Indeed, a  $h\gamma\gamma$  coupling can arise from the  $\hat{h}$ -dependence of e with constant  $s_{\theta_W}$ :

$$e(\hat{h}) = e(1 + \kappa_{\gamma\gamma} \frac{\hat{h}^2}{v^2}), \qquad s_{\theta_W}(\hat{h}) = s_{\theta_W}, \qquad (4)$$

that plugged into Eq. (3) gives, in the canonical basis, the BSM-terms:

$$\Delta \mathcal{L}^{h}_{\gamma\gamma} = \kappa_{\gamma\gamma} \left( \frac{h}{v} + \frac{h^2}{2v^2} \right) \left[ A_{\mu\nu} A^{\mu\nu} + Z_{\mu\nu} Z^{\mu\nu} + 2W^+_{\mu\nu} W^{-\mu\nu} \right], \quad (5)$$

where now and henceforth  $Z_{\mu\nu} \equiv \hat{Z}_{\mu\nu} - igc_{\theta_W}W^+_{[\mu}W^-_{\nu]}$ ,  $A_{\mu\nu} \equiv \hat{A}_{\mu\nu} - igs_{\theta_W}W^+_{[\mu}W^-_{\nu]}$  and  $W^{\pm}_{\mu\nu} \equiv \hat{W}^{\pm}_{\mu\nu} \pm igW^{\pm}_{[\mu}(s_{\theta_W}A + c_{\theta_W}Z)_{\nu]}$ . The first term of Eq. (5) contains a  $h\gamma\gamma$  coupling and corresponds to our first BSM primary effect: it defines the best observable that can be used to bound all terms in  $\Delta \mathcal{L}^h_{\gamma\gamma}$ . Indeed, from the experimental value of  $h \to \gamma\gamma$  [7] we obtain bounds on  $\kappa_{\gamma\gamma}$  at the per-mille level [4].

On the other hand, we can take, orthogonally to

Eq. (4), the direction

$$e(\hat{h}) = e, \qquad s_{\theta_W}^2(\hat{h}) = s_{\theta_W}^2(1 - \kappa_{Z\gamma} \frac{h^2}{v^2}), \qquad (6)$$

that in Eq. (3) gives the BSM-induced interactions

$$\Delta \mathcal{L}_{Z\gamma}^{h} = \kappa_{Z\gamma} \left( \frac{h}{v} + \frac{h^{2}}{2v^{2}} \right) \left[ t_{\theta_{W}} A_{\mu\nu} Z^{\mu\nu} + \frac{c_{2\theta_{W}}}{2c_{\theta_{W}}^{2}} Z_{\mu\nu} Z^{\mu\nu} + W_{\mu\nu}^{+} W^{-\mu\nu} \right].$$
(7)

The first term of Eq. (7) defines another BSM primary effect: its contribution to the  $hZ\gamma$  coupling, that is constrained by  $h \rightarrow Z\gamma$  searches. Similarly, taking the  $SU(3)_c$  coupling  $g_s(\hat{h}) = g_s(1 + \kappa_{GG}\hat{h}^2/v^2)$ , one obtains

$$\Delta \mathcal{L}_{GG}^{h} = \kappa_{GG} \left( \frac{h}{v} + \frac{h^2}{2v^2} \right) G_{\mu\nu}^{A} G^{A\,\mu\nu} \,, \qquad (8)$$

whose first term modifies the hGG coupling measured in  $GG \rightarrow h$  [7], that leads also to a per-mille bound on  $\kappa_{GG}$  [4]. Also, from  $Y_f(\hat{h})$  and  $\lambda_h(\hat{h})$ , we obtain

$$\Delta \mathcal{L}_{ff}^{h} = \delta \boldsymbol{g}_{ff}^{h} \left( h \bar{f}_{L} f_{R} + \text{h.c.} \right) \left( 1 + \frac{3h}{2v} + \frac{h^{2}}{2v^{2}} \right) ,$$
  
$$\Delta \mathcal{L}_{3h} = \delta \boldsymbol{g}_{3h} h^{3} \left( 1 + \frac{3h}{2v} + \frac{3h^{2}}{4v^{2}} + \frac{h^{3}}{8v^{3}} \right) , \qquad (9)$$

with  $\delta g_{ff}^{h} = -\frac{v}{\sqrt{2}} \partial_{h} Y_{f}(\hat{h})$  and  $\delta g_{3h} = -v^{2} \partial_{h} \lambda_{h}(\hat{h})$ , whose BSM primary effects are respectively the contributions to the hff and  $h^{3}$  interactions. Finally, from  $Z_{h}(\hat{h})$  we obtain, by going to the canonical basis, the BSM-effect

$$\Delta \mathcal{L}_{VV}^{h} = \delta \boldsymbol{g}_{\boldsymbol{V}\boldsymbol{V}}^{h} \left[ h \left( W^{+\mu} W_{\mu}^{-} + \frac{Z^{\mu} Z_{\mu}}{2c_{\theta_{W}}^{2}} \right) + \Delta_{h} \right],$$
(10)

where

$$\Delta_{h} = \left( W^{+\mu}W^{-}_{\mu} + \frac{Z^{\mu}Z_{\mu}}{2c^{2}_{\theta_{W}}} \right) \left( \frac{2h^{2}}{v} + \frac{4h^{3}}{3v^{2}} + \frac{h^{4}}{3v^{3}} \right) + \frac{m^{2}_{h}}{12m^{2}_{W}} \left( \frac{h^{4}}{v} + \frac{3h^{5}}{4v^{2}} + \frac{h^{6}}{8v^{3}} \right) + \frac{m_{f}}{4m^{2}_{W}} \left( \frac{h^{2}}{v} + \frac{h^{3}}{3v^{2}} \right) \left( \bar{f}_{L}f_{R} + \text{h.c.} \right), \quad (11)$$

 $\delta g_{VV}^{h} = m_{W}^{2} \partial_{h} Z_{h}(\hat{h})$  and we have redefined  $\delta g_{3h} \rightarrow \delta g_{3h} - m_{h}^{2}/(4m_{W}^{2}) \delta g_{VV}^{h}$  in Eq. (9) to eliminate a contribution to the  $h^{3}$  coupling from  $Z_{h}(\hat{h})$ . The

first term of Eq. (10) gives a contribution to the custodial-preserving coupling hVV (V = Z, W) and determines another BSM primary effect. This coupling can be measured, for example, in  $WW \rightarrow h$ .

It is important at this point to stress that, by construction, the different  $\Delta \mathcal{L}_i$  are orthogonally projected into different BSM primary effects, and none of the terms in a given  $\Delta \mathcal{L}_i$  contributes to other BSM primaries that is not its own (e.g., no term in  $\Delta \mathcal{L}^h_{\gamma\gamma}$ contributes to the  $hZ\gamma$ , hGG, hff,  $h^3$  or hVV coupling). The additional terms in each  $\Delta \mathcal{L}_i$ , beyond the BSM primary effect, tell us what physical processes are not independent and are instead correlated with the BSM primaries. This can be useful if a departure from the SM predictions is observed: for example, if only a deviation in  $h \to \gamma \gamma$  is measured, Eq. (5) tells us that there must also be departures in  $h \to ZZ/WW$ . Alternatively, if no deviations from the SM are found in the BSM primary effects, these relations can be used to put constraints on the size of the other terms in  $\Delta \mathcal{L}_i$ .

Having presented all possible interactions achieved by an  $\hat{h}$ -dependent shift in the SM parameters, we study next the set of possible (CP-conserving) BSM contributions that can lead to departures from gaugecoupling universality. How many effects of this type can we have? Since EM and  $SU(3)_c$  must be unbroken, only the W and Z couplings can receive deviations from the SM. Assuming for simplicity family universality we have, in principle, 9 gauge-boson couplings to fermions (the Z couplings to  $e_{L,R}$ ,  $\nu_L$ ,  $u_{L,R}$ ,  $d_{L,R}$  and the W couplings to  $e_L \nu_L$  and  $u_L d_L$ ), the Z coupling to  $\hat{h}$ , <sup>3</sup> and triple-gauge (TGC) and quarticgauge (QGC) self-couplings. We must however keep in mind that not all deviations in these couplings are independent from each other, since a linear combination of all these corresponds to the universal shift of Eq. (6).

Let us first look at the 10 gauge-boson couplings to fermions and to the Higgs-field  $\hat{h}$ . In the gauge eigenstate basis, corrections to these couplings arise from the  $\hat{h}$ -dependent interactions

$$\hat{h}^2 V^a_\mu J^{\mu a}_f$$
,  $\hat{h}^2 \eta^a V^a_\mu J^\mu_{Lf}$ ,  $\hat{h}^2 \eta^a V^a_\mu J^\mu_{Rf}$ , (12)

$$\hat{h}^{4}\eta^{a}\eta^{b}V_{\mu}^{a}V^{\mu\,b}\,,\quad \hat{h}^{4}\eta^{a}\eta^{b}V_{\mu}^{a}J_{L\,f}^{\mu\,b}\,,\tag{13}$$

where  $V^a_{\mu} \equiv W^a_{\mu} - t_{\theta_W} \delta^{a3} B_{\mu}$  (to preserve EM) and, to make the global  $SU(2)_L$  properties manifest, we have separated the interactions in which the Higgs enters as a singlet,  $\hat{h}^2$ , or as a triplet,

$$\hat{h}^2 \eta^a \in H^{\dagger} \sigma^a H$$
, with  $\eta^a \equiv (0,0,1)$ . (14)

The lepton currents are  $J_f^{\mu a} = \bar{L}_L \sigma^a \gamma^{\mu} L_L$ ,  $J_{Lf}^{\mu} = \bar{L}_L \gamma^{\mu} L_L$  and  $J_{Rf}^{\mu} = \bar{e}_R \gamma^{\mu} e_R$ , and similarly for quarks. All terms of Eq. (12) can arise from  $\mathcal{L}_6$  built as products of fermion currents and Higgs currents, these latter being, in the unitary gauge,

$$iH^{\dagger}\overset{\leftrightarrow}{D}_{\mu}H = -g\frac{\hat{h}^2}{2}\eta^a V^a_{\mu} , \ iH^{\dagger}\sigma^a\overset{\leftrightarrow}{D}_{\mu}H = g\frac{\hat{h}^2}{2}V^a_{\mu}.$$
(15)

Similarly, the first term of Eq. (13) can arise from a dimension-6 operator built by squaring the first Higgs current of Eq. (15). On the other hand, the second term in Eq. (13), containing four Higgs, can only arise from a dimension-8 operator and can then be neglected. This has the implication that, at the leading order ( $\mathcal{L}_6$ ), BSM-effects in the W couplings are not independent from those in the Z couplings.

There are many ways to connect the BSM-effects of Eq. (12) to experiments. Since the best constraints on Vff vertices come from measurement of the couplings at the Z-pole by LEP, it is convenient to parametrize the effects of Eq. (12) as modifications of the Z couplings to fermions:

$$\Delta \mathcal{L}_{ee}^{V} = \delta g_{eR}^{Z} \frac{\hat{h}^{2}}{v^{2}} Z^{\mu} \bar{e}_{R} \gamma_{\mu} e_{R}$$
(16)  
+  $\delta g_{eL}^{Z} \frac{\hat{h}^{2}}{v^{2}} \left[ Z^{\mu} \bar{e}_{L} \gamma_{\mu} e_{L} - \frac{c_{\theta_{W}}}{\sqrt{2}} (W^{+\mu} \bar{\nu_{L}} \gamma_{\mu} e_{L} + \text{h.c.}) \right]$   
+  $\delta g_{\nu L}^{Z} \frac{\hat{h}^{2}}{v^{2}} \left[ Z^{\mu} \bar{\nu}_{L} \gamma_{\mu} \nu_{L} + \frac{c_{\theta_{W}}}{\sqrt{2}} (W^{+\mu} \bar{\nu_{L}} \gamma_{\mu} e_{L} + \text{h.c.}) \right]$ 

for leptons, and similarly for quarks:

$$\Delta \mathcal{L}_{qq}^{V} = \delta \boldsymbol{g}_{\boldsymbol{u}\boldsymbol{R}}^{\boldsymbol{Z}} \frac{\hat{h}^{2}}{v^{2}} Z^{\mu} \bar{u}_{R} \gamma_{\mu} u_{R} + \delta \boldsymbol{g}_{\boldsymbol{d}\boldsymbol{R}}^{\boldsymbol{Z}} \frac{\hat{h}^{2}}{v^{2}} Z^{\mu} \bar{d}_{R} \gamma_{\mu} d_{R}$$
$$+ \delta \boldsymbol{g}_{\boldsymbol{d}\boldsymbol{L}}^{\boldsymbol{Z}} \frac{\hat{h}^{2}}{v^{2}} \left[ Z^{\mu} \bar{d}_{L} \gamma_{\mu} d_{L} - \frac{c_{\theta_{W}}}{\sqrt{2}} (W^{+\mu} \bar{u}_{L} \gamma_{\mu} d_{L} + \text{h.c.}) \right]$$
$$+ \delta \boldsymbol{g}_{\boldsymbol{u}\boldsymbol{L}}^{\boldsymbol{Z}} \frac{\hat{h}^{2}}{v^{2}} \left[ Z^{\mu} \bar{u}_{L} \gamma_{\mu} u_{L} + \frac{c_{\theta_{W}}}{\sqrt{2}} (W^{+\mu} \bar{u}_{L} \gamma_{\mu} d_{L} + \text{h.c.}) \right]$$
(17)

<sup>&</sup>lt;sup>3</sup> We do not independently count the W coupling to  $\hat{h}$ , since this is equivalent to considering the custodial-preserving combination  $\hat{h}^2(W^{\mu}W_{\mu} + Z^{\mu}Z_{\mu}/2c^2_{\theta_W})$  that has already been accounted for in Eq. (10).

Notice that, as discussed above, modifications to the W couplings are explicitly related to modifications to the Z couplings.

It remains to consider the independent effect of the first term of Eq. (13). We consider it in the following linear combination (that includes also terms of Eq. (12)):

$$- \delta g_{1}^{Z} c_{\theta_{W}}^{2} \frac{\hat{h}^{2}}{v^{2}} \left[ \frac{g^{2} \hat{h}^{2}}{2} \left( W_{\mu}^{+} W^{-\mu} + \frac{c_{2\theta_{W}}}{2 c_{\theta_{W}}^{4}} Z_{\mu} Z^{\mu} \right) + g(W_{\mu}^{-} J_{-}^{\mu} + \text{h.c.}) + \frac{g c_{2\theta_{W}}}{c_{\theta_{W}}^{3}} Z_{\mu} J_{Z}^{\mu} + 2e t_{\theta_{W}} Z_{\mu} J_{em}^{\mu} \right],$$
(18)

where  $J_Z^{\mu} = J_3^{\mu} - s_{\theta_W}^2 J_{em}^{\mu}$ . Why this particular combination? This is obtained by performing the EMpreserving shift  $s_{\theta_W}^2 \to s_{\theta_W}^2 (1 + 2\delta g_1^Z c_{\theta_W}^2 \hat{h}^2 / v^2)$ (keeping *e* constant), only in the SM gauge-couplings of the fermions and  $\hat{h}$ . In the vacuum  $\hat{h} = v$ , the effects in Eq. (18) can only be probed as a relative difference of  $s_{\theta_W}$  as measured in the fermion and  $\hat{h}$  sector (*Vff* couplings and gauge-boson masses), with respect to the value as measured in interactions involving gauge bosons only. Therefore it requires the knowledge of TGC/QGC. Indeed, by field redefinitions, the non-Higgs physics part of Eq. (18) can be rewritten as a contribution to the *VWW* coupling ( $g^V$ ) and *VV'WW* coupling ( $g^{VV'}$ ) only. This explicitly gives

$$\boldsymbol{\delta g_1^Z} = \frac{\delta g^Z}{g_{SM}^Z} = \frac{\delta g^{WW}}{2c_{\theta_W}^2 g_{SM}^{WW}} = \frac{\delta g^{ZZ}}{2g_{SM}^{ZZ}} = \frac{\delta g^{\gamma Z}}{g_{SM}^{\gamma Z}}.$$
 (19)

where  $\delta g_1^Z$  has been chosen to match the TGC definition of Ref. [8]. Eq. (18) gives however also a contribution to the custodial-preserving hVV coupling that defines one of our BSM primaries,  $\delta g_{VV}^h$ . To eliminate this, we redefine  $\delta g_{VV}^h \rightarrow \delta g_{VV}^h + g^2 v \delta g_1^Z c_{\theta_W}^2$  in Eq. (10), that gives an extra contribution proportional  $\delta g_1^Z$  to be added to Eq. (18). The final result is

$$\Delta \mathcal{L}_{g_1^Z} = \delta g_1^Z \bigg[ igc_{\theta_W} \bigg( Z^\mu (W^{+\nu} W^-_{\mu\nu} - \text{h.c.}) + Z^{\mu\nu} W^+_\mu W^-_\nu \bigg)$$
  
$$- 2gc_{\theta_W}^2 \frac{h}{v} \bigg( W^-_\mu J^\mu_- + \text{h.c.} + \frac{c_{2\theta_W}}{c_{\theta_W}^3} Z_\mu J^\mu_Z + \frac{2s_{\theta_W}^2}{c_{\theta_W}} Z_\mu J^\mu_{em} \bigg)$$
  
$$\times \bigg( 1 + \frac{h}{2v} \bigg) + \frac{e^2 v}{2c_{\theta_W}^2} h Z_\mu Z^\mu + g^2 c_{\theta_W}^2 v \Delta_h \qquad (20)$$

$$-g^{2}c_{\theta_{W}}^{2}\left(W_{\mu}^{+}W^{-\mu}+\frac{c_{2\theta_{W}}}{2c_{\theta_{W}}^{4}}Z_{\mu}Z^{\mu}\right)\left(\frac{5h^{2}}{2}+\frac{2h^{3}}{v}+\frac{h^{4}}{2v^{2}}\right)\right]$$

The interesting property of our parametrization Eqs. (16,17,20) is the following. Since BSM-effects to SM propagators can always be eliminated through the equations of motion (EOM), there is a one to one correspondence between each of the  $\delta g_f^Z$  of Eqs. (16,17) and the corresponding  $\Gamma(Z \rightarrow ff)$ partial-width measured at LEP1 [9].<sup>4</sup> Therefore all the 7 parameters  $\delta g_f^Z$  can be bounded at the permille level by Z decay-widths and asymmetries at LEP1,<sup>5</sup> using  $\alpha_{em}$ ,  $m_W$  and  $m_Z$  as the SM input parameters. This latter choice makes the phenomenological analysis particularly transparent, since these input parameters receive no corrections from BSMeffects.<sup>6</sup> On the other hand,  $\delta g_1^Z$  is constrained by TGC measurements at LEP2 [10] and LHC. Although a global analysis on the  $\mathcal{L}_6$  contributions to TGC, using all existing data, does not exist yet, we expect that bounds on  $\delta g_1^Z$  can reach at present the per-cent level [11, 12]. Eq. (20) gives also contributions to Higgs physics, such as a custodial-breaking hVV coupling or new effects to  $h \to V \bar{f} f$  that could be also used to constrain  $\delta g_1^Z$ . We believe however that when rigorous analyses of both TGC and Higgs physics (on the lines of Ref. [13]) will be available, TGC constraints will always outdo Higgs physics ones [12], so that our parametrization will remain the most convenient.

At this point the reader might be surprised by the unorthodox choice of abandoning the traditional parametrization of BSM-effects that includes the *S* and *T*-parameters [5], in favor of a description in terms of vertex corrections as given in Eqs. (16,17). As already explained, we consider the latter more suited to compare with the experimental data. Nevertheless, if desired, two of the parameters of Eqs. (16,17), e.g.,  $\delta g_{eL}^Z$  and  $\delta g_{\nu L}^Z$  [4], can always

<sup>) &</sup>lt;sup>4</sup> This is true in the limit in which  $m_f$  is neglected, so that interference with dipole-type BSM contributions vanishes.

<sup>&</sup>lt;sup>5</sup> These measurements can also be combined with the measurement of the Wud coupling as extracted from K and  $\beta$ -decays in combination with information on 4-fermion interactions from the LHC [4].

<sup>&</sup>lt;sup>6</sup> Four-fermion interactions in  $\mathcal{L}_6$  (which have no direct relation with Higgs physics and can therefore be studied separately) affect the value of  $G_F$  as extracted through the measurement of  $\mu$ -decay. For this reason the traditional choice of using  $G_F$  to fix one input parameter is less convenient than the one we propose here.

be traded with

$$\Delta \mathcal{L}_{\hat{S}} = \hat{S} \, \frac{g s_{\theta_W}^2}{c_{\theta_W}^3} \frac{\hat{h}^2}{v^2} \, Z_\mu \left[ J_Z^\mu - c_{\theta_W}^2 J_{em}^\mu + \frac{g}{c_{\theta_W}} \frac{\hat{h}^2}{4} Z^\mu \right] \,,$$
  
$$\Delta \mathcal{L}_T = -\frac{\hat{T}}{2} \frac{\hat{h}^4}{v^4} m_Z^2 Z^\mu Z_\mu \,. \tag{21}$$

It is clear that Eqs. (21) can arise from linear combinations of terms in Eq. (12) and Eq. (13), and therefore gives a different parametrization from Eqs. (16,17). To see that indeed  $\hat{\mathbf{S}}$  is the *S*-parameter (we use the *S*, *T* normalization of Ref. [14]), one must realize that the terms in brackets in  $\Delta \mathcal{L}_{\hat{S}}$  are proportional to  $J_Y^{\mu}$ , and therefore (by the EOM) can be written as  $\Delta \mathcal{L}_{\hat{S}} = -\hat{\mathbf{S}}(s_{\theta_W}/c_{\theta_W}^2)(\hat{h}^2/v^2) Z_{\mu}\partial_{\nu}B^{\mu\nu}$ . After partial integration, this gives

$$\Delta \mathcal{L}_{\hat{S}} = -\frac{\hat{S}}{2} \Big[ t_{\theta_W} \, \frac{\hat{h}^2}{v^2} B_{\mu\nu} \hat{W}_3^{\mu\nu} - t_{\theta_W}^2 \, \frac{\hat{h}^2}{v^2} B_{\mu\nu} B^{\mu\nu} \\ -2 \, \frac{\partial_\nu \hat{h}^2}{v^2} B_{\mu\nu} (t_{\theta_W} W_3^\mu - t_{\theta_W}^2 B^\mu) \Big] \,. \tag{22}$$

For  $\hat{h} = v$ , the first term coincides with the  $\hat{S}$  parameter [14], while the second term merely represents a redefinition of the SM parameter g', and the last one  $\propto \partial_{\nu} \hat{h}^2$  vanishes.

Apart from Eq. (12), there can also be BSMinduced interactions between fermion and gaugebosons of a different type than those in the SM. These include couplings of W to right-handed quarks and dipole-type interactions, that we parametrize as

$$\Delta \mathcal{L}_{R}^{W} = \delta g_{R}^{W} \frac{\hat{h}^{2}}{v^{2}} W_{\mu}^{+} \bar{u}_{R} \gamma^{\mu} d_{R} + \text{h.c.}, \qquad (23)$$

$$\Delta \mathcal{L}_{\text{dipole}}^{V} = \frac{eY_{q} \hat{h}}{m_{W}^{2}} \Big[ \delta \kappa_{q}^{G} \frac{g_{s}}{e} \bar{q}_{L} T^{A} \sigma^{\mu\nu} q_{R} G_{\mu\nu}^{A} + \delta \kappa_{q}^{A} (T_{3} \bar{q}_{L} \sigma^{\mu\nu} q_{R} A_{\mu\nu} + \frac{s_{\theta_{W}}}{\sqrt{2}} \bar{u}_{L} \sigma^{\mu\nu} d_{R} W_{\mu\nu}^{+}) + \delta \kappa_{q}^{Z} (T_{3} \bar{q}_{L} \sigma^{\mu\nu} q_{R} Z_{\mu\nu} + \frac{c_{\theta_{W}}}{\sqrt{2}} \bar{u}_{L} \sigma^{\mu\nu} d_{R} W_{\mu\nu}^{+}) + \text{h.c.} \Big],$$

for quarks q = u, d, where the coefficients are assumed to be real and  $T_3$  denotes weak isospin (and similarly for leptons). Note that the dipole interactions with W are not independent from those of Aand Z, as the term that splits these dipole interactions,  $\hat{h}^2 \eta_a W^a_{\mu\nu} \hat{h} \bar{q}_L \sigma^{\mu\nu} q_R$ , arises at dimension-8.

Let us now move to TGC and QGC. At  $O(p^4)$  there are 4 possible CP-conserving TGC couplings and 5 QGC [15], but not all can arise from  $\mathcal{L}_6$ . We already encountered one with the same Lorentz structure as in the SM:  $\Delta \mathcal{L}_{g_1^Z}$  that led to Eq. (19). Other contributions could in principle arise from  $\mathcal{L}_6$  operators containing covariant derivatives and/or fieldstrengths. However, by integration by parts and using the EOM, one can reduce them to dimension-6 operators with only field-strengths [2].<sup>7</sup> The only operators at  $O(p^4)$ , made of field-strengths and contributing to EWSB, are

$$\hat{h}^2 \eta^a W^a_{\mu\nu} B^{\mu\nu} , \quad \hat{h}^4 \eta^a \eta^b W^a_{\mu\nu} W^{b\,\mu\nu} .$$
 (24)

The second one involves four Higgs and cannot arise from dimension-6 operators, while the first one gives

$$\hat{h}^{2}\eta^{a}W^{a}_{\mu\nu}B^{\mu\nu} =$$

$$\hat{h}^{2}\left[\hat{W}^{3}_{\mu\nu}B^{\mu\nu} + 2igc_{\theta_{W}}W^{-}_{\mu}W^{+}_{\nu}(A^{\mu\nu} - t_{\theta_{W}}Z^{\mu\nu})\right].$$
(25)

The second term clearly contains a new dipole-type TGC for the W, that can be identified with  $\delta \kappa_{\gamma}$  of Ref. [8]. Since Eq. (25) also contains contributions to other BSM primaries (such as  $\hat{S}$  and  $h \to \gamma\gamma, Z\gamma$ ), we must arrange a linear combination that does not project into them. From the relation

$$t_{\theta_W} \frac{\hat{h}^2}{2v^2} W^3_{\mu\nu} B^{\mu\nu} = -\frac{\Delta \mathcal{L}_{\hat{S}}}{\hat{S}} + s^2_{\theta_W} \frac{\Delta \mathcal{L}^h_{\gamma\gamma}}{\kappa_{\gamma\gamma}} + c_{2\theta_W} \frac{\Delta \mathcal{L}^h_{Z\gamma}}{\kappa_{Z\gamma}} -\frac{\Delta \mathcal{L}_{\kappa_{\gamma}}}{\delta \kappa_{\gamma}}, \qquad (26)$$

we have that

$$\Delta \mathcal{L}_{\kappa_{\gamma}} = \frac{\delta \kappa_{\gamma}}{v^2} \Big[ i e \hat{h}^2 (A_{\mu\nu} - t_{\theta_W} Z_{\mu\nu}) W^{+\mu} W^{-\nu} \quad (27) \\ + Z_{\nu} \partial_{\mu} \hat{h}^2 (t_{\theta_W} A^{\mu\nu} - t_{\theta_W}^2 Z^{\mu\nu}) + \frac{(\hat{h}^2 - v^2)}{2} \\ \times \Big( t_{\theta_W} Z_{\mu\nu} A^{\mu\nu} + \frac{c_{2\theta_W}}{2c_{\theta_W}^2} Z_{\mu\nu} Z^{\mu\nu} + W^+_{\mu\nu} W^{-\mu\nu} \Big) \Big]$$

gives us the combination that we were looking for: it projects into a new BSM primary effect, the TGC  $\delta \kappa_{\gamma}$  [8], but not into previous ones.

The list of the  $\hat{h}$ -dependent interactions presented so far is in principle not complete. We could also have dimension-6 interaction terms of the following type:  $(\partial_{\mu}\hat{h})^{2}V^{\nu}V_{\nu}, \ \partial_{\mu}\hat{h}V^{\mu}V_{\nu}V^{\nu}, \ (\partial_{\mu}\hat{h})V_{\nu}\bar{f}_{L}\sigma^{\mu\nu}f_{R}$ 

<sup>&</sup>lt;sup>7</sup> Dimension-6 operators with Higgs and derivatives can always be written, using integration by parts, as operators of the type  $H^{\dagger}D_{\mu}HD_{\nu}F^{\mu\nu}$  and  $H^{\dagger}D_{\mu}D_{\nu}HF^{\mu\nu}$ . The former reduces to Eq. (12) and Eq. (13) by the gauge-boson EOM, while the latter can, by means of  $[D_{\mu}, D_{\nu}] = F_{\mu\nu}$ , be written as an operator made of field-strengths only. Operators built with Eq. (15) contributing to TGC/QGC are always of dimension larger than six.

and  $\hat{h}V_{\mu}V^{\mu}\bar{f}_{L}f_{R}$  (other ones can be reduced to previous ones by redefinitions or partial integration). Nevertheless, it is easy to check that none of these terms can arise from operators in  $\mathcal{L}_{6}$  independent from the ones introduced before.

The list of CP-even independent effects is completed by 4-fermion interactions  $\Delta \mathcal{L}_{4f}$  and  $O(p^6)$  interactions, both of which cannot contain the Higgsfield  $\hat{h}$  if they arise from dimension-6 operators (they preserve the SM symmetries). The relation of 4fermion interactions with experiments is straightforward and they do not interfere with the observables determining our previous BSM primaries (see footnote 6). The complete list can be found in Ref. [2]. On the other hand, we have two  $O(p^6)$  interactions,  $\epsilon_{abc} W^{a\,\nu}_{\mu} W^{b}_{\nu\rho} W^{c\,\rho\mu}$  and  $\epsilon_{ABC} G^{A\,\nu}_{\mu} G^{B\,\rho\rho}_{\nu\rho} G^{C\,\rho\mu}$ , since terms with only two gauge field-strengths can be eliminated by EOM. In the physical basis these are given by

$$\Delta \mathcal{L}_{\lambda\gamma} = \frac{i\lambda\gamma}{m_W^2} \left[ (eA^{\mu\nu} + gc_{\theta W} Z^{\mu\nu}) W_{\nu}^{-\rho} W_{\rho\mu}^+ \right],$$
  
$$\Delta \mathcal{L}_{3G} = \frac{\kappa_{3G}}{m_W^2} g_s \epsilon_{ABC} G_{\mu}^{A\nu} G_{\nu\rho}^B G^{C\,\rho\mu}.$$
(28)

The former projects onto the TGC observable  $\lambda_{\gamma}$  defined in Ref. [8] and can interfere with the extraction of  $\delta g_1^Z$  and  $\delta \kappa_{\gamma}$  from  $ff \to WW$  [11].

For completeness, we also briefly comment on CPviolating interactions  $\Delta \mathcal{L}_{CPV}$ , beyond those in  $\Delta \mathcal{L}_{4f}$ : these can easily be obtained from the CP-even ones of Eq. (9) and Eq. (23) by  $\delta g_{ff}^{h} \rightarrow i \delta \tilde{g}_{ff}^{h}$ ,  $\delta g_{R}^{W} \rightarrow$  $i \delta \tilde{g}_{R}^{W}$ ,  $\delta \kappa_{q}^{V} \rightarrow i \delta \tilde{\kappa}_{q}^{V}$ , and by substituting one of the field strengths with its dual  $(V_{\mu\nu} \rightarrow \tilde{V}_{\mu\nu})$  in each term of Eqs. (5), (7), (8), (27) and (28).

Having written the BSM effects as interactionterms allows us an easy estimate of their coefficients. The BSM primaries  $\delta g_i$  (as well as 4-fermion interactions) are expected to scale as [16]

$$\frac{\delta g_{ff}^{h}}{Y_{f}} \sim \frac{\delta g_{3h}}{\lambda_{h}v} \sim \frac{\delta g_{VV}^{h}}{gm_{W}} \sim \frac{\delta g_{f,R}^{Z,W}}{g} \sim \delta g_{1}^{Z} \sim \frac{g_{*}^{2}v^{2}}{\Lambda^{2}},$$
(29)

where  $g_*$  denotes a generic BSM coupling. Our operator expansion makes sense if  $g_*^2 v^2 / \Lambda^2 \ll 1$ . On the other hand, the BSM primaries  $\delta \kappa_i$  and  $\lambda_{\gamma}$ , associated to interactions carrying extra derivatives or gauge-fields, are expected to scale as  $g^2 v^2 / \Lambda^2$ , where g is the corresponding SM gauge-coupling. This means that for strongly coupled theories in which  $g_* \sim 4\pi$ , the BSM primaries  $\delta g_i$  are potentially more important. Also in renormalizable weakly-coupled theories  $\delta \kappa_i$  and  $\lambda_{\gamma}$  are suppressed by at least a oneloop factor [16]. In the particular case of TGC/QGC, we have that  $\delta g_1^Z$  could potentially give the most sizeable effects, having an important impact on highenergy processes if  $g_*$  is large [17].

In summary, the Lagrangian up to dimension-6 operators can be written as  $\mathcal{L} = \mathcal{L}_{SM} + \Delta \mathcal{L}_{BSM}$  with

$$\Delta \mathcal{L}_{\text{BSM}} = \Delta \mathcal{L}_{\gamma\gamma}^{h} + \Delta \mathcal{L}_{Z\gamma}^{h} + \Delta \mathcal{L}_{GG}^{h} + \Delta \mathcal{L}_{ff}^{h} + \Delta \mathcal{L}_{3h}^{h} + \Delta \mathcal{L}_{VV}^{h} + \Delta \mathcal{L}_{ee}^{V} + \Delta \mathcal{L}_{qq}^{V} + \Delta \mathcal{L}_{R}^{W} + \Delta \mathcal{L}_{\text{dipole}}^{V} + \Delta \mathcal{L}_{g_{1}^{Z}}^{Z} + \Delta \mathcal{L}_{\kappa\gamma} + \Delta \mathcal{L}_{\lambda\gamma} + \Delta \mathcal{L}_{3G} + \Delta \mathcal{L}_{4f} + \Delta \mathcal{L}_{\text{CPV}}.$$
(30)

The first term of each  $\Delta \mathcal{L}_i$  gives a BSM primary effect which sets, at present or in the near future, the most compelling constraint on the coefficient of  $\Delta \mathcal{L}_i$ . Notice that  $\Delta \mathcal{L}_{BSM}$  only includes interaction terms, as BSM contributions to propagators have been eliminated through the EOM, making the connection between BSM primaries and observables particularly transparent. The main predictions from Eq. (30) are: the Wff and Zff vertices are related (Eqs. (16,17)); the W dipole-type interaction for the fermions are related to those of A and Z (Eq. (23)); there are only 3 types of CP-conserving TGC, characterized by  $\delta g_1^Z$ .  $\delta \kappa_{\gamma}$  and  $\lambda_{\gamma}$  [8], while QGC are related to them by Eqs. (19,28); there are only 8 Higgs BSM primary effects (for one family) [16] while all other Higgs interactions (e.g. effects to  $h \to Vff$  or custodialbreaking hVV couplings) are related to BSM primaries [4, 17].

We are thankful to Eduard Masso, for discussions and precious comments. FR acknowledges support from the Swiss National Science Foundation, under the Ambizione grant PZ00P2 136932. The work of AP has been supported by the grants FPA2011-25948, 2009SGR894 and the Catalan ICREA Academia Program.

- W. Buchmuller and D. Wyler, Nucl. Phys. B 268 (1986) 621.
- [2] B. Grzadkowski, M. Iskrzynski, M. Misiak and J. Rosiek, JHEP **1010** (2010) 085.
- [3] Z. Han and W. Skiba, Phys. Rev. D 71 (2005) 075009.
- [4] A. Pomarol and F. Riva, JHEP 1401, 151 (2014).
- [5] M. E. Peskin and T. Takeuchi, Phys. Rev. Lett. 65 (1990) 964.

- [6] R. Contino *et al.*, JHEP **1307** (2013) 035; I. Brivio *et al.*, JHEP **1403** (2014) 024.
- [7] The ATLAS collaboration, ATLAS-CONF-2014-009.
- [8] K. Hagiwara, R. D. Peccei, D. Zeppenfeld and K. Hikasa, Nucl. Phys. B 282 (1987) 253.
- [9] S. Schael *et al.* [ALEPH and DELPHI and L3 and OPAL and SLD and LEP Electroweak Working Group and SLD Electroweak Group and SLD Heavy Flavour Group Collaborations], Phys. Rept. 427 (2006) 257.
- [10] S. Schael *et al.* [ALEPH and DELPHI and L3 and OPAL and LEP Electroweak Collaborations], Phys. Rept. **532** (2013) 119.

- [11] A. Falkowski et al., Les Houches 2013 Proceedings.
- [12] A. Falkowski et al., in preparation.
- [13] G. Isidori and M. Trott, JHEP **1402** (2014) 082;
   J. Ellis, V. Sanz and T. You, arXiv:1404.3667 [hep-ph].
- [14] R. Barbieri, A. Pomarol, R. Rattazzi and A. Strumia, Nucl. Phys. B 703 (2004) 127.
- [15] J. Reuter, W. Kilian and M. Sekulla, arXiv:1307.8170 [hep-ph].
- [16] J. Elias-Miro, J. R. Espinosa, E. Masso and A. Pomarol, JHEP **1311** (2013) 066.
- [17] R.S. Gupta et al., in preparation.