

# The influence of societal individualism on a century of tobacco use: modelling the prevalence of smoking

John C. Lang<sup>1</sup>, Daniel M. Abrams<sup>2</sup>, and Hans De Sterck<sup>1</sup>

<sup>1</sup>*Department of Applied Mathematics, University of Waterloo*

<sup>2</sup>*Department of Engineering Sciences and Applied Mathematics & Northwestern Institute on Complex Systems & Department of Physics and Astronomy, Northwestern University*

September 5, 2018

## Abstract

Smoking of tobacco is predicted to cause approximately six million deaths worldwide in 2014. Responding effectively to this epidemic requires a thorough understanding of how smoking behaviour is transmitted and modified. Here, we present a new mathematical model of the social dynamics that cause cigarette smoking to spread in a population. Our model predicts that more individualistic societies will show faster adoption and cessation of smoking. Evidence from a new century-long composite data set on smoking prevalence in 25 countries supports the model, with direct implications for public health interventions around the world. Our results suggest that differences in culture between societies can measurably affect the temporal dynamics of a social spreading process, and that these effects can be understood via a quantitative mathematical model matched to observations.

**Keywords:** smoking prevalence, individualism, mathematical modelling

## 1 Introduction and Motivation

In the fifty years since the first report of the Surgeon General’s Advisory Committee on Smoking and Health [1] the smoking epidemic has been responsible for more than 20 million deaths in the United States alone [2, 3], and continues to be responsible for over 6 million deaths worldwide each year [4, 5]. The strong social component of the dynamics of smoking prevalence has been modelled mathematically [6, 7, 8, 9], and examined statistically through analysis of social network data [10] and survey data [11, 12, 13]. However, whereas previous works tend to focus on the micro-level, in this paper we investigate how social aspects of smoking affect its prevalence at the societal level.

Significant inter-country differences exist in smoking prevalence [14]. For example, Fig. 1(a) shows smoking prevalence estimates over most of the past century for Sweden and the USA, obtained from surveys and cigarette consumption data (see Section 3 and Appendix A.1). In both countries, smoking prevalence increased rapidly starting from the early decades of the 20<sup>th</sup> century and reached a peak in the 1960s–1980s era when the adverse health effects of smoking became widely known [1], after which smoking prevalence declined rapidly. However, there are conspicuous differences between the curves: the rate of smoking adoption and cessation before and after the peak is much greater in the US than in Sweden, and the peak in prevalence in the US occurs much earlier than in Sweden.

In this paper we present a new mathematical model of the social spreading of smoking that incorporates the concepts of individual utility (including awareness of health effects), peer influence and social inertia. We propose an interpretation for our model in the context of societal individualism/collectivism and test the model’s predictions and interpretation in three separate phases, see Fig. 2. First, we compile smoking prevalence data spanning the past century for seven OECD countries and find good agreement between this data and the fitted model. Second, the parameter in our model that we interpret as the degree of societal collectivism (see Section 2), and that we fit to smoking prevalence data, is found to be significantly negatively correlated to an established measure of societal individualism (Hofstede’s IDV [15]). Thirdly, the central role played by societal individualism/collectivism in our model motivates us to investigate directly the role that individualism (as measured by Hofstede’s IDV) plays in observed historical tobacco use data. Specifically, we find that IDV is significantly positively correlated to the average rate of increase in smoking prevalence ( $s_x$ ) in seven OECD countries for which smoking prevalence data is available, and that it is significantly negatively correlated to the peak year of tobacco consumption ( $t_{max}$ ) for 25 countries in which tobacco consumption data are available. These findings are interpreted according to our modelling framework, which offers an explanation for the compelling effects of individualism/collectivism on smoking prevalence.

## 2 Model Specification

We begin formulating our model by observing that individuals derive utility from smoking via two mechanisms. First, they derive utility directly from the act of smoking (*individual utility*). Second, they derive utility from social interaction with other smokers (*social utility*). We note that social utility commonly manifests itself in the form of peer influence or peer pressure [16, 17]. We then proceed using a modelling framework that explicitly accounts for the effect of competition between individual and social utilities, and that was first applied to explore the temporal dynamics of language death and religious affiliation as binary choice problems [18, 19]. Specifically, we propose the model

$$\frac{dx}{dt} = b [(1-x) x^a u_x - x (1-x)^a (1-u_x)], \quad (1)$$




Figure 1: Model results and relation with tobacco consumption and individualism data. (a) Output of Eq. (1) (solid line) fitted to historical tobacco use data for the United States (dots) and Sweden (asterisks); US IDV is 91, Swedish IDV is 71. (b) Relative conformity parameter  $a$  versus IDV ( $\rho = -0.87$ ,  $p = 0.011$ ) for seven OECD countries. (c) Average slope  $s_x$  versus IDV ( $\rho = 0.85$ ,  $p = 0.015$ ) for seven OECD countries. (d) Peak year  $t_{max}$  in cigarette consumption versus IDV in a set of 25 countries ( $\rho = -0.524$ ,  $p = 0.008$ ). The seven OECD countries considered for the mathematical model (and displayed in panels (b-c)) are indicated by dots ( $\rho = -0.76$ ,  $p = 0.047$ ), and the remaining 18 countries are indicated by asterisks. For panels (b-d) the line of best fit is given by a solid line. For panel (d) the line of best fit is calculated using data from all 25 countries.




Figure 2: Model specification and testing of model predictions/interpretation in three phases.

where  $x \in [0, 1]$  is the fraction of smokers in the population (i.e., the prevalence),  $u_x \in [0, 1]$  is the individual utility from smoking, and the constant  $b > 0$  determines the timescale of the equation. The interpretation of the positive term in Eq. (1) is, therefore, that non-smokers  $1 - x$  take up smoking at a rate proportional to the *total utility* derived from smoking,  $x^a u_x$ , which is the weighted product of the individual utility from smoking  $u_x$  and the social utility from interactions with other smokers  $x$ , with weighting determined by the constant parameter  $a$ . Since societies with large  $a$  weigh social utility more heavily than individual utility when calculating total utility, we call  $a$  the *relative conformity parameter*. We therefore interpret societies with large  $a$  to be more collectivistic (or less individualistic) than societies with small  $a$ . The interpretation of the negative term in Eq. (1), which models smoking cessation, follows analogously: smokers  $x$  cease smoking at a rate proportional to the total utility derived from non-smoking,  $(1 - x)^a (1 - u_x)$ , which is the weighted product of the individual utility from non-smoking  $u_y = 1 - u_x$  and the social utility from interactions with other non-smokers  $1 - x$ , where we have normalized individual utilities from smoking  $u_x$  and from non-smoking  $u_y$  such that  $u_x + u_y = 1$ . We note that this modelling framework is conceptually consistent with the findings presented in [11]: that personal attitudes about smoking have a stronger influence on smoking behaviour in individualistic countries than in collectivistic countries.

Next, we observe that a combination of factors, including advances in our understanding of the health effects of smoking and public policy initiatives designed to curb smoking, have likely reduced individual utility from smoking ( $u_x$ ) over the past century. Thus, in a significant departure from previous work that treats individual utility as a constant [18, 19], we account for this decline in individual utility by using the cumulative number of scholarly articles on the health effects of smoking ( $n(t)$ ) as a proxy for the reduction in individual utility over the past century. Specifically, following the principle of temporal discounting [20], we assume that each additional article published is discounted by the factor  $\delta \in [0, 1]$  so that for year  $t$

$$u_x(t) = u_\infty + \delta^{n(t)} (u_0 - u_\infty), \quad (2)$$

where  $u_0$  and  $u_\infty$  are the limiting individual utilities from smoking when there is no knowledge and perfect knowledge of the adverse effects of smoking, respectively. Here,  $u_0$ ,  $u_\infty$  and  $\delta$  are parameters to be fitted to observational data. Fig. A.2 shows the cumulative number of scholarly articles on the health effects of smoking  $n(t)$  (see Appendix A.2 and ESM data file *lang\_supporting\_data1.txt*), and the resulting individual utility from smoking  $u_x(t)$ , as defined in Eq. (2), for various values of  $\delta$ . We remark that this approach leads to better fits between model output and observational data than alternatives that do not directly take into account the effect of increased scientific understanding of health effects (see Appendices A.3 and A.4).

### 3 Data

We note that Eq. (1) subject to Eq. (2) requires the fitting of four parameters per country ( $x_0 = x(t_0)$ ,  $a$ ,  $u_0$ , and  $u_\infty$ ) and two parameters  $b$  and  $\delta$  that we take equal for all countries in the data set (see Appendix A.3). We determine these parameters by fitting them to historical smoking prevalence data. Such historical data (mostly obtained by surveys) is available for a set of 24 OECD countries, but unfortunately is limited to, on average, only 21.5 observations over a period of 31.4 years spanning 1960–2012 [21]. As such, it misses much of the crucial period in the earlier parts of the 20th century during which smoking steadily gained popularity in many countries. However, historical national cigarette consumption data is available for the same 24 OECD countries plus Romania for an average of 78.4 observations over a period of 82.2 years spanning 1900–2012 [22, 23]. In order to exploit the more abundant consumption data in our prevalence model, we assume a linear relationship between national cigarette consumption and smoking prevalence for all times where both measurements are available. We then apply this relationship to the full cigarette consumption data set to produce an estimate of historical smoking prevalence ( $\hat{x}$ ) over nearly the entire past century. Sufficient data for this procedure is available in seven OECD countries (for additional details, see Appendix A.1): Australia, Canada, France, New Zealand, Sweden, the United Kingdom, and the United States.

## 4 Results: Testing the Model

### 4.1 Phase (i): Direct test

Figure 1(a) shows the fit of our model to data sets from the United States and Sweden (additional fits and parameter values are displayed in Fig. A.3 and Table A.3). The good agreement that we found with all data sets provides support for the model.

### 4.2 Phase(ii): Test of model implications for $a$

If the model and its interpretation are correct, then we expect that the fitted “relative conformity parameter”  $a$  will capture something meaningful about the individualism/collectivism of a society. To test this, we compare with Hofstede’s IDV, an established metric for societal individualism [15] that has been evaluated in most countries. Panel (b) of Fig. 1 shows the comparison. As expected, the relative conformity parameter  $a$  shows significant negative correlation with Hofstede’s IDV (negative because  $a$  increases with collectivism while IDV decreases with it). This concordance with independently assessed individualism values supports our model.

### 4.3 Phase (iii): Test of model implications for slope and peak year

Besides the correlation of  $a$  with collectivism, we note that another prediction is implicit in model (1). As the relative conformity parameter increases, the model requires that changes in smoking prevalence occur more slowly (this is true for the range of  $a$  and  $u$  values corresponding to the observational data). Put another way, societies with higher levels of individualism should experience faster changes in smoking prevalence. Intuitively, when smoking prevalence is low the lack of existing smokers inhibits smoking initiation more strongly in a collectivistic society than in an individualistic society. Thus, we expect the average rate of increase in a collectivistic society to be smaller than in an individualistic society. In contrast, when smoking prevalence is high, and once the deleterious health effects of smoking become widely known and negatively impact individual utility from smoking, the presence of existing smokers inhibits smoking cessation more strongly in a collectivistic society than in an individualistic society. In both cases collectivism acts as a brake on change in the status quo (higher cultural inertia [24, 25]). Panel (c) of Fig. 1 demonstrates that this is indeed the case: the average slope  $s_x$  of the smoking prevalence curves leading up to the peak (see Eq. (A.1) of Appendix A.5) increases with Hofstede’s IDV (shown) and decreases with  $a$  (see Fig. A.4(a)). Correlations are significant (see Table A.5).

This reasoning further suggests that the peak year for smoking prevalence  $t_{max}$  should be later in collectivistic societies and earlier in individualistic countries. As shown in Fig. 1(d), the raw observational data are consistent with this prediction:  $t_{max}$  is significantly negatively correlated with IDV (shown) and significantly positively correlated with  $a$  (see Fig. A.4(b)). Note that our assumption of a linear relationship between national cigarette consumption and smoking prevalence is not needed to establish  $t_{max}$ , so Fig. 1(d) is independent of any model assumptions.

## 5 Discussion and Conclusion

We have proposed a quantitative mathematical model of the social spreading of smoking that is derived from basic principles well-documented in the sociology and social psychology literature. The model appears to match real-world smoking prevalence data from a variety of countries well (to our knowledge, the largest historical data set of this type ever compiled), and all predictions of the model appear to be supported by the data. In particular, the model predicts that the level of individualism or collectivism of a society may significantly affect the temporal dynamics of smoking prevalence: the strong influence of the personal utility of smoking (and its decrease due to increased awareness of adverse health effects) is predicted to lead to faster adoption and cessation of smoking in individualistic societies than in more collectivistic societies.

It has previously been argued that social support mechanisms in collectivistic societies make it more likely that a person will stop smoking [26, 27] based on findings that social support (supportive counselors) can help people to

adhere to decisions to quit smoking [13]. We find that, to the contrary, cessation of smoking occurs more slowly in collectivistic societies. Our model suggests that this is so because social inertia will inhibit decisions to stop smoking more strongly in collectivistic societies than in individualistic societies.

These results have significant implications for combating the ongoing smoking epidemic. For example, they imply that interventions designed to discourage smoking should be tailored differently in societies or social groups whose cultures differ in how they value individualism versus collectivism [28]. More broadly, these results demonstrate that differences in culture can measurably affect the dynamics of a social spreading process, and that a mathematical model can help to illuminate this phenomenon.

Despite the good match between model predictions and data, a number of limitations remain. We have made an implicit “mean-field” approximation in taking social utility to be a function of the overall smoking prevalence  $x$ , rather than the local smoking prevalence among contacts in an individual’s social network. Similarly, we have taken individual utility to be uniform across the population (though not in time), whereas a more detailed model might allow for individual variation. As a mild justification for these assumptions, we point out that analysis of a similar model in another context [19] suggests that inclusion of more detail will not change qualitative predictions.

We claim that the correlation of individualism with faster societal change results from a causative influence as predicted by our model. Other factors such as income levels also correlate with individualism, and it’s possible that what we observe is ultimately also related to GDP or other variables. We certainly cannot exclude that there may be other causative factors. For example, our model in its current form is incapable of explaining differences in smoking prevalence between genders and why these inter-gender differences vary between countries [14]. Nevertheless, we remark that many previously proposed causative factors for differences in observed inter-country smoking dynamics can be accounted for within our modelling framework. In particular, beliefs about the harmful effects of smoking, the price of cigarettes, socioeconomic status and inequality, and government regulation have all been cited as potential factors affecting the differences observed in inter-country smoking dynamics [14]. Each of these factors can be interpreted within our modelling framework. For example, beliefs about the harmful effects of smoking, as well as the price of cigarettes, both likely contribute directly to individual utility derived from smoking ( $u_x$ ) and from non-smoking ( $u_y$ ). Moreover, socioeconomic status may affect individual utility from smoking indirectly by affecting an individual’s tolerance for risk and/or how they discount future rewards and costs (i.e. how they discount their future health status) [29]. Addressing the model’s inability to account for gender differences in smoking prevalence and quantifying the relationship between other causative factors and model parameters are potential areas for future work.

We also welcome future work comparing a variety of social contagion phenomena across societies. Our model suggests that the increased cultural inertia in collectivistic societies would lead to slower change across a wide spectrum of spreading processes (those where important changes occur in personal utility), a hypothesis that could



be supported or rejected by further study.

## Acknowledgments

We would like to thank Dr. James Fowler for insightful remarks on an early draft of this manuscript. DMA thanks the James S. McDonnell Foundation for support through grant #220020230. JL and HDS acknowledge support from NSERC of Canada.

## A Appendix - Materials and Methods

### A.1 Smoking prevalence and cigarette consumption data

We consider smoking prevalence  $x(t) \in [0, 1]$  for 24 OECD countries which we download from the OECD iLibrary online statistical database [21] in Excel format. We also consider manufactured cigarette consumption (in grams) per person per day  $c(t)$  for the same 24 OECD countries plus Romania (which is a non-OECD country) [22, 23]. When available, cigarette consumption data is downloaded directly from the International Smoking Statistics (Web Edition) website [22] in Excel format. Cigarette consumption data for countries not included in the International Smoking Statistics (Web Edition) are retrieved from the International Smoking Statistics (2<sup>nd</sup> Ed.) [23] by manually transferring these entries into Excel. We make these data available in CSV format in the ESM data file *lang\_supplementary\_data1.txt*, which contains four columns: country number as it appears in Table A.1, year ( $t$ ), measurement ( $x(t)$  or  $c(t)$ ), and type of measurement (0 indicates a smoking prevalence measurement, while 1 indicates a cigarette consumption measurement).

Recall that the cigarette consumption data are more dense and span a larger time interval than smoking prevalence data, see Table A.1. Thus, since our model is specified in terms of smoking prevalence, we estimate smoking prevalence from cigarette consumption in order to exploit the much richer cigarette consumption data for model fitting purposes. First, we assume a linear relationship between smoking prevalence  $x(t)$  and smoking consumption  $c(t)$

$$x(t) = Cc(t) + B.$$

Next, we calculate estimates  $\hat{C}$  and  $\hat{B}$  by regressing smoking prevalence  $x(t)$  on tobacco consumption  $c(t)$  for all years for which both measurements are available. The results of this regression are summarized in Table A.2, which illustrates that the assumption that  $x$  and  $c$  are linearly related does not hold equally well for all countries. In order to restrict ourselves to the cases where the assumption of linearity between  $x$  and  $c$  is valid we restrict ourselves to the seven OECD countries with  $R^2 \geq 0.7$ ,  $p < 0.001$ , and  $n_{obs} \geq 15$ : Australia, Canada, France, New Zealand,

Table A.1: Summary of data on smoking prevalence  $x$  and cigarette consumption (in grams) per person per day  $c$

No.	Country	Abbrev.	Smoking prevalence ( $x$ )		Cigarette consumption per person per day ( $c$ )		Source
			Obs. Period	No. of Obs.	Obs. Period	No. of Obs.	
1	Australia	AUS	1964–2010	16	1920–2010	91	[22]
2	Austria	AUT	1972–2006	5	1923–2004	82	[22]
3	Belgium	BEL	1997–2008	4	1921–2011	91	[22]
4	Canada	CAN	1964–2011	29	1920–2010	91	[22]
5	Denmark	DNK	1970–2010	41	1920–2010	91	[22]
6	Finland	FIN	1978–2011	34	1920–2009	90	[22]
7	France	FRA	1960–2010	22	1900–2010	93	[22]
8	Greece	GRE	1998–2009	6	1920–1995	76	[23]
9	Hungary	HUN	1994–2009	4	1920–2012	87	[22]
10	Iceland	ICE	1987–2012	26	1932–1995	64	[23]
11	Ireland	IRE	1973–2007	14	1920–1995	76	[23]
12	Israel	ISR	1996–2010	8	1967–1995	29	[23]
13	Italy	ITA	1980–2012	23	1905–2010	73	[22]
14	Japan	JPN	1965–2011	47	1920–2007	88	[22]
15	Netherlands	NLD	1966–2011	39	1923–1995	67	[23]
16	New Zealand	NZL	1976–2012	28	1920–2009	90	[22]
17	Norway	NOR	1973–2012	40	1927–2011	85	[22]
18	Poland	POL	1996–2009	4	1925–1995	43	[23]
19	Portugal	PRT	1987–2006	4	1940–1995	56	[23]
20	Romania	ROM	–	0	1920–1995	52	[23]
21	Spain	SPA	1985–2011	11	1920–2010	87	[22]
22	Sweden	SWE	1980–2011	32	1920–2006	87	[22]
23	Switzerland	CHE	1992–2007	4	1934–2009	76	[22]
24	United Kingdom	GBR	1960–2010	38	1905–2009	105	[22]
25	United States	USA	1965–2011	36	1920–2010	91	[22]

Sweden, the United Kingdom, and the United States. We display the raw data for these seven OECD nations in Fig. A.1. The smoking prevalence for these seven OECD countries is then estimated from tobacco consumption using the relationship

$$\hat{x}(t) = \hat{C}c(t) + \hat{B}.$$

Table A.2: Estimates  $\hat{C}$ ,  $\hat{B}$

Country	$\hat{C} \times 10^2$	$\hat{B} \times 10^2$	$R^2$	$p$	$n_{obs}$
Australia	$4.5 \pm 1.3$	$-0.3 \pm 8.8$	0.80	$3.2 \times 10^{-6}$	16
Austria	$0.0 \pm 4.9$	$24.2 \pm 32.4$	0.00	0.99	4
Belgium	$2.6 \pm 20.3$	$13.0 \pm 81.5$	0.13	0.64	4
Canada	$3.5 \pm 0.5$	$6.3 \pm 3.8$	0.87	$3.0 \times 10^{-13}$	28
Denmark	$0.0 \pm 9.2$	$40.5 \pm 44.4$	0.00	0.99	41
Finland	$2.0 \pm 0.7$	$15.8 \pm 2.8$	0.55	$1.0 \times 10^{-6}$	32
France	$1.8 \pm 0.5$	$19.1 \pm 2.5$	0.72	$6.3 \times 10^{-7}$	22
Greece	—	—	—	—	0
Hungary	$1.9 \pm 1.6$	$17.4 \pm 11.2$	0.93	$3.5 \times 10^{-2}$	4
Iceland	$4.9 \pm 1.2$	$0.9 \pm 7.0$	0.93	$2.6 \times 10^{-5}$	9
Ireland	$5.4 \pm 1.1$	$-4.0 \pm 7.4$	0.93	$1.7 \times 10^{-6}$	11
Israel	—	—	—	—	—
Italy	$4.8 \pm 2.5$	$-0.3 \pm 13.2$	0.47	$6.1 \times 10^{-4}$	21
Japan	$1.3 \pm 3.2$	$25.7 \pm 27.2$	0.02	0.43	43
Netherlands	$4.8 \pm 3.2$	$20.5 \pm 15.0$	0.32	$4.7 \times 10^{-3}$	23
New Zealand	$2.0 \pm 0.3$	$18.8 \pm 1.4$	0.86	$2.6 \times 10^{-12}$	27
Norway	$-7.2 \pm 4.3$	$50.1 \pm 10.6$	0.24	$1.6 \times 10^{-3}$	39
Poland	—	—	—	—	0
Portugal	—	—	—	—	1
Romania	—	—	—	—	0
Spain	$6.0 \pm 6.2$	$-7.4 \pm 41.7$	0.38	$5.7 \times 10^{-2}$	10
Sweden	$5.4 \pm 0.6$	$4.3 \pm 2.3$	0.92	$1.7 \times 10^{-15}$	27
Switzerland	$2.8 \pm 5.6$	$7.2 \pm 38.6$	0.69	0.17	4
United Kingdom	$5.6 \pm 0.7$	$1.6 \pm 4.5$	0.88	$5.3 \times 10^{-18}$	37
United States	$3.6 \pm 0.3$	$-0.1 \pm 2.3$	0.95	$1.1 \times 10^{-22}$	35

$\pm$  indicates 95% confidence intervals. We report  $R^2$  values for the linear regression of  $x$  on  $c$ , the  $p$ -value of the correlation between  $x$  and  $c$ , and the number of years for which both  $x$  and  $c$  measurements are available,  $n_{obs}$ .

Note that survey-based prevalence data are susceptible to noise stemming from variations in the survey methodology. In particular, prior to performing the linear regression of  $x$  on  $c$  for France, we removed the outlier  $x(1960) = 0.32$  since it is inconsistent with the rest of the data for France, see Fig. A.1(c). Specifically, the Grubbs test on  $x/\hat{x}$  indicates that the 1960 data point is a significant outlier ( $p < 0.05$ ). This can also be seen intuitively: from  $t = 1960$  until the next measurement at  $t = 1965$  smoking prevalence drops from  $x(1960) = 0.32$  to  $x(1965) = 0.25$  (a decrease of 21.9%), while cigarette consumption steadily increases from  $c(1960) = 3.6$  to  $c(1965) = 4.1$  (an increase of 13.9%). Given the population in France in 1960 (45.5 million) and in 1965 (48.6 million) [30], this would correspond to an increase in the average mass of cigarettes smoked (in grams) per smoker per day from 11.3 to 16.4 (an increase of




Figure A.1: Raw smoking prevalence ( $x$ ) and raw cigarette consumption ( $c$ ) data. (Left axis - blue asterisks) Raw smoking prevalence data and (right axis - black dots) raw cigarette consumption data (in grams per person per day). A single outlier for smoking prevalence ( $x$ ) for the country of France (panel c) is denoted with a red asterisk.

45.1%) over a short 5 year period. This is in sharp contrast with the relatively stable relationship between  $x$  and  $c$  for France’s remaining data points and justifies the exclusion of the outlier  $x(1960) = 0.32$ . With the outlier removed, France satisfies our data quality requirements for inclusion in the set of seven OECD countries ( $R^2 \geq 0.7$ ,  $p < 0.001$ , and  $n_{obs} \geq 15$ ).

## A.2 Proxy data $n(t)$ : articles published on the health effects of smoking

In order to implement Eq. (1) subject to Eq. (2) we require data on the cumulative number of articles published on the health effects of smoking  $n(t)$ . We calculate  $n(t)$  by performing a search of the online research database Scopus for papers with

- (i) tobacco, smok\*, or cigar\* in the title, and
- (ii) death, illness, mortality, risk\*, tumour\*, tumor\*, or cancer in the title, and
- (iii) medicine, dentistry, nursing, veterinary, health professions, or multidisciplinary in the subject area, and
- (iv) plant\*, mosaic, botany, smog, fog, and soot not in the title.

Items (i)-(iii) are search terms included in order to select for papers researching the health effects of smoking, whereas items (iv) are search terms excluded in order to prevent selection of papers researching the tobacco mosaic virus (plant\*, mosaic, botany) and the health effects of atmospheric smoke (smog, fog, soot). This provides us with  $n(t)$  for integer  $t$ , where time  $t$  is measured in years. We make the article data available in CSV format in the ESM data file, *lang\_supplementary\_data2.txt*, which contains three columns: year ( $t$ ), number of articles published in year  $t$ , and cumulative number of articles published up to and including year  $t$  ( $n(t)$ ). To calculate  $n(t)$  for non-integer and missing values of  $t$  we use linear interpolation, see Fig. A.2(a). Furthermore, Fig. A.2(b) displays  $u_x(t)$  from Eq. (2) using  $n(t)$  calculated above for various discount factors  $\delta$  and with  $u_0 = 0.51$  and  $u_\infty = 0.49$ . (For comparison, see Table A.3 for model-fitted values of  $\delta$ ,  $u_0$  and  $u_\infty$ .)

## A.3 Model fitting

We fit Eq. (1) to the estimated prevalence,  $\hat{x}(t)$ . To reduce the dimensionality of the optimization problem, we assume that certain *universal parameters* are constant across countries. Specifically, we assume that  $b$  and  $\delta$  are universal parameters, and that  $x_i(t_{i,0}) = x_{i,0}$ ,  $a_i$ ,  $u_{i,0}$ , and  $u_{i,\infty}$  are *local parameters* for country  $i$ , where  $t_{i,0}$  is the first year for which cigarette consumption data ( $c$ ), and hence estimated smoking prevalence data ( $\hat{x}$ ), are available. We denote the smoking prevalence estimated above for country  $i$  at time  $t$  by  $\hat{x}_i(t)$ . The time series of estimated smoking prevalences for country  $i$  is then denoted by the vector  $\widehat{X}_i$ . Analogously, we denote the time series of smoking




Figure A.2: Articles published on the health effects of smoking. Articles retrieved by Scopus with search terms (i)-(iv) and individual utility profiles from Eq. (2) for varying values of  $\delta$ . (a) (Left axis - blue, solid) Number of articles published per year and (right axis - black, dashed) cumulative number of articles published  $n(t)$ . (b) Discounted utility  $u_x(t)$  from Eq. (2) with  $u_0 = 0.51$  and  $u_\infty = 0.49$ , using cumulative number of articles published  $n(t)$ . (Solid black)  $\delta = 0.4$ , (dashed black)  $\delta = 0.6$ , (dotted black)  $\delta = 0.8$ , (solid blue)  $\delta = 0.9$ , (dashed blue)  $\delta = 0.99$ , (dotted blue)  $\delta = 0.998$ , and (solid red)  $\delta = 0.9995$ .

prevalences predicted by Eq. (1) for country  $i$  by  $\tilde{X}_i$ . We solve Eq. (1) using the Matlab differential equation solver *ode45*.

Using the Matlab function *lsqcurvefit* we now proceed as follows:

1. Holding universal parameters constant, for each country  $i$  we find the  $x_{i,0}$ ,  $a_i$ ,  $u_{i,0}$ , and  $u_{i,\infty}$  that minimize

$$E_i = \|\tilde{X}_i - \hat{X}_i\|_2^2.$$

2. Holding local parameters constant for each country  $i$ , we find the  $b$  and  $\delta$  that minimize  $E = \sum_i \|\tilde{X}_i - \hat{X}_i\|_2^2$ .

3. Repeat steps (1) and (2) until either

(a) the change in the objective function  $\sum_i \|\tilde{X}_i - \hat{X}_i\|_2^2$  is below tolerance  $tol$ , or

(b) the number of iterations exceeds a limit  $max_{itn}$ .

We perform the optimization with the initial guess  $u_{i,0} \equiv 0.51$ ,  $u_{i,\infty} \equiv 0.49$ ,  $x_{i,0} = \hat{x}_i(t_{i,0})$ ,  $a_i = 1$ ,  $b = 1$ , and  $\delta = 0.9985$ . We also provide the optimization algorithm *lsqcurvefit* with constraints

$$0 \leq a_i, b \leq 2 \text{ and}$$

$$0 \leq x_{i,0}, u_{i,0}, u_{i,\infty}, \delta \leq 1,$$

and with parameters  $tol = 10^{-6}$  and  $max_{itn} = 150$ . The fitting procedure terminates after 114 iterations, the results of which are recorded in Table A.3 and Fig. A.3. Complete model simulation code with all necessary data files are

available upon request.

Table A.3: The result of fitting Model Eq. (1) to the estimated smoking prevalence  $\hat{x}$

Universal parameters and Total Error ( $E$ )		
$b$	$\delta$	$E$
1.049	0.9981	0.163

Country ( $i$ )	Local parameters and local error ( $E_i$ )				
	$a_i$	$x_{i,0}$	$u_{i,0}$	$u_{i,\infty}$	$E_i$
Australia	1.035	0.033	0.551	0.484	0.032
Canada	1.020	0.083	0.530	0.483	0.020
France	1.121	0.198	0.543	0.524	0.004
New Zealand	1.062	0.202	0.525	0.504	0.012
Sweden	1.076	0.077	0.555	0.503	0.015
United Kingdom	0.976	0.079	0.513	0.478	0.060
United States	0.963	0.063	0.513	0.470	0.024

#### A.4 Alternative to temporal discounting of individual utility

We have found in more detailed analysis that using the discounting utility formula of Eq. (2) produces a better fit (significantly lower total error  $E$ ) than alternatives, e.g. constant utility  $u_{i,x}(t) \equiv u_{i,x}$  or step-function utility

$$u_{i,x}(t) = \begin{cases} u_{i,0} & \text{if } t < t_i^* \\ u_{i,\infty} & \text{if } t \geq t_i^* \end{cases},$$

where  $t_i^*$  is a threshold parameter whose value is determined by the fitting procedure. This is consistent with the expectation that increasing knowledge of health effects has indeed influenced the individual utility from smoking over the past century.

#### A.5 Calculation of average slope

We estimate the average slope of the estimated prevalence versus time curve in the period leading up to its peak as:

$$s_x = \frac{\hat{x}(t_{max}) - \hat{x}(t_0)}{t_{max} - t_0}, \quad (\text{A.1})$$

where  $t_0 = 1920$  is the first year for which smoking prevalence estimates are available in the subset of seven OECD countries, and where  $t_{max}$  are recorded in Table A.4. Correlation between fitted relative conformity parameter  $a$  and average slope  $s_x$  (and peak year  $t_{max}$ ) is displayed in Fig. A.4 and recorded in Table A.5.




Figure A.3: The result of fitting Eq. (1) to the estimated smoking prevalence  $\hat{x}$ . Estimated smoking prevalence values  $\hat{x}$  are given by blue dots.



Table A.4: Hofstede's Individualism Index IDV and peak year in cigarette consumption ( $t_{max}$ )

Country	IDV	Peak year ( $t_{max}$ )
Australia	90	1974
Austria	55	1979
Belgium	75	1973
Canada	80	1976
Denmark	74	1976
Finland	63	1963
France	71	1985
Greece	35	1986
Hungary	80	1980
Iceland	60	1984
Ireland	70	1974
Israel	54	1974
Italy	76	1984
Japan	46	1977
Netherlands	80	1977
New Zealand	79	1975
Norway	69	2004
Poland	60	1991
Portugal	27	1994
Romania	30	1995
Spain	51	1985
Sweden	71	1976
Switzerland	68	1972
United Kingdom	89	1973
United States	91	1963




Figure A.4: Average slope  $s_x$  and peak year of smoking prevalence  $t_{max}$  versus relative conformity parameter  $a$  for seven OECD countries. (a) Average slope  $s_x$  versus relative conformity parameter  $a$  ( $\rho = -0.92$ ,  $p = 0.003$ ). (b) Peak year  $t_{max}$  versus relative conformity parameter  $a$  ( $\rho = 0.88$ ,  $p = 0.009$ ). The line of best fit is given by a solid line.

Table A.5: Individualism index IDV and relative conformity  $a$  are significantly correlated with and  $a$ ,  $s_x$ , and  $t_{max}$

	7-country subset			25-country set
	$a$	$s_x$	$t_{max}$	$t_{max}$
IDV	-0.87 (0.011)	0.85 (0.015)	-0.76 (0.047)	-0.53 (0.006)
$a$	–	-0.92 (0.003)	0.88 (0.009)	–

Correlations between IDV,  $a$ ,  $s_x$ , and  $t_{max}$  are recorded for the seven-country subset. Correlation between IDV and  $t_{max}$  is recorded for the full set of 25 countries.  $p$ -values are in parentheses. All correlations are significant at the 95% confidence level.

## References

- [1] US Department of Health, Education, and Welfare. Smoking and Health: Report of the Advisory Committee to the Surgeon General of the Public Health Service; 1964.
- [2] US Department of Health and Human Services. The Health Consequences of Smoking—50 Years of Progress: A Report of the Surgeon General; 2014.
- [3] Warner KE. Cigarette smoking in the 1970s: the impact of the antismoking campaign on consumption. *Science*. 1981;211:729–731.
- [4] Jha P. Avoidable global cancer deaths and total deaths from smoking. *Nat Rev Cancer*. 2009;9:655–664.
- [5] Mathers CD, Loncar D. Projections of global mortality and burden of disease from 2002 to 2030. *PLoS Med*. 2006;3:e442.

- [6] Cavana RY, Tobias M. Integrative System Dynamics: Analysis of Policy Options for Tobacco Control in New Zealand. *Syst Res.* 2008;25:675–694.
- [7] Sharomi O, Gumel AB. Curtailing smoking dynamics: A mathematical modeling approach. *Appl Math Comput.* 2008;195:475–499.
- [8] Levy DT, Bauer JE, Lee H. Simulation Modeling and Tobacco Control: Creating More Robust Public Health Policies. *Am J Public Health.* 2006;96:494–498.
- [9] Rowe DC, Chassin L, Presson CC, Edwards D, Sherman SJ. An “Epidemic” Model of Adolescent Cigarette Smoking. *J Appl Soc Psychol.* 1992;22:261–285.
- [10] Christakis NA, Fowler JH. The Collective Dynamics of Smoking in a Large Social Network. *N Engl J Med.* 2008;358:2249–2258.
- [11] Hosking W, Borland R, Yong HH, Fong G, Zanna M, Laux F, et al. The effects of smoking norms and attitudes on quitting intentions in Malaysia, Thailand and four Western nations: a cross-cultural comparison. *Psychol Health.* 2009;24:95–107.
- [12] Unger JB, Rorhbach LA, Cruz TB, Baezconde-Garbanati L, Howard KA, Palmer PH, et al. Ethnic variation in peer influences on adolescent smoking. *Nicotine Tob Res.* 2001;3:167–76.
- [13] Janis IL. The Role of Social Support in Adherence to Stressful Decisions. *Am Psychol.* 2001;38:143–160.
- [14] Pampel F. Divergent Patterns of Smoking Across High-Income Nations. In: Crimmins EM, Preston SH, Cohen B, editors. *International Differences in Mortality at Older Ages: Dimensions and Sources.* Washington D.C.: National Academies Press; 2010. p. 132–163.
- [15] Hofstede G, Hofstede GJ, Minkov M. I, We, and They. In: *Cultures and organizations: software of the mind: intercultural cooperation and its importance for survival.* 3rd ed. New York: McGraw Hill; 2010. p. 89–134.
- [16] Calavó-Armengol A, Jackson MO. Peer Pressure. *J Eur Econ Assoc.* 2010;8:62–89.
- [17] Centola D, Willer R, Macy M. The Emperor’s Dilemma: A Computational Model of Self-Enforcing Norms. *Am J Sociol.* 2005;110:1009–1040.
- [18] Abrams DM, Strogatz SH. Modelling the dynamics of language death. *Nature.* 2003;424:900.
- [19] Abrams DM, Yaple HA, Wiener RJ. Dynamics of social group competition: modeling the decline of religious affiliation. *Phys Rev Lett.* 2011;107:088701.

- [20] Frederick S, Lowenstein G, O'donoghue T. Time discounting and time preference: A critical review. *J Econ Lit.* 2002;40:351–401.
- [21] OECD. OECD.Stat, Health: Non-Medical Determinants of Health: Tobacco consumption. OECDStat database. 2013;Accessed October 24, 2013.
- [22] Forey B, Hamling J, Hamling J, Thorton A, Lee P. International Smoking Statistics: A collection of worldwide historical data. Web ed. 17 Cedar Road, Sutton, SM2 5DA, UK: P N Lee Statistics & Computing Ltd; 2013. Accessed October 7, 2013. Available from: <http://www.pnlee.co.uk/ISS.htm>.
- [23] Forey B, Hamling J, Lee P, Wald N. International Smoking Statistics: A collection of historical data from 30 economically developed countries. 2nd ed. Great Clarendon Street, Oxford, OX2 6DP, UK: Oxford University Press; 2002.
- [24] Zárata MA, Shaw M, Marquez JA, Biagas D. Cultural inertia: The effects of cultural change on intergroup relations and the self-concept. *J Exp Soc Psychol.* 2012;48:634–645.
- [25] Henrich J, McElreath R. The Evolution of Cultural Evolution. *Evol Anthr.* 2003;12:123–135.
- [26] Sun T, Horn M, Merritt D. Values and lifestyles of individuals and collectivists: a study on Chinese, Japanese, British, and US consumers. *J Consum Marketing.* 2004;21:318–331.
- [27] Triandis HC, Bontempo R, Villareal MJ. Individualism and Collectivism: Cross-Cultural Perspectives on Self-Group Relationships. *J Pers Soc Psychol.* 1988;54:323–338.
- [28] Pentecostes JU. Individualism vs Collectivism: Implications for Health Promotion. *Phillippine J Psychol.* 1999;32:397–402.
- [29] Griskevicius V, Tybur JM, Delton AW, Robertson TE. The Influence of Mortality and Socioeconomic Status on Risk and Delayed Rewards: A Life History Theory Approach. *J Pers Soc Psychol.* 2011;100:1015–1026.
- [30] OECD. OECD.Stat, Demography and Population: Population Statistics: Population and Vital Statistics. OECDStat database. 2013;Accessed January 20, 2014.