LIE GROUPS AS 3-DIMENSIONAL ALMOST CONTACT B-METRIC MANIFOLDS

HRISTO MANEV, DIMITAR MEKEROV

ABSTRACT. The object of investigations are almost contact B-metric structures on 3-dimensional Lie groups considered as smooth manifolds. There are established the existence and some geometric characteristics of these manifolds in all basic classes. An example is given as a support of obtained results.

INTRODUCTION

The differential geometry of the almost contact metric manifolds is well studied (e.g. [2]). The beginning of the investigations on the almost contact manifolds with B-metric is given by [5]. These manifolds are the odd-dimensional counterpart of the almost complex manifolds with Norden metric [4, 6], where the almost complex structure acts as an anti-isometry on the metric. Almost contact B-metric manifolds are investigated and studied in [5, 12, 13, 14, 15, 16, 17, 19].

An object of special interest is the case of the lowest dimension of the considered manifolds. Curvature properties of the 3-dimensional manifolds of this type are studied in [11, 18, 20].

The basic problem considered in this work is the existence and the geometric characteristics of almost contact B-metric structures on 3-dimensional Lie algebras. The main result is a construction of the considered structures and their characterization in all basic classes of the almost contact B-metric manifolds in the classification given in [5].

The present paper is organized as follows. In Sect. 1 we recall some facts about the almost contact B-metric manifolds. In Sect. 2 we construct different types of contact structures on Lie algebras in dimension 3. In Sect. 3 we consider an example in relation with the investigations in the previous section.

1. Preliminaries

1.1. Almost contact manifolds with B-metric. Let $(M, \varphi, \xi, \eta, g)$ be an almost contact manifold with B-metric or an *almost contact B-metric manifold*, i.e. M is a (2n + 1)-dimensional differentiable manifold with an almost contact structure (φ, ξ, η) consisting of an endomorphism φ of the tangent bundle, a vector field ξ ,

¹⁹⁹¹ Mathematics Subject Classification. 53C15, 53C50, 53D15.

 $Key\ words\ and\ phrases.$ Almost contact structure, B-metric, Lie group, Lie algebra, indefinite metric.

its dual 1-form η as well as M is equipped with a pseudo-Riemannian metric g of signature (n + 1, n), such that the following algebraic relations are satisfied [5]:

(1)
$$\begin{aligned} \varphi \xi &= 0, \qquad \varphi^2 = -\operatorname{Id} + \eta \otimes \xi, \qquad \eta \circ \varphi = 0, \qquad \eta(\xi) = 1, \\ g(\varphi x, \varphi y) &= -g(x, y) + \eta(x)\eta(y), \end{aligned}$$

where Id is the identity and x, y are elements of the algebra $\mathfrak{X}(M)$ of the smooth vector fields on M.

Further x, y, z will stand for arbitrary elements of $\mathfrak{X}(M)$ or vectors in the tangent space T_pM of M at an arbitrary point p in M.

The associated metric \tilde{g} of g on M is defined by $\tilde{g}(x, y) = g(x, \varphi y) + \eta(x)\eta(y)$. The manifold $(M, \varphi, \xi, \eta, \tilde{g})$ is also an almost contact B-metric manifold. The signature of both metrics g and \tilde{g} is necessarily (n + 1, n). By ∇ and $\tilde{\nabla}$ we denote the Levi-Civita connection of g and \tilde{g} , respectively.

A classification of almost contact B-metric manifolds is given in [5]. This classification, consisting of eleven basic classes $\mathcal{F}_1, \mathcal{F}_2, \ldots, \mathcal{F}_{11}$, is made with respect to the tensor F of type (0,3) defined by

$$F(x, y, z) = g((\nabla_x \varphi) y, z).$$

The following properties are valid:

$$F(x, y, z) = F(x, z, y) = F(x, \varphi y, \varphi z) + \eta(y)F(x, \xi, z) + \eta(z)F(x, y, \xi).$$

The intersection of the basic classes is the special class \mathcal{F}_0 determined by the condition F(x, y, z) = 0. Hence \mathcal{F}_0 is the class of almost contact B-metric manifolds with ∇ -parallel structures, i.e. $\nabla \varphi = \nabla \xi = \nabla \eta = \nabla g = \nabla \tilde{g} = 0$.

If $\{e_i; \xi\}$ (i = 1, 2, ..., 2n) is a basis of $T_p M$ and (g^{ij}) is the inverse matrix of (g_{ij}) , then with F are associated 1-forms θ , θ^* , ω , called *Lee forms*, defined by:

$$\theta(z) = g^{ij}F(e_i, e_j, z), \quad \theta^*(z) = g^{ij}F(e_i, \varphi e_j, z), \quad \omega(z) = F(\xi, \xi, z).$$

Further, we use the following characteristic conditions of the basic classes [9]:

$$\begin{array}{rcl} \mathcal{F}_{1}:& F(x,y,z) = \frac{1}{2n} \Big\{ g(x,\varphi y) \theta(\varphi z) + g(\varphi x,\varphi y) \theta(\varphi^{2}z) \\ & + g(x,\varphi z) \theta(\varphi y) + g(\varphi x,\varphi z) \theta(\varphi^{2}y) \Big\}; \\ \mathcal{F}_{2}:& F(\xi,y,z) = F(x,\xi,z) = 0, \\ & F(x,y,\varphi z) + F(y,z,\varphi x) + F(z,x,\varphi y) = 0, \quad \theta = 0; \\ \mathcal{F}_{3}:& F(\xi,y,z) = F(x,\xi,z) = 0, \\ & F(x,y,z) + F(y,z,x) + F(z,x,y) = 0; \\ \mathcal{F}_{4}:& F(x,y,z) = -\frac{1}{2n} \theta(\xi) \Big\{ g(\varphi x,\varphi y) \eta(z) + g(\varphi x,\varphi z) \eta(y) \Big\}; \\ \mathcal{F}_{5}:& F(x,y,z) = -\frac{1}{2n} \theta^{*}(\xi) \Big\{ g(x,\varphi y) \eta(z) + g(x,\varphi z) \eta(y) \Big\}; \\ \mathcal{F}_{6}:& F(x,y,z) = F(x,y,\xi) \eta(z) + F(x,z,\xi) \eta(y), \\ & F(x,y,\xi) = F(y,x,\xi) = -F(\varphi x,\varphi y,\xi), \quad \theta = \theta^{*} = 0; \\ \mathcal{F}_{7}:& F(x,y,z) = F(x,y,\xi) \eta(z) + F(x,z,\xi) \eta(y), \\ & F(x,y,\xi) = -F(y,x,\xi) = -F(\varphi x,\varphi y,\xi); \\ \mathcal{F}_{8}:& F(x,y,z) = F(x,y,\xi) \eta(z) + F(x,z,\xi) \eta(y), \\ & F(x,y,\xi) = F(y,x,\xi) = F(\varphi x,\varphi y,\xi); \\ \mathcal{F}_{9}:& F(x,y,z) = F(x,y,\xi) \eta(z) + F(x,z,\xi) \eta(y), \\ & F(x,y,\xi) = -F(y,x,\xi) = F(\varphi x,\varphi y,\xi); \\ \mathcal{F}_{10}:& F(x,y,z) = F(\xi,\varphi y,\varphi z) \eta(x); \\ \mathcal{F}_{11}:& F(x,y,z) = \eta(x) \Big\{ \eta(y) \omega(z) + \eta(z) \omega(y) \Big\} \,. \end{array}$$

Let now consider the case of the lowest dimension of the manifold $(M, \varphi, \xi, \eta, g)$, i.e. dim M = 3.

Let us denote the components $F_{ijk} = F(e_i, e_j, e_k)$ of the tensor F with respect to a φ -basis $\{e_i\}_{i=0}^2 = \{e_0 = \xi, e_1 = e, e_2 = \varphi e\}$ of T_pM , which satisfies the following conditions

(3)
$$g(e_0, e_0) = g(e_1, e_1) = -g(e_2, e_2) = 1, \quad g(e_i, e_j) = 0, \ i \neq j \in \{0, 1, 2\}.$$

According to [8], the components of the Lee forms with respect to a φ -basis with conditions (3) are

$$\begin{split} \theta_0 &= F_{110} - F_{220}, & \theta_1 &= F_{111} - F_{221}, & \theta_2 &= F_{112} - F_{211}, \\ \theta_0^* &= F_{120} + F_{210}, & \theta_1^* &= F_{112} + F_{211}, & \theta_2^* &= F_{111} + F_{221}, \\ \omega_0 &= 0, & \omega_1 &= F_{001}, & \omega_2 &= F_{002}. \end{split}$$

Then, if F_s (s = 1, 2, ..., 11) are the components of F in the corresponding basic classes \mathcal{F}_s and $x = x^i e_i$, $y = y^j e_j$, $z = z^k e_k$ are arbitrary vectors we have [8]:

$$F_{1}(x, y, z) = (x^{1}\theta_{1} - x^{2}\theta_{2}) (y^{1}z^{1} + y^{2}z^{2}), \\ \theta_{1} = F_{111} = F_{122}, \qquad \theta_{2} = -F_{211} = -F_{222}; \\ F_{2}(x, y, z) = F_{3}(x, y, z) = 0; \\ F_{4}(x, y, z) = \frac{1}{2}\theta_{0} \Big\{ x^{1} (y^{0}z^{1} + y^{1}z^{0}) - x^{2} (y^{0}z^{2} + y^{2}z^{0}) \Big\}, \\ \frac{1}{2}\theta_{0} = F_{101} = F_{110} = -F_{202} = -F_{220}; \\ F_{5}(x, y, z) = \frac{1}{2}\theta_{0}^{*} \Big\{ x^{1} (y^{0}z^{2} + y^{2}z^{0}) + x^{2} (y^{0}z^{1} + y^{1}z^{0}) \Big\}, \\ \frac{1}{2}\theta_{0}^{*} = F_{102} = F_{120} = F_{201} = F_{210}; \\ F_{6}(x, y, z) = F_{7}(x, y, z) = 0; \\ F_{8}(x, y, z) = \lambda \Big\{ x^{1} (y^{0}z^{1} + y^{1}z^{0}) + x^{2} (y^{0}z^{2} + y^{2}z^{0}) \Big\}, \\ \lambda = F_{101} = F_{110} = F_{202} = F_{220}; \\ F_{9}(x, y, z) = \mu \Big\{ x^{1} (y^{0}z^{2} + y^{2}z^{0}) - x^{2} (y^{0}z^{1} + y^{1}z^{0}) \Big\}, \\ \mu = F_{102} = F_{120} = -F_{201} = -F_{210}; \\ F_{10}(x, y, z) = \nu x^{0} (y^{1}z^{1} + y^{2}z^{2}), \qquad \nu = F_{011} = F_{022}; \\ F_{11}(x, y, z) = x^{0} \Big\{ (y^{1}z^{0} + y^{0}z^{1}) \omega_{1} + (y^{2}z^{0} + y^{0}z^{2}) \omega_{2} \Big\}, \\ \omega_{1} = F_{010} = F_{001}, \qquad \omega_{2} = F_{020} = F_{002}. \\ \end{cases}$$

Obviously, the class of 3-dimensional almost contact B-metric manifolds is

(5)
$$\mathcal{F}_1 \oplus \mathcal{F}_4 \oplus \mathcal{F}_5 \oplus \mathcal{F}_8 \oplus \mathcal{F}_9 \oplus \mathcal{F}_{10} \oplus \mathcal{F}_{11}.$$

1.2. Curvature properties of almost contact B-metric manifolds. Let $R = [\nabla, \nabla] - \nabla_{[,]}$ be the curvature (1,3)-tensor of ∇ and the corresponding curvature (0,4)-tensor be denoted by the same letter: R(x, y, z, w) = g(R(x, y)z, w). The following properties are valid:

(6)
$$R(x, y, z, w) = -R(y, x, z, w) = -R(x, y, w, z),$$
$$R(x, y, z, w) + R(y, z, x, w) + R(z, x, y, w) = 0.$$

On any \mathcal{F}_0 -manifold we have

(7)
$$R(x, y, \varphi z, \varphi w) = -R(x, y, z, w).$$

Any (0,4)-tensor having the properties (6) is called a *curvature-like* tensor. If such a tensor has the Kähler property (7) it is called a *Kähler* tensor.

From [18] it is known that every Kähler tensor on a 3-dimensional almost contact B-metric manifold is zero and therefore every 3-dimensional \mathcal{F}_0 -manifold is flat, i.e. R = 0.

The Ricci tensor ρ and the scalar curvature τ for R as well as their associated quantities are defined respectively by

$$\begin{split} \rho(y,z) &= g^{ij} R(e_i, y, z, e_j), & \tau &= g^{ij} \rho(e_i, e_j), \\ \rho^*(y,z) &= g^{ij} R(e_i, y, z, \varphi e_j), & \tau^* &= g^{ij} \rho^*(e_i, e_j). \end{split}$$

4

Further, we use the notation $g \otimes h$ for the Kulkarni-Nomizu product of two (0,2)-tensors, i.e.

$$(g \otimes h) (x, y, z, w) = g(x, z)h(y, w) - g(y, z)h(x, w)$$
$$+ g(y, w)h(x, z) - g(x, w)h(y, z).$$

Obviously, $g \otimes h$ has the properties of R in (6) when g and h are symmetric.

It is well known that the curvature tensor has the following form on any 3-dimensional manifold:

(8)
$$R = -g \otimes \left(\rho - \frac{\tau}{4}g\right).$$

Every non-degenerate 2-plane β with respect to g in T_pM , $p \in M$, has the following sectional curvature

$$k(\beta; p) = -\frac{2R(x, y, y, x)}{g \oslash g},$$

where $\{x, y\}$ is a basis of β .

A 2-plane β is said to be a φ -holomorphic section (respectively, a ξ -section) if $\beta = \varphi \beta$ (respectively, $\xi \in \beta$).

2. LIE GROUPS AS 3-DIMENSIONAL ALMOST CONTACT B-METRIC MANIFOLDS

Let L be a 3-dimensional real connected Lie group, and \mathfrak{l} be its Lie algebra with a basis $\{E_0, E_1, E_2\}$ of left invariant vector fields. Let an almost contact structure (φ, ξ, η) be defined by

(9)
$$\varphi E_0 = 0, \quad \varphi E_1 = E_2, \quad \varphi E_2 = -E_1, \quad \xi = E_0, \\ \eta(E_0) = 1, \quad \eta(E_1) = \eta(E_2) = 0$$

and let g be a pseudo-Riemannian metric such that

(10)
$$g(E_0, E_0) = g(E_1, E_1) = -g(E_2, E_2) = 1,$$
$$g(E_0, E_1) = g(E_0, E_2) = g(E_1, E_2) = 0.$$

Then we have a 3-dimensional almost contact B-metric manifold $(L, \varphi, \xi, \eta, g)$.

The corresponding Lie algebra $\mathfrak l$ is determined as follows:

(11)
$$[E_i, E_j] = C_{ij}^k E_k, \quad i, j, k \in \{0, 1, 2\}.$$

Applying the Jacobi identity

$$[[E_i, E_j], E_k] + [[E_j, E_k], E_i] + [[E_k, E_i], E_j] = 0,$$

from the nine commutation coefficients C_{ij}^k remain six which could be chosen as parameters, so the three coefficients with different indices are expressed by the six parameters (if the denominators are non-zero) as follows:

$$C_{12}^{0} = \frac{C_{12}^{1}C_{01}^{0} + C_{12}^{2}C_{02}^{0}}{C_{01}^{1} + C_{02}^{2}}, \qquad C_{02}^{1} = \frac{C_{01}^{1}C_{02}^{0} - C_{12}^{1}C_{02}^{0}}{C_{01}^{0} - C_{12}^{2}},$$
$$C_{01}^{2} = \frac{C_{01}^{0}C_{02}^{2} + C_{01}^{1}C_{12}^{2}}{C_{02}^{0} + C_{12}^{1}}.$$

The Koszul equality

(12) $2g(\nabla_{E_i}E_j, E_k) = g([E_i, E_j], E_k) + g([E_k, E_i], E_j) + g([E_k, E_j], E_i)$ implies the following formula for the components $F_{ijk} = F(E_i, E_j, E_k), i, j, k \in \{0, 1, 2\}$, of the tensor F:

$$2F_{ijk} = g\left([E_i, \varphi E_j] - \varphi[E_i, E_j], E_k\right) + g\left(\varphi[E_k, E_i] - [\varphi E_k, E_i], E_j\right) + g\left([E_k, \varphi E_j] - [\varphi E_k, E_i], E_i\right).$$

Then we have

(13)

$$F_{111} = F_{122} = 2C_{12}^{1}, \qquad F_{211} = F_{222} = 2C_{12}^{2}, \\
F_{120} = F_{102} = -C_{01}^{1}, \qquad F_{020} = F_{002} = -C_{01}^{0}, \\
F_{210} = F_{201} = -C_{02}^{2}, \qquad F_{010} = F_{001} = C_{02}^{0}, \\
F_{110} = F_{101} = \frac{1}{2} \left(C_{12}^{0} - C_{01}^{2} + C_{02}^{1} \right), \\
F_{220} = F_{202} = \frac{1}{2} \left(C_{12}^{0} + C_{01}^{2} - C_{02}^{1} \right), \\
F_{011} = F_{022} = C_{12}^{0} + C_{01}^{2} + C_{02}^{1}.
\end{cases}$$

For the Lee forms we obtain

(14)
$$\begin{aligned} \theta_0 &= -C_{01}^2 + C_{02}^1, \qquad \theta_0^* = -C_{01}^1 - C_{02}^2, \qquad \omega_0 = 0, \\ \theta_1 &= 2C_{12}^1, \qquad \theta_1^* = 2C_{12}^2, \qquad \omega_1 = C_{02}^0, \\ \theta_2 &= -2C_{12}^2, \qquad \theta_2^* = 2C_{12}^1, \qquad \omega_2 = -C_{01}^0. \end{aligned}$$

Using (4), (13) and (14), we deduce the following

Theorem 2.1. The manifold $(L, \varphi, \xi, \eta, g)$ belongs to the basic class \mathcal{F}_s $(s \in \{1, 4, 5, 8, 9, 10, 11\})$ if and only if the corresponding Lie algebra \mathfrak{l} is determined by the following commutators:

$$\mathcal{F}_{1}: \quad [E_{0}, E_{1}] = [E_{0}, E_{2}] = 0, \quad [E_{1}, E_{2}] = \alpha E_{1} + \beta E_{2};$$

$$\mathcal{F}_{4}: \quad [E_{0}, E_{1}] = \alpha E_{2}, \quad [E_{0}, E_{2}] = -\alpha E_{1}, \quad [E_{1}, E_{2}] = 0;$$

$$\mathcal{F}_{5}: \quad [E_{0}, E_{1}] = \alpha E_{1}, \quad [E_{0}, E_{2}] = \alpha E_{2}, \quad [E_{1}, E_{2}] = 0;$$
(15)
$$\mathcal{F}_{8}: \quad [E_{0}, E_{1}] = \alpha E_{2}, \quad [E_{0}, E_{2}] = \alpha E_{1}, \quad [E_{1}, E_{2}] = -2\alpha E_{0};$$

$$\mathcal{F}_{9}: \quad [E_{0}, E_{1}] = \alpha E_{1}, \quad [E_{0}, E_{2}] = -\alpha E_{2}, \quad [E_{1}, E_{2}] = 0;$$

$$\mathcal{F}_{10}: \quad [E_{0}, E_{1}] = \alpha E_{2}, \quad [E_{0}, E_{2}] = \alpha E_{1}, \quad [E_{1}, E_{2}] = 0;$$

$$\mathcal{F}_{11}: \quad [E_{0}, E_{1}] = \alpha E_{0}, \quad [E_{0}, E_{2}] = \alpha E_{0}, \quad [E_{1}, E_{2}] = 0;$$

where α , β are arbitrary real parameters. Moreover, the relations of α and β with the non-zero components F_{ijk} in the different basic classes \mathcal{F}_s from (4) are the following:

$$\begin{array}{lll} \mathcal{F}_{1}: & \alpha = \frac{1}{2}\theta_{1}, & \beta = \frac{1}{2}\theta_{2}; & \mathcal{F}_{4}: & \alpha = \frac{1}{2}\theta_{0}; \\ \mathcal{F}_{5}: & \alpha = -\frac{1}{2}\theta_{0}^{*}; & \mathcal{F}_{8}: & \alpha = -\lambda; \\ \mathcal{F}_{9}: & \alpha = -\mu; & \mathcal{F}_{10}: & \alpha = \frac{1}{2}\nu; \\ \mathcal{F}_{11}: & \alpha = -\omega_{2}, & \beta = \omega_{1}. \end{array}$$

Let us remark that an \mathcal{F}_0 -manifold is obtained when the Lie algebra is Abelian, i.e. all commutators are zero. Further, we omit this special case.

2.1. Structures of special type on the considered manifolds. The metric g is called *Killing* (or, Ad-*invariant*) if

(16)
$$g([x,y],z) = g(x,[y,z]).$$

Bearing in mind (10), (11) and (16), we obtain that g is Killing if and only if the equalities $C_{12}^0 = -C_{01}^2 = -C_{02}^1$ are valid and the rest of the commutation coefficients are zero. Then, according to (13) and (14), we get the following nonzero components $-F_{011} = -F_{022} = 2F_{110} = 2F_{101} = 2F_{220} = 2F_{202} = C_{12}^0$ and $\theta = \theta^* = \omega = 0$. Therefore, using (4), we establish that $F = F_8 + F_{10}$ and $2\lambda = -\nu$. Thus, it is valid the following

Theorem 2.2. The metric g is Killing if and only if $(L, \varphi, \xi, \eta, g)$ belongs to the subclass of $\mathcal{F}_8 \oplus \mathcal{F}_{10}$ determined by the condition $2\lambda = -\nu$.

In a similar way we obtain that \tilde{g} is Killing, i.e. $\tilde{g}([x, y], z) = \tilde{g}(x, [y, z])$, if and only if $C_{12}^0 = -C_{01}^1 = C_{02}^2$ and the others C_{ij}^k are zero. Then from (13) we have zero Lee forms and $F_{120} = F_{102} = -F_{210} = -F_{201} = 2F_{110} = 2F_{101} = 2F_{220} = 2F_{202} =$ $F_{011} = F_{022} = C_{12}^0$, i.e. $2\lambda = \mu = \nu$, bearing in mind (4). Therefore, we get

Theorem 2.3. The metric \tilde{g} is Killing if and only if $(L, \varphi, \xi, \eta, g)$ belongs to the subclass of $\mathcal{F}_8 \oplus \mathcal{F}_9 \oplus \mathcal{F}_{10}$ determined by the condition $2\lambda = \mu = \nu$.

The structure φ is called *bi-invariant*, if $\varphi[x, y] = [x, \varphi y]$. This condition is satisfied if and only if $F_{011} = F_{022} = 2F_{110} = 2F_{101} = 2F_{220} = 2F_{202} = C_{12}^0$, i.e. $2\lambda = \nu$. Therefore, it is valid the following

Theorem 2.4. The structure φ is bi-invariant if and only if $(L, \varphi, \xi, \eta, g)$ belongs to the subclass of $\mathcal{F}_8 \oplus \mathcal{F}_{10}$ determined by the condition $2\lambda = \nu$.

The structure φ is called *Abelian*, if $[\varphi x, \varphi y] = [x, y]$. This condition is valid if and only if $F_{011} = F_{022} = 2F_{110} = 2F_{101} = 2F_{220} = 2F_{202} = C_{12}^0$, i.e. $2\lambda = \nu$, and $F_{111} = F_{122} = 2C_{12}^1 = \theta_1$, $F_{211} = F_{222} = 2C_{12}^2 = -\theta_2$. Therefore, we deduce

Theorem 2.5. The structure φ is Abelian if and only if $(L, \varphi, \xi, \eta, g)$ belongs to the subclass of $\mathcal{F}_1 \oplus \mathcal{F}_8 \oplus \mathcal{F}_{10}$ determined by the condition $2\lambda = \nu$.

The vector field ξ is called *Killing*, if the Lie derivative with respect to ξ of the metric g is zero, i.e. $\mathcal{L}_{\xi}g = 0$. Then the manifold $(L, \varphi, \xi, \eta, g)$ is called a *K*-contact *B*-metric manifold. This condition is equivalent to $(\nabla_x \eta) y + (\nabla_y \eta) x = 0$, which is satisfied for any manifold in $\mathcal{F}_1 \oplus \mathcal{F}_2 \oplus \mathcal{F}_3 \oplus \mathcal{F}_7 \oplus \mathcal{F}_8 \oplus \mathcal{F}_{10}$ for arbitrary dimension, according to (2). Then, bearing in mind (5) – the class of the considered manifolds of dimension 3, we get

Theorem 2.6. The vector field ξ is Killing if and only if $(L, \varphi, \xi, \eta, g) \in \mathcal{F}_1 \oplus \mathcal{F}_8 \oplus \mathcal{F}_{10}$, *i.e.* the class of the 3-dimensional K-contact B-metric manifolds is $\mathcal{F}_1 \oplus \mathcal{F}_8 \oplus \mathcal{F}_{10}$.

2.2. Curvature properties of the considered manifolds. Using (12) and (15), we obtain the covariant derivatives of E_i with respect to ∇ . The non-zero ones of

them are the following:

$$\begin{array}{ll} \mathcal{F}_{1}: & \nabla_{E_{1}}E_{1} = \alpha E_{2}, \, \nabla_{E_{1}}E_{2} = \alpha E_{1}, \, \nabla_{E_{2}}E_{1} = -\beta E_{2}, \, \nabla_{E_{2}}E_{2} = -\beta E_{1}; \\ \mathcal{F}_{4}: & \nabla_{E_{1}}E_{0} = -\alpha E_{2}, \, \nabla_{E_{2}}E_{0} = \alpha E_{1}; \\ \mathcal{F}_{5}: & \nabla_{E_{1}}E_{1} = -\nabla_{E_{2}}E_{2} = \alpha E_{0}, \, \nabla_{E_{1}}E_{0} = -\alpha E_{1}, \, \nabla_{E_{2}}E_{0} = -\alpha E_{2}; \\ \mathcal{F}_{8}: & \nabla_{E_{1}}E_{2} = -\nabla_{E_{2}}E_{1} = -\alpha E_{0}, \, \nabla_{E_{1}}E_{0} = -\alpha E_{2}, \, \nabla_{E_{2}}E_{0} = -\alpha E_{1}; \\ \mathcal{F}_{9}: & \nabla_{E_{1}}E_{1} = \nabla_{E_{2}}E_{2} = \alpha E_{0}, \, \nabla_{E_{1}}E_{0} = -\alpha E_{1}, \, \nabla_{E_{2}}E_{0} = \alpha E_{2}; \\ \mathcal{F}_{10}: & \nabla_{E_{0}}E_{1} = \alpha E_{2}, \, \nabla_{E_{0}}E_{2} = \alpha E_{1}; \\ \mathcal{F}_{11}: & \nabla_{E_{0}}E_{0} = -\alpha E_{1} + \beta E_{2}, \, \nabla_{E_{0}}E_{1} = \alpha E_{0}, \nabla_{E_{0}}E_{2} = \beta E_{0}. \end{array}$$

By virtue of the latter equalities and bearing in mind the definition equalities of the curvature tensor R, the Ricci tensor ρ , the *-Ricci tensor ρ^* , the scalar curvature τ , its *-scalar curvature τ^* and the sectional curvature k_{ij} of the non-degenerate 2-plane $\{E_i, E_j\}$

(17)
$$R_{ijkl} = g \left(\nabla_{E_i} \nabla_{E_j} E_k - \nabla_{E_j} \nabla_{E_i} E_k - \nabla_{[E_i, E_j]} E_k, E_l \right),$$
$$(17) \qquad \rho_{jk} = R_{0jk0} + R_{1jk1} - R_{2jk2}, \qquad \rho_{jk}^* = R_{1jk2} + R_{2jk1},$$
$$\tau = \rho_{00} + \rho_{11} - \rho_{22}, \qquad \tau^* = 2\rho_{12}, \qquad k_{ij} = -2R_{ijij}/(g \odot g)_{ijij},$$

from Theorem 2.1 we obtain the following

Theorem 2.7. The \mathcal{F}_{10} -manifolds $(L, \varphi, \xi, \eta, g)$ are flat and the non-zero components of R, ρ , ρ^* and the non-zero values of τ , τ^* , k_{ij} for the manifolds $(L, \varphi, \xi, \eta, g)$ from the rest of the basic classes are the following:

$$\mathcal{F}_{1}: \quad R_{1212} = \rho_{11} = -\rho_{22} = \rho_{12}^{*} = \rho_{21}^{*} = \frac{1}{2}\tau = k_{12} = \alpha^{2} - \beta^{2};$$

$$\mathcal{F}_{4}: \quad -R_{0101} = R_{0202} = \frac{1}{2}\rho_{00} = \rho_{11} = -\rho_{22}$$

$$= \frac{1}{4}\tau = k_{01} = k_{02} = \alpha^{2};$$

$$\mathcal{F}_{5}: \quad -R_{0101} = R_{0202} = R_{1212} = \frac{1}{2}\rho_{00} = \frac{1}{2}\rho_{11} = -\frac{1}{2}\rho_{22}$$

$$= \rho_{12}^{*} = \rho_{21}^{*} = \frac{1}{6}\tau = k_{12} = k_{01} = k_{02} = -\alpha^{2};$$

$$(18) \quad \mathcal{F}_{8}: \quad R_{0101} = -R_{0202} = R_{1212} = -\frac{1}{2}\rho_{00} = \rho_{12}^{*} = \rho_{21}^{*}$$

$$= -\frac{1}{2}\tau = k_{12} = -k_{01} = -k_{02} = \alpha^{2};$$

$$\mathcal{F}_{9}: \quad R_{0101} = -R_{0202} = R_{1212} = -\frac{1}{2}\rho_{00} = \rho_{12}^{*} = \rho_{21}^{*}$$

$$= -\frac{1}{2}\tau = k_{12} = -k_{01} = -k_{02} = \alpha^{2};$$

$$\mathcal{F}_{11}: \quad R_{0101} = -\rho_{11} = -k_{01} = \alpha^{2}, \quad R_{0202} = -\rho_{22} = k_{02} = \beta^{2},$$

$$R_{0120} = \rho_{12} = \frac{1}{2}\tau^{*} = -\alpha\beta, \quad \rho_{00} = \frac{1}{2}\tau = -\alpha^{2} + \beta^{2}.$$

Let us remark, on the 3-dimensional almost contact B-metric manifold with the basis $\{E_0, E_1, E_2\}$ defined by (9), we have two basic ξ -sections $\{E_0, E_1\}$, $\{E_0, E_2\}$ and one basic φ -holomorphic section $\{E_1, E_2\}$.

According to Theorem 2.7, we obtain

Theorem 2.8. For the \mathcal{F}_s -manifolds $(L, \varphi, \xi, \eta, g)$, which do not belong to \mathcal{F}_0 , the following propositions are valid:

(1) Only the \mathcal{F}_1 -manifolds have a Kähler tensor R;

9

- (2) An \mathcal{F}_1 -manifold is flat if and only if $\alpha^2 = \beta^2$;
- All F₈-manifolds and F₉-manifolds have curvature tensors of the same form;
- (4) The curvature tensor of any \mathcal{F}_s -manifold (s = 4,11) has the property $R(x, y, \varphi z, \varphi w) = 0;$
- (5) Every \mathcal{F}_s -manifold (s = 4, 11) has a vanishing *-Ricci tensor;
- (6) Every \mathcal{F}_s -manifold (s = 4, 8, 9) has a positive scalar curvature;
- An F₁-manifold (resp., F₁₁-manifold) has a positive scalar curvature if and only if α² > β² (resp., α² < β²);
- (8) Every \mathcal{F}_5 -manifold has a negative scalar curvature;
- (9) An \mathcal{F}_1 -manifold (resp., \mathcal{F}_{11} -manifold) has a negative scalar curvature if and only if $\alpha^2 < \beta^2$ (resp., $\alpha^2 > \beta^2$);
- (10) Every \mathcal{F}_s -manifold (s = 1, 4, 5, 8, 9) has a vanishing *-scalar curvature;
- (11) An \mathcal{F}_{11} -manifold has a vanishing *-scalar curvature if and only if $\alpha\beta = 0$ for $\alpha \neq \beta$;
- (12) An \mathcal{F}_{11} -manifold has a positive (resp., negative) *-scalar curvature if and only if $\alpha\beta < 0$ (resp., $\alpha\beta > 0$);
- (13) Every \mathcal{F}_1 -manifold has vanishing sectional curvatures of the ξ -sections;
- (14) Every \mathcal{F}_4 -manifold has positive sectional curvatures of the basic ξ -sections;
- (15) Every \mathcal{F}_s -manifold (s = 5,8,9) has negative sectional curvatures of the basic ξ -sections;
- (16) An \mathcal{F}_1 -manifold has a vanishing scalar curvature of the basic φ -holomorphic section if and only if the manifold is flat;
- (17) Every \mathcal{F}_s -manifold (s = 4, 11) has a vanishing scalar curvature of the basic φ -holomorphic section;
- (18) Every \mathcal{F}_s -manifold (s = 8,9) (resp., \mathcal{F}_5 -manifold) has a positive (resp., negative) scalar curvature of the basic φ -holomorphic section;
- (19) An \mathcal{F}_1 -manifold has a positive (resp., negative) scalar curvature of the basic φ -holomorphic section if and only if $\alpha^2 > \beta^2$ (resp., $\alpha^2 < \beta^2$).

Taking into account (18), we get the following

Corollary 2.9. The form of the Ricci tensor on $(L, \varphi, \xi, \eta, g)$ in the respective basic class is:

$$\begin{aligned} \mathcal{F}_{1}: \quad \rho &= \frac{\tau}{2} \left(g - \eta \otimes \eta \right); & \mathcal{F}_{4}: \quad \rho &= \frac{\tau}{4} \left(g + \eta \otimes \eta \right); \\ \mathcal{F}_{5}: \quad \rho &= \frac{\tau}{3} g; & \mathcal{F}_{8}: \quad \rho &= \tau(\eta \otimes \eta); \\ \mathcal{F}_{9}: \quad \rho &= \tau(\eta \otimes \eta); & \mathcal{F}_{11}: \quad \rho &= \rho(\varphi \cdot, \varphi \cdot) + \frac{\tau}{2} g - \tau^{*} g^{*}, \end{aligned}$$

where $g^*(\cdot, \cdot) = g(\cdot, \varphi \cdot) = \tilde{g} - \eta \otimes \eta$.

Using Corollary 2.9 and (8), we obtain for the following

Corollary 2.10. The form of the curvature tensor on $(L, \varphi, \xi, \eta, g)$ in the respective basic class is:

$$\begin{split} \mathcal{F}_{1} : & R = -\frac{\tau}{4} \left(g \otimes g \right) + \frac{\tau}{2} \left(g \otimes \left(\eta \otimes \eta \right) \right); \\ \mathcal{F}_{4} : & R = -\frac{\tau}{4} \left(g \otimes \left(\eta \otimes \eta \right) \right); \\ \mathcal{F}_{5} : & R = -\frac{\tau}{12} \left(g^{*} \otimes g^{*} \right) - \frac{\tau}{6} \left(g \otimes \left(\eta \otimes \eta \right) \right); \\ \mathcal{F}_{8} : & R = \frac{\tau}{4} \left(g^{*} \otimes g^{*} \right) - \frac{\tau}{2} \left(g \otimes \left(\eta \otimes \eta \right) \right); \\ \mathcal{F}_{9} : & R = \frac{\tau}{4} \left(g^{*} \otimes g^{*} \right) - \frac{\tau}{2} \left(g \otimes \left(\eta \otimes \eta \right) \right); \\ \mathcal{F}_{11} : & R = -\rho \otimes \left(\eta \otimes \eta \right). \end{split}$$

Let $(V, \varphi, \xi, \eta, g)$ be a vector space endowed with an almost contact structure (φ, ξ, η) . The decomposition $x = -\varphi^2 x + \eta(x)\xi$ generates the projectors h and v on (V, φ, ξ, η) determined by $h(x) = -\varphi^2 x$ and $v(x) = \eta(x)\xi$ and having the properties $h \circ h = h$, $v \circ v = v$, $h \circ v = v \circ h = 0$. Therefore, we have the orthogonal decomposition $V = h(V) \oplus v(V)$. Then we obtain the corresponding linear operators ℓ_1, ℓ_2, ℓ_3 in the space $V^* \times V^*$ of the tensors S of type (0,2) over (V, φ, ξ, η) defined in [1]:

$$\ell_1(S)(x,y) = S(h(x), h(y)), \quad \ell_2(S)(x,y) = S(v(x), v(y)), \\ \ell_3(S)(x,y) = S(v(x), y) + S(x, v(y)) - 2S(v(x), v(y)).$$

The structure group \mathcal{G} of $(M, \varphi, \xi, \eta, g)$ is determined by $\mathcal{G} = \mathcal{O}(n; \mathbb{C}) \times \mathcal{I}$, where \mathcal{I} is the identity on $\text{Span}(\xi)$ and $\mathcal{O}(n; \mathbb{C}) = \mathcal{GL}(n; \mathbb{C}) \cap \mathcal{O}(n, n)$.

It is known [10], the following orthogonal decomposition which is invariant with respect to the action of the structure group \mathcal{G} :

$$V^* \times V^* = \ell_1(V^* \times V^*) \oplus \ell_2(V^* \times V^*) \oplus \ell_3(V^* \times V^*),$$

where

$$\ell_i(V^* \times V^*) = \{ S \in (V^* \times V^*) \mid S = \ell_i(S) \}, \quad i = 1, 2, 3.$$

Since the metrics g and \tilde{g} belong to $V^* \times V^*$, then their components in the three orthogonal subspaces are:

$$\begin{split} \ell_1(g) &= -g(\varphi \cdot, \varphi \cdot) = g - \eta \otimes \eta, \qquad \ell_2(g) = \eta \otimes \eta, \qquad \ell_3(g) = 0, \\ \ell_1(\tilde{g}) &= g(\cdot, \varphi \cdot) = \tilde{g} - \eta \otimes \eta, \qquad \ell_2(\tilde{g}) = \eta \otimes \eta, \qquad \ell_3(\tilde{g}) = 0. \end{split}$$

On any almost contact B-metric manifold there are a pair of B-metrics (g, \tilde{g}) which restrictions on $H = \text{Ker}(\eta)$ coincide with their ℓ_1 -images and are the real part and the imaginary part of the complex Riemannian metric $g^{\mathbb{C}}$, i.e. $g^{\mathbb{C}} = g|_H + i\tilde{g}|_H =$ $\ell_1(g) + i\ell_1(\tilde{g})$. Moreover, the restrictions of g and \tilde{g} on $H^{\perp} = \text{Span}(\xi)$ coincide with their ℓ_2 -images and $\eta \otimes \eta$, i.e. $g|_{H^{\perp}} = \tilde{g}|_{H^{\perp}} = \ell_2(g) = \ell_2(\tilde{g}) = \eta \otimes \eta$.

It is well known that a manifold is called *Einstein* if the Ricci tensor is proportional to the metric tensor, i.e. $\rho = \lambda g$, $\lambda \in \mathbb{R}$.

Then, bearing in mind the above arguments, it is reasonable to call a manifold M whose Ricci tensor satisfies the condition

(19)
$$\rho = \lambda g + \mu \tilde{g} + \nu \eta \otimes \eta$$

an η -complex-Einstein manifold. The case of the η -complex-Einstein Sasaki-like manifolds is considered in [7]. If $\mu = 0$ we call M an η -Einstein manifold. On

the almost contact metric manifolds in [3] is studied the η -Einstein geometry as a class of distinguished Riemannian metrics on the contact metric manifolds. It is well-known [22] that λ and ν are constants if the almost contact metric manifold is K-contact and has dimension greater than 3.

It is meaningful to consider the Einstein property in the separate components $\ell_i(\rho)$ and $\ell_i(\rho^*)$, i = 1, 2, 3.

In [18] the following notions related to the Einstein condition are introduced. An almost contact B-metric manifold $(M, \varphi, \xi, \eta, g)$ is called *contact-Einstein* if the Ricci tensor has the form $\rho = \lambda g|_H + \mu \tilde{g}|_H + \nu \eta \otimes \eta$, where $\lambda, \mu, \nu \in \mathbb{R}$. A contact-Einstein manifold is called *h-Einstein* (resp., *v-Einstein*) if $\rho = \lambda g|_H + \mu \tilde{g}|_H$ (resp., $\rho = \nu \eta \otimes \eta$). An *h*-Einstein manifold is called φ -Einstein (resp., **-Einstein*) if $\rho = \lambda g|_H$ (resp., **-Einstein*) if $\rho = \lambda g|_H$ (resp., $\rho = \mu \tilde{g}|_H$).

According to Corollary 2.9, we obtain

Proposition 2.11. The manifold $(M, \varphi, \xi, \eta, g)$ is:

- (1) φ -Einstein if it belongs to \mathcal{F}_1 ;
- (2) η -Einstein if it belongs to \mathcal{F}_4 ;
- (3) Einstein if it belongs to \mathcal{F}_5 ;
- (4) v-Einstein if it belongs to $\mathcal{F}_8 \oplus \mathcal{F}_9$.

3. An Example

In [8] is introduced a real connected Lie group L as a manifold from the class $\mathcal{F}_9 \oplus \mathcal{F}_{10}$. We consider now this example for dim L = 3 and give geometrical characteristics in relation with some previous results in this work. In this case L is a 3-dimensional real connected Lie group and its associated Lie algebra with a global basis $\{E_0, E_1, E_2\}$ of left invariant vector fields on L is defined by

$$[E_0, E_1] = -a_1E_1 - a_2E_2, \quad [E_0, E_2] = -a_2E_1 + a_1E_2, \quad [E_1, E_2] = 0,$$

where a_1, a_2 are real constants. Actually, L is a family of Lie groups depending on the pair of real parameters. An invariant almost contact structure is defined by (9) and g is a pseudo-Riemannian metric such that (10) are valid. Thus, because of (1), the induced 3-dimensional manifold $(L, \varphi, \xi, \eta, g)$ is an almost contact B-metric manifold.

Let us remark that in [21] the same Lie group with the same almost contact structure but equipped with a compatible Riemannian metric is studied as an almost cosymplectic manifold.

The non-zero components F_{ijk} of the tensor F are following:

 $F_{011} = F_{022} = -2a_2, \quad F_{102} = F_{120} = -F_{201} = -F_{210} = a_1.$

Thus, it follows that $(L, \varphi, \xi, \eta, g) \in \mathcal{F}_9 \oplus \mathcal{F}_{10}$, bearing in mind (4) for $\mu = a_1$, $\nu = -2a_2$.

Further, we continue with examination of this family of almost contact B-metric manifolds.

Using (12), we obtain the covariant derivatives with respect to ∇ of E_i for the manifold $(L, \varphi, \xi, \eta, g)$:

$$\nabla_{E_0} E_1 = \frac{\nu}{2} E_2, \quad \nabla_{E_0} E_2 = \frac{\nu}{2} E_1, \quad \nabla_{E_1} E_0 = \mu E_1, \quad \nabla_{E_2} E_0 = -\mu E_2,$$

$$\nabla_{E_1} E_1 = \nabla_{E_2} E_2 = -\mu E_0.$$

By virtue of the latter equalities and bearing in mind the definition equalities (17), we get:

(20)

$$R_{0110} = -R_{0220} = R_{1221} = -\mu^2, \qquad R_{0120} = \mu\nu,$$

$$\rho_{00} = -2\rho_{22} = -2\mu^2, \qquad \rho_{00}^* = 2\rho_{12} = 2\mu\nu,$$

$$\tau = -2\mu^2, \qquad \tau^* = 2\mu\nu,$$

$$k_{01} = k_{02} = -\mu^2, \qquad k_{12} = \mu^2$$

We obtain the form of the Ricci tensor ρ :

(21)
$$\rho = \left(\tau + \frac{\tau^*}{2}\right)\eta \otimes \eta - \frac{\tau^*}{2}\tilde{g}.$$

Consequently, bearing in mind (19), $(L, \varphi, \xi, \eta, g)$ is an η -complex-Einstein manifold with $\lambda = 0$.

According to (8) and (21), we obtain the following form of the curvature tensor:

(22)
$$R = \frac{1}{4}g \oslash \left(\tau g + 2\tau^* \tilde{g} - 2\left(2\tau + \tau^*\right)\eta \otimes \eta\right).$$

If $\mu = 0$, $\nu \neq 0$, i.e. the manifold belongs to \mathcal{F}_{10} , then $\tau = \tau^* = 0$. Therefore, (22) implies R = 0.

If $\nu = 0$, $\mu \neq 0$, i.e. the manifold belongs to \mathcal{F}_9 , then $\tau^* = 0$, $\tau \neq 0$ and

$$R = \frac{\tau}{4}g \oslash \left(g - 4\eta \otimes \eta\right).$$

The latter conclusions support Corollary 2.9 and Corollary 2.10.

Taking into account (20), we obtain that the sectional curvatures of the basic ξ sections (respectively, the basic φ -holomorphic sections) are negative (respectively,
positive) for $(L, \varphi, \xi, \eta, g) \in \mathcal{F}_9 \oplus \mathcal{F}_{10}$, which support Theorem 2.8 for \mathcal{F}_9 and \mathcal{F}_{10} .

Acknowledgment. This paper is partially supported by project NI13-FMI-002 of the Scientific Research Fund, Plovdiv University, Bulgaria

References

- Alexiev, V., Ganchev, G.: On the classification of almost contact metric manifolds. In: Math. and Educ. in Math. Proc. 15th Spring Conf. of UBM. Sunny Beach, 155–161 (1986)
- Blair, D.E.: Riemannian Geometry of Contact and Symplectic Manifolds. Progress in Mathematics 203. Birkhäuser, Boston (2002)
- [3] Boyer, C.P., Galicki, K., Matzeu, P.: On η-Einstein Sasakian geometry. Commun. Math. Phys. 262, 177–208 (2006)
- [4] Ganchev, G., Borisov, A.: Note on the almost complex manifolds with a Norden metric. C. R. Acad. Bulg. Sci. 39, 31–34 (1986)
- [5] Ganchev, G., Mihova, V., Gribachev, K.: Almost contact manifolds with B-metric. Math. Balkanica (N.S.) 7 (3-4), 261–276 (1993)

- [6] Gribachev, K., Mekerov, D., Djelepov, G.: Generalized B-manifolds. C. R. Acad. Bulg. Sci. 38, 299–302 (1985)
- [7] Ivanov, S., Manev, H., Manev, M.: Sasaki-like almost contact complex Riemannian manifolds. arXiv:1402.5426
- [8] Manev, H.: On the structure tensors of almost contact B-metric manifolds. arXiv:1405.3088
- [9] Manev, M.: Almost contact B-metric hypersurfaces of Kaehlerian manifolds with B-metric. In: Dimiev, S., Sekigawa, K. (eds.) Perspectives of Complex Analysis, Differential Geometry and Mathematical Physics, pp. 159–170. World Sci. Publ., Singapore (2001)
- [10] Manev, M.: On the conformal geometry of almost contact manifolds with B-metric. Ph. D. Thesis. University of Plovdiv, (1998)
- [11] Manev, M.: Curvature properties on some classes of almost contact manifolds with B-metric.
 C. R. Acad. Bulg. Sci. 65 (3), 283–290 (2012)
- [12] Manev, M.: Contactly conformal transformations of general type of almost contact manifolds with B-metric. Applications. Math. Balkanica (N.S.) 11 (3-4), 347–357 (1997)
- [13] Manev, M.: Natural connection with totally skew-symmetric torsion on almost contact manifolds with B-metric. Int. J. Geom. Methods Mod. Phys. 9 (5), 1250044 (20 pages) (2012)
- [14] Manev, M., Gribachev, K.: Contactly conformal transformations of almost contact manifolds with B-metric. Serdica Math. J. 19, 287–299 (1993)
- [15] Manev, M., Gribachev, K.: Conformally invariant tensors on almost contact manifolds with B-metric. Serdica Math. J. 20, 133–147 (1994)
- [16] Manev, M., Ivanova, M.: Canonical type connections on almost contact manifold with Bmatric. Ann. Global Anal. Geom. 43 (4), 397–408 (2013)
- [17] Manev, M., Ivanova, M.: A classification of the torsion tensors on almost contact manifolds with B-metric. Cent. Eur. J. Math. 12 (10), 1416–1432 (2014)
- [18] Manev, M., Nakova, G.: Curvature properties of some three-dimensional almost contact B-metric manifolds. Plovdiv Univ. Sci. Works – Math. 34 (3), 51–60 (2004)
- [19] Nakova, G., Gribachev, K.: Submanifolds of some almost contact manifolds with B-metric with codimension two, I. Math. Balkanica (N.S.) 11, 255–267 (1997)
- [20] Manev, M., Nakova, G.: Curvature properties on some three-dimensional almost contact manifolds with B-metric, II. In: Mladenov, I.M., Hirshfeld, A.C. (eds.) Proc. 5th Internat. Conf. Geometry, Integrability & Quantization, V, pp. 169–177. SOFTEX, Sofia (2004)
- [21] Olszak, Z.: On almost cosymplectic manifolds. Kodai Math. J. 4 (2), 239-250 (1981)
- [22] Yano, K., Kon, M.: Structures on Manifolds. Series in Pure Mathematics 3, World Sci. Publ., Singapore (1984)

(H. Manev) MEDICAL UNIVERSITY OF PLOVDIV, FACULTY OF PHARMACY, DEPARTMENT OF PHARMACEUTICAL SCIENCES, 15-A VASIL APRILOV BLVD., PLOVDIV 4002, BULGARIA;

PAISII HILENDARSKI UNIVERSITY OF PLOVDIV, FACULTY OF MATHEMATICS AND INFORMATICS, DE-PARTMENT OF ALGEBRA AND GEOMETRY, 236 BULGARIA BLVD., PLOVDIV 4027, BULGARIA

E-mail address: hmanev@uni-plovdiv.bg

(D. Mekerov) PAISII HILENDARSKI UNIVERSITY OF PLOVDIV, FACULTY OF MATHEMATICS AND INFORMATICS, DEPARTMENT OF ALGEBRA AND GEOMETRY, 236 BULGARIA BLVD., PLOVDIV 4027, BULGARIA

 $E\text{-}mail\ address: \texttt{mircho@uni-plovdiv.bg}$