

THE DENSITY OF PRIMES DIVIDING A TERM IN THE SOMOS-5 SEQUENCE

BRYANT DAVIS, REBECCA KOTSONIS, AND JEREMY ROUSE

ABSTRACT. The Somos-5 sequence is defined by $a_0 = a_1 = a_2 = a_3 = a_4 = 1$ and $a_m = \frac{a_{m-1}a_{m-4} + a_{m-2}a_{m-3}}{a_{m-5}}$ for $m \geq 5$. We relate the arithmetic of the Somos-5 sequence to the elliptic curve $E : y^2 + xy = x^3 + x^2 - 2x$ and use properties of Galois representations attached to E to prove the density of primes p dividing some term in the Somos-5 sequence is equal to $\frac{5087}{10752}$.

1. INTRODUCTION AND STATEMENT OF RESULTS

There are many results in number theory that relate to a determination of the primes dividing some particular sequence. For example, it well-known that if p is a prime number, then p divides some term of the Fibonacci sequence, defined by $F_0 = 0$, $F_1 = 1$, and $F_n = F_{n-1} + F_{n-2}$ for $n \geq 2$. Students in elementary number theory learn that a prime p divides a number of the form $n^2 + 1$ if and only if $p = 2$ or $p \equiv 1 \pmod{4}$.

In 1966, Hasse proved in [4] that if $\pi_{\text{even}}(x)$ is the number of primes $p \leq x$ so that $p|2^n + 1$ for some n , then

$$\lim_{x \rightarrow \infty} \frac{\pi_{\text{even}}(x)}{\pi(x)} = \frac{17}{24}.$$

Note that a prime number p divides $2^n + 1$ if and only if 2 has even order in \mathbb{F}_p^\times .

A related result is the following. The Lucas numbers are defined by $L_0 = 2$, $L_1 = 1$ and $L_n = L_{n-1} + L_{n-2}$ for $n \geq 2$. In 1985, Lagarias proved (see [9] and [10]) that the density of primes dividing some Lucas number is $2/3$. Given a prime number p , let $Z(p)$ be the smallest integer m so that $p|F_m$. A prime p divides L_n for some n if and only if $Z(p)$ is even. In [2], Paul Cubre and the third author prove a conjecture of Bruckman and Anderson on the density of primes p for which $m|Z(p)$, for an arbitrary positive integer m .

In the early 1980s, Michael Somos discovered integer-valued non-linear recurrence sequences. The Somos- k sequence is defined by $c_0 = c_1 = \cdots = c_{k-1} = 1$ and

$$c_m = \frac{c_{m-1}c_{m-(k-1)} + c_{m-2}c_{m-(k-2)} + \cdots + c_{m-\lfloor \frac{k}{2} \rfloor}c_{m-\lceil \frac{k}{2} \rceil}}{c_{m-k}}$$

for $m \geq k$. Despite the fact that division is involved in the definition of the Somos sequences, the values c_m are integral for $4 \leq k \leq 7$. Fomin and Zelevinsky [3] show that the introduction of parameters into the recurrence results in the c_m being Laurent polynomials in those parameters. Also, Speyer [15] gave a combinatorial interpretation of the Somos sequences in terms of the number of perfect matchings in a family of graphs.

Somos-4 and Somos-5 type sequences are also connected with the arithmetic of elliptic curves (a connection made quite explicit by A. N. W. Hone in [5], and [6]). If a_n is the n th term in the Somos-4 sequence, $E : y^2 + y = x^3 - x$ and $P = (0, 0) \in E(\mathbb{Q})$, then the x -coordinate of the denominator of $(2n - 3)P$ is equal to a_n^2 . It follows from this that $p|a_n$ if and only if $(2n - 3)P$ reduces to the identity in $E(\mathbb{F}_p)$, and so a prime p divides a term in the Somos-4 sequence if and only if $(0, 0) \in E(\mathbb{F}_p)$ has odd order. In [8], Rafe Jones and the third author prove that the density of primes dividing some term of the Somos-4 sequence is $\frac{11}{21}$. The goal of the present paper is to prove an analogous result for the Somos-5 sequence.

Let $\pi'(x)$ denote the number of primes $p \leq x$ so that p divides some term in the Somos-5 sequence. We have the following table of data.

x	$\pi'(x)$	$\frac{\pi'(x)}{\pi(x)}$
10	3	0.750000
10^2	12	0.480000
10^3	83	0.494048
10^4	588	0.478438
10^5	4539	0.473207
10^6	37075	0.472305
10^7	314485	0.473209
10^8	2725670	0.473087
10^9	24057711	0.473134
10^{10}	215298607	0.473129

Our main result is the following.

Theorem 1. *We have*

$$\lim_{x \rightarrow \infty} \frac{\pi'(x)}{\pi(x)} = \frac{5087}{10752} \approx 0.473121.$$

The Somos-5 sequence is related to the coordinates of rational points on the elliptic curve $E : y^2 + xy = x^3 + x^2 - 2x$. This curve has $E(\mathbb{Q}) \cong \mathbb{Z} \times \mathbb{Z}/2\mathbb{Z}$ and generators are $P = (2, 2)$ (of infinite order) and $Q = (0, 0)$ (of order 2). We have (see Lemma 3) that

$$mP + Q = \left(\frac{a_{m+2}^2 - a_m a_{m+4}}{a_{m+2}^2}, \frac{4a_m a_{m+2} a_{m+4} - a_m^2 a_{m+6} - a_{m+2}^3}{a_{m+2}^3} \right).$$

It follows that a prime p divides a term in the Somos-5 sequence if and only if the reduction of Q modulo p is in $\langle P \rangle \subseteq E(\mathbb{F}_p)$. Another way of stating this is the following: there is a 2-isogeny $\phi : E \rightarrow E'$, where $E' : y^2 + xy = x^3 + x^2 + 8x + 10$ and

$$\phi(x, y) = \left(\frac{x^2 - 2}{x}, \frac{x^2 y + 2x + 2y}{x^2} \right).$$

The kernel of ϕ is $\{0, Q\}$. Letting $R = \phi(P)$ we show (see Theorem 4) that p divides some term in the Somos-5 sequence if and only if the order of P in $E(\mathbb{F}_p)$ is twice that of R in $E'(\mathbb{F}_p)$.

A result of Pink (see Proposition 3.2 on page 284 of [11]) shows that the ℓ -adic valuation of the order of a point $P \pmod{p}$ can be determined from a suitable Galois representation attached to an elliptic curve. For a positive integer k , we let K_k be the field obtained by adjoining to \mathbb{Q} the x and y coordinates of all points β_k with $2^k \beta_k = P$. There is a Galois representation $\rho_{E, 2^k} : \text{Gal}(K_k/\mathbb{Q}) \rightarrow$

$\mathrm{AGL}_2(\mathbb{Z}/2^k\mathbb{Z})$ and we relate the power of 2 dividing the order of P in $E(\mathbb{F}_p)$ to $\rho_{E,2^k}(\sigma_p)$, where σ_p is a Frobenius automorphism at p in $\mathrm{Gal}(K_k/\mathbb{Q})$. Using the isogeny ϕ we are able to relate $\rho_{E,2^k}(\sigma_p)$ and $\rho_{E',2^{k-1}}(\sigma_p)$, obtaining a criterion that indicates when p divides some term in the Somos-5 sequence. We then determine the image of $\rho_{E,2^k}$ for all k .

Once the image of $\rho_{E,2^k}$ is known, the problem of computing the fraction of elements in the image with the desired properties is quite a difficult one. We introduce a new and simple method for computing this fraction and apply it to prove Theorem 1.

Acknowledgements. *The first and second authors thank the Wake Forest Undergraduate Research and Creative Activities Center for financial support. The authors used Magma [1] version 2.20-6 for computations.*

2. BACKGROUND

If E/F is an elliptic curve given in the form $y^2 + a_1xy + a_3y = x^3 + a_2x^2 + a_4x + a_6$, the set $E(F)$ has the structure of an abelian group. Specifically, if $P, Q \in E(F)$, let $R = (x, y)$ be the third point of intersection between E and the line through P and Q . We define $P + Q = (x, -y - a_1x - a_3)$. The multiplication by m map on an elliptic curve has degree m^2 , and so if E/\mathbb{C} is an elliptic curve and $\alpha \in E(\mathbb{C})$, then there are m^2 points β so that $m\beta = \alpha$.

If K/\mathbb{Q} is a finite extension, let \mathcal{O}_K denote the ring of algebraic integers in K . A prime p ramifies in K if $p\mathcal{O}_K = \prod_{i=1}^r \mathfrak{p}_i^{e_i}$ and some $e_i > 1$, where the \mathfrak{p}_i are distinct prime ideals of \mathcal{O}_K .

Suppose K/\mathbb{Q} is Galois, p is a prime number that does not ramify in K , and $p\mathcal{O}_K = \prod_{i=1}^g \mathfrak{p}_i$. For each i , there is a unique element $\sigma \in \mathrm{Gal}(K/\mathbb{Q})$ for which

$$\sigma(\alpha) \equiv \alpha^p \pmod{\mathfrak{p}_i}$$

for all $\alpha \in \mathcal{O}_K$. This element is called the Artin symbol of \mathfrak{p}_i and is denoted $\left[\frac{K/\mathbb{Q}}{\mathfrak{p}_i}\right]$. If $i \neq j$, $\left[\frac{K/\mathbb{Q}}{\mathfrak{p}_i}\right]$ and $\left[\frac{K/\mathbb{Q}}{\mathfrak{p}_j}\right]$ are conjugate in $\mathrm{Gal}(K/\mathbb{Q})$ and $\left[\frac{K/\mathbb{Q}}{p}\right] := \left\{ \left[\frac{K/\mathbb{Q}}{\mathfrak{p}_i}\right] : 1 \leq i \leq g \right\}$ is a conjugacy class in $\mathrm{Gal}(K/\mathbb{Q})$.

The key tool we will use in the proof of Theorem 1 is the Chebotarev density theorem.

Theorem 2 ([7], page 143). *If $C \subseteq \mathrm{Gal}(K/\mathbb{Q})$ is a conjugacy class, then*

$$\lim_{x \rightarrow \infty} \frac{\#\{p \leq x : p \text{ prime}, \left[\frac{K/\mathbb{Q}}{p}\right] = C\}}{\pi(x)} = \frac{|C|}{|\mathrm{Gal}(K/\mathbb{Q})|}.$$

Roughly speaking, each element of $\mathrm{Gal}(K/\mathbb{Q})$ arises as $\left[\frac{K/\mathbb{Q}}{\mathfrak{p}}\right]$ equally often.

Let $E[m] = \{P \in E : mP = 0\}$ be the set of points of order dividing m on E . Then $\mathbb{Q}(E[m])/\mathbb{Q}$ is Galois and $\mathrm{Gal}(\mathbb{Q}(E[m])/\mathbb{Q})$ is isomorphic to a subgroup of $\mathrm{Aut}(E[m]) \cong \mathrm{GL}_2(\mathbb{Z}/m\mathbb{Z})$. Moreover, Proposition V.2.3 of [13] implies that if σ_p is a Frobenius automorphism at some prime above p and $\tau : \mathrm{Gal}(\mathbb{Q}(E[m])/\mathbb{Q}) \rightarrow \mathrm{GL}_2(\mathbb{Z}/m\mathbb{Z})$, then $\mathrm{tr} \tau(\sigma_p) \equiv p + 1 - \#E(\mathbb{F}_p) \pmod{m}$ and $\det(\tau(\sigma_p)) \equiv p \pmod{m}$. Another useful fact is the following. If K/\mathbb{Q} is a number field, \mathfrak{p} is a prime ideal in \mathcal{O}_K above p , $\mathrm{gcd}(m, p) = 1$ and $P \in E(K)[m]$ is not the identity, then P does not reduce to the identity in $E(\mathcal{O}_K/\mathfrak{p})$. This is a consequence of Proposition VII.3.1 of [13].

We will construct Galois representations attached to elliptic curves with images in $\text{AGL}_2(\mathbb{Z}/2^k\mathbb{Z}) \cong (\mathbb{Z}/2^k\mathbb{Z})^2 \rtimes \text{GL}_2(\mathbb{Z}/2^k\mathbb{Z})$. Elements of such a group can be thought of either as pairs (\vec{v}, M) , where \vec{v} is a row vector, and $M \in \text{GL}_2(\mathbb{Z}/2^k\mathbb{Z})$, or as 3×3 matrices $\begin{bmatrix} a & b & 0 \\ c & d & 0 \\ e & f & 1 \end{bmatrix}$, where $\vec{v} = [e \ f]$ and $M = \begin{bmatrix} a & b \\ c & d \end{bmatrix}$. In the former notation, the group operation is given by

$$(\vec{v}_1, M_1) * (\vec{v}_2, M_2) = (\vec{v}_1 + \vec{v}_2 M_1, M_2 M_1).$$

3. CONNECTION BETWEEN THE SOMOS-5 SEQUENCE AND E

Lemma 3. *Define $P = (2, 2)$ and $Q = (0, 0)$ on $E : y^2 + xy = x^3 + x^2 - 2x$. For all $m \geq 0$, we have the following relationship between the Somos-5 sequence and E :*

$$mP + Q = \left(\frac{a_{m+2}^2 - a_m a_{m+4}}{a_{m+2}^2}, \frac{4a_m a_{m+2} a_{m+4} - a_m^2 a_{m+6} - a_{m+2}^3}{a_{m+2}^3} \right).$$

Proof. We will prove this by strong induction. A straightforward calculation shows that the base cases $m = 0$ and $m = 1$ are true. For simplicity's sake, we will denote $a = a_m$, $b = a_{m+1}$, $c = a_{m+2}$, $d = a_{m+3}$, $e = a_{m+4}$, $f = a_{m+5}$ and $g = a_{m+6}$. Our inductive hypothesis is that

$$mP + Q = \left(\frac{c^2 - ae}{c^2}, \frac{4ace - a^2g - c^3}{c^3} \right).$$

We will now compute $(m+2)P + Q$.

To find the x and y coordinates of $(m+2)P + Q$, we add $2P = (1, -1)$ to $mP + Q$. If w is the slope and v is the y -intercept, the line between $2P$ and $mP + Q$ is $y = wx + v$ with $w = \frac{ag-4ce}{ce}$ and $v = \frac{-ag+3ce}{ce}$. Substituting this into the equation for E , we find the x -coordinate of $2P + (mP + Q)$ to be $r_x = \frac{a^2g^2 - 7aceg + ae^3 + 9c^2e^2}{c^2e^2}$. A straightforward but lengthy inductive calculation shows that if

$$F(a, c, e, g) = a^2g^2 - 7aceg + ae^3 + c^3g + 8c^2e^2,$$

then $F(a_n, a_{n+2}, a_{n+4}, a_{n+6}) = 0$ for all n . Since $F(a, c, e, g) = 0$, we know that $r_x - F(a, c, e, g) = r_x$. Therefore, we know that $r_x = \frac{-cg+e^2}{e^2}$.

Denote the y -coordinate of $(m+2)P + Q$ as r_y . We compute that $r_y = \frac{g(ag-3ce)}{e^3}$. Using that $r_y = r_y - \frac{F(a, c, e, g)}{ae^3}$, we find that $r_y = \frac{4ceg - c^2e - e^3}{e^3}$. Therefore, it is evident that

$$(m+2)P + Q = \left(\frac{a_{m+4}^2 - a_{m+2}a_{m+6}}{a_{m+4}^2}, \frac{4a_{m+2}a_{m+4}a_{m+6} - a_{m+2}^2a_{m+8} - a_{m+4}^3}{a_{m+4}^3} \right). \quad \square$$

Let E' be given by $E' : y^2 + xy = x^3 + x^2 + 8x + 10$ and let $R = (1, 4) \in E'(\mathbb{Q})$. We have a 2-isogeny $\phi : E \rightarrow E'$ given by

$$\phi(x, y) = \left(\frac{x^2 - 2}{x}, \frac{x^2y + 2x + 2y}{x^2} \right).$$

Theorem 4. *If p is a prime that divides a term in the Somos-5 sequence, the order of $P = (2, 2)$ in $E(\mathbb{F}_p)$ is twice the order of $R = (1, 4)$ in $E'(\mathbb{F}_p)$. Otherwise, their order is the same.*

Proof. If p divides a term in our sequence, say a_m , we know from our previous lemma that the denominators $(m-2)P+Q$ are divisible by p . Therefore, modulo p , $(m-2)P+Q=0$. The point Q has order 2, so adding Q to both sides we know that $(m-2)P=Q$. Therefore, we can deduce that $Q \in \langle P \rangle$. We have $\ker(\phi) = \{Q, 0\}$ (see Section 3.4 of [14]). Therefore, if ϕ is restricted to the subgroup generated by P , we have $|\ker(\phi)| = 2$. Since $\phi(P) = R$, by the first isomorphism theorem for groups, $\frac{|\langle P \rangle|}{|\ker(\phi)|} = |\langle R \rangle|$. It follows that $|P| = 2 \cdot |R|$.

Alternatively, assume p does not divide a term in the Somos-5 sequence. So, there is no m such that $mP + Q = 0$ modulo p , which implies that $Q \notin \langle P \rangle$. Therefore, the kernel of ϕ restricted to $\langle P \rangle$ is $\{0\}$ and so $|P| = |\phi(P)| = |R|$. \square

4. GALOIS REPRESENTATIONS

Denote by $E[2^r]$ the set of points on E with order dividing 2^r . Denote K_r as the field obtained by adjoining to \mathbb{Q} all x and y coordinates of points β with $2^r\beta = P$. For a prime p that is unramified in K_r , let $\sigma = \left[\frac{K_r/\mathbb{Q}}{\mathfrak{p}_i} \right]$ for some prime ideal \mathfrak{p}_i above p . Given a basis $\langle A, B \rangle$ for $E[2^r]$, for any such $\sigma \in \text{Gal}(K_r/\mathbb{Q})$, we have $\sigma(\beta) = \beta + eA + fB$. Also, $\sigma(A) = aA + bB$ and $\sigma(B) = cA + dB$. Define the map $\rho_{E,2^k} : \text{Gal}(K_r/\mathbb{Q}) \rightarrow \text{AGL}_2(\mathbb{Z}/2^k\mathbb{Z})$ by $\rho_{E,2^k}(\sigma) = (\vec{v}, M)$ where $M = \begin{bmatrix} a & b \\ c & d \end{bmatrix}$ and $\vec{v} = [e \ f]$. Let $\tau : \text{Gal}(K_r/\mathbb{Q}) \rightarrow \text{GL}_2(\mathbb{Z}/2^k\mathbb{Z})$ be given by $\tau(\sigma) = M$.

Let $S = \{\beta \in E(\mathbb{C}) : m \cdot \beta \in E(K)\}$ and let L be the field obtained by adjoining all x and y coordinates of points in S to K . Then the only primes p that ramify in L/K are those that divide m and those where E/K has bad reduction (see Proposition VIII.1.5(b) in [13]). For $E : y^2 + xy = x^3 + x^2 - 2x$, the conductor of E is $102 = 2 \cdot 3 \cdot 17$ and so the only primes that ramify in K_r/\mathbb{Q} are 2, 3 and 17.

Note that, if p is unramified, there are multiple primes \mathfrak{p}_i above p which could result in different matrices M_i and \vec{v}_i . However, properties we consider of these \vec{v}_i and M_i do not depend on the specific choice of \mathfrak{p}_i . The map depends on the choice of basis for $E[2^r]$, we choose this basis as described below in Theorem 7.

Let $\beta_r \in E(\mathbb{C})$ be a point with $2^r\beta_r = P$. We say that β_r is an r^{th} preimage of P under multiplication by 2. Let p be a prime with $p \neq 2, 3$ or 17 , $\sigma = \left[\frac{K_r/\mathbb{Q}}{\mathfrak{p}_i} \right]$, and $(\vec{v}, M) = \rho_{E,2^r}(\sigma)$. Assume that $\det(I - M) \not\equiv 0 \pmod{2^r}$. This implies that $\#E(\mathbb{F}_p) \not\equiv 0 \pmod{2^r}$.

Theorem 5. *Assume the notation above. Then 2^hP has odd order in $E(\mathbb{F}_p)$ if and only if $2^h\vec{v}$ is in the image of $I - M$.*

Proof. First, assume $2^h\vec{v}$ is in the image of $I - M$. This means that $\vec{x} = 2^h\vec{v} + \vec{x}M$ for some row vector \vec{x} with coordinates in $(\mathbb{Z}/2^r\mathbb{Z})^2$. If this is true for $\vec{x} = [e \ f]$, define $C := 2^h\beta_r + eA + fB$. We know then that $\sigma(C) = C$. We have $2^rC = 2^h(2^r\beta_r) = 2^hP$. If $|C|$ is odd, then clearly $|2^rC| = |2^hP|$ is also odd.

If $|C|$ is even, then every multiplication of C by 2 cuts the order by a factor of 2 until we arrive at a point of odd order. Since $|E(\mathbb{F}_p)| \equiv \det(I - M) \not\equiv 0 \pmod{2^r}$, the power of 2 dividing $|C|$ is also less than r , and so $|2^rC| = |2^hP|$ is odd.

Conversely, assume that $|2^hP|$ is odd. Let a be the multiplicative inverse of 2^r modulo $|2^hP|$ and define $C := a2^hP \in E(\mathbb{F}_p)$. Then $2^rC = 2^hP$ and so $2^r(C - 2^h\beta_r) = 0$. It follows that $C = 2^h\beta_r + yA + zB \in E(\mathbb{F}_p)$ for some $y, z \in \mathbb{Z}/2^r\mathbb{Z}$.

This implies that there is a Frobenius automorphism $\sigma \in \text{Gal}(K_r/\mathbb{Q})$ for which $\sigma(C) \equiv C \pmod{\mathfrak{p}_i}$ for some prime ideal \mathfrak{p}_i above p .

We claim that $\sigma(C) = C$ (as elements of $E(K_r)$). Note that $\sigma(C) - C \in E[2^r]$ and $\sigma(C) - C$ reduces to the identity modulo \mathfrak{p}_i . Since reduction is injective on torsion points of order coprime to the characteristic, and p is odd, it follows that $\sigma(C) = C$. It follows that if $\rho_{E,2^r}(\sigma) = (\vec{v}, M)$ then $2^h \vec{v} = (I - M)[y \ z]$, which implies that $2^h \vec{v}$ is in the image of $I - M$. \square

The following corollary is immediate.

Corollary 6. *Let o be the smallest positive integer so that $2^o \vec{v} = (I - M)\vec{x}$ for some \vec{x} with entries in $(\mathbb{Z}/2^r\mathbb{Z})^2$. Then 2^o is the highest power of 2 dividing $|P|$.*

The following theorem gives a convenient choice of basis for $E[2^k]$ and $E'[2^k]$.

Theorem 7. *Given a positive integer k , there are points $A_k, B_k \in E(\mathbb{C})$ that generate $E[2^k]$ and points $C_k, D_k \in E'(\mathbb{C})$ that generate $E'[2^k]$ so that $\phi(A_k) = C_k$ and $\phi(B_k) = 2D_k$. These points also satisfy the relations:*

$$2A_k = A_{k-1}, \quad 2B_k = B_{k-1}, \quad 2C_k = C_{k-1}, \quad \text{and} \quad 2D_k = D_{k-1}.$$

Proof. We will prove this by induction. Recall that $\phi : E \rightarrow E'$ is the isogeny with $\ker \phi = \{0, T\}$ where $T = (0, 0)$. Let $\phi' : E' \rightarrow E$ be the dual isogeny, and note that $\phi \circ \phi'(P) = 2P$. *Base Case:* Let $k = 1$. We want to find $\langle A_1, B_1 \rangle$ to generate $E[2]$ and $\langle C_1, D_1 \rangle$ to generate $E'[2]$ so that $\phi(A_1) = C_1$ and $\phi(B_1) = 2D_1$. We set $B_1 = (0, 0)$, and choose A_1 to be any non-identity point in $E[2]$ other than $(0, 0)$. We set $C_1 = \phi(A_1) = (-5/4, 5/8)$ and choose D_1 to be any non-identity point in $E'[2]$ other than C_1 . Note that $\phi'(D_1) = B_1$.

Inductive Hypothesis: Assume $\langle A_k, B_k \rangle = E[2^k]$ and $\langle C_k, D_k \rangle = E'[2^k]$ so that $\phi(A_k) = C_k$, $\phi(B_k) = 2D_k$, and $\phi'(D_k) = B_k$. Moreover, $D_k \notin \phi(E[2^k])$.

Since $|\ker \phi| = 2$, we have that $\phi(E[2^{k+1}]) \supseteq E'[2^k]$. Hence, we can choose B_{k+1} so that $\phi(B_{k+1}) = D_k$. Then $2B_{k+1} = \phi'(\phi(B_{k+1})) = \phi'(D_k) = B_k$. We choose D_{k+1} so that $\phi'(D_{k+1}) = B_{k+1}$. Note that $2D_{k+1} = \phi(B_{k+1}) = D_k$ and so $D_{k+1} \in E'[2^{k+1}]$. Now we pick A_{k+1} so that $2A_{k+1} = A_k$ and define $C_{k+1} = \phi(A_{k+1})$.

By our Inductive Hypothesis, $\langle A_k, B_k \rangle = E[2^k]$. This implies that $\langle A_k \rangle \cap \langle B_k \rangle = 0$, which in turn implies that $\langle 2A_{k+1} \rangle \cap \langle 2B_{k+1} \rangle = 0$. Let $C \in \langle A_{k+1} \rangle \cap \langle B_{k+1} \rangle$. Then, $C = aA_{k+1} = bB_{k+1}$. Because $|g^m| = \frac{|g|}{\gcd(m, |g|)}$, $|c| = \frac{2^{k+1}}{2^{\text{ord}_2(a)}} = \frac{2^{k+1}}{2^{\text{ord}_2(b)}}$, where $\text{ord}_2(n)$ is the highest power of 2 dividing n , it follows that either a and b are both even, or they are both odd. If a and b are even, then $C \in \langle A_k \rangle \cap \langle B_k \rangle = 0$, which is a contradiction. If a and b are odd, then $|C| = 2^{k+1}$ but $2C \in \langle A_k \rangle \cap \langle B_k \rangle = 0$, which is also a contradiction. It follows that $\langle A_{k+1} \rangle \cap \langle B_{k+1} \rangle = 0$, which gives that $E[2^{k+1}] = \langle A_{k+1}, B_{k+1} \rangle$.

Now we show that $\langle C_{k+1}, D_{k+1} \rangle = E'[2^{k+1}]$, by way of showing that $\langle C_{k+1} \rangle \cap \langle D_{k+1} \rangle = 0$. We have shown that $\langle A_{k+1}, B_{k+1} \rangle = E[2^{k+1}]$, and so $\phi(E[2^{k+1}]) = \langle C_{k+1}, 2D_{k+1} \rangle$. We want to show that $D_{k+1} \notin \phi(E[2^{k+1}])$.

If $D_{k+1} \in \phi(E[2^{k+1}])$, then $D_{k+1} = aC_{k+1} + (2b-1)D_{k+1} = 0$. Since $(2b-1)$ is odd, $(2b-1)D_{k+1}$ has order dividing 2^{k+1} . Hence, aC_{k+1} has

order dividing 2^{k+1} . We can then see that

$$\begin{aligned} 2aC_{k+1} + 2(2b-1)D_{k+1} &= 0 \\ aC_k + (2b-1)D_k &= 0 \\ \implies a &\equiv (2b-1) \equiv 0 \pmod{2^k}, \end{aligned}$$

which is a contradiction. This implies that $\phi(E[2^{k+1}])$ is an index 2 subgroup of $\langle C_{k+1}, D_{k+1} \rangle$ of order 2^{2k+1} , and so $\langle C_{k+1}, D_{k+1} \rangle = E'[2^{k+1}]$. This proves the desired claim. \square

Recall the maps $\rho_{E,2^k} : \text{Gal}(K_k/\mathbb{Q}) \rightarrow \text{AGL}_2(\mathbb{Z}/2^k\mathbb{Z})$ and $\tau : \text{Gal}(K_k/\mathbb{Q}) \rightarrow \text{GL}_2(\mathbb{Z}/2^k\mathbb{Z})$, defined at the beginning of this section. In [12], an algorithm is given to compute the image of the 2-adic Galois representation τ . Running this algorithm shows that the image of τ (up to conjugacy) is the index 6 subgroup of $\text{GL}_2(\mathbb{Z}/2^k\mathbb{Z})$ generated by $\begin{bmatrix} 1 & 1 \\ 0 & 1 \end{bmatrix}$, $\begin{bmatrix} 7 & 0 \\ 2 & 1 \end{bmatrix}$, and $\begin{bmatrix} 5 & 0 \\ 2 & 1 \end{bmatrix}$. Moreover, the subgroup generated by the aforementioned matrices is the unique conjugate that corresponds to the basis chosen in Theorem 7.

Theorem 8. *If $\rho_{E,2^k}(\sigma) = (\vec{v}, M)$ where $\vec{v} = (e, f)$, then $e \equiv 0 \pmod{2}$ if and only if $M \equiv 1, 7 \pmod{8}$.*

Proof. We will show that $e \equiv 0 \pmod{2}$ and $M \equiv 1, 7 \pmod{8}$ if and only if $\sigma(\sqrt{2}) = \sqrt{2}$.

Let β_1 be a point in $E(K_1)$ so that $2\beta_1 = (2, 2)$. We pick a basis $\langle A_1, B_1 \rangle$ according to Theorem 7. We have $\sigma(\beta_1) = \beta_1 + eA_1 + fB_1$, where $e, f \in \mathbb{Z}/2^k\mathbb{Z}$.

Let $\phi : E \rightarrow E'$ be the usual isogeny and note that $B_1 \in \ker \phi$. Thus, $\phi(\sigma(\beta_1)) = \phi(\beta_1 + eA_1 + fB_1) = \phi(\beta_1) + e\phi(A_1)$. It follows that $e \equiv 0 \pmod{2}$ if and only if $\sigma(\phi(\beta_1)) = \phi(\sigma(\beta_1)) = \phi(\beta_1)$. A straightforward computation shows that the coordinates of $\phi(\beta_1)$ generate $\mathbb{Q}(\sqrt{2})$. It follows that $e \equiv 0 \pmod{2}$ if and only if $\sigma(\sqrt{2}) = \sqrt{2}$.

Finally, suppose that σ is the Artin symbol associated to a prime ideal \mathfrak{p} above a rational prime p . By properties of the Weil pairing (see [13], Section III.8), we have that $\zeta_{2^k} = e^{2\pi i/2^k} \in \mathbb{Q}(E[2^k])$, and that $\sigma(\zeta_{2^k}) = \zeta_{2^k}^{\det(M)} = \zeta_{2^k}^p$. Since $\sqrt{2} = \zeta_8 + \zeta_8^{-1}$, it follows easily that $\sigma(\sqrt{2}) = \sqrt{2} \iff p \equiv 1, 7 \pmod{8}$ and hence $\sigma(\sqrt{2}) = \sqrt{2}$ if and only if $\det(M) \equiv 1, 7 \pmod{8}$. \square

For $k \geq 3$, define I_k to be the subgroup of $\text{AGL}_2(\mathbb{Z}/2^k\mathbb{Z})$ whose elements are ordered pairs $\{(\vec{v}, M)\}$ where $\vec{v} = [e \ f]$, the reduction of $M \pmod{8}$ is in the group generated by $\begin{bmatrix} 1 & 1 \\ 0 & 1 \end{bmatrix}$, $\begin{bmatrix} 7 & 0 \\ 2 & 1 \end{bmatrix}$, and $\begin{bmatrix} 5 & 0 \\ 2 & 1 \end{bmatrix}$, and $e \equiv 0 \pmod{2}$ if and only if $\det(M) \equiv 1$ or $7 \pmod{8}$. By Theorem 8 and the discussion preceding it, we know that the image of $\rho : \text{Gal}(K_k/\mathbb{Q}) \rightarrow \text{AGL}_2(\mathbb{Z}/2^k\mathbb{Z})$ is contained in I_k .

We now aim to show that the map $\rho_{E,2^k} : \text{Gal}(K_k/\mathbb{Q}) \rightarrow I_k$ is surjective for $k \geq 3$. By [13] (page 105), if we have an elliptic curve $E : y^2 = x^3 + Ax + B$, the division polynomial $\psi_m \in \mathbb{Z}[A, B, x, y]$ is determined recursively by:

$$\begin{aligned} \psi_1 &= 1, \psi_2 = 2y, \psi_3 = 3x^4 + 6Ax^2 + 12Bx - A^2, \\ \psi_4 &= 4y(x^6 + 5Ax^4 + 20Bx^3 - 5A^2x^2 - 4ABx - 8B^2 - A^3), \\ \psi_{2m+1} &= \psi_{m+2}\psi_m^3 - \psi_{m-1}\psi_{m+1}^3, \quad 2y\psi_{2m} = \psi_m(\psi_{m+1}\psi_{m-1}^2 - \psi_{m-2}\psi_{m+1}^2). \end{aligned}$$

We then define ϕ_m and ω_m as follows:

$$\begin{aligned}\phi_m &= x\psi_m^2 - \psi_{m+1}\psi_{m-1} \\ 4y\omega_m &= \psi_{m+2}\psi_{m-1}^2 - \psi_{m-2}\psi_{m+1}^2.\end{aligned}$$

If $\Delta = -16(4A^3 + 27B^2) \neq 0$, then $\phi_m(x)$ and $\psi_m(x)^2$ are relatively prime. This also implies that, for $P = (x_0, y_0) \in E$,

$$[m]P = \left(\frac{\phi_m(P)}{\psi_m(P)^2}, \frac{\omega_m(P)}{\psi_m(P)^3} \right).$$

Lemma 9. *The map $\rho_{E,8} : \text{Gal}(K_3, \mathbb{Q}) \rightarrow I_3$ is surjective.*

Proof. The curve E is isomorphic to $E_2 : y^2 = x^3 - 3267x + 45630$. The isomorphism that takes E to E_2 takes $P = (2, 2)$ on E to $P_2 = (87, 648)$ on E_2 .

We use division polynomials to construct a polynomial $f(x)$ whose roots are the x -coordinates of points β_3 on E_2 so that $8\beta_3 = P_2$. By the above formulas, $8P_2 = \left(\frac{\phi_8(P_2)}{\psi_8(P_2)^2}, \frac{\omega_8(P_2)}{\psi_8(P_2)^3} \right)$. Since $P_2 = (87, 648)$,

$$f(x) = \phi_8(P_2) - 87\psi_8(P_2)^2 = 0$$

will yield the equation with roots that satisfy our requirement. This is a degree 64 polynomial. By using Magma to compute the Galois Group of $f(x)$, we find the order to be 8192. A simple calculation shows that I_3 has order 8192 and since $f(x)$ splits in K_3/\mathbb{Q} , we have that $\text{Gal}(K_3/\mathbb{Q}) \cong I_3$. \square

To prove the surjectivity of $\rho_{E,2^k}$, we will consider the Frattini subgroup of I_k . This is the intersection of all maximal subgroups of I_k . Since I_k is a 2-group, every maximal subgroup is normal and has index 2. It follows from this that if $g \in I_k$, then $g^2 \in \Phi(I_k)$.

Lemma 10. *For $3 \leq k$, $\Phi(I_k)$ contains all pairs (\vec{v}, M) such that $\vec{v} \equiv \vec{0} \pmod{4}$ and $M \equiv I \pmod{8}$.*

Proof. We begin by observing that for $r = k$, $(0, I) \in \Phi(I_k)$. We prove the result by backwards induction on r .

Inductive Hypothesis: $\Phi(I_k)$ contains all pairs $(0, M)$, $M \equiv I \pmod{2^r}$. Let $g = I + 2^{r-2}N$, and let $h = I + 2^{r-1}N$. If $r \geq 5$, then a straightforward calculation shows that $(0, g) \in I_k$. So, $(0, g)^2 = (0, g^2) \in \Phi(I_k)$. Therefore, for $r > 3$,

$$g^2 = I + 2^{r-1}N + 2^{2r-4}N^2 \equiv h \pmod{2^{2r-4}}.$$

By the induction hypothesis, $(0, g^2h^{-1}) \in \Phi(I_k)$, and so $(0, h) \in \Phi(I_k)$.

So, for $k \geq r \geq 4$, all pairs $(0, M)$, $M \equiv I \pmod{2^r} \in \Phi(I_k)$. We will now construct I_4 , compute $\Phi(I_4)$, and determine if $\Phi(I_4) \in \{(\vec{v}, M) : \vec{v} \equiv 0 \pmod{8}, M \equiv I \pmod{8}\}$. A computation with Magma shows that

$$I_4 = \left\langle \left[\begin{array}{ccc} 1 & 1 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{array} \right], \left[\begin{array}{ccc} 7 & 0 & 0 \\ 2 & 1 & 0 \\ 0 & 0 & 1 \end{array} \right], \left[\begin{array}{ccc} 5 & 0 & 0 \\ 2 & 1 & 0 \\ 1 & 0 & 1 \end{array} \right] \right\rangle.$$

We then construct $\Phi(I_4)$ and then $\phi : \Phi(I_4) \rightarrow \text{GL}_3(\mathbb{Z}/8\mathbb{Z})$. We check that $\ker \phi$ has order 64 and this proves the desired claim about $\Phi(I_4)$.

Now, observe that if $\vec{v}_1 = (2x, 2y)$, then $(\vec{v}_1, I) \in I_k$ and so $(2\vec{v}_1, I) = (\vec{v}_1, I)^2 \in \Phi(I_k)$, and so $\Phi(I_k)$ contains all pairs (\vec{v}, I) with $\vec{v} \equiv \vec{0} \pmod{4}$. Finally, for any matrix $M \equiv I \pmod{8}$, we have

$$(\vec{v}_1, I) * (0, M) = (\vec{v}_1, M) \in \Phi(I_k)$$

and this proves the desired claim. \square

Finally, we prove the desired surjectivity.

Theorem 11. *The map $\rho_{E, 2^k} : \text{Gal}(K_k/\mathbb{Q}) \rightarrow I_k$ is surjective for all $k \geq 3$.*

Proof. Suppose to the contrary the map is not surjective. Lemma 10 implies that if $\rho_{E, 2^k}$ is not surjective, the image lies in a maximal subgroup M which contains the kernel of the map from $I_k \rightarrow I_3$, and so the image of $\rho_{E, 8}$ must lie in a maximal subgroup of I_3 . This contradicts Lemma 9, and shows the map is surjective. \square

Now, we indicate the relationship between $\rho_{E, 2^k}$ and $\rho_{E', 2^k}$. Let $\sigma \in \text{Gal}(K_k/\mathbb{Q})$. If β_k is chosen so $2^k \beta_k = P$, then

$$\begin{aligned}\sigma(A_k) &= aA_k + bB_k, \\ \sigma(B_k) &= cA_k + dB_k, \\ \sigma(\beta_k) &= \beta_k + eA_k + fB_k.\end{aligned}$$

Applying ϕ to these equations, we have

$$\begin{aligned}\phi(\sigma(A_k)) &= aC_k + 2bD_k = \sigma(\phi(A_k)) = \sigma(C_k), \\ \phi(\sigma(B_k)) &= cC_k + 2dD_k = \sigma(\phi(B_k)) = \sigma(2D_k), \\ \phi(\sigma(\beta_k)) &= \phi(\beta_k) + eC_k + 2fD_k = \sigma(\phi(\beta_k)) = \sigma(\beta'_k),\end{aligned}$$

where $2^k \beta'_k = R$ on E' . Using the relations from Theorem 7, we have that $2D_k = D_{k-1}$ and $2C_k = C_{k-1}$. This gives

$$\begin{aligned}\sigma(C_{k-1}) &= aC_{k-1} + 2bD_{k-1}, \\ \sigma(D_{k-1}) &= \frac{c}{2}C_{k-1} + dD_{k-1}.\end{aligned}$$

Thus, the vector-matrix pair associated with $\rho_{E', 2^{k-1}}$ is (v', M') , where $\vec{v}' = [e \quad 2f]$ and $M' = \begin{bmatrix} a & 2b \\ \frac{c}{2} & d \end{bmatrix}$.

Let (v, M) be a vector-matrix pair in I_k . Suppose that o is the smallest non-negative integer so that $2^o \vec{v}$ is in the image of $(I - M)$. Thus there are integers c_1 and c_2 (not necessarily unique) so that $2^o \vec{v} = c_1 \vec{x}_1 + c_2 \vec{x}_2$, where \vec{x}_1 and \vec{x}_2 are the first and second rows of $I - M$.

Lemma 12. *Assume that $\det(M - I) \not\equiv 0 \pmod{2^k}$. If $c_1 \vec{x}_1 + c_2 \vec{x}_2 = d_1 \vec{x}_1 + d_2 \vec{x}_2$, then $c_1 \equiv d_1 \pmod{2}$.*

Proof. The assumption on $\det(M - I)$ implies that $\ker(M - I)$ has order dividing 2^{k-1} . However, if $c_1 \vec{x}_1 + c_2 \vec{x}_2 = d_1 \vec{x}_1 + d_2 \vec{x}_2$, then $[c_1 - d_1 \quad c_2 - d_2]$ is an element of $\ker(M - I)$. If $c_1 \not\equiv d_1 \pmod{2}$, then this element has order 2^k , which is a contradiction. \square

The above lemma makes it so we can speak of $c_1 \pmod{2}$ unambiguously. We now have the following result.

Theorem 13. *Assume the notation above. Let o' be the smallest positive integer so that $2^{o'}v'$ is in the image of $I - M$. If $\det(M - I) \not\equiv 0 \pmod{2^{k-1}}$, then $o \neq o'$ if and only if c_1 is even.*

Proof. Let \vec{y}_1 and \vec{y}_2 be the first two rows of $I - M'$. A straightforward calculation shows that if $2^o\vec{v} = c_1\vec{x}_1 + c_2\vec{x}_2$, then $2^o\vec{v}' = c_1\vec{y}_1 + 2c_2\vec{y}_2$. If c_1 is even, then it follows that $2^{o-1}\vec{v}' = (c_1/2)\vec{y}_1 + c_2\vec{y}_2$ and so $o \neq o'$.

Conversely, if $o \neq o'$, then $o' = o - 1$ and so $2^{o-1}\vec{v}' = d_1\vec{y}_1 + d_2\vec{y}_2$. We have then that

$$2^o\vec{v} \equiv 2d_1\vec{x}_1 + d_2\vec{x}_2 \pmod{2^{k-1}}.$$

So if $\vec{x} = \begin{bmatrix} 2d_1 \\ d_2 \end{bmatrix}$ we have $\vec{x}(I - M) \equiv 2^o\vec{v} \pmod{2^{k-1}}$. If there is a vector \vec{x}' with $\vec{x} \not\equiv \vec{x}' \pmod{2}$ so that $\vec{x}'(I - M) \equiv 2^o\vec{v} \pmod{2^{k-1}}$, then $\vec{x} - \vec{x}'$ is in the kernel of $I - M \pmod{2^{k-1}}$. However, the order of $\vec{x} - \vec{x}'$ is 2^{k-1} and this contradicts the condition on the determinant. This proves the desired result. \square

5. PROOF OF THEOREM 1

Theorem 4 states that a prime p divides a term in the Somos-5 sequence if and only if the order of $(2, 2) \in E(\mathbb{Q})$ is different than the order of $(1, 4) \in E'(\mathbb{Q})$. Recall that o , the power of two dividing the order of P , is the smallest positive integer such that $2^o\vec{v} \in \text{im}(I - M)$, and o' is the power of two dividing the order of R .

Suppose that $\det(I - M) \not\equiv 0 \pmod{2^{k-1}}$. We have $2^o\vec{v} \in \text{im}(I - M)$ if and only if $c_1\vec{x}_1 + c_2\vec{x}_2 = 2^o\vec{v}$, where $M = \begin{bmatrix} a & b \\ c & d \end{bmatrix}$, $\vec{x}_1 = [1 - a \quad -b]$, and $\vec{x}_2 = [-c \quad 1 - d]$. We know that $o \neq o'$ if and only if c_1 is even. For the remainder of the argument,

we will consider elements of I_k as 3×3 matrices $\begin{bmatrix} a & b & 0 \\ c & d & 0 \\ e & f & 1 \end{bmatrix}$. We let $I - M =$

$\begin{bmatrix} \alpha & \beta & 0 \\ \gamma & \delta & 0 \\ e & f & 0 \end{bmatrix}$ and define $A = \gamma f - \delta e$, $B = \alpha f - \beta e$, and $C = \alpha \delta - \beta \gamma$. We define

$M_3^0(\mathbb{Z}/2^k\mathbb{Z})$ to be the set of 3×3 matrices with entries in $\mathbb{Z}/2^k\mathbb{Z}$ whose third column is zero. We will use $\text{ord}_2(r)$ to denote the highest power of 2 dividing r for $r \in \mathbb{Z}/2^k\mathbb{Z}$. If $r = 0 \in \mathbb{Z}/2^k\mathbb{Z}$, we will interpret $\text{ord}_2(r)$ to have an undefined value, but we will declare the inequality $\text{ord}_2(r) \geq k$ to be true.

Solving the equation $c_1\vec{x}_1 + c_2\vec{x}_2 = 2^o\vec{v}$ using Cramer's rule gives that $c_1C = -2^oA$ and $c_2C = 2^oB$. Assuming that c_1 is even and $o > 0$ implies that c_2 must be odd, and this implies that $\text{ord}_2(B) < \text{ord}_2(C)$. Moreover, since the power of 2 dividing c_1C must be higher than that dividing c_2C it follows that $\text{ord}_2(B) < \text{ord}_2(A)$. Conversely, if $\text{ord}_2(B) < \text{ord}_2(A)$ and $\text{ord}_2(B) < \text{ord}_2(C)$, then $o > 0$ and c_1 is even. Therefore, our goal is counting of elements of I_k with $\text{ord}_2(A) > \text{ord}_2(B)$ and $\text{ord}_2(C) > \text{ord}_2(B)$. For an $M_0 \in M_3^0(\mathbb{Z}/2^r\mathbb{Z})$, define

$$\eta(M_0, r, k) = \# \{ M \in M_3^0(\mathbb{Z}/2^k\mathbb{Z}) : M \equiv M_0 \pmod{2^r}, \text{ord}_2(A), \text{ord}_2(C) > \text{ord}_2(B) \},$$

$$\mu(M_0, r) = \lim_{k \rightarrow \infty} \frac{\eta(M_0, r, k)}{|I_3| \cdot 64^{k-3}}.$$

Roughly speaking, $\mu(M_0, r)$ is the fraction of matrices $M \equiv M_0 \pmod{2^r}$ in I_k with the property that $\rho_{E, 2^k}(\sigma_p) = M$ implies that p divides a term of the Somos-5 sequence.

Theorem 14. *We have*

$$\lim_{x \rightarrow \infty} \frac{\pi'(x)}{\pi(x)} = \sum_{M \in I_3} \mu(I - M, 3).$$

Before we start the proof, we need some lemmas. The first is straightforward, and we omit its proof.

Lemma 15. *If $a \in \mathbb{Z}/2^k\mathbb{Z}$, then the number of pairs $(x, y) \in (\mathbb{Z}/2^k\mathbb{Z})^2$ with $xy \equiv a \pmod{2^k}$ is $(\text{ord}_2(a) + 1)2^{k-1}$, where if $a \equiv 0 \pmod{2^k}$, we take $\text{ord}_2(a) = k + 1$.*

Lemma 16. *The number of matrices $M \in M_2(\mathbb{Z}/2^k\mathbb{Z})$ with $\det(M) \equiv 0 \pmod{2^k}$ is $3 \cdot 2^{3k-1} - 2^{2k-1}$.*

Proof. We counting quadruples (a, b, c, d) with $ad \equiv bc \pmod{2^k}$. By Lemma 15, this number is equal to

$$\sum_{\alpha \in \mathbb{Z}/2^k\mathbb{Z}} ((\text{ord}_2(\alpha) + 1)2^{k-1})^2,$$

which can easily be shown to equal $3 \cdot 2^{3k-1} - 2^{2k-1}$. \square

Proof of Theorem 14. For $k \geq 1$, let $G = \text{Gal}(K_k/\mathbb{Q})$ and $\sigma \in G$ have the property that $\sigma = \left[\frac{K_k/\mathbb{Q}}{\mathfrak{p}} \right]$ for some prime ideal $\mathfrak{p} \subseteq O_{K_k}$ with $\mathfrak{p} \cap \mathbb{Z} = (p)$. Assume that p is unramified in K_k/\mathbb{Q} and E/\mathbb{F}_p has good reduction at p . Let M be the 3×3 matrix corresponding to $\rho_{E, 2^k}(\sigma)$, and A, B and C be the corresponding minors of $I - M$. Then one of three alternatives occurs:

(a) $B \not\equiv 0 \pmod{2^k}$, and a higher power of 2 divides both A and C .

In this situation (the good case), previous results ensure that the order of P in $E(\mathbb{F}_p)$ is twice the order of R in $E'(\mathbb{F}_p)$, and hence p divides some term in the Somos-5 sequence.

(b) One of A or C is not congruent to 0 mod 2^k and the power of 2 dividing B is equal or higher than for A or C .

In this situation (the bad case), previous results ensure that the order of P in $E(\mathbb{F}_p)$ is equal to the order of R in $E'(\mathbb{F}_p)$ and p does not divide any term in the Somos-5 sequence.

(c) $A \equiv B \equiv C \equiv 0 \pmod{2^k}$.

In this situation (the inconclusive case), we do not have enough information to determine if p divides a term in the Somos-5 sequence or not.

Fix $\epsilon > 0$ and choose a k large enough so that both of the following conditions are satisfied:

(i) $\left| \sum_{M \in I_3} \frac{\eta(I-M, 3, k)}{|I_3|64^{k-3}} - \sum_{M \in I_3} \mu(I - M, 3) \right| < \epsilon/3$, and

(ii) the fraction of elements M in I_3 with $C \equiv \det(I - M) \equiv 0 \pmod{2^{k-1}}$ is less than $\epsilon/3$. (This fraction tends to zero by Lemma 16.)

Let $\mathcal{C} \subseteq I_k$ be the collection of “good” elements of I_k and let \mathcal{C}' be the collection of “good or inconclusive” elements.

By the statements above, we have that

$$\sum_{M \in I_3} \mu(I - M, 3) - 2\epsilon/3 < \frac{|\mathcal{C}|}{|I_k|}$$

and

$$\frac{|\mathcal{C}'|}{|I_k|} < \sum_{M \in I_3} \mu(I - M, 3) + \epsilon/3.$$

By the Chebotarev density theorem, we have

$$\lim_{x \rightarrow \infty} \frac{\#\{p \text{ prime} : p \leq x \text{ is unramified in } K_k \text{ and } [\frac{K_k/\mathbb{Q}}{p}] \subseteq \mathcal{C}\}}{\pi(x)} = \frac{|\mathcal{C}|}{|I_k|},$$

and the same with \mathcal{C}' .

Let r be the number of primes that either ramify in K_k/\mathbb{Q} or for which E/\mathbb{Q} has bad reduction. Then there is a constant N so that if $x > N$, then

$$\sum_{M \in I_3} \mu(I - M, 3) - \epsilon + \frac{r}{\pi(x)} < \frac{\#\{p \text{ prime} : p \leq x \text{ is unramified in } K_k \text{ and } [\frac{K_k/\mathbb{Q}}{p}] \subseteq \mathcal{C}\}}{\pi(x)},$$

and

$$\frac{\#\{p \text{ prime} : p \leq x \text{ is unramified in } K_k \text{ and } [\frac{K_k/\mathbb{Q}}{p}] \subseteq \mathcal{C}'\}}{\pi(x)} < \sum_{M \in I_3} \mu(I - M, 3) + \epsilon - \frac{r}{\pi(x)}.$$

It follows from these inequalities that for $x > N$, then

$$-\epsilon < \frac{\pi'(x)}{\pi(x)} - \sum_{M \in I_3} \mu(I - M, 3) < \epsilon.$$

This proves that

$$\lim_{x \rightarrow \infty} \frac{\pi'(x)}{\pi(x)} = \sum_{M \in I_3} \mu(I - M, 3).$$

□

Our goal is now to compute $\sum_{M \in I_3} \mu(I - M, 3)$. To do this, we will develop rules to compute $\mu(M, r)$ for any matrix $M \in M_3(\mathbb{Z}/2^r\mathbb{Z})$ whose third column is zero. Observe that $\mu(M_0, r) \leq \frac{\#\{M \in M_3^0(\mathbb{Z}/2^r\mathbb{Z}) : M \equiv M_0 \pmod{2^r}\}}{|I_3| \cdot 64^{r-3}} = \frac{1}{2 \cdot 64^{r-1}}$.

Also, if all the entries in M are even, then $\mu(M, r) = \frac{1}{64} \mu(\frac{M}{2}, r-1)$. This allows us to reduce to matrices where at least one entry is odd. If $M \in M_3^0(\mathbb{Z}/2\mathbb{Z})$ is the zero matrix, we have

$$\mu(M, 1) = \frac{1}{64} \mu(M/2, 0) = \frac{1}{64} \sum_{N \in M_3^0(\mathbb{Z}/2\mathbb{Z})} \mu(N, 1) = \frac{1}{64} \mu(M, 1) + \sum_{\substack{N \in M_3^0(\mathbb{Z}/2\mathbb{Z}) \\ N \neq M}} \mu(N, 1).$$

It follows that $\mu(M, 1) = \frac{1}{63} \sum_{\substack{N \in M_3^0(\mathbb{Z}/2\mathbb{Z}) \\ N \neq M}} \mu(N, 1)$.

In order to determine $\mu(M_0, r)$, it is necessary to consider a matrix $M \in M_3(\mathbb{Z}/2^k\mathbb{Z})$ and examine the behavior of matrices $M' \in M_3(\mathbb{Z}/2^{k+1}\mathbb{Z})$ with $M' \equiv M \pmod{2^k}$. We refer to these as ‘lifts’ of M . We define A , B and C to be functions defined on

a matrix $M = \begin{bmatrix} \alpha & \beta & 0 \\ \gamma & \delta & 0 \\ e & f & 0 \end{bmatrix}$, given by $A = \gamma f - \delta e$, $B = \alpha f - \beta e$ and $C = \alpha \delta - \beta \gamma$.

Theorem 17. Let $M = \begin{bmatrix} \alpha & \beta & 0 \\ \gamma & \delta & 0 \\ e & f & 0 \end{bmatrix} \in M_3^0(\mathbb{Z}/2^k\mathbb{Z})$ and suppose $A \equiv B \equiv C \equiv 0 \pmod{2^k}$.

- (1) If γ or δ is odd, then $\mu(M, k) = 0$.
- (2) If γ and δ are both even, but one of α, β, e or f is odd, then $\mu(M, k) = \frac{1}{6 \cdot 64^{k-1}}$.

Proof. Suppose that $M \in M_3^0(\mathbb{Z}/2^k\mathbb{Z})$ is a matrix with γ or δ odd and with $A \equiv C \equiv 0 \pmod{2^k}$. In the case that γ is odd, the congruences $A \equiv 0 \pmod{2^k}$ and $C \equiv 0 \pmod{2^k}$ imply that $f \equiv \frac{e\delta}{\gamma} \pmod{2^k}$ and $\beta \equiv \frac{\alpha\delta}{\gamma} \pmod{2^k}$. We then find that $B \equiv \alpha f - \beta e \equiv \alpha \left(\frac{e\delta}{\gamma}\right) - \left(\frac{\alpha\delta}{\gamma}\right)e \equiv 0 \pmod{2^k}$. It follows that none of the lifts of M have $\text{ord}_2(B) < \min\{\text{ord}_2(A), \text{ord}_2(C)\}$ and so $\mu(M, k) = 0$. A similar argument applies in the case that δ is odd.

Suppose now that γ and δ are both even. Consider M' to be a lift of $M \pmod{2^{k+1}}$. Then we have

$$M' = \begin{bmatrix} \alpha' & \beta' & 0 \\ \gamma' & \delta' & 0 \\ e' & f' & 0 \end{bmatrix} = \begin{bmatrix} \alpha + \alpha_1 2^k & \beta + \beta_1 2^k & 0 \\ \gamma + \gamma_1 2^k & \delta + \delta_1 2^k & 0 \\ e + e_1 2^k & f + f_1 2^k & 0 \end{bmatrix},$$

where $\alpha_1, \beta_1, \gamma_1, \delta_1, e_1, f_1 \in \mathbb{F}_2$. If A', B' and C' are the values of A, B , and C associated to M' , then

$$\begin{aligned} A' &\equiv A + 2^k(\gamma_1 f - \delta_1 e) \pmod{2^{k+1}} \\ B' &\equiv B + 2^k(\alpha_1 f + \alpha f_1 - \beta_1 e - \beta e_1) \pmod{2^{k+1}} \\ C' &\equiv C + 2^k(\alpha \delta_1 - \beta \gamma_1) \pmod{2^{k+1}}. \end{aligned}$$

Suppose that e or f is odd. Then the map $\mathbb{F}_2^6 \rightarrow \mathbb{F}_2^2$ given by $(\alpha_1, \beta_1, \gamma_1, \delta_1, e_1, f_1) \mapsto (\gamma_1 f - \delta_1 e, \alpha_1 f + \alpha f_1 - \beta_1 e - \beta e_1)$ is surjective. It follows that of the 64 lifts of M' , one quarter have $(A' \pmod{2^{k+1}}, B' \pmod{2^{k+1}})$ equal to each of $(2^k, 2^k)$, $(0, 2^k)$, $(2^k, 0)$ and $(0, 0)$. Moreover, if $A' \equiv 0 \pmod{2^{k+1}}$, then we must have $C' \equiv 0 \pmod{2^{k+1}}$. This is because if e' is odd, then $\delta' \equiv \frac{\gamma' f'}{e'} \pmod{2^{k+1}}$, and $\beta' \equiv \frac{\alpha f - B'}{e'} \pmod{2^{k+1}}$. Plugging these into $C' = \alpha' \delta' - \beta' \gamma'$ gives $C' \equiv \frac{B' \gamma'}{e'} \pmod{2^{k+1}}$. Since γ' is even, it follows that $C' \equiv 0 \pmod{2^{k+1}}$. A similar argument shows that $C' \equiv 0 \pmod{2^{k+1}}$ if f' is odd. As a consequence, of the 64 lifts of M , 32 have $\mu(M', k+1) = 0$, 16 have $\text{ord}_2(B') < \text{ord}_2(A')$ and $\text{ord}_2(B') < \text{ord}_2(A')$. For these, we have $\mu(M', k+1) = \frac{1}{2 \cdot 64^k}$. The remainder have $A' \equiv B' \equiv C' \equiv 0 \pmod{2^{k+1}}$. It follows that

$$\mu(M, k) = \frac{1}{2 \cdot 64^{k-1}} \cdot \frac{1}{4} + \sum_{\substack{M' \equiv M \pmod{2^{k+1}} \\ A' \equiv B' \equiv C' \equiv 0 \pmod{2^{k+1}}}} \mu(M', k+1).$$

Applying the above argument repeatedly gives

$$\mu(M, k) = \frac{1}{2 \cdot 64^{k-1}} \cdot \left(\frac{1}{4} + \frac{1}{16} + \cdots + \frac{1}{4^\ell} \right) + \sum_{\substack{M' \equiv M \pmod{2^{k+\ell}} \\ A' \equiv B' \equiv C' \equiv 0 \pmod{2^{k+\ell}}}} \mu(M', k+\ell).$$

Using the bound $0 \leq \mu(M', k + \ell) \leq \frac{1}{2 \cdot 64^{k+\ell-1}}$, noting that the sum contains 16^ℓ terms, and taking the limit as $\ell \rightarrow \infty$ yields that $\mu(M, k) = \frac{1}{2 \cdot 64^{k-1}} \sum_{r=1}^{\infty} \frac{1}{4^r} = \frac{1}{6 \cdot 64^{k-1}}$.

The case when α or β is odd is very similar. In that case, one can show that the 64 lifts M' have $(B' \bmod 2^{k+1}, C' \bmod 2^{k+1})$ divided equally between $(2^k, 2^k)$, $(0, 2^k)$, $(2^k, 0)$ and $(0, 0)$, and that $C' \equiv 0 \pmod{2^{k+1}}$ implies that $A' \equiv 0 \pmod{2^{k+1}}$. Again, one quarter of the lifts M' have $B' \equiv 2^k \pmod{2^{k+1}}$ and $A' \equiv C' \equiv 0 \pmod{2^{k+1}}$, and $\mu(M, k) = \frac{1}{6 \cdot 64^{k-1}}$. \square

Let $M \in M_3^0(\mathbb{Z}/8\mathbb{Z})$ be the zero matrix. We have that $\mu(M, 3) = \frac{1}{64^2} \mu(M, 1) = \frac{1}{63} \cdot \frac{1}{64^2} \sum_{N \in M_3^0(\mathbb{Z}/2\mathbb{Z})} \mu(N, 1)$. Of the 63 nonzero matrices in $M_3^0(\mathbb{Z}/2\mathbb{Z})$ we find that 6 have B odd and A and C even, while 36 have A or C odd. Of the remaining 21, there are 12 that have γ or δ odd, and the remaining 9 have γ and δ both even. It follows that

$$\mu(M, 3) = \frac{1}{63} \cdot \frac{1}{64^2} \cdot \frac{1}{2} \cdot \left[6 + 36 \cdot 0 + 12 \cdot 0 + 9 \cdot \frac{1}{3} \right] = \frac{1}{8192} \cdot \frac{1}{7} = \frac{1}{57344}.$$

(Note that in the denominator of $\mu(N, 1)$ we have $|I_3|64^{-2} = 8192 \cdot (1/4096) = 2$.)

For each of the 8191 non-identity elements M of I_3 , we divide $I - M$ by the highest power of 2 dividing all of the elements, say 2^r . In 3754 cases, we have $\text{ord}_2(B) < \text{ord}_2(A)$ and $\text{ord}_2(B) < \text{ord}_2(C)$. For each of these, $\mu(I - M, 3) = \frac{1}{8192}$.

In 4036 cases, we have $\text{ord}_2(B) \geq \text{ord}_2(A)$ or $\text{ord}_2(B) \geq \text{ord}_2(C)$ and not all of A , B and C are congruent to 0 modulo 2^{3-r} . For each of these, $\mu(I - M, 3) = 0$.

In 365 cases, we have $A \equiv B \equiv C \equiv 0 \pmod{2^{3-r}}$ and γ and δ are both even. In each of these cases, $\mu(I - M, 3) = \frac{1}{3 \cdot 8192}$ by Theorem 17.

In the remaining 36 cases, we have $A \equiv B \equiv 0 \pmod{2^{3-r}}$ and one of γ or δ is odd. By Theorem 17, $\mu(I - M, 3) = 0$.

It follows that

$$\sum_{M \in I_3} \mu(I - M, 3) = 3754 \cdot \frac{1}{8192} + 365 \cdot \frac{1}{3 \cdot 8192} + \frac{1}{57344} = \frac{5087}{10752}.$$

This concludes the proof of Theorem 1.

REFERENCES

- [1] W. Bosma, J. Cannon, and C. Playoust. The Magma algebra system. I. The user language. *J. Symbolic Comput.*, 24(3-4):235–265, 1997. Computational algebra and number theory (London, 1993).
- [2] Paul Cubre and Jeremy Rouse. Divisibility properties of the Fibonacci entry point. *Proc. Amer. Math. Soc.*, 142(11):3771–3785, 2014.
- [3] Sergey Fomin and Andrei Zelevinsky. The Laurent phenomenon. *Adv. in Appl. Math.*, 28(2):119–144, 2002.
- [4] Helmut Hasse. Über die Dichte der Primzahlen p , für die eine vorgegebene ganzrationale Zahl $a \neq 0$ von gerader bzw. ungerader Ordnung mod. p ist. *Math. Ann.*, 166:19–23, 1966.
- [5] A. N. W. Hone. Elliptic curves and quadratic recurrence sequences. *Bull. London Math. Soc.*, 37(2):161–171, 2005.
- [6] A. N. W. Hone. Sigma function solution of the initial value problem for Somos 5 sequences. *Trans. Amer. Math. Soc.*, 359(10):5019–5034, 2007.
- [7] Henryk Iwaniec and Emmanuel Kowalski. *Analytic number theory*, volume 53 of *American Mathematical Society Colloquium Publications*. American Mathematical Society, Providence, RI, 2004.
- [8] Rafe Jones and Jeremy Rouse. Galois theory of iterated endomorphisms. *Proc. Lond. Math. Soc. (3)*, 100(3):763–794, 2010. Appendix A by Jeffrey D. Achter.

- [9] J. C. Lagarias. The set of primes dividing the Lucas numbers has density $2/3$. *Pacific J. Math.*, 118(2):449–461, 1985.
- [10] J. C. Lagarias. Errata to: “The set of primes dividing the Lucas numbers has density $2/3$ ” [Pacific J. Math. **118** (1985), no. 2, 449–461; MR0789184 (86i:11007)]. *Pacific J. Math.*, 162(2):393–396, 1994.
- [11] Richard Pink. On the order of the reduction of a point on an abelian variety. *Math. Ann.*, 330(2):275–291, 2004.
- [12] Jeremy Rouse and David Zureick-Brown. Elliptic curves over \mathbb{Q} and 2-adic images of Galois. To appear in *Research in Number Theory*.
- [13] Joseph H. Silverman. *The arithmetic of elliptic curves*, volume 106 of *Graduate Texts in Mathematics*. Springer-Verlag, New York, 1992. Corrected reprint of the 1986 original.
- [14] Joseph H. Silverman and John Tate. *Rational points on elliptic curves*. Undergraduate Texts in Mathematics. Springer-Verlag, New York, 1992.
- [15] David E. Speyer. Perfect matchings and the octahedron recurrence. *J. Algebraic Combin.*, 25(3):309–348, 2007.

DEPARTMENT OF MATHEMATICS, WAKE FOREST UNIVERSITY, WINSTON-SALEM, NC 27109

E-mail address: `davibf11@wfu.edu`

E-mail address: `kotsrc11@wfu.edu`

E-mail address: `rouseja@wfu.edu`