

The Cycle Structure of LFSR with Arbitrary Characteristic Polynomial over Finite Fields

Zuling Chang · Martianus Frederic Ezerman ·
San Ling · Huaxiong Wang

Received: date / Accepted: date

Abstract We determine the cycle structure of linear feedback shift register with arbitrary monic characteristic polynomial over any finite field. For each cycle, a method to find a state and a new way to represent the state are proposed.

Keywords cycle structure · cyclotomic number · de Bruijn sequence · decimation · LFSR · periodic sequence

Mathematics Subject Classification (2000) 11B50 · 94A55 · 94A60

1 Introduction

Let \mathbb{F}_q , where $q = p^m$ with $m \in \mathbb{N}$, be the finite field of characteristic p and cardinality q . A linear feedback shift register (LFSR) with characteristic polynomial $f(x) \in \mathbb{F}_q[x]$ satisfying $f(0) \neq 0$ generates some periodic sequences $\mathbf{s} = (s_0, s_1, \dots, s_{N-1}) \in \mathbb{F}_q^N$ with $N \mid \text{ord}(f(x))$. All polynomials in this work are monic with positive degrees.

Let $\Omega(f(x))$ denote the set containing all sequences with $f(x)$ as their characteristic polynomial. The cycle structure of $\Omega(f(x))$ for arbitrary $f(x)$ reflects the algebraic structure and properties of its LFSR and has many potential applications. Determining the cycle structure is a crucial step in the construction of q -ary de Bruijn sequences using the *cycle joining method* discussed in, e.g., [5, 6].

Binary de Bruijn sequences have numerous well-known applications in cryptography, most notably as an ingredient in stream ciphers. Powerful tools developed from de Bruijn graphs whose vertices are length ℓ strings of 4 alphabets accelerated DNA sequencing technologies [3]. Genome assembly triggered in earnest investigations into possible applications

Z. Chang
School of Mathematics and Statistics, Zhengzhou University, Zhengzhou 450001, China
E-mail: zuling_chang@zzu.edu.cn

M. F. Ezerman · S. Ling · H. Wang
Division of Mathematical Sciences, School of Physical and Mathematical Sciences,
Nanyang Technological University, 21 Nanyang Link, Singapore 637371
E-mail: {fredezerman, lingsan, HXWang}@ntu.edu.sg

A preliminary version of this work was presented at SETA 2016.

of q -ary de Bruijn sequences and related structures, inviting us to go beyond the usual binary set-up.

Two areas where q -ary de Bruijn sequences have recently been shown to play important roles are neuroscience and biochemistry. In the former, the sequences are crucial in designing experiments to test the relationship between ordered stimuli and the neural responses that they trigger [1]. In the latter, 4-ary de Bruijn sequences help build compact libraries of RNA probes, useful to measure protein-RNA interactions [2]. These implementations require a *large number of efficiently generated sequences*. Our work lays a good foundation for further investigation in this direction.

Results on the distribution of the periods and the number of sequences having a particular period can be found in [10, Ch. 8, Sect. 5]. The exact cycle structure, *i.e.*, the list of states belonging to the sequences that generate each cycle in $\Omega(f(x))$, is known only for several limited cases. When $f(x)$ is a primitive polynomial, the cycle structure is treated in [10, Ch. 8]. For $f(x) = (1+x)^n$, it is given in [9].

Let $\{g_1(x), g_2(x), \dots, g_k(x)\}$ be a set of k pairwise distinct monic irreducible polynomials in $\mathbb{F}_q[x]$, $n_i := \deg(g_i(x))$, and $b_i \in \mathbb{N}$. Let $f(x) := \prod_{i=1}^k g_i^{b_i}(x)$. This paper settles the open problem of determining the cycle structure of $\Omega(f(x))$. For each cycle, a state belonging to the cycle is identified. We then use this knowledge to build a procedure to generate all sequences in $\Omega(f(x))$ by supplying a state belonging to each of these sequences.

2 Preliminaries

A q -ary sequence $\mathbf{s} = s_0, s_1, s_2, \dots$ has entries $s_i \in \mathbb{F}_q$. For $N \in \mathbb{N}$, if $s_{i+N} = s_i$ for all $i \geq 0$, then \mathbf{s} is N -periodic or with period N and we write $\mathbf{s} = (s_0, s_1, \dots, s_{N-1})$. The period of \mathbf{s} is not unique, so we call the smallest one the *least period* of \mathbf{s} . For brevity and when it is clear from the context, by *period* we mean *least period* in the sequel.

Let $\mathbf{s} = s_0, s_1, s_2, \dots$, $\mathbf{s}' = s'_0, s'_1, s'_2, \dots$, and $c \in \mathbb{F}_q$. Addition and scalar multiplication are given respectively by $\mathbf{s} + \mathbf{s}' = s_0 + s'_0, s_1 + s'_1, s_2 + s'_2, \dots$ and $c\mathbf{s} = cs_0, cs_1, cs_2, \dots$

It is well-known (see, *e.g.*, [7, Ch. 4]) that $\mathbf{s} = (s_0, s_1, \dots, s_{N-1})$ can be generated by a linear feedback shift register (LFSR). The entries satisfy the linear recursive relation

$$s_{k+n} + c_{n-1}s_{k+n-1} + \dots + c_1s_{k+1} + c_0s_k = 0 \text{ for } k \geq 0, \quad (1)$$

where $c_0, c_1, \dots, c_{n-1} \in \mathbb{F}_q$, $n \in \mathbb{N}$, and $c_0 \neq 0$.

A (left) shift operator L acts on $\mathbf{s} = (s_0, s_1, \dots, s_{N-1})$ by $L\mathbf{s} = (s_1, s_2, \dots, s_{N-1}, s_0)$. Thus, $L^i\mathbf{s} = (s_i, s_{i+1}, \dots, s_{i-1})$ for $i \in \mathbb{N}$ with the convention that $L^0\mathbf{s} = \mathbf{s}$. Using L , we write (1) as

$$\mathbf{0} = L^n\mathbf{s} + c_{n-1}L^{n-1}\mathbf{s} + \dots + c_1L\mathbf{s} + c_0L^0\mathbf{s} = (L^n + c_{n-1}L^{n-1} + \dots + c_1L + c_0L^0)\mathbf{s}.$$

The equation defines a *characteristic polynomial*

$$f(x) = x^n + c_{n-1}x^{n-1} + \dots + c_1x + c_0 \quad (2)$$

of \mathbf{s} over \mathbb{F}_q . The sequence \mathbf{s} may have many such polynomials. We call the monic characteristic polynomial with the least degree the *minimal polynomial* and its degree the *linear complexity* of \mathbf{s} . The (least) period of \mathbf{s} is equal to the order of its minimal polynomial.

If the minimal polynomial of \mathbf{s} is primitive with degree n , then \mathbf{s} is the corresponding m -sequence (also known as *maximal length sequence*) with period $q^n - 1$. This period is the maximal period of all sequences generated by any LFSR with minimal polynomial of degree n . We use \mathbf{m} to denote an m -sequence.

A sequence \mathbf{v} is said to be a d -decimation sequence of \mathbf{s} , denoted by $\mathbf{v} = \mathbf{s}^{(d)}$, if $v_j = s_{d \cdot j}$ for all $j \geq 0$. A d -decimation $\mathbf{m}^{(d)}$ of an m -sequence \mathbf{m} is also an m -sequence if and only if $\gcd(d, q^n - 1) = 1$. Hence, the number of distinct m -sequences of period $q^n - 1$, up to cyclic shifts, is $\frac{\phi(q^n - 1)}{n}$ where $\phi(\cdot)$ is the Euler totient function. There is a bijection between the set of all such sequences and the set of primitive polynomials of degree n in $\mathbb{F}_q[x]$. More properties of sequences in relation to their characteristic and minimal polynomials can be found in [7, Ch. 4] and [10, Ch. 8].

For $\mathbf{s} = (s_0, s_1, \dots, s_{N-1})$ with characteristic polynomial $f(x)$ defined in (2), the vector $\mathbf{s}_i = (s_i, s_{i+1}, \dots, s_{i+n-1}) \in \mathbb{F}_q^n$ is the i -th n -stage state of \mathbf{s} . In particular, we call \mathbf{s}_0 the *initial state*. Note that the states of the sequence depends on the specified characteristic polynomial.

A *state operator* T turns the state \mathbf{s}_i into \mathbf{s}_{i+1} , i.e., $\mathbf{s}_{i+1} = T\mathbf{s}_i$. If \mathbf{s}_i is a state of \mathbf{s} and e is the period of \mathbf{s} , then the e distinct states of \mathbf{s} are $\mathbf{s}_i, T\mathbf{s}_i = \mathbf{s}_{i+1}, \dots, T^{e-1}\mathbf{s}_i = \mathbf{s}_{i+e-1}$.

To $f(x)$ in (2), one associates a matrix

$$A := \begin{pmatrix} 0 & 0 & \cdots & 0 & -c_0 \\ 1 & 0 & \cdots & 0 & -c_1 \\ 0 & 1 & \cdots & 0 & -c_2 \\ \vdots & \vdots & \ddots & \vdots & \vdots \\ 0 & 0 & \cdots & 1 & -c_{n-1} \end{pmatrix}. \quad (3)$$

Using \mathbf{s}_0 as the input state, the state vectors of the resulting sequence are $\mathbf{s}_j = \mathbf{s}_0 A^j$ for $j \in \{0, 1, 2, \dots\}$. We know from [10, Thm. 8.13] that e divides the order of A in $GL(n, \mathbb{F}_q)$. Furthermore, $\mathbf{s}_{i+1} = \mathbf{s}_i A = T\mathbf{s}_i$.

For an N -periodic sequence $\mathbf{u} \in \Omega(f(x))$, define the set $[\mathbf{u}] := \{\mathbf{u}, L\mathbf{u}, L^2\mathbf{u}, \dots, L^{N-1}\mathbf{u}\}$ to be a *shift equivalent class* or a *cycle* in $\Omega(f(x))$. Sequences in a cycle are said to be *shift equivalent*. Thus, $\Omega(f)$ can be partitioned into cycles.

If $\Omega(f(x))$ consists of exactly j cycles $[\mathbf{u}_1], [\mathbf{u}_2], \dots, [\mathbf{u}_j]$ for some $j \in \mathbb{N}$, then its *cycle structure* is presented as $\Omega(f(x)) = [\mathbf{u}_1] \cup [\mathbf{u}_2] \cup \dots \cup [\mathbf{u}_j]$. To study the cycle structure, determining a state belonging to each sequence \mathbf{u}_i is crucial. Starting with the state, one easily generates the whole sequence and, hence, the corresponding cycle in $\Omega(f(x))$.

Given our setup, from [7, Sect. 4.4] we can derive the following properties.

1. $\Omega(f(x))$ can be decomposed into a direct sum of $\Omega(g_i^{b_i}(x))$ with $1 \leq i \leq k$, i.e.,

$$\Omega(f(x)) = \Omega(g_1^{b_1}(x)) \oplus \Omega(g_2^{b_2}(x)) \oplus \cdots \oplus \Omega(g_k^{b_k}(x)).$$

2. Every sequence $\mathbf{s} \in \Omega(f(x))$ can be written as

$$\mathbf{s} = \mathbf{s}_1 + \mathbf{s}_2 + \cdots + \mathbf{s}_k \text{ with } \mathbf{s}_i \in \Omega(g_i^{b_i}(x)) \text{ for } 1 \leq i \leq k.$$

3. The period of \mathbf{s} is $\text{lcm}(e_1, e_2, \dots, e_k)$ where e_i is the period of \mathbf{s}_i .

Remark 1 The polynomial $f(x)$ in [7, Thm. 4.10] is a product of distinct irreducible polynomials, i.e., $b_i = 1$ for all i . The same inductive proof works in our setup here.

3 The Cycle Structure of $\Omega(g(x))$ when $g(x)$ is Irreducible

This section discusses two ways to determine a state belonging to a sequence with irreducible minimal polynomial

$$g(x) = x^n + g_{n-1}x^{n-1} + \cdots + g_1x + g_0 \in \mathbb{F}_q[x]. \quad (4)$$

In the first subsection the state is found by using cyclotomic classes while the second subsection makes use of decimation sequences.

3.1 Using Cyclotomic Classes

The relation between sequences and cyclotomic classes can be used to find a state for each sequence with minimal polynomial $g(x)$. Let $\beta \in \mathbb{F}_{q^n}$ be a root of $g(x)$ with order e and let $t = \frac{q^n-1}{e}$. There must exist a primitive polynomial

$$q(x) = x^n + q_{n-1}x^{n-1} + \cdots + q_0 \in \mathbb{F}_q[x]$$

with a root α satisfying $\beta = \alpha^t$. While we need only one $q(x)$ to be associated with a given $g(x)$, in general the choice is not unique.

Lemma 1 *Let $g(x)$ be a non-primitive irreducible polynomial of degree n with a root β and order e . There are $\frac{\phi(q^n-1)}{\phi(e)}$ primitive polynomials that can be associated with $g(x)$. Such a polynomial $q(x)$ has degree n with a root α that satisfies $\beta = \alpha^t$.*

Proof Each of the roots of $q(x)$, say χ , satisfies $g(\chi^t) = 0$. It is then clear that a chosen $q(x)$ can be an associated primitive polynomial of only one irreducible polynomial of degree n and order e . Two distinct non-primitive irreducible polynomials, both with degree n and order e , have the same number of associated primitive polynomials. There are $\phi(q^n-1)/n$ primitive polynomials with degree n and there are $\phi(e)/n$ non-primitive irreducible polynomials with degree n and order e . Taking their ratio completes the proof. \square

Following [11], the *cyclotomic classes* $C_i \subseteq \mathbb{F}_{q^n}$, for $0 \leq i < t$, are defined by

$$C_i = \{\alpha^{i+k \cdot t} \mid 0 \leq k < e\} = \{\alpha^i \beta^k \mid 0 \leq k < e\} = \alpha^i C_0. \quad (5)$$

The classes partition \mathbb{F}_{q^n} into disjoint union

$$\mathbb{F}_{q^n} = \{0\} \cup C_0 \cup \cdots \cup C_{t-1}.$$

Using $\{1, \beta, \dots, \beta^{n-1}\}$ as a basis for \mathbb{F}_{q^n} as an \mathbb{F}_q -vector space, for $0 \leq j < q^n - 1$, one can uniquely express α^j as $\alpha^j = \sum_{i=0}^{n-1} a_{j,i} \beta^i$ with $a_{j,i} \in \mathbb{F}_q$. Hence,

$$\alpha^{j+t} = \alpha^j \beta = \beta \sum_{i=0}^{n-1} a_{j,i} \beta^i = \sum_{i=0}^{n-1} a_{j,i} \beta^{i+1} = \sum_{i=0}^{n-2} a_{j,i} \beta^{i+1} - \sum_{i=0}^{n-1} a_{j,n-1} g_i \beta^i. \quad (6)$$

Define the mapping $\varphi : \mathbb{F}_{q^n} \rightarrow \mathbb{F}_q^n$ by $\varphi(0) = \mathbf{0}$ and $\varphi(\alpha^j) = (a_{j,0}, a_{j+t,0}, \dots, a_{j+(n-1)t,0})$ with the subscripts reduced modulo $(q^n - 1)$. By the recursive relation in (6), φ is a one-to-one mapping. Let

$$\mathbf{u}_i := (a_{i,0}, a_{i+t,0}, \dots, a_{i+(e-1)t,0}). \quad (7)$$

The discussion in [8, Thm. 3] treats the case of $q = 2$. It generalizes naturally to any q . Following the same line of arguments, we deduce that, under φ , the cyclotomic class C_i corresponds to the cycle $[\mathbf{u}_i]$. In other words, \mathbf{u}_i and the sequence of states of \mathbf{u}_i , namely $((\mathbf{u}_i)_0, (\mathbf{u}_i)_1, \dots, (\mathbf{u}_i)_{e-1})$, where $(\mathbf{u}_i)_j = (a_{i+jt,0}, a_{i+(j+1)t,0}, \dots, a_{i+(j+n-1)t,0}) = \varphi(\alpha^i \beta^j)$ for $0 \leq j < e$, are shift equivalent. Hence, $\mathbf{u}_i \longleftrightarrow C_i$. Thus,

$$\Omega(g(x)) = [\mathbf{0}] \cup [\mathbf{u}_0] \cup [\mathbf{u}_1] \cup \dots \cup [\mathbf{u}_{t-1}]. \quad (8)$$

Since $\{1, \alpha, \dots, \alpha^{n-1}\}$ and $\{1, \beta, \dots, \beta^{n-1}\}$ are bases for \mathbb{F}_{q^n} as an \mathbb{F}_q -vector space, there is an invertible matrix M such that $(1, \beta, \dots, \beta^{n-1}) = (1, \alpha, \dots, \alpha^{n-1})M$.

Build M by using the representations of $1, \beta, \dots, \beta^{n-1}$ in the basis $\{1, \alpha, \dots, \alpha^{n-1}\}$ and the fact that $q(\alpha) = 0$. Conversely, with $\{1, \beta, \dots, \beta^{n-1}\}$ as the basis, $(1, \alpha, \dots, \alpha^{n-1}) = (1, \beta, \dots, \beta^{n-1})M^{-1}$, from which we derive the representation of α^i for $0 \leq i < q^n - 1$.

For $0 \leq i < t$, using the representation of α^i in the basis $\{1, \beta, \dots, \beta^{n-1}\}$ and by (6), we derive $\varphi(\alpha^i)$, which is a state of sequence \mathbf{u}_i .

3.2 Via Decimation

An alternative method to find a state for each nonzero cycle in $\Omega(g(x))$ involves decimating m -sequences. The *trace function* from \mathbb{F}_{q^n} to \mathbb{F}_q is given by $\text{Tr}(x) = x + x^q + \dots + x^{q^{n-1}}$. Recall, e.g., from [7, Sect. 4.6] that $\mathbf{m} = (m_0, m_1, \dots, m_{q^n-2})$ whose characteristic polynomial $q(x)$ is primitive of degree n with a root α can be described using

$$m_i = \text{Tr}(\gamma \alpha^i) : 0 \neq \gamma \in \mathbb{F}_{q^n} \text{ for } i = 0, 1, 2, \dots, q^n - 2.$$

Without loss of generality, one can let $\gamma = 1$, i.e., $m_i = \text{Tr}(\alpha^i)$.

From \mathbf{m} , construct the t distinct t -decimation sequences, each of period e :

$$\mathbf{u}_0 = \mathbf{m}^{(t)}, \mathbf{u}_1 = (L\mathbf{m})^{(t)}, \dots, \mathbf{u}_{t-1} = (L^{t-1}\mathbf{m})^{(t)}.$$

Observe that the entries in the resulting sequences satisfy

$$(\mathbf{u}_k)_j = \text{Tr}(\alpha^{k+t \cdot j}) \text{ for } 0 \leq k < t \text{ and } 0 \leq j < e.$$

Since $\beta = \alpha^t$, each cycle $[\mathbf{u}_i]$ is a cycle in $\Omega(g(x))$.

Starting from an arbitrary $n \cdot t$ consecutive elements of \mathbf{m} , one can derive t distinct n -stage states by the above t -decimating process. It is then straightforward to verify that each of the derived states corresponds to one nonzero cycle. Since the ordering of the cycles in $\Omega(g(x))$ has no significant bearing in our investigation, we can label these states arbitrarily.

Given an irreducible polynomial $g(x)$, finding an associated primitive polynomial $q(x)$ can become computationally expensive for large values of n and t . Decimating m -sequences is a useful tool to accomplish the task while keeping the time complexity low.

Let $\lambda := \frac{\phi(q^n-1)}{n}$. Given $g(x)$, for $1 \leq j \leq \lambda$ let $p_j(x)$ be a primitive polynomial of degree n and let \mathbf{m}_j be its corresponding m -sequence. The set \mathcal{M}_n of all shift inequivalent m -sequences with period $q^n - 1$ is given by

$$\mathcal{M}_n = \{\mathbf{m}_1, \mathbf{m}_2, \dots, \mathbf{m}_\lambda\}.$$

One can also build \mathcal{M} from an arbitrary m -sequence \mathbf{m} by simply collecting all the d_i -decimations $\{\mathbf{m}^{(d_i)}\}$ with d_i satisfying $\gcd(d_i, q^n - 1) = 1$ and, for $i \neq j$, d_i and d_j do not belong to the same conjugate coset.

We now perform a systematic check to determine which among the $p_j(x)$ s can be used as the associated primitive polynomials of $g(x)$. Derive $\mathbf{m}_j^{(t)}$ with period e and check if it shares a common string of $2n$ consecutive elements with a sequence generated by an LFSR with characteristic polynomial $g(x)$. If yes, then $p_j(x)$ can be associated to $g(x)$. The states that we want can then be determined using the method already described above. Performing the routine for all \mathbf{m}_j s guarantees a match between $g(x)$ and some $p_j(x)$. The required steps can be performed without using costly operations over \mathbb{F}_q . As q or n grows, our saving becomes more prominent since operations on polynomials over large finite fields are prohibitively expensive.

To end this section, the following example follows both approaches.

Example 1 Let $g(x) = x^3 + 2x + 2 \in \mathbb{F}_3[x]$. It is irreducible with a root β of order 13. Hence, $(e, t) = (13, 2)$ and $\Omega(g(x)) = [\mathbf{0}] \cup [\mathbf{u}_0] \cup [\mathbf{u}_1]$. Note that $q(x) = x^3 + x^2 + 2x + 1 \in \mathbb{F}_3[x]$ is a primitive polynomial whose root α satisfies $\beta = \alpha^2$. We can then express

$$(1, \alpha, \alpha^2) = (1, \beta, \beta^2) \begin{pmatrix} 1 & 2 & 0 \\ 0 & 1 & 1 \\ 0 & 1 & 0 \end{pmatrix}.$$

Using $\beta^3 = 1 + \beta$ and writing in the form of (6), $\alpha = 2 + \beta + \beta^2$, $\alpha^3 = 1 + \beta^2$, and $\alpha^5 = 1 + 2\beta$. This implies that $\mathbf{a}_1 = \varphi(\alpha) = (2, 1, 1)$ is a state of \mathbf{u}_1 and $\mathbf{a}_0 = (1, 0, 0)$ is a state of \mathbf{u}_0 . Thus, $\mathbf{u}_0 = (1, 0, 0, 1, 0, 1, 1, 1, 2, 2, 0, 1, 2)$ and $\mathbf{u}_1 = (2, 1, 1, 0, 2, 1, 2, 0, 0, 2, 0, 2, 2)$.

We can also get the respective states of \mathbf{u}_0 and \mathbf{u}_1 via decimation. The m -sequence with characteristic polynomial $q(x)$ is

$$\mathbf{m} = (1, 1, 1, 2, 1, 0, 2, 0, 2, 2, 0, 0, 1, 2, 2, 2, 1, 2, 0, 1, 0, 1, 1, 0, 0, 2).$$

The first 6 consecutive elements of \mathbf{m} can be taken to form $\mathbf{v} = (1, 1, 1, 2, 1, 0)$. Hence, $\mathbf{v}^{(2)} = (1, 1, 1)$ and $(L\mathbf{v})^{(2)} = (1, 2, 0)$ are the respective 3-stage states of \mathbf{u}_0 and \mathbf{u}_1 .

In $\mathbb{F}_3[x]$, there are 4 primitive polynomials of degree 3: $p_1(x) = x^3 + 2x + 1$, $p_2(x) = x^3 + 2x^2 + x + 1$, $p_3(x) = x^3 + x^2 + 2x + 1$, and $p_4(x) = x^3 + 2x^2 + 1$. There are also 4 non-primitive irreducible polynomials of degree 3, each having $e = 13$ and $t = 2$: $g_1(x) = x^3 + 2x^2 + 2x + 2$, $g_2(x) = x^3 + 2x + 2$, $g_3(x) = x^3 + x^2 + 2$, and $g_4(x) = x^3 + x^2 + x + 2$. By Lemma 1, each $g_i(x)$ is associated with exactly one $p_j(x)$ for $1 \leq i, j \leq 4$. We now use decimation to find the association and, subsequently, determine the respective states of the nonzero cycles in $\Omega(g_i(x))$ for all i .

The m -sequences are

$$\begin{aligned} \mathbf{m}_1 &= (1, 1, 1, 0, 0, 2, 0, 2, 1, 2, 2, 1, 0, 2, 2, 2, 0, 0, 1, 0, 1, 2, 1, 1, 2, 0), \\ \mathbf{m}_2 &= (1, 1, 1, 2, 0, 0, 1, 1, 0, 1, 0, 2, 1, 2, 2, 2, 1, 0, 0, 2, 2, 0, 2, 0, 1, 2), \\ \mathbf{m}_3 &= (1, 1, 1, 2, 1, 0, 2, 0, 2, 2, 0, 0, 1, 2, 2, 2, 1, 2, 0, 1, 0, 1, 1, 0, 0, 2), \text{ and} \\ \mathbf{m}_4 &= (1, 1, 1, 0, 2, 1, 1, 2, 1, 0, 1, 0, 0, 2, 2, 2, 0, 1, 2, 2, 1, 2, 0, 2, 0, 0). \end{aligned}$$

Their respective 2-decimation sequences are

$$\begin{aligned} \mathbf{m}_1^{(2)} &= (1, 1, 0, 0, 1, 2, 0, 2, 0, 1, 1, 1, 2), \quad \mathbf{m}_2^{(2)} = (1, 1, 0, 1, 0, 0, 1, 2, 1, 0, 2, 2, 1), \\ \mathbf{m}_3^{(2)} &= (1, 1, 1, 2, 2, 0, 1, 2, 1, 0, 0, 1, 0), \quad \mathbf{m}_4^{(2)} = (1, 1, 2, 1, 1, 1, 0, 2, 0, 2, 1, 0, 0). \end{aligned}$$

We feed the first three entries of $\mathbf{m}_j^{(2)}$ to each LFSR whose characteristic polynomial is $g_i(x)$ to establish the association and the results given in Table 1.

□

Table 1 States and Nonzero Cycles in $\Omega(g_i(x))$

$g(x)$	$q(x)$	States	Nonzero Cycles
$g_1(x)$	$p_4(x)$	(1, 1, 2), (1, 0, 1)	[(1, 1, 2, 1, 1, 1, 0, 2, 0, 2, 1, 0, 0)], [(1, 0, 1, 2, 0, 0, 2, 2, 1, 2, 2, 2, 0)]
$g_2(x)$	$p_3(x)$	(1, 1, 1), (1, 2, 0)	[(1, 1, 1, 2, 2, 0, 1, 2, 1, 0, 0, 1, 0)], [(1, 2, 0, 0, 2, 0, 2, 2, 2, 1, 1, 0, 2)]
$g_3(x)$	$p_2(x)$	(1, 1, 0), (1, 2, 0)	[(1, 1, 0, 1, 0, 0, 1, 2, 1, 0, 2, 2, 1)], [(1, 2, 0, 1, 1, 2, 2, 2, 0, 2, 0, 0, 2)]
$g_4(x)$	$p_1(x)$	(1, 1, 0), (1, 0, 2)	[(1, 1, 0, 0, 1, 2, 0, 2, 0, 1, 1, 1, 2)], [(1, 0, 2, 2, 2, 1, 2, 2, 0, 0, 2, 1, 0)]

4 The Cycle Structure of $\Omega(g^b(x))$ with $b > 1$

Our next task is to find one state of each cycle in $\Omega(g^b(x))$ with $g(x)$, β , e , and t as defined in Sect. 3 and $1 < b \in \mathbb{N}$. Since (8) already gives the cycle structure of $\Omega(g(x))$, we consider the states of the cycles in $\Omega(g^b(x)) \setminus \Omega(g(x))$. The following result follows directly from [10, Thm. 3.8 and Thm. 8.63].

Lemma 2 For $1 \leq r \leq b$, sequences in $\Omega(g^b(x))$ with minimal polynomial $g^r(x)$ can be divided into

$$\frac{q^{nr} - q^{n(r-1)}}{e \cdot p^c} = t \cdot \frac{q^{n(r-1)}}{p^c}$$

distinct cycles, each with period $e \cdot p^c$ where c is the least integer satisfying $p^c \geq r$.

We adapt the notion of *D-morphism*, first introduced by Lempel in [9], to generate cycles with minimal polynomial $g^{r+1}(x)$ from cycles with minimal polynomial $g^r(x)$. A state of a cycle with minimal polynomial $g^{r+1}(x)$ can then be derived from a state belonging to a cycle with minimal polynomial $g^r(x)$. We apply the procedure recursively to cover all cycles. The mapping $D: \mathbb{F}_q^{(r+1)n} \rightarrow \mathbb{F}_q^{rn}$ sends $\mathbf{a} = (a_0, a_1, \dots, a_{(r+1)n-1}) \mapsto \mathbf{b} = (b_0, b_1, \dots, b_{rn-1})$

$$\Leftrightarrow \begin{cases} b_0 &= g_0 a_0 + g_1 a_1 + \dots + g_{n-1} a_{n-1} + a_n, \\ b_1 &= g_0 a_1 + g_1 a_2 + \dots + g_{n-1} a_n + a_{n+1}, \\ \vdots & \vdots \\ b_{rn-1} &= g_0 a_{rn-1} + g_1 a_{rn} + \dots + g_{n-1} a_{(r+1)n-2} + a_{(r+1)n-1}. \end{cases} \quad (9)$$

Note that D is onto. Given $(a_0, a_1, \dots, a_{n-1}) \in \mathbb{F}_q^n$ and $\mathbf{b} \in \mathbb{F}_q^{rn}$, one can uniquely determine $a_n, a_{n+1}, \dots, a_{(r+1)n-1}$. Hence, for any \mathbf{b} , there are q^n distinct \mathbf{a} 's satisfying $\mathbf{b} = D(\mathbf{a})$.

Let $\mathbf{a} = (a_0, a_1, \dots, a_{(r+1)n-1}) \in \mathbb{F}_q^{(r+1)n}$ be an $n(r+1)$ -stage state of a sequence with characteristic polynomial $g(x)$. Then $g_0 a_i + g_1 a_{i+1} + \dots + a_{i+n} = 0$ implies $D(\mathbf{a}) = \mathbf{0}$ for $0 \leq i \leq nr - 1$.

Lemma 3 The mapping D is a surjective homomorphism whose kernel, denoted by $\ker(D)$, is the set $\{\mathbf{k}_0, \mathbf{k}_1, \dots, \mathbf{k}_{q^n-1}\}$ of q^n distinct $n(r+1)$ -stage states of the sequences in $\Omega(g(x))$.

Since $L(D(\mathbf{a})) = D(L(\mathbf{a}))$ for any $\mathbf{a} \in \mathbb{F}_q^{(r+1)n}$, D preserves shift relation. Whenever $\mathbf{a}_0, \mathbf{a}_1, \dots$ are consecutive states of a sequence, we know that $D(\mathbf{a}_0), D(\mathbf{a}_1), \dots$ are also consecutive states of some sequence. Hence, for a sequence $\mathbf{s} = (s_0, s_1, \dots, s_{N-1})$ with period N , sequence $\mathbf{u} := D(\mathbf{s})$ has period $N' \mid N$ and entries

$$u_i = g_0 s_i + g_1 s_{i+1} + \dots + g_{n-1} s_{i+n-1} + s_{i+n}. \quad (10)$$

Lemma 4 If $\mathbf{s} \in \Omega(g^{r+1}(x))$, then $\mathbf{u} := D(\mathbf{s}) \in \Omega(g^r(x))$. If $g^c(x)$ is the minimal polynomial of \mathbf{s} , for some $1 \leq c \leq r+1$, then $g^{c-1}(x)$ is the minimal polynomial of \mathbf{u} .

Proof For a given sequence \mathbf{s} in $\Omega(g^{r+1}(x))$, sequence \mathbf{u} in (10) can be written as

$$\mathbf{u} = g_0\mathbf{s} + g_1L\mathbf{s} + \dots + g_{n-1}L^{n-1}\mathbf{s} + L^n\mathbf{s} = g(L)\mathbf{s}.$$

Since $g^{r+1}(L)\mathbf{s} = \mathbf{0}$, we have $g^r(L)\mathbf{u} = g^r(L)g(L)\mathbf{s} = g^{r+1}(L)\mathbf{s} = \mathbf{0}$. Thus, $\mathbf{u} \in \Omega(g^r(x))$.

The second assertion can be similarly argued using the fact that $g(x)$ is irreducible. \square

Corollary 1 *Let \mathbf{s} be a sequence with minimal polynomial $h(x)$. Then the minimal polynomial of $\mathbf{u} = D(\mathbf{s})$ is*

$$\begin{cases} \frac{h(x)}{g(x)} & \text{if } g(x) \text{ divides } h(x) \text{ and} \\ h(x) & \text{if } g(x) \text{ does not divide } h(x). \end{cases}$$

Let $\mathbf{b} \in \mathbb{F}_q^{nr}$ be a state belonging to sequence \mathbf{u} with minimal polynomial $g^r(x)$. By (9), the q^n distinct vectors \mathbf{a} satisfying $\mathbf{b} = D(\mathbf{a})$ are the respective states of some sequences with minimal polynomial $g^{r+1}(x)$. Given one such \mathbf{a} , these states can be written as the sum of \mathbf{a} and the corresponding states in $\ker(D)$, i.e., as $\mathbf{a} + \mathbf{k}_i$ with $0 \leq i < q^n$.

Let \mathbf{s} be a sequence with minimal polynomial $g^{r+1}(x)$ containing \mathbf{a} as a state. Now, consider all q^n sequences: \mathbf{s} plus each of the q^n distinct sequences in $\Omega(g(x))$. We next decide whether these sequences correspond to distinct cycles with minimal polynomial $g^{r+1}(x)$. The consideration is based on whether r is a power of p .

Lemma 5 *Suppose r is not a power of p . Let \mathbf{s} be a sequence with minimal polynomial $g^{r+1}(x)$. Let \mathbf{s}_1 and \mathbf{s}_2 be two distinct sequences in $\Omega(g(x))$. Then $\mathbf{s} + \mathbf{s}_1$ and $\mathbf{s} + \mathbf{s}_2$ are shift inequivalent.*

Proof Lemma 2 says that the period of \mathbf{s} is $e \cdot p^c$ where c is the least integer with $p^c \geq r + 1$. Assume that $\mathbf{s} + \mathbf{s}_1$ and $\mathbf{s} + \mathbf{s}_2$ are shift equivalent. Then there exists an integer $0 \leq \ell < e \cdot p^c$ such that $\mathbf{s} + \mathbf{s}_1 = L^\ell(\mathbf{s} + \mathbf{s}_2) = L^\ell\mathbf{s} + L^\ell\mathbf{s}_2$. Hence, $(L^\ell - I)\mathbf{s} = \mathbf{s}_1 - L^\ell\mathbf{s}_2 \in \Omega(g(x))$. Thus, $g(x)$ must be a characteristic polynomial of $(L^\ell - I)\mathbf{s}$, implying that $g^r(x) \mid x^\ell - 1$ with $e \cdot p^c = \text{ord}(g^{r+1}(x))$. This forces $\ell \geq e \cdot p^c$, which is a contradiction. \square

With r not a power of p , let a state \mathbf{b} of a sequence \mathbf{u} with minimal polynomial $g^r(x)$ be given. Then, by (9), the q^n distinct vectors \mathbf{a} satisfying $\mathbf{b} = D(\mathbf{a})$ are states of pairwise distinct cycles, all of which has $g^{r+1}(x)$ as their minimal polynomial.

Given respective states \mathbf{b} and \mathbf{b}' from two distinct cycles, each with minimal polynomial $g^r(x)$, their D pre-image states \mathbf{a} and \mathbf{a}' belong to two distinct cycles with minimal polynomial $g^{r+1}(x)$. Thus, given all $\delta := t \cdot q^{n(r-1)} \cdot p^{-c}$ states of distinct cycles with minimal polynomial $g^r(x)$, the exact $q^n \cdot \delta$ states of all distinct cycles having minimal polynomial $g^{r+1}(x)$ are known.

In fact, given \mathbf{b} , we can let $a_0 = \dots = a_{n-1} = 0$ and construct a state \mathbf{a} by using (9). Taking the sum of \mathbf{a} and each of the $n(r+1)$ -stage states of sequences in $\Omega(g(x))$ gives us what we are after.

Lemma 6 *Let $r = p^d$ for some nonnegative integer d . For a sequence \mathbf{s} with minimal polynomial $g^{r+1}(x)$, there exists a sequence \mathbf{s}' in $\Omega(g(x))$ such that the set $\{\mathbf{s} + \eta\mathbf{s}' : \eta \in \mathbb{F}_p\}$ contains p shift equivalent sequences.*

Proof Lemma 2 says that the period of \mathbf{s} is $e \cdot p^{d+1} = p \cdot e \cdot r$.

Observe that $L^{e \cdot r}\mathbf{s} - \mathbf{s} = (L^{e \cdot r} - I)\mathbf{s} = (L^e - I)^r\mathbf{s}$. Since $g(x)$ is irreducible, $g^r(x)$ divides $(x^e - 1)^r$ but $g^{r+1}(x)$ does not divide $(x^e - 1)^r$. Hence, $g(x)$ is the minimal polynomial of $\mathbf{s}' := (L^e - I)^r\mathbf{s} \neq \mathbf{0}$ and $L^{e \cdot r}\mathbf{s} = \mathbf{s} + \mathbf{s}'$.

Let $2 \leq \eta \leq p-1$. Since \mathbf{s}' has period e ,

$$L^{\eta \cdot e \cdot r} \mathbf{s} - L^{(\eta-1) \cdot e \cdot r} \mathbf{s} = L^{(\eta-1) \cdot e \cdot r} [(L^{e \cdot r} - I) \mathbf{s}] = L^{(\eta-1) \cdot e \cdot r} \mathbf{s}' = \mathbf{s}'.$$

Proceeding inductively, one obtains $L^{\eta \cdot e \cdot r} \mathbf{s} = L^{(\eta-1) \cdot e \cdot r} \mathbf{s} + \mathbf{s}' = \mathbf{s} + \eta \mathbf{s}'$. \square

When $r = p^d$, given a state \mathbf{b} of a sequence \mathbf{u} with minimal polynomial $g^r(x)$, we combine Lemma 6 and (9) to determine the q^n distinct vectors $\mathbf{a} \in \mathbb{F}_q^{n(r+1)}$ satisfying $\mathbf{b} = D(\mathbf{a})$. For an arbitrary \mathbf{a} , let $\mathbf{a}' := T^{e \cdot r} \mathbf{a} - \mathbf{a}$. Then the p distinct states $\mathbf{a} + \eta \mathbf{a}'$ with $\eta \in \mathbb{F}_p$ correspond to one cycle with minimal polynomial $g^{r+1}(x)$.

For two arbitrary such states, say \mathbf{a}_1 and \mathbf{a}_2 , it follows that $\mathbf{a}' = T^{e \cdot r} \mathbf{a}_1 - \mathbf{a}_1 = T^{e \cdot r} \mathbf{a}_2 - \mathbf{a}_2$. Thus, given \mathbf{b} and \mathbf{a}' , we can construct the states of all the $\frac{q^n}{p}$ distinct cycles with minimal polynomial $g^{r+1}(x)$. Since there are $t \cdot \frac{q^{n(r-1)}}{r}$ possible choices for \mathbf{b} , we can provide the exact $t \cdot q^{n \cdot r} \cdot p^{-(d+1)}$ states of all the distinct cycles with minimal polynomial $g^{r+1}(x)$.

Remark 2 It may be possible to derive an exact formula for \mathbf{a}' given $g(x)$, the value $r = p^d$, and a vector \mathbf{b} . The formula is likely to be quite complicated and we leave it as an open problem. Using the LFSR to compute $T^{e \cdot r} \mathbf{a} - \mathbf{a}$ directly from an arbitrary \mathbf{a} seems to be the simplest approach.

Example 2 ([9]) Let $g(x) = x+1 \in \mathbb{F}_2[x]$. Given a state $\mathbf{b} = (b_0, b_1, \dots, b_{r-1}) \in \mathbb{F}_2^r$ of a sequence with minimal polynomial $g^r(x) = (x+1)^r$ and an element $a_0 \in \mathbb{F}_2$, the entries of the resulting $\mathbf{a} = (a_0, a_1, \dots, a_r) \in \mathbb{F}_2^{r+1}$ are $a_i = a_0 + b_0 + \dots + b_{i-1}$ for $i \in \{1, 2, \dots, r\}$. When $a_0 = 0$, we denote the resulting state by \mathbf{a}_0 . The state is \mathbf{a}_1 when $a_0 = 1$. Since the only nonzero sequence with minimal polynomial $x+1$ is the all one sequence $\mathbf{1}$, we know that $\mathbf{a}_1 = \mathbf{a}_0 + \mathbf{1}$.

If r is not a power of 2, then \mathbf{a}_0 and \mathbf{a}_1 are the respective states of two distinct sequences sharing the same minimal polynomial $(x+1)^{r+1}$. If r is a power of 2, then \mathbf{a}_0 and \mathbf{a}_1 belong to the same sequence with minimal polynomial $(x+1)^{r+1}$. \square

Example 3 Let $g(x) = x^3 + 2x + 2 \in \mathbb{F}_3[x]$, and note that $r = 1 = 3^0$. From Example 1, the respective states of the two nonzero cycles are $(1, 0, 0)$ and $(2, 1, 1)$.

Using $\mathbf{b} = (1, 0, 0)$ and letting $a_0 = a_1 = a_2 = 0$, we get a 6-stage state $\mathbf{a} = (0, 0, 0, 1, 0, 1)$ from (9) and $\mathbf{a}' = T^{13} \mathbf{a} - \mathbf{a} = (1, 2, 0, 0, 2, 0)$. The corresponding sequence \mathbf{s} of period 39 with minimal polynomial $g^2(x) = x^6 + x^4 + x^3 + x^2 + 2x + 1$ is

$$\underbrace{(0, 0, 0, 1, 0, 1)}_{\mathbf{a}}, 2, 1, 1, 1, 0, 1, 0, \underbrace{1, 2, 0, 1, 2, 1}_{\mathbf{a} + \mathbf{a}'}, 1, 1, 0, 0, 2, 1, 1, 2, \underbrace{2, 1, 0, 1, 1, 1}_{\mathbf{a} + 2\mathbf{a}'}, 0, 2, 2, 0, 2, 1, 1). \quad (11)$$

It is easy to check that $L^{13} \mathbf{s} - \mathbf{s} = L^{26} \mathbf{s} - L^{13} \mathbf{s} = (1, 2, 0, 0, 2, 0, 2, 2, 2, 1, 1, 0, 2)$. As indicated by the underbraces in (11), \mathbf{a} , $\mathbf{a} + \mathbf{a}'$ and $\mathbf{a} + 2\mathbf{a}'$ generate the same cycle $[\mathbf{s}]$.

The 27 distinct 6-stage states in $\ker(D)$ can be partitioned into 9 disjoint sets, each of the form $\mathbf{a}_i + \{\mathbf{0}, \mathbf{a}', 2\mathbf{a}'\}$ for $0 \leq i < 9$. The respective states of the cycles with minimal polynomial $g^2(x)$ are $\mathbf{a} + \mathbf{a}_i$. After some computation we arrive at

$$\{\mathbf{a}_i : 0 \leq i < 9\} = \{(\mathbf{0}), (1, 0, 0, 1, 0, 1), (0, 0, 1, 0, 1, 1), (1, 0, 1, 1, 1, 2), (1, 1, 1, 2, 2, 0), (1, 1, 2, 2, 0, 1), (1, 2, 2, 0, 1, 2), (0, 1, 2, 1, 0, 0), (1, 1, 0, 2, 1, 2)\}.$$

Obtaining $\{\mathbf{a} + \mathbf{a}_i : 0 \leq i < 9\}$ means we are halfway done.

To derive the remaining half we use $\mathbf{b} = (2, 1, 1)$ and let $a_0 = a_1 = a_2 = 0$. This yields a 6-stage state $\mathbf{a} = (0, 0, 0, 2, 1, 0)$ with $\mathbf{a}' = (1, 1, 2, 2, 0, 1)$. The corresponding sequence \mathbf{s} is

$$(0, 0, 0, 2, 1, 0, 0, 0, 1, 2, 1, 0, 2, 1, 1, 2, 1, 1, 1, 2, 1, 1, 2, 2, 0, 0, 2, 2, 1, 0, 1, 2, 1, 2, 1, 2, 0, 0, 1).$$

$\underbrace{\hspace{10em}}_{\mathbf{a}} \quad \underbrace{\hspace{10em}}_{\mathbf{a}+\mathbf{a}'} \quad \underbrace{\hspace{10em}}_{\mathbf{a}+2\mathbf{a}'}$

One eventually gets

$$\{\mathbf{a}_i : 0 \leq i < 9\} = \{(\mathbf{0}), (1, 0, 0, 1, 0, 1), (0, 0, 1, 0, 1, 1), (0, 1, 0, 1, 1, 1), (1, 0, 1, 1, 1, 2), \\ (0, 1, 1, 1, 2, 2), (1, 1, 1, 2, 2, 0), (0, 1, 2, 1, 0, 0), (2, 1, 1, 0, 2, 1)\}$$

from which we can explicitly compute elements in $\{\mathbf{a} + \mathbf{a}_i : 0 \leq i < 9\}$.

All of the 18 states we have determined above belong to their 18 respective distinct cycles in $\Omega(g^2(x)) \setminus \Omega(g(x))$. \square

Applying the approach recursively, starting from the states of distinct cycles in $\Omega(g(x))$, we determine the states of distinct cycles in $\Omega(g^b(x))$. For brevity, we present them as $n \cdot b$ -stage states. This step can be performed by using well-known properties of the LFSRs or, better still, by the method that we propose in the next section.

5 The Cycle Structure of $\Omega(f(x))$

We combine the results established in previous sections to study $\Omega(f(x))$. The first subsection considers the cycle structure of $\Omega(f(x))$. The second one develops a procedure to find a state of each cycle in $\Omega(f(x))$.

5.1 The Cycle Structure

We start by setting up the notations. Recall that $f(x) = \prod_{i=1}^k g_i^{b_i}(x)$. Let $g_i(x)$ be of degree n_i having β_i as a root. Let $e_i = \text{ord}(g_i(x)) = \text{ord}(\beta_i)$ and $t_i = \frac{q^{n_i} - 1}{e_i}$. Based on Lemma 2, we label the cycles in $\Omega(g_i^{b_i}(x))$:

$$[\mathbf{0}], [\mathbf{u}_0^i], [\mathbf{u}_1^i], \dots, [\mathbf{u}_{\sigma_i-1}^i] \text{ with } \sigma_i = \sum_{r_i=1}^{b_i} t_i \frac{q^{n_i(r_i-1)}}{p^{c_{r_i}}} \quad (12)$$

where c_{r_i} is the least integer satisfying $p^{c_{r_i}} \geq r_i$. The respective periods and states of the cycles are $1, e_0^i, e_1^i, \dots, e_{\sigma_i-1}^i$ and $\mathbf{0}, \mathbf{a}_0^i, \mathbf{a}_1^i, \dots, \mathbf{a}_{\sigma_i-1}^i \in \mathbb{F}_q^{b_i \cdot n_i}$.

The cycle structure of $\Omega(f(x))$ can be deduced from the three properties listed at the end of Section 2. We give the formal statement in the next theorem.

Theorem 1 *Let $\gamma := \gcd(e_{j_k}^k, \text{lcm}(e_{j_1}^1, \dots, e_{j_{k-1}}^{k-1}))$. Then*

$$\Omega(f(x)) = [\mathbf{0}] \bigcup_{i=1}^k \bigcup_{j=0}^{\sigma_i-1} [\mathbf{u}_j^i] \bigcup_{1 \leq i_1 < i_2 \leq k} \bigcup_{j_1=0}^{\sigma_{i_1}-1} \bigcup_{j_2=0}^{\sigma_{i_2}-1} \bigcup_{\ell_2=0}^{\gcd(e_{j_2}^{i_2}, e_{j_1}^{i_1})-1} [\mathbf{u}_{j_1}^{i_1} + L^{\ell_2} \mathbf{u}_{j_2}^{i_2}] \dots \dots \\ \bigcup_{j_1=0}^{\sigma_1-1} \dots \bigcup_{j_k=0}^{\sigma_k-1} \bigcup_{\ell_2=0}^{\gcd(e_{j_2}^2, e_{j_1}^1)-1} \dots \bigcup_{\ell_k=0}^{\gamma-1} [\mathbf{u}_{j_1}^1 + L^{\ell_2} \mathbf{u}_{j_2}^2 + \dots + L^{\ell_k} \mathbf{u}_{j_k}^k]. \quad (13)$$

If some cycles being the same is allowed, we can express $\Omega(f(x))$ more succinctly as

$$\Omega(f(x)) = \bigcup_{\substack{a_i \in \{0,1\} \\ 1 \leq i \leq k}} \bigcup_{j_1=0}^{\sigma_1-1} \cdots \bigcup_{j_k=0}^{\sigma_k-1} \bigcup_{\ell_2=0}^{\gcd(e_{j_2}^2, e_{j_1}^1)-1} \cdots \bigcup_{\ell_k=0}^{\gamma-1} \left[a_1 \mathbf{u}_{j_1}^1 + a_2 L^{\ell_2} \mathbf{u}_{j_2}^2 + \cdots + a_k L^{\ell_k} \mathbf{u}_{j_k}^k \right]. \quad (14)$$

Remark 3 Theorem 1 shows the types of nonzero cycles that we have in $\Omega(f(x))$. Cycles $[\mathbf{u}_j^i]$ come from $\Omega(g_i^{b_i}(x))$. For an r -subset $S_r := \{i_1, i_2, \dots, i_r\}$ of $\{1, 2, \dots, k\}$ with $i_j < i_\ell$ for $j < \ell$, cycles $[a_1 \mathbf{u}_{j_1}^{i_1} + a_2 L^{\ell_2} \mathbf{u}_{j_2}^{i_2} + \cdots + a_r L^{\ell_r} \mathbf{u}_{j_r}^{i_r}]$ are from $\Omega(\prod_{i \in S_r} (g_i^{b_i}(x)))$. If some of the a_1, a_2, \dots, a_k are 0, then $[a_1 \mathbf{u}_{j_1}^1 + a_2 L^{\ell_2} \mathbf{u}_{j_2}^2 + \cdots + a_k L^{\ell_k} \mathbf{u}_{j_k}^k]$ may be the same for different parameter choices.

Proof When $k = 1$, the statements hold.

Let $k = 2$ and $f(x) = g_1^{b_1}(x) g_2^{b_2}(x) \in \mathbb{F}_q[x]$. It is clear that $\Omega(f(x))$ contains $\Omega(g_1^{b_1}(x))$ and $\Omega(g_2^{b_2}(x))$ as subsets. Hence,

$$[\mathbf{0}] \bigcup_{j_1=0}^{\sigma_1-1} [\mathbf{u}_{j_1}^1] \bigcup_{j_2=0}^{\sigma_2-1} [\mathbf{u}_{j_2}^2] \subseteq \Omega(f(x)).$$

All other sequences in $\Omega(f(x))$ must be of the form

$$L^{\ell_1} \mathbf{u}_{j_1}^1 + L^{\ell_2} \mathbf{u}_{j_2}^2 = L^{\ell_1} \left(\mathbf{u}_{j_1}^1 + L^{\ell_2 - \ell_1} \mathbf{u}_{j_2}^2 \right) \quad (15)$$

with $0 \leq j_1 < \delta_1$, $0 \leq j_2 < \delta_2$, $0 \leq \ell_1 < e_{j_1}^1$, $0 \leq \ell_2 < e_{j_2}^2$, and $\ell_2 - \ell_1$ reduced modulo $e_{j_2}^2$. The period is $\text{lcm}(e_{j_1}^1, e_{j_2}^2)$.

When $j_1 \neq j_1'$ or $j_2 \neq j_2'$, sequences $L^{\ell_1} (\mathbf{u}_{j_1}^1 + L^{\ell_2 - \ell_1} \mathbf{u}_{j_2}^2)$ and $L^{\ell_1'} (\mathbf{u}_{j_1'}^1 + L^{\ell_2 - \ell_1'} \mathbf{u}_{j_2}^2)$ are shift inequivalent. For a contradiction, assume that for some ℓ_1, ℓ_1', ℓ_2 , and ℓ_2' ,

$$L^{\ell_1} \mathbf{u}_{j_1}^1 + L^{\ell_2} \mathbf{u}_{j_2}^2 = L^{\ell_1'} \mathbf{u}_{j_1'}^1 + L^{\ell_2'} \mathbf{u}_{j_2}^2 \iff L^{\ell_1} \mathbf{u}_{j_1}^1 - L^{\ell_1'} \mathbf{u}_{j_1'}^1 = -(L^{\ell_2} \mathbf{u}_{j_2}^2 - L^{\ell_2'} \mathbf{u}_{j_2}^2) = \mathbf{0}.$$

The last equality holds since

$$L^{\ell_1} \mathbf{u}_{j_1}^1 - L^{\ell_1'} \mathbf{u}_{j_1'}^1 \in \Omega(g_1^{b_1}(x)) \text{ and } L^{\ell_2} \mathbf{u}_{j_2}^2 - L^{\ell_2'} \mathbf{u}_{j_2}^2 \in \Omega(g_2^{b_2}(x)),$$

and $g_1(x)$ and $g_2(x)$ are distinct irreducible polynomials. This forces $j_1 = j_1'$ and $j_2 = j_2'$, a contradiction.

The $e_{j_1}^1 \cdot e_{j_2}^2$ distinct sequences having the form of (15) are divided into $\frac{e_{j_1}^1 \cdot e_{j_2}^2}{\text{lcm}(e_{j_1}^1, e_{j_2}^2)} = \gcd(e_{j_1}^1, e_{j_2}^2)$ distinct cycles. Let $\tau_1 := \frac{e_{j_1}^1}{\gcd(e_{j_1}^1, e_{j_2}^2)}$ and $\tau_2 := \frac{e_{j_2}^2}{\gcd(e_{j_1}^1, e_{j_2}^2)}$. We show that

$$\mathbf{u}_{j_1}^1 + L^{\ell_2} \mathbf{u}_{j_2}^2 \text{ and } \mathbf{u}_{j_1}^1 + L^{\ell_2 + \kappa \cdot \gcd(e_{j_1}^1, e_{j_2}^2)} \mathbf{u}_{j_2}^2$$

are shift equivalent for $0 \leq \ell_2 < \gcd(e_{j_1}^1, e_{j_2}^2)$ and $0 < \kappa < \tau_2$.

Since $\gcd(\tau_1, \tau_2) = 1$, there exist $v, w \in \mathbb{Z}$ such that $v \cdot \tau_1 + w \cdot \tau_2 = 1$, implying

$$\kappa \cdot v \cdot e_{j_1}^1 = \kappa (\gcd(e_{j_1}^1, e_{j_2}^2) - w \cdot e_{j_2}^2).$$

The respective periods of $\mathbf{u}_{j_1}^1$ and $\mathbf{u}_{j_2}^2$ are $e_{j_1}^1$ and $e_{j_2}^2$. Hence,

$$\mathbf{u}_{j_1}^1 + L^{\ell_2 + t \cdot \gcd(e_{j_1}^1, e_{j_2}^2)} \mathbf{u}_{j_2}^2 = L^{\kappa \cdot v \cdot e_{j_1}^1} + L^{\ell_2 + \kappa \cdot (\gcd(e_{j_1}^1, e_{j_2}^2) - w \cdot e_{j_2}^2)} \mathbf{u}_{j_2}^2 = L^{\kappa \cdot v \cdot e_{j_1}^1} (\mathbf{u}_{j_1}^1 + L^{\ell_2} \mathbf{u}_{j_2}^2).$$

Thus, a (j_1, j_2) pair corresponds to the cycle $\mathbf{u}_{j_1}^1 + L^{\ell_2} \mathbf{u}_{j_2}^2$ with $0 \leq \ell_2 \leq \gcd(e_{j_1}^1, e_{j_2}^2) - 1$.

Going through all possible (j_1, j_2) pairs gives us all the possible cycles, confirming (13) for the case of $k = 2$. The case of $k > 2$ follows by induction. \square

5.2 A State Belonging to Each Cycle

We now denote a cycle in $\Omega(f(x))$ by $[\mathbf{c}]$ with $\mathbf{c} = a_1 \mathbf{u}_{j_1}^1 + a_2 L^{\ell_2} \mathbf{u}_{j_2}^2 + \dots + a_k L^{\ell_k} \mathbf{u}_{j_k}^k$. If $a_i = 0$ for some i , then $a_i \mathbf{u}_{j_i}^i$ is the all zero sequence $\mathbf{0}$.

Let $n'_i = b_i \cdot n_i$ and $n = n'_1 + n'_2 + \dots + n'_s$. By Sections 3 and 4, to find a state of \mathbf{c} , it suffices to find an n -stage state $a_i \mathbf{v}_{j_i}^i \in \mathbb{F}_q^n$ of $a_i \mathbf{u}_{j_i}^i$ for all i . The desired state is

$$a_1 \mathbf{a}_{j_1}^1 + a_2 T^{\ell_2} \mathbf{a}_{j_2}^2 + \dots + a_k T^{\ell_k} \mathbf{a}_{j_k}^k. \quad (16)$$

The representation in (16) has several limitations. Given such a global expression, we can only use the properties of the whole cycle $[\mathbf{c}]$ without being able to utilize the properties of every *component sequence* $\mathbf{u}_{j_i}^i$. When the period of $[\mathbf{c}]$ is fairly large, determining whether a state is in this cycle may be hard to do, so we propose an alternative.

To each $g_i^{b_i}(x)$, one associates an $n'_i \times n'_i$ matrix \mathcal{A}_i similar to the one constructed in (3), *i.e.*, treating $g_i^{b_i}(x)$ as a characteristic polynomial. Construct the $n'_i \times n$ matrix \mathcal{P}_i in the following manner. The first n'_i columns form the identity matrix $I_{n'_i}$. The matrix \mathcal{A}_i occupies columns 2 to $n'_i + 1$. Next, add columns recursively by appending the last column of \mathcal{A}_i^j , starting with $j = 2$ until all n columns of \mathcal{P}_i are completed.

Given an n'_i -stage state \mathbf{a}_i of some sequence in $\Omega(g_i^{b_i}(x))$, it is evident that $\mathbf{a}_i \mathcal{P}_i$ is the corresponding n -stage state of that same sequence.

Lemma 7 *The $n \times n$ matrix*

$$\mathcal{P} = \begin{pmatrix} \mathcal{P}_1 \\ \mathcal{P}_2 \\ \vdots \\ \mathcal{P}_k \end{pmatrix} \text{ is of full rank, i.e., } \text{rank}(\mathcal{P}) = n, \text{ making } \mathcal{P} \in GL(n, \mathbb{F}_q).$$

Proof Let $\alpha_{i,j}$ denote the j -th row of \mathcal{P}_i . We show that the rows $\alpha_{i,j}$ of \mathcal{P} are linearly independent for all (i, j) with $1 \leq i \leq k$ and $1 \leq j \leq n'_i$.

Notice that $\alpha_{i,j}$ is the first n entries of the sequence from LFSR with characteristic polynomial $g_i^{b_i}(x)$ and initial n'_i -stage state $(0, \dots, 0, 1, 0, \dots, 0) \in \mathbb{F}_q^{n'_i}$, where the unique 1 is in the j -th position. For a fixed i , it is clear that the rows of \mathcal{P}_i are linearly independent.

For a contradiction, suppose that there exists a linear combination of the n rows of \mathcal{P}

$$\sum_{i=1}^k \sum_{j=1}^{n'_i} a_{i,j} \boldsymbol{\alpha}_{i,j} = \mathbf{0}$$

with not all $a_{i,j} = 0$. Without loss of generality, with not all $a_{1,j} = 0$, write

$$\sum_{j=1}^{n'_1} a_{1,j} \boldsymbol{\alpha}_{1,j} = \sum_{i=2}^k \sum_{j=1}^{n'_i} (-a_{i,j}) \boldsymbol{\alpha}_{i,j}. \quad (17)$$

The left hand side of (17) is the first n entries of a sequence in $\Omega(g_1^{b_1}(x))$ with a nonzero initial state of length n'_1 while the right hand side is the first n entries of a sequence in $\Omega(g_2^{b_2}(x) \cdots g_k^{b_k}(x))$ with a nonzero initial state of length $n - n'_1$. Since $\deg(g_1^{b_1}(x))$ and $\deg(g_2^{b_2}(x) \cdots g_k^{b_k}(x))$ are $< n$, if the first n entries of these two sequences are equal, then they must be the same sequence. Hence, there exists a sequence that simultaneously belongs to both $\Omega(g_1^{b_1}(x))$ and $\Omega(g_2^{b_2}(x) \cdots g_k^{b_k}(x))$. Since $g_1(x), \dots, g_k(x)$ are pairwise distinct, this sequence must be $\mathbf{0}$, a contradiction. \square

Suppose that we already have \mathcal{P} . Let $\mathbf{v} \in \mathbb{F}_q^n$ and $\mathbf{a}_i \in \mathbb{F}_q^{n'_i}$ with $1 \leq i \leq k$ be the respective n -stage and n'_i -stage states of the sequences in $\Omega(f(x))$ and $\Omega(g_i^{b_i}(x))$. Then \mathcal{P} provides a one-to-one correspondence between \mathbf{v} and $(\mathbf{a}_1, \mathbf{a}_2, \dots, \mathbf{a}_k)$ via $\mathbf{v} = (\mathbf{a}_1, \mathbf{a}_2, \dots, \mathbf{a}_k) \mathcal{P}$. Since

$$T\mathbf{v} = T[(\mathbf{a}_1, \mathbf{a}_2, \dots, \mathbf{a}_k) \mathcal{P}] = (T\mathbf{a}_1, T\mathbf{a}_2, \dots, T\mathbf{a}_k) \mathcal{P},$$

we know \mathcal{P} and T commute. Hence, any sequence $\mathbf{s} \in \Omega(f(x))$ with an initial state \mathbf{v} is the sum of sequences \mathbf{s}_i from $\Omega(g_i^{b_i}(x))$ with corresponding initial states \mathbf{a}_i for $1 \leq i \leq k$. When convenient, we use $(\mathbf{a}_1, \mathbf{a}_2, \dots, \mathbf{a}_k)$ to represent \mathbf{v} .

Assume that we have obtained the set $E_i := \{\mathbf{0}, \mathbf{a}_0^i, \mathbf{a}_1^i, \dots, \mathbf{a}_{\sigma_i-1}^i\}$, where \mathbf{a}_j^i is a state of cycle $[\mathbf{s}_j^i]$ and $\mathbf{0}$ the state of $[\mathbf{0}]$. This set contains all of the states corresponding to the $\sigma_i + 1$ distinct cycles in $\Omega(g_i^{b_i}(x))$. Then

$$\mathbf{v} = (\mathbf{a}_{j_1}^1, \mathbf{a}_{j_2}^2, \dots, \mathbf{a}_{j_k}^k) \mathcal{P} \text{ with } \mathbf{a}_{j_i}^i \in E_i \quad (18)$$

can be taken as an initial state of a cycle $[\mathbf{s}] \in \Omega(f(x))$. Notice that

$$\mathbf{s} = a_1 \mathbf{u}_{j_1}^1 + a_2 \mathbf{u}_{j_2}^2 + \cdots + a_k \mathbf{u}_{j_k}^k$$

with $a_i = 0$ if the corresponding component state is $\mathbf{0}$ and $a_i = 1$ otherwise.

Let ℓ_i be a nonnegative integer for $1 \leq i \leq k$. By the state obtained in (18) and by the properties of \mathcal{P} and T , the state $\mathbf{v}' = (T^{\ell_1} \mathbf{a}_{j_1}^1, T^{\ell_2} \mathbf{a}_{j_2}^2, \dots, T^{\ell_k} \mathbf{a}_{j_k}^k) \mathcal{P}$ belongs to cycle

$$\left[a_1 L^{\ell_1} \mathbf{s}_{j_1}^1 + a_2 L^{\ell_2} \mathbf{s}_{j_2}^2 + \cdots + a_k L^{\ell_k} \mathbf{s}_{j_k}^k \right] = \left[a_1 \mathbf{s}_{j_1}^1 + a_2 L^{\ell_2 - \ell_1} \mathbf{s}_{j_2}^2 + \cdots + a_k L^{\ell_k - \ell_1} \mathbf{s}_{j_k}^k \right].$$

This approach enables us to find a state belonging to any cycle in $\Omega(f(x))$.

Example 4 Consider the characteristic polynomial

$$\begin{aligned} f(x) &= \underbrace{(x^2+1)^2}_{=g_1^2(x)} \underbrace{(x^3+2x+2)^2}_{=g_2^2(x)} \underbrace{(x^3+x^2+2)^2}_{=g_3^2(x)} \in \mathbb{F}_3[x] \\ &= (x^4+2x^2+1)(x^6+x^4+x^3+x^2+2x+1)(x^6+2x^5+x^4+x^3+x^2+1) \\ &= x^{16}+2x^{15}+x^{14}+2x^{13}+x^{12}+x^{11}+x^9+x^8+x^7+x^5+x^4+2x^3+x^2+2x+1. \end{aligned}$$

The parameters are $(e_1, t_1) = (4, 2)$, $(e_2, t_2) = (e_3, t_3) = (13, 2)$, and $b_i = 2$ for all i . Our $g_2(x)$ and $g_3(x)$ here are exactly the $g_2(x)$ and $g_3(x)$ in Example 1 so we can use its relevant results. We choose $p_1(x) = x^2 + x + 2$ as the associated primitive polynomial of $g_1(x)$. The two nonzero cycles in $\Omega(g_1(x))$ are $[(1, 0, 2, 0)]$ and $[(1, 1, 2, 2)]$.

With each respective \mathcal{A}_i in the box, the matrix \mathcal{P} is given by

$$\begin{aligned} \mathcal{P} &= \begin{pmatrix} \mathcal{P}_1 \\ \mathcal{P}_2 \\ \mathcal{P}_3 \end{pmatrix} \text{ with } \mathcal{P}_1 = \begin{pmatrix} \boxed{1} & \boxed{0} & \boxed{0} & \boxed{0} & \boxed{2} & 0 & 2 & 0 & 0 & 0 & 1 & 0 & 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 & 0 & 2 & 0 & 2 & 0 & 0 & 0 & 1 & 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 & 1 & 0 & 0 & 0 & 2 & 0 & 2 & 0 & 0 & 0 & 1 & 0 \\ 0 & \boxed{0} & \boxed{0} & \boxed{1} & \boxed{0} & 1 & 0 & 0 & 0 & 2 & 0 & 2 & 0 & 0 & 0 & 1 \end{pmatrix}, \\ \mathcal{P}_2 &= \begin{pmatrix} \boxed{1} & \boxed{0} & \boxed{0} & \boxed{0} & \boxed{0} & \boxed{0} & \boxed{2} & 0 & 1 & 1 & 0 & 0 & 2 & 0 & 1 & 0 \\ 0 & 1 & 0 & 0 & 0 & 0 & 1 & 2 & 2 & 0 & 1 & 0 & 1 & 2 & 2 & 1 \\ 0 & 0 & 1 & 0 & 0 & 0 & 2 & 1 & 0 & 0 & 0 & 1 & 2 & 1 & 0 & 2 \\ 0 & 0 & 0 & 1 & 0 & 0 & 2 & 2 & 1 & 0 & 0 & 0 & 2 & 2 & 0 & 0 \\ 0 & 0 & 0 & 0 & 1 & 0 & 2 & 2 & 0 & 0 & 1 & 0 & 2 & 0 & 0 & 2 \\ 0 & \boxed{0} & \boxed{0} & \boxed{0} & \boxed{0} & \boxed{1} & \boxed{0} & 2 & 2 & 0 & 0 & 1 & 0 & 2 & 0 & 0 \end{pmatrix}, \quad \mathcal{P}_3 = \begin{pmatrix} \boxed{1} & \boxed{0} & \boxed{0} & \boxed{0} & \boxed{0} & \boxed{0} & \boxed{2} & 2 & 0 & 2 & 1 & 0 & 1 & 2 & 0 & 1 \\ 0 & 1 & 0 & 0 & 0 & 0 & 0 & 2 & 2 & 0 & 2 & 1 & 0 & 1 & 2 & 0 \\ 0 & 0 & 1 & 0 & 0 & 0 & 2 & 2 & 2 & 1 & 1 & 2 & 2 & 2 & 1 & 0 \\ 0 & 0 & 0 & 1 & 0 & 0 & 2 & 1 & 2 & 1 & 2 & 1 & 0 & 1 & 2 & 2 \\ 0 & 0 & 0 & 0 & 1 & 0 & 2 & 1 & 1 & 1 & 2 & 2 & 2 & 2 & 1 & 0 \\ 0 & \boxed{0} & \boxed{0} & \boxed{0} & \boxed{0} & \boxed{1} & \boxed{1} & 0 & 1 & 2 & 0 & 2 & 1 & 0 & 2 & 0 \end{pmatrix}. \end{aligned} \tag{19}$$

There are 8 nonzero cycles in $\Omega(g_1^2(x))$: 2 cycles of period 4 in $\Omega(g_1(x))$ and 6 cycles, each with period 12, in $\Omega(g_1^2(x)) \setminus \Omega(g_1(x))$. Performing the required computations, the first 3 cycles with period 12 can be derived using $\mathbf{b} = (1, 0)$, $\mathbf{a} = (1, 0, 0, 0)$, and $\mathbf{a}' = T^4\mathbf{a} - \mathbf{a} = (1, 0, 2, 0)$, yielding $\mathbf{a}_i \in \{(\mathbf{0}), (0, 2, 0, 1), (0, 1, 0, 2)\}$. The 4-stage states that we want are $\{\mathbf{a} + \mathbf{a}_i\}$. To get the other 3, we use $\mathbf{b} = (1, 1)$, $\mathbf{a} = (1, 1, 0, 0)$, and $\mathbf{a}' = T^4\mathbf{a} - \mathbf{a} = (1, 1, 2, 2)$, yielding $\mathbf{a}_i \in \{(\mathbf{0}), (1, 0, 2, 0), (0, 1, 0, 2)\}$. Explicitly determining elements in $\{\mathbf{a} + \mathbf{a}_i\}$ completes our task for E_1 .

Examples 1 and 3 lead us to the nonzero cycles in $\Omega(g_2^2(x))$. There are 2 cycles of period 13 in $\Omega(g_2(x))$ and 18 cycles, each with period 39, in $\Omega(g_2^2(x)) \setminus \Omega(g_2(x))$.

To determine all nonzero cycles in $\Omega(g_3^2(x))$ we perform a similar analysis. Table 1 already listed the 2 cycles of period 13 in $\Omega(g_3(x))$. The remaining 18 cycles, each with period 39, can be easily determined. The first 9 comes from $\mathbf{b} = (1, 1, 0)$, $\mathbf{a} = (0, 0, 0, 1, 0, 0)$, and $\mathbf{a}' = T^{13}\mathbf{a} - \mathbf{a} = (1, 2, 2, 2, 0, 2)$. The initial states are $\mathbf{a} + \mathbf{a}_i$ with

$$\begin{aligned} \mathbf{a}_i \in \{ & (\mathbf{0}), (1, 1, 0, 1, 0, 0), (1, 0, 1, 0, 0, 1), (0, 1, 0, 0, 1, 2), (1, 0, 0, 1, 2, 1), \\ & (0, 0, 1, 2, 1, 0), (1, 2, 1, 0, 2, 2), (0, 2, 2, 1, 1, 1), (2, 0, 1, 1, 2, 2) \}. \end{aligned}$$

The remaining 9 comes from $\mathbf{b} = (1, 2, 0)$, $\mathbf{a} = (0, 0, 0, 1, 1, 2)$, and $\mathbf{a}' = (0, 1, 2, 1, 0, 2)$. The initial states are $\mathbf{a} + \mathbf{a}_i$ with

$$\begin{aligned} \mathbf{a}_i \in \{ & (\mathbf{0}), (1, 1, 0, 1, 0, 0), (0, 1, 0, 0, 1, 2), (1, 0, 0, 1, 2, 1), (2, 1, 0, 2, 2, 1), \\ & (1, 0, 2, 2, 1, 1), (2, 2, 1, 1, 1, 0), (2, 1, 1, 1, 0, 1), (0, 1, 1, 2, 2, 2) \}. \end{aligned}$$

The complete list of nonzero cycles and their corresponding states is in Table 2.

Table 2 States and Nonzero Cycles in $\Omega(g_i^2(x))$

No.	State	Nonzero Cycle
1	$\mathbf{a}_0^1 = (1, 0)$	$\mathbf{u}_0^1 = (1, 0, 2, 0)$
2	$\mathbf{a}_1^1 = (1, 1)$	$\mathbf{u}_1^1 = (1, 1, 2, 2)$
3	$\mathbf{a}_2^1 = (1, 0, 0, 0)$	$\mathbf{u}_2^1 = (1, 0, 0, 2, 0, 2, 0, 0, 0, 1, 0)$
4	$\mathbf{a}_3^1 = (1, 2, 0, 1)$	$\mathbf{u}_3^1 = (1, 2, 0, 1, 2, 2, 2, 1, 0, 2, 1, 1)$
5	$\mathbf{a}_4^1 = (1, 1, 0, 2)$	$\mathbf{u}_4^1 = (1, 1, 0, 2, 2, 1, 2, 2, 0, 1, 1, 2)$
6	$\mathbf{a}_5^1 = (1, 1, 0, 0)$	$\mathbf{u}_5^1 = (1, 1, 0, 0, 2, 2, 2, 2, 0, 0, 1, 1)$
7	$\mathbf{a}_6^1 = (2, 1, 2, 0)$	$\mathbf{u}_6^1 = (2, 1, 2, 0, 0, 2, 1, 2, 1, 0, 0, 1)$
8	$\mathbf{a}_7^1 = (1, 2, 0, 2)$	$\mathbf{u}_7^1 = (1, 2, 0, 2, 2, 0, 2, 1, 0, 1, 1, 0)$
1	$\mathbf{a}_0^2 = (1, 1, 1)$	$\mathbf{u}_0^2 = (1, 1, 1, 2, 2, 0, 1, 2, 1, 0, 0, 1, 0)$
2	$\mathbf{a}_1^2 = (1, 2, 0)$	$\mathbf{u}_1^2 = (1, 2, 0, 0, 2, 0, 2, 2, 1, 1, 0, 2)$
3	$\mathbf{a}_2^2 = (0, 0, 0, 1, 0, 1)$	$\mathbf{u}_2^2 = (0, 0, 0, 1, 0, 1, 2, 1, 1, 0, 1, 0, 1, 2, 0, 1, 2, 1, 1, 0, 0, 2, 1, 1, 2, 2, 1, 0, 1, 1, 1, 0, 2, 2, 0, 2, 1, 1)$
4	$\mathbf{a}_3^2 = (1, 0, 0, 2, 0, 2)$	$\mathbf{u}_3^2 = (1, 0, 0, 2, 0, 2, 0, 0, 0, 2, 2, 2, 2, 0, 2, 2, 2, 1, 2, 1, 1, 2, 1, 0, 1, 0, 2, 1, 2, 1, 0, 1, 2, 2, 2, 0)$
5	$\mathbf{a}_4^2 = (0, 0, 1, 1, 1, 2)$	$\mathbf{u}_4^2 = (0, 0, 1, 1, 1, 2, 0, 0, 0, 1, 1, 0, 1, 1, 2, 1, 1, 0, 2, 2, 2, 2, 2, 0, 0, 2, 1, 1, 1, 2, 2, 2, 1, 1, 1, 0, 0, 0, 2)$
6	$\mathbf{a}_5^2 = (1, 0, 1, 2, 1, 0)$	$\mathbf{u}_5^2 = (1, 0, 1, 2, 1, 0, 1, 1, 2, 0, 1, 1, 0, 2, 2, 1, 2, 0, 0, 0, 0, 1, 1, 2, 1, 2, 0, 1, 1, 2, 2, 0, 2, 0, 2, 0, 1, 1)$
7	$\mathbf{a}_6^2 = (1, 1, 1, 0, 2, 1)$	$\mathbf{u}_6^2 = (1, 1, 1, 0, 2, 1, 0, 0, 2, 1, 0, 2, 0, 1, 0, 1, 1, 2, 2, 2, 1, 2, 1, 2, 2, 0, 2, 1, 0, 0, 1, 1, 1, 0, 0, 2, 2, 1)$
8	$\mathbf{a}_7^2 = (1, 1, 2, 0, 0, 2)$	$\mathbf{u}_7^2 = (1, 1, 2, 0, 0, 2, 1, 2, 1, 1, 1, 1, 1, 2, 0, 2, 0, 2, 2, 0, 1, 0, 2, 2, 1, 0, 0, 2, 2, 0, 1, 2, 2, 0, 2, 0, 0, 1, 2)$
9	$\mathbf{a}_8^2 = (1, 2, 2, 1, 1, 0)$	$\mathbf{u}_8^2 = (1, 2, 2, 1, 1, 0, 0, 1, 1, 2, 0, 2, 1, 2, 1, 2, 1, 0, 0, 2, 0, 0, 0, 1, 2, 0, 0, 0, 2, 1, 2, 0, 1, 2, 2, 1, 2, 2, 2)$
10	$\mathbf{a}_9^2 = (0, 1, 2, 2, 0, 1)$	$\mathbf{u}_9^2 = (0, 1, 2, 2, 0, 1, 0, 1, 2, 2, 1, 0, 2, 1, 0, 2, 2, 2, 1, 2, 0, 1, 0, 2, 0, 1, 2, 2, 2, 2, 1, 1, 1, 2, 0, 1, 0, 0, 0)$
11	$\mathbf{a}_{10}^2 = (1, 1, 0, 0, 1, 0)$	$\mathbf{u}_{10}^2 = (1, 1, 0, 0, 1, 0, 2, 1, 0, 1, 2, 0, 2, 2, 0, 0, 0, 0, 0, 1, 0, 2, 2, 0, 0, 1, 0, 2, 0, 0, 2, 0, 0, 2, 1, 0, 1, 0, 0)$
12	$\mathbf{a}_{11}^2 = (0, 0, 0, 2, 1, 0)$	$\mathbf{u}_{11}^2 = (0, 0, 0, 2, 1, 0, 0, 0, 1, 2, 1, 0, 2, 1, 1, 2, 1, 1, 2, 1, 1, 2, 2, 0, 0, 2, 2, 1, 0, 1, 2, 1, 2, 1, 2, 0, 0, 1)$
13	$\mathbf{a}_{12}^2 = (1, 0, 0, 0, 1, 1)$	$\mathbf{u}_{12}^2 = (1, 0, 0, 0, 1, 1, 1, 1, 0, 1, 1, 1, 1, 2, 1, 2, 2, 1, 2, 0, 2, 0, 1, 2, 1, 2, 0, 2, 1, 1, 1, 0, 2, 0, 0, 1, 0, 1, 0)$
14	$\mathbf{a}_{13}^2 = (0, 0, 1, 2, 2, 1)$	$\mathbf{u}_{13}^2 = (0, 0, 1, 2, 2, 1, 1, 2, 0, 2, 2, 2, 0, 1, 1, 0, 1, 2, 2, 0, 0, 0, 2, 0, 2, 1, 2, 2, 2, 0, 2, 0, 2, 1, 0, 2, 1, 2, 2)$
15	$\mathbf{a}_{14}^2 = (0, 1, 0, 0, 2, 1)$	$\mathbf{u}_{14}^2 = (0, 1, 0, 0, 2, 1, 2, 2, 1, 0, 0, 1, 2, 1, 2, 2, 2, 2, 1, 0, 1, 0, 1, 1, 0, 2, 0, 1, 1, 2, 2, 0, 0, 1, 1, 0, 2, 1, 1)$
16	$\mathbf{a}_{15}^2 = (1, 0, 1, 0, 2, 2)$	$\mathbf{u}_{15}^2 = (1, 0, 1, 0, 2, 2, 2, 0, 2, 1, 2, 0, 2, 2, 1, 0, 2, 2, 0, 1, 1, 2, 1, 0, 0, 0, 0, 2, 2, 1, 2, 1, 0, 2, 2, 1, 1, 0, 1)$
17	$\mathbf{a}_{16}^2 = (0, 1, 1, 0, 0, 2)$	$\mathbf{u}_{16}^2 = (0, 1, 1, 0, 0, 2, 0, 1, 0, 0, 1, 0, 0, 1, 2, 0, 2, 0, 2, 2, 0, 2, 0, 1, 2, 0, 2, 1, 0, 1, 1, 0, 0, 0, 0, 0, 2)$
18	$\mathbf{a}_{17}^2 = (1, 1, 1, 1, 0, 0)$	$\mathbf{u}_{17}^2 = (1, 1, 1, 1, 0, 0, 1, 2, 2, 2, 1, 1, 2, 2, 2, 0, 0, 0, 1, 0, 0, 2, 2, 2, 1, 0, 0, 0, 2, 2, 0, 2, 2, 1, 2, 2, 0, 1, 1)$
19	$\mathbf{a}_{18}^2 = (0, 1, 2, 0, 1, 0)$	$\mathbf{u}_{18}^2 = (0, 1, 2, 0, 1, 0, 1, 0, 2, 0, 2, 2, 1, 1, 2, 2, 1, 1, 0, 1, 2, 0, 0, 2, 2, 2, 0, 0, 1, 1, 2, 2, 2, 2, 0, 1, 2, 0)$
20	$\mathbf{a}_{19}^2 = (2, 1, 1, 2, 0, 1)$	$\mathbf{u}_{19}^2 = (2, 1, 1, 2, 0, 1, 1, 1, 2, 1, 0, 2, 0, 1, 0, 2, 1, 1, 1, 1, 2, 2, 2, 1, 0, 2, 0, 0, 0, 2, 1, 1, 0, 2, 0, 2, 0)$
1	$\mathbf{a}_0^3 = (1, 1, 0)$	$\mathbf{u}_0^3 = (1, 1, 0, 1, 0, 0, 1, 2, 1, 0, 2, 2, 1)$
2	$\mathbf{a}_1^3 = (1, 2, 0)$	$\mathbf{u}_1^3 = (1, 2, 0, 1, 1, 2, 2, 2, 0, 2, 0, 0, 2)$
3	$\mathbf{a}_2^3 = (0, 0, 0, 1, 0, 0)$	$\mathbf{u}_2^3 = (0, 0, 0, 1, 0, 0, 2, 1, 2, 1, 2, 1, 0, 1, 2, 2, 0, 0, 2, 2, 1, 1, 2, 1, 1, 1, 2, 1, 1, 2, 0, 1, 2, 0, 1, 2, 1, 0, 0, 0, 1, 2)$
4	$\mathbf{a}_3^3 = (1, 1, 0, 2, 0, 0)$	$\mathbf{u}_3^3 = (1, 1, 0, 2, 0, 0, 0, 0, 0, 1, 1, 0, 1, 2, 0, 2, 1, 0, 2, 0, 2, 2, 0, 0, 2, 0, 2, 1, 0, 0, 1, 0, 0, 1, 0, 2, 0, 0)$
5	$\mathbf{a}_4^3 = (1, 0, 1, 1, 0, 1)$	$\mathbf{u}_4^3 = (1, 0, 1, 1, 0, 1, 1, 2, 2, 0, 1, 2, 1, 2, 2, 0, 0, 0, 0, 1, 2, 1, 1, 0, 2, 2, 0, 1, 2, 2, 0, 2, 1, 2, 0, 2, 2, 2, 0)$
6	$\mathbf{a}_5^3 = (0, 1, 0, 1, 1, 2)$	$\mathbf{u}_5^3 = (0, 1, 0, 1, 1, 2, 0, 1, 1, 0, 0, 2, 1, 1, 0, 2, 0, 1, 1, 0, 1, 0, 1, 2, 2, 2, 2, 1, 2, 1, 0, 0, 1, 2, 2, 1, 2, 0)$
7	$\mathbf{a}_6^3 = (1, 0, 0, 2, 2, 1)$	$\mathbf{u}_6^3 = (1, 0, 0, 2, 2, 1, 2, 0, 1, 2, 0, 2, 0, 2, 2, 1, 2, 0, 2, 0, 0, 0, 2, 2, 1, 0, 1, 1, 0, 2, 2, 0, 2, 1, 1, 2, 2)$
8	$\mathbf{a}_7^3 = (0, 0, 1, 0, 1, 0)$	$\mathbf{u}_7^3 = (0, 0, 1, 0, 1, 0, 1, 0, 0, 2, 0, 1, 1, 1, 2, 0, 2, 1, 2, 1, 0, 2, 0, 2, 1, 2, 2, 1, 2, 1, 1, 1, 0, 1, 1, 1, 0)$
9	$\mathbf{a}_8^3 = (1, 2, 1, 1, 2, 2)$	$\mathbf{u}_8^3 = (1, 2, 1, 1, 2, 2, 0, 2, 0, 1, 0, 1, 0, 2, 1, 0, 2, 2, 2, 1, 1, 0, 0, 2, 2, 2, 0, 0, 2, 1, 0, 1, 1, 2)$
10	$\mathbf{a}_9^3 = (0, 2, 2, 2, 1, 1)$	$\mathbf{u}_9^3 = (0, 2, 2, 2, 1, 1, 2, 2, 2, 1, 0, 0, 1, 1, 1, 1, 1, 0, 2, 2, 1, 2, 2, 0, 2, 0, 0, 0, 1, 2, 2, 2, 0, 0, 1, 0, 0)$
11	$\mathbf{a}_{10}^3 = (2, 0, 1, 2, 2, 2)$	$\mathbf{u}_{10}^3 = (2, 0, 1, 2, 2, 2, 1, 1, 1, 1, 2, 0, 1, 0, 2, 0, 1, 2, 1, 1, 1, 0, 2, 1, 0, 2, 1, 1, 2, 0, 2, 0, 1, 1, 2, 0, 0, 0, 0)$
12	$\mathbf{a}_{11}^3 = (0, 0, 0, 1, 1, 2)$	$\mathbf{u}_{11}^3 = (0, 0, 0, 1, 1, 2, 0, 2, 2, 0, 1, 1, 1, 0, 1, 2, 2, 1, 1, 2, 0, 0, 1, 1, 2, 1, 0, 2, 1, 0, 2, 1, 0, 1, 0, 1, 1, 1, 2, 1, 0, 1)$
13	$\mathbf{a}_{12}^3 = (1, 1, 0, 2, 1, 2)$	$\mathbf{u}_{12}^3 = (1, 1, 0, 2, 1, 2, 1, 1, 0, 0, 0, 2, 1, 2, 2, 0, 1, 1, 0, 2, 1, 1, 0, 2, 1, 0, 1, 2, 1, 0, 1, 1, 1, 0, 2, 0, 2, 2, 0, 2, 2)$
14	$\mathbf{a}_{13}^3 = (0, 1, 0, 1, 2, 1)$	$\mathbf{u}_{13}^3 = (0, 1, 0, 1, 2, 1, 1, 2, 1, 2, 2, 2, 0, 2, 2, 2, 0, 0, 0, 2, 0, 2, 0, 2, 0, 0, 1, 0, 2, 2, 2, 1, 0, 1, 2, 1, 2)$
15	$\mathbf{a}_{14}^3 = (1, 0, 0, 2, 0, 0)$	$\mathbf{u}_{14}^3 = (1, 0, 0, 2, 0, 0, 0, 1, 1, 1, 2, 2, 1, 1, 1, 2, 0, 0, 2, 2, 2, 2, 2, 0, 1, 1, 2, 1, 1, 0, 1, 1, 0, 0, 0, 2, 1, 1)$
16	$\mathbf{a}_{15}^3 = (2, 1, 0, 0, 0, 0)$	$\mathbf{u}_{15}^3 = (2, 1, 0, 0, 0, 0, 1, 0, 2, 1, 1, 1, 2, 2, 2, 2, 1, 0, 2, 0, 1, 0, 2, 1, 2, 2, 2, 0, 1, 2, 0, 1, 2, 2, 1, 0, 1, 0, 2)$
17	$\mathbf{a}_{16}^3 = (1, 0, 2, 0, 2, 0)$	$\mathbf{u}_{16}^3 = (1, 0, 2, 0, 2, 0, 1, 2, 0, 0, 1, 2, 0, 0, 1, 1, 1, 1, 2, 2, 0, 0, 1, 1, 1, 0, 0, 0, 1, 2, 0, 2, 2, 1, 2, 2, 1, 1, 0)$
18	$\mathbf{a}_{17}^3 = (2, 2, 1, 2, 2, 2)$	$\mathbf{u}_{17}^3 = (2, 2, 1, 2, 2, 2, 1, 2, 2, 1, 0, 2, 1, 2, 0, 0, 0, 2, 1, 0, 0, 0, 2, 0, 0, 1, 2, 1, 2, 1, 2, 0, 2, 1, 1, 0, 0, 1, 1)$
19	$\mathbf{a}_{18}^3 = (2, 1, 1, 2, 1, 0)$	$\mathbf{u}_{18}^3 = (2, 1, 1, 2, 1, 0, 0, 0, 2, 2, 1, 0, 2, 2, 0, 0, 1, 2, 2, 0, 1, 0, 2, 2, 0, 2, 0, 2, 1, 1, 1, 1, 1, 2, 1, 2, 0, 0)$
20	$\mathbf{a}_{19}^3 = (0, 1, 1, 0, 0, 1)$	$\mathbf{u}_{19}^3 = (0, 1, 1, 0, 0, 1, 0, 1, 2, 0, 0, 2, 0, 0, 2, 0, 1, 0, 0, 2, 2, 0, 1, 0, 0, 0, 0, 2, 2, 0, 2, 1, 0, 1, 2, 0, 1, 0)$

Consulting (18), we know that

$$\begin{aligned} \mathbf{v} &= (\mathbf{a}_3^1, \mathbf{a}_9^2, \mathbf{a}_8^3) \mathcal{P} = (1, 2, 0, 1, 0, 1, 2, 2, 0, 1, 1, 2, 1, 1, 2, 2) \mathcal{P} \\ &= (2, 2, 0, 1, 1, 2, 2, 1, 2, 2, 2, 2, 0, 2, 1, 0) \end{aligned}$$

is a state of $[\mathbf{s} = \mathbf{u}_3^1 + \mathbf{u}_9^2 + \mathbf{u}_8^3] \in \Omega(f(x))$. The period of \mathbf{s} is $\text{lcm}(12, 39, 39) = 156$. One can quickly verify that using \mathbf{v} as the input state to the LFSR with characteristic polynomial $f(x)$ indeed yields \mathbf{s} since the first 32 entries of the two sequences match. \square

We now highlight the advantage of our new representation of the states of the cycles in $\Omega(f(x))$. To construct de Bruijn sequences by the cycle joining method [5] one must find the conjugate pairs between any two cycles C_1 and C_2 in $\Omega(f(x))$. We transform this problem into one that decides on whether two states belong to the same cycle. The latter can be solved by applying the state shift operator T repeatedly until one state is shown to be the other's cyclic shift or not.

The naive approach is to do exhaustive searching. If the period of the cycles are large, this is time consuming. Using the new representation, Algorithm 1 simplifies the work.

Algorithm 1 Determining if two states belong to the same cycle

Input: \mathcal{P} and two states $\mathbf{v}_1 \neq \mathbf{v}_2$.
Output: Whether $\mathbf{v}_1, \mathbf{v}_2$ are in the same cycle.

```

1: procedure MAIN( $\mathcal{P}, \mathbf{v}_1, \mathbf{v}_2$ )
2:    $(\mathbf{a}_1, \mathbf{a}_2, \dots, \mathbf{a}_k) \leftarrow \mathbf{v}_1 \mathcal{P}^{-1}$ 
3:    $(\mathbf{b}_1, \mathbf{b}_2, \dots, \mathbf{b}_k) \leftarrow \mathbf{v}_2 \mathcal{P}^{-1}$ 
4:   for  $i$  from 1 to  $k$  do ▷ Finding the period of the cycle containing state  $\mathbf{a}_i$ 
5:      $\mathbf{a} \leftarrow \mathbf{a}_i, e_i \leftarrow 0$ 
6:     while  $T\mathbf{a} \neq \mathbf{a}_i$  do
7:        $e_i \leftarrow e_i + 1$ 
8:     end while
9:   end for
10:  for  $i$  from 1 to  $k$  do ▷ Find whether  $T^{\ell_i} \mathbf{a}_i = \mathbf{b}_i$ 
11:     $\mathbf{a} \leftarrow \mathbf{a}_i, \ell_i \leftarrow 0$ 
12:    if  $\mathbf{a} \neq \mathbf{b}_i$  then
13:       $\mathbf{a} \leftarrow T\mathbf{a}, \ell_i \leftarrow 1$ 
14:    end if
15:    while  $\mathbf{a} \neq \mathbf{b}_i$  and  $\mathbf{a} \neq \mathbf{a}_i$  do
16:       $\mathbf{a} \leftarrow T\mathbf{a}, \ell_i \leftarrow \ell_i + 1$ 
17:    end while
18:    if  $\mathbf{a} \neq \mathbf{b}_i$  then
19:      Output NO and break
20:    end if
21:  end for
22:  for  $i$  from 2 to  $k$  do ▷ by Generalized CRT
23:    for  $j$  from 1 to  $i - 1$  do
24:      if  $\text{gcd}(e_i, e_j)$  does not divide  $\ell_i - \ell_j$  then
25:        Output NO and break
26:      end if
27:    end for
28:  end for
29:  Output: YES
30: end procedure

```

Theorem 2 Algorithm 1 is correct.

Proof If the input states \mathbf{v}_1 and \mathbf{v}_2 in $\Omega(f(x))$ are in the same cycle, then there exists an integer ℓ such that $\mathbf{v}_1 = T^\ell \mathbf{v}_2$. Multiplying by \mathcal{P}^{-1} , we have $\mathbf{v}_1 \mathcal{P}^{-1} = (\mathbf{a}_1, \mathbf{a}_2, \dots, \mathbf{a}_k)$ and $\mathbf{v}_2 \mathcal{P}^{-1} = (\mathbf{b}_1, \mathbf{b}_2, \dots, \mathbf{b}_k)$ with \mathbf{a}_i and \mathbf{b}_i in $\Omega(g_i^{b_i}(x))$. If \mathbf{v}_1 and \mathbf{v}_2 are in the same cycle, then \mathbf{a}_i and \mathbf{b}_i must be in the same cycle in $\Omega(g_i^{b_i}(x))$ for all $1 \leq i \leq k$.

The first part of Algorithm 1 determines the period e_i of the cycle containing the state \mathbf{a}_i . The second part tests whether \mathbf{a}_i and \mathbf{b}_i are in the same ‘‘component cycle’’ in $\Omega(g_i^{b_i}(x))$. If the test fail for at least one i then \mathbf{v}_1 and \mathbf{v}_2 belong to distinct cycles. Otherwise, we obtain integers ℓ_i s that satisfy $T^{\ell_i}\mathbf{a}_i = \mathbf{b}_i$ for each i . The last part verifies if there exists an integer ℓ that makes $\mathbf{v}_1 = T^\ell\mathbf{v}_2$. If such an ℓ exists, then $\mathbf{v}_1\mathcal{P}^{-1} = T^\ell\mathbf{v}_2\mathcal{P}^{-1}$, i.e.,

$$(T^{\ell_1}\mathbf{a}_1, T^{\ell_2}\mathbf{a}_2, \dots, T^{\ell_k}\mathbf{a}_k) = (T^\ell\mathbf{b}_1, T^\ell\mathbf{b}_2, \dots, T^\ell\mathbf{b}_k).$$

Hence, ℓ exists if and only if there exists a solution to the system of congruences:

$$\begin{cases} \ell \equiv \ell_1 \pmod{e_1} \\ \ell \equiv \ell_2 \pmod{e_2} \\ \dots \\ \ell \equiv \ell_k \pmod{e_k} \end{cases} \quad (20)$$

We know from Generalized Chinese Remainder Theorem (CRT) [4, Thm. 2.4.2] that (20) has a solution if and only if the following equations hold simultaneously:

$$\gcd(e_i, e_j) \text{ divides } \ell_i - \ell_j \text{ for all } 1 \leq i \neq j \leq k.$$

□

Our new representation allows for a faster check on whether a state \mathbf{v} is in some $[\mathbf{s}]$ by inputting \mathbf{v} and any state \mathbf{v}' belonging to $[\mathbf{s}]$ as \mathbf{v}_1 and \mathbf{v}_2 in Algorithm 1. Recall that the number of states in $[\mathbf{s}]$ is the period of \mathbf{s} . Instead of comparing \mathbf{v} with all possible states in $[\mathbf{s}]$, the representation transforms the problem into performing the verification in k ‘‘component cycles’’. Algorithm 1 requires at most $\sum_{i=1}^k e_i$ steps while an exhaustive search takes $\text{lcm}\{e_1, \dots, e_k\}$ steps to complete.

Example 5 The input consists of \mathcal{P} in (19), $\mathbf{v}_1 = (2, 2, 0, 1, 1, 2, 2, 1, 2, 2, 2, 2, 0, 2, 1, 0)$, and $\mathbf{v}_2 = (0, 1, 0, 1, 0, 2, 0, 1, 0, 1, 1, 0, 0, 1, 1, 2)$. Hence, $\mathbf{a}_1 = (1, 2, 0, 1)$, $\mathbf{a}_2 = (0, 1, 2, 2, 0, 1)$, and $\mathbf{a}_3 = (1, 2, 1, 1, 2, 2)$ while $\mathbf{b}_1 = (0, 1, 0, 2)$, $\mathbf{b}_2 = (1, 1, 0, 2, 0, 0)$, and $\mathbf{b}_3 = (2, 2, 0, 0, 0, 1)$. The sequence generated by the LFSR with characteristic polynomial $g_1^2(x)$ on input \mathbf{a}_1 never contains \mathbf{b}_1 as a state. Hence, \mathbf{v}_1 and \mathbf{v}_2 are never in a common cycle.

Keep the same \mathcal{P} and \mathbf{v}_1 but with $\mathbf{v}_2 = (1, 0, 2, 0, 2, 2, 0, 1, 1, 2, 2, 1, 2, 2, 2, 2)$. Hence, $\mathbf{b}_1 = (0, 2, 1, 1)$, $\mathbf{b}_2 = (1, 0, 0, 0, 0, 1)$, and $\mathbf{b}_3 = (0, 1, 1, 2, 1, 2)$. The sequence generated by the LFSR with characteristic polynomial $g_1^2(x)$ on input \mathbf{a}_1 is $(1, 2, 0, 1, 2, 2, 2, 1, 0, 2, 1, 1)$, making $T^8\mathbf{a}_1 = \mathbf{b}_1$. Similarly, one obtains $T^{35}\mathbf{a}_2 = \mathbf{b}_2$ and $T^{35}\mathbf{a}_3 = \mathbf{b}_3$. Thus, $\ell_1 = 8$ and $\ell_2 = \ell_3 = 35$. Since $e_1 = 12$ and $e_2 = e_3 = 39$, it is immediate to confirm that $\gcd(39, 12) = 3$ divides $\ell_3 - \ell_1 = \ell_2 - \ell_1 = 27$ and $\gcd(e_3, e_2) = 39$ divides $\ell_3 - \ell_2 = 0$, ensuring \mathbf{v}_1 and \mathbf{v}_2 belong to the same cycle. □

6 Conclusions

Theorem 1 presents the cycle structure of LFSRs with arbitrary characteristic polynomial $f(x) \in \mathbb{F}_q[x]$. This had not been previously done in the literature. We put forward a method to find a state of each cycle in $\Omega(f(x))$ by devising a new representation of the states to expedite the verification process. Finding conjugate pairs shared by two arbitrary cycles, which is crucial in the cycle joining method, can then be done more efficiently.

Acknowledgements The work of Z. Chang is supported by the Joint Fund of the National Natural Science Foundation of China under Grant U1304604, Research Grants TL-9014101684-01 and MOE2013-T2-1-041 support the research carried out by M. F. Ezerman, S. Ling, and H. Wang.

References

1. G. K. Aguirre, M. G. Mattar, and L. Magis-Weinberg, "De Bruijn cycles for neural decoding," *NeuroImage*, vol. 56, no. 3, pp. 1293–1300, 2011.
2. B. Berger and Y. Orenstein, "Efficient design of compact unstructured RNA libraries covering all k -mers," in *Proc. WABI 2015*, pp. 308–325, Atlanta, USA, Sept. 10-12, 2015, Springer Berlin Heidelberg.
3. P. Compeau, P. Pevzner, and G. Tesler, "How to apply de Bruijn graphs to genome assembly," *Nature Biotechnology*, vol. 29, no. 11, pp. 987–991, 2011.
4. C. Ding, D. Pei, and A. Salomaa, *Chinese Remainder Theorem: Applications in Computing, Coding, Cryptography*. River Edge, NJ, USA: World Scientific Publishing Co., Inc., 1996.
5. H. Fredricksen, "A survey of full length nonlinear shift register cycle algorithms," *SIAM Review*, vol. 24, no. 2, pp. 195–221, 1982.
6. S. W. Golomb, *Shift Register Sequences*. Laguna Hills: Aegean Park Press, 1981.
7. S. W. Golomb and G. Gong, *Signal Design for Good Correlation: for Wireless Communication, Cryptography, and Radar*. New York: Cambridge Univ. Press, 2004.
8. E. R. Hauge and T. Helleseth, "De Bruijn sequences, irreducible codes and cyclotomy," *Discrete Math.*, vol. 159, no. 1-3, pp. 143–154, 1996.
9. A. Lempel, "On a homomorphism of the de Bruijn graph and its applications to the design of feedback shift registers," *IEEE Trans. on Computers*, vol. C-19, no. 12, pp. 1204–1209, Dec. 1970.
10. R. Lidl and H. Niederreiter, *Finite Fields*, ser. Encyclopaedia of Mathematics and Its Applications vol. 20. New York: Cambridge Univ. Press, 1997.
11. T. Storer, *Cyclotomy and Difference Sets*, ser. Lectures in Advanced Mathematics. Chicago: Markham Pub. Co., 1967.