# An Optimal Control Problem for the Steady Nonhomogeneous Asymmetric Fluids

Exequiel Mallea-Zepeda<sup>1</sup>, Elva Ortega-Torres<sup>2</sup>, Élder J. Villamizar-Roa<sup>3</sup>

 $1$ Departamento de Matemática, Universidad de Tarapacá, Arica, Chile

 $2$ Departamento de Matemáticas, Universidad Católica del Norte, Antofagasta, Chile

 $3$ Escuela de Matemáticas, Universidad Industrial de Santander, Bucaramanga, Colombia

#### Abstract

We study an optimal boundary control problem for the two-dimensional stationary micropolar fluids system with variable density. We control the system by considering boundary controls, for the velocity vector and angular velocity of rotation of particles, on parts of the boundary of the flow domain. On the remaining part of the boundary, we consider mixed boundary conditions for the vector velocity (Dirichlet and Navier conditions) and Dirichlet boundary conditions for the angular velocity. We analyze the existence of a weak solution obtaining the fluid density as a scalar function of the stream function. We prove the existence of an optimal solution and, by using the Lagrange multipliers theorem, we state first-order optimality conditions. We also derive, through a penalty method, some optimality conditions satisfied by the optimal controls.

Keywords: Micropolar fluids system, variable density, boundary control problems. AMS Subject Classifications (2010): 49J20, 76D55, 76D05, 35Q30.

## 1 Introduction

Incompressible fluids with variable density (non homogeneous Navier-Stokes equation) correspond to a coupling between the equation for the velocity given by the conservation of momentum, the transport equation for the density provided by the mass conservation law, and the incompressibility condition. This kind of fluids are relevant to be analyzed from the mathematical and physical point of view. They can be used to model, among others, stratified fluids [\[26\]](#page-23-0), meeting of fluids coming from various regions with different densities, like the junction of pipes filled with incompressible fluids with different densities or the junction of two or more rivers [\[27\]](#page-23-1). There exists a considerable number of papers devoted to the mathematical analysis of the non homogeneous Navier-Stokes equations, principally in the non stationary case, including results when the initial density is assumed to be positive or when the initial-vacuum is allowed (see [\[13,](#page-22-0) [21,](#page-23-2) [28\]](#page-23-3) and references therein); however, not much is known about the stationary case including optimal control problems, where the state equations are given by the equations describing the motion of a viscous incompressible fluid with variable density.

An important model which generalizes the non homogeneous Navier-Stokes equation is given by the non homogeneous micropolar fluids. Non homogenous micropolar fluids refer to the micropolar fluid model with variable density; meanwhile, micropolar fluids are fluids with microstructure and asymmetric stress tensor. Physically, they represent fluids consisting of randomly oriented (or spherical) particles suspended in a viscous medium, when the deformation of fluid particles is ignored [\[10\]](#page-22-1). This model, in stationary state,

<sup>1</sup> E-mail:emallea@uta.cl

 ${}^{2}$ E-mail: eortega@ucn.cl

<sup>3</sup>E-mail: jvillami@uis.edu.co

is given by the following system of partial differential equations which expresses the balance of momentum, mass, and moment of momentum (cf. [\[11,](#page-22-2) [22\]](#page-23-4)):

<span id="page-1-0"></span>
$$
\begin{cases}\n-(\mu + \mu_r)\Delta \mathbf{u} + \rho(\mathbf{u} \cdot \nabla)\mathbf{u} + \nabla p = 2\mu_r \text{rot } \mathbf{w} + \rho \mathbf{f} \text{ in } \Omega, \\
-(c_a + c_d)\Delta \mathbf{w} - (c_0 + c_d - c_a)\nabla \text{div } \mathbf{w} + \rho(\mathbf{u} \cdot \nabla)\mathbf{w} + 4\mu_r \mathbf{w} = 2\mu_r \text{rot } \mathbf{u} + \rho \mathbf{g} \text{ in } \Omega, \\
\mathbf{u} \cdot \nabla \rho = 0 \text{ in } \Omega, \\
\text{div } \mathbf{u} = 0 \text{ in } \Omega,\n\end{cases}
$$
\n(1)

where  $\Omega$  is a connected bounded domain of  $\mathbb{R}^3$  with Lipschitz boundary,  $\mathbf{u} = (u_1, u_2, u_3)$  is the velocity field,  $\rho$  denotes the density, p represents the pressure, and w is the microrotation field interpreted as the angular velocity field of rotation of particles. The fields  $\mathbf{f} = (f_1, f_2, f_3)$  and  $\mathbf{g} = (g_1, g_2, g_3)$  represent external sources of linear and angular momentum respectively. The positive constants  $\mu, \mu_r, c_a, c_d, c_0$  characterize isotropic properties of the fluid; in particular,  $\mu$  denotes the dynamic viscosity, and  $\mu_r$ ,  $c_a$ ,  $c_d$ ,  $c_0$  are new viscosities connected with physical characteristics of the fluid. These constants satisfy  $c_0 + c_d > c_a$ . For simplicity we denote  $\mu_1 = \mu + \mu_r$ ,  $\mu_2 = c_a + c_d$  and  $\mu_3 = c_0 + c_d - c_a$ . When the microrotation viscous effects are neglected, that is  $\mu_r = 0$ , or the microrotation velocity is null, the micropolar fluid model reduces to the classical incompressible Navier-Stokes system.

From the mathematical point of view, the micropolar fluid system, with constant density, has been studied by several authors, and important results on well-posedness, large time asymptotic behavior and general qualitative analysis, have been obtained (see, for instance, [\[12,](#page-22-3) [22,](#page-23-4) [33,](#page-23-5) [34\]](#page-23-6) and references therein). However, as far as we know, the variable density stationary model [\(1\)](#page-1-0) has only been previously considered in [\[35\]](#page-23-7), where, by using the Galerkin method, the author proved the existence of weak solutions for the system [\(2\)](#page-1-1)-[\(6\)](#page-2-0) with Dirichlet boundary conditions. The main difficulty of studying model [\(1\)](#page-1-0) is due to the first-order equation  $\mathbf{u} \cdot \nabla \rho = 0$  in  $\Omega$  with  $\rho = \rho_q$  on  $\Gamma_0 \subseteq \partial \Omega$ . Even in the particular case when  $\mathbf{w} = \mathbf{0}$ , there are fewer results available in the literature related to the existence of solutions for [\(1\)](#page-1-0), and they depend on the dimension of the domain  $\Omega$  (see [\[1,](#page-22-4) [2,](#page-22-5) [3,](#page-22-6) [4,](#page-22-7) [14,](#page-22-8) [15,](#page-23-8) [27\]](#page-23-1)). In particular, in [\[14\]](#page-22-8) the author proved the existence of a solution  $[\mathbf{u}, \rho]$  for system [\(1\)](#page-1-0) in the class  $\mathbf{H}^2(\Omega) \times C^{\theta}(\Gamma_0)$  provided that  $\mathbf{f} \in \mathbf{L}^2(\Omega)$ ,  $\mathbf{u}_h \in \mathbf{C}^2(\Gamma)$ , and  $\rho_g \in C^{\theta}(\Gamma_0)$  for  $\theta > 0$ , with  $\mathbf{u}_h$  and  $\rho_g$  being the boundary data for the velocity and density respectively. This result was improved in [\[27\]](#page-23-1) where the existence of a weak solution with boundary values for the density prescribed in  $L^{\infty}$  was obtained. Still in the case 2D, but in unbounded domains, some results related to the Leray problem have been obtained in [\[1,](#page-22-4) [2,](#page-22-5) [3,](#page-22-6) [4\]](#page-22-7). The existence of solutions in the case 3D seems to be more difficult to handle and, differently to the non stationary case, we only know the paper [\[15\]](#page-23-8).

In this paper, we confine ourselves to two-dimensional flows in a bounded domain with boundary Γ of class  $C<sup>2</sup>$ . Such a flow can be interpreted as being a cross section of the three-dimensional domain  $\Omega$  by making  $x_3 = c$ , where c is a constant. In this case, it is assumed that the velocity component  $u_3$  in the  $x_3$ –direction is zero, and the axes of rotation of particles are parallel to the  $x_3$ −axis. Then, for  $\mathbf{x} = [x_1, x_2] \in \Omega \subset \mathbb{R}^2$ , the fields  $\mathbf{u}, \mathbf{w}, \rho$  and p reduce to  $\mathbf{u} = [u_1(\mathbf{x}), u_2(\mathbf{x}), 0], \mathbf{w} = [0, 0, w_3(\mathbf{x})], \rho = \rho(\mathbf{x})$ , and  $p = p(\mathbf{x})$ . Also, the external sources can be written as  $f = [f_1(\mathbf{x}), f_2(\mathbf{x}), 0]$  and  $g = [0, 0, g_3(\mathbf{x})]$ . Consequently, from now on we assume the following notations:  $\mathbf{u} = [u_1(\mathbf{x}), u_2(\mathbf{x})], \mathbf{w} = w_3(\mathbf{x}), \rho = \rho(\mathbf{x}), p = p(\mathbf{x}), \mathbf{f} = [f_1(\mathbf{x}), f_2(\mathbf{x})],$  and  $g = g_3(x)$ . Then, by observing that

$$
\text{rot } \mathbf{u} = \frac{\partial u_2}{\partial x_1} - \frac{\partial u_1}{\partial x_2}, \quad \text{rot } \mathbf{w} = [\frac{\partial \mathbf{w}}{\partial x_2}, -\frac{\partial \mathbf{w}}{\partial x_1}], \quad \text{div } \mathbf{w} = 0,
$$

and considering **u**, w,  $\rho$ ,  $p$ ,  $f$ , and g in the system [\(1\)](#page-1-0), we obtain the following two-dimensional system

<span id="page-1-1"></span>
$$
\begin{cases}\n-\mu_1 \Delta \mathbf{u} + \rho (\mathbf{u} \cdot \nabla) \mathbf{u} + \nabla p = 2\mu_r \text{rot } \mathbf{w} + \rho \mathbf{f} \text{ in } \Omega, \\
-\mu_2 \Delta \mathbf{w} + \rho (\mathbf{u} \cdot \nabla) \mathbf{w} + 4\mu_r \mathbf{w} = 2\mu_r \text{rot } \mathbf{u} + \rho \mathbf{g} \text{ in } \Omega, \\
\mathbf{u} \cdot \nabla \rho = 0 \text{ in } \Omega, \\
\text{div } \mathbf{u} = 0 \text{ in } \Omega.\n\end{cases}
$$
\n(2)

In this paper, we prove the existence of weak solutions for [\(2\)](#page-1-1) and then, we study an optimal boundary control problem where the state equations are given by the weak solutions of [\(2\)](#page-1-1). For this purpuse we consider the following boundary conditions:

<span id="page-2-1"></span>
$$
\mathbf{u} = \mathbf{u}_{g_1} := \begin{cases} \mathbf{u}_0 & \text{on} \quad \Gamma_0, \\ \mathbf{g}_1 & \text{on} \quad \Gamma_1, \end{cases} \quad [D(\mathbf{u})\mathbf{n} + \alpha \mathbf{u}]_{\text{tang}} = 0 \text{ on } \Gamma_2, \quad \mathbf{u} \cdot \mathbf{n} = 0 \text{ on } \Gamma_2,
$$
 (3)

<span id="page-2-2"></span>
$$
\int_{\Gamma} \mathbf{u} \cdot \mathbf{n} d\Gamma = 0, \quad \rho = \rho_0 > 0 \text{ on } \Gamma_0, \quad w_{g_2} = \begin{cases} w_0 & \text{on } \Gamma_0, \\ g_2 & \text{on } \Gamma_3. \end{cases}
$$
(4)

Here the boundary  $\Gamma$  of  $\Omega$  is of class  $C^2$  and  $\Gamma = \Gamma_0 \cup \Gamma_1 \cup \Gamma_2 = \Gamma_0 \cup \Gamma_3$ , where  $\Gamma_0 \cap \Gamma_i = \emptyset$ ,  $i = 1, 3$ ,  $\overline{\Gamma}_1 \cap \overline{\Gamma}_2 = \overline{\Gamma}_0 \cap \overline{\Gamma}_2 = \emptyset$ . We assume that

<span id="page-2-3"></span> $Γ_0$  is an arcwise connected closed set on Γ, with measure(Γ<sub>0</sub>) > 0. (5)

The parts  $\Gamma_1$  and  $\Gamma_3$  have nonempty interior, but  $\Gamma_2$  may be an empty set. The functions  $\rho_0$ ,  $\mathbf{u}_0$  and  $\mathbf{w}_0$  are defined on  $\Gamma_0$ , and the functions  $g_1, g_2$  describe the Dirichlet boundary control for **u** on  $\Gamma_1$  and for w on  $\Gamma_3$  respectively. The controls  $g_1, g_2$  lie in closed convex sets  $\mathcal{U}_1 \subset \mathbf{H}^{1/2}(\Gamma_1)$  and  $\mathcal{U}_2 \subset H^{1/2}(\Gamma_3)$  respectively. We assume that

<span id="page-2-0"></span> $\mathbf{u}_0 \cdot \mathbf{n} > 0$  on  $\Gamma_0$  (outflow) or  $\mathbf{u}_0 \cdot \mathbf{n} < 0$  on  $\Gamma_0$  (inflow), (6)

where n denotes the outward normal vector on Γ.



Fig. 1 Sketch of the domain  $\Omega$ 

The condition  $[D(\mathbf{u})\mathbf{n} + \alpha \mathbf{u}]_{\text{tang}} = \mathbf{0}, \mathbf{u} \cdot \mathbf{n} = 0$  on  $\Gamma_2$ , is a Navier friction boundary condition. The term  $[D(\mathbf{u})\mathbf{n} + \alpha \mathbf{u}]_{\text{tang}} := D(\mathbf{u})\mathbf{n} + \alpha \mathbf{u} - [(\underline{D}(\mathbf{u})\mathbf{n} + \alpha \mathbf{u}) \cdot \mathbf{n}] \mathbf{n}$  represents the tangential component of the vector  $D(\mathbf{u})\mathbf{n} + \alpha \mathbf{u}$ , where  $D(\mathbf{u}) := \frac{1}{2}(\nabla \mathbf{u} + \nabla^T \mathbf{u})$  is the deformation tensor, and  $\alpha \geq 0$  is the friction coefficient which measures the tendency of the fluid to slip on  $\Gamma_2$ . The Navier boundary condition was proposed by Navier [\[25\]](#page-23-9), who claimed that the tangential component of the viscous stress at the boundary should be proportional to the tangential velocity. Navier boundary condition was also derived by Maxwell [\[24\]](#page-23-10) from the kinetic theory of gases and rigorously justified as a homogenization of the no-slip condition on a rough boundary [\[19\]](#page-23-11).

We consider an objective functional given by a sum of functionals which measure, in the Lebesgue norm, the difference between the velocity vector (respectively, the density and the microrotation velocity) and a given prescribed velocity (respectively, a prescribed density and a microrotation velocity). The objective functional also measures the turbulence in the flow through a norm of the vorticity; it permits to describe the resistance in the fluid due the viscous friction (see the cost functional in [\(20\)](#page-6-0)). The state equations are given by a weak formulation of the stationary micropolar fluids equations [\(2\)](#page-1-1) with boundary conditions [\(3\)](#page-2-1)-[\(4\)](#page-2-2). The exact mathematical formulation will be given in Section 2.2 (see Definition [1\)](#page-5-0). The novelty of this paper lies in the following two aspects:

1. First, we prove the existence of a weak solution for the stationary micropolar fluids equations [\(2\)](#page-1-1) with boundary conditions [\(3\)](#page-2-1)-[\(4\)](#page-2-2). The definition of weak solution is given in Definition [1](#page-5-0) in Subsection

[2.2](#page-4-0) and he existence of weak solutions is given in Theorems [1](#page-11-0) and [2](#page-11-1) in Section [3.](#page-7-0) We look for weak solutions with  $\rho$  in the form  $\rho = \eta(\psi) := \eta \circ \psi$ , for a continuous and positive function  $\eta : \mathbb{R} \to \mathbb{R}$  and  $\psi$ the stream function associated to the velocity vector, being this the reason why we consider  $\Omega \subset \mathbb{R}^2$ .

2. Second, we prove the solvability of the optimal control problem. The existence of an optimal solution is given in Theorem [3](#page-12-0) in Section [4.](#page-12-1) Posteriorly, by using the theorem of Lagrange multipliers, we state first-order optimality conditions. The first optimality conditions are obtained in Theorem [6](#page-18-0) in Section [5.](#page-13-0) We also derive an optimality system in Remarks [4](#page-21-0) and [5.](#page-21-1)

In order to obtain the first-order optimality conditions, we will use a penalty method. This is a non standard technique which has been used previously in [\[7,](#page-22-9) [18,](#page-23-12) [20\]](#page-23-13) to derive optimality conditions for optimal control problems where the relation control-state is multivalued. To carry out this procedure, we introduce a family of penalyzed problems which approximates the initial control problem (see Theorem [4\)](#page-14-0); then, we analyze their optimality conditions (see Theorem [5\)](#page-16-0), and finally, we pass to the limit in the parameter of penalization in order to derive the optimality conditions of the original problem.

As far as we know, unlike the Navier-Stokes case, few works on optimal control problems for micropolar fluids (with constant density) are available in the literature [\[23,](#page-23-14) [29,](#page-23-15) [30,](#page-23-16) [31\]](#page-23-17). In [\[29\]](#page-23-15), a control problem for non stationary fluids in a two dimensional domains was analyzed; in that paper, a viscosity coefficient  $\lambda$ , which achieves a desired field of the microrotation velocity, is determined. In [\[30\]](#page-23-16), the author studied an optimal control problem associated with the motion of a micropolar fluid, with applications in the control of the blood pressure. In [\[31\]](#page-23-17), the author analyzed, in a two-dimensional domain, the relation between the microrotation and the vorticity of the fluid. Recently, in [\[23\]](#page-23-14), was considered an optimal boundary control problem for micropolar fluids (with constant density) equations in 3D bounded domains. Thus, the results of this paper can be seen as a 2D version of the results of [\[23\]](#page-23-14) in the case of micropolar fluids with variable density.

The outline of this paper is as follows: In Section 2, we establish the definition of weak solution and the optimal control problem to be considered. In Section 3, we prove the existence of weak solutions. In Section 4, we prove the existence of an optimal solutions. In Section 5, we derive first-order optimality conditions and, by using the Lagrange multipliers theorem, we derive an optimality system.

## <span id="page-3-0"></span>2 Statement of the Problem

### 2.1 Function Spaces

Throughout this paper we will use the Sobolev space  $H^1(\Omega)$ , and  $L^p(\Omega)$ ,  $1 \leq p \leq \infty$ , with norms  $\|\cdot\|_{H^1}$  and  $\|\cdot\|_p$  respectively. In particular, the norm and inner product in  $L^2(\Omega)$  will be represented by  $\|\cdot\|$  and  $(\cdot, \cdot)$ respectively. The norm  $L^p(\Gamma)$  will be denoted by  $\|\cdot\|_{L^p(\Gamma)}$ . Corresponding Sobolev spaces of vector valued functions will be denoted by  $\mathbf{H}^1(\Omega)$ ,  $\mathbf{L}^2(\Omega)$ , and so on. We will use the Hilbert space  $H_0^1(\Omega) = \{u \in H^1(\Omega) :$  $u = 0$  on Γ}, with the inner product  $(u, v)_{H_0^1} = (\nabla u, \nabla v)$  and the norm  $||u||_{H_0^1} = ||\nabla u||$ . We also consider the following solenoidal Banach spaces  $H_{\sigma} = \{u \in H^1(\Omega) : \text{div } u = 0 \text{ and } u \cdot n = 0 \text{ on } \Gamma_2\}$ , endowed with the usual norm of  $\mathbf{H}^1(\Omega)$ , and the space  $\tilde{\mathbf{H}}_{\sigma} = {\mathbf{u} \in \mathbf{H}_{\sigma} : \mathbf{u} = \mathbf{0} \text{ on } \Gamma \setminus \Gamma_2},$  which is a Hilbert space with the inner product  $(\mathbf{u}, \mathbf{v})_{\tilde{\mathbf{H}}_{\sigma}} = (D(\mathbf{u}), D(\mathbf{v}))$  and the norm  $\|\mathbf{u}\|_{\tilde{\mathbf{H}}_{\sigma}} := \|D(\mathbf{u})\|$ .

If X is a general Banach space, its topological dual will be denoted by  $X'$  and the duality product by  $\langle \cdot, \cdot \rangle_{X'}$  or simply by  $\langle \cdot, \cdot \rangle$ , unless this leads to ambiguity. The space  $H^{-1}$  denotes the dual of  $H_0^1(\Omega)$ ; the space  $\mathbf{H}'$  denotes the dual of  $\mathbf{H}_{\sigma}$ , and the space  $\tilde{\mathbf{H}}'$  denotes the dual of  $\tilde{\mathbf{H}}_{\sigma}$ .

If  $\Gamma_k$  is a connected subset of  $\Gamma$ , we consider the trace space  $\mathbf{H}^{1/2}(\Gamma_k) = {\mathbf{u}_{\mid \Gamma_k} : \mathbf{u} \in \mathbf{H}^1(\Omega)}$  (the restriction of the elements of  $\mathbf{H}^1(\Omega)$  to  $\Gamma_k$ ) and

$$
\mathbf{H}_{00}^{1/2}(\Gamma_k) = \{ \mathbf{v} \in \mathbf{L}^2(\Gamma_k) : \text{ there exists } \hat{\mathbf{v}} \in \mathbf{H}^{1/2}(\Gamma), \ \hat{\mathbf{v}}_{|\Gamma \backslash \Gamma_k} = \mathbf{0}, \ \hat{\mathbf{v}}_{|\Gamma_k} = \mathbf{v} \}.
$$

In the case of scalar functions, we also use the space  $H_{00}^{1/2}(\Gamma_k)$  which is defined similarly. It can be verified that  $\mathbf{H}_{00}^{1/2}(\Gamma_k)$  is a closed subspace of  $\mathbf{H}^{1/2}(\Gamma_k)$ ; moreover,  $\mathbf{H}_0^1(\Gamma_k) \hookrightarrow \mathbf{H}_{00}^{1/2}(\Gamma_k) \hookrightarrow \mathbf{L}^2(\Gamma_k)$  (cf. [\[8,](#page-22-10) 9]). For

the space  $\mathbf{H}^{1/2}(\Gamma_k)$ ,  $\mathbf{H}^{-1/2}(\Gamma_k)$  denotes its dual and  $\langle \cdot, \cdot \rangle_{\Gamma_k}$  represents its duality product. The letter C will denote diverse positive constants which may change from line to line or even within a same line.

For each function  $\mathbf{u} \in \mathbf{H}_{\sigma}$  there exists a scalar function  $\psi \in H^2(\Omega)$  (stream-function) such that

<span id="page-4-1"></span>
$$
\mathbf{u} = \operatorname{rot} \psi = \left[\frac{\partial \psi}{\partial x_2}, -\frac{\partial \psi}{\partial x_1}\right] \text{ in } \Omega. \tag{7}
$$

Let  $N: \mathbf{H}_{\sigma} \to H^2(\Omega)$  the linear operator assigning to each vector field  $\mathbf{u} \in \mathbf{H}_{\sigma}$  its stream-function  $\psi = N\mathbf{u}$ satisfying [\(7\)](#page-4-1). The assumption rot  $\psi = \mathbf{u}_0$  on  $\Gamma_0$  implies that

$$
\frac{\partial \psi}{\partial \mathbf{n}} = \mathbf{u}_0 \cdot \boldsymbol{\tau}, \quad \frac{\partial \psi}{\partial \boldsymbol{\tau}} = -(\mathbf{u}_0 \cdot \mathbf{n}) \quad \text{in} \quad \Gamma_0,
$$
\n(8)

where  $\tau$  denotes the outward tangent vector on Γ. Thus, the boundary values of  $\psi$  can be obtained by integrating with respect to the arc length, that is,

$$
\psi(\mathbf{x}) = \int_{\Gamma_0(\mathbf{x}_0, \mathbf{x})} \mathbf{u}_0 \cdot \mathbf{n} \, d\Gamma \quad \mathbf{x} \in \Gamma_0,\tag{9}
$$

where  $\mathbf{x}_0$  is the initial point of the curve  $\Gamma_0$  and  $\Gamma_0(\mathbf{x}_0, \mathbf{x})$  is the part of the curve  $\Gamma_0$  lying between the points  $\mathbf{x}_0$  and  $\mathbf{x}$  (cf. [\[14\]](#page-22-8)). Notice that since  $\psi \in H^2(\Omega)$ , then  $\psi \in H^{3/2}(\Gamma_0) \subset C^0(\Gamma_0)$ ; moreover, by [\(5\)](#page-2-3)-[\(6\)](#page-2-0),  $\psi$  is strictly monotone on  $\Gamma_0$ . Therefore, there exists  $\psi^{-1} \in C^0(\psi(\Gamma_0))$ , with  $\psi^{-1} : \psi(\Gamma_0) \subset \mathbb{R} \to \Gamma_0$ . Thus, if we assume that

$$
\rho_0 \in C^0(\Gamma_0),\tag{10}
$$

we can define the continuous function  $\tilde{\eta}(z) = \rho_0(\psi^{-1}(z)), z \in \psi(\Gamma_0) \subset \mathbb{R}$ . Since  $\rho_0(\mathbf{x}) > 0$  for all  $\mathbf{x} \in \Gamma_0$ , and  $\Gamma_0$  is an arcwise connected closed set in Γ, we can extend  $\tilde{\eta}$  to R as a strictly positive scalar function  $\eta$ such that

<span id="page-4-4"></span>
$$
\eta \in C^{0}(\mathbb{R}), \ \ \eta(y) > 0 \ \ \forall y \in \mathbb{R}, \ \eta(y) = \rho_{0}(\psi^{-1}(y)), \ \ y \in \psi(\Gamma_{0}). \tag{11}
$$

Therefore, under the above considerations, following [\[18\]](#page-23-12), we define the density  $\rho : \Omega \to \mathbb{R}$  as being

$$
\rho(\mathbf{x}) = \eta(\psi)(\mathbf{x}) = \eta(N\mathbf{u})(\mathbf{x}), \ \mathbf{x} \in \Omega.
$$

**Remark 1** *By construction of*  $\eta$  *it holds that*  $\eta(N\mathbf{u}) = \rho_0$  *on*  $\Gamma_0$ *. Moreover, for*  $\mathbf{u} \in \mathbf{H}_{\sigma}$  *and*  $\eta \in C^1(\mathbb{R})$  *it holds*

<span id="page-4-2"></span>
$$
\int_{\Omega} (\mathbf{u} \cdot \nabla \psi) \eta(N\mathbf{u}) dx = 0, \ \forall \psi \in H_0^1(\Omega).
$$
 (12)

<span id="page-4-5"></span>*If*  $\eta \in C^0(\mathbb{R})$ , *we can regularize*  $\eta$  *and the relation* [\(12\)](#page-4-2) *remains true.* 

**Remark 2** The operator  $N : H_{\sigma} \to H^2(\Omega)$  is continuous. The proof can be found in [\[18\]](#page-23-12), Lemma 2.1.

#### <span id="page-4-0"></span>2.2 Definition of Weak Solution

We consider the following operators

$$
\left\{\n \begin{array}{l}\n B: \mathbf{H}_{\sigma} \times \mathbf{H}_{\sigma} \times \mathbf{H}_{\sigma} \to \tilde{\mathbf{H}}', & F: \mathbf{H}_{\sigma} \to \tilde{\mathbf{H}}', \\
 \tilde{B}: \mathbf{H}_{\sigma} \times \mathbf{H}_{\sigma} \times H^{1}(\Omega) \to H^{-1}, & G: \mathbf{H}_{\sigma} \to H^{-1}\n \end{array}\n\right.
$$

defined by

<span id="page-4-3"></span>
$$
\begin{cases}\nB[\mathbf{u}, \mathbf{v}, \mathbf{e}] = \eta(N\mathbf{u})\mathbf{v} \cdot \nabla \mathbf{e}, & F(\mathbf{u}) = \eta(N\mathbf{u})\mathbf{f}, \\
\tilde{B}[\mathbf{u}, \mathbf{v}, \mathbf{w}] = \eta(N\mathbf{u})\mathbf{v} \cdot \nabla \mathbf{w}, & G(\mathbf{u}) = \eta(N\mathbf{u})\mathbf{g}.\n\end{cases}
$$
\n(13)

,

For  $\eta \in C^1(\mathbb{R})$  it holds that  $\text{div}(\eta(N\mathbf{u})\mathbf{u}) = \text{div}(\rho \mathbf{u}) = \mathbf{u} \cdot \nabla \rho = 0$ ; then we get

<span id="page-5-1"></span>
$$
\langle B[\mathbf{u}, \mathbf{u}, \mathbf{v}], \mathbf{v} \rangle = 0 \quad \forall \mathbf{u} \in \mathbf{H}_{\sigma}, \mathbf{v} \in \tilde{\mathbf{H}}_{\sigma}, \qquad \langle \tilde{B}[\mathbf{u}, \mathbf{u}, \mathbf{w}], \mathbf{w} \rangle = 0 \quad \forall \mathbf{u} \in \mathbf{H}_{\sigma}, \mathbf{w} \in H_0^1(\Omega). \tag{14}
$$

For  $\eta \in C^0(\mathbb{R})$ , by regularizing the function  $\eta$ , the properties [\(14\)](#page-5-1) remain true. Taking into account the operators defined in  $(13)$ , the problem  $(2)-(4)$  $(2)-(4)$  can be written as

$$
\begin{cases}\n-\mu_1 \Delta \mathbf{u} + B[\mathbf{u}, \mathbf{u}, \mathbf{u}] + \nabla p = 2\mu_r \text{rot } \mathbf{w} + F(\mathbf{u}) \text{ in } \Omega, \\
-\mu_2 \Delta \mathbf{w} + \tilde{B}[\mathbf{u}, \mathbf{u}, \mathbf{w}] + 4\mu_r \mathbf{w} = 2\mu_r \text{rot } \mathbf{u} + G(\mathbf{u}) \text{ in } \Omega, \\
\eta(N\mathbf{u}) = \rho_0 \text{ on } \Gamma_0, \\
\mathbf{u} = \mathbf{u}_{g_1} \text{ on } \Gamma \setminus \Gamma_2, \\
\mathbf{u} \cdot \mathbf{n} = 0 \text{ on } \Gamma_2, \\
[D(\mathbf{u})\mathbf{n} + \alpha \mathbf{u}]_{\text{tang}} = 0 \text{ on } \Gamma_2, \\
\mathbf{w} = \mathbf{w}_{g_2} \text{ on } \Gamma.\n\end{cases} (15)
$$

<span id="page-5-2"></span>**Lemma 1** *([\[17,](#page-23-18) 32]) Le*  $\mathbf{u} \in \mathbf{H}^2(\Omega)$  *be a divergent free vector fields verifying the Navier boundary condition* and  $\mathbf{v} \in \mathbf{H}^1(\Omega)$  *is a divergence free vector field tangent to the boundary. Then,* 

$$
-\int_{\Omega}\Delta \mathbf{u}\cdot\mathbf{v}dx=2\int_{\Omega}D(\mathbf{u}):D(\mathbf{v})dx-2\int_{\Gamma}[D(\mathbf{u})\mathbf{n}]_{\text{tang}}\cdot\mathbf{v}d\Gamma.
$$

<span id="page-5-0"></span>Through integration by parts and Lemma [1,](#page-5-2) we establish the following definition of weak solution for system  $(2)-(4).$  $(2)-(4).$  $(2)-(4).$  $(2)-(4).$ 

**Definition 1** Let  $f \in L^2(\Omega)$ ,  $g \in L^2(\Omega)$ ,  $u_{g_1}$  as in  $(3)-(4)$  $(3)-(4)$ ,  $u_0 \in H_{00}^{1/2}(\Gamma_0)$ ,  $g_1 \in H_{00}^{1/2}(\Gamma_1)$ ,  $w_0 \in H_{00}^{1/2}(\Gamma_0)$ ,  $g_2 \in H_{00}^{1/2}(\Gamma_3)$  and  $\eta \in C^0(\mathbb{R})$  defined in [\(11\)](#page-4-4). A weak solution of [\(2\)](#page-1-1)-[\(4\)](#page-2-2) is a pair  $[\mathbf{u}, \mathbf{w}] \in \mathbf{H}_{\sigma} \times H^1(\Omega)$  and  $\rho = \eta(\tilde{N} \mathbf{u})$  *satisfying* 

<span id="page-5-3"></span>
$$
\begin{cases}\n2\mu_1(D(\mathbf{u}), D(\mathbf{v})) + 2\alpha\mu_1 \int_{\Gamma_2} \mathbf{u} \cdot \mathbf{v} \, d\Gamma + \langle B[\mathbf{u}, \mathbf{u}, \mathbf{u}], \mathbf{v} \rangle = 2\mu_r(\text{rot } \mathbf{w}, \mathbf{v}) + (F(\mathbf{u}), \mathbf{v}) \,\forall \mathbf{v} \in \tilde{\mathbf{H}}_{\sigma}, \\
\mu_2(\nabla \mathbf{w}, \nabla z) + \langle \tilde{B}[\mathbf{u}, \mathbf{u}, \mathbf{w}], z \rangle + 4\mu_r(\mathbf{w}, z) = 2\mu_r(\text{rot } \mathbf{u}, z) + (G(\mathbf{u}), z) \,\forall z \in H_0^1(\Omega), \\
\eta(N\mathbf{u}) = \rho_0 \quad on \ \Gamma_0, \\
\mathbf{u} = \mathbf{u}_{g_1} \quad on \ \Gamma \setminus \Gamma_2, \\
\mathbf{w} = \mathbf{w}_{g_2} \quad on \ \Gamma.\n\end{cases} (16)
$$

In order to prove the existence of a solution to problem [\(16\)](#page-5-3), we reduce the problem to an auxiliary problem with homogeneous boundary conditions for **u** on  $\Gamma \setminus \Gamma_2$  and for w on  $\Gamma$ . For this purpose, we introduce the following result.

**Lemma 2** Let  $u_{g_1}$  as in [\(3\)](#page-2-1)–[\(4\)](#page-2-2), and assume that  $u_0 \in H_{00}^{1/2}(\Gamma_0)$ ,  $g_1 \in H_{00}^{1/2}(\Gamma_1)$ . Then, for any  $\varepsilon > 0$ , *there exists*  $\mathbf{u}^{\varepsilon} \in \mathbf{H}_{\sigma}$  *with*  $\mathbf{u}^{\varepsilon} = \mathbf{g}_1$  *on*  $\Gamma_1$ ,  $\mathbf{u}^{\varepsilon} = \mathbf{u}_0$  *on*  $\Gamma_0$  *and*  $\mathbf{u}^{\varepsilon} = \mathbf{0}$  *on*  $\Gamma_2$  *such that* 

<span id="page-5-5"></span><span id="page-5-4"></span>
$$
|(\mathbf{v} \cdot \nabla \mathbf{u}^{\varepsilon}, \mathbf{v})| \leq \varepsilon \|\mathbf{v}\|_{\tilde{\mathbf{H}}_{\sigma}}^2 \quad \forall \mathbf{v} \in \tilde{\mathbf{H}}_{\sigma}, \text{ and } \|\mathbf{u}^{\varepsilon}\|_{\mathbf{H}_{\sigma}} \leq C \|\mathbf{u}_{g_1}\|_{\mathbf{H}^{1/2}(\Gamma \backslash \Gamma_2)},\tag{17}
$$

where the constant C depends only on  $\Omega$ . Moreover, if  $w_0 \in H_{00}^{1/2}(\Gamma_0)$  and  $g_2 \in H_{00}^{1/2}(\Gamma_3)$ , then there exists  $w^e \in H^1(\Omega)$  such that  $w^e = w_0$  on  $\Gamma_0$ ,  $w^e = g_2$  on  $\Gamma_3$ , and the following estimate holds

<span id="page-5-6"></span>
$$
\|w^e\|_{H^1} \le C \|w_{g_2}\|_{H^{1/2}(\Gamma)},\tag{18}
$$

*where the constant*  $C$  *depends only on*  $\Omega$ *.* 

*Proof.* If  $u_0 \in H_{00}^{1/2}(\Gamma_0)$ ,  $g_1 \in H_{00}^{1/2}(\Gamma_1)$ , there exist  $\hat{g}_1 \in H^{1/2}(\Gamma)$ ,  $\hat{u}_0 \in H^{1/2}(\Gamma)$  such that  $\hat{u}_{0|_{\Gamma_0}} = u_0$ ,  $\hat{\mathbf{u}}_{0|_{\Gamma\setminus\Gamma_0}} = \mathbf{0}$ ,  $\hat{\mathbf{g}}_{1|_{\Gamma_1}} = \mathbf{0}$  and  $\hat{\mathbf{g}}_{1|_{\Gamma_1}} = \mathbf{g}_1$ . Thus,  $\hat{\mathbf{u}}_0 + \hat{\mathbf{g}}_1 \in \mathbf{H}^{1/2}(\Gamma)$ ; moreover, the integral in [\(4\)](#page-2-2) implies  $\int_{\Gamma} (\hat{\mathbf{u}}_0 + \hat{\mathbf{g}}_1) \cdot \mathbf{n} d\Gamma = 0$ . Then, by Lemma IX.4.2 of [\[16\]](#page-23-20), p. 610, there exists  $\mathbf{u}^{\varepsilon} \in \mathbf{H}_{\sigma}$  with  $\mathbf{u}^{\varepsilon} = \hat{\mathbf{u}}_0 + \hat{\mathbf{g}}_1$  on  $\Gamma$ verifying [\(17\)](#page-5-4); in particular we have that  $\mathbf{u}_{\vert_{\Gamma_1}}^{\varepsilon} = \mathbf{g}_1, \mathbf{u}_{\vert_{\Gamma_0}}^{\varepsilon} = \mathbf{u}_0$  and  $\mathbf{u}_{\vert_{\Gamma_2}}^{\varepsilon} = \mathbf{0}$ . The existence of  $w^e \in H^1(\Omega)$ is well-known from the lifting theorem.

Rewriting  $[\mathbf{u}, w] \in \mathbf{H}_{\sigma} \times H^1(\Omega)$  in the form  $\mathbf{u} = \mathbf{u}^{\varepsilon} + \hat{\mathbf{u}}$  and  $w = w^{\varepsilon} + \hat{w}$  with  $[\hat{\mathbf{u}}, \hat{w}] \in \widetilde{\mathbf{H}}_{\sigma} \times H^1_0(\Omega)$  new unknown functions, from [\(16\)](#page-5-3) we obtain the following nonlinear system: Find  $[\hat{\mathbf{u}}, \hat{\mathbf{w}}] \in \tilde{\mathbf{H}}_{\sigma} \times H_0^1(\Omega)$  such that

<span id="page-6-2"></span>
$$
\begin{cases}\n2\mu_1(D(\hat{\mathbf{u}}), D(\mathbf{v})) + 2\alpha\mu_1 \int_{\Gamma_2} \hat{\mathbf{u}} \cdot \mathbf{v} \, d\Gamma + \langle B[\hat{\mathbf{u}} + \mathbf{u}^\varepsilon, \hat{\mathbf{u}} + \mathbf{u}^\varepsilon, \hat{\mathbf{u}}], \mathbf{v} \rangle + \langle B[\hat{\mathbf{u}} + \mathbf{u}^\varepsilon, \hat{\mathbf{u}}, \mathbf{u}^\varepsilon], \mathbf{v} \rangle \\
= 2\mu_r(\text{rot}(\hat{\mathbf{w}} + \mathbf{w}^\varepsilon), \mathbf{v}) + (F(\hat{\mathbf{u}} + \mathbf{u}^\varepsilon), \mathbf{v}) - 2\mu_1(D(\mathbf{u}^\varepsilon), D(\mathbf{v})) - \langle B[\hat{\mathbf{u}} + \mathbf{u}^\varepsilon, \mathbf{u}^\varepsilon], \mathbf{v} \rangle, \\
\mu_2(\nabla \hat{\mathbf{w}}, \nabla z) + \langle \tilde{B}[\hat{\mathbf{u}} + \mathbf{u}^\varepsilon, \hat{\mathbf{u}} + \mathbf{u}^\varepsilon, \hat{\mathbf{w}}], z \rangle + \langle \tilde{B}[\hat{\mathbf{u}} + \mathbf{u}^\varepsilon, \hat{\mathbf{u}}, \mathbf{w}^\varepsilon], z \rangle + 4\mu_r(\hat{\mathbf{w}}, z) \\
= 2\mu_r(\text{rot}(\hat{\mathbf{u}} + \mathbf{u}^\varepsilon), z) + (G(\hat{\mathbf{u}} + \mathbf{u}^\varepsilon), z) - \mu_2(\nabla \mathbf{w}^\varepsilon, \nabla z) - \langle \tilde{B}[\hat{\mathbf{u}} + \mathbf{u}^\varepsilon, \mathbf{u}^\varepsilon, \mathbf{w}^\varepsilon], z \rangle - 4\mu_r(\mathbf{w}^\varepsilon, z),\n\end{cases} \tag{19}
$$

for all  $[\mathbf{v}, \mathbf{z}] \in \tilde{\mathbf{H}}_{\sigma} \times H_0^1(\Omega)$ .

<span id="page-6-3"></span> $\overline{\mathcal{L}}$ 

#### 2.3 Statement of Boundary Control Problem

In order to establish the statement of the boundary control problem, we suppose that  $U_1 \subset H_{00}^{1/2}(\Gamma_1)$  and  $\mathcal{U}_2 \subset H_{00}^{1/2}(\Gamma_3)$  are nonempty sets. We consider that  $\mathbf{f} \in \mathbf{L}^2(\Omega)$ ,  $g \in L^2(\Omega)$ ,  $\mathbf{u}_0 \in \mathbf{H}_{00}^{1/2}(\Gamma_0)$ ,  $w_0 \in H_{00}^{1/2}(\Gamma_0)$ , and the controls  $g_1 \in U_1$ ,  $g_2 \in U_2$ . For simplicity, we denote  $\mathbb{X} = \mathbf{H}_{\sigma} \times H^1(\Omega) \times U_1 \times U_2$  and consider the following objective functional  $J : \mathbb{X} \to \mathbb{R}$  defined by:

<span id="page-6-0"></span>
$$
J[\mathbf{u}, \mathbf{w}, \mathbf{g}_1, g_2] = \frac{\beta_1}{2} ||\text{rot } \mathbf{u}||^2 + \frac{\beta_2}{2} ||\mathbf{u} - \mathbf{u}_d||^2 + \frac{\beta_3}{2} ||\mathbf{w} - \mathbf{w}_d||^2 + \frac{\beta_4}{2} ||\eta(N\mathbf{u}) - \rho_d||^2
$$
  
+  $\frac{\beta_5}{2} ||\mathbf{g}_1||_{\mathbf{H}^{1/2}(\Gamma_1)}^2 + \frac{\beta_6}{2} ||g_2||_{H^{1/2}(\Gamma_3)}^2,$  (20)

where the constants  $\beta_k$  measure the cost of the control and satisfy the following conditions:

- $\sqrt{ }$  $\int$  $i) \beta_i \ge 0$  for  $i = 1, ..., 6$ , (not all zero),  $\mathcal{U}_1 \subset \mathbf{H}_{00}^{1/2}(\Gamma_1)$  and  $\mathcal{U}_2 \subset H_{00}^{1/2}(\Gamma_3)$  are bounded closed and convex sets;
	- $ii) \beta_i \ge 0$  for  $i = 1, ..., 4, \beta_i > 0$  for  $i = 5, 6, \mathcal{U}_1 \subset \mathbf{H}_{00}^{1/2}(\Gamma_1)$  and  $\mathcal{U}_2 \subset H_{00}^{1/2}(\Gamma_3)$  are closed and convex sets. (21)

In the functional [\(20\)](#page-6-0), the prescribed functions  $\mathbf{u}_d \in \mathbf{L}^2(\Omega)$ ,  $\mathbf{w}_d \in L^2(\Omega)$  and  $\rho_d \in L^2(\Omega)$ , correspond to the desired states for the velocity, the microrotation velocity and the density, respectively. Then we study the following constrained minimization problem related to system  $(2)-(4)$  $(2)-(4)$ :

<span id="page-6-1"></span> $\left\{\begin{array}{c}\text{Find } [\mathbf{u}, \mathbf{w}, \mathbf{g}_1, g_2] \in \mathbb{X} \text{ such that, the functional } J[\mathbf{u}, \mathbf{w}, \mathbf{g}_1, g_2] \text{ reaches its minimum over the weak solutions of system (2)-(4).}\end{array}\right.$  $\left\{\begin{array}{c}\text{Find } [\mathbf{u}, \mathbf{w}, \mathbf{g}_1, g_2] \in \mathbb{X} \text{ such that, the functional } J[\mathbf{u}, \mathbf{w}, \mathbf{g}_1, g_2] \text{ reaches its minimum over the weak solutions of system (2)-(4).}\end{array}\right.$  $\left\{\begin{array}{c}\text{Find } [\mathbf{u}, \mathbf{w}, \mathbf{g}_1, g_2] \in \mathbb{X} \text{ such that, the functional } J[\mathbf{u}, \mathbf{w}, \mathbf{g}_1, g_2] \text{ reaches its minimum over the weak solutions of system (2)-(4).}\end{array}\right.$  $\left\{\begin{array}{c}\text{Find } [\mathbf{u}, \mathbf{w}, \mathbf{g}_1, g_2] \in \mathbb{X} \text{ such that, the functional } J[\mathbf{u}, \mathbf{w}, \mathbf{g}_1, g_2] \text{ reaches its minimum over the weak solutions of system (2)-(4).}\end{array}\right.$  $\left\{\begin{array}{c}\text{Find } [\mathbf{u}, \mathbf{w}, \mathbf{g}_1, g_2] \in \mathbb{X} \text{ such that, the functional } J[\mathbf{u}, \mathbf{w}, \mathbf{g}_1, g_2] \text{ reaches its minimum over the weak solutions of system (2)-(4).}\end{array}\right.$  (22)

The set of admissible solutions of problem [\(22\)](#page-6-1) is defined by

 $\mathcal{S}_{ad} = \{ \mathbf{s} = [\mathbf{u}, \mathbf{w}, \boldsymbol{g}_1, g_2] \in \mathbb{X} \text{ such that } J(\mathbf{s}) < \infty \text{ and } \mathbf{s} \text{ satisfies (16)} \}.$  $\mathcal{S}_{ad} = \{ \mathbf{s} = [\mathbf{u}, \mathbf{w}, \boldsymbol{g}_1, g_2] \in \mathbb{X} \text{ such that } J(\mathbf{s}) < \infty \text{ and } \mathbf{s} \text{ satisfies (16)} \}.$  $\mathcal{S}_{ad} = \{ \mathbf{s} = [\mathbf{u}, \mathbf{w}, \boldsymbol{g}_1, g_2] \in \mathbb{X} \text{ such that } J(\mathbf{s}) < \infty \text{ and } \mathbf{s} \text{ satisfies (16)} \}.$ 

## <span id="page-7-0"></span>3 Existence of Weak Solutions

#### 3.1 Linearized Problem

For  $\tilde{\mathbf{u}} \in \tilde{\mathbf{H}}_{\sigma}$  fixed, we consider the following linear problem: Find  $\hat{\mathbf{w}} \in H_0^1(\Omega)$  such that

<span id="page-7-1"></span>
$$
\begin{cases}\n\mu_2(\nabla \hat{\mathbf{w}}, \nabla z) + \langle \tilde{B}[\tilde{\mathbf{u}} + \mathbf{u}^\varepsilon, \tilde{\mathbf{u}} + \mathbf{u}^\varepsilon, \hat{\mathbf{w}}], z \rangle + 4\mu_r(\hat{\mathbf{w}}, z) \\
= 2\mu_r(\text{rot}(\tilde{\mathbf{u}} + \mathbf{u}^\varepsilon), z) + (G(\tilde{\mathbf{u}} + \mathbf{u}^\varepsilon), z) - \mu_2(\nabla \mathbf{w}^\varepsilon, \nabla z) - 4\mu_r(\mathbf{w}^\varepsilon, z) \\
-\langle \tilde{B}[\tilde{\mathbf{u}} + \mathbf{u}^\varepsilon, \mathbf{u}^\varepsilon, \mathbf{w}^\varepsilon], z \rangle - \langle \tilde{B}[\tilde{\mathbf{u}} + \mathbf{u}^\varepsilon, \tilde{\mathbf{u}}, \mathbf{w}^\varepsilon], z \rangle \,\forall z \in H_0^1(\Omega),\n\end{cases} (23)
$$

where  $\mathbf{u}^{\varepsilon}$  and  $\mathbf{w}^{\varepsilon}$  are given by Lemma [2.](#page-5-5) For problem [\(23\)](#page-7-1) we have the following result.

**Lemma 3** If  $g \in L^2(\Omega)$ , then the problem [\(23\)](#page-7-1) has a unique solution  $\hat{w} \in H_0^1(\Omega)$ . Moreover, the following *inequality holds:*

<span id="page-7-4"></span>
$$
\mu_2 \|\hat{\mathbf{w}}\|_{H_0^1} \leq (\tilde{C}_{\eta} C \|\mathbf{w}^e\|_{H^1} + \mu_r C) \|\tilde{\mathbf{u}}\|_{\tilde{\mathbf{H}}_{\sigma}} + \tilde{C}_{\eta} C \|\mathbf{u}^e\|_{\mathbf{H}_{\sigma}} \|\mathbf{w}^e\|_{H^1} + \tilde{C}_{\eta} \|\mathbf{g}\| \n+ \mu_r C (\|\mathbf{u}^e\|_{\mathbf{H}_{\sigma}} + \|\mathbf{w}^e\|_{H^1}) + \mu_2 C \|\mathbf{w}^e\|_{H^1},
$$
\n(24)

 $where \ \tilde{C}_\eta > 0 \ \text{is a constant satisfying } ||\eta(N(\tilde{\mathbf{u}} + \mathbf{u}^{\varepsilon}))||_{\infty} \leq \tilde{C}_\eta, \ [ \mathbf{u}^{\varepsilon}, \mathbf{w}^{\varepsilon} ] \ \text{is given by Lemma 2, and } C > 0 \ \text{is}$  $where \ \tilde{C}_\eta > 0 \ \text{is a constant satisfying } ||\eta(N(\tilde{\mathbf{u}} + \mathbf{u}^{\varepsilon}))||_{\infty} \leq \tilde{C}_\eta, \ [ \mathbf{u}^{\varepsilon}, \mathbf{w}^{\varepsilon} ] \ \text{is given by Lemma 2, and } C > 0 \ \text{is}$  $where \ \tilde{C}_\eta > 0 \ \text{is a constant satisfying } ||\eta(N(\tilde{\mathbf{u}} + \mathbf{u}^{\varepsilon}))||_{\infty} \leq \tilde{C}_\eta, \ [ \mathbf{u}^{\varepsilon}, \mathbf{w}^{\varepsilon} ] \ \text{is given by Lemma 2, and } C > 0 \ \text{is}$ *a constant that depends only on* Ω*.*

*Proof.* We define the bilinear form  $a_{\tilde{\mathbf{u}}} : H_0^1(\Omega) \times H_0^1(\Omega) \to \mathbb{R}$  and the linear functional  $f_{\tilde{\mathbf{u}}} : H_0^1(\Omega) \to \mathbb{R}$  by

<span id="page-7-2"></span>
$$
a_{\tilde{\mathbf{u}}}([\hat{\mathbf{w}}, \mathbf{z}]) = \mu_2(\nabla \hat{\mathbf{w}}, \nabla \mathbf{z}) + \langle \tilde{B}[\tilde{\mathbf{u}} + \mathbf{u}^\varepsilon, \tilde{\mathbf{u}} + \mathbf{u}^\varepsilon, \hat{\mathbf{w}}], \mathbf{z} \rangle + 4\mu_r(\hat{\mathbf{w}}, \mathbf{z}) \,\forall \mathbf{z} \in H_0^1(\Omega),
$$
  
\langle f\_{\tilde{\mathbf{u}}}, \mathbf{z} \rangle = 2\mu\_r(\text{rot}(\tilde{\mathbf{u}} + \mathbf{u}^\varepsilon), \mathbf{z}) + (G(\tilde{\mathbf{u}} + \mathbf{u}^\varepsilon), \mathbf{z}) - \mu\_2(\nabla \mathbf{w}^\varepsilon, \nabla \mathbf{z}) - 4\mu\_r(\mathbf{w}^\varepsilon, \mathbf{z})\n(25)

$$
-\langle \tilde{B}[\tilde{\mathbf{u}} + \mathbf{u}^{\varepsilon}, \mathbf{u}^{\varepsilon}, \mathbf{w}^{\mathrm{e}}], \mathbf{z} \rangle - \langle \tilde{B}[\tilde{\mathbf{u}} + \mathbf{u}^{\varepsilon}, \tilde{\mathbf{u}}, \mathbf{w}^{\mathrm{e}}], \mathbf{z} \rangle \ \forall \mathbf{z} \in H_0^1(\Omega). \tag{26}
$$

Then, from [\(25\)](#page-7-2)-[\(26\)](#page-7-2), problem [\(23\)](#page-7-1) is equivalent to find  $\hat{\mathbf{w}} \in H_0^1(\Omega)$  such that

<span id="page-7-7"></span><span id="page-7-3"></span>
$$
a_{\tilde{\mathbf{u}}}([\hat{\mathbf{w}}, \mathbf{z}]) = \langle \mathbf{f}_{\tilde{\mathbf{u}}}, \mathbf{z} \rangle \ \forall \mathbf{z} \in H_0^1(\Omega). \tag{27}
$$

The bilinear form  $a_{\tilde{u}}$  is continuous and coercive on  $H_0^1(\Omega) \times H_0^1(\Omega)$ , and the linear functional  $f_{\tilde{u}}$  is continuous on  $H_0^1(\Omega)$ . Then, by Lax-Milgram Theorem, it follows that there is a unique  $\hat{w} \in H_0^1(\Omega)$  such that [\(27\)](#page-7-3) is satisfied, and therefore the problem [\(23\)](#page-7-1) has a unique solution.

Now, in order to obtain inequality [\(24\)](#page-7-4), by replacing  $z = \hat{w}$  in [\(23\)](#page-7-1) and taking into account [\(14\)](#page-5-1), we have

<span id="page-7-5"></span>
$$
\mu_2 \|\nabla \hat{\mathbf{w}}\|^2 + 4\mu_r \|\hat{\mathbf{w}}\|^2 \leq 2\mu_r |(\text{rot}(\tilde{\mathbf{u}} + \mathbf{u}^{\varepsilon}), \hat{\mathbf{w}})| + |(G(\hat{\mathbf{u}} + \mathbf{u}^{\varepsilon}), \hat{\mathbf{w}})| + \mu_2 |(\nabla \mathbf{w}^{\mathrm{e}}, \nabla \hat{\mathbf{w}})| \n+ 4\mu_r |(\mathbf{w}^{\mathrm{e}}, \hat{\mathbf{w}})| + |\langle \tilde{B}[\tilde{\mathbf{u}} + \mathbf{u}^{\varepsilon}, \mathbf{u}^{\varepsilon}, \mathbf{w}^{\mathrm{e}}], \hat{\mathbf{w}}\rangle| \n+ |\langle \tilde{B}[\tilde{\mathbf{u}} + \mathbf{u}^{\varepsilon}, \tilde{\mathbf{u}}, \mathbf{w}^{\mathrm{e}}], \hat{\mathbf{w}}\rangle|.
$$
\n(28)

Now, we find estimates for the terms on the right hand side of  $(28)$ . By applying the Hölder and Poincaré inequalities we get

<span id="page-7-6"></span>
$$
2\mu_r |(\text{rot}(\tilde{\mathbf{u}}+\mathbf{u}^{\varepsilon}), \hat{\mathbf{w}})| \leq 2\sqrt{2}\mu_r C \|\nabla(\tilde{\mathbf{u}}+\mathbf{u}^{\varepsilon})\| \|\hat{\mathbf{w}}\|_{H_0^1} \leq \mu_r C (\|\tilde{\mathbf{u}}\|_{\tilde{\mathbf{H}}_{\sigma}} + \|\mathbf{u}^{\varepsilon}\|_{\mathbf{H}_{\sigma}}) \|\hat{\mathbf{w}}\|_{H_0^1},
$$
(29)

$$
|(G(\tilde{\mathbf{u}} + \mathbf{u}^{\varepsilon}), \hat{\mathbf{w}})| \leq \|\eta(N(\tilde{\mathbf{u}} + \mathbf{u}^{\varepsilon}))\|_{\infty} \|g\| \|\hat{\mathbf{w}}\|_{H_0^1} \leq \tilde{C}_\eta \|g\| \|\hat{\mathbf{w}}\|_{H_0^1},
$$
\n(30)

$$
\mu_2 |(\nabla w^e, \nabla \hat{w})| \leq \mu_2 \|\nabla w^e\| \|\nabla \hat{w}\| \leq \mu_2 \|w^e\|_{H^1} \|\hat{w}\|_{H_0^1},\tag{31}
$$

$$
4\mu_r |(\mathbf{w}^e, \hat{\mathbf{w}})| \le 4\mu_r \|\mathbf{w}^e\| \|\hat{\mathbf{w}}\| \le 4\mu_r C \|\mathbf{w}^e\|_{H^1} \|\nabla \hat{\mathbf{w}}\| \le \mu_r C \|\mathbf{w}^e\|_{H^1} \|\hat{\mathbf{w}}\|_{H_0^1},\tag{32}
$$

$$
|\langle \tilde{B}[\tilde{\mathbf{u}} + \mathbf{u}^{\varepsilon}, \mathbf{u}^{\varepsilon}, \mathbf{w}^{\varepsilon}], \hat{\mathbf{w}} \rangle| \leq \|\eta(N(\tilde{\mathbf{u}} + \mathbf{u}^{\varepsilon}))\|_{\infty} \|\mathbf{u}^{\varepsilon}\|_{3} \|\nabla \mathbf{w}^{\varepsilon}\| \|\hat{\mathbf{w}}\|_{6} \leq \tilde{C}_{\eta} C \|\nabla \mathbf{u}^{\varepsilon}\| \|\nabla \hat{\mathbf{w}}^{\varepsilon}\| \|\nabla \hat{\mathbf{w}}\|
$$
\n
$$
\leq \tilde{C}_{\eta} C \|\mathbf{u}^{\varepsilon}\|_{\mathbf{H}_{\sigma}} \|\mathbf{w}^{\varepsilon}\|_{\mathbf{H}^{1}} \|\hat{\mathbf{w}}\|_{H^{1}_{0}}, \tag{33}
$$

$$
|\langle \tilde{B}[\tilde{\mathbf{u}} + \mathbf{u}^{\varepsilon}, \tilde{\mathbf{u}}, \mathbf{w}^e], \hat{\mathbf{w}} \rangle| \leq \|\eta(N(\tilde{\mathbf{u}} + \mathbf{u}^{\varepsilon}))\|_{\infty} \|\tilde{\mathbf{u}}\|_{3} \|\nabla \mathbf{w}^e\| \|\hat{\mathbf{w}}\|_{6} \leq \tilde{C}_{\eta} C \|\nabla \tilde{\mathbf{u}}\| \|\nabla \mathbf{w}^e\| \|\nabla \hat{\mathbf{w}}\|
$$
\n
$$
\leq \tilde{C}_{\eta} C \|\tilde{\mathbf{u}}\|_{\tilde{\mathbf{H}}_{\sigma}} \|\mathbf{w}^e\|_{H^1} \|\hat{\mathbf{w}}\|_{H^1_0}.
$$
\n(34)

By substituting  $(29)-(34)$  $(29)-(34)$  in  $(28)$  we obtain

$$
\mu_2 \|\hat{\mathbf{w}}\|_{H_0^1}^2 \leq \tilde{C}_\eta C(\|\tilde{\mathbf{u}}\|_{\tilde{\mathbf{H}}_\sigma} + \|\mathbf{u}^\varepsilon\|_{\mathbf{H}_\sigma}) \|\mathbf{w}^\varepsilon\|_{H^1} \|\hat{\mathbf{w}}\|_{H_0^1} + \tilde{C}_\eta \|g\| \|\hat{\mathbf{w}}\|_{H_0^1} + \mu_r C(\|\tilde{\mathbf{u}}\|_{\tilde{\mathbf{H}}_\sigma} + \|\mathbf{u}^\varepsilon\|_{\mathbf{H}_\sigma} + \|\mathbf{w}^\varepsilon\|_{H^1}) \|\hat{\mathbf{w}}\|_{H_0^1} + \mu_2 C \|\mathbf{w}^\varepsilon\|_{H^1} \|\hat{\mathbf{w}}\|_{H_0^1},
$$

that is,

$$
\mu_2 \|\hat{\mathbf{w}}\|_{H_0^1} \leq \tilde{C}_\eta C(\|\tilde{\mathbf{u}}\|_{\tilde{\mathbf{H}}_\sigma} + \|\mathbf{u}^\varepsilon\|_{\mathbf{H}_\sigma}) \|\mathbf{w}^\varepsilon\|_{H^1} + \tilde{C}_\eta \|\mathbf{g}\| \n+ \mu_r C(\|\tilde{\mathbf{u}}\|_{\tilde{\mathbf{H}}_\sigma} + \|\mathbf{u}^\varepsilon\|_{\mathbf{H}_\sigma} + \|\mathbf{w}^\varepsilon\|_{H^1}) + \mu_2 C \|\mathbf{w}^\varepsilon\|_{H^1},
$$
\n(35)

which implies [\(24\)](#page-7-4). ⋄

#### 3.2 Weak Solutions

In order to prove the existence of solutions to the problem [\(19\)](#page-6-2), we define the linear operator  $T : \tilde{H}_{\sigma} \to \tilde{H}_{\sigma}$ as follows: For each  $\tilde{\mathbf{u}} \in \tilde{\mathbf{H}}_{\sigma}$ , let  $T\tilde{\mathbf{u}} = \hat{\mathbf{u}}$ , where  $\hat{\mathbf{u}}$  satisfies the following system

<span id="page-8-0"></span>
$$
2\mu_1(D(\hat{\mathbf{u}}), D(\mathbf{v})) + 2\alpha\mu_1 \int_{\Gamma_2} \hat{\mathbf{u}} \cdot \mathbf{v} d\Gamma + \langle B[\tilde{\mathbf{u}} + \mathbf{u}^\varepsilon, \tilde{\mathbf{u}} + \mathbf{u}^\varepsilon, \hat{\mathbf{u}}], \mathbf{v} \rangle + \langle B[\tilde{\mathbf{u}} + \mathbf{u}^\varepsilon, \hat{\mathbf{u}}, \mathbf{u}^\varepsilon], \mathbf{v} \rangle
$$
  
= 
$$
2\mu_r(\text{rot}(\hat{\mathbf{w}} + \mathbf{w}^\text{e}), \mathbf{v}) + (F(\tilde{\mathbf{u}} + \mathbf{u}^\varepsilon), \mathbf{v}) - 2\mu_1(D(\mathbf{u}^\varepsilon), D(\mathbf{v})) - \langle B[\tilde{\mathbf{u}} + \mathbf{u}^\varepsilon, \mathbf{u}^\varepsilon, \mathbf{u}^\varepsilon], \mathbf{v} \rangle,
$$
(36)

<span id="page-8-5"></span>for all  $\mathbf{v} \in \tilde{\mathbf{H}}_{\sigma}$ , and  $\hat{\mathbf{w}}$  being the unique solution of the linear problem [\(23\)](#page-7-1).

**Lemma 4** *The operator*  $T : \tilde{H}_{\sigma} \to \tilde{H}_{\sigma}$  *defined by* [\(36\)](#page-8-0) *is compact.* 

*Proof.* Let  ${\{\tilde{\mathbf{u}}^m\}}_{m \geq 1} \subset \tilde{\mathbf{H}}_{\sigma}$  a sequence weakly convergent to  $\tilde{\mathbf{u}} \in \tilde{\mathbf{H}}_{\sigma}$ . Since the embedding  $\tilde{\mathbf{H}}_{\sigma} \hookrightarrow \mathbf{L}^p(\Omega)$  is compact for  $1 \leq p < \infty$  we have

$$
\tilde{\mathbf{u}}^m \to \tilde{\mathbf{u}} \text{ strongly in } \mathbf{L}^p(\Omega).
$$

From Remark [2](#page-4-5) and since the embedding  $H^2(\Omega) \hookrightarrow C^0(\overline{\Omega})$  is compact, we get that, for some subsequence of  ${\{\tilde{\mathbf{u}}^m\}}_{m\geq1}$ , still denoted by  ${\{\tilde{\mathbf{u}}^m\}}_{m\geq1}$ , it holds that  $N\tilde{\mathbf{u}}^m \to N\tilde{\mathbf{u}}$  in  $C^0(\overline{\Omega})$ ; moreover, taking into account that  $\eta \in \overline{C}^0(\mathbb{R})$ , we have  $\eta(N\tilde{\mathbf{u}}^m) \to \eta(N\tilde{\mathbf{u}})$  strongly in  $C^0(\overline{\Omega})$ , which implies that there exists a constant  $C_n > 0$ , independent of m, such that

<span id="page-8-4"></span>
$$
\|\eta(N(\tilde{\mathbf{u}}^m + \mathbf{u}^\varepsilon))\|_{\infty} \le C_{\eta}.\tag{37}
$$

Then, denoting  $\hat{\mathbf{u}}^m = T(\tilde{\mathbf{u}}^m)$ , for all  $\mathbf{v} \in \tilde{\mathbf{H}}_{\sigma}$  we get

<span id="page-8-1"></span>
$$
2\mu_1(D(\hat{\mathbf{u}}^m), D(\mathbf{v})) + 2\alpha\mu_1 \int_{\Gamma_2} \hat{\mathbf{u}}^m \cdot \mathbf{v} \, d\Gamma + \langle B[\tilde{\mathbf{u}}^m + \mathbf{u}^\varepsilon, \tilde{\mathbf{u}}^m + \mathbf{u}^\varepsilon, \hat{\mathbf{u}}^m], \mathbf{v} \rangle + \langle B[\tilde{\mathbf{u}}^m + \mathbf{u}^\varepsilon, \hat{\mathbf{u}}^m, \mathbf{u}^\varepsilon], \mathbf{v} \rangle
$$
  
=  $2\mu_r(\text{rot}(\hat{\mathbf{w}} + \mathbf{w}^\varepsilon), \mathbf{v}) + (F(\tilde{\mathbf{u}}^m + \mathbf{u}^\varepsilon), \mathbf{v}) - 2\mu_1(D(\mathbf{u}^\varepsilon), D(\mathbf{v})) - \langle B[\tilde{\mathbf{u}}^m + \mathbf{u}^\varepsilon, \mathbf{u}^\varepsilon], \mathbf{v} \rangle.$  (38)

Taking the difference between [\(38\)](#page-8-1) and [\(36\)](#page-8-0), we have

<span id="page-8-2"></span>
$$
2\mu_1(D(\hat{\mathbf{u}}^m - \hat{\mathbf{u}}), D(\mathbf{v})) + 2\alpha\mu_1 \int_{\Gamma_2} (\hat{\mathbf{u}}^m - \hat{\mathbf{u}}) \cdot \mathbf{v} \, d\Gamma + \langle B[\tilde{\mathbf{u}}^m + \mathbf{u}^\varepsilon, \tilde{\mathbf{u}}^m + \mathbf{u}^\varepsilon, \hat{\mathbf{u}}^m - \hat{\mathbf{u}}], \mathbf{v} \rangle
$$
  
+  $\langle B[\tilde{\mathbf{u}}^m + \mathbf{u}^\varepsilon, \hat{\mathbf{u}}^m - \hat{\mathbf{u}}, \mathbf{u}^\varepsilon], \mathbf{v} \rangle$   
=  $\langle B[\tilde{\mathbf{u}} + \mathbf{u}^\varepsilon, \tilde{\mathbf{u}} + \mathbf{u}^\varepsilon, \hat{\mathbf{u}}], \mathbf{v} \rangle - \langle B[\tilde{\mathbf{u}}^m + \mathbf{u}^\varepsilon, \tilde{\mathbf{u}}^m + \mathbf{u}^\varepsilon, \hat{\mathbf{u}}], \mathbf{v} \rangle + \langle B[\tilde{\mathbf{u}} + \mathbf{u}^\varepsilon, \hat{\mathbf{u}} + \mathbf{u}^\varepsilon], \mathbf{v} \rangle$   
-  $\langle B[\tilde{\mathbf{u}}^m + \mathbf{u}^\varepsilon, \hat{\mathbf{u}} + \mathbf{u}^\varepsilon, \mathbf{u}^\varepsilon], \mathbf{v} \rangle + (F(\tilde{\mathbf{u}}^m + \mathbf{u}^\varepsilon) - F(\tilde{\mathbf{u}} + \mathbf{u}^\varepsilon), \mathbf{v}).$  (39)

Replacing  $\mathbf{v} = \hat{\mathbf{u}}^m - \hat{\mathbf{u}}$  in [\(39\)](#page-8-2) and using [\(14\)](#page-5-1), we deduce

<span id="page-8-3"></span>
$$
2\mu_1 \|\hat{\mathbf{u}}^m - \hat{\mathbf{u}}\|_{\tilde{\mathbf{H}}_{\sigma}}^2 \leq |\langle B[\tilde{\mathbf{u}}^m + \mathbf{u}^\varepsilon, \hat{\mathbf{u}}^m - \hat{\mathbf{u}}, \mathbf{u}^\varepsilon], \hat{\mathbf{u}}^m - \hat{\mathbf{u}}\rangle| + |\langle B[\tilde{\mathbf{u}} + \mathbf{u}^\varepsilon, \tilde{\mathbf{u}} + \mathbf{u}^\varepsilon, \hat{\mathbf{u}}] - B[\tilde{\mathbf{u}}^m + \mathbf{u}^\varepsilon, \tilde{\mathbf{u}}^m + \mathbf{u}^\varepsilon, \hat{\mathbf{u}}], \hat{\mathbf{u}}^m - \hat{\mathbf{u}}\rangle| + |\langle B[\tilde{\mathbf{u}} + \mathbf{u}^\varepsilon, \hat{\mathbf{u}} + \mathbf{u}^\varepsilon, \mathbf{u}^\varepsilon] - B[\tilde{\mathbf{u}}^m + \mathbf{u}^\varepsilon, \hat{\mathbf{u}} + \mathbf{u}^\varepsilon, \hat{\mathbf{u}}^\varepsilon], \hat{\mathbf{u}}^m - \hat{\mathbf{u}}\rangle| + |\langle F(\tilde{\mathbf{u}}^m + \mathbf{u}^\varepsilon) - F(\tilde{\mathbf{u}} + \mathbf{u}^\varepsilon), \hat{\mathbf{u}}^m - \hat{\mathbf{u}}\rangle|.
$$
(40)

Now we will bound the terms on the right hand side of [\(40\)](#page-8-3). Using [\(17\)](#page-5-4) and [\(37\)](#page-8-4) we obtain

<span id="page-9-0"></span>
$$
\left| \langle B[\tilde{\mathbf{u}}^m + \mathbf{u}^\varepsilon, \hat{\mathbf{u}}^m - \hat{\mathbf{u}}, \mathbf{u}^\varepsilon], \hat{\mathbf{u}}^m - \hat{\mathbf{u}} \rangle \right| \leq \left| \|\eta(N(\tilde{\mathbf{u}}^m + \mathbf{u}^\varepsilon))\|_{\infty} |\langle(\hat{\mathbf{u}}^m - \hat{\mathbf{u}}) \cdot \nabla \mathbf{u}^\varepsilon, \hat{\mathbf{u}}^m - \hat{\mathbf{u}} \rangle|
$$
  

$$
\leq C_\eta \varepsilon \|\hat{\mathbf{u}}^m - \hat{\mathbf{u}}\|_{\tilde{\mathbf{H}}_\sigma}^2.
$$
 (41)

Now, notice that

$$
B[\tilde{\mathbf{u}} + \mathbf{u}^{\varepsilon}, \tilde{\mathbf{u}} + \mathbf{u}^{\varepsilon}, \hat{\mathbf{u}}] - B[\tilde{\mathbf{u}}^m + \mathbf{u}^{\varepsilon}, \tilde{\mathbf{u}}^m + \mathbf{u}^{\varepsilon}, \hat{\mathbf{u}}]
$$
  
=  $(\eta(N(\tilde{\mathbf{u}} + \mathbf{u}^{\varepsilon})) - \eta(N(\tilde{\mathbf{u}}^m + \mathbf{u}^{\varepsilon})))(\tilde{\mathbf{u}} + \mathbf{u}^{\varepsilon}) \cdot \nabla \hat{\mathbf{u}} - \eta(N(\tilde{\mathbf{u}}^m + \mathbf{u}^{\varepsilon}))(\tilde{\mathbf{u}}^m - \tilde{\mathbf{u}}) \cdot \nabla \hat{\mathbf{u}}.$ 

Then, by the Hölder inequality and  $(37)$  we obtain

$$
|\langle B[\tilde{\mathbf{u}} + \mathbf{u}^{\varepsilon}, \tilde{\mathbf{u}} + \mathbf{u}^{\varepsilon}, \hat{\mathbf{u}}] - B[\tilde{\mathbf{u}}^m + \mathbf{u}^{\varepsilon}, \tilde{\mathbf{u}}^m + \mathbf{u}^{\varepsilon}, \hat{\mathbf{u}}], \hat{\mathbf{u}}^m - \hat{\mathbf{u}} \rangle|
$$
  
\n
$$
\leq ||\eta(N(\tilde{\mathbf{u}} + \mathbf{u}^{\varepsilon})) - \eta(N(\tilde{\mathbf{u}}^m + \mathbf{u}^{\varepsilon}))||_{\infty} |\langle(\tilde{\mathbf{u}} + \mathbf{u}^{\varepsilon}) \cdot \nabla \hat{\mathbf{u}}, \hat{\mathbf{u}}^m - \hat{\mathbf{u}} \rangle|
$$
  
\n
$$
+ ||\eta(N(\tilde{\mathbf{u}}^m + \mathbf{u}^{\varepsilon}))||_{\infty} |\langle(\tilde{\mathbf{u}}^m - \tilde{\mathbf{u}}) \cdot \nabla \hat{\mathbf{u}}, \hat{\mathbf{u}}^m - \hat{\mathbf{u}} \rangle|
$$
  
\n
$$
\leq ||\eta(N(\tilde{\mathbf{u}} + \mathbf{u}^{\varepsilon})) - \eta(N(\tilde{\mathbf{u}}^m + \mathbf{u}^{\varepsilon}))||_{\infty} ||(\tilde{\mathbf{u}} + \mathbf{u}^{\varepsilon}) \cdot \nabla \hat{\mathbf{u}}||_{\tilde{\mathbf{H}}'} ||\hat{\mathbf{u}}^m - \hat{\mathbf{u}}||_{\tilde{\mathbf{H}}_{\sigma}}
$$
  
\n
$$
+ C_{\eta} ||\tilde{\mathbf{u}}^m - \tilde{\mathbf{u}}||_3 ||\hat{\mathbf{u}}||_{\tilde{\mathbf{H}}_{\sigma}} ||\tilde{\mathbf{u}}^m - \hat{\mathbf{u}}||_6
$$
  
\n
$$
\leq C ||\eta(N(\tilde{\mathbf{u}} + \mathbf{u}^{\varepsilon})) - \eta(N(\tilde{\mathbf{u}}^m + \mathbf{u}^{\varepsilon})) ||_{\infty} ||\hat{\mathbf{u}}^m - \hat{\mathbf{u}}||_{\tilde{\mathbf{H}}_{\sigma}}
$$
  
\n
$$
+ CC_{\eta} ||\tilde{\math
$$

Again, by using the Hölder inequality we have

$$
\begin{split} |\langle B[\tilde{\mathbf{u}}+\mathbf{u}^{\varepsilon},\hat{\mathbf{u}}+\mathbf{u}^{\varepsilon},\mathbf{u}^{\varepsilon}] - B[\tilde{\mathbf{u}}^{m}+\mathbf{u}^{\varepsilon},\hat{\mathbf{u}}+\mathbf{u}^{\varepsilon},\mathbf{u}^{\varepsilon}],\hat{\mathbf{u}}^{m} - \hat{\mathbf{u}} \rangle| \\ \leq & \|\eta(N(\tilde{\mathbf{u}}+\mathbf{u}^{\varepsilon})) - \eta(N(\tilde{\mathbf{u}}^{m}+\mathbf{u}^{\varepsilon}))\|_{\infty} \|(\hat{\mathbf{u}}+\mathbf{u}^{\varepsilon}) \cdot \nabla \mathbf{u}^{\varepsilon}\|_{\tilde{\mathbf{H}}'} \|\hat{\mathbf{u}}^{m} - \hat{\mathbf{u}}\|_{\tilde{\mathbf{H}}_{\sigma}} \\ \leq & C \|\eta(N(\tilde{\mathbf{u}}+\mathbf{u}^{\varepsilon})) - \eta(N(\tilde{\mathbf{u}}^{m}+\mathbf{u}^{\varepsilon}))\|_{\infty} \|\hat{\mathbf{u}}^{m} - \hat{\mathbf{u}}\|_{\tilde{\mathbf{H}}_{\sigma}}, \end{split} \tag{43}
$$

and

<span id="page-9-1"></span>
$$
|(F(\tilde{\mathbf{u}}^{m} + \mathbf{u}^{\varepsilon}) - F(\tilde{\mathbf{u}} + \mathbf{u}^{\varepsilon}), \hat{\mathbf{u}}^{m} - \hat{\mathbf{u}})|
$$
  
\n
$$
\leq ||\eta(N(\tilde{\mathbf{u}}^{m} + \mathbf{u}^{\varepsilon})) - \eta(N(\tilde{\mathbf{u}} + \mathbf{u}^{\varepsilon}))||_{\infty} ||\mathbf{f}|| ||\hat{\mathbf{u}}^{m} - \hat{\mathbf{u}}||_{\tilde{\mathbf{H}}_{\sigma}}
$$
  
\n
$$
\leq C ||\eta(N(\tilde{\mathbf{u}}^{m} + \mathbf{u}^{\varepsilon})) - \eta(N(\tilde{\mathbf{u}} + \mathbf{u}^{\varepsilon})) ||_{\infty} ||\hat{\mathbf{u}}^{m} - \hat{\mathbf{u}}||_{\tilde{\mathbf{H}}_{\sigma}}.
$$
\n(44)

Replacing the inequalities  $(41)-(44)$  $(41)-(44)$  in  $(40)$ , and taking into account  $(37)$ , we obtain

$$
(2\mu_1 - \varepsilon C_\eta) \|\hat{\mathbf{u}}^m - \hat{\mathbf{u}}\|_{\tilde{\mathbf{H}}_{\sigma}}^2 \leq C \|\eta(N(\tilde{\mathbf{u}} + \mathbf{u}^{\varepsilon})) - \eta(N(\tilde{\mathbf{u}}^m + \mathbf{u}^{\varepsilon}))\|_{\infty} \|\hat{\mathbf{u}}^m - \hat{\mathbf{u}}\|_{\tilde{\mathbf{H}}_{\sigma}} + CC_\eta \|\tilde{\mathbf{u}}^m - \tilde{\mathbf{u}}\|_{3} \|\hat{\mathbf{u}}^m - \hat{\mathbf{u}}\|_{\tilde{\mathbf{H}}_{\sigma}}.
$$

Thus, for  $\varepsilon$  small enough such that  $2\mu_1-\varepsilon C_\eta>0$  we deduce that

<span id="page-9-2"></span>
$$
\|\hat{\mathbf{u}}^m - \hat{\mathbf{u}}\|_{\tilde{\mathbf{H}}_{\sigma}} \le C(\|\eta(N(\tilde{\mathbf{u}} + \mathbf{u}^{\varepsilon})) - \eta(N(\tilde{\mathbf{u}}^m + \mathbf{u}^{\varepsilon}))\|_{\infty} + C_{\eta} \|\tilde{\mathbf{u}}^m - \tilde{\mathbf{u}}\|_{3}).
$$
\n(45)

Passing to the limit in [\(45\)](#page-9-2), when  $m \to \infty$ , and considering the strong convergences of  ${\{\tilde{\mathbf{u}}^m\}}_{m\geq 1}$  in  $\mathbf{L}^3(\Omega)$ , and  $\{\eta(N\tilde{\mathbf{u}}^m)\}_{m\geq 1}$  in  $C^0(\overline{\Omega})$ , we have

$$
\|T\tilde{\mathbf{u}}^m - T\tilde{\mathbf{u}}\|_{\tilde{\mathbf{H}}_{\sigma}} = \|\hat{\mathbf{u}}^m - \hat{\mathbf{u}}\|_{\tilde{\mathbf{H}}_{\sigma}} \to 0, \text{ as } m \to \infty.
$$

Thus, we conclude that T is a compact operator.  $\Diamond$ 

<span id="page-9-3"></span>Lemma 5 *Let* T *be the operator defined by [\(36\)](#page-8-0), and consider the set*

$$
M_{\lambda} = \{ \tilde{\mathbf{u}} \in \tilde{\mathbf{H}}_{\sigma} : \tilde{\mathbf{u}} = \lambda T \tilde{\mathbf{u}} \text{ for some } \lambda \in [0, 1] \}. \tag{46}
$$

*If*  $\mu_1$  *and*  $\mu_2$  *are large enough such that* 

<span id="page-10-4"></span>
$$
\min\{2\mu_1 - \tilde{C}_\eta C \|\mathbf{w}_{g_2}\|_{H^{1/2}(\Gamma)} - \mu_r C, \mu_2 - \mu_r C\} > 0,
$$
\n(47)

*where*  $\tilde{C}_\eta$  *is the constant defined in Lemma [3](#page-7-7) and*  $C > 0$  *is a constant that depends only on*  $\Omega$ *, then, the set*  $M_{\lambda}$  *is bounded in*  $\tilde{\mathbf{H}}_{\sigma}$ . Moreover, for  $\lambda \in [0,1]$ , all functions  $\tilde{\mathbf{u}} = \tilde{\mathbf{u}}(\lambda) \in M_{\lambda}$  are contained, independently *of*  $\lambda$ *, in the open ball*  $B(0; R) \subset \widetilde{H}_{\sigma}$ *, with* 

<span id="page-10-5"></span>
$$
R = \tilde{C}_{\eta} C_{1} (\|\mathbf{u}_{g_{1}}\|_{\mathbf{H}^{1/2}(\Gamma\backslash\Gamma_{2})} \|\mathbf{w}_{g_{2}}\|_{H^{1/2}(\Gamma)} + \|\mathbf{u}_{g_{1}}\|_{\mathbf{H}^{1/2}(\Gamma\backslash\Gamma_{2})}^{2} + \|f\| + \|g\|)
$$
  
+  $C_{1} (\|\mathbf{u}_{g_{1}}\|_{\mathbf{H}^{1/2}(\Gamma\backslash\Gamma_{2})} + \|\mathbf{w}_{g_{2}}\|_{H^{1/2}(\Gamma)}),$  (48)

*where the constant*  $C_1 > 0$  *depends only on*  $\Omega$ ,  $\mu_1$  *and*  $\mu_2$ *.* 

*Proof.* Assuming  $\lambda > 0$ , for any  $\tilde{\mathbf{u}} \in M_{\lambda}$  we can write  $T\tilde{\mathbf{u}} = \frac{1}{\lambda}\tilde{\mathbf{u}}$ ; then, substituting  $\hat{\mathbf{u}} = \frac{1}{\lambda}\tilde{\mathbf{u}}$  and  $\mathbf{v} = \lambda\tilde{\mathbf{u}}$  in [\(36\)](#page-8-0), and taking into account [\(14\)](#page-5-1), we obtain

<span id="page-10-0"></span>
$$
2\mu_1 \|\tilde{\mathbf{u}}\|_{\tilde{\mathbf{H}}_{\sigma}}^2 \leq |\langle B[\tilde{\mathbf{u}} + \mathbf{u}^{\varepsilon}, \tilde{\mathbf{u}}, \mathbf{u}^{\varepsilon}], \tilde{\mathbf{u}}\rangle| + 2\mu_r \lambda |(\text{rot}(\hat{\mathbf{w}} + \mathbf{w}^{\mathrm{e}}), \tilde{\mathbf{u}})| + \lambda |(F(\tilde{\mathbf{u}} + \mathbf{u}^{\varepsilon}), \tilde{\mathbf{u}})| + 2\mu_1 \lambda |(D(\mathbf{u}^{\varepsilon}), D(\tilde{\mathbf{u}}))| + \lambda |\langle B[\tilde{\mathbf{u}} + \mathbf{u}^{\varepsilon}, \mathbf{u}^{\varepsilon}], \tilde{\mathbf{u}}\rangle|.
$$
(49)

Now we will bound the terms on the right hand side of  $(49)$ . Using the Hölder inequality and  $(17)$  we obtain

<span id="page-10-1"></span>
$$
|\langle B[\tilde{\mathbf{u}}+\mathbf{u}^{\varepsilon},\tilde{\mathbf{u}},\mathbf{u}^{\varepsilon}],\tilde{\mathbf{u}}\rangle| \leq \varepsilon \|\tilde{\mathbf{u}}\|_{\tilde{\mathbf{H}}_{\sigma}}^2 \|\eta(N(\tilde{\mathbf{u}}+\mathbf{u}^{\varepsilon}))\|_{\infty} \leq \varepsilon \tilde{C}_{\eta} \|\tilde{\mathbf{u}}\|_{\tilde{\mathbf{H}}_{\sigma}}^2.
$$
 (50)

Applying the Hölder and Poincaré inequalities we obtain

<span id="page-10-2"></span>
$$
2\mu_r \lambda |(\text{rot}(\hat{\mathbf{w}} + \mathbf{w}^e), \tilde{\mathbf{u}})| \leq 2\mu_r \lambda ||\text{rot}(\hat{\mathbf{w}} + \mathbf{w}^e)|| \|\tilde{\mathbf{u}}\| \leq \lambda \mu_r C ||\nabla(\hat{\mathbf{w}} + \mathbf{w}^e)|| \|\tilde{\mathbf{u}}\|_{\tilde{\mathbf{H}}_{\sigma}} \leq \lambda \mu_r C (||\hat{\mathbf{w}}||_{H_0^1} + ||\mathbf{w}^e||_{H^1}) ||\tilde{\mathbf{u}}||_{\tilde{\mathbf{H}}_{\sigma}},
$$
\n(51)

$$
\lambda |(F(\tilde{\mathbf{u}} + \mathbf{u}^{\varepsilon}), \tilde{\mathbf{u}})| \leq \lambda \|\eta(N(\tilde{\mathbf{u}} + \mathbf{u}^{\varepsilon}))\|_{\infty} \|\mathbf{f}\| \|\tilde{\mathbf{u}}\|_{\tilde{\mathbf{H}}_{\sigma}} \leq \lambda \tilde{C}_{\eta} \|\mathbf{f}\| \|\tilde{\mathbf{u}}\|_{\tilde{\mathbf{H}}_{\sigma}},
$$
\n(52)

$$
2\mu_1\lambda|(D(\mathbf{u}^{\varepsilon}),D(\tilde{\mathbf{u}}))|\leq 2\mu_1\lambda\|D(\mathbf{u}^{\varepsilon})\|\|\tilde{\mathbf{u}}\|_{\tilde{\mathbf{H}}_{\sigma}}\leq 2\mu_1\lambda\|\mathbf{u}^{\varepsilon}\|_{\mathbf{H}_{\sigma}}\|\tilde{\mathbf{u}}\|_{\tilde{\mathbf{H}}_{\sigma}},\tag{53}
$$

$$
\lambda |\langle B[\tilde{\mathbf{u}} + \mathbf{u}^{\varepsilon}, \mathbf{u}^{\varepsilon}, \mathbf{u}^{\varepsilon}], \tilde{\mathbf{u}} \rangle| \leq \lambda \|\eta(N(\tilde{\mathbf{u}} + \mathbf{u}^{\varepsilon}))\|_{\infty} \|\mathbf{u}^{\varepsilon}\|_{3} \|\nabla \mathbf{u}^{\varepsilon}\| \|\tilde{\mathbf{u}}\|_{6} \leq \lambda \tilde{C}_{\eta} C \|\mathbf{u}^{\varepsilon}\|_{\mathbf{H}_{\sigma}}^{2} \|\tilde{\mathbf{u}}\|_{\tilde{\mathbf{H}}_{\sigma}}. \tag{54}
$$

Replacing [\(50\)](#page-10-1)-[\(54\)](#page-10-2) in [\(49\)](#page-10-0) and taking into account that  $\lambda \leq 1$ , we have

$$
(2\mu_1 - \varepsilon \tilde{C}_{\eta}) \|\tilde{\mathbf{u}}\|_{\tilde{\mathbf{H}}_{\sigma}}^2 \leq \mu_r C(\|\hat{\mathbf{w}}\|_{H_0^1} + \|\mathbf{w}^{\varepsilon}\|_{H^1}) \|\tilde{\mathbf{u}}\|_{\tilde{\mathbf{H}}_{\sigma}} + \tilde{C}_{\eta} \|\mathbf{f}\| \|\tilde{\mathbf{u}}\|_{\tilde{\mathbf{H}}_{\sigma}} + 2\mu_1 \|\mathbf{u}^{\varepsilon}\|_{\mathbf{H}_{\sigma}} \|\tilde{\mathbf{u}}\|_{\tilde{\mathbf{H}}_{\sigma}} + C\tilde{C}_{\eta} \|\mathbf{u}^{\varepsilon}\|_{\mathbf{H}_{\sigma}}^2 \|\tilde{\mathbf{u}}\|_{\tilde{\mathbf{H}}_{\sigma}},
$$

which implies

<span id="page-10-3"></span>
$$
(2\mu_1 - \varepsilon \tilde{C}_{\eta}) \|\tilde{\mathbf{u}}\|_{\tilde{\mathbf{H}}_{\sigma}} \leq \mu_r C(\|\hat{\mathbf{w}}\|_{H_0^1} + \|\mathbf{w}^e\|_{H^1}) + \tilde{C}_{\eta} \|\mathbf{f}\| + 2\mu_1 \|\mathbf{u}^{\varepsilon}\|_{\mathbf{H}_{\sigma}} + \tilde{C}_{\eta} C \|\mathbf{u}^{\varepsilon}\|_{\mathbf{H}_{\sigma}}^2. \tag{55}
$$

Adding  $(24)$  and  $(55)$ , and taking into account  $(17)-(18)$  $(17)-(18)$ , we get

$$
(2\mu_1 - \varepsilon \tilde{C}_{\eta}) \|\tilde{\mathbf{u}}\|_{\tilde{\mathbf{H}}_{\sigma}} + \mu_2 \|\hat{\mathbf{w}}\|_{H_0^1} \leq (\tilde{C}_{\eta} C \|\mathbf{w}_{g_2}\|_{H^{1/2}(\Gamma)} + \mu_r C) \|\tilde{\mathbf{u}}\|_{\tilde{\mathbf{H}}_{\sigma}} + \mu_r C \|\hat{\mathbf{w}}\|_{H_0^1} + \tilde{C}_{\eta} C \|\mathbf{u}_{g_1}\|_{\mathbf{H}^{1/2}(\Gamma \setminus \Gamma_2)} \|\mathbf{w}_{g_2}\|_{H^{1/2}(\Gamma)} + \mu_r C (\|\mathbf{u}_{g_1}\|_{\mathbf{H}^{1/2}(\Gamma \setminus \Gamma_2)} + \|\mathbf{w}_{g_2}\|_{H^{1/2}(\Gamma)}) + \mu_2 C \|\mathbf{w}_{g_2}\|_{H^{1/2}(\Gamma)} + 2\mu_1 C \|\mathbf{u}_{g_1}\|_{\mathbf{H}^{1/2}(\Gamma \setminus \Gamma_2)} + \tilde{C}_{\eta} C \|\mathbf{u}_{g_1}\|_{\mathbf{H}^{1/2}(\Gamma \setminus \Gamma_2)} + \tilde{C}_{\eta} (\|\mathbf{f}\| + \|\mathbf{g}\|),
$$

and thus,

<span id="page-11-2"></span>
$$
(2\mu_1 - \varepsilon \tilde{C}_{\eta} - \tilde{C}_{\eta} C \|\mathbf{w}_{g_2}\|_{H^{1/2}(\Gamma)} - \mu_r C) \|\tilde{\mathbf{u}}\|_{\tilde{\mathbf{H}}_{\sigma}} + (\mu_2 - \mu_r C) \|\hat{\mathbf{w}}\|_{H_0^1}
$$
  
\n
$$
\leq \tilde{C}_{\eta} C \|\mathbf{u}_{g_1}\|_{\mathbf{H}^{1/2}(\Gamma \backslash \Gamma_2)} \|\mathbf{w}_{g_2}\|_{H^{1/2}(\Gamma)} + \mu_r C (\|\mathbf{u}_{g_1}\|_{\mathbf{H}^{1/2}(\Gamma \backslash \Gamma_2)} + \|\mathbf{w}_{g_2}\|_{H^{1/2}(\Gamma)})
$$
  
\n
$$
+ \mu_2 C \|\mathbf{w}_{g_2}\|_{H^{1/2}(\Gamma)} + 2\mu_1 C \|\mathbf{u}_{g_1}\|_{\mathbf{H}^{1/2}(\Gamma \backslash \Gamma_2)} + \tilde{C}_{\eta} C \|\mathbf{u}_{g_1}\|_{\mathbf{H}^{1/2}(\Gamma \backslash \Gamma_2)}
$$
  
\n
$$
+ \tilde{C}_{\eta} (\|\mathbf{f}\| + \|\mathbf{g}\|).
$$
 (56)

By using [\(47\)](#page-10-4),  $\mu_2 - \mu_r C > 0$  and  $\delta = 2\mu_1 - \varepsilon \tilde{C}_\eta - \tilde{C}_\eta C ||\mathbf{w}_{g_2}||_{H^{1/2}(\Gamma)} - \mu_r C > 0$ , for  $\varepsilon$  small enough. Then, from [\(56\)](#page-11-2) it follows

<span id="page-11-3"></span>
$$
\delta \|\tilde{\mathbf{u}}\|_{\tilde{\mathbf{H}}_{\sigma}} \leq \tilde{C}_{\eta} C \|\mathbf{u}_{g_1}\|_{\mathbf{H}^{1/2}(\Gamma \backslash \Gamma_2)} \|\mathbf{w}_{g_2}\|_{H^{1/2}(\Gamma)} + \mu_r C (\|\mathbf{u}_{g_1}\|_{\mathbf{H}^{1/2}(\Gamma \backslash \Gamma_2)} + \|\mathbf{w}_{g_2}\|_{H^{1/2}(\Gamma)}) + \mu_2 C \|\mathbf{w}_{g_2}\|_{H^{1/2}(\Gamma)} + 2\mu_1 C \|\mathbf{u}_{g_1}\|_{\mathbf{H}^{1/2}(\Gamma \backslash \Gamma_2)} + \tilde{C}_{\eta} C \|\mathbf{u}_{g_1}\|_{\mathbf{H}^{1/2}(\Gamma \backslash \Gamma_2)}^2 + \tilde{C}_{\eta} (\|\mathbf{f}\| + \|\mathbf{g}\|),
$$
\n(57)

which implies that  $M_{\lambda}$  is bounded in  $\tilde{H}_{\sigma}$  for  $\lambda > 0$ . For  $\lambda = 0$  the result is trivial. The radius R in [\(48\)](#page-10-5) follows from [\(57\)](#page-11-3). ⋄

<span id="page-11-0"></span>With the previous results, we have the following theorem of existence of solutions for the system [\(19\)](#page-6-2).

**Theorem 1** Let  $f \in L^2(\Omega)$ ,  $g \in L^2(\Omega)$ ,  $\eta \in C^0(\mathbb{R})$ , and  $\mu_1$ ,  $\mu_2$  satisfying [\(47\)](#page-10-4). Then, the operator  $T: \tilde{H}_{\sigma} \to \tilde{H}_{\sigma}$  defined in *(36),* with  $\hat{w}$  a solution of *(23),* has a fixed point  $\hat{u} \in \tilde{H}_{\sigma}$  and the pair of functions  $[\hat{u}, \hat{w}]$  *is a solution of system [\(19\)](#page-6-2). Furthermore, the solution*  $[\hat{u}, \hat{w}]$  *satisfies the inequality* 

<span id="page-11-5"></span>
$$
\|\hat{\mathbf{u}}\|_{\tilde{\mathbf{H}}_{\sigma}} + \|\hat{\mathbf{w}}\|_{H_0^1} \le C\Theta,
$$
\n<sup>(58)</sup>

 $where \ \Theta = \|\mathbf{u}_{g_1}\|_{\mathbf{H}^{1/2}(\Gamma \backslash \Gamma_2)} \|\mathbf{w}_{g_2}\|_{H^{1/2}(\Gamma)} + \|\mathbf{u}_{g_1}\|_{\mathbf{H}^{1/2}(\Gamma \backslash \Gamma_2)} + \|\mathbf{w}_{g_2}\|_{H^{1/2}(\Gamma)} + \|\mathbf{u}_{g_1}\|_{\mathbf{H}^{1/2}(\Gamma \backslash \Gamma_2)}^2 + \|\mathbf{f}\| + \|g\|,$  and C positive constant that depends only on  $\Omega$ ,  $\mu_1$ ,  $\mu_2$ ,  $\mu_r$ ,  $\tilde{C}_\eta$ .

*Proof.* From Lemmas [4](#page-8-5) and [5,](#page-9-3) it follows that the operator T and the set  $M_{\lambda}$  satisfy the conditions of the Leray-Schauder theorem (cf. Theorem 1.2.4, p. 42 of [\[22\]](#page-23-4)); therefore the operator T has a fixed point, that is, there exists  $\hat{\mathbf{u}} \in \tilde{\mathbf{H}}_{\sigma}$  such that  $T\hat{\mathbf{u}} = \hat{\mathbf{u}}$ . Then, by the definition of T it follows that  $\tilde{\mathbf{u}} = \hat{\mathbf{u}}$  which satisfies [\(36\)](#page-8-0), and the auxiliary equation [\(23\)](#page-7-1). Thus, we concluded that  $[\hat{\mathbf{u}}, \hat{\mathbf{w}}]$  is a solution of system [\(19\)](#page-6-2).

Now, from [\(56\)](#page-11-2) with  $\tilde{\mathbf{u}} = \hat{\mathbf{u}}$  we have

<span id="page-11-4"></span>
$$
(2\mu_1 - \varepsilon \tilde{C}_{\eta} - \tilde{C}_{\eta} C \|\mathbf{w}_{g_2}\|_{H^{1/2}(\Gamma)} - \mu_r C) \|\hat{\mathbf{u}}\|_{\tilde{\mathbf{H}}_{\sigma}} + (\mu_2 - \mu_r C) \|\hat{\mathbf{w}}\|_{H_0^1}
$$
  
\n
$$
\leq \tilde{C}_{\eta} C \|\mathbf{u}_{g_1}\|_{\mathbf{H}^{1/2}(\Gamma \backslash \Gamma_2)} \|\mathbf{w}_{g_2}\|_{H^{1/2}(\Gamma)} + \mu_r C (\|\mathbf{u}_{g_1}\|_{\mathbf{H}^{1/2}(\Gamma \backslash \Gamma_2)} + \|\mathbf{w}_{g_2}\|_{H^{1/2}(\Gamma)})
$$
  
\n
$$
+ \mu_2 C \|\mathbf{w}_{g_2}\|_{H^{1/2}(\Gamma)} + 2\mu_1 C \|\mathbf{u}_{g_1}\|_{\mathbf{H}^{1/2}(\Gamma \backslash \Gamma_2)} + \tilde{C}_{\eta} C \|\mathbf{u}_{g_1}\|_{\mathbf{H}^{1/2}(\Gamma \backslash \Gamma_2)}^2
$$
  
\n
$$
+ \tilde{C}_{\eta} (\|\mathbf{f}\| + \|\mathbf{g}\|).
$$
  
\n(59)

From [\(47\)](#page-10-4) we have that  $\hat{\delta} = \min\{2\mu_1 - \varepsilon \tilde{C}_\eta - \tilde{C}_\eta C ||\mathbf{w}_{g_2}||_{H^{1/2}(\Gamma)} - \mu_r C, \mu_2 - \mu_r C\} > 0$ , for  $\varepsilon$  small enough. Then, from [\(59\)](#page-11-4) we deduce that

$$
\hat{\delta}(\|\hat{\mathbf{u}}\|_{\tilde{\mathbf{H}}_{\sigma}} + \|\hat{\mathbf{w}}\|_{H_0^1}) \leq C(\|\mathbf{u}_{g_1}\|_{\mathbf{H}^{1/2}(\Gamma \backslash \Gamma_2)} \|\mathbf{w}_{g_2}\|_{H^{1/2}(\Gamma)} + \|\mathbf{u}_{g_1}\|_{\mathbf{H}^{1/2}(\Gamma \backslash \Gamma_2)} + \|\mathbf{w}_{g_2}\|_{H^{1/2}(\Gamma)} + \|\mathbf{u}_{g_1}\|_{\mathbf{H}^{1/2}(\Gamma \backslash \Gamma_2)} + \|\mathbf{u}_{g_1}\|_{\mathbf{H}^{1/2}(\Gamma \backslash \Gamma_2)} + \|\mathbf{f}\| + \|g\|),
$$

which implies inequality [\(58\)](#page-11-5). ⋄

As a consequence of Theorem [1,](#page-11-0) we have the following result.

<span id="page-11-1"></span>

**Theorem 2** Let **f**  $\in$  **L**<sup>2</sup>( $\Omega$ ),  $g \in L^2(\Omega)$ , **u**<sub>0</sub>  $\in$  **H**<sub>00</sub><sup>1</sup>/<sub>2</sub>( $\Gamma$ <sub>0</sub>), **g**<sub>1</sub>  $\in$  **H**<sub>00</sub><sup>1</sup>/<sub>2</sub>( $\Gamma$ <sub>0</sub>), *g*<sub>2</sub>  $\in$  *H*<sub>00</sub><sup>1</sup>/<sub>2</sub>( $\Gamma$ <sub>3</sub>), **g**<sub>2</sub>  $\in$  *H*<sub>00</sub><sup>1</sup>/<sub>2</sub>( $\Gamma$ <sub>3</sub>),  $\eta \in C^0(\mathbb{R})$ ,  $\mu_1$  and  $\mu_2$  satisfying the condition  $(47)$  given in Theorem [1.](#page-11-0) Then, there exists functions  $[\mathbf{u}, \mathbf{w}] \in \mathbf{H}_{\sigma} \times H^1(\Omega)$  and  $\rho = \eta(N\mathbf{u})$  satisfying the nonhomogeneous system [\(16\)](#page-5-3). Moreover, the solution [u, w] *satisfies the following inequality*

<span id="page-12-2"></span>
$$
\|\mathbf{u}\|_{\mathbf{H}_{\sigma}} + \|\mathbf{w}\|_{H^1} \le C\Theta,\tag{60}
$$

*where C is a positive constant depending on*  $\Omega$ ,  $\mu_1$ ,  $\mu_2$ , and  $\Theta$  *is defined as in* [\(58\)](#page-11-5)*.* 

*Proof.* Considering the solution  $[\hat{\mathbf{u}}, \hat{\mathbf{w}}] \in \tilde{\mathbf{H}}_{\sigma} \times H_0^1(\Omega)$  of system [\(19\)](#page-6-2) given by Theorem [1,](#page-11-0) we deduce that there exists a solution  $[\mathbf{u}, \mathbf{w}] \in \mathbf{H}_{\sigma} \times H^1(\Omega)$  for system [\(16\)](#page-5-3), where  $\mathbf{u} = \hat{\mathbf{u}} + \mathbf{u}^{\varepsilon}$  and  $\mathbf{w} = \hat{\mathbf{w}} + \mathbf{w}^e$ . Moreover, by using triangle inequality we have

$$
\begin{array}{rcl}\|\mathbf{u}\|_{\mathbf{H}_{\sigma}}+\|\mathbf{w}\|_{H^1} & \leq & \|\hat{\mathbf{u}}\|_{\tilde{\mathbf{H}}_{\sigma}}+\|\hat{\mathbf{w}}\|_{H_0^1}+\|\mathbf{u}^{\varepsilon}\|_{\mathbf{H}_{\sigma}}+\|\mathbf{w}^e\|_{H^1} \\
& \leq & C\Theta+C(\|\mathbf{u}_{\mathbf{g}_1}\|_{\mathbf{H}^{1/2}(\Gamma\backslash\Gamma_2)}+\|\mathbf{w}_{g_2}\|_{H^{1/2}(\Gamma)}),\end{array}
$$

which implies inequality [\(60\)](#page-12-2).  $\Diamond$ 

## <span id="page-12-1"></span>4 Existence of an Optimal Solution

In this section, we will prove the existence of an optimal solution for problem [\(22\)](#page-6-1). We remember that the set of admissible solutions of problem [\(22\)](#page-6-1) is defined by

<span id="page-12-0"></span>
$$
\mathcal{S}_{ad} = \{ \mathbf{s} = [\mathbf{u}, \mathbf{w}, \mathbf{g}_1, g_2] \in \mathbb{X} \text{ such that } J(\mathbf{s}) < \infty \text{ and } \mathbf{s} \text{ satisfies (16)} \},
$$

where  $\mathbb{X} = \mathbf{H}_{\sigma} \times H^1(\Omega) \times \mathcal{U}_1 \times \mathcal{U}_2$ . We have the following result.

Theorem 3 *Under the conditions of Theorem [2,](#page-11-1) if one the conditions* (i) *or* (ii) *in [\(21\)](#page-6-3) is satisfied, then the problem* [\(22\)](#page-6-1) has at least one optimal solution  $\tilde{\mathbf{s}} = [\tilde{\mathbf{u}}, \tilde{\mathbf{w}}, \tilde{\mathbf{g}}_1, \tilde{\mathbf{g}}_2] \in S_{ad}$ .

*Proof.* From Theorem [2](#page-11-1) we have that  $S_{ad}$  is nonempty, and since the functional  $J$  is bounded below, there exists a minimizing sequence  $\{s^m = [\mathbf{u}^m, \mathbf{w}^m, g_1^m, g_2^m]\}_{m \ge 1} \in \mathcal{S}_{ad}, m \in \mathbb{N}$ , such that  $\lim_{m \to \infty} J(\mathbf{s}^m) = \inf_{\mathbf{s} \in \mathcal{S}_{ad}} J(\mathbf{s})$ . Moreover,  $\mathbf{s}^m$  satisfies the system [\(16\)](#page-5-3), that is,

<span id="page-12-4"></span>
$$
\begin{cases}\n2\mu_1(D(\mathbf{u}^m), D(\mathbf{v})) + 2\alpha\mu_1 \int_{\Gamma_2} \mathbf{u}^m \cdot \mathbf{v} \, d\Gamma + \langle B[\mathbf{u}^m, \mathbf{u}^m, \mathbf{u}^m], \mathbf{v} \rangle = 2\mu_r(\text{rot } \mathbf{w}^m, \mathbf{v}) + (F(\mathbf{u}^m), \mathbf{v}), \\
\mu_2(\nabla \mathbf{w}^m, \nabla z) + \langle \tilde{B}[\mathbf{u}^m, \mathbf{u}^m, \mathbf{w}^m], z \rangle + 4\mu_r(\mathbf{w}^m, z) = 2\mu_r(\text{rot } \mathbf{u}^m, z) + (G(\mathbf{u}^m), z), \\
\mathbf{u}^m = \mathbf{u}_{g_1^m} \quad \text{on } \Gamma \setminus \Gamma_2, \\
\mathbf{w}^m = \mathbf{w}_{g_2^m} \quad \text{on } \Gamma, \\
\eta(N\mathbf{u}^m) = \rho_0 \quad \text{on } \Gamma_0,\n\end{cases} \tag{61}
$$

for all  $[\mathbf{v}, \mathbf{z}] \in \tilde{\mathbf{H}}_{\sigma} \times H_0^1(\Omega)$ , and  $\eta \in C^0(\mathbb{R})$ .

If one of the conditions  $(i)$  or  $(ii)$  in  $(21)$  is satisfied, we have that there exists a constant C independent of m such that  $\|\boldsymbol{g}_1^m\|_{\mathbf{H}^{1/2}(\Gamma_1)}^2 + \|g_2^m\|_{H^{1/2}(\Gamma_3)}^2 \leq C$ ; consequently, [\(60\)](#page-12-2) implies  $\|\mathbf{u}^m\|_{\mathbf{H}^{\sigma}} + \|\mathbf{w}^m\|_{H^1} \leq C$ . Therefore, since  $\mathcal{U}_1 \times \mathcal{U}_2$  is closed convex subset of  $\mathbf{H}_{00}^{1/2}(\Gamma_1) \times H_{00}^{1/2}(\Gamma_3)$ , there exists  $\tilde{\mathbf{s}} = [\tilde{\mathbf{u}}, \tilde{\mathbf{w}}, \tilde{\mathbf{g}}_1, \tilde{\mathbf{g}}_2] \in \mathbb{X}$  and some subsequence of  $\{\mathbf s^m\}_{m\geq 1}$ , still denoted by  $\{\mathbf s^m\}_{m\geq 1}$ , such that when  $m\to\infty$  we have

> <span id="page-12-3"></span> $\mathbf{u}^m \to \tilde{\mathbf{u}}$  weakly in  $\mathbf{H}_{\sigma}$  and strongly in  $\mathbf{L}^2(\Omega)$ ,  $w^m \to \tilde{w}$  weakly in  $H^1(\Omega)$  and strongly in  $L^2(\Omega)$ ,  $g_1^m \to \tilde{g}_1$  weakly in  $\mathbf{H}_{00}^{1/2}(\Gamma_1)$  and strongly in  $\mathbf{L}^2(\Gamma_1)$ ,  $g_2^m \to \tilde{g}_2$  weakly in  $H_{00}^{1/2}(\Gamma_3)$  and strongly in  $L^2(\Gamma_3)$ . (62)

Moreover, since  $\mathbf{u}^m = \mathbf{u}_{\mathbf{g}_1^m}$  on  $\Gamma \setminus \Gamma_2$ ,  $\mathbf{w}_2^m = \mathbf{w}_{\mathbf{g}_2^m}$  on  $\Gamma$ , and  $\eta(N\mathbf{u}^m) = \rho_0$  on  $\Gamma_0$ , from [\(62\)](#page-12-3) it follows that  $\tilde{\mathbf{u}} = \mathbf{u}_{\tilde{\mathbf{g}}_1}$  on  $\Gamma \setminus \Gamma_2$ ,  $\tilde{\mathbf{w}} = \mathbf{w}_{\tilde{\mathbf{g}}_2}$  on  $\Gamma$ , and  $\eta(N\tilde{\mathbf{u}}) = \rho_0$  on  $\Gamma_0$ ; thus,  $\tilde{\mathbf{s}}$  satisfies the boundary conditions in [\(16\)](#page-5-3). A standard procedure permits to pass to the limit in [\(61\)](#page-12-4) when m goes to  $\infty$ , and then,  $\tilde{s}$  is solution of the system [\(16\)](#page-5-3). Thus, we have  $\tilde{\mathbf{s}} \in \mathcal{S}_{ad}$  and

<span id="page-13-1"></span>
$$
\lim_{m \to \infty} J(\mathbf{s}^m) = \inf_{\mathbf{s} \in S_{ad}} J(\mathbf{s}) \le J(\tilde{\mathbf{s}}). \tag{63}
$$

Also, since the functional J is weakly lower semicontinuous on  $\mathcal{S}_{ad}$ , we get that  $J(\tilde{s}) \leq \lim_{m\to\infty} \inf J(\mathbf{s}^m)$ . Therefore, from [\(63\)](#page-13-1) and the last inequality, we conclude that  $J[\tilde{\mathbf{u}}, \tilde{\mathbf{w}}, \tilde{\mathbf{g}}_1, \tilde{g}_2] = J(\tilde{\mathbf{s}}) = \min_{\mathbf{s} \in S_{ad}} J(\mathbf{s})$ , which implies the existence of an optimal solution for the control problem [\(22\)](#page-6-1). ⋄

## <span id="page-13-0"></span>5 Necessary Optimality Conditions and an Optimality System

This section is devoted to obtain an optimality system to problem [\(22\)](#page-6-1). We shall use the theorem of Lagrange multipliers to turn the constrained optimization problem [\(22\)](#page-6-1) into an unconstrained one. In order to prove the existence of Lagrange multipliers, we use a penalty method. This method consists in introducing a family of penalized problems  $(P)_{\varepsilon}$ ,  $\varepsilon > 0$ , whose solutions converge towards a solution to problem [\(22\)](#page-6-1); then we derive the optimality conditions for problems  $(P)_{\varepsilon}$  and finally, we pass to the limit in these optimality conditions. This method has been previously used in [\[5,](#page-22-11) [18,](#page-23-12) [20\]](#page-23-13) in the context of Navier-Stokes and Boussinesq equations.

#### 5.1 Penalized Problem

For an optimal solution  $\tilde{\mathbf{s}} = [\tilde{\mathbf{u}}, \tilde{\mathbf{w}}, \tilde{\mathbf{g}}_1, \tilde{g}_2]$  of the optimal control problem [\(22\)](#page-6-1) we consider the following family of auxiliary extremal problems: Find

<span id="page-13-3"></span>
$$
\min J_{\varepsilon}[\mathbf{u}, \mathbf{w}, \mathbf{g}_1, g_2], \qquad [\mathbf{u}, \mathbf{w}, \mathbf{g}_1, g_2] \in \mathbb{X}, \tag{64}
$$

where for any  $\varepsilon > 0$  the functional  $J_{\varepsilon} : \mathbb{X} \to \mathbb{R}$  is defined by

<span id="page-13-2"></span>
$$
J_{\varepsilon}[\mathbf{u}, \mathbf{w}, \mathbf{g}_{1}, g_{2}] = J[\mathbf{u}, \mathbf{w}, \mathbf{g}_{1}, g_{2}] + \frac{1}{2} ||\mathbf{u} - \tilde{\mathbf{u}}||_{\mathbf{H}\sigma}^{2} + \frac{1}{2} ||\mathbf{w} - \tilde{\mathbf{w}}||_{H^{1}}^{2} + \frac{1}{2} ||\mathbf{g}_{1} - \tilde{\mathbf{g}}_{1}||_{\mathbf{H}^{1/2}(\Gamma_{1})}^{2} + \frac{1}{2} ||g_{2} - \tilde{g}_{2}||_{H^{1/2}(\Gamma_{3})}^{2} + \frac{1}{2\varepsilon} ||\mu_{1}A\mathbf{u} + B[\mathbf{u}, \mathbf{u}, \mathbf{u}] - 2\mu_{r} \text{rot } \mathbf{w} - F(\mathbf{u})||_{\mathbf{H}^{1}}^{2} + \frac{1}{2\varepsilon} ||\mu_{2}\tilde{A}\mathbf{w} + \tilde{B}[\mathbf{u}, \mathbf{u}, \mathbf{w}] + 4\mu_{r}\mathbf{w} - 2\mu_{r} \text{rot } \mathbf{u} - G(\mathbf{u})||_{H^{-1}}^{2} + \frac{1}{2\varepsilon} ||\mathbf{u} - \mathbf{u}_{\mathbf{g}_{1}}||_{\mathbf{H}^{1/2}(\Gamma\backslash\Gamma_{2})}^{2} + \frac{1}{2\varepsilon} ||\mathbf{w} - \mathbf{w}_{g_{2}}||_{H^{1/2}(\Gamma)}^{2}.
$$
 (65)

In [\(65\)](#page-13-2),  $J[\mathbf{u}, \mathbf{w}, \mathbf{g}_1, g_2]$  is the functional defined in [\(22\)](#page-6-1), the operators B,  $\tilde{B}$ , F, and G are defined in [\(13\)](#page-4-3), and the operators  $A: \mathbf{H}_{\sigma} \to \tilde{\mathbf{H}}'$  and  $\tilde{A}: H^1(\Omega) \to H^{-1}(\Omega)$  are defined by

<span id="page-13-5"></span>
$$
\langle A\mathbf{u}, \mathbf{v} \rangle = 2(D(\mathbf{u}), D(\mathbf{v})) + 2\alpha \int_{\Gamma_2} \mathbf{u} \cdot \mathbf{v} \, d\Gamma \quad \forall \mathbf{v} \in \tilde{\mathbf{H}}_{\sigma}, \qquad \langle \tilde{A}\mathbf{w}, \mathbf{z} \rangle = (\nabla \mathbf{w}, \nabla \mathbf{z}) \quad \forall \mathbf{z} \in H_0^1(\Omega), \tag{66}
$$

where  $\alpha$  is given in [\(3\)](#page-2-1).

**Remark 3** *Since*  $\tilde{\mathbf{s}}$  *satisfies* [\(16\)](#page-5-3)*, then by definition of*  $J_{\varepsilon}$  *it holds* 

<span id="page-13-4"></span>
$$
J_{\varepsilon}(\tilde{\mathbf{s}}) = J(\tilde{\mathbf{s}}). \tag{67}
$$

Following the proof of Theorem [3,](#page-12-0) and recalling that the functional  $J_{\varepsilon}$  is weakly lower semincontinuous, we can prove that there exists an optimal solution of problem [\(64\)](#page-13-3)-[\(65\)](#page-13-2), that is, for each  $\varepsilon > 0$ , there exists  $\mathbf{s}^{\varepsilon} = [\mathbf{u}^{\varepsilon}, \mathbf{w}^{\varepsilon}, \mathbf{g}^{\varepsilon}_1, g^{\varepsilon}_2] \in \mathbb{X}$  such that

$$
J_{\varepsilon}(\mathbf{s}^{\varepsilon}) = \min_{\mathbf{s} \in \mathbb{X}} J_{\varepsilon}(\mathbf{s}).
$$

<span id="page-14-0"></span>**Theorem 4** *Assume the hypotheses of Theorem [3.](#page-12-0) For each*  $\varepsilon > 0$ , let  $\mathbf{s}^{\varepsilon} = [\mathbf{u}^{\varepsilon}, \mathbf{w}^{\varepsilon}, \mathbf{g}_1^{\varepsilon}, \mathbf{g}_2^{\varepsilon}]$  *a solution of problem* [\(64\)](#page-13-3)<sup>*-*</sup>[\(65\)](#page-13-2)*.* Then, as  $\varepsilon \to 0$ , the sequence  $\{s^{\varepsilon} = [\mathbf{u}^{\varepsilon}, \mathbf{w}^{\varepsilon}, \mathbf{g}_1^{\varepsilon}, g_2^{\varepsilon}]\}_{\varepsilon > 0}$  satisfies the following convergences

<span id="page-14-5"></span> $[\mathbf{u}^{\varepsilon}, \mathbf{w}^{\varepsilon}, \mathbf{g}_{1}^{\varepsilon}, \tilde{g}_{2}^{\varepsilon}] \longrightarrow [\tilde{\mathbf{u}}, \tilde{\mathbf{w}}, \tilde{\mathbf{g}}_{1}, \tilde{g}_{2}] \text{ strongly in } \mathcal{S}_{ad},$ (68)

$$
J_{\varepsilon}[\mathbf{u}^{\varepsilon}, \mathbf{w}^{\varepsilon}, \mathbf{g}_1^{\varepsilon}, g_2^{\varepsilon}] \longrightarrow J[\tilde{\mathbf{u}}, \tilde{\mathbf{w}}, \tilde{\mathbf{g}}_1, \tilde{g}_2],
$$
\n(69)

where  $\tilde{\mathbf{s}} = [\tilde{\mathbf{u}}, \tilde{\mathbf{w}}, \tilde{\mathbf{g}}_1, \tilde{\mathbf{g}}_2] \in \mathcal{S}_{ad}$  *is an optimal solution of problem [\(22\)](#page-6-1).* 

*Proof.* Since  $\tilde{s} \in S_{ad} \subset \mathbb{X}$  and the functional  $J_{\varepsilon}$  attains its minimum in  $[\mathbf{u}^{\varepsilon}, \mathbf{w}^{\varepsilon}, \mathbf{g}_{1}^{\varepsilon}, \mathbf{g}_{2}^{\varepsilon}]$ , we get that  $J_{\varepsilon}(\mathbf{s}^{\varepsilon}) \leq J_{\varepsilon}(\tilde{\mathbf{s}})$ ; thus, equality [\(67\)](#page-13-4) implies

<span id="page-14-1"></span>
$$
J_{\varepsilon}(\mathbf{s}^{\varepsilon}) \le J(\tilde{\mathbf{s}}). \tag{70}
$$

Observing that  $J(\mathbf{s}^{\varepsilon}) \geq 0$ , from [\(65\)](#page-13-2) we obtain

$$
\frac{1}{2} \|\mathbf{u}^{\varepsilon}-\tilde{\mathbf{u}}\|_{\mathbf{H}\sigma}^2 + \frac{1}{2} \|\mathbf{w}^{\varepsilon}-\tilde{\mathbf{w}}\|_{H^1}^2 + \frac{1}{2} \|\boldsymbol{g}_1^{\varepsilon}-\tilde{\boldsymbol{g}}_1\|_{\mathbf{H}^{1/2}(\Gamma_1)}^2 + \frac{1}{2} \|\boldsymbol{g}_2^{\varepsilon}-\tilde{\boldsymbol{g}}_2\|_{H^{1/2}(\Gamma_3)}^2 \leq J_{\varepsilon}(\mathbf{s}^{\varepsilon}).
$$

Therefore,

$$
\begin{split} \|\mathbf{u}^{\varepsilon}\|_{\mathbf{H}_{\sigma}}^{2}+\|\mathbf{w}^{\varepsilon}\|_{H^{1}}^{2}+\|g_{1}^{\varepsilon}\|_{\mathbf{H}^{1/2}(\Gamma_{1})}^{2}+\|g_{2}^{\varepsilon}\|_{H^{1/2}(\Gamma_{3})}^{2} \\ &\leq 2(\|\mathbf{u}^{\varepsilon}-\tilde{\mathbf{u}}\|_{\mathbf{H}_{\sigma}}^{2}+\|\mathbf{w}^{\varepsilon}-\tilde{\mathbf{w}}\|_{H^{1}}^{2}+\|g_{1}^{\varepsilon}-\tilde{g}_{1}\|_{\mathbf{H}^{1/2}(\Gamma_{1})}^{2}+\|g_{2}^{\varepsilon}-\tilde{g}_{2}\|_{H^{1/2}(\Gamma_{3})}^{2}) \\ &\quad+2(\|\tilde{\mathbf{u}}\|_{\mathbf{H}_{\sigma}}^{2}+\|\tilde{\mathbf{w}}\|_{H^{1}}^{2}+\|\tilde{g}_{1}\|_{\mathbf{H}^{1/2}(\Gamma_{1})}^{2}+\|\tilde{g}_{2}\|_{H^{1/2}(\Gamma_{3})}^{2}) \\ &\leq 4J_{\varepsilon}(\mathbf{s}^{\varepsilon})+2(\|\tilde{\mathbf{u}}\|_{\mathbf{H}_{\sigma}}^{2}+\|\tilde{\mathbf{w}}\|_{H^{1}}^{2}+\|\tilde{g}_{1}\|_{\mathbf{H}^{1/2}(\Gamma_{1})}^{2}+\|\tilde{g}_{2}\|_{H^{1/2}(\Gamma_{3})}^{2}), \end{split}
$$

and using [\(70\)](#page-14-1) we obtain

<span id="page-14-2"></span>
$$
\|\mathbf{u}^{\varepsilon}\|_{\mathbf{H}_{\sigma}}^{2} + \|\mathbf{w}^{\varepsilon}\|_{H^{1}}^{2} + \|g_{1}^{\varepsilon}\|_{\mathbf{H}^{1/2}(\Gamma_{1})}^{2} + \|g_{2}^{\varepsilon}\|_{H^{1/2}(\Gamma_{3})}^{2} \le 4J(\tilde{\mathbf{s}}) + C \le C,
$$
\n(71)

where C is a constant independent of  $\varepsilon$ . Thus, [\(71\)](#page-14-2) implies that there exists  $\mathbf{s} = [\mathbf{u}, \mathbf{w}, \mathbf{g}_1, g_2] \in \mathbb{X}$  and a subsequence of  $\{\mathbf{s}^{\varepsilon} = [\mathbf{u}^{\varepsilon}, \mathbf{w}^{\varepsilon}, \mathbf{g}^{\varepsilon}_1, \mathbf{g}^{\varepsilon}_2]\}_{\varepsilon > 0}$ , still denoted by  $\{\mathbf{s}^{\varepsilon}\}_{\varepsilon > 0}$ , such that

<span id="page-14-3"></span>
$$
[\mathbf{u}^{\varepsilon}, \mathbf{w}^{\varepsilon}, \mathbf{g}_1^{\varepsilon}, g_2^{\varepsilon}] \longrightarrow [\mathbf{u}, \mathbf{w}, \mathbf{g}_1, g_2] \text{ weakly in } \mathbb{X}, \text{ as } \varepsilon \to 0.
$$
 (72)

Since the embeddings  $H_{\sigma} \hookrightarrow L^2(\Omega)$ ,  $H^1(\Omega) \hookrightarrow L^2(\Omega)$ ,  $\mathcal{U}_1 \hookrightarrow L^2(\Gamma_1)$  and  $\mathcal{U}_2 \hookrightarrow L^2(\Gamma_3)$  are compact, we have

<span id="page-14-4"></span>
$$
[\mathbf{u}^{\varepsilon}, \mathbf{w}^{\varepsilon}, \mathbf{g}_1^{\varepsilon}, g_2^{\varepsilon}] \longrightarrow [\mathbf{u}, \mathbf{w}, \mathbf{g}_1, g_2] \text{ strongly in } \mathbf{L}^2(\Omega) \times L^2(\Omega) \times L^2(\Gamma_1) \times L^2(\Gamma_3). \tag{73}
$$

Following the proof of Theorem [3,](#page-12-0) by considering [\(72\)](#page-14-3) and [\(73\)](#page-14-4), we obtain that  $\mathbf{s} = [\mathbf{u}, \mathbf{w}, \mathbf{g}_1, g_2] \in \mathcal{S}_{ad}$ . Furthermore, since  $s^{\varepsilon} \in \mathbb{X}$ , by the definition of  $J_{\varepsilon}$  given in [\(65\)](#page-13-2), we obtain

$$
J(\mathbf{s}^{\varepsilon}) \leq J_{\varepsilon}(\mathbf{s}^{\varepsilon}),
$$

and by using [\(70\)](#page-14-1), we get

<span id="page-14-6"></span>
$$
J(\mathbf{s}^{\varepsilon}) \le J_{\varepsilon}(\mathbf{s}^{\varepsilon}) \le J(\tilde{\mathbf{s}}). \tag{74}
$$

Since  $s \in S_{ad}$  and J is weakly lower semicontinuous on  $S_{ad}$ , from [\(72\)](#page-14-3) and [\(73\)](#page-14-4) we obtain

$$
J[\mathbf{u}, \mathbf{w}, \mathbf{g}_1, g_2] \leq \lim_{\varepsilon \to 0} \inf J(\mathbf{s}^{\varepsilon}) \leq J[\tilde{\mathbf{u}}, \tilde{\mathbf{w}}, \tilde{\mathbf{g}}_1, \tilde{g}_2] \leq J[\mathbf{u}, \mathbf{w}, \mathbf{g}_1, g_2],
$$

and taking into account that  $\tilde{\mathbf{s}} = [\tilde{\mathbf{u}}, \tilde{\mathbf{w}}, \tilde{\mathbf{g}}_1, \tilde{g}_2]$  is an optimal solution for control problem [\(22\)](#page-6-1), we conclude that  $[\mathbf{u}, \mathbf{w}, \mathbf{g}_1, g_2] = [\tilde{\mathbf{u}}, \tilde{\mathbf{w}}, \tilde{\mathbf{g}}_1, g_2]$ . Then, from [\(72\)](#page-14-3)-[\(73\)](#page-14-4), as  $\varepsilon \to 0$ , we get

<span id="page-15-0"></span>
$$
[\mathbf{u}^{\varepsilon}, \mathbf{w}^{\varepsilon}, \mathbf{g}_1^{\varepsilon}, \mathbf{g}_2^{\varepsilon}] \longrightarrow [\tilde{\mathbf{u}}, \tilde{\mathbf{w}}, \tilde{\mathbf{g}}_1, \tilde{\mathbf{g}}_2] \text{ weakly in } \mathcal{S}_{ad} \subset \mathbb{X},
$$
\n(75)

$$
[\mathbf{u}^{\varepsilon}, \mathbf{w}^{\varepsilon}, \mathbf{g}_1^{\varepsilon}, \mathbf{g}_2^{\varepsilon}] \longrightarrow [\tilde{\mathbf{u}}, \tilde{\mathbf{w}}, \tilde{\mathbf{g}}_1, \tilde{\mathbf{g}}_2] \text{ strongly in } \mathbf{L}^2(\Omega) \times L^2(\Omega) \times L^2(\Gamma_1) \times L^2(\Gamma_3). \tag{76}
$$

Now, we observe that

 $\overline{\phantom{a}}$ 

<span id="page-15-1"></span>
$$
\|\mathbf{u}^{\varepsilon}\|_{\mathbf{H}_{\sigma}}^{2} - \|\tilde{\mathbf{u}}\|_{\mathbf{H}_{\sigma}}^{2} = \|\mathbf{u}^{\varepsilon}\|^{2} - \|\tilde{\mathbf{u}}\|^{2} + (\nabla \mathbf{u}^{\varepsilon}, \nabla \mathbf{u}^{\varepsilon} - \nabla \tilde{\mathbf{u}}) + (\nabla \tilde{\mathbf{u}}, \nabla \mathbf{u}^{\varepsilon} - \nabla \tilde{\mathbf{u}}),\tag{77}
$$

$$
|\mathbf{w}^{\varepsilon}\|_{H^1}^2 - \|\tilde{\mathbf{w}}\|_{H^1}^2 = \|\mathbf{w}^{\varepsilon}\|^2 - \|\tilde{\mathbf{w}}\|^2 + (\nabla \mathbf{w}^{\varepsilon}, \nabla \mathbf{w}^{\varepsilon} - \nabla \tilde{\mathbf{w}}) + (\nabla \tilde{\mathbf{w}}, \nabla \mathbf{w}^{\varepsilon} - \nabla \tilde{\mathbf{w}}),\tag{78}
$$

where  $\nabla \mathbf{u}^{\varepsilon}$ ,  $\nabla \tilde{\mathbf{u}} \in \mathbf{L}^2(\Omega)$  and  $\nabla \mathbf{w}^{\varepsilon}$ ,  $\nabla \tilde{\mathbf{w}} \in L^2(\Omega)$ . Therefore, from [\(75\)](#page-15-0)-[\(76\)](#page-15-0), as  $\varepsilon \to 0$ , we get

$$
(\nabla \mathbf{u}^{\varepsilon}, \nabla \mathbf{u}^{\varepsilon} - \nabla \tilde{\mathbf{u}}) + (\nabla \tilde{\mathbf{u}}, \nabla \mathbf{u}^{\varepsilon} - \nabla \tilde{\mathbf{u}}) \longrightarrow 0 \text{ and } ||\mathbf{u}^{\varepsilon}|| \longrightarrow ||\tilde{\mathbf{u}}||, (\nabla \mathbf{w}^{\varepsilon}, \nabla \mathbf{w}^{\varepsilon} - \nabla \tilde{\mathbf{w}}) + (\nabla \tilde{\mathbf{w}}, \nabla \mathbf{w}^{\varepsilon} - \nabla \tilde{\mathbf{w}}) \longrightarrow 0 \text{ and } ||\mathbf{w}^{\varepsilon}|| \longrightarrow ||\tilde{\mathbf{w}}||, \tag{80}
$$

which, together with  $(77)-(78)$  $(77)-(78)$ , implies that

$$
\|\mathbf{u}^{\varepsilon}\|_{\mathbf{H}_{\sigma}}^{2} \longrightarrow \|\tilde{\mathbf{u}}\|_{\mathbf{H}_{\sigma}}^{2} \text{ and } \|\mathbf{w}^{\varepsilon}\|_{H^{1}}^{2} \longrightarrow \|\tilde{\mathbf{w}}\|_{H^{1}}^{2}.
$$

In particular,

<span id="page-15-2"></span> $\limsup_{\varepsilon \to 0} \|\mathbf{u}^{\varepsilon}\|_{\mathbf{H}_{\sigma}} = \|\tilde{\mathbf{u}}\|_{\mathbf{H}_{\sigma}} \text{ and } \limsup_{\varepsilon \to 0} \|\mathbf{w}^{\varepsilon}\|_{H^{1}} = \|\tilde{\mathbf{w}}\|_{H^{1}}.$  (81)

Since  $H_{\sigma} \times H^{1}(\Omega)$  is a Hilbert space, it is uniformly convex; then from [\(81\)](#page-15-2) and Proposition 3.32 of [\[6\]](#page-22-12) p. 78, as  $\varepsilon \to 0$ , we obtain

<span id="page-15-5"></span>
$$
[\mathbf{u}^{\varepsilon}, \mathbf{w}^{\varepsilon}] \longrightarrow [\tilde{\mathbf{u}}, \tilde{\mathbf{w}}] \text{ strongly in } \mathbf{H}_{\sigma} \times H^{1}(\Omega). \tag{82}
$$

Now, we observe that

<span id="page-15-3"></span>
$$
\|\boldsymbol{g}_1^{\varepsilon}\|_{\mathbf{H}^{1/2}(\Gamma_1)}^2 - \|\tilde{\boldsymbol{g}}_1\|_{\mathbf{H}^{1/2}(\Gamma_1)}^2 = \langle \boldsymbol{g}_1^{\varepsilon} - \tilde{\boldsymbol{g}}_1, \boldsymbol{g}_1^{\varepsilon}\rangle_{\Gamma_1} + \langle \tilde{\boldsymbol{g}}_1, \boldsymbol{g}_1^{\varepsilon} - \tilde{\boldsymbol{g}}_1 \rangle_{\Gamma_1},
$$
\n(83)

$$
\|\hat{g}_2\|_{H^{1/2}(\Gamma_3)}^2 - \|\tilde{g}_2\|_{H^{1/2}(\Gamma_3)}^2 = \langle g_2^{\varepsilon} - \tilde{g}_2, g_2^{\varepsilon} \rangle_{\Gamma_3} + \langle \tilde{g}_2, g_2^{\varepsilon} - \tilde{g}_2 \rangle_{\Gamma_3}.
$$
 (84)

Taking into account that  $U_1 \subset \mathbf{H}_{00}^{1/2}(\Gamma_1) \subset \mathbf{L}^2(\Gamma_1)$  and  $U_2 \subset H_{00}^{1/2}(\Gamma_3) \subset L^2(\Gamma_3)$ , then from [\(75\)](#page-15-0)-[\(76\)](#page-15-0), as  $\varepsilon \to 0$ , we deduce that

$$
\langle g_1^{\varepsilon}-\tilde{g}_1, g_1^{\varepsilon}\rangle_{\Gamma_1}+\langle \tilde{g}_1, g_1^{\varepsilon}-\tilde{g}_1\rangle_{\Gamma_1}\to 0, \quad \langle g_2^{\varepsilon}-\tilde{g}_2, g_2^{\varepsilon}\rangle_{\Gamma_3}+\langle \tilde{g}_2, g_2^{\varepsilon}-\tilde{g}_2\rangle_{\Gamma_3}\to 0,
$$

which, together with  $(83)-(84)$  $(83)-(84)$ , implies that

$$
\|\mathbf{g}_1^{\varepsilon}\|_{\mathbf{H}^{1/2}(\Gamma_1)}^2 \to \|\tilde{\mathbf{g}}_1\|_{\mathbf{H}^{1/2}(\Gamma_1)}^2 \text{ and } \|\mathbf{g}_2^{\varepsilon}\|_{H^{1/2}(\Gamma_3)}^2 \to \|\tilde{g}_2\|_{H^{1/2}(\Gamma_3)}^2.
$$

In particular,

<span id="page-15-4"></span>
$$
\limsup_{\varepsilon \to 0} \|\boldsymbol{g}_1^{\varepsilon}\|_{\mathbf{H}^{1/2}(\Gamma \backslash \Gamma_2)} = \|\tilde{\boldsymbol{g}}_1\|_{\mathbf{H}^{1/2}(\Gamma \backslash \Gamma_2)} \text{ and } \limsup_{\varepsilon \to 0} \|\boldsymbol{g}_2^{\varepsilon}\|_{H^{1/2}(\Gamma)} = \|\tilde{g}_2\|_{H^{1/2}(\Gamma)}. \tag{85}
$$

Since  $\mathcal{U}_1 \times \mathcal{U}_2$  is a closed set of  $\mathbf{H}_{00}^{1/2}(\Gamma_1) \times H^{1/2}(\Gamma_3)$ , from [\(85\)](#page-15-4) and Proposition 3.32 of [\[6\]](#page-22-12) p.78, as  $\varepsilon \to 0$ , we obtain that

<span id="page-15-6"></span>
$$
[\mathbf{g}_1^{\varepsilon}, g_2^{\varepsilon}] \longrightarrow [\tilde{\mathbf{g}}_1, \tilde{g}_2] \text{ strongly in } \mathcal{U}_1 \times \mathcal{U}_2. \tag{86}
$$

Therefore, from [\(82\)](#page-15-5) and [\(86\)](#page-15-6), we conclude [\(68\)](#page-14-5).

In order to prove [\(69\)](#page-14-5), from [\(74\)](#page-14-6)-[\(75\)](#page-15-0) and, since that J is weakly lower semicontinuous on  $S_{ad}$ , we obtain

$$
J[\tilde{\mathbf{u}}, \tilde{\mathbf{w}}, \tilde{\mathbf{g}}_1, \tilde{g}_2] \le \liminf_{\varepsilon \to 0} J(\mathbf{s}^{\varepsilon}) \le \limsup_{\varepsilon \to 0} J(\mathbf{s}^{\varepsilon}) \le J[\tilde{\mathbf{u}}, \tilde{\mathbf{w}}, \tilde{\mathbf{g}}_1, \tilde{g}_2],
$$

which implies

<span id="page-16-1"></span>
$$
\lim_{\varepsilon \to 0} J(\mathbf{s}^{\varepsilon}) = J[\tilde{\mathbf{u}}, \tilde{\mathbf{w}}, \tilde{\mathbf{g}}_1, \tilde{g}_2].
$$
\n(87)

Then, from [\(74\)](#page-14-6) and [\(87\)](#page-16-1), we obtain

$$
J[\tilde{\mathbf{u}}, \tilde{\mathbf{w}}, \tilde{\boldsymbol{g}}_1, \tilde{g}_2] = \lim_{\varepsilon \to 0} J[\mathbf{u}^{\varepsilon}, \mathbf{w}^{\varepsilon}, \boldsymbol{g}_1^{\varepsilon}, \tilde{g}_2^{\varepsilon}] \leq \lim_{\varepsilon \to 0} J_{\varepsilon}[\mathbf{u}^{\varepsilon}, \mathbf{w}^{\varepsilon}, \boldsymbol{g}_1^{\varepsilon}, \tilde{g}_2^{\varepsilon}] \leq J[\tilde{\mathbf{u}}, \tilde{\mathbf{w}}, \tilde{\boldsymbol{g}}_1, \tilde{g}_2],
$$

which implies [\(69\)](#page-14-5).  $\Diamond$ 

#### 5.2 Existence of Lagrange Multipliers and Adjoint Equations

For simplicity, we consider the following operators

$$
\begin{cases}\nK: \mathbf{H}_{\sigma} \times \mathbf{H}_{\sigma} \times \mathbf{H}_{\sigma} \times \tilde{\mathbf{H}}_{\sigma} \to \mathbf{H}', & \tilde{K}: \mathbf{H}_{\sigma} \times H^{1}(\Omega) \times H^{1}_{0}(\Omega) \to \mathbf{H}', \nB^{T}: \mathbf{H}_{\sigma} \times \mathbf{H}_{\sigma} \times \tilde{\mathbf{H}}_{\sigma} \to \mathbf{H}', & E: \mathbf{H}_{\sigma} \times H^{1}(\Omega) \times H^{1}_{0}(\Omega) \to \mathbf{H}', \n\tilde{E}: \mathbf{H}_{\sigma} \times \mathbf{H}_{\sigma} \times H^{1}_{0}(\Omega) \to (H^{1}(\Omega))', & F': \mathbf{H}_{\sigma} \times \tilde{\mathbf{H}}_{\sigma} \to \mathbf{H}', \nG': \mathbf{H}_{\sigma} \times H^{1}_{0}(\Omega) \to \mathbf{H}',\n\end{cases}
$$
\n(88)

defined by

<span id="page-16-4"></span><span id="page-16-0"></span>
$$
\begin{cases}\n\langle K[\mathbf{u}, \mathbf{u}, \mathbf{u}, \lambda], \mathbf{v}\rangle = (\eta'(N\mathbf{u})(N\mathbf{v})\mathbf{u}\cdot\nabla\mathbf{u}, \lambda) \quad \forall \mathbf{v} \in \mathbf{H}_{\sigma}, \\
\langle \tilde{K}[\mathbf{u}, \mathbf{u}, \mathbf{w}, \kappa], \mathbf{v}\rangle = (\eta'(N\mathbf{u})(N\mathbf{v})\mathbf{u}\cdot\nabla\mathbf{w}, \kappa) \quad \forall \mathbf{v} \in \mathbf{H}_{\sigma}, \\
\langle B^T[\mathbf{u}, \mathbf{u}, \lambda], \mathbf{v}\rangle = (\eta(N\mathbf{u})(\mathbf{u}\cdot\nabla\mathbf{v} + \mathbf{v}\cdot\nabla\mathbf{u}), \lambda) \quad \forall \mathbf{v} \in \mathbf{H}_{\sigma}, \\
\langle E[\mathbf{u}, \mathbf{w}, \kappa], \mathbf{v}\rangle = (\eta(N\mathbf{u})\mathbf{v}\cdot\nabla\mathbf{w}, \kappa) \quad \forall \mathbf{v} \in \mathbf{H}_{\sigma}, \\
(\tilde{E}[\mathbf{u}, \mathbf{u}, \kappa], \mathbf{z}) = (\eta(N\mathbf{u})\mathbf{u}\cdot\nabla\mathbf{z}, \kappa) \quad \forall \mathbf{z} \in H^1(\Omega), \\
\langle F'[\mathbf{u}, \lambda], \mathbf{v}\rangle = (\eta'(N\mathbf{u})(N\mathbf{v})\mathbf{f}, \lambda) \quad \forall \mathbf{v} \in \mathbf{H}_{\sigma}, \\
\langle G'[\mathbf{u}, \kappa], \mathbf{v}\rangle = (\eta'(N\mathbf{u})(N\mathbf{v})\mathbf{g}, \kappa) \quad \forall \mathbf{v} \in \mathbf{H}_{\sigma},\n\end{cases}
$$
\n(89)

where  $\eta'$  denotes the first derivative of  $\eta$ .

**Theorem 5** Let  $f \in L^2(\Omega)$ ,  $g \in L^2(\Omega)$ ,  $u_0 \in H_{00}^{1/2}(\Gamma_0)$ ,  $g_1 \in \mathcal{U}_1$ ,  $w_0 \in H_{00}^{1/2}(\Gamma_0)$ ,  $g_2 \in \mathcal{U}_2$  and  $\eta \in C^1(\mathbb{R})$ . Then, for any optimal solution  $[\mathbf{u}^{\varepsilon}, \mathbf{w}^{\varepsilon}, \mathbf{g}^{\varepsilon}_1, g^{\varepsilon}_2] \in \mathbb{X}$  of problem  $(64)$ - $(65)$  there exist Lagrange multipliers  $[\lambda^{\varepsilon}, \phi^{\varepsilon}, \xi^{\varepsilon}, \vartheta^{\varepsilon}] \in \tilde{\mathbf{H}}_{\sigma} \times H_0^1(\Omega) \times \mathbf{H}_{00}^{-1/2}(\Gamma \setminus \Gamma_2) \times H^{-1/2}(\Gamma)$  given by

<span id="page-16-2"></span>
$$
\lambda^{\varepsilon} = \frac{1}{\varepsilon} A^{-1} (\mu_1 A \mathbf{u}^{\varepsilon} + B[\mathbf{u}^{\varepsilon}, \mathbf{u}^{\varepsilon}, \mathbf{u}^{\varepsilon}] - 2\mu_r \text{rot } \mathbf{w}^{\varepsilon} - F(\mathbf{u}^{\varepsilon})) \in \tilde{\mathbf{H}}_{\sigma},\tag{90}
$$

$$
\phi^{\varepsilon} = \frac{1}{\varepsilon} \tilde{A}^{-1} (\mu_2 \tilde{A} \mathbf{w}^{\varepsilon} + \tilde{B}[\mathbf{u}^{\varepsilon}, \mathbf{u}^{\varepsilon}, \mathbf{w}^{\varepsilon}] + 4\mu_r \mathbf{w}^{\varepsilon} - 2\mu_r \text{rot } \mathbf{u}^{\varepsilon} - G(\mathbf{u}^{\varepsilon})) \in H_0^1(\Omega), \tag{91}
$$

$$
\boldsymbol{\xi}^{\varepsilon} = \frac{1}{\varepsilon} (\mathbf{u}_{|\Gamma \backslash \Gamma_2}^{\varepsilon} - \mathbf{u}_{\boldsymbol{g}_1^{\varepsilon}}) \in \mathbf{L}^2(\Gamma \setminus \Gamma_2) \subset \mathbf{H}_{00}^{-1/2}(\Gamma \setminus \Gamma_2), \tag{92}
$$

$$
\vartheta^{\varepsilon} = \frac{1}{\varepsilon} (\mathbf{w}_{|\Gamma}^{\varepsilon} - \mathbf{w}_{g_2^{\varepsilon}}) \in L^2(\Gamma) \subset H^{-1/2}(\Gamma), \tag{93}
$$

*which satisfies the following system*

<span id="page-16-3"></span>
$$
\mu_1 \langle A\mathbf{\lambda}^{\varepsilon}, \mathbf{v} \rangle + \langle K[\mathbf{u}^{\varepsilon}, \mathbf{u}^{\varepsilon}, \mathbf{u}^{\varepsilon}, \mathbf{\lambda}^{\varepsilon}], \mathbf{v} \rangle + \langle B^T[\mathbf{u}^{\varepsilon}, \mathbf{u}^{\varepsilon}, \mathbf{\lambda}^{\varepsilon}], \mathbf{v} \rangle + \langle \tilde{K}[\mathbf{u}^{\varepsilon}, \mathbf{u}^{\varepsilon}, \mathbf{v}^{\varepsilon}, \phi^{\varepsilon}], \mathbf{v} \rangle \n+ \langle E[\mathbf{u}^{\varepsilon}, \mathbf{w}^{\varepsilon}, \phi^{\varepsilon}], \mathbf{v} \rangle + \langle J'_{\mathbf{u}}(\mathbf{u}^{\varepsilon}), \mathbf{v} \rangle + \langle \xi^{\varepsilon}, \mathbf{v} \rangle_{\Gamma \setminus \Gamma_2} + (\mathbf{u}^{\varepsilon} - \tilde{\mathbf{u}}, \mathbf{v})_{\mathbf{H}_{\sigma}} \n= 2\mu_r(\text{rot}\,\phi^{\varepsilon}, \mathbf{v}) + \langle F'[\mathbf{u}^{\varepsilon}, \mathbf{\lambda}^{\varepsilon}], \mathbf{v} \rangle + \langle G'[\mathbf{u}^{\varepsilon}, \phi^{\varepsilon}], \mathbf{v} \rangle \quad \forall \mathbf{v} \in \mathbf{H}_{\sigma},
$$
\n(94)

$$
\mu_2(\tilde{A}\phi^{\varepsilon},z) + \langle \tilde{E}[\mathbf{u}^{\varepsilon},\mathbf{u}^{\varepsilon},\phi^{\varepsilon}],z\rangle + 4\mu_r(\phi^{\varepsilon},z) + (\mathbf{w}^{\varepsilon} - \tilde{\mathbf{w}},z)_{H^1} + \beta_3(\mathbf{w}^{\varepsilon} - \mathbf{w}_d,z) + \langle \vartheta^{\varepsilon},z \rangle_{\Gamma} \n= 2\mu_r(\text{rot}\,\mathbf{\lambda}^{\varepsilon},z) \quad \forall z \in H^1(\Omega),
$$
\n(95)

$$
\beta_5 \langle \boldsymbol{g}_1^{\varepsilon}, \boldsymbol{g}_1 - \boldsymbol{g}_1^{\varepsilon} \rangle_{\Gamma_1} + \langle \boldsymbol{g}_1^{\varepsilon} - \tilde{\boldsymbol{g}}_1, \boldsymbol{g}_1 - \boldsymbol{g}_1^{\varepsilon} \rangle_{\Gamma_1} - \langle \boldsymbol{\xi}^{\varepsilon}, \boldsymbol{g}_1 - \boldsymbol{g}_1^{\varepsilon} \rangle_{\Gamma_1} \ge 0 \quad \forall \boldsymbol{g}_1 \in \mathcal{U}_1,
$$
\n(96)

$$
\beta_6 \langle g_2^{\varepsilon}, g_2 - g_2^{\varepsilon} \rangle_{\Gamma_3} + \langle g_2^{\varepsilon} - \tilde{g}_2, g_2 - g_2^{\varepsilon} \rangle_{\Gamma_3} - \langle \vartheta^{\varepsilon}, g_2 - g_2^{\varepsilon} \rangle_{\Gamma_3} \ge 0 \quad \forall g_2 \in \mathcal{U}_2,
$$
\n
$$
(97)
$$

*where*

<span id="page-17-4"></span>
$$
\langle J'_{\mathbf{u}}(\mathbf{u}^{\varepsilon}), \mathbf{v} \rangle := \beta_1(\text{rot } \mathbf{u}^{\varepsilon}, \text{rot } \mathbf{v}) + \beta_2(\mathbf{u}^{\varepsilon} - \mathbf{u}_d, \mathbf{v}) + \beta_4(\eta(N\mathbf{u}^{\varepsilon}) - \rho_d, \eta'(N\mathbf{u}^{\varepsilon})N\mathbf{v}). \tag{98}
$$

*Moreover, there exists a constant*  $C > 0$ *, independent of*  $\varepsilon$ *, such that* 

<span id="page-17-1"></span>
$$
\|\boldsymbol{\xi}^{\varepsilon}\|_{\mathbf{H}_{00}^{-1/2}(\Gamma\backslash\Gamma_{2})} \leq C + C \|\left[\boldsymbol{\lambda}^{\varepsilon},\phi^{\varepsilon}\right]\|_{\tilde{\mathbf{H}}_{\sigma}\times H_{0}^{1}},\tag{99}
$$

$$
\|\vartheta^{\varepsilon}\|_{H^{-1/2}(\Gamma)} \leq C + C \|\left[\mathbf{\lambda}^{\varepsilon}, \phi^{\varepsilon}\right]\|_{\mathbf{\tilde{H}}_{\sigma} \times H_0^1}.
$$
\n(100)

*Proof.* We introduce the function  $\mathcal{F} : [0,1] \times [0,1] \times [0,1] \times [0,1] \longrightarrow \mathbb{R}$  defined by

$$
\mathcal{F}[\zeta_1, \zeta_2, \zeta_3, \zeta_4] = J_{\varepsilon}[\mathbf{u}^{\varepsilon} + \zeta_1 \mathbf{v}, \mathbf{w}^{\varepsilon} + \zeta_2 \mathbf{z}, \mathbf{g}_1^{\varepsilon} + \zeta_3 (\mathbf{g}_1 - \mathbf{g}_1^{\varepsilon}), \mathbf{g}_2^{\varepsilon} + \zeta_4 (\mathbf{g}_2 - \mathbf{g}_2^{\varepsilon})],
$$
(101)

where  $[\mathbf{v}, \mathbf{z}, \mathbf{g}_1, g_2] \in \mathbb{X}$ .

Since the function F attains its minimum at  $\mathbf{0} = [0, 0, 0, 0]$  and  $\mathcal{U}_1 \times \mathcal{U}_2$  is convex, we have

<span id="page-17-0"></span>
$$
\frac{\partial \mathcal{F}}{\partial \zeta_1}(\mathbf{0}) = 0, \quad \frac{\partial \mathcal{F}}{\partial \zeta_2}(\mathbf{0}) = 0, \quad \frac{\partial \mathcal{F}}{\partial \zeta_3}(\mathbf{0}) \ge 0, \quad \frac{\partial \mathcal{F}}{\partial \zeta_4}(\mathbf{0}) \ge 0.
$$
\n(102)

Therefore, from [\(102\)](#page-17-0), and the definitions of  $\lambda^{\varepsilon}, \phi^{\varepsilon}, \xi^{\varepsilon}, \vartheta^{\varepsilon}$  given in [\(90\)](#page-16-2)-[\(93\)](#page-16-2), we obtain the system [\(94\)](#page-16-3)-[\(97\)](#page-16-3). Now we will prove inequality [\(99\)](#page-17-1). From [\(94\)](#page-16-3) we get

<span id="page-17-2"></span>
$$
|\langle \xi^{\varepsilon}, \mathbf{v} \rangle_{\Gamma \backslash \Gamma_{2}}| \leq \mu_{1} |\langle A\lambda^{\varepsilon}, \mathbf{v} \rangle| + |\langle K[\mathbf{u}^{\varepsilon}, \mathbf{u}^{\varepsilon}, \mathbf{u}^{\varepsilon}, \lambda^{\varepsilon}], \mathbf{v} \rangle| + |\langle B^{T}[\mathbf{u}^{\varepsilon}, \mathbf{u}^{\varepsilon}, \lambda^{\varepsilon}], \mathbf{v} \rangle| + |\langle \tilde{K}[\mathbf{u}^{\varepsilon}, \mathbf{u}^{\varepsilon}, \mathbf{w}^{\varepsilon}, \phi^{\varepsilon}], \mathbf{v} \rangle| + |\langle E[\mathbf{u}^{\varepsilon}, \mathbf{w}^{\varepsilon}, \phi^{\varepsilon}], \mathbf{v} \rangle| + |\langle J'_{\mathbf{u}}(\mathbf{u}^{\varepsilon}), \mathbf{v} \rangle| + |(\mathbf{u}^{\varepsilon} - \tilde{\mathbf{u}}, \mathbf{v})_{\mathbf{H}_{\sigma}}| + 2\mu_{r}|(\text{rot}\,\phi^{\varepsilon}, \mathbf{v})| + |\langle F'[\mathbf{u}^{\varepsilon}, \lambda^{\varepsilon}], \mathbf{v} \rangle| + |\langle G'[\mathbf{u}^{\varepsilon}, \phi^{\varepsilon}], \mathbf{v} \rangle|.
$$
\n(103)

We shall find bounds for the terms on right hand side of  $(103)$ . By using the Hölder inequality and observing that  $\|\mathbf{v}\|_{\mathbf{H}_{\sigma}} \leq C \|\mathbf{v}\|_{\tilde{\mathbf{H}}_{\sigma}}$ , we obtain

<span id="page-17-5"></span>
$$
\mu_1|\langle A\mathbf{\lambda}^{\varepsilon}, \mathbf{v}\rangle| \leq 2\mu_1|(D(\mathbf{\lambda}^{\varepsilon}), D(\mathbf{v}))| + 2\alpha\mu_1 \left| \int_{\Gamma_2} \mathbf{\lambda}^{\varepsilon} \cdot \mathbf{v} d\Gamma \right|
$$
  
\n
$$
\leq 2\mu_1 \|D(\mathbf{\lambda}^{\varepsilon})\| \|D(\mathbf{v})\| + 2\alpha\mu_1 \|\mathbf{\lambda}^{\varepsilon}\|_{\mathbf{L}^2(\Gamma)} \|\mathbf{v}\|_{\mathbf{L}^2(\Gamma)}
$$
  
\n
$$
\leq C \|\mathbf{v}\|_{\mathbf{H}_{\sigma}} \|\mathbf{\lambda}^{\varepsilon}\|_{\tilde{\mathbf{H}}_{\sigma}} + 2\mu_1 \alpha C \|\mathbf{\lambda}^{\varepsilon}\|_{\tilde{\mathbf{H}}_{\sigma}} \|\mathbf{v}\|_{\mathbf{H}_{\sigma}} \leq C \|\mathbf{v}\|_{\mathbf{H}_{\sigma}} \|\mathbf{\lambda}^{\varepsilon}\|_{\tilde{\mathbf{H}}_{\sigma}}, \qquad (104)
$$
  
\n
$$
2\mu_r |(\mathrm{rot}\,\phi^{\varepsilon}, \mathbf{v})| \leq 2\mu_r |\mathrm{rot}\,\phi^{\varepsilon}| \|\mathbf{v}\| \leq C\mu_r \|\phi^{\varepsilon}\|_{H_0^1} \|\mathbf{v}\|_{\mathbf{H}_{\sigma}} \leq C \|\mathbf{v}\|_{\mathbf{H}_{\sigma}} \|\phi^{\varepsilon}\|_{H_0^1}.
$$

Now, by using the Hölder inequality and the fact that  $\|\mathbf{u}^{\varepsilon}\|_{\mathbf{H}_{\sigma}} \leq C$ , where C is independent of  $\varepsilon$ , we obtain

<span id="page-17-6"></span>
$$
|(\mathbf{u}^{\varepsilon} - \tilde{\mathbf{u}}, \mathbf{v})_{\mathbf{H}_{\sigma}}| \leq C \|\mathbf{u}^{\varepsilon} - \tilde{\mathbf{u}}\|_{\mathbf{H}_{\sigma}} \|\mathbf{v}\|_{\mathbf{H}_{\sigma}} \leq C \|\mathbf{v}\|_{\mathbf{H}_{\sigma}}.
$$
\n(106)

From [\(68\)](#page-14-5), the definition of N and  $\eta$  given in Subsection [2.1,](#page-3-0) we deduce that

<span id="page-17-3"></span>
$$
\|\eta(N(\mathbf{u}^{\varepsilon})\|_{\infty} \leq C, \ \|\eta'(N(\mathbf{u}^{\varepsilon})\|_{\infty} \leq C, \ \|N\mathbf{v}\|_{\infty} \leq C\|N\mathbf{v}\|_{H^{2}} \leq C\|\mathbf{v}\|_{\mathbf{H}_{\sigma}},\tag{107}
$$

where C is a constant independent of  $\varepsilon$ . By the Hölder inequality, [\(107\)](#page-17-3) and the definition of  $\langle J'_{\mathbf{u}}(\mathbf{u}^{\varepsilon}), \mathbf{v} \rangle$ given in [\(98\)](#page-17-4), we obtain

<span id="page-17-7"></span>
$$
\begin{array}{rcl}\n|\langle J'_{\mathbf{u}}(\mathbf{u}^{\varepsilon}), \mathbf{v} \rangle| & \leq & \beta_{1}|\langle \text{rot } \mathbf{u}^{\varepsilon}, \text{rot } \mathbf{v} \rangle| + \beta_{2}|\langle \mathbf{u}^{\varepsilon} - \mathbf{u}_{d}, \mathbf{v} \rangle| + \beta_{4}|\langle \eta(N(\mathbf{u}^{\varepsilon})) - \rho_{d}, \eta'(N(\mathbf{u}^{\varepsilon}))N\mathbf{v} \rangle| \\
& \leq & \beta_{1}\|\text{rot } \mathbf{u}^{\varepsilon}\|\|\text{rot } \mathbf{v}\| + \beta_{2}\|\mathbf{u}^{\varepsilon} - \mathbf{u}_{d}\|\|\mathbf{v}\| + \beta_{4}\|\eta(N\mathbf{u}^{\varepsilon}) - \rho_{d}\|_{\infty}\|\eta'(N\mathbf{u}^{\varepsilon})\|\|N\mathbf{v}\| \\
& \leq & \beta_{1}C\|\mathbf{v}\|_{\mathbf{H}_{\sigma}} + \beta_{2}C\|\mathbf{v}\|_{\mathbf{H}_{\sigma}} + \beta_{4}C\|\mathbf{v}\|_{\mathbf{H}_{\sigma}} \leq & C\|\mathbf{v}\|_{\mathbf{H}_{\sigma}}.\n\end{array} \tag{108}
$$

Also, by the Hölder and Poincaré inequalities, [\(107\)](#page-17-3) and the definition of the operators  $B^T$ , E, K,  $\tilde{K}$ , F', and  $G'$  given in [\(89\)](#page-16-4), we obtain

<span id="page-18-2"></span>
$$
|\langle B^{T}[\mathbf{u}^{\varepsilon}, \mathbf{u}^{\varepsilon}, \boldsymbol{\lambda}^{\varepsilon}], \mathbf{v}\rangle| = |\langle B[\mathbf{u}^{\varepsilon}, \mathbf{u}^{\varepsilon}, \mathbf{v}] + B[\mathbf{u}^{\varepsilon}, \mathbf{v}, \mathbf{u}^{\varepsilon}], \boldsymbol{\lambda}^{\varepsilon}\rangle|
$$
  
\n
$$
\leq |(\eta(N\mathbf{u}^{\varepsilon})\mathbf{u}^{\varepsilon} \cdot \nabla \mathbf{v}, \boldsymbol{\lambda}^{\varepsilon})| + |(\eta(N\mathbf{u}^{\varepsilon})\mathbf{v} \cdot \nabla \mathbf{u}^{\varepsilon}, \boldsymbol{\lambda}^{\varepsilon})|
$$
  
\n
$$
\leq ||\eta(N\mathbf{u}^{\varepsilon})||_{\infty} (||\mathbf{u}^{\varepsilon}||_{3}||\nabla \mathbf{v}|| + ||\mathbf{v}||_{3}||\nabla \mathbf{u}^{\varepsilon}||) ||\boldsymbol{\lambda}^{\varepsilon}||_{6}
$$
  
\n
$$
\leq C||\mathbf{u}^{\varepsilon}||_{\mathbf{H}_{\sigma}} ||\mathbf{v}||_{\mathbf{H}_{\sigma}} ||\boldsymbol{\lambda}^{\varepsilon}||_{\tilde{\mathbf{H}}_{\sigma}} \leq C||\mathbf{v}||_{\mathbf{H}_{\sigma}} ||\boldsymbol{\lambda}^{\varepsilon}||_{\tilde{\mathbf{H}}_{\sigma}}, \qquad (109)
$$

$$
|\langle E[\mathbf{u}^{\varepsilon}, \mathbf{w}^{\varepsilon}, \phi^{\varepsilon}], \mathbf{v}\rangle| = |(\eta(N\mathbf{u}^{\varepsilon})\mathbf{v} \cdot \nabla \mathbf{w}^{\varepsilon}, \phi^{\varepsilon})|
$$
  
\n
$$
\leq \|\eta(N\mathbf{u}^{\varepsilon})\|_{\infty} \|\mathbf{v}\|_{3} \|\nabla \mathbf{w}^{\varepsilon}\| \|\phi^{\varepsilon}\|_{6}
$$
  
\n
$$
\leq C \|\mathbf{v}\|_{\mathbf{H}_{\sigma}} \|\mathbf{w}^{\varepsilon}\|_{H^{1}} \|\phi^{\varepsilon}\|_{H^{1}_{0}} \leq C \|\mathbf{v}\|_{\mathbf{H}_{\sigma}} \|\phi^{\varepsilon}\|_{H^{1}_{0}}, \tag{110}
$$

$$
|\langle K[\mathbf{u}^{\varepsilon}, \mathbf{u}^{\varepsilon}, \mathbf{u}^{\varepsilon}, \boldsymbol{\lambda}^{\varepsilon}], \mathbf{v}\rangle| = |(\eta'(N\mathbf{u}^{\varepsilon})(N\mathbf{v})\mathbf{u}^{\varepsilon} \cdot \nabla \mathbf{u}^{\varepsilon}, \boldsymbol{\lambda}^{\varepsilon})|
$$
  
\n
$$
\leq \|\eta'(N\mathbf{u}^{\varepsilon})\|_{\infty} \|N\mathbf{v}\|_{\infty} \|\mathbf{u}^{\varepsilon}\|_{3} \|\nabla \mathbf{u}^{\varepsilon}\| \|\boldsymbol{\lambda}^{\varepsilon}\|_{6}
$$
  
\n
$$
\leq C \|\mathbf{v}\|_{\mathbf{H}_{\sigma}} \|\mathbf{u}^{\varepsilon}\|_{\mathbf{H}_{\sigma}}^2 \|\boldsymbol{\lambda}^{\varepsilon}\|_{\mathbf{H}_{\sigma}} \leq C \|\mathbf{v}\|_{\mathbf{H}_{\sigma}} \|\boldsymbol{\lambda}^{\varepsilon}\|_{\mathbf{H}_{\sigma}}, \tag{111}
$$

$$
\begin{array}{rcl}\n|\langle \tilde{K}[\mathbf{u}^{\varepsilon}, \mathbf{u}^{\varepsilon}, \mathbf{w}^{\varepsilon}, \phi^{\varepsilon}], \mathbf{v}\rangle| & = & |(\eta'(N\mathbf{u}^{\varepsilon})(N\mathbf{v})\mathbf{u}^{\varepsilon} \cdot \nabla \mathbf{w}^{\varepsilon}, \phi^{\varepsilon})| \\
& \leq & ||\eta'(N\mathbf{u}^{\varepsilon})||_{\infty} ||N\mathbf{v}||_{\infty} ||\mathbf{u}^{\varepsilon}||_{3} ||\nabla \mathbf{w}^{\varepsilon}|| ||\phi^{\varepsilon}||_{6} \\
& \leq & C ||\mathbf{v}||_{\mathbf{H}_{\sigma}} ||\mathbf{u}^{\varepsilon}||_{\mathbf{H}_{\sigma}} ||\mathbf{w}^{\varepsilon}||_{H^{1}} ||\phi^{\varepsilon}||_{H_{0}^{1}} \leq & C ||\mathbf{v}||_{\mathbf{H}_{\sigma}} ||\phi^{\varepsilon}||_{H_{0}^{1}},\n\end{array} \tag{112}
$$

$$
\begin{array}{rcl}\n|\langle F'[\mathbf{u}^{\varepsilon}, \boldsymbol{\lambda}^{\varepsilon}], \mathbf{v} \rangle| & = & |(\eta' (N \mathbf{u}^{\varepsilon}) (N \mathbf{v}) \mathbf{f}, \boldsymbol{\lambda}^{\varepsilon})| \\
& \leq & \|\eta' (N \mathbf{u}^{\varepsilon})\|_{\infty} \|N \mathbf{v}\|_{\infty} \| \mathbf{f} \| \|\boldsymbol{\lambda}^{\varepsilon}\| \leq C \| \mathbf{v} \|_{\mathbf{H}_{\sigma}} \|\boldsymbol{\lambda}^{\varepsilon}\|_{\tilde{\mathbf{H}}_{\sigma}},\n\end{array} \tag{113}
$$

<span id="page-18-1"></span>
$$
|\langle G'[\mathbf{u}^{\varepsilon},\phi^{\varepsilon}],\mathbf{v}\rangle| = |(\eta'(N\mathbf{u}^{\varepsilon})(N\mathbf{v})g,\phi^{\varepsilon})|
$$
  
\n
$$
\leq \|\eta'(N\mathbf{u}^{\varepsilon})\|_{\infty} \|N\mathbf{v}\|_{\infty} \|g\| \|\phi^{\varepsilon}\| \leq C \|\mathbf{v}\|_{\mathbf{H}_{\sigma}} \|\phi^{\varepsilon}\|_{H_{0}^{1}}.
$$
\n(114)

By substituting inequalities  $(104)-(106)$  $(104)-(106)$  and  $(108)-(114)$  $(108)-(114)$  in  $(103)$ , we obtain

$$
|\langle \xi^\varepsilon, \mathbf{v} \rangle_{\Gamma \backslash \Gamma_2}| \leq \ C \|\mathbf{v}\|_{\mathbf{H}_\sigma} + C \|\mathbf{v}\|_{\mathbf{H}_\sigma} (\|\boldsymbol{\lambda}^\varepsilon\|_{\tilde{\mathbf{H}}_\sigma} + \|\phi^\varepsilon\|_{H_0^1}),
$$

and then, we deduce that

<span id="page-18-0"></span>
$$
\|\pmb \xi^\varepsilon\|_{{\bf H}_{00}^{-1/2}(\Gamma\backslash \Gamma_2)}\leq C+C(\|\pmb \lambda^\varepsilon\|_{\tilde{\bf H}_\sigma}+\|\phi^\varepsilon\|_{H_0^1})\leq C+\sqrt{2}C\|[\pmb \lambda^\varepsilon,\phi^\varepsilon]\|_{\tilde{\bf H}_\sigma\times H_0^1},
$$

which implies [\(99\)](#page-17-1). Analogously we can obtain [\(100\)](#page-17-1). ⋄

#### 5.3 Optimality System

This subsection is devoted to obtain an optimality system to problem [\(22\)](#page-6-1). We first show the existence of Lagrange multipliers.

**Theorem 6** Let  $f \in L^2(\Omega)$ ,  $g \in L^2(\Omega)$ ,  $\mathbf{u}_0 \in \mathbf{H}_{00}^{1/2}(\Gamma_0)$ ,  $g_1 \in \mathcal{U}_1$ ,  $w_0 \in H_{00}^{1/2}(\Gamma_0)$ ,  $g_2 \in \mathcal{U}_2$  and  $\eta \in C^1(\mathbb{R})$ . *Then, for any optimal solution*  $\tilde{\mathbf{s}} = [\tilde{\mathbf{u}}, \tilde{\mathbf{w}}, \tilde{\mathbf{g}}_1, \tilde{g}_2] \in \mathcal{S}_{ad}$  *of problem [\(22\)](#page-6-1) there exist Lagrange multipliers*  $[\lambda_0, \lambda, \phi, \xi, \vartheta] \in (\mathbb{R}^+ \cup \{0\}) \times \tilde{H}_{\sigma} \times H_0^1(\Omega) \times H_{00}^{-1/2}(\Gamma \setminus \Gamma_2) \times H^{-1/2}(\Gamma),$  *not all zero, satisfying the following system:*

<span id="page-19-4"></span>
$$
\mu_1 \langle A \mathbf{\lambda}, \mathbf{v} \rangle + \langle K[\tilde{\mathbf{u}}, \tilde{\mathbf{u}}, \tilde{\mathbf{u}}, \mathbf{\lambda}], \mathbf{v} \rangle + \langle B^T[\tilde{\mathbf{u}}, \tilde{\mathbf{u}}, \mathbf{\lambda}], \mathbf{v} \rangle + \langle \tilde{K}[\tilde{\mathbf{u}}, \tilde{\mathbf{u}}, \tilde{\mathbf{w}}, \phi], \mathbf{v} \rangle + \langle E[\tilde{\mathbf{u}}, \tilde{\mathbf{w}}, \phi], \mathbf{v} \rangle + \lambda_0 \langle J'_{\mathbf{u}}(\tilde{\mathbf{u}}), \mathbf{v} \rangle + \langle \xi, \mathbf{v} \rangle_{\Gamma \setminus \Gamma_2} - 2\mu_r(\text{rot}\phi, \mathbf{v}) = \langle F'[\tilde{\mathbf{u}}, \mathbf{\lambda}], \mathbf{v} \rangle + \langle G'[\tilde{\mathbf{u}}, \phi], \mathbf{v} \rangle \quad \forall \mathbf{v} \in \mathbf{H}_\sigma,
$$
 (115)

$$
\mu_2\langle\tilde{A}\phi,z\rangle+\langle\tilde{E}[\tilde{\mathbf{u}},\tilde{\mathbf{u}},\phi],\text{z}\rangle+4\mu_r(\phi,z)+\lambda_0\beta_3(\tilde{\mathbf{w}}-\mathbf{w}_d,\text{z})+\langle\vartheta,\text{z}\rangle_\Gamma
$$

$$
=2\mu_r(\operatorname{rot}\lambda,z)\quad\forall z\in H^1(\Omega),\tag{116}
$$

$$
\lambda_0 \beta_5 \langle \tilde{\boldsymbol{g}}_1, \boldsymbol{g}_1 - \tilde{\boldsymbol{g}}_1 \rangle_{\Gamma_1} - \langle \boldsymbol{\xi}, \boldsymbol{g}_1 - \tilde{\boldsymbol{g}}_1 \rangle_{\Gamma_1} \ge 0 \quad \forall \boldsymbol{g}_1 \in \mathcal{U}_1,
$$
\n(117)

$$
\lambda_0 \beta_6 \langle \tilde{g}_2, g_2 - \tilde{g}_2 \rangle_{\Gamma_3} - \langle \vartheta, g_2 - \tilde{g}_2 \rangle_{\Gamma_3} \ge 0 \quad \forall g_2 \in \mathcal{U}_2. \tag{118}
$$

*Proof.* From [\(99\)](#page-17-1)-[\(100\)](#page-17-1) we have

<span id="page-19-2"></span>
$$
\|\boldsymbol{\xi}^{\varepsilon}\|_{\mathbf{H}_{00}^{-1/2}(\Gamma\backslash\Gamma_2)} + \|\vartheta^{\varepsilon}\|_{H^{-1/2}(\Gamma)} \leq C + C \|\left[\boldsymbol{\lambda}^{\varepsilon}, \phi^{\varepsilon}\right]\|_{\mathbf{\tilde{H}}_{\sigma}\times H_0^1}.
$$
\n(119)

For  $\|[\lambda^{\varepsilon}, \phi^{\varepsilon}]\|_{\tilde{\mathbf{H}}_{\sigma} \times H_0^1}$  we have the following cases

$$
\|[\boldsymbol{\lambda}^{\varepsilon},\phi^{\varepsilon}]\|_{\tilde{\mathbf{H}}_{\sigma}\times H_0^1} \leq C \quad \text{ or } \quad \|[\boldsymbol{\lambda}^{\varepsilon},\phi^{\varepsilon}]\|_{\tilde{\mathbf{H}}_{\sigma}\times H_0^1} \longrightarrow \infty, \text{ when } \varepsilon \to 0.
$$

**Case 1:**  $I_{\varepsilon} = ||\mathbf{\hat{A}}^{\varepsilon}, \phi^{\varepsilon}||_{\mathbf{\tilde{H}}_{\sigma} \times H_0^1} \leq C$ , where C is independent of  $\varepsilon$ .

Since the sequence  $\{[\mathcal{X}^{\varepsilon}, \phi^{\varepsilon}]\}_{\varepsilon>0}$  is bounded in  $\tilde{\mathbf{H}}_{\sigma} \times H_0^1(\Omega)$ , there exists  $[\mathcal{X}, \phi] \in \tilde{\mathbf{H}}_{\sigma} \times H_0^1(\Omega)$  and a subsequence of  $\{[\mathcal{X}^{\varepsilon}, \phi^{\varepsilon}]\}_{\varepsilon>0}$ , still denoted by  $\{[\mathcal{X}^{\varepsilon}, \phi^{\varepsilon}]\}_{\varepsilon>0}$ , such that as  $\varepsilon \to 0$ ,

<span id="page-19-0"></span>
$$
[\lambda^{\varepsilon}, \phi^{\varepsilon}] \longrightarrow [\lambda, \phi] \text{ weakly in } \tilde{\mathbf{H}}_{\sigma} \times H_0^1(\Omega) \text{ and strongly in } \mathbf{L}^2(\Omega) \times L^2(\Omega). \tag{120}
$$

Then, taking into account the convergences [\(68\)](#page-14-5) and [\(120\)](#page-19-0), as  $\varepsilon \to 0$ , we can obtain

<span id="page-19-1"></span>
$$
\begin{cases}\n\langle K[\mathbf{u}^{\varepsilon}, \mathbf{u}^{\varepsilon}, \mathbf{v}] \rangle \rightarrow \mu_{1} \langle A\lambda, \mathbf{v} \rangle, \\
\langle K[\mathbf{u}^{\varepsilon}, \mathbf{u}^{\varepsilon}, \mathbf{v}^{\varepsilon}], \mathbf{v} \rangle \rightarrow \langle K[\tilde{\mathbf{u}}, \tilde{\mathbf{u}}, \tilde{\mathbf{w}}, \lambda], \mathbf{v} \rangle, \\
\langle \tilde{K}[\mathbf{u}^{\varepsilon}, \mathbf{u}^{\varepsilon}, \mathbf{v}^{\varepsilon}, \phi^{\varepsilon}], \mathbf{v} \rangle \rightarrow \langle \tilde{K}[\tilde{\mathbf{u}}, \tilde{\mathbf{u}}, \tilde{\mathbf{w}}, \phi], \mathbf{v} \rangle, \\
\langle B^{T}[\mathbf{u}^{\varepsilon}, \mathbf{u}^{\varepsilon}, \lambda^{\varepsilon}], \mathbf{v} \rangle \rightarrow \langle B^{T}[\tilde{\mathbf{u}}, \tilde{\mathbf{u}}, \tilde{\mathbf{w}}, \phi], \mathbf{v} \rangle, \\
\langle E[\mathbf{u}^{\varepsilon}, \mathbf{w}^{\varepsilon}, \phi^{\varepsilon}], \mathbf{v} \rangle \rightarrow \langle E[\tilde{\mathbf{u}}, \tilde{\mathbf{w}}, \phi], \mathbf{v} \rangle, \\
\langle F'[\mathbf{u}^{\varepsilon}, \lambda^{\varepsilon}], \mathbf{v} \rangle \rightarrow \langle F'[\tilde{\mathbf{u}}, \tilde{\mathbf{w}}, \phi], \mathbf{v} \rangle, \\
\langle G'[\mathbf{u}^{\varepsilon}, \phi^{\varepsilon}], \mathbf{v} \rangle \rightarrow \langle G'[\tilde{\mathbf{u}}, \phi], \mathbf{v} \rangle, \\
\langle G'[\mathbf{u}^{\varepsilon}, \phi^{\varepsilon}], \mathbf{v} \rangle \rightarrow \langle B[\tilde{\mathbf{u}}, \tilde{\mathbf{u}}, \phi], \mathbf{v} \rangle, \\
\langle E[\mathbf{u}^{\varepsilon}, \mathbf{u}^{\varepsilon}, \phi^{\varepsilon}], \mathbf{z} \rangle \rightarrow \langle E[\tilde{\mathbf{u}}, \tilde{\mathbf{u}}, \phi], \mathbf{z} \rangle, \\
\langle E[\mathbf{u}^{\varepsilon}, \mathbf{u}^{\varepsilon}, \phi^{\varepsilon}], \mathbf{z} \rangle \rightarrow
$$

for all  $[\mathbf{v}, \mathbf{z}] \in \mathbf{H}_{\sigma} \times H^1(\Omega)$ . In [\(121\)](#page-19-1), the operators  $A, \tilde{A}, K, \tilde{K}, B^T, E, \tilde{E}, F', G'$ , and  $J'_{\mathbf{u}}$  are defined in [\(66\)](#page-13-5), [\(89\)](#page-16-4) and [\(98\)](#page-17-4).

Since  $\|[\mathbf{\lambda}^{\varepsilon}, \phi^{\varepsilon}]\|_{\mathbf{\tilde{H}}_{\sigma} \times H_0^1} \leq C$ , from [\(119\)](#page-19-2), we have that the sequence  $\{[\xi^{\varepsilon}, \vartheta^{\varepsilon}]\}_{\varepsilon > 0}$  is bounded in  $\mathbf{H}_{00}^{-1/2}(\Gamma \setminus$  $(\Gamma_2) \times H^{-1/2}(\Gamma)$ . Then, there exist  $[\xi, \vartheta] \in \mathbf{H}_{00}^{-1/2}(\Gamma \setminus \Gamma_2) \times H^{-1/2}(\Gamma)$  and a subsequence of  $\{[\xi^{\varepsilon}, \vartheta^{\varepsilon}]\}_{\varepsilon > 0}$ , still denoted by  $\{[\xi^{\varepsilon}, \vartheta^{\varepsilon}]\}_{\varepsilon>0}$ , such that as  $\varepsilon \to 0$ , we have

$$
[\xi^{\varepsilon}, \vartheta^{\varepsilon}] \longrightarrow [\xi, \vartheta] \text{ weakly in } \mathbf{H}_{00}^{-1/2}(\Gamma \setminus \Gamma_2) \times H^{-1/2}(\Gamma),
$$

that is,

<span id="page-19-3"></span>
$$
\langle \xi^{\varepsilon}, \mathbf{v} \rangle_{\Gamma \backslash \Gamma_2} \longrightarrow \langle \xi, \mathbf{v} \rangle_{\Gamma \backslash \Gamma_2} \ \forall \mathbf{v} \in \mathbf{H}_{\sigma}, \quad \langle \vartheta^{\varepsilon}, z \rangle_{\Gamma} \longrightarrow \langle \vartheta, z \rangle_{\Gamma} \ \forall z \in H^1(\Omega). \tag{122}
$$

Thus, observing [\(121\)](#page-19-1)-[\(122\)](#page-19-3) and passing to the limit in [\(96\)](#page-16-3)-[\(97\)](#page-16-3), as  $\varepsilon \to 0$ , we obtain the system [\(115\)](#page-19-4)-[\(118\)](#page-19-4) with  $\lambda_0 = 1$ .

**Case 2:**  $I_{\varepsilon} = ||[\boldsymbol{\lambda}^{\varepsilon}, \phi^{\varepsilon}]||_{\tilde{\mathbf{H}}_{\sigma} \times H_0^1} \to \infty \text{ as } \varepsilon \to 0.$ By denoting

<span id="page-20-0"></span>
$$
\tilde{\lambda}^{\varepsilon} = \frac{\lambda^{\varepsilon}}{I_{\varepsilon}}, \quad \tilde{\phi}^{\varepsilon} = \frac{\phi^{\varepsilon}}{I_{\varepsilon}}, \tag{123}
$$

for all  $\varepsilon > 0$ , we have

<span id="page-20-5"></span>
$$
\|[\tilde{\lambda}^{\varepsilon}, \tilde{\phi}^{\varepsilon}]\|_{\tilde{\mathbf{H}}_{\sigma} \times H_0^1} = 1. \tag{124}
$$

Thus, the sequence  $\{[\tilde{\boldsymbol{\lambda}}^{\varepsilon}, \tilde{\phi}^{\varepsilon}]\}_{\varepsilon>0}$  is bounded in  $\tilde{\mathbf{H}}_{\sigma} \times H_0^1(\Omega)$ . Then there exist  $[\boldsymbol{\lambda}, \phi] \in \tilde{\mathbf{H}}_{\sigma} \times H_0^1(\Omega)$  and a subsequence of  $\{[\tilde{\lambda}^{\varepsilon}, \tilde{\phi}^{\varepsilon}]\}_{\varepsilon>0}$ , still denoted by  $\{[\tilde{\lambda}^{\varepsilon}, \tilde{\phi}^{\varepsilon}]\}_{\varepsilon>0}$ , such that, as  $\varepsilon \to 0$ , we have

<span id="page-20-2"></span>
$$
[\tilde{\boldsymbol{\lambda}}^{\varepsilon}, \tilde{\phi}^{\varepsilon}] \longrightarrow [\boldsymbol{\lambda}, \phi] \text{ weakly in } \tilde{\mathbf{H}}_{\sigma} \times H_0^1(\Omega) \text{ and strongly in } \mathbf{L}^2(\Omega) \times L^2(\Omega). \tag{125}
$$

Moreover, by denoting

<span id="page-20-1"></span>
$$
\tilde{\xi}^{\varepsilon} = \frac{\xi}{I_{\varepsilon}}, \quad \tilde{\vartheta}^{\varepsilon} = \frac{\vartheta^{\varepsilon}}{I_{\varepsilon}}, \tag{126}
$$

from [\(99\)](#page-17-1) and [\(100\)](#page-17-1), we obtain that  $\|\tilde{\xi}^{\varepsilon}\|_{\mathbf{H}_{00}^{-1/2}(\Gamma\setminus\Gamma_2)} + \|\tilde{\vartheta}^{\varepsilon}\|_{H^{-1/2}(\Gamma)} \leq \frac{C}{I_{\varepsilon}} + C \leq C$ , which implies that  $\{[\tilde{\xi}^{\varepsilon}, \tilde{\vartheta}^{\varepsilon}]\}_{\varepsilon>0}$  is bounded in  $\mathbf{H}_{00}^{-1/2}(\Gamma \setminus \Gamma_2) \times \tilde{H}^{-1/2}(\Gamma)$ . Then, there exist  $[\xi, \vartheta] \in \mathbf{H}_{00}^{-1/2}(\Gamma \setminus \Gamma_2) \times H^{-1/2}(\Gamma)$ and a subsequence of  $\{[\tilde{\xi}^{\varepsilon}, \tilde{\vartheta}^{\varepsilon}]\}_{\varepsilon>0}$ , still denoted by  $\{[\tilde{\xi}^{\varepsilon}, \tilde{\vartheta}^{\varepsilon}]\}_{\varepsilon>0}$ , such that, as  $\varepsilon \to 0$ , we have

<span id="page-20-3"></span>
$$
\langle \tilde{\xi}^{\varepsilon}, \mathbf{v} \rangle_{\Gamma \backslash \Gamma_2} \longrightarrow \langle \xi, \mathbf{v} \rangle_{\Gamma \backslash \Gamma_2} \ \forall \mathbf{v} \in \mathbf{H}_{\sigma}, \qquad \langle \tilde{\vartheta}^{\varepsilon}, z \rangle_{\Gamma} \longrightarrow \langle \vartheta, z \rangle_{\Gamma} \ \forall z \in H^1(\Omega). \tag{127}
$$

Observing [\(123\)](#page-20-0), [\(126\)](#page-20-1), and dividing the terms of the system [\(94\)](#page-16-3)-[\(97\)](#page-16-3) by  $I_{\varepsilon}$ , we obtain

<span id="page-20-4"></span>
$$
\mu_1 \langle A\tilde{\lambda}^{\varepsilon}, \mathbf{v} \rangle + \langle K[\mathbf{u}^{\varepsilon}, \mathbf{u}^{\varepsilon}, \mathbf{\tilde{\lambda}}^{\varepsilon}], \mathbf{v} \rangle + \langle B^T[\mathbf{u}^{\varepsilon}, \mathbf{u}^{\varepsilon}, \tilde{\lambda}^{\varepsilon}], \mathbf{v} \rangle + \langle \tilde{K}[\mathbf{u}^{\varepsilon}, \mathbf{u}^{\varepsilon}, \mathbf{w}^{\varepsilon}, \tilde{\phi}^{\varepsilon}], \mathbf{v} \rangle \n+ \langle E[\mathbf{u}^{\varepsilon}, \mathbf{w}^{\varepsilon}, \tilde{\phi}^{\varepsilon}], \mathbf{v} \rangle + \frac{1}{I_{\varepsilon}} \langle J'_{\mathbf{u}}(\mathbf{u}^{\varepsilon}), \mathbf{v} \rangle + \langle \tilde{\xi}^{\varepsilon}, \mathbf{v} \rangle_{\Gamma \setminus \Gamma_2} + \frac{1}{I_{\varepsilon}} (\mathbf{u}^{\varepsilon} - \tilde{\mathbf{u}}, \mathbf{v})_{\mathbf{H}_{\sigma}} \n= 2\mu_r(\text{rot } \tilde{\phi}^{\varepsilon}, \mathbf{v}) + \langle F'[\mathbf{u}^{\varepsilon}, \tilde{\lambda}^{\varepsilon}], \mathbf{v} \rangle + \langle G'[\mathbf{u}^{\varepsilon}, \tilde{\phi}^{\varepsilon}], \mathbf{v} \rangle, \n\mu_2 \langle \tilde{A} \tilde{\phi}^{\varepsilon}, z \rangle + \langle \tilde{E}[\mathbf{u}^{\varepsilon}, \mathbf{u}^{\varepsilon}, \tilde{\phi}^{\varepsilon}], z \rangle + 4\mu_r(\tilde{\phi}^{\varepsilon}, z) + \frac{1}{I_{\varepsilon}} (\mathbf{w}^{\varepsilon} - \tilde{\mathbf{w}}, z)_{H^1} + \beta_3 \frac{1}{I_{\varepsilon}} (\mathbf{w}^{\varepsilon} - \mathbf{w}_d, z)
$$
\n(128)

$$
+\langle \tilde{\vartheta}^{\varepsilon}, z \rangle_{\Gamma} = 2\mu_r(\operatorname{rot} \tilde{\boldsymbol{\lambda}}^{\varepsilon}, z), \tag{129}
$$

$$
\frac{\beta_5}{I_{\varepsilon}}\langle \boldsymbol{g}_1^{\varepsilon}, \boldsymbol{g}_1 - \boldsymbol{g}_1^{\varepsilon}\rangle_{\Gamma_1} + \frac{1}{I_{\varepsilon}}\langle \boldsymbol{g}_1^{\varepsilon} - \tilde{\boldsymbol{g}}_1, \boldsymbol{g}_1 - \boldsymbol{g}_1^{\varepsilon}\rangle_{\Gamma_1} - \langle \tilde{\boldsymbol{\xi}}^{\varepsilon}, \boldsymbol{g}_1 - \boldsymbol{g}_1^{\varepsilon}\rangle_{\Gamma_1} \ge 0, \tag{130}
$$

$$
\frac{\beta_6}{I_{\varepsilon}} \langle g_2^{\varepsilon}, g_2 - g_2^{\varepsilon} \rangle_{\Gamma_3} + \frac{1}{I_{\varepsilon}} \langle g_2^{\varepsilon} - \tilde{g}_2, g_2 - g_2^{\varepsilon} \rangle_{\Gamma_3} - \langle \tilde{\vartheta}^{\varepsilon}, g_2 - g_2^{\varepsilon} \rangle_{\Gamma_3} \ge 0,
$$
\n(131)

for all  $[\mathbf{v}, \mathbf{z}, \mathbf{g}_1, g_2] \in \mathbf{H}_{\sigma} \times H^1(\Omega) \times \mathcal{U}_1 \times \mathcal{U}_2$ .

Therefore, observing the convergences [\(121\)](#page-19-1), [\(125\)](#page-20-2), [\(127\)](#page-20-3), and passing to the limit in [\(128\)](#page-20-4)-[\(131\)](#page-20-4) when  $\varepsilon \to 0$ , we obtain the system [\(115\)](#page-19-4)-[\(118\)](#page-19-4) with  $\lambda_0 = 0$ .

Now we only need to verify that  $[0, \lambda, \phi, \xi, \vartheta] \neq \mathbf{0}$ . Observing that  $\langle A\tilde{\lambda}^{\varepsilon}, \tilde{\lambda}^{\varepsilon}\rangle = 2\|\tilde{\lambda}^{\varepsilon}\|^2_{\tilde{\mathbf{H}}_{\sigma}} + 2\alpha\|\dot{\tilde{\lambda}}^{\varepsilon}\|^2_{\mathbf{L}^2(\Gamma_2)}$  and  $\langle \tilde{A}\tilde{\phi}^{\varepsilon}, \tilde{\phi}^{\varepsilon}\rangle = \|\tilde{\phi}^{\varepsilon}\|^2_{H_0^1}$ , and replacing  $\mathbf{v} = \tilde{\lambda}^{\varepsilon}$  in [\(128\)](#page-20-4) and  $z = \tilde{\phi}^{\varepsilon}$  in [\(129\)](#page-20-4), we obtain

<span id="page-21-2"></span>
$$
2\mu_1(\|\tilde{\boldsymbol{\lambda}}^{\varepsilon}\|_{\tilde{\mathbf{H}}_{\sigma}}^2 + \alpha \|\tilde{\boldsymbol{\lambda}}^{\varepsilon}\|_{\mathbf{L}^2(\Gamma_2)}^2) = -\langle K[\mathbf{u}^{\varepsilon}, \mathbf{u}^{\varepsilon}, \tilde{\boldsymbol{\lambda}}^{\varepsilon}], \tilde{\boldsymbol{\lambda}}^{\varepsilon}\rangle - \langle B^T[\mathbf{u}^{\varepsilon}, \mathbf{u}^{\varepsilon}, \tilde{\boldsymbol{\lambda}}^{\varepsilon}], \tilde{\boldsymbol{\lambda}}^{\varepsilon}\rangle
$$
  
\n
$$
- \langle \tilde{K}[\mathbf{u}^{\varepsilon}, \mathbf{u}^{\varepsilon}, \mathbf{w}^{\varepsilon}, \tilde{\phi}^{\varepsilon}], \tilde{\boldsymbol{\lambda}}^{\varepsilon}\rangle - \langle E[\mathbf{u}^{\varepsilon}, \mathbf{w}^{\varepsilon}, \tilde{\phi}^{\varepsilon}], \tilde{\boldsymbol{\lambda}}^{\varepsilon}\rangle
$$
  
\n
$$
- \frac{1}{I_{\varepsilon}} \langle J'_{\mathbf{u}}(\mathbf{u}^{\varepsilon}), \tilde{\boldsymbol{\lambda}}^{\varepsilon}\rangle - \langle \tilde{\boldsymbol{\xi}}^{\varepsilon}, \tilde{\boldsymbol{\lambda}}^{\varepsilon}\rangle_{\Gamma\backslash\Gamma_2} - \frac{1}{I_{\varepsilon}} (\mathbf{u}^{\varepsilon} - \tilde{\mathbf{u}}, \tilde{\boldsymbol{\lambda}}^{\varepsilon})_{\mathbf{H}_{\sigma}}
$$
  
\n
$$
+ 2\mu_r(\text{rot}\,\tilde{\phi}^{\varepsilon}, \tilde{\boldsymbol{\lambda}}^{\varepsilon}) + \langle F'[\mathbf{u}^{\varepsilon}, \tilde{\boldsymbol{\lambda}}^{\varepsilon}], \tilde{\boldsymbol{\lambda}}^{\varepsilon}\rangle + \langle G'[\mathbf{u}^{\varepsilon}, \tilde{\phi}^{\varepsilon}], \tilde{\boldsymbol{\lambda}}^{\varepsilon}\rangle, \tag{132}
$$
  
\n
$$
\mu_2 \|\tilde{\phi}^{\varepsilon}\|_{H_0^1}^2 + 4\mu_r \|\tilde{\phi}^{\varepsilon}\|^2 = -\langle \tilde{E}[\mathbf{u}^{\varepsilon}, \mathbf{u}^{\varepsilon}, \tilde{\phi}^{\varepsilon
$$

If  $[\lambda, \phi] = [0, 0]$ , considering [\(120\)](#page-19-0)-[\(121\)](#page-19-1) and passing to the limit in [\(132\)](#page-21-2)-[\(133\)](#page-21-2) as  $\varepsilon \to 0$ , we obtain that  $\mu_1 \|\tilde{\boldsymbol{\lambda}}^{\varepsilon}\|^2_{\tilde{\mathbf{H}}_{\sigma}} \longrightarrow 0$  and  $\mu_2 \|\tilde{\phi}^{\varepsilon}\|^2_{H_0^1} \longrightarrow 0$ ; then it follows

$$
\|[\tilde{\boldsymbol{\lambda}}^\varepsilon,\tilde{\phi}^\varepsilon]\|^2_{\tilde{\mathbf{H}}_\sigma\times H^1_0}=\|\tilde{\boldsymbol{\lambda}}^\varepsilon\|^2_{\tilde{\mathbf{H}}_\sigma}+\|\tilde{\phi}^\varepsilon\|^2_{H^1_0}\rightarrow0,
$$

which contradicts the equality given in [\(124\)](#page-20-5). Therefore, we conclude that  $[\lambda, \phi] \neq [0, 0]$  and consequently  $[0, \lambda, \phi, \xi, \vartheta] \neq 0$ . Thus, the proof of the theorem is finished.  $[0, \lambda, \phi, \xi, \vartheta] \neq 0$ . Thus, the proof of the theorem is finished.

<span id="page-21-0"></span>Remark 4 *From [\(16\)](#page-5-3) and [\(115\)](#page-19-4)-[\(118\)](#page-19-4), we obtain the following optimality system for problem [\(22\)](#page-6-1) constituted by the state equations [\(16\)](#page-5-3), the adjoint equations [\(115\)](#page-19-4)-[\(116\)](#page-19-4) and the optimality conditions [\(117\)](#page-19-4)-[\(118\)](#page-19-4).*

**Corollary 1** *Suppose that the assumptions of Theorem* [6](#page-18-0) are satisfied and let  $[\tilde{\mathbf{u}}, \tilde{\mathbf{w}}, \tilde{\mathbf{g}}_1, \tilde{g}_2] \in S_{ad}$  an optimal *solution to problem* [\(22\)](#page-6-1)*. Let*  $\mu_1$  *and*  $\mu_2$  *large enough such that* 

<span id="page-21-3"></span>
$$
\tilde{\delta} > C(\|[\tilde{\mathbf{u}}, \tilde{\mathbf{w}}]\|_{\mathbf{H}_{\sigma} \times H^1} + \|[\tilde{\mathbf{u}}, \tilde{\mathbf{w}}]\|_{\mathbf{H}_{\sigma} \times H^1}^2 + \|\mathbf{f}\| + \|g\| + \mu_r),\tag{134}
$$

where  $\tilde{\delta} = \min\{2\mu_1, \mu_2\}$  and C is a positive constant depending only on  $\Omega$ ,  $\|\eta(N\tilde{\mathbf{u}})\|_{\infty}$ , and  $\|\eta'(N\tilde{\mathbf{u}})\|_{\infty}$ . Then, there exists a unique  $[\lambda_0, \lambda, \phi, \xi, \vartheta] \in (\mathbb{R}^+ \cup \{0\}) \times \tilde{H}_{\sigma} \times H_0^1(\Omega) \times H_{00}^{-1/2}(\Gamma \setminus \Gamma_2) \times H^{-1/2}(\Gamma)$  satisfying  $(115)$ <sup>2</sup> $(118)$ *, with*  $\lambda_0 = 1$ .

*Proof.* We assume that  $\lambda_0 = 0$ . Then, by setting  $\mathbf{v} = \lambda$  in [\(115\)](#page-19-4) and  $z = \phi$  in [\(116\)](#page-19-4), taking into account that  $\langle \xi, \lambda \rangle_{\Gamma \backslash \Gamma_2} = 0$  and  $\langle \vartheta, \phi \rangle_{\Gamma} = 0$ , observing the estimates given in [\(105\)](#page-17-5), [\(109\)](#page-18-2)-[\(114\)](#page-18-1), and definition of A,  $\tilde{A}$ , we obtain

$$
2\mu_1 \|\lambda\|^2 \leq C\mu_r \|\[\lambda, \phi\]\|_{\tilde{\mathbf{H}}_{\sigma} \times H_0^1}^2 + C\|\eta(N\tilde{\mathbf{u}})\|_{\infty} (\|\tilde{\mathbf{u}}\|_{\mathbf{H}_{\sigma}} + \|\tilde{\mathbf{w}}\|_{H^1} + \|\tilde{\mathbf{u}}\|_{\mathbf{H}_{\sigma}}^2) \|[\lambda, \phi]\|_{\tilde{\mathbf{H}}_{\sigma} \times H_0^1}^2
$$
  
+
$$
C\|\eta'(N\tilde{\mathbf{u}})\|_{\infty} (\|\tilde{\mathbf{u}}\|_{\mathbf{H}_{\sigma}} \|\tilde{\mathbf{w}}\|_{H^1} + \|f\| + \|g\|) \|[\lambda, \phi]\|_{\tilde{\mathbf{H}}_{\sigma} \times H_0^1}^2,
$$
  

$$
\mu_2 \|\phi\|^2 \leq C\mu_r \|\[\lambda, \phi\]\|_{\tilde{\mathbf{H}}_{\sigma} \times H_0^1}^2.
$$

Then, by adding the above inequalities, we deduce

$$
\tilde{\delta} \Vert [\lambda, \phi] \Vert^{2}_{\tilde{\mathbf{H}}_{\sigma} \times H^{1}_{0}} \leq C (\Vert [\tilde{\mathbf{u}}, \tilde{\mathbf{w}}] \Vert_{\mathbf{H}_{\sigma} \times H^{1}} + \Vert [\tilde{\mathbf{u}}, \tilde{\mathbf{w}}] \Vert^{2}_{\mathbf{H}_{\sigma} \times H^{1}} + \Vert \mathbf{f} \Vert + \Vert \mathbf{g} \Vert + \mu_{r}) \Vert [\lambda, \phi] \Vert^{2}_{\tilde{\mathbf{H}}_{\sigma} \times H^{1}_{0}},
$$

<span id="page-21-1"></span>which, by applying condition [\(134\)](#page-21-3) implies that  $\|\left[\lambda, \phi\right]\|_{\tilde{\mathbf{H}}_{\sigma} \times H_0^1}^2 = 0$ , that is,  $\lambda = 0$  and  $\phi = 0$ . In this case, the equations [\(115\)](#page-19-4) and [\(116\)](#page-19-4) can be rewritten as  $\langle \xi, v \rangle_{\Gamma \backslash \Gamma_2} = 0$  for any  $v \in H_{\sigma}$  and  $\langle \vartheta, z \rangle_{\Gamma} = 0$  for any <sup>z</sup> <sup>∈</sup> <sup>H</sup><sup>1</sup> (Ω), respectively. Hence we have ξ = 0 and ϑ = 0, which contradicts Theorem [6.](#page-18-0) ⋄ **Remark 5** If the Lagrange multiplier  $\lambda_0 = 1$ , then the optimality conditions are equivalent to

<span id="page-22-13"></span>
$$
\langle \beta_5 \tilde{g}_1 - \xi, g_1 - \tilde{g}_1 \rangle_{\Gamma_1} \ge 0 \quad \text{and} \quad \langle \beta_6 \tilde{g}_2 - \vartheta, g_2 - \tilde{g}_2 \rangle_{\Gamma_3} \ge 0. \tag{135}
$$

*Since the set of controls*  $U_1 \times U_2$  *is convex, from inequalities* [\(135\)](#page-22-13) we obtain

$$
\tilde{\boldsymbol{g}}_1 = \Pr_{\mathcal{U}_1} \left( \frac{\boldsymbol{\xi}}{\beta_5} \right) \quad on \; \Gamma_1, \qquad \tilde{g}_2 = \Pr_{\mathcal{U}_2} \left( \frac{\vartheta}{\beta_6} \right) \quad on \; \Gamma_3.
$$

Acknowledgments: The first author was supported by proyecto UTA-Mayor, 4738-17. The third author was supported by Fondo Nacional de Financiamiento para la Ciencia, la Tecnología y la Innovación Francisco José de Caldas, contrato Colciencias FP 44842-157-2016.

## <span id="page-22-4"></span>References

- [1] Ammar-Khodja F., Santos M.M., 2D density-dependent, Leray problem with a discontinuous density. Methods Appl. Anal. 13, 321-335, (2006).
- <span id="page-22-5"></span>[2] Ammar-Khodja F., Santos M.M., 2D, Ladyzhenskaya-Solonnikov problem for inhomogeneous fluids. Contributions to nonlinear analysis. Progr. Nonlinear Differential Equations Appl. Birkhäuser, 66, 351-364, (2006).
- <span id="page-22-6"></span>[3] Ammar-Khodja F., Santos M.M., The Leray problem for 2D inhomogeneous fluids. Regularity and other aspects of the Navier-Stokes equations. Banach Center Publ., Polish Acad. Sci., Warsaw, (2005).
- <span id="page-22-7"></span>[4] Ammar-Khodja F., Santos M.M., 2D density-dependent Leray's problem for inhomogeneous fluids. Mat. Contemp. 27, 19-35, (2004).
- <span id="page-22-11"></span>[5] Arbergel F., Casas E., Some optimal control problems of multistate equations appearing in fluid mechanics. Math. Model. Numer. Anal. 27, 2, 223-247, (1993).
- <span id="page-22-12"></span><span id="page-22-9"></span>[6] Brézis H., Functional Analysis, Sobolev spaces and Partial differential Equations, Springer, (2010).
- [7] Casas E., Mateos M., Raymond J.P., Error estimates for the numercal approximation of a distributed control problem for the steady-state Navier-Stokes equations. SIAM J. Control Optim. 46, 952-982, (2007).
- <span id="page-22-10"></span>[8] Dautray R., Lions J.L., Mathematical Analysis and Numerical Methods for Science and Technology, Vol. 2, Springer, Berlin, (2000).
- [9] De los Reyes J.C., Kunisch K., A semi-smooth Newton method for control constrained boundary optimal control of the Navier-Stokes equations. Nonlinear Analysis, 62, 1289-1316, (2005).
- <span id="page-22-2"></span><span id="page-22-1"></span>[10] Eringen A.C., Simple microfluids. Int. J. Engng. Sci. 2, 205-217, (1964).
- <span id="page-22-3"></span>[11] Eringen A.C., Theory of micropolar fluids, J. Math. Mech. 16, 1-18, (1966).
- [12] Ferreira L.C.F., Villamizar-Roa E.J., Micropolar fluid system in a space of distributions and large time behavior. J. Math. Anal. Appl. 332, 1425-1445, (2007).
- <span id="page-22-0"></span>[13] Ferreira L.C.F., Planas G., Villamizar-Roa E.J., On the nonhomogeneous Navier-Stokes system with Navier friction boundary conditions. SIAM J. Math. Anal. 45, 2576-2595, (2013).
- <span id="page-22-8"></span>[14] Frolov N.N., Solvability of a boundary problem of motion of an inhomogeneous fluid, Mat. Zametki 53, 130-140, (1993).
- <span id="page-23-20"></span><span id="page-23-8"></span>[15] Frolov N.N., Boundary value problem describing the motion of an inhomogeneous fluid, Siberian Math. J. 37, 433-451, (1995).
- [16] Galdi G.P., An introduction to the mathematical theory of the Navier-Stokes equations. Second Edition, Springer, (2011).
- <span id="page-23-18"></span><span id="page-23-12"></span>[17] Iftimie D., Planas G., Inviscid limits for the Navier-Stokes equations with Navier friction boundary conditions. Nonlinearity 19, 899-918, (2006).
- <span id="page-23-11"></span>[18] Illarionov A.I., Optimal boundary control of steady-state flow of a viscous inhomogeneous incompressible fluid. Mathematical Notes, 69, 5, 614-624, (2001).
- [19] Jägger W., Mikelić A., On the roughness-induced effective boundary conditions for an incompressible viscous flow. J. Diff. Eqns, 170, 96-122, (2001).
- <span id="page-23-13"></span>[20] Lee H.-C., Imanuvilov O.Y., Analysis of Newmann boundary optimal control problems for the stationary Boussinesq equations including solid media. SIAM J. Control Optim. 2, 39, 457-477, (2000).
- <span id="page-23-4"></span><span id="page-23-2"></span>[21] Lions P.L., Mathematical Topics in Fluid Mechanics, Clarendon Press, Oxford, (1996).
- <span id="page-23-14"></span>[22] Lukaszewicz G., Micropolar Fluids: Theory and Applications Birkhäuser, Boston, (1999).
- [23] Mallea-Zepeda E., Ortega-Torres E., Villamizar-Roa E.J., A boundary control problem for micropolar fluids. J. Optim. Theory Appl. 169, 349-369, (2016).
- <span id="page-23-10"></span>[24] Maxwell J.C., On stressed in rariffed gases arising from inequalities of temperature. Phil. Trans. Royal society, 704-712, (1879).
- <span id="page-23-9"></span>[25] Navier C.L.M.H., Sur le lois de l'équilibrie et du mouvement des corps élastiques, Mem. and Acad. R. Sci. Inst. France, 369, (1827).
- <span id="page-23-1"></span><span id="page-23-0"></span>[26] Pedlosky J., Geophysical Fluid Dynamics, Springer-Verlag, (1984).
- [27] Santos M.M., Stationary solution of the Navier-Stokes equations in a 2D bounded domain for incompressible flow with discontinuous density. Z. Angew. Math. Phys. 53, 661-675, (2002).
- <span id="page-23-3"></span>[28] Simon J., Nonhomogeneous viscous incompressible fluids: existence of velocity, density, and pressure. SIAM J. Math. Anal., 21, 1093-1117, (1990).
- <span id="page-23-15"></span>[29] Stavre R., A distributed control problem for micropolar fluids. In honour of Academician Nicolae Dan Cristescu on his 70th birthday. Rev. Roumaine Math. Pures Appl. 45, 2, 353-358, (2001).
- <span id="page-23-16"></span>[30] Stavre R., The control of the pressure for a micropolar fluid. Dedicated to Eugen Sos. Z. Angew. Math. Phys. 53, 6, 912-922, (2002).
- <span id="page-23-17"></span>[31] Stavre R., Optimization and numerical approximation for micropolar fluids. Numer. Funct. Anal. Optim. 24, 3-4, 223-241, (2003).
- <span id="page-23-19"></span>[32] Verfüth R., Finite element approximation of incompresible Navier-Stokes equations with slip boundary condition. Numer. Math. 50, 697-721, (1987).
- <span id="page-23-5"></span>[33] Villamizar-Roa E.J. and Rodríguez-Bellido M.A., Global existence and exponential stability for the micropolar fluid system. Z. Angew. Math. Phys. 59, 5, 790-809, (2008).
- <span id="page-23-6"></span>[34] Villamizar-Roa E.J., Rodríguez-Bellido M.A., Rojas-Medar M.A., Some properties of a class of abstract stationary equations. Nonlinear Anal. 64, 10, 2203-2214, (2006).
- <span id="page-23-7"></span>[35] Vitoriano F., On the steady viscous flow of a nonhomegeneuos asymmetric fluid. Annali di Matematica. 192, 665-672, (2013).