REGULARIZED WEIGHTED DISCRETE LEAST SQUARES APPROXIMATION USING GAUSS QUADRATURE POINTS

Congpei An^{*} and Hao-Ning Wu[†]

Abstract

We consider polynomial approximation over the interval [-1,1] by regularized weighted discrete least squares methods with ℓ_2- or ℓ_1 -regularization, respectively. As the set of nodes we use Gauss quadrature points (which are zeros of orthogonal polynomials). The number of Gauss quadrature points is N + 1. For $2L \leq 2N + 1$, with the aid of Gauss quadrature, we obtain approximation polynomials of degree L in closed form without solving linear algebra or optimization problems. In fact, these approximation polynomials can be expressed in the form of the barycentric interpolation formula (Berrut & Trefethen, 2004) when an interpolation condition is satisfied. We then study the approximation quality of the ℓ_2 -regularized approximation polynomial in terms of Lebesgue constants, and the sparsity of the ℓ_1 -regularized approximation polynomial. Finally, we give numerical examples to illustrate these theoretical results and show that a well-chosen regularization parameter can lead to good performance, with or without contaminated data.

Keywords. regularized least squares approximation, Gauss quadrature, Lebesgue constants, sparsity, barycentric interpolation.

AMS subject classifications. 41A05, 65D99, 65D32, 94A99

1 Introduction

In this paper, we are interested in approximating or recovering a function (possibly noisy) $f \in C([-1,1])$ by a polynomial

$$p_L(x) = \sum_{\ell=0}^{L} \beta_{\ell} \tilde{\Phi}_{\ell}(x) \in \mathbb{P}_L, \quad x \in [-1, 1],$$
(1.1)

where \mathbb{P}_L is a linear space of polynomials of degree at most L, and $\tilde{\Phi}_{\ell}(x)$, $\ell = 0, 1, \ldots, L$, are normalized orthogonal polynomials (Szegö, 1939; Gautschi, 2004). As long as the basis $\{\tilde{\Phi}_{\ell}(x)\}_{\ell=0}^{L}$ for \mathbb{P}_L is given, the next step is to determine coefficients β_{ℓ} , $\ell = 0, \ldots, L$. We will consider the ℓ_2 -regularized approximation problem

$$\min_{\beta_{\ell} \in \mathbb{R}} \left\{ \sum_{j=0}^{N} \omega_j \left(\sum_{\ell=0}^{L} \beta_{\ell} \tilde{\Phi}_{\ell}(x_j) - f(x_j) \right)^2 + \lambda \sum_{\ell=0}^{L} (\mu_{\ell} \beta_{\ell})^2 \right\}, \quad \lambda > 0,$$
(1.2)

and the ℓ_1 -regularized approximation problem

$$\min_{\beta_{\ell} \in \mathbb{R}} \left\{ \sum_{j=0}^{N} \omega_j \left(\sum_{\ell=0}^{L} \beta_{\ell} \tilde{\Phi}_{\ell}(x_j) - f(x_j) \right)^2 + \lambda \sum_{\ell=0}^{L} |\mu_{\ell} \beta_{\ell}| \right\}, \quad \lambda > 0,$$
(1.3)

where f is a given continuous function with values (possibly noisy) taken at N + 1 distinct points x_0, x_1, \ldots, x_N over the interval [-1, 1]; $2L \le 2N + 1$; $\mathbf{w} = [\omega_0, \omega_1, \ldots, \omega_N]^T$ is a vector of positive

^{*}School of Economic Mathematics, Southwestern University of Finance and Economics, Chengdu 611130, China. Email address: andbach1984@hotmail.com, ancp@swufe.edu.cn

 $^{^\}dagger \mathrm{Department}$ of Mathematics, The University of Hong Kong, Hong Kong, China.

Email address: haoning.wu@outlook.com, hnwu@hku.hk

Gauss quadrature weights (Gautschi, 2004); $\{\mu_\ell\}_{\ell=0}^L$ is a nonnegative nondecreasing sequence, which penalizes coefficients $\{\beta_\ell\}_{\ell=0}^L$; and $\lambda > 0$ is the regularization parameter.

It is known that approximation schemes (1.2) and (1.3) are special cases of classical penalized least squares methods, see Powell (1967), Golitschek & Schumaker (1990), Gautschi (2004), Lazarov *et al.* (2007), Cai *et al.* (2009), Kim *et al.* (2009), An *et al.* (2012), Xiang & Zou (2013) and Zhou & Chen (2018). Some optimization methods or iterative algorithms are presented to find minimizers. However, we will concentrate on the aspect of constructing minimizers to problems (1.2) and (1.3) by means of orthogonal polynomials and Gauss quadrature (Kress, 1998; Gautschi, 2004, 2012; Trefethen, 2013). In this paper, Gauss quadrature will play an important role. We assume that the weight function $w: (-1, 1) \rightarrow \mathbb{R}$ is positive, such that

$$\int_{-1}^{1} w(x)dx < \infty, \quad \int_{-1}^{1} x^{i}w(x)dx < \infty, \ i = 1, 2, \dots.$$

Definition 1.1 A quadrature formula

$$\int_{-1}^{1} w(x) f(x) dx \approx \sum_{j=0}^{N} \omega_j f(x_j)$$

with N + 1 distinct quadrature points x_0, x_1, \ldots, x_N is called a Gauss quadrature formula if it integrates all polynomials $p \in \mathbb{P}_{2N+1}$ exactly, i.e., if

$$\sum_{j=0}^{N} \omega_j p(x_j) = \int_{-1}^{1} w(x) p(x) dx \quad \forall p \in \mathbb{P}_{2N+1}.$$
(1.4)

 x_0, x_1, \ldots, x_N are called Gauss quadrature points.

Without causing confusion, we call $\mathcal{X}_{N+1} = \{x_j\}_{j=0}^N$ Gauss quadrature points. Throughout this paper, we always assume that \mathcal{X}_{N+1} are Gauss quadrature points. It is well known (see, for example, Powell (1981), Kress (1998) and Gautschi (2012)) that Gauss quadrature points \mathcal{X}_{N+1} are the zeros of the orthogonal polynomial of degree N + 1. The orthogonality is with respect to the L_2 inner product

$$(f,g)_{L_2} := \int_{-1}^1 w(x)f(x)g(x)dx$$

This inner product induces the standard L_2 norm

$$||f||_{L_2} := \sqrt{(f,f)_{L_2}} = \left(\int_{-1}^1 |f(x)|^2 dx\right)^{1/2}.$$

Given a continuous function f defined on [-1, 1], sampling on \mathcal{X}_{N+1} generates

$$\mathbf{f} := \mathbf{f}(\mathcal{X}_{N+1}) = [f(x_0), f(x_1), \dots, f(x_N)]^T \in \mathbb{R}^{N+1}$$

Let $\mathbf{A}_L := \mathbf{A}_L(\mathcal{X}_{N+1}) \in \mathbb{R}^{(N+1) \times (L+1)}$ be a matrix of orthogonal polynomials evaluated at \mathcal{X}_{N+1} :

$$\mathbf{A}_L = \left[\tilde{\Phi}_\ell(x_j)\right] \in \mathbb{R}^{(N+1) \times (L+1)}, \quad j = 0, 1, \dots, N, \quad \ell = 0, 1, \dots, L.$$

By subtracting the structure (1.1) of the approximation polynomial into the ℓ_2 -regularized approximation problem (1.2), the problem (1.2) transforms into the following problem

$$\min_{\boldsymbol{\beta} \in \mathbb{R}^{L+1}} \| \mathbf{W}^{\frac{1}{2}} (\mathbf{A}_L \boldsymbol{\beta} - \mathbf{f}) \|_2^2 + \lambda \| \mathbf{R}_L \boldsymbol{\beta} \|_2^2, \quad \lambda > 0,$$
(1.5)

where

$$\mathbf{W} = \operatorname{diag}(\omega_0, \omega_1, \dots, \omega_N) \in \mathbb{R}^{(N+1) \times (N+1)}$$

and the diagonal matrix $\mathbf{R}_L := \operatorname{diag}(\mu_0, \mu_1, \dots, \mu_L) \in \mathbb{R}^{(L+1) \times (L+1)}$ is a semi-definite positive matrix.

With the same basis and weight vector as the ℓ_2 -regularized approximation problem (1.2) above, the ℓ_1 -regularized approximation problem (1.3) transforms into

$$\min_{\boldsymbol{\beta} \in \mathbb{R}^{L+1}} \| \mathbf{W}^{\frac{1}{2}} (\mathbf{A}_L^T \boldsymbol{\beta} - \mathbf{f}) \|_2^2 + \lambda \| \mathbf{R}_L \boldsymbol{\beta} \|_1, \quad \lambda > 0.$$
(1.6)

Now the next step is to fix $\boldsymbol{\beta} = [\beta_0, \beta_1, \dots, \beta_L]^T \in \mathbb{R}^{L+1}$.

The goal of this paper is to construct approximation polynomials in the form of (1.1). We specify coefficients $\{\beta_\ell\}_{\ell=0}^L$ by solving problems (1.5) and (1.6) directly. This immediately yields the ℓ_2 -regularized approximation polynomial $p_{L,N+1}^{\ell_2}$ (2.3) and the ℓ_1 -regularized approximation polynomial $p_{L,N+1}^{\ell_1}$ (2.7).

As is known to all, Gauss quadrature goes hand in hand with the theory and computation of orthogonal polynomials, see Gautschi (2004) and Trefethen (2013) and references therein. Orthogonal polynomials occur in a wide range of applications and act as a remarkable role in pure and applied mathematics. Chebyshev polynomials and Legendre polynomials are two excellent factors in the family of orthogonal polynomials. Many polynomial approximation textbooks introduce fruitful results of Chebyshev and Legendre polynomials (Szegö, 1939; Powell, 1981; Gautschi, 2012; Trefethen, 2013). In particular, we take these two orthogonal polynomials (Chebyshev and Legendre) as representative examples in the choice of basis and Gauss quadrature points.

In the next section, we introduce some necessary notations and terminologies. The construction of the ℓ_2- and ℓ_1 -regularized minimizers to problems (1.2) and (1.3) are presented, respectively. The crucial fact is that both $p_{L,N+1}^{\ell_2}$ and $p_{L,N+1}^{\ell_1}$ could be presented in the barycentric form under an interpolation condition, see the ℓ_2 -regularized barycentric interpolation formula (2.13) and the ℓ_1 -regularized barycentric interpolation formula (2.17). It is worth noting that the Wang-Xiang formula (Wang & Xiang, 2012) is a special case of $p_{L,N+1}^{\ell_2}$ when we set Legendre polynomials as the basis, see Section 2.3. In Section 3, we study the quality of the approximation polynomial $p_{L,N+1}^{\ell_2}$ in terms of Lebesgue constants. We illustrate that Lebesgue constants decay when the regularization parameter increases. Section 4 analyses the ℓ_1 -regularized approximation problem (1.3) in the view of sparsity. In particular, we derive a sharp upper bound of nonzero entries in the solution to the ℓ_1 -regularized approximation problem (1.6). We consider, in Section 5, numerical experiments containing approximation with exact and contaminated data.

All numerical results^{*} in this paper are carried out by using MATLAB R2017A on a desktop (8.00 GB RAM, Intel(R) Processor 5Y70 at 1.10 GHz and 1.30 GHz) with the Windows 10 operating system.

2 Regularized weighted least squares approximation

The construction of minimizers to problems (1.2) and (1.3) is presented in this section.

2.1 ℓ_2 -regularized approximation problem

First, we consider solving the ℓ_2 -regularized weighted discrete least squares approximation problem (the ℓ_2 -regularized approximation problem) (1.2). The problem can be transformed into a convex and differential optimization problem (1.5).

Taking the first derivative of the objective function in the ℓ_2 -regularized approximation problem (1.5) in the matrix form with respect to β leads to the first order condition

$$\left(\mathbf{A}_{L}^{T}\mathbf{W}\mathbf{A}_{L}+\lambda\mathbf{R}_{L}^{T}\mathbf{R}_{L}\right)\boldsymbol{\beta}=\mathbf{A}_{L}^{T}\mathbf{W}\mathbf{f}, \quad \lambda>0.$$
(2.1)

One may solve the first order condition (2.1) using methods of numerical linear algebra; however, in this paper we concentrate on how to obtain the solution to the first order condition (2.1) in a closed form.

^{*}All codes are available at

https://github.com/HaoNingWu/Regularized-Least-Squares-Approximation-using-Orthogonal-Polynomials.

Lemma 2.1 Let $\{\tilde{\Phi}_{\ell}\}_{\ell=0}^{L}$ be a class of normalized orthogonal polynomials with the weight function w(x) over [-1,1], and $\mathcal{X}_{N+1} = \{x_0, x_1, \ldots, x_N\}$ be the set of zeros of $\tilde{\Phi}_{N+1}$. Assume $2L \leq 2N+1$ and \mathbf{w} is a vector of weights satisfying the Gauss quadrature formula (1.4). Then

$$\mathbf{H}_L := \mathbf{A}_L^T \mathbf{W} \mathbf{A}_L = \mathbf{I}_L \in \mathbb{R}^{(L+1) \times (L+1)},$$

where \mathbf{I}_L is the identity matrix.

Proof. By the structure of the matrix \mathbf{H}_L and the definition of Gauss quadrature formula (see (1.4)), we obtain

$$(\mathbf{H}_L)_{\ell,\ell'} = \sum_{j=0}^N \omega_j \tilde{\Phi}_\ell(x_j) \tilde{\Phi}_{\ell'}(x_j) = \int_{-1}^1 w(x) \tilde{\Phi}_\ell(x) \tilde{\Phi}_{\ell'}(x) dx = \delta_{\ell,\ell'},$$

where $\delta_{\ell,\ell'}$ is the Kronecker delta. The middle equality holds from $\tilde{\Phi}_{\ell}(x)\tilde{\Phi}_{\ell'}(x) \in \mathbb{P}_{2L} \subset \mathbb{P}_{2N+1}$, and the last equality holds because of the orthonormality of $\{\tilde{\Phi}_{\ell}\}_{\ell=0}^{L}$.

Theorem 2.1 Under the condition of Lemma 2.1, the optimal solution to the ℓ_2 -regularized approximation problem (1.5) in the matrix form can be expressed by

$$\beta_{\ell} = \frac{1}{1 + \lambda \mu_{\ell}^2} \sum_{j=0}^N \omega_j \tilde{\Phi}_{\ell}(x_j) f(x_j), \quad \ell = 0, 1, \dots, L, \quad \lambda > 0.$$
(2.2)

Consequently, the ℓ_2 -regularized approximation polynomial is

$$p_{L,N+1}^{\ell_2}(x) = \sum_{\ell=0}^{L} \frac{\tilde{\Phi}_{\ell}(x)}{1 + \lambda \mu_{\ell}^2} \sum_{j=0}^{N} \omega_j \tilde{\Phi}_{\ell}(x_j) f(x_j).$$
(2.3)

Proof. This is immediately obtained from the first order condition (2.1) of the problem (1.5) and Lemma 2.1. \Box

In the limiting case $N \to \infty$, we obtain the following simple but significant result.

Theorem 2.2 Adopt the notation and assumptions of Lemma 2.1. Let $f \in C([-1,1])$, and let $L \ge 0$ be given. Then the polynomial $p_{L,N+1}^{\ell_2}$ (2.3) has the uniform limit p_L^* as $N \to \infty$, that is

$$\lim_{N \to \infty} \|p_{L,N+1}^{\ell_2} - p_L^*\|_{\infty} = 0,$$

where

$$p_L^*(x) = \sum_{\ell=0}^L \frac{\tilde{\Phi}_\ell(x)}{1 + \lambda \mu_\ell^2} \int_{-1}^1 w(x) \tilde{\Phi}_\ell(x) f(x) dx.$$
(2.4)

Proof. Since the interval [-1, 1] is a compact set, and since the sums over ℓ in (2.2) and (2.4) are finite, to prove the theorem, it is sufficient to prove that

$$\lim_{N \to \infty} \sum_{j=0}^{N} \omega_j \tilde{\Phi}_\ell(x_j) f(x_j) = \int_{-1}^1 w(x) \tilde{\Phi}_\ell(x) f(x) dx, \quad 0 \le \ell \le L.$$
(2.5)

Since $w(x)\Phi_{\ell}(x)f(x) \in \mathcal{C}([-1, 1])$, the result follows from that, the sequence of Gauss quadrature formulae is convergent (Kress, 1998, Chapter 9). Hence (2.5) holds, proving the whole theorem. \Box

2.2 ℓ_1 -regularized approximation problem

Now we discuss the ℓ_1 -regularized approximation problem (1.3), but we still convert it into solving the problem (1.6) in a matrix form. To solve this problem, we first define the soft threshold operator $S_k(a)$.

Definition 2.1 (Donoho & Johnstone, 1994) The soft threshold operator, denoted as $S_k(a)$, is defined as

$$S_k(a) = \max(0, a - k) + \min(0, a + k).$$

Theorem 2.3 Adopt the notation and assumptions of Lemma 2.1. Then the ℓ_1 -regularized approximation problem (1.6) in the matrix form has the unique closed-form solution

$$\beta_{\ell} = \frac{1}{2} S_{\lambda \mu_{\ell}}(2\alpha_{\ell}), \quad \ell = 0, 1, \dots, L,$$
(2.6)

where $\alpha_{\ell} = \sum_{j=0}^{N} \omega_j \tilde{\Phi}_{\ell}(x_j) f(x_j)$. Consequently, the ℓ_1 -regularized approximation polynomial is

$$p_{L,N+1}^{\ell_1}(x) = \frac{1}{2} \sum_{\ell=0}^N S_{\lambda\mu_\ell} (2\alpha_\ell) \tilde{\Phi}_\ell(x).$$
(2.7)

The method of the proof is similar to the proof of Zhou & Chen (2018, Theorem 5.1). But we explain that our regularized least squares approximation problem (1.3) is over the interval [-1, 1] rather than over the unit sphere.

Proof. Since \mathbf{H}_L is non-singular, for the ℓ_1 -regularized approximation problem (1.6) in the matrix form, its first order condition is

$$\mathbf{0} \in 2\mathbf{H}_L \boldsymbol{\beta} - 2\mathbf{A}_L^T \mathbf{W} \mathbf{f} + \lambda \partial (\|\mathbf{R}_L \boldsymbol{\beta}\|_1),$$
(2.8)

where $\partial(\cdot)$ denotes the subgradient (Clarke, 1990). Since $\mathbf{H}_L = \mathbf{I}_L$ is an identity matrix and \mathbf{R}_L is a diagonal matrix, $\boldsymbol{\beta}$ is the solution to the ℓ_1 -regularized approximation problem (1.6) in the matrix form if and only if

$$0 \in 2\beta_{\ell} - 2\alpha_{\ell} + \lambda \mu_{\ell} \partial |\beta_{\ell}|, \quad \ell = 0, 1, \dots, L,$$

$$(2.9)$$

where $-1 \leq \partial |\beta_{\ell}| \leq 1$. Let β_{ℓ}^* be the optimal solution to the problem (2.9), then

$$\beta_{\ell}^* = \frac{1}{2} (2\alpha_{\ell} - \lambda \mu_{\ell} \partial |\beta_{\ell}^*|), \quad \ell = 0, 1, \dots, L.$$

Next there exist three cases to be considered:

1. If $2\alpha_{\ell} > \lambda \mu_{\ell}$, then $2\alpha_{\ell} - \lambda \mu_{\ell} \partial |\beta_{\ell}^*| > 0$, thus $\beta_{\ell}^* > 0$, yielding $\partial |\beta_{\ell}^*| = 1$, then

$$\beta_{\ell}^* = \frac{1}{2}(2\alpha_{\ell} - \lambda\mu_{\ell}) > 0.$$

2. If $2\alpha_{\ell} < -\lambda \mu_{\ell}$, then $2\alpha_{\ell} + \lambda \mu_{\ell} \partial |\beta_{\ell}^*| < 0$, thus $\beta_{\ell}^* < 0$, giving $\partial |\beta_{\ell}^*| = -1$, then

$$\beta_{\ell}^* = \frac{1}{2}(2\alpha_{\ell} + \lambda\mu_{\ell}) < 0.$$

3. If $-\lambda \mu_{\ell} \leq 2\alpha_{\ell} \leq \lambda \mu_{\ell}$, then on the one hand, $\beta_{\ell}^* > 0$ leads to $\partial |\beta_{\ell}| = 1$, and thus $\beta_{\ell}^* \leq 0$, on the other hand, $\beta_{\ell}^* < 0$ produces $\partial |\beta_{\ell}| = -1$, and hence $\beta_{\ell}^* \geq 0$. Thus

$$\beta_{\ell}^* = 0$$

As we hoped, with the aid of soft threshold operator, we obtain

$$\beta_{\ell}^{*} = \frac{1}{2} (\max(0, 2\alpha_{\ell} - \lambda\mu_{\ell}) + \min(0, 2\alpha_{\ell} + \lambda\mu_{\ell})) \\ = \frac{1}{2} S_{\lambda\mu_{\ell}}(2\alpha_{\ell}).$$

2.3 Regularized barycentric interpolation formulae

In this subsection, we focus on the condition of L = N and the required interpolation conditions

$$p(x_j) = f(x_j), \quad j = 0, 1, \dots, N,$$

where $p \in \mathbb{P}_L$ is the interpolant of f. As pointed out by Wang, Huybrechs, & Vandewalle (2014), "barycentric interpolation is arguably the method of choice for numerical polynomial interpolation". It is significant to express the ℓ_2 -regularized approximation polynomial (2.3) and the ℓ_2 -regularized approximation polynomial (2.7) in a barycentric form (Berrut & Trefethen, 2004; Higham, 2004)

$$p(x) = \frac{\sum_{j=0}^{N} \frac{\Omega_j}{x - x_j} f(x_j)}{\sum_{j=0}^{N} \frac{\Omega_j}{x - x_j}}, \quad \Omega_j = \frac{1}{\prod_{k \neq j} (x_k - x_j)},$$

respectively. $\{\Omega_j\}_{j=0}^N$ are called barycentric weights. The study of the $\{\Omega_j\}_{j=0}^N$ for roots and extrema of classical orthogonal polynomials is well developed, see Salzer (1972), Schwarz & Waldvogel (1989), Berrut & Trefethen (2004), Wang & Xiang (2012) and Wang *et al.* (2014).

We first derive the ℓ_2 -regularized barycentric interpolation formula. Recall (2.3), we have

$$p_{L,N+1}^{\ell_2}(x) = \sum_{\ell=0}^{N} \frac{\sum_{j=0}^{N} \omega_j \tilde{\Phi}_{\ell}(x_j) f(x_j)}{1 + \lambda \mu_{\ell}^2} \tilde{\Phi}_{\ell}(x)$$
$$= \sum_{j=0}^{N} \omega_j f(x_j) \sum_{\ell=0}^{N} \frac{\tilde{\Phi}_{\ell}(x_j) \tilde{\Phi}_{\ell}(x)}{1 + \lambda \mu_{\ell}^2}.$$
(2.10)

From the orthonormality of $\tilde{\Phi}_{\ell}(x)$, $\ell = 0, 1, ..., N$, by letting $f(x) \equiv 1$ we have

$$\sum_{j=0}^{N} \omega_j \sum_{\ell=0}^{N} \frac{\tilde{\Phi}_{\ell}(x_j) \tilde{\Phi}_{\ell}(x)}{1 + \lambda \mu_{\ell}^2} = \sum_{\ell=0}^{N} \left(\sum_{j=0}^{N} \omega_j \tilde{\Phi}_{\ell}(x_j) \cdot 1 \right) \frac{\tilde{\Phi}_{\ell}(x)}{1 + \lambda \mu_{\ell}^2}$$
$$= \|\tilde{\Phi}_0(x)\|_{L_2} \frac{\tilde{\Phi}_0(x)}{1 + \lambda \mu_0^2} = \frac{1}{1 + \lambda \mu_0^2}.$$

Note that

$$1 \neq \sum_{j=0}^{N} \omega_j \sum_{\ell=0}^{N} \frac{\tilde{\Phi}_{\ell}(x_j)\tilde{\Phi}_{\ell}(x)}{1+\lambda\mu_{\ell}^2} \quad \text{when} \quad \lambda\mu_{\ell}^2 \neq 0,$$

due to the existence of regularization. Then the ℓ_2 -regularized approximation polynomial (2.10) can be expressed as

$$p_{L,N+1}^{\ell_2}(x) = \frac{\sum_{j=0}^{N} \left(\omega_j \sum_{\ell=0}^{N} \frac{\tilde{\Phi}_{\ell}(x_j) \tilde{\Phi}_{\ell}(x)}{1 + \lambda \mu_{\ell}^2} \right) f(x_j)}{(1 + \lambda \mu_0^2) \sum_{j=0}^{N} \omega_j \sum_{\ell=0}^{N} \frac{\tilde{\Phi}_{\ell}(x_j) \tilde{\Phi}_{\ell}(x)}{1 + \lambda \mu_{\ell}^2}}.$$
(2.11)

Without loss of generality, assume $\mu_{\ell} = 1$ for $\ell \geq N + 1$. Note that under this assumption, $\left\{\frac{\tilde{\Phi}_{\ell}(x)}{\sqrt{1+\lambda\mu_{\ell}^{2}}}\right\}_{\ell\in\mathbb{N}}$ is still a sequence of orthogonal polynomials. By the Christoffel-Darboux formula (Gautschi, 2004, Section 1.3.3), we rewrite $\sum_{\ell=0}^{N} \frac{\tilde{\Phi}_{\ell}(x_{\ell})\tilde{\Phi}_{\ell}(x)}{1+\lambda\mu_{\ell}^{2}}$ in the form

$$\sum_{\ell=0}^{N} \frac{\tilde{\Phi}_{\ell}(x)\tilde{\Phi}_{\ell}(x_{j})}{1+\lambda\mu_{\ell}^{2}} = \frac{k_{N}}{h_{N}k_{N+1}} \frac{\frac{\tilde{\Phi}_{N+1}(x)}{\sqrt{1+\lambda\mu_{N+1}^{2}}} \frac{\tilde{\Phi}_{N}(x_{j})}{\sqrt{1+\lambda\mu_{N}^{2}}} - \frac{\tilde{\Phi}_{N+1}(x_{j})}{\sqrt{1+\lambda\mu_{N+1}^{2}}} \frac{\tilde{\Phi}_{N}(x)}{\sqrt{1+\lambda\mu_{N}^{2}}} = \frac{k_{N}}{h_{N}k_{N+1}} \frac{\tilde{\Phi}_{N+1}(x)\tilde{\Phi}_{N}(x_{j})}{\sqrt{1+\lambda\mu_{N+1}^{2}}\sqrt{1+\lambda\mu_{N}^{2}}(x-x_{j})},$$
(2.12)

where k_{ℓ} and h_{ℓ} denote the leading coefficient and the L_2 norm of $\frac{\tilde{\Phi}_{\ell}(x)}{\sqrt{1+\lambda\mu_{\ell}^2}}$, respectively. Let us combine (2.12) with the other expression (2.11) of the ℓ_2 -regularized approximation polynomial and cancel the factor $\frac{k_N}{h_N k_{N+1}} \frac{\tilde{\Phi}_{N+1}(x)}{\sqrt{1+\lambda\mu_{N+1}^2}\sqrt{1+\lambda\mu_N^2}}$ from both the numerator and the denomi-

nator. Together with (2.11) and (2.12) we obtain the solution to the ℓ_2 -regularized approximation problem in a barycentric form, and we name it the ℓ_2 -regularized barycentric interpolation formula:

$$p_N^{\ell_2 - \text{bary}}(x) = \frac{\sum_{j=0}^N \frac{\Omega_j}{x - x_j} f(x_j)}{(1 + \lambda \mu_0^2) \sum_{j=0}^N \frac{\Omega_j}{x - x_j}},$$
(2.13)

where $\Omega_j = \omega_j \Phi_N(x_j)$ is the corresponding barycentric weight at x_j . This relation between barycentric weights and Gauss quadrature weights is revealed by Wang, Huybrechs, & Vandewalle (2014); however, this relation does not lead to the fast computation since it still requires evaluating orthogonal polynomials on \mathcal{X}_{N+1} . From the relation they also find the explicit barycentric weights for all classical orthogonal polynomials.

Secondly, we induce the ℓ_1 -regularized barycentric interpolation formula. The ℓ_1 -regularized approximation polynomial (2.7) can be expressed as the sum of two terms:

$$p_{L,N+1}^{\ell_1}(x) = \sum_{\ell=0}^N \frac{S_{\lambda\mu_\ell} \left(2\sum_{j=0}^N \omega_j \tilde{\Phi}_\ell(x_j) f(x_j)\right)}{2} \tilde{\Phi}_\ell(x)$$
$$= \sum_{\ell=0}^N \left(\sum_{j=0}^N \omega_j \tilde{\Phi}_\ell(x_j) f(x_j)\right) \tilde{\Phi}_\ell(x) + \sum_{\ell=0}^N c_\ell \tilde{\Phi}_\ell(x), \tag{2.14}$$

where

$$c_{\ell} = \frac{S_{\lambda\mu_{\ell}}(2\alpha_{\ell})}{2} - \alpha_{\ell}, \quad \ell = 0, 1, \dots, N.$$

The first term in (2.14) can be directly written in the barycentric form by letting $\lambda = 0$ using the ℓ_2 -regularized barycentric formula. Then let the basis $\{\tilde{\Phi}_\ell\}_{\ell=0}^N$ transform into Lagrange polynomials $\{\ell_j(x)\}_{j=0}^N$. By the basis-transformation relation between orthogonal polynomials and Lagrange polynomials (Gander, 2005), the second term in (2.14) can be represented by Lagrange polynomials in the form

$$\sum_{\ell=0}^{N} c_{\ell} \tilde{\Phi}_{\ell}(x) = \sum_{j=0}^{N} \left(\sum_{\ell=0}^{N} c_{\ell} \tilde{\Phi}_{\ell}(x_j) \right) \ell_j(x).$$
(2.15)

With the same procedure of obtaining the barycentric formula from the classical Lagrange interpolation formula in Berrut & Trefethen (2004), the second term (2.15) in (2.14) can be rewritten as

$$\sum_{\ell=0}^{N} c_{\ell} \tilde{\Phi}_{\ell}(x) = \frac{\sum_{j=0}^{N} \frac{\Omega_j}{x - x_j} \left(\sum_{\ell=0}^{N} c_{\ell} \tilde{\Phi}_{\ell}(x_j)\right)}{\sum_{j=0}^{N} \frac{\Omega_j}{x - x_j}}.$$
(2.16)

Together with (2.14) and (2.16), we obtain the ℓ_1 -regularized barycentric interpolation formula:

$$p_{N}^{\ell_{1}-\text{bary}}(x) = \frac{\sum_{j=0}^{N} \frac{\Omega_{j}}{x-x_{j}} \left(f(x_{j}) + \sum_{\ell=0}^{N} c_{\ell} \tilde{\Phi}_{\ell}(x_{j}) \right)}{\sum_{j=0}^{N} \frac{\Omega_{j}}{x-x_{j}}}.$$
(2.17)

If $\lambda = 0$, the basis is normalized Legendre polynomials (i.e., interpolation nodes are Legendre points) and $\Omega_j = (-1)^j \sqrt{(1-x_j^2)\omega_j}$ where ω_j is the Gauss quadrature weight at x_j (Wang & Xiang, 2012), then both the ℓ_2 -regularized barycentric interpolation formula (2.13) and the ℓ_1 -regularized barycentric interpolation formula (2.17) reduce to the Wang-Xiang formula (Wang & Xiang, 2012; Trefethen, 2013). Inspired by the work of Higham (2004), we will conduct numerical studies on both regularized barycentric interpolation formulae (2.13) and (2.17), such as numerical stability, see the next paper (An & Wu, 2019).

3 Quality of ℓ_2 -regularized weighted least squares approximation

In this section, we study the quality of the ℓ_2 -regularized weighted least squares approximation in terms of Lebesgue constants. As is known to all, the Lebesgue constant is a tool for quantifying the divergence or convergence of polynomial approximation. From 1910, a lot of works have been done on Lebesgue constants (Fejér, 1910; Szegö, 1939; Rivlin, 2003; Powell, 1981; Wang & Xiang, 2012; Trefethen, 2013). This paper considers Lebesgue constants in the case of regularization. The Lebesgue constant is the ∞ -norm of the linear mapping from data to approximation polynomial:

$$\Lambda_L := \sup_{f \neq 0} \frac{\|p_L\|_{\infty}}{\|f\|_{\infty}}.$$

In particular, we have the following estimation on the Λ_L as a consequence of (2.3).

Proposition 3.1 Adopt the notation and assumptions of Theorem 2.1. We have

$$\Lambda_{L} = \max_{x \in [-1, 1]} \sum_{j=1}^{N} w_{j} \Big| \sum_{\ell=0}^{L} \frac{1}{1 + \lambda \mu_{\ell}^{2}} \tilde{\Phi}_{\ell}(x) \tilde{\Phi}_{\ell}(x_{j}) \Big| \\ \leq \max_{x \in [-1, 1]} \sum_{j=0}^{N} w_{j} \sum_{\ell=0}^{L} \frac{1}{1 + \lambda \mu_{\ell}^{2}} \Big| \tilde{\Phi}_{\ell}(x) \tilde{\Phi}_{\ell}(x_{j}) \Big|.$$
(3.1)

3.1 Lebesgue constants with the basis of Chebyshev polynomials of the first kind

We mimic the discussion of the least squares approximation without regularization in Rivlin (2003, Section 2.4). We shall treat the case of normalized Chebyshev polynomials of the first kind $T_{\ell}(x), \ell = 0, 1, \ldots$, as the basis for \mathbb{P}_L . Primary results are also available in Powell (1967). Consider a weighted Fourier series of a given continuous function $g(\theta)$ over $[-\pi, \pi]$:

$$q_L(\theta) = \frac{\rho_{0,L}}{2}a_0 + \sum_{\ell=1}^L \rho_{\ell,L}(a_\ell \cos \ell \theta + b_\ell \sin \ell \theta),$$

where $a_0, a_1, \ldots, a_L, b_1, \ldots, b_L$ are Fourier coefficients defined as

$$a_{\ell} = \frac{1}{\pi} \int_{-\pi}^{\pi} g(t) \cos \ell t dt, \quad \ell = 0, 1, \dots, L,$$

and

$$b_{\ell} = \frac{1}{\pi} \int_{-\pi}^{\pi} g(t) \sin \ell t dt, \quad \ell = 1, \dots, L,$$

and weights $\rho_{\ell,L} = 1/(1 + \lambda \mu_{\ell}^2), \ \ell = 0, 1, ..., L.$

Lemma 3.1 (Rivlin, 2003) If $g(\theta)$ is continuous on $[-\pi, \pi]$ with period 2π , then

$$q_L(\theta) = \frac{1}{\pi} \int_{-\pi}^{\pi} g(t+\theta) u_L(t) dt,$$

where

$$u_L(t) = \frac{\rho_{0,L}}{2} + \sum_{\ell=1}^L \rho_{\ell,L} \cos \ell t.$$

Definition 3.1 Lebesgue constants for ℓ_2 -regularized least squares approximation using Chebyshev polynomials of the first kind are defined as

$$\Lambda_L := \frac{1}{\pi} \int_{-\pi}^{\pi} |u_L(t)| dt$$

The case of $\lambda = 0$ leads to Lebesgue constants for Fourier series (without regularization) (Rivlin, 2003, Section 2.4) in the form of

$$\Lambda_L = \frac{1}{\pi} \int_0^{\pi} \frac{|\sin(L + \frac{1}{2})t|}{\sin\frac{t}{2}} dt = \frac{1}{2\pi} \int_{-\pi}^{\pi} \left| \frac{\sin(L + \frac{1}{2})t}{\sin\frac{t}{2}} \right| dt,$$

where the last integrand is the famous Dirichlet kernel. For estimation of Λ_L , we have the following lemma.

Lemma 3.2 Let D_n denote the Dirichlet kernel (Stein & Shakarchi, 2011)

$$D_n(x) := \sum_{k=-n}^n e^{ikx} = \frac{\sin(n+\frac{1}{2})x}{\sin\frac{x}{2}}.$$

Then for $n \geq 2$,

$$\frac{1}{2\pi} \int_{-\pi}^{\pi} |D_n(x)| dx = \frac{4}{\pi^2} \log n + \mathcal{O}(1), \tag{3.2}$$

and

$$\frac{1}{2\pi} \int_{-\pi}^{\pi} |D_n(x)| dx \le \frac{4}{\pi^2} \log(n-1) + \eta,$$
(3.3)

where $\eta = \frac{4}{\pi^2} + \frac{2}{\pi} (1 + \int_0^\pi \frac{\sin x}{x} dx) = 2.220884...$

(3.2) is a known result given by Fejér (1910), one might find in Lorentz (1966, p.5) and Stein & Shakarchi (2011, Section 2.7.2). (3.3) is sharper than the known result $\Lambda_L < \frac{4}{\pi^2} \log n + 3$, $n \ge 2$ (Rivlin, 2003, Lemma 2.2). For completeness, the proof of Lemma 3.2 is given in Appendix: Proof of Lemma 3.2.

Theorem 3.1 Suppose f is continuous on [-1, 1], and normalized Chebyshev polynomials constitute a basis for \mathbb{P}_L . Then Lebesgue constants Λ_L for the ℓ_2 -regularized least squares approximation of degree L ($L \ge 2$) on [-1, 1] satisfy

$$\Lambda_L = \frac{4\log L/\pi^2}{1+\lambda\tilde{\mu}^2} + \mathcal{O}(1),$$
(3.4)

and

$$\Lambda_L \le \frac{4\log(L-1)/\pi^2 + \eta}{1 + \lambda \underline{\mu}^2},\tag{3.5}$$

where $\underline{\mu} = \min\{\mu_0, \mu_1, \dots, \mu_L\}, \ \tilde{\mu} \in \mathbb{R}$ which satisfies $\min\{\mu_0, \mu_1, \dots, \mu_L\} \le \tilde{\mu} \le \max\{\mu_0, \mu_1, \dots, \mu_L\}$ and $\eta = 2.220884...$

Proof. Since f is continuous on [-1, 1], then $g(\theta) = f(\cos \theta)$ is continuous on $[0, \pi]$. If $g(-\theta) = g(\theta)$, then g is continuous on $[-\pi, \pi]$. The even function g gives $b_{\ell} = 0$ for all $\ell = 1, \ldots, L$, and then

$$q_{L}(\theta) = \frac{\rho_{0,L}}{2} a_{0} + \sum_{\ell=1}^{L} \rho_{\ell,L} a_{\ell} \cos \ell \theta$$

$$= \frac{\rho_{0,L}}{2} a_{0} + \sum_{\ell=1}^{L} \rho_{\ell,L} a_{\ell} T_{\ell}(x)$$
(3.6)
$$= \frac{\sqrt{\pi/2} \rho_{0,L}}{2} a_{0} \tilde{T}_{0} + \sum_{\ell=1}^{L} \sqrt{\pi} \rho_{\ell,L} a_{\ell} \tilde{T}_{\ell}(x),$$

which reveals that this is the ℓ_2 -regularized least squares approximation of degree L with the basis for \mathbb{P}_L being normalized Chebyshev polynomials of the first kind. Since $g(\theta)$ is continuous on $[-\pi, \pi]$ with period 2π , there must exist $M \ge 0$ such that

$$|g(t+\theta)| \le M, \quad t, \theta \in [-\pi, \pi].$$

By Lemma 3.1,

$$\max_{\theta \in [-\pi,\pi]} |q_L(\theta)| \le M\Lambda_L.$$

When $\lambda = 0$, one may easily verify that

$$u_L(t) = \frac{1}{2} + \sum_{\ell=0}^{L} \cos \ell t = \frac{\sin(L + \frac{1}{2})t}{2\sin\frac{t}{2}},$$

then by Lemma 3.2,

$$\Lambda_L = \frac{1}{2\pi} \int_{-\pi}^{\pi} \left| \frac{\sin(L + \frac{1}{2})t}{\sin\frac{t}{2}} \right| dt \le \frac{4}{\pi^2} \log(L - 1) + \eta,$$

and

$$\Lambda_L = \frac{4}{\pi^2} \log L + \mathcal{O}(1).$$

Let $\underline{\mu} = \min\{\mu_0, \mu_1, \dots, \mu_L\}$ and $\overline{\mu} = \max\{\mu_0, \mu_1, \dots, \mu_L\}$. And let $\tilde{\mu}$ be a real number satisfying $\underline{\mu} \leq \underline{\tilde{\mu}} \leq \overline{\mu}$. If $\lambda > 0$, then

$$u_L(t) \le \frac{1}{1+\lambda\underline{\mu}^2} \left(\frac{1}{2} + \sum_{\ell=0}^L \cos\ell t\right) = \frac{1}{1+\lambda\underline{\mu}^2} \frac{\sin(L+\frac{1}{2})t}{2\sin\frac{t}{2}},$$

and

$$u_L(t) = \frac{1}{1 + \lambda \tilde{\mu}^2} \left(\frac{1}{2} + \sum_{\ell=0}^L \cos \ell t \right) = \frac{1}{1 + \lambda \tilde{\mu}^2} \frac{\sin(L + \frac{1}{2})t}{2\sin\frac{t}{2}},$$

which gives the asymptotic result (3.4) and the inequality (3.5).

Take the family of normalized Chebyshev polynomials of the first kind $\{\tilde{T}_{\ell}(x)\}_{\ell=0}^{L}$ as the basis for \mathbb{P}_{L} and the set of zeros of $\tilde{T}_{L+1}(x)$ as the set of nodes. Setting L = N and $\lambda = 10^{-1}$. Fig. 1 illustrates Lebesgue constants with respect to different choices of regularization parameter λ .

3.2 Lebesgue constants with the basis of Legendre polynomials

In this subsection, we will derive asymptotic bounds of Lebesgue constants of the ℓ_2 - regularized approximation by using Legendre polynomials. Consider the kernel

$$K_L(x,y) := \sum_{\ell=0}^{L} \frac{2\ell+1}{1+\lambda\mu_{\ell}^2} P_{\ell}(x) P_{\ell}(y),$$

where $P_{\ell}(\cdot)$ denotes the Legendre polynomial of degree ℓ . The case of $\lambda = 0$ gives a simple kernel

$$T_L(x,y) = \sum_{\ell=0}^{L} (2\ell+1)P_\ell(x)P_\ell(y) = (L+1)\frac{P_L(x)P_{L+1}(y) - P_{L+1}(x)P_L(y)}{y-x}$$

where the last equality is due to Gronwall (1913, (4)). Thus,

$$K_L(x) \triangleq K_L(x,1) := \sum_{\ell=0}^{L} \frac{2\ell+1}{1+\lambda\mu_{\ell}^2} P_{\ell}(x),$$



Figure 1: The Lebesgue constant of ℓ_2 -regularized approximation with L = N and $\mu_{\ell} = 1$ for $\ell = 0, 1, \ldots, L$ using Chebyshev polynomials of the first kind

and

$$T_L(x) \triangleq T_L(x,1) := \sum_{\ell=0}^{L} (2\ell+1)P_\ell(x) = (L+1)\frac{P_L(x) - P_{L+1}(x)}{1-x}$$

where the rightmost equality is due to Gronwall (1913, (5)). The reader may note that

$$|K_L(x)| \le \frac{1}{1+\lambda\mu^2} |T_L(x)|,$$
(3.7)

where $\underline{\mu} = \min\{\mu_0, \mu_1, \dots, \mu_L\}.$

Definition 3.2 Lebesgue constants for the ℓ_2 -regularized approximation using Legendre polynomials are defined as

$$\Lambda_L := \frac{1}{2} \int_{-1}^1 |K_L(x)| dx.$$

The case of $\lambda = 0$ leads to

$$\Theta_L := \frac{1}{2} \int_{-1}^{1} |T_L(x)| dx = \frac{L+1}{2} \int_{-1}^{1} \left| \frac{P_L(x) - P_{L+1}(x)}{1-x} \right| dx,$$
(3.8)

which is the definition of Lebesgue constant of Legendre truncation of degree L (Gronwall, 1913).

Lemma 3.3 (Gronwall, 1913 and Szegö, 1934) Let Θ_L be defined as (3.8). Then

$$\lim_{L \to \infty} \frac{\Theta_L}{\sqrt{L}} = 2\sqrt{\frac{2}{\pi}}.$$
(3.9)

Combining Lemma (3.3) with the inequality (3.7), we obtain the estimation of Λ_L in the case of Legendre polynomials.

Theorem 3.2 Suppose f is continuous on [-1,1], and Legendre polynomials constitute the basis for \mathbb{P}_L . Then Lebesgue constants Λ_L for the ℓ_2 -regularized least squares approximation of degree L ($L \ge 2$) on [-1,1] satisfy

$$\Lambda_L \le \frac{1}{1 + \lambda \underline{\mu}^2} \left(\frac{2^{3/2}}{\sqrt{\pi}} L^{1/2} + o(L^{1/2}) \right),$$

where $\underline{\mu} = \min\{\mu_0, \mu_1, \dots, \mu_L\}.$

The proof for Theorem 3.2 is based on the above discussion.

4 Sparsity of solution to the ℓ_1 -regularized approximation problem

Some real-world problems, such as signal processing, often have sparse solutions. One may seek the sparsest solution to a problem, that is, the solution containing zero elements at most. However, a vector of real data would rarely contains many strict zeros. One may introduce other methods to seek sparsity, such as $\min_x ||x||_p$, where $||x||_p = (\sum_i |x_i|^p)^{1/p}$, 0 . Nevertheless, these optimization problems mentioned above are nonconvex and nondifferentiable (Clarke, 1990; Bruckstein*et al.* $, 2009). Regularized methods, especially <math>\ell_1$ -regularized methods, also produce sparse solutions, according to our examples. One may find a relatively sparse solution by minimizing ℓ_1 norm, because such an optimization problem is a convex optimization problem, and its solution is the closest one to the sparsest solution for all $p \ge 1$ in $||x||_p$. For topics on sparsity, we refer to Bruckstein *et al.* (2009). We consider the sparsity of the solution β to the ℓ_1 -regularized approximation problem (1.6). The sparsity is measured by the number of nonzero elements of β , denoted as $||\beta||_0$, also known as the zero "norm" (it is not a norm actually) of β .

Before discussing the upper bound of $\|\beta\|_0$, we offer a quick glimpse into the zero elements distribution of the ℓ_1 -regularized approximation solution. From Definition 2.1 of the soft threshold operator, we have the following result.

Proposition 4.1 (Zero elements distribution of the ℓ_1 -regularized approximation solution) Adopt the notation and assumptions of Lemma 2.1. If μ_ℓ satisfies

$$-\lambda \mu_{\ell} \le 2 \sum_{j=0}^{N} \omega_j \tilde{\Phi}_{\ell}(x_j) f(x_j) \le \lambda \mu_{\ell},$$
(4.1)

then its corresponding β_{ℓ} is zero, $\ell = 0, 1, \ldots, L$.

If $\lambda > 0$, then $\|\mathbf{A}_{L}^{T}\mathbf{W}\mathbf{f}\|_{0}$ becomes an upper bound of the number of nonzero elements of $\boldsymbol{\beta}$. Furthermore, we obtain the exact number of nonzero elements of $\boldsymbol{\beta}$ with the help of the information of $\boldsymbol{\beta}$.

Theorem 4.1 Let $\boldsymbol{\beta} = [\beta_0, \beta_1, \dots, \beta_L]^T$ be the solution to the ℓ_1 -regularized problem (1.6). If $\lambda > 0$, then the number of nonzero elements of $\boldsymbol{\beta}$ satisfies

$$\|\boldsymbol{\beta}\|_0 \le \|\mathbf{A}_L^T \mathbf{W} \mathbf{f}\|_0, \tag{4.2}$$

and

$$\|\boldsymbol{\beta}\|_{0} = \|\mathbf{A}_{L}^{T}\mathbf{W}\mathbf{f}\|_{0} - \#\{occurrences \ of \ \beta_{\ell} = 0 \ but \ \alpha_{\ell} \neq 0\},$$
(4.3)

where $\#\{$ occurrences of $\beta_{\ell} = 0$ but $\alpha_{\ell} \neq 0 \}$ denotes the number of occurrences of $\beta_{\ell} = 0$ but $\alpha_{\ell} \neq 0$ for $\ell = 0, 1, ..., L$.

Proof. The first order condition (2.8) of the ℓ_1 -regularized approximation problem (1.6) in the matrix form can be rewritten as

$$oldsymbol{eta} \in \mathbf{A}_L^T \mathbf{W} \mathbf{f} - rac{\lambda \partial (\|\mathbf{R}_L oldsymbol{eta}\|_1)}{2}$$

To obtain a solution, there must exist an L + 1 vector $\mathbf{h} = [h_0, h_1, \dots, h_L]^T \in \partial(||\mathbf{R}_L \boldsymbol{\beta}||_1)$ such that

$$\boldsymbol{\beta} = \mathbf{A}_L^T \mathbf{W} \mathbf{f} - \frac{\lambda \mathbf{h}}{2}.$$
(4.4)

Since $\mu_{\ell} > 0$ for all $\ell = 0, 1, \dots, L$, elements of **h** satisfy

$$h_{\ell} = \begin{cases} \mu_{\ell}, & \mu_{\ell}\beta_{\ell} > 0, \text{ i.e., } \beta_{\ell} > 0\\ -\mu_{\ell}, & \mu_{\ell}\beta_{\ell} < 0, \text{ i.e., } \beta_{\ell} < 0\\ r\mu_{\ell} \ \forall r \in [-1, 1], & \mu_{\ell}\beta_{\ell} = 0, \text{ i.e., } \beta_{\ell} = 0, \end{cases}$$

yielding $\|\mathbf{h}\|_0 \ge \|\boldsymbol{\beta}\|_0$. Expression (4.4) gives

$$\left| \boldsymbol{\beta} + \frac{\lambda \mathbf{h}}{2} \right\|_{0} = \left\| \mathbf{A}_{L}^{T} \mathbf{W} \mathbf{f} \right\|_{0}.$$

If $\beta_{\ell} > (\text{or } <)0$, then $h_{\ell} > (\text{or } <)0$. If $\beta_{\ell} = 0$, whereas h_{ℓ} may not be zero. Thus,

$$\left\|\boldsymbol{\beta} + \frac{\lambda \mathbf{h}}{2}\right\|_0 = \|\mathbf{h}\|_0.$$

Hence,

$$\|\boldsymbol{\beta}\|_0 \leq \|\mathbf{h}\|_0 = \|\mathbf{A}_L^T \mathbf{W} \mathbf{f}\|_0.$$

Let β_{ℓ}^* denote the optimal solution to the problem (2.9). With the aid of the closed-form solution to the ℓ_1 -regularized approximation problem (see (2.6)), expression (4.4) gives birth to

$$\frac{h_{\ell}}{\mu_{\ell}} = \frac{2}{\lambda \mu_{\ell}} (\alpha_{\ell} - \beta_{\ell}^*) = \begin{cases} 1, & \beta_{\ell}^* > 0, \\ -1, & \beta_{\ell}^* < 0, \\ \frac{2\alpha_{\ell}}{\lambda \mu_{\ell}}, & \beta_{\ell}^* = 0. \end{cases}$$

Due to $\|\mathbf{h}\|_0 = \left\|\frac{\mathbf{h}}{\mu}\right\|_0$, where $\frac{\mathbf{h}}{\mu}$ denotes the pointwise division between \mathbf{h} and μ , the difference between $\|\mathbf{h}\|_0$ and $\|\boldsymbol{\beta}\|_0$ is expressed as

$$\|\mathbf{h}\|_{0} - \|\boldsymbol{\beta}\|_{0} = \left\|\frac{\mathbf{h}}{\boldsymbol{\mu}}\right\|_{0} - \|\boldsymbol{\beta}\|_{0} = \#\{\text{occurrences of } \boldsymbol{\beta}_{\ell} = 0 \text{ but } \boldsymbol{\alpha}_{\ell} \neq 0\}.$$
(4.5)

Together with $\|\mathbf{h}\|_0 = \|\mathbf{A}_{\mathrm{L}}^{\mathrm{T}} \mathbf{W} \mathbf{f}\|_0$ and (4.5), we obtain the exact number (4.3) of nonzero elements of $\boldsymbol{\beta}$.

Corollary 4.1 If $\lambda = 0$, then the number of nonzero elements of β satisfies

$$\|\boldsymbol{\beta}\|_0 = \|\mathbf{A}_L^T \mathbf{W} \mathbf{f}\|_0. \tag{4.6}$$

REMARK. Together, Theorem 4.1 and Corollary 4.1 state that the regularized minimization is better than minimization without regularization in terms of sparsity.

Let the basis for \mathbb{P}_L be the family of normalized Chebyshev polynomials of the first kind $\{\tilde{T}_{\ell}(x)\}_{\ell=0}^L$ and the set of nodes be the set of zeros of $\tilde{T}_{N+1}(x)$. With degree L of approximation polynomial ranging from 1 to 60, λ evaluated 10^{-1} and μ_{ℓ} evaluated 1 for all $\ell = 0, 1, \ldots, L$, Fig. 2 gives four examples on the bounds and estimations given above.

5 Numerical experiments

In this section, we report numerical results to illustrate the theoretical results derived above and test the efficiency of the ℓ_2 -regularized approximation polynomial $p_{L,N+1}^{\ell_2}$ (2.3) and the ℓ_1 -regularized approximation polynomial $p_{L,N+1}^{\ell_1}$ (2.7). The choices of the basis for \mathbb{P}_L and the set of nodes \mathcal{X}_{N+1} are primary when using both models. We choose Chebyshev polynomials of the first kind and the corresponding Chebyshev points. Certainly, choosing other orthogonal polynomials such as Legendre polynomials is also possible. All computations are performed in MATLAB in double precision arithmetic. Some related commands, for instance, obtaining quadrature points and weights, are included in CHEBFUN 5.7.0 (Trefethen *et al.*, 2017).

To test the efficiency of approximation, we define the uniform error and the L_2 error to measure the approximation error:

• The uniform error of the approximation is estimated by

$$\|f(x) - p_{L,N+1}(x)\|_{\infty} := \max_{x \in [-1,1]} |f(x) - p_{L,N+1}(x)|$$
$$\simeq \max_{x \in \mathcal{X}} |f(x) - p_{L,N+1}(x)|,$$

where \mathcal{X} is a large but finite set of well distributed points (for example, clustered grids, see Trefethen (2000, Chapter 5)) over the interval [-1, 1].



Figure 2: Examples on bounding the number of nonzero elements, where # denotes #{occurrences of $\beta_{\ell} = 0$ but $\alpha_{\ell} \neq 0$ }

• The L_2 error of the approximation is estimated by a proper Gauss quadrature rule:

$$\|f(x) - p_{L,N+1}(x)\|_{L_2} = \left(\int_{-1}^1 w(x)(f(x) - p_{L,N+1}(x))^2 dx\right)^{1/2}$$
$$\simeq \left(\sum_{j=0}^N \omega_j(f(x_j) - p_{L,N+1}(x_j))^2\right)^{1/2}.$$

5.1 Regularized approximation models for exact data

The fact should always stick in readers' mind that regularization is introduced to solve ill-posed problems or to prevent overfitting. When approximation applies to functions without noise, regularization parameter $\lambda = 0$ (no regularization) contributes to the best choice of approximating. Fig. 3 reports the efficiency and errors for approximating function

$$f_1(x) = \tanh(20\sin(12x)) + 0.02e^{3x}\sin(300x),$$

with or without regularization over [-1, 1]. The test function is given in Trefethen (2013). Let $N = 600, L = 200, \lambda = 10^{-1}$ and $\mu_{\ell} = 1$ for all $\ell = 0, 1, ..., L$. Fig. 3 illustrates that regularization is beyond use in this well-posed approximation problem, and ℓ_2 -regularization is better than ℓ_1 -regularization in approximating smooth functions.

5.2 Regularized approximation models for contaminated data

We consider

$$f_2(x) = \begin{cases} \frac{5\sin(5\pi x)}{5\pi x}, & x \neq 0, \\ 5, & x = 0, \end{cases}$$
(5.1)

which is the Fourier transform of the gate signal

$$g(t) = \begin{cases} 1, & |t| \le 5/2, \\ 0, & |t| > 5/2, \end{cases}$$



Figure 3: Approximation of function $f_1(x) = \tanh(20\sin(12x)) + 0.02e^{3x}\sin(300x)$ with exact values on the Chebyshev polynomials of the first kind

see Bracewell (1965). We use regularized least squares models to reduce Gaussian white noise added to the function (5.1) with the signal-noise ratio (SNR) 10 dB. The choice of λ is critical in these models, so we first consider the relation between λ and approximation errors to choose the optimal λ . Let L = 30 and N = 100. We take $\lambda = 10^{-15}$, $10^{-14.5} \ 10^{-14}$, $\dots, 10^{4.5}$, 10^5 to choose the best regularization parameter. Here we choose $\lambda = 10^{-1}$. For more advanced methods to choose the parameter λ , we refer to Lazarov *et al.* (2007) and Pereverzyev *et al.* (2015) for a further discussion.

Fig. 4 shows that the ℓ_2 - and ℓ_1 -regularized approximation models with $\lambda = 10^{-1}$ are effective in recovering the noisy function. In the case we let

$$\mu_{\ell} = \frac{1}{F(\ell/L)}, \quad \ell = 0, 1, \dots, L,$$

where the filter function F is defined as (An *et al.*, 2012)

$$F(x) := \begin{cases} 1, & x \in [0, 1/2], \\ \sin^2 \pi x, & x \in [1/2, 1], \\ 0, & x \in [1, +\infty]. \end{cases}$$

In this case, $\{\mu_\ell\}_{\ell=0}^L$ is a sequence of nonnegative nondecreasing parameters.

Results in Fig. 5 illustrate that the ℓ_1 -regularized approximation model is the best choice when recovering a contaminated function, which accords with the known facts (Lu & Pereverzev, 2009). Both Fig. 4 and Fig. 5 show that regularized models are better than those without regularization ($\lambda = 0$).

Besides this, consider the highly oscillatory function

$$f_3(x) = \operatorname{Airy}(40x), \quad x \in [-1, 1],$$

with 12dB Gauss white noise added (noisy function is shown in Fig. 6). We use regularized barycentric formulae (2.13) and (2.17) to conduct this experiment. Let L = N = 500 and $\{\mu_\ell\}_{\ell=0}^L$ be the same as above. Different values of λ , say 10^{-1} , 10^{-2} , 10^{-5} , lead to different results, see Fig. 6. This experiment indicates that one could apply a simple formula to reduce noise, rather than employ an iterative scheme.

These numerical examples illustrate that for some problems, ℓ_2 -regularization also can be better than ℓ_1 -regularization. For example, $\lambda = 10^{-1}$ suits ℓ_2 -regularization, but almost straightens



Figure 4: Regularized approximation models with L = 30 and N = 100 to recover the function (5.1) from contaminated data on the Chebyshev polynomials of the first kind



Figure 5: Errors for regularized approximation model to recover $f_2(x)$ with fixed N = 100 on the Chebyshev polynomials of the first kind



Figure 6: Denoising by regularized barycentric formulae with L = N = 500 on Chebyshev points of the first kind: function $f_3(x) = \text{Airy}(40x)$ with 12dB Gauss white noise added

the function with ℓ_1 -regularization. Besides this, we can see that ℓ_2 -regularization contains lower sensitivity than ℓ_1 -regularization.

6 Concluding remarks

In this paper, we have investigated minimizers to the ℓ_2- and ℓ_1 -regularized least squares approximation problems with the aid of Gauss quadrature points and orthogonal polynomials on the interval [-1, 1]. Based on those explicit constructed approximation polynomials (2.3) and (2.7), the ℓ_2 -regulariarized barycentric interpolation formula (2.13) and the ℓ_1 -regulariarized barycentric interpolation formula (2.13) and the ℓ_1 -regulariarized barycentric interpolation formula (2.17) have been derived. In addition, Lebesgue constants are studied in the case of using Legendre polynomials and normalized Chebyshev polynomials of the first kind as the basis for the polynomial space \mathbb{P}_L . A bound for sparsity of the solution to ℓ_1 -regularized approximation polynomial and the ℓ_1 -regularized approximation polynomial and the ℓ_1 -regularized approximation polynomial are practicable and efficient. Regularization parameter choice strategies and error bounds of approximation should be studied in future work. These results provide new insights into the ℓ_2- and ℓ_1 -regularized approximation, and can be adapted to many practical applications such as noise reduction by using the barycentric interpolation scheme on Gauss quadrature points.

References

- AN, C., CHEN, X., SLOAN, I. H. & WOMERSLEY, R. S. (2012) Regularized least squares approximations on the sphere using spherical designs. *SIAM Journal on Numerical Analysis*, 50, 1513–1534.
- AN, C. & WU, H.-N. (2019) The numerical stability of regularized barycentric interpolation formulae for interpolation and extrapolation. arXiv preprint arXiv:1904.07187.
- BERRUT, J.-P. & TREFETHEN, L. N. (2004) Barycentric Lagrange interpolation. SIAM Review, 46, 501–517.

BRACEWELL, R. (1965) The Fourier Transform and Its Applications. New York: McGraw-Hill.

- BRUCKSTEIN, A. M., DONOHO, D. L. & ELAD, M. (2009) From sparse solutions of systems of equations to sparse modeling of signals and images. *SIAM Review*, **51**, 34–81.
- CAI, J.-F., OSHER, S. & SHEN, Z. (2009) Split Bregman methods and frame based image restoration. *Multiscale Modeling & Simulation*, 8, 337–369.
- CLARKE, F. H. (1990) Optimization and nonsmooth analysis, vol. 5. Philadelphia: SIAM.
- DONOHO, D. L. & JOHNSTONE, J. M. (1994) Ideal spatial adaptation by wavelet shrinkage. Biometrika, 81, 425–455.
- FEJÉR, L. (1910) Lebesguessche konstanten und divergente fourierreihen. Journal f
 ür die Reine und Angewandte Mathematik, 138, 22–53.
- GANDER, W. (2005) Change of basis in polynomial interpolation. Numerical Linear Algebra with Applications, 12, 769–778.
- GAUTSCHI, W. (2004) Orthogonal Polynomials: Computation and Approximation. Oxford: Oxford University Press on Demand.
- GAUTSCHI, W. (2012) Numerical analysis 2nd edition. Basel: Birkhäuser Basel.
- GOLITSCHEK, M. & SCHUMAKER, L. (1990) Data fitting by penalized least squares. Algorithms for Approximation II. New York: Springer-Verlag, pp. 210–227.
- GRONWALL, T. H. (1913) Über die Laplacesche reihe. Mathematische Annalen, 74, 213–270.
- HIGHAM, N. J. (2004) The numerical stability of barycentric Lagrange interpolation. IMA Journal of Numerical Analysis, 24, 547–556.
- KIM, S.-J., KOH, K., BOYD, S. & GORINEVSKY, D. (2009) ℓ_1 trend filtering. *SIAM Review*, **51**, 339–360.
- KRESS, R. (1998) Numerical analysis. Graduate Texts in Mathematics, vol. 181. New York: Springer-Verlag.
- LAZAROV, R. D., LU, S. & PEREVERZEV, S. V. (2007) On the balancing principle for some problems of numerical analysis. *Numerische Mathematik*, **106**, 659–689.
- LORENTZ, G. G. (1966) Approximation of functions. Athena Series: Selected Topics in Mathematics. New York Chicago San Francisco Toronto London: Holt, Rinehart and Winston.
- LU, S. & PEREVERZEV, S. V. (2009) Sparse recovery by the standard Tikhonov method. Numerische Mathematik, 112, 403–424.
- PEREVERZYEV, S. V., SLOAN, I. H. & TKACHENKO, P. (2015) Parameter choice strategies for least-squares approximation of noisy smooth functions on the sphere. SIAM Journal on Numerical Analysis, 53, 820–835.
- POWELL, M. J. D. (1967) On the maximum errors of polynomial approximations defined by interpolation and by least squares criteria. *The Computer Journal*, **9**, 404–407.
- POWELL, M. J. D. (1981) Approximation theory and methods. Cambridge: Cambridge University Press.
- RIVLIN, T. J. (2003) An introduction to the approximation of functions. North Chelmsford: Courier Corporation.
- SALZER, H. E. (1972) Lagrangian interpolation at the Chebyshev points xn, $\nu \equiv \cos (\nu \pi/n)$, $\nu = 0$ (1) n; some unnoted advantages. The Computer Journal, 15, 156–159.
- SCHWARZ, H. R. & WALDVOGEL, J. (1989) Numerical analysis: a comprehensive introduction. New York: Wiley.

- STEIN, E. M. & SHAKARCHI, R. (2011) *Fourier analysis: an introduction*, vol. 1. Princeton: Princeton University Press.
- SZEGÖ, G. (1934) Über einige asymptotische entwicklungen der Legendreschen funktionen. Proceedings of the London Mathematical Society, 2, 427–450.
- SZEGÖ, G. (1939) Orthogonal polynomials. Colloquium Publications Volume XXIII, vol. 23. Providence, Rhode Island: American Mathematical Society.
- TREFETHEN, L. N. (2000) Spectral methods in MATLAB, vol. 10. Philadelphia: SIAM.
- TREFETHEN, L. N. (2013) Approximation theory and approximation practice, vol. 128. Philadelphia: SIAM.
- TREFETHEN, L. N. et al. (2017) Chebfun Version 5.7.0. http://www/maths.ox.ac.uk/chebfun/: Chebfun Development Team, Chebfun Development Team.
- WANG, H., HUYBRECHS, D. & VANDEWALLE, S. (2014) Explicit barycentric weights for polynomial interpolation in the roots or extrema of classical orthogonal polynomials. *Mathematics of Computation*, 83, 2893–2914.
- WANG, H. & XIANG, S. (2012) On the convergence rates of Legendre approximation. Mathematics of Computation, 81, 861–877.
- XIANG, H. & ZOU, J. (2013) Regularization with randomized SVD for large-scale discrete inverse problems. *Inverse Problems*, 29, 085008.
- ZHOU, Y. & CHEN, X. (2018) Spherical t_{ϵ} -designs for approximations on the sphere. Mathematics of Computation, 87, 2831–2855.

Appendix A. Proof of Lemma 3.2

Proof. We adopt the proof of Rivlin (2003, Lemma 2.2), but bring up a sharper bound of this inequality:

$$\frac{1}{2\pi} \int_{-\pi}^{\pi} |D_n(x)| dx = \frac{1}{2\pi} \int_{-\pi}^{\pi} \left| \frac{\sin(n + \frac{1}{2})x}{\sin\frac{x}{2}} \right| dx < \frac{4}{\pi^2} \log n + 3, \qquad n \ge 1.$$
(.1)

When n = 1, $\frac{1}{2\pi} \int_{-\pi}^{\pi} |D_n(x)| dx = 1$. Suppose $n \ge 2$, then

$$\frac{1}{2\pi} \int_{-\pi}^{\pi} \left| \frac{\sin(n+\frac{1}{2})x}{\sin\frac{x}{2}} \right| dx = \frac{1}{\pi} \int_{0}^{\pi} \frac{|\sin(n+\frac{1}{2})x|}{\sin\frac{x}{2}} dx$$
$$= \frac{1}{\pi} \int_{0}^{\pi} \left| \frac{\sin nx}{\tan\frac{x}{2}} + \cos nx \right| dx$$
$$\leq \frac{1}{\pi} \int_{0}^{\pi} \frac{|\sin nx|}{\tan\frac{x}{2}} dx + \frac{1}{\pi} \int_{0}^{\pi} |\cos nx| dx. \tag{.2}$$

We have

$$\frac{1}{\pi} \int_0^\pi |\cos nx| dx = \frac{2}{\pi},$$
(.3)

and

$$\int_{0}^{\pi} \frac{|\sin nx|}{\tan \frac{x}{2}} dx \le 2 \int_{0}^{\pi} \frac{\sin x}{x} dx + \frac{4}{\pi} [1 + \log(n-1)].$$
(.4)

Together with (.2), (.3) and (.4), we obtain

$$\frac{1}{2\pi} \int_{-\pi}^{\pi} \Big| \frac{\sin(n+\frac{1}{2})x}{\sin\frac{x}{2}} \Big| dx \le \frac{2}{\pi} \int_{0}^{\pi} \frac{\sin x}{x} dx + \frac{4}{\pi^{2}} [1 + \log(n-1)] + \frac{2}{\pi}.$$

Thus,

$$\frac{1}{2\pi} \int_{-\pi}^{\pi} |D_n(x)| dx \le \frac{4}{\pi^2} \log(n-1) + \eta, \tag{.5}$$

where

$$\eta := \frac{4}{\pi^2} + \frac{2}{\pi} + \frac{2\int_0^\pi \frac{\sin x}{x} dx}{\pi}.$$

The value of η can be calculated by MATLAB with the function quadgk as

$$\eta := \frac{4}{\pi^2} + \frac{2}{\pi} + \frac{2\int_0^\pi \frac{\sin x}{x} dx}{\pi} = 2.220884\dots$$

It is not difficult to see inequality (.5) is sharper than inequality (.1) for $n \ge 2$.