## Analysis of nonsmooth stochastic approximation: the differential inclusion approach \*

SZYMON MAJEWSKI<sup>†</sup>, BLAZEJ MIASOJEDOW<sup>‡</sup>, AND ERIC MOULINES <sup>§</sup>

Abstract. In this paper we address the convergence of stochastic approximation when the functions to be minimized are not convex and nonsmooth. We show that the "mean-limit" approach to the convergence which leads, for smooth problems, to the ODE approach can be adapted to the non-smooth case. The limiting dynamical system may be shown to be, under appropriate assumption, a differential inclusion. Our results expand earlier works in this direction by [11] and provide a general framework for proving convergence for unconstrained and constrained stochastic approximation problems, with either explicit or implicit updates. In particular, our results allow us to establish the convergence of stochastic subgradient and proximal stochastic gradient descent algorithms arising in a large class of deep learning and high-dimensional statistical inference with sparsity inducing penalties.

Key words. Stochastic Approximation, Subgradient algorithm, Stochastic Proximal Gradient, Proximal operator, Differential Inclusions

**1. Introduction.** Stochastic approximation algorithms are stochastic processes defined iteratively as

(1) 
$$x_k = x_{k-1} + \gamma_k Y_k$$

where  $x_k$  takes value in  $\mathbb{R}^d$ ,  $\{Y_k, k \in \mathbb{N}\}$  is a sequence of random variable and  $\{\gamma_k, k \in \mathbb{N}^*\}$  is a sequence of stepsizes satisfying  $\gamma_k > 0$ ,  $\lim_{k\to\infty} \gamma_k = 0$  and  $\sum_{k=1}^{\infty} \gamma_k = \infty$ . The decreasing stepsizes imply that the rate of change of the parameter decreases as k goes to infinity, thus providing an implicit averaging. The value  $x_k$  might represent the current fit of a parameter or the state of a system and  $Y_k = \Phi(x_{k-1}, \xi_k)$  is a (measurable) function of the past fit  $x_{k-1}$  of the parameter and a new information observed at time k. Such problems have been considered in the early work by [35] and since found numerous applications, especially in machine learning and computational statistics.

A powerful method to analyze stochastic gradient algorithm, introduced in the early works by [28] and [26] is the ordinary differential equation (ODE) method. The ODE method has led to an enormous literature; see for example [10], [27] and the references therein. The ODE method can be informally summarized as follows: first we rewrite

$$Y_k = F(x_{k-1}) + \eta$$

where  $F : \mathbb{R}^d \to \mathbb{R}^d$  is a locally-Lipschitz vector field defined by an appropriate averaging and  $\eta_k = Y_k - F(x_{k-1})$ . If  $\{\xi_k, k \in \mathbb{N}\}$  is an i.i.d. sequence and  $\mathbb{E}[\|\Phi(x,\xi)\|] < \infty$  for all  $x \in \mathbb{R}^d$ , one may take for example  $F(x) = \mathbb{E}[\Phi(x,\xi_1)]$  and  $\eta_k = Y_k - F(x_{k-1})$  which then is a martingale increment sequence. The situation becomes more complex when the noise  $\{\xi_k, k \in \mathbb{N}\}$  is no longer i.i.d.. Of particular importance is the case where the conditional distribution of  $\xi_k$  given the past  $\{(x_j, \xi_j) : j \in \{0, \dots, k-1\}\}$  is Markovian (see for example [3]). The ODE approach to analyze the asymptotic behavior of the sequence  $\{x_k, k \in \mathbb{N}\}$  is to consider them as approximated solutions of the ODE  $\dot{x} = F(x)$ .

In this paper, we are primarily interested by the application of stochastic approximation algorithms to minimize a function  $f : \mathbb{R}^d \to \mathbb{R}$ . If the function f is differentiable and the

<sup>\*</sup> This work was funded by the project BayesScale - Pierre Laffitte -, the Data Science Initiative of Ecole Polytechnique and by Polish National Science Center grant no. 2015/17/D/ST1/01198.

<sup>&</sup>lt;sup>†</sup>Institute of Mathematics, Polish Academy of Sciences (smajewski@impan.pl).

<sup>&</sup>lt;sup>‡</sup>Institute of Applied Mathematics and Mechanics, University of Warsaw, Institute of Mathematics, Polish Academy of Sciences, (bmia@mimuw.edu.pl)

<sup>&</sup>lt;sup>§</sup>Ecole Polytechnique, Paris, National Research University Higher School of Economics (HSE), Moscow, (eric.moulines@polytechnique.edu).

gradient  $\nabla f$  is known, a classical method to minimize f consists in performing a gradient descent. If only a noise corrupted  $H_k$  version of the gradient  $\nabla f(x_{k-1})$  is available, a popular algorithm is the stochastic gradient descent (SGD) algorithm whose iterations are given by  $x_k = x_{k-1} - \gamma_k H_k$ . In this context  $F(x) = -\nabla f$  and the associated ODE is  $\dot{x} = -\nabla f(x)$ .

If the function f is not differentiable, it is no longer possible to use the SGD algorithm. However, if the function f is locally Lipschitz, the Clarke generalized gradient  $\bar{\partial} f(x)$  can still be defined (see Definition 2.6 and [16, Section 1.2]). The Clarke generalized gradient  $\bar{\partial} f(x)$ is a point-to-set map: for any  $x \in \mathbb{R}^d$ ,  $\bar{\partial} f(x)$  is a nonempty convex compact subset of  $\mathbb{R}^d$ . When f is continuously differentiable at x,  $\bar{\partial} f(x)$  reduces to the singleton  $\{\nabla f(x)\}$ . When fis convex, then  $\bar{\partial} f(x)$  coincides with the subdifferential of convex analysis. As above, if the Clarke gradient cannot be computed but a noise corrupted version of a selection of  $\bar{\partial} f(x_{k-1})$ is available, we may consider a generalization of the stochastic subgradient algorithm

$$x_k = x_{k-1} - \gamma_k \{v_{k-1} + \eta_k\}$$
, where  $v_{k-1} \in \partial f(X_{k-1})$ 

which we sometimes denote more concisely

$$x_k \in x_{k-1} - \gamma_k \{ \partial f(x_{k-1}) + \eta_k \}$$
.

The classical stochastic approximation algorithm update rule is replaced by a stochastic recursive inclusion:

$$x_k \in x_{k-1} + \gamma_k \{F(x_{k-1}) + \eta_k\}$$

where *F* is a point-to-set map and  $\eta_k$  is defined in (1). Such algorithms play an important role in game theory, as illustrated in [8, 9] where numerous examples of stochastic recursive inclusions are introduced.

[11] have shown that the "mean-limit" approach leading to the ODE method in the smooth case can be extended to the analysis of stochastic recursive inclusion. In this case, the limit ODE is replaced by a solution of the differential inclusion

$$\dot{x} \in F(x)$$

i.e. an absolutely continuous mapping  $x : \mathbb{R} \to \mathbb{R}^d$  such that  $\dot{x}(t) \in F(x(t))$  for almost every  $t \in \mathbb{R}$ . Such differential inclusions play a key role in the analysis of nonsmooth dynamical systems; see for example [16, Chapter 4] or [5, Section 2.1] and the references therein.<sup>1</sup>

In this paper, we will also consider proximal algorithms, which have become an important tool in nonsmooth optimization problems; the literature in this field is also huge, see for example [4, 7, 31, 33, 13]). Proximal algorithms with stochastic updates have been proposed and studied in recent years. One such algorithm is Proximal Stochastic Gradient Descent (proxSGD), that optimizes composite convex function P = f + g where f is a continuously differentiable function with Lipschitz-gradients and g is a "proximable" function (e.g. g is lower semi-continuous and convex, but this notion can be extended to nonconvex functions). The proxSGD algorithm alternates between stochastic gradient update for f and deterministic proximal step for g. This above optimization problem plays a fundamental role in many machine learning problems, ranging from convex optimization such as convex regression problem with sparsity inducing penalties like LASSO to highly nonconvex problem such as optimizing the weights of deep neural networks. Numerous papers have been devoted to the

<sup>&</sup>lt;sup>1</sup>Just before submitting this paper, [19] published an analysis of stochastic subgradient algorithms. We have developed our results completely independently; our work is based on results obtained earlier in [11] and even if some statements are similar, the proofs in this paper are all original.

case when f and g are both convex, f gradient Lipschitz and g lower semi-continuous; see for example [37, 32, 40, 18]. Atchade et al. [3] have extended these results in the Markovian noise case. In recent years, triggered by the surge deep learning, the nonconvex case has started to attract many research efforts, at least in the smooth case (f gradient Lipschitz and  $g \equiv 0$ ); see for example [23, 1] and the references therein. For the nonsmooth and nonconvex case, the results are still partial. Ghadimi et al. [24] considered the case where f is differentiable but possibly nonconvex and g is non-differentiable but convex. They have analyzed the deterministic proximal gradient algorithm (where the full gradient is computed at each iteration). They have also extended their results to the stochastic case; Reddi et al. [34] provides rates of convergence.

We consider in this paper nonconvex and nonsmooth minimization problems. We establish the convergence of stochastic inclusion equation generalizing (1) by allowing implicit steps and projections on a closed compact convex set at each iteration. Our results generalize [11]. We also discuss the stability of the limit differential inclusion by means of locally Lipschitz continuous and regular Lyapunov functions (see Definition 2.8). We in particular establish a characterization of the possible limit point of the stochastic approximation algorithm as the set of zeros of an upper-bound of the set-valued Lie derivative (see Definition 3.3) of the Lyapunov function. We then apply our results to the analysis of the proximal stochastic gradient descent for the composite minimization problem P = f + g, under assumptions on the noise sequence analogous to those commonly used for the SGD in the smooth nonconvex case. We also show that V = f + g can play the role of a Lyapunov function. We finally analyse a projected version of stochastic subgradient algorithm.

The paper is organized as follows. In Section 2, we introduce our main assumptions and notations and introduce the proximal stochastic gradient and projected subgradient algorithms. In Section 3, we state and prove our main convergence results under the assumption that the iterates are stable. In Section 4, we extend these convergence results to the case where the updates are projected on a compact convex set. In Section 5, we consider applications of our main results to ProxSGD and projected stochastic subgradient. Finally in Section 6 we present postponed proofs.

### 2. Assumptions and Notations. In this section we introduce definitions and notations.

DEFINITION 2.1 (Perturbed approximate discretization (PAD) and projected perturbed approximate discretization (PPAD) ). Let X be an open subset of  $\mathbb{R}^d$  and F be a set-valued function mapping each point  $x \in X$  to a set  $F(x) \subset X$ . We say that the sequence  $\{x_k, k \in \mathbb{N}\} \subset X$  is a Perturbed Approximate Discretization with noise  $\{\eta_k, k \in \mathbb{N}\}$  and step sizes  $\{\gamma_k, k \in \mathbb{N}\}$  if an only if there exists  $\{y_k, k \in \mathbb{N}\}$  such that

(3) 
$$x_k \in x_{k-1} + \gamma_k \{F(y_k) + \eta_k\} .$$

Let K a compact convex set. We say that the sequence  $\{x_k, k \in \mathbb{N}\}$  is a K-Projected Perturbed Approximate Discretization (K-PPAD) with noise  $\{\eta_k, k \in \mathbb{N}\}$  and step sizes  $\{\gamma_k, k \in \mathbb{N}\}$  if and only if there exists  $\{y_k, k \in \mathbb{N}\}$  such that

(4) 
$$x_k \in \Pi_K \left( x_{k-1} + \gamma_k \{ F(y_k) + \eta_k \} \right) .$$

By convention for a convex closed set *K* and a given set *A*, by  $\Pi_K(A)$  we denote the projection of *A* onto *K*, defined as  $\Pi_K(A) = {\Pi_K(a), a \in A}$ .

*Remark* 2.2. The condition  $x_k \in \Pi_K(x_{k-1} + \gamma_k(F(y_k) + \eta_k))$  is satisfied if and only if there exists  $v_k \in F(y_k)$  such that

(5) 
$$\langle x_k - z, x_k - x_{k-1} - \gamma_k (v_k + \eta_k) \rangle \le 0$$

for all  $z \in K$ .

*Example* 2.3 (Benaïm *et al.* discretization). [11, Definition III] deal with sequence satisfying a recursion of the form

$$x_k \in x_{k-1} + \gamma_k \{F(x_{k-1}) + \eta_k\}$$
.

Such a sequence clearly is a PAD with noise  $\{\eta_k, k \in \mathbb{N}\}$ , step sizes  $\{\gamma_k, k \in \mathbb{N}\}$ . In such case  $y_k = x_{k-1}$ .

We now show that the PAD and *K*-PPAD formalism cover the proximal stochastic gradient descent (ProxSGD) and the stochastic (sub)gradient algorithms for nonsmooth and nonconvex minimization problems. First we introduce some additional definitions and notations.

DEFINITION 2.4 (Generalized directional derivative, after [16, Chapter 2, Section 1]). Let  $f : \mathbb{R}^d \to \mathbb{R}$  be a locally Lipschitz function at  $x_0$ . The generalized directional derivative of f at  $x_0$  in the direction  $h \in \mathbb{R}^d$  is:

$$f^{0}(x_{0},h) = \lim_{\substack{\delta \downarrow 0^{+} \\ 0 < \lambda < \delta}} \sup_{\substack{\delta \downarrow 0^{+} \\ 0 < \lambda < \delta}} \frac{f(x_{0} + \tilde{h} + \lambda h) - f(x_{0} + \tilde{h})}{\lambda}$$

*Remark* 2.5. Contrary to the standard directional derivative, the generalized directional derivative  $f^0(x_0, h)$  is always well defined for any interior point  $x_0$  of the domain  $Dom(f) = \{x \in \mathbb{R}^n : |f(x)| < \infty\}$ .

It is shown in [16, Proposition 2.1.1] that the generalized directional derivative at  $x_0$ ,  $h \mapsto f(x_0,h)$  is positively homogeneous and subadditive and  $|f^0(x_0,h)| \le L ||h||$ , where *L* is a Lipschitz constant on some neighborhood of  $x_0$ . Denote by  $\langle \cdot, \cdot \rangle$  the scalar product on  $\mathbb{R}^d$ .

DEFINITION 2.6 (Clarke generalized gradient). Let  $f : \mathbb{R}^d \to \mathbb{R}$  be a function and  $x_0$  a point in the interior of Dom(f). If f is locally Lipschitz function at  $x_0$ , the Clarke generalized gradient of f at  $x_0$  is the set defined by:

$$\overline{\partial} f(x_0) = \left\{ \zeta \in \mathbb{R}^d : f^0(x,h) \ge \langle h, \zeta \rangle, \quad \text{for all } h \in \mathbb{R}^d \right\}$$

Similarly to subgradient, the Clarke generalized gradient is a set-valued generalization of the gradient. In particular, when function f is continuously differentiable at some point  $x_0$ , then we have  $\overline{\partial} f(x_0) = \{\nabla f(x_0)\}$ . Furthermore, if the function f is convex and locally Lipschitz and  $x_0$  belongs to the interior of its domain, then the Clarke generalized gradient of f coincides with the subgradient.

We say that the set-valued map  $F : X \to \mathbb{R}^d$  is convex-compact if for any point  $x \in X$  the set F(x) is convex and compact. We say that the set-valued function F is locally bounded if for any compact set  $K \subset X$ ,  $\bigcup_{x \in K} F(x)$  is bounded. We also define upper hemicontinuity.

DEFINITION 2.7 (Upper Hemicontinuity). Let X be an open subset of  $\mathbb{R}^d$ . A set-valued map  $F: X \to \mathscr{P}(\mathbb{R}^d)$  is said to be upper hemicontinuous at  $x \in X$ , if and only if F(x) is nonempty, and for every open neighborhood U of F(x), there exist an open neighborhood V of x, such that:

$$\left(\bigcup_{z\in V}F(z)\right)\subseteq U.$$

It is shown in [16, Propositions 2.1.2] that if *f* is Lipschitz on a neighborhood of  $x_0$  with Lipschitz constant  $||f||_{\text{Lip},x_0}$  then  $\overline{\partial}f(x_0)$  is a non-empty set, compact, convex, and for any  $u \in \overline{\partial}f(x_0)$ ,  $||u|| \leq ||f||_{\text{Lip},x_0}$ . By [16, Proposition 2.1.5],  $\overline{\partial}f$  is upper hemicontinuous at  $x_0$ .

For any compact set  $K \subset X$ ,  $\sup_{x \in K} ||f||_{\operatorname{Lip},x} < \infty$  showing that  $\overline{\partial} f$  is locally bounded. For proofs of those results and additional properties of Clarke generalized gradient, we refer the reader to [16].

DEFINITION 2.8 (Regular function). Let f be a function and  $x_0 \in \text{Dom}(f)$ . Assume that f is locally Lipschitz at  $x_0$ . We say that f is regular at  $x_0$ , if and only if for any  $h \in \mathbb{R}^d$ 

$$f^{0}(x_{0},h) = \lim_{\lambda \downarrow 0^{+}} \frac{f(x_{0} + \lambda h) - f(x_{0})}{\lambda} ,$$

where  $f^0$  is the generalized directional derivatives (see Definition 2.4).

In words, f is regular at  $x_0$  if the directional derivatives exist for all directions  $d \in \mathbb{R}^d$  and coincide with the generalized directional derivatives.

*Remark* 2.9. It may happen that the usual directional derivative exists, but does not coincide with the generalized directional derivative. The classical example is f(x) = -|x|. As shown in [16, Proposition 2.3.6] every convex locally Lipschitz function is regular. The same property obviously holds for any continuously differentiable function. Note finally that if the functions  $f_1, \ldots, f_p$  are regular at  $x_0$ , then for any nonnegative weights  $\alpha_1, \ldots, \alpha_p, \sum_{i=1}^p \alpha_i f_i$ is also regular at  $x_0$ , see [16, Proposition 2.3.6].

Now we are ready to discuss the proximal gradient descent and projected subgradient algorithms.

*Example* 2.10 (Perturbed proximal gradient descent algorithm). Consider the problem of minimizing a composite function P = f + g define on an open subset of  $X \subseteq \mathbb{R}^d$  where f is continuously differentiable and g is locally Lipschitz, bounded from below and regular (see Definition 2.8). The proximal gradient algorithm is defined by the following recursion:

$$x_k \in \operatorname{prox}_{\gamma_k,g}(x_{k-1} - \gamma_k \nabla f(x_{k-1})), \quad k \ge 1,$$

where  $\{\gamma_k, k \in \mathbb{N}^*\}$  is a sequence of stepsizes and  $\operatorname{prox}_{\gamma,g}$  stands for the proximal operator defined by

(6) 
$$\operatorname{prox}_{\gamma,g}(x) \in \operatorname*{arg\,min}_{y \in \mathsf{X}} \left\{ g(y) + (2\gamma)^{-1} \left\| y - x \right\|^2 \right\} \,.$$

In our settings the function g is lower bounded and under this condition the set appearing in the right-hand side of (6) is nonempty. Necessary and sufficient conditions for this algorithm to be well-defined can be found in [36, Excersise 1.24]. In the perturbed version of the proximal gradient algorithm, for any  $k \in \mathbb{N}^*$  we replace the gradient  $\nabla f(x_{k-1})$  by a noise corrupted version  $\nabla f(x_{k-1}) + \zeta_k$  which leads to the recursion

$$x_k \in \operatorname{prox}_{\gamma_k \varrho}(x_{k-1} - \gamma_k \{ \nabla f(x_{k-1}) + \zeta_k \})$$
.

The characterization of the minimum by the Clarke generalized gradient (see [16, Proposition 2.3.2]) yields

$$0 \in \gamma_k^{-1}(x_k - x_{k-1}) + \nabla f(x_{k-1}) + \zeta_k + \overline{\partial}g(x_k)$$

Setting, for any  $k \in \mathbb{N}^*$ ,  $\eta_k = -\zeta_k + \nabla f(x_k) - \nabla f(x_{k-1})$  we get that for any  $k \in \mathbb{N}^*$ 

$$x_k \in x_{k-1} + \gamma_k \left[ -\nabla f(x_k) - \overline{\partial} g(x_k) + \eta_k \right]$$

Therefore perturbed proximal gradient is a PAD in the sense of Definition 2.1 with  $F = -\nabla f - \overline{\partial}g$ ,  $y_k = x_k$ , noise sequence  $\{\eta_k, k \in \mathbb{N}^*\}$ , and step sizes  $\{\gamma_k, k \in \mathbb{N}^*\}$ .

*Example* 2.11 (Projected stochastic (sub)gradient). We consider projected subgradient algorithm framework introduced in the convex case by [31] to solve the constrained minimization problem  $\arg\min_{x \in K} f(x)$ . Let *K* be a bounded closed convex set and  $f: K \to \mathbb{R}$  be a locally Lipshitz function. Assume that for each iteration an oracle returns a perturbed element of the subgradient, i.e.  $H_k := v_k + \zeta_k$  with  $v_k \in \partial f(x_{k-1})$  and with  $\zeta_k$  a perturbation.

The projected stochastic subgradient algorithm generates iteratively the sequence  $\{x_k, k \in \mathbb{N}^*\}$  as follows

$$x_k = \prod_K (x_{k-1} - \gamma_k H_k)$$

where  $\{\gamma_k, k \in \mathbb{N}^*\}$  is a sequence of stepsizes and  $\Pi_K$  is the projection on the set *K*. The projected subgradient algorithm is a *K*-PPAD with field  $F = -\partial f$ ,  $y_k = x_{k-1}$  and noise  $\eta_k = -\zeta_k$ .

- We will analyse the convergence of PAD (see (3)) under the following assumptions:
- (A1) X is an open subset of  $\mathbb{R}^d$  and  $F : X \to \mathscr{P}(\mathbb{R}^d)$  is a set-valued map, that is upper hemicontinuous, cf. Definition 2.7, convex-compact valued and locally bounded.
- (A2) The sequence of step sizes  $\{\gamma_k, k \in \mathbb{N}^*\}$  satisfies  $\gamma_k > 0$ ,  $\sum_{k=0}^{\infty} \gamma_k = \infty$ , and  $\lim_{k \to \infty} \gamma_k = 0$ .
- (A3) The perturbation sequence  $\{\eta_k, k \in \mathbb{N}^*\}$  can be decomposed as  $\eta_k = e_k + r_k$ , where  $\{e_k, k \in \mathbb{N}^*\}$  and  $\{r_k, k \in \mathbb{N}^*\}$  are two sequences satisfying  $\lim_{k\to\infty} ||r_k|| = 0$ , and  $\sum_{k=1}^{\infty} \gamma_k e_k$  converges.
- (A4) The approximation sequence  $\{y_k, k \in \mathbb{N}^*\}$  belongs to X and satisfies

$$\lim_{k\to\infty}\|x_k-y_k\|=0.$$

Condition (A1) is a rather mild regularity condition. Upper hemicontinuity replaces the continuity of the vector field which plays a key role in the classical theory of stochastic approximation [27]. The requirement for F to be convex-compact valued and locally bounded might be less obvious, but this assumption is commonly used in nonsmooth analysis. This is not a serious limitation for the minimization problems we have primarily in mind.

The assumptions (A2, A3) are usual in stochastic approximation literature [2, 22]. It is worth noting, that condition (A3) allows perturbations sequences which have random and deterministic components and hence our results can be used for proving almost sure convergence for ProxSGD for which the proximal operator is computed numerically and is therefore inexact (although in our framework the deterministic noise should vanish asymptotically faster than step size). The assumptions (A4) allows to cover both explicit and implicit discretization of differential inclusions as illustrated in Example 2.10.

As in classical ODE method for stochastic approximation, establishing convergence results first requires to show that algorithm is stable in the sense that the sequence  $\{x_k, k \in \mathbb{N}^*\}$ remains in some compact set. This issue is non-trivial even in the noiseless case and might be challenging to establish in the stochastic case. One of possible solution to overcome this difficulty is to introduce a projection on convex compact set *K*. This case is considered in Section 4, however first in Section 3 we consider the standard version of stochastic approximation, where we establish the convergence of PAD assuming that the sequence  $\{x_k, k \in \mathbb{N}^*\}$ remains in some compact set.

**3.** Convergence of Perturbed Approximate Discretisation. In this section, we state our main convergence results for PAD. First in Theorem 3.2 we show that a translated and interpolated version of the PAD converges to a solution of a differential inclusion. Further in Theorem 3.5 we combine these results with Lyapunov stability conditions to obtain convergence of the iterates to the set of stationary points of the differential inclusion.

DEFINITION 3.1 (Solution of Differential Inclusion). Let  $X \subseteq \mathbb{R}^d$  be an open subset,  $F : X \to \mathscr{P}(\mathbb{R}^d)$  be a set-valued map, and  $I \subseteq \mathbb{R}$  be an interval. A function  $x : I \to X$  is a solution of the differential inclusion  $\dot{x} \in F(x)$  if it is absolutely continuous and for almost every  $t \in I, \dot{x}(t) \in F(x(t))$ .

Let us define the piecewise linear interpolation  $X_0 : \mathbb{R} \to \mathbb{R}^d$  of the sequence  $\{x_k, k \in \mathbb{N}\}$  with positive stepsize  $\{\gamma_k, k \in \mathbb{N}\}$ :

(7) 
$$X_0(t) = \begin{cases} x_0 & \text{if } t \le 0\\ x_k \frac{(t-t_{k-1})}{\gamma_k} + x_{k-1} \frac{(t_k-t)}{\gamma_k} & \text{if } t \in [t_{k-1}, t_k], k \ge 1 \end{cases}$$

where  $t_0 = 0$  and for  $k \ge 1$ ,

(8) 
$$t_k = t_{k-1} + \gamma_k = \sum_{i=1}^k \gamma_i .$$

Let  $\{s_k, k \in \mathbb{N}^*\}$  be an increasing sequence of positive real numbers, such that  $\lim_{k\to\infty} s_k = \infty$ . Let us define shifted linear interpolation of  $\{x_k, k \in \mathbb{N}\}$  by

(9) 
$$X_k(t) = X_0(t+s_k), t \ge 0$$

Consider the PAD sequence  $\{x_k, k \in \mathbb{N}\}$  defined in (3): for  $k \in \mathbb{N}^*$ ,

(10) 
$$x_k = x_{k-1} + \gamma_k (v_k + e_k + r_k) ,$$

where  $\{e_k, k \in \mathbb{N}\}$  and  $\{r_k, k \in \mathbb{N}\}$  are defined in A3 and  $v_k \in F(y_k)$ . Let us define piecewise constant functions  $\hat{v}, \hat{r}, \hat{e}$  on  $[0, \infty)$  as follows

$$\hat{v}(t) = \sum_{k=1}^{\infty} v_k \mathbb{1}_{[t_{k-1}, t_k)}(t) , \hat{r}(t) = \sum_{k=1}^{\infty} r_k \mathbb{1}_{[t_{k-1}, t_k)}(t) , \hat{e}(t) = \sum_{k=1}^{\infty} e_k \mathbb{1}_{[t_{k-1}, t_k)}(t) ,$$

where  $\{t_k, k \in \mathbb{N}^*\}$  is defined in (8). Denote

(11) 
$$\hat{V}_0(t) = \int_0^t \hat{v}(s) ds, \quad \hat{R}_0(t) = \int_0^t \hat{r}(s) ds, \quad \hat{E}_0(t) = \int_0^t \hat{e}(s) ds.$$

Analogously to (9), for any  $k \in \mathbb{N}^*$  we denote by  $\hat{V}_k, \hat{E}_k, \hat{R}_k$  the shifts of  $\hat{V}_0, \hat{E}_0, \hat{R}_0$  respectively. With this notation for any  $k \in \mathbb{N}^*$  we can decompose  $X_k(t)$  as follows

(12) 
$$X_k(t) = X_0(t+s_k) = x_0 + \hat{V}_k(t) + \hat{R}_k(t) + \hat{E}_k(t) .$$

Without loss of generality we can assume that  $\sum_{k=1}^{\infty} \gamma_k e_k = 0$  (if this is not true, we can just modify  $r_1$  and  $e_1$ ).

THEOREM 3.2. Let  $\{x_k, k \in \mathbb{N}^*\}$  be the PAD (10). Assume that conditions (A1–4) hold, and there exists a compact set  $K \subset X$  such that  $x_k \in K$  for any  $k \ge 0$ . Then

(i) the family of functions {X<sub>k</sub>, k ∈ N\*}, defined by (9) is precompact in the topology of compact convergence: for any increasing sequence {n<sub>k</sub>, k ∈ N\*} of positive integers, there exist a subsequence {n<sub>k</sub>, k ∈ N\*} and an absolutely continuous function X<sub>∞</sub>: [0,+∞) → K such that for any T > 0, lim<sub>k→∞</sub> sup<sub>t∈[0,T]</sub> ||X<sub>n<sub>k</sub></sub>(t) − X<sub>∞</sub>(t)|| = 0 and X<sub>∞</sub> is a solution of differential inclusion

$$\dot{x} \in F(x)$$
.

(ii) In addition, for any  $t \ge 0$ ,  $X_{\infty}(t)$  is a limiting point of the sequence  $\{x_k, k \in \mathbb{N}^*\}$ .

*Proof.* We divide the proof into three steps.

STEP 1. The family of functions  $\{X_k, k \in \mathbb{N}^*\}$ , defined by (9) is precompact in the topology of compact convergence.

*Proof.* First we prove equicontinuity, cf. Definition 6.4, of the sequence of functions  $\{X_k, k \in \mathbb{N}\}$ . The proof follows essentially by the same arguments as in [27, Theorem 2.3.1], but we included it for completeness. Obviously equicontinuity (see Definition 6.4) of all terms on the RHS of (12) implies pointwise equicontinuity of the family of functions  $\{X_k, k \in \mathbb{N}^*\}$ . Assumption (A3) implies boundedness of the sequence  $\{r_k, k \in \mathbb{N}^*\}$ . Therefore the sequence of functions  $\{\hat{R}_k, k \in \mathbb{N}^*\}$  is equicontinuous since for each  $k \in \mathbb{N}^*$ ,  $\hat{R}_k$  is Lipschitz continuous with constant  $\|\hat{R}_k\|_{\text{Lip}} = \sup_{\ell \in \mathbb{N}^*} \|r_\ell\|$ . Consider now  $\{\hat{V}_k, k \in \mathbb{N}^*\}$ . Since the sequence  $\{x_k, k \in \mathbb{N}\}$  belongs to the compact set

Consider now  $\{V_k, k \in \mathbb{N}^*\}$ . Since the sequence  $\{x_k, k \in \mathbb{N}\}$  belongs to the compact set K, under (A4) the sequence  $\{y_k, k \in \mathbb{N}\}$  belongs to a compact neighborhood of K. By (A1) F is locally bounded so the sequence  $\{v_k, k \in \mathbb{N}\}$  is also bounded and functions  $\{V_k, k \in \mathbb{N}^*\}$  are Lipschitz continuous with  $\|\hat{V}_k\|_{\text{Lip}} = \sup_{\ell \in \mathbb{N}^*} \|v_\ell\|$ , which implies equicontinuity.

Consider finally  $\{\hat{E}_k, k \in \mathbb{N}^*\}$ . For any arbitrarily chosen  $t \in \mathbb{R}_+, k, n \in \mathbb{N}^*$ , define  $b_{k,n}^t$ 

$$b_{k,n}^{t} = \sup_{\{s:|s-t|<1/n\}} \left\| \hat{E}_{k}(s) - \hat{E}_{k}(t) \right\|$$

and consider the sequence  $\{a_n^t, n \in \mathbb{N}\}$  given for  $n \in \mathbb{N}^*$  by

$$a_n^t = \sup_{k\geq 1} b_{k,n}^t$$
.

By construction  $\{a_n^t, n \in \mathbb{N}^*\}$  is nonincreasing and nonnegative. Hence  $\{a_n^t, n \in \mathbb{N}^*\}$  converges to some limit. Moreover, by assumption (A3) the series  $\sum_{k=1}^{\infty} \gamma_k e_k$  converges, so for any  $\varepsilon > 0$ , there exists  $N \in \mathbb{N}^*$  such that

(13) 
$$\sup_{m\geq l\geq N} \left\| \sum_{i=l}^{m} \gamma_i e_i \right\| \leq \varepsilon$$

First we observe that, since  $\lim_{k\to\infty} s_k = \infty$ , the set  $\mathscr{E}_N^t = \{k : s_k < t_N - t + 1\}$  is finite. Therefore, since for each  $\ell \in \mathbb{N}^*$  the functions  $\hat{E}_\ell$  is continuous, there exists  $N' \ge N$  such that for any  $k \in \mathscr{E}_N^t$  and  $n \ge N'$  we have  $b_{k,n}^t \le \varepsilon$ . Now assume that  $k \notin \mathscr{E}_N^t$ . Then for all  $n \ge 1$  and ssuch that  $|t-s| \le 1/n$ , we get  $s + s_k \ge t_N$  and by (13) for all  $n \ge 1$  and  $k \notin \mathscr{E}_N^t$  we get

$$b_{k,n}^{t} = \sup_{\{s:|s-t|<1/n\}} \left\| \hat{E}_{k}(s) - \hat{E}_{k}(t) \right\| \leq 3 \sup_{m \geq l \geq N} \left\| \sum_{i=l}^{m} \gamma_{i} e_{i} \right\| \leq 3\varepsilon$$

Therefore for  $n \ge N'$  we have  $a_n^t \le 3\varepsilon$ . Since  $\varepsilon$  was arbitrary positive number, we get that  $\lim_{n\to\infty} a_n^t = 0$ . T Hence, for all  $\varepsilon > 0$ , there exists N'' > 0 such that

for all 
$$s \in \mathbb{R}$$
,  $|t-s| < 1/N'' \Rightarrow \sup_{k \ge 1} \left\| \hat{E}_k(t) - \hat{E}_k(s) \right\| \le \varepsilon$ 

That is the family  $\{\hat{E}_k, k \in \mathbb{N}^*\}$  is pointwise equicontinuous at an arbitrary point *t*. Together with equicontinuity of  $\{\hat{V}_k, k \in \mathbb{N}^*\}$  and  $\{\hat{R}_k, k \in \mathbb{N}^*\}$  it give us pointwise equicontinuity of  $\{X_k, k \in \mathbb{N}^*\}$ . Since by assumption  $\{x_k, k \in \mathbb{N}^*\}$  remains in the compact set *K* so functions  $\{X_k, k \in \mathbb{N}^*\}$  are uniformly bounded and we can apply Arzela-Ascoli theorem (Theorem 6.5), showing that, from every subsequence of  $\{X_k, k \in \mathbb{N}^*\}$ , we can choose a further subsequence that converges uniformly on compact intervals, to some continuous limit  $X_{\infty}$ . STEP 2. Any limit  $X_{\infty}$  of converging subsequence is absolutely continuous and for almost every  $t \ge 0$  there exists subsequence  $\{n_k, k \in \mathbb{N}\}$  such that

$$X_{\infty}(t) = \lim_{k \to \infty} x_{n_k} = \lim_{k \to \infty} y_{n_k}$$

*Proof.* Let  $\{X_{n_k}, k \in \mathbb{N}\}\$  be a subsequence that converges compactly to  $X_{\infty}$ . We start by proving that  $X_{\infty}$  is absolutely continuous on compact intervals. It is clear that  $\{\hat{E}_{n_k}, k \in \mathbb{N}\}\$  converges compactly to a function, that is equal to  $\sum_{k=1}^{\infty} \gamma_k e_k = 0$  everywhere. That means, that the sequence

(14) 
$$M_k = x_0 + \hat{V}_{n_k} + \hat{R}_{n_k}$$

converges compactly to the same limit as  $X_{n_k}$ . Recall that  $\hat{V}_k$  and  $\hat{R}_k$  are Lipschitz continuous with constant independent on k. Therefore all functions  $M_k$  are Lipschitz continuous with common constant and  $X_{\infty}$ , which is equal to the limit of  $\{M_k, k \in \mathbb{N}\}$ , is also Lipschitz continuous and hence absolutely continuous on compact intervals.

For any  $t \ge 0$  and  $k \in \mathbb{N}$  let us define m(k, t) by

(15) 
$$m(k,t) = \min\{n \in \mathbb{N} : t_n > t + s_k\},$$

where  $t_n$  is defined in (8). By assumption (A2) m(k,t) is well defined and converges to  $\infty$  as  $k \to \infty$ .

By construction, for each  $k \in \mathbb{N}$  we get that

(16) 
$$X_{n_k}(t_{m(n_k,t)} - s_{n_k}) = x_{m(n_k,t)} .$$

Moreover, since  $\lim_{k\to\infty} \gamma_k = 0$  and  $\lim_{k\to\infty} s_k = \infty$ , by (15) we have

(17) 
$$\lim_{k \to \infty} \{ t_{m(n_k,t)} - s_{n_k} - t \} = 0.$$

By the triangle inequality we get that

(18) 
$$\|y_{m(n_k,t)} - X_{\infty}(t)\|$$
  
 $\leq \|y_{m(n_k,t)} - x_{m(n_k,t)}\| + \|x_{m(n_k,t)} - X_{\infty}(t_{m(n_k,t)} - s_{n_k})\| + \|X_{\infty}(t_{m(n_k,t)} - s_{n_k}) - X_{\infty}(t)\|$ 

By assumption  $s_k$  converges to  $\infty$ , so also  $m(n_k, t)$  goes to  $\infty$  as  $k \to \infty$ . Therefore by assumption (A4) the first part of the RHS of (18) converges to 0. By (16), the second term in the RHS of (18) is equal to

$$\|X_{n_k}(t_{m(n_k,t)}-s_{n_k})-X_{\infty}(t_{m(n_k,t)}-s_{n_k})\|$$

which goes to zero by uniform convergence of  $X_{n_k}$  to  $X_{\infty}$ . Finally, continuity of  $X_{\infty}$  implies that the last term of (18) also converges to 0. All together we have therefore established that

(19) 
$$\lim_{t \to \infty} y_{m(n_k,t)} = X_{\infty}(t) . \qquad \Box$$

STEP 3. The limit  $X_{\infty}$  is a solution of differential inclusion  $\dot{x}(t) \in F(x(t))$ .

*Proof.* We denote by *G* a weak derivative of  $X_{\infty}$ . We will prove that  $G(t) \in F(X_{\infty}(t))$  for almost every  $t \in \mathbb{R}_+$ . By the definition (11), for each  $k \in \mathbb{N}^*$  and almost every  $t \in \mathbb{R}_+$ , the weak derivatives of  $M_k$  (see (14)) at *t* is equal to  $\dot{M}_k(t) = \hat{v}(t+s_{n_k}) + \hat{r}(t+s_{n_k})$ . Because  $\sup_k(||v_k|| + ||r_k||) < \infty$ , the functions  $\{\dot{M}_k, k \in \mathbb{N}\}$  are uniformly integrable on finite intervals. Thus,

from Lemma 6.6, for any  $0 < a < b < \infty$  the sequence  $\{\mathbb{1}_{[a,b]}\dot{M}_k, k \in \mathbb{N}\}$  converges weakly to  $\mathbb{1}_{[a,b]}G$  in  $L_1([a,b])$ . From Lemma 6.8 there exists  $\{\dot{M}_k^w, k \in \mathbb{N}\}$  a convex combination subsequence (see Definition 6.7) of  $\{\dot{M}_k, k \in \mathbb{N}\}$  that converges to *G* almost everywhere on [a,b], i.e. for almost every  $t \in [a,b]$ ,  $\lim_{k\to\infty} \dot{M}_k^w(t) = G(t)$ . By construction for any  $t \in \mathbb{R}_+$ , we get  $\dot{M}_k^w(t) = \hat{v}^w(t + s_{n_k}) + \hat{r}^w(t + s_{n_k})$  and

$$\hat{v}^{w}(t+s_{n_{k}}) = \sum_{j=1}^{\infty} w_{n,j} v_{m(n_{j},t)}$$
 and  $\hat{r}^{w}(t+s_{n_{k}}) = \sum_{j=1}^{\infty} w_{n,j} r_{m(n_{j},t)}$ .

By assumption (A3) for any  $t \in \mathbb{R}_+$ ,  $\lim_{k\to\infty} ||r_{m(n_k,t)}|| = 0$  and hence  $\lim_{k\to\infty} \hat{r}^w(t+s_{n_k}) = 0$  It follows, that the for almost every  $t \in [a,b] \lim_{k\to\infty} \hat{v}^w(t+s_{n_k}) = G(t)$ . But we have for all  $t \in \mathbb{R}_+$ ,  $v_{m(n_k,t)} \in F(y_{m(n_k,t)})$  and by (19) we know that  $\lim_{k\to\infty} y_{m(n_k,t)} = X_{\infty}(t)$ . Since F is upper hemicontinous and closed convex, we apply Lemma 6.9 to conclude that  $G(t) \in F(X_{\infty}(t))$  for almost every  $t \in [a,b]$ .

We have proven that  $G(t) \in F(X_{\infty}(t))$  for almost all  $t \in [a,b]$ , where  $[a,b] \subset \mathbb{R}_+$  is an arbitrary compact interval. We can cover the real line  $\mathbb{R}_+$  by a countable family of compact intervals of form  $[0,\ell]$  for  $\ell \in \mathbb{N}^*$ . Let  $\{n_k, k \in \mathbb{N}^*\}$  be a sequence. For each  $\ell \in \mathbb{N}^*$  can extract subsequence  $\{n_k^{\ell}, k \in \mathbb{N}^*\} \subseteq \{n_k^{\ell-1}, k \in \mathbb{N}^*\}$  such that  $\{X_{n_k^{\ell}}, k \in \mathbb{N}^*\}$  converges uniformly on  $[0,\ell]$ . Setting  $\tilde{n}_k = n_k^k$  we get that there exists function  $X_{\infty} : \mathbb{R} \mapsto \mathbb{R}^d$  such that that for any T > 0,  $\lim_{k \to \infty} \sup_{t \in [0,T]} ||X_{\tilde{n}_k}(t) - X_{\infty}(t)|| = 0$ . and  $\dot{X}_{\infty}(t) \in F(X_{\infty}(t))$  for almost all  $t \in \mathbb{R}$ . This is equivalent to saying, that  $X_{\infty}$  is a solution of the differential inclusion  $\dot{x} \in F(x)$ .

Combining Theorem 3.2 with stability properties of underlying differential inclusion  $\dot{x} \in F(x)$  we establish convergence of PADs. To state the result we need to define a set valued Lie derivative, introduced in [6].

DEFINITION 3.3. Let  $X \subseteq \mathbb{R}^d$  be an open subset,  $F : X \to \mathbb{R}^n$  be a set-valued map, and  $V : \mathbb{R}^d \to \mathbb{R}_+$  be a locally Lipschitz function. The set-valued Lie derivative of V with respect to F at x is defined by

$$\mathscr{L}_F V(x) = \left\{ a \in \mathbb{R} : \exists v \in F(x) \text{ such that } \langle v, w \rangle = a , \forall w \in \overline{\partial} V(x) \right\},\$$

where  $\bar{\partial}V$  is the Clarke generalized gradient of V, cf. Definition 2.6.

The Lie derivative plays important role in analysis of stability of solution of differential inclusions. We in particular will used an important property stated in [6, Lemma 1].

LEMMA 3.4. Let  $F : X \to \mathscr{P}(\mathbb{R}^d)$  be a set-valued map on an open domain X,  $I \in \mathbb{R}$  be an interval, and assume that there exists  $\phi : I \to X$  a solution of the differential inclusion  $\dot{x} \in F(x)$ . Let V be a locally Lipschitz regular function defined on X. Then  $\frac{d}{dt}V(\phi(t))$  exists for almost all  $t \in I$ , and for almost all  $t \in I$  we have:

$$\frac{d}{dt}V(\phi(t)) \in \mathscr{L}_F V(\phi(t))$$

where  $\mathscr{L}_F V$  is the set-valued Lie derivative of V with respect to F, cf. Definition 3.3.

THEOREM 3.5. Let  $X \subseteq \mathbb{R}^d$  be an open subset and  $F : X \mapsto \mathscr{P}(\mathbb{R}^d)$  be a set valued map,  $\{x_k, k \in \mathbb{N}^*\}$  be a PAD of F with step sizes  $\{\gamma_k, k \in \mathbb{N}^*\}$  and perturbations  $\{\eta_k, k \in \mathbb{N}^*\}$ . Assume that conditions (A1-4) hold, and there exists a compact set  $K \subset X$  such that  $x_k \in K$ for any  $k \ge 1$ . Let  $V : X \to \mathbb{R}$  be a locally Lipschitz, regular function. Suppose that there exists an upper semicontinuous function  $U : X \to \mathbb{R}$ , such that for all  $x \in K$ 

$$\sup \mathscr{L}_F V(x) \le U(x) \le 0,$$

and set  $\mathscr{S} := \{x \in \mathsf{X} : U(x) = 0\}.$ 

- (i) The image by V of the set of limiting points of  $\{x_k, k \in \mathbb{N}^*\}$  is a compact interval in  $V(\mathscr{S} \cap K)$ .
- (ii) If  $V(\mathscr{S} \cap K)$  has empty interior, then  $\{x_k, k \in \mathbb{N}^*\}$  converges to  $K \cap \mathscr{S}$ .

Proof. The proof is divided into four steps.

STEP 1. The set  $K \cap \mathscr{S}$  is nonempty.

*Proof.* By Theorem 3.2, there exists a solution  $X : \mathbb{R}_+ \to \mathbb{R}^d$  of the differential inclusion  $\dot{x} \in F(x)$  satisfying  $X(t) \in K$  for all  $t \in \mathbb{R}_+$ . By Lemma 3.4 for almost all  $t \in \mathbb{R}_+ \frac{d}{dt}V(X(t))$  is well-defined and we have:

$$\frac{d}{dt}V(X(t)) \in \mathscr{L}_F V(X(t))$$

If the  $K \cap \mathscr{S} = \emptyset$  by upper semicontinuity of U(x) and compactness of K we would have  $\sup_{x \in K} U(x) = -\delta$  for some  $\delta > 0$ . Therefore function  $V \circ X$  must decrease at a rate at least  $\delta$ , and thus  $\lim_{t \to \infty} V(X(t)) = -\infty$ . But this is a contradiction with the assumption that V is bounded from below.

Since V is continuous and K is compact,

(20) 
$$L = \liminf_{k \in \mathbb{N}} V(x_k) > -\infty,$$

and there exists  $x_* \in K$  and a subsequence  $\{x_{n_k}, k \in \mathbb{N}^*\}$  such that  $\lim_{k\to\infty} x_{n_k} = x_*$  and  $V(x_*) = L$ .

STEP 2. If  $\{x_{n_k}, k \in \mathbb{N}^*\}$  is a subsequence such that  $\lim_{k\to\infty} x_{n_k} = x_*$  and  $V(x_*) = L$  then  $x^* \in \mathscr{S} \cap K$ .

*Proof.* The proof is by contradiction. Assume that  $x_* \notin \mathscr{S} \cap K$ . Therefore, we can find disjoint open neighborhoods  $X \supset A \supset \mathscr{S} \cap K$  and by  $X \supset B \ni x_*$ . Also, there exists r > 0, such that  $\overline{B}(x_*, r) := \{y \in \mathbb{R}^d : ||y - x^*|| \le r\} \subseteq B$ . Define

(21) 
$$v_{\max} = \sup_{x \in K} \sup_{z \in F(x)} \|z\|,$$

which is finite by local boundedness of *F*. We denote by  $\Delta t = r/v_{\text{max}}$ . From Theorem 3.2 it follows, that on the interval  $[0,\Delta t]$  there exists a subsequence  $\{\tilde{n}_k, k \in \mathbb{N}^*\} \subseteq \{n_k, k \in \mathbb{N}^*\}$  such that  $\lim_{k\to\infty} \sup_{t\in[0,\Delta t]} ||X_{\tilde{n}_k} - X_{\infty}|| = 0$ , where  $X_{\infty}$  is a solution of  $\dot{x} \in F(x)$ . Let  $X_0$  be defined by (7) and set, for all  $k \in \mathbb{N}$ ,  $s_k = t_{\tilde{n}_k}$ , where  $\{t_k, k \in \mathbb{N}\}$  is defined in (8). By (A2)  $\lim_{k\to\infty} s_k = \infty$ . Consider  $\{X_k, k \in \mathbb{N}\}$  defined by (9). By definition of  $\{s_k, k \in \mathbb{N}\}$ , for all k, we have  $X_k(0) = x_{\tilde{n}_k}$  and since  $\lim_{k\to\infty} x_{\tilde{n}_k} = x_*$  we have  $X_{\infty}(0) = x_*$ . For almost every  $t \in [0,\Delta t]$  we have  $\dot{X}_{\infty}(t) \in F(X_{\infty}(t))$  and hence by (21) we get  $||\dot{X}_{\infty}(t)|| \leq v_{\text{max}}$ . Thus, for all  $t \in [0,\Delta t]$ ,

$$\|X_{\infty}(t) - X_{\infty}(0)\| = \left\|\int_0^t \dot{X}_{\infty}(s) \mathrm{d}s\right\| \le \int_0^t v_{\max} \mathrm{d}s \le v_{\max} \Delta t = r\,,$$

and hence  $X_{\infty}(t) \in B(x_*, r)$ . By upper semicontinuity of U and compactness of  $\overline{B}(x_*, r)$  we get that there exists  $\delta > 0$  such that  $\sup_{x \in \overline{B}(x_*, r)} U(x) = -\delta$ , and, using Lemma 3.4, we conclude that for almost every  $t \in [0, \Delta t]$ ,  $\frac{d}{dt}V(X_{\infty}(t)) \leq -\delta$ . This means, that

(22) 
$$V(X_{\infty}(\Delta t)) \leq V(X_{\infty}(0)) - \delta \Delta t = L - \delta \Delta t ,$$

where *L* is defined in (20). But, by Theorem 3.2 for almost every  $t \in [0, \Delta t]$ ,  $X_{\infty}(t)$  is a accumulation point of sequence  $\{x_k, k \in \mathbb{N}^*\}$  so (22) contradicts with  $\liminf_{k\to\infty} V(x_k) = L.\Box$ 

STEP 3. If  $\hat{x}$  is an accumulation point of the sequence  $\{x_k, k \in \mathbb{N}^*\}$ , then  $V(\hat{x}) \in V(\mathscr{S} \cap K)$ .

*Proof.* The proof is by contradiction. Assume that there exists a subsequence  $\{x_{n_k}, k \in \mathbb{N}^*\}$  such that  $\lim_{k\to\infty} x_{n_k} = \hat{x}$  and  $V(\hat{x}) \notin V(\mathscr{S} \cap K)$ . Then we have  $\hat{x} \notin \mathscr{S} \cap K$  as well. By Step 2, we know that there exists another subsequence  $\{m_l, l \in \mathbb{N}^*\}$  such that  $\lim_{l\to\infty} x_{m_l} = x_*$ ,  $V(x_*) = \liminf_k V(x_k) = L$  and  $x_* \in \mathscr{S} \cap K$ , where *L* is defined in (20). Note that  $V(x_*) < V(\hat{x})$ . So, there exist  $a_1 < a_2 < b_1 < b_2$  such that intervals  $(a_1, a_2) \ni V(x_*)$ ,  $(b_1, b_2) \ni V(\hat{x})$  are disjoint and  $(a_2, b_2) \cap V(\mathscr{S} \cap K) = \emptyset$ . We denote by  $A = V^{-1}((a_1, a_2))$  and by  $B = V^{-1}((b_1, b_2))$ . Observe that sets *A* and *B* are also disjoint.

Since  $\hat{x} \in B$  and  $x_* \in A$ , the function  $X_0$ , defined in (7), must go from A to B infinitely often. More precisely, for  $j \in \mathbb{N}^*$  we can define three increasing sequences  $\{l_j, j \in \mathbb{N}\}$ ,  $\{\tilde{l}_j, j \in \mathbb{N}\}$ , and  $\{r_j, j \in \mathbb{N}\}$  by recurrence as follows:  $r_0 = l_0 = \tilde{l}_0 = 0$  and for  $j \ge 1$ ,  $\tilde{l}_j = \min\{t \ge r_{j-1} : X_0(t) \in A\}$ ,  $r_j = \min\{t \ge \tilde{l}_j : X_0(t) \in B\}$  and  $l_j = \max\{t \le r_j : X_0(t) \in A\}$ . Because  $X_0$  is continuous, the sequences  $\{l_j, j \in \mathbb{N}\}$ ,  $\{r_j, j \in \mathbb{N}\}$  are well defined. Since, sets A and B are disjoint and both contain accumulation points of  $\{x_k, k \in \mathbb{N}^*\}$ , by construction  $\lim_{j\to\infty} l_j = \infty$ , in addition by continuity of V we get that  $V(X_0(l_j)) = a_2$ ,  $V(X_0(r_j)) = b_1$ . Furthermore, for all  $t \in (l_j, r_j)$  we have  $V(X_0(t)) \in (a_1, b_2)$  and  $X_0(t) \in K \setminus (A \cup B)$ .

Consider the sequence  $\{r_j - l_j : j \in \mathbb{N}^*\}$  of positive numbers. Set  $S = \limsup_{j \to \infty} \{r_j - l_j\} \in [0,\infty]$  and let  $\{m_j, j \in \mathbb{N}\}$  be a sequence such that  $\lim_{j\to\infty} \{r_{m_j} - l_{m_j}\} = S$ .

Let  $s_k = l_{m_k}$ , T > 0 and consider  $\{X_k, k \in \mathbb{N}^*\}$  defined in (9). By Theorem 3.2 there exists  $\{\tilde{n}_k, k \in \mathbb{N}^*\}$  such that  $\lim_{k\to\infty} \sup_{t\in[0,T]} ||X_{\tilde{n}_k} - X_{\infty}|| = 0$ , where  $X_{\infty}$  is a solution of  $\dot{x} \in F(x)$ . First assume that S = 0. By construction for all  $k \in \mathbb{N}^*$ ,  $V(X_k(0)) = a_2$  and  $V(X_k(r_{m_k} - l_{m_k})) = b_1$ . But this contradicts the continuity of  $V \circ X_{\infty}$ . Therefore we must have S > 0.

Consider now the case: S > 0. By construction, we may find  $\tilde{S} \in (0, S)$  such that for large enough k for all  $t \in (0, \tilde{S}]$  we have  $X_k(t) \in K \setminus (A \cup B)$  and  $V(X_k(t)) > a_2$ . Therefore, for all  $t \in (0, \tilde{S}]$  the limit  $X_{\infty}$  also satisfies  $X_{\infty}(t) \in K \setminus (A \cup B)$ ,  $V(X_{\infty}(t)) > a_2$ ,  $V(X_{\infty}(0)) = a_2$ . On the other hand, by upper semicontinuity of U and compactness of  $K \setminus (A \cup B)$  we get that there exists  $\delta > 0$  such that  $\sup_{x \in K \setminus A} U(x) = -\delta$  and by Lemma 3.4  $V \circ X_{\infty}$  is strictly decreasing on  $[0, \tilde{S}]$ , which contradicts with  $V(X_{\infty}(t)) > a_2$ ,  $V(X_{\infty}(0)) = a_2$ . Hence  $V(\hat{x}) \in V(\mathscr{S} \cap K)$ .

STEP 4. If  $\hat{x}$  is an accumulation point of  $\{x_k, k \in \mathbb{N}\}$ , then  $V(\hat{x}) = L$ , where L is defined in (20).

*Proof.* Consider  $X_0$ , defined in (7). Suppose there were two different points x', x'', that are accumulation points of  $\{x_k, k \in \mathbb{N}^*\}$ , with V(x') < V(x''). Then the sequence  $\{V(x_k), k \in \mathbb{N}^*\}$  must oscillate between between V(x') and V(x''). Let  $v \in [V(x'), V(x'')]$ . We define the sequence  $\{s_k, k \in \mathbb{N}\}$  by  $s_0 = 0$  and for  $k \ge 1$ ,  $s_k = \min\{t > s_{k-1} : V(X_0(t)) = v\}$ , which is well defined by continuity of  $V \circ X_0$ . Let  $X_k$  be defined by (9). For any T > 0 by Theorem 3.2 there exists  $\{\tilde{n}_k, k \in \mathbb{N}^*\}$  such that  $\lim_{k\to\infty} \sup_{t\in[0,T]} ||X_{\tilde{n}_k}(t) - X_{\infty}(t)|| = 0$ , where by construction  $V(X_{\infty}(0)) = v$  and  $x''' = X_{\infty}(0)$  is accumulation point of  $\{x_k, k \in \mathbb{N}^*\}$ .

Using Step 3,  $V(x'), V(x''), V(x''') \in V(\mathscr{S} \cap K)$ . Since  $v \in [V(x'), V(x'')]$  is arbitrary, therefore, the whole interval [V(x'), V(x'')] must be contained in  $V(\mathscr{S} \cap K)$ . But this contradicts our assumption, that  $V(\mathscr{S} \cap K)$  has empty interior. Therefore, for any point  $\hat{x}$  that is an accumulation point of the sequence  $\{x_k, k \in \mathbb{N}\}$  we must have  $V(\hat{x}) = L$ .

We can now conclude the proof of Theorem 3.5. Combining Step 2 with Step 4 we get that any accumulation point of  $\{x_k, k \in \mathbb{N}^*\}$  belongs to  $\mathscr{S} \cap K$ , and hence  $\lim_{k\to\infty} d(x_k, \mathscr{S} \cap K) = 0.\square$ 

4. Convergence of Projected Perturbed Approximate Discretisation. Let  $K \in \mathbb{R}^d$  be a compact convex set . For any  $x \in K$  the tangent and the normal cone are set valued maps

defined as

(23) 
$$\mathbf{T}_K(x) = \operatorname{cl}\{d \in \mathbb{R}^d : \exists \varepsilon > 0 \text{ such that } x + \varepsilon d \in K\},\$$

(24) 
$$\mathbf{N}_{K}(x) = \{g \in \mathbb{R}^{d} : \langle g, z - x \rangle \leq 0, \text{ for all } z \in K\},\$$

where cl(*A*) denotes closure of set *A*. Let *F* be a convex compact set valued map defined on  $X \subseteq \mathbb{R}^d$ . Consider the *K*-PPAD sequence  $\{x_k, k \in \mathbb{N}\}$  defined in (4): for  $k \in \mathbb{N}^*$ ,

(25) 
$$x_k = \Pi_K \left( x_{k-1} + \gamma_k \{ v_k + e_k + r_k \} \right) \,,$$

where  $\{e_k, k \in \mathbb{N}\}$  and  $\{r_k, k \in \mathbb{N}\}$  are defined in A3 and  $v_k \in F(y_k)$ .

- THEOREM 4.1. Let  $\{x_k, k \in \mathbb{N}^*\}$  be the K-PPAD given in (25). Assume that conditions (A1–4) hold and  $\sup_k ||e_k|| < \infty$ . Then
- (i) The sequence of functions  $\{X_k, k \in \mathbb{N}^*\}$ , defined in (9), is precompact in the topology of compact convergence.
- (ii) For every convergent subsequence of  $\{X_k, k \in \mathbb{N}^*\}$  the limit  $X_{\infty}$  is a solution of a projected differential inclusion  $\dot{x} \in \Pi_{T_K(x)}(F(x))$ , where  $\Pi_{T_K(x)}$  is the projection onto the closed convex cone  $T_K(x)$ .

*Proof.* As in the proof of Theorem 3.2, we assume  $\sum_{k=1}^{\infty} \gamma_k e_k = 0$ . We denote by

(26) 
$$p_k = \frac{x_k - x_{k-1}}{\gamma_k} - v_k - e_k - r_k$$

and we define the functions for any  $t \ge 0$ 

$$\hat{p}(t) = \sum_{k=1}^{\infty} p_k \mathbb{1}_{[t_{k-1}, t_k)}(t) , \quad \hat{P}_0(t) = \int_0^t \hat{p}(s) ds ,$$

where  $t_k$  is defined in (8). Let  $\{s_k, k \in \mathbb{N}^*\}$  be an increasing sequence of positive real numbers, such that  $\lim_{k\to\infty} s_k = \infty$ . For any  $k \in \mathbb{N}^*$  and  $t \in \mathbb{R}_+$  we define

(27) 
$$\hat{P}_k(t) = \hat{P}_0(t+s_k) - \hat{P}_0(s_k)$$

The proof is divided into three steps.

STEP 1. The family of functions  $\{(X_k, \hat{P}_k) : k \in \mathbb{N}\}$ , defined by (9) and (27), respectively, is precompact in the topology of compact convergence.

*Proof.* The boundedness of  $\{X_k, k \in \mathbb{N}\}$  is trivial due to the fact, that  $x_k \in K$  for all  $k \ge 0$ . Using (26) and noting that, since projection is a contraction,

(28) 
$$\gamma_k^{-1} \|x_k - x_{k-1}\| \le \|v_k\| + \|r_k\| + \|e_k\|$$

we get that  $\sup_k ||p_k|| < \infty$ . Next, since

$$\hat{P}_k(t) = \int_{s_k}^{t+s_k} \hat{p}(s) \mathrm{d}s \; ,$$

and for every  $s \ge 0$ ,  $\|\hat{p}(s)\| \le \sup_k \|p_k\|$  we get  $\|\hat{P}_k(t)\| \le \sup_k \|p_k\| t$ .

In addition since  $\sup_k ||p_k|| < \infty$  and by (28)  $\sup_k \gamma_k^{-1} ||x_k - x_{k-1}|| < \infty$ , for all  $k \in \mathbb{N}$  the functions  $\hat{P}_k$ , and  $X_k$  are Lipschitz continuous with a Lipschitz constant, which does not depend on k. Hence  $\{(X_k, P_k), k \in \mathbb{N}^*\}$  is equicontinuous.

By Arzela-Ascoli theorem thee exists subsequence  $\{n_k, k \in \mathbb{N}\}$  such that  $\{X_{n_k}, k \in \mathbb{N}\}$  converges uniformly on compact subsets to a limit  $X_{\infty}$  and  $\{\hat{P}_{n_k}, k \in \mathbb{N}\}$  converges uniformly on compact subsets to  $P_{\infty}$ .

STEP 2. Any limit  $(X_{\infty}, P_{\infty})$  of converging subsequence is Lipschitz continuous and for almost every  $t \ge 0$  there exists subsequence  $\{n_k, k \in \mathbb{N}\}$  such that

$$X_{\infty}(t) = \lim_{k \to \infty} x_{n_k} = \lim_{k \to \infty} y_{n_k} \, .$$

*Proof.* Since  $\{X_k, k \in \mathbb{N}^*\}$  and  $\{\hat{P}_k, k \in \mathbb{N}^*\}$  are Lipschitz continuous, so the limits  $X_{\infty}$ ,  $P_{\infty}$  are also Lipschitz continuous. Along the same lines as in Step 2 of proof of Theorem 3.2, i.e. from (16), (17) and (18), we get that

(29) 
$$\lim_{k \to \infty} y_{m(n_k,t)} = X_{\infty}(t) ,$$

where  $m(n_k, t)$  is defined in (15).

STEP 3. Any limit  $X_{\infty}$  of converging subsequence is a solution of the projected differential inclusion  $\dot{x} \in \prod_{F_K(x)} (F(x))$ .

We use the following characterization of the solution of projected differential inclusion  $\dot{x} \in \Pi_{T_K(x)}(F(x))$  given in [5, Chapter 5, Section 6, Propositions 1 and 2]:

- (i)  $X_{\infty}$  is absolutely continuous
- (ii) For all  $t \ge 0$  we have  $X_{\infty}(t) \in K$ .
- (iii) For almost every  $t \ge 0$  there exists  $w(X_{\infty}(t)) \in F(X_{\infty}(t))$  such that,

$$\dot{X}_{\infty}(t) - w(X_{\infty}(t)) \in -\mathbf{N}_K(X_{\infty}(t)).$$

The condition (i) is already proved in Step 2. First we show (ii). Since *K* is a convex and by construction for all  $k \in \mathbb{N}$  and  $t \ge 0$ ,  $X_k(t)$  is convex combination of two elements of *K*,  $X_k(t) \in K$ . Because *K* is compact and  $X_\infty$  is the limit of convergent subsequence  $\{X_{n_k}, k \in \mathbb{N}\}$  also for all  $t \ge 0$ ,  $X_\infty(t) \in K$ .

It remains to show the condition (iii). We choose a compact interval  $[a,b] \subset \mathbb{R}_+$ . Let G and Q denote the weak derivatives of  $X_{\infty}$  and  $P_{\infty}$ , respectively. Recall that by assumption (A3) the family of functions  $\{\hat{E}_k, k \in \mathbb{N}\}$  is uniformly bounded on  $\mathbb{R}$ , and converges uniformly to 0 on compact intervals. Hence  $\{(X_{n_k} - \hat{E}_{n_k}, \hat{P}_{n_k}), k \in \mathbb{N}\}$  converges uniformly on compact sets to the limit  $(X_{\infty}, P_{\infty})$ . Because the functions  $\{(X_{n_k} - \hat{E}_{n_k}, \hat{P}_{n_k}), k \in \mathbb{N}\}$  are Lipschitz continuous with the same constant, their weak derivatives  $\{(G_{n_k}, Q_{n_k}), k \in \mathbb{N}\}$  are uniformly integrable. Hence, applying Lemma 6.6 we get that  $\{(G_{n_k}, Q_{n_k}), k \in \mathbb{N}\}$  converges in the weak topology of  $L_1([a,b])$  to (G,Q). By Lemma 6.8, we conclude that there exists a convex combination subsequence  $\{(G_{n_k}^w, Q_{n_k}^w), k \in \mathbb{N}\}$  that converges to (G,Q) for almost every  $t \in [a,b]$ .

Let  $t \in [a,b]$  be such, that  $\lim_{k\to\infty} (G_{n_k}^w(t), Q_{n_k}^w(t)) = (G(t), Q(t))$ . Since  $Q_{n_k}(t) = p_{m(n_k,t)}$ , where  $m(n_k,t)$  is defined in (15),  $Q(t) = \lim_{k\to\infty} p_{m(n_k,t)}^w$ .

Inequality (5) shows that for any  $l \in \mathbb{N}$  and  $z \in K$ ,  $\langle x_l - z, p_l \rangle \leq 0$ , or equivalently  $-p_l \in N_K(x_l)$ , see definition (24). By [5, Chapter 5, Section 1, Theorem 1] the normal cone has closed graph. On the other hand, since  $\sup_l ||p_l|| = r < \infty$ , by [5, Chapter 1, Section 1, Theorem 1] the map  $x \mapsto N_K(x) \cap B(0, r)$  is upper hemicontinuous. Therefore, since  $\lim_{k\to\infty} x_{m(n_k,t)} = X_{\infty}(t)$ , we can apply Lemma 6.9 to conclude that  $Q(t) = \lim_{k\to\infty} Q_{n_k}^w(t)$  belongs to the minus normal cone of K at  $X_{\infty}(t)$ , i.e.  $-Q(t) \in N_K(X_{\infty}(t))$ .

On the other hand,  $\lim_{k\to\infty} (G_{n_k}^w(t) - Q_{n_k}^w(t)) = G(t) - Q(t)$  and since for any  $m \in \mathbb{N}$ 

$$\frac{x_{m+1} - x_m}{\gamma_{m+1}} - \left(\frac{x_{m+1} - x_m}{\gamma_{m+1}} - v_m - e_m - r_m\right) - e_m = v_m + r_m$$

we get  $G_{n_k}^w(t) - Q_{n_k}^w(t) = v_{m(n_k,t)}^w + r_{m(n_k,t)}^w$ .

For all  $t \ge 0$ , since  $\lim_{k\to\infty} ||r_{m(n_k,t)}|| = 0$ , we have  $\lim_{k\to\infty} ||r_{m(n_k,t)}^w|| = 0$ . Therefore, for almost every  $t \in [a,b]$ , we get that  $\lim_{k\to\infty} v_{m(n_k,t)}^w = G(t) - Q(t)$ . On the other hand, for all  $t \in \mathbb{R}_+$ ,  $v_{m(n_k,t)} \in F(y_{m(n_k,t)})$  and by (29),  $\lim_{k\to\infty} y_{m(n_k,t)} = X_{\infty}(t)$ . Since *F* is upper hemicontinuous closed convex we apply Lemma 6.9 to show that  $G(t) - Q(t) \in F(X_{\infty}(t))$ . We denote  $w(X_{\infty}(t)) = G(t) - Q(t)$ . Hence for almost every  $t \in [a,b]$  we have:

(30) 
$$\dot{X}_{\infty}(t) - w(X_{\infty}(t)) = G(t) - (G(t) - Q(t)) = Q(t) \in -N_K(X_{\infty}(t)).$$

We therefore proved, that for almost all  $t \in [a, b] \subset \mathbb{R}_+$  there exists  $w(\dot{X}_{\infty}(t)) \in F(X_{\infty}(t))$ , such that the (30) holds. Since there is countable cover of real line by compact intervals, by diagonal extraction argument we can define  $w(\dot{X}_{\infty}(t))$ , which satisfies (30) for almost all  $t \in \mathbb{R}_+$ . Hence condition (iii) holds, and that concludes the proof.

THEOREM 4.2. Let  $K \subset \mathbb{R}^d$  be a convex and compact,  $X \subseteq \mathbb{R}^d$  be an open subset and  $F : X \mapsto \mathscr{P}(\mathbb{R}^d)$  be a set valued map,  $\{x_k, k \in \mathbb{N}^*\}$  be a K-PPAD of F with step sizes  $\{\gamma_k, k \in \mathbb{N}^*\}$  and perturbations  $\{\eta_k, k \in \mathbb{N}^*\}$ . Assume that conditions (A1–4) hold and  $\sup_k ||e_k|| < \infty$ . Let  $V : X \to \mathbb{R}$  be a locally Lipschitz, regular function. Suppose that there exists an upper semicontinuous function  $U : X \to \mathbb{R}$ , such that for all  $x \in K$ 

$$\sup \mathscr{L}_{\Pi_{\mathcal{T}_{K}}(F)}V(x) \leq U(x) \leq 0,$$

and set  $\mathscr{S} := \{x \in \mathsf{X} : U(x) = 0\}.$ 

- (i) The image by V of the set of limiting points of  $\{x_k, k \in \mathbb{N}^*\}$  is a compact interval in  $V(\mathscr{S} \cap K)$ .
- (ii) If  $V(\mathscr{S} \cap K)$  has empty interior, then  $\{x_k, k \in \mathbb{N}^*\}$  converges to  $K \cap \mathscr{S}$ .

*Proof.* The proof is along the same lines as the proof of Theorem 3.5.

**5. Applications.** In this section we apply the result from previous sections to projected ProxSGDand projected subgradient descent algorithm.

**5.1. Proximal stochastic gradient descent algorithm.** Stochastic proximal gradient is a natural extension of Proximal Gradient algorithm to the case where the gradient cannot be computed exactly and is therefore affected by some errors. More specifically, we want to optimize a composite function of form

$$P(x) = f(x) + g(x)$$

where *f* is a continuously differentiable function on some open set  $X \subseteq \mathbb{R}^d$ , and *g* is locally Lipschitz, bounded from below, regular function (see Definition 2.8).

In many Machine Learning applications f is the empirical risk of the model, and g a sparsity inducing penalties like LASSO [39], MCP [41], or SCAD [20]. We consider applications in which the gradient  $\nabla f(x_k)$  cannot be computed but for which noisy estimates  $H_k$  of  $\nabla f(x_k)$  are available. It is well known that, even in the noiseless case, additional conditions are needed to guarantee that the successive iterates remains in compact set and further converge to stationary points. To overcome this issue we consider a projected version of algorithm. For a predefined compact set  $K \subseteq X$ , we choose a sequence of step sizes  $\{\gamma_k, k \in \mathbb{N}\}$  satisfying (A2) and a starting point  $x_0 \in K$ .

Denote by  $\mathbb{I}_K$  the convex indicator function of set K ( $\mathbb{I}(x) = 0$  if  $x \in K$  and  $\mathbb{I}(x) = \infty$  otherwise) and by prox the proximal operator (see (6)).

We consider two versions of the projected ProxSGD algorithm which are given by

(31) 
$$x_k \in \operatorname{prox}_{\gamma_k(g+\mathbb{I}_K)}(x_{k-1} - \gamma_k H_k)$$

or

(32) 
$$x_k \in \Pi_K \left( \operatorname{prox}_{\gamma_k g} (x_{k-1} - \gamma_k H_k) \right)$$

Those two approaches to projection are not equivalent, and depending on g and K one might be easier to compute than the other. Consider the following assumptions:

- (P1)  $X \subset \mathbb{R}^d$  be open set,  $f : X \to \mathbb{R}$  be a continuously differentiable function, and  $g : X \to \mathbb{R}$  be a locally Lipschitz, bounded from below, regular function (see Definition 2.8).
- (P2) The sequence of step sizes  $\{\gamma_k, k \in \mathbb{N}^*\}$  satisfies  $\gamma_k > 0$ ,  $\sum_{k=0}^{\infty} \gamma_k = \infty$ , and  $\lim_{k \to \infty} \gamma_k = 0$ .

Denote by  $\{\delta_k, k \in \mathbb{N}^*\}$  the gradient perturbation

$$\delta_k = H_k - \nabla f(x_{k-1})$$

(P3) The sequence  $\{\delta_k, k \in \mathbb{N}^*\}$  can be decomposed as  $\delta_k = e_k^{\delta} + r_k^{\delta}$  where  $\{e_k^{\delta}, k \in \mathbb{N}^*\}$  and  $\{r_k^{\delta}, k \in \mathbb{N}^*\}$  are two sequences satisfying  $\lim_{k \to \infty} ||r_k^{\delta}|| = 0$  and the series  $\sum_{k=1}^{\infty} \gamma_k e_k^{\delta}$  converges.

To illustrate our derivations, we consider now two possible choices of sparsity inducing penalties g.

*Example* 5.1 (MCP penalty). The MCP penalty, introduced in [41] is given for  $x = (x_1, \ldots, x_d) \in \mathbb{R}^d$  by  $g(x) = \sum_{i=1}^d p_{\lambda,\kappa}(x_i)$  where for  $z \in \mathbb{R}$ ,  $\lambda > 0$  and  $\kappa > 1$ ,

$$p_{\lambda,\kappa}(z) = \begin{cases} \lambda |z| - z^2 / (2\kappa) & \text{if } |z| \le \kappa \lambda \\ 1 / (2\kappa\lambda^2) & \text{otherwise.} \end{cases}$$

The function g is Lipschitz on  $\mathbb{R}^d$  and nonnegative. If  $z \neq 0$ ,  $p_{\lambda,\kappa}$  is differentiable at z and hence  $p_{\lambda,\kappa}$  is regular at z by [16, Proposition 2.3.6]. The function  $z \mapsto \lambda |z|$  is convex and therefore regular on  $\mathbb{R}$ , applying again [16, Proposition 2.3.6]. The function  $z \mapsto -z^2/(2\kappa)$ is differentiable on  $\mathbb{R}$  and therefore regular. The sum of regular functions being regular by [16, Proposition 2.3.6],  $z \mapsto \lambda |z| - z^2/(2\kappa)$  is regular on  $\mathbb{R}$ . Therefore,  $p_{\lambda,\kappa}$  is regular at 0. By Lemma 6.2, the function g is also regular and satisfy (P1). The proximal operator for the MCP penalty  $p_{\lambda,\kappa}$  is the function given (see for example [14]), for all  $\gamma \in (0, \kappa)$  by

$$\operatorname{prox}_{\gamma p_{\lambda,\kappa}}(z) = \begin{cases} S(z,\gamma\lambda)/(1-\gamma/\kappa) & \text{for } |z| < \lambda \kappa \\ z & \text{otherwise }, \end{cases}$$

where  $S(z, \lambda) = \text{sign}(z)(|z| - \lambda)_+$  is the soft thresholding operator (here  $(x)_+ = x \lor 0$  is the positive part of *x*). The proximal operator for *g* is given by (see [33, Section 2.1])

$$\operatorname{prox}_{\gamma g}(x_1,\ldots,x_d) = \left(\operatorname{prox}_{\gamma p_{\lambda,\kappa}}(x_1),\ldots,\operatorname{prox}_{\gamma p_{\lambda,\kappa}}(x_d)\right) \,.$$

The SCAD penalty ([20]) can be handled along the same lines (the expression for the proximal function can be found in [14, Section 2]).

PROPOSITION 5.2. Assume (P1)–(P3) is satisfied. Then, the sequence  $\{x_k, k \in \mathbb{N}\}$  defined by (31) is K-PPAD (see Definition 2.1-(4)) with  $F = -\nabla f - \bar{\partial}g$ , noise sequence  $\{\eta_k, k \in \mathbb{N}\}$  where for each  $k \in \mathbb{N}^*$ ,  $\eta_k = -\delta_k + \{\nabla f(x_k) - \nabla f(x_{k-1})\}$  and  $y_k = x_k$ . Moreover, Assumptions (A1–4) are satisfied and, if  $\sup_{k \in \mathbb{N}^*} \|\delta_k\| < \infty$  then  $\sup_{k \in \mathbb{N}^*} \|\eta_k\| < \infty$ .

*Proof.* We denote by F the set-valued map  $-\nabla f - \overline{\partial}g$ . The Clarke gradient of a locally Lipschitz function is convex-compact valued and locally bounded (see [16, Proposition 2.1.2]) and is upper hemi-continuous (see [16, Proposition 2.1.5]. Since  $\nabla f$  is continuous, this implies that F satisfies (A1). By [16, Corollary 2.3.2],  $\overline{\partial}(-f-g) = -\nabla f - \overline{\partial}g$ .

First, we need to show that  $\{x_k, k \in \mathbb{N}\}$  generated by (31) can be seen as PPAD. Suppose that  $\{x_k, k \in \mathbb{N}\}$  is generated by iteration (31). Denote by

(34) 
$$w_k = x_{k-1} - \gamma_k \{ \nabla f(x_{k-1}) + \delta_k \}.$$

With this notation (31) can be written as

$$x_k = \operatorname{prox}_{\gamma_k(g+\mathbb{I}_K)}(w_k)$$
.

By [16, corollary of Proposition 2.4.3, p. 52],  $0 \in \gamma_k^{-1}(x_k - w_k) + \bar{\partial}g(x_k) + N_K(x_k)$ . Therefore there exists

$$(35) u_k \in \bar{\partial}g(x_k)$$

such that

(36) 
$$x_k - w_k + \gamma_k u_k \in -\gamma_k \mathbf{N}_K(x_k) .$$

The normal cone to *K* at  $x_k$  consists of vectors  $v_c$ , such that for all  $z \in K$  we have  $\langle v_c, x_k - z \rangle \ge 0$ . Since  $\gamma_k N_K(x_k) = N_K(x_k)$  and using (34), (36) implies that for all  $z \in K$ ,

(37) 
$$\langle x_k - w_k - \gamma_k u_k, x_k - z \rangle = \langle x_k - x_{k-1} + \gamma_k (\nabla f(x_{k-1}) + \delta_k + u_k), x_k - z \rangle \le 0,$$

where  $u_k$  is defined in (35). Setting  $v_k = -\nabla f(x_k) - u_k \in F(x_k)$  and  $\eta_k = -\delta_k + (\nabla f(x_k) - \nabla f(x_{k-1}))$  we get

$$\langle x_k-z, x_k-x_{k-1}-\gamma_k(v_k+\eta_k)\rangle \leq 0$$

which is (5), but with noise sequence  $\{\eta_k, k \in \mathbb{N}\}$ .

We finally have to check that (A3). Since  $\{\delta_k, k \in \mathbb{N}\}$  satisfies condition (P3) and

(38) 
$$\eta_k = \nabla f(x_k) - \nabla f(x_{k-1}) - \delta_k = \nabla f(x_k) - \nabla f(x_{k-1}) - r_k^{\delta} - e_k^{\delta}$$

it is enough to show that  $\lim_{k\to\infty} \|\nabla f(x_{k-1}) - \nabla f(x_k)\| = 0$ . Plugging  $z = x_{k-1}$  into (37), we get that:

$$\langle x_k - w_k + \gamma_k u_k, x_k - x_{k-1} \rangle \leq 0$$
,

Hence

$$\begin{aligned} \|x_k - x_{k-1}\|^2 &\leq \langle x_k - x_{k-1} - (x_k - w_k + \gamma_k u_k), x_k - x_{k-1} \rangle \\ &= \langle -x_{k-1} + w_k - \gamma_k u_k, x_k - x_{k-1} \rangle , \end{aligned}$$

and using the Cauchy-Schwartz and triangle inequalities and (34), we obtain

(39) 
$$\|x_k - x_{k-1}\| \le \|-x_{k-1} + w_k - \gamma_k u_k\| \le \gamma_k \|\nabla f(x_{k-1})\| + \gamma_k \|\delta_k\| + \gamma_k \|u_k\| .$$

Since  $x_k \in K$  for any  $k \in \mathbb{N}$  and  $\nabla f$  is continuous under (P1),  $\sup_{k \in \mathbb{N}} ||\nabla f(x_{k-1})|| < \infty$ . On the other hand, under (P3),  $\lim_{k\to\infty} \gamma_k ||\delta_k|| = 0$ . Finally, using again (P1) and  $x_k \in K$  for any  $k \in \mathbb{N}^*$ , [16, Proposition 2.1.2] shows that  $\sup_{k \in \mathbb{N}^*} ||u_k|| < \infty$ . Therefore,  $\lim_{k\to\infty} ||x_k - x_{k-1}|| = 0$  under (P2) and, the continuity of  $\nabla f$  combined with the decomposition (38) shows that (A3) holds.

PROPOSITION 5.3. Assume (P1)–(P3) is satisfied and that  $g: X \to \mathbb{R}$  is Lipschitz. Then, the sequence  $\{x_k, k \in \mathbb{N}\}$  defined by (32) is K-PPAD with  $F = -\nabla f - \overline{\partial}g$  (where  $\overline{\partial}$  denotes the Clarke generalized gradient) and noise  $\{\eta_k, k \in \mathbb{N}\}$  where for each  $k \in \mathbb{N}^*$ ,  $\eta_k = -\delta_k + \{\nabla f(x_k) - \nabla f(x_{k-1})\}$ . Moreover, Assumptions (A1–4) are satisfied and, if  $\sup_{k \in \mathbb{N}^*} ||\delta_k|| < \infty$ then  $\sup_{k \in \mathbb{N}^*} ||\eta_k|| < \infty$ .

Proof. Denote by

$$y_k = \operatorname{prox}_{\gamma_k g} (x_{k-1} - \gamma_k \nabla f(x_{k-1}) - \gamma_k \delta_k)$$
.

Using the definitions of proximal and [16, Proposition 2.3.2], for each  $k \in \mathbb{N}^*$  there exists  $u_k \in \bar{\partial}g(y_k)$  such that

$$\mathbf{y}_k = \mathbf{x}_{k-1} - \mathbf{\gamma}_k \nabla f(\mathbf{x}_{k-1}) - \mathbf{\gamma}_k \mathbf{\delta}_k - \mathbf{\gamma}_k \mathbf{u}_k$$

Denoting by  $v_k = -\nabla f(y_k) - u_k$  and by  $\eta_k = -\delta_k - \nabla f(x_{k-1}) + \nabla f(y_k)$  we get that

$$y_k = x_{k-1} + \gamma_k(v_k + \eta_k) .$$

We now chack (A3). The perturbation  $\eta_k$  may be decomposed as  $\eta_k = e_k + r_k$  where  $e_k = -e_k^{\delta}$ and  $r_k = -r_k^{\delta} + \nabla f(y_k) - \nabla f(x_{k-1})$ . Since  $\nabla f$  is continuous, Assumption (A3) is satisfied if

$$\lim_{k\to\infty}\|y_k-x_{k-1}\|=0$$

Note that, since g is Lipschitz, [16, Proposition 2.1.2-(a)] shows that for all  $u \in \bar{\partial}g(y)$ ,  $||u|| \le ||g||_{\text{Lip}} < \infty$ . Boundedness of  $\nabla f$  on the set K and assumptions (P2) and (P3) implies that

$$\limsup_{k \to \infty} \|x_{k-1} - y_k\| \le \limsup_{k \to \infty} \gamma_k \|\nabla f(x_{k-1}) + \delta_k\| + \limsup_{k \to \infty} \gamma_k \|u_k\| = 0$$

Because  $x_k$  is a projection of  $y_k$  on the set K we have

$$||y_k - x_{k-1}|| \ge ||y_k - x_k||$$
,

showing that (A4) is satisfied.

Applying our results from Section 4, we now show that both versions of the projected proximal gradient algorithms converge.

THEOREM 5.4. Assume (P1-3) and denote

(40) 
$$\mathscr{S} := \left\{ x \in K : 0 \in \nabla f(x) + \bar{\partial}g(x) - \mathcal{N}_K(x) \right\},$$

where  $\overline{\partial}g$  is the Clarke gradient of g (see Definition 2.6), and  $N_K(x)$  is the normal cone to set K at  $x \in K$  (see (24)). Suppose  $(f+g)(\mathscr{S})$  has empty interior and  $\sup_{k \in \mathbb{N}^*} \|\delta_k\| < \infty$ , where  $\delta_k$  is defined in (33).

- (i) The sequence  $\{x_k, k \in \mathbb{N}\}$  generated by iterations (31) converges to the  $\mathscr{S}$ .
- (ii) Assume in addition that g is Lipschitz. Then the sequence  $\{x_k, k \in \mathbb{N}\}$  generated by iterations (32) converges to the  $\mathscr{S}$ .

*Proof.* Using Propositions 5.2 and 5.3, the sequences  $\{x_k, k \in \mathbb{N}\}$  defined by (31) and (32) are *K*-PPAD satisfying (A1–4) with

$$F = -\nabla f - \bar{\partial}g$$

To apply Theorem 4.2, we show that V = f + g is a Lyapunov function for  $F_K(x) := \prod_{T_K(x)} (F(x))$ ,  $x \in X$ . Under the stated assumptions, V is locally Lipschitz regular and by [16, Corollary 2 of Proposition 2.3.3]

$$\bar{\partial}V(x) = \nabla f(x) + \bar{\partial}g(x), \quad x \in \mathsf{X}.$$

We now compute the Lie derivative of *V* with respect to the field  $F_K$  see Definition 3.3). Let  $x \in K$ . Suppose that there exists  $a \in \mathscr{L}_{F_K}V(x)$ . Then there exists  $v \in \Pi_{T_K(x)}(F(x))$ , such that for all  $w \in \bar{\partial}V(x)$ ,  $\langle v, w \rangle = a$ . Let  $u \in F(x)$  be such that  $\Pi_{T_K(x)}(u) = v$ . Note that  $-u \in \bar{\partial}V(x)$ , which implies  $\langle \Pi_{T_K(x)}(u), -u \rangle = a$  and

$$a = - \|\Pi_{\mathbf{T}_{K}(x)}(u)\|^{2} + \langle \Pi_{\mathbf{T}_{K}(x)}(u), \Pi_{\mathbf{T}_{K}(x)}(u) - u \rangle$$

Applying [5, Proposition 0.6.2], we get that  $\langle \Pi_{T_K(x)}(u), \Pi_{T_K(x)}(u) - u \rangle = 0$ . Therefore, if  $a \in \mathscr{L}_{F_K}V(x)$ , then  $a = -\|\Pi_{T_K(x)}(u)\|^2$  for some  $u \in F(x)$ . Hence, for any  $x \in K$ ,  $\mathscr{L}_{F_K}V(x)$  is either empty, or contains only non-positive elements.

For any  $x \in X$ , [16, Proposition 2.1.2] shows that F(x) is non-empty, convex and compact. Define U for any  $x \in K$  as follows

$$U(x) = -\min_{u \in F(x)} \|\Pi_{\mathbf{T}_{K}(x)}(u)\|^{2}.$$

By [5, Proposition 0.6.4] we get for any  $x \in K$ ,

(41) 
$$U(x) = -\min_{v \in F(x) - N_K(x)} ||v||^2 .$$

Under (A1), *F* is upper hemicontinuous compact-convex valued. On the other hand, by [5, Chapter 5, Section 1, Theorem 1], N<sub>K</sub> has closed graph. Hence, the map  $F - N_K$  has closed graph by [5, Proposition 1.1.2, p. 41]. By Lemma 6.1 the function *U* is upper semicontinuous. Thus we have an upper semicontinuous function *U*, such that for all  $x \in K$ :

$$\sup \mathscr{L}_{F_K} V(x) \leq U(x) \leq 0.$$

Using (41), we get that

$$\{x \in K \colon U(x) = 0\} = \{x \in K \colon 0 \in -\nabla f(x) - \overline{\partial}g(x) - N_K(x)\} = \mathscr{S},$$

which concludes the proof

**5.2.** Online proximal stochastic gradient descent algorithm. Let X be an open subset of  $\mathbb{R}^d$  and  $K \subset \mathbb{R}^d$  be a nonempty bounded closed convex set. We first consider the following composite minimization problem

(42) 
$$\min_{x \in K} \{ f(x) + g(x) \}$$

where  $f : \mathbb{R}^d \to \mathbb{R}$  is continuously differentiable on a neighborhood of *K* and  $g : X \to \mathbb{R}^d$  is locally Lipschitz, bounded from below, regular function.

In the online learning case, the gradient  $\nabla f$  of the function f cannot be computed but that a noisy version of the gradient is available To make the discussion simple, we assume that

(i) an i.i.d. sequence  $\{\xi_k, k \in \mathbb{N}\}$  with common distribution  $\pi$  on a measurable space  $(\mathsf{Z}, \mathscr{Z})$ 

(ii) there is an oracle which for a given input point (x, ξ) ∈ X × Z returns a stochastic gradient, a vector H(x, ξ) such that E[Φ(x, ξ)] = ∫<sub>Z</sub>Φ(x, z)π(dz) is well defined and is equal to ∇f(x).

We are considering the two following stochastic approximation procedures

(43) 
$$x_{k} = \operatorname{prox}_{\gamma_{k}(g+\mathbb{I}_{K})} \{ x_{k-1} - \gamma_{k} \Phi(x_{k-1}, \xi_{k}) \}$$

or

(44) 
$$x_k = \Pi_K \left( \operatorname{prox}_{\gamma_k g} \{ x_{k-1} - \gamma_k \Phi(x_{k-1}, \xi_k) \} \right)$$

where  $\{\gamma_k, k \in \mathbb{N}\}$  is a sequence of step sizes satisfying  $\sum_{k=1}^{\infty} \gamma_k^{1+\varepsilon} < \infty$  for some  $\varepsilon > 0$ . Clearly such  $\{\gamma_k, k \in \mathbb{N}\}$  satisfies (P2). Assume for simplicity that the essential supremum of  $\sup_{x \in K} \|\Phi(x, \cdot)\|$  is finite

(45) 
$$\inf \left\{ a \in \mathbb{R} : \pi \left( \sup_{x \in K} \| \Phi(x, z) \| > a \right) = 0 \right\} < \infty.$$

Setting, for all  $k \in \mathbb{N}^*$ ,  $\delta_k = \Phi(x_{k-1}, \xi_k) - \nabla f(x_{k-1})$ , we get that  $\sup_k ||\delta_k|| < \infty \mathbb{P}$ -almost surely and since  $\{\delta_k, k \in \mathbb{N}\}$  is a bounded martingale increment sequence the conditional version of the Kolmogorov three-series theorem shows that (see [25, Theorem 2.16] shows that  $\sum_{k=1}^{\infty} \gamma_k \delta_k < \infty$  almost surely. Therefore (P3) holds. Hence, if  $(f + g)(\mathscr{S})$  has an empty interior (where  $\mathscr{S}$  is the set of stationary point defined in (40)), Theorem 5.4 implies that  $\{x_k, k \in \mathbb{N}\}$  generated by (43) almost surely converge to the set of stationary points (40) If in addition the function g is Lipschitz, then the sequence  $\{x_k, k \in \mathbb{N}^*\}$  generated by (44) also converges almost surely to  $\mathscr{S}$ .

**5.3. Monte Carlo Proximal stochastic gradient descent algorithm.** In this section, we still consider the composite minimization problem (42). We assume that *f* is continuously differentiable and that for all  $x \in K \nabla f(x)$  satisfies

(46) 
$$\nabla f(x) = \int_{\mathsf{Z}} \Phi(x, z) \pi_x(\mathrm{d} z) \;,$$

for some probability measure  $\pi_x$  and an integrable function  $(x,z) \mapsto \Phi(x,z)$  from  $K \times Z$  to  $\mathbb{R}^d$ . Note the dependence of  $\pi_x$  on x which makes the situation trikier. To approximate  $\nabla f(x)$ , several options are available. Of course, when the dimension of the state space Z is small to moderate, it is always possible to perform a numerical integration using either Gaussian quadratures or low-discrepancy sequences. Such approximations necessarily introduce some bias, which might be difficult to control. In addition, these techniques are not applicable when the dimension of Z becomes large. In this paper, we rather consider some form of Monte Carlo approximation.

When sampling directly  $\pi_x$  is doable, then an obvious choice is to use a naive Monte Carlo estimator which amounts to sample a batch  $\{z_k^{(j)}, 1 \le j \le m\}$  independently of the past values of the parameters  $\{x_j, j \le k-1\}$  and of the past draws i.e. independently of the  $\sigma$ -algebra

(47) 
$$\mathscr{F}_n := \sigma(x_0, z_k^{(j)}, 0 \le k \le n, 0 \le j \le m) .$$

We then form

$$Y_k = m^{-1} \sum_{j=0}^{m-1} \Phi(x_{k-1}, z_k^{(j)})$$

Conditionally to  $\mathscr{F}_{k-1}$ ,  $Y_k$  is an unbiased estimator of  $\nabla f(x_{k-1})$ .

When direct sampling from  $\pi_x$  is not an option, we may still construct a Markov kernel  $P_x$  with invariant distribution  $\pi_x$ . Monte Carlo Markov Chains (MCMC) provide a set of principled tools to sample from complex distributions over large dimensional spaces. In such case, conditional to the past,  $\{z_k^{(j)}, 1 \le j \le m\}$  is a realization of a Markov chain with transition kernel  $M_{x_{k-1}}$  and started from  $z_{k-1}^{(m)}$  (the last sample draws in the previous minibatch).

Recall that a Markov kernel *P* is an application on  $Z \times \mathscr{Z}$ , taking values in [0,1] such that for any  $z \in Z$ ,  $P(z, \cdot)$  is a probability measure on  $\mathscr{Z}$ ; and for any  $A \in \mathscr{Z}$ ,  $z \mapsto P(z,A)$  is measurable. Furthermore, if *P* is a Markov kernel on *Z*, we denote by  $P^k$  the *k*-th iterate of *P* defined recursively as  $P^0(z,A) := \mathbb{1}_A(z)$ , and  $P^k(z,A) := \int P^{k-1}(z,dz)P(z,A)$ ,  $k \ge 1$ . Finally, the kernel *P* acts on probability measure: for any probability measure  $\mu$  on  $\mathscr{Z}$ ,  $\mu P$  is a probability measure defined by

$$\mu P(A) := \int \mu(\mathrm{d}z) P(z,A), \qquad A \in \mathscr{Z};$$

and *P* acts on positive measurable functions: for a measurable function  $f : Z \to \mathbb{R}_+$ , *Pf* is a function defined by

$$Pf(z) := \int f(y) P(z, \mathrm{d}y)$$

We refer the reader to [29] for the definitions and basic properties of Markov chains.

In this section, we assume that  $Y_k$  is a Monte Carlo approximation of the expectation  $\nabla f(x_{k-1})$ :

$$Y_k = m^{-1} \sum_{j=0}^{m-1} \Phi(x_{k-1}, z_k^{(j)});$$

for all  $k \ge 1$ , conditionally to the past,  $\{z_k^{(j)}, 1 \le j \le m\}$  is a Markov chain started from  $z_{k-1}^{(m)}$ and with transition kernel  $M_{x_{k-1}}$  (we set  $z_0^{(m)} = x_{\star} \in X$ ), where for all  $x \in X$ ,  $M_x$  is a Markov kernel with invariant distribution  $\pi_x$ .

From a mathematical standpoint, the Markovian setting is trickier than the fixed batch size, because  $Y_k$  is no longer an unbiased estimator of  $\nabla f(x_{k-1})$ , i.e. the bias  $B_k$  defined by

(48)  

$$B_{k} := \mathbb{E}\left[Y_{k} \mid \mathscr{F}_{k-1}\right] - \nabla f(x_{k-1}) = m^{-1} \sum_{j=0}^{m-1} \mathbb{E}\left[\Phi(x_{k-1}, z_{k}^{(j)}) \mid \mathscr{F}_{k-1}\right] - \nabla f(x_{k-1})$$

$$= m^{-1} \sum_{j=0}^{m-1} M_{x_{k-1}}^{j} \Phi(x_{k-1}, z_{k}^{(0)}) - \nabla f(x_{k-1}),$$

does not vanish.

Let  $W : Z \to [1,\infty)$  be a measurable function. The *W*-norm of a measurable function  $h : Z \to \mathbb{R}^{\ell}$  is defined as  $|h|_W = \sup_{z \in Z} ||h(z)|| / W(z)$ . The *W*-variation of a finite signed measure  $\mu$  on the measurable space  $(Z, \mathscr{Z})$  as  $||\mu||_W := \sup_{|h|_W \leq 1} \mu(h)$  where supremum is taken over all measurable functions  $h : Z \to \mathbb{R}$  satisfying  $|h|_W \leq 1$ . Denote by  $D_W(x, x')$  the *W*-variation of the kernels  $M_x$  and  $M_{x'}$ 

$$D_W(x,x') = \sup_{z \in \mathsf{Z}} \frac{\|M_x(z,\cdot) - M_{x'}(z,\cdot)\|_W}{W(z)}.$$

Consider the following assumptions

(B1) There exists  $\lambda \in [0,1)$ ,  $b < \infty$  and a measurable function  $W : Z \rightarrow [1, +\infty)$  such that

$$\sup_{x\in K} \|\Phi(x,\cdot)\|_{W^{1/2}} < \infty, \quad \sup_{x\in K} M_x W \le \lambda W + b.$$

In addition for any  $\ell \in (0,1]$  there exists  $C < \infty$  and  $\rho \in (0,1)$  such that for any  $z \in Z$ ,

$$\sup_{x\in K} \|M_x^n(z,\cdot)-\pi_x\|_{W^\ell} \le C\rho^n W^\ell(z).$$

Sufficient conditions for the uniform-in-*x* ergodic behavior are given e.g. in [22, Lemma 2.3], in terms of aperiodicity, irreducibility and minorization conditions on the kernels  $\{M_x : x \in X\}$ . Examples of MCMC kernels  $M_x$  satisfying this assumption can be found in [2, Proposition 12], [38, Proposition 15].

(B2) The kernels  $M_x$  and the stationary distributions  $\pi_x$  are locally Lipschitz with respect to x, i.e. for any compact set K and any  $x, x' \in K$  there exists  $C < \infty$  such that

$$\left\|\Phi(x,\cdot) - \Phi(x',\cdot)\right\|_{W^{1/2}} + D_{W^{1/2}}(x,x') \le C \left\|x - x'\right\|$$

PROPOSITION 5.5. Let us consider  $\{x_k, k \in \mathbb{N}^*\}$  defined by (43) or (44) with Markovian dynamic  $\{z_k, k \in \mathbb{N}^*\}$  and with  $\Phi$  satisfying (46). Assume (B1–2), (P1)–(P2),  $\mathbb{E}[W(z_1)] < \infty$ , and

$$\sum_{k=1}^{\infty} |\gamma_{k-1} - \gamma_k| < \infty, \quad \sum_{k=1}^{\infty} \gamma_k^2 < \infty \ .$$

### Then (P3) is satisfied.

*Proof.* The proof follows along the same lines as [30, Proof of Lemma 27]. However for completeness we give a detailed proof in Appendix A.

**5.4. Projected stochastic subgradient descent algorithm.** We consider the projected subgradient descent algorithm framework introduced in [31], for constrained minimization of a possibly nonsmooth convex objective function. Let X be an open set of  $\mathbb{R}^d$  and  $K \subset X$  be a convex compact. Consider the constrained minimization problem  $\arg \min_{x \in K} f(x)$ , where *f* is locally Lipschitz regular function (see Definition 2.8). The projected stochastic subgradient algorithm generates iteratively the sequence  $\{x_k, k \in \mathbb{N}^*\}$  as follows

(49) 
$$x_k = \Pi_K (x_{k-1} - \gamma_k Y_k) ,$$

where  $\{\gamma_k, k \in \mathbb{N}^*\}$  is a sequence of positive step sizes,  $\Pi_K$  is the projection on the set *K* and  $Y_k$  is a noisy version of Clarke generalized gradient, i.e.  $Y_k = v_k + \delta_k$  with  $v_k \in \overline{\partial} f(x_{k-1})$ .

In [19], the stochastic subgradient defined recursively by  $x_k = x_{k-1} - \gamma_k Y_k$  (without projection) is analyzed, under the assumption that the iterates  $\{x_k, k \in \mathbb{N}\}$  stay in the compact set K and that the noise  $\{\delta_k, k \in \mathbb{N}^*\}$  is bounded. This paper establishes the almost-sure convergence of the iterates  $\{x_k, k \in \mathbb{N}^*\}$  to the stationary set  $\mathscr{S} := \{x \in K : 0 \in \overline{\partial} f(x)\}$ , under a descent condition on f. Specifically, it is assumed in [19] that if  $z : \mathbb{R}_+ \to \mathbb{R}^d$  is a solution of the differential inclusion  $\dot{z}(t) \in -\overline{\partial} f(z(t))$  and  $z(0) \notin \mathscr{S}$  (i.e. z(0) is not a critical point of f), then there exists a T > 0 such that  $f(z(T)) < \sup_{t \in [0,T)} f(z(t)) \leq f(z(0))$ . It is also proved in this paper that this condition is satisfied for two classes of functions: subdifferential regularity is equivalent to our condition of regularity (see Lemma 6.3), while the class of Whitney stratifiable functions, and contains for example the class of semialgebraic and semianalytic functions.

The convergence of stochastic subgradient algorithm for regular functions f can be easily deduced from Section 3. Here, we show that projected stochastic subgradient descent algorithm also fits into our framework and its convergence can be established based on the results of Section 4.

Consider the following assumptions:

- (**P**1)  $X \subset \mathbb{R}^d$  is an open set,  $K \subset X$  is a convex compact set,  $f : X \to \mathbb{R}$  is a locally Lipschitz, regular function (see Definition 2.8).
- (**\hat{P}2**) The sequence of step sizes  $\{\gamma_k, k \in \mathbb{N}^*\}$  satisfies  $\gamma_k > 0$ ,  $\sum_{k=0}^{\infty} \gamma_k = \infty$ , and  $\lim_{k\to\infty} \gamma_k = 0$ .
- (**P3**) The sequence  $\{\delta_k, k \in \mathbb{N}^*\}$  can be decomposed as  $\delta_k = e_k^{\delta} + r_k^{\delta}$  where  $\{e_k^{\delta}, k \in \mathbb{N}^*\}$  and  $\{r_k^{\delta}, k \in \mathbb{N}^*\}$  are two sequences satisfying  $\lim_{k\to\infty} ||r_k^{\delta}|| = 0$  and the series  $\sum_{k=1}^{\infty} \gamma_k e_k^{\delta}$  converges.

PROPOSITION 5.6. Assume  $(\hat{P}^{1}-3)$  is satisfied. Then, the sequence  $\{x_k, k \in \mathbb{N}\}$  defined by (49) is K-PPAD with  $F = -\bar{\partial}f$  (where  $\bar{\partial}$  denotes the Clarke generalized gradient) and noise  $\{\eta_k, k \in \mathbb{N}\}$  where for each  $k \in \mathbb{N}^*$ ,  $\eta_k = -\delta_k$ . Moreover, Assumptions (A1-4) are satisfied.

*Proof.* Set  $F = -\bar{\partial}f$ . The Clarke gradient of a locally Lipschitz function is convexcompact valued and locally bounded (see [16, Proposition 2.1.2]) and is upper hemi-continuous (see [16, Proposition 2.1.5]. Hence (A1) is satisfied. By construction the  $\{x_k, k \in \mathbb{N}\}$  generated by (49) is a K-PPAD with stepsizes  $\{\gamma_k, k \in \mathbb{N}^*\}$ , noise  $\eta_k = -\delta_k$  and  $y_k = x_{k-1}$ . Hence, assumptions ( $\hat{P}2-3$ ) implies (A2-3). We need to check only (A4). Denote for any  $k \in \mathbb{N}^*$  by  $w_k = x_{k-1} - \gamma_k Y_k = x_{k-1} - \gamma_k v_k - \gamma_k \delta_k$ , where  $v_k \in \bar{\partial}f(x_{k-1})$ . Since  $x_k$  is projection of  $w_k$  on the compact convex set K, the triangle inequality implies

(50) 
$$\|x_k - y_k\| = \|x_k - x_{k-1}\| \le \|w_k - x_{k-1}\| \le \gamma_k \|v_k\| + \gamma_k \|\delta_k\| .$$

Since  $\{x_k, k \in \mathbb{N}^*\}$  remains in the compact set *K* and  $\bar{\partial} f$  is locally bounded we obtain that  $\sup_k ||v_k|| < \infty$ . Thereofore by ( $\hat{P}2-3$ ) and (50) we get that  $\lim_{k\to\infty} ||x_k - y_k|| = 0$ , and that completes the proof.

Applying our results from Section 4, we now show that projected stochastic subgradient algorithms converge.

THEOREM 5.7. Assume  $(\hat{P}_{1-3})$  and denote

$$\mathscr{S} := \left\{ x \in K : 0 \in \overline{\partial} f(x) - \mathcal{N}_K(x) \right\},\$$

where  $\overline{\partial} f$  is the Clarke gradient of g (see Definition 2.6), and  $N_K(x)$  is the normal cone to set K at  $x \in K$  (see (24)). Suppose  $f(\mathscr{S})$  has empty interior and  $\sup_{k \in \mathbb{N}^*} ||\delta_k|| < \infty$ . Then the sequence  $\{x_k, k \in \mathbb{N}\}$  generated by the iterations (49) converges to the  $\mathscr{S}$ .

*Proof.* The proof follows along the sime lines as proof of Theorem 5.4.

Note, that adaptation of results from Subsection 5.2 and Subsection 5.3 to the case of projected stochastic subgradient descent algorithm is straightforward.

**6. Proofs.** In this section we introduce some notations and preliminary facts used in the proofs of results from Section 3 and Section 4, as well as some auxiliary definitions and theorems.

LEMMA 6.1. Let K be a compact subset of  $\mathbb{R}^d$ , and  $G: K \to \mathscr{P}(\mathbb{R}^d)$  be a nontrivial closed set-valued map (i.e. for any  $x \in K$ ,  $G(x) \neq \emptyset$  and the graph of the function G is closed;

see [16, Section 2]). Then  $W: K \to \mathbb{R}$  defined by

$$W(x) = -\min_{v \in G(x)} ||v||^2$$

is upper semicontinuous.

*Proof.* Let  $x \in K$  and  $\{x_n, n \in \mathbb{N}\} \subset K$  be any sequence such that  $\lim_{n\to\infty} x_n = x$ . Consider a subsequence  $\{x_{n_k}, k \in \mathbb{N}\}$  such that  $\lim_k W(x_{n_k}) = \limsup_n W(x_n)$ . For any  $\tilde{x} \in K$ , there exists  $\tilde{w} \in G(\tilde{x})$  such that  $-\|\tilde{w}\|^2 = W(\tilde{x})$  (since  $G(\tilde{x})$  is closed and  $\|\cdot\|$  is continuous). We may therefore define a sequence  $\{w_{n_k}, k \in \mathbb{N}\}$  such that for all  $k \in \mathbb{N}$ ,  $-\|w_{n_k}\|^2 = W(x_{n_k})$ . Because  $\lim_{k\to\infty} W(x_{n_k})$  is finite, the subsequence  $\{w_{n_k}, k \in \mathbb{N}\}$  is bounded. We may hence extract a subsequence  $\{w_{\tilde{n}_k}, k \in \mathbb{N}\} \subseteq \{w_{n_k}, k \in \mathbb{N}\}$  such that  $\lim_{k\to\infty} w_{\tilde{n}_k} = w$  for some  $w \in \mathbb{R}^d$ . Since *G* has closed graph we have  $w \in G(x)$ . By continuity of norm we get

$$\limsup_{n} W(x_{n}) = \lim_{k \to \infty} W(x_{\tilde{n}_{k}}) = -\lim_{k \to \infty} ||w_{\tilde{n}_{k}}||^{2} = -||w||^{2} \le W(x) ,$$

which means that W is upper semicontinuous.

LEMMA 6.2. Let  $d \in \mathbb{N}^*$ ,  $p_i : \mathbb{R} \to \mathbb{R}$ ,  $i \in \{1, ..., d\}$ , be functions and  $x^0 = (x_1^0, ..., x_d^0) \in \mathbb{R}^d$ . If for any  $i \in \{1, ..., d\}$  the functions  $p_i$  are regular at  $x_i^0$ , then the function  $g(x_1, ..., x_d) = \sum_{i=1}^d p_i(x_i)$  is regular at  $x^0$ .

*Proof.* By [16, Proposition 2.3.6] a finite linear combination (by nonnegative scalars) of functions regular at  $x^0$  is regular at  $x^0$ . The proof then follows by noting that, for any  $i \in \{1, ..., d\}$ , the function  $p^i(x_1, ..., x_d) = p_i(x_i)$  is regular at  $x^0$ .

LEMMA 6.3. Let  $f : \mathbb{R}^d \to \mathbb{R}$  be a locally Lipschitz function. Then f is regular at x if and only if for all  $v \in \overline{\partial} f(x)$  we have:

(51) 
$$f(y) \ge f(x) + \langle v, y - x \rangle + o(||x - y||).$$

*Proof.* Let  $\partial_D f(x)$  be the D-subdifferential of f at x (see [17, chapter 3.4, subsection D-differential] for definition). Then according to [17, Proposition 4.10], the inequality (51) is satisfied for v if and only if  $v \in \partial_D f(x)$ . On the other hand, [17, Proposition 4.8, part (b)] implies that for a locally Lipschitz function we have  $\overline{\partial} f(x) = \partial_D f(x)$  if and only if f is regular. Combined, these two facts conclude the proof.

DEFINITION 6.4 (Equicontinuity). A sequence of functions  $\{f_n, n \in \mathbb{N}\}$ , from  $\mathbb{R}$  to  $\mathbb{R}^d$ , is said to be equicontinuous at  $t_0$ , if for all  $\varepsilon > 0$  there exists  $\eta > 0$  such that for all  $|t - t_0| \le \eta$ ,  $||f_n(t) - f_n(t_0)|| \le \varepsilon$  for all  $n \in \mathbb{N}$ . A sequence of functions  $\{f_n, n \in \mathbb{N}\}$  from  $\mathbb{R}$  to  $\mathbb{R}^d$  is said to be equicontinuous, if and only if it is equicontinuous at every point of  $t_0 \in \mathbb{R}$ .

# THEOREM 6.5 (Arzela-Ascoli theorem).

Let  $\{f_n, n \in \mathbb{N}\}$  from  $\mathbb{R}$  to  $\mathbb{R}^k$  be a sequence of functions. Assume that the sequence  $\{f_n, n \in \mathbb{N}\}$  is equicontinuous and pointwise bounded (meaning that  $\sup_{n \in \mathbb{N}} ||f_n(x)||$  is finite for all  $x \in \mathbb{R}$ ). Then the sequence  $\{f_n, n \in \mathbb{N}\}$  is precompact in the topology of compact convergence.

Proof. See [21, Theorem 4.44].

LEMMA 6.6. Let  $\{f_n, n \in \mathbb{N}\}\$  be a sequence of functions from  $\mathbb{R}$  to  $\mathbb{R}^d$ , that are absolutely continuous on compact intervals, and converge pointwise to a function f, which is also absolutely continuous on compact intervals. For each  $n \in \mathbb{N}$ , let  $g_n$  be a weak derivative of  $f_n$  and g be a weak derivative of f. Also assume that  $\{g_n, n \in \mathbb{N}\}\$  are uniformly integrable on bounded intervals. Then for every interval  $[a,b], 0 \leq a < b < \infty$ , the sequence

25

 $\{g_n \mathbb{1}_{[a,b]}, n \in \mathbb{N}\}\$  converges in the weak topology of  $L_1([a,b])$  to  $g\mathbb{1}_{[a,b]}$ , i.e. we get, for all  $\varphi \in L_{\infty}([a,b])$ ,

$$\lim_{n\to\infty}\int_a^b g_n(t)\varphi(t)\mathrm{d}t = \int_a^b g(t)\varphi(t)\mathrm{d}t$$

*Proof.* We prove the result for d = 1, the extension for d > 1 is straightforward. Denote by  $h_n = \mathbb{1}_{[a,b]}g_n$ , and  $h = \mathbb{1}_{[a,b]}g$ . Under the stated assumptions, for each  $n \in \mathbb{N}$ ,  $h_n \in L_1([a,b])$  and the sequence  $\{h_n, n \in \mathbb{N}\}$  is uniformly integrable. Using the Dunford-Pettis theorem [12, Corollary 4.7.19], we conclude that for every subsequence  $\{h_n, n \in \mathbb{N}\}$  there exists a further subsequence  $\{h_{n_l}, l \in \mathbb{N}\}$  which converges in the weak topology of  $L_1[a,b]$ ) to  $h_*$ . We know that for any c such that  $a \le c \le b$  we have:

$$\int_{a}^{c} h_{*} dx = \lim_{l \to \infty} \int_{a}^{c} h_{n_{l}} dx = \lim_{l \to \infty} (f_{n_{l}}(c) - f_{n_{l}}(a)) = f(c) - f(a)$$

Hence  $h_*$  is a weak derivative of f (restricted to [a,b]). Since a weak derivative is unique up to a set of measure zero, we conclude that  $h_* = h$  in  $L_1([a,b])$ . Since from every subsequence of  $\{h_n, n \in \mathbb{N}\}$  we can choose a further subsequence converging weakly to h, the weak limit of  $\{h_n, n \in \mathbb{N}\}$  exists and is equal to h. This concludes the proof.

DEFINITION 6.7 (Convex combination subsequence). Let  $\{z_n, n \in \mathbb{N}\}$  be a sequence belonging to a linear subspace over  $\mathbb{R}$ . Let  $\{w_{n,k}, k \in \mathbb{N}, n \in \mathbb{N}\}$  be a sequence of weights satisfying:

- (*i*) For all  $k, n \in \mathbb{N}$ ,  $0 \le w_{n,k} \le 1$ .
- (ii) For all  $n \in \mathbb{N}$  we have  $\sum_{k=1}^{\infty} w_{n,k} = 1$ ,  $w_{n,k} > 0$  only for a finite number of indices k, and

$$\lim_{n \to \infty} \left( \inf\{k : w_{n,k} > 0\} \right) = \infty$$

The sequence  $\{z_n^w, n \in \mathbb{N}\}$  defined for each  $n \in \mathbb{N}$  by  $z_n^w = \sum_{k=1}^{\infty} w_{n,k} z_k$  is said to be a convex combination of  $\{z_n, n \in \mathbb{N}\}$  with weights  $\{w_{n,k}, k \in \mathbb{N}, n \in \mathbb{N}\}$ .

In the sequel, we will consider convex combinations of elements of  $\mathbb{R}^d$  and of the Banach space of integrable functions over some intervals [a,b] of  $\mathbb{R}$ ,  $L_1([a,b])$ .

LEMMA 6.8. Let  $0 \le a < b < \infty$ . Let  $\{f_n, n \in \mathbb{N}\}$  be a sequence of functions from [a,b] to  $\mathbb{R}^d$  which are integrable. Suppose the sequence  $\{f_n, n \in \mathbb{N}\}$  converges in the weak topology of  $L_1([a,b])$  to a limit f. Then there exists a convex combination subsequence of  $\{f_n, n \in \mathbb{N}\}$  that converges almost everywhere to f.

*Proof.* The space  $L_1([a,b])$  is a Banach space, so from Mazur's Lemma [15, Corollary 3.8] it follows that there exists a convex combination subsequence of  $\{f_n, n \in \mathbb{N}\}$  that converges strongly in  $L_1([a,b])$  to f, i.e. there exists a sequence of weights  $\{w_{n,k}, n, k \in \mathbb{N}\}$  satisfying the conditions of Definition 6.7 such that

$$\lim_{n\to\infty}\int_a^b |f_n^w(t)-f(t)|\mathrm{d}t=0\;.$$

A strongly convergent sequence in  $L_1([a,b])$  has an almost everywhere convergent subsequence. Since a subsequence of a convex combination subsequence is a convex combination subsequence that  $\{f_n, n \in \mathbb{N}\}$  has a convex combination subsequence that converges almost everywhere to f.

LEMMA 6.9. Let  $G : X \to \mathscr{P}(\mathbb{R}^d)$  be a set-valued function, define on a open subset  $X \subset \mathbb{R}^d$ . Assume that G is upper hemicontinuous and convex-closed valued. Let  $\{x_k, k \in \mathbb{N}\} \subset X$  be a sequence that converges to  $x \in X$ , and  $\{v_k, k \in \mathbb{N}\}$  be a sequence such that  $v_k \in G(x_k)$  for any  $k \in \mathbb{N}$ . Suppose a convex combination subsequence of  $\{v_k, k \in \mathbb{N}\}$  converges to a limit v. Then  $v \in G(x)$ .

*Proof.* For a closed set  $A \subseteq \mathbb{R}^d$  and  $z \in \mathbb{R}^d$  we denote by  $d(z,A) = \inf_{y \in A} ||z - y||$ . Let  $\{v_k^w, k \in \mathbb{N}\}$  be a convex combination subsequence of  $\{v_k, k \in \mathbb{N}\}$  with weights  $\{w_{n,k}, n, k \in \mathbb{N}\}$  (see Definition 6.7). Since *G* is upper hemicontinuity (see Definition 2.7), for any  $\varepsilon > 0$ , there exists an integer  $k_{\varepsilon}$ , such that for  $k \ge k_{\varepsilon}$ ,  $d(v_k, G(x)) \le \varepsilon$ . Since *G*(*x*) is convex, then function  $d(\cdot, G(x))$  is convex, and hence from Jensen's inequality we have  $d(v_n^w, G(x)) \le \varepsilon$  for *n* large enough. Hence for any  $\varepsilon > 0$  and *n* large enough we have  $d(v, G(x)) \le ||v - v_n^w|| + d(v_n^w, G(x)) \le 2\varepsilon$ .

### REFERENCES

- Z. ALLEN-ZHU AND E. HAZAN, Variance reduction for faster non-convex optimization, in International Conference on Machine Learning, 2016, pp. 699–707.
- [2] C. ANDRIEU AND E. MOULINES, On the ergodicity properties of some adaptive MCMC algorithms, Ann. Appl. Probab., 16 (2006), pp. 1462–1505.
- [3] Y. F. ATCHADÉ, G. FORT, AND E. MOULINES, On perturbed proximal gradient algorithms, J. Mach. Learn. Res., 18 (2017), pp. Paper No. 10, 33.
- [4] H. ATTOUCH AND J. BOLTE, On the convergence of the proximal algorithm for nonsmooth functions involving analytic features, Mathematical Programming, 116 (2007), pp. 5–16, https://doi.org/10.1007/ s10107-007-0133-5.
- [5] J.-P. AUBIN AND A. CELLINA, Differential inclusions : set-valued maps and viability theory, vol. 264 of Grundlehren der mathematischen Wissenschaften, Springer-Verlag, Berlin, Heidelberg, 1984.
- [6] A. BACCIOTTI AND F. CERAGIOLI, Stability and stabilization of discontinuous systems and nonsmooth Lyapunov functions, ESAIM: Control, Optimisation and Calculus of Variations, 4 (1999), pp. 361–376.
- [7] A. BECK AND M. TEBOULLE, A fast iterative shrinkage-thresholding algorithm for linear inverse problems, SIAM Journal on Imaging Sciences, 2 (2009), pp. 183–202.
- [8] M. BENAÏ M, J. HOFBAUER, AND S. SORIN, Stochastic approximations and differential inclusions. II. Applications, Math. Oper. Res., 31 (2006), pp. 673–695, https://doi.org/10.1287/moor.1060.0213, https://doi.org/10.1287/moor.1060.0213.
- [9] M. BENAÏ M, J. HOFBAUER, AND S. SORIN, Perturbations of set-valued dynamical systems, with applications to game theory, Dyn. Games Appl., 2 (2012), pp. 195–205, https://doi.org/10.1007/ s13235-012-0040-0, https://doi.org/10.1007/s13235-012-0040-0.
- [10] M. BENAÏM, Dynamics of stochastic approximation algorithms, in Seminaire de probabilites XXXIII, Springer, 1999, pp. 1–68.
- [11] M. BENAÏM, J. HOFBAUER, AND S. SORIN, Stochastic approximations and differential inclusions, SIAM Journal on Control and Optimization, 44 (2005), pp. 328–348.
- [12] V. I. BOGACHEV, Measure Theory, Vol I, Springer, 2006, http://www.ebook.de/de/product/5848086/vladimir\_ i\_bogachev\_measure\_theory.html.
- [13] J. BOLTE, S. SABACH, AND M. TEBOULLE, Proximal alternating linearized minimization for nonconvex and nonsmooth problems, Math. Program., 146 (2014), pp. 459–494, https://doi.org/10.1007/ s10107-013-0701-9, https://doi.org/10.1007/s10107-013-0701-9.
- [14] P. BREHENY AND J. HUANG, Coordinate descent algorithms for nonconvex penalized regression, with applications to biological feature selection, Ann. Appl. Stat., 5 (2011), pp. 232–253, https://doi.org/10.1214/ 10-AOAS388, https://doi.org/10.1214/10-AOAS388.
- [15] H. BREZIS, Functional Analysis, Sobolev Spaces and Partial Differential Equations, Springer-Verlag GmbH, 2010, http://www.ebook.de/de/product/9223754/haim\_brezis\_functional\_analysis\_sobolev\_spaces\_and\_ partial\_differential\_equations.html.
- [16] F. H. CLARKE, Optimization and nonsmooth analysis, SIAM, 1990.
- [17] F. H. CLARKE, Y. S. LEDYAEV, R. J. STERN, AND P. R. WOLENSKI, Nonsmooth Analysis and Control Theory, Springer-Verlag New York, Inc., Secaucus, NJ, USA, 1998.
- [18] P. L. COMBETTES AND J.-C. PESQUET, Stochastic approximations and perturbations in forward-backward splitting for monotone operators, Pure Appl. Funct. Anal., 1 (2016), pp. 13–37.
- [19] D. DAVIS, D. DRUSVYATSKIY, S. KAKADE, AND J. D. LEE, Stochastic subgradient method converges on tame functions, ArXiv e-prints, (2018), https://arxiv.org/abs/1804.07795.
- [20] J. FAN AND R. LI, Variable selection via nonconcave penalized likelihood and its oracle properties, J. Amer. Statist. Assoc., 96 (2001), pp. 1348–1360, https://doi.org/10.1198/016214501753382273, https://doi.org/ 10.1198/016214501753382273.
- [21] G. B. FOLLAND, *Real analysis*, Pure and Applied Mathematics (New York), John Wiley & Sons, Inc., New York, second ed., 1999.
- [22] G. FORT, E. MOULINES, AND P. PRIOURET, Convergence of adaptive and interacting Markov chain Monte

27

Carlo algorithms, Ann. Statist., 39 (2011), pp. 3262-3289.

- [23] S. GHADIMI AND G. LAN, Stochastic first- and zeroth-order methods for nonconvex stochastic programming, SIAM J. Optim., 23 (2013), pp. 2341–2368, https://doi.org/10.1137/120880811, https://doi.org/10.1137/ 120880811.
- [24] S. GHADIMI, G. LAN, AND H. ZHANG, Mini-batch stochastic approximation methods for nonconvex stochastic composite optimization, Math. Program., 155 (2016), pp. 267–305, https://doi.org/10.1007/ s10107-014-0846-1, https://doi.org/10.1007/s10107-014-0846-1.
- [25] P. HALL AND C. HEYDE, Martingale Limit Theory and its Application, Academic Press, 1980.
- [26] H. J. KUSHNER AND D. S. CLARK, Stochastic approximation methods for constrained and unconstrained systems, Applied mathematical sciences, Springer, New York, 1978.
- [27] H. J. KUSHNER AND G. YIN, Stochastic approximation and recursive algorithms and applications, Applications of mathematics, Springer, New York, 2003.
- [28] L. LJUNG, Analysis of recursive stochastic algorithms, IEEE Transactions On Automatic Control, 22 (1977), pp. 551–575, https://doi.org/10.1109/TAC.1977.1101561.
- [29] S. MEYN AND R. TWEEDIE, Markov chains and stochastic stability, Cambridge University Press, Cambridge, second ed., 2009. With a prologue by Peter W. Glynn.
- [30] B. MIASOJEDOW, E. MOULINES, AND M. VIHOLA, An adaptive parallel tempering algorithm, Journal of Computational and Graphical Statistics, 22 (2013), pp. 649–664, https://doi.org/10.1080/10618600. 2013.778779.
- [31] A. NEMIROVSKI, A. JUDITSKY, G. LAN, AND A. SHAPIRO, Robust stochastic approximation approach to stochastic programming, SIAM Journal on Optimization, 19 (2009), pp. 1574–1609.
- [32] A. NITANDA, Stochastic proximal gradient descent with acceleration techniques, in Advances in Neural Information Processing Systems, 2014, pp. 1574–1582.
- [33] N. PARIKH AND S. BOYD, Proximal algorithms, Foundations and Trends in Optimization, 1 (2014), pp. 127– 239.
- [34] S. J. REDDI, A. HEFNY, S. SRA, B. POCZOS, AND A. SMOLA, Stochastic variance reduction for nonconvex optimization, in International conference on machine learning, 2016, pp. 314–323.
- [35] H. ROBBINS AND S. MONRO, A stochastic approximation method, The Annals of Mathematical Statistics, (1951), pp. 400–407.
- [36] R. T. ROCKAFELLAR, R. J. B. WETS, AND R. J.-B. WETS, Variational Analysis, Springer, 2009, http:// www.ebook.de/de/product/2835226/r\_tyrrell\_rockafellar\_roger\_j\_b\_wets\_roger\_j\_b\_wets\_variational\_ analysis.html.
- [37] L. ROSASCO, S. VILLA, AND B. C. VŨ, Convergence of stochastic proximal gradient algorithm, arXiv preprint arXiv:1403.5074, (2014).
- [38] E. SAKSMAN AND M. VIHOLA, On the ergodicity of the adaptive Metropolis algorithm on unbounded domains, Ann. Appl. Probab., 20 (2010), pp. 2178–2203.
- [39] R. TIBSHIRANI, Regression shrinkage and selection via the lasso: a retrospective, J. R. Stat. Soc. Ser. B Stat. Methodol., 73 (2011), pp. 273–282, https://doi.org/10.1111/j.1467-9868.2011.00771.x, https://doi.org/ 10.1111/j.1467-9868.2011.00771.x.
- [40] L. XIAO AND T. ZHANG, A proximal stochastic gradient method with progressive variance reduction, SIAM J. Optim., 24 (2014), pp. 2057–2075, https://doi.org/10.1137/140961791, https://doi.org/10.1137/ 140961791.
- [41] C.-H. ZHANG, Nearly unbiased variable selection under minimax concave penalty, Ann. Statist., 38 (2010), pp. 894–942, https://doi.org/10.1214/09-AOS729, https://doi.org/10.1214/09-AOS729.

### Appendix A. Proof of Proposition 5.5.

LEMMA A.1. Under assumptions of Proposition 5.5, there exists  $C < \infty$  such that for any  $k \in \mathbb{N}^*$ ,

$$||x_k - x_{k-1}|| \le C \gamma_k W^{1/2}(z_k)$$

*Proof.* First consider  $\{x_k, k \in \mathbb{N}^*\}$  generated by (43). By (39) we get

$$|x_k - x_{k-1}|| \le \gamma_k \|\nabla f(x_{k-1})\| + \gamma_k \|\delta_k\| + \gamma_k \|u_k\|$$

where  $\delta_k = \Phi(x_{k-1}, z_k) - \nabla f(x_{k-1})$  and  $u_k \in \overline{\partial}g(x_k)$ . Since  $\{x_k, k \in \mathbb{N}^*\} \subset K$ , where *K* is compact, by (P1) we obtain  $\sup_{k \in \mathbb{N}^*} \{ \|u_k\| + \|\nabla f(x_{k-1}\|) < \infty$ . In addition by (B2), there exists  $\tilde{C} < \infty$  such that for any  $z \in \mathbb{Z}$ ,  $\sup_{k \in \mathbb{N}^*} \|\Phi(x_{k-1}, z)\| \leq \tilde{C}W^{1/2}(z)$  which concludes the proof.

Now consider  $\{x_k, k \in \mathbb{N}^*\}$  generated by (44). Denote by

$$y_k = \operatorname{prox}_{\gamma_k g} \left( x_{k-1} - \gamma_k \nabla f(x_{k-1}) - \gamma_k \delta_k \right) ,$$

where  $\delta_k = \Phi(x_{k-1}, z_k) - \nabla f(x_{k-1})$ . Using the definitions of the proximal operator (see (6)) and [16, Proposition 2.3.2], for each  $k \in \mathbb{N}^*$  there exists  $u_k \in \overline{\partial}g(y_k)$  such that

$$y_k = x_{k-1} - \gamma_k \nabla f(x_{k-1}) - \gamma_k \delta_k - \gamma_k u_k .$$

Since  $x_k$  is projection of  $y_k$  on set K, by the triangle inequality we get

$$||x_k - x_{k-1}|| \le ||y_k - x_{k-1}|| \le \gamma_k ||\nabla f(x_{k-1})|| + \gamma_k ||\delta_k|| + \gamma_k ||u_k|| .$$

Note that, since g is Lipschitz, [16, Proposition 2.1.2-(a)] shows that for all  $u \in \bar{\partial}g(y)$ ,  $||u|| \le ||g||_{\text{Lip}} < \infty$ . Boundedness of  $\nabla f$  on the set K and assumption (B2) concludes the proof.

*Proof of Proposition* 5.5. In this proof, *C* is a constant whose value may change upon each appearance. Observe that it is enough to show  $\sum_{k=1}^{\infty} \gamma_k \delta_k < \infty$  almost surely, where by construction

$$\delta_k = \Phi(x_{k-1}, z_k) - \nabla f(x_{k-1}) = \Phi(x_{k-1}, z_k) - \pi_{x_{k-1}}(\Phi(x_{k-1}, \cdot)) .$$

Geometric ergodicity (B1) in turn implies the existence of a solution of the Poisson equation, and also provide bounds on the growth of this solution; see [3, Lemma 13]. For  $z \in Z$ , we set  $\Phi_x(z) = \Phi(x, z)$ . For any  $x \in K$  there exists a solution  $\hat{\Phi}_x$  to the Poisson equation

$$\hat{\Phi}_x - M_x \hat{\Phi}_x = \Phi_x - \pi_x (\Phi_x)$$

and there exists a constant  $C < \infty$  such that for any  $x \in K$  and  $z \in Z$ 

(52) 
$$\left| \hat{\Phi}_x(z) \right|_{W^{1/2}} \le C W^{1/2}(z) \text{ and } \left| M_x \hat{\Phi}_x(z) \right|_{W^{1/2}} \le C W^{1/2}(z) .$$

Hence, for any  $k \in \mathbb{N}^*$  we can decompose  $\delta_k = \delta M_k + \kappa_k$  where

$$\begin{split} \delta M_k &:= \hat{\Phi}_{x_{k-1}}(z_k) - M_{x_{k-1}} \hat{\Phi}_{x_{k-1}}(z_{k-1}) \\ \kappa_k &:= M_{x_{k-1}} \hat{\Phi}_{x_{k-1}}(z_{k-1}) - M_{x_{k-1}} \hat{\Phi}_{x_{k-1}}(z_k). \end{split}$$

By construction, the sequence  $\{\delta M_k, k \in \mathbb{N}\}$  is a martingale increment sequence and Doob's inequality implies that there exists a constant  $C < \infty$  such that for all  $z_0 \in \mathbb{Z}$ ,

(53) 
$$\mathbb{E}\left[\left(\sup_{k>n}\left|\sum_{l=n}^{k}\gamma_{l}\delta M_{l}\right|\right)^{2}\right] \leq C\sum_{l=n}^{\infty}\gamma_{l}^{2}\mathbb{E}\left[\left\|\hat{\Phi}_{x_{l-1}}(z_{l})-M_{x_{l-1}}\hat{\Phi}_{x_{l-1}}(z_{l-1})\right\|^{2}\right].$$

By construction the sequence  $\{x_k, k \in \mathbb{N}\}$  remains in the compact set K. Using (52), we obtain that,

$$\mathbb{E}\left[\left(\sup_{k>n}\left|\sum_{l=n}^{k}\gamma_{l}\delta M_{l}\right|\right)^{2}\right] \leq C\sup_{k\geq 1}\mathbb{E}[W(z_{k})]\sum_{l=n}^{\infty}\gamma_{l}^{2}.$$

Geometric ergodicity together with  $\mathbb{E}W(z_1) < \infty$  implies that (see. [30, Lemma 21] or [3, Lemma 14])

$$\sup_{k} \mathbb{E}W(z_k) < \infty.$$

Therefore, since  $\sum_{k=1}^{\infty} \gamma_k^2 < \infty$  we conclude that  $\sum_k \gamma_k \delta M_k$  converges almost surely.

Decompose  $\sum_{l=n}^{k} \gamma_l \kappa_l = R_{n,k}^1 + R_{n,k}^2 + R_{n,k}^3$  with  $R_{n,k}^{1} := \sum_{l=k-1}^{n-1} \gamma_{l+1} \left[ M_{x_{l}} \hat{\Phi}_{x_{l}}(z_{l}) - M_{x_{l-1}} \hat{\Phi}_{x_{l}}(z_{l}) \right]$  $R_{n,k}^2 := \gamma_{n-1} M_{x_{n-2}} \hat{\Phi}_{x_{n-2}}(z_{n-1}) - \gamma_k M_{x_{k-1}} \hat{\Phi}_{x_{k-1}}(z_k)$  $R_{n,k}^3 := \sum_{l=n-1}^{k-1} (\gamma_{l+1} - \gamma_l) M_{x_{l-1}} \hat{\Phi}_{x_{l-1}}(z_l)$ 

Applying [22, Lemma 4.2] we get that

$$|R_{n,k}^{1}| \leq C \sum_{l=k-1}^{n-1} \gamma_{l+1} W^{1/2}(z_{l}) \left[ D_{W^{1/2}}(x_{l}, x_{l-1}) + \|\Phi(x_{l}, \cdot) - \Phi(x_{l-1}, \cdot)\|_{W^{1/2}} \right]$$

By assumption (B2) and Lemma A.1 we obtain  $\sup_{k} [\mathbb{E}[R_{n,k}^{1}]] \leq C \sup_{k} \mathbb{E}[W(z_{k})] \sum_{l>n} \gamma_{l}^{2}$  and

this converges to zero by  $\sum_{k=1}^{\infty} \gamma_k^2 < \infty$ . Finally, from (52), (P2) and  $\sum_k |\gamma_k - \gamma_{k-1}| < \infty$  we deduce that  $R_{n,k}^2$  and  $R_{n,k}^3$  also converges to zero almost surely and that completes the proof.