

Gauge enhancement of super M-branes via parametrized stable homotopy theory

Vincent Braunack-Mayer*, Hisham Sati†, Urs Schreiber‡

March 5, 2019

Abstract

A key open problem in M-theory is to explain the mechanism of “gauge enhancement” through which M-branes exhibit the nonabelian gauge degrees of freedom seen perturbatively in the limit of 10d string theory. In fact, since only the twisted K-theory classes represented by nonabelian Chan–Paton gauge fields on D-branes have an invariant meaning, the problem is really the understanding the M-theory lift of the classification of D-brane charges by twisted K-theory.

Here we show that this problem has a solution by universal constructions in rational super homotopy theory. We recall how double dimensional reduction of super M-brane charges is described by the cyclification adjunction applied to the 4-sphere, and how M-theory degrees of freedom hidden at ADE singularities are induced by the suspended Hopf action on the 4-sphere. Combining these, we demonstrate that, in the approximation of rational homotopy theory, gauge enhancement in M-theory is exhibited by lifting against the fiberwise stabilization of the unit of this cyclification adjunction on the A-type orbispace of the 4-sphere. This explains how the fundamental D6 and D8 brane cocycles can be lifted from twisted K-theory to a cohomology theory for M-brane charge, at least rationally.

Contents

1	Introduction	2
2	Two universal constructions in homotopy theory	10
2.1	Fiberwise stabilization	10
2.2	The Ext/Cyc-adjunction	19
3	The A-type orbispace of the 4-sphere	26
3.1	Rational homotopy type of the A-type orbispace	29
3.2	Fiberwise stabilized Ext/Cyc-unit of the A-type orbispace	32
4	Gauge enhancement of M-branes	40
4.1	The mechanism without supersymmetry	40
4.2	The mechanism with local supersymmetry	44


*Institute of Mathematics, University of Zurich, Winterthurerstrasse 190, CH-8057 Zürich, Switzerland.

†Division of Science and Mathematics, New York University, Abu Dhabi, UAE.

‡Division of Science and Mathematics, New York University, Abu Dhabi, UAE, on leave from Czech Academy of Science.

1 Introduction

The conjecture of *M-theory* [To95a, Wi95] (see [Du99, BBS06]) says, roughly, that there exists a non-perturbative physical theory, which makes the following schematic diagram commute:




Here both *perturbative string theory* as well as *higher-dimensional quantum supergravity* may, with some effort, be well-defined as *perturbative S-matrix theories* (e.g. [Po01, Sec. 12.5][Wi12] and [Do95][BFR13]); and the conjecture is that there is a joint *non-perturbative* completion of 11-dimensional supergravity and of IIA string theory, hence of any effective quantum field theory approximating the latter. Even though the actual nature of M-theory has remained an open problem [Mo14, Sec. 12]¹, there is a huge and steadily growing network of hints supporting the M-theory conjecture. This should be regarded in light of the situation of perturbative quantum field theory as used in the Standard Model of particle physics, where the identification of any aspect of its non-perturbative completion is a crucial but wide-open problem, referred to as one of the “millennium problems” [ClayMP].

In a series of articles [FSS13, FSS16a, FSS16b, HS17], we have shown that a systematic analysis of the *Green–Schwarz sigma-models* (which define *fundamental* super p -branes, such as the fundamental membrane that gives M-theory its name) from the point of view of *super homotopy theory* provides a concrete handle on some previously elusive aspects of M-theory; see [Sc17c, FSS19] for exposition of this perspective. Specifically, in the companion article [HSS18] it is shown that the existence and classification of *black M-branes at real ADE singularities* can be systematically derived and analyzed in the supergeometric enhancement of *equivariant homotopy theory* (see [B17]). However, the M-theory folklore suggests ([Se97, Sec. 2], see e.g. [IU12, Sec. 6.3.3], also [AG04]) that understanding black branes at ADE-singularities also holds the key to the all-important, widely expected yet still mysterious phenomenon of *gauge enhancement* in M-theory. This suggests that the super homotopy theoretic analysis may also shed light on the true nature of gauge enhancement in M-theory. Here we show that this is indeed the case.

In the remainder of this introduction we review the issue of gauge enhancement in various guises, survey what is known, what is conjectured, and which problems remain essentially unsolved. After establishing some mathematical results in Sec. 2 and Sec. 3, we explain our solution to the gauge enhancement problem in Sec. 4.


Double dimensional reduction. At the heart of the matter is *double dimensional reduction*, originally due to [DHI⁺87, To95b] and whose rigorous formulation from [FSS16a, Sec. 3] [FSS16b, Sec. 3] we recall and expand upon below in Sec. 4. Going back to ideas of Kaluza and Klein a century ago, in ordinary *dimensional reduction*, we take spacetime to be a pseudo-Riemannian S^1 -fibration over a $(D - 1)$ -dimensional base space and consider the limit in which the circle fiber becomes infinitesimal. In this limit, we obtain a field theory on the $(D - 1)$ -dimensional base space of the circle fibration, hence in a lower-dimensional spacetime, which has a larger space of field species including the Fourier modes of the original fields on the circle fiber.

¹At least in mathematics it is not uncommon that a theory is conjectured to exist before its actual nature is known—famous examples of this include the theory of *motives*, which has meanwhile been discovered, and the *field with one element*.



Secondly, one finds higher-dimensional analogs of black holes in higher-dimensional supergravity, with $(p + 1)$ -dimensional singularities, called *black* (or “solitonic”) *p-branes* [DIP⁺88]. Under dimensional reduction of the ambient supergravity theory, the singularity of a black *p*-brane may or may not extend along the circle fiber that is being shrunk away. If it does not, then the result is again a black *p*-brane solution, now in the lower-dimensional supergravity theory. But if it does, then along with reduction in spacetime dimension from D to $(D - 1)$, the black *p*-brane singularity effectively appears as a black $(p - 1)$ -brane solution in the lower-dimensional supergravity theory; whence “double dimensional reduction”.

Chan–Paton gauge enhancement. In its most immediate (albeit naive) form, the formulation of the problem of *gauge enhancement* in M-theory proceeds from the expected double dimensional reduction (2) of the black branes of M-theory (see [HSS18] for a precise account), to those of type IIA string theory, which for black branes looks as follows [To95a, To95b, FOS02a, FOS02b]:



From perturbative string theory one finds that open fundamental strings ending on the D-branes behave as quanta for an abelian $U(1)$ -gauge theory (i.e. electromagnetism) on the worldvolume of the D-brane. A widely accepted but informal² argument [Wi96, Sec. 3] indicates that if N such D-branes are *coincident* then the gauge group *enhances* from the abelian group $(U(1))^N$ to the non-abelian group $U(N)$. The idea is that massless open fundamental strings stretch in $N \times N$ possible ways between the N coincident D-branes, thus constituting gauge bosons that organize in $N \times N$ unitary matrices, called “Chan–Paton factors”. However, it is non-trivial to check that scattering of these open strings reproduces the scattering amplitudes of gauge bosons for non-abelian gauge theory (Yang–Mills theory). First approximate numerical checks of this idea are due to [CST03] and similar numerical checks as well as an exact derivation under simplifying assumptions are given in [BS03]; a full derivation was claimed in [Le17].

This phenomenon of *gauge enhancement on D-branes* is of paramount importance for string theory, in particular as a candidate for a theory of realistic physics. The fundamental gauge fields observed in nature, per the Standard Model of particle physics, do of course involve non-abelian gauge groups corresponding to the weak and strong nuclear forces. While Kaluza–Klein dimensional reduction from 11-dimensional supergravity may exhibit such non-abelian gauge groups, this happens in a manner incompatible with realistic coupling of fermionic matter fields [Wi81]. Instead, realistic gauge fields must arise from gauge enhancement on coincident D-branes (see [IU12, Sec. 10]). Moreover, all discussion of modern string theoretic topics such as *AdS/CFT duality* (see [AGM⁺99]) or *geometric engineering of gauge theories* (going back to [HW97], see e.g. [Fa17]), such as for classification of 6d SCFTs (as in [DZH⁺15]), depend crucially on gauge enhancement on coincident D-branes. But, under the M-theory conjecture, double

² In [Wi96, first line on p. 8] the argument was introduced as an “obvious guess”. Most subsequent references cite this as a fact, e.g. the review [My03, Sec. 3], despite the lack of a formal argument.


dimensional reduction should exhibit an equivalence (“duality”) between (strongly coupled) type IIA string theory and M-theory, in particular between the full non-perturbative theory of D-branes and their M-brane pre-images. Hence if M-theory exists, then gauge enhancement on coincident D-branes must correspond to, and is potentially explained by, a corresponding phenomenon on M-branes. The most immediate incarnation of the problem of gauge enhancement in M-theory can, therefore, be succinctly phrased as:

Open Problem, version 1:

What is the lift to M-theory of the non-abelian Chan–Paton gauge field degrees of freedom on coincident D-branes?

Since the string theory literature tends to blur the distinction between what is known and what is conjectured, we briefly highlight what the folklore on this problem (e.g. [IU12, Sec. 6.3.3], [AG04]) does and does not achieve:

- A celebrated recent result (see [BLM⁺13] for a review) shows the existence of a class of non-abelian gauge field theories that are plausible candidates for the worldvolume theories expected to live on black M2-branes sitting at ADE singularities (the “BLG-model” [BL08, Gus09] and, more generally, the “ABJM-model” [ABJM08]). However, a derivation of these field theories from M-theoretic degrees of freedom is missing; the argument works just by consistency checks.
- On the other hand, a conjectural sketch of an explicit derivation does exist for the M-brane species MK6, whose image in the low-energy approximation provided by 11-dimensional supergravity is supposed to be the *Kaluza–Klein monopole spacetime* and which becomes the black D6-brane under dimensional reduction [To95a], [Se97, Sec. 1]. For an *abelian* gauge field on the D6-brane, a straightforward analysis shows that it is sourced by doubly dimensionally reduced M2-branes ending on the KK-monopole [GM03, Sec. 2], [Ma04, Sec. 5.4]. But more generally, KK monopoles may be argued to be the fixed point loci of spacetime orbifolds locally of the form $\mathbb{R}^{6,1} \times \mathbb{C}^2 // G_{\text{ADE}}$ for a finite subgroup $G_{\text{ADE}} \subset SU(2)$. A classical theorem [duV34] (see [Re87] for a review) implies that such singularities are canonically resolved by spheres touching along the shape of a simply-laced Dynkin diagram:



If one here imagines that M2-branes wrap these *vanishing 2-cycles* then, under double dimensional reduction, this situation looks like an M-theoretic lift of the strings stretching between several coincident D6-branes, as indicated in the figure above, and hence looks like an M-theoretic lift of the Chan–Paton gauge enhancement mechanism discussed above.

This argument goes back to [Se97, Sec. 2]; for review see [IU12, Sec. 6.3.3]. While this story is appealing, it is unsatisfactory that it has to treat the membrane in 11-dimensional spacetime as a direct analogue of the fundamental string in 10-dimensional spacetime; after all, the very term “M-theory” instead of “Membrane theory” was chosen as a reminder that this direct analogy is too naive.³

In short, the traditional attempts to understand gauge enhancement on M-branes suffer from the lack of any handle on the actual nature of M-theory. But what is worse, these stories argue a point that has meanwhile come to be thought as being inaccurate:

K-Theoretic gauge enhancement. A well-known series of arguments [MM97, Wi98, FW99, FH00, MW00] shows that the gauge fields carried by D-branes do not actually have well-defined existence in


³[HW06]: “As it has been proposed that [this] theory is a supermembrane theory but there are some reasons to doubt that interpretation, we will non-committedly call it the M-theory, leaving to the future the relation of M to membranes.”

themselves. Instead, one must view all the Dp -branes taken together and then their gauge fields serve as representatives for classes in *twisted K-theory* [BM00, Wi00, Fr00, BCM⁺02, BEV03, MS04, ES06, Ev06]. It is only these twisted K-theory classes that are supposed to have an intrinsic meaning. In other words, non-abelian gauge fields on separate D-brane species are much like coordinate charts on spacetime: a convenient but non-invariant means of presenting a specific structure. In actual reality, the D-brane species Dp , $p \in \{0, 2, 4, 6, 8\}$ only have a unified joint existence and gauge enhancement to on separate D-brane species is only one presentation, out of many, of a unified *higher gauge field*: a cocycle in twisted K-theory. In view of this, the problem of gauge enhancement in M-theory is really the following:⁴

Open Problem, version 2:

What is the cohomology theory classifying M-brane charge, and how does double dimensional reduction reduce it to the classification of D-brane charge in twisted K-theory?

Hence the refined perspective of twisted K-theory shifts the focus away from Chan–Paton-like gauge enhancement on one particular D-brane species (which seems to have no invariant meaning in the full theory) and instead highlights the problem of how the full list of D-brane species arises and carries a unified charge in twisted K-theory. However, this is in conflict with the M-theoretic origin of the D-branes in the traditional story recalled above: only the M2-brane and M5-brane exist as *fundamental branes*, meaning that they have corresponding Green–Schwarz type sigma-models (we recall this in detail below in Sec. 4.2). The double dimensional reduction of these yields the fundamental brane species in type IIA string theory, *except* the D6 and the D8 (while the D0 now encodes the circle fibration itself):




On the other hand, up to now the M-theoretic origin of the D6-brane has only been argued in its *black brane* incarnation, which is supposed to be given by the MK6 as recalled above. The nature of the D8-brane is yet more subtle [BRG⁺96]—an M-theory lift has been proposed in [Hu98], but the proposal is not among the usual list of expected black M-branes. On the other hand, the reduction of the MO9-brane, whose existence in M-theory is solid (see [HSS18]), is not the black D8-brane, but rather the O8-plane (e.g. [GKS⁺01, Sec. 3]). This highlights that the core of the open problem, from the refined K-theoretic perspective, is really in the appearance of the D6- and D8-brane:

Open Problem, version 3:

What is the lift to M-theory of the fundamental D6-branes and D8-branes in type IIA string theory, such that the unified Dp -branes are jointly classified by twisted K-theory?

K-Theoretic gauge enhancement in rational approximation. Twisted K-theory is a comparatively complicated structure, and the fine detail of which of its various variants really applies to D-branes is still the subject of discussion ([KS05, DFM09, Sa10, GS19]). Of course the glaring problem here is, once more, that the non-perturbative theory that ought to answer this question is missing. It is worth highlighting that the issue of gauge enhancement is visible, and has remained unresolved, already in the *rational* approximation (i.e. ignoring all torsion-group effects), where a cocycle in twisted K-theory reduces to a cocycle in *twisted de Rham cohomology*. Associated to each brane species is a differential form on spacetime—its *flux form*—which corresponds to the brane in analogy to the correspondence between the *Faraday tensor* and charged particles (0-brane) in electromagnetism. The double dimensional reduction of these flux forms is as follows (see [MS04, Sec 4.2]), parallel to the pattern of the double dimensional reduction of the fundamental branes.

⁴ While a *derivation of K-theory from M-theory* is suggested by the title of [DMW03], that article only checks that the behavior of the partition function of the 11d supergravity C -field is compatible with the *a priori* K-theory classification of D-branes. Seeking a generalized cohomology describing the M-field and M-branes was originally advocated for in [Sa05a, Sa05b, Sa06, Sa10].



Here the right hand sides have been recognized in [Sa13, Sec 2.5] [FSS15b] for the top part and derived in [FSS16a, FSS16b] for the bottom part; we discuss this in detail below in Sec. 4.1. Notice that the F_2 -contribution does arise from plain double dimensional reduction, not directly from the (G_4, G_7) , but as a rational image of the first Chern-class of the circle fibration itself. Moreover, these differential forms satisfy relations (“twisted Bianchi identities”) that identify them, via the de Rham theorem, as cocycles in the rationalization of a generalized cohomology theory:

Flux forms	Twisted Bianchi identity	Rational cocycle in
M-theory	$dG_4 = 0$ $dG_7 = -\frac{1}{2}G_4 \wedge G_4$	cohomotopy in degree 4
Type IIA string theory	$dH_3 = 0$ $dF_{2p+4} = H_3 \wedge F_{2p+2}$	twisted K-theory in even degree

In this rational approximation, the core of the gauge enhancement problem is still fully visible:


Open Problem, rational version 1:

What is the origin of the RR-flux forms F_8 and F_{10} in M-theory, such that these unify with the double dimensional reduction of the M-flux (G_4, G_7) to an un-truncated cocycle in rational twisted K-theory (i.e. in twisted de Rham cohomology)?

We present a solution to this version of the problem in Sec. 4.1. Though working in the rationalized setting means that we are disregarding all torsion⁵ information for the time being, it has the striking advantage that these rationalized relations follow rigorously from a *first principles*-definition of M-branes (recalled below in Sec. 4.2), and hence serve as a starting point for a systematic analysis of the problem of gauge enhancement.

To properly take local supersymmetry into account, one has to consider the refinement of the plain flux forms to super-flux forms on super-spacetime. The torsion-freeness constraints of supergravity geometrically require [Lo90, EE12] the bifermionic component of these super-flux forms to be covariantly constant on each super tangent space (see [FSS18, Sec. 1]), where they correspond to those cocycles μ_{p+2} in the supersymmetry super Lie algebra cohomology defining the Green–Schwarz-type sigma-models for the fundamental p -branes [AETW87, AT89]. This identification locates the problem in the precise context of *super homotopy theory* of super-spacetimes. The beauty of this is that homotopy theory is governed by *universal constructions*, which, roughly, means that it exhibits the emergence of “god-given” structures from a minimum of input. In fact, in super homotopy theory the super-cocycles μ_{M_2} and μ_{M_5} witnessing the fundamental M-branes emerge by a universal construction (a kind of equivariant Whitehead tower) from nothing but the superpoint [FSS13, HS17], and their double dimensional reduction is reflected by another universal construction [FSS16a, Sec. 3] [FSS16b, Sec. 3]—the *Ext/Cyc-adjunction* (discussed in detail in Sec. 2.2 below).

⁵This torsion is in the sense of cohomology or homotopy classes. In the following paragraph we use torsion in the sense of differential (super)geometry. We hope that the distinction will be clear from the context.



Fundamental brane super-cocycle	Cocycle condition	in rational cohomology theory
M-theory	$d\mu_{M2} = 0$ $d\mu_{M5} = -\frac{1}{2}\mu_{M2} \wedge \mu_{M2}$	cohomology in degree 4
Type IIA string theory	$d\mu_{F1} = 0$ $d\mu_{2p+2} = \mu_{F1} \wedge \mu_{2p}$	twisted K-theory in even degree

Here (rational) *cohomotopy* in degree 4 is the generalized *non-abelian* cohomology theory represented by the (rationalized) 4-sphere, meaning that the joint M2/M5-brane cocycle is a morphism in the rational super homotopy category of the form [FSS15b, FSS16a]

$$\mathbb{R}^{10,1|\mathbf{32}} \xrightarrow{\mu_{M2/M5}} S^4 \in \text{Ho}(\text{SuperSpaces}_{\mathbb{R}}).$$

That cohomotopy governs the M-brane charges this way, at least rationally, was first proposed and highlighted in [Sa13, Sec. 2.5]. *A priori* there are many homotopy types that look like the 4-sphere in the rational approximation; however in [HSS18] further precise evidence was provided to demonstrate the sense in which the 4-sphere is the correct coefficient for M-brane charge. Namely, the 4-sphere coefficient is naturally identified with the 4-sphere around a black M5-brane singularity in $D = 11$ supergravity and this identification induces a real structure on the 4-sphere together with actions of the ADE subgroups of $SU(2)$ that are compatible with the corresponding BPS actions on super-spacetime. It is therefore natural to ask for enhancements of the M2/M5-brane cocycle to *equivariant* cohomotopy

$$\begin{array}{ccc} G_{\text{ADE}} \times G_{\text{HW}} & & G_{\text{ADE}} \times G_{\text{HW}} \\ \curvearrowright & & \curvearrowright \\ \mathbb{R}^{10,1|\mathbf{32}} & \xrightarrow{\hat{\mu}_{M2/M5}} & S^4 \end{array} \in \text{Ho}((G_{\text{ADE}} \times G_{\text{HW}})\text{-SuperSpaces}_{\mathbb{R}}).$$

Here $G_{\text{ADE}} \subset SU(2)$ is a finite subgroup as per the ADE-classification that acts by orientation-preserving super-spacetime automorphisms, while $G_{\text{HW}} = \mathbb{Z}_2$ is an orientation-reversing reflection as in Hořava–Witten theory.

It is shown in [HSS18] that such an equivariant enhancement exists and makes the *black branes* at ADE singularities appear. This results in a unified framework for black and fundamental M-branes. In particular, the corresponding A-series actions on the 4-sphere factor through the $U(1)$ -action

$$\begin{array}{ccc} S^1 & & U(1) & & SU(2) \\ \curvearrowright & & \curvearrowright & & \curvearrowright \\ S^4 & := & S(\mathbb{R} \oplus \mathbb{C}^2) & \subset & S(\mathbb{R} \oplus \mathbb{C}^2) \end{array}$$

obtained as the suspension of the circle action on the complex Hopf fibration $H_C: S^3 \rightarrow S^2$. The projection to the corresponding homotopy quotient is identified with the M-theory S^1 -fibration in the near horizon geometry of an M5-brane:

$$\begin{array}{ccccc}
& & \boxed{\text{M2/M5-brane cocycle}} & & \\
& & \mathbb{R}^{10,1|\mathbf{32}} \xrightarrow{\mu_{\text{M2/M5}}} S^4 & & \boxed{\text{4-sphere coefficient}} \\
\boxed{\text{11d super-spacetime}} & & \downarrow & & \downarrow \\
& & \mathbb{R}^{9,1|\mathbf{16}+\overline{\mathbf{16}}} \simeq \mathbb{R}^{10,1|\mathbf{32}} // S^1 & & S^4 // S^1 \\
\boxed{\text{10d super-spacetime}} & & & & \boxed{\text{A-type orbispace of the 4-sphere}}
\end{array} \tag{5}$$

In conclusion, the problem of gauge enhancement in M-theory in the rational approximation, but otherwise proceeding from *first principles*, reads as follows:

Open Problem, rational version 2:

Which universal construction in rational super homotopy theory enhances the cyclification of the M2/M5-cocycle from 6-truncated to un-truncated rational twisted K-theory?


Gauge enhancement explained. It is this version of the problem to which we present a solution. First we explain and analyze two relevant universal constructions in homotopy theory:

- (i) *Fiberwise stabilization* (in Sec. 2.1) and
- (ii) *the Ext/Cyc-adjunction* (in Sec. 2.2).

We then consider (in Sec. 3.1) the rational homotopy type of the *A-type orbispace of the 4-sphere*, as in (5) above, and we apply the two aforementioned universal constructions to it (in Sec. 3.2). Our main result, Theorem 3.19, shows that rational untruncated twisted K-theory appears as a direct summand in the fiberwise stabilization of the Ext/Cyc-unit on the A-type orbispace of the 4-sphere.


Since homotopy theory is immensely rich and computationally demanding (see. e.g. [Ra03, HHR09]) one often simplifies calculations by working in successive approximations, such as in the filtration by *chromatic layers*. The first of these approximations is *rational homotopy theory* (e.g. [He06]) obtained by disregarding all torsion elements in homotopy and cohomology groups. The model for *rational parametrized stable homotopy theory* that we use in our computations had been conjectured in [FSS16a, p. 20] and was subsequently worked out in [BM18]. This model allows us to reveal a deeper meaning behind a curious dg-algebraic observation due to [RS00] (recalled as Prop. 3.13 below), culminating in our main Theorem 3.19, below. However, since the gauge enhancement mechanism that we present is obtained by *universal constructions* (specifically the *derived adjunctions* discussed in Sec. 2), the lift of the mechanism beyond the rational approximation certainly exists, but is just much harder to analyze.

Finally, in Sec. 4.2 we apply Theorem 3.19 to the double dimensional reduction of the fundamental M-brane cocycles. We show that this solves **Open Problem, rational version 2** by making the D6- and D8-brane cocycles appear and by exhibiting a single unified super-cocycle in rational un-truncated twisted K-theory:



Notice how all the folkloric ingredients recalled above do appear in this rigorous result, albeit in a somewhat subtle way. First of all, the fact that fundamental branes and black branes are closely related, while still

crucially different (particularly in the matter of gauge enhancement), is reflected by how super-cocycles interact with spacetime ADE singularities in the data specifying a real ADE-equivariant cohomotopy class [HSS18]. Second, the claim that gauge enhancement in M-theory is connected to the appearance of ADE singularities in spacetime is reflected here in the fact that the untruncated rational twisted K-theory spectrum (which, as we have discussed, is the true rational coefficient for the gauge enhanced brane charges), only appears from fiberwise stabilization of the *equivariant* 4-sphere coefficient. This equivariant coefficient also induces the appearance of singular fixed point strata in spacetime via equivariant enhancement:



Of course it remains to lift our result beyond rational homotopy theory. However, we suggest that the rational derivation of gauge enhancement in Theorem 3.19 and Sec. 4.2 points to its own non-rational refinement. The reason is that the universal constructions that we have used also make sense non-rationally – they are just much harder to compute. More concretely, we observe that the manner in which the rational version of twisted K-theory appears below is via a twisted, rational version of *Snaith’s theorem* (see Rem. 3.14). This theorem says that the K-theory spectrum KU is obtained from the suspension spectrum $\Sigma_+^\infty BS^1$ of the classifying space BS^1 by adjoining a multiplicative inverse of the Bott generator β :

$$\Sigma_+^\infty BS^1[\beta^{-1}] \simeq_{\text{swhe}} KU.$$

Rationally, Snaith’s theorem is rather immediate, as is its rational twisted version (Ex. 2.28 below) that underlies our identification of rational twisted K-theory in Theorem 3.19. Since rationalization is the coarsest non-trivial approximation to full homotopy theory (and in this regard quite similar to taking the first derivative of a non-linear function at a single point) it loses plenty of information.

A priori, what looks like twisted K-theory in the rational approximation could correspond non-rationally to different twisted cohomology theories (see [Sa10, GS17, GS19] for discussions in this context). However, we do not just see rational twisted K-theory in isolation, but rather appearing after applying universal constructions to the A-type orbispace of the 4-sphere. For our main theorem on gauge enhancement, these universal constructions are what really matter. Therefore, any non-rational lift of our gauge enhancement mechanism should arise by the same universal construction, applied non-rationally, possibly in conjunction with other universal constructions that are rationally invisible. This considerably constrains the possibilities. The conclusion we draw is that (fiberwise) inversion of the Bott generator is a good candidate for lifting our gauge enhancement mechanism beyond the rational approximation.

It is quite plausible that the gauge enhancement mechanism presented in Sec. 4.2 generalizes beyond the rational approximation to a derivation of full twisted K-theory from degree 4 cohomotopy. We may state this as the remaining part of the problem of gauge enhancement in M-theory:

Open Problem, remaining part:

What is the non-rational lift of the gauge enhancement mechanism, by universal constructions in super homotopy theory, from Sec. 4.2?

We will return to this open problem elsewhere.


2 Two universal constructions in homotopy theory

Here we discuss two universal constructions in homotopy theory (see e.g. [Sc17a, Sc17b]), which when applied to the A-type orbispace of the 4-sphere (Sec. 3) reveal the mechanism of gauge enhancement on M-branes (Sec. 4). Firstly, in Sec. 2.1 we recall *parametrized stable homotopy theory* and recall an algebraic model for its rationalization from [BM18, BM19b], which enables us to effectively compute in this setting. The main results of this section are Theorem 2.19, which establishes differential-graded modules as rational models for parametrized spectra, and Prop. 2.21 characterizing the fiberwise stabilization adjunction in terms of these models. In Sec. 2.2 we demonstrate that forming cyclic loop spaces is one part of a homotopy-theoretic adjunction, and we characterize the unit of the adjunction (Theorem 2.44). This universal construction is used in our formulation of double dimensional reduction.


This section may be read independently of the rest of the article and is of interest beside its application to M-brane phenomena. Conversely, readers interested only in the application to M-theory and willing to accept our homotopy-theoretic machinery as a black box may be inclined to skip this section.

2.1 Fiberwise stabilization

In [FSS16a, p. 20], we found that the super L_∞ -algebraic F1/Dp-brane cocycles organize into a diagram as shown in (b) below. We further indicated that this ought to be thought of as the image in supergeometric rational homotopy theory of a cocycle in twisted K-theory realized as a morphism of parametrized spectra, as shown in (a) below.



(a) A twisted K-theory cocycle according to [ABG10, NSS12].



(b) The descended IIA F1/Dp-brane cocycle according to [FSS16a, Theorem 4.16].

Roughly, a *spectrum* is a kind of *linearized* or *abelianized* version of a topological space. By the classical Brown Representability Theorem, maps into spectra represent cocycles in generalized cohomology theories, such as K-theory. A *parametrized spectrum* is a family of spectra that is parametrized in a homotopy-coherent manner by a topological space (see [MS06]). Roughly, these are equivalent to bundles of spectra over the parameter space, where maps into the total space of such a bundle represent cocycles in a *twisted* generalized cohomology theory.

$$\text{Spectra} \hookrightarrow \begin{array}{c} \xrightarrow{\text{Parametrized}} \\ \text{over the point} \end{array} \text{Parametrized spectra} \xrightarrow{\begin{array}{c} \text{Underlying} \\ \text{parameter space} \end{array}} \text{Spaces}$$

Under Koszul duality, the conjecture of [FSS16a] means, roughly, that there ought to be highlighted entries as in the following table. These entries unify Quillen–Sullivan’s DG-models for rational homotopy theory of topological spaces (the central result is recalled as Prop. 2.11 below) with chain complex models for stable rational homotopy theory (recalled as Prop. 2.17 below).

Homotopy theory	Stable	Parametrized stable	Plain
Plain	Spectra	Parametrized spectra	Spaces
Rational	Cochain complexes	<i>DG-modules</i>	DG-algebras
Super rational	Super cochain complexes	<i>Super DG-modules</i>	Super DG-algebras (“FDA”s)

This conjecture has been proven recently in [BM18] (see also the forthcoming articles [BM19a, BM19b]). In this paper we review those parts of the resulting *rational parametrized stable homotopy theory* that we

need for the proof of Theorem 3.19. The main results in this section are Theorem 2.19, together with Prop. 2.21, which provide differential graded models for fiberwise suspension spectra.

To fix notation and conventions, we briefly recall some background on homotopy theory:

Definition 2.1 (Classical homotopy theory (see e.g. [Sc17a])). We write

- (i) $\text{Ho}(\text{Spaces})$ for the homotopy category of topological spaces, called the *classical homotopy category*, which is the localization of the category of topological spaces at the *weak homotopy equivalences* (those maps inducing isomorphisms on all homotopy groups, which we will denote by \simeq_{whe}).
- (ii) $\text{Ho}(\text{Spaces})_{\mathbb{Q}, \text{ft}}$ for the full subcategory on homotopy types of *finite rational type*, namely those spaces X for which the homotopy groups $\pi_{k \geq 1}(X)$ are uniquely divisible (i.e., torsion-free and divisible), and $H^1(X, \mathbb{Q})$ and $\pi_{k \geq 2}(X) \otimes \mathbb{Q}$ are finite-dimensional \mathbb{Q} -vector spaces;
- (iii) $\text{Ho}(\text{Spaces})_{\mathbb{Q}, \text{nil}, \text{ft}} \subset \text{Ho}(\text{Spaces})_{\mathbb{Q}, \text{ft}}$ for the full subcategory of homotopy types that are moreover *nilpotent*, meaning those X that are connected and whose fundamental group $\pi_1(X)$ is a nilpotent group with each $\pi_n(X)$ a nilpotent $\pi_1(X)$ -module.

A nilpotent action is one whose iteration a certain number of times is the identity. Being simply-connected is a special case. Similar constructions hold in the parametrized case (see [CJ98, MS06]):

Definition 2.2 (Parametrized homotopy theory (e.g. [BM18, Sec. 1.1])). For any topological space X , we write

- (i) $\text{Ho}(\text{Spaces}/_X)$ for the homotopy category of spaces over X , that is, of spaces equipped with a map to X . We denote such objects by $[Y \xrightarrow{\pi} X]$; this is the *X -parametrized classical homotopy category*;
- (ii) $\text{Ho}(\text{Spaces} // _X)$ for the homotopy category of spaces over X that are equipped with a section. We denote such objects by $[X \xrightarrow{\sigma} Y \xrightarrow{\pi} X]$, where it is understood that the composite $\pi \circ \sigma = \text{id}_X$; morphisms are maps of spaces over X respecting the sections up to homotopy.

Remark 2.3. In the special case that $X \simeq *$ is the point, we simply have that

- (i) $\text{Ho}(\text{Spaces}/_*) \simeq \text{Ho}(\text{Spaces})$ is the classical homotopy category (Def. 2.1); and
- (ii) $\text{Ho}(\text{Spaces} // _*)$ is the homotopy category of *pointed* spaces.

In general, it is sensible to think of $\text{Ho}(\text{Spaces} // _X)$ as the homotopy category of *X -parametrized pointed spaces*.

Recall that there are two fundamental constructions associated to any pointed space Y : its based loop space $\Omega_* Y$ and its (reduced) suspension $\Sigma_* Y$. There are analogous constructions in the parametrized setting over a fixed base space X (see [CJ98, MS06]). These constructions compute loop spaces and reduced suspensions fiberwise over X :

Proposition 2.4 (Looping and suspension). *If Y is parametrized over a space X (Def. 2.2), we can form its fiberwise loop space $\Omega_X Y$. This construction is functorial and admits a left adjoint Σ_X , called the fiberwise reduced suspension:*

$$\text{Ho}(\text{Spaces} // _X) \begin{array}{c} \xleftarrow{\Omega_X} \\ \xrightarrow[\Sigma_X]{\tau} \end{array} \text{Ho}(\text{Spaces} // _X).$$

A fiberwise loop space $\Omega_X Y$ carries the structure of a fiberwise homotopical group (or, more precisely, a fiberwise *grouplike A_∞ -space*) given by concatenating loops. In the case of a fiberwise *double* loop space $\Omega_X^2 Y = \Omega_X \Omega_X Y$, the Eckmann–Hilton argument implies an additional fiberwise first-order homotopy commutativity structure given by twisting based loops around each other. Increasing the fiberwise loop order, as n goes to infinity the fiberwise homotopical group structure on $\Omega_X^n Y$ becomes increasingly homotopy-commutative. The $n \rightarrow \infty$ limit therefore provides a useful heuristic for obtaining *X -parametrized homotopical abelian groups*.

One way to formalize the idea that increasing loop order *stabilizes* to abelian homotopy theory is to exhibit a homotopy category for which the adjunction of Prop. 2.4 is an equivalence. In the unparametrized setting, the homotopy category obtained in this manner is the *stable homotopy category*, the objects of which are called *spectra*. By definition, a spectrum P is a sequence of pointed topological spaces $\{P_n\}_{n \in \mathbb{N}}$ equipped with structure maps $\Sigma P_n \rightarrow P_{n+1}$. To each spectrum P is naturally assigned a sequence

of abelian groups $\{\pi_k(P)\}_{k \in \mathbb{Z}}$ called the *stable homotopy groups* of P . There is a natural notion of a morphism of spectra with respect to which the assignment of stable homotopy groups is functorial. A map of spectra is a *stable weak equivalence* if it induces an isomorphism on all stable homotopy groups, and localizing the category of spectra at the class of stable weak equivalences produces the *stable homotopy category*. Every spectrum is stably weakly equivalent to a spectrum P for which the adjoints of the structure maps are weak homotopy equivalences $P_n \cong_{\text{whe}} \Omega P_{n+1}$. Spectra of this special type are called Ω -spectra, and for an Ω -spectrum P there are weak homotopy equivalences $P_k \cong \Omega^n P_{n+k}$ for all $n, k \geq 0$, exhibiting each P_k as an infinite loop space (or “homotopical abelian group”). Finally, the looping and suspension operations on spaces prolong to spectra and for any spectrum P we have natural isomorphisms

$$\pi_{k+1}(\Sigma P) \cong \pi_k(P) \cong \pi_{k-1}(\Omega P)$$

for all $k \in \mathbb{Z}$. Spectra are primarily of interest since they represent generalized cohomology theories on topological spaces. It is a consequence of the previously-mentioned Brown representability theorem that *all* generalized cohomology theories arise from spectra in this way. See [Sc17b] for a review of classical stable homotopy theory.

The above notions for spaces have analogues for spectra.

Definition 2.5 (Stable homotopy theory). We write

- (i) $\text{Ho}(\text{Spectra})$ for the *homotopy category of spectra*, also called the *stable homotopy category*;
- (ii) $\text{Ho}(\text{Spectra})_{\mathbb{Q}}$ for the *rational stable homotopy category*, hence the localization of the category of spectra at the maps inducing isomorphisms on rationalized stable homotopy groups $\pi_* \otimes \mathbb{Q}$;
- (iii) $\text{Ho}(\text{Spectra})_{\mathbb{Q}, \text{ft}}$ for the full subcategory on those spectra P which are of *finite rational type*, meaning that $\pi_k(P) \otimes \mathbb{Q}$ is a finite-dimensional \mathbb{Q} -vector space for all $k \in \mathbb{Z}$;
- (iv) $\text{Ho}(\text{Spectra})_{\mathbb{Q}, \text{bbl}}$ for the full subcategory of those spectra which are *rationally bounded below*, hence whose rationalized stable homotopy groups all vanish below some given dimension.

We are mainly concerned here with *parametrized* spectra, which are families of spectra parametrized by a topological space. Equivalently, these are bundles of spectra over that base space. Given a family of spectra P parametrized by a topological space X , for any point $x \in X$ we can extract a *stable homotopy fiber* $\mathbb{R}x^*P$ in a way that depends functorially on P (and on x in the appropriate homotopy-coherent sense). There is a natural notion of a map of X -parametrized spectra, and we declare a map $P \rightarrow Q$ to be a *fiberwise stable equivalence* if the induced map $\mathbb{R}x^*Q \rightarrow \mathbb{R}x^*P$ is a stable weak equivalence for all $x \in X$. For a rigorous approach to the theory using simplicial homotopy theory see [BM18, BM19a].

Definition 2.6 (Parametrized stable homotopy theory). For a fixed parameter space X , we write

- (i) $\text{Ho}(\text{Spectra}_X)$ for the *homotopy category of spectra parametrized by X* , hence the localization of the category of X -parametrized spectra at the fiberwise stable equivalences;
- (ii) $\text{Ho}(\text{Spectra}_X)_{\mathbb{Q}}$ for the *rational homotopy category of spectra parametrized by X* , hence the localization of category of X -parametrized spectra at the maps inducing isomorphisms on rationalized stable homotopy groups on all homotopy fiber spectra;
- (iii) $\text{Ho}(\text{Spectra}_X)_{\mathbb{Q}, \text{ft}, \text{bbl}} \subset \text{Ho}(\text{Spectra}_X)_{\mathbb{Q}}$ for the full subcategory on those X -spectra P whose homotopy fiber spectra $\mathbb{R}x^*(P)$ are of *finite rational type* and are *rationally bounded below* for all $x \in X$.

A key point about parametrized spectra (Def. 2.6) is that they represent *twisted* generalized cohomology theories, generalizing the fact that plain spectra represent generalized cohomology theories (e.g. [ABG⁺14]). We will be particularly interested in the (rational version of the) twisted cohomology theory called twisted K-theory (Lemma 2.31 below). The fundamental relation between unstable and stable homotopy theory in the parametrized setting is captured by the following:

Proposition 2.7 (Fiberwise stabilization adjunction). *For any space X there are pairs of adjoint functors*

$$\begin{array}{ccc}
 & \xleftarrow{\Omega_X^\infty} & \\
 \text{Ho}(\text{Spaces}_{/X}) & \xleftarrow{\perp} \text{Ho}(\text{Spaces}_{//X}) \xrightarrow{\perp} & \text{Ho}(\text{Spectra}_X) \\
 & \xrightarrow{(-)_{+,X}} & \xrightarrow{\Sigma_X^\infty} \\
 & \xrightarrow{\Sigma_{+,X}^\infty} &
 \end{array} \tag{6}$$

between the classical parametrized homotopy categories (Def. 2.2) and the parametrized stable homotopy category (Def. 2.6). Here $(-)+_{+,X}$ adjoins a copy of X , e.g. $[Y \rightarrow X] \mapsto [X \rightarrow X \coprod Y \rightarrow X]$, while Ω_X^∞ sends an X -parametrized spectrum to its fiberwise infinite loop space, and the operations Σ_X^∞ and/or $\Sigma_{+,X}^\infty$ are called forming fiberwise suspension spectra.

Moreover, the adjunction (6) stabilizes the looping/suspension adjunction from Prop. 2.4 in that there is a diagram, commuting up to natural isomorphism, as follows

$$\begin{array}{ccc}
 \text{Ho}(\text{Spaces}_{//X}) & \begin{array}{c} \xleftarrow{\Omega_X} \\ \xrightarrow{\Sigma_X} \end{array} & \text{Ho}(\text{Spaces}_{//X}) \\
 \begin{array}{c} \downarrow \Sigma_X^\infty \\ \uparrow \Omega_X^\infty \end{array} \dashv \! \! \dashv & & \begin{array}{c} \downarrow \Sigma_X^\infty \\ \uparrow \Omega_X^\infty \end{array} \dashv \! \! \dashv \\
 \text{Ho}(\text{Spectra}_X) & \begin{array}{c} \xleftarrow{\Omega_X} \\ \xrightarrow{\Sigma_X} \end{array} & \text{Ho}(\text{Spectra}_X)
 \end{array}$$

Remark 2.8 (Units). We denote the unit morphism of the adjunction (6) on $Y \in \text{Ho}(\text{Spaces}_{/X})$ by

$$\begin{array}{ccc}
 Y & \xrightarrow{\text{st}_X(Y)} & \Omega_X^\infty \Sigma_{+,X}^\infty(Y) \\
 & \searrow & \swarrow \\
 & & X
 \end{array} \quad (7)$$

Both the classical and stable homotopy categories are extremely rich mathematical settings. In order to get a better handle on these categories, one may filter them in various ways so as to study (stable) homotopy types in controlled approximations. A particularly useful approximation of this sort is provided by rational homotopy theory, which discards all torsion information carried by the homotopy groups. The main reason that rational homotopy theory is so tractable is that both the unstable and stable variants can be completely described in terms of algebraic data. We recall this as Prop. 2.11 and Prop. 2.17 below, but first we must recall some terminology:

Definition 2.9 (DG-algebraic homotopy theory). We write

- (i) $\text{Ho}(\text{DGCAlg})$ for the *homotopy category of connective differential graded (unital) commutative algebras (DG-algebras)* over \mathbb{Q} . The connectivity condition means that the underlying cochain complex vanishes identically in negative degree, and working in the homotopy category means that we localize the category of DG-algebras with respect to the class of quasi-isomorphisms;
- (ii) $\text{Ho}(\text{DGCAlg})_{\text{cn}}$ for the full subcategory on the *(cohomologically) connected* DG-algebras; those A for which the algebra unit $\mathbb{Q} \rightarrow A$ induces an isomorphism $\mathbb{Q} \simeq H^0(A)$;
- (iii) $\text{Ho}(\text{DGCAlg})_{\text{ft}}$ for the full subcategory of DG-algebras A of *finite type*, so that A is cohomologically connected and quasi-isomorphic to a DG-algebra that is degreewise finitely generated.

Remark 2.10 (Differentials). (i) We write “DG” throughout to indicate that we are working with cohomological grading conventions, so that in particular all differentials increase degree by +1.

(ii) We will write free graded commutative algebras (without differentials) as polynomial algebras

$$\mathbb{Q}[\alpha_{k_1}, \beta_{k_2}, \dots],$$

where the subscript on the generator will always indicate its degree. Differentials on such free algebras are fully determined by their actions on generators by the graded Leibniz rule, and so we will denote DG-algebras obtained this way by

$$\mathbb{Q}[\alpha_{k_1}, \beta_{k_2}, \dots] \left/ \left(\begin{array}{l} d\alpha_{k_1} = \dots \\ d\beta_{k_2} = \dots \\ \vdots \end{array} \right) \right.$$

The main result of Sullivan’s approach to rational homotopy theory [Su77] (a detailed treatment is the subject of the monograph [BG76]) is a characterization of certain well-behaved rational homotopy types in terms of DG-algebras:

Proposition 2.11 (DG-models for rational homotopy theory (e.g. [BG76, He06])).

(i) *There is an adjunction*

$$\mathrm{Ho}(\mathrm{Spaces}) \begin{array}{c} \xrightarrow{\mathcal{O}} \\ \perp \\ \xleftarrow{S} \end{array} \mathrm{Ho}(\mathrm{DGAlg})^{\mathrm{op}}$$

between the classical homotopy category of topological spaces (Def. 2.1) and the opposite of the homotopy category of DG-algebras (Def. 2.9), where \mathcal{O} denotes the derived functor of forming the DG-algebra of rational polynomial differential forms.

(ii) *This adjunction restricts to an equivalence of categories*

$$\mathrm{Ho}(\mathrm{Spaces})_{\mathbb{Q}, \mathrm{nil}, \mathrm{ft}} \begin{array}{c} \xrightarrow{\mathcal{O}} \\ \simeq \\ \xleftarrow{S} \end{array} \mathrm{Ho}(\mathrm{DGAlg})_{\mathrm{ft}}^{\mathrm{op}} \quad (8)$$

between the rational homotopy category of nilpotent spaces of finite type (Def. 2.1) and the homotopy category of DG-algebras of finite type (Def. 2.9).

(iii) *The rational cohomology of a space is computed by the cochain cohomology of any one of its DG-algebra models:*

$$H^\bullet(X, \mathbb{Q}) \simeq H^\bullet(\mathcal{O}(X)).$$

(iv) *Under the equivalence of (8), every space X on the left has a model by a minimal DG-algebra, whose underlying graded algebra is the free graded commutative algebra on the dual rational homotopy groups of X :*

$$\mathcal{O}(X) \simeq \mathbb{Q}[(\pi_\bullet(X) \otimes \mathbb{Q}^*)] / (d(\cdots) = (\cdots)).$$

Minimal models are unique up to isomorphism, with the isomorphism between any two minimal models unique up to homotopy.

Example 2.12 (Minimal model for the 3-sphere). The minimal model for the 3-sphere is given by $\mathcal{O}(S^3) \simeq \mathbb{Q}[h_3] / (dh_3 = 0)$. Observe that S^3 is rationally indistinguishable from a $K(\mathbb{Z}, 3)$. This is true for all odd-dimensional spheres, since $\pi_*(S^{2k+1}) \otimes \mathbb{Q}$ is a one-dimensional graded vector space concentrated in dimension $2k + 1$.

Example 2.13 (Minimal model for the 4-sphere). In contrast, the minimal model for the 4-sphere is not free, and is given by

$$\mathcal{O}(S^4) \simeq \mathbb{Q}[\omega_4, \omega_7] / \left(\begin{array}{l} d\omega_4 = 0 \\ d\omega_7 = -\frac{1}{2}\omega_4 \wedge \omega_4 \end{array} \right).$$

Example 2.14 (Minimal model for BS^1). The minimal model for the classifying space BS^1 of the circle is given by $\mathcal{O}(BS^1) \simeq \mathbb{Q}[\omega_2] / (d\omega_2 = 0)$. Note that this is decidedly not a minimal DGC-algebra model for the 2-sphere, since $\pi_2(S^2) \cong \mathbb{Z} \cong \pi_3(S^2)$.

We will model the rational homotopy theory of parametrized spectra in terms of DG-modules over DG-algebras:

Definition 2.15 (DG-modular homotopy theory). Given a DG-algebra A (Def. 2.9), we write

- (i) $A\text{-Mod}$ for the category of unbounded DG-modules over A ;
- (ii) $A\text{-Mod}_{\mathrm{ft}}$ for the subcategory of those DG- A -modules of *finite type*, hence those whose cochain cohomology is finite-dimensional in each degree;
- (iii) $A\text{-Mod}_{\mathrm{bbl}}$ for the subcategory of *bounded below* DG- A -modules, namely those whose cochain cohomology vanishes identically below some degree;
- (iv) $\mathrm{DGAlg}^{A/}$ for the slice category of A -algebras. The objects of this category are simply DG-algebras equipped with an algebra morphism from A , which we will frequently denote $B \leftarrow A: \pi^*$.
- (v) $\mathrm{DGAlg}_{//A}$ for the category of *augmented A -algebras*, whose objects are diagrams of DG-algebras

$$A \xleftarrow{\sigma^*} B \xleftarrow{\pi^*} A$$

such that $\sigma^* \circ \pi^*$ is the identity on A . The morphism σ^* is called the *augmentation*, and its kernel $\ker(\sigma^*) \in A\text{-Mod}$ is the *augmentation ideal*.

(vi) Passing to the homotopy category in any of the above cases (i)-(v) (e.g. $A\text{-Mod} \mapsto \text{Ho}(A\text{-Mod})$) means that we localize with respect to the class of quasi-isomorphisms.

Remark 2.16. We emphasise that we always work with *connective* DG-algebras, however modules over these algebras may be unbounded.

The stable analogue of Sullivan’s rational homotopy theory equivalence Prop. 2.11 (i) in the unparametrized context is the following:

Proposition 2.17 (DG-models for rational stable homotopy theory). *There is an equivalence*

$$\text{Ho}(\text{Spectra})_{\mathbb{Q},\text{ft}} \xrightleftharpoons[\simeq]{} \text{Ho}(\text{Ch}(\mathbb{Q}))_{\text{ft}}^{\text{op}}$$

between the rational homotopy category of spectra of finite type (Def. 2.1) and the opposite homotopy category of rational cochain complexes of finite type (as in Def. 2.15 for $A = \mathbb{Q}$).

Sketch of proof. This is a well-known fact in stable homotopy theory (and used extensively in differential cohomology, see [GS17]), but we sketch a proof for completeness. Passing to the rational stable homotopy category is implemented by smashing with the Eilenberg-MacLane spectrum $H\mathbb{Q}$, so that $\text{Ho}(\text{Spectra})_{\mathbb{Q}} \cong \text{Ho}(H\mathbb{Q}\text{-Mod})$. But the latter homotopy category is equivalent to the homotopy category of rational chain complexes $\text{Ho}(H\mathbb{Q}\text{-Mod}) \cong \text{Ho}(\text{ch}(\mathbb{Q}))$ (see [Sh07]). Under the assumption of finite type, dualizing then gives

$$\text{Ho}(\text{Spectra})_{\mathbb{Q},\text{ft}} \cong \text{Ho}(\text{Ch}(\mathbb{Q}))_{\text{ft}}^{\text{op}}.$$

An alternative proof which does not appeal to dualization is given in [BM18, Sec. 2.7.4]; see also [BM19b]. \square

Remark 2.18 (Operations). Under the above equivalence, suspension of spectra corresponds to shifting the corresponding cochain complex up by one. Looping corresponds to the cochain complex down by one.

The unification of Prop. 2.11 with Prop. 2.17 established in [BM18, BM19b] is:

Theorem 2.19 (DG-models for rational parametrized spectra). *For any connective DG-algebra A (Def. 2.9), there is a pseudo-natural transformation*

$$\text{Ho}(\text{Spectra}_{\mathcal{S}(A)}) \xrightarrow{\mathcal{M}_A} \text{Ho}(A\text{-Mod})^{\text{op}}$$

from the homotopy category of parametrized spectra (Def. 2.6) parametrized by the rational space $\mathcal{S}(A)$ (Prop. 2.11) to the opposite homotopy category of DG-modules over A (Def. 2.15) with the following properties:

1. There is a factorization over the rational homotopy category of parametrized spectra:

$$\begin{array}{ccc} \text{Ho}(\text{Spectra}_{\mathcal{S}(A)}) & \longrightarrow & \text{Ho}(\text{Spectra}_{\mathcal{S}(A)})_{\mathbb{Q}} \\ & \searrow \mathcal{M}_A & \downarrow \\ & & \text{Ho}(A\text{-Mod})^{\text{op}}. \end{array}$$

2. If the space $\mathcal{S}(A)$ is simply-connected, then \mathcal{M}_A restricts to an equivalence of rational homotopy categories

$$\text{Ho}(\text{Spectra}_{\mathcal{S}(A)})_{\mathbb{Q},\text{ft},\text{bbl}} \xrightarrow[\simeq]{\mathcal{M}_A} \text{Ho}(A\text{-Mod})_{\text{ft},\text{bbl}}^{\text{op}}$$

between finite-type, bounded below objects.

3. For $A = \mathbb{Q}$, this extends to the equivalence of rational stable homotopy theory from Prop. 2.17:

$$\text{Ho}(\text{Spectra})_{\mathbb{Q},\text{ft}} \xrightarrow[\simeq]{\mathcal{M}_A} \text{Ho}(\text{Ch}(\mathbb{Q}))_{\text{ft}}^{\text{op}}.$$

Sketch of proof. The functor \mathcal{M}_A is constructed by stabilizing the Sullivan–de Rham adjunction of Prop. 2.11 (i). The various properties of the stabilized functor are established in [BM18, BM19b]. Specifically, pseudo-naturality is [BM18, Cor. 2.7.26], the first item is [BM18, Cor. 2.7.31], the second item is [BM18, Theorem 2.7.42], and the final item is [BM18, Rem. 2.7.44]. \square

Remark 2.20 (Fiberwise operations). The obvious parametrized analogue of Rem. 2.18 holds, namely *fiberwise* suspension of a parametrized spectrum corresponds to shifting the corresponding DG-module up by one, and forming fiberwise loop spaces corresponds to shifting the DG-module down by one.

In the proof of Theorem 3.19 below, we will need to know explicitly how the functor \mathcal{M}_A behaves on fiberwise suspension spectra.

Proposition 2.21 (DG-models for fiberwise suspension spectra [BM18, BM19b]). *Let A be a DG-algebra (Def. 2.9) and let*

$$[Y \xrightarrow{\pi} \mathcal{S}(A)] \in \text{Ho}(\text{Spaces}/_{\mathcal{S}(A)})_{\mathbb{Q}}$$

be a space over (Def. 2.2) the rational space $\mathcal{S}(A)$ determined by A via Prop. 2.11. Then, after passing to DG-models via \mathcal{M}_A (Theorem 2.19), the fiberwise suspension spectrum (Prop. 2.7) is modeled by $\mathcal{O}(Y)$ (according to Prop. 2.11), regarded as an A -module via π^ ; that is,*

$$\mathcal{M}_A(\Sigma_{+, \mathcal{S}(A)}^{\infty}(Y)) \simeq \mathcal{O}(Y). \quad (9)$$

Sketch of proof. In the proof of Theorem 2.19 (see [BM18, Lemma 2.7.25], also [BM19b]), we encounter the commutative diagram of left Quillen functors:

$$\begin{array}{ccccc} & & \Sigma_{+, \mathcal{S}(A)}^{\infty} & & \\ & \xrightarrow{(-)_{+, \mathcal{S}(A)}} & & \xrightarrow{\Sigma_{\mathcal{S}(A)}^{\infty}} & \\ \text{Spaces}/_{\mathcal{S}(A)} & \xrightarrow{\quad} & \text{Spaces}/_{/\mathcal{S}(A)} & \xrightarrow{\quad} & \text{Spectra}_{\mathcal{S}(A)} \\ \mathcal{O}_A \downarrow & & \mathcal{O}_A \downarrow & & \downarrow \mathcal{M}_A \\ (\text{DGCAlg}^{A/})^{\text{op}} & \xrightarrow{(-) \oplus A} & (\text{DGCAlg}/_{/A})^{\text{op}} & \xrightarrow{\text{aug}_A} & A\text{-Mod}^{\text{op}} \end{array}$$

Here the left and bottom horizontal functors send

$$\begin{array}{c} \left[\begin{array}{c} Y \\ \downarrow \pi \\ \mathcal{S}(A) \end{array} \right] \xrightarrow{\mathcal{O}_A} \left[\begin{array}{c} \mathcal{O}(Y) \\ \uparrow \pi^* \\ \mathcal{O}(\mathcal{S}(A)) \\ \uparrow \eta \\ A \end{array} \right] \xrightarrow{(-) \oplus A} \left[\begin{array}{c} A \\ \uparrow 0 \oplus \text{id} \\ \mathcal{O}(Y) \oplus A \\ \uparrow (\pi^* \circ \eta) \oplus \text{id} \\ A \end{array} \right] \xrightarrow{\text{aug}_A} \ker(0 \oplus \text{id}) = \mathcal{O}(Y), \end{array}$$

from which the assertion follows. \square

Theorem 2.19 allows us to make use of the established theory of *minimal DG-modules* in order to obtain models for parametrized spectra in the rational approximation:

Definition 2.22 (Minimal DG-modules (see [Ro94][RS00, Sec. 1])). Let A be a DG-algebra. Write

$$A[n] \simeq A \otimes \langle c_n \rangle \in A\text{-Mod}$$

for the DG-module over A that is freely generated by a single generator c_n in degree $n \in \mathbb{Z}$. The underlying graded vector space is simply A shifted up (or down, if n is negative) in degree by $|n|$ and c_n is identified with shifted algebra unit. The differential is same as A , shifted in degree, and the module structure is given by left multiplication with elements in A .

(i) Given $N \in A\text{-Mod}$ and an element $\alpha \in N$ of degree $n + 1$ such that $d\alpha = 0$, we construct a new DG-module

$$N \oplus_{\alpha} (A \otimes \langle c_n \rangle) \in A\text{-Mod}$$

whose underlying graded vector space is the direct sum $N \oplus (A \otimes \langle c_n \rangle)$, equipped with the evident A -module structure. The differential is induced from the differentials of N and A , with the additional condition that $dc_n := \alpha$. Hence, the differential is specified by

$$d(n + a \otimes c_n) = d_N n + (d_A a) \otimes c_n + (-1)^{\deg(a)} a \cdot \alpha$$

for all $n \in N$ and $a \in A$. An A -module of the form $N \oplus_{\alpha} (A \otimes \langle c_n \rangle)$ is called an *n -cell attachment* of N .

(ii) A *relative cell complex over A* is an inclusion $N \hookrightarrow \widehat{N}$ of A -modules such that \widehat{N} arises from N via a countable sequence of cell attachments as in (i). If $N = 0$ is the zero module, then such a \widehat{N} is called a *cell complex over A* . A cell complex over A is therefore equivalent to the data of

- a graded vector space V equipped with a countable, ordered linear basis $\{e^{(i)}\}_{i \in \mathbb{N}}$; and
- a differential on $A \otimes V$ making this graded vector space a DG- A -module and such that

$$de^{(i)} \in A \otimes \langle e^{(j)} \rangle_{j \leq i}.$$

(iii) Furthermore, a cell complex over A is *minimal* the ordered basis $\{e^{(i)}\}_{i \in \mathbb{N}}$ satisfies the additional condition that $i \leq j$ if and only if $\deg(e^{(i)}) \leq \deg(e^{(j)})$.

Remark 2.23 (Existence of minimal models [Ro94]). For a (necessarily connective) DG-algebra A , any A -module M admits a *minimal model*. This means that there is a minimal A -module N together with a quasi-isomorphism of A -modules $N \rightarrow M$. Any two minimal models of M are unique up to isomorphism, and this isomorphism is unique up to homotopy.

Example 2.24 (Minimal cochain complexes have vanishing differential). If $A = \mathbb{Q}$, so that $A\text{-Mod} \simeq \text{Ch}(\mathbb{Q})$ is the category of rational cochain complexes, then the minimal DG-modules (Def. 2.22) are precisely the cochain complexes of finite type with *vanishing* differential. This is because a non-vanishing differential would have to take a generator $e^{(i)}$ in some degree n to a \mathbb{Q} -linear multiple of a generator $e^{(<i)}$ of degree $n + 1$, but this is ruled out by the degree condition on the generators.

According to Theorem 2.19, minimal DG-modules over DG-algebra A determine rational $\mathcal{S}(A)$ -spectra. Just as for minimal DG-algebras, the fiberwise rational stable homotopy groups of a parametrized spectrum can be read off directly from a minimal module:

Lemma 2.25. *For $E \in \text{Ho}(\text{Spectra})_{\mathbb{Q}, \text{ft}}$ any spectrum of finite rational type, the minimal DG-module model for its rationalization (Rem. 2.23) is the cochain complex*

$$(\pi_{\bullet}(E) \otimes \mathbb{Q})^*$$

equipped with vanishing differential.

Lemma 2.26 (Minimal models for fiber spectra (see [BM18, Rem. 2.7.35]; also [BM19b])). *Let $X \in \text{Ho}(\text{Spaces})$ be simply-connected of finite rational type, and let $E \in \text{Ho}(\text{Spectra}_X)_{\mathbb{Q}, \text{ft}, \text{bbl}}$ be a bounded-below X -spectrum of finite rational type (Def. 2.6). If, moreover,*

$$A \simeq \mathcal{O}(X) \in \text{DGCAlg}$$

is any DG-algebra model for X under Prop. 2.11, and if

$$A \otimes V \simeq \mathcal{M}(E) \in A\text{-Mod}$$

is a minimal DG-module model (Def. 2.22) for E under Theorem 2.19, then for every $x \in X$ the cochain complex with vanishing differential

$$V \simeq \mathcal{M}_{\mathbb{Q}}(E_x) \in \text{Ch}(\mathbb{Q})$$

is a minimal model for the fiber spectrum E_x .

Example 2.27 (Minimal model for suspension spectrum of BS^1). The minimal model for the suspension spectrum (Prop. 2.7) of the classifying space BS^1 (viewed as a parametrized spectrum over the point) is the graded vector space spanned by one generator in every non-negative even degree:

$$\mathcal{M}_{\mathbb{Q}}(\Sigma_+^{\infty} BS^1) \simeq \mathbb{Q}[\beta_2] = \langle 1, \beta_2, \beta_2^2, \dots \rangle \in \text{Ch}(\mathbb{Q}). \quad (10)$$

Indeed, a minimal DG-algebra model for BS^1 is the symmetric graded algebra on a single generator in degree 2:

$$\mathbb{Q}[\beta_2] \simeq \mathcal{O}(BS^1) \in \text{DGCAlg},$$

which necessarily has trivial differential; this implies the claim with Prop. 2.21. Note that in (10) the generator in degree $2n$ is β_2^n (however, this notation should be taken with a grain of salt since we have forgotten the algebra structure at this point).

Example 2.28 (Rational Snaith theorem). Let $KU \in \text{Ho}(\text{Spectra})$ be the spectrum representing complex K-theory and write $ku \in \text{Spectra}$ for its connective cover (obtained by killing negative-dimensional homotopy groups). Minimal DG-module models for KU and ku are, by Lemma 2.25, given by

$$\mathcal{M}_{\mathbb{Q}}(ku) \simeq \mathbb{Q}[\beta_2] \in \text{Ch}(\mathbb{Q}) \quad \text{and} \quad \mathcal{M}_{\mathbb{Q}}(KU) \simeq \mathbb{Q}[\beta_2, \beta_2^{-1}] \in \text{Ch}(\mathbb{Q}).$$

In particular, rationally there is no difference between ku and $\Sigma_+^{\infty} BS^1$:

$$\Sigma_+^{\infty} BS^1 \simeq_{\mathbb{Q}} ku.$$

On the other hand, if we remember the algebra structure on $\mathbb{Q}[\beta_2]$, the full non-connective K-theory is obtained by multiplicatively inverting the element β_2

$$\Sigma_+^{\infty} BS^1[\beta_2^{-1}] \simeq_{\mathbb{Q}} KU.$$

Remark 2.29 (Full Snaith theorem). As a matter of fact, the last statement in the previous example is still true *non-rationally*: this is the content of Snaith’s theorem [Sn81]. After rationalization, Snaith’s theorem essentially reduces to a triviality, however keeping in mind that this is a “rational shadow” may help identify the non-rational situation approximated by our main Theorem 3.19 below. We conjecture that Theorem 3.19 remains true non-rationally by a generalization of Snaith’s theorem to twisted K-theory obtained by *fiberwise* inversion of the Bott generator. We will return to this point elsewhere.

We have seen in Prop. 2.21 how stabilization – the process of passing from spaces to spectra – works in terms of rational models by taking augmentation ideals. Conversely, the *destabilization* process that extracts an infinite loop space from a spectrum also has a straightforward incarnation in terms of algebraic models. For connective parametrized spectra, extracting fiberwise infinite loop spaces is represented by taking the free algebra of the corresponding DG-module:

Proposition 2.30 (Rational models for fiberwise infinite loop spaces (see [BM18, Sec. 2.7]; [BM19b])). *Let A be a DG-algebra of finite type such that $\mathcal{S}(A)$ is simply-connected. If M is a connective A -module, then under the inverse equivalence of*

$$\text{Ho}(\text{Spectra}_{\mathcal{S}(A)})_{\mathbb{Q}, \text{ft}, \text{bbl}} \xrightarrow[\simeq]{\mathcal{M}_A} \text{Ho}(A\text{-Mod})_{\text{ft}, \text{bbl}}^{\text{op}},$$

the fiberwise infinite loop space (Prop. 2.7) is modelled by the augmented A -algebra $\text{Sym}_A(N)$ where N is a minimal model of M (Rem. 2.23).

Lemma 2.31 (Minimal model for twisted connective K-theory). *Denote the parametrized spectrum representing general twisted connective K-theory (e.g. [ABG10]) by*

$$ku//GL_1(ku) \longrightarrow BGL_1(ku)$$

*and its restriction to the twist by ordinary degree-3 cohomology by*⁶

$$\begin{array}{ccc} ku//BS^1 & \longrightarrow & ku//GL_1(ku) \\ \downarrow & \text{(pb)} & \downarrow \\ K(\mathbb{Z}, 3) & \longrightarrow & BGL_1(ku). \end{array}$$

The minimal DG-algebra model for the base space is the graded symmetric algebra freely generated by a single generator in degree h_3 with vanishing differential:

$$\mathcal{O}(K(\mathbb{Z}, 3)) \simeq \mathbb{Q}[h_3].$$

The corresponding minimal DG-module model (Def. 2.22) for the rationalization of $ku//BS^1$ is

$$\mathcal{M}_{\mathbb{Q}[h_3]}(ku//BS^1) \simeq \mathbb{Q}[h_3] \otimes \langle \omega_{2k} \mid k \in \mathbb{N} \rangle / \left(\begin{array}{l} d\omega_0 = 0 \\ d\omega_{2k+2} = h_3 \wedge \omega_{2k} \end{array} \right). \quad (11)$$

The module structure over $\mathbb{Q}[h_3]$ is given by the evident action on this tensor factor.

Moreover, for each $n \geq 0$ the fiberwise infinite loop space of the fiberwise suspension $\Sigma_{B^3\mathbb{Q}}^{2n}(ku//BS^1)$ has minimal DG-model

$$\mathcal{O}(\Omega_{B^3\mathbb{Q}}^{\infty-2n}(ku//BS^1)) \simeq \mathbb{Q}[h_3, \omega_{2k+2n} \mid k \in \mathbb{N}] / \left(\begin{array}{l} d\omega_{2n} = 0 \\ d\omega_{2k+2n+2} = h_3 \wedge \omega_{2k+2n} \end{array} \right). \quad (12)$$

⁶Here and elsewhere, “(pb)” denotes a (homotopy-)pullback square.

Proof. For the structure as a $\mathbb{Q}[h_3]$ -module, we appeal to Lemma 2.26 and Ex. 2.28. It only remains to determine the differential. By [AS05, Prop. 3.9] the degree-3 twist on complex K-theory is non-trivial in every degree. But by the degrees of the generators in (11), the given differential is degreewise and up to isomorphism the only possible non-trivial differential. The minimal models (12) are obtained by using Rem. 2.20 and Prop. 2.30. \square

A form of this statement also appears as [BN14, Ex. 12.5], [GS17, Sec. 3.1].

2.2 The Ext/Cyc-adjunction

Our mathematical formalization of double dimensional reduction and gauge enhancement in Sec. 4 involves at its core particular universal construction – the *Ext/Cyc-adjunction*. While we ultimately work with the rational homotopy theory version of this construction, we would like to amplify that this adjunction can be formulated much more generally. In particular, for \mathbf{H} any “good” homotopy theory (for instance an ∞ -topos [Lu09]) and for G a strong homotopy group in \mathbf{H} (namely, a grouplike A_∞ -monoid) with delooping $\mathbf{B}G$, there is a duality (an ∞ -adjunction)

$$\mathbf{H} \begin{array}{c} \xleftarrow{\text{Ext}_G} \\ \perp \\ \xrightarrow{\text{Cyc}_G} \end{array} \mathbf{H}/\mathbf{B}G$$

between

- (i) the operation Ext_G of forming G -extensions; and
- (ii) the operation of *G -cyclification*; the result of first forming the space of maps out of G and then taking the homotopy quotient by the G -action rigidly reparametrizing these maps.

In terms of abstract homotopy theory, this adjunction turns out to be right base change along the essentially unique point inclusion map $* \rightarrow \mathbf{B}G$. For the reader familiar with abstract homotopy theory this fully defines the adjunction, and the only point to check is that this right adjoint is indeed obtained by forming cyclifications as claimed.

For any object $X \in \mathbf{H}$, specifying a G -principal bundle on X is equivalent to the data of a map

$$\tau: X \longrightarrow \mathbf{B}G.$$

The G -principal bundle associated to τ is obtained by computing the homotopy fiber at the essentially unique point in $\mathbf{B}G$:

$$\begin{array}{ccc} \text{Ext}_G(\tau) & \longrightarrow & * \\ \downarrow & \text{(pb)} & \downarrow \\ X & \xrightarrow{\tau} & \mathbf{B}G, \end{array}$$

(see [NSS12] for an exposition of the general theory). Importantly for our purposes, for each map τ the Ext/Cyc-adjunction provides us with a natural morphism – the unit of the adjunction – which fits into a (homotopy) commutative diagram:

$$\begin{array}{ccc} X & \longrightarrow & \text{Cyc}_G \text{Ext}_G(\tau) \\ & \searrow \tau & \swarrow \\ & \mathbf{B}G & \end{array}$$

The map $X \rightarrow \text{Cyc}_G \text{Ext}_G(\tau)$ is the operation that takes any point in X to the map $G \rightarrow \text{Ext}_G(\tau)$ which winds identically around the extension fiber over that point. This is only well-defined up to a choice of base point in the fiber, but this is precisely the ambiguity that this quotiented out in the definition of Cyc_G .

Remark 2.32 (Notation). We will often abuse notation and write $\text{Ext}_G(X)$ instead of $\text{Ext}_G(\tau)$ when it is understood that we are considering a particular map $\tau: X \rightarrow \mathbf{B}G$.

We now describe the Ext/Cyc-adjunction in some detail in the setting of classical homotopy theory (Def. 2.1). More precisely, for any strict topological group G we exhibit an ordinary adjunction (Theorem 2.44 below) between categories of topological spaces which for $G = S^1$ presents Ext/Cyc-adjunction in homotopy (see Rem. 2.45). Let us first recall some preliminaries to establish our conventions:

Definition 2.33 (Group action on topological space). For X a topological space and G a topological group, a (*right*) *action* of G on X is a continuous function

$$\begin{aligned} X \times G &\longrightarrow X \\ (x, g) &\longmapsto x \cdot g \end{aligned}$$

such that $x \cdot e = x$ and $(x \cdot g_1) \cdot g_2 = x \cdot (g_1 \cdot g_2)$ for all $x \in X$ and $g_1, g_2 \in G$. One also refers to this situation by saying that X is a G -space.

Definition 2.34 (Quotients by group actions). For a G -space X , we write

$$X/G := X/(x \sim x \cdot g)$$

for the (ordinary) quotient space, which comes with the quotient projection

$$X \xrightarrow{\pi_G} X/G. \quad (13)$$

For G -spaces X and Y , we write

$$X \times_G Y := (X \times Y)/G := (X \times Y)/((x, y) \sim (x \cdot g, y \cdot g)) \quad (14)$$

for the quotient by the diagonal action.

Remark 2.35 (Comparison map for free actions). Recall that the G -action on X is called *free* if for every pair of points $(x_1, x_2) \in X \times X$ there is at most one $g \in G$ such that $x_2 = x_1 \cdot g$. For a free action there is a well-defined *comparison map* which we suggestively write as

$$\begin{aligned} X \times_{X/G} X &\longrightarrow G \\ [x_1, x_2] &\longmapsto x_1^{-1} \cdot x_2 \end{aligned} \quad (15)$$

such that $y_1 \cdot (x_1^{-1} \cdot x_2) = y_2$ whenever $[y_1, y_2] = [x_1, x_2]$. The comparison map determines a homeomorphism $X \times_{X/G} X \rightarrow X \times G$ via $[x_1, x_2] \mapsto (x_1, x_1^{-1} \cdot x_2)$.

Definition 2.36 (G -Extension functor). For any topological group G , there exists a topological space EG such that EG is a free G -space which is weakly contractible: the map $EG \rightarrow *$ is a weak homotopy equivalence. The quotient space

$$BG := (EG)/G \quad (16)$$

is the *classifying space* of G , and the quotient projection

$$EG \xrightarrow{\pi_G} BG \quad (17)$$

this is called the *universal G -principal bundle*. The G -bundle $EG \rightarrow BG$ is determined by this specification uniquely up to homotopy equivalence.

Given any topological space X equipped with a map $X \xrightarrow{\phi} BG$, we can pull back the universal bundle (17) to obtain a space $X \times_{BG} EG$ with free G -action whose quotient space is X :

$$\begin{array}{ccc} X \times_{BG} EG & \longrightarrow & EG \\ \pi_G \downarrow & \text{(pb)} & \downarrow \pi_G \\ X & \xrightarrow{\phi} & BG. \end{array} \quad (18)$$

This is the G -principal bundle *classified by ϕ* . For our purposes, it is useful to think of $X \times_{BG} EG$ as the *extension* that is classified by the *cocycle ϕ* . Therefore, we write Ext_G for the functor that computes these fiber products:

$$\begin{aligned} \text{Ext}_G: \text{Spaces}_{/BG} &\longrightarrow \text{Spaces} \\ (Y \rightarrow BG) &\longmapsto Y \times_{BG} EG. \end{aligned} \quad (19)$$

Definition 2.37 (Homotopy quotient). Given a G -space X (Def. 2.33), write $X//G$ for the *homotopy quotient space*. This is specified up to weak homotopy equivalence by the *Borel construction*

$$X//G := X \times_G EG, \quad (20)$$

which we take as our definition.

Example 2.38 (Homotopy quotient of trivial G -actions). The homotopy quotient of the (unique) G -action on the point $*$ is the classifying space (16):

$$*//G := (* \times EG)/G = (EG)/G = BG.$$

More generally, for a *trivial* G -space X (so that $x \cdot g = x$ for all g), the homotopy quotient is simply

$$X//G \simeq X \times BG.$$

Remark 2.39 (Maps related to the homotopy quotient). The homotopy quotient (Def. 2.37) is naturally equipped with the following maps of interest:

(i) The ordinary quotient projection (13) factors canonically up to homotopy via the homotopy quotient as

$$\pi_G : X \longrightarrow X//G \longrightarrow X/G.$$

Indeed, choosing any point $p \in EG$ (which is unique up to homotopy since EG is contractible), the factorization is obtained by the sequence of maps on the left-hand side of the diagram

$$\begin{array}{ccc} X & \xrightarrow{x \mapsto (x,p)} & X \times EG \\ \downarrow x \mapsto [x,p] & & \downarrow \\ X//G & \xlongequal{\quad} & X \times_G EG \\ \downarrow & & \downarrow \\ X/G & \xlongequal{\quad} & X \times_G * \end{array} \quad (21)$$

On the right-hand side we first take the quotient projection by the diagonal G -action on $X \times EG$ and then project out the EG factor via the (G -equivariant) map $EG \rightarrow *$. If the G -action on X is free, the comparison map $X//G \rightarrow X/G$ is a weak homotopy equivalence (see e.g. [Kö18]).

(ii) The homotopy quotient $X//G$ is equipped with a canonical map to the classifying space (16)

$$X//G \longrightarrow BG, \quad (22)$$

obtained via the map $X \rightarrow *$ as

$$X//G = X \times_G EG \longrightarrow * \times_G EG = (EG)/G = BG. \quad (23)$$

Proposition 2.40 (Extension of homotopy quotient). *Any G -space X is weakly homotopy equivalent to the G -extension (Def. 2.36) of its homotopy quotient $X//G$ (Def. 2.37) along the canonical map (22):*

$$\text{Ext}_G(X//G) \xrightarrow{\simeq_{\text{whe}}} X. \quad (24)$$

Proof. Unwinding the definitions, the extension in question is obtained as the pullback

$$\text{Ext}_G(X//G) = (X \times_G EG) \times_{BG} EG, \quad (25)$$

which is homeomorphic as a G -space to $X \times EG$, via the map

$$\begin{aligned} (X \times_G EG) \times_{BG} EG &\longrightarrow X \times EG \\ ([x, e_1], e_2) &\longmapsto (x \cdot (e_1^{-1} \cdot e_2), e_2) \end{aligned} \quad (26)$$

where we have used the comparison map (15) fiberwise over BG . Thus we have a map

$$\text{Ext}_G(X//G) \cong X \times EG \longrightarrow X$$

which is a weak homotopy equivalence by weak contractibility of EG . \square

Remark 2.41 ($\text{Ext}_G(X//G)$ as a free resolution). The induced action on $X//G \times_{BG} EG$ in (25) is always free, even if the action on X is not. Additionally, the isomorphism (26) is manifestly G -equivariant for the diagonal G -action on $X \times EG$. Hence Prop. 2.40 is saying that $\text{Ext}_G(X//G)$ is a *resolution* of X by a free G -space. For example, if $X = *$ then we have $*//G \times_{BG} EG \cong EG$.

We now turn to the description of the G -cyclification functor, which extends the cyclification functor from [FSS17, FSS18].

Definition 2.42 (Mapping space out of G). For a topological space Y , write $\text{Maps}(G, Y)$ for the space of continuous maps $G \rightarrow Y$.⁷ This mapping space is regarded as equipped with the G -action

$$\begin{aligned} \text{Maps}(G, Y) \times G &\longrightarrow \text{Maps}(G, Y) \\ (f, g) &\longmapsto [(f \cdot g): h \mapsto f(hg^{-1})], \end{aligned}$$

which, equivalently, is the conjugation action on maps of G -spaces where Y is regarded as having the trivial G -action.

Definition 2.43 (G -Cyclification). For G a topological group and Y a topological space, the G -cyclification of Y is the map

$$\text{Maps}(G, Y)//G \longrightarrow *//G = BG$$

obtained by forming the homotopy quotient (Def. 2.37) of the mapping space $\text{Maps}(G, Y)$ (Def. 2.42).

(i) This assignment extends to a functor

$$\begin{aligned} \text{Cyc}_G: \text{Spaces} &\longrightarrow \text{Space}/_{BG} \\ Y &\longmapsto \text{Maps}(G, Y) \times_G EG. \end{aligned}$$

(ii) For the special case that $G = S^1$ is the circle group we omit the subscript “ G ” and write simply

$$\text{Cyc}(X) = \mathcal{L}(X) := \text{Maps}(S^1, X)//S^1. \quad (27)$$

This is the homotopy quotient of the *free loop space* of X by the rigid rotation action on loops. The cohomology of $\text{Cyc}(X)$ is the *cyclic cohomology* of X , whence the terminology and notation.

The key fact relating the functors Ext_G and Cyc_G in this setting is the following:

Theorem 2.44 (The Ext/Cyc -adjunction). *Let G be a topological group. Then*

(i) *The functors Ext_G (Def. 2.36) and Cyc_G (Def. 2.43) are adjoints, with Ext_G the left and Cyc_G the right adjoint:*

$$\text{Spaces} \begin{array}{c} \xleftarrow{\text{Ext}_G} \\ \perp \\ \xrightarrow{\text{Cyc}_G} \end{array} \text{Spaces}/_{BG}.$$

(ii) *The unit of the adjunction*

$$X \xrightarrow{\eta_x} \text{Cyc}_G(\text{Ext}_G(X))$$

is the map that sends $x \in X$ to the equivalence class of the map $G \rightarrow \text{Ext}_G(X)|_x$ obtained by choosing any image of the neutral element $e \in G$ and extending to all of G by the group action:

$$\eta_x : x \longmapsto [G \xrightarrow{\simeq} \text{Ext}_G(X)|_x].$$

(iii) *The counit of the adjunction*

$$\begin{array}{ccc} \text{Ext}_G(\text{Cyc}_G(Y)) & \xrightarrow{\epsilon_Y} & Y \\ & \searrow \simeq_{\text{whe}} & \nearrow \text{ev}_e \\ & & \text{Maps}(G, Y) \end{array}$$

is the composite of a weak homotopy equivalence to $\text{Maps}(G, Y)$ followed by evaluation at the neutral element.

⁷we will always assume that all topological spaces are compactly generated, so that $\text{Maps}(G, Y)$ is the exponential object in the category of compactly generated spaces—this completely specifies the topology.

Proof. To show **(i)**, given $(c: X \rightarrow BG) \in \text{Spaces}_{BG}$ and $Y \in \text{Spaces}$, we must produce a natural bijection of sets

$$\text{Hom}\left(X \times_{BG} EG, Y\right) \cong \text{Hom}_{/BG}(X, \text{Maps}(G, Y) \times_G EG). \quad (28)$$

On the one hand, given a map

$$\begin{array}{ccc} X & \xrightarrow{\phi} & \text{Maps}(G, Y) \times_G EG \\ & \searrow c & \swarrow \\ & & BG \end{array}$$

that sends $x \mapsto [f_x, p_x]$, we define the map $\Phi(\phi): X \times_{BG} EG \rightarrow Y$ via $(x, p) \mapsto f_x(p^{-1} \cdot p_x)$. It is easy to check that this assignment does not depend on the choice of representative of the class $[f_x, p_x]$, and the argument of f_x is determined by the comparison map (15), since p, p_x lie in the same fiber of EG over BG .

Conversely, given a map

$$\begin{aligned} \Psi: X \times_{BG} EG &\longrightarrow Y \\ (x, p) &\longmapsto y_{x,p}, \end{aligned}$$

we define the map $\psi(\Psi): X \rightarrow \text{Maps}(G, Y) \times_G EG$ via the assignment $x \mapsto [y_{x, p_x \cdot (-)^{-1}}, p_x]$, where $p_x \in EG$ is such that $(x, p_x) \in X \times_{BG} EG$ and $y_{x, p_x \cdot (-)^{-1}}$ denotes the map $G \rightarrow Y$ given by the assignment $g \mapsto y_{x, p_x \cdot g^{-1}}$. It is straightforward to see that $\psi(\Psi)$ is well-defined and indeed determines a map of spaces over BG .

We now check that that the assignments $\phi \mapsto \Phi(\phi)$ and $\Psi \mapsto \psi(\Psi)$ are inverses of each other. If $\phi: x \mapsto [f_x, p_x]$ is as above, then we have

$$\psi(\Phi(\phi)): x \longmapsto [f_x((-) \cdot (p'_x)^{-1} \cdot p_x), p'_x] = [f_x, p_x],$$

where $p'_x \in EG$ is any point such that $(x, p_x) \in X \times_{BG} EG$. Similarly, $\Phi(\psi(\Psi)) = \Psi$ for all maps of spaces $\Psi: X \rightarrow \text{Cyc}_G(Y)$ over BG . Indeed, writing $\Psi: (x, p) \mapsto y_{x,p}$ as above then

$$\Phi(\psi(\Psi)): (x, p) \longmapsto y_{x, p_x \cdot (p_x^{-1} \cdot p)} = y_{x,p}.$$

This gives us a bijection on hom-sets of the desired form (28), which is manifestly natural in $(c: X \rightarrow BG)$ and Y . This completes the proof of **(i)**.

As to **(ii)**, we recall that the component of the unit at $(X \rightarrow BG)$ is the adjunct of the identity map on $\text{Ext}_G(X)$. By the above, this map is described by the assignment

$$x \longmapsto [(x, p_x \cdot (-)^{-1}), p_x],$$

where $p_x \in EG$ is any point such that $(x, p_x) \in X \times_{BG} EG$. But this is precisely of the claimed form: for each x we choose a point (x, p_x) in the fiber of $\text{Ext}_G(X)$ over X which is the image of the neutral element of G . Then an arbitrary element $g \in G$ is sent to $(x, p_x \cdot g^{-1})$, which determines a homeomorphism $G \cong \text{Ext}_G(X)|_x$. Passing to the homotopy quotient removes the choice ambiguity.

For **(iii)**, the component of the counit at Y is the adjunct of the identity on $\text{Cyc}_G(Y)$. By the above, this is simply the map

$$\begin{aligned} (\text{Maps}(G, Y) \times_G EG) \times_{BG} EG &\longrightarrow Y \\ ([f, p], p') &\longmapsto f((p')^{-1} \cdot p). \end{aligned}$$

In the proof of Prop. 2.40, we saw that

$$\begin{aligned} \kappa: (\text{Maps}(G, Y) \times_G EG) \times_{BG} EG &\longrightarrow \text{Maps}(Y, G) \times EG \\ ([f, p], p') &\longmapsto (f \cdot (p^{-1} \cdot p'), p') \end{aligned}$$

is a homeomorphism of G -spaces, so that the counit factors as

$$(\text{Maps}(G, Y) \times_G EG) \times_{BG} EG \xrightarrow{\simeq_{\text{whe}}} \text{Maps}(G, Y) \xrightarrow{\text{ev}_e} Y,$$

where we have used that $EG \rightarrow *$ is a weak homotopy equivalence. This completes the proof. \square

Remark 2.45 (Extension to the homotopy categories). The result we have just proven establishes Ext/Cyc-adjunction as an ordinary adjunction between categories of topological spaces. However, as we are interested in the corresponding adjunction between the *homotopy* categories, some additional points are in order.

- Since the universal G -principal bundle $EG \rightarrow BG$ is always a (Serre) fibration, taking fiber products with this map preserves weak homotopy equivalences. In particular, the functor Ext_G is *homotopical* and so descends to functor on derived categories $\text{Ho}(\text{Spaces}_{/BG}) \rightarrow \text{Ho}(\text{Spaces})$.
- The homotopical properties of the cyclification functor are more involved. Indeed, Cyc_G may fail to be a homotopical functor in general since $Y \mapsto \text{Maps}(G, Y)$ need not preserve weak homotopy equivalences, though this problem evaporates if G is a CW complex. In this article, we are primarily interested in $G = S^1$, in which case $\text{Cyc} = \text{Cyc}_{S^1}$ is a homotopical functor and so does determine a functor between the corresponding homotopy categories.

In summary, we have that Theorem 2.44 presents the adjunction between homotopy categories

$$\text{Ho}(\text{Spaces}) \begin{array}{c} \xleftarrow{\text{Ext}} \\ \perp \\ \xrightarrow{\text{Cyc}} \end{array} \text{Ho}(\text{Spaces}_{/BS^1})$$

and, therefore (upon further localization), between *rational* homotopy categories, which is our primary focus in this article. For more general G , the adjunction of Theorem 2.44 may fail to descend to homotopy categories. There are various ways of remedying this issue, but this takes us beyond the scope of the present article.

Below we will be mainly concerned with the Ext/Cyc-adjunction in the *rational* approximation. To see how this works, we first establish good models for the rationalization of the cyclification functor:

Remark 2.46 (Minimal models for cyclic loop spaces [VB85]). Let X be a simply-connected topological space of finite rational type (Def. 2.1) and let $(\mathbb{Q}[\{\omega_i\}], d_X)$ be a corresponding minimal DG-algebra (Prop. 2.11 (iv)). Then

(i) a minimal DG-algebra model for the free loop space $\mathcal{L}(X) := \text{Maps}(S^1, X)$ is obtained by adjoining a second copy $\{s\omega_i\}$ of the generators, where the degrees of these additional “looped generators” satisfy $\deg(s\omega_i) = \deg(\omega_i) - 1 = i - 1$, and with differential given by

$$d_{\mathcal{L}(X)}\omega_i := d_X\omega_i, \quad d_{\mathcal{L}(X)}s\omega_i := -s(d_X\omega_i),$$

where on the right s is uniquely extended from a linear map on generators to a graded derivation of degree -1 .

(ii) a minimal DG-algebra model for the cyclic loop space $\text{Cyc}(X)$ (Def. 2.43) is obtained from this by adjoining one more generator $\tilde{\omega}_2$ in degree 2, and taking the differential to be given by

$$d_{\text{Cyc}(X)}\tilde{\omega}_2 = 0, \quad d_{\text{Cyc}(X)}w = d_{\mathcal{L}(X)}w + \tilde{\omega}_2 \wedge sw,$$

where on the right $w \in \{\omega_i, s\omega_i\}$.

Example 2.47 (Minimal model for $\text{Cyc}(S^4)$ [FSS16a, Example 3.3]). By Remark 2.46, a minimal DG-algebra model for the cyclification of the 4-sphere is given by

$$\mathcal{O}(\text{Cyc}(S^4)) \simeq \mathbb{Q}[h_3, h_7, \omega_2, \omega_4, \omega_6] \left/ \begin{array}{l} dh_3 = 0 \\ dh_7 = -\frac{1}{2}\omega_4 \wedge \omega_4 + \omega_6 \wedge \omega_2 \\ d\omega_2 = 0 \\ d\omega_4 = h_3 \wedge \omega_2 \\ d\omega_6 = h_3 \wedge \omega_4 \end{array} \right.$$

This exhibits the structure of a DG-algebra over $\mathbb{Q}[h_3]/(dh_3 = 0)$, hence exhibiting a rational model for a map

$$\text{Cyc}(S^4) \longrightarrow S^3. \tag{29}$$

Example 2.48 ($\text{Cyc}(S^4)$ covers 6-truncated twisted K-theory, rationally). Ex. 2.47 reveals a close relationship between $\text{Cyc}(S^4)$ and the 6-truncation of $\Omega_{B^3\mathbb{Q}}^{\infty-2}(\text{ku}/BS^1)$ in the rational approximation. In

terms of the minimal models of Lemma 2.31, the 6-truncation $\Omega_{B^3\mathbb{Q}}^{\infty-2}(\mathrm{ku}/BS^1)\langle 6 \rangle$ is obtained simply by setting all $\omega_{\bullet>6}$ to zero. We then have the following morphisms in the rational homotopy category:

$$\begin{array}{ccccc} \Omega_{B^3\mathbb{Q}}^{\infty-2}(\mathrm{ku}/BS^1) & \xrightarrow{\tau_6} & \Omega_{B^3\mathbb{Q}}^{\infty-2}(\mathrm{ku}/BS^1)\langle 6 \rangle & \xleftarrow{p} & \mathrm{Cyc}(S^4) \\ h_3 & \longleftarrow & h_3 & \longrightarrow & h_3 \\ \omega_2 & \longleftarrow & \omega_2 & \longrightarrow & \omega_2 \\ \omega_4 & \longleftarrow & \omega_4 & \longrightarrow & \omega_4 \\ \omega_6 & \longleftarrow & \omega_6 & \longrightarrow & \omega_6 \\ \omega_8 & & & & h_7 \\ \vdots & & & & \end{array}$$

In Theorem 3.19 we encounter lifts $\widehat{\phi}$ of morphisms of rational homotopy types $\phi: X \xrightarrow{\phi} \mathrm{Cyc}(S^4)$ through this zig-zag, i.e., maps $\widehat{\phi}$ making the following diagram of maps of rational homotopy types commute:

$$\begin{array}{ccc} & & \Omega_{B^3\mathbb{Q}}^{\infty-2}(\mathrm{ku}/BS^1) \\ & \nearrow \widehat{\phi} & \downarrow \tau_6 \\ & & \Omega_{B^3\mathbb{Q}}^{\infty-2}(\mathrm{ku}/BS^1)\langle 6 \rangle \\ X_{\mathbb{Q}} & \xrightarrow{\phi} & \mathrm{Cyc}(S^4) \\ & & \uparrow p \end{array} \quad (30)$$

Proposition 2.49 (Minimal DG-module for fiberwise stabilization of $\mathrm{Cyc}(S^4)$).

(i) A minimal DG-module (Def. 2.22) modelling the fiberwise stabilization (Prop. 2.7) of the cyclic loop space of the 4-sphere (Ex. 2.47) over S^3 (via (29)) is

$$\mathcal{M}_{\mathbb{Q}[S^3]}(\Sigma_{+,S^3}^{\infty}\widehat{\mathrm{Cyc}}(S^4)) \simeq \frac{\mathbb{Q}[h_3, \omega_2, \omega_4, \omega_6]}{(\omega_6 \wedge \omega_2 - \frac{1}{2}\omega_4 \wedge \omega_4)} \Big/ \begin{pmatrix} dh_3 = 0 \\ d\omega_2 = 0 \\ d\omega_4 = h_3 \wedge \omega_2 \\ d\omega_6 = h_3 \wedge \omega_4 \end{pmatrix} \in \mathbb{Q}[h_3]\text{-Mod}.$$

Here the module structure is the evident one induced by multiplication in $\mathbb{Q}[h_3]$.

(ii) There is a quasi-isomorphism from this minimal model to the DG-module underlying the minimal DG-algebra from Ex. 2.47

$$\frac{\mathbb{Q}[h_3, \omega_2, \omega_4, \omega_6]}{(\omega_6 \wedge \omega_2 - \frac{1}{2}\omega_4 \wedge \omega_4)} \Big/ \begin{pmatrix} dh_3 = 0 \\ d\omega_2 = 0 \\ d\omega_4 = h_3 \wedge \omega_2 \\ d\omega_6 = h_3 \wedge \omega_4 \end{pmatrix} \xrightarrow{\simeq_{\mathrm{qi}}} \mathbb{Q}[h_3, h_7, \omega_2, \omega_4, \omega_6] \Big/ \begin{pmatrix} dh_3 = 0 \\ dh_7 = -\frac{1}{2}\omega_4 \wedge \omega_4 + \omega_2 \wedge \omega_6 \\ d\omega_2 = 0 \\ d\omega_4 = h_3 \wedge \omega_2 \\ d\omega_6 = h_3 \wedge \omega_4 \end{pmatrix}$$

given by any choice of linear splitting of the underlying quotient map of graded algebras, for example, by the map sending equivalence classes on the left to their unique representatives on the right that are at most linear in ω_4 .

Proof. For (i): by Prop. 2.21 and Lemma 2.26 the underlying graded $\mathbb{Q}[h_3]$ -module is the free $\mathbb{Q}[h_3]$ -module on the cohomology of the homotopy cofiber of

$$\mathbb{Q}[h_3] \longrightarrow \mathbb{Q}[h_3, h_7, \omega_2, \omega_4, \omega_6] \Big/ \begin{pmatrix} dh_3 = 0 \\ dh_7 = -\frac{1}{2}\omega_4 \wedge \omega_4 + \omega_2 \wedge \omega_6 \\ d\omega_2 = 0 \\ d\omega_4 = h_3 \wedge \omega_2 \\ d\omega_6 = h_3 \wedge \omega_4 \end{pmatrix},$$

where the minimal DG-algebra on the right is from Ex. 2.47. This cofiber cohomology is evidently the graded-commutative algebra

$$\frac{\mathbb{Q}[\omega_2, \omega_4, \omega_6]}{(\omega_6 \wedge \omega_2 - \frac{1}{2}\omega_4 \wedge \omega_4)},$$

obtained as the quotient by the two-sided tensor ideal generated by $\omega_6 \wedge \omega_2 - \frac{1}{2}\omega_4 \wedge \omega_4$. The graded vector space underlying the minimal DG-module is therefore

$$\mathbb{Q}[h_3] \otimes \frac{\mathbb{Q}[\omega_2, \omega_4, \omega_6]}{(\omega_6 \wedge \omega_2 - \frac{1}{2}\omega_4 \wedge \omega_4)} \simeq \frac{\mathbb{Q}[h_3, \omega_2, \omega_4, \omega_6]}{(\omega_6 \wedge \omega_2 - \frac{1}{2}\omega_4 \wedge \omega_4)}.$$

The differential on this must be such that fiberwise stabilization does not change the cohomology (by Prop. 2.21). This completely determines the differential, fixing it as claimed. The second point (ii) now follows at once. \square

3 The A-type orbispace of the 4-sphere

In this section we consider a particular circle action on the 4-sphere, as well as the induced homotopy quotient, which we call the *A-type orbispace of the 4-sphere* (see Def. 3.1 and Rem. 3.2 below). We first provide an informal string-theoretic motivation for considering this space in Rem. 3.4, and then substantiate this by a more formal mathematical analysis. After establishing some results on the rational homotopy type of the A-type orbispace in Sec. 3.1, our main result Theorem 3.19 shows that, rationally, there is a copy of twisted K-theory in the fiberwise stabilization of the A-type orbispace, fibered over the 3-sphere. In Sec. 4, we demonstrate how this result witnesses the phenomenon of gauge enhancement of M-branes.

Definition 3.1 (The A-type orbispace of the 4-sphere). Writing S^4 as the unit sphere in $\mathbb{R}^5 = \mathbb{R} \oplus \mathbb{C}^2$, the identification

$$S^4 = S(\mathbb{R} \oplus \mathbb{C}^2) \begin{array}{c} \text{SU}(2)_L \\ \curvearrowright \end{array} \quad (31)$$

shows that S^4 inherits an action of $SU(2)$. Specifically, on the right-hand side above we are referring to the defining linear representation of $SU(2)$ on \mathbb{C}^2 , regarded as a *left* action. This restricts along the canonical inclusion $S^1 \simeq U(1) \hookrightarrow SU(2)$ to define an S^1 -action on S^4 . We refer to the corresponding homotopy quotient (20)

$$S^4 // S^1 \simeq S^4 \times_{S^1} ES^1 \quad (32)$$

as the *A-type orbispace of the 4-sphere*.

Remark 3.2 (A-series vs. S^1). The terminology in Def. 3.1 is motivated as follows: the finite subgroups of $SU(2)$ have a famous ADE-classification, corresponding to the simply-laced Dynkin diagrams. The finite subgroups in the A-series are cyclic and, up to conjugation, are all subgroups of the canonical copy of S^1 inside $SU(2)$:

$$\mathbb{Z}_{n+1} \hookrightarrow S^1 \simeq U(1) \hookrightarrow SU(2).$$

The S^1 -action considered in Def. 3.1 is thus the limiting case (as $n \rightarrow \infty$) of the A-series actions. Now the homotopy quotient of the smooth 4-sphere such a finite group action is an *orbifold*, hence an *A-type orbifold* for an A-series group action (see [HSS18] for further discussion). More generally, homotopy quotients by (possibly non-finite) topological groups are *orbispaces* [HG07], whence our terminology.

The following result is immediate:

Proposition 3.3 (Quotient and fixed points of the A-type orbispace). *For the A-type S^1 -action on S^4 (Def. 3.1), it holds that:*

(i) *The ordinary quotient space is the 3-sphere: $S^4/S^1 \simeq S^3$. Hence, via (21) there is a canonical map from the A-type orbispace of the 4-sphere to the 3-sphere:*

$$\begin{array}{ccc} S^4 // S^1 & \longrightarrow & S^3 \\ \parallel & & \parallel \\ S^4 \times_{S^1} ES^1 & \xrightarrow{\text{id} \times_{S^1} p} & S^4 \times_{S^1} * \end{array} \quad (33)$$

(ii) *The space of S^1 -fixed points is the 0-sphere, included as two antipodal points*

$$S^0 = (S^4)^{S^1} \hookrightarrow S^4.$$

In summary, we have the following system of spaces over S^3 :

$$\begin{array}{ccccccc}
\text{Fixed} & & \text{4-sphere} & & \text{Homotopy} & & \text{Naive} \\
\text{points} & & & & \text{quotient} & & \text{quotient} \\
\overbrace{S^0 = (S^4)^{S^1}} & \hookrightarrow & \overbrace{S^4} & \longrightarrow & \overbrace{S^4 // S^1} & \longrightarrow & \overbrace{S^4 / S^1} \\
& & & & \downarrow & & \\
& & & & S^3 & &
\end{array} \tag{34}$$

Below in Sec. 4 we regard the (rationalization of the) A-type orbispace of the 4-sphere as the *coefficient* of a generalized cohomology theory. However, as explained in [HSS18, Sec. 2.2], the 4-sphere coefficient here ultimately originates as a factor in a black M5-brane spacetime $\sim \text{AdS}_7 \times S^4$. With this in mind, the spaces appearing in (34) readily explain those spaces appearing in the string theory literature.

Remark 3.4 (The A-type orbispace from black M5-brane geometry).

(i) The near-horizon geometries of black M2-brane and black M5-brane solutions of 11-dimensional supergravity are given by $\text{AdS}_4 \times S^7$ and $\text{AdS}_7 \times S^4$, respectively [Gu92]. Both of the spherical factors admit natural maps to the four-sphere, namely the quaternionic Hopf fibration $H_{\mathbb{H}}: S^7 \rightarrow S^4$ and the identity map $S^4 \rightarrow S^4$, and these maps generate the torsion-free homotopy of S^4 . It is natural to posit that S^4 is the coefficient for a *nonabelian* cohomology theory (in this case *cohomotopy*) that measures M-brane charge in the spirit of Dirac charge quantization [Fr00], at least rationally [Sa13, FSS15b, FSS16a].

$$\begin{array}{ccc}
\begin{array}{c} \text{one unit of} \\ \text{M2-brane charge} \end{array} & \curvearrowright & \\
\left[\underbrace{\text{AdS}_4 \times S^7}_{\text{black M2-brane spacetime}} \xrightarrow{\text{pr}_2} \underbrace{S^7}_{\text{sphere around M2-brane singularity}} \xrightarrow{H_{\mathbb{H}}} S^4 \right] & \in [Y, S^4] & \\
\text{cohomotopy classes} & & \\
\text{of } Y \text{ in degree 4} & & \tag{35} \\
\begin{array}{c} \text{one unit of} \\ \text{M5-brane charge} \end{array} & \curvearrowright & \\
\left[\underbrace{\text{AdS}_7 \times S^4}_{\text{black M5-brane spacetime}} \xrightarrow{\text{pr}_2} \underbrace{S^4}_{\text{sphere around M5-brane singularity}} \xrightarrow{\text{id}} S^4 \right] & \in [Y, S^4] &
\end{array}$$

(ii) More generally, the black M5-brane may sit inside an MK6, which itself is located at the singular locus of a global orbifold

$$\text{AdS}_7 \times S^4 // G_{\text{ADE}}$$

(see [HSS18, Ex. 2.7] for a precise statement and for pointers to the literature), where $G_{\text{ADE}} \subset \text{SU}(2)$ is a finite subgroup acting on the 4-sphere via the identification


$$\begin{array}{ccc}
G_{\text{ADE}} & & G_{\text{ADE}} \\
\curvearrowright & & \curvearrowright \\
S^4 & \simeq & S(\mathbb{R} \oplus \mathbb{C}^2).
\end{array}$$

In order for the 4-sphere charge coefficient to be able to measure the unit charges of such M5-branes sitting at ADE singularities in a manner generalizing (35), it must be equipped with that same group action. The resulting *equivariant* cohomotopy theory for M-branes is the subject of [HSS18].

(iii) Our current focus is on the A-series subgroups, which up to conjugation are the cyclic subgroups $\mathbb{Z}_{n+1} \hookrightarrow S^1 = \text{U}(1) \hookrightarrow \text{SU}(2)$, as in Rem. 3.2. By analogy with the case for M2-branes as in [ABJM08, p. 3], we may interpret the S^1 -action as being that of the M-theory circle fibration over 10d type IIA supergravity. With this interpretation in mind, passage to the finite A-type orbifold quotient

$$\text{AdS}_7 \times S^4 // \mathbb{Z}_{n+1}$$

corresponds to shrinking the M-theory circle fiber, and hence the coupling constant of non-perturbative type IIA string theory, by the factor $n + 1$. The limit $n \rightarrow \infty$, in which the cyclic groups \mathbb{Z}_{n+1} exhausts the group S^1 , corresponds to the limit of perturbative type IIA string theory. Via the maps of (21):



Applying the same logic as before, we might expect that the A-type orbispace $S^4//S^1$ serves as the charge quantization coefficient when M-branes are identified with their dual incarnations as D-branes in type IIA string theory. That this is indeed the case, up to a subtlety related to fiberwise stabilization, is essentially our result on gauge enhancement.

(iv) While the A-type orbispace $S^4//S^1$ has not previously featured in the string theory literature, the *ordinary* quotient space $S^4/S^1 \simeq S^3$ has been considered in this context. We briefly survey the related literature:

- The dimensional reduction of 11-dimensional supergravity on the 4-sphere factor yields a maximal $SO(5)$ -gauged supergravity in seven dimensions [PNT84]. The consistency of this reduction is established in [NVvN00] and a systematic classification of such reductions is given in [FOS04]. On the other hand, the reduction of type IIA supergravity on S^3 leads to an $SO(4)$ -gauged supergravity in seven dimensions. To compare these two gauged supergravity theories, one needs a means of breaking the $SO(5)$ gauge symmetry. In [CLP⁺00] the comparison between the two reductions is achieved using the singular scaling limit of S^4 opening up to $S^3 \times \mathbb{R}$, based on earlier arguments [HW88, CLL⁺00]. The consistency of such reductions is studied and established in [CLP00].
- Reductions with less symmetry are also possible, for instance by gauging only a left-acting $SU(2)$ subgroup of $\text{Spin}(4) \cong SU(2)_L \times SU(2)_R$ [CS00]. In [NV00], this was achieved using a singular limit of the S^4 reduction of 11d supergravity. In [HSS18, Sec. 2.2] it is explained how the distinction between these actions relates to the 4-sphere detecting black M5-branes as well as black M2-branes at singularities.

Remark 3.5 (Other circle actions on the 4-sphere). Circle actions on spheres form one of the most important problems in the theory of transformation groups. For a compact connected topological group G acting non-trivially on the 4-sphere, requiring orbits to be of dimension ≤ 1 immediately forces G to be the circle group. However, the action may not be equivalent to a differentiable one [MZ54]. Furthermore, there are infinitely many nonlinear circle actions on S^4 [Pa78]. Since S^4 is compact, it follows (by applying [FOT08, Theorem 7.33]) that in any case the fixed point set F will have the same Euler characteristic as S^4 , namely 2. Since the sum of dimensions of the cohomology groups of F is always at most as large as the corresponding sum for the space S^4 [FOT08, Theorem 7.37], this forces $\dim H^{\text{ev}}(F; \mathbb{Q}) = 2$ and $\dim H^{\text{odd}}(F; \mathbb{Q}) = 0$. Away from the trivial case ($F \simeq_{\mathbb{Q}} S^4$), this implies that there are only two possibilities for the rational homotopy type of the fixed point space: either F is rationally a 0-sphere (union of two points), or F is a rational homology 2-sphere. The latter case is described in [FOT08, Ex. 7.39]; in this article we deal exclusively with the former case.

3.1 Rational homotopy type of the A-type orbispace

In this section we study the rational homotopy type of the A-type orbispace of the 4-sphere (Def. 3.1) and apply the Ext/Cyc-adjunction to it. The main result is Prop. 3.12 below, but first we establish some preliminary results:

Lemma 3.6 (Rational homotopy and cohomology of $S^4//S^1$). *For every S^1 -action on the 4-sphere, the resulting homotopy quotient $S^4//S^1$ has the following properties:*

(i) *Its rational homotopy groups are*

$$\pi_{\bullet}^{\mathbb{Q}}(S^4//S^1) := \pi_{\bullet}(S^4//S^1) \otimes \mathbb{Q} \simeq \begin{cases} \mathbb{Q} & \text{in dimensions 2, 4 and 7} \\ 0 & \text{otherwise.} \end{cases} \quad (36)$$

(ii) *Its rational cohomology groups are*

$$H^{\bullet}(S^4//S^1, \mathbb{Q}) \simeq \langle \tilde{\omega}_0 \rangle \oplus \langle \omega_2 \rangle \oplus \bigoplus_{k \in \mathbb{N}} \langle \omega_2^{\wedge(k+2)}, \omega_4 \wedge \omega_2^{\wedge k} \rangle, \quad (37)$$

so that $\dim H^{2k}(S^4//S^1, \mathbb{Q})$ is 1 for $k = 0, 1$, is 2 for $k \geq 2$, and the odd cohomology vanishes.

Proof. The first statement follows with the long exact sequence of rational homotopy groups induced by the homotopy fiber sequence

$$S^1 \longrightarrow S^4 \longrightarrow S^4//S^1.$$

The second statement follows with the corresponding multiplicative Serre spectral sequence in rational cohomology (though we make no claims regarding the *algebra* structure on cohomology—the notation is merely suggestive of the manner in which these classes arise in the Serre spectral sequence). \square

Lemma 3.7 (DG-algebra model for general $S^4//S^1$). *For every S^1 -action on the 4-sphere, the minimal DG-algebra (via Prop. 2.11) of the resulting homotopy quotient (Def. 2.37) is of the form*

$$\mathcal{O}(S^4//S^1) \simeq \mathbb{Q}[\omega_2, \omega_4, h_7] \Big/ \left(\begin{array}{l} d\omega_2 = 0 \\ d\omega_4 = 0 \\ dh_7 = -\frac{1}{2}\omega_4 \wedge \omega_4 \\ \quad + c_1 \omega_2^{\wedge 4} + c_2 \omega_2^{\wedge 2} \wedge \omega_4 \end{array} \right) \in \text{DGCAlg} \quad (38)$$

for some coefficients $c_1, c_2 \in \mathbb{R}$.

Proof. By Prop. 2.11, the minimal DG-algebra model of $S^4//S^1$ has the following properties:

1. as a graded algebra, it is generated by the rational homotopy groups;
2. the differential on the minimal model is such that the cochain cohomology reproduces the rational cohomology of $S^4//S^1$.

By Lemma 3.6, we therefore have that the underlying graded commutative algebra of the minimal model of *any* $S^4//S^1$ is $\mathbb{Q}[\omega_2, \omega_4, h_7]$. By the second item in Lemma 3.6 and for degree reasons, the differential is necessarily of the form

$$d\omega_2 = 0, \quad d\omega_4 = 0, \quad dh_7 \neq 0.$$

There is a homotopy fiber sequence

$$S^4 \longrightarrow S^4//S^1 \longrightarrow BS^1,$$

which in the rational models is reflected by the requirement that setting ω_2 to zero in $\mathcal{O}(S^4//S^1)$ produces a (necessarily minimal) DG-algebra model for S^4 . Comparing with Ex. 2.13, this means that

$$dh_7 = -\frac{1}{2}\omega_4 \wedge \omega_4 + \text{terms of degree 8 at most linear in } \omega_4,$$

which completes the proof. \square

Lemma 3.8 (Rational Ext/Cyc-adjunction unit at $S^4//S^1$). *For any S^1 -action on S^4 , the composite of the Ext/Cyc-adjunction unit (Theorem 2.44) at $S^4//S^1$ with the cyclification of the equivalence of Prop. 2.40 is presented by the map of minimal DG-algebra models*

$$\begin{array}{ccc}
S^4//S^1 & \xrightarrow{\eta_{S^4//S^1}} & \text{Cyc Ext}(S^4//S^1) \xrightarrow{\simeq_{\text{whe}}} \text{Cyc}(S^4) \\
\omega_2 \longleftarrow & \xrightarrow{\quad} & \longleftarrow \omega_2 \\
\omega_4 \longleftarrow & \xrightarrow{\quad} & \longleftarrow \omega_4 \\
c_1 \omega_2^{\wedge 3} + c_2 \omega_2 \wedge \omega_4 \longleftarrow & \xrightarrow{\quad} & \longleftarrow \omega_6 \\
0 \longleftarrow & \xrightarrow{\quad} & \longleftarrow h_3 \\
h_7 \longleftarrow & \xrightarrow{\quad} & \longleftarrow h_7
\end{array} \tag{39}$$

where $c_1, c_2 \in \mathbb{R}$ are the constants as in (38).

Proof. We determine what the map $S^4//S^1 \rightarrow \text{Cyc}(S^4)$ does on generators of the rational homotopy groups. With this we can posit a map on minimal models, which turns out to be uniquely specified for degree reasons. To begin, we observe that the degree-3 generator h_3 must be sent to zero (since $S^4//S^1$ has no free homotopy in dimension 3). The degree-2 generator ω_2 on the right of (39) is sent to the generator of the same name on the left: all morphisms considered are over BS^1 , and ω_2 generates the minimal model of this space (Ex. 2.14). The zig-zag identity of the Ext/Cyc-adjunction gives us a commuting diagram

$$\begin{array}{ccccc}
S^4 \simeq_{\text{whe}} \text{Ext}(S^4//S^1) & \xrightarrow{\text{Ext}(\eta_{S^4//S^1})} & \text{Ext Cyc Ext}(S^4//S^1) & \simeq_{\text{whe}} & \text{Maps}(S^1, S^4) \\
& \searrow \text{id} & \downarrow \epsilon_{\text{Ext}(S^4//S^1)} & & \downarrow \text{ev}_0 \\
& & \text{Ext}(S^4//S^1) & \simeq_{\text{whe}} & S^4
\end{array}$$

with weak homotopy equivalences as shown due to Prop. 2.40. Examining the DG-algebra model of the free loop space (Remark 2.46), this means that the unit η sends non-shifted (non-looped) algebra generators to themselves. These generators are ω_4 and h_7 .

So far we have defined the desired map of DG-algebras on the generators ω_2 , ω_4 , h_3 , and h_7 . This map respects the differentials on ω_2 , ω_4 and h_3 , whereas respect for the differential on h_7

$$\begin{array}{ccc}
-\frac{1}{2}\omega_4 \wedge \omega_4 & \xleftarrow{\quad} & -\frac{1}{2}\omega_4 \wedge \omega_4 \\
+\omega_2 \wedge (c_1\omega_2^{\wedge 3} + c_2\omega_2 \wedge \omega_4) & \xleftarrow{\quad} & +\omega_2 \wedge \omega_6 \\
\uparrow d & & \uparrow d \\
h_7 & \xleftarrow{\quad} & h_7
\end{array}$$

forces $\omega_6 \mapsto c_1\omega_2^{\wedge 3} + c_2\omega_2 \wedge \omega_4$. This completely determines the map of minimal models (39). \square

Example 3.9 (DG-algebra model for homotopy quotient of the trivial action). The homotopy quotient of the trivial S^1 -action on S^4 is $S^4//S^1 \simeq S^4 \times BS^1$ (Ex.2.38). The minimal DG-algebra model of this product space is obtained by setting $c_1 = c_2 = 0$ in Lemma 3.7:

$$\mathcal{O}(S^4 \times BS^1) \simeq \mathbb{Q}[\omega_2, \omega_4, h_7] \Big/ \left(\begin{array}{l} d\omega_2 = 0 \\ d\omega_4 = 0 \\ dh_7 = -\frac{1}{2}\omega_4 \wedge \omega_4 \end{array} \right) \in \text{DGCAlg}.$$

Lemma 3.10 (A-type action on 4-sphere is rationally trivial). *The A-type circle action on the 4-sphere (Def. 3.1) is rationally trivial. That is, the action is represented in rational homotopy theory by the coprojection map of DG-algebras*

$$\begin{aligned}
\mathcal{O}(S^4) &\longrightarrow \mathcal{O}(S^1) \otimes \mathcal{O}(S^4) \\
\eta &\longmapsto 1 \otimes \eta.
\end{aligned}$$

In particular, the A-type orbispace $S^4//S^1$ is equivalent to the rationalization of the trivial action (Ex. 3.9).

Proof. It is sufficient to argue on minimal models: the action $\mu: S^1 \times S^4 \rightarrow S^4$ determines a dual map in the category of DG-algebras $\mu^*: \mathcal{O}(S^4) \rightarrow \mathcal{O}(S^1) \otimes \mathcal{O}(S^4)$ via Prop. 2.11. Since minimal models are cofibrant and fibrant in the model structure on DG-algebras (see [He06] for a review), the map μ^* is homotopic to a map between minimal DG-algebra models:

$$\mathbb{Q}[\omega_4, \omega_7] \Big/ \left(\begin{array}{l} d\omega_4 = 0 \\ dh_7 = -\frac{1}{2}\omega_4 \wedge \omega_4 \end{array} \right) \xrightarrow{\nu} \mathbb{Q}[\theta_1, \omega_4, \omega_7] \Big/ \left(\begin{array}{l} d\theta_1 = 0 \\ d\omega_4 = 0 \\ dh_7 = -\frac{1}{2}\omega_4 \wedge \omega_4 \end{array} \right).$$

The source of this map is the minimal DG-algebra model of the 4-sphere (Ex. 2.13), and the extra degree-1 generator θ_1 appearing in the target corresponds to $\pi_1(S^1) = \mathbb{Z}$. The map ν is completely determined by the images of the generators ω_4 and h_7 .

Now, $SU(2)$ acts on S^4 via the inclusion $SU(4) \hookrightarrow SO(5)$ (compare with (31)). In particular, the $SU(2)$ -action preserves the round volume form on S^4 . Restricting along $S^1 \hookrightarrow SU(2)$ and observing that the generator ω_4 represents the round volume form in cohomology forces $\nu(\omega_4) = \omega_4$. Up to non-zero scaling, the only way to define ν on the degree-7 generator h_7 that respects the differential is $\nu(h_7) = h_7$. \square

Remark 3.11. Some remarks on the above results are in order:

(i) In the above proof, we refer to the degree-7 generator in the minimal model of S^4 as h_7 , in line with the notation used throughout this section and in Sec. 4. This generator was called ω_7 in Ex. 2.13.

(ii) In interpreting expression (38) it may be worthwhile to view passing to rational homotopy theory as a homotopical analogue of forming first derivatives: that the A-type action on the 4-sphere is rationally trivial is analogous to finding that the derivative of some non-trivial function on the real line vanishes at the origin. That is to say, the A-type orbispace does *not* itself split as a product, but it does in the rational approximation. This turns out to be crucial for our gauge enhancement mechanism. In the companion article [HSS18] we go further and work in *equivariant* rational homotopy theory, which captures a great deal more information.

In summary, we have the following:

Proposition 3.12 (Minimal DG-algebra model for the A-type orbispace). *The minimal DG-algebra model of the A-type orbispace of the 4-sphere (Def. 3.1) is*

$$\mathcal{O}(S^4 // S^1) \simeq \mathbb{Q}[\omega_2, \omega_4, h_7] \Big/ \left(\begin{array}{l} d\omega_2 = 0 \\ d\omega_4 = 0 \\ dh_7 = -\frac{1}{2}\omega_4 \wedge \omega_4 \end{array} \right) \in \text{DGCAlg}.$$

Furthermore, the unit of the Ext \dashv Cyc-adjunction (Theorem 2.44) on the A-type orbispace, composed with the equivalence (23) from Prop. 2.40, pulls back the generators of the DG-algebra model for $\text{Cyc}(S^4)$ (Ex. 2.47) as follows:

$$\begin{array}{ccc} S^4 // S^1 & \xrightarrow{\eta_{S^4 // S^1}} & \text{Cyc Ext}(S^4 // S^1) \xrightarrow[\simeq_{\text{whe}}]{\text{Cyc}(\kappa)} \text{Cyc}(S^4) \\ \omega_2 \leftarrow & \xlongequal{\hspace{10em}} & \omega_2 \\ \omega_4 \leftarrow & \xlongequal{\hspace{10em}} & \omega_4 \\ 0 \leftarrow & \xlongequal{\hspace{10em}} & \omega_6 \\ 0 \leftarrow & \xlongequal{\hspace{10em}} & h_3 \\ h_7 \leftarrow & \xlongequal{\hspace{10em}} & h_7 \end{array} \tag{40}$$

Proof. By Lemma 3.10, the minimal DG-algebra model for A-type homotopy quotient $S^4 // S^1$ coincides with that of the trivial action, given by Ex. 3.9, hence is given by setting $c_1 = c_2 = 0$ in (38). We conclude with Lemma 3.8. \square

This concludes our discussion of the rational homotopy type of the A-type orbispace of the 4-sphere. Next, we discuss the fiberwise stabilization of $S^4 // S^1 \rightarrow S^4 / S^1 = S^3$ in rational homotopy theory.

3.2 Fiberwise stabilized Ext/Cyc-unit of the A-type orbispace

In Prop. 3.12 we described the rationalization of the unit $\eta_{S^4//S^1}$ of the Ext/Cyc-adjunction (Theorem 2.44) on the A-type orbispace of the 4-sphere (Def. 3.1). In the rational approximation, we may regard this map as lying over the classifying space $B^2S^1 \simeq B^3\mathbb{Z} \simeq_{\mathbb{Q}} B^3\mathbb{Q} \simeq_{\mathbb{Q}} S^3$ and discuss its *fiberwise stabilization* (Prop. 2.7) in rational parametrized stable homotopy theory (Theorem 2.19).

The main result of this section is Theorem 3.19, which states that the fiberwise stabilization of the A-type orbispace contains two summands of twisted connective K-theory, and characterizes lifts through the fiberwise stabilization adjunction unit in terms of lifting from 6-truncated to untruncated twisted K-theory (cf. Ex. 2.48). We interpret these lifts as being the rational image of gauge enhancement of M-branes in Sec. 4 below.

The next result appears in [RS00]. We spell out the proof in some detail, since we will need certain details in the proof of our main Theorem 3.19. In order to make certain features more apparent, we use a different naming convention for algebra generators than is used in [RS00]:

$$\begin{array}{c} \text{[RS00]:} \\ \text{here:} \end{array} \left\| \begin{array}{ccccccc} a & 1 & c_{2n} & c_{2n+1} & e & \gamma_{2n} & \gamma_{2n+1} \\ h_3 & \tilde{\omega}_0 & \omega_{2n+2}^L & \omega_{2n+4}^R & \omega_2 & \tilde{\omega}_{2n} & \omega_{2n} \end{array} \right.$$

Proposition 3.13 (Minimal DG-module for fiberwise stabilization of the A-type orbispace). *We have the following table of minimal DG-module models (Def. 2.22) describing fiberwise stabilizations of the spaces (34) over the 3-sphere:*

Fibration	Vector space underlying minimal DG-model	Differential of minimal DG-model
$\begin{array}{c} S^0 = (S^4)^{S^1} \\ \downarrow \\ S^3 \end{array}$	$\mathbb{Q}[h_3] \otimes \langle \omega_{2p}^L, \omega_{2p}^R \mid p \in \mathbb{N} \rangle$	$d : \left\{ \begin{array}{l} \omega_0^L \mapsto 0 \\ \omega_{2p+2}^L \mapsto h_3 \otimes \omega_{2p}^L \\ \omega_0^R \mapsto 0 \\ \omega_{2p+2}^R \mapsto h_3 \otimes \omega_{2p}^R \end{array} \right\} \begin{array}{l} (\text{ku} // BS^1) \\ (\text{ku} // BS^1) \end{array}$
$\begin{array}{c} S^4 \\ \downarrow \\ S^3 \end{array}$	$\mathbb{Q}[h_3] \otimes \langle \tilde{\omega}_{2p}, \omega_{2p+4} \mid p \in \mathbb{N} \rangle$	$d : \left\{ \begin{array}{l} \tilde{\omega}_0 \mapsto 0 \\ \tilde{\omega}_{2p+2} \mapsto h_3 \otimes \tilde{\omega}_{2p} \\ \omega_4 \mapsto 0 \\ \omega_{2p+6} \mapsto h_3 \otimes \omega_{2p+4} \end{array} \right\} \begin{array}{l} (\text{ku} // BS^1) \\ (\Sigma^4 \text{ku} // BS^1) \end{array}$
$\begin{array}{c} S^4 // S^1 \\ \downarrow \\ S^3 \end{array}$	$\mathbb{Q}[h_3, \omega_2] \otimes \langle \tilde{\omega}_{2p}, \omega_{2p+4} \mid p \in \mathbb{N} \rangle$	$d : \left\{ \begin{array}{l} \tilde{\omega}_0 \mapsto 0 \\ \tilde{\omega}_{2p+2} \mapsto h_3 \otimes \tilde{\omega}_{2p} \\ \omega_2 \mapsto 0 \\ \omega_4 \mapsto h_3 \wedge \omega_2 \otimes \tilde{\omega}_0 \\ \omega_{2p+6} \mapsto h_3 \otimes \omega_{2p+4} \end{array} \right\} \begin{array}{l} (\text{ku} // BS^1) \\ (\Sigma^2 \text{ku} // BS^1) \end{array}$

Beware of the special placement of the element ω_2 in the last line, as a generator of (the graded vector space underlying) a whole graded-commutative algebra. On the far right of the table we are highlighting that there are two sequences of differentials in each case, differing only in the degrees in which they start, and that each are of the same form as those for the minimal DG-model of the labelled shifted twisted K-theory spectrum (Lemma 2.31). Observe, moreover, that the second sequence in the last line really starts with the element $\omega_2 \otimes \tilde{\omega}_0$.

Proof. To determine the minimal DG-models, in each case we use that:

- by the particular form of the minimal DG-algebra model of S^3 (Ex. 2.12), the minimal DG-modules in question are necessarily *free* modules over $\mathbb{Q}[h_3]$; and
- according to (9), the cochain cohomology of the minimal DG-module must coincide with the rational cohomology of the total space of the corresponding fibration.

In all present cases of interest, these two constraints have a unique solution.

We begin by adjoining additional closed generators ω_k to capture the cohomology of the total space. But by the free module structure, this also makes the element $h_3 \otimes \omega_k$ appear, which must be killed off in cohomology. To do this, we introduce new generators with prescribed differential, which produce new spurious elements that need to be killed off, and so on. Explicitly, we have:

(i) We start with the case $S^0 \rightarrow S^3$. Write the 0-sphere as the disjoint union of a “left” and a “right” point

$$S^0 = *^L \amalg *^R.$$

In cohomology, the map $S^0 \xrightarrow{\pi} S^3$ is

$$\begin{array}{ccccc} H^\bullet(S^0, \mathbb{Q}) = \langle [\omega_0^L], [\omega_0^R] \rangle & & [\omega_0^L] + [\omega_1^R] & & 0 \\ \uparrow \pi^* & & \uparrow & & \uparrow \\ H^\bullet(S^3, \mathbb{Q}) = \langle [1], [h_3] \rangle & & [1] & & [h_3]. \end{array}$$

Thus, if the DG-model for the fiberwise stabilization of S^0 over S^3 is to be of the form

$$\begin{array}{ccccc} \langle 1, h_3 \rangle \otimes \langle \omega_0^L, \omega_0^R, \dots \rangle & & 1 \otimes (\omega_0^L + \omega_0^R) & & h_3 \otimes (\omega_0^L + \omega_0^R) \\ \uparrow \pi^* & & \uparrow & & \uparrow \\ \underbrace{\langle 1, h_3 \rangle}_{=\mathbb{Q}[h_3]} & & 1 & & h_3 \end{array}$$

with

$$d1 = 0, \quad dh_3 = 0, \quad d\omega_0^{L/R} = 0,$$

then arguing by induction proves the claim; firstly, there must be additional generators $\omega_2^{L/R}$ in order to remove the elements $h_3 \otimes \omega_0^{L/R}$ from cohomology:

$$d\omega_2^{L/R} = h_3 \otimes \omega_0^{L/R}.$$

But this means that the $h_3 \otimes \omega_2^{L/R}$ are closed, since

$$\begin{aligned} d(h_3 \otimes \omega_2^{L/R}) &= \underbrace{(dh_3)}_{=0} \otimes \omega_2^{L/R} - h_3 \otimes d\omega_2^{L/R} \\ &= \underbrace{-h_3 \wedge h_3}_{=0} \otimes \omega_0^{L/R}. \end{aligned}$$

In order to remove these elements from cohomology, we need to introduce new elements $\omega_4^{L/R}$ with differential

$$d\omega_4^{L/R} = h_3 \otimes \omega_2^{L/R},$$

and so on.

(ii) We now consider the case $S^4 \xrightarrow{\pi} S^3$, which in cohomology is the assignment

$$\begin{array}{ccccc} H^\bullet(S^4, \mathbb{Q}) = \langle [\tilde{\omega}_0], [\omega_4] \rangle & & [\tilde{\omega}_0] & & 0 \\ \uparrow \pi^* & & \uparrow & & \uparrow \\ H^\bullet(S^3, \mathbb{Q}) = \langle [1], [h_3] \rangle & & [1] & & [h_3]. \end{array}$$

Hence, if the minimal DG-model is to be of the form

$$\begin{array}{ccccc} \langle 1, h_3 \rangle \otimes \langle \tilde{\omega}_0, \omega_4, \dots \rangle & & 1 \otimes \tilde{\omega}_0 & & h_3 \otimes \tilde{\omega}_0 \\ \uparrow \pi^* & & \uparrow & & \uparrow \\ \langle 1, h_3 \rangle & & 1 & & h_3 \end{array}$$

with

$$d1 = 0, \quad d\tilde{\omega}_0 = 0, \quad d\omega_4 = 0,$$

then there need to be elements $\tilde{\omega}_2$ and ω_6 that remove $h_3 \otimes \tilde{\omega}_0$ and $h_3 \otimes \omega_4$ from cohomology via

$$d\tilde{\omega}_2 = h_3 \otimes \tilde{\omega}_0, \quad d\omega_6 = h_3 \otimes \omega_4.$$

But this implies that

$$d(h_3 \otimes \tilde{\omega}_2) = 0, \quad d(h_3 \otimes \omega_6) = 0,$$

so that to we must introduce additional generators $\tilde{\omega}_4$ and ω_8 to make these elements exact. But then $h_3 \otimes \tilde{\omega}_4$ and $h_3 \otimes \omega_8$ are closed, so that we must add $\tilde{\omega}_6$ and ω_{10} to remove them from cohomology. The result follows by induction.

(iii) Now consider the main case of interest, the A-type orbispace $S^4//S^1$. The cohomology of the total space was determined in Lemma 3.6, so that in cohomology the map $S^4//S^1 \rightarrow S^3$ is of the form

$$\begin{array}{ccc} H^\bullet(S^4//S^1, \mathbb{Q}) = \langle [\tilde{\omega}_0], [\omega_2], [\omega_4], [\omega_2 \wedge \omega_2], \dots \rangle & \begin{array}{c} [\tilde{\omega}_0] \\ \uparrow \\ [1] \end{array} & \begin{array}{c} 0 \\ \uparrow \\ [h_3] \end{array} \\ \uparrow \pi^* & & \\ H^\bullet(S^3, \mathbb{Q}) = \langle [1], [h_3] \rangle & & \end{array}$$

First consider $\tilde{\omega}_0$ and of $\omega_2 (\equiv \omega_2 \otimes \tilde{\omega}_0)$: analogously to (ii) above, these induce sequences of additional generators $\tilde{\omega}_{2p}$ and ω_{2p+4} with differentials as claimed. But adding in these generators already implies the existence of a further cohomology class in degree four, exhibited by the cocycle

$$\hat{\omega}_4 := 1 \otimes \omega_4 - \omega_2 \otimes \tilde{\omega}_2 \quad (41)$$

By degree reasons, the only potential primitives of $\hat{\omega}_4$ are non-zero multiples of h_3 , but this is closed. Thus we have already found the correct cohomology in dimensions ≤ 5 . Extending in powers of ω_2 , we find that the cocycles

$$\omega_2^{\wedge(k+2)} \otimes \tilde{\omega}_0 \quad \text{and} \quad \omega_2^{\wedge k} \otimes \hat{\omega}_4$$

recover the correct $2k+4$ -dimensional cohomology of $S^4//S^1$, and no new non-trivial cocycles arise in odd degrees. This completes the proof. \square

At this point, let us pause to provide some intuition for what is going on in Prop. 3.13:

Remark 3.14 (Interpretation of fiberwise stabilization of A-type orbispace). Due to the rational homotopy equivalence $S^3 \simeq_{\text{whe}, \mathbb{Q}} B^2S^1$, the homotopy fiber of any point inclusion $* \rightarrow S^3$ is, rationally, the classifying space BS^1 (16):

$$\text{hofib}(* \longrightarrow S^3) \simeq_{\text{whe}, \mathbb{Q}} BS^1 .$$

Accordingly, the homotopy fiber of $S^0 \rightarrow S^3$ is, rationally, the disjoint union of two copies of BS^1 :


$$\text{hofib}\left((S^4)^{S^1} = * \amalg * \longrightarrow S^3 \right) \simeq_{\text{whe}, \mathbb{Q}} BS^1 \amalg BS^1 .$$

Forming fiberwise suspension spectra (Prop. 2.7) really means that we stabilize these homotopy fibers, which means that the fiberwise suspension spectrum of $S^0 \rightarrow S^3$ is a parametrized spectrum whose fiber over any point is

$$\Sigma_+^\infty BS^1 \oplus \Sigma_+^\infty BS^1 \simeq_{\text{swhe}, \mathbb{Q}} \text{ku} \oplus \text{ku}$$

(cf. Ex. 2.28). Hence, the fiberwise suspension spectrum $\Sigma_{+, S^3}^\infty S^0$ is rationally equivalent to the direct sum of two copies of twisted connective K-theory, which by comparison in Lemma 2.31 is precisely what item (i) in Prop. 3.13 asserts.

The second item in Prop. 3.13 can also be heuristically understood in similar terms: as the cartoon picture



for the suspended Hopf action from Def. 3.1 indicates, the homotopy fiber now is some mixture of the two copies of BS^1 attached to the fixed points, and a copy of S^1 attached to all the other points. Prop. 3.16 below shows that, accordingly, there is one unshifted copy of $\Sigma_+^\infty BS^1$ in the fiber spectrum of $\Sigma_{+,S^3}^\infty S^4$ that pulls back diagonally into the direct sum of the suspension spectra associated with the two fixed points.

We record the following useful consequences of Prop. 3.13:

Lemma 3.15 (Comparison map between DG-module models of A-type orbispace). *A quasi-isomorphism of DG-modules over $\mathbb{Q}[h_3]$ from the minimal DG-model of $\Sigma_{+,S^3}^\infty (S^4 // S^1)$ (Prop. 3.13) to the (non-minimal) DG-model underlying the DG-algebra model of Prop. 3.12 is determined on generators as follows:*

$$\begin{array}{ccc}
\mathbb{Q}[\omega_2, \omega_4, h_7] / \left(\begin{array}{l} d\omega_2 = 0 \\ d\omega_4 = 0 \\ dh_7 = -\frac{1}{2}\omega_4 \wedge \omega_4 \end{array} \right) & \xleftarrow{\simeq_{\text{qi}}} & \mathbb{Q}[h_3, \omega_2] \otimes \langle \tilde{\omega}_{2p}, \omega_{2p+4} \rangle / \left(\begin{array}{l} d\tilde{\omega}_0 = 0 \\ d\tilde{\omega}_{2p+2} = h_3 \otimes \tilde{\omega}_{2p} \\ d\omega_2 = 0 \\ d\omega_4 = h_3 \wedge \omega_2 \otimes \tilde{\omega}_0 \\ d\omega_{2p+6} = h_3 \otimes \omega_{2p+4} \end{array} \right) \\
0 & \longleftarrow & h_3 \\
1 & \longleftarrow & \tilde{\omega}_0 \\
0 & \longleftarrow & \tilde{\omega}_{2p+2} \\
\omega_2 & \longleftarrow & \omega_2 \\
\omega_4 & \longleftarrow & \omega_4 \\
0 & \longleftarrow & \omega_{2p+6}
\end{array} \tag{42}$$

where both wedge products as well as tensor products (including powers of ω_2) on the right are sent to wedge products on the left.

Proof. The only non-evident point to note is that this map is indeed an isomorphism on cohomology in degree 4. But this follows from (41), since $\widehat{\omega}_4 \mapsto \omega_4$. \square

Proposition 3.16 (DG-model for S^0 - S^3 - S^4 system). *The commutative diagram (34)*

$$\begin{array}{ccc}
S^0 & \hookrightarrow & S^4 \\
& \searrow & \swarrow \\
& S^3 &
\end{array}$$

is represented on rational fiberwise suspension spectra in terms of the DG-modules of Prop. 3.13 by the commuting diagram

$$\begin{array}{ccc}
\left(\begin{array}{l} d\omega_0^{L/R} = 0 \\ d\omega_{2p+2}^{L/R} = h_3 \otimes \omega_{2p}^{L/R} \end{array} \right) & \xleftarrow{\left(\begin{array}{l} \tilde{\omega}_{2p} \mapsto (\omega_{2p}^L + \omega_{2p}^R) \\ \omega_{2p+4} \mapsto 0 \end{array} \right)} & \left(\begin{array}{l} d\tilde{\omega}_0 = 0, d\tilde{\omega}_{2p+2} = h_3 \otimes \tilde{\omega}_{2p} \\ d\omega_4 = 0, d\omega_{2p+6} = h_3 \otimes \omega_{2p+4} \end{array} \right) \\
\left(\begin{array}{l} 1 \mapsto 1 \otimes (\omega_0^L + \omega_0^R) \\ h_3 \mapsto h_3 \otimes (\omega_0^L + \omega_0^R) \end{array} \right) & \swarrow & \left(\begin{array}{l} 1 \mapsto 1 \otimes \tilde{\omega}_0 \\ h_3 \mapsto h_3 \otimes \tilde{\omega}_0 \end{array} \right) \\
& \left(\begin{array}{l} d1 = 0 \\ dh_3 = 0 \end{array} \right) &
\end{array}$$

Proof. We know that the various cohomology generators are mapped as follows:

$$\begin{array}{ccc}
1 \otimes ([\omega_0^L] + [\omega_0^R]) & \longleftarrow & 1 \otimes [\tilde{\omega}_0] \\
& \searrow & \swarrow \\
& [1] &
\end{array}$$

This implies the statement along the lines of the proof of Prop. 3.13. \square

So far, we have shown that copies of rational twisted K-theory appear as summands in the rational fiberwise stabilization of the A-type orbispace. Our next aim is to analyze how the Ext/Cyc-adjunction unit compares these copies to the copy of rational 6-truncated twisted K-theory to be found inside the rational cyclification of the 4-sphere (Ex. 2.48). For this, we will need to know how the adjunction unit is represented in terms of minimal DG-modules:

Proposition 3.17 (Fiberwise stabilization of Ext/Cyc-unit at A-type orbispace via minimal DG-models). *The fiberwise stabilization of the component of the Ext/Cyc-unit at the A-type orbispace of the 4-sphere*

$$\Sigma_{+,S^3}^\infty (S^4 // S^1) \xrightarrow{\Sigma_{+,S^3}^\infty(\eta)} \Sigma_{+,S^3}^\infty \text{Cyc}(S^4)$$

is represented on minimal DG-modules (Prop. 2.49 and Prop. 3.13) by a homomorphism of DG-modules over $\mathbb{Q}[h_3]$, shown as a dotted map in (44), which has the following properties:

(i) The generators ω_2, ω_4 and ω_6 from Prop. 2.49 are sent to the generators of the same name in Prop. 3.13:

$$\begin{array}{ccc} \omega_2 \otimes \tilde{\omega}_0 & \xleftarrow{\Sigma_{+,S^3}^\infty(\eta)^*} & \omega_2 \\ 1 \otimes \omega_4 & \xleftarrow{\quad} & \omega_4 \\ 1 \otimes \omega_6 & \xleftarrow{\quad} & \omega_6 \end{array} \quad (43)$$

(ii) The elements

$$1 \otimes \omega_{2p+8}, \quad p \in \mathbb{N}$$

are not in its image.

(iii) It is a minimal relative DG-module inclusion (Def. 2.22).

Proof. A general representative for $\Sigma_{+,S^3}^\infty(\eta)$ on minimal DG-models is a dotted morphism that makes the following diagram commute *up to homotopy*, but we claim that a representative with the stated properties exists that makes this diagram even *strictly* commutative:

$$\begin{array}{ccc} \Sigma_{+,S^3}^\infty (S^4 // S^1) \xrightarrow{\Sigma_{+,S^3}^\infty(\eta_{S^4 // S^1})} \Sigma_{+,S^3}^\infty \text{Cyc Ext} (S^4 // S^1) \xrightarrow[\simeq_{\text{whe}}]{\Sigma_{S^3}^\infty \text{Cyc}(\kappa)} \Sigma_{+,S^3}^\infty \text{Cyc}(S^4) & & (44) \\ \\ \mathbb{Q}[\omega_2, \omega_4, h_7] / \left(\begin{array}{l} d\omega_2 = 0 \\ d\omega_4 = 0 \\ dh_7 = -\frac{1}{2}\omega_4 \wedge \omega_4 \end{array} \right) \xleftarrow[\begin{array}{l} \omega_2 \mapsto \omega_2 \\ \omega_4 \mapsto \omega_4 \\ \omega_6 \mapsto 0 \end{array}]{\begin{array}{l} h_3 \mapsto 0 \\ h_7 \mapsto h_7 \end{array}} \mathbb{Q}[h_3, h_7, \omega_2, \omega_4, \omega_6] / \left(\begin{array}{l} dh_3 = 0 \\ dh_7 = -\frac{1}{2}\omega_4 \wedge \omega_4 + \omega_2 \wedge \omega_6 \\ d\omega_2 = 0 \\ d\omega_4 = h_3 \wedge \omega_2 \\ d\omega_6 = h_3 \wedge \omega_4 \end{array} \right) \\ \uparrow \simeq_{\text{qi}} \begin{array}{l} h_3 \mapsto 0 \\ \tilde{\omega}_0 \mapsto 1 \\ \tilde{\omega}_{2p+2} \mapsto 0 \\ \omega_2 \mapsto \omega_2 \\ \omega_4 \mapsto \omega_4 \\ \omega_{2p+6} \mapsto 0 \end{array} \uparrow \simeq_{\text{qi}} \begin{array}{l} h_3 \mapsto h_3 \\ \omega_2 \mapsto \omega_2 \\ \omega_4 \mapsto \omega_4 \\ \omega_6 \mapsto \omega_6 \end{array} \\ \mathbb{Q}[h_3, \omega_2] \otimes \langle \tilde{\omega}_{2p}, \omega_{2p+4} \rangle / \left(\begin{array}{l} dh_3 = 0 \\ d\tilde{\omega}_0 = 0 \\ d\tilde{\omega}_{2p+2} = h_3 \otimes \tilde{\omega}_{2p} \\ d\omega_2 = 0 \\ d\omega_4 = h_3 \wedge \omega_2 \otimes \tilde{\omega}_0 \\ d\omega_{2p+6} = h_3 \otimes \omega_{2p+4} \end{array} \right) \xleftarrow[\begin{array}{l} \omega_4 \mapsto 1 \otimes \omega_4 \\ \omega_6 \mapsto 1 \otimes \omega_6 \\ \vdots \end{array}]{\begin{array}{l} h_3 \mapsto h_3 \\ \omega_2 \mapsto \omega_2 \otimes \tilde{\omega}_0 \end{array}} \mathbb{Q}[h_3, \omega_2, \omega_4, \omega_6] / \left(\begin{array}{l} dh_3 = 0 \\ d\omega_2 = 0 \\ d\omega_4 = h_3 \wedge \omega_2 \\ d\omega_6 = h_3 \wedge \omega_4 \end{array} \right) \end{array}$$

Here the top horizontal morphism is the map from Prop. 3.12, regarded as a map between underlying DG-modules, while in the bottom row we have the corresponding minimal models. The left-hand vertical morphism is (42) from Lemma 3.15, while the right-hand vertical morphism is provided by Prop. 2.49.

Firstly, we show that if such a dotted morphism exists, then it must satisfy (i): To start with, the dotted morphism necessarily sends $h_3 \mapsto h_3$, since it is a homomorphism of free $\mathbb{Q}[h_3]$ -modules. Next, in order for the underlying linear maps to commute, the dotted morphism needs to send

$$\omega_2 \mapsto \omega_2 \otimes \tilde{\omega}_0 + \dots,$$

where the ellipsis indicates a term of degree 2 that vanishes under the left vertical map. The only possibility for such a term is a scalar multiple of $1 \otimes \tilde{\omega}_2$, but then the respect for the differential

$$\begin{array}{ccc} c h_3 \otimes \tilde{\omega}_0 & \longleftarrow & 0 \\ d \uparrow & & \uparrow d \\ \omega_2 \otimes \tilde{\omega}_0 + c 1 \otimes \tilde{\omega}_2 & \longleftarrow & \omega_2 \end{array}$$

enforces $c = 0$.

A similar argument applies to ω_4 : we must have

$$\omega_4 \mapsto 1 \otimes \omega_4 + \cdots$$

where the ellipsis indicates a term of degree 4 that vanishes under the left vertical map, which must be of the form $c_1 \omega_2 \wedge \tilde{\omega}_2 + c_2 1 \otimes \tilde{\omega}_4$ for some $c_1, c_2 \in \mathbb{R}$. But then respect for the differentials

$$\begin{array}{ccc} h_3 \wedge \omega_2 \otimes \tilde{\omega}_0 + c_1 \omega_2 \wedge h_3 \otimes \tilde{\omega}_0 + c_2 h_3 \otimes \tilde{\omega}_2 & \longleftarrow & h_3 \wedge \omega_2 \\ d \uparrow & & \uparrow d \\ 1 \otimes \omega_4 + c_1 \omega_2 \otimes \tilde{\omega}_2 + c_2 1 \otimes \tilde{\omega}_4 & \longleftarrow & \omega_4 \end{array}$$

implies $c_1 = 0$ and $c_2 = 0$.

Finally, we must have that

$$\omega_6 \mapsto 0 + \cdots,$$

where the ellipsis is a term of degree 6 that vanishes under the left vertical map. Hence

$$\omega_6 \mapsto c_1 1 \otimes \omega_6 + c_2 1 \otimes \tilde{\omega}_6 + c_3 \omega_2 \otimes \tilde{\omega}_4 + c_4 \omega_2^{\wedge 2} \otimes \tilde{\omega}_2$$

for some $c_1, c_2, c_3, c_4 \in \mathbb{R}$. Now the respect for the differentials

$$\begin{array}{ccc} c_1 h_3 \otimes \omega_4 + c_2 h_3 \otimes \tilde{\omega}_4 + c_3 \omega_2 \wedge h_3 \otimes \tilde{\omega}_2 + c_4 \omega_2^{\wedge 2} \wedge h_3 \otimes \tilde{\omega}_0 & \longleftarrow & h_3 \wedge \omega_4 \\ \uparrow & & \uparrow \\ c_1 1 \otimes \omega_6 + c_2 1 \otimes \tilde{\omega}_6 + c_3 \omega_2 \otimes \tilde{\omega}_4 + c_4 \omega_2 \wedge \omega_2 \otimes \tilde{\omega}_2 & \longleftarrow & \omega_6 \end{array}$$

implies $c_1 = 1$ and $c_2 = c_3 = c_4 = 0$. This establishes (i), provided that the dotted morphism actually exists.

We now prove that a dotted morphism in (44) does exist as claimed. Indeed, we have just seen that the images of the generators h_3, ω_2, ω_4 , and ω_6 are predetermined. Extending this map as a $\mathbb{Q}[h_3, \omega_2]$ -module homomorphism, we have defined the dotted morphism on all terms that are at most linear in ω_4 and ω_6 . We must now extend this map to higher order terms in these generators—but by the relation $\frac{1}{2}\omega_4 \wedge \omega_4 = \omega_2 \wedge \omega_6$, which holds in the minimal DG-module in the bottom right of (44), we may consider those unique representatives which are at most linear in ω_4 . Hence we need only find consistent images for all elements of the form

$$(\omega_6)^{\wedge n}, \quad (\omega_6)^{\wedge(n-1)} \wedge \omega_4 \quad \text{for } n \geq 2.$$

We claim that there are unique coefficients $a_n, b_n \in \mathbb{N} \subset \mathbb{Q}$ such that the assignment

$$\begin{array}{ccc} b_n \omega_2^{\wedge(n-1)} \otimes \omega_{4n} & \longleftarrow & (\omega_6)^{\wedge(n-1)} \wedge \omega_4 \\ a_n \omega_2^{\wedge(n-1)} \otimes \omega_{4n+2} & \longleftarrow & (\omega_6)^{\wedge n} \end{array} \quad \text{for } n \geq 2. \quad (45)$$

respects the differentials and makes the underlying linear maps of (44) commute. We argue by induction: for the initial case $n = 2$, respect for the differentials on $\omega_6 \wedge \omega_4$ means that the following square has to commute

$$\begin{array}{ccc} 3h_3 \wedge \omega_2 \otimes \omega_6 & \longleftarrow & 3h_3 \wedge \omega_2 \wedge \omega_6 \\ d \uparrow & & \uparrow d \\ b_2 \omega_2 \otimes \omega_8 & \longleftarrow & \omega_6 \wedge \omega_4, \end{array} \quad (46)$$

where the top left entry shows the image of the top right element. To see this, notice that:

- (a) in computing the differential on the right of (46), we are using the relation $\frac{1}{2}\omega_4 \wedge \omega_4 = \omega_2 \wedge \omega_6$ in the minimal DG-module model from Prop. 2.49, to uniquely represent all differentials by representatives that are at most linear in ω_4 ; and
- (b) the definition (45) applies only to terms at least quadratic in ω_6, ω_4 . For the image of the top right element in (46) we instead use (43) and respect for the $\mathbb{Q}[h_3, \omega_2]$ -module structure.

But the differential of the bottom left element in (46) is $b_2 \omega_2 \wedge h_3 \otimes \omega_6$, and hence the square (46) commutes precisely if $b_2 = 3$. In an analogous manner, we see that the respect for the differential on $\omega_6 \wedge \omega_6$ means that the following square has to commute:

$$\begin{array}{ccc} 2b_2 h_3 \wedge \omega_2 \otimes \omega_8 & \longleftarrow & 2h_3 \wedge \omega_6 \wedge \omega_4 \\ \uparrow d & & \uparrow d \\ a_2 \omega_2 \otimes \omega_{10} & \longleftarrow & \omega_6 \wedge \omega_6, \end{array}$$

where again the top left element shown is the image of the top right element, now obtained via (45). We see that the differential on the left has this same image precisely if

$$a_2 = 2b_2 = 6.$$

For the inductive argument, we assume that the claim (45) holds for $(n-1)$, with $n \geq 3$. Then we need to assure the commutativity of the squares

$$\begin{array}{ccc} nb_n h_3 \wedge \omega_2^{\wedge(n-1)} \otimes \omega_{4n} & \longleftarrow & n h_3 \wedge (\omega_6)^{\wedge(n-1)} \wedge \omega_4 \\ \uparrow d & & \uparrow d \\ a_n \omega_2^{\wedge(n-1)} \otimes \omega_{4n+2} & \longleftarrow & (\omega_6)^{\wedge n} \end{array}$$

and

$$\begin{array}{ccc} (2n-1) a_{n-1} h_3 \wedge \omega_2^{\wedge(n-2)} \otimes \omega_{4n-2} & \longleftarrow & (2n-1) h_3 \wedge \omega_2 \wedge (\omega_6)^{\wedge(n-1)} \\ \uparrow d & & \uparrow d \\ b_n \omega_2^{\wedge(n-1)} \otimes \omega_{4n} & \longleftarrow & (\omega_6)^{\wedge(n-1)} \wedge \omega_4. \end{array}$$

By the inductive hypothesis, the second square above implies that $b_n = (2n-1)a_{n-1}$, whereas the first square implies that $a_n = n \cdot b_n$. By induction, this gives a map of DG- $\mathbb{Q}[h_3, \omega_2]$ -modules as per the dotted arrow in (44) via the specification (45). Let us note for completeness that the coefficients in (45) are given recursively by

$$\frac{a_n}{a_{n-1}} = 2n^2 - n, \quad b_n = \frac{a_n}{n} \quad \text{for } n > 2,$$

with $a_2 = 6, b_2 = 3$. This completes the construction of the dotted morphism in (44), and it is now easy to see that that diagram commutes.

By construction, it is clear that the elements $1 \otimes \omega_{2p+8}, p \geq 0$ are not in the image of the dotted morphism. This proves item (ii).

Finally, the construction of the dotted morphism via (i) and (45) is a map between minimal DG-modules (over $\mathbb{Q}[h_3]$) which is moreover manifestly an injective cell map in that it injectively maps generators to generators. It follows that this map is a relative cell complex inclusion, in fact a minimal DG-module inclusion. This completes the proof. \square

Remark 3.18. The proof of (iii) depends crucially upon the fact that the elements $\omega_2^{\wedge n}$ for varying n are *distinct* as generators for the minimal DG-modules appearing along the bottom of Diagram (44). Despite being a potential source of great confusion, we persist in using this notation because of the utility of being able to work “multiplicatively in ω_2 ” in the argument above.

We are now in a position to state and prove our main technical result:

Theorem 3.19 (K-theory “detruncation” by lifting against fiberwise stabilization of A-type orbispace). *Regarding the A-type orbispace of the 4-sphere (Def. 3.1) as fibered over the 3-sphere via the canonical map (34) from Prop. 3.3,*

$$S^4 // S^1 \longrightarrow S^3 \xrightarrow{\simeq_Q} B^3\mathbb{Q}, \quad (47)$$

we have that:

(i) *Rationally, the fiberwise stabilization (Prop. 2.7) of (47) contains a copy of the 2-fold suspension of twisted connective K-theory as a direct summand:*

$$\Sigma^2 \text{ku} // BS^1 \hookrightarrow \Sigma_{+,S^3}^\infty (S^4 // S^1).$$

(ii) *A map ϕ of rational homotopy types over $B^3\mathbb{Q}$*

$$\begin{array}{ccc} X & \xrightarrow{\phi} & \text{Cyc}(S^4) \xrightarrow{p} \Omega_{B^3\mathbb{Q}}^{\infty-2}(\text{ku} // BS^1) \langle 6 \rangle \\ & \searrow \mu_3 & \swarrow \mu_3 \\ & & B^3\mathbb{Q} \end{array}$$

lifts from rational 6-truncated twisted K-theory to rational untruncated twisted K-theory if and only if ϕ admits a lift through the fiberwise stabilization of the Ext/Cyc-unit $\eta_{S^4 // S^1}$:

$$\begin{array}{ccccc} & & \widehat{\text{st} \circ \phi} & & \\ & \nearrow \text{dotted} & \text{dotted} & \searrow \text{dotted} & \\ & \Omega_{B^3\mathbb{Q}}^{\infty-2}(\text{ku} // BS^1) & \hookrightarrow & \Omega_{S^3}^\infty \Sigma_{+,S^3}^\infty (S^4 // S^1) & \\ & \downarrow \tau_6 & & \downarrow \Omega_{S^3}^\infty \Sigma_{+,S^3}^\infty (\eta_{S^4 // S^1}) & \\ X & \xrightarrow{\phi} & \text{Cyc}(S^4) \xrightarrow{p} \Omega_{B^3\mathbb{Q}}^{\infty-2}(\text{ku} // BS^1) \langle 6 \rangle & \xrightarrow{\text{st}} & \Omega_{S^3}^\infty \Sigma_{+,S^3}^\infty \text{Cyc}(S^4) \xrightarrow{\simeq} \Omega_{S^3}^\infty \Sigma_{+,S^3}^\infty \text{Cyc Ext}(S^4 // S^1) \\ & \searrow \mu_3 & \swarrow \mu_3 & & \\ & & B^3\mathbb{Q} & & \end{array}$$

That is, specifying a dotted lift $\widehat{\phi}$ is equivalent to specifying a dotted lift $\widehat{\text{st} \circ \phi}$, where st is the unit of the fiberwise stabilization adjunction.

Proof. The rational parametrized spectra appearing on the right-hand side of the diagram have minimal DG-module models as in Prop. 2.47 and Prop. 3.13. On these minimal models, the vertical map on the right-hand side is represented by the dotted morphism in (44). The fiberwise stabilization map st is represented in algebra by the right-hand vertical map in (44). The claim (i) was already established in Prop. 3.13.

To prove (ii), we argue as follows. Recall that by Prop. 2.30, rational models for fiberwise infinite loop spaces are obtained by taking free relative algebras, which allows us to argue on module generators (equivalently, we argue on $\Sigma_{+,S^3}^\infty \dashv \Omega_{S^3}^\infty$ -adjuncts). By item (i) of Prop. 3.17, a strict lifting of $\text{st} \circ \phi$ through the stabilized Ext/Cyc-unit implies a strict lifting of ϕ through the zig-zag morphism τ_6 —this is since the images of ω_2 , ω_4 and ω_6 are already specified in the rational model of X . Since untruncated twisted K-theory includes as a direct summand via ι , the converse is also true: a strict lifting of ϕ through the zig-zag τ_6 extends by zero to the complementary direct summand in $\Omega_{S^3}^\infty \Sigma_{+,S^3}^\infty (S^4 // S^1)$.

To complete the proof, we argue that strict lifts of $p \circ \phi$ and $\text{st}_{S^3} \circ \phi$ are equivalent to up-to-homotopy lifts. Ultimately, this follows from item (iii) in Prop. 3.17. by using a standard argument in homotopy theory. For completeness, let us recall how this works: homotopy commutative squares in a homotopical category

$$\begin{array}{ccc} X_1 & \xrightarrow{f} & Y_1 \\ p \downarrow & \swarrow & \downarrow q \\ X_2 & \xrightarrow{f} & Y_2 \end{array} \quad (48)$$

are equivalent to morphisms in the homotopy category of the corresponding arrow category. When working with a cofibrantly generated model category \mathcal{C} , the homotopy theory of the arrow category can be presented by the projective model structure on functors $\text{Fun}(\Delta[1], \mathcal{C})$ (see e.g. [Lu09, Sec. A.2.8]). General model category theory then implies that *strictly* commutative squares are equivalent to homotopy commutative squares (48) as soon as

- the source morphism $p: X_1 \rightarrow X_2$ is projectively cofibrant; and
- the target morphism $q: Y_1 \rightarrow Y_2$ is projectively fibrant.

In terms of maps of DG- $\mathbb{Q}[h_3]$ -modules, the fibrancy condition is always satisfied, whereas a map of DG-modules is projectively cofibrant if and only if it is a cofibration between cofibrant objects. In the case at hand, observe that minimal DG-modules are cofibrant, and the dotted morphism in (44) is a cofibration by Prop. 3.17 item (iii), as is the 6-truncation map of Ex. 2.48. Working with our rational models, this means that up-to-homotopy lifts of $p \circ \phi$ through τ_6 are equivalent to strict lifts, and likewise for lifts of $\text{st} \circ \phi$ through the stabilization of the adjunction unit. This completes the proof. \square

4 Gauge enhancement of M-branes

In this section, we explain how Theorem 3.19 provides a mechanism implementing gauge enhancement of fundamental M-branes. Firstly, in Sec. 4.1, we explain the construction without local supersymmetry taken into account. We present a solution to **Open Problem, rational version 1** within the framework of rational homotopy theory that explains the core of the gauge enhancement mechanism via double dimensional reduction of flux forms as in [MS04]. This focuses the problem on the enhancement from 6-truncated to untruncated twisted de Rham cohomology but falls short of exhibiting the existence and uniqueness of cocycles and their lifts, which requires local supersymmetry. Then, in Sec. 4.2, we pass to rational *super* homotopy theory to explain the gauge enhancement of super M-branes from first principles. To be self-contained, we first recall how super-spacetimes and super p -branes propagating in them are incarnated as super-cocycles in super-homotopy theory. We then explain how Theorem 3.19 implements gauge enhancement of fundamental M-branes under double dimensional reduction to type IIA D-branes in the rational approximation.

4.1 The mechanism without supersymmetry

We first discuss the aspects of the gauge enhancement mechanism that are already apparent in rational homotopy, disregarding for the moment the crucial interaction with local supersymmetry. Being that local supersymmetry is a necessary ingredient for the very existence and classification of fundamental p -branes and their charges, we will incorporate it into our discussion in Sec. 4.2 below. For the moment, however, we will simply assume that a collection of flux-like forms is provided to us on some spacetime without commenting on their origin in string or M-theory.

We take as input data a smooth manifold Y (to be thought of as our 11-dimensional background spacetime), equipped with a map to the 4-sphere

$$Y \xrightarrow{(G_4, G_7)} S^4, \quad (49)$$

to be thought of as measuring M-brane charge in Y . In the approximation of rational homotopy theory (Def. 2.1), the datum of such a map (49) is encoded by a differential 4-form G_4 and a differential 7-form G_7 on Y satisfying the relations

$$(G_4, G_7) \sim \begin{cases} dG_4 = 0 \\ dG_7 = -\frac{1}{2}G_4 \wedge G_4. \end{cases} \quad (50)$$

This follows from the equivalence Prop. 2.11, using the minimal DG-algebra model for the 4-sphere from Ex. 2.13. The relations (50) have the form of the “topological sector” of the equations of motion in 11-dimensional supergravity for the *C-field strength* G_4 and its dual flux G_7 , which we have identified with a rational homotopy map to the 4-sphere (this was first highlighted in [Sa13, Sec. 2.5]).

Observe that the contribution of the “dual flux” G_7 can be disentangled from that of the “C-field strength” G_4 by taking the homotopy pullback along the quaternionic Hopf fibration $H_{\mathbb{H}}: S^7 \rightarrow S^4$. This

is an $SU(2)$ -principal bundle over S^4 , hence classified by a map $\phi_{\mathbb{H}}: S^4 \rightarrow BSU(2)$ to the classifying space, as in (18). The cohomotopy cocycle (49) decomposes along the quaternionic Hopf fibration as follows:

$$\begin{array}{ccc}
 \widehat{Y} & \xrightarrow{G_7} & S^7 \\
 \downarrow & \text{(homotopy pullback)} & \downarrow H_{\mathbb{H}} \\
 Y & \xrightarrow{(G_4, G_7)} & S^4 \\
 \searrow G_4 & & \downarrow \phi_{\mathbb{H}} \\
 & & BSU(2) \\
 & \swarrow c_2 & \\
 & B^4\mathbb{Q}, &
 \end{array} \tag{51}$$

where $c_2: BSU(2) \rightarrow B^4\mathbb{Q}$ is the second Chern class of the universal $SU(2)$ -bundle. We are interested in the *double dimensional reduction* of these data, assuming that Y is a circle fibration as in (2). This means that we assume Y sits in a homotopy fiber sequence

$$\begin{array}{ccc}
 Y \simeq_{\text{whe}} \text{Ext}(X) & & \\
 \pi \downarrow & & \\
 X := Y // S^1 & \searrow & BS^1,
 \end{array} \tag{52}$$

hence that Y is an S^1 -extension of X as in Sec. 2.2. Consequently, our cohomotopy cocycle as in (51) is now exhibited equivalently as a map out of a space in the image of the Ext-functor (19):

$$\begin{array}{ccc}
 \text{Ext}(X) & \xrightarrow{(G_4, G_7)} & S^4 \\
 \searrow G_4 & & \swarrow c_2(\phi_{\mathbb{H}}) \\
 & B^4\mathbb{Q}. &
 \end{array}$$

By the adjunction of Theorem 2.44, maps out of the image of Ext are equivalent to maps into the image of the right adjoint Cyc, where this equivalence is exhibited by first applying Cyc and then precomposing with the adjunction unit map $\eta_X: X \rightarrow \text{Cyc Ext}(X)$. (e.g. [Bo94, Sec. 3]). In summary, this means that the cohomotopy cocycle on Y (with rational components (G_4, G_7)) is naturally identified with a $\text{Cyc}(S^4)$ -valued cocycle on X :

$$\begin{array}{ccc}
 & \text{double dimensional reduction} & \\
 & \text{of } (G_4, G_7) & \\
 X & \xrightarrow{(\widetilde{G_4, G_7})} & \text{Cyc}(S^4), \\
 \eta_X \downarrow & \text{Cyc}(G_4, G_7) \longrightarrow & \\
 \text{Cyc Ext}(X) & \longrightarrow & \\
 & \searrow & \swarrow \\
 & \text{Cyc}(B^4\mathbb{Q}) & \\
 & \downarrow & \\
 & B^3\mathbb{Q} &
 \end{array} \tag{53}$$

where the map $\text{Cyc}(B^4\mathbb{Q}) \rightarrow B^3\mathbb{Q}$ is obtained using the infinite loop space structure of $B^4\mathbb{Q}$; in terms of minimal DG-algebra models it is simply the map

$$\begin{array}{ccc}
 \text{Cyc}(B^4\mathbb{Q}) & \longrightarrow & B^3\mathbb{Q}. \\
 \omega_2 & & \\
 \omega_3 & \longleftarrow & \omega_3 \\
 \omega_4 & &
 \end{array}$$

Using the minimal DG-algebra model for $\text{Cyc}(S^4)$ (Ex. 2.47), we have that the Ext/Cyc-adjunct of the cohomotopy cocycle (G_4, G_7) is rationally determined by a collection of differential forms on X satisfying the following relations:

$$(\widetilde{G}_4, \widetilde{G}_7) = \begin{cases} dH_3 = 0, & dH_7 = F_2 \wedge F_6 - \frac{1}{2}F_4 \wedge F_4, \\ dF_2 = 0, \\ dF_4 = H_3 \wedge F_2, \\ dF_6 = H_3 \wedge F_4. \end{cases} \quad (54)$$

These differential forms can also be seen as arising from the data (G_4, G_7) via the Gysin sequence [MS04, Sec 4.2]. Interpreting G_4 and G_7 as M-brane flux forms in 11-dimensional supergravity, corresponding to the M2-brane and the M5-brane respectively, then the forms in (54) are naturally interpreted as flux forms of brane species in 10-dimensional type IIA supergravity, together with their Bianchi identities. We notice that not all of the expected brane flux forms appear: we have the NS1-brane flux H_3 , the RR-flux forms $F_{(2p+2 \leq 6)}$ for the D($2p \leq 4$)-branes, as well the NS5-brane flux H_7 ,⁸ but the flux forms for the D6 and D8 branes are missing.

So far we have merely recapitulated the mechanism of double dimensional reduction formalized via the Ext/Cyc-adjunction in rational homotopy theory [FSS13, FSS16b]. We are now interested in how the collection of flux forms in (54) may be naturally enhanced so as to contain flux forms F_8 and F_{10} . As discussed in Rem. 3.4, operations on spacetime Y should be accompanied by corresponding operations on the 4-sphere brane charge coefficient. This means that we should consider the A-type circle action on the 4-sphere:

$$\begin{array}{ccc} \text{Ext}(S^4 // S^1) \simeq_{\text{whe}} S^4 & & \\ \downarrow & & \\ S^4 // S^1 & & \\ \swarrow & & \\ BS^1 & & \end{array} \quad (55)$$

Diagram (53) can now be cast in a more symmetric form:

$$\begin{array}{ccccc} X & \xrightarrow{\quad (\widetilde{G}_4, \widetilde{G}_7) \quad} & S^4 // S^1 & & \\ \eta_X \downarrow & & \downarrow \eta_{S^4 // S^1} & & \\ \text{Cyc Ext}(X) & \xrightarrow{\quad \text{Cyc}(G_4, G_7) \quad} & \text{Cyc}(\text{Ext}(S^4 // S^1)) & & \\ & \searrow & \swarrow & & \\ & \text{Cyc}(B^4 \mathbb{Q}) & & & \\ & \downarrow & & & \\ & B^3 \mathbb{Q} & & & \end{array} \quad (56)$$

Diagram (56) poses the natural question of whether we can find a lift through the Ext/Cyc-unit:

$$\begin{array}{ccccc} & \xrightarrow{\quad \text{enhanced} \\ \quad \quad \quad \text{double dimensional reduction} \\ \quad \quad \quad \text{of } (G_4, G_7) \text{ ??} \quad} & & & & \\ X & \xrightarrow{\quad \quad \quad} & S^4 // S^1 & & \\ \eta_X \downarrow & & \downarrow \eta_{S^4 // S^1} & & \\ \text{Cyc Ext}(X) & \xrightarrow{\quad \text{Cyc}(G_4, G_7) \quad} & \text{Cyc}(\text{Ext}(S^4 // S^1)) & & \\ & \searrow \quad H_3 & \swarrow & & \\ & B^3 \mathbb{Q} & & & \end{array} \quad (57)$$

⁸The identity $dH_7 = F_2 \wedge F_6 - \frac{1}{2}F_4 \wedge F_4$ for the type IIA NS5-brane flux does not hold after fiberwise stabilization in (60)—indeed, the flux form H_7 is part of the obstruction to completing the zig-zag truncation map τ_6 in Ex. 2.48 to an actual homomorphism.

In general, there are strong obstructions to such a lift: Prop. 3.12 implies, via (40), that for such a lift to exist, the cocycle data in (54) needs to satisfy the conditions that $F_6 = 0$ and $H_3 \wedge F_2 = 0$ (so that $dF_4 = 0$). Once we take local supersymmetry into account, we will see that these conditions are too restrictive and that a direct lift as in (57) simply does not exist.

Remark 4.1 (Goodwillie calculus). At this point it behooves us to highlight that similarly to string theory and quantum field theory, homotopy theory also has a concept of *perturbative approximation*. As we have touched upon at varied points, homotopy theory is an immensely rich and computationally demanding area of mathematics (see [Ra03, HHR09] for good examples of this). However, in striking analogy to Taylor series expansions in differential calculus, mapping spaces between homotopy types can often be approximated by a sequence of increasingly accurate approximations called the *Goodwillie–Taylor tower* (see [Ku04] and [Lu17, Ch. 6]). The first-order approximation in this *Goodwillie perturbation theory* is provided by stabilization adjunction between spaces and spectra, as discussed in Sec. 2.1.

In view of Rem. 4.1, as we encounter strong obstructions in homotopy theory as in (57), we are led to ask whether the obstruction persists “perturbatively”, hence stage-wise in the Goodwillie–Taylor approximation. To first order in Goodwillie perturbation theory, this means that we should ask whether enhanced double dimensional reduction exists after *fiberwise stabilization* (Prop. 2.7) over $S^3 \simeq_{\mathbb{Q}} B^3\mathbb{Q}$:

$$\begin{array}{ccc}
 & \text{perturbatively enhanced} & \\
 & \text{double dimensional reduction} & \\
 & \text{of } (G_4, G_7) \text{ ??} & \\
 X & \xrightarrow{\quad} & \Omega_{S^3}^{\infty} \Sigma_{+, S^3}^{\infty} (S^4 // S^1) \\
 \eta_X \downarrow & & \downarrow \Omega_{S^3}^{\infty} \Sigma_{+, S^3}^{\infty} (\eta_{S^4 // S^1}) \\
 \text{Cyc Ext}(X) & \xrightarrow{\text{Cyc}(G_4, G_7)} & \text{Cyc}(S^4) \xrightarrow{\text{st}} \Omega_{S^3}^{\infty} \Sigma_{+, S^3}^{\infty} \text{Cyc}(S^4) \\
 & \searrow H_3 & \swarrow \\
 & & B^3\mathbb{Q}
 \end{array} \tag{58}$$

But this is precisely the question that is addressed by Theorem 3.19: the obstruction to a dotted morphism as in (58) is precisely the lift to untruncated twisted K-theory as in the diagram:

$$\begin{array}{ccc}
 & \text{perturbatively enhanced} & \\
 & \text{double dimensional reduction} & \\
 & \text{of } (G_4, G_7) \text{ !} & \\
 X & \xrightarrow{\widehat{(G_4, G_7)}} & \Omega_{B^3\mathbb{Q}}^{\infty-2} (\text{ku} // BS^1) \hookrightarrow \Omega_{S^3}^{\infty} \Sigma_{+, S^3}^{\infty} S^4 // S^1 \\
 \eta_X \downarrow & \searrow \widetilde{(G_4, G_7)} & \downarrow \Omega_{S^3}^{\infty} \Sigma_{+, S^3}^{\infty} (\eta_{S^4 // S^1}) \\
 \text{Cyc Ext}(X) & \xrightarrow{\text{Cyc}(G_4, G_7)} & \text{Cyc}(S^4) \xrightarrow{\text{st}} \Omega_{S^3}^{\infty} \Sigma_{+, S^3}^{\infty} \text{Cyc}(S^4) \\
 & \searrow H_3 & \swarrow \\
 & & B^3\mathbb{Q}
 \end{array} \tag{59}$$

That is, exhibiting a *perturbatively* enhanced double dimensional reduction cocycle is equivalent to specifying an extension of the cocycle data (54) to a cocycle of the form

$$\widehat{(G_4, G_7)} = \begin{cases} dH_3 = 0 \\ dF_2 = 0 \\ dF_4 = H_3 \wedge F_2 \\ dF_6 = H_3 \wedge F_4 \\ dF_8 = H_3 \wedge F_6 \\ dF_{10} = H_3 \wedge F_8 \\ \vdots \end{cases} \tag{60}$$

According to the discussion in Sec. 1, such a cocycle exhibits the full gauge enhancement mechanism in rational homotopy theory. That is,

A **solution to Open Problem, rational version 1** (p. 6) is obtained as follows:


- Double dimensional reduction of the M-theory flux data (G_4, G_7) of (50) is given by the Ext/Cyc-adjunct along the M-theory spacetime extension (52); and
- Its *perturbative gauge enhancement*, making all RR flux forms appear, is exhibited by lifting against the fiberwise stabilization of the Ext/Cyc-unit on the A-type orbispace of the 4-sphere (59)(60).

Remark 4.2 (Copies of twisted K-theory). Let us briefly pause to highlight a curious aspect of our result on gauge enhancement. By Prop. 3.13, there are in fact *two* direct summands of rationalized twisted K-theory inside the rational fiberwise stabilization of the A-type orbispace. Theorem 3.19 assigns a special role to the shifted copy $\Sigma^2 \text{ku} // BS^1$, since it is this copy that obstructs the lifting problem. The 2-fold suspension here reflects the fact that RR-flux forms in type IIA string theory ordinarily start with D0-flux F_2 in degree 2. But there is also a rational copy of *unshifted* twisted K-theory; a cocycle landing in this this copy corresponds to a twisted de Rham cocycle whose lowest component is a 0-form instead of a 2-form. The subtle issue of considering this possibility in the context of the K-theory classification of RR-flux is discussed in [MS03, MS04, BV05] and revisited systematically, and in a refined form, in [GS19].

4.2 The mechanism with local supersymmetry

In this section, we finally present our solution to the problem of gauge enhancement for super M-branes in its formulation as **Open Problem, rational version 2** (p. 8). We proceed as per the solution to **Open Problem, rational version 1** detailed in the previous section, with the important caveat that we now work in the proper context for super p -branes, namely locally supersymmetric supergeometry. Before presenting our solution, we briefly review some of the necessary background material (for a detailed introduction with an emphasis on the aspects of relevance here, see [Sc16]).

While there is no known mechanism in string theory that would enforce – or even prefer – *global* supersymmetry,⁹ what is predicted by perturbative string theory, and what is at the heart of its non-perturbative description as in [HSS18], is *local* supersymmetry, in particular *supergravity*. An elegant way to understand supergravity is via *Cartan geometry*, a powerful generalization of Riemannian geometry (see [ČS09, Ch. 1] for an introduction). In Cartan geometry, one fixes a local model space V and a subgroup $G \subset \text{Aut}(V)$ of its automorphism group, and then studies spaces that look *locally*, namely on the infinitesimal neighborhood of every point, like V equipped with a reduction of the structure group to G :




In the physics literature, this is known as the method of *local gauging via moving frames* (following [Ca23]), while mathematically this is the study of (*torsion-free*) G -structures [Gu65]. If one chooses the local model to be Minkowski spacetime

Local model for off-shell gravity	$\begin{array}{c} \text{SO}(d,1) \\ \curvearrowright \\ \mathbb{R}^{d,1} \end{array}$	(61)
--------------------------------------	---	------

then the resulting Cartan geometries are precisely the pseudo-Riemannian manifolds modelling configurations in general relativity. However, not all of these configurations are physically admissible (they are not all “on-shell”) and we need to impose the Einstein field equations to pick out the physically admissible

⁹Models of particle physics obtained from dimensional reductions of M-theory on singular manifolds of G_2 -holonomy (see [Ka17]) are among the globally supersymmetric extensions of the Standard Model of particle physics that are, so far, still consistent with experimental constraints [BGK18]. If and when supersymmetric extensions of the Standard Model are ruled out, then this will also rule out dimensional reductions of M-theory on singular fiber manifolds of G_2 -holonomy as realistic models for particle physics. However, this particular type of dimensional reduction is in no way dictated by the theory, and are certainly not generic amongst all possibilities, but were motivated by the expectation of global supersymmetry in the first place. What *is* dictated by the theory is *local* supersymmetry, which is already present as soon as fermions are.

geometries. In this Cartan-geometric formulation, the passage to supergravity is immediate: taking the local model space to be instead *super* Minkowski spacetime

Local model for off-shell supergravity	$\text{Spin}(d,1)$  $\mathbb{R}^{d,1 \mathbf{N}}$
---	---

(62)

then the resulting Cartan geometries are supergravity super-spacetimes. This is spelled out explicitly in [Lo90, EE12], and is implicit in much of the physics literature on supergravity, following [CDF91].

In the spirit of Kleinian geometry [Kl72], the group G acting on the local model space V reflects extra structure on V . For plain Minkowski spacetime (61) with its Lorentz group action, this extra structure is the Minkowski metric of special relativity. For the super-Minkowski spacetime (62) with its Lorentzian Spin group action, this structure is the translational *supersymmetry* super Lie algebra structure on $\mathbb{R}^{d,1|\mathbf{N}}$. Recall that the only non-trivial component of the super Lie bracket is the odd-odd spinor-to-vector pairing:

$$\begin{array}{ccc} \mathbf{N} \otimes \mathbf{N} & \xrightarrow{\overline{(-)}\Gamma(-)} & \mathbb{R}^{d,1} \\ \psi_1, \psi_2 \dagger & \longrightarrow & (\overline{\psi}_1 \Gamma^a \psi_2)_{a=0}^d . \end{array}$$

The Chevalley–Eilenberg algebra of this super-Minkowski super Lie algebra is a super DG-algebra

$$\text{CE}(\mathbb{R}^{d,1|\mathbf{N}}) := \left(\mathbb{R} \left[\underbrace{(e^a)_{a=0}^d}_{\text{deg}=(1,\text{even})}, \underbrace{(\psi^\alpha)_{\alpha=1}^{\mathbf{N}}}_{(1,\text{odd})} \right] / \left(\begin{array}{l} de^a = \overline{\psi} \Gamma^a \psi \\ d\psi^\alpha = 0 \end{array} \right) \right) \in \text{DGCSuperAlg}, \quad (63)$$

like those used in rational homotopy theory in Def. 2.9, but equipped with additional \mathbb{Z}_2 -grading. In view of the equivalence of Prop. 2.11 we may think of *super* DG-algebras of finite type as the formal duals of rational *superspaces* (see [HSS18, Sec. 3.2] for details in this context):

$$\text{Ho}(\text{SuperSpaces}_{\mathbb{R},\text{cn},\text{ft},\text{nil}}) := \text{Ho}(\text{DGCSuperAlg}_{\text{cn},\text{ft}}^{\text{op}}). \quad (64)$$

All of ordinary (henceforth “bosonic”) rational homotopy theory embeds into rational super homotopy theory by

- replacing the ground field by \mathbb{R} via $A \mapsto A \otimes_{\mathbb{Q}} \mathbb{R}$,¹⁰ and
- regarding ordinary DG-algebras as having an additional \mathbb{Z}_2 -grading in which all elements have trivial degree.


For instance, a morphism of the form

$$\begin{array}{ccc} \text{Spin}(d,1) & & \\ \downarrow \curvearrowright & & \\ \mathbb{R}^{d,1|\mathbf{N}} & \xrightarrow{\mu_p} & B^{p+1}S^1 \end{array}$$

in rational super homotopy theory is, equivalently, a $\text{Spin}(d,1)$ -invariant super Lie algebra cocycle of $\mathbb{R}^{p,1|\mathbf{N}}$ in degree $p+2$. The requirement of $\text{Spin}(d,1)$ -invariance forces such a cocycle to be a product of elements of the form

$$\mu_p := (\overline{\psi} \Gamma_{a_1 \dots a_p} \psi) \wedge e^{a_1} \wedge \dots \wedge e^{a_p} \in \text{CE}(\mathbb{R}^{d,1|\mathbf{N}}).$$

Remarkably, when the local model space is taken to be the $D=11, \mathcal{N}=1$ super-Minkowski spacetime

Local model for <i>on-shell</i> $D=11, \mathcal{N}=1$ supergravity	$\text{Spin}(10,1)$  $\mathbb{R}^{10,1 \mathbf{32}}$
---	--

the condition of (super-)torsion freeness is already equivalent to the supergravity equations of motion [CL94, Ho97, FOS17]. That is, (super-)torsion-free $D=11, \mathcal{N}=1$ (super-)Cartan geometries are precisely on-shell configurations of supergravity. In fact, more is true: the supergravity equations of motion imply that the bifermionic components of the supergravity C -field curvature 4-form G_4 and its

¹⁰From the point of view of homotopy theory, there is little difference between working over \mathbb{Q} or over \mathbb{R} .

Hodge dual 7-form G_7 are covariantly constant on each super tangent space $\mathbb{R}^{10,1|32}$ and constrained there to be of the form

$$(G_4, G_7)_{\text{fermionic}} = \left(\begin{array}{l} \mu_{M2} := \frac{i}{2} (\overline{\Psi} \Gamma_{a_1 a_2} \Psi) \wedge E^{a_1} \wedge E^{a_2}, \\ \mu_{M5} := \frac{1}{5!} (\overline{\Psi} \Gamma_{a_1 \dots a_5} \Psi) \wedge E^{a_1} \wedge \dots \wedge E^{a_5} \end{array} \right). \quad (65)$$


In this expression, (E^a, Ψ^α) is the *super-vielbein* field on super-spacetime, hence the supergeometric analog of the vielbein field determining a Cartan geometry. Torsion-freeness implies that on the infinitesimal neighborhood of each point of super-spacetime, the super-vielbein reproduces the super left-invariant 1-form generators (e^a, ψ^α) of $\text{CE}(\mathbb{R}^{10,1|32})$ as in (63). Due to the *Fierz identities* of Spin-representation theory [D'AF82], the super-forms of (65) satisfy the relations

$$\mu_{M2/M5} = \begin{cases} d\mu_{M2} = 0 \\ d\mu_{M5} = -\frac{1}{2} \mu_{M2} \wedge \mu_{M2} \end{cases} \quad (66)$$


on each super tangent space $\mathbb{R}^{10,1|32}$. Moreover, this is precisely the super tangent space-wise condition that witnesses propagation of fundamental super M2-branes [BST87, (15)] and the fundamental super M5-branes [LT96, (5)], [BLN⁺97, (6)] in the background super-spacetime (see [HSS18, Sec. 2.1] for further background on fundamental super p -branes).

After comparing the M2/M5-brane Fierz identity (66) with the minimal DG-algebra model for the 4-sphere (Ex. 2.13, also (50)), we may summarize the state of affairs using the language of (rational) super homotopy theory as follows:

11-dimensional super-Minkowski spacetime carries an exceptional super-cocycle in rational cohomotopy of degree four [FSS16a, FSS15b], and 11-dimensional supergravity super-spacetimes together with fundamental M2/M5-branes propagating in them are the *higher* Cartan geometries [Sc15, We17] locally modelled on the higher geometric data:

Local model for fundamental M2/M5-branes in $D = 11, \mathcal{N} = 1$ supergravity	$\text{Spin}(10,1)$  $\mathbb{R}^{10,1 32} \xrightarrow{\mu_{M2/M5}} S^4$	(67)
--	---	------

Analogous statements hold for all fundamental branes that appear in string theory [FSS13]. In particular, the super-spacetimes of $D = 10, \mathcal{N} = (1, 1)$ supergravity (that is, type IIA supergravity) are Cartan geometries locally modelled on the type IIA super-Minkowski spacetime

Local model for $D = 10, \mathcal{N} = (1, 1)$ supergravity	$\text{Spin}(9,1)$  $\mathbb{R}^{9,1 16+16}$
--	--

The presence of the fundamental F1- and Dp -branes propagating in these super-spacetimes is captured, as for the M2/M5-brane cocycles above, by the non-trivial super Lie algebra cocycles [CGN⁺97, (3.9)]:

$$\begin{aligned} \mu_{F1}^{\text{IIA}} &= i (\overline{\Psi} \Gamma_a \Gamma_{10} \Psi) \wedge E^a \\ \mu_{D0} &= (\overline{\Psi} \Gamma_{10} \Psi) \\ \mu_{D2} &= \frac{i}{2} (\overline{\Psi} \Gamma_{a_1 a_2} \Psi) \wedge E^{a_1} \wedge E^{a_2} \\ \mu_{D4} &= \frac{1}{4!} (\overline{\Psi} \Gamma_{a_1 \dots a_4} \Gamma_{10} \Psi) \wedge E^{a_1} \wedge \dots \wedge E^{a_4} \\ \mu_{D6} &= \frac{i}{6!} (\overline{\Psi} \Gamma_{a_1 \dots a_6} \Psi) \wedge E^{a_1} \wedge \dots \wedge E^{a_6} \\ \mu_{D8} &= \frac{1}{8!} (\overline{\Psi} \Gamma_{a_1 \dots a_8} \Gamma_{10} \Psi) \wedge E^{a_1} \wedge \dots \wedge E^{a_8} \\ \mu_{D10} &= \frac{i}{10!} (\overline{\Psi} \Gamma_{a_1 \dots a_{10}} \Psi) \wedge E^{a_1} \wedge \dots \wedge E^{a_{10}} \\ (\mu_{NS5}) &= \frac{1}{5!} (\overline{\Psi} \Gamma_{a_1 \dots a_5} \Psi) \wedge E^{a_1} \wedge \dots \wedge E^{a_5}, \end{aligned} \quad (68)$$

where we have also included here the NS5-brane cocycle μ_{NS5} for completeness. Using the Fierz identities for $\text{Spin}(9, 1)$ -representations, one finds that these expressions satisfy the following relations:

$$(\mu_{F1/Dp}) = \begin{cases} d\mu_{F1} = 0 & (d\mu_{NS5} = \mu_{D0} \wedge \mu_{D4} - \frac{1}{2} \mu_{D2} \wedge \mu_{D2}) \\ d\mu_{D(2p+4)} = \mu_{F1} \wedge \mu_{D(p+2)}, \end{cases} \quad (69)$$

([CGN⁺97, App. A], [CAI⁺00, (6.8), (6.9)], see also [FSS16a, Theorem 4.16], [FSS16b, Prop. 4.8]). This is the supersymmetric version of the cocycle (60) that we had encountered in the previous section.

By analogy with (67), using Lemma 2.31 we may concisely summarize this in the language of rational super homotopy theory as follows:

10-dimensional type IIA super-Minkowski spacetime carries an exceptional cocycle in rational 2-shifted twisted connective K-theory and super-spacetimes of 10-dimensional type IIA supergravity together with fundamental F1/D p -branes propagating inside them are the *higher* Cartan geometries locally modelled on the higher geometric data:

<p style="text-align: center;">Local model for fundamental F1/Dp-branes in $D = 10$, $\mathcal{N} = (1, 1)$ supergravity</p>	$ \begin{array}{ccc} & \text{Spin}(9,1) & \\ & \curvearrowright & \\ \mathbb{R}^{9,1 \mathbf{16}+\overline{\mathbf{16}}} & \xrightarrow{\mu_{F1/Dp}} & \text{ku} // BS^1 \\ & \searrow \mu_{F1} & \swarrow \\ & B^3\mathbb{Q} & \end{array} $ (70)
---	--

Working with higher Cartan geometry in this way, one finds large parts of the string/M-theory literature appearing in cocycle incarnations (see [HSS18, Sec. 2] for a more detailed account). The upshot is that all information about brane species and their behavior in string/M-theory is encoded in *cohomological data*, depending on the local model space. The actual brane dynamics are provided by the higher-super-Cartan-geometric globalization of these local data.

The problem of gauge enhancement for M-branes, therefore, reduces to the question of how double dimensional reduction turns the local model (67) for $D = 11$, $\mathcal{N} = 1$ superspacetime with its M2/M5-brane cocycle into the local model (70) for $D = 10$, $\mathcal{N} = (1, 1)$ super-spacetime with its unified D-brane cocycle. This works via the mechanism of Sec. 4.1: our input datum (49) is now specifically the fundamental M2/M5-brane super cocycle (67)

$$\mathbb{R}^{10,1|\mathbf{32}} \xrightarrow{\mu_{M2/M5}} S^4. \quad (71)$$

The supersymmetric version of the spacetime circle extension (52) is the extension of type IIA super-Minkowski spacetime to $D = 11$ super-Minkowski spacetime classified by the D0-brane cocycle [FSS13, Prop. 4.5] (this extension implements “D0-brane condensation”):

$$\begin{array}{ccc}
 \mathbb{R}^{10,1|\mathbf{32}} & \simeq & \text{Ext} \left(\mathbb{R}^{9,1|\mathbf{16}+\overline{\mathbf{16}}} \right) \\
 \downarrow & & \\
 \mathbb{R}^{9,1|\mathbf{16}+\overline{\mathbf{16}}} & & \\
 & \searrow \mu_{D0} & \\
 & & BS^1.
 \end{array} \quad (72)$$

Thus, (71) can be recast in the form

$$\begin{array}{ccc}
 \text{Ext} \left(\mathbb{R}^{9,1|\mathbf{16}+\overline{\mathbf{16}}} \right) & \xrightarrow{\mu_{M2/M5}} & S^4, \\
 & \searrow \mu_{M2} & \swarrow \\
 & B^4\mathbb{Q} &
 \end{array}$$

so that its double dimensional reduction is given as in (53) by the Ext/Cyc-adjunct as in the following

diagram:

$$\begin{array}{ccc}
& \text{double dimensional reduction} \\
& \text{of } \mu_{M2/M5} \\
\mathbb{R}^{9,1|16+\overline{16}} & \xrightarrow{\tilde{\mu}_{M2/M5}} & \text{Cyc}(S^4) \\
\eta \downarrow & & \\
\text{CycExt}(\mathbb{R}^{9,1|16+\overline{16}}) & \xrightarrow{\text{Cyc}(\mu_{M2/M5})} & \text{Cyc}(S^4) \\
& \searrow & \swarrow \\
& \text{Cyc}(B^4\mathbb{Q}) & \\
& \downarrow & \\
& B^3\mathbb{Q} &
\end{array} \quad (73)$$

In [FSS16a, Prop. 3.8] and [FSS16b, Theorem 3.8], this double dimensional reduction cocycle $\tilde{\mu}_{M2/M5}$ was computed as:

$$\tilde{\mu}_{M2/M5} = \begin{cases} d\mu_{F1} = 0 & d\mu_{NS5} = \mu_{D0} \wedge \mu_{D4} - \frac{1}{2}\mu_{D2} \wedge \mu_{D2} \cdot \\ d\mu_{D0} = 0 & \\ d\mu_{D2} = \mu_{F1} \wedge \mu_{D0} & \\ d\mu_{D4} = \mu_{F1} \wedge \mu_{D2} \cdot & \end{cases} \quad (74)$$

We, therefore, obtain the truncation of (68), (69) that contains the fundamental F1-brane cocycle μ_{F1} , as well as the fundamental Dp-brane cocycles μ_{Dp} for $p \in \{0, 2, 4\}$, together with Bianchi identities. We also obtain the NS5-brane cocycle μ_{NS5} . To enhance this double dimensional reduction picture, we could ask for a lift as in (57):

$$\begin{array}{ccc}
& \text{enhanced} \\
& \text{double dimensional reduction} \\
& \text{of } \mu_{M2/M5} \text{ ??} \\
X & \dashrightarrow & S^4 // S^1 \\
\eta_{\mathbb{R}^{9,1|16+\overline{16}}} \downarrow & & \downarrow \eta_{S^4 // S^1} \\
\text{CycExt}(\mathbb{R}^{9,1|16+\overline{16}}) & \xrightarrow{\text{Cyc}(\mu_{M2/M5})} & \text{Cyc}(\text{Ext}(S^4 // S^1)) \\
& \searrow \mu_{F1} & \swarrow \\
& B^3\mathbb{Q} &
\end{array} \quad (75)$$

But a dashed lift in (75) does *not* exist: Prop. 3.12 requires that for such a lift to exist, the double dimensionally reduced cocycle data in (74) needs to satisfy the extra conditions

- (i) $\mu_{D4} = 0$; and
- (ii) $\mu_{F1} \wedge \mu_{D0} = 0$,

both of which fail, the first by (68) the second by (69). If we now approach the problem with homotopy theoretic perturbation theory (Rem. 4.1), we might instead ask for a *first-order* lift as in (58), hence a lift in the following diagram:

$$\begin{array}{ccc}
& \text{perturbatively enhanced} \\
& \text{double dimensional reduction} \\
& \text{of } \mu_{M2/M5} \text{ ??} \\
\mathbb{R}^{9,1|16+\overline{16}} & \dashrightarrow & \Omega_{S^3}^\infty \Sigma_{+,S^3}^\infty(S^4 // S^1) \\
\eta_{\mathbb{R}^{9,1|16+\overline{16}}} \downarrow & & \downarrow \Omega_{S^3}^\infty \Sigma_{+,S^3}^\infty(\eta_{S^4 // S^1}) \\
\text{CycExt}(\mathbb{R}^{9,1|16+\overline{16}}) & \xrightarrow{\text{Cyc}(\mu_{M2/M5})} & \text{Cyc}(S^4) \xrightarrow{\text{st}} \Omega_{S^3}^\infty \Sigma_{+,S^3}^\infty \text{Cyc}(S^4) \\
& \searrow \mu_{F1} & \swarrow \\
& B^3\mathbb{Q} &
\end{array} \quad (76)$$

As was the case for (59), Theorem 3.19 says that the lift in (76) exists precisely if the lift $\widehat{\mu}_{M2/M5}$ in the following diagram exists:

$$\begin{array}{ccc}
& \text{perturbatively enhanced} \\
& \text{double dimensional reduction} \\
& \text{of } \mu_{M2/M5} \text{ !} \\
\mathbb{R}^{9,1|16+\overline{16}} & \xrightarrow{\widehat{\mu}_{M2/M5}} & \Omega_{B^3\mathbb{Q}}^{\infty-2}(\text{ku} // BS^1) \xrightarrow{\iota} \Omega_{S^3\Sigma_{+,S^3}}^{\infty}(S^4 // S^1) \\
\downarrow \eta_{\mathbb{R}^{9,1|16+\overline{16}}} & \searrow \widetilde{\mu}_{M2/M5} & \downarrow \Omega_{S^3\Sigma_{+,S^3}}^{\infty}(\eta_{S^4 // S^1}) \\
\text{Cyc Ext}(\mathbb{R}^{9,1|16+\overline{16}}) & \xrightarrow{\text{Cyc}(\mu_{M2/M5})} & \text{Cyc}(S^4) \xrightarrow{\text{st}} \Omega_{S^3\Sigma_{+,S^3}}^{\infty}\text{Cyc}(S^4) \\
& \searrow \mu_{F1} & \downarrow \tau_6 \\
& & B^3\mathbb{Q}
\end{array} \tag{77}$$

As in (60), such a lift $\widehat{\mu}_{M2/M5}$ is equivalent to specifying an extension of the truncated data in (74) to a cocycle:

$$\widehat{\mu}_{M2/M5} = \begin{cases} d\mu_{F1} = 0 \\ d\mu_{D0} = 0 \\ d\mu_{D2} = \mu_{F1} \wedge \mu_{D0} \\ d\mu_{D4} = \mu_{F1} \wedge \mu_{D2} \\ d\mu_{D6} = \mu_{F1} \wedge \mu_{D4} \\ d\mu_{D8} = \mu_{F1} \wedge \mu_{D6} \\ d\mu_{D10} = \mu_{F1} \wedge \mu_{D8} \end{cases} \tag{78}$$

(after discarding the NS5-brane cocycle μ_{NS5}). By (68) and (69), such an extension exists and is precisely the required enhancement of the double dimensional reduction of the fundamental M2/M5-brane cocycle by the missing D($p \geq 6$)-brane cocycles to the full F1/D p -brane cocycle of type IIA string theory with coefficients in rational twisted K-theory. This is exactly the required gauge enhancement:

A solution to Open Problem, rational version 2 (p. 8) is given as follows:

- While double dimensional reduction of the fundamental M2/M5-brane cohomotopy 4-cocycle $\mu_{M2/M5}$ (66) is obtained as the Ext/Cyc-adjunct (74) along the M-theoretic super-spacetime extension (72);
- Its *perturbative gauge enhancement*, making the full combined F1/D p -brane cocycle appear is obtained by lifting through the fiberwise stabilization of the Ext/Cyc-unit on the A-type orbispace of the 4-sphere (76).

Remark 4.3 (Higher-dimensional branes). An often neglected point is that, with local supersymmetry taken into account, the system of relations (78) is indeed non-trivial up to the degree shown. The purely bosonic part of the D8-flux form F_{10} is necessarily closed (being a 10-form on a 10-dimensional manifold). However, its fermionic component which is proportional to $(\overline{\Psi}\Gamma_{a_1 \dots a_8}\Gamma_{10}\Psi) \wedge E^{a_1} \wedge \dots \wedge E^{a_8}$ is only of bosonic degree 8, so that its differential as a super-form need not vanish (and, indeed, does not vanish [CAI⁺00, (6.9)]). For the same reason, we include in (68) and (78) the non-trivial component μ_{D10} which ought to correspond to a D10-brane, were it not for the fact that there is no bosonic aspect of a 10-brane in a 10-dimensional spacetime. One way to see the ‘‘D10-brane contribution’’ μ_{D10} arise is to consider the type IIB D-brane super cocycles $\mu_{D1}, \mu_{D3}, \mu_{D5}, \mu_{D7}, \mu_{D9}$ [Sa00, Sec. 2] and then apply super-geometric T-duality [FSS16b, Theorem 5.3]. The existence of the charge structure for would-be D10 branes was also noticed in [CS09, p. 30] by different means.

Acknowledgements. We are grateful to Augustí Roig and Martintxo Saralegi-Aranguren for discussion of [RS00], as well as to David Corfield, Ted Erler, Domenico Fiorenza, and David Roberts for useful comments. We also thank the anonymous referee for their careful reading and helpful suggestions. VBM acknowledges partial support of SNF Grant No. 200020_172498/1. This research was partly supported by the NCCR SwissMAP, funded by the Swiss National Science Foundation, and by the COST Action MP1405 QSPACE, supported by COST (European Cooperation in Science and Technology).

References

- [AG04] B. S. Acharya and S. Gukov, *M theory and Singularities of Exceptional Holonomy Manifolds*, Phys. Rept. **392** (2004), 121-189, [[arXiv:hep-th/0409191](#)].
- [AETW87] A. Achúcarro, J. Evans, P. Townsend, and D. Wiltshire, *Super p-Branes*, Phys. Lett. **B 198** (1987), 441-446, [[spire:22286](#)].
- [ABJM08] O. Aharony, O. Bergman, D. Jafferis, and J. Maldacena, *$N = 6$ superconformal Chern-Simons-matter theories, M2-branes and their gravity duals*, J. High Energy Phys. **0810** (2008), 091, [[arXiv:0806.1218](#)].
- [AGM⁺99] O. Aharony, S. S. Gubser, J. Maldacena, H. Ooguri, and Y. Oz, *Large N Field Theories, String Theory and Gravity*, Phys. Rept. **323** (2000), 183-386, [[arXiv:hep-th/9905111](#)].
- [ABG10] M. Ando, A. Blumberg, and D. Gepner, *Twists of K-theory and TMF*, in *Superstrings, geometry, topology, and C*-algebras*, 27–63, Proc. Symp. Pure Math. 81, Amer. Math. Soc., Providence, RI 2010 [[arXiv:1002.3004](#)].
- [ABG⁺14] M. Ando, A. Blumberg, D. Gepner, M. J. Hopkins, and C. Rezk, *An ∞ -categorical approach to R-line bundles, R-module Thom spectra, and twisted R-homology*, J. Topol. **7** (2014), 869-893, [[arXiv:1403.4325](#)].
- [AS05] M. Atiyah and G. Segal, *Twisted K-theory and cohomology*, In *Inspired By SS Chern*, 5–43, Nankai Tracts Math., 11, World Sci. Publ., Hackensack, NJ, 2006, [[arXiv:math/0510674](#)].
- [D'AF82] R. D'Auria and P. Fré, *Geometric supergravity in $D = 11$ and its hidden supergroup*, Nucl. Phys. **B 201** (1982), 101–140, [[ncatlab.org/nlab/files/GeometricSupergravity.pdf](#)].
- [AT89] J. de Azcárraga and P. Townsend, *Superspace geometry and the classification of supersymmetric extended objects*, Phys. Rev. Lett. **62** (1989), 2579-2582, [[spire:284635](#)].
- [BL08] J. Bagger and N. D. Lambert, *Gauge symmetry and supersymmetry of multiple M2- branes*, Phys. Rev. D **77** (2008) 065008, [[arXiv:0711.0955](#)] [[hep-th](#)].
- [BLM⁺13] J. Bagger, N. Lambert, S. Mukhi, and C. Papageorgakis, *Multiple Membranes in M-theory*, Phys. Rep. **527** (2013), 1-100, [[arXiv:1203.3546](#)].
- [BLN⁺97] I. Bandos, K. Lechner, A. Nurmagambetov, P. Pasti, D. Sorokin, and M. Tonin, *Covariant action for the super-five-brane of M-theory*, Phys. Rev. Lett. **78** (1997) 4332-4334, [[arXiv:hep-th/9701149](#)].
- [BBS06] K. Becker, M. Becker, J. Schwarz, *String theory and M-theory: a modern introduction*, Cambridge University Press, 2006.
- [BS03] N. Berkovits and M. Schnabl, *Yang-Mills Action from Open Superstring Field Theory*, J. High Energy Phys. **0309** (2003), 022, [[arXiv:hep-th/0307019](#)].
- [BV05] A. Bergman and U. Varadarajan, *Loop Groups, Kaluza-Klein Reduction and M-Theory*, J. High Energy Phys. **0506** (2005), 043, [[arXiv:hep-th/0406218](#)].
- [BRG⁺96] E. Bergshoeff, M. de Roo, M. Green, G. Papadopoulos, and P. Townsend, *Duality of Type II 7-branes and 8-branes*, Nucl. Phys. **B470** (1996), 113-135, [[arXiv:hep-th/9601150](#)].
- [BST87] E. Bergshoeff, E. Sezgin, and P. K. Townsend, *Supermembranes and eleven-dimensional supergravity*, Phys. Lett. **B189** (1987), 75-78, [[spire:248230](#)].
- [Bl17] A. Blumberg *Equivariant homotopy theory*, lecture notes 2017, [[github.com/adebray/equivariant_homotopy_theory](#)].
- [Bo94] F. Borceux, *Basic Category Theory*, Vol. 1 of *Handbook of Categorical Algebra*, Cambridge University Press, 1995.
- [BG76] A. Bousfield and V. Gugenheim, *On PL deRham theory and rational homotopy type*, Memoirs of the AMS, vol. 179, Amer. Math. Soc., Providence, RI, 1976.
- [BCM⁺02] P. Bouwknegt, A. L. Carey, V. Mathai, M. K. Murray, and D. Stevenson, *Twisted K-theory and K-theory of bundle gerbes*, Commun. Math. Phys. **228** (2002), 17-49, [[arXiv:hep-th/0106194](#)].
- [BEV03] P. Bouwknegt, J. Evslin, and V. Mathai, *T-Duality: Topology Change from H-flux*, Commun. Math. Phys. **249** (2004), 383-415, [[arXiv:hep-th/0306062](#)].

- [BM00] P. Bouwknegt and V. Mathai, *D-branes, B-fields and twisted K-theory*, J. High Energy Phys. **0003** (2000), 007, [arXiv:hep-th/0002023].
- [BM18] V. Braunack-Mayer, *Rational parametrised stable homotopy theory*, PhD thesis, Zurich University, 2018, [ncatlab.org/schreiber/show/thesis+Braunack-Mayer].
- [BM19a] V. Braunack-Mayer, *Strict algebraic models for rational parametrised spectra I*, in preparation.
- [BM19b] V. Braunack-Mayer, *Strict algebraic models for rational parametrised spectra II*, in preparation.
- [BFR13] R. Brunetti, K. Fredenhagen, and K. Rejzner, *Quantum gravity from the point of view of locally covariant quantum field theory*, Commun. Math. Phys. **345** (2016), 741-779, [arXiv:1306.1058].
- [BN14] U. Bunke and T. Nikolaus, *Twisted differential cohomology*, [arXiv:1406.3231].
- [BGK18] M. Buschmann, E. Gonzalez, and G. L. Kane, *Revisiting Gluinos at LHC*, [arXiv:1803.04394].
- [CS09] A. K. Callister and D. J. Smith, *Topological charges in $SL(2, \mathbb{R})$ covariant massive 11-dimensional and Type IIB SUGRA*, Phys. Rev. **D80** (2009), 125035, [arXiv:0907.3614].
- [CL94] A. Candiello and K. Lechner, *Duality in supergravity theories*, Nucl. Phys. **B412** (1994) 479-501, [arXiv:hep-th/9309143].
- [ČS09] A. Čap and J. Slovák, *Parabolic Geometries I: Background and General Theory*, Amer. Math. Soc., Providence, RI, 2009.
- [Ca23] É. Cartan, *Sur les variétés à connexion affine et la théorie de la relativité généralisée (première partie)* Ann. scient. de l'Ecole Normale Supérieure, Sér. 3, **40** (1923), 325-412.
- [CDF91] L. Castellani, R. D'Auria, and P. Fré, *Supergravity and Superstrings – A geometric perspective*, World Scientific, Singapore, 1991.
- [CGN⁺97] M. Cederwall, A. von Gussich, B.E.W. Nilsson, P. Sundell, and A. Westerberg, *The Dirichlet Super-p-Branes in Ten-Dimensional Type IIA and IIB Supergravity*, Nucl. Phys. **B490** (1997), 179-201, [hep-th/9611159].
- [CS00] A. H. Chamseddine and W. A. Sabra, *$D = 7$ $SU(2)$ gauged supergravity from $D = 10$ supergravity*, Phys. Lett. **B 476** (2000), 415-419, [arXiv:hep-th/9911180].
- [CAI⁺00] C. Chryssomalakos, J. de Azcárraga, J. Izquierdo, and C. Pérez Bueno, *The geometry of branes and extended superspaces*, Nucl. Phys. **B 567** (2000), 293-330, [arXiv:hep-th/9904137].
- [ClayMP] Clay Mathematics Institute, *Millennium Problems – Yang–Mills and Mass Gap* [claymath.org/millennium-problems/yang-mills-and-mass-gap].
- [CST03] E. Coletti, I. Sigalov, and W. Taylor, *Abelian and nonabelian vector field effective actions from string field theory*, J. High Energy Phys. **0309** (2003), 050, [arXiv:hep-th/0306041].
- [CJ98] M. Crabb and I. James, *Fibrewise homotopy theory*, Springer-Verlag London, Ltd., London, 1998.
- [CLL⁺00] M. Cvetič, J. T. Liu, H. Lü, and C. N. Pope, *Domain-wall supergravities from sphere reduction*, Nucl. Phys. **B 560** (1999), 230-256, [arXiv:hep-th/0005137].
- [CLP00] M. Cvetič, H. Lü, and C. N. Pope, *Consistent Kaluza-Klein sphere reductions*, Phys. Rev. **D 62** (2000) 064028, [arXiv:hep-th/0003286].
- [CLP⁺00] M. Cvetič, H. Lü, C. N. Pope, A. Sadrzadeh, and T. A. Tran, *S^3 and S^4 reductions of type IIA supergravity*, Nucl. Phys. **B 590** (2000), 233-251, [arXiv:hep-th/0005137].
- [DZH⁺15] M. Del Zotto, J. Heckman, A. Tomasiello, and C. Vafa, *6d Conformal Matter*, J. High Energy Phys. **2015** (2015), 54, [arXiv:1407.6359].
- [DMW03] D. Diaconescu, G. Moore, and E. Witten, *E_8 -gauge theory and a derivation of K-theory from M-theory*, Adv. Theor. Math. Phys **6** (2003), 1031–1134, [arXiv:hep-th/0005090].
- [DFM09] J. Distler, D. Freed, and G. Moore, *Orientifold Précis* in: H. Sati, U. Schreiber (eds.), *Mathematical Foundations of Quantum Field and Perturbative String Theory*, Proc. Symp. Pure Math., AMS, 2011, [arXiv:0906.0795].
- [Do95] J. F. Donoghue, *Introduction to the Effective Field Theory Description of Gravity*, [arXiv:gr-qc/9512024].
- [Du94] M. Duff, *Kaluza-Klein Theory in Perspective*, Proc. of the Symposium: *The Oskar Klein Centenary*, World Scientific, Singapore, 1994, [arXiv:hep-th/9410046].

- [Du99] M. Duff (ed.) *The World in Eleven Dimensions: Supergravity, Supermembranes and M-theory*, IoP, Bristol, 1999.
- [DIP⁺88] M. Duff, T. Inami, C. Pope, E. Sezgin, and K. Stelle, *Semiclassical Quantization of the Supermembrane*, Nucl. Phys. **B297** (1988), 515-538, [[spire:247064](#)].
- [DHI⁺87] M. Duff, P. Howe, T. Inami, and K. Stelle, *Superstrings in $D = 10$ from Supermembranes in $D = 11$* , Phys. Lett. **B191** (1987), 70-74 (reprinted in [Du99]), [[spire:245249](#)].
- [EE12] M. Egeileh and F. El Chami, *Some remarks on the geometry of superspace supergravity*, J. Geom. Phys. **62** (2012), 53-60.
- [Ev06] J. Evslin, *What Does(n't) K-theory Classify?*, Second Modave Summer School in Mathematical Physics [[arXiv:hep-th/0610328](#)].
- [ES06] J. Evslin and H. Sati, *Can D-Branes Wrap Nonrepresentable Cycles?*, J. High Energy Phys. **0610** (2006), 050, [[arXiv:hep-th/0607045](#)].
- [Fa17] M. Fazzi, *Higher-dimensional field theories from type II supergravity*, [[arXiv:1712.04447](#)].
- [FOT08] Y. Félix, J. Oprea, and D. Tanré, *Algebraic models in geometry*, Oxford University Press, Oxford, 2008.
- [FOS17] J. Figueroa-O'Farrill and A. Santi, *Spencer cohomology and eleven-dimensional supergravity*, Comm. Math. Phys. **349** (2017), 627-660, [[arXiv:1511.08737](#)].
- [FOS02a] J. Figueroa-O'Farrill and J. Simón, *Supersymmetric Kaluza-Klein reductions of M2 and M5-branes*, Adv. Theor. Math. Phys. **6** (2003), 703-793, [[arXiv:hep-th/0208107](#)].
- [FOS02b] J. Figueroa-O'Farrill and J. Simón, *Supersymmetric Kaluza-Klein reductions of M-waves and MKK-monopoles*, Class. Quant. Grav. **19** (2002), 6147-6174, [[hep-th/0208108](#)].
- [FOS04] J. Figueroa-O'Farrill and J. Simón, *Supersymmetric Kaluza-Klein reductions of AdS backgrounds*, Adv. Theor. Math. Phys. **8** (2004), 217-317, [[arXiv:hep-th/0401206](#)].
- [FSS13] D. Fiorenza, H. Sati, and U. Schreiber, *Super Lie n -algebra extensions, higher WZW models, and super p -branes with tensor multiplet fields*, Intern. J. Geom. Meth. Mod. Phys. **12** (2015) 1550018, [[arXiv:1308.5264](#)].
- [FSS15b] D. Fiorenza, H. Sati, and U. Schreiber, *The WZW term of the M5-brane and differential cohomotopy*, J. Math. Phys. **56** (2015), 102301 [[arXiv:1506.07557](#)].
- [FSS16a] D. Fiorenza, H. Sati, and U. Schreiber, *Rational sphere valued supercocycles in M-theory and type IIA string theory*, J. Geom. Phys. **114** (2017) 91-108 [[arXiv:1606.03206](#)].
- [FSS16b] D. Fiorenza, H. Sati, and U. Schreiber, *T-Duality from super Lie n -algebra cocycles for super p -branes*, Adv. Theor. Math. Phys. **22** (2018) no. 5, [[arXiv:1611.06536](#)].
- [FSS17] D. Fiorenza, H. Sati, and U. Schreiber, *T-duality in rational homotopy theory via L_∞ -algebras*, Geom. Topol. Math. Phys. **1** (2018), special volume in tribute of Jim Stasheff and Dennis Sullivan, [[arXiv:1712.00758](#)] [[math-ph](#)].
- [FSS18] D. Fiorenza, H. Sati, and U. Schreiber, *Higher T-duality of M-branes*, [[arXiv:1803.05634](#)].
- [FSS19] D. Fiorenza, H. Sati, and U. Schreiber, *The rational higher structure of M-theory*, Proceedings of Higher Structures in M-Theory, Durham Symposium 2018, Fortsch. Phys. 2019
- [Fr00] D. Freed, *Dirac charge quantization and generalized differential cohomology*, Surveys in Differential Geometry, Int. Press, Somerville, MA, 2000, pp. 129-194, [[arXiv:hep-th/0011220](#)].
- [FH00] D. Freed and M. Hopkins, *On Ramond-Ramond fields and K-theory*, J. High Energy Phys. 0005 (2000) 044 [[arXiv:hep-th/0002027](#)].
- [FW99] D. Freed and E. Witten, *Anomalies in String Theory with D-Branes*, Asian J. Math. **3** (1999), 819-852, [[arXiv:hep-th/9907189](#)].
- [GM03] C. Gómez and J. J. Manjarín, *A note on the dyonic D6-brane*, 6th International Workshop on Conformal Field Theory and Integrable Models, Landau Institute, Sept. 2002, [[arXiv:hep-th/0302096](#)].
- [GKS⁺01] E. Gorbatov, V.S. Kaplunovsky, J. Sonnenschein, S. Theisen, and S. Yankielowicz, *On Heterotic Orbifolds, M Theory and Type I' Brane Engineering*, J. High Energy Phys. **0205** (2002), 015, [[arXiv:hep-th/0108135](#)].
- [GS15] D. Grady and H. Sati, *Massey products in differential cohomology via stacks*, J. Homotopy Relat. Struct. **13** (2017), 169-223, [[arXiv:1510.06366](#)].

- [GS17] D. Grady and H. Sati, *Twisted differential generalized cohomology theories and their Atiyah-Hirzebruch spectral sequence*, Alg. Geom. Topol. (2019), [[arXiv:1711.06650](#)].
- [GS19] D. Grady and H. Sati, *Ramond-Ramond fields and twisted differential K-theory*, preprint.
- [Gu92] R. Gueven, *Black p-brane solutions of D = 11 supergravity theory*, Phys. Lett. **B276** (1992), 49-55, (reprinted in [Du99]), [[spire:338203](#)].
- [Gu65] V. Guillemin, *The integrability problem for G-structures*, Trans. Amer. Math. Soc. **116** (1965), 544-560, [[jstor:1994134](#)].
- [Gus09] A. Gustavsson, *Algebraic structures on parallel M2-branes*, Nucl. Phys. **B 811** (2009) 66–76, [[arXiv:0709.1260](#)] [[hep-th](#)].
- [HW97] A. Hanany and E. Witten, *Type IIB Superstrings, BPS Monopoles, And Three-Dimensional Gauge Dynamics*, Nucl. Phys. **B492** (1997), 152-190, [[arXiv:hep-th/9611230](#)].
- [HG07] A. Henriques and D. Gepner, *Homotopy Theory of Orbispaces*, [[arXiv:math/0701916](#)].
- [He06] K. Hess, *Rational homotopy theory: a brief introduction*, Interactions between homotopy theory and algebra, Contemp. Math **436**, 175-202, [[arXiv:math.AT/0604626](#)].
- [HHR09] M. A. Hill, M. J. Hopkins, and D. C. Ravenel, *On the non-existence of elements of Kervaire invariant one*, Ann. Math. **184** (2016), 1-262, [[arXiv:0908.3724](#)].
- [HW06] P. Hořava and E. Witten, *Heterotic and Type I string dynamics from eleven dimensions*, Nucl. Phys. **B460** (1996) 506–524, [[arXiv:hep-th/9510209](#)].
- [Ho97] P. Howe, *Weyl superspace*, Phys. Lett. **B 415** (1997), 149-155, [[arXiv:hep-th/9707184](#)].
- [HS17] J. Huerta and U. Schreiber, *M-Theory from the superpoint*, Lett. Math. Phys. **108** (2018), 2695–2727, [[arXiv:1702.01774](#)].
- [HSS18] J. Huerta, H. Sati, and U. Schreiber, *Real ADE-equivariant (co)homotopy of super M-branes*, [[arXiv:1805.05987](#)].
- [Hu98] C. M. Hull, *Massive string theories from M-theory and F-theory*, J. High Energy Phys. **11** (1998) 027, [[arXiv:hep-th/9811021](#)].
- [HW88] C. M. Hull and N. P. Warner, *Non-compact gaugings from higher dimensions*, Classical Quantum Grav. **5** (1988), 1517-1530.
- [IU12] L. Ibáñez and A. Uranga, *String Theory and Particle Physics: An Introduction to String Phenomenology*, Cambridge University Press, Cambridge, 2012.
- [Ka17] G. Kane, *String theory and the real world*, Morgan & Claypool, 2017.
- [Kl72] F. Klein, *Vergleichende Betrachtungen über neuere geometrische Forschungen* (1872) translation by M. W. Haskell, *A comparative review of recent researches in geometry*, Bull. New York Math. Soc. **2**, (1892-1893), 215-249.
- [Kö18] A. Körschgen, *A Comparison of two Models of Orbispaces*, Homol., Homotpy Appl. **20** (2018), 329-358, [[arXiv:1612.04267](#)] [[math.AT](#)].
- [KS05] I. Kriz and H. Sati, *Type IIB String Theory, S-Duality, and Generalized Cohomology*, Nucl. Phys. **B715** (2005) 639-664, [[arXiv:hep-th/0410293](#)].
- [Ku04] N. J. Kuhn, *Goodwillie towers and chromatic homotopy: an overview* Geom. Topol. Monogr. **10** (2007), 245-279, [[arXiv:math/0410342](#)].
- [LT96] K. Lechner and M. Tonin, *Worldvolume and target space anomalies in the D = 10 super-fivebrane sigma-model*, Nucl. Phys. **B475** (1996), 545-561, [[hep-th/9603094](#)].
- [Le17] T. Lee, *Covariant open bosonic string field theory on multiple D-branes in the proper-time gauge*, J. Korean Phys. Soc. **71** (2017), 886-903, [[arXiv:1609.01473](#)].
- [Lo90] J. Lott, *The geometry of supergravity torsion constraints*, Comm. Math. Phys. **133** (1990), 563-615; see [[arXiv:math/0108125](#)].
- [Lu09] J. Lurie, *Higher Topos Theory*, Princeton University Press, 2009, [[arXiv:math/0608040](#)].
- [Lu17] J. Lurie, *Higher Algebra*, [[math.harvard.edu/~lurie/papers/HA.pdf](#)].
- [Ma04] J. J. Manjarín, *Topics on D-brane charges with B-fields*, Int. J. Geom. Meth. Mod. Phys. **1** (2004), 545-602, [[arXiv:hep-th/0405074](#)].

- [MS04] V. Mathai and H. Sati, *Some relations between twisted K-theory and E_8 gauge theory*, J. High Energy Phys. **03** (2004), 016, [[arXiv:hep-th/0312033](#)].
- [MS06] J. P. May and J. Sigurdsson, *Parametrized homotopy theory*, Amer. Math. Soc., Providence, RI, 2006.
- [MM97] R. Minasian and G. Moore, *K-theory and Ramond-Ramond charge*, J. High Energy Phys. **9711** (1997), 002, [[arXiv:hep-th/9710230](#)].
- [MY60] D. Montgomery and C. T. Yang, *Groups on S^n with principal orbits of dimension $n-3$, I*, Illinois J. Math. **4** (1960), 507-517.
- [MZ54] D. Montgomery and L. Zippin, *Examples of transformation groups*, Proc. Amer. Math. Soc. **5** (1954), 460-465.
- [Mo14] G. Moore, *Physical Mathematics and the Future*, talk at Strings 2014 [[physics.rutgers.edu/~gmoore/PhysicalMathematicsAndFuture.pdf](#)].
- [MS03] G. Moore and N. Saulina, *T-duality, and the K-theoretic partition function of type IIA superstring theory*, Nucl. Phys. **B670** (2003) 27-89 [[arXiv:hep-th/0206092](#)].
- [MW00] G. Moore and E. Witten, *Self-duality, Ramond-Ramond Fields, and K-theory*, J. High Energy Phys. **0005** (2000), 032, [[hep-th/9912279](#)].
- [My03] R. C. Myers, *Nonabelian Phenomena on D-branes*, Class. Quant.Grav. **20** (2003), S347-S372, [[arXiv:hep-th/0303072](#)].
- [NV00] H. Nastase and D. Vaman, *On the nonlinear KK reductions on spheres of supergravity theories*, Nucl. Phys. **B 583** (2000), 211-236, [[arXiv:hep-th/0002028](#)].
- [NVvN00] H. Nastase, D. Vaman, and P. van Nieuwenhuizen, *Consistency of the $AdS_7 \times S^4$ reduction and the origin of self-duality in odd dimensions*, Nucl. Phys. **B 581** (2000) 179-239, [[arXiv:hep-th/9911238](#)].
- [NSS12] T. Nikolaus, U. Schreiber, and D. Stevenson, *Principal ∞ -bundles – general theory*, J. Homotopy Related Structr. **10** (2015), 749–801, [[arXiv:1207.0248](#)].
- [Pa78] P. S. Pao, *Nonlinear circle actions on the 4-sphere and twisting spun knots*, Topology **17** (1978), no. 3, 291-296.
- [PNT84] K. Pilch, P. van Nieuwenhuizen, and P.K. Townsend, *Compactification of $D = 11$ supergravity on S^4 (or $11 = 7 + 4$, too)*, Nucl. Phys. **B 242** (1984), 377-392.
- [Po01] J. Polchinski, *String theory*, Cambridge University Press, Cambridge, 2001.
- [Ra03] D. Ravenel, *Complex cobordism and stable homotopy groups of spheres*, Amer. Math. Soc., Providence, RI, 2003, [[web.math.rochester.edu/people/faculty/doug/mu.html](#)].
- [Re87] M. Reid, *Young person's guide to canonical singularities*, in S. Bloch (ed.), *Algebraic geometry – Bowdoin 1985, Part 1*, Proc. Sympos. Pure Math. 46 Part 1, Amer. Math. Soc., Providence, RI, 1987, pp. 345-414.
- [Ro93] A. Roig, *Minimal resolutions and other minimal models*, Publ. Matemáticas **37** (1993), 285-303.
- [Ro94] A. Roig, *Formalizability of dg modules and morphisms of cdg algebras*, Illinois J. Math. **38** (1994), 434-451, [[euclid:1255986724](#)].
- [RS00] A. Roig and M. Saralegi-Aranguren, *Minimal models for non-free circle actions*, Illinois J. Math. **44** (2000), no. 4, 784-820, [[arXiv:math/0004141](#)].
- [Sa00] M. Sakaguchi, *IIB-branes and new spacetime superalgebras*, J. High Energy Phys. **0004** (2000), 019, [[arXiv:hep-th/9909143](#)].
- [Sa05a] H. Sati, *M-theory and Characteristic Classes*, J. High Energy Phys. **0508** (2005) 020, [[arXiv:hep-th/0501245](#)].
- [Sa05b] H. Sati, *Flux Quantization and the M-Theoretic Characters*, Nucl. Phys. **B727** (2005) 461-470, [[arXiv:hep-th/0507106](#)].
- [Sa06] H. Sati, *Duality symmetry and the form fields of M-theory*, J. High Energy Phys. **0606** (2006) 062, [[arXiv:hep-th/0509046](#)].
- [Sa10] H. Sati, *Geometric and topological structures related to M-branes*, Superstrings, geometry, topology, and C^* -algebras, 181-236, Proc. Sympos. Pure Math., 81, Amer. Math. Soc., Providence, RI, 2010, [[arXiv:1001.5020](#)].

- [Sa13] H. Sati, *Framed M-branes, corners, and topological invariants*, J. Math. Phys. **59** (2018), 062304, [arXiv:1310.1060].
- [SS18] H. Sati and U. Schreiber, *Higher T-duality of M-branes via local supersymmetry*, Phys. Lett. **B 781** (2018), 694–698, [arXiv:1805.00233].
- [Sc15] U. Schreiber, *Higher Cartan Geometry*, lecture notes, Prague 2015, [ncatlab.org/schreiber/show/Higher+Cartan+Geometry].
- [Sc16] U. Schreiber, *From the Superpoint to T-Folds*, lecture notes, Prague 2016, [ncatlab.org/schreiber/show/From+the+Superpoint+to+T-Folds].
- [Sc17a] U. Schreiber, *Introduction to Homotopy Theory*, lecture notes, Bonn 2017, [ncatlab.org/nlab/show/Introduction+to+Homotopy+Theory].
- [Sc17b] U. Schreiber, *Introduction to Stable Homotopy Theory*, lecture notes, Bonn 2017, [ncatlab.org/nlab/show/Introduction+to+Stable+homotopy+theory+---+1].
- [Sc17c] U. Schreiber, *Super p-Brane Theory emerging from Super Homotopy Theory*, talk at String Math 17, Hamburg, 2017, [ncatlab.org/schreiber/show/StringMath2017].
- [SS03] S. Schwede and B. Shipley, *Stable model categories are categories of modules*, Topology **42** (2003), 103-153.
- [Se97] A. Sen, *A Note on Enhanced Gauge Symmetries in M- and String Theory*, J. High Energy Phys. **9709** (1997), 001, [arXiv:hep-th/9707123].
- [Sh07] B. Shipley, *HZ-algebra spectra are differential graded algebras*, Amer. J. Math. **129** (2007), 351-379, [arXiv:math/0209215].
- [Sn81] V. Snaitch, *Localized stable homotopy of some classifying spaces*, Math. Proc. Cambridge Philos. Soc. **89** (1981), no. 2, 325-330.
- [Su77] D. Sullivan, *Infinitesimal computations in topology*, Publ. Math. de I.H.É.S. **47** (1977), 269-331.
- [To95a] P. Townsend, *The eleven-dimensional supermembrane revisited*, Phys. Lett. **B350** (1995), 184-187, [arXiv:hep-th/9501068].
- [To95b] P. Townsend, *D-branes from M-branes*, Phys. Lett. **B373** (1996) 68-75, [arXiv:hep-th/9512062].
- [duV34] P. du Val, *On isolated singularities of surfaces which do not affect the conditions of adjunction*, I, II and III, Proc. Camb. Phil. Soc. **30** (1934), 453–459, 460–465, 483–491.
- [VB85] M. Vigué-Poirrier and D. Burghelea, *A model for cyclic homology and algebraic K-theory of 1-connected topological spaces*, J. Differential Geom. **22** (1985), 243-253,
- [We17] F. Wellen, *Formalizing Cartan Geometry in Modal Homotopy Type Theory*, PhD Thesis, KIT 2017, [ncatlab.org/schreiber/show/thesis+Wellen].
- [Wi81] E. Witten, *Search for a realistic Kaluza-Klein theory*, Nucl. Phys. **B 186** (1981), 412-428.
- [Wi95] E. Witten, *String Theory Dynamics In Various Dimensions*, Nucl. Phys. **B443** (1995), 85-126, [arXiv:hep-th/9503124].
- [Wi96] E. Witten, *Bound States of Strings and p-Branes*, Nucl. Phys. **B460** (1996), 335-350, [arXiv:hep-th/9510135].
- [Wi98] E. Witten, *D-Branes And K-Theory*, J. High Energy Phys. **9812** (1998), 019, [arXiv:hep-th/9810188].
- [Wi00] E. Witten, *Overview Of K-Theory Applied To Strings*, Int. J. Mod. Phys. **A16** (2001), 693-706, [arXiv:hep-th/0007175].
- [Wi12] E. Witten, *Superstring Perturbation Theory Revisited*, [arXiv:1209.5461].