# HOFFMANN-OSTENHOF'S CONJECTURE FOR CLAW-FREE CUBIC GRAPHS

### MILAD AHANJIDEH AND ELHAM ABOOMAHIGIR

ABSTRACT. Hoffmann-Ostenhof's Conjecture states that the edge set of every connected cubic graph can be decomposed into a spanning tree, a matching and a 2-regular subgraph. In this paper, we show that the conjecture holds for claw-free cubic graphs.

#### 1. Introduction

Graphs are finite without loops and multiple edges throughout this paper. Let G be a finite undirected graph with the vertex set V(G) and the edge set E(G). For a vertex  $v \in V(G)$ , the degree of v in G is denoted by  $d_G(v)$ . An edge joining vertices u and v is denoted by uv. Here,  $N_G(v)$  denotes the set of all neighbours of v. The complete graph of order n is denoted by  $K_n$ . The complete graph on four vertices minus one edge is called a diamond. A graph is cubic in which all vertices have degree three and a subcubic graph is a graph in which each vertex has degree at most three. A graph is claw-free if it has no induced subgraph isomorphic to  $K_{1,3}$ . A cycle C is called chordless if C has no cycle chord. For every  $S \subseteq V(G)$ , the graph obtained by removing all vertex in S and all associated incident edges is denoted by  $G \setminus S$ . If X is a subset of the edges, then  $G \setminus X$  is the graph obtained by removing all edges in X from G. A cut-edge of a connected graph G is an edge  $e \in E(G)$  such that  $G \setminus e$  is disconnected.

Now, we adopt some terminology from [6] in which we need for the proof of the main theorem. For convenience, we repeat their definitions here.

Let  $k \geq 2$  be an integer, a diamond-necklace  $N_k$  with k diamonds is a connected cubic graph constructed as follows. Take k disjoint copies  $D_1, D_2, \ldots, D_k$  of a diamond, where  $V(D_i) = \{a_i, b_i, c_i, d_i\}$  and  $a_i d_i$  is the missing edge in  $D_i$ . Let  $N_k$  be obtained from the disjoint union of these k diamonds by adding the edges  $\{d_i a_{i+1} | i = 1, 2, \ldots, k-1\}$  and adding the edge  $d_k a_1$ , see Figure 1.

For  $k \geq 1$ , a diamond-bracelet  $B_k$  with k diamonds is defined as follows. Let  $B_k$  be obtained from a diamond-necklace  $N_{k+1}$  with k+1 diamonds  $D_1, D_2, \ldots, D_{k+1}$  by removing the diamond  $D_{k+1}$  and adding a triangle T with  $V(T) = \{a, b, c\}$ , and adding two edges  $ba_1$  and  $ad_k$ , see Figure 1.

For  $k \geq 1$ , a diamond-chain  $L_k$  with k diamonds is defined as follows. Let  $L_k$  be obtained from k disjoint copies  $D_1, D_2, \ldots, D_k$  of a diamond by adding the edges  $\{d_i a_{i+1} | i = 1, \ldots, k-1\}$ . After that we add two disjoint triangles  $T_1$  and  $T_2$  and add one edge joining  $a_1$  to a vertex of  $T_1$ , and one edge joining  $d_k$  to a vertex of  $T_2$ , see Figure 1.

 $<sup>1991\</sup> Mathematics\ Subject\ Classification.\ 05C70,\ 05C05.$ 

Key words and phrases. Cubic graph, Hoffmann-Ostenhof's Conjecture, Claw-free graph.




FIGURE 1. From the left side: A diamond-necklace  $N_7$ ; A diamond-bracelet  $B_6$ ; A diamond-chain  $L_2$ .

The following conjecture was posed in [7] and appeared as a problem in BCC22 [5]:

**Hoffmann-Ostenhof's Conjecture.** Let G be a connected cubic graph. Then the edges of G can be decomposed into a spanning tree, a matching and a 2-regular subgraph.

Note that the spanning tree or the 2-regular subgraph cannot be empty, however the matching may be empty. An edge decomposition of a graph G is called a *good decomposition*, if the edges of G can be decomposed into a spanning tree, a matching and a 2-regular subgraph. A graph is called *good* if it has a good decomposition. Throughout, we use  $\{T, M, O\}$  to denote the spanning tree, the matching and the 2-regular subgraph of the good decomposition of G, respectively.

Hoffmann-Ostenhof's Conjecture is known to be true for some families of cubic graphs. Kostochka [9] showed that the Petersen graph, the prism over cycles, and many other graphs are good. Bachstein [4] proved that every 3-connected cubic graph embedded in torus or Klein-bottle is good. Furthermore, Ozeki and Ye [10] proved that 3-connected cubic plane graphs are good. Akbari et. al. [3] showed that hamiltonian cubic graphs are good. Also, it has been proved that traceable cubic graphs are good [1]. In 2017, Hoffmann-Ostenhof et. al. [8] proved that planar cubic graphs are good. Recently in [2] the authors proved that claw-free subcubic graphs and 4-chordal subcubic graphs have the same property. In this paper, we are interested in finding a good decomposition of claw-free cubic graphs. We prove the following theorem.

**Main Theorem.** Let G be a connected claw-free cubic graph. Then G is good.

# 2. Preliminary Results

In order to prove the main theorem, we need to know a few basic properties about claw-free cubic graphs.

**Lemma 2.1.** [6, Claim A] If  $G \neq K_4$  is a connected claw-free cubic graph, then the vertex set V(G) can be uniquely partitioned into sets each of which induces a triangle or a diamond in G.

In the next two lemmas, we discuss some properties of a good decomposition of a claw-free cubic graph.

**Lemma 2.2.** Let G be a connected claw-free cubic graph which has a good decomposition and let C be a cycle in  $G \setminus T$ , where T is a spanning tree of the good decomposition of G. Then C is either a triangle or the following statements hold for C:

- (i) C is a chordless cycle.
- (ii) There is no vertex of C in a diamond of G.
- (iii) C is an even cycle.

*Proof.* Assume that C is not a triangle.

- (i) Suppose that C contains two vertices a and b such that ab is a chord in C. Since  $d_G(a) = d_G(b) = 3$ , the spanning tree T does not contain a and b, a contradiction.
- (ii) Suppose that (ii) is false. Since G is cubic, the cycle C should pass through of 3 or 4 vertices of a diamond. It is easy to see that the spanning tree T does not contain some vertices of this diamond, a contradiction.
- (iii) Clearly, every vertex of a claw-free cubic graph lies in a triangle. Assume that  $a_1$ ,  $a_2$  and  $a_3$  are three consecutive vertices in C such that  $a_1a_2a_3$  is a path. By (i),  $a_1$  and  $a_3$  are not adjacent. Thus it is easy to see that two consecutive vertices of every three consecutive vertices of C lie in a triangle. Note that there is no vertex in C lying in a diamond. So there are |V(C)|/2 edges belonging to triangles of C and hence |V(C)| is even.

**Lemma 2.3.** Let G be a connected claw-free cubic graph with the good decomposition  $\{T, M, O\}$ . If  $C \neq K_3$  is a cycle in the 2-regular subgraph O, then for the fix edge xy of C, we can find another good decomposition such that xy belongs to either T or M.

*Proof.* Assume that C is a cycle in O containing xy. By Lemma 2.2, C is an even cycle. For the convenience, let us recall x by  $b_1$  and y by  $a_2$  and call every triangle  $a_i t_i b_i$  with one edge on C by  $C_i$ . If  $b_1 a_2$  does not lie in a triangle, then there are two distinct triangles  $C_1$  and  $C_2$  such that  $b_1$  belongs to  $C_1$  and  $a_2$  belongs to  $C_2$ . Note that  $t_i$  is not adjacent to  $t_j$  for i, j, where  $1 \le i, j \le |V(C)|/2$ , otherwise we have a spanning forest instead of a spanning tree in G. Now, assume that the number of triangles with one edge on C is 2k. We change the decomposition of G such that  $b_1a_2$  lies in the spanning tree as follows. We consider  $T' = T \cup \{b_1 a_2, b_{2k} a_1, a_{2i} t_{2i}, t_{2i} b_{2i}, b_j a_{j+1} | i = 1, \dots, k, j = 1, \dots, 2k - 1\},$  $M' = M \cup \{a_{2i}b_{2i} | i = 1, ..., k\}$  and  $O' = (O \setminus \{b_1a_2\}) \cup \{C_{2i-1} | i = 1, ..., k\}$  as a new good decomposition into a spanning tree, a matching and a 2-regular subgraph of G. If the number of triangles with one edge on C is 2k-1, then similar to the above argument, we change the decomposition of G such that  $b_1a_2$  lies in a matching. We consider  $T' = T \cup \{a_1b_1, t_1b_1, a_2t_2, a_2b_2, b_{2k-1}a_1, a_{2i}t_{2i}, t_{2i}b_{2i}, b_ja_{j+1} | i = 1\}$  $2, \ldots, k-1, j=2, \ldots, 2k-2\}, M'=M \cup \{a_1t_1, b_1a_2, t_2b_2, a_{2i}b_{2i} | i=2, \ldots, k-1\}$ and  $O' = (O \setminus \{b_1 a_2\}) \cup \{C_{2i-1} | i = 2, ..., k\}$  as a new good decomposition of G. See Figure 2(b).

Now, Assume that  $b_1a_2$  lies in a triangle. Then consider one of incident edges with

xy on cycle C and fix it. Then we apply the above method. We can see at once that xy belongs to T or M.




FIGURE 2. From the left side: a, b.

# 3. Proof of the Main Theorem

*Proof.* We proceed by induction on the order |V(G)| = n of G. Obviously, both  $K_4$  and the Cartesian product  $C_3 \square K_2$  have the desired decompositions. Now, we divide the proof into two steps:

**Step 1.** Let G be a claw-free cubic graph of order n with no diamond.

Let  $n \geq 8$  and the hypothesis holds for all claw-free cubic graphs of order less than n. Assume that  $v_1v_2v_3$  is a triangle in G. Let  $u_1$  and  $u_2$  be the other neighbours of  $v_1$  and  $v_2$ , respectively. If  $u_1$  and  $u_2$  are adjacent, then since G is claw-free,  $u_1$  and  $u_2$  have a common neighbour  $u_3$ . If  $u_3$  and  $v_3$  are adjacent, then G is the Cartesian product  $C_3 \square K_2$  and it has a good decomposition. So, assume that  $x \neq u_3$  is a neighbour of  $v_3$  and  $y \neq v_3$  is a neighbour of  $u_3$ . It is easy to see that  $x \neq y$ , otherwise there is a claw in x. If x and y are not adjacent, then we put  $A = \{u_1, u_2, u_3, v_1, v_2, v_3\}$  and consider  $G \setminus A$ . Join x to y and call the resulting claw-free cubic graph  $G^A$ . Since  $|V(G^A)| < |V(G)|$ , by induction hypothesis  $G^A$  has a good decomposition  $\{T^A, M^A, O^A\}$ , where  $T^A$  is a spanning tree of  $G^A$ ,  $M^A$  is a matching and  $O^A$  is a 2-regular subgraph of  $G^A$ . We extend this decomposition of  $G^A$  to G. It is clear that  $G^A$  is connected, so we have the following three cases:

- i. If xy lies in the spanning tree  $T^A$ , then since the spanning tree  $T^A$  saturates vertices x and y, we can consider  $T = (T^A \setminus \{xy\}) \cup \{xv_3, v_3v_1, v_1v_2, v_2u_2, u_2u_1, u_1u_3, u_3y\}$  and  $M = M^A \cup \{v_3v_2, v_1u_1, u_2u_3\}$  as a spanning tree and a matching of G, respectively and we keep the cycles in the good decomposition of  $G^A$ , see Figure 3 (a).
- ii. If xy lies in the matching  $M^A$ , then we define  $T = T^A \cup \{xv_3, v_1v_3, v_2v_3, u_1u_3, u_2u_3, u_3y\}$ ,  $M = M^A \setminus \{xy\}$  and  $O = O^A \cup \{v_1v_2u_2u_1\}$  as a good decomposition of G, see Figure 3 (b).

Before we discuss the third case, we need the following claim.

Claim 1. Let C be a cycle in  $G^A$  containing xy, then |V(C)| > 3.

Proof. Assume that |V(C)| = 3, then there is  $t_1 \in N_{G^A}(x) \cap N_{G^A}(y)$ . Let  $a_1 \in N_{G^A}(x) \setminus \{t_1\}$  and  $a_2 \in N_{G^A}(y) \setminus \{t_1, a_1\}$ . Note that if  $a_1$  and  $a_2$  are same, then there are claws in G. Since G is cubic,  $t_1$  can be adjacent to at most one of the two vertices  $a_1$  and  $a_2$ . Therefore G has at least one claw containing x or y, a contradiction. So |V(C)| > 3 and in particular xy is not in a triangle in  $G^A$ .  $\square$ 

iii. If xy lies in the cycle, then by Claim 1, |V(C)| > 3 and hence by Lemma 2.3, this case will be converted to either Case i or Case ii.



FIGURE 3. From the left side: a; b.

Now, if x and y are adjacent, then for avoiding of a claw, there is  $b \in N_G(x) \cap N_G(y)$ , see Figure 4. Note that triangle  $v_1v_2v_3$  join to another triangle xyb by the edge  $v_3x$ . We investigate this case in the below by replacing x with  $u_3$ , y with  $u_1$  and b with  $u_2$ .

Now, assume that triangle  $v_1v_2v_3$  join to another triangle  $u_1u_2u_3$  by the edge  $u_3v_3$ .



FIGURE 4. If x and y are adjacent.

Let x, y, w and z be the neighbours of  $v_1, v_2, u_1$  and  $u_2$ , respectively. If either  $x = u_1$  or  $x = u_2$ , then we have done before. Without loss of generality, we can assume that x, y, w and z are different from  $u_1, u_2, v_1$  and  $v_2$ . Since there is no diamond in  $G, x \neq y$  and  $w \neq z$ . If x and y or w and z are adjacent, then we have done before. So, assume that x and y (w and z) are not adjacent. If there is  $t \in N_G(x) \cap N_G(y)$ , then there is a claw in vertices x or y. If x = w or x = z, then we have a claw in x. So, assume that x, y, w and z are not adjacent. Put  $B = \{u_1, u_2, u_3, v_1, v_2, v_3\}$  and consider  $G \setminus B$ . Join x to y and w to z, we denote the new graph  $G^B$ . It is obvious that  $G^B$  is a claw-free cubic graph and  $|V(G^B)| < |V(G)|$ .

If G is 2-edge connected, then  $G^B$  is connected. So by induction hypothesis  $G^B$  has a good decomposition  $\{T^B, M^B, O^B\}$ , where  $T^B$  is a spanning tree of  $G^B$ ,  $M^B$  is a matching and  $O^B$  is a 2-regular subgraph. We extend this decomposition of  $G^B$  to G. The proof falls into six cases:

Case 1. If xy and wz belong to  $M^B$ , then since the spanning tree  $T^B$  saturates vertices x, y, w and z, consider  $T = T^B \cup \{xv_1, v_1v_2, v_2v_3, v_3u_3, wu_1, u_2z\}$ ,  $M = (M^B \setminus \{xy, wz\}) \cup \{v_1v_3, v_2y\}$  and  $O = O^B \cup \{u_1u_2u_3\}$  as a good decomposition of G, see Figure 5 (a).

- Case 2. If xy and wz belong to  $T^B$ , then we can consider  $T = (T^B \setminus \{xy, wz\}) \cup \{xv_1, v_1v_3, v_3v_2, v_2y, wu_1, u_1u_3, u_3u_2, u_2z\}, M = M^B \cup \{v_1v_2, u_1u_2, u_3v_3\}$  and  $O = O^B$  as a good decomposition of G, see Figure 5 (b).
- Case 3. If  $xy \in T^B$  and  $wz \in M^B$ , then we consider  $T = (T^B \setminus \{xy\}) \cup \{xv_1, v_1v_2, v_2v_3, v_2y, v_3u_3, wu_1, u_2z\}$ ,  $M = (M^B \setminus \{wz\}) \cup \{v_1v_3\}$  and  $O = O^B \cup \{u_1u_2u_3\}$  as a good decomposition of G.
- Case 4. If  $xy \in T^B$  and  $wz \in O^B$ , then we can consider  $T = (T^B \setminus \{xy\}) \cup \{xv_1, v_1v_3, v_3v_2, v_2y, v_3u_3, u_3u_1, u_3u_2\}$ ,  $M = M^B \cup \{v_1v_2\}$  and  $O = (O^B \setminus \{wz\}) \cup \{wu_1, u_1u_2, u_2z\}$  as a good decomposition of G, see Figure 5 (c).
- Case 5. If  $xy \in M^B$  and  $wz \in O^B$ , then we have  $T = T^B \cup \{xv_1, v_1v_2, v_2v_3, v_3u_3, u_3u_1, u_3u_2\}$ ,  $M = (M^B \setminus \{xy\}) \cup \{v_1v_3, v_2y\}$  and  $O = (O^B \setminus \{wz\}) \cup \{wu_1, u_1u_2, u_2z\}$  as a good decomposition of G, see Figure 5 (d).




FIGURE 5. From the left side: a, b, c, d.

Case 6. If xy and wz belong to  $O^B$ , then similar to that in the proof of Claim 1 we have  $C \neq K_3$ . So by Lemma 2.3, this case will be converted to either Case 4 or Case 5.

Now, assume that  $G^B$  is not connected. Let  $G_1^B$  and  $G_2^B$  be the connected components of  $G^B$ . So by induction hypothesis  $G_1^B$  and  $G_2^B$  have good decompositions  $\{T_1^B, M_1^B, O_1^B\}$  and  $\{T_2^B, M_2^B, O_2^B\}$ . From this we want to find a new decomposition  $\{T^B, M^B, O_1^B\}$  of  $G^B$ , where  $T^B$  is a spanning forest, M is a matching and  $O^B$  is a 2-regular subgraph. Define  $T^B = T_1^B \cup T_2^B$ ,  $M^B = M_1^B \cup M_2^B$  and  $O^B = O_1^B \cup O_2^B$ . For extending this decomposition of  $G^B$  to G, we make slight changes in some cases for when  $G^B$  is connected. Now, we investigate those cases here. We remove xy and xy from the decomposition of xy. In Case 1, a good decomposition of yy is obtained by adding the set of edges xy, xy, xy, xy, xy, xy, xy, xy, xy, yy, y

Note that by applying the proof of Claim 1, we can deduce that  $G^B$  has no diamond.

**Step 2.** Let G be a claw-free cubic graph of order  $n \geq 8$  with at least one string which contains  $k \geq 1$  diamonds. First, assume that G is a diamond-necklace with  $k \geq 2$  diamonds. Because G is hamiltonian, according to the proof of Theorem 9 in [3], G has the desired decomposition. Now, assume that G is a claw-free cubic

graph of order n containing a string of at least two consecutive diamonds. Without loss of generality, consider a string of the maximum number of diamonds with starting and ending vertices w and z. It is easy to check that  $w \neq z$ . By Lemma 2.1, w and z belong to triangles or diamonds. If at least one of w or z belongs to a diamond, then it leads to a contradiction with our assumption on the maximum number of diamonds in string. Now, we consider the following cases:

- a. Let w and z belong to a common triangle. That is, we have a diamond-bracelet  $B_k$  in G.
- b. Let w and z belong to two distinct triangles.

Since the proof of both cases are similar, we prove only part (a) and the proof of part (b) is left to the reader. we consider  $G \setminus \{a_i, b_i, c_i, d_i \mid i = 2, \dots, k\}$  and then join  $d_1$  to z. Let us call this new graph G'. By induction hypothesis, G' has a good decomposition  $\{T', M', O'\}$ . We extend this good decomposition to G. If  $d_1z$  lies in the spanning tree T', then we define  $T = T' \cup \{a_ib_i, b_ic_i, c_id_i \mid 2 \leq i \leq k\} \cup \{d_ja_{j+1} \mid 1 \leq j \leq k-1\} \cup \{d_kz\}, M = M' \cup \{a_ic_i, b_id_i \mid 2 \leq i \leq k\}$  and we keep the cycles in the good decomposition of G'. If  $d_1z$  lies in the matching M', then we define  $T = T' \cup \{d_kz, b_2d_2, c_2d_2\} \cup \{a_ib_i, b_ic_i, c_id_i \mid 3 \leq i \leq k\} \cup \{d_ja_{j+1} \mid 1 \leq j \leq k-1\}, M = M' \cup \{a_ic_i, b_id_i \mid 3 \leq i \leq k\}$  and  $O = O' \cup \{a_2b_2c_2\}$ , see Figure 6. Now, let us consider the case in which  $d_1z$  lies in a cycle. Since G is cubic, the edge  $wa_1$  lies in every possible cycle containing  $d_1z$  in the decomposition of G'. It is easy to check that of  $a_1, b_1, c_1$  and  $d_1$  are not covered by the spanning tree of G', a contradiction.

Here, we examine the case in which G has only one diamond. First, assume that w




FIGURE 6. From the left side: a)  $d_1z$  lies in the spanning tree T'; b)  $d_1z$  lies in the matching M'.

and z lie in two distinct triangles. Remove the diamond and the edges  $wa_1, d_1z$ , then join w to z and call this new cubic graph G'. The graph G' is a claw-free cubic graph and |V(G')| < |V(G)|. So by Step 1, G' has a good decomposition  $\{T', M', O'\}$ . If wz lies in the spanning tree T', then we define  $T = T' \cup \{wa_1, a_1b_1, b_1c_1, c_1d_1, d_1z\}$ ,  $M = M' \cup \{a_1c_1, b_1d_1\}$  and we keep the cycles in the good decomposition of G'. If wz lies in the matching M', then we define  $T = T' \cup \{wa_1, a_1b_1, a_1c_1, d_1z\}$ ,  $O = O' \cup \{b_1c_1d_1\}$  and we keep the matching in the good decomposition of G', see Figure 7. Assume that wz lies in a cycle C, by Step 1, the graph G' has a good decomposition  $\{T', M', O'\}$ . It is obvious that |V(C)| > 3. So by Lemma 2.3, we

can convert this case to previous cases, that is the cases that wz lies in either a spanning tree or a matching.




FIGURE 7. From the left side: a) wz lies in the spanning tree T'; b) wz lies in the matching M'.




FIGURE 8. Only one diamond, w and z lie in a common triangle.

Now, assume that w and z are on a common triangle. Assume that u is a common neighbour of w and z. Let another adjacent vertex to u be x. It is obvious that xu is a cut-edge of G. Assume that  $G^x$  is a connected component of  $G \setminus \{xu\}$  containing x. Since  $G^x$  is a connected claw-free subcubic graph, by [2, Theorem 2.1], the graph  $G^x$  has a good decomposition  $\{T^x, M^x, O^x\}$ . Now, we extend this good decomposition to G. Add the set of edges  $\{xu, uw, uz, wa_1, zd_1, d_1c_1, d_1b_1\}$  to  $T^x$ , wz to  $M^x$  and the triangle  $a_1b_1c_1$  to  $O^x$  to obtain a good decomposition into the spanning tree, the matching and the 2-regular subgraph of G, see Figure 8.  $\square$ 

**Remark 3.1.** Note that in the main theorem, the matching M can be empty. We can see at once that the main theorem does not hold for subcubic graphs, a cycle is a counterexample.

Acknowledgements. The authors would like to thank Professor Saieed Akbari for a discussion on this topic and for his comments.

# References

- F. Abdolhosseini, S. Akbari, H. Hashemi, M.S. Moradian, Hoffmann-Ostenhofs conjecture for traceable cubic graphs, (2016), arXiv:1607.04768v1.
- [2] E. Aboomahigir, M. Ahanjideh, S. Akbari, Decomposing claw-free subcubic graphs and 4chordal subcubic graphs, (2018), arXiv:1806.11009v1.
- [3] S. Akbari, T.R. Jensen, M. Siggers, Decomposition of graphs into trees, forests, and regular subgraphs, Discrete Math. 338(8) (2015), 1322–1327.

- [4] A.C. Bachstein, Decomposition of Cubic Graphs on The Torus and Klein Bottle, A Thesis Presented to the Faculty of the Department of Mathematical Sciences Middle Tennessee State University, 2015.
- [5] P.J. Cameron, Research problems from the BCC22, Discrete Math. 311 (2011), 1074-1083.
- [6] M.A. Henning, C. Lwenstein, Locating-total domination in claw-free cubic graphs. Discrete Math. 312(21) (2012), 3107–3116.
- [7] A. Hoffmann-Ostenhof, Nowhere-zero Flows and Structures in Cubic Graphs, Ph.D. thesis (2011), University of Vienna.
- [8] A. Hoffmann-Ostenhof, T. Kaiser, K. Ozeki, Decomposition planar cubic graphs, J. Graph Theory. 88 (2018), 631–640.
- [9] A. Kostochka, Spanning trees in 3-regular graphs, REGS in Combinatorics, University of Illinois at Urbana Champaign, (2009). http://www.math.uiuc.edu/~west/regs/span3reg.html. 52 (2016), 40–46.
- [10] K. Ozeki, D. Ye, Decomposing plane cubic graphs, European J. Combin. 52 (2016), 40–46.
- [11] D.B. West, Introduction to Graph Theory, Prentice-Hall (2001).

Department of Mathematics, Faculty of Mathematical Sciences, Tarbiat Modares University, Tehran, Iran.

 $E ext{-}mail\ address: ahanjidm@gmail.com}$ 

DEPARTMENT OF MATHEMATICS, FACULTY OF MATHEMATICAL SCIENCES, TARBIAT MODARES UNIVERSITY, TEHRAN, IRAN.

E-mail address: mahigir.elham@gmail.com