Markov-Induced CM Model

Reza Rezaie and X. Rong Li Department of Electrical Engineering University of New Orleans New Orleans, LA 70148 rrezaie@uno.edu and xli@uno.edu

Abstract

Conditionally Markov (CM) sequences are powerful mathematical tools for modeling random phenomena. There are several classes of CM sequences one of which is the reciprocal sequence. Reciprocal sequences have been widely used in many areas including image processing, intelligent systems, and acausal systems. To use them in application, we need not only their applicable dynamic models, but also some general approaches to designing parameters of dynamic models. Dynamic models governing two important classes of nonsingular Gaussian (NG) CM sequences (called CM_L and CM_F models), and a dynamic model governing the NG reciprocal sequence (called reciprocal CM_L model) were presented in our previous work. In this paper, these models are studied in more detail and general approaches are presented for their parameter design. It is shown that every reciprocal CM_L model can be induced by a Markov model and parameters of the reciprocal CM_L model can be obtained from those of the Markov model. Also, it is shown how NG CM sequences can be represented in terms of a NG Markov sequence and an independent NG vector. This representation provides a general approach for parameter design of CM_L and CM_F models. In addition, it leads to a better understanding of CM sequences, including the reciprocal sequence.

Keywords: Conditionally Markov (CM) sequence, reciprocal sequence, Markov sequence, Gaussian sequence, dynamic model, characterization.

1 Introduction

Reciprocal processes have been applied to many different areas, including stochastic mechanics, image processing, acausal systems, trajectory modeling and intent inference, and intelligent systems (e.g., [1]–[13]). In [1]–[2], reciprocal processes were discussed in the context of stochastic mechanics. [3]–[4] considered the application of reciprocal processes in image processing. In [5] the behavior of acausal systems was described using reciprocal processes. Based on quantized state space, [6]–[9] used finite-state reciprocal sequences for detection of anomalous trajectory pattern, intent inference, and tracking. [10] used the idea of reciprocal processes for intent inference in intelligent interactive displays of vehicles. [12]–[13] proposed some classes of CM sequences (including reciprocal) for trajectory modeling.

Since they are desirable for many applications, reciprocal processes should be studied more to develop easily applicable tools for their application. This is the main goal of this paper.

Consider stochastic sequences defined over the interval $[0, N] = \{0, 1, ..., N\}$. For convenience, let the index be time. A sequence is Markov iff conditioned on the state at any time k, the subsequences before and after k are independent. A sequence is reciprocal iff conditioned on the states at any two times k_1 and k_2 , the subsequences inside and outside the interval $[k_1, k_2]$ are independent. In other words, inside and outside are independent given the boundaries. A sequence is CM over [0, N] iff conditioned on the state at time 0(N), the sequence is Markov over (0, N] ([0, N)). The Markov sequence and the reciprocal sequence are two important classes of CM sequences.

CM processes were introduced in [14] for Gaussian processes based on mean and covariance functions. Also, stationary Gaussian CM processes were characterized, and construction of some nonstationary Gaussian CM processes was discussed. [15] extended the definition of Gaussian CM processes (presented in [14]) to the general (Gaussian/non-Gaussian) case. The relationship between the CM process and the reciprocal process was studied in [16], [15], [17]. Moreover, it was shown how continuous time Gaussian CM processes can be represented in terms of a Wiener process and an uncorrelated Gaussian random vector [15], [14]. In [18], different (Gaussian/non-Gaussian) CM sequences were defined based on conditioning at the first or the last index (time) of the CM interval, and (stationary/non-stationary) NG CM sequences were studied. Also, dynamic models and characterizations of NG CM sequences were presented.

Reciprocal processes were introduced in [19] in connection to a problem posed by E. Schrodinger [20]–[21]. Later, they were studied more in [22]–[48] and others. A dynamic model and a characterization of the NG reciprocal sequence were presented in [34]. It was shown that the evolution of a reciprocal sequence can be described by a second-order nearest-neighbor model driven by locally correlated dynamic noise [34]. That model is interesting since it can be considered a generalization of the Markov model. However, due to the correlation of the dynamic noise as well as the nearest-neighbor structure, it is not necessarily easy to apply [12]–[13]. A dynamic model (governing the NG reciprocal sequence) was presented in [16] from the viewpoint of the CM sequence. This dynamic model, called the reciprocal CM_L dynamic model, is driven by white noise and is easily applicable.

The contributions of this paper are as follows. In [16], it was not discussed how the parameters of the reciprocal CM_L model can be designed in application. In this paper, an approach is presented for designing these parameters. Also, inspired by [15], a representation of NG CM sequences in terms of a NG Markov sequence and an uncorrelated NG vector is obtained. In addition, construction of a NG CM sequence based on a NG Markov sequence and an uncorrelated NG vector is presented. This representation and construction of CM sequences can be used for designing parameters of CM_L and CM_F dynamic models based on those of the Markov model. Besides providing a better insight into CM and reciprocal sequences, the results of this paper make these sequences more easily applicable.

The paper is organized as follows. Section 2 reviews some definitions and results required for later sections. In Section 3, an approach is presented for design of parameters of the reciprocal CM_L model. In Section 4, a representation (construction) of a NG CM sequence as a sum of a NG Markov sequence and an uncorrelated NG vector is discussed. Section 5 contains a summary and conclusions.

2 Definitions and Preliminaries

i, j

2.1 Conventions

Throughout the paper we consider stochastic sequences defined over the interval [0, N], which is a general discrete index interval. For convenience this discrete index is called time. The following conventions are used throughout the paper:

$$\begin{split} [i,j] &\triangleq \{i,i+1,\cdots,j-1,j\} \\ (i,j) &\triangleq \{i+1,i+2,\cdots,j-2,j-1\} \\ [x_k]_i^j &\triangleq \{x_k,k \in [i,j]\} \\ [x_k] &\triangleq [x_k]_0^N \\ [x_k]_J &\triangleq \{x_k,k \in J\}, \quad J \subset [0,N] \\ ,k_1,k_2,l_1,l_2 \in [0,N], k_1 < k_2, i < j \end{split}$$

where k in $[x_k]_i^j$ is a dummy variable. We use "\" for set subtraction. C_{l_1,l_2} is a covariance function. C is the covariance matrix of the whole sequence $[x_k]$. Also, 0 may denote a zero scalar, vector, or matrix, as is clear from the context. $F(\cdot|\cdot)$ denotes the conditional cumulative distribution function (CDF).

The abbreviations ZMNG and NG are used for "zero-mean nonsingular Gaussian" and "nonsingular Gaussian", respectively.

2.2 Definitions and Notations

CM sequences are defined as follows. A sequence $[x_k]$ is $CM_c, c \in \{0, N\}$, iff conditioned on the state at time 0 (or N), the sequence is Markov over (0, N] ([0, N)). The above definition is equivalent to the following lemma [18].

Lemma 2.1. $[x_k]$ is $CM_c, c \in \{0, N\}$, iff

$$F(\xi_k | [x_i]_0^j, x_c) = F(\xi_k | x_j, x_c)$$
⁽¹⁾

for every $j, k \in [0, N], j < k$, and every $\xi_k \in \mathbb{R}^d$, where d is the dimension of x_k .

Remark 2.2. For the forward direction, we have

$$CM_c = \begin{cases} CM_F & ifc = 0\\ CM_L & ifc = N \end{cases}$$

where the subscript "F" or "L" is used because the conditioning is at the first or last time.

Forward and backward directions are parallel. So we only consider the forward one.

The reciprocal sequence is defined as follows. A sequence is reciprocal iff the subsequences inside and outside the interval $[k_1, k_2]$ are independent conditioned on the boundaries x_{k_1} and x_{k_2} $(\forall k_1, k_2 \in [0, N])$. The above definition is equivalent to the following lemma [16], [23].

Lemma 2.3. $[x_k]$ is reciprocal iff

$$F(\xi_k | [x_i]_0^j, [x_i]_l^N) = F(\xi_k | x_j, x_l)$$
⁽²⁾

(3)

for every $j, k, l \in [0, N]$ (j < k < l), and every $\xi_k \in \mathbb{R}^d$, where d is the dimension of x_k . Lemma 2.4. $[x_k]$ is Markov iff

$$F(\xi_k | [x_i]_0^j) = F(\xi_k | x_j)$$

for every $j, k \in [0, N], j < k$, and every $\xi_k \in \mathbb{R}^d$, where d is the dimension of x_k .

2.3 Preliminaries

We review some results for later sections [18], [16], [34], [49].

Definition 2.5. A symmetric positive definite matrix is called CM_L if it has form (4) and CM_F if it has form (5).

$$\begin{bmatrix} A_{0} & B_{0} & 0 & \cdots & 0 & 0 & D_{0} \\ B'_{0} & A_{1} & B_{1} & 0 & \cdots & 0 & D_{1} \\ 0 & B'_{1} & A_{2} & B_{2} & \cdots & 0 & D_{2} \\ \vdots & \vdots & \vdots & \vdots & \vdots & \vdots & \vdots \\ 0 & \cdots & 0 & B'_{N-3} & A_{N-2} & B_{N-2} & D_{N-2} \\ 0 & \cdots & 0 & 0 & B'_{N-2} & A_{N-1} & B_{N-1} \\ D'_{0} & D'_{1} & D'_{2} & \cdots & D'_{N-2} & B'_{N-1} & A_{N} \end{bmatrix}$$

$$\begin{bmatrix} A_{0} & B_{0} & D_{2} & \cdots & D_{N-2} & D_{N-1} & D_{N} \\ B'_{0} & A_{1} & B_{1} & 0 & \cdots & 0 & 0 \\ D'_{2} & B'_{1} & A_{2} & B_{2} & \cdots & 0 & 0 \\ \vdots & \vdots & \vdots & \vdots & \vdots & \vdots & \vdots \\ D'_{N-2} & \cdots & 0 & B'_{N-3} & A_{N-2} & B_{N-2} & 0 \\ D'_{N-1} & \cdots & 0 & 0 & B'_{N-2} & A_{N-1} & B_{N-1} \\ D'_{N} & 0 & 0 & \cdots & 0 & B'_{N-1} & A_{N} \end{bmatrix}$$

$$(4)$$

Definition 2.6. A matrix is called cyclic tri-diagonal if it has form (6).

$$\begin{bmatrix} A_0 & B_0 & 0 & \cdots & 0 & 0 & D_0 \\ B'_0 & A_1 & B_1 & 0 & \cdots & 0 & 0 \\ 0 & B'_1 & A_2 & B_2 & \cdots & 0 & 0 \\ \vdots & \vdots & \vdots & \vdots & \vdots & \vdots & \vdots \\ 0 & \cdots & 0 & B'_{N-3} & A_{N-2} & B_{N-2} & 0 \\ 0 & \cdots & 0 & 0 & B'_{N-2} & A_{N-1} & B_{N-1} \\ D'_0 & 0 & 0 & \cdots & 0 & B'_{N-1} & A_N \end{bmatrix}$$
(6)

Definition 2.7. A matrix is called tri-diagonal if it has form (7).

$$\begin{bmatrix} A_0 & B_0 & 0 & \cdots & 0 & 0 & 0 \\ B'_0 & A_1 & B_1 & 0 & \cdots & 0 & 0 \\ 0 & B'_1 & A_2 & B_2 & \cdots & 0 & 0 \\ \vdots & \vdots & \vdots & \vdots & \vdots & \vdots & \vdots \\ 0 & \cdots & 0 & B'_{N-3} & A_{N-2} & B_{N-2} & 0 \\ 0 & \cdots & 0 & 0 & B'_{N-2} & A_{N-1} & B_{N-1} \\ 0 & 0 & 0 & \cdots & 0 & B'_{N-1} & A_N \end{bmatrix}$$

$$(7)$$

 A_k , B_k , and D_k are matrices in general.

Remark 2.8. We use CM_c to mean either CM_L or CM_F or both. **Theorem 2.9.** (i) A NG sequence with covariance matrix C is CM_c iff C^{-1} has the CM_c form. (ii) A NG sequence with covariance matrix C is reciprocal iff C^{-1} is cyclic tri-diagonal. (iii) A NG sequence with covariance matrix C is Markov iff C^{-1} is tri-diagonal.

The following corollary follows from the characterizations presented in Theorem 2.9 [16]. Corollary 2.10. A NG sequence is reciprocal iff it is both CM_L and CM_F .

A CM_c dynamic model is as follows.

Theorem 2.11. A ZMNG $[x_k]$ with covariance function C_{l_1,l_2} is CM_c iff it is governed by

$$x_k = G_{k,k-1}x_{k-1} + G_{k,c}x_c + e_k, \quad k \in (0,N] \setminus \{c\}$$
(8)

where $[e_k]$ is a zero-mean white Gaussian sequence with nonsingular covariances G_k , along with boundary condition¹

$$x_0 = e_0, \quad x_c = G_{c,0}x_0 + e_c \text{ (for } c = N)$$
(9)

or equivalently

$$x_c = e_c, \quad x_0 = G_{0,c} x_c + e_0 \text{ (for } c = N)$$
 (10)

The following theorem presents the reciprocal CM_c models [16].

Theorem 2.12. A ZMNG $[x_k]$ with covariance function C_{l_1,l_2} is reciprocal iff it is governed by (8) along with (9) or (10), and

$$G_k^{-1}G_{k,c} = G'_{k+1,k}G_{k+1}^{-1}G_{k+1,c}$$
(11)

 $\forall k \in (0, N-1)$ for c = N, and $\forall k \in (1, N)$ for c = 0. Moreover, $[x_k]$ is Markov iff additionally we have, for c = N,

$$G_N^{-1}G_{N,0} = G_{1,N}'G_1^{-1}G_{1,0}$$
(12)

for (9) or

$$G_0^{-1}G_{0,N} = G_{1,0}'G_1^{-1}G_{1,N}$$
(13)

for (10); for c = 0, we have

$$G_{N,0} = 0 \tag{14}$$

 $^{{}^{1}}e_{0}$ and e_{N} in (9) are not necessarily the same as e_{0} and e_{N} in (10). Just for simplicity we use the same notation.

3 A Dynamic Model of Reciprocal Sequences

Based on Theorem 2.12, we can determine whether a given CM_c model governs a reciprocal sequence or not. Also, Theorem 2.12 gives the required conditions on the parameters of a reciprocal CM_c model for their design. Theorem 3.1 below provides an approach to parameter design of the reciprocal CM_L model. We have a similar result for a backward CM_L model (which is useful for the CM_F sequence, because a CM_F sequence is backward CM_L).

Theorem 3.1. A ZMNG sequence $[x_k]$ with covariance function C_{l_1,l_2} is reciprocal iff it is governed by (8) (for c = N) along with (9) or (10), where $(G_{k,k-1}, G_{k,N}, G_k)$, $k \in (0, N)$, are given by

$$G_{k,k-1} = M_{k,k-1} - G_{k,N}M_{N|k-1}$$
(15)

$$G_{k,N} = G_k M'_{N|k} C_{N|k}^{-1}$$
(16)

$$G_k = (M_k^{-1} + M_{N|k}' C_{N|k}^{-1} M_{N|k})^{-1}$$
(17)

$$M_{N|k} = M_{N,N-1} \cdots M_{k+1,k}, \quad k \in (0,N)$$

and $M_{N|N} = I$. Also,

$$C_{N|k} = \sum_{n=k}^{N-1} M_{N|n+1} M_{n+1} M'_{N|n+1}, \quad k \in (0, N)$$

where $M_{k,k-1}$, $k \in (0, N]$, are square matrices, and M_k , $k \in (0, N]$, are positive definite of the dimension of x_k .

Proof. We only show how (15)–(17) are obtained. More detail can be found in [50].

Given the square matrices $M_{k,k-1}$ and the positive definite matrices $M_k, k \in (0, N]$, there exists a ZMNG Markov sequence $[y_k]$ governed by

$$y_k = M_{k,k-1}y_{k-1} + e_k^M, \quad k \in (0,N]$$
(18)

where $[e_k^M]_1^N$ is a zero-mean white Gaussian sequence with nonsingular covariances M_k , uncorrelated with y_0 with nonsingular covariance C_0^y .

Since every Markov sequence is CM_L , one can obtain the following model for $[y_k]$

$$y_k = G_{k,k-1}y_{k-1} + G_{k,N}y_N + e_k^y, \quad k \in (0,N)$$
(19)

where $[e_k^y]$ is a zero-mean white Gaussian sequence with nonsingular covariances G_k , and boundary condition

$$y_N = e_N^y, \quad y_0 = G_{0,N}^y y_N + e_0^y$$
 (20)

Parameters of model (19) can be obtained as follows. Since

$$p(y_k|y_{k-1}) = \mathcal{N}(y_k; M_{k,k-1}y_{k-1}, M_k)$$

by the Markov property, we have

$$p(y_k|y_{k-1}, y_N) = \frac{p(y_k|y_{k-1})p(y_N|y_k)}{p(y_N|y_{k-1})}$$

$$= \mathcal{N}(y_k; G_{k,k-1}y_{k-1} + G_{k,N}y_N; G_k)$$
(21)

 $\forall k \in (0, N)$, and $G_{k,k-1}$, $G_{k,N}$, and G_k are calculated by (15)–(17), where for $i \in [0, N)$ we have $p(y_N|y_i) = \mathcal{N}(y_N; M_{N|i}y_i, C_{N|i})$

Now, we construct a sequence $[x_k]$ governed by the same model (19) as

$$x_k = G_{k,k-1}x_{k-1} + G_{k,N}x_N + e_k, \quad k \in (0,N)$$
(22)

and boundary condition

$$x_N = e_N, \quad x_0 = G_{0,N} x_N + e_0 \tag{23}$$

Note that parameters of (19) and (22) are the same (i.e., $G_{k,k-1}, G_{k,N}, G_k, k \in (0, N)$), but parameters of (20) (i.e., $G_{0,N}^y, G_0^y, G_N^y$) and (23) (i.e., $G_{0,N}, G_0, G_N$) are different. By Theorem 2.11, $[x_k]$ is a ZMNG CM_L sequence. Then, based on the above results, necessity and sufficiency of the theorem are proved [50].

By Theorem 3.1, we can first design $M_{k,k-1}$ and M_k , $k \in (0, N]$, and then parameters of the reciprocal CM_L model. This approach is useful because design of the parameters of a Markov model is much easier (than of a reciprocal CM_L model) and $M_{k,k-1}$ and M_k can be seen as parameters of a Markov model. It is explained for the problem of motion trajectory modeling with destination information as follows. Let $[y_k]$ be a Markov sequence governed by (18) (for example, a nearly constant velocity motion model without considering destination information) with parameters $(M_{k,k-1}, M_k), k \in (0, N]$ (and the initial covariance C_0^y), and $[x_k]$ be a CM_L sequence governed by (8) with parameters $(G_{k,k-1}, G_{k,N}, G_k), k \in (0, N)$ given by (15)–(17) (and any boundary condition $(G_0, G_{N,0}, G_N)$ or $(G_N, G_{0,N}, G_0)$). It can be seen that $[x_k]$ and $[y_k]$ have the same CM_L evolution law ((19) and (22)) while $[x_k]$ can have any joint endpoint distribution. Therefore, $[x_k]$ can model trajectories with any origin and destination distributions, where parameters of its CM_L model are designed based on those of the Markov model. In the above example, on the one hand it is assumed that the moving object follows the Markov model, and on the other hand it is assumed that the destination (distribution) is known (this is the case in a real problem of trajectory modeling with destination information). Considering these two assumptions (i.e., motion Markov model and destination), the corresponding CM_L model (with (15)–(17)) describes the behavior of the moving object satisfying both the above assumptions (also, see (21)).

The dynamic model (22) is called a CM_L model *induced* by a Markov model. By Theorem 3.1, every CM_L model induced by a Markov model is a reciprocal CM_L model. Also, every reciprocal CM_L model can be induced by a Markov model.

4 Representations of CM and Reciprocal Sequences

A representation of continuous time Gaussian CM processes by a Wiener process and an uncorrelated Gaussian vector was presented in [15]. Inspired by [15], we show how a NG CM sequence can be represented by a NG Markov sequence and an uncorrelated NG vector. Also, it is shown how to use a NG Markov sequence and an uncorrelated NG vector to construct a NG CM sequence. This representation not only gives more insight into the CM sequence, but also provides an approach for designing CM sequences in application.

Proposition 4.1. A ZMNG $[x_k]$ is CM_c iff it can be represented as

$$x_k = y_k + \Gamma_k x_c, \quad k \in [0, N] \setminus \{c\}$$
(24)

where $[y_k]_{[0,N]\setminus\{c\}}$ is a ZMNG Markov sequence, x_c is a ZMNG vector uncorrelated with $[y_k]_{[0,N]\setminus\{c\}}$, and Γ_k are some matrices.

Proof. We only show that given a ZMNG Markov sequence $[y_k]_0^{N-1}$ uncorrelated with a ZMNG vector x_N , $[x_k]$ constructed by $x_k = y_k + \Gamma_k x_N$, $k \in [0, N)$ is a ZMNG CM_L sequence, where Γ_k are some matrices. More detail can be found in [50]. It suffices to show that $[x_k]$ is governed by (8) and (10) (c = N).

Since $[y_k]_0^{N-1}$ is a ZMNG Markov sequence, we have

$$y_k = M_{k,k-1}y_{k-1} + e_k^M, \quad k \in (0,N)$$
(25)

where $[e_k^M]_1^{N-1}$ is a zero-mean white Gaussian sequence with nonsingular covariances M_k , uncorrelated with y_0 with nonsingular covariance C_0^y . We have

$$x_0 = y_0 + \Gamma_0 x_N$$

So, we consider $G_{0,N} = \Gamma_0$ and $e_0 = y_0$. Then, for $k \in (0, N)$, we have
$$x_k = y_k + \Gamma_k x_N$$
$$= M_{k,k-1}y_{k-1} + e_k^M + \Gamma_k x_N$$
$$= M_{k,k-1}x_{k-1} + (\Gamma_k - M_{k,k-1}\Gamma_{k-1})x_N + e_k^M$$

Thus, for $k \in (0, N)$, we consider

$$G_{k,k-1} = M_{k,k-1}$$

$$G_{k,N} = \Gamma_k - M_{k,k-1}\Gamma_{k-1}$$

$$e_k = e_k^M$$

Nonsingularity of $[x_k]$ can be proved based on the nonsingularity of $[y_k]_0^{N-1}$ and x_N .

Proposition 4.1 makes a key concept behind the CM_c sequence clear. Also, it provides an approach for design of CM_c models in application. Below we explain the idea for designing a CM_L model for motion trajectory modeling with destination information. The CM_L model is more general than the reciprocal CM_L model. Consequently, the following approach for parameter design is more general and flexible than that of Theorem 3.1. For example, it can incorporate other available information in the problem. The approach is as follows. First, a Markov model (e.g., a nearly constant velocity model) with the given origin distribution (without considering other information) is considered. The sequence governed by this model is $[y_k]_0^{N-1}$ in (24) (let c = N)). Assuming the destination, distribution of x_N , is known. Then, the matrices Γ_k are designed according to the information available in the problem. In other words, based on Γ_k , the Markov sequence $[y_k]_0^{N-1}$ is modified to satisfy the available information leading to the desired trajectories which end up at the destination. A direct attempt for designing parameters of a CM_L model for this problem is hard. However, the above approach based on Proposition 4.1 provides some guidelines which make parameter design easier and intuitive.

A representation of the reciprocal sequence is as follows.

Proposition 4.2. A ZMNG $[x_k]$ is reciprocal iff it can be represented as both

$$x_k = y_k^L + \Gamma_k^L x_N, \quad k \in [0, N)$$
(26)

$$x_k = y_k^F + \Gamma_k^F x_0, \quad k \in (0, N]$$

$$\tag{27}$$

where $[y_k^L]_0^{N-1}$ and $[y_k^F]_1^N$ are ZMNG Markov sequences, x_N and x_0 are ZMNG vectors uncorrelated with $[y_k^L]_0^{N-1}$ and $[y_k^F]_1^N$, respectively, and Γ_k^L and Γ_k^F are some matrices.

5 Summary and Conclusions

An approach has been presented for designing parameters of reciprocal CM_L dynamic models. This approach makes the nonsingular Gaussian (NG) reciprocal sequence more easily applicable. The approach is based on inducing a CM_L model from a Markov model. It was shown that every CM_L model induced by a Markov model is a reciprocal CM_L model. Also, every reciprocal CM_L model can be induced by a Markov model.

It has been shown how a NG CM (or reciprocal) sequence can be represented as a sum of a NG Markov sequence and an uncorrelated NG vector. In addition, construction of a NG CM sequence based on a NG Markov sequence and an uncorrelated NG vector is discussed. This representation and construction can be used for designing parameters of the CM dynamic models in application (based on designing parameters of a NG Markov model and an uncorrelated NG vector). The results of this paper lead to more insight into CM and reciprocal sequences, provide some tools for application of CM (and reciprocal) sequences, and demonstrate the benefits of studying the reciprocal sequence from the viewpoint of the CM sequence.

Acknowledgments

Research was supported by NASA Phase03-06 through grant NNX13AD29A.

References

- B. Levy and A. J. Krener. Dynamics and Kinematics of Reciprocal Diffusions," *Journal of Mathematical Physics*. Vol. 34, No. 5, pp. 1846-1875, 1993.
- [2] B. Levy and A. J. Krener. Stochastic Mechanics of Reciprocal Diffusions. *Journal of Mathematical Physics*. Vol. 37, No. 2, pp. 769-802, 1996.
- [3] A. Chiuso, A. Ferrante, and G. Picci. Reciprocal Realization and Modeling of Textured Images. 44th IEEE Conference on Decision and Control, Seville, Spain, Dec. 2005.
- [4] G. Picci and F. Carli. Modelling and Simulation of Images by Reciprocal Processes. *Tenth International Conference on Computer Modeling and Simulation*, Cambridge, UK, Apr. 2008.

- [5] A. J. Krener. Reciprocal Processes and the Stochastic Realization Problem for Acausal Systems. Modeling, Identification, and Robust Control, C. I. Byrnes and A. Lindquist (editors), Elsevier, 1986.
- [6] M. Fanaswala, V. Krishnamurthy, and L. White. Destination-aware Target Tracking via Syntactic Signal Processing. *IEEE International Conference on Acoustics, Speech and Signal Processing (ICASSP)*, Prague, Czech Republic, May 2011.
- [7] M. Fanaswala and V. Krishnamurthy. Spatiotemporal trajectory models for metalevel target tracking. IEEE Aerospace and Electronic Systems Magazine, Vol. 30, No. 1, pp. 16-31, 2015.
- [8] M. Fanaswala and V. Krishnamurthy. Detection of Anomalous Trajectory Patterns in Target Tracking via Stochastic Context-Free Grammer and Reciprocal Process Models. *IEEE Journal of Selected Topics in Signal Processing*, Vol. 7, No. 1, pp. 76-90, 2013.
- [9] G. Stamatescu, L. B. White, and R. Bruce-Doust. Track Extraction With Hidden Reciprocal Chains. *IEEE Trans. on Automatic Control*, Vol. 63, No. 4, pp. 1097-1104, 2018.
- [10] B. I. Ahmad, J. K. Murphy, S. J. Godsill, P. M. Langdon, and R. Hardy. Intelligent Interactive Displays in Vehicles with Intent Prediction: A Bayesian framework. *IEEE Signal Processing Magazine*, Vol. 34, No. 2, 2017.
- [11] B. I. Ahmad, J. Murphy, P. M. Langdon, R. Hardy, and S. J. Godsill. Destination Inference Using Bridging Distributions. *IEEE International Conference on Acoustics, Speech and Signal Processing (ICASSP)*, Brisbane, QLD, Australia, Apr. 2015.
- [12] R. Rezaie and X. R. Li. Destination-Directed Trajectory Modeling and Prediction Using Conditionally Markov Sequences. *IEEE Western New York Image and Signal Processing Workshop*, Rochester, NY, USA, Oct. 2018, pp. 1-5.
- [13] R. Rezaie and X. R. Li. Trajectory Modeling and Prediction with Waypoint Information Using a Conditionally Markov Sequence. 56th Allerton Conference on Communication, Control, and Computing, Monticello, IL, USA, Oct. 2018, pp. 486-493.
- [14] C. B. Mehr and J. A. McFadden, Certain Properties of Gaussian Processes and their First-Passage Times. *Journal of Royal Statistical Society (B)*, Vol. 27, pp. 505-522, 1965.
- [15] J. Abraham and J. Thomas. Some Comments on Conditionally Markov and Reciprocal Gaussian Processes. *IEEE Trans. on Information Theory*. Vol. 27, No. 4, July 1981.
- [16] R. Rezaie and X. R. Li. Gaussian Reciprocal Sequences from the Viewpoint of Conditionally Markov Sequences. *Inter. Conference on Vision, Image and Signal Processing*, Las Vegas, NV, USA, Aug. 2018., pp. 33:1-33:6.
- [17] J-P Carmichael, J-C Masse, and R. Theodorescu. Representations for Multivariate Reciprocal Gaussian Processes. *IEEE Trans. on Inform. Theory*, Vol. 34, No. 1, pp. 155-157, 1988.
- [18] R. Rezaie and X. R. Li. Nonsingular Gaussian Conditionally Markov Sequences. *IEEE West. New York Image and Signal Processing Workshop*. Rochester, NY, USA, Oct. 2018, pp. 1-5.
- [19] S. Bernstein. Sur les liaisons entre les grandeurs aleatoires. *Verh. des intern. Mathematikerkongr I*, Zurich, 1932.
- [20] E. Schrodinger. Uber die Umkehrung der Naturgesetze. Sitz. Ber. der Preuss. Akad. Wissen., Berlin Phys. Math. 144, 1931.
- [21] E. Schrodinger. Theorie relativiste de l'electron et l'interpretation de la mechanique quantique. *Ann. Inst. H. Poincare* 2, 269-310, 1932.
- [22] D. Slepian. First Passage Time for a Particular Gaussian Process. Annals of Mathematical Statistics, Vol. 32, pp. 610-612, 1961.
- [23] B. Jamison. Reciprocal Processes: The Stationary Gaussian Case. Annals of Mathematical Statistics, Vol. 41, No. 5, pp. 1624-1630, 1970.
- [24] S. C. Chay, On Quasi-Markov Random Fields, Journal of Multivariate Analysis, Vol. 2, pp. 14-76, 1972.
- [25] J-P Carmichael, J-C Masse, and R. Theodorescu, Processus Gaussiens Stationnaires Reciproques sur un Intervalle, C. R. Acad. Sc. Paris, t. 295 (27 Sep 1982).

- [26] B. Jamison. Reciprocal Processes. Z. Wahrscheinlichkeitstheorie verw. Gebiete, vol. 30, pp. 65-86, 1974.
- [27] C. Leonard, S. Rœlly, and J-C Zambrini. Reciprocal Processes. A Measure-theoretical Point of View. *Probability Surveys*, Vol. 11, pp. 237–269, 2014.
- [28] G. Conforti, P. Dai Pra, and S. Roelly. Reciprocal Class of Jump Processes. *Journal of Theoretical Probability*, June 2014.
- [29] R. Murr, *Reciprocal Classes of Markov Processes. An Approach with Duality Formulae*, Ph.D. thesis, Universitat Potsdam, 2012.
- [30] S. Rœlly. Reciprocal Processes. A Stochastic Analysis Approach. In V. Korolyuk, N. Limnios, Y. Mishura, L. Sakhno, and G. Shevchenko, editors, Modern Stochastics and Applications, volume 90 of Optimization and Its Applications, pp. 53–67. Springer, 2014.
- [31] J. Chen and H. L. Weinert, A New Characterization of Multivariate Gaussian Reciprocal Processes. *IEEE Trans. on Automatic Control*, Vol. 38, No. 10, pp. 1601-1602, 1993.
- [32] F. P. Carli, A. Ferrante, M. Pavon, G. Picci. A Maximum Entropy Solution of the Covariance Extension Problem for Reciprocal Processes. *IEEE Trans. on Automatic Control*, Vol. 56, No. 9, pp. 1999-2012, 2011.
- [33] B. C. Levy, A. Beghi, Discrete-time Gauss-Markov Processes with Fixed Reciprocal Dynamics. *Journal of Mathematical Systems, Estimation, and Control*, Vol. 4, No. 3, pp. 1-25, 1994.
- [34] B. C. Levy, R. Frezza, and A. Krener. Modeling and Estimation of Discrete-Time Gaussian Reciprocal Processes. *IEEE Trans. on Automatic Control.* Vol. 35, No. 9, pp. 1013-1023, 1990.
- [35] A. Beghi, Continuous-time Gauss-Markov Processes with Fixed Reciprocal Dynamics. Journal of Mathematical Systems, Estimation, and Control, Vol. 4, No. 4, pp. 1-24, 1994.
- [36] A. J. Krener. Reciprocal Diffusions and Stochastic Differential Equations of Second Order. *Stochastics*, Vol. 24, No. 4, pp. 393-422, 1988.
- [37] A. J. Krener, R. Frezza, and B. C. Levy. Gaussian Reciprocal Processes and Self-adjoint Stochastic Differential Equations of Second Order. *Stochastics and Stochastic Reports*, Vol. 34, No. 1-2, pp. 29-56, 1991.
- [38] F. Carravetta and L. B. White. Modelling and Estimation for Finite State Reciprocal Processes. *IEEE Trans. on Automatic Control*, Vol. 57, No. 9, pp. 2190-2202, 2012.
- [39] F. Carravetta. Nearest-neighbour Modelling of Reciprocal Chains. An International Journal of Probability and Stochastic Processes, Vol. 80, No. 6, pp. 525-584, 2008.
- [40] L. B. White and F. Carravetta. Optimal Smoothing for Finite State Hidden Reciprocal Processes. *IEEE Trans. on Automatic Control*, Vol. 56, No. 9, pp. 2156-2161, 2011.
- [41] L B. White and H. X. Vu. Maximum Likelihood Sequence Estimation for Hidden Reciprocal Processes. *IEEE Trans. on Automatic Control*, Vol. 58, No. 10, pp. 2670-2674, 2013.
- [42] E. Baccarelli and R. Cusani. Recursive Filtering and Smoothing for Gaussian Reciprocal Processes with Dirichlet Boundary Conditions. *IEEE Trans. on Signal Processing*, Vol. 46, No. 3, pp. 790-795, 1998.
- [43] E. Baccarelli, R. Cusani, and G. Di Blasio. Recursive filtering and smoothing for reciprocal Gaussian processes-pinned boundary case. *IEEE Trans. on Information Theory*, Vol. 41, No. 1, pp. 334-337, 1995.
- [44] D. Vats and J. M. F. Moura. Recursive Filtering and Smoothing for Discrete Index Gaussian Reciprocal Processes. 43rd Annual Conference on Information Sciences and Systems, Baltimore, MD, USA, Mar. 2009.
- [45] R. Rezaie and X. R. Li. Gaussian Conditionally Markov Sequences: Algebraically Equivalent Dynamic Models. *IEEE Trans. on Aerospace and Electronic Systems*, 2019, DOI: 10.1109/TAES.2019.2951188.
- [46] R. Rezaie and X. R. Li. Gaussian Conditionally Markov Sequences: Singular/Nonsingular. *IEEE Trans.* on Automatic Control, 2019, DOI: 10.1109/TAC.2019.2944363.
- [47] R. Rezaie. Gaussian Conditionally Markov Sequences: Theory with Application. Ph.D. Dissertation, Dept of Electrical Engineering, University of New Orleans, July 2019.

- [48] R. Rezaie and X. R. Li. Explicitly Sample-Equivalent Dynamic Models for Gaussian Conditionally Markov, Reciprocal, and Markov Sequences. *Inter. Conf. on Control, Automation, Robotics, and Vision Engineering*, New Orleans, LA, USA, Nov. 2018, pp. 1-6.
- [49] R. Ackner, T. Kailath. Discrete-Time Complementary Models and Smoothing. International Journal of Control, Vol. 49, No. 5, pp. 1665-1682, May 1989.
- [50] R. Rezaie and X. R. Li. Gaussian Conditionally Markov Sequences: Dynamic Models and Representations of Reciprocal and other Classes. *IEEE Transactions on Signal Processing*, 2019, DOI: 10.1109/TSP.2019.2919410.