HOLLOW POLYTOPES OF LARGE WIDTH

GIULIA CODENOTTI AND FRANCISCO SANTOS

ABSTRACT. We construct the first known hollow lattice polytopes of width larger than dimension: a hollow lattice polytope (resp. a hollow lattice simplex) of dimension 14 (resp. 404) and of width 15 (resp. 408). We also construct a hollow (non-lattice) tetrahedron of width $2 + \sqrt{2}$ and conjecture that this is the maximum width among 3-dimensional hollow convex bodies.

We show that the maximum lattice width grows (at least) additively with d. In particular, the constructions above imply the existence of hollow lattice polytopes (resp. hollow simplices) of arbitrarily large dimension d and width $\simeq 1.14d$ (resp. $\simeq 1.01d$).

1. INTRODUCTION

A convex body $K \subset \mathbb{R}^d$ is called *hollow* or *lattice-free* with respect to a lattice $\Lambda \cong \mathbb{Z}^d$ if $\text{int}(K) \cap \Lambda = \emptyset$. The *width* of K with respect to a linear functional $f \in (\mathbb{R}^d)^*$ is the length of the segment $f(K)$. We denote it width (K, f) . The *lattice* width of K is

$$
\mathrm{width}_{\Lambda}(K) := \inf_{f \in \Lambda^* \setminus 0} \mathrm{width}(K, f).
$$

We omit Λ and write just width (K) when this creates no ambiguity. We also say width of K meaning lattice width.

The celebrated flatness theorem states that hollow bodies in fixed dimension d have bounded lattice width. That is, for each fixed d , the supremum width among all hollow convex bodies in \mathbb{R}^d is a certain constant $w_c(d) < \infty$. We are also interested in the following specializations of w_c . We call $w_p(d)$ the maximum width among all hollow *lattice d-polytopes*, $w_s(d)$ the maximum width among hollow lattice d-simplices and $w_e(d)$ the maximum width among empty d-simplices; here a lattice simplex is empty if its only lattice points are its vertices. Observe that these specializations take integer values. Obviously,

$$
w_e(d) \le w_s(d) \le w_p(d) \le w_c(d).
$$

Much work has been done in finding upper bounds for $w_c(d)$ (see references, e.g., in the introductions to [\[6,](#page-14-0) [12\]](#page-14-1)). The current best upper bound is $w_c(d) \in$ $O^*(d^{4/3})$ [\[14\]](#page-14-2), where the notation $O($)^{*} denotes that a polylog factor is neglected. Better upper bounds are known for restricted classes of convex bodies. For example, it is known that the maximum width of hollow (not necessarily lattice) simplices [\[6\]](#page-14-0)

Date: June 7, 2019.

²⁰¹⁰ Mathematics Subject Classification. Primary 52C07, 52B20; Secondary 52C17.

The authors were supported by the Einstein Foundation Berlin under grant EVF-2015-230 and, while they were in residence at the Mathematical Sciences Research Institute in Berkeley, California during the Fall 2017 semester, by the Clay Institute and the National Science Foundation (Grant No. DMS-1440140). Work of F. Santos is also supported by project MTM2017-83750-P of the Spanish Ministry of Science (AEI/FEDER, UE).

and of centrally symmetric hollow bodies $[5]$ is in $O(n \log n)$. But work on lower bounds is very scarce. To the best of our knowledge, can be summarized as follows:

- Since the d-th dilation of a unimodular d-simplex is hollow and has width $d, w_s(d) \geq d$.
- Sebő [\[15\]](#page-14-4) showed $w_e(d) \geq d-2$.
- Conway and Thompson (see $[13,$ Theorem I.9.5]) showed a lower bound of $\Omega(d)$ for the maximum width of hollow ellipsoids.
- Dash et al. [\[7\]](#page-14-6) (Theorem 3.2 and the paragraphs before it) show that

$$
3.1547\ldots = 2 + \frac{2}{\sqrt{3}} \le w_s(3) \le w_c(3) \le 1 + \frac{2}{\sqrt{3}} + \left(\frac{90}{\pi}\right)^{1/3} = 4.2439\ldots
$$

• The following exact values are known for small d:

In this paper we establish some new lower bounds, both for specific dimensions (Sections [4](#page-5-0) to [6\)](#page-10-0) and in the asymptotic sense (Section [7\)](#page-11-0).

Equals 4 to 0) and in the asymptotic sense (Section 7).
More precisely, in Section [5](#page-7-0) we show that $w_c(3) \geq 2 + \sqrt{2}$:

Theorem 1.1. There is a hollow (non-lattice) tetrahedron of width $2+\sqrt{2} \simeq 3.4142$.

The tetrahedron of Theorem [1.1](#page-1-1) is symmetric with respect to the fcc-cubic lattice The tetranearon of Theorem 1.1 is symmetric with respect to the rcc-cubic lattice and has width $2 + \sqrt{2}$ for seven different linear functionals (the three coordinates and four diagonals of the cube). To certify that no integer functional gives smaller width to it we develop in Section [3](#page-4-0) a method which may be of independent interest, based on existence of long piecewise-linear paths of rational directions.

This tetrahedron maximizes width among a two-parameter family of hollow tetrahedra that contains *two* of the five existing hollow 3-polytopes of width 3 (see Theorem [5.1\)](#page-8-0), in much the same way as the value of $w_c(2) = 1 + 2/\sqrt{3} \approx 2.1547$ in the table above is attained by optimizing a perturbation of the second dilation of the unimodular triangle (see details in Section [4\)](#page-5-0). This makes us conjecture that:

Conjecture 1.2. The tetrahedron in Theorem [1.1](#page-1-1) is the convex 3-body of largest **Conjecture 1.2.** The tetraned width; that is, $w_c(3) = 2 + \sqrt{2}$.

Similarly, in Sections [4](#page-5-0) and [6,](#page-10-0) we show $w_p(14) \ge 15$ and $w_s(404) \ge 408$:

Theorem 1.3. There is a hollow lattice 14-polytope of width 15 and a hollow lattice 404-simplex of width 408.

We do not know of any hollow lattice d -polytope of width larger than d in previous literature.

Our main technical tool, both in Theorem [1.3](#page-1-2) and for the asymptotic results, is to use dilated *direct sums* of polytopes and convex bodies. Let $C_i \subset \mathbb{R}^{d_i}$, $i = 1, ..., m$,

¹As Sebő points out, for even d the bound can be increased to $d-1$.

be convex bodies containing the origin. Their *direct sum* $[9]$ (sometimes called *free* sum $[1]$) is defined as

$$
C_1 \oplus \cdots \oplus C_m := \left\{ (\lambda_1 x_1, \ldots, \lambda_m x_m) : x_i \in C_i, \lambda_i \geq 0, \sum_{i=1}^m \lambda_i = 1 \right\}.
$$

For a constant $k \in \mathbb{R}_{\geq 0}$, kC denotes the dilation of C by a factor of k. For a given lattice polytope or convex body C containing the origin (not necessarily in the interior) let us denote $C^{\oplus m} = \bigoplus_{i=1}^{m} mC$, the m-fold direct sum of mC with itself. The following proposition is a particular case of Theorem [2.2](#page-3-0) in Section [2.](#page-3-1)

Proposition 1.4. (1) width $(C^{\oplus m}) = m$ width (C) . (2) If C is hollow then $C^{\oplus m}$ is hollow.

As a consequence we have the following statement, proved in Section [7.](#page-11-0)

Theorem 1.5. Let $w_* : \mathbb{N} \to \mathbb{R}$ denote any of the functions w_s , w_p , or w_c . Then

$$
\lim_{d \to \infty} \frac{w_*(d)}{d} = \sup_{d \in \mathbb{N}} \frac{w_*(d)}{d}.
$$

This, in turn, implies our main asymptotic result:

Theorem 1.6.

$$
\lim_{d \to \infty} \frac{w_p(d)}{d} = \lim_{d \to \infty} \frac{w_c(d)}{d} \ge \frac{2 + \sqrt{2}}{3} = 1.138...
$$

$$
\lim_{d \to \infty} \frac{w_s(d)}{d} \ge \frac{102}{101} = 1.0099...
$$

Proof. From Theorem [1.5,](#page-2-0) together with the explicit lower bounds $w_c(3) \geq 2 + \sqrt{2}$ (Theorem [1.1\)](#page-1-1) and $w_s(404) \geq 408$ (Theorem [1.3\)](#page-1-2), we get

$$
\lim_{d \to \infty} \frac{w_c(d)}{d} \ge \frac{2}{3} + \frac{\sqrt{2}}{3}, \qquad \lim_{d \to \infty} \frac{w_s(d)}{d} \ge \frac{102}{101}.
$$

Thus, we only need to show the equality

$$
\sup_{d \in \mathbb{N}} \frac{w_c(d)}{d} = \sup_{d \in \mathbb{N}} \frac{w_p(d)}{d}.
$$

The " \geq " is obvious. For the " \leq ", let C be a hollow convex body such that width $(C)/\dim(C)$ is very close to $\sup_d w_c(d)/d$. We can approximate C arbitrarily by a hollow rational polytope P , and choose an integer m such that mP is a lattice polytope. By Proposition [1.4](#page-2-1) we have that $P^{\oplus m}$ is a hollow lattice polytope of dimension $m \dim(C)$ and

$$
\frac{\text{width}(P^{\oplus m})}{\dim(P^{\oplus m})} = \frac{\text{width}(P)}{\dim(P)} \simeq \frac{\text{width}(C)}{\dim(C)}.
$$

Another implication of our analysis of direct sums together with the values $w_c(2) = 2.1547...$ and $w_c(3) \geq 3.4142...$ is

Proposition 1.7 (Section [7\)](#page-11-0). For every d we have

$$
w_c(d+1) \ge w_c(d) + 1,
$$

\n
$$
w_c(d+2) \ge w_c(d) + 2.1547...,
$$

\n
$$
w_c(d+3) \ge w_c(d) + 3.4142...,
$$

As a consequence,

$$
\frac{w_c(d)}{d} \ge \frac{1}{2} \cdot 2.1547 \cdots = 1.0773 \ldots \qquad \forall d \ge 2.
$$

In Section [8](#page-12-0) we study the width of empty simplices. We do not know whether $\lim_{d\to\infty}\frac{w_e(d)}{d}$ exists. However, we can prove the following slightly weaker result:

Theorem 1.8 (Section [8\)](#page-12-0). For every $d, m \in \mathbb{N}$ we have

$$
w_e(dm) \ge (m-3) w_e(d).
$$

In particular,

$$
\limsup_{d \to \infty} \frac{w_e(d)}{d} = \sup_{d \in \mathbb{N}} \frac{w_e(d)}{d} \ge 1
$$

We do not know whether there is an empty simplex of width larger than its dimension. Yet, Theorem [1.8](#page-3-2) disproves the following guess from $[15, p. 403]$: "it seems to be reasonable to think that the maximum width of an empty integer simplex in \mathbb{R}^n is $n + constant$ " (unless the constant is zero).

We believe our results are a first step towards the main goal concerning flatness lower bounds, which would be to show that $\sup_d w_*(d)/d = \infty$, at least for w_c .

Acknowledgement: We thank Gennadiy Averkov, Benjamin Nill, and an anonymous referee for useful comments on the first version of this paper.

2. Hollow direct sums

Since we will often be using direct sums of polytopes, let us remind the reader of their combinatorial structure.

Lemma 2.1. Let $P = P_1 \oplus \cdots \oplus P_m$ be a direct sum of polytopes. Then:

- (1) If F_i is a face of P_i that does not contain the origin for each i then the join $F_1 * \cdots * F_m$ of them is a face of P that does not contain the origin of dimension $\sum_i \dim(F_i) + m - 1$. All faces of P that do not contain the origin arise in this way.
- (2) If F_i is a face of P_i that contains the origin for each i then the direct sum $F_1\oplus\cdots\oplus F_m$ of them is a face of P that contains the origin of dimension $\sum_i \dim(F_i)$. All faces of P that contain the origin arise in this way.

In particular, the non-zero vertices of P are the points of the form $(0, \ldots, 0, v, \ldots)$ $(0, \ldots, 0)$, with v a non-zero vertex of the corresponding P_i , and 0 is a vertex of P if and only if it is a vertex of every P_i .

Our main technical result is the following theorem. Proposition [1.4](#page-2-1) is the case $C_1 = \cdots = C_m$ and $k_i = m$ of it. Part (4) of Theorem [2.2](#page-3-0) is equivalent to Corollary $5.5(a)$ in [\[1\]](#page-14-12).

Theorem 2.2. Let C_1, \ldots, C_m be convex bodies containing the origin and let $k_1, \ldots, k_m > 0$ be dilation factors. Let

$$
C := \bigoplus_i k_i C_i = k_1 C_1 \oplus \cdots \oplus k_m C_m.
$$

Then:

(1) If k_iC_i is a lattice polytope for every i then C is a lattice polytope.

- (2) If C_i is a simplex with a vertex at the origin for every i then C is a simplex with a vertex at the origin.
- (3) The width of C equals $\min_i \{k_i \text{width}(C_i)\}.$
- (4) If C_i is hollow for every i and $\sum_i \frac{1}{k_i} \ge 1$ then C is hollow.

Proof. Part (1) is obvious, from the description of the vertices of direct sums in Lemma [2.1.](#page-3-3) For part (2) let d_i be the dimension of C_i . Each C_i has d_i non-zero vertices plus the origin so, by the same Lemma, C has $d_1 + \cdots + d_m$ vertices plus the origin. Since C lives in dimension $d_1 + \cdots + d_m$, it must be a simplex.

To prove (3), first note that width $(k_iC_i) = k_i$ width (C_i) , so we can assume w.l.o.g. $k_i = 1$ for all i. Let $f_i \in \Lambda_i^*$ be a lattice direction for which width (C_i) is obtained. Then

$$
\begin{aligned} \text{width}(P) &\leq \text{width}(C, (0, \dots, 0, f_i, 0, \dots, 0)) \\ &= \text{width}(C_i, f_i) = \text{width}(C_i). \end{aligned}
$$

This proves that $\text{width}(C) \leq \min_i \{ \text{width}(C_i) \}.$ For the other inequality, given any lattice functional $g = (g_1, \ldots, g_m) \in \Lambda^* \setminus \{0\} = \bigoplus_i \Lambda_i^* \setminus \{0\}$, we want to show that width $(C, g) \geq$ width (C_i) for some i. For this, let us choose any i with $g_i \neq 0$. Then:

$$
\begin{split} \text{width}(C, g) &= \max_{c, c' \in C} |g^\mathsf{T} c - g^\mathsf{T} c'| \\ &\ge \max_{c_i, c'_i \in C_i} |g^\mathsf{T} (0, \dots, 0, c_i, 0, \dots, 0) - g^\mathsf{T} (0, \dots, 0, c'_i, 0, \dots, 0)| \\ &= \text{width}(C_i, g_i) \ge \text{width}(C_i) \ge \min_j \text{width}(C_j). \end{split}
$$

Finally, to prove part (4) , suppose by contradiction that C is not hollow, and let $c \in \text{int } C \cap \Lambda$. Since $c \in \text{int } C$, we can write $c = (\lambda_1 k_1 c_1, \dots, \lambda_m k_m c_m)$ with $c_i \in \text{int } C_i$ and $\lambda_i > 0$ with $\sum \lambda_i = 1$. On the other hand, since $c \in \Lambda$, we know that each $\lambda_i k_i c_i \in \Lambda_i$. Since C_i is hollow and $c_i \in \text{int } C_i$, we have that $\lambda_i k_i > 1$. This implies $\sum \frac{1}{k_i} < \sum \lambda_i = 1$, contradicting our assumption.

Observe that the assumption that the C_i s contain the origin is no loss of generality: lattice polytopes can be translated to have the origin as a vertex; convex bodies can first be enlarged so that they have lattice points in the boundary, then translated. In both cases, the direct sum C of Theorem [2.2](#page-3-0) can be constructed using these modified C_i s.

3. A certificate for width

In Sections [4–](#page-5-0)[6,](#page-10-0) we construct explicit examples of polytopes of width larger than their dimension. Before that, we show a heuristic method to certify the width of a convex body. This method indirectly takes advantage of the fact that in our examples the width is attained with respect to several different functionals.

By a rational path Γ in \mathbb{R}^d , with respect to a certain lattice Λ , we mean a concatenation of segments in rational directions. That is, Γ is given as a sequence p_0, p_1, \ldots, p_t of points in \mathbb{R}^d such that for every i the vector $p_{i+1} - p_i$ is parallel to a lattice vector. This allows us to define the *lattice length* of each segment $[p_i, p_{i+1}]$ as the scalar $\lambda > 0$ such that $\frac{1}{\lambda}(p_{i+1} - p_i)$ is *primitive*, meaning that it is the shortest

integer vector in its direction. The *lattice length* of the rational path Γ is the sum of the lattice lengths of the individual segments; we denote this by length_Λ(Γ).

We say that a functional f is *strictly increasing* along Γ if

$$
f(p_0) < f(p_1) < \cdots < f(p_t).
$$

The open polar cone of Γ, denoted cone (Γ)[°], is the set of functionals $f \in (\mathbb{R}^d)^*$ that are strictly increasing along Γ ^{[2](#page-5-1)}

Lemma 3.1. Let $P \subset \mathbb{R}^d$ be a convex body. Let Γ be a rational path of lattice length w for a certain lattice Λ , with the first and last points of Γ in P. Then any lattice functional in the open polar cone of Γ gives width at least w to P.

Proof. If $f \in \text{cone}(\Gamma)^\circ$ then

$$
\text{width}(P, f) \ge \text{length}_{\Lambda}(\Gamma) = w,
$$

since f takes an integer positive value in the primitive vector parallel to each segment of Γ. \Box

Remark 3.2. As a consequence of the lemma, if $\Gamma_1, \ldots, \Gamma_k$ is a collection of rational paths with end-points in P , all of length at least w , and with the property that

$$
\bigcup_{i=1}^k \text{cone}(\Gamma_i)^\circ = \mathbb{R}^d \setminus \{0\},\
$$

then the lattice width of P is at least w .

Example 3.3. The necessity for using the *open* polar cone cone $(\Gamma)^\circ$ and not the closed one in Lemma [3.1](#page-5-2) can be illustrated by considering P to be the square $[0, 1]^2$. The two boundary paths between opposite vertices in P have lattice length two and by Lemma 3.1 this guarantees that the width of P with respect to any functional in $\Lambda^* \setminus (\langle e_1^* \rangle \cup \langle e_2^* \rangle)$ is at least two. But, of course, the width of P with respect to the functionals e_1^* and e_2^* is 1, and these two functionals are weakly increasing along the boundary paths.

4. A hollow lattice 14-polytope of width 15

Let A, B and C be the vertices of an equilateral triangle Δ in the plane; without loss of generality, $A = (0,0), B = (1,0), C = \left(\frac{1}{2}, \frac{\sqrt{3}}{2}\right)$. Let Λ be the lattice they generate:

$$
\Lambda := \left\{ (a, b\sqrt{3}) \in \mathbb{R}^2 : 2a, 2b, a+b \in \mathbb{Z} \right\}.
$$

We consider the family $\{T(x, y)\}\$ of equilateral triangles circumscribed around Δ , where (x, y) denotes, by convention, the vertex lying between A and C. A point (x, y) defines such a triangle if and only if it lies outside Δ and along the circle

$$
S_1 := \left\{ (x,y) : x^2 + \left(y - \frac{\sqrt{3}}{3}\right)^2 = \frac{1}{3} \right\}.
$$

It is easy to see that every triangle in the family is hollow. For example, $T(-\frac{1}{2}, \frac{\sqrt{3}}{2})$ is a hollow lattice triangle of width two, unimodularly equivalent to

²The notation cone (Γ)[°] comes from the fact that this cone is the (open) polar, in the standard sense, of the cone generated by the vectors $p_{i+1} - p_i$, $i = 1, ..., t$.

FIGURE 1. Left: the hollow equilateral triangle $T(x, y)$ circumscribed to the triangle Δ , depending on the position of (x, y) along the circle S_1 (Hurkens position, maximizing width, is shown in the picture). Right: the refinement of the lattice by a factor of seven creates a lattice point in the circle and close to Hurkens position.

the second dilation of Δ . The triangle $T\left(-\frac{\sqrt{3}}{3}, \frac{\sqrt{3}}{3}\right)$, pictured in Figure [1](#page-6-0) (left) maximizes the width of the family and was shown by Hurkens [\[10\]](#page-14-7) to maximize width among all hollow convex 2-bodies (see also Averkov and Wagner [\[4\]](#page-14-13)).

We now consider the seventh refinement $\Lambda' := \frac{1}{7}\Lambda$ of Λ . The circle S_1 contains, apart from the points A, B, C , additional points of Λ' . In particular, if we fix $T := T(D)$ for the point $D = \left(-\frac{4}{7}, \frac{2\sqrt{3}}{7}\right) \in \Lambda' \cap S_1$ we get a triangle with vertices in $Λ'$ and of width close to the maximum, since D is close to Hurken's point $\left(-\frac{\sqrt{3}}{3}, \frac{\sqrt{3}}{3}\right)$ (See Figure [1,](#page-6-0) again). Specifically:

$$
T := T(D) = \text{conv}\left(\left(-\frac{4}{7}, \frac{2\sqrt{3}}{7} \right), \left(\frac{17}{14}, \frac{9\sqrt{3}}{14} \right), \left(\frac{6}{7}, -\frac{3\sqrt{3}}{7} \right) \right).
$$

Lemma 4.1. The triangle T defined above is hollow and of width $2 + 1/7 = 2.1419$ with respect to Λ . It is also rational, with its seventh dilation being a lattice triangle.

Proof. It is clear by our construction that T is hollow with respect to Λ , and since it has its vertices in Λ' , its seventh dilation is a lattice triangle of Λ .

We now claim that the width of T in Λ' is 15. It is easy to check that it has width 15 with respect to the three functionals f_0 , f_1 and f_2 that define edges of Δ . We call E and F the vertices of T in clockwise order from D . To show that the width of T is at least 15, we apply Lemma [3.1](#page-5-2) to each of the three paths DHE (drawn in red in Figure [1\)](#page-6-0), EIF and FGD . These paths have lattice length equal to 15. It is easy to see that the polar cones of the paths are cone (f_0, f_1) , cone (f_0, f_2) and cone (f_1, f_2) , so by Lemma [3.1,](#page-5-2) functionals in the interior of any of these cones give width at least 15 to T. The only (primitive) functionals not in the open cones are precisely f_0, f_1 and f_2 which, as said above, yield width 15.

We can now prove the first half of Theorem 1.3 :

Theorem 4.2. $T^{\oplus 7}$ is a 14-dimensional hollow lattice polytope of width 15. It has 21 vertices and $2^7 + 7$ facets $(2^7 \text{ simplices and seven combinatorially of the form})$ $segment \oplus triangle^{\oplus 6}$).

Proof. The first claim follows from Proposition [1.4](#page-2-1) and Lemma [4.1.](#page-6-1) $T^{\oplus 7}$ has 21 vertices by the description of vertices of direct sums in Lemma [2.1.](#page-3-3) The same lemma implies the following description of the facets:

- (1) Facets of $T^{\oplus 7}$ that do not contain the origin are the joins of edges of T that do not contain the origin. Since there are two such edges to chose from in T and joins of simplices are simplices, we obtain the $2⁷$ stated simplices.
- (2) Facets of $T^{\oplus 7}$ that contain the origin are of the form

$$
T \oplus \cdots \oplus T \oplus F \oplus T \oplus \cdots \oplus T
$$

where F is the edge of T that contains the origin. Since F can be placed anywhere in the sum, we have seven such facets. \Box

5. A hollow 3-simplex of width 3.4142

Consider the (dilated) face-centered cubic lattice

$$
\Lambda := \{ (a, b, c) : a, b, c \in 2\mathbb{Z}, a + b + c \in 4\mathbb{Z} \},
$$

with dual

$$
\Lambda^* = \left\{ (a, b, c) \in \frac{1}{4} \mathbb{Z} : a + b, a + c, b + c \in \frac{1}{2} \mathbb{Z}^3 \right\}.
$$

Here and in what follows we use the standard coordinates in $(\mathbb{R}^3)^*$, so that (a, b, c) denotes the functional $(x, y, z) \mapsto ax + by + cz$.

For the sake of symmetry, all constructions in this section are with respect to the following *affine* lattice, which is a translation of Λ :

$$
\Lambda_{-1} := \Lambda - (1,1,1) = \left\{ (a,b,c) : a,b,c \in 1 + 2\mathbb{Z}, a+b+c \in 1 + 4\mathbb{Z} \right\}.
$$

By lattice width with respect to Λ_{-1} we mean the lattice width with respect to Λ .

Consider the following lattice tetrahedron (see Figure [2\)](#page-8-1) of width three in Λ_{-1} :

$$
\Delta_0 := \operatorname{conv}\left\{ (3,1,5), (-1,3,-5), (-3,-1,5), (1,-3,-5) \right\}.
$$

 Δ_0 is (modulo unimodular transformation) the hollow 3-simplex of normalized volume 25 and width 3 that appears in $[2,$ Figure 2 and $[3,$ Figure 1(h). We want to modify Δ_0 to a non-hollow simplex of larger width, in the spirit of the previous section. We chose this Δ_0 because it achieves its lattice width only with respect to two lattice functionals, namely $x/4$ and $y/4$. This gives a certain freedom to scale down the z coordinate and enlarge the other two, thus increasing the minimum width. We can simultaneously rotate the whole tetrahedron around the z axis.

To formalize this, we consider the family of tetrahedra that share the following properties with Δ_0 : they are circumscribed around the unimodular simplex conv $\{(-1, 1, 1), (-1, -1, -1), (1, -1, 1), (1, 1, -1)\}$, and they are invariant under the order four isometry $(x, y, z) \mapsto (-y, x, -z)$. Put differently, for each $(x, y, z) \in$ \mathbb{R}^3 we define $\Delta(x, y, z)$ to be the tetrahedron with vertices

$$
A = (x, y, z),
$$
 $B = (-y, x, -z),$ $C = (-x, -y, z),$ $D = (y, -x, -z).$

We constrain (x, y, z) to satisfy that $(-1, 1, 1), (-1, -1, -1), (1, -1, 1)$ and $(1, 1, -1)$ lie, respectively, in the planes containing ABC, BCD, ACD and ABD. By symmetry, these four conditions are equivalent to one another and an easy computation

FIGURE 2. Projection along the z-axis of the hollow lattice 3simplex Δ_0 of width three in the lattice $\Lambda_{-1} := \{(a, b, c) : a, b, c \in$ $1+2\mathbb{Z}, a+b+c \in 1+4\mathbb{Z}$. Dots represent (projections of) vertical lattice lines. For those that intersect Δ_0 , next to the dot we show the interval of values of z in the intersection. For example, the interval $[-1, 3]$ next to the dot with coordinates $(1, 1)$ indicates that the points $(1, 1, -1)$ and $(1, 1, 3)$ are in the boundary of Δ_0 . At the four vertices of Δ_0 the interval degenerates to a point.

shows that they translate to the equality

(1)
$$
z = \frac{x^2 + y^2}{x^2 + y^2 - 2x - 2y}.
$$

In the rest of this section we show the following, which implies Theorem [1.1:](#page-1-1)

Theorem 5.1. Let $(x, y, z) \in \mathbb{R}^3$ be a point satisfying the constraint of Equa-**Theorem 3.1.** Let $(x, y, z) \in \mathbb{R}$ be a point satisfying the constraint of Equation [\(1\)](#page-8-2). Then, the width of $\Delta(x, y, z)$ with respect to Λ is at most $2 + \sqrt{2}$, with equality if and only if

$$
(x, y, z) \in \left\{ \left(2 + \sqrt{2}, \sqrt{2}, 2 + \sqrt{2}\right), \left(\sqrt{2}, 2 + \sqrt{2}, 2 + \sqrt{2}\right)\right\}.
$$

Proof of the upper bound in Theorem [5.1.](#page-8-0) Let us consider the functionals $(\frac{1}{2}, 0, 0)$, $(0, \frac{1}{2}, 0)$, and $(0, 0, \frac{1}{2})$, which are in Λ^* . The width of $\Delta(x, y, z)$ with respect to first two equals max $\{|x|, |y|\}$, and with respect to the third equals |z|. We are going to two equals $\max\{|x|, |y|\}$, and with respect to the third equals $|z|$. We show that whenever $\max\{|x|, |y|\} \geq 2 + \sqrt{2}$ we have $|z| \leq 2 + \sqrt{2}$. Let

$$
f(x,y) := \frac{x^2 + y^2}{x^2 + y^2 - 2x - 2y}
$$

be the function giving z in terms of x and y. The assumption max $\{|x|, |y|\} \geq 2+\sqrt{2}$ implies that the denominator of f is positive, since it is only negative (or zero) inside miques that the denominator of f is positive, since it is only negative (or zero) inside
(or on) the circle with center (1, 1) and radius $\sqrt{2}$. The numerator is also obviously positive, and thus z is positive. The equation

$$
f(x,y) = \frac{x^2 + y^2}{x^2 + y^2 - 2x - 2y} = 2 + \sqrt{2}
$$

defines again a circle, with center $(\sqrt{2},$ √ 2) and radius 2. Outside the circle z is dennes again a circle, with center $(\sqrt{2}, \sqrt{2})$ and radius 2. Outside the smaller than $2 + \sqrt{2}$ and inside the circle at least one of |x| and |y| is.

 \Box

Thus, for the rest of the section we fix $\Delta = \Delta(2 + \sqrt{2})$, $\sqrt{2}$, $2 + \sqrt{2}$), which has the following vertices and is depicted in Figure [3:](#page-9-0)

$$
A = \begin{pmatrix} 2 + \sqrt{2}, & \sqrt{2}, & 2 + \sqrt{2} \end{pmatrix}, \qquad B = \begin{pmatrix} -\sqrt{2}, & 2 + \sqrt{2}, & -2 - \sqrt{2} \end{pmatrix},
$$

\n
$$
C = \begin{pmatrix} -2 - \sqrt{2}, & -\sqrt{2}, & 2 + \sqrt{2} \end{pmatrix}, \qquad D = \begin{pmatrix} \sqrt{2}, & -2 - \sqrt{2}, & -2 - \sqrt{2} \end{pmatrix}.
$$

FIGURE 3. The hollow 3-simplex Δ of width $2 + \sqrt{2}$, drawn with FIGURE 3. The notions as in Figure [2.](#page-8-1) We abbreviate $\alpha = 1 + \sqrt{2}$.

Observe that the width of Δ with respect to the following 14 lattice functionals Observe that
equals $2 + \sqrt{2}$:

(2)
$$
\pm \frac{1}{2}(1,0,0), \pm \frac{1}{2}(0,1,0), \pm \frac{1}{2}(0,0,1), \frac{1}{4}(\pm 1, \pm 1, \pm 1).
$$

We now prove that this is the width of Δ .

Proof of the equality in Theorem [5.1.](#page-8-0) In Figure [3](#page-9-0) we have written, next to each vertical lattice line $\ell = (x_0, y_0) \times \mathbb{R}$ intersected by Δ , the interval $\{z \in \mathbb{R} : (x_0, y_0, z) \in$ ∆}. Hollowness follows from this information, since the intervals do not contain points of Λ in their interior. To check correctness of these computations observe that the facet-defining inequality for triangle ABC is

$$
z \le \frac{x-y}{\sqrt{2}} - y + 2 + \sqrt{2}.
$$

Plugging in the coordinates $(x_0, y_0) \in \{(-3, -1), (-1, 1), (-1, 3), (1, 1)\}$ of the four vertical lines meeting the triangle ABC we get that the highest points of Δ on each are indeed

$$
(-3,-1,3), (-1,1,1), (-1,3,-1-\sqrt{2}), (1,1,1+\sqrt{2}).
$$

The rest of upper and lower bounds for the intervals in Figure [3](#page-9-0) follow by symmetry.

To show that the width is at least $2 + \sqrt{2}$ we apply Lemma [3.1](#page-5-2) to various paths. For example, the expression

$$
D = C + \left(\frac{1}{2} + \frac{\sqrt{2}}{2}\right)(4,0,0) + \frac{1}{2}(0,-4,0) + \left(1 + \frac{\sqrt{2}}{2}\right)(0,0,-4).
$$

gives a rational path from vertex C to vertex D with directions $(4, 0, 0)$, $(0, -4, 0)$ and $(0, 0, -4)$ and of length

$$
\left(\frac{1}{2} + \frac{\sqrt{2}}{2}\right) + \frac{1}{2} + \left(1 + \frac{\sqrt{2}}{2}\right) = 2 + \sqrt{2}.
$$

The open polar cone of this rational path is the octant $\{(a, b, c) \in (\mathbb{R}^3)^* : a > 0, b <$ $0, c < 0$, so all lattice functionals in the interior of the octant give width at least $2 + \sqrt{2}$ to Δ . The same path in reverse implies the same for the opposite octant, and the symmetry of order 4 in Δ implies it for the eight open octants.

We now define a second family of paths whose open polar cones are the connected components of $(\mathbb{R}^3)^* \setminus \{(a, b, c) : a = \pm b\}.$ (Observe that these are non-pointed cones). The first one goes from B to D based on the equality

$$
D = B + (-2, -2, 0) + (1 + \sqrt{2})(2, -2, 0).
$$

Its length is $1 + (1 + \sqrt{2}) = 2 + \sqrt{2}$ and its open polar cone is

$$
\{(a,b,c) : a+b < 0, a-b > 0\}.
$$

Again, symmetry of Δ gives paths for the other three connected components of $(\mathbb{R}^3)^* \setminus \{ (a, b, c) : a = \pm b \}.$

To $\setminus \{a, b, c\}$: $a = \pm b$.
Together, these two sets of paths show width $\geq 2 + \sqrt{2}$ for all lattice functionals except for the integer multiples of $\frac{1}{2}(0,0,1)$, $\frac{1}{2}(1,1,0)$ and $\frac{1}{2}(1,-1,0)$. These three except for the integer multiples of $\frac{1}{2}(0, 0, 1)$, $\frac{1}{2}(1, 1, 0)$ and $\frac{1}{2}(1, -1, 0)$. These three
give widths $2 + \sqrt{2}$, $2 + 2\sqrt{2}$ and $2 + 2\sqrt{2}$ to Δ , respectively.

Remark 5.2. The family of tetrahedra $\Delta(x, y, z)$ also contains

$$
\Delta(3,3,3) = \text{conv}\{(3,3,3), (3,-3,-3), (-3,3,-3), (-3,3,-3)\},\
$$

which is the third dilation of a unimodular simplex. In this sense, $\Delta(x, y, z)$ is a common generalization of two of the three existing lattice tetrahedra of maximal width [\[2\]](#page-14-9). This is further motivation for Conjecture [1.2.](#page-1-3)

6. A hollow lattice 404-simplex of width 408

We now want to construct a lattice *simplex* of width larger than its dimension. To do this via Theorem [2.2,](#page-3-0) we need a rational hollow simplex with the origin as a vertex and of width larger than its dimension, which can be found in dimension four. We do not know whether one exists in dimension three.

Lemma 6.1. There is a rational hollow 4-simplex S of width $4 + 4/101$ and with a lattice vertex whose 101-th dilation is a lattice simplex.

Proof. It is known that the following lattice 4-simplex is *empty*, that is, it has no latticepoints other than its vertices, and it has width four $([8, 11])$ $([8, 11])$ $([8, 11])$ $([8, 11])$ $([8, 11])$:

 $S_0 := \text{conv } \{(0, 0, 0, 0), (1, 0, 0, 0), (0, 1, 0, 0), (0, 0, 1, 0), (6, 14, 17, 101)\}.$

Observe that the facet of S_0 opposite the origin lies in the hyperplane $101x_1 +$ $101x_2 + 101x_3 - 36x_4 = 101$. Since 101 is coprime with 36, dilating S_0 by a factor of $102/101$ gives a hollow simplex S: apart of the five vertices of S_0 (which lie in the boundary of S) all other lattice points must be in the facet-defining hyperplane $101x_1 + 101x_2 + 101x_3 - 36x_4 = 102.$

Applying Proposition [1.4](#page-2-1) to the hollow simplex S, we obtain that $S^{\oplus 101}$ is a 404-dimensional lattice simplex of width 408. This proves the second half of Theorem [1.3.](#page-1-2)

Remark 6.2. Any dilation of S_0 by a factor strictly greater than $102/101$ is not hollow anymore, since the point

$$
(1,2,3,14) = \frac{17}{101}(1,0,0,0) + \frac{6}{101}(0,1,0,0) + \frac{65}{101}(0,0,1,0) + \frac{14}{101}(6,14,17,101)
$$

lies in the relative interior of the facet of S opposite the origin.

7. General lower bounds

In this section, we apply Theorem [2.2](#page-3-0) to the explicit examples from Sections [4–](#page-5-0) [6](#page-10-0) to obtain lower bounds for $w_c(d)$, $w_p(d)$ and $w_s(d)$ in general dimension d. In particular, we prove Theorem [1.5](#page-2-0) and Proposition [1.7.](#page-2-2)

Corollary 7.1. For $w_* = w_c$, w_p or w_s we have that

$$
w_*(md) \ge m w_*(d), \quad \forall m \in \mathbb{N}.
$$

For w_c we have the more general result

(3)
$$
w_c(d_1 + \cdots + d_m) \ge w_c(d_1) + \cdots + w_c(d_m), \quad \forall d_1, \ldots, d_m \in \mathbb{N}.
$$

Proof. For the first inequality, let K be a hollow convex d-body (resp., a lattice d polytope, a lattice d-simplex) achieving $w_c(d)$ (resp. $w_p(d)$, $w_s(d)$ and, in the case of a lattice polytope, assume without loss of generality that the origin is a vertex of K). Then, apply Theorem [2.2](#page-3-0) with $C_i = K$ and $k_i = m$ for all i. This gives a dm-dimensional hollow convex body (resp., a lattice polytope, a lattice simplex) of width $m w_*(d)$.

For the case of w_c we have more freedom, since we do not need the k_i s to be integers. Thus, if for each $i = 1, \ldots, m$ we let the C_i s in Theorem [2.2](#page-3-0) be hollow convex bodies of width $w_c(d_i)$ and we take $k_i = (\sum_j w_c(d_j)) / w_c(d_i)$ for each i, we obtain a hollow convex body C of width $\sum_j w_c(d_j)$ and dimension $\sum_j d_j$.

Inequality [\(3\)](#page-11-1) implies that w_c is strictly increasing. For w_p and w_s we can only prove weak monotonicity:

Corollary 7.2.

 $w_p(d+1) \geq w_p(d), \quad w_s(d+1) \geq w_s(d).$

Proof. Let $C_1 = [0, 1]$ and let C_2 be a lattice polytope (resp. a hollow simplex) of dimension d and with width(C) = $w_*(d)$. Apply Theorem [2.2](#page-3-0) with $k_1 > w_*(d)$ and $k_2 = 1.$

Question 7.3. Are w_p , w_s or w_e strictly increasing? Since these w_* take only integer values, strict monotonicity is equivalent to the inequality

$$
w_*(d+1) \ge w_*(d) + 1.
$$

(For w_e even non-strict monotonicity is unclear, due to its more arithmetic nature).

We can now prove Theorem [1.5](#page-2-0) and Proposition [1.7:](#page-2-2)

Proof of Theorem [1.5.](#page-2-0) By Corollaries [7.1](#page-11-2) and [7.2,](#page-11-3) the three sequences $w_c(d)$, $w_n(d)$ and $w_s(d)$ satisfy the conditions of the following elementary statement:

If a sequence $(a_d)_{d\in\mathbb{N}}$ satisfies $a_{d+1} \ge a_d$ and $a_{md} \ge m a_d \ \forall d, m \in \mathbb{N}$, then

$$
\lim_{d \to \infty} \frac{a_d}{d} = \sup_{d \in \mathbb{N}} \frac{a_d}{d}.
$$

Proof of Proposition [1.7.](#page-2-2) The inequalities

$$
w_c(d+1) \ge w_c(d) + 1,
$$

\n
$$
w_c(d+2) \ge w_c(d) + 1 + \frac{2}{\sqrt{3}},
$$

\n
$$
w_c(d+3) \ge w_c(d) + 2 + \sqrt{2},
$$

follow from applying Equation [\(3\)](#page-11-1) of Corollary [7.1](#page-11-2) with $w_c(1) = 1$ $w_c(2) = 1 +$ $\frac{2}{ }$ now from applying Equation (3) or C
 $\frac{1}{3}$ [\[10\]](#page-14-7) and $w_c(3) \geq 2 + \sqrt{2}$ (Section [5\)](#page-7-0).

Any integer $d \geq 2$ can be written as $d = 2a + 3b$ for some nonnegative integers a, b. Then for all $d \geq 2$, the inequalities above yield

$$
w_c(d) \ge w_c(2)a + w_c(3)b \ge \left(1 + \frac{2}{\sqrt{3}}\right)a + \left(2 + \sqrt{2}\right)b
$$

$$
\ge \left(\frac{1}{2} + \frac{1}{\sqrt{3}}\right)(2a + 3b) = \left(\frac{1}{2} + \frac{1}{\sqrt{3}}\right)d.
$$

8. Lower bound for empty simplices

To prove the asymptotic lower bound of Theorem [1.8,](#page-3-2) we use the following lemma:

Lemma 8.1. Let $P = \text{conv}(0, v_1, \dots, v_d) \subset \mathbb{R}^d$ be an empty d-simplex of width w and let $m \geq 2$ be an integer. For each $i \in [d]$ and $j \in [m]$, let

$$
v_i^{(j)} := 0 \oplus \cdots \oplus v_i \oplus \cdots \oplus 0 \in \mathbb{R}^{md}
$$

with v_i in the j-th summand, and define

 (1)

$$
w_i^{(j)} := (m-2)v_i^{(j)} + v_{i+1}^{(j+1)} \in \mathbb{R}^{md},
$$

with i taken modulo d and j modulo m. Let

$$
P_m := \text{conv}\left(\{0\} \cup \left\{w_i^{(j)} : (i,j) \in [d] \times [m]\right\}\right).
$$

Then: (1) width $(P_m) \ge (m-3)w$; and (2) P_m is empty.

Proof. Observe that P_m is contained in $P^{\oplus m} = \bigoplus_{j=1}^m mP$ and tries to approximate it: the vertices of $P^{\oplus m}$ are 0 and $\{mv_i^{(j)} : (i,j) \in [d] \times [m]\}$, and the vertices $w_i^{(j)}$ of P_m are close to them.

To prove (1), let $f = f_1 \oplus \cdots \oplus f_m$ be an integer functional. Assume without loss of generality that

$$
\max_{j} \{\text{width}(P, f_j)\} = \text{width}(P, f_1).
$$

Let us denote $v_0 = 0$ and let $i^+, i^- \in \{0, \ldots, d\}$ be indices such that $f_1(v_{i^+})$ and $f_1(v_i-)$ are the maximum and minimum values of f_1 on P, respectively. Then,

$$
|f(w_{i^{+}}^{(1)} - w_{i^{-}}^{(1)})| \ge |f_1((m-2)v_{i^{+}} - (m-2)v_{i^{-}})| - |f_2(v_{i^{+}+1} - v_{i^{-}+1})|
$$

\n
$$
\ge (m-2)|f_1(v_{i^{+}} - v_{i^{-}})| - |f_1(v_{i^{+}} - v_{i^{-}})|
$$

\n
$$
= (m-3)|f_1(v_{i^{+}} - v_{i^{-}})| \ge (m-3)w.
$$

For part (2), to search for a contradiction assume P_m is not empty. Let $z \in \mathbb{R}$ $P_m \cap \mathbb{Z}^{dm}$ be an integer point different from 0 and from the $w_i^{(j)}$ s. We can then write z as a convex combination of the vertices of P_m . That is:

(4)
$$
z = \sum_{j=1}^{m} \sum_{i=1}^{d} \lambda_i^{(j)} w_i^{(j)} = \sum_{j=1}^{m} \sum_{i=1}^{d} ((m-2)\lambda_i^{(j)} + \lambda_{i-1}^{(j-1)}) v_i^{(j)},
$$

with $\lambda_i^{(j)} \geq 0$ and $\sum_{i,j} \lambda_i^{(j)} \leq 1$.

But since $P_m \subset P^{\oplus m}$, we can also write

(5)
$$
z = \mu_1 z_1 \oplus \cdots \oplus \mu_m z_m,
$$

with each $z_j \in P$, $\mu_j z_j \in \mathbb{Z}^d$, $\mu_j \geq 0$ and $\sum_j \mu_j \leq m$. Comparing Equations [\(4\)](#page-13-0) and (5) we obtain

(6)
$$
\mu_j z_j = \sum_{i=1}^d ((m-2)\lambda_i^{(j)} + \lambda_{i-1}^{(j-1)}) v_i.
$$

Claim: $\sum_i \lambda_i^{(j)} \neq 0$ *for every j.* Indeed, if there is a j where this sum is zero, assume without loss of generality that $\sum_i \lambda_i^{(j-1)} \neq 0$. Then Equation [\(6\)](#page-13-2) gives

$$
\mu_j z_j = \sum_{i=1}^d \lambda_{i-1}^{(j-1)} v_i,
$$

which is a nonzero point in P. Since P is empty and $\mu_j z_j \in \mathbb{Z}^d$, we conclude that one of the $\lambda_i^{(j-1)}$ s equals 1, so that $z = w_i^{(j-1)}$, a contradiction because z was assumed not to be a vertex of P_m .

From the claim and Equation [\(6\)](#page-13-2) it follows that $\mu_i z_i \neq 0$ for all j. In order for $\mu_j z_j$ to be a lattice point we need $\mu_j \geq 1$ (because $0 < \mu_j < 1$ implies $\mu_j z_j$ to be a lattice point in P but not a vertex of P , which is not possible). Since on the other hand $\sum_j \mu_j \leq m$, we conclude that $\mu_j = 1$ for every j. This implies that every z_j is a non-zero lattice point of P; that is, for each j there is an i_j such that $z_j = v_{i_j}$. Equation (6) now becomes

$$
v_{i_j} = \sum_{i=1}^{d} ((m-2)\lambda_i^{(j)} + \lambda_{i-1}^{(j-1)})v_i.
$$

Since the v_i s are independent, we have

$$
1 = (m-2)\lambda_{i_j}^{(j)} + \lambda_{i_j-1}^{(j-1)}, \qquad \forall j.
$$

Summing over j we get the contradiction

$$
m = \sum_{j} (m-2)\lambda_{i_j}^{(j)} + \sum_{j} \lambda_{i_j-1}^{(j-1)} \le (m-2) + 1 = m - 1.
$$

Remark 8.2. Lemma [8.1](#page-12-1) and its proof generalize Sebő's construction of empty msimplices of width $m-2$ [\[15\]](#page-14-4). Indeed, letting $P = [0, 1]$ our lemma gives an empty m-simplex of width (at least) $m-3$. Sebő's $m-2$ is obtained with an additional argument that works for $[0, 1]$ but not (as far as we can see) for an arbitrary P.

Proof of Theorem [1.8.](#page-3-2) Let P be an empty d-simplex of maximum width; that is, with width(P) = $w_e(d)$. Applying Lemma [8.1](#page-12-1) to P we obtain a sequence $(P_m)_{m \in \mathbb{N}}$ of empty md-simplices of width $(m-3)w_e(d)$, which implies $w_e(dm) \geq (m-3)w_e(d)$.

From this fact, combined with $w_e(1) = 1$, we obtain

$$
\limsup_{d \to \infty} \frac{w_e(d)}{d} = \sup_{d \in \mathbb{N}} \frac{w_e(d)}{d} \ge 1.
$$

REFERENCES

- 1. G. Averkov, A. Basu. Lifting properties of maximal lattice-free polyhedra, Math. Program., Ser. B 154 (2015) 81–111. <http://dx.doi.org/10.1007/s10107-015-0865-6>
- 2. G. Averkov, J. Krümpelmann and S. Weltge, Notions of maximality for integral lattice-free polyhedra: the case of dimension three. Math. Oper. Res., 42:4 (2017), 1035–1062. [http:](http://dx.doi.org/10.1287/moor.2016.0836) [//dx.doi.org/10.1287/moor.2016.0836](http://dx.doi.org/10.1287/moor.2016.0836)
- 3. G. Averkov, C. Wagner and R. Weismantel. Maximal lattice-free polyhedra: finiteness and an explicit description in dimension three, Math. Oper. Res. 36 (2011), no. 4, 721–742.
- 4. G. Averkov and C. Wagner. Inequalities for the lattice width of lattice-free convex sets in the plane, Beitr. Algebra Geom. (2012) 53:1–23. [http://dx.doi.org/10.1007/](http://dx.doi.org/10.1007/s13366-011-0028-8) [s13366-011-0028-8](http://dx.doi.org/10.1007/s13366-011-0028-8)
- 5. W. Banaszczyk. Inequalities for convex bodies and polar reciprocal lattices in \mathbb{R}^n II: Application of K-convexity. Discrete Comput. Geom. 16. (1996) 305–311.
- 6. W. Banaszczyk, A. E. Litvak, A. Pajor, and S. J. Szarek, The flatness theorem for nonsymmetric convex bodies via the local theory of Banach spaces., Math. Oper. Res. 24 (1999), no. 3, 728–750.
- 7. S. Dash, N. B. Dobbs, O. Günlük, T. J. Nowicki, G. M. Świrszcz, Lattice-free sets, multibranch split disjunctions, and mixed-integer programming, Math. Program. 145:1–2 (2014) 483–508. https://doi.org/10.1007/s10107-013-0654-z
- 8. C. Haase and G. M. Ziegler, On the maximal width of empty lattice simplices. Eur. J. Combin. 21 (2000), 111–119.
- 9. M. Henk, J. Richter-Gebert, G. M. Ziegler, Basic Properties of Convex Polytopes. In Handbook of Discrete and Computational Geometry, Third Edition, edited by Jacob E. Goodman, Joseph O'Rourke, and Csaba D. Tóth, CRC Press, November 2017, pp 383-413.
- 10. C. A. J. Hurkens. Blowing up convex sets in the plane, Linear Algebra Appl. 134 (1990), 121–128.
- 11. O. Iglesias Valiño and F. Santos, Classification of empty lattice 4-simplices of width larger than two, Trans. Amer. Math. Soc., to appear. arXiv:1704.07299
- 12. R. Kannan and L. Lovász, Covering minima and lattice-point-free convex bodies, Ann. of Math. (2) 128 (1988), no. 3, 577-602.
- 13. J. Milnor, D. Husemoller. Symmetric Bilinear Forms. Springer-Verlag, 1973.
- 14. M. Rudelson. Distances between nonsymmetric convex bodies and the MM∗-estimate. Positivity 4:2 (2000), 161–178.
- 15. A. Sebő. An introduction to empty lattice simplices, Integer programming and combinatorial optimization (Graz, 1999), Lecture Notes in Comput. Sci., vol. 1610, Springer, Berlin, 1999, pp. 400–414.
- 16. G. K. White. Lattice tetrahedra. Canadian J. Math. 16 (1964), 389–396.

(G. Codenotti) INSTITUT FÜR MATHEMATIK, FREIE UNIVERSITÄT BERLIN, GERMANY E-mail address: giulia.codenotti@fu-berlin.de

(F. Santos) Dept. of Mathematics, Statistics and Comp. Sci., Univ. of Cantabria, Spain E-mail address: francisco.santos@unican.es