

# A class of narrow-sense BCH codes over $\mathbb{F}_q$ of length $\frac{q^m-1}{2}$

Xin Ling · Sihem Mesnager · Yanfeng Qi ·  
Chunming Tang

Received: date / Accepted: date

**Abstract** BCH codes with efficient encoding and decoding algorithms have many applications in communications, cryptography and combinatorics design. This paper studies a class of linear codes of length  $\frac{q^m-1}{2}$  over  $\mathbb{F}_q$  with special trace representation, where  $q$  is an odd prime power. With the help of the inner distributions of some subsets of association schemes from bilinear forms associated with quadratic forms, we determine the weight enumerators of these codes. From determining some cyclotomic coset leaders  $\delta_i$  of cyclotomic cosets modulo  $\frac{q^m-1}{2}$ , we prove that narrow-sense BCH codes of length  $\frac{q^m-1}{2}$  with designed distance  $\delta_i = \frac{q^m-q^{m-1}}{2} - 1 - \frac{q^{\lfloor \frac{m-3}{2} \rfloor + i} - 1}{2}$  have the corresponding trace representation, and have the minimal distance  $d = \delta_i$  and the Bose distance  $d_B = \delta_i$ , where  $1 \leq i \leq \lfloor \frac{m+3}{4} \rfloor$ .

**Keywords** Linear codes · BCH codes · association schemes · the weight distribution · quadratic forms

**Mathematics Subject Classification (2010)** 94B05 · 94B15 · 05E30 · 15A63

## 1 Introduction

As an important class of cyclic codes, BCH codes with efficient encoding and decoding algorithms have many applications in communications, cryptography and combinatorics de-

---

X. Ling

School of Mathematics and Information, China West Normal University, Nanchong 637002, China. E-mail: xinlingcwnu@163.com.

S. Mesnager

the Department of Mathematics, University of Paris VIII, 93526 Saint-Denis, France, with LAGA UMR 7539, CNRS, Sorbonne Paris Cité, University of Paris XIII, 93430 Paris, France, and also with Telecom ParisTech, 75013 Paris, France. E-mail: smesnager@univ-paris8.fr.

Y. Qi

School of Science, Hangzhou Dianzi University, Hangzhou, Zhejiang, 310018, China. E-mail: qiyanfeng07@163.com.

C. Tang

School of Mathematics and Information, China West Normal University, Nanchong 637002, China, and Department of Mathematics, The Hong Kong University of Science and Technology, Clear Water Bay, Kowloon, Hong Kong, China. E-mail: tangchunmingmath@163.com.

sign. They were independently discovered by Hocquenghem [17] and Bose, Ray-Chaudhuri [3], and were generalized from binary fields to finite fields by Gorenstein and Zierler [15]. The determination of parameters of BCH codes is an interesting and difficult problem. There are many results on BCH codes [1, 2, 4, 5, 6, 7, 8, 9, 10, 12, 13, 14, 16, 19, 22, 23, 24, 26, 28, 30, 31, 33, 34, 38, 39, 40, 41, 42].

Let  $m$  be a positive integer and  $\mathbb{F}_{q^m}$  be a finite field. Let  $\beta$  be an element in  $\mathbb{F}_{q^m}$  with order  $n$ , where  $n \mid (q^m - 1)$ . A BCH code  $C$  over  $\mathbb{F}_q$  with designed distance  $\delta$  is a cyclic code with a generator polynomial  $g(x)$ , where  $g(x)$  is determined by its  $\delta - 1$  consecutive roots  $\beta^l, \beta^{l+1}, \dots, \beta^{l+\delta-1}$ . The minimal distance of the BCH code  $C$  is greater than  $\delta$ . When  $n = q^m - 1$ , the code  $C$  is called primitive. When  $l = 1$ , this code  $C$  is called narrow-sense. Two narrow-sense BCH codes with different designed distances may be the same. The Bose distance  $d_B$  of a BCH code is the largest designed distance. The Bose distance satisfies that  $d_B \leq d \leq d_B + 4$  [5], where  $d$  is the minimum distance. The determination of the minimum distance  $d$  and the Bose distance of a narrow-sense BCH code attracts much interest [1, 4, 5, 9, 20, 21, 22, 32, 33, 40]. Some results on  $[n, k, d]$  narrow-sense BCH codes over  $\mathbb{F}_q$  such that  $d = d_B$  are listed:

- the primitive case:  $n = q^m - 1$ 
  - $d = d_B = q^i - 1$ , where  $1 \leq i \leq m - 1$  [34];
  - $q = 2, d = d_B = 2^{m-1} - 2^i - 1$ , where  $\frac{m-2}{2} \leq i \leq m - \lfloor \frac{m}{3} \rfloor - 1$  [2];
  - $q = 2, d = d_B = 2^{m-1-s} - 2^{m-1-i-s}$ , where  $1 \leq i \leq m - s - 2$  and  $0 \leq s \leq m - 2i$  [20];
  - $d = d_B = q^i + 1$ , where  $(m - i) \mid i$  or  $2i \mid m$  [9];
  - $d = d_B = (q - l_0)q^{m-l_0-1} - 1$ , where  $0 \leq l_0 \leq q - 2$  and  $0 \leq l_1 \leq m - 1$  [7];
  - $d = d_B = q^m - q^{m-1} - q^i - 1$ , where  $\frac{m-2}{2} \leq i \leq m - \lfloor \frac{m}{3} \rfloor - 1$  [Theorem 1.2, [25]].
- $n = \frac{q^m - 1}{2}$ 
  - $q = 3, d = d_B = \frac{q^m - q^{m-1}}{2} - 1 - \frac{q^{\lfloor \frac{m-3}{2} \rfloor + i} - 1}{2}$  for  $i = 1, 2$  [27];
  - $d = d_B = \frac{q^m - q^{m-1}}{2} - 1 - \frac{q^{\lfloor \frac{m-3}{2} \rfloor + i} - 1}{2}$  for  $i = 1, 2$  [43];
  - $d = d_B = \frac{q^m - q^{m-1}}{2} - 1 - \frac{q^{\lfloor \frac{m-3}{2} \rfloor + i} - 1}{2}$ , where  $1 \leq i \leq \lfloor \frac{m+3}{4} \rfloor$  [this paper].

To determine the minimal distance  $d$  and the Bose distance  $d_B$  of narrow-sense primitive BCH codes with designed distance  $q^m - q^{m-1} - q^i - 1$ , Li [25] employed the framework of association schemes from bilinear forms associated with quadratic forms and presented the weight enumerators of these BCH codes, where these quadratic forms are corresponding to the trace representation of BCH codes. In [36], Tang et al. studied a class of ternary linear codes with trace representation, determined the weight distributions of some shortened codes and punctured codes of these three-weight subcodes, and presented some 2-designs from these codes.

Motivated by these papers and their ideas, this paper studies the following linear codes  $C_{1,h}$  (resp.  $C_{2,h}$ ) of length  $n = \frac{q^m - 1}{2}$  with special trace representation over finite field  $\mathbb{F}_q$  for  $m$  odd (resp. even), where  $q$  is odd prime power and

$$C_{1,h} = \left\{ \left( \text{Tr}_1^n \left( \sum_{j=h}^{\frac{m-1}{2}} a_j \alpha^{(q^j+1)l} \right) + a \right)_{l=0}^{n-1} : a_h, \dots, a_{\frac{m-1}{2}} \in \mathbb{F}_{q^m}, a \in \mathbb{F}_q \right\}$$

and

$$C_{2,h} = \left\{ \left( \text{Tr}_1^m \left( a_{\frac{m}{2}} \alpha^{(q^{\frac{m}{2}+1)l} + \sum_{j=h}^{\frac{m-2}{2}} a_j \alpha^{(q^j+1)l} \right) + a \right)_{l=0}^{n-1} : a_h, \dots, a_{\frac{m-2}{2}} \in \mathbb{F}_{q^m}, a_{\frac{m}{2}} \in \mathbb{F}_{q^{\frac{m}{2}}}, a \in \mathbb{F}_q \right\}.$$

These codes correspond to quadratic forms. With the help of the theory of association schemes from bilinear forms associated with quadratic forms, we determine the weight enumerators of these codes. Determining some largest cyclotomic coset leaders of cyclotomic cosets modulo  $n = \frac{q^m-1}{2}$ , we prove that a narrow-sense BCH code with designed distance  $\delta_i = \frac{q^m-q^{m-1}}{2} - 1 - q^{\lfloor \frac{m-3}{2} \rfloor + i} - 1$  has the above trace representation and has the minimal distance  $d$  such that  $d = d_B = \delta_i$ , where  $1 \leq i \leq \lfloor \frac{m+3}{4} \rfloor$ .

The rest of the paper is organized as follows. Section 2 introduces some basic results on BCH codes, quadratic forms and association schemes. Section 3 studies some linear codes with trace representation, uses association schemes to determine the weight enumerators of these codes, and presents the minimum distance  $d$  and the Bose distance of the narrow-sense BCH codes of length  $n = \frac{q^m-1}{2}$  with designed distance  $\delta_i$ . Section 4 makes a conclusion.

## 2 Preliminaries

In this section, some results on BCH codes, quadratic forms and association schemes are introduced.

### 2.1 BCH codes

Let  $q$  be a prime power. An  $[n, k, d]$  linear code  $C$  over the finite field  $\mathbb{F}_q$  is a  $k$ -dimensional subspace of  $\mathbb{F}_q^n$ , where  $d$  is the minimal distance of  $C$ . A linear code is called a cyclic code if  $(c_0, c_1, \dots, c_{n-1}) \in C$  implies that  $(c_{n-1}, c_0, \dots, c_{n-2}) \in C$ . We also write a cyclic code as a principal ideal of the ring  $\mathbb{F}_q[x]/(x^n - 1)$ , since there is a map

$$\begin{aligned} \Psi: \mathbb{F}_q^n &\longrightarrow \mathbb{F}_q[x]/(x^n - 1) \\ (c_0, c_1, \dots, c_{n-1}) &\longmapsto c_1 + c_1x + \dots + c_{n-1}x^{n-1}. \end{aligned}$$

Then  $C = \langle g(x) \rangle$ , where  $g(x)$  is a monic polynomial and is called the generator polynomial of  $C$ . Note that  $g(x)|(x^n - 1)$ . The polynomial  $h(x) = g(x)|(x^n - 1)$  is called the parity-check polynomial of  $C$ . The code  $C$  is called a cyclic code with  $s$  zeros (resp.  $s$  nonzeros) if  $g(x)$  (resp.  $h(x)$ ) can be factored into a product of  $s$  irreducible polynomials over  $\mathbb{F}_q$ .

Let  $n$  be a positive integer and  $m$  be the smallest positive integer such that  $n|(q^m - 1)$ . Let  $\alpha$  be a primitive element of  $\mathbb{F}_{q^m}$  and  $\beta = \alpha^{\frac{q^m-1}{n}}$ . Let  $m_i(x)$  be the minimal polynomial of  $\beta^i$  over  $\mathbb{F}_q$ , where  $0 \leq i \leq n - 1$ . For each  $2 \leq \delta \leq n$ , define two polynomials

$$g_{(n,q,m,\delta)}(x) = lcm(m_1(x), \dots, m_{\delta-1}(x)) \text{ and } \tilde{g}_{(n,q,m,\delta)}(x) = (x-1)g_{(n,q,m,\delta)}(x),$$

where  $lcm$  denotes the least common multiple of the polynomials. Let  $C_{(n,q,m,\delta)} = \langle g_{(n,q,m,\delta)}(x) \rangle$  and  $\tilde{C}_{(n,q,m,\delta)} = \langle \tilde{g}_{(n,q,m,\delta)}(x) \rangle$ . Then  $C_{(n,q,m,\delta)}$  is called a narrow-sense BCH code with designed distance  $\delta$  and  $\tilde{C}_{(n,q,m,\delta)}$  is the even-like subcode of  $C_{(n,q,m,\delta)}$ . Note that  $dim(\tilde{C}_{(n,q,m,\delta)}) = dim(C_{(n,q,m,\delta)}) - 1$  and the minimal distances of  $C_{(n,q,m,\delta)}$  and  $\tilde{C}_{(n,q,m,\delta)}$  are at least  $\delta$  and  $\delta + 1$ , respectively. If  $n = q^m - 1$ , the code  $C_{(n,q,m,\delta)}$  is called a narrow-sense primitive BCH code. Note that two BCH codes with different designed distances may be the same. For a narrow-sense BCH code  $C_{(n,q,m,\delta)}$ , the Bose distance [31] of this code is defined by the largest designed distance  $d_B = \max(\delta')$ , where  $C_{(n,q,m,\delta')} = C_{(n,q,m,\delta)}$ .

The narrow-sense BCH code  $C_{(n,q,m,\delta)}$  has close relation with  $q$ -cyclotomic cosets modulo  $n$ . The  $q$ -cyclotomic coset of  $s$  modulo  $n$  is defined by

$$C_s = \{s, sq, sq^2, \dots, sq^{l_s-1}\} \pmod n \subseteq \mathbb{Z}_n,$$

where  $0 \leq s \leq n-1$  and  $l_s$  called the size of the  $q$ -cyclotomic coset is the smallest positive integer such that  $s \equiv sq^{l_s} \pmod n$ . Note that  $l_s \mid m$ . The coset leader of  $C_s$  is the smallest integer in  $C_s$ . Let  $\Gamma_{n,q}$  be the set of all the coset leaders. Then  $C_s \cap C_t = \emptyset$  and  $\mathbb{Z}_n = \bigcup_{s \in \Gamma_{n,q}} C_s$ , where  $s, t \in \Gamma_{n,q}$  and  $s \neq t$ .

Let  $m_s(x)$  be the minimal polynomial of  $\beta^s$  over  $\mathbb{F}_q$ . Then

$$m_s(x) = \prod_{i \in C_s} (x - \beta^i) \in \mathbb{F}_q[x],$$

$\deg(m_s(x)) = l_s$  and

$$x^n - 1 = \prod_{s \in \Gamma_{n,q}} m_s(x).$$

Hence, the generator polynomial  $g_{(n,q,m,\delta)}(x) = \prod_{s \in \bigcup_{j=1}^{\delta-1} C_j} m_s(x)$ , the parity-check polynomial of  $C_{(n,q,m,\delta)}$  is

$$h(x) = (x-1) \prod_{s \geq \delta, s \in \Gamma_{n,q}} m_s(x),$$

and the dimension of  $C_{(n,q,m,\delta)}$  is

$$\sum_{s \geq \delta, s \in \Gamma_{n,q}} l_s + 1.$$

The following lemma gives the relation between the Bose distance  $d_B$  of  $C_{(n,q,m,\delta)}$  and coset leaders.

**Lemma 1 (Proposition 4, [27])** *The Bose distance  $d_B$  of the narrow-sense BCH code  $C_{(n,q,m,\delta)}$  is a coset leader of a  $q$ -cyclotomic coset modulo  $n$ . Furthermore, if  $\delta$  is a coset leader, then  $d_B = \delta$ .*

From Delsarte's theorem [11], a cyclic code has the following trace representation.

**Proposition 1** *Let  $q$  be a prime power,  $n$  be a positive integer,  $m$  be the smallest positive integer such that  $n \mid (q^n - 1)$ , and  $\beta$  be an  $n$ -th primitive element of  $\mathbb{F}_{q^m}$ . Let  $C$  be a cyclic code of length  $n$  over  $\mathbb{F}_q$  with  $s$  nonzeros. Let  $\beta^{i_1}, \dots, \beta^{i_s}$  be  $s$  roots of its parity-check polynomial  $h(x)$ , which are not conjugate with each other. Then the code  $C$  has the following trace representation*

$$C = \{c(a_1, \dots, a_s) : a_j \in \mathbb{F}_{q^{l_j}}, \text{ for } 1 \leq j \leq s\},$$

where  $l_j$  is the size of the  $q$ -cyclotomic coset  $C_{i_j}$ ,  $\text{Tr}_1^{l_j}$  is the trace function from  $\mathbb{F}_{q^{l_j}}$  to  $\mathbb{F}_q$ , and  $c(a_1, \dots, a_s) = \left( \sum_{j=1}^s \text{Tr}_1^{l_j}(a_j \beta^{-li_j}) \right)_{l=0}^{n-1}$ .

## 2.2 Quadratic forms

This subsection introduces some results on quadratic forms [29]. Let  $q$  be an odd prime power and  $V$  be an  $n$ -dimensional vector space over  $\mathbb{F}_q$ . A quadratic form  $Q$  on  $V$  is a function from  $V$  to  $\mathbb{F}_q$  satisfying  $Q(\lambda x) = \lambda^2 Q(x)$ , where  $x \in V$  and  $\lambda \in \mathbb{F}_q$ . A symmetric bilinear form  $B_Q(x, y) = \frac{1}{2}(Q(x+y) - Q(x) - Q(y))$  on  $V$  is associated with  $Q$ . The radical of  $Q$  is  $Rad(Q) = Q^{-1}(0) \cap Rad(B_Q)$ , where  $Rad(B_Q) = \{y \in V : B(x, y) = 0, \forall x \in V\}$  is the radical of the symmetric bilinear form  $B_Q$ . The radical of  $Q$  is a vector space over  $\mathbb{F}_q$ . The rank of  $Q$  is  $rank(Q) = n - dim(Rad(Q))$  and the rank of  $B_Q$  is  $rank(B_Q) = n - dim(Rad(B_Q))$ .

**Lemma 2 (Lemma 3.6, [25])** *Let  $q$  be an odd prime power,  $Q$  be a quadratic form on the  $n$ -dimensional vector space  $V$  over  $\mathbb{F}_q$  and  $B_Q$  be its associated bilinear form. Then  $rank(Q) = rank(B_Q)$ .*

Let  $V = \mathbb{F}_q^n$  and  $Q$  be a quadratic form on  $V$ . Then  $Q$  has the following expression  $Q(x) = \sum_{1 \leq i, j \leq n} c_{ij} x_i x_j$ . Two quadratic forms  $Q$  and  $Q'$  are equivalent if there is an  $n \times n$  nonsingular matrix  $A$  such that  $Q(x_1, \dots, x_n) = Q'((x_1, \dots, x_n)A)$ . Every quadratic form  $Q$  is equivalent to  $\sum_{i=1}^r a_i x_i^2$ , where  $a_i \in \mathbb{F}_q^*$  and  $r = rank(Q)$ . The type of  $Q$  is  $\tau = \eta(\prod_{i=1}^r a_i)$ , where  $\eta$  is the quadratic character of  $\mathbb{F}_q$ .

**Lemma 3 (Lemma 5.1, [35])** *Let  $q$  be an odd prime power and  $Q$  be quadratic form of rank  $r$  and type  $\tau$ . Let  $N(b)$  be the number of solutions  $Q(x) = b$ , where  $b \in \mathbb{F}_q$ . Then*

$$N(b) = \begin{cases} q^{n-1} + \tau \eta(-1)^{\frac{r-1}{2}} \eta(b) q^{n-\frac{r+1}{2}}, & \text{if } r \text{ is odd;} \\ q^{n-1} + \tau \eta(-1)^{\frac{r}{2}} v(b) q^{n-\frac{r+2}{2}}, & \text{if } r \text{ is even.} \end{cases}$$

where  $\eta(0) = 0$ ,  $v(x) = -1$  for  $x \in \mathbb{F}_q^*$  and  $v(0) = q - 1$ .

Let  $V = \mathbb{F}_q^n$ . Choose a basis  $\alpha_1, \dots, \alpha_n$  of  $V$  over  $\mathbb{F}_q$ . We have a bijection between  $\mathbb{F}_q^n$  and  $\mathbb{F}_q^n$ . Note that the rank  $r$  and the type  $\tau$  of a quadratic form  $Q$  on  $V$  are independent of the choice of the basis.

## 2.3 Association schemes

This subsection introduces some results on association schemes [31, 18, 37]. Let  $X$  be a finite set and  $R_0, R_1, \dots, R_n$  be a partition of  $X \times X$ . Then a pair  $(X, (R_i))$  is called an association scheme with  $n$  classes if it satisfies

- $R_0 = \{(x, x) : x \in X\}$ ;
- for each  $i$ , there exists  $j$  such that the inverse of  $R_i$  equals  $R_j$ ;
- if  $(x, y) \in R_k$ , the number of the set  $\{z \in X : (x, z) \in R_i, (z, y) \in R_j\}$  is a constant  $p_{ij}^k$  depending on only  $i, j$ , and  $k$ , but not on the particular choice of  $x$  and  $y$ .

An association scheme  $(X, (R_i))$  is symmetric if for all  $i$  the inverse of  $R_i$  equals  $R_i$ . It is commutative if for all  $i, j$ , and  $k$ ,  $p_{ij}^k = p_{ji}^k$ .

Let  $(X, (R_i))$  be a commutative association scheme with  $n$  classes and  $A_i$  be the adjacency matrix of the diagraph  $(X, R_i)$ . The matrices  $A_0, A_1, \dots, A_n$  span a vector space over the complex numbers called the Bose-Mesner algebra of  $(X, (R_i))$  with dimension  $n + 1$ . This vector space has another uniquely defined basis consisting of minimal idempotent matrices  $E_0, E_1, \dots, E_n$ . Then

$$A_i = \sum_{k=0}^n P_i(k) E_k \text{ and } E_k = \frac{1}{|X|} \sum_{i=0}^n Q_k(i) A_i.$$

The uniquely defined numbers  $P_i(k)$  and  $Q_k(i)$  are called the  $P$ -numbers and the  $Q$ -numbers of  $(X, (R_i))$ , respectively.

Let  $V$  be a vector space over the finite field  $\mathbb{F}_q$  of dimension  $m$  and  $X(m, q)$  be the set of symmetric bilinear forms on  $V$ , where  $q$  is an odd prime power. Let  $\alpha_1, \dots, \alpha_m$  be a basis of  $V$ . Then a symmetric bilinear form  $B \in X(m, q)$  has the  $m \times m$  symmetric matrix

$$(B(\alpha_i, \alpha_j))_{1 \leq i, j \leq m}.$$

The rank of  $B$  is the rank of this matrix, which is independent of the choice of the basis. This matrix is congruent to a diagonal matrix, whose diagonal is either zero or  $[z, 1, \dots, 1, 0, \dots, 0]$  for some nonzero  $z \in \mathbb{F}_q$ . The type of  $B$  is  $\eta(z)$ , where  $\eta$  is the quadratic character of  $\mathbb{F}_q$ .

Let  $X_{r,\tau}$  be the set of all the symmetric bilinear forms with rank  $r$  and type  $\tau$ . Define

$$R_{r,\tau} = \{(A, B) \in X(m, q) \times X(m, q) : A - B \in X_{r,\tau}\}.$$

Then  $(X(m, q), (R_{r,\tau}))$  is an association scheme with  $2m$  classes.

Let  $Y$  be a subset of  $X(m, q)$  and the inner distribution of  $Y$  be the sequence of numbers  $(a_{r,\tau})$ , where

$$a_{r,\tau} = \frac{|(Y \times Y) \cap R_{r,\tau}|}{|Y|}.$$

Let  $Q_{k,\varepsilon}(r, \tau)$  be the  $Q$ -numbers of  $(X(m, q), (R_{r,\tau}))$ . The dual inner distribution of  $Y$  is the sequence of  $(a'_{k,\varepsilon})$ , where

$$a'_{k,\varepsilon} = \sum_{r,\tau} Q_{k,\varepsilon}(r, \tau) a_{r,\tau}.$$

Note that  $\tau \in \{1, -1\}$ . When  $r = 0$ , we write  $a_0$ .

**Definition 1** Let  $Y$  be a subset of  $X(m, q)$ . The set  $Y$  is a  $d$ -code if  $a_{i,1} = a_{i,-1} = 0$  for each  $i \in \{1, 2, \dots, d-1\}$ . The set  $Y$  is a proper  $d$ -code if it is a  $d$ -code and it is not a  $(d+1)$ -code. The set  $Y$  is a  $t$ -design if  $a'_{i,1} = a'_{i,-1} = 0$  for each  $i \in \{1, 2, \dots, t\}$ . The set  $Y$  is a  $(2t+1, \varepsilon)$ -design if it is a  $(2t+1)$ -design and  $a'_{2t+2,\varepsilon} = 0$ .

Note that the designs involved in associate schemes are not the usual  $t$ -designs studied in combinatorial design theory. Define  $q^2$ -analogs of binomial coefficients

$$\begin{bmatrix} n \\ k \end{bmatrix} = \prod_{i=1}^k \frac{q^{2n-2i+2} - 1}{q^{2i} - 1}$$

for integers  $n$  and  $k \geq 0$ . Note that  $\begin{bmatrix} n \\ 0 \end{bmatrix} = 1$ . The inner distribution of  $Y$  can be given by  $q^2$ -analogs of binomial coefficients in the following theorem and proposition.

**Theorem 1 (Theorem 3.9, [35])** *If  $Y$  is a  $(2\delta - 1)$ -code and a  $(2n - 2\delta + 3)$ -design in  $X(2n + 1, q)$ , then the inner distribution  $(a_{r,\tau})$  of  $Y$  satisfies*

$$a_{2i-1,\tau} = \frac{1}{2} \begin{bmatrix} n \\ i-1 \end{bmatrix} \sum_{j=0}^{i-\delta} (-1)^j q^{j(j-1)} \begin{bmatrix} i \\ j \end{bmatrix} \left( \frac{|Y|}{q^{(2n+1)(n+1+j-i)}} - 1 \right)$$

$$a_{2i,\tau} = \frac{1}{2} (q^{2i} + \tau \eta(-1)^i q^i) \begin{bmatrix} n \\ i \end{bmatrix} \sum_{j=0}^{i-\delta} (-1)^j q^{j(j-1)} \begin{bmatrix} i \\ j \end{bmatrix} \left( \frac{|Y|}{q^{(2n+1)(n+1+j-i)}} - 1 \right)$$

for  $i > 0$ . If  $Y$  is a  $(2\delta - 1)$ -code and a  $(2n - 2\delta + 2)$ -design in  $X(2n, q)$ , then the inner distribution  $(a_{r,\tau})$  of  $Y$  satisfies

$$\begin{aligned} a_{2i-1,\tau} &= \frac{1}{2}(q^{2i} - 1) \begin{bmatrix} n \\ i \end{bmatrix} \sum_{j=0}^{i-\delta} (-1)^j q^{j(j-1)} \begin{bmatrix} i-1 \\ j \end{bmatrix} \frac{|Y|q^{2j}}{q^{(2n+1)(n+1+j-i)}} \\ a_{2i,\tau} &= \frac{1}{2} \begin{bmatrix} n \\ i \end{bmatrix} \sum_{j=0}^{i-\delta+1} (-1)^j q^{j(j-1)} \begin{bmatrix} i \\ j \end{bmatrix} \left( \frac{|Y|q^{2j}}{q^{(2n+1)(n+j-i)}} - 1 \right) \\ &\quad + \frac{\tau}{2} \eta (-1)^i q^i \begin{bmatrix} n \\ i \end{bmatrix} \sum_{j=0}^{i-\delta} (-1)^j q^{j(j-1)} \begin{bmatrix} i \\ j \end{bmatrix} \left( \frac{|Y|}{q^{(2n-1)(n+j-i)} q^{2n}} - 1 \right) \end{aligned}$$

for  $i > 0$ .

**Proposition 2 (Proposition 3.10, [35])** If  $Y$  is a  $(2\delta)$ -code and a  $(2n - 2\delta + 1)$ -design in  $X(2n, q)$ , then the inner distribution  $(a_{r,\tau})$  of  $Y$  satisfies

$$\begin{aligned} a_{2i-1,\tau} &= \frac{1}{2}(q^{2i} - 1) \begin{bmatrix} n \\ i \end{bmatrix} \sum_{j=0}^{i-\delta-1} (-1)^j q^{j(j-1)} \begin{bmatrix} i-1 \\ j \end{bmatrix} \frac{|Y|q^{2j}}{q^{(2n+1)(n+1+j-i)}} \\ a_{2i,\tau} &= \frac{1}{2} \begin{bmatrix} n \\ i \end{bmatrix} \sum_{j=0}^{i-\delta} (-1)^j q^{j(j-1)} \begin{bmatrix} i \\ j \end{bmatrix} \left( \frac{|Y|q^{2j}}{q^{(2n+1)(n+j-i)}} - 1 \right) \\ &\quad + \frac{\tau}{2} \eta (-1)^i q^i \begin{bmatrix} n \\ i \end{bmatrix} \sum_{j=0}^{i-\delta} (-1)^j q^{j(j-1)} \begin{bmatrix} i \\ j \end{bmatrix} \left( \frac{|Y|}{q^{(2n-1)(n+j-i)} q^{2n}} - 1 \right) \end{aligned}$$

for  $i > 0$ . If  $Y$  is a  $(2\delta)$ -code and a  $(2n - 2\delta + 1, \eta(-1)^{n-\delta+1})$ -design in  $X(2n + 1, q)$ , then the inner distribution  $(a_{r,\tau})$  of  $Y$  satisfies

$$\begin{aligned} a_{2i-1,\tau} &= \frac{1}{2} \begin{bmatrix} n \\ i-1 \end{bmatrix} \sum_{j=0}^{i-\delta} (-1)^j q^{j(j-1)} \begin{bmatrix} i \\ j \end{bmatrix} \left( \frac{|Y|}{q^{(2n+1)(n+1+j-i)}} - 1 \right) \\ &\quad + \frac{1}{2} (-1)^{i-\delta} q^{(i-\delta)(i-\delta-1)} \begin{bmatrix} n \\ \delta-1 \end{bmatrix} \left( \frac{|Y|}{q^{(2n+1)(n-\delta+1)}} - 1 \right) \left( \begin{bmatrix} n-\delta \\ n-i+1 \end{bmatrix} (q^{n-\delta+1} + 1) - \begin{bmatrix} n-\delta+1 \\ n-i+1 \end{bmatrix} \right) \\ a_{2i,\tau} &= \frac{1}{2} (q^{2i} + \tau \eta (-1)^i q^i) \begin{bmatrix} n \\ i \end{bmatrix} \sum_{j=0}^{i-\delta} (-1)^j q^{j(j-1)} \begin{bmatrix} i \\ j \end{bmatrix} \left( \frac{|Y|}{q^{(2n+1)(n+1+j-i)}} - 1 \right) \\ &\quad + \frac{1}{2} \eta (-1)^{i-\delta} q^{(i-\delta+1)(i-\delta)} \begin{bmatrix} n \\ \delta-1 \end{bmatrix} \begin{bmatrix} n-\delta \\ n-i \end{bmatrix} (q^{n-\delta+1} + 1) \left( \frac{|Y|}{q^{(2n+1)(n-\delta+1)}} - 1 \right) \end{aligned}$$

for  $i > 0$ .

### 3 A class of linear codes of length $n = \frac{q^m-1}{2}$

In this section, let  $q$  be an odd prime power,  $m$  be a positive integer,  $\alpha$  be a generator of  $\mathbb{F}_{q^m}^*$  and  $\beta = \alpha^2$ . Then  $\beta$  is a primitive  $n$ -th root of unity in  $\mathbb{F}_{q^m}$ , where  $n = \frac{q^m-1}{2}$ .

Let  $h$  be a positive integer. When  $m$  is odd, define the codes

$$C_{1,h} = \left\{ \left( \text{Tr}_1^m \left( \sum_{j=h}^{\frac{m-1}{2}} a_j \alpha^{(q^j+1)l} \right) + a \right)_{l=0}^{n-1} : a_h, \dots, a_{\frac{m-1}{2}} \in \mathbb{F}_{q^m}, a \in \mathbb{F}_q \right\}$$

and

$$\tilde{C}_{1,h} = \left\{ \left( \text{Tr}_1^m \left( \sum_{j=h}^{\frac{m-1}{2}} a_j \alpha^{(q^j+1)l} \right) \right)_{l=0}^{n-1} : a_h, \dots, a_{\frac{m-1}{2}} \in \mathbb{F}_{q^m}, a \in \mathbb{F}_q \right\}.$$

When  $m$  is even, define the codes

$$C_{2,h} = \left\{ \left( \text{Tr}_1^m \left( a_{\frac{m}{2}} \alpha^{(q^{\frac{m}{2}+1)l} + \sum_{j=h}^{\frac{m-2}{2}} a_j \alpha^{(q^j+1)l} \right) + a \right)_{l=0}^{n-1} : a_h, \dots, a_{\frac{m-2}{2}} \in \mathbb{F}_{q^m}, a_{\frac{m}{2}} \in \mathbb{F}_{q^{\frac{m}{2}}}, a \in \mathbb{F}_q \right\}$$

and

$$\tilde{C}_{2,h} = \left\{ \left( \text{Tr}_1^m \left( a_{\frac{m}{2}} \alpha^{(q^{\frac{m}{2}+1)l} + \sum_{j=h}^{\frac{m-2}{2}} a_j \alpha^{(q^j+1)l} \right) \right)_{l=0}^{n-1} : a_h, \dots, a_{\frac{m-2}{2}} \in \mathbb{F}_{q^m}, a_{\frac{m}{2}} \in \mathbb{F}_{q^{\frac{m}{2}}}, a \in \mathbb{F}_q \right\}.$$

Let  $Q(x)$  be a quadratic form defined by

$$Q(x) = \text{Tr}_1^m \left( \sum_{j=1}^l a_j x^{q^{k_j}+1} \right),$$

where  $k_j \geq 0$  and  $a_j \in \mathbb{F}_{q^m}$ . The symmetric bilinear form associated with  $Q(x)$  is

$$\begin{aligned} B_Q(x, y) &= \frac{1}{2} (Q(x+y) - Q(x) - Q(y)) \\ &= \frac{1}{2} \sum_{j=0}^l \text{Tr}_1^m \left( a_j (x^{q^{k_j}} + y^{q^{k_j}})(x+y) - a_j x^{q^{k_j}+1} - a_j y^{q^{k_j}+1} \right) \\ &= \frac{1}{2} \sum_{j=1}^l \text{Tr}_1^m (a_j x^{q^{k_j}} y + a_j x y^{q^{k_j}}) \\ &= \frac{1}{2} \sum_{j=1}^l \text{Tr}_1^m (a_j x^{q^{k_j}} y) + \frac{1}{2} \sum_{j=1}^l \text{Tr}_1^m (a_j x y^{q^{k_j}}) \\ &= \sum_{j=1}^l \text{Tr}_1^m \left( \left( \frac{a_j}{2} \right) x^{q^{k_j}} y \right) + \sum_{j=1}^l \text{Tr}_1^m \left( \left( \frac{a_j}{2} \right) q^{-k_j} x^{q^{-k_j}} y \right) \\ &= \sum_{j=0}^l \text{Tr}_1^m \left( \left( \left( \frac{a_j}{2} \right) x^{q^{k_j}} + \left( \frac{a_j}{2} \right) q^{-k_j} x^{q^{-k_j}} \right) y \right). \end{aligned}$$

Let  $h \geq 1$ . Define the set of quadratic forms:

$$Q_1 = \left\{ \text{Tr}_1^m \left( \sum_{j=h}^{\frac{m-1}{2}} a_j x^{q^j+1} \right) : a_j \in \mathbb{F}_{q^m} \text{ for } h \leq j \leq \frac{m-1}{2} \right\},$$

where  $m$  is odd. Define the set of quadratic forms:

$$Q_2 = \left\{ \text{Tr}_1^m \left( a_{\frac{m}{2}} x^{q^{\frac{m}{2}+1}} + \sum_{j=h}^{\frac{m-2}{2}} a_j x^{q^j+1} \right) : a_{\frac{m}{2}} \in \mathbb{F}_{q^{\frac{m}{2}}}, a_j \in \mathbb{F}_{q^m} \text{ for } h \leq j \leq \frac{m-2}{2} \right\},$$



where  $m$  is even. Then we have the following two sets of bilinear forms associated with  $Q_1$  and  $Q_2$  respectively.

$$\begin{aligned} S_1 &= \left\{ \text{Tr}_1^m \left( \sum_{j=h}^{\frac{m-1}{2}} \left( \left( \frac{a_j}{2} \right) x^{q^j} + \left( \frac{a_j}{2} \right)^{q^{-j}} x^{q^{-j}} \right) y \right) : a_j \in \mathbb{F}_{q^m} \text{ for } h \leq j \leq \frac{m-1}{2} \right\} \\ &= \left\{ \text{Tr}_1^m \left( \sum_{j=h}^{\frac{m-1}{2}} \left( a_j x^{q^j} + a_j^{q^{-j}} x^{q^{-j}} \right) y \right) : a_j \in \mathbb{F}_{q^m} \text{ for } h \leq j \leq \frac{m-1}{2} \right\} \end{aligned}$$

and

$$\begin{aligned} S_2 &= \left\{ \text{Tr}_1^m \left( \frac{a_{\frac{m}{2}}}{2} x^{\frac{m}{2}} + \sum_{j=h}^{\frac{m-2}{2}} \left( \left( \frac{a_j}{2} \right) x^{q^j} + \left( \frac{a_j}{2} \right)^{q^{-j}} x^{q^{-j}} \right) y \right) : a_{\frac{m}{2}} \in \mathbb{F}_{q^{\frac{m}{2}}}, a_j \in \mathbb{F}_{q^m} \text{ for } h \leq j \leq \frac{m-2}{2} \right\} \\ &= \left\{ \text{Tr}_1^m \left( a_{\frac{m}{2}} x^{\frac{m}{2}} + \sum_{j=h}^{\frac{m-2}{2}} \left( a_j x^{q^j} + a_j^{q^{-j}} x^{q^{-j}} \right) y \right) : a_{\frac{m}{2}} \in \mathbb{F}_{q^{\frac{m}{2}}}, a_j \in \mathbb{F}_{q^m} \text{ for } h \leq j \leq \frac{m-2}{2} \right\}. \end{aligned}$$

Then  $|S_1| = |S_2| = q^{m(\frac{m+1}{2}-h)}$ . The inner distributions of  $S_1$  and  $S_2$  are given in the following proposition.

**Proposition 3** *Let  $(a_0, a_{1,1}, a_{1,-1}, \dots, a_{m,1}, a_{m,-1})$  be the inner distribution of  $S_i$ . If  $m$  is odd, then  $S_1$  is a proper  $(2h+1)$ -code and  $(m+1-2h)$ -design in  $X(m, q)$ , and the inner distribution of  $S_1$  is given in Theorem 1. If  $m$  is even, then  $S_2$  is a proper  $(2h)$ -code and  $(m+1-2h)$ -design in  $X(m, q)$ , and the inner distribution of  $S_2$  is given in Proposition 2.*

*Proof* We first prove that  $S_1$  is a  $(m+1-2h)$ -design in  $X(m, q)$ , where  $m$  is odd.

Let  $U$  be a  $t$ -dimensional subspace of  $\mathbb{F}_{q^m}$  and  $A$  be a symmetric bilinear form on  $U$ , where  $t = m+1-2h$ . For  $a = (a_0, a_1, \dots, a_{t-1}) \in \mathbb{F}_{q^m}^t$ , define a bilinear form  $B_a$  on  $\mathbb{F}_{q^m}^t$

$$B_a(x, y) = \text{Tr}_1^m \left( \sum_{j=0}^{t-1} a_j x^{q^{j+h}} y \right) = \text{Tr}_1^m \left( \sum_{j=h}^{m-h} a_{j-h} x^{q^{sj}} y \right).$$

From Lemma 4.6 in [35], the set  $\{B_a \mid U : a \in \mathbb{F}_{q^m}^t\}$  is a multiset in which each bilinear form on  $U$  occurs a constant number (depending only on  $t$ ) of times. From Lemma 4.3 in [25], the number of elements in the multiset  $\{D+D' : D \in \{B_a : a \in \mathbb{F}_{q^m}^t\}\}$  that are an extension of  $A$  is a constant independent of  $U$  and  $A$ .

For  $a = (a_0, a_1, \dots, a_{t-1}) \in \mathbb{F}_{q^m}^t$ , we have

$$\begin{aligned}
B_a(x, y) + B'_a(x, y) &= \text{Tr}_1^m \left( \sum_{j=h}^{m-h} a_{j-h} (x^{q^{sj}} y + xy^{q^{sj}}) \right) \\
&= \text{Tr}_1^m \left( \sum_{j=h}^{\frac{m-1}{2}} a_{j-h} (x^{q^{sj}} y + xy^{q^{sj}}) \right) + \text{Tr}_1^m \left( \sum_{j=\frac{m+1}{2}}^{m-h} a_{j-h} (x^{q^{sj}} y + xy^{q^{sj}}) \right) \\
&= \text{Tr}_1^m \left( \sum_{j=h}^{\frac{m-1}{2}} a_{j-h} (x^{q^{sj}} y + xy^{q^{sj}}) \right) + \text{Tr}_1^m \left( \sum_{j=h}^{\frac{m-1}{2}} a_{m-j-h} (x^{q^{s(m-j)}} y + xy^{q^{s(m-j)}}) \right) \\
&= \text{Tr}_1^m \left( \sum_{j=h}^{\frac{m-1}{2}} a_{j-h} (x^{q^{sj}} y + xy^{q^{sj}}) \right) + \text{Tr}_1^m \left( \sum_{j=h}^{\frac{m-1}{2}} a_{m-j-h}^{sj} (x^{q^{sm}} y^{q^{sj}} + x^{q^{sj}} y^{q^{sm}}) \right) \\
&= \text{Tr}_1^m \left( \sum_{j=h}^{\frac{m-1}{2}} a_{j-h} (x^{q^{sj}} y + xy^{q^{sj}}) \right) + \text{Tr}_1^m \left( \sum_{j=h}^{\frac{m-1}{2}} a_{m-j-h}^{sj} (xy^{q^{sj}} + x^{q^{sj}} y) \right) \\
&= \text{Tr}_1^m \left( \sum_{j=h}^{\frac{m-1}{2}} (a_{j-h} + a_{m-j-h}^{sj}) (x^{q^{sj}} y + xy^{q^{sj}}) \right).
\end{aligned}$$

Then

$$\{D + D' : D \in \{B_a : a \in \mathbb{F}_{q^m}^t\}\} = \left\{ \text{Tr}_1^m \left( \sum_{j=h}^{\frac{m-1}{2}} (a_{j-h} + a_{m-j-h}^{sj}) (x^{q^{sj}} y + xy^{q^{sj}}) \right) : a \in \mathbb{F}_{q^m}^t \right\}.$$

Note that  $\{a_{j-h}, a_{m-j-h} : h \leq j \leq \frac{m-1}{2}\} = \{a_j : 0 \leq j \leq t-1\}$ . Then  $a_{j-h} + a_{m-j-h}^{sj}$  ranges over  $\mathbb{F}_{q^m}$  for  $q^m$  times when  $a_{j-h}$  and  $a_{m-j-h}$  ranges over  $\mathbb{F}_{q^m}$ . Hence, the number of elements in  $S_1$  which are an extension of  $A$  is a constant independent of  $U$  and  $A$ . From Theorem 3.11 in [35], the set  $S_1$  is a  $t$ -design in  $X(m, q)$ .

We then prove that  $S_1$  is a proper  $(2h+1)$ -code. For a bilinear form

$$B(x, y) = \text{Tr}_1^m \left( \sum_{j=h}^{\frac{m-1}{2}} (a_j x^{q^j} + a_j^{q^{-j}} x^{q^{-j}}) y \right) \in S_1,$$

we have

$$\begin{aligned}
\text{Rad}(B(x, y)) &= \left\{ x \in \mathbb{F}_{q^m} : \text{Tr}_1^m \left( \sum_{j=h}^{\frac{m-1}{2}} (a_j x^{q^j} + a_j^{q^{-j}} x^{q^{-j}}) y \right) = 0, \forall y \in \mathbb{F}_{q^m} \right\} \\
&= \left\{ x \in \mathbb{F}_{q^m} : \text{Tr}_1^m \left( \sum_{j=h}^{\frac{m-1}{2}} (a_j x^{q^j} + a_j^{q^{m-j}} x^{q^{m-j}}) y \right) = 0, \forall y \in \mathbb{F}_{q^m} \right\} \\
&= \left\{ x \in \mathbb{F}_{q^m} : \sum_{j=h}^{\frac{m-1}{2}} (a_j x^{q^j} + a_j^{q^{m-j}} x^{q^{m-j}}) y = 0, \forall y \in \mathbb{F}_{q^m} \right\} \\
&= \left\{ x \in \mathbb{F}_{q^m} : \sum_{j=h}^{\frac{m-1}{2}} (a_j^{-h} x^{q^{j-h}} + a_j^{q^{m-j-h}} x^{q^{m-j-h}}) y = 0, \forall y \in \mathbb{F}_{q^m} \right\}.
\end{aligned}$$

Since the degree of the linearized polynomial  $\sum_{j=h}^{\frac{m-1}{2}} (a_j^{-h} x^{q^{j-h}} + a_j^{q^{m-j-h}} x^{q^{m-j-h}})y$  over  $\mathbb{F}_{q^m}$  is at most  $m-2h$ , the dimension of the vector space  $\text{Rad}(B(x,y))$  is at most  $m-2h$ . Hence,  $\text{rank}(B(x,y)) = m - \dim(\text{Rad}(B(x,y))) \geq 2h$ . Hence,  $S_1$  is a  $(2h)$ -code in  $X(m,q)$ . Since  $S_1$  is a  $(2m+1-2h)$ -design, then  $S_1$  is a  $(2m-2h, \tau)$ -design. From Proposition 2 with  $\delta = h$ , we have  $a_{2h, \tau} = 0$  for  $\tau = \{1, -1\}$ . Hence,  $S_1$  is a  $(2h+1)$ -code. From Theorem 1, we have  $a_{2h+1, \tau} > 0$ . Hence,  $S_1$  is a proper  $(2h+1)$ -code.

From a similar discussion, we have the corresponding results for  $S_2$  when  $m$  is even.

**Lemma 4** *Let  $q$  be odd and  $Q$  be a quadratic form of rank  $r \geq 1$  and type  $\tau$  on  $\mathbb{F}_{q^m}$ . Then the weight enumerator  $W_{r, \tau}$  of  $(Q(\alpha^l) + a)_{l=0}^{n-1}$  is*

$$Z^{\frac{q^m - q^{m-1}}{2}} + \frac{q-1}{2} Z^{\frac{1}{2}} (q^m - q^{m-1} - \tau \eta(-1)^{\frac{r-1}{2}} q^{m - \frac{r+1}{2}} - 1) + \frac{q-1}{2} Z^{\frac{1}{2}} (q^m - q^{m-1} + \tau \eta(-1)^{\frac{r-1}{2}} q^{m - \frac{r+1}{2}} - 1)$$

if  $r$  is odd, or

$$Z^{\frac{q^m - q^{m-1} - \tau \eta(-1)^{\frac{r}{2}} (q-1) q^{m - \frac{r+2}{2}}}{2}} + (q-1) Z^{\frac{1}{2}} (q^m - q^{m-1} + \tau \eta(-1)^{\frac{r}{2}} q^{m - \frac{r+2}{2}} - 1)$$

if  $r$  is even.

*Proof* Let  $f(x) = Q(x) + a$ . We just compute the weight  $wt(f)$ . Let  $N(f=0)$  be the number of solutions of  $f(x) = 0$ .

When  $a = 0$ , from Lemma 3, we have

$$N(f=0) = \begin{cases} q^{m-1}, & \text{if } r \text{ is odd;} \\ q^{m-1} + \tau \eta(-1)^{\frac{r}{2}} (q-1) q^{m - \frac{r+2}{2}}, & \text{if } r \text{ is even.} \end{cases}$$

Then

$$wt(f) = \frac{q^m - N(f=0)}{2} = \begin{cases} \frac{q^m - q^{m-1}}{2}, & \text{if } r \text{ is odd;} \\ \frac{q^m - q^{m-1} - \tau \eta(-1)^{\frac{r}{2}} (q-1) q^{m - \frac{r+2}{2}}}{2}, & \text{if } r \text{ is even.} \end{cases}$$

When  $a \neq 0$  and  $\eta(-a) = 1$ , from Lemma 3, we have

$$N(f=0) = \begin{cases} q^{m-1} + \tau \eta(-1)^{\frac{r-1}{2}} q^{m - \frac{r+1}{2}}, & \text{if } r \text{ is odd;} \\ q^{m-1} - \tau \eta(-1)^{\frac{r}{2}} q^{m - \frac{r+2}{2}}, & \text{if } r \text{ is even.} \end{cases}$$

Then

$$wt(f) = \frac{q^m - 1 - N(f=0)}{2} = \begin{cases} \frac{1}{2} (q^m - q^{m-1} - \tau \eta(-1)^{\frac{r-1}{2}} q^{m - \frac{r+1}{2}} - 1), & \text{if } r \text{ is odd;} \\ \frac{1}{2} (q^m - q^{m-1} + \tau \eta(-1)^{\frac{r}{2}} q^{m - \frac{r+2}{2}} - 1), & \text{if } r \text{ is even.} \end{cases}$$

When  $a \neq 0$  and  $\eta(-a) = -1$ , from Lemma 3, we have

$$N(f=0) = \begin{cases} q^{m-1} - \tau \eta(-1)^{\frac{r-1}{2}} q^{m - \frac{r+1}{2}}, & \text{if } r \text{ is odd;} \\ q^{m-1} - \tau \eta(-1)^{\frac{r}{2}} q^{m - \frac{r+2}{2}}, & \text{if } r \text{ is even.} \end{cases}$$

Then

$$wt(f) = \frac{q^m - 1 - N(f=0)}{2} = \begin{cases} \frac{1}{2} (q^m - q^{m-1} + \tau \eta(-1)^{\frac{r-1}{2}} q^{m - \frac{r+1}{2}} - 1), & \text{if } r \text{ is odd;} \\ \frac{1}{2} (q^m - q^{m-1} + \tau \eta(-1)^{\frac{r}{2}} q^{m - \frac{r+2}{2}} - 1), & \text{if } r \text{ is even.} \end{cases}$$

Hence, this lemma follows.

**Lemma 5** Let  $q$  be odd and  $Q$  be a quadratic form of rank  $r \geq 1$  and type  $\tau$  on  $\mathbb{F}_q^m$ . Then the weight enumerator  $W_{r,\tau}$  of  $(Q(\alpha^l))_{l=0}^{n-1}$  is  $Z^{\frac{q^m - q^{m-1}}{2}}$  (resp.  $Z^{\frac{q^m - q^{m-1} - \tau(-1)^{\frac{m}{2}}(q-1)q^{\frac{m-r-2}{2}}}{2}}$ ) if  $r$  is odd (resp. even).

*Proof* From results of  $a = 0$  in Lemma 4, this lemma follows.

We determine the weight enumerators of  $C_{1,h}$ ,  $\tilde{C}_{1,h}$ ,  $C_{1,h}$ , and  $\tilde{C}_{1,h}$  in the following theorem.

**Theorem 2** When  $m$  is odd, the weight enumerator of the code  $C_{1,h}$  is

$$1 + (q-1)Z^{\frac{q^m-1}{2}} + \sum_{r=2h+1}^m \sum_{\tau \in \{1,-1\}} a_{r,\tau} W_{r,\tau},$$

where  $a_{r,\tau}$  is given in Theorem 1 and  $W_{r,\tau}$  is given in Lemma 4. The weight enumerator of the code  $\tilde{C}_{1,h}$  is

$$1 + \sum_{r=2h+1}^m \sum_{\tau \in \{1,-1\}} a_{r,\tau} W_{r,\tau}$$

where  $a_{r,\tau}$  is given in Theorem 1 and  $W_{r,\tau}$  is given in Lemma 5. When  $m$  is even, the weight enumerator of the code  $C_{2,h}$  is

$$1 + (q-1)Z^{\frac{q^m-1}{2}} + \sum_{r=2h}^m \sum_{\tau \in \{1,-1\}} a_{r,\tau} W_{r,\tau}$$

where  $a_{r,\tau}$  is given in Proposition 2 and  $W_{r,\tau}$  is given in Lemma 4. The weight enumerator of the code  $\tilde{C}_{2,h}$  is

$$1 + \sum_{r=2h}^m \sum_{\tau \in \{1,-1\}} a_{r,\tau} W_{r,\tau},$$

where  $a_{r,\tau}$  is given in Proposition 2 and  $W_{r,\tau}$  is given in Lemma 5.

*Proof* We first give the weight enumerator of  $C_{1,h}$ . When  $m$  is odd, we have

$$C_{1,h} = \bigcup_{Q \in Q_1} (Q(\alpha^l) + a)_{l=0}^{n-1}.$$

The weight enumerator of  $\{(a)_{l=0}^{n-1} : a \in \mathbb{F}_q\}$  is  $1 + (q-1)Z^{\frac{q^m-1}{2}}$ . From Proposition 3,  $S_1$  is a proper  $(2h+1)$ -code. For any nonzero  $Q(x) \in Q_1$ , we have  $\text{rank}(Q(x)) = r \geq 2h+1$ . The weight enumerator  $W_{r,\tau}$  of  $(Q(\alpha^l) + a)_{l=0}^{n-1}$  with  $\text{rank } r > 0$  is given in Lemma 4. The inner distribution of  $Q_1$  is just the inner distribution of  $S_1$ , which is given in Theorem 1. Hence, we have the weight enumerator of  $C_{1,h}$

$$1 + (q-1)Z^{\frac{q^m-1}{2}} + \sum_{r=2h+1}^m \sum_{\tau \in \{1,-1\}} a_{r,\tau} W_{r,\tau}.$$

From the similar method, we have the other results. Hence, this theorem follows.

**Lemma 6** When  $m \geq 2$  and  $1 \leq i \leq \lfloor \frac{m+3}{4} \rfloor$ , then  $\delta_i = \frac{q^m - q^{m-1}}{2} - 1 - \frac{q^{\lfloor \frac{m-3}{2} \rfloor + i - 1}}{2}$  is the  $i$ -th largest coset leader module  $n$ .

*Proof* When  $q = 3$ , from [27], this lemma holds. Suppose that  $q > 3$ . When  $m$  is odd, then the  $q$ -adic expansion of  $\delta_i$  is

$$\delta_i = \left(\frac{q-3}{2}\right)q^{m-1} + (q-1) \sum_{j=\frac{m-3}{2}+i}^{m-2} q^j + \left(\frac{q-1}{2}\right) \sum_{j=0}^{\frac{m-3}{2}+i-1} q^j.$$

and its cyclotomic coset  $C_{\delta_i}$  is

$$\left\{ \frac{q^m - 1 - q^{l-1} - q^{l+\frac{m-3}{2}+i}}{2} : 1 \leq l \leq \frac{m+1}{2} - i \right\} \cup \left\{ \frac{q^m - 1 - q^{l-1} - q^{l-\frac{m+3}{2}+i}}{2} : \frac{m+3}{2} - i \leq l \leq m \right\}.$$

From Lemma 9 in [43],  $\delta_1$  is the largest coset leader module  $n$ . Then we just prove that for  $i > 1$ , there does not exist a coset leader  $s$  satisfying  $\delta_i < s < \delta_{i-1}$ .

Suppose such a coset leader  $s$  satisfying  $\delta_i < s < \delta_{i-1}$  exists. From Lemma 8 in [43], then the  $q$ -adic expansion of  $s$  is

$$s = \left(\frac{q-3}{2}\right)q^{m-1} + (q-1) \sum_{j=k+1}^{m-2} q^j + \sum_{j=0}^k s_j q^j,$$

where  $k = \frac{m-3}{2} + i - 1$ ,  $\frac{q-3}{2} \leq s_j \leq q-1$  and  $\left(\frac{q-1}{2}\right) \sum_{j=0}^k q^j < \sum_{j=0}^k s_j q^j < (q-1)q^k + \left(\frac{q-1}{2}\right) \sum_{j=0}^{k-1} q^j$ .

When  $s_k = \frac{q-1}{2}$ , from  $t > \delta_i$ , we have  $(q^{m-2-t} s \bmod n) < s$ , where  $0 \leq t < k$  and  $t$  is the largest index such that  $s_t > \frac{q-1}{2}$ . It is a contradiction to the coset leader  $s$ .

When  $\frac{q-1}{2} < s_k < q-1$ , then  $(q^{m-1-k} s \bmod n) < s$ , which is a contradiction to the coset leader  $s$ .

When  $s_k = q-1$ , from  $s < \delta_{i-1}$ , there exists a  $t$  satisfying  $(q^{m-1-t} s \bmod n) < s$  or  $(q^{m-2-t} s \bmod n) < s$ , where  $0 \leq t < k$ . It is a contradiction to the coset leader  $s$ .

Hence,  $\delta_i$  is the  $i$ -th largest coset leader module  $n$  when  $m$  is odd. From the similar method, it holds when  $m$  is even.

The following theorem gives parameters of the BCH code  $C_{(n,q,m,\delta_i)}$ .

**Theorem 3** Let  $m \geq 2$ ,  $1 \leq i \leq \lfloor \frac{m+3}{4} \rfloor$ , and  $h = \lfloor \frac{m}{2} \rfloor - i + 1$ . Then the BCH code  $C_{(n,q,m,\delta_i)}$  is  $C_{1,h}$  (resp.  $C_{2,h}$ ) for  $m$  odd (resp. even) and it has parameters  $[n, (i+1 + \frac{m-1}{2} - \lfloor \frac{m}{2} \rfloor)m, \delta_i]$  and the Bose distance  $d_B = \delta_i$ . The code  $\tilde{C}_{(n,q,m,\delta_i)}$  is  $\tilde{C}_{1,h}$  (resp.  $\tilde{C}_{2,h}$ ) for  $m$  odd (resp. even) and it has dimension  $(i + \frac{m-1}{2} - \lfloor \frac{m}{2} \rfloor)m$ .

*Proof* Note that  $\delta_i$  is the  $i$ -th largest coset leader module  $n$ . When  $m$  is odd, we have  $l_i = |C_{\delta_i}| = m$  and

$$\begin{aligned} \text{Tr}_1^{l_i}(b\beta^{-\delta_i l}) &= \text{Tr}_1^m \left( b\alpha^{(-q^m + q^{m-1} + 1 + q^{\frac{m-3}{2}+i})l} \right) \\ &= \text{Tr}_1^m \left( b\alpha^{(q^{m-1} + q^{\frac{m-3}{2}+i})l} \right) \\ &= \text{Tr}_1^m \left( b^{-\frac{m-3}{2}-i} \alpha^{(q^{\frac{m+1}{2}-i+1})l} \right). \end{aligned}$$

Hence the code  $C_{(n,q,m,\delta_i)} = C_{1,h}$  and the dimension of  $C_{(n,q,m,\delta_i)}$  is  $(i+1 + \frac{m-1}{2} - \lfloor \frac{m}{2} \rfloor)m$ . When  $r = 2h+1$ , the  $(Q(\alpha^l) + a)_{l=0}^{n-1}$  has a codeword of weight  $\delta_i$ . From Lemma 1, the code

$C_{(n,q,m,\delta_i)}$  has Bose distance  $\delta_i$ . Hence, the code  $C_{(n,q,m,\delta_i)}$  has parameters  $[n, (i+1 + \frac{m-1}{2} - \lfloor \frac{m}{2} \rfloor)m, \delta_i]$  and Bose distance  $\delta_i$ . From the similar discussion, we have the other results of this proposition.

*Example 1* Let  $q = 3$ ,  $h = \lfloor \frac{m}{2} \rfloor$ , and  $i = 1$ . When  $m$  is odd (resp. even), the code  $C_{(n,3,m,\delta_1)}$  is  $C_{1, \frac{m-1}{2}}$  (resp.  $C_{2, \frac{m}{2}}$ ), and it has parameters  $[\frac{3^m-1}{2}, m+1, \delta_1]$  (resp.  $[\frac{3^m-1}{2}, \frac{m+2}{2}, \delta_1]$ ) and the weight distribution in Table II (resp. Table I) in Theorem 19 [27]. When  $m$  is odd (resp. even), the code  $\tilde{C}_{(n,3,m,\delta_1)}$  is  $\tilde{C}_{1, \frac{m-1}{2}}$  (resp.  $\tilde{C}_{2, \frac{m}{2}}$ ), and it has parameters  $[\frac{3^m-1}{2}, m, 3^{m-1}]$  (resp.  $[\frac{3^m-1}{2}, \frac{m}{2}, 3^{m-1} + 3^{\frac{m-2}{2}}]$ ) and the weight enumerator  $1 + (3^m - 1)Z^{3^{m-1}}$  (resp.  $1 + (3^{\frac{m}{2}} - 1)Z^{3^{m-1} + 3^{\frac{m-2}{2}}}$ ) in Theorem 22 [27].

*Example 2* Let  $q = 3$ ,  $h = \lfloor \frac{m}{2} \rfloor - 1$ , and  $i = 2$ . When  $m$  is odd (resp. even), the code  $C_{(n,3,m,\delta_2)}$  is  $C_{1, \frac{m-3}{2}}$  (resp.  $C_{2, \frac{m-2}{2}}$ ), and it has parameters  $[\frac{3^m-1}{2}, 2m+1, \delta_2]$  (resp.  $[\frac{3^m-1}{2}, \frac{3m+2}{2}, \delta_2]$ ) and the weight distribution in Table VI (resp. Table V) in Theorem 29 [27]. When  $m$  is odd (resp. even), the code  $\tilde{C}_{(n,3,m,\delta_2)}$  is  $\tilde{C}_{1, \frac{m-3}{2}}$  (resp.  $\tilde{C}_{2, \frac{m-2}{2}}$ ), and it has parameters  $[\frac{3^m-1}{2}, 2m, 3^{m-1} - 3^{\frac{m-1}{2}}]$  (resp.  $[\frac{3^m-1}{2}, \frac{3m}{2}, 3^{m-1} - 3^{\frac{m-2}{2}}]$ ) and the weight distribution in Table IV (resp. Table III) in Theorem 26 [27].

*Example 3* When  $h = \lfloor \frac{m}{2} \rfloor$  and  $i = 1$ , the code  $C_{(n,q,m,\delta_1)}$  (resp.  $\tilde{C}_{(n,q,m,\delta_1)}$ ) is given in Theorem 5 (resp. Theorem 4) in [43]. When  $h = \lfloor \frac{m}{2} \rfloor - 1$  and  $i = 2$ , the code  $C_{(n,q,m,\delta_2)}$  (resp.  $\tilde{C}_{(n,q,m,\delta_2)}$ ) is given in Theorem 7 (resp. Theorem 6) in [43].

*Remark 1* Let  $q = 3$ ,  $h = \lfloor \frac{m}{2} \rfloor - 1$ , and  $i = 2$ . The code  $\tilde{C}_{(n,3,m,\delta_2)}$  is a three-weight code with the weight distribution in Table 1 of [36]. Then these codes can be used to construct 2-designs.

## 4 Conclusion

This paper studies a class of linear codes of length  $\frac{q^m-1}{2}$  over  $\mathbb{F}_q$  with special trace representation, uses association schemes from bilinear forms associated with quadratic forms, and determines the weight enumerators of these codes. From determining some cyclotomic coset leaders  $\delta_i$  of cyclotomic cosets modulo  $\frac{q^m-1}{2}$ , we prove that narrow-sense BCH codes of length  $\frac{q^m-1}{2}$  with designed distance  $\delta_i = \frac{q^m-q^{m-1}}{2} - 1 - q^{\lfloor \frac{m-3}{2} \rfloor + i - 1}$  have the corresponding trace representation and have the minimal distance  $d = \delta_i$  and the Bose distance  $d_B = \delta_i$ , where  $1 \leq i \leq \lfloor \frac{m+3}{4} \rfloor$ . It is interesting to determine parameters of more BCH codes.

**Acknowledgement.** This work was supported by SECODE project and the National Natural Science Foundation of China (Grant No. 11871058, 11531002, 11701129). C. Tang also acknowledges support from 14E013, CXTD2014-4 and the Meritocracy Research Funds of China West Normal University. Y. Qi also acknowledges support from Zhejiang provincial Natural Science Foundation of China (LQ17A010008, LQ16A010005).

## References

1. S. A. Aly, A. Klappenecker, and P. K. Sarvepalli, On quantum and classical BCH codes, IEEE Trans. Inf. Theory, vol. 53, no. 3, pp. 1183-1188, Mar. 2007.
2. E. R. Berlekamp, The weight enumerators for certain subcodes of the second order binary Reed-Muller codes, Inform. and Control, 17, pp. 485-500, 1970.

3. R. C. Bose and D. K. Ray-Chaudhuri, On a class of error correcting binary group codes, *Information and control*, vol. 3, no. 1, pp. 68-79, 1960.
4. P. Charpin, On a class of primitive BCH codes, *IEEE Trans. Inf. Theory*, vol. 36, no. 1, pp. 222-228, Jan. 1990.
5. P. Charpin, Open problems on cyclic codes, in *Handbook of Coding Theory*, Vol. I, NorthHolland, Amsterdam, pp. 963-1063, 1998.
6. P. Charpin, T. Helleseeth, and V. A. Zinoviev, The coset distribution of triple-error-correcting binary primitive BCH codes, *IEEE Trans. Inf. Theory*, vol. 52, no. 4, pp. 1727-1732, Apr. 2006.
7. C. Ding, Parameters of several classes of BCH codes, *IEEE Trans. Inf. Theory*, vol. 61, no. 10, pp. 5322-5330, Oct. 2015.
8. C. Ding: *Designs from linear codes*. World Scientific, Singapore, 2018.
9. C. Ding, X. Du, and Z. Zhou, The Bose and minimum distance of a class of BCH codes, *IEEE Trans. Inf. Theory* vol. 61, no. 5, pp. 2351-2356, May 2015.
10. C. Ding, C. Fan, and Z. Zhou, The dimension and minimum distance of two classes of primitive BCH codes, *Finite Fields Appl.*, vol. 45, pp. 237-263, 2017.
11. P. Delsarte, On subfield subcodes of modified Reed-Solomon codes (Corresp.), *IEEE Trans. Inf. Theory*, vol. 21, no. 5, pp. 575-576, Sept. 1975.
12. Y. Desaki, T. Fujiwara, and T. Kasami, The weight distributions of extended binary primitive BCH codes of length 128, *IEEE Trans. Inf. Theory*, vol. 43, no. 4, pp. 1364-1371, Jul. 1997.
13. K. Feng and J. Luo, Weight distribution of some reducible cyclic codes, *Finite Fields Appl.*, vol. 14, no. 2, pp. 390-409, 2008.
14. T. Fujiwara, T. Takata, T. Kasami, and S. Lin, An approximation to the weight distribution of binary primitive BCH codes with designed distances 9 and 11 (Corresp.), *IEEE Trans. Inf. Theory*, vol. 32, no. 5, pp. 706-709, Sept. 1986.
15. D. C. Gorenstein and N. Zierler, A class of error-correcting codes in  $p^m$  symbols, *J. SIAM* 9 (1961), 207-214.
16. H. Helgert and R. Stinaff, Shortened BCH codes (Corresp.), *IEEE Trans. Inf. Theory*, vol. 19, no. 6, pp. 818-820, Nov. 1973.
17. A. Hocquenghem, Codes correcteurs d'erreurs, *Chiffres*, vol. 2, no. 2, pp. 147-156, 1959.
18. Y.J. Huo, Z.X. Wan, Nonsymmetric association schemes of symmetric matrices. *Acta Math. Appl. Sinica (English Ser.)* 9(3), 236-255, 1993.
19. T. Kasami, Weight distributions of Bose-Chaudhuri-Hocquenghem codes, in: R.C. Bose, T.A. Dowlings (Eds.), *Combinatorial Mathematics and Applications*, Univ. North Carolina Press, Chapel Hill, NC, 1969, Ch.20.
20. T. Kasami and S. Lin, Some results on the minimum weight of primitive BCH codes, *IEEE Trans. Inf. Theory*, vol. 18, no. 6, pp. 824-825, Nov. 1972.
21. T. Kasami, S. Lin, and W. W. Peterson, Linear codes which are invariant under the affine group and some results on minimum weights in BCH codes, *Electron. Commun. Japan*, vol. 50, no. 9, pp. 100-106, 1967.
22. T. Kasami and N. Tokura, Some remarks on BCH bounds and minimum weights of binary primitive BCH codes, *IEEE Trans. Inf. Theory*, vol. 15, no. 3, pp. 408-413, May 1969.
23. O. Keren and S. Litsyn, More on the distance distribution of BCH codes, *IEEE Trans. Inf. Theory*, vol. 45, no. 1, pp. 251-255, Jan. 1999.
24. I. Krasikov and S. Litsyn, On the distance distributions of BCH codes and their duals, *Des. Codes Cryptogr.*, vol. 23, no. 2, pp. 223-232, 2001.
25. S. Li, The minimum distance of some narrow-sense primitive BCH codes, *SIAM Journal on Discrete Mathematics*, vol. 31, no. 4, 2530-2569, 2017.
26. C. Li, C. Ding, and S. Li, LCD cyclic codes over finite fields, *IEEE Trans. Inf. Theory*, vol. 63, no. 7, pp. 4344-4356, Jul. 2017.
27. S. Li, C. Ding, M. Xiong, and G. Ge, Narrow-Sense BCH codes over  $GF(q)$  with length  $n = \frac{q^m-1}{q-1}$ , *IEEE Trans. Inf. Theory*, vol. 63, no. 11, pp. 7219-7236, Nov. 2017.
28. S. Li, C. Li, C. Ding, and H. Liu, Two Families of LCD BCH Codes, *IEEE Trans. Inf. Theory*, vol. 63, no. 9, pp. 5699-5717, Sept. 2017.
29. R. Lidl and H. Niederreiter, *Finite fields*, Cambridge University Press, Cambridge, 1997.
30. H. Liu, C. Ding, and C. Li, Dimensions of three types of BCH codes over  $GF(q)$ , *Discrete Mathematics*, vol. 240, no. 8, pp. 1910-1927, 2017.
31. F. J. MacWilliams and N. J. A. Sloane, *The theory of error-correcting codes*, North-Holland Pub. Co, 1977.
32. D. Mandelbaum, Two applications of cyclotomic cosets to certain BCH codes (Corresp.), *IEEE Trans. Inf. Theory*, vol. 26, no. 6, pp. 737-738, Nov. 1980.
33. H. B. Mann, On the number of information symbols in Bose-Chaudhuri codes, *Inf. Control*, vol. 5, no. 2, pp. 153-162, 1962.

34. W. W. Peterson, Some new results on finite fields and their application to the theory of BCH codes, in *Combinatorial Mathematics and Its Applications* (Proc. Conf., Univ. North Carolina, Chapel Hill, NC, 1967), University of North Carolina Press, Chapel Hill, NC, 1969, pp. 329-334.
35. K.-U. Schmidt, Symmetric bilinear forms over finite fields with applications to coding theory, *J. Algebraic Combin.*, 42 (2015), pp. 635-670.
36. C. Tang, C. Ding, M. Xiong, Steiner systems  $S(2, 4, \frac{3^m-1}{2})$  and 2-designs from ternary linear codes of length  $\frac{3^m-1}{2}$ , <https://arxiv.org/abs/1901.09228>
37. Y. Wang, C. Wang, C. Ma, J. Ma, Association schemes of quadratic forms and symmetric bilinear forms. *J. Algebr. Comb.* 17(2), 149-161, 2003.
38. J. K. Wolf, Adding two information symbols to certain nonbinary BCH codes and some applications, *Bell Labs Technical Journal*, vol. 48, no. 7, pp. 2405-2424, 1969.
39. H. Yan, A class of primitive BCH codes and their weight distribution, *AAECC*, Volume 29, Issue 1, pp 1-11, January 2018.
40. D. Yue and G. Feng, Minimum cyclotomic coset representatives and their applications to BCH codes and Goppa codes, *IEEE Trans. Inf. Theory*, vol. 46, no. 7, pp. 2625-2628, 2000.
41. D. Yue and Z. Hu, On the dimension and minimum distance of BCH codes over  $GF(q)$ , *J. of Electronics*, vol. 13, no. 3, pp. 216-221, 1996.
42. Z. Zhou and C. Ding, A class of three-weight cyclic codes, *Finite Fields Appl.*, vol. 25, pp. 79-93, 2014.
43. S. Zhu, Z. Sun, and X. Kai, A Class of Narrow-Sense BCH Codes. [arXiv:1902.04372v1](https://arxiv.org/abs/1902.04372v1), 12 Feb 2019.