

COMPLEX ASTEROSEISMOLOGY OF SX PHOENICIS

J. Daszyńska-Daszkiewicz¹, P. Walczak¹, A. Pamyatnykh², W. Szewczuk¹ and W. Dziembowski²

Abstract. We present seismic analysis of the prototype SX Phoenicis that aims at fitting the two radial-mode frequencies and the corresponding values of the bolometric flux amplitude (the parameter f), whose empirical values are derived from multi-colour photometric observations. Seismic model that meets these conditions is of low mass, $M = 1.05M_{\odot}$, has moderately effective convection in the outer layers, described by the mixing length parameter $\alpha_{\text{MLT}} \approx 0.7$, and the microturbulent velocity in the atmospheres of about $\xi_t \approx 8 \text{ km s}^{-1}$. Such seismic studies of stars like SX Phe are very important for deriving constraints on outer-layer convection, because the object is on the border between very effective and ineffective convection.

Keywords: Stars: evolution, Stars: atmospheres, Asteroseismology, Convection

1 Introduction

SX Phoenicis (HD 223065) is A3-type star of Population II discovered to be variable almost seven decades ago by Eggen (1952a,b). This is a prototype for the whole class of high-amplitude and usually metal-poor pulsators located inside the δ Scuti instability region. SX Phe was a target of several studies based on photometric and spectroscopic observations. The analysis of photometric data revealed two frequencies and their combinations (e.g., by Coates et al. 1979; Rolland et al. 1991; Garrido & Rodriguez 1996). The frequency ratio indicated that SX Phe pulsates in the two radial modes; fundamental and first overtone ones. These two periodicities are present also in the radial velocity variations (Kim et al. 1993). Interestingly, the recent analysis of the high-precision data from the TESS satellite confirmed the old results that only these two frequencies dominate the light curve of SX Phe (Antoci et al. 2019). The values of frequencies extracted from the TESS light curve are $\nu_1 = 18.193565(6) \text{ d}^{-1}$ and $\nu_2 = 23.37928(2) \text{ d}^{-1}$.

There is also some evidence that both pulsation periods change in a timescale of decades (Landes et al. 2007). Moreover, for the dominant pulsational period the effective temperature varies in a huge range from 7210 to 8120 K and the surface gravity from 3.63 to 4.23. The corresponding mean values are 7640 K and 3.89 (Kim et al. 1993). The recent determination of the effective temperature from spectroscopy amounts to $T_{\text{eff}} = 7500 \pm 150 \text{ K}$ and the luminosity derived from the Gaia DR2 data is $\log L/L_{\odot} = 0.844 \pm 0.009$ (Antoci et al. 2019). The metallicity of SX Phe is low and amounts to about $[\text{Fe}/\text{H}] = -1.4$ (McNamara 1997).

SX Phe has been also a subject of seismic modelling. In the first attempts, Dziembowski & Kozłowski (1974) indicated a small mass of about $0.2 M_{\odot}$. The modeling with the recomputed OPAL opacity data (Iglesias et al. 1992) by Petersen & Christensen-Dalsgaard (1996) showed that the period ratio of the two radial modes is best reproduced by the model with parameters: a mass $M = 1.0M_{\odot}$, metallicity $Z = 0.001$, initial hydrogen abundance $X_0 = 0.70$ and age 4.07 Gyr.

The aim of this paper is to repeat seismic modelling of SX Phe and extend it by employing the bolometric flux amplitude (the parameter f) which is very sensitive to physical conditions in subphotospheric layers.

2 Models fitting the two radial-mode frequencies

Using the Warsaw-New Jersey evolutionary code (e.g., Pamyatnykh et al. 1998; Pamyatnykh 1999) and non-adiabatic pulsational code (Dziembowski 1977), we computed models of SX Phe with the aim to fit ν_1 and ν_2 as

¹ Astronomical Institute, Wrocław University, ul. Kopernika 11, 51-622 Wrocław, Poland

² Nicolaus Copernicus Astronomical Center, Polish Academy of Sciences, Bartycka 18, 00-716 Warsaw, Poland

the radial fundamental and first overtone modes, respectively. Part of these results have been already published in Antoci et al. (2019).

We searched the whole range of the effective temperature derived for SX Phe in the literature, i.e., from about 7000 to 9000 K. The OPAL opacity tables and the solar element mixture of Asplund et al. (2009) were adopted. We considered different values of the metallicity, Z , and initial hydrogen abundance, X_0 , from the range (0.0005, 0.003) and (0.66, 0.74), respectively. Different values of the mixing length parameter were investigated $\alpha_{\text{MLT}} \in (0, 2.5)$ and overshooting from the convective core was not included. At such low metallicity and effective temperatures $T_{\text{eff}} > 7000$ K, the effect of α_{MLT} on mode frequencies is negligible but it strongly affects the values of the parameter f , discussed in the next section.

We started from re-calculation of the seismic model of Petersen & Christensen-Dalsgaard (1996), whose constraints were: a mass $M = 1.0 M_{\odot}$, metallicity $Z = 0.001$ and initial hydrogen $X_0 = 0.70$. Our best seismic model for these parameters has the effective temperature $\log T_{\text{eff}} = 3.88427$, luminosity $\log L/L_{\odot} = 0.811$ and the frequency ratio of the radial fundamental mode to the first overtone amounts to $\nu_1/\nu_2 = 0.780014$. The observed counterpart is 0.778192. The theoretical values of the frequencies are $\nu_1 = 18.193565 \text{ d}^{-1}$ (the model was interpolated on this frequency) and $\nu_2 = 23.324654 \text{ d}^{-1}$. Both radial modes are in this model excited. However, as one can see, the difference between the theoretical and observed value of ν_1/ν_2 is significant and amounts to 0.001822. The age of the model is about 3.93 Gyr, which is slightly less than for the model of Petersen & Christensen-Dalsgaard (1996) who obtained about 4.07 Gyr. This difference may result from different versions of the OPAL opacity tables as well as from the usage of various evolutionary and pulsational codes.

In the next step, we changed the value of the metallicity and found the model for $Z = 0.0014$ with a mass $M = 1.15 M_{\odot}$ which fits better the observed frequency ratio. The model has the following parameters: $\log T_{\text{eff}} = 3.91770$, luminosity $\log L/L_{\odot} = 0.984$. The frequency ratio is $\nu_1/\nu_2 = 0.778412$ and the individual frequencies are $\nu_1 = 18.193565 \text{ d}^{-1}$ and $\nu_2 = 23.372681 \text{ d}^{-1}$. The model is much younger than the previous one with the age of about 2.45 Gyr. The fundamental and first overtone radial modes are in this model both stable. Increasing slightly metallicity, to $Z = 0.002$ allowed to find the model with a mass $M = 1.05 M_{\odot}$, $\log T_{\text{eff}} = 3.87152$, $\log L/L_{\odot} = 0.770$ and with the frequency ratio of the two considered radial modes $\nu_1/\nu_2 = 0.778073$. This model is about 3.42 Gyr old and has both radial fundamental and first overtone modes excited. As one can see the models with higher metallicity abundance give much better fit of frequencies but still there is a room for an improvement. Therefore, in the next step we changed also the initial abundance of hydrogen.

We have managed to find the model that reproduces the observed frequency ratio up to the fifth decimal place which we adopt as the numerical accuracy. The model has the following parameters: $M = 1.05 M_{\odot}$, $X_0 = 0.67$, $Z = 0.002$, $\log T_{\text{eff}} = 3.88979$, $\log L/L_{\odot} = 0.844$ and it was interpolated on the dominant frequency $\nu_1 = 18.193565 \text{ d}^{-1}$. The value of the second frequency is $\nu_2 = 23.379761 \text{ d}^{-1}$ which differs by 0.00048 d^{-1} from the observed value. Thus, the theoretical value of the frequency ratio is 0.778176 compared to the observed value 0.778192. Taking into account the numerical accuracy, which is not better than five decimal places, we can conclude that this model perfectly reproduces the observed frequencies of the two radial modes of SX Phe. Moreover, this model predicts instability (excitation) of both radial fundamental and first overtone modes. The age of the model amounts to about 2.85 Gyr. Another advantage of this model is the fact that it has the luminosity which agrees with the value derived from the Gaia DR2 data.

The lower hydrogen abundance (higher helium abundance) is also quite probable because, as other SX Phoenicis variables, SX Phe itself can be a blue straggler. Such objects are presumably formed by the merger of two stars or by interactions in a binary system and, as a consequence, they may have enhanced helium abundance (e.g., McNamara 2011; Nemeč et al. 2017).

In the left panel of Fig. 1, we show the evolution of the frequency ratio as a function of the dominant frequency (the Petersen diagram) for the four best seismic models we have found. The observed value is marked as a solid square. In the right panel of Fig. 1, we depicted the corresponding evolutionary tracks on the HR diagram and the position of SX Phe.

3 Complex seismic modeling

In the next step we extended our seismic analysis by including the parameter which describes the relative amplitude of the radiative flux perturbations at the photosphere level. This amplitude is the so-called nonadiabatic parameter f and it is defined as

$$\frac{\delta \mathcal{F}_{\text{bol}}}{\mathcal{F}_{\text{bol}}} = \text{Re}\{\varepsilon f Y_{\ell}^m(\theta, \varphi) e^{-i\omega t}\}, \quad (3.1)$$




Fig. 1. Left: The Petersen diagram for the four best seismic models suitable for SX Phoenicis. The solid square marks the observed values. The theoretical values of ν_1/ν_2 are given in the legend along with the mass and chemical composition (X, Z). **Right:** The corresponding evolutionary tracks of the seismic models with the observed position of SX Phe.

where \mathcal{F}_{bol} is the bolometric flux, ε is the mode intrinsic amplitude, Y_{ℓ}^m is the spherical harmonic with the degree ℓ and the azimuthal order m . Other quantities have their usual meaning. The value of f is complex and it is associated with a given mode. The theoretical values of f are very sensitive to the subphotospheric layers where the driving of pulsation occurs. In the case of B-type pulsators, these values depend mainly on the adopted opacity data and their modifications around the Z-bump (Daszyńska-Daszkiewicz et al. 2005a). In case of cooler pulsators, like δ Scuti or SX Phoenicis stars, the values of f are affected by convection in the envelope (Daszyńska-Daszkiewicz et al. 2003).

Therefore, in this paper we aim at obtaining some constraints on the efficiency of convective transport in the outer layers of SX Phe. The semi-empirical values of f can be derived from multi-colour light variations and radial velocity measurements (e.g., Daszyńska-Daszkiewicz et al. 2003, 2005a). To this end, we used the Strömgen amplitudes and phases for the two radial mode frequencies as determined by Rolland et al. (1991). The atmospheric flux derivatives over the effective temperature and gravity as well as limb darkening were derived from the Vienna model atmospheres (NEMO, see, e.g., Nendwich et al. 2004).

In Fig. 2 we show a comparison of the theoretical and empirical values of f for our best seismic model with the parameters: $M = 1.05 M_{\odot}$, $\log T_{\text{eff}} = 3.889793$, $\log L/L_{\odot} = 0.84375$ and $X_0 = 0.67$ and $Z = 0.002$. The theoretical values were computed for different values of the mixing length parameter α_{MLT} whereas the empirical counterparts for different values of the microturbulent velocity ξ_t . In the left panel, we plotted the results for the dominant frequency ν_1 (the radial fundamental mode) and in the right panel for the frequency ν_2 (the first overtone radial mode).

As one can see, in the case of the frequency ν_1 there is a pretty good agreement for the MLT parameter of $\alpha_{\text{MLT}} \approx 0.7$ and the microturbulent velocity of $\xi_t \approx 8 \text{ km s}^{-1}$. The results for the second mode frequency ν_2 are less unambiguous. While the imaginary part of f indicates lower values of the MLT parameter ($\alpha_{\text{MLT}} < 1.5$), the real part of f does not agree with any theoretical value. It can result from much smaller light amplitude of ν_2 which is determined with much lower accuracy. For example, the amplitude in the Strömgen v filter is almost three times lower for the frequency ν_2 than for the frequency ν_1 .

4 Conclusions

We made seismic analysis of the prototype SX Phoenicis. We started from the fitting of the two radial mode frequencies and constrained the mass, luminosity and chemical composition. Our best seismic model, that reproduces these two frequencies has the parameters: $M = 1.05 M_{\odot}$, $\log T_{\text{eff}} = 3.889793$, $\log L/L_{\odot} = 0.84375$ and the chemical composition: $X_0 = 0.67$ and $Z = 0.002$. The effective temperature is within the allowed observational range and luminosity perfectly agrees with the determination from the Gaia DR2 data. The age of other seismic models found by us is in the range from 2.5 to 3.9 Gyr. Further studies are needed to estimate




Fig. 2. A comparison of the theoretical and empirical values of f on the complex plane for the radial fundamental mode (the left panel) and for the first overtone mode (the right panel). The theoretical values were computed for our best seismic model (see text for details) considering different values of the mixing length parameter α_{MLT} . The (semi)empirical counterparts were determined from the Strömgen photometry and the Vienna model atmospheres assuming different values of the microturbulent velocity ξ_t .

the age more accurately, because it would give a clue to the star's evolutionary past and its origin.

In the next step, we tried to reproduce also the bolometric flux amplitude (the parameter f) corresponding to each mode. The aim was to get further constraints, e.g., on convection or atmospheric conditions. The (semi)empirical values of f were derived from the Strömgen amplitudes and phases adopting the Vienna model atmospheres. We found a very strong effect of the microturbulent velocity, ξ_t , on the empirical values of f which was already announced by Daszyńska-Daszkiewicz et al. (2005b) and Daszyńska-Daszkiewicz (2007). In the case of the fundamental radial mode, the empirical and theoretical values of f agree if the MLT parameter is about $\alpha_{\text{MLT}} = 0.7$ and the microturbulent velocities in the atmospheres is about $\xi_t = 8 \text{ km s}^{-1}$. It would mean that the efficiency of convective transport in the outer layers of SX Phe is rather moderate. In the case of the first overtone mode the agreement is poor and further studies are needed. It can result from some interaction between the two modes or a need of additional modification in pulsational and/or atmospheric models. Another reason could be much smaller photometric amplitudes determined with much lower accuracy.

We are going to extend these studies by re-computations of atmospheric models for a higher abundance of helium and higher microturbulent velocities. Moreover, the effect of modification of the mean opacities will be examined.

Definitely, new simultaneous multi-colour photometric and spectroscopic time series observations would lead to more plausible seismic constraints on the parameters of the model and theory.

The work was financially supported by the Polish NCN grant 2018/29/B/ST9/02803.

References

- Antoci, V., Cunha, M. S., Bowman, D. M., & et al. 2019, MNRAS, 490, 4040
 Asplund, M., Grevesse, N., Sauval, A. J., & Scott, P. 2009, Annu.Rev.Astron.Astrophys., 47, 481
 Coates, D. W., Dale, M., Halprin, L., & et al. 1979, MNRAS, 187, 83
 Daszyńska-Daszkiewicz, J. 2007, Comm. in Asteroseismology, 150, 32
 Daszyńska-Daszkiewicz, J., Dziembowski, W. A., & Pamyatnykh, A. A. 2003, A&A, 407, 999
 Daszyńska-Daszkiewicz, J., Dziembowski, W. A., & Pamyatnykh, A. A. 2005a, A&A, 441, 641
 Daszyńska-Daszkiewicz, J., Dziembowski, W. A., Pamyatnykh, A. A., & et al. 2005b, A&A, 438, 653
 Dziembowski, W. & Kozłowski, M. 1974, Acta Astr., 24, 245
 Dziembowski, W. A. 1977, Acta Astr., 27, 95
 Eggen, O. J. 1952a, PASP, 64, 31
 Eggen, O. J. 1952b, PASP, 64, 305

- Garrido, R. & Rodriguez, E. 1996, MNRAS, 281, 696
- Iglesias, C. A., Rogers, F. J., & Wilson, B. G. 1992, ApJ, 397, 717
- Kim, C., McNamara, D. H., & Christensen, C. G. 1993, AJ, 106, 2493
- Landes, H., Bamberg, K. R., Coates, D. W., & et al. 2007, PASAustral., 24, 41
- McNamara, D. H. 1997, PASP, 109, 1221
- McNamara, D. H. 2011, AJ, 142, 110
- Nemec, J. M., Balona, L. A., Murphy, S. J., & et al. 2017, MNRAS, 466, 1290
- Nendwich, J., Heiter, U., Kupka, F., & et al. 2004, Comm. in Asteroseismology, 144, 43
- Pamyatnykh, A. A. 1999, Acta Astr., 49, 119
- Pamyatnykh, A. A., Dziembowski, W. A., Handler, G., & et al. 1998, A&A, 333, 141
- Petersen, J. O. & Christensen-Dalsgaard, J. 1996, A&A, 312, 463
- Rolland, A., Rodriguez, E., Lopez de Coca, P., & et al. 1991, A&A Suppl., 91, 347