DISPERSIVE DECAY ESTIMATES FOR DIRAC EQUATIONS WITH A DOMAIN WALL

JOSEPH KRAISLER, AMIR SAGIV, AND MICHAEL I. WEINSTEIN

ABSTRACT. Dispersive time-decay estimates are proved for a one-parameter family of one-dimensional Dirac Hamiltonians with dislocations; these are operators which interpolate between two phase-shifted massive Dirac Hamiltonians at $x = +\infty$ and $x = -\infty$. This family of Hamiltonians arises in the theory of topologically protected states of one-dimensional quantum materials. For certain values of the phase-shift parameter, τ , the Dirac Hamiltonian has a *threshold resonance* at the endpoint of its essential spectrum. Such resonances are known to influence the time-decay rate. Our main result explicitly displays the transition in time-decay rate as τ varies between resonant and non-resonant values. Our results appear to be the first dispersive time-decay estimates for Dirac Hamiltonians which are not a relatively compact perturbation of a free Dirac operator.

1. INTRODUCTION

We study the time-dynamics for a family of Dirac Hamiltonians, $\{\mathcal{D}(\tau)\}_{\tau \in S^1}$, which arise in the theory of topologically protected states in one-dimensional asymptotically periodic quantum systems with a dislocation defect (domain wall). The parameter $\tau \in S^1 = \mathbb{R}/2\pi\mathbb{Z}$ specifies the size of the dislocation. We now introduce the model and in Section 1.1.1 below, we discuss a context in which this class of models plays a central role.

Consider the following initial-value problem, describing the evolution of complex vector amplitude, $\alpha(t, x) : \mathbb{R}_t \times \mathbb{R}_x \to \mathbb{C}^2$:

(1.1a)
$$i\partial_t \alpha = \mathscr{D}(\tau)\alpha, \qquad \alpha(0, x) = \alpha_0(x) \in L^2(\mathbb{R}; \mathbb{C}^2).$$

For each $\tau \in \mathbb{R}/2\pi\mathbb{Z}$, the Dirac Hamiltonian, $\mathscr{D}(\tau)$, is given by:

(1.1b)
$$\mathscr{D}(\tau) \equiv i\sigma_3\partial_x + \sigma_1 \mathbb{1}_{(-\infty,0)}(x) + \sigma_\star(\tau)\mathbb{1}_{[0,\infty)}(x) .$$

Here, σ_j denote standard Pauli matrices (see (1.8)),

$$\sigma_{\star}(\tau) \equiv \begin{pmatrix} 0 & e^{-i\tau} \\ e^{i\tau} & 0 \end{pmatrix} \ ,$$

and $\mathbb{1}_{S}(x)$ denote the indicator function of a set $S \subseteq \mathbb{R}$. The curve $\tau \in S^{1} \mapsto \mathscr{D}(\tau)$ sweeps out a periodic family of self-adjoint operators on $L^{2}(\mathbb{R}; \mathbb{C}^{2})$ with common domain $H^{1}(\mathbb{R}; \mathbb{C}^{2})$.

The goal of this paper is to provide a detailed understanding of the time-dynamics (1.1) as the dislocation parameter τ varies; in particular the dispersive decay of solutions, as $t \to \infty$, for initial data projected onto the continuous spectral part of $\mathcal{P}(\tau)$.

Note that

$$\mathcal{D}(\tau) = \begin{cases} \mathcal{D}_+(\tau) \equiv i\sigma_3\partial_x + \sigma_\star(\tau) & \text{for } x > 0\\ \mathcal{D}_-(\tau) \equiv i\sigma_3\partial_x + \sigma_1 & \text{for } x < 0. \end{cases}$$

If $\tau = 0$ (or $\tau = 2\pi$), $\mathcal{D}_+(0) = \mathcal{D}_-(0)$, and $\mathcal{D}(0)$ is a constant coefficient (i.e., translation invariant) massive Dirac Hamiltonian. In contrast, for all $\tau \notin \{0, 2\pi\}$, the operator $\mathcal{D}_+(\tau)$ is non-trivially "phase-shifted" from $\mathcal{D}_-(\tau)$. Such a defect is called a *domain wall* or a *dislocation*.

Since $\mathcal{D}(0) = \mathcal{D}(2\pi)$ commutes with spatial translations, its spectrum consists entirely of essential (continuous) spectrum, with a gap consisting of all energies in (-1, +1). For $\tau \neq 0, 2\pi$, however, $\mathcal{D}(\tau)$ has a defect-mode: an eigenvalue in this spectral gap, spatially localized around x = 0. The spectral properties of $\mathcal{D}(\tau)$ are well-known (see [9, 10, 19]) and are summarized in the following (also a consequence of Proposition 3.6 below):

Theorem 1.1. (1) For all $\tau \in [0, 2\pi]$, the essential spectrum of $\mathcal{D}(\tau)$ is given by:

$$\sigma_{\rm ess}(\mathcal{D}(\tau)) = (-\infty, -1] \cup [1, \infty).$$

(2) For $\tau \in [0, 2\pi]$,

$$\sigma(\mathcal{B}(\tau)) = \begin{cases} (-\infty, -1] \cup \{\omega_{\tau}\} \cup [1, \infty), \quad \tau \in (0, 2\pi) \\ (-\infty, -1] \cup [1, \infty), \quad \tau = 0, 2\pi \end{cases}$$

Here, ω_{τ} is an eigenvalue of multiplicity one within the spectral gap (the gap within the essential spectrum) and is given by the expression:

$$\omega_{\tau} = \cos(\tau/2) \, .$$

with corresponding one-dimensional eigenspace spanned by

(1.2)
$$\psi_{\tau}(x) = \frac{1}{2} \sqrt{\sin\left(\frac{\tau}{2}\right)} e^{-\sin\left(\frac{\tau}{2}\right)|x|} \begin{pmatrix} 1\\ -e^{i\frac{\tau}{2}} \end{pmatrix}, \quad \|\psi_{\tau}\|_{L^{2}} = 1.$$

That ω_{τ} traverses the spectral gap as τ varies from 0 to 2π has an underlying topological explanation, which we discuss in Section 1.1.1.

For $\tau = 0, 2\pi$, the Hamiltonian $\mathcal{D}(\tau)$ is a translation invariant (constant coefficient) differential operator. Each component of α satisfies a Klein-Gordon equation: $\partial_t^2 U(x,t) = (\partial_x^2 - 1)U(x,t)$. Hence, for $\tau = 0, 2\pi$, all sufficiently smooth and spatially localized initial conditions disperse to zero (spread and decay) as time advances under the evolution (1.1) [3].

In contrast, by Theorem 1.1, if $\tau \in (0, 2\pi)$, one expects dispersive time-decay only after projecting onto the continuous spectral subspace of $\mathcal{D}(\tau)$. We denote this projection by $P_{ac}(\mathcal{D}(\tau))$. By Theorem 1.1

$$P_{ac}(\mathcal{D}(\tau)) = I - \langle \psi_{\tau}, \cdot \rangle \ \psi_{\tau}.$$

Our main result is the following time-decay estimate, with a τ - dependent rate of decay.

Theorem 1.2. Let $\tau \in [0, 2\pi]$ and $\mathcal{D}(\tau)$ be as defined in (1.1b). For any $\epsilon > 0$, there exists $C_{\epsilon} > 0$, which is independent of τ , such that for all we have

(1.3)
$$\|\langle x \rangle^{-2} e^{-i \mathcal{D}(\tau) t} P_{ac}(\mathcal{D}(\tau)) \langle \mathcal{D}(\tau) \rangle^{-3/2 - \epsilon} \langle x \rangle^{-2} \|_{L^1 \to L^\infty} \le C_\epsilon \frac{1}{\langle t \rangle^{1/2}} \frac{1}{1 + \sin^2(\tau/2)t}.$$

From the bound (1.3) we note the non-uniformity of the time scale on which $t^{-\frac{3}{2}}$ occurs for τ or $2\pi - \tau \ll 1 \pmod{2\pi}$; indeed the $t^{-3/2}$ rate of decay only becomes visible on a time scale $t \gg \tau^{-2}$. Furthermore, while there is discontinuity in the $t \to \infty$ decay rate at $\tau = 0$, for a fixed time (1.3) shows that the decay rate is continuous with respect to τ even at $\tau = 0$.

A dispersive decay bound which displays a similar transition was obtained for the time-dependent Schrödinger equation with highly oscillatory potential in [11].

$$(1.4) \qquad \|\langle x\rangle^{-2}e^{-i\mathcal{P}(\tau)t}P_{ac}(\mathcal{P}(\tau))\alpha_0\|_{L^{\infty}(\mathbb{R})} \leq C_{\epsilon}\frac{1}{\langle t\rangle^{1/2}}\frac{1}{1+\sin^2(\tau/2)t}\|\langle x\rangle^2\langle \mathcal{P}(\tau)\rangle^{3/2+\epsilon}\alpha_0\|_{L^1(\mathbb{R})}.$$

This time-decay bound requires both smoothness and spatial localization of the initial condition. The smoothness required on the right hand side of eq. (1.4) arises from high-energy considerations; for the high-energy part of the data, the dynamics are comparable to those of the Klein-Gordon equation, whose dispersion relation $\omega(k) = \sqrt{1+k^2}$ has a dispersion rate, $\omega''(k)$, which tends to zero as $k \to \infty$.¹ Smoothing provides an effective truncation of the spectral support of the data α_0 , leading to an effective lower bound on the dispersion rate.

Further, the localization in x arises from seeking a upper bound for the time-decay rate which holds uniformly in $\tau \in [0, 2\pi]$, i.e., the constant C_{ϵ} in (1.3) should be independent of τ . This boils down to an analysis of the behavior of the resolvent of $\mathcal{D}(\tau)$ for energies near the endpoints (thresholds) of the essential spectrum, for which we require spatial weights which are at least quadratic in x. It is an open problem whether the spatial weights can be reduced to $\langle x \rangle^{-1}$.

Remark 1.4 (Threshold resonance phenomena). While our initial consideration of $\mathcal{D}(\tau)$ was motivated by its role in applications to periodic structures with dislocations (see Section 1.1 below), Theorem 1.2 also turns out to be a useful and simple paradigm for illustrating dispersive wave phenomena: The transition in decay rate in the estimate (1.3) reflects the appearance of a threshold resonance just as the eigenvalue ω_{τ} , when $\tau \equiv 0 \pmod{2\pi}$, reaches the endpoint of the essential spectrum. The threshold resonance phenomenon, associated with the emergence of point spectrum from (or disappearance into) the continuous spectrum, is well-known to impact the rate of local energy decay in wave equations. This is best known in the context of the Schrödinger operator with a spatially localized potential; see, for example, [25, 29, 30, 37]. The effect of threshold resonances for one-dimensional massive Dirac equations with a spatially localized potential is discussed in [17]. Our family of Dirac operators is, to the best our knowledge, the only example of Dirac operators where the transition in decay rate, due to a threshold resonance, is explicitly displayed. Such a transition in decay rates has displayed as well for a class of Schrödinger Hamiltonians with rapidly oscillatory potentials in [11].

Remark 1.5 (Relation to previous time-decay results for Dirac Hamiltonians). Time-decay estimates for massive Dirac operators for classes of potentials, which decay to zero as $x \to \pm \infty$, were obtained by Erdogan and Green one space dimension in [17]. For results in two and three space dimensions, see e.g., [7, 13, 14, 16, 15, 18, 31]. Pelinovsky and Stefanov derived estimates for the linearized evolution about standing waves of a one-dimensional semilinear Dirac equation in [34].

In this remark, we make two contrasts of the results and methods of this article with those in [17]. First, the operators we study, $\mathcal{D}(\tau)$, are not spatially localized perturbations of a constant coefficient operator. Second, our strategy of proof is different in several ways (e.g., in developing the scattering theory of $\mathcal{D}(\tau)$), see Section 2.

The following theorem states alternative variations of time-decay upper bounds, which hold for various relaxations of the assumptions on spatial localization of the initial conditions.

Theorem 1.6. (1) For any $\tau \in [0, 2\pi]$ and $\varepsilon > 0$, there exists a constant, $C_{\varepsilon}(\tau) > 0$, such that

(1.5)
$$\|e^{-i\mathcal{D}(\tau)t}P_{ac}(\mathcal{D}(\tau))\langle\mathcal{D}(\tau)\rangle^{-3/2-\varepsilon}\|_{L^1\to L^\infty} \le C_{\varepsilon}(\tau)\frac{1}{\langle t\rangle^{1/2}}.$$

¹For the Klein-Gordon equation k is simply the Fourier variable.

(2) Let $K \subset (0, 2\pi)$ be a compact set and fix $\varepsilon > 0$. Then there exists a constant $C_1(K) > 0$ such that for any $\tau \in K$ and t > 0 we have

(1.6)
$$\|\langle x\rangle^{-1}e^{-i\mathcal{D}(\tau)t}P_{ac}(\mathcal{D}(\tau))\langle \mathcal{D}(\tau)\rangle^{-3/2-\varepsilon}\langle x\rangle^{-1}\|_{L^1\to L^\infty} \le C_1(K)\frac{1}{\langle t\rangle^{3/2}}$$

and $C_1(K_n) \to \infty$ for every sequence of compact sets such that $K_n \to (0, 2\pi)$.

Remark 1.7 (Generalizations of Theorems 1.2 and 1.6). It is natural to consider extending Theorems 1.2 and 1.6 to the case where $\mathcal{D}(\tau)$ in (1.1b) is replaced by an operator which continuously interpolates between the asymptotic operators \mathcal{D}_{-} and \mathcal{D}_{+} . In this general setting we expect $\mathcal{D}(\tau)$ to have, for each fixed τ , a finite number of eigenvalues in the spectral gap and also possible threshold resonance energies at the edge of the essential spectrum. Analogous decay estimates to those presented in Theorems 1.2 and 1.6 would hold with (a) $P_{ac}(\mathcal{D}(\tau))$ defined as the projection orthogonal to all bound states corresponding to the energies in the spectral gap and (b) decay rates adjusted to transition from the faster $\langle t \rangle^{-\frac{3}{2}}$ rate, for typical values of τ , to the slower $\langle t \rangle^{-\frac{1}{2}}$ rate arising from those values of τ for which a threshold resonance occurs. See the further discussion in Section 1.2 below.

Remark 1.8. In the resonant case $\tau = 0, 2\pi$, which is in fact the standard massive Dirac equation, a $t^{-3/2}$ decay rate holds true for a variation of (1.6). Essentially, if one subtracts a finite-rank operator from the evolution equation, one can counteract the effect of resonance and obtain a nonresonant type order of decay. For details, see [17, Theorem 1.2].

1.1. Motivation; topologically protected states and radiation damping for Floquet systems.

1.1.1. Topologically protected states in dislocated systems. Since the experimental observation of the integer quantum Hall effect [4, 28], and its subsequent explanation using the topological phases of matter [43], there has been a great deal of fundamental and applied interest in wave systems having properties which are "topologically protected", i.e., systems in which energy transport remains unchanged under continuous localized deformations, due to a topological invariant; see, for example, the reviews [24, 33, 47] and references therein.

The Dirac operators which this article studies, $\{\mathcal{D}(\tau)\}_{\tau \in \mathbb{R}/2\pi\mathbb{Z}}$, play a central role in a class of one-dimensional PDE models of physical media which exhibit topological properties. We briefly describe these models, and refer to [9, 10, 19, 20] for details. For a photonic realization of such systems, see [32]. Consider a Schrödinger operator, $H^{(0)}$, with a one-periodic potential, V(x). Discrete translation-invariance of $H^{(0)}$ implies a band structure; its spectrum is characterized by a sequence of eigenvalue dispersion curves: $k \in [0, 2\pi] \mapsto E(k)$ (Floquet-Bloch band functions). A scenario of great is interest is when two such curves meet at a pair (k_D, E_D) and form a linear crossing, a Dirac point [19]. In this case, E_D is a two-fold degenerate eigenvalue of $H^{(0)}$. The effective (or homogenized) Hamiltonian which describes a blowup of the band structure near such linear crossings is $H_{\text{eff}}^{(0)} = i\sigma_3\partial_X$, a massless Dirac Hamiltonian.

Next, we introduce a *dislocation*, a perturbing potential which is also asymptotically one-periodic but, as $x \to +\infty$ is phase-shifted by an amount $\tau/2\pi$ from its behavior as $x \to -\infty$. The parameter $\tau \in [0, 2\pi]$ measures the amount of dislocation. The corresponding Schrödinger operator, $H^{(\tau)}$, has a gap in its essential spectrum about energy E_D (where two bands of spectra of H^0 cross linearly). Typically, $H^{(\tau)}$ will have a finite number of eigenvalues, $E_i^{(\tau)}$, which "flow" within the spectral gap

4

as τ varies. Figure 1 displays several examples where eigenvalues traverse the gap as τ varies from 0 to 2π .

The spectral flow of the family of operators $\{H^{(\tau)}\}_{\tau \in S^1}$ is an integer which measures the net number of edge state curves which traverse the spectral gap. The spectral flow is a topological invariant; it does not change under continuous deformations of the Hamiltonian under which the gap remains open [44]. In [9], $H^{(\tau)}$ is embedded in a continuously varying family of Hamiltonians $s \in (0, 1] \mapsto H^{(\tau, s)}$, where $H^{(\tau, 1)} = H^{(\tau)}$. In the asymptotic regime, $s \ll 1$, the operators $H^{(\tau, s)}$ encode a small and adiabatic dislocation. By a multiple-scales analysis [10, 20, 19], the study of the spectral flow of $\{H^{(\tau, s)}\}_{\tau \in S^1}$ can be reduced to that of Dirac operators of the form:

(1.7)
$$\mathcal{D}(m\tau) \equiv i\sigma_3\partial_X + \sigma_1\mathbb{1}_{(-\infty,0)}(X) + \begin{pmatrix} 0 & e^{-im\tau} \\ e^{im\tau} & 0 \end{pmatrix} \mathbb{1}_{[0,\infty)}(X) \ .$$

The integer m denotes the winding number about zero of a complex-valued "effective mass" parameter which emerges from the multiple scale analysis. By Theorem 1.1, the spectral flow of the family $\{\mathcal{D}(m\tau)\}_{\tau\in\mathbb{R}/(2\pi/m)}$ is equal to 1 and hence the spectral flow of the family $\{\mathcal{D}(m\tau)\}_{\tau\in\mathbb{R}/(2\pi/m)}$ is equal to 1 and hence the spectral flow of the family $\{\mathcal{D}(m\tau)\}_{\tau\in\mathbb{R}/(2\pi/m)}$ is equal to 1 and hence the spectral flow of the family $\{\mathcal{D}(m\tau)\}_{\tau\in\mathbb{R}/(2\pi/m)}$ is equal to 1 and hence the spectral flow of the family $\{\mathcal{D}(m\tau)\}_{\tau\in\mathbb{R}/(2\pi/m)}$ is equal to m. We conclude that, provided that a gap remains open throughout the deformation, the net number of eigenvalue curves of $H^{(\tau)}$ which traverse the spectral gap is equal to m (see Figure 1). Each operator family $\{\mathcal{D}(m\tau)\}_{\tau\in S^1}$ is a canonical operator (or normal form) to which all operator families $\{H^{(\tau)}\}_{\tau\in S^1}$ in the same topological class can be deformed; see right panel of Figure 1.

Finally, Drouot proves in [9] that the spectral flow, m, can also be identified as the first Chern number, $C_1(\mathcal{E})$, associated with a vector bundle over $\mathbb{T}^2_{k,\tau}$, whose fibers are the k-pseudoperiodic eigenspaces of the periodic Schrödinger operator at infinity $H^{(\tau)}_+$ associated with all bands below the energy E_D , which lies in a gap. This equality of bulk and edge indices is a variant of the bulk-edge correspondence principle.

1.1.2. Radiative decay and metastability of edge modes. In [23], motivated by modeling and experiments in photonic waveguides [2, 27], parametric forcing perturbations of the dynamics $i\partial_t \alpha = \mathcal{D}(\pi)\alpha$ is studied (i.e., linear and weakly time-dependent perturbations of $\mathcal{D}(\pi)$). In particular, it is shown via an asymptotic expansion of solutions and detailed comparison with computer simulations, that the zero energy eigenstate of $\mathcal{D}(\pi)$ is only meta-stable under sufficiently rapid forcing; we demonstrate it decays exponentially in t for $0 < t \leq \beta^{-2}$, for $0 < \beta \ll 1$, the amplitude of the forcing.²

To establish a systematic multiple-scale expansion of the solution, we required a weak form of the local energy decay estimate of Theorem 1.2. The approach we present here implies such an estimate by a more direct and generalizable method than in [23]. The latter involved first relating the problem to a Schrödinger evolution, by squaring the Dirac Hamiltonian, and then applying results on the boundedness of wave operators associated with Schrödinger equation [45, 49].

1.2. Open questions and future directions.

(1) We believe that the time-decay estimates in Theorems 1.2 and 1.6 can be extended to the case where $\mathcal{P}(\tau)$ in (1.1b) is replaced by a Dirac operator with a domain wall, which continuously interpolates between the asymptotic operators \mathcal{P}_{-} and \mathcal{P}_{+} (as opposed to the discontinuity present in the definition of $\mathcal{P}(\tau)$, (1.1b)); see also the discussion in Remark 1.7. One approach to such a result would be to modify the present analysis by working with

²It is conjectured that for times $t \gg \beta^{-2}$, the decay is at an algebraic rate. See analogous results concerning the Schrödinger equation in [39].



FIGURE 1. Spectral flow illustrated (see Section 1.1.1). All figures show energy (spectrum) as a function of the dislocation parameter τ : continuous spectrum (red stripes) and curves of point spectra (solid blue). Left: $\mathcal{D}(m\tau)$ with m = 1, see (1.7). Center: same, with m = 2. Right: a continuous deformation of the m = 1 case in which τ is replaced by a curve $\theta(\tau)$, which does not close the gap. The curve ω_{τ} , the energy of the eigenfunction of $\mathcal{D}(\theta(\tau))$, is non-monotonic and the spectral flow is 1.

the distorted plane waves derivable from appropriate linear integral equations; in the present work we used explicit expressions for distorted plane waves, obtainable because $\mathcal{D}(\tau)$ has piecewise constant coefficients. Alternatively, if $\tilde{\mathcal{D}}(\tau)$ is a Dirac operator with continuously varying coefficients, and such that $\tilde{\mathcal{D}}(\tau) \to \mathcal{D}(\tau)$ sufficiently rapidly as $x \to \pm \infty$, then $\tilde{\mathcal{D}}(\tau)$ is a relatively compact perturbation of $\mathcal{D}(\tau)$, and one can seek to derive time-decay estimates for $\exp(-it\tilde{\mathcal{D}}(\tau))P_{\rm ac}(\tilde{\mathcal{D}}(\tau))$ by a perturbation argument.

- (2) A fully rigorous justification of the expansion constructed in [23], which describes the radiation damping of the localized mode (see Section 1.1.2), is an interesting open problem. We believe that such a proof would require time-decay estimates for the *full Floquet (time-periodic) Hamiltonian* with a domain wall, where the time-dependent forcing is not spatially localized. Such estimates present a challenging open analytical problem, and do not follow from known results on decay estimates for time-periodic Hamiltonians, which impose spatial localization conditions on the forcing [1, 21, 40].
- (3) Section 1.1.1 outlines a context in which Dirac operators with domain wall / dislocation defects arise and their role in determining topological properties. In physical systems, such as in optics (e.g. [32]), nonlinear corrections to the underlying linear wave equations are needed to model effects at high intensities. For the systems described in Section 1.1.1, the corresponding effective Dirac equations have a nonlinearity, analogous to the Kerr (cubic) nonlinearity of the nonlinear Schrödinger equation. A natural agenda of analytical questions for the semi-linear Dirac models with domain walls ranges from low energy scattering for systems without a domain wall defect, nonlinear scattering and asymptotic stability theory for nonlinear bound states which bifurcate from the linear spectrum of Dirac operators with

domain walls, to radiation damping and metastability of states. This agenda connects naturally with the extensive literature on these topics for nonlinear Schrödinger and nonlinear Klein Gordon type equations; ; see, for example, [46, 38]. Concerning the subtle effects on large time weakly nonlinear dynamics, which are caused by threshold resonances, see for example [5, 36].

1.3. Notation, conventions, remarks.

(1) Pauli-type matrices:

(1.8)
$$\sigma_1 = \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}, \qquad \sigma_3 = \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}, \qquad \sigma_\star(\tau) = \begin{pmatrix} 0 & e^{-i\tau} \\ e^{i\tau} & 0 \end{pmatrix}.$$

(2) For an operator $T: L^2(\mathbb{R}; \mathbb{C}^2) \to L^2(\mathbb{R}; \mathbb{C}^2)$, which arises from an integral kernel, we denote its kernel by T(x, y), i.e., for $f \in L^2(\mathbb{R}; \mathbb{C}^2)$,

$$(Tf)(x) = \int_{\mathbb{R}} T(x,y)f(y) \, dy$$
.

(3) Let a, b, c > 0 be such that $a \leq b$ and $a \leq c$. Then,

(1.9)
$$a \le \frac{2}{b^{-1} + c^{-1}} ,$$

- (4) $\langle x \rangle = \sqrt{1+x^2}$
- (5) $\lambda(\omega) = \sqrt{1-\omega^2}$ for all $\omega \in (-1,1)$, and as the unique analytic continuation of that root for all $\omega \in U$ where $U = \mathbb{C} \setminus \{(-\infty, -1] \cup [1, \infty)\}$ (see Lemma 3.1).
- (6) The resolvent operator is defined by $\Re(\omega; \tau) \equiv (\mathcal{D}(\tau) \omega)^{-1}$ for every $\omega \in \mathbb{C}$, and for $k \in \mathbb{R}$ we define

$$\mathcal{R}_{\pm}(k,\tau) \equiv \Re(\sqrt{1+k^2\pm i0},\tau)\,.$$

1.4. Structure of the article. We first provide a rough outline of the proof in Section 2. We then develop the spectral theory of the Dirac operators $\mathcal{D}(\tau)$: the resolvent kernel is constructed in closed form in Section 3; In Section 4, we the limits of the resolvent kernel as the spectral parameter approaches the essential spectrum; In Section 5, we solve the associated scattering problem and relate it to the spectral measure. Finally, the main result (Theorem 1.2), as well as its variation (Theorem 1.6), are proved in Section 6.

Acknowledgements. This research was supported in part by National Science Foundation grants DMS-1908657 and DMS-1937254 (MIW), Simons Foundation Math + X Investigator Award #376319 (MIW, JK, AS), the Binational Science Foundation Research Grant #2022254 (MIW, AS), and the AMS-Simons Travel Grant (AS). This research was initiated while JK was a postdoctoral fellow and AS was an associate research Scientist, both in the Department of Applied Physics and Applied Mathematics at Columbia University.

2. Strategy and overview of the proof of Theorem 1.2

For any $\alpha_0 \in L^2(\mathbb{R}, \mathbb{C}) \cap L^1(\mathbb{R}, \mathbb{C})$ data, by the $L^2(\mathbb{R})$ -functional calculus and Stone's formula for the spectral measure associated with $\mathcal{D}(\tau)$ [22, 30, 42], we have that:

$$e^{-i\mathcal{B}(\tau)t}P_{ac}(\mathcal{D}(\tau))\langle\mathcal{D}(\tau)\rangle^{-3/2-\varepsilon}\alpha_{0}$$

$$(2.1) \qquad \qquad = \frac{1}{2\pi i}\int_{\sigma_{ac}(\mathcal{D}(\tau))}e^{-i\omega t}\left[\Re(\omega+i0,\tau)-\Re(\omega-i0,\tau)\right]\alpha_{0}\langle\omega\rangle^{-3/2-\varepsilon}d\omega ,$$

where $\Re(\omega, \tau)$ is the resolvent

$$\Re(\omega,\tau) = (\mathscr{D}(\tau) - \omega)^{-1},$$

and where $\sigma_{\rm ac} = (-\infty, 1] \cup [1, \infty)$ is the continuous spectrum of $\mathcal{D}(\tau)$.

Remark 2.1. Since the continuous spectrum is made up of the two disjoint intervals, the integral (2.1) breaks up into two disjoint integration domains. Throughout the proof, we treat only the positive part of the spectrum; the arguments for the negative part follow analogously. In particular, treating $[1, \infty)$ independently allows us to use the change of variables (2.3), which simplifies the analysis considerably.

Alternatively, in Appendix B we establish the invariance of the equation $\mathcal{D}(\tau)\alpha = \omega \alpha$ under the transformation

$$(\omega, \tau, \alpha(x)) \mapsto (-\omega, -\tau \mod(2\pi), \sigma_3 S\alpha(-x))$$

Hence, proving decay-rates for data in $P_{ac}\left(\mathcal{D}(\tau)\mathbb{1}_{[1,\infty)}\right)$ for all $\tau \in [0, 2\pi)$ immediately leads to the same estimates for data in $P_{ac}\left(\mathcal{D}(\tau)\mathbb{1}_{(-\infty,-1]}\right)$.

Define

$$P(\mathcal{D}(\tau)) \equiv P_{ac} \left(\mathcal{D}(\tau) \mathbb{1}_{[1,\infty)} \right) \,,$$

to be the projection onto the positive part of the continuous spectrum. Then replacing P_{ac} in (2.1) with P yields

(2.2)
$$\alpha(x,t) = e^{-i\mathcal{B}(\tau)t} P(\mathcal{D}(\tau)) \langle \mathcal{D}(\tau) \rangle^{-3/2-\varepsilon} \alpha_0$$
$$= \frac{1}{2\pi i} \int_1^\infty e^{-i\omega t} \left[\Re(\omega + i0, \tau) - \Re(\omega - i0, \tau) \right] \alpha_0 \ \langle \omega \rangle^{-3/2-\varepsilon} \, d\omega \,.$$

Next we make the change of variables

(2.3)
$$\omega \equiv \sqrt{1+k^2}$$

and define

$$\mathcal{R}_{\pm}(k,\tau) \equiv \Re(\sqrt{1+k^2} \pm i0,\tau) = (\mathcal{D}(\tau) - \sqrt{1+k^2} \pm i0)^{-1}.$$

Then,

(2.4)
$$\alpha(x,t) = \frac{1}{2\pi i} \int_0^\infty dk \, \langle k \rangle^{-3/2-\varepsilon} \frac{k}{\sqrt{1+k^2}} e^{-i\sqrt{1+k^2}t} \left[\mathcal{R}_+(k,\tau) - \mathcal{R}_-(k,\tau) \right] \alpha_0 \,,$$

This last representation of the dynamics is the main object of our analysis.

Remark 2.2. When carefully substituting $\omega = \sqrt{1+k^2}$ into (2.2), one really gets

$$\langle \omega \rangle = \langle \sqrt{1+k^2} \rangle = (1+(\sqrt{1+k^2})^2)^{1/2} = (2+k^2)^{1/2} \,,$$

in (2.4), instead of $\langle k \rangle$. However, we allow this small abuse of notations in (2.4) since the analysis is unchanged by this difference.

The strategy for proving Theorem 1.2 from here on is as follows:

- In Section 3 we construct the resolvent kernel, $\Re_{\pm}(\omega, \tau)(x, y)$ for spectral parameter, ω , in the complement of $\sigma(\mathcal{D}(\tau))$, by the variation of parameters method for ODEs (Proposition 3.5).
- Results in Sections 3, 4, and 5 lead to explicit expression for the solution $\alpha(t, x)$ in (2.4) as an oscillatory integral.
- The oscillatory integral is divided into a high-energy and low-energy components (see (6.4)).
- The high energy component is bounded from above uniformly in τ using a stationary phase argument (Lemma 6.2).
- The low energy component is bounded from above in two steps: first a uniform $t^{-1/2}$ bound is proved (Proposition 6.6). Then a more careful analysis of the transmition coefficient leads to a $t^{-3/2}$ decay rate for non-resonant τ .

3. The resolvent operator $\Re(\omega; \tau)$

In this section we construct the resolvent, $\Re(\omega, \tau) = (\mathscr{D}(\tau) - \omega)^{-1}$, by providing an explicit, closed-form expression for its kernel. For any $\omega \in \mathbb{C}$, the resolvent operator is defined in such a way that if $\Re(\omega; \tau)f = \beta$ for $\beta, f \in L^2(\mathbb{R}; \mathbb{C}^2)$, then

(3.1)
$$(\mathcal{D}(\tau) - \omega) \beta = f.$$

Hence, the resolvent acting on f returns an L^2 solution to the inhomogeneous ODE forced by f. Our first goal is to solve this ODE for ω outside of the spectrum i.e., $\omega \in U$ where

(3.2)
$$U \equiv \mathbb{C} \setminus \{(-\infty, -1] \cup [1, \infty)\}.$$

To solve the ODE (3.1) in $L^2(\mathbb{R}; \mathbb{C}^2)$ (and hence to construct the resolvent kernel), we will use the method of variation of parameters: in Section 3.1 we first solve the corresponding homogeneous ODE $(\mathcal{D}(\tau) - \omega)\eta = 0$ to find asymptotically decaying solutions, known as the *Jost solution* and are denoted by $\eta_{\pm}(x; \omega, \tau)$. Afterwards, in Section 3.2, we use the Jost solutions to construct the kernel, see Proposition 3.5.

3.1. The homogeneous equation. We begin by considering general solutions of the homogeneous ordinary differential equation

 $(\not\!\!\!D(\tau)-\omega)\beta=0.$

This may be written as:

$$\partial_x \beta(x) = M(x; \omega, \tau) \beta(x)$$

where

(3.

(3.4)
$$M(x;\omega,\tau) \equiv \begin{cases} M_{+}(\omega,\tau), & x > 0, \\ M_{-}(\omega,\tau), & x < 0, \end{cases}$$

and

(3.5)
$$M_{+}(\omega,\tau) \equiv \begin{pmatrix} -i\omega & ie^{-i\tau} \\ -ie^{i\tau} & i\omega \end{pmatrix}, \qquad M_{-}(\omega,\tau) \equiv \begin{pmatrix} -i\omega & i \\ -i & i\omega \end{pmatrix}.$$

We observe that

det
$$(M_{\pm}(\omega, \tau)) = \omega^2 - 1$$
 and Tr $(M_{\pm}(\omega, \tau)) = 0$.

Hence, for a fixed ω the eigenvalues of M_+ and M_- are solutions, λ , of

$$\lambda^2 = 1 - \omega^2.$$

The following lemma ensures that, on an appropriate domain, U, we may choose a solution of (3.6), $\lambda(\omega)$, with strictly positive real part.

Lemma 3.1. There is a holomorphic function $\omega \mapsto \lambda(\omega)$, defined on the domain U (see (3.2)), such that for all $\omega \in U$

$$\lambda^2(\omega) = 1 - \omega^2 \text{ and } \operatorname{Re}(\lambda(\omega)) > 0.$$

Furthermore, $\overline{\lambda(\omega)} = \lambda(\overline{\omega})$ for all $\omega \in U$.

Proof. We claim that the desired mapping is $\omega \in U \mapsto \lambda(\omega) = \sqrt{1 - \omega^2} \in \mathbb{C}$ (we take the positive square root) analytically continued from the interval $-1 < \omega < 1$ to the domain U. We need only check that $\operatorname{Re}(\lambda(\omega))$ is strictly positive on U. If not, then by continuity of $\lambda(\omega)$ on U there is some $\omega_0 \in U$ where $\operatorname{Re}(\lambda(\omega_0)) = 0$. Then, $\lambda(\omega_0) = i\zeta$, for some $\zeta \in \mathbb{R}$. Squaring both sides of this equation yields $1 - \omega_0^2 = -\zeta^2$ or equivalently $\omega_0^2 = 1 + \zeta^2 > 1$, which contradicts that assumption that $\omega_0 \in U$.

For all $\omega \in U$, the matrices $M_{+}(\omega, \tau)$ and $M_{-}(\omega, \tau)$ (see (3.5)) both have eigenvalues $\pm \lambda(\omega)$. The eigenpairs of M_{\pm} are given by

(3.7a)
$$M_{-}(\omega, \tau)v_{-}^{(+)} = +\lambda(\omega) v_{-}^{(+)}, \qquad v_{-}^{(+)} = \begin{pmatrix} 1\\ \omega - i\lambda(\omega) \end{pmatrix}$$

(3.7b)
$$M_{-}(\omega, \tau)v_{-}^{(-)} = -\lambda(\omega) \ v_{-}^{(-)}, \qquad v_{-}^{(-)} = \begin{pmatrix} 1\\ \omega + i\lambda(\omega) \end{pmatrix}$$

(3.7c)
$$M_{+}(\omega,\tau)v_{+}^{(+)} = +\lambda(\omega) v_{+}^{(+)}, \qquad v_{+}^{(+)} = \begin{pmatrix} e^{-i\tau} \\ \omega - i\lambda(\omega) \end{pmatrix}$$

(3.7d)
$$M_{+}(\omega,\tau)v_{+}^{(-)} = -\lambda(\omega) v_{+}^{(-)}, \qquad v_{+}^{(-)} = \begin{pmatrix} e^{-i\tau} \\ \omega + i\lambda(\omega) \end{pmatrix}.$$

Any solution of (3.3), $\partial_x \beta = M\beta$, can be constructed by piecing together the states $e^{\pm \lambda(\omega)x} v_{-}^{(\pm)}$ for x < 0 and $e^{\pm \lambda(\omega)x} v_{+}^{(\pm)}$ for x > 0 across the equation's discontinuity at x = 0.

We seek a construction of the resolvent $(\mathcal{D}(\tau) - \omega)^{-1}$ which for all $\omega \in U$, is a bounded linear operator on $L^2(\mathbb{R}; \mathbb{C}^2)$. Our construction uses the method of variation of parameters [6], which produces the resolvent kernel by appropriate modulation of two linearly independent solutions, decaying at $x \to \pm \infty$, respectively. These are introduced in the following proposition.

Definition 3.2. For any $\tau \in [0, 2\pi]$ and $\omega \in U$ (see (3.2)), we define the solutions $\eta_{\pm}(x; k, \tau)$, to be the unique solutions of

$$(\mathscr{D}(\tau) - \omega)\eta_{\pm}(x;\omega,\tau) = 0, \quad -\infty < x < \infty,$$

10

with the asymptotic behavior

(3.8a)
$$\lim_{x \to +\infty} e^{\lambda(\omega)x} \eta_+(x;\omega,\tau) = v_+^{(-)}(\omega,\tau) \,,$$

(3.8b)
$$\lim_{x \to -\infty} e^{-\lambda(\omega)x} \eta_{-}(x;\omega,\tau) = v_{-}^{(+)}(\omega,\tau) \,,$$

where $\lambda(\omega)$ is given by Lemma 3.1. For $\omega \in U$, $\eta_+(x; \omega, \tau)$ is exponentially decaying as $x \to \infty$ and $\eta_-(x; \omega, \tau)$ is exponentially decaying as $x \to -\infty$.

Remark 3.3. In Proposition 3.4 we display explicit expressions for the homogeneous solutions, η_{\pm} , of $(\mathcal{D}(\tau) - \omega)\eta = 0$ for a piecewise constant domain wall. For the case of general domain walls, the existence of a unique solution satisfying (3.8a) (respectively (3.8b)) follows from a standard equivalent formulation as a fixed point problem for a Volterra (integral) operator of the type that arise in the construction of Jost solutions for Schrödinger operators [35].

The solutions $\eta_{\pm}(x; \omega, \tau)$, introduced in Definition 3.2, are displayed in the following proposition, whose proof is presented in Appendix A.

Proposition 3.4. Assume $\omega \in U = \mathbb{C} \setminus \{(-\infty, -1] \cup [1, \infty)\}$ so that $\operatorname{Re}(\lambda(\omega)) > 0$, and $\tau \in [0, 2\pi]$.

(1) The solutions $\eta_{\pm}(x;\omega,\tau)$ have the explicit expressions:

(3.9a)
$$\eta_{+}(x;\omega,\tau) = \begin{cases} v_{+}^{(-)} e^{-\lambda(\omega)x} & x > 0\\ A v_{-}^{(+)} e^{\lambda(\omega)x} + B v_{-}^{(-)} e^{-\lambda(\omega)x} & x < 0 \end{cases}$$

(3.9b)
$$\eta_{-}(x;\omega,\tau) = \begin{cases} C \ v_{+}^{(+)}e^{\lambda(\omega)x} + D \ v_{+}^{(-)}e^{-\lambda(\omega)x} & x > 0\\ v_{-}^{(+)}e^{\lambda(\omega)x} & x < 0 \end{cases}$$

Here, the vectors $v_{\pm}^{(\pm)} = v_{\pm}^{(\pm)}(\omega, \tau)$ are displayed in (3.7) and A, B, C, D, which depend on ω and τ , are given by

(3.10a)
$$A(\omega,\tau) = \frac{(e^{-i\tau} - 1)(\omega + i\lambda(\omega))}{2i\lambda(\omega)},$$

(3.10b)
$$B(\omega,\tau) = \frac{\omega(1-e^{-i\tau})+i\lambda(\omega)(1+e^{-i\tau})}{2i\lambda(\omega)},$$

(3.10c)
$$C(\omega,\tau) = \frac{\omega(e^{i\tau}-1) + i\lambda(\omega)(e^{i\tau}+1)}{2i\lambda(\omega)},$$

(3.10d)
$$D(\omega,\tau) = \frac{(1-e^{i\tau})(\omega-i\lambda(\omega))}{2i\lambda(\omega)}.$$

(2) Moreover, we have the algebraic relations

(3.11a)
$$\overline{\eta_+(x;\omega,\tau)} = S(\tau)\eta_-(-x;\overline{\omega},\tau) ,$$

(3.11b)
$$\overline{\eta_{-}(x;\omega,\tau)} = S(\tau)\eta_{+}(-x;\overline{\omega},\tau) ,$$

where $S(\tau)$ is the 2 × 2 matrix defined

$$S(\tau) = \begin{pmatrix} e^{i\tau} & 0\\ 0 & 1 \end{pmatrix}.$$

3.2. Constructing the resolvent kernel. In this section we give an explicit construction of the resolvent kernel in terms of the solutions η_{\pm} of Proposition 3.4. For a fixed τ , we omit the τ dependence and write $\eta_{\pm}(x;\omega,\tau) = \eta_{\pm}(x;\omega)$ for brevity.

Proposition 3.5. The resolvent kernel of $\Re(\omega, \tau) = (\mathcal{D}(\tau) - \omega)^{-1}$ is given by

$$\Re(\omega,\tau)(x,y) = \frac{i}{\varphi(\omega,\tau)} \begin{pmatrix} \eta_{+,1}(x;\omega) & \eta_{-,1}(x;\omega) \\ \eta_{+,2}(x;\omega) & \eta_{-,2}(x;\omega) \end{pmatrix} \begin{pmatrix} \mathbbm{1}_{(-\infty,x]}(y)\eta_{-,2}(y;\omega) & \mathbbm{1}_{(-\infty,x]}(y)\eta_{-,1}(y;\omega) \\ \mathbbm{1}_{[x,\infty)}(y)\eta_{+,2}(y;\omega) & \mathbbm{1}_{[x,\infty)}(y)\eta_{+,1}(y;\omega) \end{pmatrix}$$

where

(3.12)
$$\varphi(\omega,\tau) := i\lambda(\omega)(e^{-i\tau}+1) - \omega(e^{-i\tau}-1)$$

 $\lambda(\omega)$ is given by Lemma 3.1, and $\eta_{\pm,j}$ for j = 1,2 are the *j*-th coordinates of the Jost solutions, given which are given by Proposition 3.4.

Proof. To obtain the resolvent operator, we need to solve the inhomogeneous ODE for $\gamma(x)$

$$(\mathcal{D}(\tau) - \omega) \gamma(x) = f(x) +$$

for general $f \in L^2(\mathbb{R}; \mathbb{C}^2)$, which we do by the method of variations of parameters. This problem can be rewritten as

(3.13)
$$(\partial_x - M(x,\omega))\gamma(x) = -i\sigma_3 f(x) ,$$

where $M(x,\omega) = M(x;\omega,\tau)$ is given by (3.4). Let $X(x;\omega) = X(x;\omega,\tau)$ be the fundamental matrix (of the homogeneous problem) with columns given by $\eta_{\pm}(x;\omega) = (\eta_{\pm,1},\eta_{\pm,2})^{\top}$,

$$X(x;\omega) = \begin{pmatrix} \eta_+(x;\omega) & \eta_-(x;\omega) \end{pmatrix}$$

A particular solution of the inhomogeneous system (3.13) can be found in the form

(3.14)
$$\gamma(x;\omega) = X(x;\omega)c(x;\omega) ,$$

where $c(\cdot; \omega) : \mathbb{R} \to \mathbb{C}^2$ is to be determined. Substitution into (3.13) yields

(3.15)
$$X(x;\omega)c'(x;\omega) = -i\sigma_3 f(x).$$

To invert $X(x;\omega)$ in (3.15), note that since $\operatorname{Tr} M(x;\omega) \equiv 0$ for all $x \in \mathbb{R}$, we have by Liouville's formula

$$\det X(x;\omega) = e^{\int_0^x \operatorname{Tr} M(y;\omega) \, dy} \det X(0;\omega) = \det X(0;\omega)$$
$$= \omega(e^{-i\tau} - 1) - i\lambda(\omega)(e^{-i\tau} + 1).$$

We set, as in (3.12),

$$\varphi(\omega,\tau) \equiv i\lambda(\omega)(e^{-i\tau}+1) - \omega(e^{-i\tau}-1).$$

Solving (3.15) for $c'(x;\omega)$, we get

(3.16)
$$c'(x;\omega) = -iX(x;\omega)^{-1}\sigma_{3}f(x)$$
$$= \frac{i}{\varphi(\omega,\tau)} \begin{pmatrix} \eta_{-,2}(x;\omega) & \eta_{-,1}(x;\omega) \\ -\eta_{+,2}(x;\omega) & -\eta_{+,1}(x;\omega) \end{pmatrix} \begin{pmatrix} f_{1}(x) \\ f_{2}(x) \end{pmatrix}$$

When constructing $c(x;\omega)$ by integration of (3.16), we must choose limits of integration of each component. Informed by the exponential decay of η_{\pm} as $x \to \pm \infty$, we find that in order to ensure that $\gamma(x;\omega) = X(x;\omega)c(x;\omega)$ decays as $x \to \pm \infty$ we must take:

$$c(x,\omega) = \frac{i}{\varphi(\omega,\tau)} \int_{\mathbb{R}} \begin{pmatrix} \mathbbm{1}_{(-\infty,x]}(y)\eta_{-,2}(y;\omega) & \mathbbm{1}_{(-\infty,x]}(y)\eta_{-,1}(y;\omega) \\ \mathbbm{1}_{[x,\infty)}(y)\eta_{+,2}(y;\omega) & \mathbbm{1}_{[x,\infty)}(y)\eta_{+,1}(y;\omega) \end{pmatrix} \begin{pmatrix} f_1(y) \\ f_2(y) \end{pmatrix} dy$$

With this choice we can write the resolvent in kernel form:

(3.17)
$$\Re(\omega,\tau)f(x) = \int_{\mathbb{R}} \Re(\omega,\tau)(x,y)f(y)dy,.$$

where

$$\begin{aligned} \Re(\omega,\tau)(x,y) &= \\ \frac{i}{\varphi(\omega,\tau)} \begin{pmatrix} \eta_{+,1}(x;\omega) & \eta_{-,1}(x;\omega) \\ \eta_{+,2}(x;\omega) & \eta_{-,2}(x;\omega) \end{pmatrix} \begin{pmatrix} \mathbb{1}_{(-\infty,x]}(y)\eta_{-,2}(y;\omega) & \mathbb{1}_{(-\infty,x]}(y)\eta_{-,1}(y;\omega) \\ \mathbb{1}_{[x,\infty)}(y)\eta_{+,2}(y;\omega) & \mathbb{1}_{[x,\infty)}(y)\eta_{+,1}(y;\omega) \end{pmatrix}. \end{aligned}$$

Our construction of $\Re(\omega, \tau)(x, y)$ implies the following result, which encompasses Theorem 1.1:

Proposition 3.6. (1) Let ω_{τ} be defined as

$$\omega_{\tau} \equiv \cos(\tau/2) \, .$$

For $\omega \in \mathbb{C} \setminus \{(-\infty, -1] \cup \{\omega_{\tau}\} \cup [1, \infty)\}\)$, the mapping $f(x) \mapsto \Re(\omega, \tau) f(x)$, given in (3.17), defines a bounded operator on a dense subspace of $L^2(\mathbb{R}; \mathbb{C}^2)$.

- (2) The resolvent kernel, $\Re(\omega, \tau)(x, y)$, has a pole at $\omega = \omega_{\tau}$.
- (3) Correspondingly, $\mathcal{D}(\tau)$ has an eigenvalue $\tau \in [0, 2\pi] \mapsto \omega_{\tau} = \cos(\tau/2)$, which traverses the spectral gap (-1, 1).

Proof. The energy, ω , is a pole of the resolvent kernel if and only if the Wronskian det $X(0; \omega, \tau) = \varphi(\omega, \tau)$ vanishes. Thus, $\omega(e^{-i\tau} - 1) - i\lambda(\omega)(e^{-i\tau} + 1) = 0$, which has the unique solution

(3.18)
$$\omega_{\tau} = \cos(\tau/2)$$

The corresponding eigenstate can be deduced as follows. Consider the expression (3.9a) for the function $\eta_+(x;\omega,\tau)$, satisfying $\mathcal{D}(\tau)\eta_+ = \omega\eta_+$. By (3.10b) and the expression (3.12) for $\varphi(\omega,\tau)$

$$B = -\frac{1}{2i\lambda(\omega)}\varphi(\omega,\tau)$$

Since $\varphi(\omega_{\tau}, \tau) = 0$, then B = 0 and so we have that

$$\eta_+(x;\omega_\tau,\tau) = \begin{cases} v_+^{(-)}(\omega_\tau,\tau) \times e^{-\lambda(\omega_\tau)x} & x > 0\\ A(\omega_\tau,\tau) v_-^{(+)}(\omega_\tau,\tau) \times e^{\lambda(\omega_\tau)x} & x < 0 \end{cases},$$

which decays exponentially as $x \to \pm \infty$ since $\lambda(\omega_{\tau}) > 0$ (Lemma 3.1). Finally, define $\psi_{\tau}(x)$ to be a constant multiple of $\eta_{+}(x;\omega_{\tau},\tau)$ for which $\|\psi_{\tau}\|_{L^{2}(\mathbb{R})} = 1$. This yields the expression displayed in (1.2). 4. Limits of the resolvent kernel as the energy approaches $\sigma_{\rm ess}(\mathcal{D}(\tau))$

A representation of the time-evolution for the Hamiltonian, $\mathscr{D}(\tau)$, is given in (2.4) based on the functional calculus and Stone's formula for the spectral measure of $\mathscr{D}(\tau)$. To work with this representation we need to evaluate the $\mathcal{R}_+(k,\tau) - \mathcal{R}_-(k,\tau)$, where we recall the definition

(4.1)
$$\mathcal{R}_{\pm}(k,\tau) := \lim_{\varepsilon \to 0^+} \Re(\sqrt{1+k^2} \pm i\varepsilon,\tau) = \lim_{\varepsilon \to 0^+} (\mathcal{D}(\tau) - (\sqrt{1+k^2} \pm i\varepsilon))^{-1},$$

for $k \in \mathbb{R}$ (corresponding to the spectral parameter, ω , being in the essential spectrum of $\mathcal{D}(\tau)$). In this section we shall study the limits (4.1) by studying the resolvent kernel. Then, in Section 5 we study the properties of the difference $\mathcal{R}_+(k,\tau) - \mathcal{R}_-(k,\tau)$, which are summarized in Proposition 5.4.

In order to calculate the limits (4.1), we remark that by our choice of $\lambda(\omega)$ (Lemma 3.1) for $\omega \in [1, \infty)$ we have

(4.2)
$$\lim_{\varepsilon \to 0^+} \lambda(\omega \pm i\varepsilon) = \mp i \sqrt{\omega^2 - 1}.$$

Let $k \in \mathbb{R}$ and define Jost solutions, $\xi_{\pm}(x; k, \tau)$, which satisfy

$$(\mathscr{D}(\tau) - \sqrt{1+k^2})\xi_{\pm}(x;k,\tau) = 0 , \quad -\infty < x < \infty,$$

with asymptotic behavior

$$\lim_{x \to +\infty} e^{-ikx} \xi_+(x;k,\tau) = \begin{pmatrix} e^{-i\tau} \\ \sqrt{1+k^2} + k \end{pmatrix},$$
$$\lim_{x \to -\infty} e^{+ikx} \xi_-(x;k,\tau) = \begin{pmatrix} 1 \\ \sqrt{1+k^2} - k \end{pmatrix}.$$

The following result states that the Jost solutions arise as limits of the decaying solutions $\eta_{\pm}(x;\omega,\tau)$ for ω approaching the essential spectrum.

Proposition 4.1. The Jost solutions $\xi_{\pm}(x;k,\tau)$ are related to the decaying solutions $\eta_{\pm}(x;\omega,\tau)$ through the following limits.

$$\begin{split} \xi_+(x;\pm|k|,\tau) &= \lim_{\varepsilon \to 0^+} \eta_+(x;\sqrt{1+k^2} \pm i\varepsilon,\tau), \\ \xi_-(x;\pm|k|,\tau) &= \lim_{\varepsilon \to 0^+} \eta_-(x;\sqrt{1+k^2} \pm i\varepsilon,\tau). \end{split}$$

In order to prove the proposition, one uses the identity

$$\lim_{\varepsilon \to 0^+} \lambda(\sqrt{1+k^2} \pm i\varepsilon) = \mp i|k|,$$

which follows from (4.2) and the substitution $\omega = \sqrt{1+k^2}$. Henceforth we assume that $k \ge 0$. An immediate consequence of this is that the limiting resolvents $\mathcal{R}_{\pm}(k,\tau)$ may be expressed in terms

of the Jost solutions $\xi_{\pm}(x; k, \tau)$ (4.3a)

$$\mathcal{R}_{\pm}(k,\tau)(x,y) = \begin{cases} \frac{-i}{\varphi(\pm k,\tau)} \begin{pmatrix} \xi_{-,1}(x;\pm k,\tau)\xi_{+,2}(y;\pm k,\tau) & \xi_{-,1}(x;\pm k,\tau)\xi_{+,1}(y;\pm k,\tau) \\ \xi_{-,2}(x;\pm k,\tau)\xi_{+,2}(y;\pm k,\tau) & \xi_{-,2}(x;\pm k,\tau)\xi_{+,1}(y;\pm k,\tau) \end{pmatrix} & x < y \\ \frac{-i}{\varphi(\pm k,\tau)} \begin{pmatrix} \xi_{+,1}(x;\pm k,\tau)\xi_{-,2}(y;\pm k,\tau) & \xi_{+,1}(x;\pm k,\tau)\xi_{-,1}(y;\pm k,\tau) \\ \xi_{+,2}(x;\pm k,\tau)\xi_{-,2}(y;\pm k,\tau) & \xi_{+,2}(x;\pm k,\tau)\xi_{-,1}(y;\pm k,\tau) \end{pmatrix} & x > y \end{cases}$$

$$(4.3b) = \begin{cases} \frac{-i}{\varphi(\pm k,\tau)}\xi_{-}(x;\pm k,\tau)\xi_{+}(y;\pm k,\tau)^{\top}\sigma_{1} & x < y \\ \frac{-i}{\varphi(\pm k,\tau)}\xi_{+}(x;\pm k,\tau)\xi_{-}(y;\pm k,\tau)^{\top}\sigma_{1} & x > y \end{cases}$$

For explicitly formulae of the resolvent kernel, see Appendix C.

5. Scattering Theory

In this section we construct the scattering theory for $\mathcal{D}(\tau)$ and find explicit formulas for the transmission and reflection coefficients. For a fixed τ , These coefficients relate the Jost solutions $\xi_{\pm}(x;k)$ to $\xi_{\pm}(x;-k)$, in the following way:

Lemma 5.1. Let $k \in \mathbb{R} \setminus \{0\}$ be such that $\varphi(k,\tau) \neq 0$ (see (3.12)). Then, there exist unique coefficients $T_1(k,\tau), R_1(k,\tau), T_2(k,\tau)$, and $R_2(k,\tau)$ such that

- (5.1a) $\xi_+(x;-k,\tau) = T_1(k,\tau)\xi_-(x;k,\tau) R_1(k,\tau)\xi_+(x;k,\tau) ,$
- (5.1b) $\xi_{-}(x;-k,\tau) = T_{2}(k,\tau)\xi_{+}(x;k,\tau) R_{2}(k,\tau)\xi_{-}(x;k,\tau).$

We call the T_j 's and the R_j 's the transmission and reflection coefficients, respectively.

Proof. The existence and uniqueness of the coefficients T_1, T_2, R_1, R_2 follow immediately from two facts

- (1) $\xi_{\pm}(x;\pm k,\tau)$ solve the same differential equation as $\xi_{\pm}(x;\pm k,\tau)$.
- (2) For $k \neq 0$, the Jost solutions $\xi_+(x;k,\tau)$ and $\xi_-(x;k,\tau)$ are linearly independent.

The first is clear as the equation is preserved with $k \to -k$. One may observe the second by a direct calculation of the Wronskian at x = 0:

$$W[\xi_{+}(x;k,\tau),\xi_{-}(x;k,\tau)] = \varphi(k,\tau)$$

= $k(e^{-i\tau}+1) - \sqrt{1+k^{2}}(e^{-i\tau}-1) \neq 0.$

We can even go further and solve for the coefficients:

Proposition 5.2. For every $k \in \mathbb{R} \setminus \{0\}$ and $\tau \in (0, 2\pi)$,

(5.2a)
$$T(k,\tau) := T_1(k,\tau) = e^{-i\tau}T_2(k,\tau) ,$$

(5.2b)
$$\frac{R_1(k,\tau)}{T_1(k,\tau)} = \frac{-1}{e^{-i\tau}} \frac{R_2(-k,\tau)}{T_2(-k,\tau)} ,$$

where

$$T(k,\tau) = \frac{2k}{k(e^{i\tau}+1) + \sqrt{1+k^2}(e^{i\tau}-1)} = \frac{2e^{-i\tau}}{\varphi(k,\tau)} \,.$$

Proof. Applying Cramer's rule to the definition of the transmition and reflection coefficients, and having already computed the relevant Wronskians in the proof of Lemma 5.1, we have

$$\begin{split} T_1(k,\tau) &= \frac{W[\xi_+(x;-k,\tau),\xi_+(x;k,\tau)]}{W[\xi_-(x;k,\tau),\xi_+(x;k,\tau)]} = \frac{2e^{-i\tau}k}{\varphi(k,\tau)} \ ,\\ R_1(k,\tau) &= -\frac{W[\xi_-(x;k,\tau),\xi_+(x;-k,\tau)]}{W[\xi_-(x;k,\tau),\xi_+(x;k,\tau)]} = \frac{W[\xi_+(x;-k,\tau),\xi_-(x;k,\tau)]}{\varphi(k,\tau)} \ ,\\ T_2(k,\tau) &= \frac{W[\xi_-(x;-k,\tau),\xi_-(x;k,\tau)]}{W[\xi_+(x;k,\tau),\xi_-(x;k,\tau)]} = \frac{2k}{\varphi(k,\tau)} \ ,\\ R_2(k,\tau) &= -\frac{W[\xi_+(x;k,\tau),\xi_-(x;-k,\tau)]}{W[\xi_+(x;k,\tau),\xi_-(x;k,\tau)]} = \frac{W[\xi_+(x;k,\tau),\xi_-(x;-k,\tau)]}{\varphi(k,\tau)} .\end{split}$$

This leads to the relations (5.2).

The explicit formula for the transmission coefficient is given by

$$T(k,\tau) = \frac{2k}{k(e^{i\tau}+1) + \sqrt{1+k^2}(e^{i\tau}-1)}$$

Moreover, the square magnitude and its derivatives are given by

(5.3)
$$|T(k,\tau)|^2 = \frac{k^2}{k^2 + \sin^2(\tau/2)}$$

(5.4)
$$\partial_k |T(k,\tau)|^2 = \frac{2k\sin^2(\tau/2)}{(k^2 + \sin^2(\tau/2))^2} ,$$

(5.5)
$$\partial_k^2 |T(k,\tau)|^2 = \frac{2\sin^2(\tau/2)(\sin^2(\tau/2) - 3k^2)}{(k^2 + \sin^2(\tau/2))^3}.$$

From this we have the estimates

Proposition 5.3. We have the following bounds on $\partial_k^j |T(k,\tau)|^2$ for j = 0, 1, 2.

$$|T(k,\tau)|^{2} \leq \min\left(1, \frac{k^{2}}{k^{2} + \sin^{2}(\tau/2)}\right) ,$$

$$\partial_{k}|T(k,\tau)|^{2} \leq \min\left(\frac{1}{k}, \frac{2k}{k^{2} + \sin^{2}(\tau/2)}, \frac{2}{k^{3}}\right) ,$$

$$\partial_{k}^{2}|T(k,\tau)|^{2} \leq \min\left(\frac{3}{k^{2}}, \frac{8}{k^{2} + \sin^{2}(\tau/2)}, \frac{7}{k^{4}}\right).$$

Proof. By direct computation and elementary bounds $\sin^2(\theta) \in [0,1]$ and $k^2, k^4 \ge 0$.

We now state a proposition which gives several alternative ways to calculate the difference across the continuous spectrum of the two operators defined by the limiting absorption principle above.

Proposition 5.4 (Jump in the resolvent kernel across the real axis). For any $k \in \mathbb{R}$

$$\begin{aligned} \mathcal{R}_{+}(k,\tau)(x,y) &- \mathcal{R}_{-}(k,\tau)(x,y) \\ &= \frac{|T(k,\tau)|^{2}}{2ie^{-i\tau}k} \left(\xi_{+}(x;k,\tau)\xi_{+}(y;-k,\tau)^{\top}\sigma_{1} + e^{-i\tau}\xi_{-}(x;k,\tau)\xi_{-}(y;-k,\tau)^{\top}\sigma_{1} \right) \end{aligned}$$

Proof. We start with x > y. Using (5.1) along with the properties of the transmission and reflection coefficients we find

$$\begin{aligned} &\mathcal{R}_{+}(k,\tau)(x,y) - \mathcal{R}_{-}(k,\tau)(x,y) \\ &= \frac{-i}{\varphi(k,\tau)} \xi_{+}(x;k,\tau) \xi_{-}(y;k,\tau)^{\top} \sigma_{1} - \frac{-i}{\varphi(-k,\tau)} \xi_{+}(x;-k,\tau) \xi_{-}(y;-k,\tau)^{\top} \sigma_{1} \\ &= \frac{-iT(k,\tau)}{2e^{-i\tau}k} \xi_{+}(x;k,\tau) \left(e^{i\tau}T(-k,\tau) \xi_{+}(x;-k,\tau) \right) - R_{2}(-k,\tau) \xi_{-}(x;-k,\tau) \right)^{\top} \sigma_{1} \\ &- \frac{iT(-k,\tau)}{2e^{-i\tau}k} \left(T(k,\tau) \xi_{-}(x;k,\tau) - R_{1}(k,\tau) \xi_{+}(x;k,\tau) \right) \xi_{-}(y,-k)^{\top} \sigma_{1} \\ &= \frac{|T(k,\tau)|^{2}}{2ie^{-i\tau}k} \left(\xi_{+}(x;k,\tau) \xi_{+}(y;-k,\tau)^{\top} \sigma_{1} + e^{-i\tau} \xi_{-}(x;k,\tau) \xi_{-}(y;-k,\tau)^{\top} \sigma_{1} \right) \end{aligned}$$

The case where x < y is similar.

Now that we have developed the requisite scattering theory we may prove the main theorem.

6. Proof of Theorem 1.2, the main theorem

We begin by remarking that it suffices to prove the following weaker bound which, in contrast to (1.3) (equivalently (1.4)), is singular as $t \downarrow 0$:

(6.1)
$$\|\langle x \rangle^{-2} e^{-i \mathscr{D}(\tau) t} P_{ac}(\mathscr{D}(\tau)) \alpha_0 \|_{L^{\infty}(\mathbb{R})} \le C_{\epsilon} \frac{1}{t^{1/2}} \frac{1}{1 + \sin^2(\tau/2) t} \|\langle x \rangle^2 \langle \mathscr{D}(\tau) \rangle^{3/2 + \epsilon} \alpha_0 \|_{L^1(\mathbb{R})}$$

Indeed, from the expression for the resolvent kernel in Appendix C, we note that for all x and y, the resolvent kernel satisfies: $|k\mathcal{R}_{\pm}(k,\tau)(x,y)| \leq C_{\tau}$, for some constant C_{τ} . Therefore, directly from the representation of $\alpha(t,x)$ as an oscillatory integral of the data α_0 in (2.4), we have:

(6.2)
$$\|\alpha(t,x)\|_{L^{\infty}(\mathbb{R}_{x})} \lesssim \|\alpha_{0}(y)\|_{L^{1}(\mathbb{R}_{y})} \quad \text{for all } t \in \mathbb{R}$$

We next derive an estimate reflecting (6.1) for t bounded away from zero and (6.2) for t near zero by interpolation. Denote

$$\begin{split} a &= \|\langle x \rangle^{-2} e^{-i \mathscr{D}(\tau) t} P_{ac}(\mathscr{D}(\tau)) f\|_{L^{\infty}(\mathbb{R})} \,, \qquad b \sim \|\langle x \rangle^{2} \langle \mathscr{D}(\tau) \rangle^{3/2 + \epsilon} f\|_{L^{1}(\mathbb{R})} \,, \\ c &\sim \frac{1}{t^{1/2}} \frac{1}{1 + \sin^{2}(\tau/2) t} \|\langle x \rangle^{2} \langle \mathscr{D}(\tau) \rangle^{3/2 + \epsilon} f\|_{L^{1}(\mathbb{R})} \,. \end{split}$$

Then, using the bounds (6.1) and (6.2), we have that $a \leq b$ and $a \leq c$. A bound which is nonsingular as $t \to 0$ now follows from the algebraic interpolation inequality (1.9). We therefore focus on proving the decay-estimate (6.1).

We now embark on the proof. Our strategy is to decompose the representation of the evolution operator, given in (2.4), into "high k" (or "high energy") and "low k" (or "low energy") integrals. To do so, fix $k_0 > 0$ and define a smooth cutoff function χ , defined on \mathbb{R} , such that

(6.3)
$$\chi(k) = \begin{cases} 1 & \text{if } k < k_0 \\ 0 & \text{if } k > 2k_0 \end{cases}$$

Then, for every $\alpha_0 \in L^2(\mathbb{R}; \mathbb{C}^2)$, we have from (2.4)) that

(6.4)
$$\alpha(t,x) = e^{-i\mathcal{D}(\tau)t} P(\mathcal{D}(\tau)) \langle \mathcal{D}(\tau) \rangle^{-3/2-\varepsilon} \alpha_0 := \alpha_l(t,x) + \alpha_h(t,x) ,$$

where

(6.5)

$$\alpha_{l}(t,x) := \frac{1}{2\pi i} \int_{0}^{\infty} e^{-i\sqrt{1+k^{2}t}} \chi(k) \left[\left(\mathcal{R}_{+}(k,\tau) - \mathcal{R}_{-}(k,\tau) \right) \alpha_{0} \right](x) \left\langle k \right\rangle^{-3/2-\varepsilon} \frac{k}{\sqrt{1+k^{2}}} \, dk \,,$$
(6.6)
$$\alpha_{l}(t,x) := \frac{1}{2\pi i} \int_{0}^{\infty} e^{-i\sqrt{1+k^{2}t}} (1-\chi(k)) \left[\left(\mathcal{R}_{+}(k,\tau) - \mathcal{R}_{-}(k,\tau) \right) \alpha_{0} \right](x) \left\langle k \right\rangle^{-3/2-\varepsilon} \frac{k}{\sqrt{1+k^{2}t}} \, dk \,,$$

$$\alpha_h(t,x) := \frac{1}{2\pi i} \int_0^\infty e^{-i\sqrt{1+k^2}t} (1-\chi(k)) \left[\left(\mathcal{R}_+(k,\tau) - \mathcal{R}_-(k,\tau) \right) \alpha_0 \right](x) \langle k \rangle^{-3/2-\varepsilon} \frac{k}{\sqrt{1+k^2}} dk \, .$$

Note that $\alpha(t, x)$ in (6.4) is the solution of the time-dependent Dirac equation (1.1) with the smoothed initial data $P(\mathcal{D}(\tau))\langle \mathcal{D}(\tau)\rangle^{-3/2-\varepsilon}\alpha_0$. The functions $\alpha_l(t, x)$ and $\alpha_l(t, x)$ are, respectively, the low k and high k components of $\alpha(t, x)$. We estimate these separately in the following sections.

6.1. High Energy Estimates. In this section we prove high-energy the following τ -independent bounds on $\alpha_h(t, x)$, defined in (6.4).

Theorem 6.1. There exist positive constants $C_{h,1}$ and $C_{h,2}$, such that for any $\tau \in [0, 2\pi)$ and $t \ge 0$

(6.7)
$$\|\langle x \rangle^{-1} \alpha_h(t,x)\|_{L^{\infty}(\mathbb{R}_x)} \le C_{h,1} \frac{1}{|t|^{3/2}} \|\langle y \rangle \alpha_0(y)\|_{L^1(\mathbb{R}_y)}$$

(6.8)
$$\|\alpha_h(t,x)\|_{L^{\infty}(\mathbb{R}_x)} \le C_{h,2} \frac{1}{|t|^{1/2}} \|\alpha_0(y)\|_{L^1(\mathbb{R}_y)}$$

Proof. To bound α_h , the high energy part of α , it is sufficient to bound the contributions from \mathcal{R}_+ and \mathcal{R}_- (see (4.1)) individually. Write

(6.9a)
$$\alpha_h(t,x) = \alpha_h^{(+)}(t,x) - \alpha_h^{(-)}(t,x)$$

where

(6.9b)
$$\alpha_h^{(\pm)}(t,x) \equiv \frac{1}{2\pi i} \int_0^\infty e^{-i\sqrt{1+k^2}t} (1-\chi(k)) \left[\mathcal{R}_{\pm}(k,\tau)\alpha_0\right](x) \langle k \rangle^{-3/2-\varepsilon} \frac{k}{\sqrt{1+k^2}} dk$$

We will estimate $\alpha_h^{(+)}$ explicitly; the estimates for $\alpha_h^{(-)}$ follow analogously. The operator $\mathcal{R}_+(k,\tau)$, see (4.1), acting on functions $\beta \in L^2(\mathbb{R}; \mathbb{C}^2)$, can be written via its kernel representation, i.e.,

$$[\mathcal{R}_+(k,\tau)\beta](x) = \int_{\mathbb{R}} \mathcal{R}_+(k,\tau)(x,y)\beta(y) \, dy \; ,$$

where the resolvent kernel $\mathcal{R}_+(k,\tau)(x,y)$, is displayed in (6.9). Substitution into (6.9) yields, via the Fubini-Tonelli theorem, that for any $x \in \mathbb{R}$

(6.10)
$$\alpha_h^{(+)}(t,x) = \int_{\mathbb{R}} A_t(x,y)\alpha_0(y)dy$$

where

$$A_t(x,y) \equiv \frac{1}{2\pi i} \int_0^\infty e^{-i\sqrt{1+k^2}t} (1-\chi(k)) \mathcal{R}_+(k,\tau)(x,y) \langle k \rangle^{-3/2-\varepsilon} \frac{k}{\sqrt{1+k^2}} dk$$

Hence,

(6.11)
$$\|\alpha_h(t,x)\|_{L^{\infty}(\mathbb{R}_x)} \leq \sup_{z,z' \in \mathbb{R}} |A_t(z,z')| \times \|\alpha_0(y)\|_{L^1(\mathbb{R}_y)},$$

and (by inserting a $1 = \langle y \rangle^{-1} \langle y \rangle$ terms into the integrand)

(6.12)
$$\|\langle x \rangle^{-1} \alpha_h(t,x)\|_{L^{\infty}(\mathbb{R}_x)} \leq \sup_{z,z' \in \mathbb{R}} \left(\langle z \rangle^{-1} |A_t(z,z')| \langle z' \rangle^{-1} \right) \times \|\langle y \rangle \alpha_0\|_{L^1(\mathbb{R}_y)}.$$

Theorem 6.1 will follow if we can establish the following kernel bounds:

(6.13)
$$\sup_{x,y \in \mathbb{R}} |A_t(x,y)| \le C|t|^{-\frac{1}{2}}$$

(6.14)
$$\sup_{x,y\in\mathbb{R}} \langle x \rangle^{-1} |A_t(x,y)| \langle y \rangle^{-1} \le C' |t|^{-\frac{3}{2}}$$

We focus on the proof of bound (6.14) and its use in the proof of (6.7). The proof of (6.13) and its application to the proof of (6.8) is simpler.

 $A_t(x,y)$ is defined as an integral with respect to $k \in \text{supp}(1-\chi) = [k_0,\infty)$. To bound $A_t(x,y)$, we express $[k_0,\infty)$ as union of overlapping intervals of exponentially increasing size: $[k_0,\infty) = \bigcup_{j>0} [k_0 2^j, k_0 2^{j+2}], j \ge 0$, and use the corresponding smooth partition of unity

(6.15)
$$\mathbb{1}_{(k_0,\infty)}(x) = \sum_{j\geq 0} \chi_j(x), \quad \chi_j \in C_c^\infty\left([k_0 2^j, k_0 2^{j+2}]\right).$$

By the triangle inequality, for every $x, y \in \mathbb{R}$

(6.16)
$$|A_t(x,y)| \le \sum_{j=0}^{\infty} I_j(x,y)$$

where

(6.17)
$$I_j(x,y) = \left| \frac{1}{2\pi i} \int_{k_0}^{\infty} e^{-it\sqrt{1+k^2}} \mathcal{R}_+(k,\tau)(x,y) \langle k \rangle^{-3/2-\varepsilon} \chi_j(k) \frac{k}{\sqrt{1+k^2}} \, dk \right|.$$

The kernel, $\mathcal{R}_+(k,\tau)(x,y)$, is a 2×2 matrix whose elements are displayed in Appendix C. Based on the precise expressions for these matrix elements, I_j can be written as a linear combination of expressions of the following type:

(6.18)
$$\left| \frac{1}{2\pi i} \int_{k_0}^{\infty} e^{-it\sqrt{1+k^2} \pm ik(x\pm y)} L(k) \langle k \rangle^{-3/2-\varepsilon} \chi_j(k) \frac{k}{\sqrt{1+k^2}} \, dk \right| ,$$

where L(k) is a rational function of k and $\sqrt{1+k^2}$, which can be read off the formulae in Appendix C. To obtain upper bounds on (6.18) we make use of the following general result on oscillatory integrals, whose proof we give in Appendix D:

Lemma 6.2. Let $\psi(k)$ be a smooth function supported in $[2^j, 2^{j+2}] \cup [-2^{j+2}, -2^j]$ for $j \ge 1$, and for j = 0 let $\psi(k)$ be supported in a neighborhood of the origin. Then, there is a constant C > 0, which is independent of ψ , such that for all $r \in \mathbb{R}$ and all $t \in \mathbb{R}$:

$$\begin{split} & \left| \int_{\mathbb{R}} e^{-it\sqrt{1+k^2} + ikr} \psi(k) dk \right| \\ & \leq C \min\left(\|\psi\|_{L^1}, |t|^{-\frac{1}{2}} 2^{\frac{3}{2}j} \|\partial_k \psi\|_{L^1}, |t|^{-\frac{3}{2}} 2^{\frac{3}{2}j} \left\| (\partial_{kk} + ir\partial_k) \left(\frac{\sqrt{1+k^2}}{k} \psi \right) \right\|_{L^1} \right) \,. \end{split}$$

In order to bound (6.17) via (6.18), we apply Lemma 6.2, for each $j \ge 0$ to

(6.19)
$$\psi = \psi_j(k) = \chi_j(k)L(k)\frac{k}{\sqrt{1+k^2}}\langle k \rangle^{-3/2-\varepsilon}$$

Substituting (6.19) into the third norm expression in the upper bound of Lemma 6.2, we have that there is a constant, C_{ε} , which is independent of k, such that

$$\left\| \left(\partial_{kk} \pm i(x\pm y)\partial_k\right) \left(\frac{\sqrt{1+k^2}}{k}\psi_j\right) \right\|_{L^1} = \left\| \left(\partial_{kk} \pm i(x\pm y)\partial_k\right) \left(\langle k \rangle^{-3/2-\varepsilon} L(k)\chi_j\right) \right\|_{L^1} \\ \leq C_{\varepsilon} \left[\mathcal{I}_j^1 + \mathcal{I}_j^2 + \mathcal{I}_j^3\right] \,,$$

where the $\mathcal{I}_j(x, y)$ are given by

$$\begin{aligned} &(6.20a)\\ \mathcal{I}_{j}^{1} \equiv \left\| \langle k \rangle^{-3/2 - \varepsilon} \left[L''(k) \chi_{j}(k) + 2L'(k) \chi_{j}(k)' + L(k) \chi_{j}''(k) \pm (x \pm y) \left(L'(k) \chi_{j}(k) + L(k) \chi_{j}'(k) \right) \right] \right\|_{L_{k}^{1}}, \\ &(6.20b)\\ \mathcal{I}_{j}^{2} \equiv \left\| k \langle k \rangle^{-7/2 - \varepsilon} \left[2L'(k) \chi_{j}(k) + 2L(k) \chi_{j}'(k) \pm (x \pm y) L(k) \chi_{j}(k) \right] \right\|_{L_{k}^{1}}, \\ &(6.20c)\\ \mathcal{I}_{j}^{3} \equiv \left\| \left(\langle k \rangle^{-7/2 + \varepsilon} + k^{2} \langle k \rangle^{-11/2 - \varepsilon} \right) L(k) \chi_{j}(k) \right\|_{L_{k}^{1}}. \end{aligned}$$

Proposition 6.3. There is a constant C > 0 such that $\mathcal{I}_{i}^{1}, \mathcal{I}_{i}^{2}, \mathcal{I}_{i}^{3} \leq C2^{-(\frac{3}{2}+\varepsilon)j} \langle x \rangle \langle y \rangle$.

Proposition 6.3 is proved in Appendix D using Van der Corput's Lemma on oscillatory integrals. We finally apply the last upper bound in Lemma 6.2 with $\psi = \psi_j$ given by (6.19), to bound I_j defined in (6.17). We obtain

$$I_j(x,y) \lesssim \left(|t|^{-\frac{3}{2}} 2^{\frac{3}{2}j} \right) \cdot \left[\mathcal{I}_j^1 + \mathcal{I}_j^2 + \mathcal{I}_j^3 \right] \lesssim \left(|t|^{-\frac{3}{2}} 2^{\frac{3}{2}j} \right) \cdot \left(2^{-(\frac{3}{2} + \varepsilon)j} \langle x \rangle \langle y \rangle \right) = 2^{-\varepsilon j} |t|^{-\frac{3}{2}} \langle x \rangle \langle y \rangle.$$

Therefore, by (6.17) we have for any $x, y \in \mathbb{R}$

$$\langle x \rangle^{-1} |A_t(x,y)| \langle y \rangle^{-1} \le \sum_{j=0}^{\infty} \langle x \rangle^{-1} I_j(x,y) \langle y \rangle^{-1} \lesssim |t|^{-\frac{3}{2}},$$

since ε is strictly positive. This proves the kernel bound (6.14) and therewith the high energy time-decay bound (6.7) of Theorem 6.1. As remarked above, the bound (6.8) follows by a closely related, but simpler, argument, involving the second upper bound in Lemma 6.2. This completes the proof of Theorem 6.1.

6.2. Low Energy Estimates. Concerning $\alpha_l(x,t)$, the low energy part of $\alpha(x,t)$, we have the following time-decay estimate:

Theorem 6.4. Let $\alpha_l(x,t)$ be defined as in (6.5). There is a constant C > 0 such that for $\tau \in [0, 2\pi]$ we have the following weighted $L^1 \to L^\infty$ bound

(6.21)
$$\|\langle x \rangle^{-2} \alpha_l(x,t)\|_{L^{\infty}(\mathbb{R}_x)} \le \frac{C}{|t|^{1/2}} \frac{1}{1+\sin^2(\tau/2)t} \|\langle y \rangle^2 \alpha_0(y)\|_{L^1(\mathbb{R}_y)}.$$

In contrast to the high-energy time-decay bound (Theorem 6.1), the low-energy bound (Theorem 6.4) depends on τ , and in particular for (threshold) resonant values of τ , the decay rate is only $\lesssim t^{-1/2}$.

The proof of Theorem 6.4 proceeds in the following way:

- (1) Proposition 6.6 establishes a $t^{-1/2}$ upper bound for all τ values.
- (2) Proposition 6.7 establishes a $t^{-3/2}$ upper bound for all $\tau \neq 0, 2\pi$.

(3) Finally, we use these upper bounds together with the interpolation inequality (1.9) to get the upper bound in Theorem 6.4.

To capture the subtleties involving k near zero (equivalently - energies near the band edge), we must carefully account for cancellations in $\mathcal{R}_+(k,\tau) - \mathcal{R}_-(k,\tau)$. The key is the jump formula of Proposition 5.4, which we use to reexpress the low energy representation formula (6.5):

$$\alpha_{l}(t,x) = \frac{1}{2\pi i} \int_{0}^{\infty} e^{-i\sqrt{1+k^{2}t}} \chi(k) \ \langle k \rangle^{-3/2-\varepsilon} \frac{k}{\sqrt{1+k^{2}}} \left[\left(\mathcal{R}_{+}(k,\tau) - \mathcal{R}_{-}(k,\tau) \right) \alpha_{0} \right](x) \ dk$$

By Proposition 5.4, we have

where

(6.23)
$$f(x,y;k,\tau) = \left(\xi_+(x;k,\tau)\xi_+(y;-k,\tau)^\top \sigma_1 + e^{-i\tau}\xi_-(x;k,\tau)\xi_-(y;-k,\tau)^\top \sigma_1\right).$$

Substitution of (6.22) into the definition of $\alpha_l(t, x)$ (and cancelling factors of k), we obtain

(6.24)
$$\alpha_l(t,x) = \frac{1}{-4\pi e^{-i\tau}} \int_0^\infty e^{-i\sqrt{1+k^2}t} \chi(k) \frac{|T(k,\tau)|^2}{\sqrt{1+k^2}} F(x;k,\tau) dk,$$

where

(6.25)
$$F(x;k,\tau) = \langle k \rangle^{-3/2-\varepsilon} \int_{\mathbb{R}} f(x,y;k,\tau) \ \alpha_0(y) dy \ .$$

We derive the desired low energy bound in two steps: first, in Proposition 6.6, we prove a $|t|^{-\frac{1}{2}}$ decay-estimate for $\alpha_l(t,x)$ which holds uniformly in τ . Second, in Proposition 6.7, we prove a τ -dependent $|t|^{-\frac{3}{2}}$ decay-estimate, as in Theorem 1.2.

Remark 6.5. In order to prove the aforementioned bounds, we must split the integral representation of $\alpha_l(x,t)$ further into two subdomains: k values in a neighborhood of zero, and k values which are small but bounded away from zero. This is necessary as there are no uniform in τ bounds near k = 0 for $\partial_k |T(k,\tau)|^2$; see (5.4). Therefore, one cannot apply a stationary phase argument for small values of k. For those values of k, a cruder estimate is used, and therefore this interval is chosen to be vanishingly small as $t \to \infty$.

Proposition 6.6 (τ -independent estimate). Let $\alpha_l(t, x)$ be defined as in (6.5) or equivalently (6.24). There exists a constant $C_l > 0$ such that for every $\tau \in [0, 2\pi]$ and every t > 0

(6.26)
$$\|\langle x\rangle^{-1}\alpha_l(x,t)\|_{L^{\infty}(\mathbb{R}_x)} \leq \frac{C_l}{|t|^{1/2}} \|\langle y\rangle\alpha_0(y)\|_{L^1(\mathbb{R}_y)}$$

Proof. It is useful to decompose $\alpha_l(x,t)$ further into its spectral contributions from: $0 \le k \le |t|^{-\frac{1}{2}}$ and $|t|^{-\frac{1}{2}} \le k < \infty$ as follows

(6.27)
$$\alpha_{l}(x,t) = \frac{1}{-4\pi e^{-i\tau}} \int_{0}^{\infty} e^{-i\sqrt{1+k^{2}t}} \chi(k)\chi(k\sqrt{t}) \frac{|T(k,\tau)|^{2}}{\sqrt{1+k^{2}}} F(x;k,\tau) dk + \frac{1}{-4\pi e^{-i\tau}} \int_{0}^{\infty} e^{-i\sqrt{1+k^{2}t}} \chi(k)(1-\chi(k\sqrt{t})) \frac{|T(k,\tau)|^{2}}{\sqrt{1+k^{2}}} F(x;k,\tau) dk := \alpha_{l}^{(1)}(x,t) + \alpha_{l}^{(2)}(x,t).$$

To bound the first term, we note that since $|T(k,\tau)| \leq 1$ (Lemma 5.3), and since the integral is over an interval of length $\sim t^{-1/2}$, we get

(6.28)
$$|\alpha_l^{(1)}(x,t)| \le \frac{1}{4\pi} \int_0^{2k_0/\sqrt{t}} \frac{|T(k,\tau)|^2}{\sqrt{1+k^2}} |F(x;k,\tau)| dk \le \frac{k_0}{2\pi\sqrt{t}} \sup_k |F(x;k,\tau)|.$$

To bound the second term, we rewrite

$$|\alpha_l^{(2)}(x,t)| = \left| \frac{1}{4\pi e^{-i\tau}} \int_0^\infty \frac{\sqrt{1+k^2}}{ik} \partial_k \left(e^{-i\sqrt{1+k^2}t} \right) \chi(k) (1-\chi(k\sqrt{t})) \frac{|T(k,\tau)|^2}{\sqrt{1+k^2}} F(x;k,\tau) dk \right| \,.$$

Since the support of $\chi(k)(1 - \chi(k\sqrt{t}))$ is bounded away from k = 0 and $k = +\infty$, integration by parts and the triangle inequality yields

$$\begin{aligned} |\alpha_{l}^{(2)}(x,t)| &= \left| \frac{1}{4\pi t} \int_{0}^{\infty} e^{-i\sqrt{1+k^{2}t}} \partial_{k} \left[\chi(k)(1-\chi(k\sqrt{t})) \frac{|T(k,\tau)|^{2}}{k} F(x;k,\tau) \right] dk \right| \\ &\leq \frac{1}{4\pi t} \int_{0}^{\infty} \left| e^{-i\sqrt{1+k^{2}t}} \partial_{k} \left[\chi(k)(1-\chi(k\sqrt{t})) \frac{|T(k,\tau)|^{2}}{k} F(x;k,\tau) \right] \right| dk , \\ &\leq \frac{1}{4\pi t} \int_{k_{0}/\sqrt{t}}^{2k_{0}} \left| \frac{\partial_{k} \left(|T(k,\tau)|^{2} F(x;k,\tau) \right)}{k} \right| + \left| \partial_{k} \left(\frac{\chi(k)(1-\chi(k\sqrt{t}))}{k} \right) |T(k,\tau)|^{2} F(x;k,\tau) \right| dk \\ (6.29) \qquad := (\mathbf{I}) + (\mathbf{II}). \end{aligned}$$

For (II) we estimate as follows, since the transmission coefficient satisfies $|T(k,\tau)| \leq 1$,

$$\begin{aligned} \text{(II)} &= \frac{1}{4\pi t} \int_{k_0/\sqrt{t}}^{2k_0} \left| \partial_k \left(\frac{\chi(k)(1-\chi(k\sqrt{t}))}{k} \right) |T(k,\tau)|^2 F(x;k,\tau) \right| dk \\ &= \frac{1}{4\pi t} \sup_{k \in [0,2k_0]} |F(x;k,\tau)| \int_{k_0/\sqrt{t}}^{2k_0} \left| \partial_k \left(\frac{\chi(k)(1-\chi(k\sqrt{t}))}{k} \right) \right| dk \\ &= \frac{1}{4\pi t} \sup_{k \in [0,2k_0]} |F(x;k,\tau)| \int_{k_0/\sqrt{t}}^{2k_0} \left| \left(\frac{\chi'(k)(1-\chi(k\sqrt{t})) + \sqrt{t}\chi(k)\chi'(k\sqrt{t})}{k} - \frac{\chi(k)(1-\chi(k\sqrt{t}))}{k^2} \right) \right| dk \end{aligned}$$

(6.20)

(6.30)

$$\leq \frac{1}{4\pi t} \sup_{k \in [0,2k_0]} |F(x;k,\tau)| \int_{k_0/\sqrt{t}}^{2k_0} \frac{|\chi'(k)|}{k} + \frac{\left|\sqrt{t}\chi'(k\sqrt{t})\right|}{k} + \frac{1}{k^2} dk.$$

To bound (6.30), we use that $\chi'(k) = 0$ for $k \in [0, k_0)$ and $k \in (2k_0, \infty)$ to get

$$\int_{k_0/\sqrt{t}}^{2k_0} \frac{|\chi'(k)|}{k} dk \leq \sup |\chi'| \int_{k_0}^{2k_0} \frac{1}{k} dk = \sup |\chi'| \cdot \ln(2) ,$$

$$\int_{k_0/\sqrt{t}}^{2k_0} \frac{|\sqrt{t}\chi'(k\sqrt{t})|}{k} dk = \sqrt{t} \int_{k_0}^{2k_0\sqrt{t}} \frac{|\chi'(z)|}{z} dz \leq \sup |\chi'| \cdot \ln(2)\sqrt{t}$$

$$\int_{k_0/\sqrt{t}}^{2k_0} \frac{1}{k^2} dk = (\sqrt{t} - 1/2)k_0^{-1} \leq \sqrt{t}k_0^{-1}$$

Substituting these bounds into (6.30), we obtain

(6.31) (II)
$$\leq \frac{C'(1+k_0^{-1})}{\sqrt{t}} \sup_{k \in [0,2k_0]} |F(x;k,\tau)|.$$

We now bound the expression (I) in (6.29). By applying the upper bounds on the transmission coefficient from Proposition 5.3, we get

$$(\mathbf{I}) \leq \frac{1}{4\pi t} \int_{k_0/\sqrt{t}}^{2k_0} |\partial_k F(x;k,\tau)| \frac{|T(k,\tau)|^2}{k} + |\partial_k|T(k,\tau)|^2 |\frac{|F(x;k,\tau)|}{k} dk$$

$$\leq \frac{1}{4\pi t} \sup_{k \in [0,2k_0]} |\partial_k F(x;k,\tau)| \int_{k_0/\sqrt{t}}^{2k_0} \frac{1}{k} dk + \frac{1}{4\pi t} \sup_{k \in [0,2k_0]} |F(x;k,\tau)| \int_{k_0/\sqrt{t}}^{2k_0} \frac{1}{k^2} dk$$

$$\leq \frac{1}{4\pi t} \sup_{k \in [0,2k_0]} |\partial_k F(x;k,\tau)| (1+\ln(t)) + \frac{1}{4\pi t} \sup_{k \in [0,2k_0]} |F(x;k,\tau)| (1+\sqrt{t})$$

$$(6.32) \qquad \leq \frac{C''}{\sqrt{t}} \left(\sup_{k \in [0,2k_0]} |\partial_k F(x;k,\tau)| + \sup_{k \in [0,2k_0]} |F(x;k,\tau)| \right)$$

By the upper bound on $|\alpha_l^{(1)}(x,t)|$, (6.28), and the upper bound on $|\alpha_l^{(2)}(x,t)| \leq (I) + (II)$, there is a constant C > 0 (independent of α_l , t, and τ), such that

$$|\alpha_l(x,t)| \le \frac{C}{\sqrt{t}} \left(\sup_{k \in [0,2k_0]} |F(x;k,\tau)| + \sup_{k \in [0,2k_0]} |\partial_k F(x;k,\tau)| \right).$$

Hence, to complete the proof of Proposition 6.6, we need to bound F and $\partial_k F$ in terms of the L^1 norm of α_0 . These bounds are displayed in Proposition E.1 of Appendix E.

Next we prove a more subtle τ -dependent decay estimate, which requires stronger localization of α_0 . We shall make crucial use of the small |k| behavior of the transmission coefficient (see (5.3)):

(6.33)
$$|T(k,\tau)|^2 = \frac{k^2}{k^2 + \sin^2(\tau/2)}.$$

Proposition 6.7 (τ -dependent estimate). Let $\alpha_l(x,t)$ denote low-energy part of $\alpha(x,t)$; see (6.5) or equivalently (6.24). There is a τ -independent constant D > 0 such that for t > 0 and $\tau \in (0, 2\pi)$ (no threshold resonance) we have:

$$\|\langle x \rangle^{-2} \alpha_l(x,t) \|_{L^{\infty}(\mathbb{R}_x)} \le \frac{D}{\sin^2(\tau/2) |t|^{\frac{3}{2}}} \|\langle y \rangle^2 \alpha_0(y) \|_{L^1(\mathbb{R}_y)} ,$$

Proof. Assume $\tau \in (0, 2\pi)$. We study the low-energy solution $\alpha_l(t, x)$ as displayed in (6.24). Using integration by parts we have

$$\begin{split} \alpha_l(t,x) &= \frac{1}{-4\pi e^{-i\tau}} \int_0^\infty e^{-i\sqrt{1+k^2}t} \chi(k) \frac{|T(k,\tau)|^2}{\sqrt{1+k^2}} F(x;k,\tau) dk \\ &= \frac{1}{-4\pi e^{-i\tau}} \int_0^\infty \frac{\sqrt{1+k^2}}{-ikt} \partial_k \left(e^{-i\sqrt{1+k^2}t} \right) \chi(k) \frac{|T(k,\tau)|^2}{\sqrt{1+k^2}} F(x;k,\tau) dk \\ &= \frac{1}{4\pi i t e^{-i\tau}} \int_0^\infty \partial_k \left(e^{-i\sqrt{1+k^2}t} \right) \chi(k) \frac{|T(k,\tau)|^2}{k} F(x;k,\tau) dk \\ &= -\frac{1}{4\pi i t e^{-i\tau}} \int_0^\infty e^{-i\sqrt{1+k^2}t} \partial_k \left(\chi(k) \frac{|T(k,\tau)|^2}{k} F(x;k,\tau) \right) dk \\ &+ e^{-i\sqrt{1+k^2}t} \chi(k) \frac{|T(k,\tau)|^2}{k} F(x;k,\tau) \Big|_{k=0}^{k=\infty} \end{split}$$

Now note that the boundary terms vanish; At $k = \infty$, this is due to the vanishing of $\chi(k)$ as $k \to +\infty$. At k = 0, the boundary term vanishes since $|T(k, \tau)|^2 = \mathcal{O}(k^2)$ as $|k| \to 0$ for $0 < \tau < 2\pi$, by (6.33). Since the boundary terms vanish, it follows that

(6.34)
$$\alpha_l(t,x) = \frac{1}{-4\pi i t e^{-i\tau}} \int_0^\infty e^{-i\sqrt{1+k^2}t} G(x;k,\tau) dk \,,$$

where

(6.35)
$$G(x;k,\tau) \equiv \partial_k \left[\chi(k) \frac{|T(k,\tau)|^2}{k} F(x;k,\tau) \right] \,.$$

Remark 6.8. Here we see why Proposition 6.7 holds only for $\tau \in (0, 2\pi)$: for the boundary terms to vanish at k = 0, we need that $T(0, \tau) = 0$. This is not true when $\tau = 0, 2\pi$, since the transmission coefficient there is always 1.

The new representation of α_l given in (6.34) can be decomposed, in analogy with (6.27), into k near zero $(k < t^{-1/2}k_0)$ and k bounded away from zero $(k \in (t^{-1/2}k_0, 2k_0))$. For every $\tau \in (0, 2\pi)$,

(6.36)
$$\alpha_{l}(t,x) = \frac{1}{-4\pi i t e^{-i\tau}} \int_{0}^{\infty} e^{-i\sqrt{1+k^{2}t}} \chi(\sqrt{t}k)G(x;k,\tau)dk + \frac{1}{-4\pi i t e^{-i\tau}} \int_{0}^{\infty} e^{-i\sqrt{1+k^{2}t}} (1-\chi(\sqrt{t}k))G(x;k,\tau)dk = \alpha_{l}^{(i)}(x,t) + \alpha_{l}^{(ii)}(x,t).$$

We estimate each term in (6.36). First,

(6.37)
$$|\alpha_l^{(i)}(t,x)| \le \frac{1}{4\pi t} \int_0^{2k_0/\sqrt{t}} |G(x;k,\tau)| dk \le \frac{k_0}{2\pi t^{3/2}} \sup_k |G(x;k,\tau)|.$$

To estimate $\alpha_l^{(ii)}$ we first use integration by parts,

$$\begin{split} \alpha_l^{(ii)}(t,x) &= \frac{1}{-4\pi i t e^{-i\tau}} \int_0^\infty e^{-i\sqrt{1+k^2}t} (1-\chi(\sqrt{t}k)) G(x;k,\tau) dk \\ &= \frac{1}{-4\pi i t e^{-i\tau}} \int_0^\infty \frac{\sqrt{1+k^2}}{-ikt} \partial_k \left(e^{-i\sqrt{1+k^2}t} \right) (1-\chi(\sqrt{t}k)) G(x;k,\tau) dk \\ &= \frac{1}{4\pi t^2 e^{-i\tau}} \int_0^\infty e^{-i\sqrt{1+k^2}t} \partial_k \left(\frac{\sqrt{1+k^2}}{k} (1-\chi(\sqrt{t}k)) G(x;k,\tau) \right) dk \,. \end{split}$$

It follows that

$$\begin{aligned} |\alpha_l^{(ii)}(t,x)| &\leq \frac{1}{4\pi t^2} \int_{k_0/\sqrt{t}}^{2k_0} \left| \partial_k \left(\frac{\sqrt{1+k^2}}{k} (1-\chi(\sqrt{t}k)) \right) G(x;k,\tau) \right| dk \\ &+ \frac{1}{4\pi t^2} \int_{k_0/\sqrt{t}}^{2k_0} \left| \left(\frac{\sqrt{1+k^2}}{k} (1-\chi(\sqrt{t}k)) \partial_k G(x;k,\tau) \right) \right| dk \\ &\leq \text{Term}_1 + \text{Term}_2 + \text{Term}_3, \end{aligned}$$

where

$$\begin{aligned} \text{Term}_1 &\equiv \frac{1}{4\pi t^2} \sup_k |G(x;k,\tau)| \int_{k_0/\sqrt{t}}^{2k_0} \left| \frac{\sqrt{1+k^2}}{k} \sqrt{t} |\chi'(\sqrt{t}k)\rangle| \right| \, dk \,, \\ \text{Term}_2 &\equiv \frac{1}{4\pi t^2} \sup_k |G(x;k,\tau)| \int_{k_0/\sqrt{t}}^{2k_0} \frac{1}{k^2\sqrt{1+k^2}} \, dk \,, \quad \text{and} \\ \text{Term}_3 &\equiv \frac{1}{4\pi t^2} \int_{k_0/\sqrt{t}}^{2k_0} \frac{\sqrt{1+k^2}}{k} |\partial_k G(x;k,\tau)| \, dk \,. \end{aligned}$$

Recall that $G(x; k, \tau)$ is defined in (6.35) in terms of $F(x; k, \tau)$, which depends on the initial data, α_0 ; see (6.25).

We now provide upper bounds for each of the three terms above. These upper bounds rely on Proposition E.1, which provides bounds on G and its derivatives. By changing variables $(z := k\sqrt{t})$, and using that $\chi'(z)$ is compactly supported and vanishes in a neighborhood of z = 0, we obtain (with positive constants D_1 , D_2 , etc., all independent of k and τ):

(6.38)
$$\operatorname{Term}_{1} \leq \frac{D_{1}}{t^{3/2}} \frac{\langle x \rangle}{\sin^{2}(\tau/2)} \|\langle y \rangle \alpha_{0}(y)\|_{L^{1}(\mathbb{R}_{y})} \int_{k_{0}/\sqrt{t}}^{2k_{0}} \frac{\sqrt{1+k^{2}}}{k} \left| \chi'(\sqrt{t}k) \right| dk$$
$$= \frac{D_{1}}{t^{3/2}} \frac{\langle x \rangle}{\sin^{2}(\tau/2)} \|\langle y \rangle \alpha_{0}(y)\|_{L^{1}(\mathbb{R}_{y})} \int_{k_{0}}^{2k_{0}\sqrt{t}} \frac{\sqrt{1+z^{2}/t}}{z} |\chi'(z)| dz$$
$$\leq \frac{D_{2}}{t^{3/2}} \frac{\langle x \rangle}{\sin^{2}(\tau/2)} \|\langle y \rangle \alpha_{0}(y)\|_{L^{1}(\mathbb{R}_{y})}.$$

Similarly,

(6.39)
$$\operatorname{Term}_{2} \leq \frac{D_{3}}{t^{2}} \frac{\langle x \rangle}{\sin^{2}(\tau/2)} \|\langle y \rangle \alpha_{0}(y)\|_{L^{1}(\mathbb{R}_{y})} \int_{k_{0}/\sqrt{t}}^{2k_{0}} \frac{1}{k^{2}\sqrt{1+k^{2}}} dk$$
$$\leq \frac{D_{3}}{t^{2}} \frac{\langle x \rangle}{\sin^{2}(\tau/2)} \|\langle y \rangle \alpha_{0}(y)\|_{L^{1}(\mathbb{R}_{y})} \int_{k_{0}/\sqrt{t}}^{2k_{0}} \frac{1}{k^{2}} dk$$
$$\leq \frac{D_{4}}{t^{3/2}} \frac{\langle x \rangle}{\sin^{2}(\tau/2)} \|\langle y \rangle \alpha_{0}(y)\|_{L^{1}(\mathbb{R}_{y})}.$$

Finally, using the change of variables $k = (k_0/\sqrt{t})l$,

$$\operatorname{Term}_{3} \leq \frac{D_{5}}{t^{2}} \langle x \rangle^{2} \| \langle y \rangle^{2} \alpha_{0}(y) \|_{L^{1}(\mathbb{R}_{y})} \int_{k_{0}/\sqrt{t}}^{2k_{0}} \frac{\sqrt{1+k^{2}}}{k} \frac{1}{k(k^{2}+\sin^{2}(\tau/2))} dk$$

$$= \frac{D_{5}}{t^{2}} \langle x \rangle^{2} \| \langle y \rangle^{2} \alpha_{0}(y) \|_{L^{1}(\mathbb{R}_{y})} \int_{k_{0}/\sqrt{t}}^{2k_{0}} \frac{\sqrt{1+k^{2}}}{k} \frac{1}{k(k^{2}+\sin^{2}(\tau/2))} dk$$

$$= \frac{D_{6}}{t^{1/2}} \langle x \rangle^{2} \| \langle y \rangle^{2} \alpha_{0}(y) \|_{L^{1}(\mathbb{R}_{y})} \int_{1}^{2\sqrt{t}} \frac{1}{l^{2}} \frac{1}{(k_{0}^{2}l^{2}+\sin^{2}(\tau/2)t)} dl$$

$$\leq \frac{D_{6}}{t^{1/2}} \langle x \rangle^{2} \| \langle y \rangle^{2} \alpha_{0}(y) \|_{L^{1}(\mathbb{R}_{y})} \frac{1}{k_{0}^{2}+\sin^{2}(\tau/2)t} \int_{1}^{2\sqrt{t}} \frac{1}{l^{2}} dl$$

$$\leq \frac{D_{6}}{\sin^{2}(\tau/2) t^{3/2}} \langle x \rangle^{2} \| \langle y \rangle^{2} \alpha_{0}(y) \|_{L^{1}(\mathbb{R}_{y})}.$$

To conclude, we combine the upper bound (6.37) for $\alpha_l^{(i)}$, and the three upper bounds which (6.38), (6.39), and (6.40) together bound $\alpha_l^{(ii)}$, to get by (6.36) and the triangle inequality

$$|\alpha_l(t,x)| \leq \frac{D_7}{\sin^2(\tau/2) t^{3/2}} \langle x \rangle^2 \| \langle y \rangle \alpha_0(y) \|_{L^2(\mathbb{R}_y)}.$$

Finally, to complete the proof of Theorem 6.4, we interpolate via inequality (1.9), using the bounds of Propositions 6.6 and 6.7. This yields

$$\|\langle x \rangle^{-2} \alpha_l(t, x)\|_{L^{\infty}(\mathbb{R}_x)} \le \frac{D}{t^{1/2}} \frac{1}{1 + \sin^2(\tau/2)t} \|\langle y \rangle \alpha_0(y)\|_{L^1(\mathbb{R}_y)}$$

which implies the bound of Theorem 6.4.

6.3. Remarks on the proof of Theorem 1.6. By the interpolation argument at the beginning of Section 6, it suffices to prove both results in Theorem 1.6 in a weaker version, where $\langle t \rangle$ replaced by |t|. Moreover, these bounds are established with τ -independent constants for high energies in Theorem 6.1. Thus, it only remains to prove the desired inequalities with τ -dependent coefficients for the low energy part of the solution $\alpha_l(x, t)$, displayed in (6.24).

The desired inequalities follow from an application of Lemma 6.2 similar to that in Theorem 6.1. The middle bound provided by Lemma 6.2 provides the necessary inequality for (1.5) while the third bound in Lemma 6.2 provides the inequality for (1.6).

Appendix A. Proof of Proposition 3.4

Proof. The expressions (3.9a) for ξ_+ and (3.9b) for ξ_- are solutions with the desired asymptotic behavior at infinity if and only if each of these expressions is continuous at x = 0. This imposes linear inhomogeneous systems of algebraic equations for (A, B) and for (C, D). The solutions are displayed (3.10b). This proves Part 1 of Proposition 3.4.

Next we turn our attention to the relations (3.11) between ξ_+ and ξ_- . Note that (3.11b) is equivalent to (3.11a). Indeed, (3.11a) implies

$$\overline{S(\tau)\xi_+(x;\omega,\tau)} = \xi_-(-x,\overline{\omega},\tau).$$

Making the replacements $x \mapsto -x$ and $\omega \to \overline{\omega}$ gives (3.11b). Reversing the steps recovers (3.11a) from (3.11b).

The last item in Proposition 3.4 is the identity (3.11b), rewritten as:

(A.1)
$$\overline{S(\tau)}\xi_+(-x;\overline{\omega},\tau) = \xi_-(x;\omega,\tau).$$

We will first show that the expressions on either side of the equality (A.1) satisfy the same differential equation and asymptotic condition at $+\infty$. Once we do that, uniqueness then implies the equality of these expressions. First, to verify the equality of asymptotic behaviors at infinity, recall by (3.8b) that

(A.2)
$$e^{-\lambda(\omega)x}\xi_{-}(x;\omega,\tau) \longrightarrow v_{-}^{(+)} = \begin{pmatrix} 1\\ \omega - i\lambda(\omega) \end{pmatrix} \text{ as } x \to -\infty.$$

Concerning the left hand side of (A.1), note that by (3.8a)

$$S(\tau)e^{\lambda(\omega)(-x)}\xi_+(-x,\omega,\tau) \to \begin{pmatrix} 1\\ \omega+i\lambda(\omega) \end{pmatrix}$$
 as $x \to -\infty$.

Hence,

$$\overline{S(\tau)e^{\lambda(\omega)(-x)}\xi_+(-x,\omega,\tau)} \to \begin{pmatrix} 1\\ \bar{\omega}-i\lambda(\bar{\omega}) \end{pmatrix} \text{ as } x \to -\infty$$

or

$$e^{-\lambda(\bar{\omega})x}\overline{S(\tau)\xi_+(-x,\omega,\tau)} \to \begin{pmatrix} 1\\ \bar{\omega}-i\lambda(\bar{\omega}) \end{pmatrix}$$
 as $x \to -\infty$

or, replacing ω by $\bar{\omega}$, we obtain

(A.3)
$$e^{-\lambda(\omega)x}\overline{S(\tau)}\xi_+(-x,\bar{\omega},\tau) \to \begin{pmatrix} 1\\ \omega - i\lambda(\omega) \end{pmatrix} \text{ as } x \to -\infty$$

Therefore, by (A.2) and (A.3), $\overline{S(\tau)\xi_+(-x;\overline{\omega},\tau)}$ and $\xi_-(x;\omega,\tau)$ satisfy the same asymptotic condition as $x \to -\infty$.

To complete the proof we now verify that both expressions in the equality (A.1), $\overline{S(\tau)}\xi_+(-x;\overline{\omega},\tau)$ and $\xi_-(x;\omega,\tau)$, satisfy the same differential equation. We begin by first noting the following identities:

(A.4)
$$S\sigma_3 = \sigma_3 S, \quad S\sigma_1 = \overline{\sigma_\star} S, \quad S\sigma_\star = \sigma_1 S,$$

where we have used the abbreviated notation $S = S(\tau)$, $\sigma_{\star} = \sigma_{\star}(\tau)$, and $\mathcal{D} = \mathcal{D}(\tau)$. Using (A.4) we have

Changing variables: x = -y gives

$$\left[-i\sigma_{3}\partial_{y}+\sigma_{1}\mathbbm{1}_{y<0}+\overline{\sigma_{\star}}\mathbbm{1}_{y>0}\right]S\xi_{+}(-y;\omega,\tau)=\omega\ S\xi_{+}(-y;\omega,\tau),$$

taking the complex conjugate gives

$$[i\sigma_3\partial_y + \sigma_1 \mathbb{1}_{y<0} + \sigma_\star \mathbb{1}_{y>0}]\overline{S\xi_+(-y;\omega,\tau)} = \overline{\omega}\ \overline{S\xi_+(-y;\omega,\tau)}$$

and finally making the replacement $\omega \to -\overline{\omega}$ gives

$$\mathcal{D}\overline{S\xi_{+}(-y;\overline{\omega},\tau)} = [i\sigma_{3}\partial_{y} + \sigma_{1}\mathbb{1}_{y<0} + \sigma_{\star}\mathbb{1}_{y>0}]\overline{S\xi_{+}(-y;\overline{\omega},\tau)} = \omega \ \overline{S\xi_{+}(-y;\overline{\omega},\tau)}.$$

Since $\xi_{-}(x;\omega,\tau)$ satisfies the same equation and boundary condition at $-\infty$, the proof of the symmetry properties (3.11) is now complete.

Appendix B. Spectral symmetry of the operator $\mathscr{D}(\tau)$

We will show that there is a symmetry of the eigenvalue equation

(B.1)
$$\mathcal{D}(\tau)\alpha = \omega \alpha$$
.

Let $\beta(x) = \sigma_3 S(\tau) \alpha(-x)$. Then we claim that β satisfies

$$\mathscr{D}(2\pi - \tau)\alpha = -\omega\beta$$
.

To see this, replace $x \mapsto -x$ in the original equation, multiply by $\sigma_3 S(\tau)$, and use the anticommutation relations satisfied by Pauli matrices to observe

(B.2)
$$\omega \sigma_3 S(\tau) \alpha(-x) = \sigma_3 S(\tau) (-i\sigma_3 \partial_x + \sigma_1 \mathbb{1}_{(-\infty,0)}(-x) + \sigma_\star(\tau) \mathbb{1}_{[0,\infty)}(-x)) \alpha(-x)$$

(B.3)
$$= \sigma_3 S(\tau) (-i\sigma_3 \partial_x + \sigma_1 \mathbb{1}_{(0,\infty)}(x) + \sigma_\star(\tau) \mathbb{1}_{(-\infty,0]}(x)) \alpha(-x)$$

(B.4)
$$= \sigma_3(-i\sigma_3\partial_x + \sigma_\star(2\pi - \tau)\mathbb{1}_{(0,\infty)}(x) + \sigma_1\mathbb{1}_{(-\infty,0]}(x))S(\tau)\alpha(-x)$$

(B.5)
$$= (-i\sigma_3\partial_x - \sigma_\star(2\pi - \tau)\mathbb{1}_{(0,\infty)}(x) - \sigma_1\mathbb{1}_{(-\infty,0]}(x))\sigma_3S(\tau)\alpha(-x)$$

(B.6)
$$= -\mathcal{D}(2\pi - \tau)\sigma_3 S(\tau)\alpha(-x) .$$

Making the definition $\beta(x) = \sigma_3 S(\tau) \alpha(-x)$ and multiplying by minus 1, we obtain the desired equation.

Appendix C. Limiting Resolvent matrix elements

For $k \in \mathbb{R}$, the limiting resolvents $\mathcal{R}_{\pm}(k, \tau)$ are given by

(C.1)
$$\mathcal{R}_{\pm}(k,\tau) = \begin{cases} \frac{-i}{\varphi(\pm k,\tau)} \xi_{-}(x;\pm k,\tau) \xi_{+}(y;\pm k,\tau)^{\top} \sigma_{1} & x < y \\ \frac{-i}{\varphi(\pm k,\tau)} \xi_{+}(x;\pm k,\tau) \xi_{-}(y;\pm k,\tau)^{\top} \sigma_{1} & x > y \end{cases}$$

Using the explicit formulas for the Jost solutions (3.9), we can write the Resolvent entries in closed form. Let $\mathcal{R}(k,\tau)$ be defined such that

(C.2)
$$\mathcal{R}_{\pm}(k,\tau) = \mathcal{R}(\pm k,\tau).$$

Then the entries of the kernel of $\mathcal{R}(k,\tau)$ are, (where the τ -dependent coefficients are determined by (3.10b) as $\tilde{A}(k) = A(\sqrt{1+k^2})$, and similarly for $\tilde{A}, \tilde{D}, \tilde{C}$):

$$\begin{cases} \frac{i}{\varphi(k,\tau)} e^{-i\tau} \left[\tilde{D}(\sqrt{1+k^2}+k)e^{ik(x+y)} + \tilde{C}(\sqrt{1+k^2}-k)e^{ik(x-y)} \right] & x > y > 0\\ \frac{i}{\tau(k-\tau)} e^{-i\tau}(\sqrt{1+k^2}-k)e^{ik(x-y)} & x > 0 > y \end{cases}$$

$$(\mathcal{R}(k,\tau)(x,y))_{1,1} = \begin{cases} \frac{i}{\varphi(k,\tau)} \begin{bmatrix} \tilde{B}(\sqrt{1+k^2}-k)e^{ik(x-y)} + \tilde{A}(\sqrt{1+k^2}-k)e^{-ik(x+y)} \end{bmatrix} & 0 > x > y \\ \frac{i}{\varphi(k,\tau)} \begin{bmatrix} \tilde{B}(\sqrt{1+k^2}+k)e^{-i\tau}e^{ik(x+y)} + \tilde{C}(\sqrt{1+k^2}+k)e^{-i\tau}e^{-ik(x-y)} \end{bmatrix} & y > x > 0 \\ \frac{i}{\varphi(k,\tau)}(\sqrt{1+k^2}+k)e^{-ik(x-y)} & y > 0 > x \\ \frac{i}{\varphi(k,\tau)} \begin{bmatrix} \tilde{B}(\sqrt{1+k^2}+k)e^{-ik(x-y)} + \tilde{A}(\sqrt{1+k^2}-k)e^{-ik(x+y)} \end{bmatrix} & 0 > y > x \end{cases}$$

$$(C.4) \qquad (\mathcal{R}(k,\tau)(x,y))_{1,2} = \begin{cases} \frac{i}{\varphi(k,\tau)}e^{-2i\tau} \begin{bmatrix} \tilde{D}e^{ik(x+y)} + \tilde{C}e^{ik(x-y)} \end{bmatrix} & x > y > 0\\ \frac{i}{\varphi(k,\tau)}e^{-i\tau}e^{ik(x-y)} & x > 0 > y\\ \frac{i}{\varphi(k,\tau)}\tilde{B}e^{ik(x-y)} + \tilde{A}e^{-ik(x+y)} & 0 > x > y\\ \frac{i}{\varphi(k,\tau)}e^{-2i\tau} \begin{bmatrix} \tilde{D}e^{ik(x+y)} + \tilde{C}e^{-ik(x-y)} \end{bmatrix} & y > x > 0 \end{cases}, \\ \frac{i}{\varphi(k,\tau)}e^{-i\tau}e^{-ik(x-y)} & y > 0 > x\\ \frac{i}{\varphi(k,\tau)}e^{-2i\tau} \begin{bmatrix} \tilde{D}e^{ik(x+y)} + \tilde{C}e^{ik(x-y)} \end{bmatrix} & 0 > y > x\\ \frac{i}{\varphi(k,\tau)}e^{-2i\tau} \begin{bmatrix} \tilde{D}e^{ik(x+y)} + \tilde{C}e^{ik(x-y)} \end{bmatrix} & 0 > y > x \end{cases}$$

DIRAC DECAY ESTIMATES

$$(C.5) \qquad (\mathcal{R}(k,\tau)(x,y))_{2,1} = \begin{cases} \frac{i}{\varphi(k,\tau)} \tilde{D}(\sqrt{1+k^2}+k)^2 e^{ik(x+y)} + \tilde{C}e^{ik(x-y)} & x > y > 0\\ \frac{i}{\varphi(k,\tau)} e^{ik(x-y)} & x > 0 > y\\ \frac{i}{\varphi(k,\tau)} \tilde{B}e^{ik(x-y)} + \tilde{A}(\sqrt{1+k^2}-k)^2 e^{-ik(x+y)} & 0 > x > y\\ \frac{i}{\varphi(k,\tau)} \tilde{D}(\sqrt{1+k^2}+k)^2 e^{ik(x+y)} + \tilde{C}e^{-ik(x-y)} & y > x > 0\\ \frac{i}{\varphi(k,\tau)} e^{-ik(x-y)} & y > 0 > x\\ \frac{i}{\varphi(k,\tau)} \tilde{B}e^{-ik(x-y)} + \tilde{A}(\sqrt{1+k^2}-k)^2 e^{-ik(x+y)} & 0 > y > x \end{cases}$$

(C.6)

$$(\mathcal{R}(k,\tau)(x,y))_{2,2} = \begin{cases} \frac{i}{\varphi(k,\tau)} \tilde{D}e^{-i\tau} (\sqrt{1+k^2}+k)e^{ik(x+y)} + \tilde{C}e^{-i\tau} (\sqrt{1+k^2}+k)e^{ik(x-y)} & x > y > 0\\ \frac{i}{\varphi(k,\tau)} (\sqrt{1+k^2}+k)e^{ik(x-y)} & x > 0 > y\\ \frac{i}{\varphi(k,\tau)} \left[\tilde{B}(\sqrt{1+k^2}+k)e^{ik(x-y)} + \tilde{A}(\sqrt{1+k^2}-k)e^{-ik(x+y)} \right] & 0 > x > y\\ \frac{i}{\varphi(k,\tau)} e^{-i\tau} \left[\tilde{D}(\sqrt{1+k^2}+k)e^{ik(x+y)} + \tilde{C}(\sqrt{1+k^2}-k)e^{-ik(x-y)} \right] & y > x > 0 \\ \frac{i}{\varphi(k,\tau)} e^{-i\tau} (\sqrt{1+k^2}-k)e^{-ik(x-y)} & y > 0 > x\\ \frac{i}{\varphi(k,\tau)} \left[\tilde{B}(\sqrt{1+k^2}-k)e^{-ik(x-y)} + \tilde{A}(\sqrt{1+k^2}-k)e^{-ik(x+y)} \right] & 0 > y > x \end{cases}$$

where as in (3.12), substituting $\omega = \sqrt{1+k^2}$ for real values of k, we get

(C.7)
$$\varphi(k,\tau) := k(e^{-i\tau} + 1) - \sqrt{1 + k^2}(e^{-i\tau} - 1) ,$$

APPENDIX D. PROOFS OF LEMMAS FROM SECTION 6.1

D.1. Proof of Lemma 6.2. The first term is a direct consequence of the triangle inequality

$$\left| \int_{\mathbb{R}} e^{-it\sqrt{1+k^2} + ikr} \psi_j(k) \, dk \right| \le \int_{\mathbb{R}} \left| e^{-it\sqrt{1+k^2} + ikr} \right| \cdot |\psi_j(k)| \, dk = \|\psi_j\|_{L^1} \, .$$

The next two upper bounds are the result of Van der Corput's Lemma for oscillatory integrals:

Theorem D.1 (Van der Corput Lemma [41]). Let $\phi : \mathbb{R} \to \mathbb{R}$ be a smooth and let $\psi \in C_c^{\infty}(\mathbb{R})$. If $|\partial_k^2 \phi(k)| \ge \lambda \ge 0$ for all $k \in \operatorname{supp}(\psi)$, then there exists a constant C > 0 such that

(D.1)
$$\left| \int_{\mathbb{R}} e^{i\phi(k)} \psi(k) \, dk \right| \le C \lambda^{-\frac{1}{2}} \left\| \partial_k \psi(k) \right\|_{L^1(\mathbb{R}_k)}$$

.

By (D.1), we only need to compute the second derivative in k of the phase term, i.e.,

$$\left|\partial_{kk}\left[-t\sqrt{1+k^2}+kr\right]\right| = \frac{t}{(1+k^2)^{\frac{3}{2}}} = t\langle k \rangle^{-3} \gtrsim t 2^{-3(j+2)},$$

since $\psi_j(k)$ is supported on $[2^j, 2^{j+2}]$. Hence, substituting this lower bound as λ in (D.1), we get the second upper bound in Lemma 6.2.

To obtain the third upper bound in Lemma 6.2, we first extract more time-decay using integration by parts, and then apply Van der Corput's Lemma. Noting that the domain of integration is

bounded away from k = 0, we have

$$\begin{split} &\int_{\mathbb{R}} e^{-it\sqrt{1+k^2}+ikr}\psi_j(k)\,dk = \int_{\mathbb{R}} \partial_k [e^{-it\sqrt{1+k^2}}]e^{ikr}\frac{1}{-it}\frac{\sqrt{1+k^2}}{k}\psi_j(k)\,dk \\ &= \left[e^{-it\sqrt{1+k^2}}e^{ikr}\frac{1}{-it}\frac{\sqrt{1+k^2}}{k}\psi_j(k)\right]_{k=-\infty}^{k=\infty} + \frac{1}{it}\int_{\mathbb{R}} e^{-it\sqrt{1+k^2}}\partial_k \left[e^{ikr}\frac{\sqrt{1+k^2}}{k}\psi_j(k)\right]\,dk \\ &= \frac{1}{it}\int_{\mathbb{R}} e^{-it\sqrt{1+k^2}+ikr}(ir+\partial_k)\left[\frac{\sqrt{1+k^2}}{k}\psi_j(k)\right]\,dk\,, \end{split}$$

where the boundary terms vanish since ψ_j is compactly supported. Now we can apply Van der Corput Lemma (D.1) on the last integral, with the lower bound on the phase term as before, and get the last upper bound in Lemma 6.2.

D.2. **Proof of Proposition 6.3.** We prove $\mathcal{I}_j^1, \mathcal{I}_j^2, \mathcal{I}_j^3 \leq C2^{-(\frac{3}{2}+\varepsilon)j} \langle x \rangle \langle y \rangle$, for some C > 0. Each term L(k) is based on the explicit expressions for the resolvent kernel Appendix C. It requires the definitions of A, B, C, D as they appear in (3.10) (as well as that of $\tilde{B}, \tilde{A}, \tilde{D}, \tilde{C}$) and of $\varphi(k, \tau)$ as defined in (C.7). We find that L(k) is a rational function where both the numerator and the denominator are quadratic in k and $\sqrt{1-k^2}$, and we can write, for some nonzero $a, b, c, d \in \mathbb{C}$:

$$L(k) = \frac{(ak+b\sqrt{1+k^2})(ck+d\sqrt{1+k^2})}{k\varphi(k,\tau)}$$

Note that, using the asymptotics of φ , $\tilde{B}, \tilde{A}, \tilde{D}, \tilde{C}$ as $k \to \infty$, we have that

(D.2)
$$|L(k)| \lesssim \text{const}, \quad |L'(k)|, |L''(k)| \lesssim k^{-3}, \quad \text{as } k \to \infty.$$

We first bound \mathcal{I}_j^1 from above since, as we shall see, it is the largest term (or, the slowest to decay as $k \to \infty$), and will yield the overall upper bound on I_j . Note that by the triangle inequality

(D.3)
$$\mathcal{I}_{j}^{1} \lesssim \int_{[k_{0}2^{j},k_{0}2^{j+1}]} \langle k \rangle^{-3/2-\varepsilon} \left(|x \pm y| \cdot |L'(k)| + |L''(k)| \right) dk$$

(D.4)
$$+ \int_{[k_0 2^j, k_0 2^{j+1}]} \langle k \rangle^{-3/2 - \varepsilon} \left[\left(|x \pm y| \cdot |L(k)| + 2|L'(k)| \right) |\chi'_j(k)| + |L(k)\chi''_j(k)| \right] dk \,.$$

Since $\langle k \rangle^{-\frac{3}{2}-\varepsilon}$ is strictly decreasing, and using the upper bounds in (D.2), we have

$$|(\mathbf{D}.3)| \lesssim 2^{-(\frac{3}{2}+\varepsilon)j} \int_{[k_0 2^j, k_0 2^{j+1}]} k^{-3} dk \langle x \rangle \langle y \rangle,$$

which vanishes as $j \to \infty$, and so (D.3) is uniformly bounded for all $j \in \mathbb{N}$. To bound (D.4), we note that the L'(k) vanishes similarly, and so we only have to consider the $L(k)\chi'_j(k)$ and $L(k)\chi''_j(k)$.

Since χ_j is a smooth identifier function, χ'_j and χ''_j are compactly supported, localized around the edges of $\operatorname{supp}(\chi_j)$. Hence the value of the integral in (D.4) does not depend on the length of $\operatorname{supp}\chi_j$. Since L(k) tends to a constant (see (D.2)), the integral of the terms $|L(k)||\chi'_k(k)|$ and $|L(k)|^2|\chi''_j(x)|$ also tend to a constant as $j \to \infty$. Therefore, (D.4) is uniformly bounded. Hence, we obtained the desired bound for \mathcal{I}_i^1 . The remaining two terms, \mathcal{I}_j^2 and \mathcal{I}_j^3 (see (6.20)), are bounded in a similar manner. In fact, both terms decay even faster, esentially like $k^{-5/2-\varepsilon}$, and so the overall decay of $I_j \leq \sum_{\ell=1,2,3} \mathcal{I}_j^\ell$ is dominated, in high *j* values (high energy), by that of \mathcal{I}_j^1 .

Appendix E. Bounds on $F(x; k, \tau)$ and $G(x; k, \tau)$

Recall the definitions of $F(x; k, \tau)$ in (6.25) and $G(x; k, \tau)$ in (6.35)

$$\begin{split} F(x;k,\tau) &\equiv \int_{\mathbb{R}} f(x,y;k,\tau) \langle k \rangle^{-3/2-\varepsilon} \alpha_0(y) dy ,\\ f(x,y;k,\tau) &\equiv \begin{pmatrix} \xi_{+,1}(x;k,\tau)\xi_{+,2}(y;-k,\tau) & \xi_{+,1}(x;k,\tau)\xi_{+,1}(y;-k,\tau) \\ \xi_{+,2}(x;k,\tau)\xi_{+,2}(y;-k,\tau) & \xi_{+,2}(x;k,\tau)\xi_{+,1}(y;-k,\tau) \end{pmatrix} \\ &\quad + \begin{pmatrix} e^{-i\tau}\xi_{-,1}(x;k,\tau)\xi_{-,2}(y;-k,\tau) & e^{-i\tau}\xi_{-,1}(x;k,\tau)\xi_{-,1}(y;-k,\tau) \\ e^{-i\tau}\xi_{-,2}(x;k,\tau)\xi_{-,2}(y;-k,\tau) & e^{-i\tau}\xi_{-,2}(x;k,\tau)\xi_{-,1}(y;-k,\tau) \end{pmatrix} ,\\ G(x;k,\tau) &\equiv \partial_k \left(\chi(k) \frac{|T(k,\tau)|^2}{k} F(x;k,\tau) \right) . \end{split}$$

Proposition E.1 (Bounds on $\partial_k^j F(x; k, \tau)$ and $\partial_k^j G(x; k, \tau)$). There exist constants A_j , for j = 0, 1, 2 and B_l for l = 0, 1, which depend on k_0 , such that for every $x \in \mathbb{R}$, $k \in (0, 2k_0]$ and $\tau \in (0, \tau)$:

(E.1)
$$|\partial_k^j F(x;k,\tau)| \le A_j \langle x \rangle^j \| \langle y \rangle^j \alpha_0(y) \|_{L^1(\mathbb{R}_y)}$$

(E.2)
$$|\partial_k^l G(x;k,\tau)| \le B_l \frac{\langle x \rangle^{l+1}}{k^l (k^2 + \sin^2(\tau/2))} \|\langle y \rangle^{l+1} \alpha_0(y)\|_{L^1(\mathbb{R}_y)}.$$

Proof. To bound $F(x; k, \tau)$ and its derivatives, we first bound the Jost solutions (see Proposition 4.1) $\xi_{\pm}(x; k, \tau)$ and their derivatives with respect to k. Every coordinate of f can be written as a sum of product of Jost solutions. Thus, $F(x; k, \tau)$ can be written as a finite sum of the form

$$F(x;k,\tau) = \sum_{j} \int_{\mathbb{R}} \phi_{j,1}(x;k,\tau) \phi_{j,2}(y;k,\tau) \langle k \rangle^{-3/2-\varepsilon} \alpha(y) \, dy$$

where $\phi_{j,\ell}(z,k,\tau)$ is an expression of the form

(E.3)
$$\frac{(ak+b\sqrt{1+k^2})(ck+d\sqrt{1+k^2})}{k}e^{\pm ikz},$$

for some real parameters a, b, c, d. Moreover, uniformly in z = x or y, we have $|\phi(z, k, \tau)| \leq \langle k \rangle$ by the explicit form of the resolvent kernel entries (see Appendix C), and it follows that

$$|F(x;k,\tau)| \le A_0 \langle k \rangle^{\frac{1}{2}-\varepsilon} \int_{\mathbb{R}} |\alpha_0(y)| \, dy \, .$$

Since we have constrained k to vary in $[0, 2k_0]$, we have the desired upper bound for $F(x; k, \tau)$.

Turning now to k- derivatives of $F(x; k, \tau)$, we note that $k \mapsto F(x; k, \tau)$ is smooth functions of k and hence for each j, $\partial_k^j F(x; k, \tau)$ is uniformly bounded on for $k \in [0, 2k_0]$. The bounds on $\partial_k^j F(x; k, \tau)$ involve the weights $\langle x \rangle^j$ and $\langle y \rangle^j$, due to the appearance of factors of e^{ikx} and e^{iky} in $\phi_{j,l}(x; k, \tau)$; see (E.3). Hence, $\partial_k^j F(x; k, \tau)$ gives rise, via $\phi_{j,1}(x; k, \tau)\phi_{j,2}(y; k, \tau)$, to terms whose behavior is bounded by $\langle x \rangle^j \langle y \rangle^j$, leading to the asserted bounds for the derivatives of F. We now turn to the bounds on $G(x; k, \tau)$. Expanding the expression, we have

(E.4)
$$G(x;k,\tau) = \frac{k\chi'(k) - \chi(k)}{k^2} |T(k,\tau)|^2 F(x;k,\tau) + \frac{\chi(k)}{k} |T(k,\tau)|^2 \partial_k F(x;k,\tau)$$
(E.5)
$$+ \frac{\chi(k)}{k^2} \partial_k |T(k,\tau)|^2 F(x;k,\tau)$$

(E.5)
$$+ \frac{\chi(k)}{k} \partial_k |T(k,\tau)|^2 F(x;k,\tau).$$

Applying the above bounds for $F(x; k, \tau)$ as well as those for $T(k, \tau)$ in Proposition 5.3, we obtain (E.6) $|G(x; k, \tau)|$

(E.7)
$$\lesssim \frac{1}{k^2} \frac{k^2}{k^2 + \sin^2(\tau/2)} \|\alpha_0\|_{L^1(\mathbb{R})} + \langle x \rangle \|\langle y \rangle \alpha_0(y)\|_{L^1(\mathbb{R}_y)} + \frac{1}{k} \frac{2k}{k^2 + \sin^2(\tau/2)} \|\alpha_0\|_{L^1(\mathbb{R})}$$

(E.8)
$$\lesssim \frac{\langle x \rangle}{k^2 + \sin^2(\tau/2)} \| \langle y \rangle \alpha_0(y) \|_{L^1(\mathbb{R}_y)},$$

with implied constants which are independent of k and τ . The argument for $\partial_k G(x; k, \tau)$ follows similarly using the product rule as well as the bounds on $\partial_k^2 F(x; k, \tau)$ and $\partial_k^2 |T(k, \tau)|^2$. This completes the proof of Proposition E.1.

References

- M. Beceanu. "New estimates for a time-dependent Schrödinger equation". Duke Mathematical Journal 159.3 (2011), pp. 417–477.
- [2] M. Bellec et al. "Non-diffracting states in one-dimensional Floquet photonic topological insulators". *Europhysics Letters* 119.1 (2017), p. 14003.
- [3] P. Brenner. "On scattering and everywhere defined scattering operators for nonlinear Klein-Gordon equations". Journal of differential equations 56.3 (1985), pp. 310–344.
- [4] M. E. Cage et al. *The quantum Hall effect*. Springer Science & Business Media, 2012.
- G. Chen and F. Pusateri. "The 1-dimensional nonlinear Schrödinger equation with a weighted L1 potential". Analysis & PDE 15.4 (2022), pp. 937–982.
- [6] E. A. Coddington and N. Levinson. Theory of ordinary differential equations. Florida: Krieger, 1955.
- [7] P. D'Ancona and L. Fanelli. "Decay estimates for the wave and Dirac equations with a magnetic potential". Communications on Pure and Applied Mathematics 60.3 (2007), pp. 357– 392.
- [8] A. Drouot. "Scattering resonances for highly oscillatory potentials." In: Annales Scientifiques de l'Ecole Normale Supérieure. Vol. 51. 4. 2018.
- [9] A. Drouot. "The bulk-edge correspondence for continuous dislocated systems". 71.3 (2021), pp. 1185–1239.
- [10] A. Drouot, C. L. Fefferman, and M. I. Weinstein. "Defect modes for dislocated periodic media". *Communications in Mathematical Physics* 377.3 (2020), pp. 1637–1680.
- [11] V. Duchene, I. Vukicevic, and M. I. Weinstein. "Scattering and Localization Properties of Highly Oscillatory Potentials". *Communications on Pure and Applied Mathematics* 67.1 (2014), pp. 83–128.
- [12] B. Erdogan, M. Goldberg, and W. R. Green. "The massless Dirac equation in two dimensions: zero-energy obstructions and dispersive estimates". *Journal of Spectral Theory* (2021).

REFERENCES

- [13] B. M. Erdoğan, M. Goldberg, and W. R. Green. "Limiting absorption principle and Strichartz estimates for Dirac operators in two and higher dimensions". *Communications in Mathematical Physics* 367.1 (2019), pp. 241–263.
- [14] M. B. Erdoğan, M. Goldberg, and W. R. Green. "The massless Dirac equation in two dimensions: zero-energy obstructions and dispersive estimates". *Journal of Spectral Theory* 11.3 (2021), pp. 935–979.
- [15] M. B. Erdoğan, W. R. Green, and E. Toprak. "Dispersive estimates for massive Dirac operators in dimension two". Journal of Differential Equations 264.9 (2018), pp. 5802–5837.
- [16] M. B. Erdoğan, W. R. Green, and E. Toprak. "Dispersive estimates for Dirac operators in dimension three with obstructions at threshold energies". *American Journal of Mathematics* 141.5 (2019), pp. 1217–1258.
- [17] M. B. Erdoğan and W. R. Green. "On the one dimensional Dirac equation with potential". Journal de Mathématiques Pures et Appliquées 151 (2021), pp. 132–170.
- [18] M. B. Erdoğan, W. R. Green, and E. Toprak. "Dispersive estimates for massive Dirac operators in dimension two". *Journal of Differential Equations* 264.9 (2018), pp. 5802–5837.
- [19] C. Fefferman, J. Lee-Thorp, and M. I. Weinstein. Topologically protected states in one-dimensional systems. Vol. 247. 1173. American Mathematical Society, 2017.
- [20] C. L. Fefferman, J. P. Lee-Thorp, and M. I. Weinstein. "Topologically protected states in one-dimensional continuous systems and Dirac points". *Proceedings of the National Academy* of Sciences 111.24 (2014), pp. 8759–8763.
- [21] A. Galtbayar, A. Jensen, and K. Yajima. "Local time-decay of solutions to Schrödinger equations with time-periodic potentials". *Journal of Statistical Physics* 116.1-4 (2004), pp. 231– 282.
- [22] B. C. Hall. Quantum theory for mathematicians. Springer, 2013.
- [23] S. N. Hameedi, A. Sagiv, and M. I. Weinstein. "Radiative decay of edge states in Floquet media". arXiv preprint arXiv:2201.11219, to appear in Multiscale Modeling and Simulations.
- [24] M. Z. Hasan and C. L. Kane. "Colloquium: topological insulators". Reviews of modern physics 82.4 (2010), p. 3045.
- [25] A. Jensen and T. Kato. "Spectral properties of Schrödinger operators and time-decay of the wave functions". Duke Math. J. 46.1 (1979), pp. 583–611.
- [26] L. Jezequel and P. Delplace. "Nonlinear edge modes from topological one-dimensional lattices". Phys. Rev. B 105 (3 2022), p. 035410.
- [27] M. Jürgensen, S. Mukherjee, and M. C. Rechtsman. "Quantized nonlinear Thouless pumping". *Nature* 596.7870 (2021), pp. 63–67.
- [28] K. v. Klitzing, G. Dorda, and M. Pepper. "New method for high-accuracy determination of the fine-structure constant based on quantized Hall resistance". *Physical review letters* 45.6 (1980), p. 494.
- [29] A. Komech and E. Kopylova. "Weighted energy decay for 1D Klein–Gordon equation". Communications in Partial Differential Equations 35.2 (2010), pp. 353–374.
- [30] E. Kopylova and A. Komech. Dispersion Decay and Scattering Theory. Wiley, 2014.
- [31] H. Kovarik. "Spectral properties and time decay of the wave functions of Pauli and Dirac operators in dimension two". Advances in Mathematics 398 (2022), p. 108244.
- [32] J. P. Lee-Thorp et al. "Photonic realization of topologically protected bound states in domainwall waveguide arrays". *Phys. Rev. A* 93 (2016), p. 033822.
- [33] T. Ozawa et al. "Topological photonics". Reviews of Modern Physics 91.1 (2019), p. 015006.

REFERENCES

- [34] D. E. Pelinovsky and A. Stefanov. "Asymptotic stability of small gap solitons in nonlinear Dirac equations". Journal of mathematical physics 53.7 (2012).
- [35] M. Reed and B. Simon. Scattering theory. Methods of modern mathematical physics, Vol. III. 1979.
- [36] W. Schlag and J. Luermann. "On codimension one stability of the soliton for the 1D focusing cubic Klein-Gordon equation". https://arxiv.org/abs/2302.05273 (2023).
- [37] W. Schlag. "Dispersive estimates for Schrödinger operators: a survey". Mathematical aspects of nonlinear dispersive equations 163 (2005), pp. 255–285.
- [38] A. Soffer and M. Weinstein. "Resonance and Radiation damping..." Inventiones Math. (1999).
- [39] A. Soffer and M. I. Weinstein. "Nonautonomous Hamiltonians". Journal of statistical physics 93 (1998), pp. 359–391.
- [40] A. Soffer and X. Wu. "L^p Boundedness of the Scattering Wave Operators of Schroedinger Dynamics with Time-dependent Potentials and applications". arXiv preprint arXiv:2012.14356 (2020).
- [41] E. M. Stein and T. S. Murphy. Harmonic analysis: real-variable methods, orthogonality, and oscillatory integrals. Vol. 3. Princeton University Press, 1993.
- [42] G. Teschl. Mathematical methods in quantum mechanics. Vol. 157. American Mathematical Soc., 2014.
- [43] D. J. Thouless et al. "Quantized Hall conductance in a two-dimensional periodic potential". *Physical review letters* 49.6 (1982), p. 405.
- [44] N. Waterstraat. "Fredholm operators and spectral flow". arXiv preprint arXiv:1603.02009 (2016).
- [45] R. Weder. "Inverse scattering on the line for the nonlinear Klein–Gordon equation with a potential". *Journal of mathematical analysis and applications* 252.1 (2000), pp. 102–123.
- [46] M. I. Weinstein. "Localized States and Dynamics in the Nonlinear Schrödinger/Gross-Pitaevskii Equation". In: *Dynamics of Partial Differential Equations*. Cham: Springer International Publishing, 2015, pp. 41–79.
- [47] H. Xue, Y. Yang, and B. Zhang. "Topological acoustics". Nature Reviews Materials (2022), pp. 1–17.
- [48] D. Yafaev. Mathematical scattering theory. General theory. 1992.
- [49] K. Yajima. "The W^{k,p}-continuity of wave operators for Schrödinger operators". Journal of the Mathematical Society of Japan 47.3 (1995), pp. 551–581.

Department of Mathematics and Statistics, Amherst College, 31 Quadrangle Dr, Amherst, MA 01002, USA

Email address: jkraisler@amherst.edu

FACULTY OF MATHEMATICS, TECHNION - ISRAEL INSTITUTE OF TECHNOLOGY, HAIFA 32000, ISRAEL *Email address*: mmirsagiv@technion.ac.il

DEPARTMENT OF APPLIED PHYSICS AND APPLIED MATHEMATICS AND DEPARTMENT OF MATHEMATICS, COLUMBIA UNIVERSITY, 500 W120 STREET, NEW YORK, NY 10027, USA

Email address: miw2103@columbia.edu

34