

Riemann-Stieltjes operators on some general spaces on the unit ball and their essential norm

Y. Estaremi and M. Hassanlou

ABSTRACT. The objective of this paper is to find an estimation of the essential norm of Riemann-Stieltjes operator from the spaces $F(p, q, s)$ (general function space) and $H(p, q, \phi)$ (mixed-norm space) into Zygmund-type space in the unit ball of \mathbb{C}^n .

1. Introduction and Preliminaries

Let \mathbb{B}_n be the unit ball in \mathbb{C}^n and $H(\mathbb{B}_n)$ be the class of all holomorphic functions on \mathbb{B}_n . For $f \in H(\mathbb{B}_n)$, with the Taylor expansion $f(z) = \sum_{|\beta| \geq 0} a_\beta z^\beta$, let $\mathcal{R}f(z) = \sum_{|\beta| \geq 0} |\beta| a_\beta z^\beta$ be the radial derivative of f . Here $z = (z_1, z_2, \dots, z_n) \in \mathbb{B}_n$, $\beta = (\beta_1, \beta_2, \dots, \beta_n)$ is a multi-index, $z^\beta = z_1^{\beta_1} \cdots z_n^{\beta_n}$ and $|\beta| = \beta_1 + \beta_2 + \cdots + \beta_n$. It is easy to see that

$$\mathcal{R}f(z) = \sum_{j=1}^n z_j \frac{\partial f}{\partial z_j}(z).$$

For more information one can see [21]. The iterates of \mathcal{R} is defined by $\mathcal{R}^m f(z) = \mathcal{R}(\mathcal{R}^{m-1} f(z))$, for $m \in \mathbb{N} \setminus \{1\}$.

Let $0 < p < \infty$, $0 \leq s < \infty$ and $-n - 1 < q < \infty$. The space $F(p, q, s)$ is the space of all analytic functions in \mathbb{B}_n for which

$$\|f\|_{F(p,q,s)}^p = |f(0)|^p + \sup_{a \in \mathbb{B}_n} \int_{\mathbb{B}_n} |\mathcal{R}f(z)|^p (1 - |z|^2)^q g^s(z, a) dV(z) < \infty.$$

Here dV is normalized area measure on \mathbb{B}_n , $g(z, a) = \log \left| \frac{1}{\varphi_a(z)} \right|$, $\varphi_a(z) = \frac{a-z}{1-\bar{a}z}$. Using complex gradient ∇ and complex invariant gradient $\tilde{\nabla}$, there are some equivalent norms for the functions in this space, see [19] for more details. The above space, introduced in [17], is called general function space which include some classes of

2020 *Mathematics Subject Classification.* 47B38, 30H30.

Key words and phrases. Riemann-Stieltjes operator, Essential norm, Zygmund type space.

* Corresponding author.

functions like Bergman, Bloch, Hardy, BMOA and Q_p spaces. See [17] for more details.

Beside the general space $F(p, q, s)$, we use another general class of analytic functions $H(p, q, \phi)$ which is called mixed-norm space.

The function ϕ on $[0, 1)$ is called normal if it is a positive continuous function for which there are two constants $b > a > 0$ such that

- (i) $\frac{\phi(r)}{(1-r)^a}$ is non-increasing in $[0, 1)$ and $\frac{\phi(r)}{(1-r)^a} \downarrow 0$
- (ii) $\frac{\phi(r)}{(1-r)^b}$ is non-decreasing in $[0, 1)$ and $\frac{\phi(r)}{(1-r)^b} \uparrow \infty$

as $r \rightarrow 1^-$. By a normal function $\phi : \mathbb{B}_n \rightarrow [0, \infty)$ we mean that $\phi(|z|)$ is normal and also ϕ is radial, $\phi(z) = \phi(|z|)$.

For $0 < p, q < \infty$ and normal function ϕ on $[0, 1)$, the mixed-norm space $H(p, q, \phi)$ consists of all analytic functions in \mathbb{B}_n for which

$$\|f\|_{p,q,\phi}^p = \int_0^1 M_q^p(f, r) \frac{\phi^p(r)}{1-r} dr < \infty$$

where

$$M_q(f, r) = \left(\frac{1}{2\pi} \int_0^{2\pi} |f(re^{i\theta})|^q d\theta \right)^{1/q}.$$

If $1 \leq p < \infty$, then $H(p, q, \phi)$ becomes a Banach space equipped with the norm $\|\cdot\|_{p,q,\phi}$. In the case $0 < p < 1$, $\|\cdot\|_{p,q,\phi}$ is a quasi-norm on $H(p, q, \phi)$ and this space is a Fréchet space but is not a Banach space. For $\alpha > -1$, $\phi(r) = (1-r)^{(\alpha+1)/p}$ is a normal function. If $p = q$, then $H(p, p, (1-r)^{(\alpha+1)/p})$ is weighted Bergman space A_α^p which is defined by

$$A_\alpha^p = \{f \in H(\mathbb{B}_n) : \|f\|_{A_\alpha^p}^p = \int_{\mathbb{B}_n} |f(z)|^p (1-|z|^2)^\alpha dV(z) < \infty\}.$$

Let μ be a positive continuous function on $[0, 1)$, such a function is called weight. The Bloch-type space \mathcal{B}_μ and Zygmund-type space \mathcal{Z}_μ are defined as

$$\begin{aligned} \mathcal{B}_\mu &= \{f \in H(\mathbb{B}_n) : \|f\|_{\mathcal{B}_\mu} = |f(0)| + \sup_{z \in \mathbb{B}_n} \mu(|z|) |\mathcal{R}f(z)| < \infty\} \\ \mathcal{Z}_\mu &= \{f \in H(\mathbb{B}_n) : \|f\|_{\mathcal{Z}_\mu} = |f(0)| + \sup_{z \in \mathbb{B}_n} \mu(|z|) |\mathcal{R}^2 f(z)| < \infty\}. \end{aligned}$$

When $\mu(r) = (1-r^2)$ then we have classical Bloch and Zygmund space, \mathcal{B} and \mathcal{Z} . If $\mu(r) = (1-r^2)^\alpha$, $\alpha > 0$, then α -Bloch and Zygmund spaces are obtained, \mathcal{B}^α and \mathcal{Z}^α . Let $g \in H(\mathbb{B}_n)$. The Riemann-Stieltjes operator L_g is defined by

$$L_g f(z) = \int_0^1 \mathcal{R}f(tz) g(tz) \frac{dt}{t}, \quad f \in H(\mathbb{B}_n), \quad z \in \mathbb{B}_n.$$

Bounded and compact Riemann-Stieltjes operator from general space of analytic functions to Zygmund-type spaces and little Zygmund-type spaces on the unit ball

are characterized in [10]. Liu and Yu have investigated boundedness and compactness of L_g from mixed-norm spaces to Zygmund-type spaces in [9]. Essential norm of Riemann-Stieltjes operator on weighted Bergman spaces with doubling weights is estimated in [1]. Volterra-type operators from $F(p, q, s)$ space to Bloch-Orlicz and Zygmund-Orlicz spaces are investigated in [6]. For more results on integral operators or generalized integral operators on some spaces of analytic functions one can see [3–5, 7, 8, 11–13, 15–18, 20] and references therein.

Motivated by [9, 10], we are going to find an estimation of the essential norm of the Riemann-Stieltjes operator from the general space of analytic functions, $F(p, q, s)$, and mixed-norm space, $H(p, q, \phi)$, into Zygmund-type spaces in the unit ball.

In this paper, for real scalars A and B , the notation $A \preceq B$ means $A \leq CB$ for some positive constant C . Also, the notation $A \approx B$ means $A \preceq B$ and $B \preceq A$. The essential norm of an operator T , $\|T\|_e$, is the distance of T from the space of all compact operators.

2. Main results

In this section, first we recall some basic lemmas from the literature for later use in the paper. Then we estimate the essential norm of Riemann-Stieltjes operator L_g from the general function space and mixed-norm space into Zygmund-type space in the unit ball of \mathbb{C}^n .

Lemma 2.1. [14] *For every $f, g \in H(\mathbb{B}_n)$,*

$$\mathcal{R}L_g(f)(z) = \mathcal{R}f(z)g(z) \quad z \in \mathbb{B}_n.$$

Lemma 2.2. [18] *Let $0 < p, s < \infty$, $\max\{-n - 1, -s - 1\} < q < \infty$ and $f \in F(p, q, s)$. Then $f \in \mathcal{B}^{\frac{n+1+q}{p}}$ and*

$$\|f\|_{\mathcal{B}^{\frac{n+1+q}{p}}} \leq C\|f\|_{F(p,q,s)}.$$

According to the Proposition 2.1 of [2], for $\alpha > 0$ and $m \in \mathbb{N}$, we have $f \in \mathcal{B}^\alpha$ if and only if

$$\sup_{z \in \mathbb{B}_n} |\mathcal{R}^m f(z)|(1 - |z|^2)^{\alpha+m-1} < \infty. \quad (1)$$

Lemma 2.3. [15] *Assume that $m \in \mathbb{N}$, $p, q \in (0, \infty)$, ϕ is normal and $f \in H(p, q, \phi)$. Then there exists a positive constant C independent of f such that*

$$|\mathcal{R}^m f(z)| \leq \frac{C|z|}{\phi(|z|)(1 - |z|^2)^{m+n/q}} \|f\|_{p,q,\phi}, \quad z \in \mathbb{B}_n.$$

Lemma 2.4. [3] *Let $0 < p, s < \infty$, $\max\{-n - 1, -s - 1\} < q < \infty$, μ be a weight and $g \in H(\mathbb{B}_n)$. Then $L_g : F(p, q, s) \rightarrow \mathcal{Z}_\mu$ is compact if and only if $L_g : F(p, q, s) \rightarrow \mathcal{Z}_\mu$ is bounded and for any bounded sequence $\{f_k\}$ in $F(p, q, s)$,*

which converges uniformly to zero on compact subsets of \mathbb{B}_n , as $k \rightarrow \infty$, we have $\|L_g f_k\|_{\mathcal{Z}_\mu} \rightarrow 0$, as $k \rightarrow \infty$.

Lemma 2.5. [4] *Let $0 < p, q < \infty$, μ be a weight, ϕ be a normal function and $g \in H(\mathbb{B}_n)$. Then $L_g : H(p, q, \phi) \rightarrow \mathcal{Z}_\mu$ is compact if and only if $L_g : H(p, q, \phi) \rightarrow \mathcal{Z}_\mu$ is bounded and for any bounded sequence $\{f_k\}$ in $H(p, q, \phi)$ which converges uniformly to zero on compact subsets of \mathbb{B}_n as $k \rightarrow \infty$, we have $\|L_g f_k\|_{\mathcal{Z}_\mu} \rightarrow 0$ as $k \rightarrow \infty$.*

By the above mentioned lemmas, in the next theorem we find an estimation of the essential norm of the Riemann-Stieltjes operator $L_g : F(p, q, s) \rightarrow \mathcal{Z}_\mu$. For simplicity, put $\alpha = \frac{n+1+q}{p}$.

Theorem 2.6. *Let $0 < p, s < \infty$, $\max\{-n-1, -s-1\} < q < \infty$, μ be a weight and $g \in H(\mathbb{B}_n)$. If $L_g : F(p, q, s) \rightarrow \mathcal{Z}_\mu$ is bounded, then*

$$\|L_g\|_{e, F(p, q, s) \rightarrow \mathcal{Z}_\mu} \approx \max \left\{ \limsup_{|z| \rightarrow 1} \frac{\mu(|z|)|g(z)|}{(1-|z|^2)^{\alpha+1}}, \limsup_{|z| \rightarrow 1} \frac{\mu(|z|)|\mathcal{R}g(z)|}{(1-|z|^2)^\alpha} \right\}.$$

PROOF. Since L_g is bounded, then there exists $C > 0$ such that

$$\|L_g f\|_{\mathcal{Z}_\mu} \leq C \|f\|_{F(p, q, s)}, \quad \text{for all } f \in F(p, q, s).$$

By applying this inequality to $f_1(z) = z$, we obtain

$$\sup_{z \in \mathbb{B}_n} \mu(|z|)|\mathcal{R}g(z)| < \infty. \quad (2)$$

Also, by applying the inequality to $f_2(z) = z^2$, we have

$$\sup_{z \in \mathbb{B}_n} \mu(|z|)|g(z)| < \infty. \quad (3)$$

Let $\{z_i\}_{i \in \mathbb{N}}$ be a sequence in \mathbb{D} , such that $|z_i| \rightarrow 1$. We define the sequence of functions as follows

$$f_i(z) = \frac{(1-|z_i|^2)^{\alpha+1}}{(1-\bar{z}_i z)^{2\alpha}} - 2 \frac{1-|z_i|^2}{(1-\bar{z}_i z)^\alpha}, \quad i \in \mathbb{N}.$$

Direct calculations shows that for every $i \in \mathbb{N}$, $f_i \in F(p, q, s)$, $\sup_i \|f_i\|_{F(p, q, s)} \leq C$ and $f_i \rightarrow 0$, uniformly on compact subsets of \mathbb{B}_n . Also,

$$\mathcal{R}f_i(z_i) = 0 \quad \text{and} \quad \mathcal{R}^2 f_i(z_i) = 2\alpha^2 \frac{|z_i|^4}{(1-|z_i|^2)^{\alpha+1}}.$$

So for any compact operator $K : F(p, q, s) \rightarrow \mathcal{Z}_\mu$ we have

$$\lim_{i \rightarrow \infty} \|K f_i\|_{\mathcal{Z}_\mu} = 0.$$

By the above observations we have

$$\begin{aligned}
\|L_g - K\|_{F(p,q,s) \rightarrow \mathcal{Z}_\mu} &\succeq \limsup_{i \rightarrow \infty} \|L_g f_i - K f_i\|_{\mathcal{Z}_\mu} \\
&\geq \limsup_{i \rightarrow \infty} \|L_g f_i\|_{\mathcal{Z}_\mu} - \limsup_{i \rightarrow \infty} \|K f_i\|_{\mathcal{Z}_\mu} \\
&= \limsup_{i \rightarrow \infty} \mu(|z_i|) |\mathcal{R}^2(L_g f_i)(z_j)| \\
&\succeq \limsup_{i \rightarrow \infty} \frac{\mu(|z_i|) |g(z_i)| |z_i|^4}{(1 - |z_i|^2)^{\alpha+1}}.
\end{aligned}$$

Therefore,

$$\|L_g\|_{e, F(p,q,s) \rightarrow \mathcal{Z}_\mu} = \inf_K \|L_g - K\|_{F(p,q,s) \rightarrow \mathcal{Z}_\mu} \succeq \limsup_{|z| \rightarrow 1} \frac{\mu(|z|) |g(z)|}{(1 - |z|^2)^{\alpha+1}}. \quad (4)$$

Now we define the sequence $\{h_i\}$ as follows

$$h_i(z) = \frac{(1 - |z_i|^2)^{\alpha+1}}{(1 - \bar{z}_i z)^{2\alpha}} - \frac{2 + 4\alpha}{\alpha + 1} \frac{1 - |z_i|^2}{(1 - \bar{z}_i z)^\alpha}.$$

Then $\{h_j\}$ is a bounded sequence in $F(p, q, s)$ which converges to 0 uniformly on compact subsets of \mathbb{B}_n . Also

$$\mathcal{R}h_i(z_i) = \mathcal{R}^2 h_i(z_i) = -\frac{2\alpha^2}{\alpha + 1} \frac{|z_j|^2}{(1 - |z_i|^2)^\alpha}.$$

For any compact operator $K : F(p, q, s) \rightarrow \mathcal{Z}_\mu$, using the definition of the operator norm and the norm in Zygmund-type space and (4), we have

$$\begin{aligned}
\|L_g - K\|_{F(p,q,s) \rightarrow \mathcal{Z}_\mu} &\succeq \limsup_{i \rightarrow \infty} \|L_g h_i - K h_i\|_{\mathcal{Z}_\mu} \\
&\geq \limsup_{i \rightarrow \infty} \|L_g h_i\|_{\mathcal{Z}_\mu} - \limsup_{i \rightarrow \infty} \|K h_i\|_{\mathcal{Z}_\mu} \\
&= \limsup_{i \rightarrow \infty} \mu(|z_i|) |\mathcal{R}^2(L_g h_i)(z_i)| \\
&= \limsup_{i \rightarrow \infty} \mu(|z_i|) |\mathcal{R}^2 h_i(z_i) g(z_i) + \mathcal{R}h_i(z_i) \mathcal{R}g(z_i)| \\
&= \limsup_{i \rightarrow \infty} \mu(|z_i|) \left| -\frac{2\alpha^2}{\alpha + 1} \frac{g(z_i) |z_i|^2}{(1 - |z_i|^2)^\alpha} - \frac{2\alpha^2}{\alpha + 1} \frac{\mathcal{R}g(z_i) |z_i|^2}{(1 - |z_i|^2)^\alpha} \right| \\
&\geq \limsup_{i \rightarrow \infty} \frac{2\alpha^2}{\alpha + 1} \frac{\mu(|z_i|) |\mathcal{R}g(z_i)| |z_i|^2}{(1 - |z_i|^2)^\alpha} - \limsup_{i \rightarrow \infty} \frac{2\alpha^2}{\alpha + 1} \frac{\mu(|z_i|) |g(z_i)| |z_i|^2}{(1 - |z_i|^2)^\alpha} \\
&= \limsup_{|z| \rightarrow 1} \frac{\mu(|z|) |\mathcal{R}g(z)|}{(1 - |z|^2)^\alpha} - \limsup_{|z| \rightarrow 1} \frac{\mu(|z|) |g(z)|}{(1 - |z|^2)^\alpha} \\
&\geq \limsup_{|z| \rightarrow 1} \frac{\mu(|z|) |\mathcal{R}g(z)|}{(1 - |z|^2)^\alpha} - \limsup_{|z| \rightarrow 1} \frac{\mu(|z|) |g(z)|}{(1 - |z|^2)^{\alpha+1}} \\
&\succeq \limsup_{|z| \rightarrow 1} \frac{\mu(|z|) |\mathcal{R}g(z)|}{(1 - |z|^2)^\alpha} - \|L_g\|_{e, F(p,q,s) \rightarrow \mathcal{Z}_\mu}.
\end{aligned}$$

Hence

$$\begin{aligned} \|L_g\|_{e, F(p, q, s) \rightarrow \mathcal{Z}_\mu} &= \inf_K \|L_g - K\|_{F(p, q, s) \rightarrow \mathcal{Z}_\mu} \\ &\succeq \limsup_{|z| \rightarrow 1} \frac{\mu(|z|)|\mathcal{R}g(z)|}{(1 - |z|^2)^\alpha} - \|L_g\|_{e, F(p, q, s) \rightarrow \mathcal{Z}_\mu}, \end{aligned}$$

and so

$$\|L_g\|_{e, F(p, q, s) \rightarrow \mathcal{Z}_\mu} \succeq \limsup_{|z| \rightarrow 1} \frac{\mu(|z|)|\mathcal{R}g(z)|}{(1 - |z|^2)^\alpha}. \quad (5)$$

From (4) and (5), the lower estimate of the essential norm is achieved.

For the upper estimate, let $\{r_i\} \subset (0, 1)$ be a sequence such that $r_i \rightarrow 1$, as $i \rightarrow \infty$. Then as $f_{r_i} \rightarrow f$ uniformly on compact subsets of \mathbb{B}_n , as $i \rightarrow \infty$, in which $f_r(z) = f(rz)$. Define the operators K_r on $F(p, q, s)$, $K_r f(z) = f_r(z)$, where $0 < r < 1$. Then it is clear that K_r is a compact operator on $F(p, q, s)$. Hence for any positive integer i , the operator $L_g K_{r_i} : F(p, q, s) \rightarrow \mathcal{Z}_\mu$ is compact and so

$$\|L_g\|_{e, F(p, q, s) \rightarrow \mathcal{Z}_\mu} \leq \limsup_{i \rightarrow \infty} \|L_g - L_g K_{r_i}\|_{F(p, q, s) \rightarrow \mathcal{Z}_\mu}. \quad (6)$$

Therefore for every $f \in F(p, q, s)$, with $\|f\|_{F(p, q, s)} \leq 1$ we have

$$\begin{aligned} \|L_g f - L_g K_{r_i} f\|_{\mathcal{Z}_\mu} &= \sup_{z \in \mathbb{B}_n} \mu(|z|) |\mathcal{R}^2(L_g f)(z) - \mathcal{R}^2(L_g K_{r_i} f)(z)| \\ &= \sup_{z \in \mathbb{B}_n} \mu(|z|) |\mathcal{R}^2 f(z)g(z) + \mathcal{R}f(z)\mathcal{R}g(z) \\ &\quad - r_i \mathcal{R}^2 f_{r_i}(z)g(z) - \mathcal{R}f_{r_i}(z)\mathcal{R}g(z)| \\ &\leq \sup_{z \in \mathbb{B}_n} \mu(|z|) |\mathcal{R}^2 f(z) - r_i \mathcal{R}^2 f_{r_i}(z)| |g(z)| \\ &\quad + \sup_{z \in \mathbb{B}_n} \mu(|z|) |\mathcal{R}f(z) - \mathcal{R}f_{r_i}(z)| |\mathcal{R}g(z)| \\ &= \sup_{|z| \leq \delta} \mu(|z|) |\mathcal{R}^2 f(z) - r_i \mathcal{R}^2 f_{r_i}(z)| |g(z)| \\ &\quad + \sup_{|z| > \delta} \mu(|z|) |\mathcal{R}^2 f(z) - r_i \mathcal{R}^2 f_{r_i}(z)| |g(z)| \\ &\quad + \sup_{|z| \leq \delta} \mu(|z|) |\mathcal{R}f(z) - \mathcal{R}f_{r_i}(z)| |\mathcal{R}g(z)| \\ &\quad + \sup_{|z| > \delta} \mu(|z|) |\mathcal{R}f(z) - \mathcal{R}f_{r_i}(z)| |\mathcal{R}g(z)|, \end{aligned}$$

where $\delta \in (0, 1)$ is fixed. By Weierstrass theorem, $\mathcal{R}f_{r_i} \rightarrow \mathcal{R}f$ and $\mathcal{R}^2 f_{r_i} \rightarrow \mathcal{R}^2 f$ uniformly on compact subsets of \mathbb{B}_n , as $i \rightarrow \infty$. Since the ball $\{z : |z| \leq \delta\}$ is a compact subset of \mathbb{B}_n , then we have

$$\begin{aligned} \limsup_{i \rightarrow \infty} \sup_{|z| \leq \delta} \mu(|z|) |\mathcal{R}^2 f(z) - r_i \mathcal{R}^2 f_{r_i}(z)| |g(z)| &= 0 \\ \limsup_{i \rightarrow \infty} \sup_{|z| \leq \delta} \mu(|z|) |\mathcal{R} f(z) - \mathcal{R} f_{r_i}(z)| |\mathcal{R} g(z)| &= 0. \end{aligned}$$

Therefore using Lemma 2.2 and (1) we have

$$\begin{aligned} \|L_g f - L_g K_{r_i} f\|_{\mathcal{Z}_\mu} &\leq \sup_{|z| > \delta} \mu(|z|) |\mathcal{R}^2 f(z) - r_i \mathcal{R}^2 f_{r_i}(z)| |g(z)| \\ &\quad + \sup_{|z| > \delta} \mu(|z|) |\mathcal{R} f(z) - \mathcal{R} f_{r_i}(z)| |\mathcal{R} g(z)| \\ &\preceq \sup_{|z| > \delta} \frac{\mu(|z|) |g(z)|}{(1 - |z|^2)^{\alpha+1}} + \sup_{|z| > \delta} \frac{r_i \mu(|z|) |g(z)|}{(1 - |r_i z|^2)^{\alpha+1}} \\ &\quad + \sup_{|z| > \delta} \frac{\mu(|z|) |\mathcal{R} g(z)|}{(1 - |z|^2)^\alpha} \|f\|_{\mathcal{B}^\alpha} + \sup_{|z| > \delta} \frac{r_i \mu(|z|) |\mathcal{R} g(z)|}{(1 - |r_i z|^2)^\alpha} \|f\|_{\mathcal{B}^\alpha} \\ &\leq \sup_{|z| > \delta} \frac{\mu(|z|) |g(z)|}{(1 - |z|^2)^{\alpha+1}} + \sup_{|z| > \delta} \frac{r_i \mu(|z|) |g(z)|}{(1 - |r_i z|^2)^{\alpha+1}} \\ &\quad + \sup_{|z| > \delta} \frac{\mu(|z|) |\mathcal{R} g(z)|}{(1 - |z|^2)^\alpha} \|f\|_{F(p,q,s)} + \sup_{|z| > \delta} \frac{r_i \mu(|z|) |\mathcal{R} g(z)|}{(1 - |r_i z|^2)^\alpha} \|f\|_{F(p,q,s)}, \end{aligned}$$

letting $i \rightarrow \infty$, we obtain

$$\|L_g f - L_g K_{r_i} f\|_{\mathcal{Z}_\mu} \preceq 2 \sup_{|z| > \delta} \frac{\mu(|z|) |g(z)|}{(1 - |z|^2)^{\alpha+1}} + 2 \sup_{|z| > \delta} \frac{\mu(|z|) |\mathcal{R} g(z)|}{(1 - |z|^2)^\alpha}.$$

When $\delta \rightarrow 1$, from (6) we have

$$\|L_g\|_{e, F(p,q,s) \rightarrow \mathcal{Z}_\mu} \preceq \limsup_{|z| \rightarrow 1} \frac{\mu(|z|) |g(z)|}{(1 - |z|^2)^{\alpha+1}} + \limsup_{|z| \rightarrow 1} \frac{\mu(|z|) |\mathcal{R} g(z)|}{(1 - |z|^2)^\alpha},$$

and consequently we have an upper estimate of the essential norm. \square

In the sequel we find an estimate for the essential norm of the Riemann-Stieltjes operator $L_g : H(p, q, \phi) \rightarrow \mathcal{Z}_\mu$.

Theorem 2.7. *Let $0 < p, q < \infty$, μ be a weight, ϕ is a normal function and $g \in H(\mathbb{B}_n)$. If $L_g : H(p, q, \phi) \rightarrow \mathcal{Z}_\mu$ is bounded, then*

$$\|L_g\|_{e, H(p,q,\phi) \rightarrow \mathcal{Z}_\mu} \approx \max \left\{ \limsup_{|z| \rightarrow 1} \frac{\mu(|z|) |g(z)|}{\phi(|z|) (1 - |z|^2)^{2 + \frac{n}{q}}}, \limsup_{|z| \rightarrow 1} \frac{\mu(|z|) |\mathcal{R} g(z)|}{\phi(|z|) (1 - |z|^2)^{1 + \frac{n}{q}}} \right\}.$$

PROOF. Let $\{z_i\}_{i \in \mathbb{N}}$ be a sequence in \mathbb{D} such that $|z_i| \rightarrow 1$. Define the sequence of functions as follows

$$f_i(z) = (\gamma + 1) \frac{(1 - |z_i|^2)^{b+1}}{\phi(|z_i|) (1 - \bar{z}_i z)^\beta} - \gamma \frac{(1 - |z_i|^2)^{b+2}}{\phi(|z_i|) (1 - \bar{z}_i z)^{\beta+1}}, \quad 1 \in \mathbb{N},$$

where $\gamma = b + 1 + \frac{n}{q}$ and b comes from the definition of the normal function ϕ . It is clear that for each $i \in \mathbb{N}$, $f_i \in H(p, q, \phi)$, $\sup_j \|f_i\|_{p, q, \phi} \leq C$ and $f_i \rightarrow 0$ uniformly on compact subsets of \mathbb{B}_n . Also

$$\mathcal{R}f_i(z_i) = 0, \quad \mathcal{R}^2 f_i(z_i) = -\gamma(\gamma + 1) \frac{|z_i|^4}{\phi(|z_i|)(1 - |z_i|^2)^{2 + \frac{n}{q}}} \quad i \in \mathbb{N}.$$

Hence for any compact operator $K : H(p, q, \phi) \rightarrow \mathcal{Z}_\mu$ we have $\lim_{n \rightarrow \infty} \|Kf_i\|_{\mathcal{Z}_\mu} = 0$ and so

$$\begin{aligned} \|L_g - K\|_{H(p, q, \phi) \rightarrow \mathcal{Z}_\mu} &\succeq \limsup_{i \rightarrow \infty} \|L_g f_i - Kf_i\|_{\mathcal{Z}_\mu} \\ &\succeq \limsup_{i \rightarrow \infty} \|L_g f_i\|_{\mathcal{Z}_\mu} - \limsup_{i \rightarrow \infty} \|Kf_i\|_{\mathcal{Z}_\mu} \\ &\succeq \limsup_{i \rightarrow \infty} \mu(|z_j|) |\mathcal{R}^2(L_g f_i)(z_i)| \\ &\succeq \limsup_{i \rightarrow \infty} \frac{\mu(|z_j|) |g(z_i)| |z_i|^4}{\phi(|z_i|)(1 - |z_i|^2)^{2 + \frac{n}{q}}}. \end{aligned}$$

Therefore we get that

$$\|L_g\|_{e, H(p, q, \phi) \rightarrow \mathcal{Z}_\mu} = \inf_K \|L_g - K\|_{H(p, q, \phi) \rightarrow \mathcal{Z}_\mu} \succeq \limsup_{|z| \rightarrow 1} \frac{\mu(|z|) |g(z)|}{\phi(|z|)(1 - |z|^2)^{2 + \frac{n}{q}}}. \quad (7)$$

Now we define the sequence $\{h_j\}$ as follows

$$h_i(z) = (\beta + 2) \frac{(1 - |z_i|^2)^{b+1}}{\phi(|z_i|)(1 - \bar{z}_i z)^\beta} - \beta \frac{(1 - |z_i|^2)^{b+2}}{(1 - \bar{z}_i z)^{\beta+1}}.$$

It is easy to see that $\{h_i\}$ is a bounded sequence in $H(p, q, \phi)$, which converges to 0, uniformly on compact subsets of \mathbb{B}_n . Also

$$\mathcal{R}h_i(z_i) = \mathcal{R}^2 h_i(z_i) = \beta \frac{|z_i|^2}{\phi(|z_i|)(1 - |z_i|^2)^{1 + \frac{n}{q}}}.$$

Using (7) we get that

$$-\limsup_{|z| \rightarrow 1} \frac{\mu(|z|) |g(z)|}{\phi(|z|)(1 - |z|^2)^{1 + \frac{n}{q}}} \geq -\limsup_{|z| \rightarrow 1} \frac{\mu(|z|) |g(z)|}{\phi(|z|)(1 - |z|^2)^{2 + \frac{n}{q}}} \succeq -\|L_g\|_{e, H(p, q, \phi) \rightarrow \mathcal{Z}_\mu},$$

and then for any compact operator $K : H(p, q, \phi) \rightarrow \mathcal{Z}_\mu$ we have

$$\begin{aligned}
\|L_g - K\|_{H(p,q,\phi) \rightarrow \mathcal{Z}_\mu} &\succeq \limsup_{i \rightarrow \infty} \|L_g h_i - K h_i\|_{\mathcal{Z}_\mu} \\
&\geq \limsup_{i \rightarrow \infty} \|L_g h_i\|_{\mathcal{Z}_\mu} - \limsup_{i \rightarrow \infty} \|K h_i\|_{\mathcal{Z}_\mu} \\
&= \limsup_{i \rightarrow \infty} \mu(|z_i|) |\mathcal{R}^2(L_g h_i)(z_j)| \\
&= \limsup_{j \rightarrow \infty} \mu(|z_j|) |\mathcal{R}^2 h_i(z_i) g(z_j) + \mathcal{R} h_j(z_j) \mathcal{R} g(z_j)| \\
&= \limsup_{j \rightarrow \infty} \mu(|z_i|) \left| \gamma \frac{g(z_j) |z_j|^2}{\phi(|z_i|)(1 - |z_i|^2)^{1 + \frac{n}{q}}} + \gamma \frac{\mathcal{R} g(z_j) |z_j|^2}{\phi(|z_i|)(1 - |z_j|^2)^{1 + \frac{n}{q}}} \right| \\
&\geq \limsup_{i \rightarrow \infty} \gamma \frac{\mu(|z_i|) |\mathcal{R} g(z_i)| |z_i|^2}{\phi(|z_i|)(1 - |z_i|^2)^{1 + \frac{n}{q}}} - \limsup_{i \rightarrow \infty} \gamma \frac{\mu(|z_i|) |g(z_i)| |z_i|^2}{\phi(|z_i|)(1 - |z_i|^2)^{1 + \frac{n}{q}}} \\
&\succeq \limsup_{|z| \rightarrow 1} \frac{\mu(|z|) |\mathcal{R} g(z)|}{\phi(|z|)(1 - |z|^2)^{1 + \frac{n}{q}}} - \limsup_{|z| \rightarrow 1} \frac{\mu(|z|) |g(z)|}{\phi(|z|)(1 - |z|^2)^{1 + \frac{n}{q}}} \\
&\succeq \limsup_{|z| \rightarrow 1} \frac{\mu(|z|) |\mathcal{R} g(z)|}{\phi(|z|)(1 - |z|^2)^{1 + \frac{n}{q}}} - \|L_g\|_{e, H(p,q,\phi) \rightarrow \mathcal{Z}_\mu}.
\end{aligned}$$

Hence

$$\begin{aligned}
\|L_g\|_{e, H(p,q,\phi) \rightarrow \mathcal{Z}_\mu} &= \inf_K \|L_g - K\|_{H(p,q,\phi) \rightarrow \mathcal{Z}_\mu} \\
&\succeq \limsup_{|z| \rightarrow 1} \frac{\mu(|z|) |\mathcal{R} g(z)|}{\phi(|z|)(1 - |z|^2)^{1 + \frac{n}{q}}} - \|L_g\|_{e, H(p,q,\phi) \rightarrow \mathcal{Z}_\mu},
\end{aligned}$$

and so

$$\|L_g\|_{e, H(p,q,\phi) \rightarrow \mathcal{Z}_\mu} \succeq \limsup_{|z| \rightarrow 1} \frac{\mu(|z|) |\mathcal{R} g(z)|}{\phi(|z|)(1 - |z|^2)^{1 + \frac{n}{q}}}. \quad (8)$$

Therefor we have a lower estimate. In order to prove the upper estimate, let $\{r_i\} \subset (0, 1)$ be a sequence such that $r_i \rightarrow 1$ as $i \rightarrow \infty$. Then $f_{r_i} \rightarrow f$ uniformly on compact subsets of \mathbb{B}_n , as $i \rightarrow \infty$. We define the operators T_r on $H(p, q, \phi)$, $T_r f(z) = f_r(z)$, where $0 < r < 1$. Then T_r is a compact operator on $H(p, q, \phi)$. So, for any positive integer i , the operator $L_g T_{r_i} : H(p, q, \phi) \rightarrow \mathcal{Z}_\mu$ is compact. Thus

$$\|L_g\|_{e, H(p,q,\phi) \rightarrow \mathcal{Z}_\mu} \leq \limsup_{j \rightarrow \infty} \|L_g - L_g T_{r_i}\|_{H(p,q,\phi) \rightarrow \mathcal{Z}_\mu}. \quad (9)$$

It will be sufficient to compute $\|L_g - L_g T_{r_i}\|_{H(p,q,\phi) \rightarrow \mathcal{Z}_\mu}$. Let $f \in H(p,q,\phi)$, with $\|f\|_{p,q,\phi} \leq 1$. Then similar to the proof of Theorem 2.6 we have

$$\begin{aligned} \|L_g f - L_g T_{r_i} f\|_{\mathcal{Z}_\mu} &\leq \sup_{|z| \leq \delta} \mu(|z|) |\mathcal{R}^2 f(z) - r_i \mathcal{R}^2 f_{r_i}(z)| |g(z)| \\ &\quad + \sup_{|z| > \delta} \mu(|z|) |\mathcal{R}^2 f(z) - r_i \mathcal{R}^2 f_{r_i}(z)| |g(z)| \\ &\quad + \sup_{|z| \leq \delta} \mu(|z|) |\mathcal{R} f(z) - \mathcal{R} f_{r_i}(z)| |\mathcal{R} g(z)| \\ &\quad + \sup_{|z| > \delta} \mu(|z|) |\mathcal{R} f(z) - \mathcal{R} f_{r_i}(z)| |\mathcal{R} g(z)|, \end{aligned}$$

where $\delta \in (0, 1)$ is fixed. By Weierstrass theorem, $\mathcal{R} f_{r_i} \rightarrow \mathcal{R} f$ and $\mathcal{R}^2 f_{r_i} \rightarrow \mathcal{R}^2 f$ uniformly on compact subsets of \mathbb{B}_n , as $i \rightarrow \infty$. So

$$\begin{aligned} \limsup_{i \rightarrow \infty} \sup_{|z| \leq \delta} \mu(|z|) |\mathcal{R}^2 f(z) - r_i \mathcal{R}^2 f_{r_i}(z)| |g(z)| &= 0 \\ \limsup_{i \rightarrow \infty} \sup_{|z| \leq \delta} \mu(|z|) |\mathcal{R} f(z) - \mathcal{R} f_{r_i}(z)| |\mathcal{R} g(z)| &= 0. \end{aligned}$$

Here we use the fact that $\sup_{z \in \mathbb{B}_n} \mu(|z|) |\mathcal{R} g(z)| < \infty$ and $\sup_{z \in \mathbb{B}_n} \mu(|z|) |g(z)| < \infty$ which is obtained from boundedness of the operator. Using Lemma 2.3 we get that

$$\begin{aligned} \|L_g f - L_g T_{r_i} f\|_{\mathcal{Z}_\mu} &\leq \sup_{|z| > \delta} \mu(|z|) |\mathcal{R}^2 f(z) - r_i \mathcal{R}^2 f_{r_i}(z)| |g(z)| \\ &\quad + \sup_{|z| > \delta} \mu(|z|) |\mathcal{R} f(z) - \mathcal{R} f_{r_i}(z)| |\mathcal{R} g(z)| \\ &\preceq \sup_{|z| > \delta} \frac{\mu(|z|) |g(z)| |z|}{\phi(|z|)(1 - |z|^2)^{2 + \frac{n}{q}}} \|f\|_{p,q,\phi} + \sup_{|z| > \delta} \frac{r_i \mu(|z|) |g(z)| |r_i z|}{\phi(|r_i z|)(1 - |r_i z|^2)^{2 + \frac{n}{q}}} \|f\|_{p,q,\phi} \\ &\quad + \sup_{|z| > \delta} \frac{\mu(|z|) |\mathcal{R} g(z)| |z|}{\phi(|z|)(1 - |z|^2)^{1 + \frac{n}{q}}} \|f\|_{p,q,\phi} + \sup_{|z| > \delta} \frac{r_i \mu(|z|) |\mathcal{R} g(z)| |r_i z|}{\phi(|r_i z|)(1 - |r_i z|^2)^{1 + \frac{n}{q}}} \|f\|_{p,q,\phi} \\ &\leq \sup_{|z| > \delta} \frac{\mu(|z|) |g(z)| |z|}{\phi(|z|)(1 - |z|^2)^{2 + \frac{n}{q}}} + \sup_{|z| > \delta} \frac{r_i \mu(|z|) |g(z)| |r_i z|}{\phi(|r_i z|)(1 - |r_i z|^2)^{2 + \frac{n}{q}}} \\ &\quad + \sup_{|z| > \delta} \frac{\mu(|z|) |\mathcal{R} g(z)| |z|}{\phi(|z|)(1 - |z|^2)^{1 + \frac{n}{q}}} + \sup_{|z| > \delta} \frac{r_i \mu(|z|) |\mathcal{R} g(z)| |r_i z|}{\phi(|r_i z|)(1 - |r_i z|^2)^{1 + \frac{n}{q}}}. \end{aligned}$$

If $i \rightarrow \infty$, then

$$\|L_g f - L_g T_{r_i} f\|_{\mathcal{Z}_\mu} \preceq 2 \sup_{|z| > \delta} \frac{\mu(|z|) |g(z)| |z|}{\phi(|z|)(1 - |z|^2)^{2 + \frac{n}{q}}} + 2 \sup_{|z| > \delta} \frac{\mu(|z|) |\mathcal{R} g(z)| |z|}{\phi(|z|)(1 - |z|^2)^{1 + \frac{n}{q}}}.$$

If $\delta \rightarrow 1$, then (9) implies that

$$\|L_g\|_{e, H(p,q,\phi) \rightarrow \mathcal{Z}_\mu} \preceq \limsup_{|z| \rightarrow 1} \frac{\mu(|z|) |g(z)|}{\phi(|z|)(1 - |z|^2)^{2 + \frac{n}{q}}} + \limsup_{|z| \rightarrow 1} \frac{\mu(|z|) |\mathcal{R} g(z)|}{\phi(|z|)(1 - |z|^2)^{1 + \frac{n}{q}}}.$$

□

References

- [1] L. Hu, S. Li and R. Yang, Essential norm of Riemann-Stieltjes operator on weighted Bergman spaces with doubling weights, *Georgian Mathematical Journal*, 2024. <https://doi.org/10.1515/gmj-2023-2110>
- [2] B. Li and C. Ouyang, Higher radial derivative of Bloch type functions, *Acta Math. Sci. Ser. B* 22(4) (2002), 433-445.
- [3] S. Li and S. Stević, Riemann-Stieltjes operators on Hardy spaces in the unit ball of \mathbb{C}^n , *Bull. Belg. Math. Soc. Simon Stevin* 14(4) (2007), 621-628.
- [4] S. Li and S. Stević, Riemann-Stieltjes-type integral operators on the unit ball in \mathbb{C}^n , *Complex Var. Elliptic Equ.* 52(6) (2007), 495-517.
- [5] S. Li and S. Stević, Riemann-Stieltjes operators between different weighted Bergman spaces, *Bull. Belg. Math. Soc. Simon Stevin* 15(4) (2008), 677-686.
- [6] Y. Liang, H. Zeng and Ze-Hua Zhou, Volterra-type operators from $F(p, q, s)$ space to Bloch-Orlicz and Zygmund-Orlicz spaces, *Filomat* 34(4) (2020), 1359-1381.
- [7] Y. Liang and Z. H. Zhou, Some integral-type operators from $F(p, q, s)$ spaces to mixed-norm spaces on the unit ball, *Math. Nachr.* 287 (11-12) (2014), 1298-1311.
- [8] Y. Liang, Integral-type operators from $F(p, q, s)$ space to α -Bloch-Orlicz and β -Zygmund-Orlicz spaces, *Complex Anal. Oper. Theory* 12 (2018), 169-194.
- [9] Y. Liu and Y. Yu, Riemann-Stieltjes operator from mixed norm spaces to Zygmund-type spaces on the unit ball, *Taiwanese J. Math.* 17(5) (2013), 1751-1764.
- [10] Y. Liu, Y. Yu and X. Liu, Riemann-Stieltjes operator from the general space to Zygmund-type spaces on the unit ball, *Complex Anal. Oper. Theory* 9 (2015), 985-997.
- [11] S. Miihkinen, P. J. Nieminen and W. Xu, Essential norms of integration operators on weighted Bergman spaces, *J. Math. Anal. Appl.* 450(1) (2017), 229-243.
- [12] J. Pau and R. Zhao, Carleson measures, Riemann-Stieltjes and multiplication operators on a general family of function spaces, *Integr. Equ. Oper. Theory* 78 (2014), 483-514.
- [13] R. Peng Y. Wu and F. Deng, Carleson measures, Riemann-Stieltjes operators and multipliers on $F(p, q, s)$ spaces in the unit ball of \mathbb{C}^n . *Acta Math Sci.* 36B (2016), 635-654.
- [14] S. Stević, On an integral operator between Bloch-type spaces on the unit ball. *Bull. Sci. Math.* 134(4) (2010), 329-339.
- [15] S. Stević, Weighted iterated radial composition operators between some spaces of holomorphic functions on the unit ball, *Abstr. Appl. Anal.* 2010, Art. ID 801264, 14 pp., doi:10.1155/2010/801264.
- [16] C. L. Yang, Integral-type operators from $F(p, q, s)$ spaces to Zygmund-type spaces on the unit ball, *J. Inequal. Appl.* 2010 (2010), Article ID 789285, 14 pages.
- [17] R. Zhao, On a general family of function spaces, *Ann. Acad. Sci. Fenn. Math. Diss.* 105 (1996), 56.
- [18] X. Zhang, Multipliers on some holomorphic function spaces (in Chinese). *Chin. Ann. Math. Ser. A* 26(4) (2005), 477-486.
- [19] X. Zhang, C. He and F. Cao, The equivalent norms of $F(p, q, s)$ space in \mathbb{C}^n . *J. Math. Anal. Appl.* 401(2) (2013), 601-610.
- [20] R. Zhao, On $F(p, q, s)$ spaces, *Acta Mathematica Scientia*, 2021, 41B(6) (2021), 1985-2020.
- [21] K. Zhu, *Spaces of holomorphic functions in the unit ball. Graduate Text in Mathematics 226.* Springer, New York (2005)

Y. ESTAREMI

Email address: `y.estaremi@gu.ac.ir`

DEPARTMENT OF MATHEMATICS, FACULTY OF SCIENCES, GOLESTAN UNIVERSITY, GORGAN, IRAN.

M. HASSANLOU

Email address: `m.hassanlou@urmia.ac.ir`

ENGINEERING FACULTY OF KHOY, URMIA UNIVERSITY OF TECHNOLOGY, URMIA, IRAN