

a -POTENT SCHWARZ MATRICES AND BESSEL-LIKE JACOBI POLYNOMIALS

ALEXANDER DYACHENKO, CARLOS M. DA FONSECA, AND MIKHAIL TYAGLOV

ABSTRACT. We consider the problem of the reconstruction of a Schwarz matrix from exactly one given eigenvalue. This inverse eigenvalue problem leads to the Jacobi orthogonal polynomials $\{P_k^{(-n,n)}\}_{k=0}^{n-1}$ that can be treated as a discrete finite analogue of Bessel polynomials.

1. INTRODUCTION

The matrix inverse eigenvalue problem consists in reconstructing a specific structured matrix from a prescribed spectral data. This topic is of great interest in different branches of mathematics, continually providing new and surprising results (see for example [10, 22, 33, 34] and the references there).

From the celebrated work [36] by Hubert S. Wall, it follows that there exists a *unique* matrix

$$(1.1) \quad J_n = \begin{bmatrix} -b_0 & 1 & & & \\ -b_1 & 0 & 1 & & \\ & -b_2 & \ddots & \ddots & \\ & & \ddots & \ddots & 1 \\ & & & -b_{n-1} & 0 \end{bmatrix}$$

with $b_k > 0$, for $k = 0, 1, \dots, n-1$, whose characteristic polynomial is $(x+1)^n$ (see also [32, 35, 37] for more details). Wall's argumentation is based on the stability of the characteristic polynomial, i.e. on that all its zeros have negative real parts — which is of course true for $(x+1)^n$.

By analogy with nilpotent matrices, one introduces *unipotent* matrices: an $n \times n$ matrix is called unipotent if its characteristic polynomial is $(x-1)^n$, or, equivalently, if all its eigenvalues are equal to 1. Various disciplines deal with unipotent matrices, cf. e.g. [6, 21, 28, 38]. We consider an extension of this definition, namely matrices with only a single eigenvalue labelled by a . In brief, we are looking for an “ a -potent” matrix of the form (1.1) where $a \in \mathbb{C} \setminus \{0\}$.

Matrices of the form (1.1) are called Schwarz matrices. Such matrices are very important in automatic control theory and signal processing (see, for example, [8]), and several general inverse problems have been previously considered in [3, 7, 11, 13, 16]. So, here we consider the inverse eigenvalue problem for a Schwarz matrix whose characteristic

Date: June 18, 2024.

2020 Mathematics Subject Classification. 15B05, 33C45, 15A18, 15A29, 15A15.

Key words and phrases. Inverse eigenvalue problem, Schwarz matrices, unipotent matrices, orthogonal polynomials.

polynomial is

$$(1.2) \quad p(z) = (z - a)^n = \sum_{i=0}^n (-1)^{n-i} \binom{n}{i} a^{n-i} z^i,$$

where a is a given nonzero complex number. That is, we want to construct a Schwarz matrix (1.1) with only one eigenvalue a .

Since J_k is the leading principal $k \times k$ submatrix of J_n , with $k \in \{1, \dots, n\}$, the polynomials

$$P_k(z) = \det(zI_k - J_k), \quad \text{for } k = 1, \dots, n,$$

where I_k is the identity matrix of order k and $P_n(z) = p(z)$, satisfy the following three-term recurrence relation:

$$(1.3) \quad \begin{aligned} P_0(z) &= 1, \\ P_1(z) &= z + b_0, \\ P_{k+1}(z) &= zP_k(z) + b_k P_{k-1}(z), \quad \text{for } k = 1, \dots, n-1. \end{aligned}$$

By the Favard theorem, see¹ [9, pp. 21–22], the sequence $\{P_k(z)\}_{k=0}^{n-1}$ is orthogonal w.r.t. a linear functional. Observe that, given arbitrary $n > 1$ and $a \in \mathbb{C} \setminus \{0\}$, this functional is not positive and cannot be written as an integral w.r.t. a measure on the real line (otherwise J_n would have n distinct real eigenvalues). Accordingly, for $n > 1$ the numbers b_1, \dots, b_{n-1} in (1.3) cannot be chosen negative while b_0 remains real. As is seen from Section 3.2, the support of the orthogonality functional is natural to take consisting of a unique point a .

We find the aforementioned matrix J_n , see the values of b_0 and b_1, \dots, b_{n-1} in, resp., (2.9) and (2.23). The sequence $\{P_k(z)\}_{k=0}^{n-1}$ satisfying (1.3) reduces to a finite chunk of the Jacobi polynomials with non-standard parameters. The explicit expression of the polynomial $P_k(z)$ for $a = 1$ is given in (3.3), while the relation of the polynomials with distinct a is in (3.2).

It turns out that three-term recurrence relations for the Bessel polynomials [24] have a form similar to (1.3), so Bessel polynomials are also characteristic polynomials of Schwarz matrices, and for $a = -\frac{1}{n}$ the polynomials $P_k(z)$ approximate Bessel polynomials uniformly on the compact sets of the complex plane as $n \rightarrow +\infty$. Moreover, the support of the functional of orthogonality for Bessel polynomials on the real line may also be treated as consisting of one point of essential singularity. Thus, we can treat polynomials $\{P_k(z)\}_{k=0}^{n-1}$ as a discrete finite analogue of Bessel polynomials.

The paper is organised as follows. Section 2 is devoted to finding the entries of the matrix J_n . In Sections 3.1 and 3.2, we investigate properties of the orthogonal polynomials $P_k(z)$ and find their functional of orthogonality on the real line. In Section 3.3, we compare our polynomials with Bessel polynomials and provide some calculations showing that the zeroes of $P_k(z)$ approach the zeroes of Bessel polynomials for $a = -\frac{1}{n}$ as $n \rightarrow +\infty$. In Section 3.4, we calculate zeroes of $P_k(z)$ for varying a and k to estimate numerically their asymptotic behaviour as n is large.

¹In [27], this extended version is attributed to Chihara, although in essence it already existed in [17].

2. EXPLICIT FORMULA FOR SCHWARZ MATRIX

To find the entries of the matrix J_n or, similarly, the coefficients of the three-term recurrence relations (1.3), we consider the rational function

$$(2.1) \quad R(z) = b_0((zI_n - J_n)^{-1}e_1, e_1) = \frac{q(z)}{p(z)},$$

where e_1 is the first coordinate vector. Here the polynomial $q(z)$, multiplied by b_0 , is the characteristic polynomial of the principal submatrix of J_n (1.1) constructed from J_n by deleting its first row and the first column:

$$q(z) = b_0 \det \begin{bmatrix} z & -1 & 0 & & \\ b_2 & z & -1 & & \\ & b_3 & \ddots & \ddots & \\ & & \ddots & \ddots & -1 \\ & & & b_{n-1} & z \end{bmatrix}.$$

Observe that the polynomial $p(z)$, the characteristic polynomial of the matrix J_n , can be represented as follows

$$(2.2) \quad p(z) = Q(z) + q(z),$$

where

$$Q(z) = \det \begin{bmatrix} z & -1 & 0 & & \\ b_1 & z & -1 & & \\ & b_2 & \ddots & \ddots & \\ & & \ddots & \ddots & -1 \\ & & & b_{n-1} & z \end{bmatrix}.$$

It is easy to see that $q(z)$ is an even polynomial whenever n is odd, and is an odd polynomial otherwise, while the polynomial $Q(z)$ is odd if n is odd, and is even if n is even. Therefore, by (2.2), the polynomial $q(z)$ is the even (odd) part of the polynomial $p(z)$ whenever n is odd (even). This means that $q(z)$ can be obtained from $p(z)$ in the following way

$$q(z) = \frac{p(z) - (-1)^n p(-z)}{2},$$

so that

$$(2.3) \quad \begin{aligned} q(z) &= -\binom{n}{1}az^{n-1} - \binom{n}{3}a^3z^{n-3} - \binom{n}{5}a^5z^{n-5} - \dots \\ &= -\sum_{j=0}^{\lfloor \frac{n-1}{2} \rfloor} \binom{n}{2j+1} a^{2j+1} z^{n-2j-1}. \end{aligned}$$

For the calculation of the entries of the matrix J_n we first study the rational function $R(z)$ defined in (2.1).

Theorem 2.1. *The rational function $R(z) = \frac{q(z)}{p(z)}$ has the form*

$$R(z) = \frac{\alpha_1}{z-a} + \frac{\alpha_1}{(z-a)^2} + \dots + \frac{\alpha_n}{(z-a)^n} = \frac{s_0}{z} + \frac{s_1}{z^2} + \frac{s_2}{z^3} + \dots,$$

where

$$(2.4) \quad \alpha_k = -2^{k-1}a^k \binom{n}{k}, \quad \text{for } k = 1, \dots, n,$$

and

$$(2.5) \quad s_m = \sum_{k=1}^n a^{m-k+1} \binom{m}{k-1} \alpha_k = -a^{m+1} \sum_{k=1}^n 2^{k-1} \binom{m}{k-1} \binom{n}{k}.$$

Proof. Expanding the polynomial $q(z)$ into the Taylor series at the point a , we obtain

$$R(z) = \frac{q(a)}{(z-a)^n} + \frac{q'(a)}{1!(z-a)^{n-1}} + \frac{q''(a)}{2!(z-a)^{n-2}} + \dots + \frac{q^{(n-1)}(a)}{(n-1)!(z-a)},$$

so

$$\alpha_k = \frac{q^{(n-k)}(a)}{(n-k)!}, \quad \text{for } k = 1, \dots, n.$$

From (2.3), we have

$$\frac{q^{(n-k)}(z)}{(n-k)!} = - \sum_{j=1}^{\lfloor \frac{k-1}{2} \rfloor} \binom{n}{2j+1} \binom{n-2j-1}{n-k} a^{2j+1} z^{k-2j-1}, \quad \text{for } k = 1, \dots, n,$$

which implies

$$(2.6) \quad \alpha_k = \frac{q^{(n-k)}(a)}{(n-k)!} = -a^k \binom{n}{k} \sum_{j=0}^{\lfloor \frac{k-1}{2} \rfloor} \binom{k}{2j+1}, \quad \text{for } k = 1, \dots, n.$$

Now substituting

$$2^{k-1} = \sum_{j=0}^{\lfloor \frac{k-1}{2} \rfloor} \binom{k}{2j+1}$$

into (2.6), we get (2.4).

To establish (2.5), we recall that, for any nonnegative integer number ℓ ,

$$\frac{1}{(z-a)^\ell} = \sum_{m=0}^{\infty} \binom{m+\ell-1}{\ell-1} \frac{a^m}{z^{m+\ell}},$$

so

$$\begin{aligned} R(z) &= \sum_{k=1}^n \frac{\alpha_k}{(z-a)^k} = \sum_{m=0}^{\infty} \sum_{k=1}^n \frac{\binom{m+k-1}{k-1} a^m \alpha_k}{z^{m+k}} \\ &= \sum_{m=0}^{\infty} \frac{1}{z^{m+1}} \sum_{k=1}^n a^{m-k+1} \binom{m}{k-1} \alpha_k. \end{aligned}$$

This formula together with (2.4) gives us (2.5). □

Remark 2.2. Note that the numbers s_m satisfy the following recurrence relation

$$(1+m)a^2 s_m + 2nas_{m+1} - (3+m)s_{m+2} = 0.$$

A careful consideration shows that the moments s_m are polynomials in n , and can be represented as

$$s_m(n) = -\frac{a^{m+1}n}{(m+1)!} p_m(n), \quad \text{for } m = 0, 1, 2, \dots,$$

where $p_m(x)$ are the polynomials satisfying the following three-term recurrence relations:

$$p_{m+1}(x) = xp_m(x) + m(m+1)p_{m-1}(x), \quad \text{for } m = 1, 2, \dots$$

with the initial conditions $p_0(x) = 1$, $p_1(x) = x$. These recurrence relations generate the following continued fraction (see, e.g., [26, p.231]):

$$\Phi(x) = x \int_0^\infty e^{-xt} \tanh(t) dt = \frac{1}{x + \frac{1 \cdot 2}{x + \frac{2 \cdot 3}{x + \dots}}}$$

convergent for $x > 0$. In fact, the integral itself converges in the right half-plane, so it generates a measure of orthogonality for the polynomials $p_m(x)$ with the support on the imaginary axis.

It is not hard to identify this system of orthogonal polynomials: on comparing the recurrence coefficients one observes that they are the Meixner–Pollaczek polynomials $P_m^{(1)}(\cdot; \frac{\pi}{2})$ up to rotation and renormalisation. Namely,

$$p_m(n) = -\frac{(m+1)!}{a^{m+1}n} s_m(n) = (m+1)! {}_2F_1 \left(\begin{matrix} -m, 1-n \\ 2 \end{matrix} \middle| 2 \right) = \frac{m!}{i^m} P_m^{(1)} \left(in; \frac{\pi}{2} \right),$$

so the corresponding orthogonality measure is $|\Gamma(1-x)|^2$ with $x \in i\mathbb{R}$, see² [9, p. 180].

The entries b_m of the matrix J_n defined in (1.1) can be found by the formula (see, e.g., [20, 35] and references therein)

$$(2.7) \quad b_m = -\frac{D_{m-1}(R)D_{m+1}(R)}{D_m^2(R)}, \quad \text{for } m = 1, 2, \dots, n-1,$$

where $D_0(R) = 1$, and $D_m(R)$, for $m = 1, \dots, n$, are the Hankel determinants defined as follows

$$(2.8) \quad D_m(R) = \begin{vmatrix} s_0 & s_1 & s_2 & \cdots & s_{m-1} \\ s_1 & s_2 & s_3 & \cdots & s_m \\ \vdots & \vdots & \vdots & \cdots & \vdots \\ s_{m-1} & s_m & s_{m+1} & \cdots & s_{2m-2} \end{vmatrix}, \quad \text{for } m = 1, 2, \dots, n.$$

Recall that

$$D_m(R) = 0, \quad \text{for } m = n+1, n+2, \dots,$$

since the rational function $R(z)$ has exactly n poles, counting multiplicities (see, e.g., [15, Ch. XV, §10, Theorem 8] or, alternatively, [20, Theorem 1.2]).

Moreover,

$$(2.9) \quad b_0 = -an,$$

since $-b_0$ equals the trace of J_n .

To find the exact formula for the minors $D_m(R)$ we need the following general result that can be obtained using elementary determinant properties.

²Note a typo in formula (3.22) on that page: the third parameter of ${}_2F_1$ is actually 2λ , not 2.

Lemma 2.3. *The minors $D_m(R)$ can be represented in the following form:*

$$(2.10) \quad D_m(R) = \begin{vmatrix} \alpha_1 & \alpha_2 & \cdots & \alpha_m \\ \alpha_2 & \alpha_3 & \cdots & \alpha_{m+1} \\ \vdots & \vdots & \ddots & \vdots \\ \alpha_m & \alpha_{m+1} & \cdots & \alpha_{2m-1} \end{vmatrix},$$

for $m = 1, 2, \dots, n$, where we set $\alpha_j = 0$, for $j > n$.

Proof. Indeed, the k th column \mathbf{C}_k of the minor $D_m(R)$ has the form

$$(2.11) \quad \mathbf{C}_k = \begin{bmatrix} s_{k-1} \\ \vdots \\ s_{k+m-2} \end{bmatrix}, \quad \text{for } k = 1, \dots, m.$$

If we change columns as follows

$$(2.12) \quad \tilde{\mathbf{C}}_k = \sum_{j=1}^k (-a)^{k-j} \binom{k-1}{j-1} \mathbf{C}_j,$$

for $k = 1, \dots, m$, where $\tilde{\mathbf{C}}_k$ is the k th new column, then the minor $D_m(R)$ does not change its value. From (2.11)–(2.12), we obtain that the ℓ -entry $(\tilde{\mathbf{C}}_k)_\ell$ of the new k th column has the form

$$\begin{aligned} (\tilde{\mathbf{C}}_k)_\ell &= \sum_{j=1}^k (-a)^{k-j} \binom{k-1}{j-1} s_{j+\ell-2} \\ &= \sum_{j=1}^k (-a)^{k-j} \binom{k-1}{j-1} \sum_{i=1}^n a^{j+\ell-1-i} \binom{j+\ell-2}{i-1} \alpha_i \\ &= \sum_{i=1}^n a^{k+\ell-1-i} \alpha_i \sum_{j=0}^{k-1} (-1)^{k-1-j} \binom{k-1}{j} \binom{j+\ell-1}{i-1} \\ &= \sum_{i=1}^n a^{k+\ell-1-i} \binom{\ell-1}{i-k} \alpha_i = \sum_{i=1}^{\ell} a^{\ell-i} \binom{\ell-1}{i-1} \alpha_{k+i-1}, \end{aligned}$$

where $k, \ell = 1, \dots, m$ and $\alpha_j = 0$, for $j > n$. Here we used the identity

$$\sum_{k=0}^m (-1)^{m-k} \binom{m}{k} \binom{k+\ell}{i} = \binom{\ell}{i-m}$$

which is true for any nonnegative integer numbers m, ℓ, i (cf., e.g., [31, Section 4.2.5, no. 55] or [18, Chapter 5, Table 169]).

Thus, we get the following formula for the minors $D_m(R)$:

$$(2.13) \quad D_m(R) = \begin{vmatrix} \alpha_1 & \alpha_2 & \cdots & \alpha_m \\ a\alpha_1 + \alpha_2 & a\alpha_2 + \alpha_3 & \cdots & a\alpha_m + \alpha_{m+1} \\ \vdots & \vdots & \ddots & \vdots \\ \sum_{r=1}^m a^{m-r} \binom{m}{r} \alpha_r & \sum_{r=1}^m a^{m-r} \binom{m}{r} \alpha_{r+1} & \cdots & \sum_{r=1}^m a^{m-r} \binom{m}{r} \alpha_{r+m-1} \end{vmatrix},$$

for $m = 1, 2, \dots, n$.

Let us consider now the ℓ th row of the matrix (2.13). It has the form

$$(2.14) \quad \mathbf{R}_\ell = [v_{\ell,1}, v_{\ell,2}, \dots, v_{\ell,m}] \quad , \quad \text{for } \ell = 1, \dots, m,$$

where

$$(2.15) \quad v_{\ell k} = \sum_{i=1}^{\ell} a^{\ell-i} \binom{\ell-1}{i-1} \alpha_{i+k-1}, \quad \text{for } \ell, k = 1, 2, \dots, m.$$

Now we change the rows as follows

$$(2.16) \quad \tilde{\mathbf{R}}_\ell = \sum_{j=1}^{\ell} (-a)^{\ell-j} \binom{\ell-1}{j-1} \mathbf{R}_j, \quad \text{for } \ell = 1, \dots, m,$$

where $\tilde{\mathbf{R}}_\ell$ is the ℓ th new row. This operation does not change the value of the minor $D_m(R)$. From (2.14)–(2.16), we conclude that the k th entry $(\tilde{\mathbf{R}}_\ell)_k$ of the new ℓ th row has the form

$$(2.17) \quad \begin{aligned} (\tilde{\mathbf{R}}_\ell)_k &= \sum_{j=1}^{\ell} (-a)^{\ell-j} \binom{\ell-1}{j-1} v_{j,k} \\ &= \sum_{j=1}^{\ell} \sum_{i=1}^j a^{\ell-i} (-1)^{\ell-j} \binom{\ell-1}{j-1} \binom{j-1}{i-1} \alpha_{i+k-1} \\ &= \sum_{i=1}^{\ell} a^{\ell-i} \alpha_{i+k-1} \sum_{j=i}^{\ell} (-1)^{\ell-j} \binom{\ell-1}{j-1} \binom{j-1}{i-1} \\ &= \sum_{i=1}^{\ell} a^{\ell-i} \alpha_{i+k-1} \delta_{\ell i} = \alpha_{k+\ell-1}, \end{aligned}$$

for $k, \ell = 1, \dots, m$. Now (2.17) implies (2.10). □

Now we are in a position to calculate the values of the Hankel minors $D_m(R)$.

Theorem 2.4. *The minors $D_m(R)$ have the form*

$$(2.18) \quad D_m(R) = (-1)^{\frac{m(m+1)}{2}} a^{m^2} 2^{m(m-1)} \binom{n}{m} \prod_{k=0}^{m-1} \frac{\binom{n+k}{n-k}}{\binom{m+k}{m-k}},$$

for $m = 1, 2, \dots, n$.

Proof. From (2.4) and (2.10) we have

$$\begin{aligned}
D_m(R) &= \begin{vmatrix} -a\binom{n}{1} & -2a^2\binom{n}{2} & \cdots & -2^{m-1}a^m\binom{n}{m} \\ -2a^2\binom{n}{2} & -2^2a^3\binom{n}{3} & \cdots & -2^m a^{m+1}\binom{n}{m+1} \\ \vdots & \vdots & \ddots & \vdots \\ -2^{m-1}a^m\binom{n}{m} & -2^m a^{m+1}\binom{n}{m+1} & \cdots & -2^{2m-2}a^{2m-1}\binom{n}{2m-1} \end{vmatrix} \\
&= (-1)^m 2^{\frac{m(m-1)}{2}} a^{\frac{m(m+1)}{2}} \begin{vmatrix} \binom{n}{1} & 2a\binom{n}{2} & \cdots & 2^{m-1}a^{m-1}\binom{n}{m} \\ \binom{n}{2} & 2a\binom{n}{3} & \cdots & 2^{m-1}a^{m-1}\binom{n}{m+1} \\ \vdots & \vdots & \ddots & \vdots \\ \binom{n}{m} & 2a\binom{n}{m+1} & \cdots & 2^{m-1}a^{m-1}\binom{n}{2m-1} \end{vmatrix} \\
&= (-1)^m 2^{\frac{m(m-1)}{2}} a^{\frac{m(m+1)}{2}} (2a)^{\frac{m(m-1)}{2}} \begin{vmatrix} \binom{n}{1} & \binom{n}{2} & \cdots & \binom{n}{m} \\ \binom{n}{2} & \binom{n}{3} & \cdots & \binom{n}{m+1} \\ \vdots & \vdots & \ddots & \vdots \\ \binom{n}{m} & \binom{n}{m+1} & \cdots & \binom{n}{2m-1} \end{vmatrix} \\
(2.19) \quad &= (-1)^m 2^{m(m-1)} a^{m^2} T_m,
\end{aligned}$$

where

$$\begin{aligned}
T_m &= \begin{vmatrix} \binom{n}{1} & \binom{n}{2} & \cdots & \binom{n}{m} \\ \binom{n}{2} & \binom{n}{3} & \cdots & \binom{n}{m+1} \\ \vdots & \vdots & \ddots & \vdots \\ \binom{n}{m} & \binom{n}{m+1} & \cdots & \binom{n}{2m-1} \end{vmatrix} \\
(2.20) \quad &= \begin{vmatrix} \frac{n}{1!} & \frac{n(n-1)}{2!} & \cdots & \frac{n(n-1)\cdots(n-m+1)}{m!} \\ \frac{n(n-1)}{2!} & \frac{n(n-1)(n-2)}{3!} & \cdots & \frac{n(n-1)\cdots(n-m)}{(m+1)!} \\ \vdots & \vdots & \ddots & \vdots \\ \frac{n(n-1)\cdots(n-m+1)}{m!} & \frac{n(n-1)\cdots(n-m)}{(m+1)!} & \cdots & \frac{n(n-1)\cdots(n-2m+2)}{(2m-1)!} \end{vmatrix}
\end{aligned}$$

for $m = 1, 2, 3, \dots, n$. Taking into account that the j th, $(j+1)$ th, ..., m th rows have the common factor $n(n-1)\cdots(n-j+1)$, for $j = 1, \dots, m$, we get

$$\begin{aligned}
T_m &= n^m (n-1)^{m-1} \cdots (n-m)^2 (n-m+1) \begin{vmatrix} \frac{1}{1!} & \frac{n-1}{2!} & \cdots & \frac{(n-1)\cdots(n-m+1)}{m!} \\ \frac{1}{2!} & \frac{n-2}{3!} & \cdots & \frac{(n-2)\cdots(n-m)}{(m+1)!} \\ \vdots & \vdots & \ddots & \vdots \\ \frac{1}{m!} & \frac{n-m}{(m+1)!} & \cdots & \frac{(n-m)\cdots(n-2m+2)}{(2m-1)!} \end{vmatrix} \\
&= \prod_{k=1}^m k! \binom{n}{k} \begin{vmatrix} \frac{1}{1!} & \frac{n-1}{2!} & \cdots & \frac{(n-1)\cdots(n-m+1)}{m!} \\ \frac{1}{2!} & \frac{n-2}{3!} & \cdots & \frac{(n-2)\cdots(n-m)}{(m+1)!} \\ \vdots & \vdots & \ddots & \vdots \\ \frac{1}{m!} & \frac{n-m}{(m+1)!} & \cdots & \frac{(n-m)\cdots(n-2m+2)}{(2m-1)!} \end{vmatrix}
\end{aligned}$$

Now we add the $(m-1)$ th column to the m th column, then the $(m-2)$ th column to the $(m-1)$ th column, etc. The identity

$$(2.21) \quad \begin{aligned} & \frac{(n-i)(n-i-1)\cdots(n-i-j+3)}{(i+j-2)!} + \frac{(n-i)(n-i-1)\cdots(n-i-j+2)}{(i+j-1)!} = \\ & = (n+1) \frac{(n-i)(n-i-1)\cdots(n-i-j+3)}{(i+j-1)!} \end{aligned}$$

is used repeatedly for each element of the j th column, for $j = m, m-1, \dots, 2$. Thus, we conclude that the determinant does not change its value, while its 2nd, 3rd, ..., m th columns have the common factor $n+1$. Therefore, we have

$$T_m = (n+1)^{m-1} \prod_{k=1}^m k! \binom{n}{k} \begin{vmatrix} \frac{1}{1!} & \frac{1}{2!} & \cdots & \frac{(n-1)\cdots(n-m+2)}{m!} \\ \frac{1}{2!} & \frac{1}{3!} & \cdots & \frac{(n-2)\cdots(n-m+1)}{(m+1)!} \\ \vdots & \vdots & \ddots & \vdots \\ \frac{1}{m!} & \frac{1}{(m+1)!} & \cdots & \frac{(n-m)\cdots(n-2m+3)}{(2m-1)!} \end{vmatrix}$$

Now we use the similar procedure for the 3rd, 4th, ..., $(m+1)$ th columns. Namely, we add the $(m-1)$ th column to the m th, then the $(m-2)$ th column to the $(m-1)$ th, etc., and use the identity

$$\begin{aligned} & \frac{(n-i)(n-i-1)\cdots(n-i-j+4)}{(i+j-2)!} + \frac{(n-i)(n-i-1)\cdots(n-i-j+3)}{(i+j-1)!} = \\ & = (n+2) \frac{(n-i)(n-i-1)\cdots(n-i-j+4)}{(i+j-1)!} \end{aligned}$$

repeatedly for each element of the j th column, for $j = 3, \dots, m$. Again we conclude that the determinant does not change its value, while its 3rd, 4th, ..., m th columns have the common factor $n+2$. So we have

$$T_m = (n+1)^{m-1} (n+2)^{m-2} \prod_{k=1}^m k! \binom{n}{k} \begin{vmatrix} \frac{1}{1!} & \frac{1}{2!} & \cdots & \frac{(n-1)\cdots(n-m+3)}{m!} \\ \frac{1}{2!} & \frac{1}{3!} & \cdots & \frac{(n-2)\cdots(n-m+2)}{(m+1)!} \\ \vdots & \vdots & \ddots & \vdots \\ \frac{1}{m!} & \frac{1}{(m+1)!} & \cdots & \frac{(n-m)\cdots(n-2m+4)}{(2m-1)!} \end{vmatrix}$$

Consequently, from a straightforward inductive argument, we get

$$\begin{aligned}
T_m &= \prod_{\ell=1}^{m-1} (n+\ell)^{m-\ell} \prod_{k=1}^m k! \binom{n}{k} \begin{vmatrix} \frac{1}{1!} & \frac{1}{2!} & \cdots & \frac{1}{m!} \\ \frac{1}{2!} & \frac{1}{3!} & \cdots & \frac{1}{(m+1)!} \\ \vdots & \vdots & \ddots & \vdots \\ \frac{1}{m!} & \frac{1}{(m+1)!} & \cdots & \frac{1}{(2m-1)!} \end{vmatrix} \\
&= \prod_{k=1}^m k! \binom{n}{k} \prod_{j=1}^{m-1} j! \binom{n+j}{j} \begin{vmatrix} \frac{1}{1!} & \frac{1}{2!} & \cdots & \frac{1}{m!} \\ \frac{1}{2!} & \frac{1}{3!} & \cdots & \frac{1}{(m+1)!} \\ \vdots & \vdots & \ddots & \vdots \\ \frac{1}{m!} & \frac{1}{(m+1)!} & \cdots & \frac{1}{(2m-1)!} \end{vmatrix} \\
&= m! \binom{n}{m} \prod_{k=1}^{m-1} (2k)! \binom{n+k}{n-k} \begin{vmatrix} \frac{1}{1!} & \frac{1}{2!} & \cdots & \frac{1}{m!} \\ \frac{1}{2!} & \frac{1}{3!} & \cdots & \frac{1}{(m+1)!} \\ \vdots & \vdots & \ddots & \vdots \\ \frac{1}{m!} & \frac{1}{(m+1)!} & \cdots & \frac{1}{(2m-1)!} \end{vmatrix}.
\end{aligned}$$

Note that the determinant T_m equals $(-1)^{\frac{m(m-1)}{2}}$, if $n = m$, as it follows from (2.20). Therefore,

$$(2.22) \quad \begin{vmatrix} \frac{1}{1!} & \frac{1}{2!} & \cdots & \frac{1}{m!} \\ \frac{1}{2!} & \frac{1}{3!} & \cdots & \frac{1}{(m+1)!} \\ \vdots & \vdots & \ddots & \vdots \\ \frac{1}{m!} & \frac{1}{(m+1)!} & \cdots & \frac{1}{(2m-1)!} \end{vmatrix} = \frac{(-1)^{\frac{m(m-1)}{2}}}{m! \prod_{k=1}^{m-1} (2k)! \binom{m+k}{m-k}},$$

and, consequently,

$$\begin{aligned}
T_m &= (-1)^{\frac{m(m-1)}{2}} \frac{m! \binom{n}{m} \prod_{k=1}^{m-1} (2k)! \binom{n+k}{n-k}}{m! \prod_{k=1}^{m-1} (2k)! \binom{m+k}{m-k}} \\
&= (-1)^{\frac{m(m-1)}{2}} \binom{n}{m} \prod_{k=1}^{m-1} \frac{\binom{n+k}{n-k}}{\binom{m+k}{m-k}}.
\end{aligned}$$

Substituting this expression for T_m into (2.19), we obtain (2.18), as required. \square

Remark 2.5. The determinant (2.22) is connected to the Padé approximations to the exponential, and its value can be found, e.g., in [2, Section 1.2, p.9].

Remark 2.6. On summing up the matrix transformations behind Theorem 2.4, one additionally arrives at the following factorisation of the matrix from (2.20):

$$\left[\binom{n}{i+j-1} \right]_{i,j=1}^m = \text{diag} [(n-j)_{j+1}]_{j=0}^{m-1} \cdot \left[\frac{1}{(i+j-1)!} \right]_{i,j=1}^m \cdot \text{diag} [(n)_j]_{j=0}^{m-1} \cdot \left[(-1)^{i+j} \binom{j}{i} \right]_{i,j=0}^{m-1},$$

where

$$\left[(-1)^{i+j} \binom{j}{i} \right]_{i,j=0}^{m-1} = \left(\left[\binom{j}{i} \right]_{i,j=0}^{m-1} \right)^{-1}$$

is the inverse of the upper-triangular Pascal matrix. This inverse of the Pascal matrix is a product of $m-1$ bidiagonal terms stemming from column operations and application of the Pascal identity starting (2.21). The Pascal matrix itself also has an analogous bidiagonal factorisation (but the terms are unsigned). The remaining term, whose determinant appears on the left-hand side of (2.22), can be expressed from the matrix $\left[\binom{m}{i+j-1}\right]_{i,j=1}^m$ by the reverse transformations with n set to be m , that is

$$\left[\frac{1}{(i+j-1)!}\right]_{i,j=1}^m = \text{diag} \left[\frac{1}{(m-j)_{j+1}}\right]_{j=0}^{m-1} \cdot \left[\binom{m}{i+j-1}\right]_{i,j=1}^m \cdot \text{diag} \left[\frac{1}{(m)_j}\right]_{j=0}^{m-1} \cdot \left[\binom{j}{i}\right]_{i,j=0}^{m-1}.$$

At the same time, the Hankel matrix $\left[\binom{m}{i+j-1}\right]_{i,j=1}^m$ is built from the coefficients of the polynomial $(x+1)^m$, and hence it may be further factorised into $m \times m$ factors

$$\left[\binom{m}{i+j-1}\right]_{i,j=1}^m = \begin{bmatrix} 0 & \dots & 0 & 0 & 1 \\ \vdots & \ddots & 0 & 1 & 0 \\ \vdots & \ddots & 1 & 0 & 0 \\ 0 & \ddots & \ddots & \ddots & \vdots \\ 1 & 0 & 0 & \dots & 0 \end{bmatrix} \cdot \begin{bmatrix} 1 & 0 & 0 & \dots & 0 \\ 1 & 1 & 0 & \ddots & \vdots \\ 0 & 1 & 1 & \ddots & 0 \\ \vdots & \ddots & \ddots & \ddots & 0 \\ 0 & \dots & 0 & 1 & 1 \end{bmatrix}^m.$$

Now (2.7) and (2.18) give us the exact formula for the entries of the matrix (1.1) whose characteristic polynomial is (1.2).

Corollary 2.7. *The coefficients b_m of the relations (1.3) have the form*

$$(2.23) \quad b_m = a^2 \frac{n^2 - m^2}{4m^2 - 1}, \quad \text{for } m = 1, 2, \dots, n-1.$$

Proof. Let us consider first the relation

$$\begin{aligned} \frac{D_{m+1}(R)}{D_m(R)} &= \frac{(-1)^{\frac{(m+1)(m+2)}{2}} 2^{(m+1)m} a^{(m+1)^2} \binom{n}{m+1} \prod_{k=1}^m \frac{\binom{n+k}{n-k}}{\binom{m+1+k}{m+1-k}}}{(-1)^{\frac{m(m+1)}{2}} 2^{m(m-1)} a^{m^2} \binom{n}{m} \prod_{k=1}^{m-1} \frac{\binom{n+k}{n-k}}{\binom{m+k}{m-k}}} \\ &= (-1)^{m+1} 4^m a^{2m+1} \frac{n-m}{m+1} \frac{\binom{n+m}{n-m}}{2m+1} \prod_{k=1}^{m-1} \frac{\binom{m+k}{m-k}}{\binom{m+1+k}{m+1-k}} \\ &= (-1)^{m+1} 4^m a^{2m+1} \frac{n-m}{m+1} \frac{\binom{n+m}{n-m}}{2m+1} \prod_{k=1}^{m-1} \frac{m+1-k}{m+1+k} \\ (2.24) \quad &= (-1)^{m+1} 4^m a^{2m+1} \frac{n-m}{(2m+1)!} \binom{n+m}{n-m} (m!)^2. \end{aligned}$$

Now, from (2.7) and (2.24), we have

$$\begin{aligned}
b_m &= -\frac{D_{m+1}(R)/D_m(R)}{D_m(R)/D_{m-1}(R)} \\
&= -\frac{(-1)^{m+1}4^m a^{2m+1} \frac{n-m}{(2m+1)!} \binom{n+m}{n-m} (m!)^2}{(-1)^m 4^{m-1} a^{2m-1} \frac{n-m+1}{(2m-1)!} \binom{n+m-1}{n-m+1} ((m-1)!)^2} \\
&= 4m^2 a^2 \frac{n-m}{2m(2m+1)} \frac{(n+m)!}{(n-m)!(2m)!} \frac{(n-m)!(2m-2)!}{(n+m-1)!} \\
&= a^2 \frac{n-m}{(2m-1)(2m+1)} \frac{(n+m)!}{(n+m-1)!} = a^2 \frac{n^2 - m^2}{4m^2 - 1}.
\end{aligned}$$

□

As a conclusion, up to similarity, the only Schwarz matrix J_n whose characteristic polynomial is $(z-a)^n$ is

$$(2.25) \quad J_n = \begin{bmatrix} an & 1 & & & \\ -a^2 \frac{n^2-1}{3} & 0 & 1 & & \\ & -a^2 \frac{n^2-4}{15} & \ddots & \ddots & \\ & & \ddots & \ddots & 1 \\ & & & -a^2 \frac{1}{2n-3} & 0 \end{bmatrix}.$$

Clearly, the spectrum of the matrix

$$\tilde{J}_n = a \begin{bmatrix} n & \frac{n-1}{1} & & & \\ -\frac{n+1}{3} & 0 & \frac{n-2}{3} & & \\ & -\frac{n+2}{5} & \ddots & \ddots & \\ & & \ddots & \ddots & \frac{1}{2n-3} \\ & & & -1 & 0 \end{bmatrix}$$

is reduced to a as well.

3. ORTHOGONAL POLYNOMIALS

In this section, we consider the sequence $\{P_k(z)\}_{k=0}^{n-1}$ of the characteristic polynomials of the leading principal minors of the matrix (2.25) satisfying the three-term recurrence relation (1.1)–(1.3):

$$(3.1) \quad \begin{aligned}
P_0(z) &= 1, \quad P_1(z) = z - an, \\
P_{k+1}(z) &= zP_k(z) + \frac{a^2(n^2 - k^2)}{(2k-1)(2k+1)} P_{k-1}(z), \quad \text{for } k = 1, \dots, n-1.
\end{aligned}$$

We give their explicit representations and study their even and odd parts. Then we show that $\{P_k(z)\}_{k=0}^{n-1}$ are orthogonal w.r.t. a linear functional supported at the only point $z = a$.

We also find the inverse Fourier transform of the corresponding distributional weight and depict the zeroes of the polynomials $P_k(z)$ for various a and relations between k and n as $n \rightarrow +\infty$.

3.1. Basic properties. For the current subsection we assume $a = 1$ keeping in mind that the general case $a \neq 0$ may be obtained by

$$(3.2) \quad a^{-k} P_k(az) = P_k(z)|_{a=1},$$

which immediately follows from (3.1). Observe that the monic Jacobi polynomials

$$\widehat{P}_k^{(-n,n)}(z) = \frac{2^k}{\binom{2k}{k}} P_k^{(-n,n)}(z) \quad \left(= \frac{(-2)^k}{\binom{2k}{k}} P_k^{(n,-n)}(-z) \right)$$

satisfy for $k \geq 1$ the three-term recurrence relation [9, p. 153, eq. (2.29)]

$$\widehat{P}_{k+1}^{(-n,n)}(z) = z \widehat{P}_k^{(-n,n)}(z) - \frac{4k(k-n)(k+n)k}{(2k-1)(2k+1)(2k)^2} \widehat{P}_{k-1}^{(-n,n)}(z),$$

which on reduction of similar terms turns into (3.1) with $a = 1$. Similarly, $\widehat{P}_0^{(-n,n)}(z) \equiv 1$ and $\widehat{P}_1^{(-n,n)}(z) = z - n$. Therefore, our $P_k(z)$ are exactly $\widehat{P}_k^{(-n,n)}(z)$, namely, for all $k = 0, \dots, n$,

$$(3.3) \quad P_k(z) = \widehat{P}_k^{(-n,n)}(z) = (-2)^k \frac{(n+1)_k}{(k+1)_k} {}_2F_1 \left(\begin{matrix} -k, k+1 \\ n+1 \end{matrix} \middle| \frac{1+z}{2} \right).$$

Also,

$$P_k(z)|_{a=1} = \frac{(-1)^k}{(k+1)_k} \left(\frac{1-z}{1+z} \right)^n \frac{d^k}{dz^k} [(1-z)^{n-k} (1+z)^{n+k}],$$

$$(1-z^2)P_k''(z) + 2(n-z)P_k'(z) + k(k+1)P_k(z) = 0.$$

Although the three-term recurrence degenerates to a two-term recurrence for $k = n$, the sequence $\{P_k\}_{k=0}^n$ may be further extended to an infinite sequence. Properties of infinite sequences of the monic polynomials $\{\widehat{P}_k^{(-n,\beta)}\}_{k=0}^\infty$ with $\beta \notin \{-1, -2, \dots\}$ were studied in [1], where orthogonality was introduced via a bilinear form.

Now, on introducing two auxiliary families of polynomials via

$$(3.4) \quad f_0(z) = 1, \quad f_1(z) = z, \quad \text{and} \quad f_{k+1}(z) = z f_k(z) + \frac{n^2 - k^2}{4k^2 - 1} f_{k-1}(z)$$

and

$$(3.5) \quad g_0(z) = 0, \quad g_1(z) = 1, \quad \text{and} \quad g_{k+1}(z) = z g_k(z) + \frac{n^2 - k^2}{4k^2 - 1} g_{k-1}(z)$$

with $k = 1, \dots, n-1$, one immediately obtains that the decomposition

$$P_k(z) = f_k(z) - n g_k(z)$$

is valid for all k . The formula (3.4) determines monic orthogonal polynomials supported on the imaginary axis, and (3.5) gives exactly the corresponding monic numerator polynomials. In particular, for each k the zeroes of f_k and g_k are all simple and pure imaginary, they also interlace on the imaginary axis, see [9, p. 86]. Therefore, the Hermite-Biehler theorem implies that all zeroes of the polynomial $P_k(z) = f_k(z) - n g_k(z)$ have positive real parts, [25] (see also [20] and references there). That is to say, all zeroes of the polynomial P_k for arbitrary $a \neq 0$ lie in the half-plane $\{z \in \mathbb{C} : \operatorname{Re}(z/a) > 0\}$, which is clearly seen on the figures in Sections 3.3 and 3.4.

In fact, the polynomials f_k and g_k may also be expressed via classical families of orthogonal polynomials. The three-term relations (3.4) and (3.5) are known to determine monic versions of the so-called associated Jacobi polynomials $\widehat{P}_k^{(\alpha,\beta)}(z; c)$, namely,

$$f_k(z) = \widehat{P}_k^{(-n,-n)}(z; n) \quad \text{and} \quad g_k(z) = \widehat{P}_{k-1}^{(-n,-n)}(z; n+1).$$

The explicit representation of the associated Jacobi polynomials is due to Wimp [23, p. 182], however our specific case allows to write these polynomials using the simple expressions

$$f_k(z) = \frac{(-1)^k P_k(-z) + P_k(z)}{2} \quad \text{and} \quad g_k(z) = \frac{(-1)^k P_k(-z) - P_k(z)}{2n}.$$

Note that the sequence $\{f_k\}_{k=0}^{n-1}$ consists of the so-called (monic) exceptional Jacobi polynomials [23, p. 97], although the parameter $\alpha = -n$ is non-standard for them. Observe also that f_k and $-ng_k$ turn into Q and q from (2.2) on letting $k = n$.

3.2. Orthogonality. Now, let us introduce the following linear functional $\mathcal{L}_n^a(f(x))$ acting on $(n-1)$ th differentiable functions f and depending on the positive integer number n and the (complex) nonzero number a :

$$\mathcal{L}_n^a(f(x)) = \sum_{k=1}^n \frac{f^{(k-1)}(a) \alpha_k}{(k-1)!} = \sum_{k=1}^n \frac{f^{(k-1)}(a) q^{(n-k)}(a)}{(k-1)! (n-k)!} = \frac{1}{(n-1)!} \frac{d^{n-1}[f(x)q(x)]}{dx^{n-1}} \Big|_{x=a}.$$

where the numbers α_k are defined in (2.4).

Alternatively, assuming that f is analytic near the point a this functional may be expressed in a more usual form — as an integral with a continuous (complex-valued) Jacobi weight $(z-a)^{-n}(z+a)^n$ over the circle $|z-a| = 2|a|$, which is typical for the Jacobi polynomials with such parameters, cf. [17, p. 222]. The weight is analytic in $\mathbb{C} \setminus \{0\}$, so the circle may be replaced with any Jordan curve enclosing the point $z = a$. Due to Cauchy's integral formula, this integral will essentially express the same functional \mathcal{L}_n^a , namely:

$$\begin{aligned} \mathcal{L}_n^a(f) &= \frac{1}{(n-1)!} \frac{d^{n-1}[f(z)q(z)]}{dz^{n-1}} \Big|_{z=a} = \frac{1}{2\pi i} \oint_{|z-a|=2|a|} \frac{f(z)q(z)}{(z-a)^n} dz \\ &= \frac{1}{2\pi i} \oint_{|z-a|=2|a|} f(z) \cdot \frac{1}{2} \left(\left(\frac{z-a}{z-a} \right)^n - \left(\frac{z+a}{z-a} \right)^n \right) dz \\ &= \frac{i}{4\pi} \oint_{|z-a|=2|a|} f(z) \left(\frac{z+a}{z-a} \right)^n dz. \end{aligned}$$

With this functional we can represent the rational function $R(z)$ defined in (2.1) as follows

$$(3.6) \quad R(z) = \mathcal{L}_n^a \left(\frac{1}{z-x} \right) = \sum_{k=1}^n \frac{\alpha_k}{(z-a)^k}.$$

Moreover, from (3.2) we have

$$\mathcal{L}_n^a(x^m) = s_m, \quad \text{for } m = 0, 1, \dots,$$

where s_m are the moments defined in (2.5). In fact,

$$\begin{aligned}\mathcal{L}_n^a(x^m) &= \sum_{k=1}^n \frac{(x^m)^{(k-1)} \Big|_{x=a}}{(k-1)!} \alpha_k = a^m \alpha_1 + \sum_{k=2}^n \frac{m(m-1) \cdots (m-k+2) a^{m-k+1}}{(k-1)!} \alpha_k \\ &= \sum_{k=1}^n \binom{m}{k-1} a^{m-k+1} \alpha_k = s_m.\end{aligned}$$

We now prove that the polynomials $P_k(z)$, for $k = 0, 1, \dots, n-1$, defined in (3.1) are orthogonal w.r.t. the functional (3.2), that is,

$$\mathcal{L}_n^a(P_m(x)P_k(x)) = C_m \delta_{km} \quad \text{for } m, k = 0, 1, \dots, n-1,$$

where δ_{km} is the Kronecker symbol. To this end, we find the three-term recurrence relations for monic polynomials $Q_m(z)$ orthogonal w.r.t. the functional $\mathcal{L}_n^a(f)$ and compare the obtained relations with (3.1). Firstly, let us note that all the minors $D_m(R)$, $m = 1, \dots, n$, defined in (2.8) are nonzero, i.e.,

$$D_m(R) \neq 0, \quad \text{for } m = 1, \dots, n,$$

by (2.18), so such a sequence exists (see, e.g., [9, Theorem 3.1]). These orthogonal polynomials have the form [23, p. 21]

$$(3.7) \quad Q_m(z) = \frac{1}{D_m(R)} \begin{vmatrix} s_0 & s_1 & s_2 & \cdots & s_m \\ s_1 & s_2 & s_3 & \cdots & s_{m+1} \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ s_{m-1} & s_m & s_{m+1} & \cdots & s_{2m-1} \\ 1 & z & z^2 & \cdots & z^m \end{vmatrix},$$

for $m = 1, \dots, n$. The roots of the last polynomial $Q_n(z)$ (partially) define the functional $\mathcal{L}_n^a(f)$. If $\mathcal{L}_n^a(f)$ is a positive or negative functional, $Q_n(z)$ would have exactly n real distinct roots that are the growth point of the measure generated by $\mathcal{L}_n^a(f)$, see, e.g., [23, Section 2.2]. But the formula (2.18) shows that $\mathcal{L}_n^a(f)$ is neither a positive nor a negative functional, since the minors $D_m(R)$ are not all positive nor of alternating signs ($\text{sgn } D_m(R) \neq (-1)^m$, for $m = 1, \dots, n$).

The polynomials $Q_m(z)$ satisfy a three-term recurrence relation

$$Q_{m+1}(z) = (z - c_{m+1})Q_m(z) - d_{m+1}Q_{m-1}(z).$$

It is well known, e.g., [9, 20, 23, 37], that

$$(3.8) \quad d_{m+1} = \frac{D_{m-1}(R)D_{m+1}(R)}{D_m^2(R)} = -b_m,$$

so the polynomials $Q_m(z)$ satisfy the recurrence relation

$$(3.9) \quad Q_{m+1}(z) = (z - c_{m+1})Q_m(z) + b_m Q_{m-1}(z), \quad \text{for } m = 0, 1, \dots, n-1,$$

where $Q_{-1}(z) = 0$ and $Q_0(z) = 1$.

Furthermore, if

$$Q_m(z) = z^m + \sum_{i=1}^m A_{im} z^{m-i},$$

then (cf., e.g., [9])

$$(3.10) \quad c_1 = A_{11}, \quad c_{m+1} = A_{1m} - A_{1,m+1}, \quad \text{for } m = 1, \dots, n-1,$$

and, from (3.7), we have

$$(3.11) \quad A_{1,m} = -\frac{D'_m(R)}{D_m(R)}, \quad \text{for } m = 1, \dots, n.$$

where

$$D'_m(R) = \begin{vmatrix} s_0 & s_1 & s_2 & \cdots & s_{m-2} & s_m \\ s_1 & s_2 & s_3 & \cdots & s_{m-1} & s_{m+1} \\ \vdots & \vdots & \vdots & \cdots & \vdots & \vdots \\ s_{m-1} & s_m & s_{m+1} & \cdots & s_{2m-3} & s_{2m-1} \end{vmatrix}, \quad \text{for } m = 1, \dots, n.$$

To find the formula for the minors $D'_m(R)$, we need the following lemma which can be proved in the same way as Lemma 2.3.

Lemma 3.1. *The minors $D'_m(R)$ can be represented in the following form:*

$$(3.12) \quad D'_m(R) = \begin{vmatrix} \alpha_1 & \alpha_2 & \alpha_3 & \cdots & \alpha_m \\ \alpha_2 & \alpha_3 & \alpha_4 & \cdots & \alpha_{m+1} \\ \vdots & \vdots & \vdots & \cdots & \vdots \\ \alpha_{m-1} & \alpha_m & \alpha_{m+1} & \cdots & \alpha_{2m-2} \\ \alpha_{m+1} & \alpha_{m+2} & \alpha_{m+3} & \cdots & \alpha_{2m} \end{vmatrix} + am \begin{vmatrix} \alpha_1 & \alpha_2 & \alpha_3 & \cdots & \alpha_m \\ \alpha_2 & \alpha_3 & \alpha_4 & \cdots & \alpha_{m+1} \\ \vdots & \vdots & \vdots & \cdots & \vdots \\ \alpha_{m-1} & \alpha_m & \alpha_{m+1} & \cdots & \alpha_{2m-2} \\ \alpha_m & \alpha_{m+1} & \alpha_{m+2} & \cdots & \alpha_{2m-1} \end{vmatrix},$$

for $m = 1, \dots, n$, and setting $\alpha_j = 0$, for $j > n$.

Furthermore, by the same technique that we used in proof of Theorem 2.4, we may conclude the next theorem (here, we omit the proof to avoid long and unnecessary repetitions).

Theorem 3.2. *The following identity holds for $m = 1, \dots, n$,*

$$(3.13) \quad \begin{vmatrix} \alpha_1 & \alpha_2 & \alpha_3 & \cdots & \alpha_m \\ \alpha_2 & \alpha_3 & \alpha_4 & \cdots & \alpha_{m+1} \\ \vdots & \vdots & \vdots & \cdots & \vdots \\ \alpha_{m-1} & \alpha_m & \alpha_{m+1} & \cdots & \alpha_{2m-2} \\ \alpha_{m+1} & \alpha_{m+2} & \alpha_{m+3} & \cdots & \alpha_{2m} \end{vmatrix} = a(n-m) \begin{vmatrix} \alpha_1 & \alpha_2 & \alpha_3 & \cdots & \alpha_m \\ \alpha_2 & \alpha_3 & \alpha_4 & \cdots & \alpha_{m+1} \\ \vdots & \vdots & \vdots & \cdots & \vdots \\ \alpha_{m-1} & \alpha_m & \alpha_{m+1} & \cdots & \alpha_{2m-2} \\ \alpha_m & \alpha_{m+1} & \alpha_{m+2} & \cdots & \alpha_{2m-1} \end{vmatrix},$$

where we set $\alpha_j = 0$, for $j > n$, and for $m = 1$ we mean that the left determinant equals α_2 .

So, from (3.12) and (3.13) we have

$$D'_m(R) = an \begin{vmatrix} \alpha_1 & \alpha_2 & \alpha_3 & \cdots & \alpha_m \\ \alpha_2 & \alpha_3 & \alpha_4 & \cdots & \alpha_{m+1} \\ \vdots & \vdots & \vdots & \cdots & \vdots \\ \alpha_{m-1} & \alpha_m & \alpha_{m+1} & \cdots & \alpha_{2m-2} \\ \alpha_m & \alpha_{m+1} & \alpha_{m+2} & \cdots & \alpha_{2m-1} \end{vmatrix},$$

i.e.,

$$D'_m(R) = anD_m(R).$$

Now, (2.10) and (3.11) imply

$$A_{1,m} = -an, \quad \text{for } m = 1, 2, \dots, n.$$

Hence

$$(3.14) \quad c_1 = an \quad \text{and} \quad c_m = 0, \quad \text{for } m = 2, \dots, n.$$

by (3.10).

Thus, from (3.8)–(3.9) and (3.14) we have

$$Q_{m+1}(z) = zQ_m(z) + \frac{a^2(n^2 - m^2)}{(2m - 1)(2m + 1)}Q_{m-1}(z), \quad \text{with } m = 1, \dots, n,$$

with initial conditions $Q_0(z) = 1$, and $Q_1(z) = z - an$. Therefore, the monic polynomials $P_k(z)$ coincide with the monic polynomials $Q_m(z)$ which are orthogonal with respect to the functional $\mathcal{L}_n^a(f(x))$ defined in (3.2) and

$$\begin{aligned} C_m &= \mathcal{L}_a^n(P_m^2(x)) = b_0 \prod_{k=1}^m (-b_k) = (-1)^{m+1} a^{2m+1} n \prod_{k=1}^m \frac{(n^2 - k^2)}{4k^2 - 1} \\ &= (-1)^{m+1} a^{2m+1} \binom{n+m}{2m+1} \frac{(2m)!!}{(2m-1)!!}, \end{aligned}$$

or, equivalently,

$$C_m = (-1)^{m+1} a^{2m+1} \sqrt{\pi} \binom{n+m}{2m+1} \frac{\Gamma(m+1)}{\Gamma(m+\frac{1}{2})},$$

for $m = 0, 1, \dots, n-1$.

For example, setting $m = n-1$, we have the following interesting fact:

$$C_{n-1} = \mathcal{L}_a^n(P_{n-1}^2(x)) (-1)^n a^{2n-1} \sqrt{\pi} \frac{\Gamma(n)}{\Gamma(n-\frac{1}{2})} \sim (-1)^n a^{2n-1} \sqrt{\pi n} \quad \text{as } n \rightarrow +\infty.$$

Having the moments s_m defined in (2.5) which are related to the orthogonal polynomials $\{P_m(z)\}_{m=0}^{n-1}$, one can introduce the distributional weight function w_a characterised by those moments (see [29]) such that

$$(3.15) \quad \langle w_a, f \rangle = \sum_{m=0}^{\infty} \frac{s_m f^{(m)}(0)}{m!}$$

for infinitely differentiable function f with a compact support. Therefore, the distributional weight function ω_a of this functional has the form [29]

$$(3.16) \quad \omega_a(x) = \sum_{m=0}^{\infty} \frac{(-1)^m s_m \delta^{(m)}(x)}{m!},$$

where $\delta(x) = \delta^{(0)}(x)$ is the Dirac delta-function, and $\delta^{(m)}(x)$, for $m = 1, 2, \dots$, are its derivatives. From (2.5) it easily follows that the moments s_m satisfy the inequality

$$|s_m| \leq C |a|^m m!,$$

for certain constants $C = C(a, n) > 0$, and the distributional weight $w_a(x)$ has the inverse Fourier transform [29]

$$F^{-1}w_a(t) = \frac{1}{2\pi} \int_{-\infty}^{+\infty} e^{-ixt} \omega_a(x) dx = \frac{1}{2\pi} \sum_{m=0}^{+\infty} \frac{s_m(-it)^m}{m!}.$$

The sum of this series can be found explicitly.

Theorem 3.3. *The inverse Fourier transform of the distributional weight (3.16) corresponding to the functional w_a has the form*

$$F^{-1}w_a(t) = \frac{-ae^{-iat}}{2\pi} L_{n-1}^{(1)}(2iat), \quad t \in \mathbb{C},$$

where $L_n^{(\alpha)}(x)$ denotes the (generalised) Laguerre polynomial.

Proof. From (2.5) one has

$$\begin{aligned} F^{-1}w_a(t) &= \frac{-a}{2\pi} \sum_{m=0}^{\infty} \sum_{k=1}^n 2^{k-1} a^m \binom{m}{k-1} \binom{n}{k} \frac{(-it)^m}{m!} \\ &= \frac{-a}{2\pi} \sum_{k=1}^n \frac{2^{k-1}}{(k-1)!} \binom{n}{k} \sum_{m=0}^{\infty} \frac{(-ait)^m}{(m-k+1)!} \\ &= \frac{-ae^{-ait}}{2\pi} \sum_{k=0}^{n-1} \frac{1}{k!} \binom{n}{k+1} (-2ait)^k = \frac{-ae^{-iat}}{2\pi} L_{n-1}^{(1)}(2iat), \end{aligned}$$

where the series converges for $t \in \mathbb{C}$. □

In (3.15) we defined the functional w_a for infinitely differentiable functions with compact support. In fact it will be extended to a larger class of functions. Indeed, let us find the functional w_a by applying the Fourier transform to $F^{-1}w_a(t)$.

$$\begin{aligned} F(F^{-1}w_a(t)) &= -\frac{a}{2\pi} F\left(e^{-iat} L_{n-1}^{(1)}(2iat)\right) \\ &= \sum_{k=1}^n \frac{-2^{k-1} a^k \binom{n}{k}}{(k-1)!} \frac{(-i)^{k-1} F(e^{-iat} t^{k-1})}{2\pi} = \sum_{k=1}^n \frac{\alpha_k \delta^{(k-1)}(x-a)}{(k-1)!}, \end{aligned}$$

where α_k are defined in (2.4). Thus, we can continuously extend the functional w_a to the set of all $(n-1)$ -times differentiable functions. From (3.2) it follows that the continuously extended functional w_a coincides with \mathcal{L}_a^n .

Note that since the matrix J_n defined in (1.1) is tridiagonal, it is non-derogatory. Thus, the normalized (by the first unit entry) eigenvector corresponding to the unique eigenvalue of J_n is $\mathbf{u}_0 := \mathbf{u}(a)$ where $\mathbf{u}(z) = (P_0(z), P_1(z), \dots, P_{n-2}(z), P_{n-1}(z))^T$. The corresponding generalised eigenvectors are $\mathbf{u}_k := \left. \frac{d^k \mathbf{u}(z)}{dz^k} \right|_{z=a}$, $k = 1, \dots, n-1$.

3.3. Relation to Bessel polynomials. The Bessel polynomials $\{B_n(x)\}_{n=0}^{\infty}$ is a sequence of polynomials satisfying the differential equation

$$x^2 \frac{d^2 y}{dx^2} + (2x+2) \frac{dy}{dx} = n(n+1)y.$$

They occur naturally in the theory of traveling spherical waves [24]. These (monic) polynomials satisfy the following three-term recurrence relations

$$(3.17) \quad \begin{aligned} B_0(z) &= 1, & B_1(z) &= z + 1, \\ B_{k+1}(z) &= zB_k(z) + \frac{1}{4k^2 - 1}B_{k-1}(z), & \text{for } k &= 1, 2, \dots \end{aligned}$$

The Bessel polynomials were introduced by H.L. Krall and O. Frink in [24]. They satisfy complex orthogonality conditions

$$\int_C B_n(z)B_m(z)e^{-2/z}dz = 0, \quad m \neq n,$$

where C is a closed path in the complex plane enclosing the origin. Their orthogonality on the real axis was a subject of discussion for many years until this problem was resolved in a few different ways, see [12, 14, 29]. We note that on the real line, the support of the functional of orthogonality for the Bessel polynomials consists of a unique point $z = 0$ which is an essential singularity. Moreover, from the recurrence relations (3.17), it follows that every monic Bessel polynomial is the characteristic polynomial of a Schwarz matrix defined by these relations and, therefore, is stable. Since the polynomials $\{P_k\}_{k=0}^{n-1}$ satisfying (3.1) and orthogonal w.r.t. the functional \mathcal{L}_a^n defined in (3.2) are characteristic polynomials of Schwarz matrices stable for $a < 0$, and the support of the functional \mathcal{L}_a^n consists of one point, we may say that $\{P_k\}_{k=0}^{n-1}$ can be treated as a finite discrete analogue of Bessel polynomials. Additionally, if one sets $a = -\frac{1}{n}$, then the three-term recurrence relations (3.1) for the polynomials $P_k(z)$ take the form

$$\begin{aligned} P_0(z) &= 1, & P_1(z) &= z + 1, \\ P_{k+1}(z) &= zP_k(z) + \frac{P_{k-1}(z)}{4k^2 - 1} - \frac{1}{n^2} \frac{k^2}{4k^2 - 1}P_{k-1}(z), & \text{for } k &= 1, \dots, n-1, \end{aligned}$$

from which it follows that as $n \rightarrow +\infty$, these recurrence relations are close to (3.17). Thus, the polynomials $P_k(z)$ approach $B_k(z)$ coefficient-wise, and, therefore, on the compact subsets of the complex plane. So for $a = -\frac{1}{n}$, sufficiently large n , and relatively small k (comparing with n) one can consider the polynomials $P_k(z)$ as an approximation of the monic Bessel polynomials $B_k(z)$. Thus, we can expect that for a fixed k and large n the zeroes of $P_k(z)$ will approximate zeroes of $B_k(z)$, and as $k \rightarrow +\infty$ and $n \rightarrow +\infty$ with $\frac{k}{n} \rightarrow 0$ the zeroes of $P_k(z)$ will approach the same curve as the zeroes of $B_k(z)$ for $k \rightarrow +\infty$, see [4, 5] and references there. Figures 3.1 and 3.2 show the zeroes of the polynomials $B_{50}(z)$ and $B_{75}(z)$, respectively, and the zeroes of $P_{50}(z)$ and $P_{75}(z)$ approaching them as $n \rightarrow +\infty$.

Figures 3.3 and 3.4 represent the zeroes of the polynomials $P_k(z)$ and $B_k(z)$ for $n = k^2$ and $k = 5, \dots, 24$. There is a visibly good approximation of $B_k(z)$ by $P_k(z)$ in such a case.

Note also that if $a = -\frac{1}{n}$, then as $n \rightarrow +\infty$ the limits of the moments $s_m(n)$, $m = 0, 1, 2, \dots$, are close to the moments of the Bessel polynomials (but not the same). At the same time, the inverse Fourier transform of the distributional weight function




FIGURE 3.1. Zeroes of B_{50} and P_{50} for $n = 50, \dots, 150$




FIGURE 3.2. Zeroes of B_{75} and P_{75} for $n = 75, \dots, 175$




FIGURE 3.3. Zeroes of P_k and B_k for $n = k^2$ and $k = 5, \dots, 19$




FIGURE 3.4. Zeroes of P_k and B_k for $n = k^2$ and $k = 20, \dots, 24$

corresponding to the Bessel polynomials has the form

$$F^{-1}w(t) = -\frac{1}{2\pi} \sum_{k=0}^{+\infty} \frac{2^{k+1}(it)^k}{k!(k+1)!} = -\frac{I_1(\sqrt{8it})}{\pi\sqrt{8it}},$$

where $I_1(x)$ is the modified Bessel function [29].

3.4. Zeroes of polynomials $P_k(z)$ for fixed a . Apart of approximation of Bessel polynomials, it is interesting to study how the zeroes the polynomials P_k behave for a fixed $a \in \mathbb{C} \setminus \{0\}$. As it was already observed in Section 3.1, the results on the polynomials P_k for a certain fixed $a \in \mathbb{C} \setminus \{0\}$ immediately extend to other values of a due to

$$a^{-k}P_k(az) = P_k(z)|_{a=1} \quad \text{or, equivalently,} \quad (-a)^{-k}P_k(-az) = P_k(z)|_{a=-1}.$$

Accordingly, varying values of a only scales and/or rotates the zeroes of these polynomials in \mathbb{C} , see Figure 3.5. We shall concentrate on the case $a = -1$ which is related to the previous section.

Figures 3.7 and 3.8 represent the zeroes of the polynomials $P_{n/10}(z)$ and $P_{n-1}(z)$ for $a = -1$. According to calculations, the zeroes tend to a certain curve in the left half-plane that depends on the limiting ratio $\frac{k}{n}$ as $k \rightarrow \infty$, $n > k$.




FIGURE 3.5. Zeros of P_k for $a = e^{i\varphi}$ with $n = k + 1 = 71$, and $\varphi = -\pi, \dots, -\frac{\pi}{2}$




FIGURE 3.6. Zeros of P_k for $a = -1$, $k = 17, \dots, 50$, and $n = 50$




FIGURE 3.7. Zeros of P_k for $a = -1$, $n = 10k$, and $k = 10, \dots, 60$




FIGURE 3.8. Zeros of P_k for $a = -1$, $n = k + 1$, and $k = 50, \dots, 150$

If one changes linear dependence between n and k to higher powers, the zeroes of P_k move to infinity as $k \rightarrow +\infty$, see Figures 3.9 and 3.10. This phenomenon may be explained as follows. As $\frac{k}{n} \rightarrow 0$ we can rescale the polynomials $P_k(z)$ to have the zeroes bounded. The proper rescaling may be achieved by letting $a = -\frac{1}{n}$ and study the asymptotic behaviour of the zeroes of $P_k\left(\frac{z}{k}\right)$, which is equivalent to letting $a = -\frac{k}{n}$ and studying the zeroes of $P_k(z)$, see Figures 3.11 and 3.12. From the discussion in Section 3.3 it follows that the zeroes of $P_k\left(\frac{z}{k}\right)$ for $a = -\frac{1}{n}$ should approximate those of $B_k\left(\frac{z}{k}\right)$ as




FIGURE 3.9. Zeros of P_k for $a = -1$, $n = k^2$, and $k = 9, \dots, 144$




FIGURE 3.10. Zeros of P_k for $a = -1$, $n = \lfloor k^{5/4} \rfloor$, and $k = 9, \dots, 144$

$k \rightarrow \infty$, $n \rightarrow \infty$, $\frac{k}{n} \rightarrow 0$. At the same time, the latter zeroes are known to remain bounded [19, 30], see also [4, 5].




FIGURE 3.11. Zeros of P_k for $a = -k/n$, $n = k^2$, and $k = 9, \dots, 144$




FIGURE 3.12. Zeros of P_k for $a = -k/n$, $n = \lfloor k^{5/4} \rfloor$, and $k = 9, \dots, 144$

ACKNOWLEDGEMENT

The results of Theorems 2.1 and 2.4, Corollary 2.7 and Sections 3.2 and 3.3 (except Theorem 3.2) were obtained with the support of the Russian Science Foundation grant 19-71-30002. The work on Remarks 2.2 and 2.6, Section 3.1 and all numerical calculations were supported by the state assignment, registration number 122041100132-9.

REFERENCES

- [1] M. Alfaro, T. E. Pérez, M. A. Piñar, M. L. Rezola, Sobolev orthogonal polynomials: the discrete-continuous case, *Methods Appl. Anal.*, **6**, no. 4, 1999, 593–616.
- [2] G. Baker, Jr. and P. Graves-Morris, *Padé approximants*, Vol. 1, Encyclopedia of Mathematics and Applications 13, Addison-Wesley Publ. Co., 1980.
- [3] A. Behn, K. R. Driessel, I. R. Hentzel, K. A. Vander Velden, and J. Wilson, Some nilpotent, tridiagonal matrices with a special sign pattern, *Linear Algebra Appl.*, **436**, no. 12, 2012, 4446–4450.

- [4] M.G. de Bruin, E.B. Saff, and R.S Varga, On the generalized Bessel polynomials. I, *Indag. Math.*, **84**, no. 1, 1981, 1–13.
- [5] M.G. de Bruin, E.B. Saff, and R.S Varga, On the generalized Bessel polynomials. II, *Indag. Math.*, **84**, no. 1, 1981, 14–25.
- [6] M.B. Can, R. Howe, and M. Joyce, Unipotent invariant matrices, *Linear Algebra Appl.*, **439**, 2013, 196–210.
- [7] K. Castillo, C.M. da Fonseca, and J. Petronilho, On Chebyshev polynomials and the inertia of certain matrices, *Appl. Math. Comput.*, **467**, 2024, no. 128497.
- [8] V. Černý and J. Hrušák, On some new similarities between non-linear observer and filter design, *Nonlinear Control Syst.*, **6**, 2004, 465–470.
- [9] T.S. Chihara, *An Introduction to Orthogonal Polynomials*, Gordon and Breach, New York, 1978.
- [10] M.T. Chu, Inverse eigenvalue problems, *SIAM Rev.*, **40**, 1998, 1–39.
- [11] J.H. Drew, C.R. Johnson, D.D. Olesky, and P. van den Driessche, Spectrally arbitrary patterns, *Linear Algebra Appl.*, **308**, 2000, 121–137.
- [12] A.J. Duran, Functions with given moments and weight functions for orthogonal polynomials, *Rocky Mountain J. Math.*, **23**, no. 1, 1993, 87–104.
- [13] L. Elsner and D. Hershkowitz, On the spectra of close-to-Schwarz matrices, *Linear Algebra Appl.*, **363**, 2003, 81–88.
- [14] W.D. Evans, W.N. Everitt, K.H. Kwon, and L.L. Littlejohn, Real orthogonalizing weights for Bessel polynomials, *J. Comput. Appl. Math.*, **49**, 1993, 51–57.
- [15] F.R. Gantmacher, *The Theory of Matrices*, vol. II, Translated by K. A. Hirsch, Chelsea Publishing Co., New York, 1959.
- [16] C. Garnett and B.L. Shader, A proof of the conjecture: Centralizers, Jacobians and spectrally arbitrary sign patterns, *Linear Algebra Appl.*, **436**, no. 12, 2012, 4451–4458.
- [17] J. Geronimus, Sur les polynômes orthogonaux relatifs à une suite des nombres donnée et sur le théorème de W. Hahn, *Izv. Akad. Nauk SSSR, Ser. Mat.*, **4**, no. 2, 1940, 215–228.
- [18] R. Gragham, D. Knuth, and O. Patashnik, *Concrete Mathematics*, Advanced Book Program (First ed.), Reading, MA: Addison-Wesley Publ. Co., 1998, pp. xiv+625.
- [19] E. Grosswald, *Bessel polynomials*, Lecture Notes in Mathematics, **698**, Springer-Verlag, New York, 1978, pp. 182.
- [20] O. Holtz and M. Tyaglov, Structured matrices, continued fractions, and root localization of polynomials, *SIAM Rev.*, **54**, 2012, 421–509.
- [21] X. Hou, Z. Xiao, Y. Hao, and Q. Yuan, Decomposition of symplectic matrices into products of symplectic unipotent matrices of index 2, *Electron. J. Linear Algebra*, **35**, 2019, 497–502.
- [22] C.R. Johnson, B.D. Sutton, and A.J. Witt, Implicit construction of multiple eigenvalues for trees, *Linear Multilinear Algebra*, **57**, 2009, 409–420.
- [23] M.E.H. Ismail, *Classical and quantum orthogonal polynomials in one variable*, with two chapters by W. Van Assche, with a foreword by R. Askey, Encyclopedia of Mathematics and its Applications **98**, Cambridge University Press, Cambridge, 2005.
- [24] H.L. Krall and O. Frink, A new class of orthogonal polynomials: The Bessel polynomials, *Trans. Amer. Math. Soc.*, **65**, 1949, 100–115.
- [25] M.G. Krein and M.A. Naimark, The method of symmetric and Hermitian forms in the theory of the separation of the roots of algebraic equations, *Linear and Multilinear Algebra*, 1981, **10**, 265–308.
- [26] L. Lorentzen and H. Waadeland, *Continued Fractions with Applications*, North-Holland Publishing Co., Amsterdam, 1992.
- [27] F. Marcellán and R. Álvarez-Nodarse, On the “Favard theorem” and its extensions, *J. Comput. Appl. Math.*, **127**, 2001, 231–254.
- [28] A.W. Mason, Unipotent matrices, modulo elementary matrices, in SL_2 over a coordinate ring, *J. Algebra*, **203**, 1998, 134–155.
- [29] R.D. Morton and A.M. Krall, Distributional weight functions for orthogonal polynomials, *SIAM J. Math. Anal.*, **9**, no. 4, 1978, 604–626.
- [30] F.W.J. Olver, The asymptotic expansions of Bessel functions of large order, *Phil. Trans. Roy. Soc. London Ser. A*, **247**, 1954, 338–368.
- [31] A. Prudnikov, Yu. Brychkov, and O. Marichev, *Integrals and Series, Vol. 1, Elementary functions*, Phys. and Math. Literature, Moscow, 2002, p. 632.
- [32] H.R. Schwarz, Ein Verfahren zur Stabilitätsfrage bei Matrizen-Eigenwertproblemen, *Zeitschrift für angew. Math. Phys.*, **7**, no. 6, 1956, 473–500.

- [33] R.L. Soto, A.I. Julio, and J.H. Alfaro, Permutative universal realizability, *Spec. Matrices*, **9**, 2021, 66–77.
- [34] B.D. Sutton, Numerical construction of structured matrices with given eigenvalues, *Spec. Matrices*, **7**, 2019, 263–271.
- [35] M. Tyaglov, Sign patterns of the Schwarz matrices and generalized Hurwitz polynomials, *Electron. J. Linear Algebra*, **24**, 2012, 215–236.
- [36] H.S. Wall, Polynomials whose zeros have negative real parts, *Amer. Math. Monthly*, **52**, no. 6, 1945, 308–322.
- [37] H.S. Wall, *Analytic Theory of Continued Fractions*, D. Van Nostrand Company Inc., New York, 1948.
- [38] H. Zassenhaus, Characterization of unipotent matrices, *J. Number Theory*, **1**, 1969, 222–230.

KELDYSH INSTITUTE OF APPLIED MATHEMATICS, RUSSIAN ACADEMY OF SCIENCES, 125047, MOSCOW, RUSSIA

Email address: `diachenko@sfedu.ru`

KUWAIT COLLEGE OF SCIENCE AND TECHNOLOGY, DOHA DISTRICT, SAFAT 13133, KUWAIT

CHAIR OF COMPUTATIONAL MATHEMATICS, UNIVERSITY OF DEUSTO, 48007 BILBAO, SPAIN

Email address: `c.dafonseca@kcst.edu.kw`

DEPARTMENT OF MATHEMATICS AND COMPUTER SCIENCES, ST.PETERSBURG STATE UNIVERSITY, ST.PETERSBURG, 199178, RUSSIA

Email address: `tyaglov@mail.ru`