# COMPLEMENTABLE OPERATORS AND THEIR SCHUR COMPLEMENTS

SACHIN MANJUNATH NAIK AND P. SAM JOHNSON

Abstract. In this paper, we characterize complementable operators and provide more precise expressions for the Schur complement of these operators using a single Douglas solution. We demonstrate the existence of subspaces where the given operator is invariably complementable. Additionally, we investigate the range-Hermitian property of these operators.

### 1. INTRODUCTION

Let T be an  $(n + m) \times (n + m)$  matrix over an arbitrary field which is partitioned in the form

<span id="page-0-0"></span>
$$
T = \begin{pmatrix} A & B \\ C & D \end{pmatrix}
$$

with a nonsingular matrix  $D$  of order  $m$ . Then, the classical Schur complement of D in T is the  $n \times n$  matrix defined by

(1.1) 
$$
T_{/D} = A - BD^{-1}C.
$$

The formula [\(1.1\)](#page-0-0) was first used by Schur [\[24\]](#page-16-0). However, this terminology is due to Haynsworth [\[15\]](#page-16-1). Matrices of the form [\(1.1\)](#page-0-0) have undoubtedly been encountered from the time matrices were first used, particularly in Gaussian elimination [\[12,](#page-15-0) [26\]](#page-16-2) and minor determinants [\[25\]](#page-16-3) and others. We remark that the Schur complement in this classical sense is a matrix of lower order, while the Schur complement in our sense is always a matrix of the order of T and it is henceforth denoted by

$$
T_{/A} = \begin{pmatrix} A - BD^{-1}C & 0 \\ 0 & 0 \end{pmatrix}.
$$

The Schur complement has been extended to the case where the block matrix D is rectangular and singular, by replacing the regular inverse by the Moore-Penrose inverse or any other generalized inverse[\[5\]](#page-15-1). This notion has proved to be a fundamental idea in many applications like numerical analysis, statistics, operator inequalities, electrical network theory, discrete-time regulator problem and to name a few.

<sup>2010</sup> Mathematics Subject Classification. 47A64, 47B65.

Key words and phrases. Complementable operators, weakly complementable operators, Schur complement.

Let H and K be real or complex Hilbert spaces. We denote  $\mathcal{B}(\mathcal{H},\mathcal{K})$  the space of all linear bounded operators from H to K and we abbreviate  $\mathcal{B}(\mathcal{H}) =$  $\mathcal{B}(\mathcal{H},\mathcal{H})$ . For  $T \in \mathcal{B}(\mathcal{H},\mathcal{K})$ , we denote by  $T^*, \mathcal{N}(T)$  and  $\mathcal{R}(T)$ , respectively, the adjoint, the null-space and the range-space of T. An operator  $T \in \mathcal{B}(\mathcal{H})$ is said to be a projection if  $T^2 = T$ . A projection, T is said to be orthogonal if  $T = T^*$ . An operator  $T \in \mathcal{B}(\mathcal{H})$  is called positive (denoted  $T \geq 0$ ) if  $T = T^*$  and  $\langle Tx, x \rangle \geq 0$  for all  $x \in \mathcal{H}$ . The cone of positive operators in  $\mathcal{B}(\mathcal{H})$  is denoted by  $\mathcal{B}(\mathcal{H})^+$ . The operator  $|T| := (T^*T)^{1/2}$  is called the modulus of T.

Anderson [\[1\]](#page-15-2) first introduced the definition of the Schur complement for positive operators on Hilbert spaces. Given  $T \in \mathcal{B}(\mathcal{H})^+$ , the shorted operator (or Schur complement) of T with respect to a closed subspace M of  $H$ is defined as

$$
T_{/M} = \max_{\leq} \{ X \in \mathcal{B}(\mathcal{H})^+ : X \leq T \text{ and } \mathcal{R}(X) \subseteq M^{\perp} \}.
$$

The existence of maximum has been studied by Krein [\[18\]](#page-16-4) in the study of symmetric operator extensions. Ando [\[2\]](#page-15-3) introduced the definition of the Schur complement for M-complementable bounded operators. Since then, the Schur complement of bounded linear operators has been attracted extensively by many researchers. See, for example [\[8,](#page-15-4) [9,](#page-15-5) [10,](#page-15-6) [13,](#page-15-7) [22,](#page-16-5) [23\]](#page-16-6). Ando's definition opened the door to various extensions and generalizations, as observed by Butler and Morley [\[6\]](#page-15-8), who compared six different notions of Schur complements for finite dimensions.

Mitra [\[19\]](#page-16-7) generalized the concept of complementable operators to the context of operators between distinct Hilbert spaces, introducing bilateral shorted operators and exploring their properties. Subsequently, Antezana et al. [\[3\]](#page-15-9) formulated the concept of complementability and weak complementability for bounded operators  $T : \mathcal{H} \to \mathcal{K}$  relative to closed subspaces  $M \subseteq \mathcal{H}, N \subseteq \mathcal{K}$ , which gave rise to the bilateral shorted operator and its applications in extending parallel sums of bounded linear operators. A comprehensive survey of the Schur complement can be found in [\[4,](#page-15-10) [8,](#page-15-4) [11,](#page-15-11) [21\]](#page-16-8).

We now describe the contents of the paper. In the next section, we provide a brief background for the rest of the material in the paper. In Section 3, we comprehend complementable bounded operators with more characterizations. Furthermore, for complementable operators relative to closed subspaces  $M$  and  $N$ , we present a new approach to derive the Schur complement using a single Douglas solution. In Section 4, we show that this methodology provides insights into the closed range of complementable operators and their Schur complements relative to the associated subspaces. Also, we provide an extension of a characterization of EP-matrix and its Schur complement given in [\[20\]](#page-16-9) to a hypo-EP operator  $T : \mathcal{H} \to \mathcal{H}$  relative to closed subspaces  $M \subseteq \mathcal{H}$  and  $N \subseteq \mathcal{H}$ .

### 2. Preliminaries

Let M and N be closed subspaces of the Hilbert spaces  $\mathcal H$  and  $\mathcal K$ , respectively. A bounded linear operator  $T \in \mathcal{B}(\mathcal{H}, \mathcal{K})$  can be expressed as

$$
T = \begin{pmatrix} A & B \\ C & D \end{pmatrix}
$$

with regard to the decompositions  $\mathcal{H} = M \oplus M^{\perp}$  and  $\mathcal{K} = N \oplus N^{\perp}$ .

**Definition 2.1.** [\[3\]](#page-15-9) Let  $P_r \in \mathcal{B}(\mathcal{H})$  and  $P_\ell \in \mathcal{B}(\mathcal{K})$  be projections. An operator  $T \in \mathcal{B}(\mathcal{H}, \mathcal{K})$  is called  $(P_r, P_\ell)$ -complementable if there exist operators  $M_r \in \mathcal{B}(\mathcal{H})$  and  $M_\ell \in \mathcal{B}(\mathcal{K})$  such that

(1) 
$$
(I - P_r)M_r = M_r
$$
 and  $(I - P_\ell)TM_r = (I - P_\ell)T$ ,  
(2)  $M_\ell(I - P_\ell) = M_\ell$  and  $M_\ell T(I - P_r) = T(I - P_r)$ .

<span id="page-2-0"></span>**Proposition 2.2.** [\[3\]](#page-15-9) *Let*  $P_r \in \mathcal{B}(\mathcal{H})$  *and*  $P_\ell \in \mathcal{B}(\mathcal{K})$  *be projections with their ranges* M *and* N, *respectively. Then* T *is*  $(P_r, P_\ell)$ -complementable *if and only if*  $\mathcal{R}(C) \subseteq \mathcal{R}(D)$  *and*  $\mathcal{R}(B^*) \subseteq \mathcal{R}(D^*)$ *.* 

It is observed from Proposition [2.2](#page-2-0) that  $(P_r, P_\ell)$ -complementable depends only on the ranges of  $P_r$  and  $P_\ell$ .

**Definition 2.3.** [\[3\]](#page-15-9) An operator  $T \in \mathcal{B}(\mathcal{H}, \mathcal{K})$  is called  $(M, N)$ complementable if it is  $(P_r, P_\ell)$ -complementable for some projections  $P_r$  and  $P_\ell$  with  $\mathcal{R}(P_r) = M$  and  $\mathcal{R}(P_\ell) = N$ .

**Definition 2.4.** [\[3\]](#page-15-9) An operator  $T \in \mathcal{B}(\mathcal{H}, \mathcal{K})$  is called  $(M, N)$ -weakly complementable if

$$
\mathcal{R}(C) \subseteq \mathcal{R}(|D^*|^{\frac{1}{2}}) \text{ and } \mathcal{R}(B^*) \subseteq \mathcal{R}(|D|^{\frac{1}{2}}).
$$

We know that  $\mathcal{R}(S) \subseteq \mathcal{R}(|S^*|^{1/2})$ , for any bounded operator S. By Proposition [2.2,](#page-2-0) it is easy to see that if T is  $(M, N)$ -complementable, then it is  $(M, N)$ -weakly complementable. Both notions of complementability coincide if  $\mathcal{R}(D)$  is closed from the following facts :

(i)  $\mathcal{R}(D)$  is closed if and only if  $\mathcal{R}(D^*)$  is closed;

(ii)  $\mathcal{R}(D) = \mathcal{R}(D^{1/2})$  if and only if  $\mathcal{R}(D)$  is closed;

(iii)  $\mathcal{R}(D) = \mathcal{R}(|D^*|) = \mathcal{R}(|D^*|^{1/2}).$ 

From the above observations, it is noted that if  $T = \begin{pmatrix} A & B \\ C & D \end{pmatrix}$  is  $(M, N)$ complementable with  $\mathcal{R}(D)$  closed, then

$$
T_{/(M,N)} = \begin{pmatrix} A - BD^{\dagger}C & 0 \\ 0 & 0 \end{pmatrix}.
$$

<span id="page-2-1"></span>**Theorem 2.5.** *Let*  $A, B \in \mathcal{B}(\mathcal{H})$ *. Then the following are equivalent:* 

- $(1)$   $\mathcal{R}(A) \subseteq \mathcal{R}(B)$ ;
- (2)  $AA^* \leq \lambda BB^*$ , for some  $\lambda > 0$ ;
- (3) *There exists a bounded operator*  $C \in \mathcal{B}(\mathcal{H})$  *such that*  $A = BC$ *.*

*Moreover, if these equivalent conditions hold, then there is a unique operator*  $C \in \mathcal{B}(\mathcal{H})$  *such that* 

(i)  $||C|| = \inf{\lambda > 0 : AA^* \leq \lambda BB^*};$ (ii)  $\mathcal{N}(A) = \mathcal{N}(B)$ ; (iii)  $\mathcal{R}(C) \subseteq \mathcal{N}(B)^{\perp}$ .

**Definition 2.6.** [\[3\]](#page-15-9) Let T be  $(M, N)$ -weakly complementable. Let F and E be the reduced solutions of the equations  $C = |D^*|^{\frac{1}{2}}UX$  and  $B^* = |D|^{\frac{1}{2}}X$ , respectively, where  $D = U|D|$  is the polar decomposition of D. Then the bilateral shorted operator (the Schur complement) of T with respect to closed subspaces M and N is  $\begin{pmatrix} A - E^*F & 0 \\ 0 & 0 \end{pmatrix}$  and is denoted by  $T_{/(M,N)}$ . If  $\mathcal{H} = \mathcal{K}$  and  $M = N$ , then we denote  $T_{/(M,N)}$  by  $T_{/M}$ .

The above definition relies on the existence of two reduced solutions to define the Schur complement of weakly complementable operators. But for an  $(M, N)$ -complementable operator  $T = \begin{pmatrix} A & B \\ C & D \end{pmatrix}$ , the Schur complement can be elegantly represented by employing either one of the Douglas solutions pertaining to the conditions  $\mathcal{R}(C) \subseteq \mathcal{R}(D)$  or  $\mathcal{R}(B^*) \subseteq \mathcal{R}(D^*)$ .

Consider the reduced Douglas solution Z for the inclusion  $\mathcal{R}(C) \subseteq \mathcal{R}(D)$ . As a result, we can express  $C = DZ = U|D|Z = U|D|^{\frac{1}{2}}|D|^{\frac{1}{2}}Z =$  $|D^*|^{\frac{1}{2}}U|D|^{\frac{1}{2}}Z$ . Therefore,  $|D|^{\frac{1}{2}}Z$  serves as a solution for the equation  $C = |D^*|^{\frac{1}{2}}UX$ . It is also important to note that  $\mathcal{R}(|D|^{\frac{1}{2}}Z) \subseteq \overline{\mathcal{R}(|D|^{\frac{1}{2}})}$ .

Therefore,  $|D|^{\frac{1}{2}}Z$  stands as the reduced solution for the equation  $C =$  $|D^*|^{\frac{1}{2}}UX$ . Furthermore, let  $B = E^*|D|^{\frac{1}{2}}$ . This allows us to express  $BZ =$  $E^*|D|^{\frac{1}{2}}Z = E^*F$  and hence

$$
T_{/(M,N)} = \begin{pmatrix} A - BZ & 0 \\ 0 & 0 \end{pmatrix}.
$$

Moreover, if Y is a solution for the equation  $B = XD$ , then  $BZ = YC$ . Hence, the Schur complement can also be represented as

$$
T_{/(M,N)} = \begin{pmatrix} A - YC & 0 \\ 0 & 0 \end{pmatrix}.
$$

In this context, the condition  $\mathcal{R}(C) \subseteq \mathcal{R}(D)$  is essential for establishing the existence of the operator Z such that  $C = DZ$ . Furthermore, through the following example, we can observe that the condition  $\mathcal{R}(B^*) \subseteq \mathcal{R}(D^*)$ ensures the Schur complement is well-defined. Indeed, if we consider two solutions,  $Z_1$  and  $Z_2$ , such that  $C = DZ_1$  and  $C = DZ_2$ , we find that  $D(Z_1 - Z_2) = 0$  if and only if  $\mathcal{R}(Z_1 - Z_2) \subseteq \mathcal{N}(D) \subseteq \mathcal{N}(B)$ , which implies  $BZ_1 = BZ_2$ . Therefore, it becomes evident that the definition of the Schur complement is independent of the specific choice of Z.

<span id="page-3-0"></span>Theorem 2.7. [\[3\]](#page-15-9) *Let* T *be* (M, N)*-complementable. Then the following statements hold good :*

(1)  $T^*$  *be*  $(N, M)$ *-complementable* ;  $(2) \ \mathcal{R}(T_{/(M,N)}) = \mathcal{R}(T) \cap N;$ (3)  $\mathcal{N}(T_{/(M,N)}) = M^{\perp} + \mathcal{N}(T);$ (4) If  $\mathcal{R}(D)$  is closed, then  $T_{/(M,N)} = \begin{pmatrix} A - BD^{\dagger}C & 0 \ 0 & 0 \end{pmatrix}$ .

<span id="page-4-0"></span>**Theorem 2.8.** [\[14\]](#page-16-10) *Let*  $T$  *be*  $(M, N)$ -complementable with  $\mathcal{R}(C^*) \subseteq$  $\mathcal{R}((T_{/(M,N)})^*), \mathcal{R}(B) \subseteq \mathcal{R}(T_{/(M,N)})$  and  $\mathcal{R}(T)$  closed. Then

$$
T^\dagger = \begin{pmatrix} T_{/M}{}^\dagger & -T_{/M}{}^\dagger B D^\dagger \\ -D^\dagger C T_{/M}{}^\dagger & D^\dagger + D^\dagger C T_{/M}{}^\dagger B D^\dagger \end{pmatrix}.
$$

For a complementable operator, the Schur complement is established through the Douglas solution of the inclusion  $\mathcal{R}(C) \subseteq \mathcal{R}(D)$ . Similarly, in the context of an operator satisfying  $\mathcal{R}(C) \subseteq \mathcal{R}(D)$  and  $\mathcal{R}(B^*) \nsubseteq \mathcal{R}(D^*),$ a Schur complement expression can be formulated, albeit not uniquely. We conclude this section by an illustrative example demonstrating these assertions.

**Example 2.9.** Let  $T = \begin{pmatrix} A & B \\ C & D \end{pmatrix} : \ell_2 \oplus \ell_2 \to \ell_2 \oplus \ell_2$  where A, B, C and D are defined from  $\ell_2$  to  $\ell_2$  as follows :

$$
A(x_1, x_2, \ldots) = \left(0, x_1, \frac{x_2}{2}, \ldots\right);
$$
  
\n
$$
B(x_1, x_2, \ldots) = \left(x_1, x_2, x_3, \ldots\right);
$$
  
\n
$$
C(x_1, x_2, \ldots) = \left(x_1, \frac{x_2}{2}, \frac{x_3}{3} \ldots\right);
$$
  
\n
$$
D(x_1, x_2, \ldots) = \left(x_2, x_3, x_4, \ldots\right).
$$

It is evident that  $\mathcal{R}(D^*) = \ell_2 \setminus \text{span}\{e_1\}, \ \mathcal{R}(C) \subseteq \mathcal{R}(D) = \ell_2$  and  $\mathcal{R}(B^*) \nsubseteq \mathcal{R}(D^*)$ . Define the operators  $Z_1(x_1, x_2, \ldots) = (0, x_1, \frac{x_2}{2})$  $(\frac{c_2}{2}, \dots)$  and  $Z_2(x_1, x_2, \ldots) = (x_1, x_1, \frac{x_2}{2})$  $\left(\frac{x_2}{2}, \ldots\right)$ . It is evident that  $C = DZ_1 = DZ_2$ . Now, consider

$$
(T_{/M_1})(x_1, x_2, x_3, \ldots) = (A - BZ_1)(x_1, x_2, x_3, \ldots) = (x_1, 0, 0, \ldots)
$$

and

$$
(T_{/M_2})(x_1, x_2, x_3, \ldots) = (A - BZ_2)(x_1, x_2, x_3, \ldots) = (0, 0, 0, \ldots).
$$

In accordance with the definition of the Schur complement, both  $(T_M)_1$ and  $(T_M)_2$  qualify as Schur complements; however, it is important to note that  $(T_M)_1$  is not equal to  $(T_M)_2$ .

#### 3. Complementable Operators and Schur Complements

Butler and Moorley's [\[6\]](#page-15-8) pioneering work illuminates a geometric pathway to comprehend complementable operators within finite dimensions. Drawing inspiration from their groundbreaking insights, our research embarks on an exploration into the vast domain of infinite-dimensional operators. By pushing the boundaries of traditional understanding, we aim to unveil new perspectives and insights that can reshape the landscape of operator theory and its applications.

<span id="page-5-0"></span>**Theorem 3.1.** Let  $T \in \mathcal{B}(\mathcal{H}, \mathcal{K})$  with  $\mathcal{R}(D)$  closed. Then T is  $(M, N)$ *complementable if and only if for each*  $x \in M$ *, there exists unique*  $z \in M$ *such that*  $\{T(x,0) + T(M^{\perp})\} \cap N = \{(z,0)\}\$ *. Moreover,*  $T_{/(M,N)}(x,0) =$  $(z, 0)$ .

*Proof.* Let Z be such that  $C = DZ$ . Then  $T(x, 0) + T(0, -Zx) = ((A BZ(x,0)$ . Therefore,  $\{T(x,0) + T(M^{\perp})\} \cap N$  is non-empty.

Now, assume  $(z_1, 0), (z_2, 0) \in \{T(x, 0) + T(M^{\perp})\} \cap N$ . There exist  $y_1, y_2 \in$  $M^{\perp}$  such that  $T(x, y_1) = (z_1, 0)$  and  $T(x, y_2) = (z_2, 0)$ . Hence  $Ax + By_1 =$  $z_1, Cx + Dy_1 = 0, Ax + By_2 = z_2, and Cx + Dy_2 = 0.$  Thus,  $y_1 - y_2 \in$  $\mathcal{N}(D) \subseteq \mathcal{N}(B)$ , leading to  $By_1 = By_2$ . This implies  $(z_1, 0) = (z_2, 0)$ .

Conversely, assume that  $\mathcal{R}(D)$  is closed. For each  $x \in M$ , let  $\{T(x, 0) +$  $T(M^{\perp})\}\cap N = \{(z,0)\}\)$  for some  $z \in M$ . Then there exists  $y \in M^{\perp}$  such that  $T(x, y) = (z, 0)$ , implying  $Cx + Dy = 0$ . Thus  $\mathcal{R}(C) \subseteq \mathcal{R}(D)$ .

Let  $y \in \mathcal{N}(D)$ . Then  $T(0,0) + T(0,y) = (By, 0) \in N$ . Since  $T(0,0)$  +  $T(0, 0) = (0, 0) \in N$  and  $\{T(x, 0) + T(M^{\perp})\} \cap N$  is a singleton, we have  $By = 0$ . Therefore  $\mathcal{N}(D) \subseteq \mathcal{N}(B)$ , leading to  $\overline{\mathcal{R}(B^*)} \subseteq \overline{\mathcal{R}(D^*)}$ . Since  $\mathcal{R}(D)$ is closed, we obtain  $\mathcal{R}(B^*) \subseteq \mathcal{R}(D^*)$ .

As observed earlier, we have  $T(x, 0) + T(M^{\perp}) \cap N = z = ((A - BZ)x, 0).$ Thus  $T_{/(M,N)}(x,0) = (z,0).$ 

In the following example, we find Schur complement of an operator using our characterization.

**Example 3.2.** Let  $T: \ell_2 \to \ell_2$  be defined by

$$
T(x_1, x_2,...) = (x_3 + x_2 - x_1, x_1 + x_2, x_5 + x_4 - x_3, \frac{x_3}{3} + x_4,...).
$$

Consider the subspaces

$$
M = \{x = (x_1, 0, x_3, 0, \ldots) : x \in \ell_2\}
$$

and

$$
M^{\perp} = \{x = (0, x_2, 0, x_4, 0, \ldots) : x \in \ell_2\}.
$$

Let  $x = (x_1, 0, x_3, 0, ...) \in M$ . Then  $Tx = (x_3 - x_1, x_1, x_5 - x_3, \frac{x_3}{3})$  $\frac{c_3}{3}, \ldots$ ). Moreover,  $T(M^{\perp}) = \{(x_2, x_2, x_4, x_4, \ldots) : (0, x_2, 0, x_4, \ldots) \in M^{\perp}\}.$ Thus,  $Tx + T(M^{\perp}) = \{(x_3 + x_2 - x_1, x_1 + x_2, x_5 + x_4 - x_3, \frac{x_3}{3} + x_4)\}$  $x_4, \ldots$ ) :  $(0, x_2, 0, x_4, \ldots) \in M^{\perp}$ .

Let  $z \in (T(x) + T(M^{\perp})) \cap M$ . Then  $z = Tx + Ty$  for some  $y =$  $(0, x_2, 0, x_4, 0, ...) \in M^{\perp}$ . Therefore,  $z = (x_3 + x_2 - x_1, x_1 + x_2, x_5 + x_4$  $x_3, \frac{x_3}{3} + x_4, \ldots$ ). Since  $z \in M$ , we have  $\frac{x_n}{n} + x_{n+1} = 0$  for all odd integers n. Hence  $x_{n+1} = -\frac{x_n}{n}$  $\frac{x_n}{n}$  for all odd integers n. So,  $z = (x_3 - 2x_1, 0, x_5 - \frac{4x_3}{3})$  $\frac{x_3}{3}, 0, \ldots$ ). Therefore  $T(x) + T(M^{\perp})$  is a singleton and

$$
T_{/M}(x_1, x_2, \ldots) = z = \left(x_3 - 2x_1, 0, x_5 - \frac{4x_3}{3}, 0, \ldots\right).
$$

We proved that the 'singleton condition' in the above theorem is a necessary and sufficient for the complementability of an operator  $T = \begin{pmatrix} A & B \\ C & D \end{pmatrix}$  with  $\mathcal{R}(D)$  closed. Without the assumption of the closed range of the operator D, the 'singleton condition' is only necessary but not sufficient, which is illustrated in the following example.

**Example 3.3.** Let  $T: \ell_2 \to \ell_2$  be defined by

$$
T(x_1, x_2,...) = \left(x_1 + \frac{x_2}{2}, x_1 + \frac{x_2}{2}, x_3 + \frac{x_4}{4}, x_3 + \frac{x_4}{4}, \dots\right).
$$

Consider the subspaces

$$
M = \{x = (x_1, 0, x_3, 0, \ldots) : x \in \ell_2\}
$$

and

$$
M^{\perp} = \{x = (0, x_2, 0, x_4, 0, \ldots) : x \in \ell_2\}.
$$

Therefore

$$
A(x_1, 0, x_3, 0, \ldots) = (x_1, 0, x_3, 0, \ldots);
$$
  
\n
$$
B(0, x_2, 0, x_4, 0, \ldots) = (\frac{x_2}{2}, 0, \frac{x_4}{4}, 0, \ldots);
$$
  
\n
$$
C(x_1, 0, x_3, 0, \ldots) = (0, x_1, 0, x_3, 0, \ldots);
$$
  
\n
$$
D(0, x_2, 0, x_4, 0, \ldots) = (0, \frac{x_2}{2}, 0, \frac{x_4}{4}, 0, \ldots).
$$

Clearly  $\mathcal{R}(D)$  is not closed and  $\mathcal{R}(C) \nsubseteq \mathcal{R}(D)$ . So T is not Mcomplementable.

Let  $x = (x_1, 0, x_3, 0, ...) \in M$ . Then  $Tx = (x_1, x_1, x_3, x_3, ...)$ . Thus  $Tx + T(M^{\perp}) = \{(x_1 + \frac{x_2}{2})\}$  $\frac{x_2}{2}, x_1 + \frac{x_2}{2}$  $\frac{x_2}{2}, x_3 + \frac{x_4}{4}$  $\frac{x_4}{4}$ ,  $x_3 + \frac{x_4}{4}$  $\left( \begin{matrix} \frac{c_4}{4}, \ldots \end{matrix} \right) : (0, x_2, 0, x_4, \ldots) \in$  $M^{\perp}$ . Suppose  $z \in (T(x) + T(M^{\perp})) \cap M$ . Then  $z = Tx + Ty$  for some  $y = (0, x_2, 0, x_4, 0, ...) \in M^{\perp}$ . Therefore,  $z = (x_1 + \frac{x_2}{2})$  $\frac{x_2}{2}, x_1 + \frac{x_2}{2}$  $\frac{x_2}{2}, x_3 + \frac{x_4}{4}$  $y = (0, x_2, 0, x_4, 0, ...) \in M^{\perp}$ . Therefore,  $z = (x_1 + \frac{x_2}{2}, x_1 + \frac{x_2}{2}, x_3 + \frac{x_4}{4}, x_3 + \frac{x_5}{24})$ <br>Since  $z \in M$  we have  $\frac{x_{n+1}}{2} + x_{n+1} = 0$  for all odd integers n So we  $\frac{x_4}{4}, \ldots$ ). Since  $z \in M$ , we have  $\frac{x_{n+1}}{n+1} + x_n = 0$  for all odd integers n. So, we get  $z = 0$ . Therefore,  $T(x) + T(M^{\perp})$  is a singleton.

It is natural to ask, given  $T \in \mathcal{B}(\mathcal{H})$ , whether there exist closed subspaces M and N such that T is  $(M, N)$  complementable. We affirmatively answer the question in the following results.

<span id="page-6-0"></span>**Theorem 3.4.** *Let*  $T \in \mathcal{B}(\mathcal{H}, \mathcal{K})$  *and* N *be a closed subspace of* K. *Let*  $M =$  $(T^{-1}(N^{\perp}))^{\perp}$ . Then T is  $(M, N)$ -complementable and  $T_{/(M, N)}(x) = P_N Tx$ .

*Proof.* Let  $T = \begin{pmatrix} A & B \\ C & D \end{pmatrix}$ . Given that  $T(M^{\perp}) = N^{\perp}$ , it follows that  $\mathcal{R}(D) =$  $\mathcal{R}(P_{N^{\perp}}TP_{M^{\perp}})=P_{N^{\perp}}T(M^{\perp})=P_{N^{\perp}}(N^{\perp})=N^{\perp},$  so  $\mathcal{R}(D)$  is closed. For each  $x \in M$ , choose  $y \in \{Tx + T(M^{\perp})\} \cap N = \{Tx + N^{\perp}\} \cap N$ . This implies that  $y = Tx + z = P<sub>N</sub> Tx + P<sub>N</sub> \perp Tx + z$ , for some  $z \in N^{\perp}$ , so  $y \in N$ . As  $z \in$  $N^{\perp}$  and  $y \in N$ , we obtain  $P_{N^{\perp}} Tx = -z$ . Therefore,  $y = P_N Tx + P_{N^{\perp}} Tx +$   $z = P<sub>N</sub> Tx$ . Hence  $\{Tx + T(M^{\perp})\} \cap N = \{P<sub>N</sub> Tx\}$ , a singleton set. So by Theorem [3.1,](#page-5-0) T is  $(M, N)$ -complementable and  $T_{/(M, N)}(x) = P_N T x$ .

**Corollary 3.5.** *Let*  $T \in \mathcal{B}(\mathcal{H}, \mathcal{K})$  *and*  $M$  *be a closed subspace of*  $\mathcal{H}$ *. Let*  $N = ((T^*)^{-1}(M^{\perp}))^{\perp}$ . Then T is  $(M, N)$ -complementable.

*Proof.* By Theorem [3.4,](#page-6-0) we have  $T^*$  is  $(N, M)$ -complementable. Hence T is  $(M, N)$ -complementable by Theorem [2.7.](#page-3-0)

**Proposition 3.6.** *Let*  $T \in \mathcal{B}(\mathcal{H}, \mathcal{K})$ *. Then there exist closed subspaces* M *and* N *such that* T *is* (M, N)*-complementable.*

*Proof.* Let  $x_1$  be a unit vector in  $H$  such that  $T(x_1) \neq 0$ . Consider an orthonormal basis  $\{x_{\lambda}\}\$  of H containing  $x_1$ , and let  $\{y_{\beta}\}\$  be an orthonormal basis of K. Also, there exist non-zero coefficients  $\beta_1, \beta_2, \ldots$ , and vectors  $y_1, y_2, \ldots$  in  $\{y_\lambda\}$  such that  $T(x_1) = \sum_{m=1}^{\infty} \beta_i y_i$ . Define  $M = (\text{span}\{x_1\})^{\perp}$ and  $N = (\text{span}\{y_1\})^{\perp}$ . The operator is expressed as  $T = \begin{pmatrix} A & B \\ C & D \end{pmatrix}$  with respect to the decomposition  $M \oplus M^{\perp}$  and  $N \oplus N^{\perp}$ . Since  $x_1 \in M^{\perp}$ , we have  $TP_{M^{\perp}}x_1 = \sum_{m=1}^{m}$  $\frac{i=1}{i}$  $\beta_i y_i$ . This implies  $P_{N^{\perp}} T P_{M^{\perp}} x_1 = \beta_1 y_1$ , making  $\mathcal{R}(P_{N^{\perp}}TP_{M^{\perp}})$  is an one-dimensional space. So,  $\mathcal{R}(D)$  is a closed subspace.

Let 
$$
x \in M
$$
. Then  $x = \sum_{i=1}^{\infty} \langle x, x_i \rangle x_i$ . Therefore,  $Tx = \sum_{i=1}^{\infty} \langle x, x_i \rangle Tx_i$ . Since  $Tx_i = \sum_{i=1}^{\infty} \langle Tx_i, y_i \rangle y_i$ , we have

$$
Tx = \sum_{i=1}^{\infty} (\langle x, x_i \rangle \Big( \sum_{j=1}^{\infty} \langle Tx_i, y_j \rangle y_j) \Big) = \sum_{j=1}^{\infty} \Big( \sum_{i=1}^{\infty} \langle x, x_i \rangle \langle Tx_i, y_j \rangle \Big) y_j.
$$

Let  $w \in \{Tx + T(M^{\perp})\} \cap N$ . This implies that  $w = Tx + Ty$  for some  $y \in M^{\perp}$ . Since  $y \in M^{\perp}$ ,  $y = \alpha_0 x_1$  for some  $\alpha_0$ . Therefore

$$
Ty = \alpha_0 Tx_1 = \alpha_0 \sum_{k=1}^{\infty} \langle Tx_1, y_k \rangle y_k.
$$

Consequently,  $Tx + Ty = \sum_{j=1}^{\infty} \left( \sum_{i=1}^{\infty} \right)$  $\langle x, x_i \rangle \langle Tx_i, y_j \rangle \bigg) y_j + \alpha_0 \sum_{i=1}^{\infty}$  $k=1$  $\langle Tx_1, y_k\rangle y_k.$ Since  $Tx + Ty \in N$ , the coefficient of  $y_1 = 0$ . Therefore

$$
\alpha_0 = -\frac{\sum_{i=1}^{\infty} \langle x, x_i \rangle \langle Tx_i, y_1 \rangle}{\langle Tx_1, y_1 \rangle}
$$

.

This says that there is only one  $y \in M^{\perp}$  such that  $Tx + Ty \in N$ . Therefore  ${T x + T(M^{\perp})}\cap N = {T x + T(\alpha_0 x_1)}$ , a singleton set. Hence, by Theorem

[3.1,](#page-5-0) T is  $(M, N)$ -complementable and

$$
T_{/(M,N)}(x) = Tx + T(\alpha_0 x_1)
$$
  
= 
$$
\sum_{j=1}^{\infty} \left( \sum_{i=1}^{\infty} \langle x, x_i \rangle \langle Tx_i, y_j \rangle \right) y_j + \alpha_0 Tx_1
$$
  
= 
$$
\sum_{j=1}^{\infty} \left( \sum_{i=1}^{\infty} \langle x, x_i \rangle \langle Tx_i, y_j \rangle \right) y_j - \left( \frac{\sum_{i=1}^{\infty} \langle x, x_i \rangle \langle Tx_i, y_1 \rangle}{\langle Tx_1, y_1 \rangle} \right) Tx_1.
$$
  
nis completes the proof

This completes the proof.

**Example 3.7.** Let  $T: \ell_2 \to \ell_2$  be defined by

 $T(x_1, x_2, \ldots) = (x_1 + x_2, x_1 + x_2 + x_3, x_2 + x_3 + x_4, x_3 + x_4 + x_5, \ldots).$ and  $M = N = (\{e_1\})^{\perp}$ . With respect to the decompositions  $M \oplus M^{\perp}$  and  $N \oplus N^{\perp}$ , we have the following operators :

$$
A(0, x_2, x_3, \ldots) = (0, x_2 + x_3, x_2 + x_3 + x_4, x_3 + x_4 + x_5, \ldots);
$$
  
\n
$$
B(x_1, 0, 0, \ldots) = (0, x_1, 0, 0, \ldots);
$$
  
\n
$$
C(0, x_2, x_3, \ldots) = (x_2, 0, 0, \ldots);
$$
  
\n
$$
D(x_1, 0, 0, \ldots) = (x_1, 0, 0, \ldots).
$$

It is evident that T is  $(M, N)$ -complementable, and all of the operators A,  $B, C,$  and  $D$  are non-zero. This serves as a demonstration that the operators  $A, B, C$ , and  $D$  in the previous proposition may not necessarily be zero.

The current conceptualization and various characterizations of complementable operators are rooted in algebraic considerations, contingent upon the range of these operators. In this context, we introduce an analytical characterization for complementable operators. This characterization pivots on the properties of the image of the unit ball under these operators.

For a subspace M of a Hilbert H, we denote the set  $\{x \in M : ||x|| \leq 1\}$ as  $\mathcal{B}_M$ .

<span id="page-8-0"></span>**Theorem 3.8.** *Let*  $T \in \mathcal{B}(\mathcal{H}, \mathcal{K})$ *. Then*  $T$  *is*  $(M, N)$ *-complementable if and only if there exists*  $\lambda > 0$  *such that*  $C(\mathcal{B}_M) \subseteq \lambda D(\mathcal{B}_{M^{\perp}})$  *and*  $B^*(\mathcal{B}_M) \subseteq$  $\lambda D^*(\mathcal{B}_{M^{\perp}})$ .

*Proof.* Theorem [2.5](#page-2-1) says that the inclusion  $\mathcal{R}(C) \subseteq \mathcal{R}(D)$  gives the existence of an operator  $Z \in \mathcal{B}(\mathcal{H})$  such that  $C = DZ$ . Suppose C is non-zero. It follows that Z is also non-zero, leading to the conclusion that  $||Z|| \neq 0$ . Let  $x \in \mathcal{B}_{\mathcal{H}}$ . We then observe that

$$
Cx = DZx = ||Z|| \left( D\left(\frac{Zx}{||Z||}\right) \right).
$$

Now we consider  $y = \frac{Zx}{\|Z\|}$  $\frac{Zx}{\|Z\|}$ . So we have  $\|y\| = \left\|\frac{Zx}{\|Z\|}\right\|$  $\frac{Zx}{\|Z\|}\| \leq 1$ , and  $y \in \mathcal{R}(Z) \subseteq \mathcal{H}$ . Therefore, we establish  $C(\mathcal{B}_{\mathcal{H}}) \subseteq ||Z||D(\mathcal{B}_{\mathcal{H}})$ .

Conversely, let us assume the existence of a positive constant  $\lambda$  such that  $C(\mathcal{B}_{\mathcal{H}}) \subseteq \lambda D(\mathcal{B}_{\mathcal{H}})$ . Let  $w \in \mathcal{R}(C)$ , represented as  $w = Cx$  for some  $x \in \mathcal{H}$ . We observe that  $C\left(\frac{x}{\ln x}\right)$  $\left( \frac{x}{\|x\|} \right) \in C(\mathcal{B}_{\mathcal{H}}) \subseteq \lambda D(\mathcal{B}_{\mathcal{H}}).$  Accordingly,  $C\left( \frac{x}{\|x\|} \right)$  $\frac{x}{\|x\|}\bigg) =$  $\lambda D(y)$  for some  $y \in \mathcal{B}_{\mathcal{H}}$ . This implies  $Cx = \lambda ||x|| D(y) = D(\lambda ||x||y)$ . Hence we conclude  $\mathcal{R}(C) \subseteq \mathcal{R}(D)$ .

Moreover,  $||Z|| = \inf\{\lambda > 0 : C(\mathcal{B}_{\mathcal{H}}) \subseteq \lambda D(\mathcal{B}_{\mathcal{H}})\}\.$  Indeed, suppose that  $C(\mathcal{B}_{\mathcal{H}}) \subseteq \lambda D(\mathcal{B}_{\mathcal{H}})$  for some  $\lambda > 0$ . For each  $x \in \mathcal{B}_{\mathcal{H}}$ , there exists  $y \in \mathcal{B}_{\mathcal{H}} \cap \mathcal{N}(D)^{\perp}$  such that  $Cx = \lambda Dy$  and also  $Cx = DZx$ with  $Zx \in \mathcal{N}(D)^{\perp}$ . Therefore, we have  $D(\lambda y - Zx) = 0$ , leading to  $\lambda y - Zx \in \mathcal{N}(D) \cap \mathcal{N}(D)^{\perp}$ . Therefore,  $Zx = \lambda y$ . As  $||y|| \leq 1$ , we deduce  $||Zx|| = \lambda ||y|| \leq \lambda$  for all  $x \in \mathcal{B}_{\mathcal{H}}$ . Hence, we establish  $||Z|| \leq \lambda$ .

Expanding upon previous arguments, we establish the existence of positive constants  $\lambda_1$  and  $\lambda_2$  such that  $C(\mathcal{B}_M) \subseteq \lambda_1 D(\mathcal{B}_{M^{\perp}})$  and  $B^*(\mathcal{B}_M) \subseteq$  $\lambda_2 D^*(\mathcal{B}_{M^{\perp}})$ . Let  $\lambda = \max{\lambda_1, \lambda_2} > 0$ . Now, we have  $\lambda_1 D(\mathcal{B}_{M^{\perp}}) \subseteq$  $\lambda D(\mathcal{B}_{M^{\perp}})$  and  $\lambda_2 D^*(\mathcal{B}_{M^{\perp}}) \subseteq \lambda D^*(\mathcal{B}_{M^{\perp}})$ . Hence we have  $C(\mathcal{B}_M) \subseteq$  $\lambda D(\mathcal{B}_{M^{\perp}})$  and  $B^*(\mathcal{B}_M) \subseteq \lambda D^*(\mathcal{B}_{M^{\perp}})$ .

Let  $T \in \mathcal{B}(\mathcal{H}, \mathcal{K})$  be an operator. The reduced minimum modulus of T is defined as

$$
\gamma(T) = \inf \{ ||Tx|| : x \in \mathcal{H} \cap \mathcal{N}(T)^{\perp} \}.
$$

Moreover,  $\mathcal{R}(T)$  is closed if and only if  $\gamma(T) > 0$  [\[16\]](#page-16-11). We now characterize algebraically  $(M, N)$ -complementable operator  $T = \begin{pmatrix} A & B \\ C & D \end{pmatrix}$  with respect to the reduced minimum modulus of the operator D.

**Theorem 3.9.** Let  $T \in \mathcal{B}(\mathcal{H}, \mathcal{K})$  with  $\mathcal{R}(D)$  closed. Then T is  $(M, N)$ *complementable if and only if*  $C(\mathcal{B}_M) \subseteq \frac{\|C\|}{\gamma(D)}D(\mathcal{B}_{M^{\perp}})$  *and*  $B^*(\mathcal{B}_M) \subseteq$  $\frac{\|B\|}{\gamma(D^*)}D^*(\mathcal{B}_{M^\perp}).$ 

*Proof.* Let T be  $(M, N)$ -complementable and  $x \in \mathcal{B}_{\mathcal{H}}$ . Since  $\mathcal{R}(C) \subseteq \mathcal{R}(D)$ , there exists  $y \in (\mathcal{N}(D)^{\perp}) \cap M^{\perp}$  such that  $Cx = Dy$ . As  $\mathcal{R}(D)$  is closed, we have

$$
||y||\gamma(D) \le ||y|| ||D(\frac{y}{||y||})|| = ||Dy|| = ||Cx|| \le ||C||.
$$

Therefore the norm of y is bounded by  $\frac{\|C\|}{\gamma(D)}$ . Therefore, y can be expressed as  $y = ky_0$ , where  $0 \leq k \leq \frac{||C||}{\gamma(D)}$  $\frac{\|C\|}{\gamma(D)}$ , and  $y_0 = \frac{y}{\|y\|}$  $\frac{y}{\|y\|}$ . Thus,  $y = \frac{\|C\|}{\gamma(D)}$  $\frac{\|C\|}{\gamma(D)}\left(k\frac{\gamma(D)}{\|C\|}\right)$  $\frac{\gamma(D)}{\|C\|}y_0\Big).$ Since  $k \frac{\gamma(D)}{\|C\|} \leq 1$ , it follows that  $y = \frac{\|C\|}{\gamma(D)}$  $\frac{\|C\|}{\gamma(D)}z$  for some  $||z|| \leq 1$ . Hence,  $C(\mathcal{B}_M)$ is contained in  $\frac{\|C\|}{\gamma(D)}D(\mathcal{B}_{M^{\perp}})$ . In a similar way, the same argument can be made for  $B^*$  and  $D^*$ , leading to the inclusion

$$
B^*(\mathcal{B}_M)\subseteq \frac{\|B^*\|}{\gamma(D^*)}D^*(\mathcal{B}_{M^\perp}).
$$

Conversely, if  $C(\mathcal{B}_M) \subseteq \frac{\|C\|}{\gamma(D)}D(\mathcal{B}_{M^{\perp}})$  and  $B^*(\mathcal{B}_M) \subseteq \frac{\|B^*\|}{\gamma(D^*)}D^*(\mathcal{B}_{M^{\perp}})$ , then by Theorem [3.8,](#page-8-0) T is  $(M, N)$ -complementable.

## 4. Schur Complements of EP and Hypo-EP Operators

Motivated by the Schur determinant formula

$$
det(T) = det(A) \, det(T_A)
$$

for matrices, where  $T_{/A}$  is given in [\(1.1\)](#page-0-0), we analyze the closedness of the range of  $(M, N)$ -complementable operator.

<span id="page-10-0"></span>**Theorem 4.1.** *Let*  $T \in \mathcal{B}(\mathcal{H}, \mathcal{K})$  *be*  $(M, N)$ *-complementable. Then*  $\mathcal{R}(T)$  *is closed if and only if*  $\mathcal{R}(T_{/(M,N)})$  *and*  $\mathcal{R}(D)$  *are closed.* 

*Proof.* Let Z and Y be such that  $C = DZ$  and  $B = YD$ . This implies the following matrix equality:

$$
\begin{pmatrix} A & B \\ C & D \end{pmatrix} = \begin{pmatrix} I & Y \\ 0 & I \end{pmatrix} \begin{pmatrix} T_{/(M,N)} & 0 \\ 0 & D \end{pmatrix} \begin{pmatrix} I & 0 \\ Z & I \end{pmatrix}
$$

where  $\begin{pmatrix} I & Y \\ 0 & I \end{pmatrix}$  $0$  *I* ) and  $\begin{pmatrix} I & 0 \\ Z & I \end{pmatrix}$  are invertible.

Assuming  $\mathcal{R}(T)$  is closed, we can express the above equality as:

$$
\begin{pmatrix} T_{/(M,N)} & 0 \ 0 & D \end{pmatrix} = \begin{pmatrix} I & -Y \\ 0 & I \end{pmatrix} \begin{pmatrix} A & B \\ C & D \end{pmatrix} \begin{pmatrix} I & 0 \\ -Z & I \end{pmatrix}
$$

.

Let  $P = \begin{pmatrix} I & Y \\ 0 & I \end{pmatrix}$  $0$  *I*  $\left( \begin{matrix} I & 0 \\ Z & I \end{matrix} \right)$  and  $U = \begin{pmatrix} T_{/(M,N)} & 0 \\ 0 & D \end{pmatrix}$  $0$   $D$  . Now, analyzing the range of  $U$ , we have

$$
\mathcal{R}(U) = U(\mathcal{H})
$$
  
=  $(P^{-1}TQ^{-1})(\mathcal{H})$   
=  $P^{-1}T(Q^{-1}(\mathcal{H}))$   
=  $(P^{-1}T)(\mathcal{H})$   
=  $P^{-1}(\mathcal{R}(T)).$ 

Since  $\mathcal{R}(T)$  is closed and P is bounded,  $\mathcal{R}(U)$  is closed. However,  $\mathcal{R}(U)$  =  $\mathcal{R}(T_{/(M,N)})\oplus \mathcal{R}(D)$ . Therefore, both  $\mathcal{R}(T_{/(M,N)})$  and  $\mathcal{R}(D)$  are closed.

Conversely, assuming  $\mathcal{R}(T_{/(M,N)})$  and  $\mathcal{R}(D)$  are closed, we observe that  $\mathcal{R}(T) = PUQ(\mathcal{H}) = PU(\mathcal{H}).$  Now  $U(\mathcal{H}) = \mathcal{R}(T_{/(M,N)}) \oplus \mathcal{R}(D).$  Given the closure of  $\mathcal{R}(T_{/(M,N)})$  and  $\mathcal{R}(D), U(\mathcal{H}) = \mathcal{R}(T_{/(M,N)}) \oplus \mathcal{R}(D)$  is closed. Because  $P^{-1}$  is bounded,  $P(U(\mathcal{H}))$  is closed. Therefore,  $\mathcal{R}(T)$  is closed.  $\Box$ 

The following example provides a  $(M, N)$ -complementable operator whose range is not closed.

**Example 4.2.** Let  $T: \ell_2 \to \ell_2$  be defined by

$$
T(x_1, x_2,...) = \left(x_1, x_2, \frac{x_3}{3}, x_4, \frac{x_5}{5}, x_6,...\right).
$$

Consider the subspaces

$$
M = \{(x_1, x_1, x_3, x_3, \ldots)\} \text{ and } M^{\perp} = \{(x_2, -x_2, x_4, -x_4, \ldots)\}.
$$

We have  $T = \begin{pmatrix} A & B \\ C & D \end{pmatrix}$  where A, B, C and D are given by

$$
A(x_1, x_1, x_3, x_3 \dots) = \frac{1}{2} \Big( 2x_1, 2x_1, \frac{4}{3}x_3, \frac{4}{3}x_3, \dots \Big);
$$
  
\n
$$
B(x_1, -x_1, x_3, -x_3 \dots) = \frac{1}{2} \Big( 0, 0, -\frac{2}{3}x_3, -\frac{2}{3}x_3, -\frac{4}{5}x_5, -\frac{4}{5}x_5, \dots \Big);
$$
  
\n
$$
C(x_1, x_1, x_3, x_3 \dots) = \frac{1}{2} \Big( 0, 0, -\frac{2}{3}x_3, \frac{2}{3}x_3, -\frac{4}{5}x_5, \frac{4}{5}x_5, \dots \Big);
$$
  
\n
$$
D(x_1, -x_1, x_3, -x_3 \dots) = \frac{1}{2} \Big( 2x_1, -2x_1, \frac{4}{3}x_3, -\frac{4}{3}x_3, \dots \Big).
$$

Clearly,  $\mathcal{R}(T)$  is not closed and  $\mathcal{R}(D)$  is closed. Now by definition we have

$$
T_{/M}(x_1,x_2,x_3,x_4,\ldots)=\frac{1}{2}\Big(0,0,x_3,x_3,\frac{4}{6}x_5,\frac{4}{6}x_5,\frac{4}{8}x_7,\frac{4}{8}x_7,\ldots\Big).
$$

So  $\mathcal{R}(T_M)$  is not closed.

It is established in [\[20\]](#page-16-9) that under certain conditions a Schur complement of an EP-matrix is EP again. Motivated by this work, we now discuss Schur complements of EP and hypo-EP operators. An operator  $T \in \mathcal{B}(\mathcal{H})$  is said to be EP (hypo-EP) if  $\mathcal{R}(T)$  is closed and  $\mathcal{R}(T) = \mathcal{R}(T^*) (\mathcal{R}(T) \subseteq \mathcal{R}(T^*))$ [\[7,](#page-15-12) [17\]](#page-16-12).

<span id="page-11-0"></span>**Theorem 4.3.** *Let*  $T \in \mathcal{B}(\mathcal{H})$  *be a*  $(M, N)$ *-complementable operator. If*  $T$ *is hypo-EP*, then  $T_{/(M,N)}$  *is hypo-EP*.

*Moreover, if*  $\mathcal{R}(C^*) \subseteq \mathcal{R}((T_{/(M,N)})^*)$ , then the following are equivalent:

(1) T *is hypo-*EP*;* (2)  $T_{/(M,N)}$  and D are hypo-EP and  $\mathcal{R}(B) \subseteq \mathcal{R}(T_{/(M,N)})$ ; (3)  $\begin{pmatrix} T/(M,N) & 0 \\ C & D \end{pmatrix}$  and  $\begin{pmatrix} T/(M,N) & B \\ 0 & D \end{pmatrix}$  $0$  D  $\bigg\}$  are hypo-EP.

*Proof.* Suppose that T is hypo-EP. Theorem [2.7](#page-3-0) establishes that  $\mathcal{R}(T_{/(M,N)}) = \mathcal{R}(T) \cap N$  and  $\mathcal{R}((T^*_{/(M,N)})) = \mathcal{R}((T_{/(M,N)})^*) = \mathcal{R}(T^*) \cap N$ . The hypo-EP property of T implies  $\mathcal{R}(T_{/(M,N)}) \subseteq \mathcal{R}((T_{/(M,N)})^*)$ . Furthermore, given the closed nature of both  $\mathcal{R}(T)$  and N, it follows that  $\mathcal{R}(T_{/(M,N)})$  is closed. Therefore,  $T_{/(M,N)}$  is hypo-EP.

(1)  $\implies$  (2): Already we have proved that  $T_{/(M,N)}$  is hypo-EP, if T is hypo-EP. Considering the matrix operators  $P = \begin{pmatrix} I & BD^{\dagger} \\ 0 & I \end{pmatrix}$  $0 \quad I$  $\bigg),$  $Q = \begin{pmatrix} I & 0 \\ C T & 1 & I \end{pmatrix}$  $CT_{/(M,N)}$ † I ), and  $L = \begin{pmatrix} T/(M,N) & 0 \\ 0 & T \end{pmatrix}$  $0$   $D$ ), we can express  $PQL$  as  $PQL = \begin{pmatrix} I + BD^{\dagger}C(T_{/(M,N)})^{\dagger} & BD^{\dagger} \\ C(T & \lambda^{\dagger})^{\dagger} & I \end{pmatrix}$ C(T/(M,N) ) † I  $\bigcap_{l\in (M,N)}$  0  $0$  D  $\setminus$  $=\begin{pmatrix} T_{/(M,N)}+BD^\dagger C(T_{/(M,N)})^\dagger T_{/(M,N)} & BD^\dagger D \ 2T & (T_{/(M,N)})^\dagger T_{/(M,N)} & DD^\dagger D \end{pmatrix}$  $CT_{/(M,N)}(T_{/(M,N)})^{\dagger}$  D  $\setminus$  $= \begin{pmatrix} A & B \\ C & D \end{pmatrix}.$ 

This implies that  $\mathcal{N}(L) = \mathcal{N}(T)$ . Taking the orthogonal complement on both sides yields  $\mathcal{R}(T^*) = \mathcal{R}(L^*)$ . Since T is hypo-EP, we have  $\mathcal{R}(T) \subseteq$  $\mathcal{R}(L^*)$ . Additionally,  $\mathcal{R}(L)$  is closed. Thus,

$$
T^* = T^* L^{\dagger} L = \begin{pmatrix} A^*(T_{/(M,N)})^{\dagger} T_{/(M,N)} & C^* D^{\dagger} D \\ B^*(T_{/(M,N)})^{\dagger} T_{/(M,N)} & D^* D^{\dagger} D \end{pmatrix}.
$$

This leads to the relationships  $B^* = B^* (T_{/(M,N)})^{\dagger} T_{/(M,N)}$  and  $D^* = D^* D^{\dagger} D$ . As a result,  $\mathcal{R}(B) \subseteq \mathcal{R}(T_{/(M,N)})$  and  $\mathcal{R}(D) \subseteq \mathcal{R}(D^*)$ .

 $(2) \implies (1)$ : Given that  $\mathcal{R}(T_{/(M,N)})$  and  $\mathcal{R}(D)$  are closed, as established by Theorem [4.1,](#page-10-0) it follows that  $\mathcal{R}(T)$  is also closed. By applying Theorem [2.8,](#page-4-0) we express  $T^{\dagger}$  as

$$
T^\dagger = \begin{pmatrix} (T_{/(M,N)})^\dagger & -(T_{/(M,N)})^\dagger B D^\dagger \\ - D^\dagger C (T_{/(M,N)})^\dagger & D^\dagger + D^\dagger C (T_{/(M,N)})^\dagger B D^\dagger \end{pmatrix}.
$$

Utilizing the conditions  $\mathcal{R}(C) \subseteq \mathcal{R}(D)$ ,  $\mathcal{R}(B^*) \subseteq \mathcal{R}(D^*)$ ,  $\mathcal{R}(C^*) \subseteq$  $\mathcal{R}(T_{/(M,N)}^*)$  and  $\mathcal{R}(B) \subseteq \mathcal{R}(T_{/(M,N)}^*)$ , we deduce

$$
TT^{\dagger} = \begin{pmatrix} T_{/(M,N)}(T_{/(M,N)})^{\dagger} & 0 \\ 0 & DD^{\dagger} \end{pmatrix} \text{ and } T^{\dagger}T = \begin{pmatrix} (T_{/(M,N)})^{\dagger}T_{/(M,N)} & 0 \\ 0 & D^{\dagger}D \end{pmatrix}.
$$

Given that  $T_{/(M,N)}$  and D are hypo-EP, it follows that  $\mathcal{R}(TT^{\dagger}) \subseteq \mathcal{R}(T^{\dagger}T)$ . Accordingly,  $\mathcal{R}(T) \subseteq \mathcal{R}(T^*)$ . Thus T is hypo-EP.

(2)  $\iff$  (3): Given that T is  $(M, N)$ -complementable. The first implication indicates that  $\begin{pmatrix} T/(M,N) & 0 \ C & D \end{pmatrix}$  and  $\begin{pmatrix} T/(M,N) & B \ 0 & D \end{pmatrix}$  $0$   $D$  are also  $(M, N)$ -complementable.  $\begin{array}{cc} T_{/(M,N)} & 0 \ C & D \end{array} \biggr) \ \ {\rm and} \ \ \begin{pmatrix} T_{/(M,N)} & B \ 0 & D \end{pmatrix}$  $0$  D  $\Big)$  satisfy all the conditions established for  $T$  in the hypothesis. Additionally,

$$
(T_{/(M,N)})_{/(M,N)} = T_{/(M,N)}.
$$
 Thus, by applying (1)  $\iff$  (2) for operators  $\begin{pmatrix} T_{/(M,N)} & 0 \\ C & D \end{pmatrix}$  and  $\begin{pmatrix} T_{/(M,N)} & B \\ 0 & D \end{pmatrix}$ , we obtain that  $\begin{pmatrix} T_{/(M,N)} & 0 \\ C & D \end{pmatrix}$  and  $\begin{pmatrix} T_{/(M,N)} & B \\ 0 & D \end{pmatrix}$  are hypo-*EP*.

**Corollary 4.4.** *Let*  $T \in \mathcal{B}(\mathcal{H})$  *be a*  $(M, N)$ *-complementable operator.* If  $T$ is EP, then  $T_{/(M,N)}$  is EP. *Moreover, if*  $\mathcal{R}(C^*) \subseteq \mathcal{R}((T_{/(M,N)})^*)$  *then the following are equivalent:* (1) T *is* EP *;* (2)  $T_{/(M,N)}$  and D are EP and  $\mathcal{R}(B) \subseteq \mathcal{R}(T_{/(M,N)})$ ;

$$
(3) \begin{pmatrix} T_{/(M,N)} & 0 \\ C & D \end{pmatrix} and \begin{pmatrix} T_{/(M,N)} & B \\ 0 & D \end{pmatrix} are EP.
$$

*Proof.* By Theorem [4.3,](#page-11-0) it is easy to see that  $T_{/(M,N)}$  is EP.  $(1) \iff (2)$ : It is easy to see that T is EP if and only if both T and T<sup>\*</sup> are hypo-EP. Hence by Theorem [4.3,](#page-11-0) T is EP if and only if  $T_{/(M,N)}$  and D are EP, and hence  $\mathcal{R}(B) \subseteq \mathcal{R}(T_{/(M,N)}).$ 

$$
(2) \iff (3): \quad \text{Both } \begin{pmatrix} T_{/(M,N)} & 0 \\ C & D \end{pmatrix} \text{ and } \begin{pmatrix} T_{/(M,N)} & B \\ 0 & D \end{pmatrix} \text{ are } (M,N)\text{-complementable, and } (T_{/(M,N)})/(M,N) = T_{/(M,N)}.
$$
\n
$$
\text{Applying } (1) \iff (2) \text{ for operators } \begin{pmatrix} T_{/(M,N)} & 0 \\ C & D \end{pmatrix} \text{ and } \begin{pmatrix} T_{/(M,N)} & B \\ 0 & D \end{pmatrix},
$$
\nwe conclude that 
$$
\begin{pmatrix} T_{/(M,N)} & 0 \\ C & D \end{pmatrix} \text{ and } \begin{pmatrix} T_{/(M,N)} & B \\ 0 & D \end{pmatrix} \text{ are } EP.
$$

Now we illustrate Theorem [4.3](#page-11-0) with the following example.

**Example 4.5.** Let  $T: \ell_2 \oplus \ell_2 \to \ell_2 \oplus \ell_2$  be defined by

$$
T((x_1,x_2,\ldots)\oplus(y_1,y_2,\ldots))=\left(\left(x_1,x_1+\frac{x_2}{2}+y_1,x_2+\frac{2x_3}{3}+y_2,\ldots\right)\oplus\left(0,x_1+y_1,x_2+y_2,\ldots\right)\right).
$$

 $0$  D

 $\Box$ 

Consider the subspaces

$$
M = \ell_2 \oplus 0 \text{ and } M^{\perp} = 0 \oplus \ell_2.
$$

We have  $T = \begin{pmatrix} A & B \\ C & D \end{pmatrix}$  where A, B, C and D are given by

$$
A((x_1, x_2,...) \oplus 0) = \left( \left( x_1, x_1 + \frac{x_2}{2}, x_2 + \frac{2x_3}{3}, ... \right) \oplus 0 \right);
$$
  
\n
$$
B(0 \oplus (y_1, y_2,...)) = \left( (0, y_1, y_2,...) \oplus 0 \right);
$$
  
\n
$$
C((x_1, x_2,...) \oplus 0) = \left( 0 \oplus (0, x_1, x_2,...) \right);
$$
  
\n
$$
D(0 \oplus (y_1, y_2,...)) = \left( 0 \oplus (0, y_1, y_2,...) \right).
$$

Clearly, T is M-complementable and

$$
T_{/M}((x_1,x_2,x_3,\ldots)\oplus 0)=\bigg(\bigg(x_1,\frac{x_2}{2},\frac{2x_3}{3}\ldots\bigg)\oplus 0\bigg).
$$

Also,  $T_{/M}^* = T_{/M}$ ,  $\mathcal{R}(D)$  is closed and  $\mathcal{R}(C^*) \subseteq \mathcal{R}(T_{/M}^*)$ . Now let  $x \oplus y \in$  $\mathcal{N}(T)$ , then  $x = y = 0$ . Then  $0 = \mathcal{N}(T) \subseteq \mathcal{N}(T^*)$ . Now  $T_{/M} : \ell_2 \to \ell_2$ is invertible, so  $\mathcal{R}(T_M)$  is closed. Also,  $0 = \mathcal{N}(T_M) \subseteq \mathcal{N}(T_M^{'})$ . Also by Theorem [4.1,](#page-10-0)  $\mathcal{R}(T)$ ,  $\mathcal{R}(T_1)$ , and  $\mathcal{R}(T_2)$  are closed, where  $T_1 = \begin{pmatrix} T_{/M} & 0 \\ C & D \end{pmatrix}$ and  $T_2 = \begin{pmatrix} T_{/M} & B \\ 0 & D \end{pmatrix}$ ). So we have T,  $T_{/M}$ , D,  $T_1$  and  $T_2$  are hypo-EP and

 $0$  D  $\mathcal{R}(B) \subseteq \mathcal{R}(T_{/M}).$ 

The next example shows that the condition  $\mathcal{R}(C^*) \nsubseteq \mathcal{R}((T_M)^*)$  is necessary for the implication (1) if and only if (2) in Theorem [4.3.](#page-11-0)

**Example 4.6.** Let  $T: \ell_2 \to \ell_2$  be defined by

 $T(x_1, x_2, \ldots) = (x_1 + x_2 + x_3 + x_4, 0, x_2 + x_3 + x_4, x_2 + x_3 + x_4, x_5, x_6, \ldots).$ Consider the subspaces

$$
M = \{(x_1, x_2, 0, 0, \ldots)\} \text{ and } M^{\perp} = \{(0, 0, x_3, x_4, \ldots)\}.
$$
  
We have  $T = \begin{pmatrix} A & B \\ C & D \end{pmatrix}$  where A, B, C and D are given by  
 $A(x_1, x_2, 0, 0, \ldots) = (x_1 + x_2, 0, 0, 0 \ldots);$   
 $B(0, 0, x_3, x_4, \ldots) = (x_3 + x_4, 0, 0, 0 \ldots);$   
 $C(x_1, x_2, 0, 0, \ldots) = (0, 0, x_2, x_2, 0, 0, \ldots);$   
 $D(0, 0, x_3, x_4, \ldots) = (0, 0, x_3 + x_4, x_3 + x_4, x_5, x_6 \ldots).$ 

Clearly, T is M-complementable and

 $T_{/M}(x_1, x_2, 0, 0, \ldots) = (x_1, 0, 0, \ldots).$ 

But  $\mathcal{R}(C^*) \nsubseteq \mathcal{R}((T_{/M})^*)$ . Now, D and  $T_{/M}$  are hypo-EP (EP), but T is not hypo- $EP$  ( $EP$ ). This shows that the range space conditions are necessary for Theorem [4.3.](#page-11-0)

The next example shows that the condition  $\mathcal{R}(C^*) \nsubseteq \mathcal{R}((T_{/M})^*)$  is necessary for the implication (2) if and only if (3) in Theorem [4.3.](#page-11-0)

**Example 4.7.** Let  $T: \ell_2 \to \ell_2$  be defined by

$$
T(x_1, x_2,...) = (x_1+x_2+x_3+x_4, x_1+x_3+x_4, x_1+x_2+x_3+x_4, x_1+x_2+x_3+x_4, x_5, x_6,...).
$$
  
with respect to the subspaces

$$
M = \{(x_1, x_2, 0, 0, \ldots)\} \text{ and } M^{\perp} = \{(0, 0, x_3, x_4, \ldots)\}.
$$
  
We have  $T = \begin{pmatrix} A & B \\ C & D \end{pmatrix}$  where A, B, C and D are given by

$$
A(x_1, x_2, 0, 0, \ldots) = (x_1 + x_2, x_1, 0, 0, \ldots);
$$
  
\n
$$
B(0, 0, x_3, x_4, \ldots) = (x_3 + x_4, x_3 + x_4, 0, 0, \ldots);
$$
  
\n
$$
C(x_1, x_2, 0, 0, \ldots) = (0, 0, x_1 + x_2, x_1 + x_2, 0, 0, \ldots);
$$
  
\n
$$
D(0, 0, x_3, x_4, \ldots) = (0, 0, x_4 + x_4, x_3 + x_4, x_5, x_6 \ldots).
$$

Clearly, T is M-complementable and

$$
T_{/M}(x_1, x_2, 0, 0, \ldots) = (0, -x_2, 0, \ldots).
$$

But  $\mathcal{R}(B) \nsubseteq \mathcal{R}(T_M)$  and  $\mathcal{R}(C^*) \nsubseteq \mathcal{R}((T_M)^*)$ . Clearly, D,  $T_M$  and T are hypo-EP (EP), but  $\begin{pmatrix} T/M & 0 \\ C & D \end{pmatrix}$  and  $\begin{pmatrix} T/M & B \\ 0 & D \end{pmatrix}$  $0$  D are not hypo- $EP(EP)$ .

# Acknowledgments

The first author thanks the National Institute of Technology Karnataka (NITK), Surathkal for the financial support. The present work of the second author was partially supported by Science and Engineering Research Board (SERB), Department of Science and Technology, Government of India (Reference Number: MTR/2023/000471) under the scheme "Mathematical Research Impact Centric Support (MATRICS)".

#### **REFERENCES**

- <span id="page-15-3"></span><span id="page-15-2"></span>[1] W. N. Anderson, Jr., Shorted operators. SIAM J. Appl. Math., 20:520–525, 1971.
- <span id="page-15-9"></span>[2] T. Ando., Generalized Schur complements Linear Algebra Appl., 27:173–186, 1979.
- [3] J. Antezana, G. Corach and D. Stojanoff., Bilateral shorted operators and parallel sums. Linear Algebra Appl., 414:570–588, 2006.
- <span id="page-15-10"></span>[4] M. L. Arias, G. Corach and A. Maestripieri., On complementable operators in the sense of T. Ando. Linear Algebra Appl., 594:158–176, 2020.
- <span id="page-15-1"></span>[5] A. Arthur., Conditions for positive and nonnegative definiteness in terms of pseudoinverses. SIAM J. Appl. Math., 17:434–440, 1969.
- <span id="page-15-8"></span>[6] C. A. Butler and T.D. Morley., Six generalized Schur complements Linear Algebra Appl., 106:259–269, 1988.
- <span id="page-15-12"></span>[7] S. L. Campbell and C. D. Meyer., EP operators and generalized inverses. Canad. Math. Bull., 18(3):327–333, 1975.
- <span id="page-15-4"></span>[8] D. Carlson., What are Schur complements, anyway?. Linear Algebra Appl., 74:257– 275, 1986.
- <span id="page-15-5"></span>[9] G. Corach, A. Maestripieri and D. Stojanoff., Generalized Schur complements and oblique projections. Linear Algebra Appl., 341, 259–272, 2002.
- <span id="page-15-6"></span>[10] G. Corach, A. Maestripieri and D. Stojanoff., Oblique projections and Schur complements. Acta Sci. Math. (Szeged), 67, 337–356, 2001.
- <span id="page-15-11"></span>[11] R. W. Cottle., On manifestations of the Schur complement. Rend. Sem. Mat. Fis. Milano , 45, 31–40, 1975.
- <span id="page-15-0"></span>[12] F. R. Gantmacher., The theory of matrices. Vol. 1. AMS Chelsea Publishing, Providence, RI, 1998.
- <span id="page-15-7"></span>[13] J. I. Giribet, A. Maestripieri and F. M. Pería., Shorting selfadjoint operators in Hilbert spaces. Linear Algebra Appl., 428, 1899–1911, 2008.
- <span id="page-16-10"></span>[14] N. C. González, M. F. Martínez-Serrano and J. Robles., Expressions for the Moore-Penrose inverse of block matrices involving the Schur complement. Linear Algebra Appl., 471:353–368, 2015.
- <span id="page-16-11"></span><span id="page-16-1"></span>[15] E. V. Haynsworth., Determination of the inertia of a partitioned Hermitian matrix Linear Algebra Appl., 1:73–81, 1968.
- [16] I. S. Hwang and W. Y. Lee., The reduced minimum modulus of operators. J. Math. Anal. Appl., 267, 679–694, 2002.
- <span id="page-16-12"></span><span id="page-16-4"></span>[17] M. Itoh., On some EP operators. Canad. Math. Bull., 16(1):49–56, 2005.
- [18] M. Krein., The theory of self-adjoint extensions of semi-bounded Hermitian transformations and its applications I. Rec. Math. [Mat. Sbornik] N.S., 20(62):431–495, 1947.
- <span id="page-16-9"></span><span id="page-16-7"></span>[19] S. K. Mitra and M. L. Puri., Shorted matrices—an extended concept and some applications. Linear Algebra Appl., 42:57–79, 1982.
- <span id="page-16-8"></span>[20] Ar. Meenakshi., On Schur complements in an EP matrix. Period. Math. Hungar., 16:193–200, 1985.
- [21] D. V. Ouellette., Schur complements and statistics. Linear Algebra Appl.,36, 187- 295,1981.
- <span id="page-16-5"></span>[22] E. L. Pekarev., Shorts of operators and some extremal problems. Acta Sci. Math. (Szeged) , 56:1-2, 147-163, 1992.
- <span id="page-16-6"></span>[23] V. Ptak., Extremal operators and oblique projections. Casopis pro pestovani Matematiky , 110, 343–350, 1985.
- <span id="page-16-0"></span>[24] J. Schur., Über Potenzreihen, die im Innern des Einheitskreises beschränkt sind. J. Reine Angew. Math., 147:205–232, 1917.
- <span id="page-16-3"></span>[25] J. J. Sylvester., On the relation between the minor determinants of linearly equivalent quadratic functions. The London, Edinburgh, and Dublin Philosophical Magazine and Journal of Science, 1(4), 295–305, 1851.
- <span id="page-16-2"></span>[26] F. Zhang., The Schur Complement and Its Applications. Springer New York, NY, 1571-5698.

(Sachin Manjunath Naik) Department of Mathematical and Computational Sciences, National Institute of Technology Karnataka (NITK), Surathkal, Mangaluru 575 025, India

Email address: sachinmaths46@gmail.com

(P. Sam Johnson) Department of Mathematical and Computational Sciences, National Institute of Technology Karnataka (NITK), Surathkal, Mangaluru 575 025, India

Email address: sam@nitk.edu.in