

On some 3-dimensional almost η -Ricci solitons with diagonal metrics

Adara M. Blaga

Abstract

We study some properties of the 3-dimensional manifold \mathbb{R}^3 with a diagonal Riemannian metric as an almost η -Ricci soliton from the following points of view: under certain assumptions, we determine the potential vector field if η is given; we get obstructions on the metric when the potential vector field has a particular expression; we compute the defining functions of the soliton when both the potential vector field and the 1-form are prescribed. Moreover, we find conditions for the manifold to be flat or Ricci-flat. Based on the theoretical results, we provide examples.

1 Introduction

The solitons' theory has been lately intensively investigated. The stationary solutions of the Ricci flow, namely, the *Ricci solitons*, still represent a very actual topic to be studied from various points of view. Problems like linear stability of compact Ricci solitons, curvature estimates and rigidity results for gradient Ricci solitons, etc. have been recently treated by Huai-Dong Cao *et. al.* in [9, 8, 7]. As a generalization of the notion of Ricci soliton given by Hamilton in [12], an *almost η -Ricci soliton* [4, 5, 10] is a complete Riemannian manifold (M, \tilde{g}) with a smooth vector field V which satisfies the following equation

$$\frac{1}{2}\mathcal{L}_V\tilde{g} + \text{Ric} + \lambda\tilde{g} + \mu\eta \otimes \eta = 0 \quad (1)$$

2010 *Mathematics Subject Classification.* 35Q51, 53B25, 53B50.

Key words and phrases. Almost η -Ricci soliton; Killing vector field; Ricci-flat manifold.

for λ and μ two smooth functions on M ($\mu \neq 0$), where Ric is the Ricci curvature tensor field, $\mathcal{L}_V \tilde{g}$ is the Lie derivative of the metric \tilde{g} in the direction of V and η is a 1-form on M . If V is a Killing vector field, i.e., $\mathcal{L}_V \tilde{g} = 0$, then the soliton is called *trivial*, and in this case, (M, \tilde{g}) is just a quasi-Einstein manifold. On the other hand, if $\eta = 0$, then $(M, \tilde{g}, V, \lambda)$ is called an *almost Ricci soliton* [13]. In this case, the soliton is said to be *shrinking*, *steady* or *expanding* according as λ is negative, zero or positive, respectively [11]. In the above cases, if λ and μ are real numbers, then we drop "almost".

The aim of the present paper is to describe the 3-dimensional manifold \mathbb{R}^3 endowed with a diagonal Riemannian metric as an almost η -Ricci soliton. More precisely, under certain assumptions, we determine the potential vector field V when η is given, we find the conditions that must be satisfied by the Riemannian metric when the potential vector field has a particular expression, and we compute the defining functions λ and μ when both the potential vector field and the 1-form are prescribed. Based on the theoretical results, we construct examples, among which the 3-dimensional Sol_3 and $\mathbb{H}^2 \times \mathbb{R}$ Lie groups. An analogous study for the canonical metric has been previously done by the present author in [6]. More recently, some particular Riemannian manifolds, namely, the 3-dimensional Sol_3 and $\mathbb{H}^2 \times \mathbb{R}$ Lie groups, have been described as Ricci solitons by Belarbi, Atashpeykara and Haji-Badali in [3, 2, 1] from a similar point of view.

2 Flatness and Ricci-flatness conditions

We consider the 3-dimensional manifold \mathbb{R}^3 endowed with a diagonal Riemannian metric \tilde{g} given by:

$$\tilde{g} = \frac{1}{f_1^2} dx^1 \otimes dx^1 + \frac{1}{f_2^2} dx^2 \otimes dx^2 + dx^3 \otimes dx^3,$$

where f_1 and f_2 are two smooth functions nowhere zero on \mathbb{R}^3 , and x^1, x^2, x^3 are the standard coordinates in \mathbb{R}^3 . Let

$$\left\{ E_1 := f_1 \frac{\partial}{\partial x^1}, \quad E_2 := f_2 \frac{\partial}{\partial x^2}, \quad E_3 := \frac{\partial}{\partial x^3} \right\}$$

be a local orthonormal frame field, and, for the sake of simplicity, we will make the following notations:

$$\frac{f_2}{f_1} \cdot \frac{\partial f_1}{\partial x^2} =: a, \quad \frac{1}{f_1} \cdot \frac{\partial f_1}{\partial x^3} =: b, \quad \frac{f_1}{f_2} \cdot \frac{\partial f_2}{\partial x^1} =: c, \quad \frac{1}{f_2} \cdot \frac{\partial f_2}{\partial x^3} =: d.$$

Computing the Lie brackets $[X, Y] := X \circ Y - Y \circ X$, we get:

$$[E_1, E_2] = -aE_1 + cE_2, \quad [E_1, E_3] = -bE_1, \quad [E_2, E_3] = -dE_2.$$

The Levi-Civita connection ∇ of \tilde{g} deduced from the Koszul's formula

$$\begin{aligned} 2\tilde{g}(\nabla_X Y, Z) &= X(\tilde{g}(Y, Z)) + Y(\tilde{g}(Z, X)) - Z(\tilde{g}(X, Y)) \\ &\quad - \tilde{g}(X, [Y, Z]) + \tilde{g}(Y, [Z, X]) + \tilde{g}(Z, [X, Y]) \end{aligned}$$

is given by:

$$\nabla_{E_1} E_1 = aE_2 + bE_3, \quad \nabla_{E_1} E_2 = -aE_1, \quad \nabla_{E_1} E_3 = -bE_1, \quad \nabla_{E_2} E_1 = -cE_2,$$

$$\nabla_{E_2} E_2 = cE_1 + dE_3, \quad \nabla_{E_2} E_3 = -dE_2, \quad \nabla_{E_3} E_1 = 0, \quad \nabla_{E_3} E_2 = 0, \quad \nabla_{E_3} E_3 = 0,$$

and the Riemannian and Ricci curvature tensor fields

$$R(X, Y)Z := \nabla_X \nabla_Y Z - \nabla_Y \nabla_X Z - \nabla_{[X, Y]} Z, \quad \text{Ric}(Y, Z) := \sum_{k=1}^3 \tilde{g}(R(E_k, Y)Z, E_k)$$

are:

$$R(E_1, E_2)E_2 = [E_1(c) + E_2(a) - a^2 - c^2 - bd]E_1 + [E_3(c) - cd]E_3,$$

$$R(E_2, E_1)E_1 = [E_1(c) + E_2(a) - a^2 - c^2 - bd]E_2 + [E_3(a) - ab]E_3,$$

$$R(E_1, E_3)E_3 = [E_3(b) - b^2]E_1,$$

$$R(E_2, E_3)E_3 = [E_3(d) - d^2]E_2,$$

$$R(E_3, E_1)E_1 = [E_3(a) - ab]E_2 + [E_3(b) - b^2]E_3,$$

$$R(E_3, E_2)E_2 = [E_3(c) - cd]E_1 + [E_3(d) - d^2]E_3,$$

$$\begin{aligned} R(E_1, E_2)E_3 &= \tilde{g}(R(E_2, E_1)E_1, E_3)E_1 - \tilde{g}(R(E_1, E_2)E_2, E_3)E_2 \\ &= [E_3(a) - ab]E_1 - [E_3(c) - cd]E_2, \end{aligned}$$

$$\begin{aligned} R(E_2, E_3)E_1 &= \tilde{g}(R(E_3, E_2)E_2, E_1)E_2 - \tilde{g}(R(E_2, E_3)E_3, E_1)E_3 \\ &= [E_3(c) - cd]E_2, \end{aligned}$$

$$\begin{aligned} R(E_3, E_1)E_2 &= -\tilde{g}(R(E_3, E_1)E_1, E_2)E_1 + \tilde{g}(R(E_1, E_3)E_3, E_2)E_3 \\ &= -[E_3(a) - ab]E_1, \end{aligned}$$

$$\text{Ric}(E_1, E_1) = E_1(c) + E_2(a) + E_3(b) - a^2 - b^2 - c^2 - bd,$$

$$\text{Ric}(E_2, E_2) = E_1(c) + E_2(a) + E_3(d) - a^2 - c^2 - d^2 - bd,$$

$$\text{Ric}(E_3, E_3) = E_3(b) + E_3(d) - b^2 - d^2,$$

$$\text{Ric}(E_1, E_2) = 0, \quad \text{Ric}(E_1, E_3) = E_3(c) - cd, \quad \text{Ric}(E_2, E_3) = E_3(a) - ab.$$

We shall determine the conditions which the two functions f_1 and f_2 must satisfy for the manifold to be flat or Ricci-flat, exploring all the possible cases when each of the two functions depend on a single variable. In all the rest of this paper, if a function h on \mathbb{R}^3 depends only on some of its variables, then we will write in its argument only that variables in order to emphasize this fact. For example, $h(x^j)$, $h(x^i, x^j)$.

PROPOSITION 2.1. *If $f_i = f_i(x^i)$ for $i \in \{1, 2\}$, then $(\mathbb{R}^3, \tilde{g})$ is a flat Riemannian manifold.*

PROOF. In this case, $a = b = c = d = 0$ and $R(E_i, E_j)E_k = 0$ for any $i, j, k \in \{1, 2, 3\}$. □

PROPOSITION 2.2. *If $f_i : \mathbb{R}^2 \times I \rightarrow \mathbb{R}$, $f_i = f_i(x^3)$ for $i \in \{1, 2\}$ and $I \subset \mathbb{R}$ an open interval, then the following assertions are equivalent:*

- (1) $(\mathbb{R}^2 \times I, \tilde{g}|_{\mathbb{R}^2 \times I})$ is a flat Riemannian manifold;
- (2) $(\mathbb{R}^2 \times I, \tilde{g}|_{\mathbb{R}^2 \times I})$ is a Ricci-flat Riemannian manifold;
- (3) f_1 and f_2 are constant, or $f_i = k_i \in \mathbb{R} \setminus \{0\}$ and $f_j(x^3) = \frac{c_1}{x^3 - c_2}$ for $c_1 \in \mathbb{R} \setminus \{0\}$, $c_2 \in \mathbb{R} \setminus I$ and $i \neq j$.

PROOF. In this case, $a = 0$, $b = \frac{f'_1}{f_1}$, $c = 0$, $d = \frac{f'_2}{f_2}$, and

$$R(E_1, E_2)E_2 = -bdE_1, \quad R(E_2, E_1)E_1 = -bdE_2,$$

$$R(E_1, E_3)E_3 = (b' - b^2)E_1, \quad R(E_3, E_1)E_1 = (b' - b^2)E_3,$$

$$R(E_2, E_3)E_3 = (d' - d^2)E_2, \quad R(E_3, E_2)E_2 = (d' - d^2)E_3,$$

$$\text{Ric}(E_1, E_1) = b' - b^2 - bd, \quad \text{Ric}(E_2, E_2) = d' - d^2 - bd, \quad \text{Ric}(E_3, E_3) = b' + d' - b^2 - d^2,$$

$$\text{Ric}(E_1, E_2) = \text{Ric}(E_1, E_3) = \text{Ric}(E_2, E_3) = 0.$$

Then, $R = 0$ if and only if

$$\begin{cases} bd = 0 \\ b' = b^2 ; \\ d' = d^2 \end{cases}$$

$\text{Ric} = 0$ if and only if

$$\begin{cases} b' - b^2 = bd = d' - d^2 \\ b' - b^2 = -(d' - d^2) \end{cases} ;$$

that is, $R = 0$ if and only if $\text{Ric} = 0$, i.e.,

$$\begin{cases} f_1' f_2' = 0 \\ \left(\frac{f_1'}{f_1}\right)' = \left(\frac{f_1'}{f_1}\right)^2 \\ \left(\frac{f_2'}{f_2}\right)' = \left(\frac{f_2'}{f_2}\right)^2 \end{cases}.$$

Therefore,

- $f_1' = 0$ and $f_2' = 0$, or
- $f_1' = 0$ and $-\frac{1}{\frac{f_2'(x^3)}{f_2(x^3)}} = x^3 + c_0$ (i.e., $f_2(x^3) = \frac{c_1}{x^3 - c_2}$ with $c_1 \in \mathbb{R} \setminus \{0\}$, $c_2 \in \mathbb{R} \setminus I$),

or, similarly

- $f_1(x^3) = \frac{c_1}{x^3 - c_2}$ with $c_1 \in \mathbb{R} \setminus \{0\}$, $c_2 \in \mathbb{R} \setminus I$ and $f_2' = 0$. □

COROLLARY 2.3. *If $f_i = f_i(x^3)$ for $i \in \{1, 2\}$, then the following assertions are equivalent:*

- (1) $(\mathbb{R}^3, \tilde{g})$ is a flat Riemannian manifold;
- (2) $(\mathbb{R}^3, \tilde{g})$ is a Ricci-flat Riemannian manifold;
- (3) f is constant.

PROOF. It follows immediately from Proposition 2.2. □

PROPOSITION 2.4. *If $f_1, f_2 : \mathbb{R}^2 \times I \rightarrow \mathbb{R}$, $f_1 = f_1(x^1)$, $f_2 = f_2(x^3)$ for $I \subset \mathbb{R}$ an open interval, then the following assertions are equivalent:*

- (1) $(\mathbb{R}^2 \times I, \tilde{g}|_{\mathbb{R}^2 \times I})$ is a flat Riemannian manifold;
- (2) $(\mathbb{R}^2 \times I, \tilde{g}|_{\mathbb{R}^2 \times I})$ is a Ricci-flat Riemannian manifold;
- (3) f_2 is constant, or $f_2(x^3) = \frac{c_1}{x^3 - c_2}$ for $c_1 \in \mathbb{R} \setminus \{0\}$, $c_2 \in \mathbb{R} \setminus I$.

PROOF. In this case, $a = b = c = 0$, $d = \frac{f_2'}{f_2}$, and

$$R(E_2, E_3)E_3 = (d' - d^2)E_2, \quad R(E_3, E_2)E_2 = (d' - d^2)E_3,$$

$$R(E_1, E_2)E_2 = R(E_1, E_3)E_3 = R(E_2, E_1)E_1 = R(E_3, E_1)E_1 = 0,$$

$$\text{Ric}(E_2, E_2) = \text{Ric}(E_3, E_3) = d' - d^2,$$

$$\text{Ric}(E_1, E_1) = \text{Ric}(E_1, E_2) = \text{Ric}(E_1, E_3) = \text{Ric}(E_2, E_3) = 0.$$

Then, $R = 0$ if and only if $\text{Ric} = 0$, i.e.,

$$d' = d^2,$$

that is,

$$\left(\frac{f_2'}{f_2}\right)' = \left(\frac{f_2'}{f_2}\right)^2.$$

Therefore,

- $f_2' = 0$, or
- $-\frac{1}{\frac{f_2'(x^3)}{f_2(x^3)}} = x^3 + c_0$ (i.e., $f_2(x^3) = \frac{c_1}{x^3 - c_2}$ with $c_1 \in \mathbb{R} \setminus \{0\}$, $c_2 \in \mathbb{R} \setminus I$). □

COROLLARY 2.5. *If $f_1 = f_1(x^1)$, $f_2 = f_2(x^3)$, then the following assertions are equivalent:*

- (1) $(\mathbb{R}^3, \tilde{g})$ is a flat Riemannian manifold;
- (2) $(\mathbb{R}^3, \tilde{g})$ is a Ricci-flat Riemannian manifold;
- (3) f_2 is constant.

PROOF. It follows immediately from Proposition 2.4. □

PROPOSITION 2.6. *If $f_i = f_i(x^2)$ for $i \in \{1, 2\}$, then the following assertions are equivalent:*

- (1) $(\mathbb{R}^3, \tilde{g})$ is a flat Riemannian manifold;
- (2) $(\mathbb{R}^3, \tilde{g})$ is a Ricci-flat Riemannian manifold;
- (3) f_1 and f_2 satisfy the equation

$$\frac{f_1' f_2'}{f_1 f_2} + \frac{f_1''}{f_1} = 2 \left(\frac{f_1'}{f_1}\right)^2;$$

- (4) f_1 is constant, or f_1' is nowhere zero and $f_2 = c_0 \frac{f_1^2}{f_1'}$ with $c_0 \in \mathbb{R} \setminus \{0\}$.

PROOF. In this case, $a = f_2 \frac{f_1'}{f_1}$, $b = c = d = 0$, and

$$\begin{aligned} R(E_1, E_2)E_2 &= [E_2(a) - a^2]E_1, & R(E_2, E_1)E_1 &= [E_2(a) - a^2]E_2, \\ R(E_1, E_3)E_3 &= R(E_3, E_1)E_1 = R(E_2, E_3)E_3 = R(E_3, E_2)E_2 = 0, \\ \text{Ric}(E_1, E_1) &= \text{Ric}(E_2, E_2) = E_2(a) - a^2, \\ \text{Ric}(E_1, E_2) &= \text{Ric}(E_1, E_3) = \text{Ric}(E_2, E_3) = \text{Ric}(E_3, E_3) = 0. \end{aligned}$$

Then, $R = 0$ if and only if $\text{Ric} = 0$, i.e.,

$$f_2 a' = a^2,$$

that is,

$$f_2 f_2' \frac{f_1'}{f_1} + f_2^2 \frac{f_1'' f_1 - (f_1')^2}{f_1^2} = f_2^2 \left(\frac{f_1'}{f_1} \right)^2,$$

which is equivalent to

$$\frac{f_1' f_2'}{f_1 f_2} + \frac{f_1''}{f_1} = 2 \left(\frac{f_1'}{f_1} \right)^2.$$

If $f_1' = 0$, then f_1 is a constant function. If $f_1' \neq 0$ (hence, if it is nowhere zero), then the previous relation can be written as

$$\frac{f_2'}{f_2} = \frac{2 \left(\frac{f_1'}{f_1} \right)^2 - \frac{f_1''}{f_1}}{\frac{f_1'}{f_1}}.$$

Let us notice that

$$\frac{f_2'}{f_2} = \frac{\left(\frac{f_1'}{f_1} \right)^2 - \left(\frac{f_1'}{f_1} \right)'}{\frac{f_1'}{f_1}} = \frac{f_1'}{f_1} - \frac{\left(\frac{f_1'}{f_1} \right)'}{\frac{f_1'}{f_1}},$$

which, by integration, gives

$$\ln |f_2| = \ln |f_1| - \ln \left| \frac{f_1'}{f_1} \right| + k = \ln \left(e^k \frac{f_1^2}{|f_1'|} \right),$$

where $k \in \mathbb{R}$; therefore, $f_2 = c_0 \frac{f_1^2}{f_1'}$ with $c_0 \in \mathbb{R} \setminus \{0\}$. □

COROLLARY 2.7. *If $f_1 = f_2 =: f(x^2)$, then the following assertions are equivalent:*

- (1) $(\mathbb{R}^3, \tilde{g})$ is a flat Riemannian manifold;
- (2) $(\mathbb{R}^3, \tilde{g})$ is a Ricci-flat Riemannian manifold;
- (3) $f(x^2) = c_1 e^{c_2 x^2}$ for $c_1 \in \mathbb{R} \setminus \{0\}$, $c_2 \in \mathbb{R}$.

PROOF. (1) and (2) are equivalent to $f f'' - (f')^2 = 0$, that is, $\left(\frac{f'}{f} \right)' = 0$, with the solution $f(x^2) = c_1 e^{c_2 x^2}$ for $c_1 \in \mathbb{R} \setminus \{0\}$, $c_2 \in \mathbb{R}$, hence we get the conclusion. □

PROPOSITION 2.8. *If $f_1 = f_1(x^2)$ and $f_2 = f_2(x^1)$, then the following assertions are equivalent:*

- (1) $(\mathbb{R}^3, \tilde{g})$ is a flat Riemannian manifold;
- (2) $(\mathbb{R}^3, \tilde{g})$ is a Ricci-flat Riemannian manifold;
- (3) f_1 and f_2 satisfy the equation

$$\frac{f_1'' f_1 - 2(f_1')^2}{f_1^4} = -\frac{f_2'' f_2 - 2(f_2')^2}{f_2^4} = \text{constant}.$$

PROOF. In this case, $a = f_2 \frac{f_1'}{f_1}$, $b = 0$, $c = f_1 \frac{f_2'}{f_2}$, $d = 0$, and

$$R(E_1, E_2)E_2 = [E_1(c) + E_2(a) - a^2 - c^2]E_1, \quad R(E_2, E_1)E_1 = [E_1(c) + E_2(a) - a^2 - c^2]E_2,$$

$$R(E_1, E_3)E_3 = R(E_3, E_1)E_1 = R(E_2, E_3)E_3 = R(E_3, E_2)E_2 = 0,$$

$$\text{Ric}(E_1, E_1) = \text{Ric}(E_2, E_2) = E_1(c) + E_2(a) - a^2 - c^2,$$

$$\text{Ric}(E_1, E_2) = \text{Ric}(E_1, E_3) = \text{Ric}(E_2, E_3) = \text{Ric}(E_3, E_3) = 0.$$

Then, $R = 0$ if and only if $\text{Ric} = 0$, i.e.,

$$f_1 \frac{\partial c}{\partial x^1} + f_2 \frac{\partial a}{\partial x^2} = a^2 + c^2,$$

that is,

$$f_1^2 \frac{f_2'' f_2 - (f_2')^2}{f_2^2} + f_2^2 \frac{f_1'' f_1 - (f_1')^2}{f_1^2} = f_2^2 \left(\frac{f_1'}{f_1} \right)^2 + f_1^2 \left(\frac{f_2'}{f_2} \right)^2,$$

which is equivalent to

$$\frac{f_1'' f_1 - 2(f_1')^2}{f_1^4} = -\frac{f_2'' f_2 - 2(f_2')^2}{f_2^4}.$$

Since f_1 depends only on x^2 and f_2 depends only on x^1 , we deduce that the above ratio must be a constant. \square

REMARK 2.9. We shall now look on the condition (3) from Proposition 2.8 satisfied by the two functions, namely:

$$\frac{f'' f - 2(f')^2}{f^4} = k \in \mathbb{R},$$

when f is a real function defined on a real interval. Let us notice that

$$\frac{f'' f - 2(f')^2}{f^4} = -\left(\frac{1}{f} \right)'' \frac{1}{f}.$$

Denoting by $h := \frac{1}{f}$, we have

$$h''h = -k.$$

Let $r \in \mathbb{R}$ and let $J \subset \mathbb{R}$ be an open interval such that $0 \notin J$ and $-2k \ln |y| + r > 0$ for any $y \in J$. Let F be an antiderivative on J of the function

$$y \mapsto \frac{1}{\sqrt{-2k \ln |y| + r}}.$$

Then, $F'(y) > 0$ for any $y \in J$; therefore, F is strictly increasing on J , hence, it is invertible onto its image. Let $\varepsilon \in \{\pm 1\}$ and let $I := \varepsilon(F(J) - c_0)$, where $c_0 \in \mathbb{R}$. Then,

$$h : I \rightarrow \mathbb{R}, \quad h(x) := F^{-1}(\varepsilon x + c_0)$$

satisfies $h''(x)h(x) = -k$ for any $x \in I$, and we get

$$f : I \rightarrow \mathbb{R}, \quad f(x) = \frac{1}{F^{-1}(\varepsilon x + c_0)}.$$

PROPOSITION 2.10. *If $f_1, f_2 : \mathbb{R} \times I_2 \times I_3 \rightarrow \mathbb{R}$, $f_1 = f_1(x^2)$, $f_2 = f_2(x^3)$ for $I_2, I_3 \subset \mathbb{R}$ two open intervals, then the following assertions are equivalent:*

- (1) $(\mathbb{R} \times I_2 \times I_3, \tilde{g}|_{\mathbb{R} \times I_2 \times I_3})$ is a flat Riemannian manifold;
- (2) $(\mathbb{R} \times I_2 \times I_3, \tilde{g}|_{\mathbb{R} \times I_2 \times I_3})$ is a Ricci-flat Riemannian manifold;
- (3) f_1 and f_2 are constant, or $f_1 = k_1 \in \mathbb{R} \setminus \{0\}$ and $f_2(x^3) = \frac{c_1}{x^3 - c_2}$ for $c_1 \in \mathbb{R} \setminus \{0\}$, $c_2 \in \mathbb{R} \setminus I_3$, or $f_2 = k_2 \in \mathbb{R} \setminus \{0\}$ and $f_1(x^2) = \frac{c_1}{x^2 - c_2}$ for $c_1 \in \mathbb{R} \setminus \{0\}$, $c_2 \in \mathbb{R} \setminus I_2$.

PROOF. In this case, $a = f_2 \frac{f_1'}{f_1}$, $b = c = 0$, $d = \frac{f_2'}{f_2}$, and

$$R(E_1, E_2)E_2 = [E_2(a) - a^2]E_1,$$

$$R(E_2, E_3)E_3 = [E_3(d) - d^2]E_2, \quad R(E_3, E_2)E_2 = [E_3(d) - d^2]E_3,$$

$$R(E_2, E_1)E_1 = [E_2(a) - a^2]E_2 + E_3(a)E_3,$$

$$R(E_1, E_3)E_3 = R(E_2, E_3)E_1 = 0,$$

$$R(E_3, E_1)E_1 = E_3(a)E_2, \quad R(E_1, E_2)E_3 = E_3(a)E_1, \quad R(E_3, E_1)E_2 = -E_3(a)E_1,$$

$$\text{Ric}(E_1, E_1) = E_2(a) - a^2, \quad \text{Ric}(E_2, E_2) = E_2(a) - a^2 + E_3(d) - d^2, \quad \text{Ric}(E_3, E_3) = E_3(d) - d^2,$$

$$\text{Ric}(E_1, E_2) = \text{Ric}(E_1, E_3) = 0, \quad \text{Ric}(E_2, E_3) = E_3(a).$$

Then, $R = 0$ if and only if $\text{Ric} = 0$, i.e.,

$$\begin{cases} f_1' f_2' = 0 \\ \left(\frac{f_1'}{f_1}\right)' = \left(\frac{f_1'}{f_1}\right)^2 \\ \left(\frac{f_2'}{f_2}\right)' = \left(\frac{f_2'}{f_2}\right)^2 \end{cases}.$$

Therefore,

- $f_1' = 0$ and $f_2' = 0$, or
- $f_1' = 0$ and $-\frac{1}{\frac{f_2'(x^3)}{f_2(x^3)}} = x^3 + c_0$ (i.e., $f_2(x^3) = \frac{c_1}{x^3 - c_2}$ with $c_1 \in \mathbb{R} \setminus \{0\}$, $c_2 \in \mathbb{R} \setminus I_3$),

or, similarly

- $f_1(x^2) = \frac{c_1}{x^2 - c_2}$ with $c_1 \in \mathbb{R} \setminus \{0\}$, $c_2 \in \mathbb{R} \setminus I_2$ and $f_2' = 0$. □

COROLLARY 2.11. *If $f_1 = f_1(x^2)$, $f_2 = f_2(x^3)$, then the following assertions are equivalent:*

- (1) $(\mathbb{R}^3, \tilde{g})$ is a flat Riemannian manifold;
- (2) $(\mathbb{R}^3, \tilde{g})$ is a Ricci-flat Riemannian manifold;
- (3) f_1 and f_2 are constant.

PROOF. It follows immediately from Proposition 2.10. □

3 \mathbb{R}^3 as an almost η -Ricci soliton

Let $V = \sum_{k=1}^3 V^k E_k$ and $\eta = \sum_{k=1}^3 \eta^k e_k$, where e_k is the dual 1-form of E_k for $k \in \{1, 2, 3\}$. From (1) we get the equations that define an almost η -Ricci soliton $(\mathbb{R}^3, \tilde{g}, V, \lambda, \mu)$:

$$\begin{aligned} \frac{1}{2} \left\{ E_i(V^j) + E_j(V^i) + \sum_{k=1}^3 V^k [\tilde{g}(\nabla_{E_i} E_k, E_j) + \tilde{g}(E_i, \nabla_{E_j} E_k)] \right\} \\ + \text{Ric}(E_i, E_j) + \lambda \delta^{ij} + \mu \eta^i \eta^j = 0 \end{aligned}$$

for any $(i, j) \in \{(1, 1), (2, 2), (3, 3), (1, 2), (1, 3), (2, 3)\}$, equivalent to:

$$\begin{cases} E_1(V^1) - aV^2 - bV^3 + \text{Ric}(E_1, E_1) + \lambda + \mu(\eta^1)^2 = 0 \\ E_2(V^2) - cV^1 - dV^3 + \text{Ric}(E_2, E_2) + \lambda + \mu(\eta^2)^2 = 0 \\ E_3(V^3) + \text{Ric}(E_3, E_3) + \lambda + \mu(\eta^3)^2 = 0 \\ \frac{1}{2}[E_1(V^2) + E_2(V^1) + aV^1 + cV^2] + \mu\eta^1\eta^2 = 0 \\ \frac{1}{2}[E_1(V^3) + E_3(V^1) + bV^1] + \text{Ric}(E_1, E_3) + \mu\eta^1\eta^3 = 0 \\ \frac{1}{2}[E_2(V^3) + E_3(V^2) + dV^2] + \text{Ric}(E_2, E_3) + \mu\eta^2\eta^3 = 0 \end{cases} \quad (2)$$

We shall further consider the cases when the potential vector field of the almost η -Ricci soliton $(\mathbb{R}^3, \tilde{g}, \lambda, \mu)$ is $V = \frac{\partial}{\partial x^3}$ or when $\eta = dx^3$.

3.1 Almost η -Ricci solitons with $V = \frac{\partial}{\partial x^3}$

If $V^1 = V^2 = 0$ and $V^3 = 1$, then the system (2) becomes

$$\begin{cases} -b + \text{Ric}(E_1, E_1) + \lambda + \mu(\eta^1)^2 = 0 \\ -d + \text{Ric}(E_2, E_2) + \lambda + \mu(\eta^2)^2 = 0 \\ \text{Ric}(E_3, E_3) + \lambda + \mu(\eta^3)^2 = 0 \\ \mu\eta^1\eta^2 = 0 \\ \text{Ric}(E_1, E_3) + \mu\eta^1\eta^3 = 0 \\ \text{Ric}(E_2, E_3) + \mu\eta^2\eta^3 = 0 \end{cases} \quad (3)$$

PROPOSITION 3.1. *Let $(\mathbb{R}^3, \tilde{g}, V, \lambda, \mu)$ be an almost η -Ricci soliton with $V = \frac{\partial}{\partial x^3}$. If $f_i = f_i(x^i)$ for $i \in \{1, 2\}$ and μ is nowhere zero, then $\eta = 0$ and V is a Killing vector field.*

PROOF. In this case, we have $a = b = c = d = 0$ and $\text{Ric}(E_i, E_j) = 0$ for any $i, j \in \{1, 2, 3\}$ from Proposition 2.1, and the system (3) becomes

$$\begin{cases} \lambda + \mu(\eta^1)^2 = 0 \\ \lambda + \mu(\eta^2)^2 = 0 \\ \lambda + \mu(\eta^3)^2 = 0 \\ \mu\eta^1\eta^2 = \mu\eta^1\eta^3 = \mu\eta^2\eta^3 = 0 \end{cases} \quad (4)$$

Since μ is nowhere zero, we get $(\eta^1)^2 = (\eta^2)^2 = (\eta^3)^2$, hence, $\eta^i = 0$ for $i \in \{1, 2, 3\}$, and $\lambda = 0$ from (4); therefore, $\mathcal{L}_V \tilde{g} = 0$ by means of (1). \square

PROPOSITION 3.2. Let $(\mathbb{R}^3, \tilde{g}, V, \lambda, \mu)$ be an almost η -Ricci soliton with $V = \frac{\partial}{\partial x^3}$. If $f_i = f_i(x^3)$ for $i \in \{1, 2\}$, $\eta^i =: f$ for $i \in \{1, 2, 3\}$, and μ is nowhere zero, then $\eta = 0$. Moreover,

- (i) if one of the functions f_1 and f_2 is constant, then the other one is constant, too; in this case, $(\mathbb{R}^3, \tilde{g})$ is a flat Riemannian manifold, $\lambda = 0$ and V is a Killing vector field;
 (ii) if $\frac{f_1}{f_2}$ is constant, then $f_i(x^3) = c_i e^{c_0 e^{-x^3}}$, $i \in \{1, 2\}$ for $c_0 \in \mathbb{R}$, $c_1, c_2 \in \mathbb{R} \setminus \{0\}$.

PROOF. In this case, we have $a = 0$, $b = \frac{f'_1}{f_1}$, $c = 0$, $d = \frac{f'_2}{f_2}$, and

$$\text{Ric}(E_1, E_1) = b' - b^2 - bd, \quad \text{Ric}(E_2, E_2) = d' - d^2 - bd, \quad \text{Ric}(E_3, E_3) = b' + d' - b^2 - d^2,$$

$$\text{Ric}(E_1, E_2) = \text{Ric}(E_1, E_3) = \text{Ric}(E_2, E_3) = 0$$

from Proposition 2.2, and the system (3) becomes

$$\begin{cases} -b + b' - b^2 - bd + \lambda + \mu(\eta^1)^2 = 0 \\ -d + d' - d^2 - bd + \lambda + \mu(\eta^2)^2 = 0 \\ b' + d' - b^2 - d^2 + \lambda + \mu(\eta^3)^2 = 0 \\ \mu\eta^1\eta^2 = \mu\eta^1\eta^3 = \mu\eta^2\eta^3 = 0 \end{cases} \quad (5)$$

Since μ is nowhere zero, (5) is equivalent to

$$\begin{cases} b(d+1) = d^2 - d' \\ d(b+1) = b^2 - b' \\ \lambda = b^2 - b' + d^2 - d' \\ f^2 = 0 \end{cases} \quad (6)$$

which implies $\eta = 0$.

(i) f_1 is constant if and only if $b = 0$, and from (6) we deduce that $b = 0$ if and only if $d = 0$, i.e., if and only if f_2 is constant. In this case, $R = 0$, $\lambda = 0$ and $\mathcal{L}_V \tilde{g} = 0$.

(ii) If $\frac{f_1}{f_2}$ is constant, then $b = d$ and (6) becomes

$$\begin{cases} b' = -b \\ \lambda = 2(b^2 - b') \end{cases} \quad (7)$$

If $b = d = 0$, then f_1 and f_2 are constant and $\lambda = 0$. If $b = d \neq 0$ (hence, if they are nowhere zero), then

$$\begin{cases} \frac{b'}{b} = -1 \\ \lambda = 2(b^2 - b') \end{cases}, \quad (8)$$

from where we get

$$\frac{f'_i(x^3)}{f_i(x^3)} = b(x^3) = c_0 e^{-x^3}, \quad c_0 \in \mathbb{R} \setminus \{0\}, i \in \{1, 2\},$$

which, by integration, gives $f_i(x^3) = c_i e^{c_0 e^{-x^3}}$ for $c_i \in \mathbb{R} \setminus \{0\}$, $i \in \{1, 2\}$. \square

COROLLARY 3.3. *Under the hypotheses of Proposition 3.2, if $f_1 = f_2 =: f(x^3)$, then $f(x^3) = c_1 e^{c_2 e^{-x^3}}$ for $c_1 \in \mathbb{R} \setminus \{0\}$, $c_2 \in \mathbb{R}$.*

PROOF. It follows from Proposition 3.2 (ii). \square

PROPOSITION 3.4. *Let $(\mathbb{R}^3, \tilde{g}, V, \lambda, \mu)$ be an almost η -Ricci soliton with $V = \frac{\partial}{\partial x^3}$. If $f_1 = f_1(x^1)$, $f_2 = f_2(x^3)$, $\eta^i =: f$ for $i \in \{1, 2, 3\}$, and μ is nowhere zero, then $\eta = 0$. Moreover, $\lambda = 0$, f_2 is constant and V is a Killing vector field.*

PROOF. In this case, we have $a = b = c = 0$, $d = \frac{f'_2}{f_2}$, and

$$\text{Ric}(E_2, E_2) = \text{Ric}(E_3, E_3) = d' - d^2,$$

$$\text{Ric}(E_1, E_1) = \text{Ric}(E_1, E_2) = \text{Ric}(E_1, E_3) = \text{Ric}(E_2, E_3) = 0$$

from Proposition 2.4, and the system (3) becomes

$$\begin{cases} \lambda + \mu(\eta^1)^2 = 0 \\ -d + d' - d^2 + \lambda + \mu(\eta^2)^2 = 0 \\ d' - d^2 + \lambda + \mu(\eta^3)^2 = 0 \\ \mu\eta^1\eta^2 = \mu\eta^1\eta^3 = \mu\eta^2\eta^3 = 0 \end{cases}. \quad (9)$$

Since μ is nowhere zero, (9) is equivalent to

$$\begin{cases} d = d' - d^2 = 0 \\ \lambda = -\mu f^2 \\ f^2 = 0 \end{cases}.$$

It follows that $\eta = 0$, $\lambda = 0$ and $\mathcal{L}_V \tilde{g} = 0$. \square

PROPOSITION 3.5. *Let $(\mathbb{R}^3, \tilde{g}, V, \lambda, \mu)$ be an almost η -Ricci soliton with $V = \frac{\partial}{\partial x^3}$. If $f_i = f_i(x^2)$ for $i \in \{1, 2\}$, $\eta^i =: f$ for $i \in \{1, 2, 3\}$, and μ is nowhere zero, then $\eta = 0$. Moreover, $\lambda = 0$ and*

$$\frac{f'_1 f'_2}{f_1 f_2} + \frac{f''_1}{f_1} = 2 \left(\frac{f'_1}{f_1} \right)^2.$$

PROOF. In this case, we have $a = f_2 \frac{f'_1}{f_1}$, $b = c = d = 0$, and

$$\text{Ric}(E_1, E_1) = \text{Ric}(E_2, E_2) = E_2(a) - a^2,$$

$$\text{Ric}(E_1, E_2) = \text{Ric}(E_1, E_3) = \text{Ric}(E_2, E_3) = \text{Ric}(E_3, E_3) = 0$$

from Proposition 2.10, and the system (3) becomes

$$\begin{cases} f_2 f'_2 \frac{f'_1}{f_1} + \frac{f_2^2}{f_1^2} [f''_1 f_1 - 2(f'_1)^2] + \lambda + \mu(\eta^1)^2 = 0 \\ f_2 f'_2 \frac{f'_1}{f_1} + \frac{f_2^2}{f_1^2} [f''_1 f_1 - 2(f'_1)^2] + \lambda + \mu(\eta^2)^2 = 0 \\ \lambda + \mu(\eta^3)^2 = 0 \\ \mu\eta^1\eta^2 = \mu\eta^1\eta^3 = \mu\eta^2\eta^3 = 0 \end{cases} \quad (10)$$

Since μ is nowhere zero, (10) is equivalent to

$$\begin{cases} f_2 f'_2 \frac{f'_1}{f_1} + \frac{f_2^2}{f_1^2} [f''_1 f_1 - 2(f'_1)^2] + \lambda + \mu f^2 = 0 \\ \lambda = -\mu f^2 \\ f^2 = 0 \end{cases}.$$

It follows that $\eta = 0$, $\lambda = 0$, and, from the first equation of the previous system, we get the relation between f_1 and f_2 . \square

COROLLARY 3.6. *Under the hypotheses of Proposition 3.5, if $f_1 = f_2 =: f(x^2)$, then $f(x^2) = c_1 e^{c_2 x^2}$ for $c_1 \in \mathbb{R} \setminus \{0\}$, $c_2 \in \mathbb{R}$.*

PROOF. It follows from Proposition 3.5 that $f f'' - (f')^2 = 0$, that is, $\left(\frac{f'}{f}\right)' = 0$, with the solution $f(x^2) = c_1 e^{c_2 x^2}$ for $c_1 \in \mathbb{R} \setminus \{0\}$, $c_2 \in \mathbb{R}$. \square

PROPOSITION 3.7. *Let $(\mathbb{R}^3, \tilde{g}, V, \lambda, \mu)$ be an almost η -Ricci soliton with $V = \frac{\partial}{\partial x^3}$. If $f_1 = f_1(x^2)$, $f_2 = f_2(x^1)$, $\eta^i =: f$ for $i \in \{1, 2, 3\}$, and μ is nowhere zero, then $\eta = 0$. Moreover, $\lambda = 0$ and*

$$\frac{f''_1 f_1 - 2(f'_1)^2}{f_1^4} = -\frac{f''_2 f_2 - 2(f'_2)^2}{f_2^4} = \text{constant}.$$

PROOF. In this case, we have $a = f_2 \frac{f'_1}{f_1}$, $b = 0$, $c = f_1 \frac{f'_2}{f_2}$, $d = 0$, and

$$\text{Ric}(E_1, E_1) = \text{Ric}(E_2, E_2) = E_1(c) + E_2(a) - a^2 - c^2,$$

$$\text{Ric}(E_1, E_2) = \text{Ric}(E_1, E_3) = \text{Ric}(E_2, E_3) = \text{Ric}(E_3, E_3) = 0$$

from Proposition 2.8, and the system (3) becomes

$$\begin{cases} f_1 \frac{\partial c}{\partial x^1} + f_2 \frac{\partial a}{\partial x^2} - a^2 - c^2 + \lambda + \mu(\eta^1)^2 = 0 \\ f_1 \frac{\partial c}{\partial x^1} + f_2 \frac{\partial a}{\partial x^2} - a^2 - c^2 + \lambda + \mu(\eta^2)^2 = 0 \\ \lambda + \mu(\eta^3)^2 = 0 \\ \mu\eta^1\eta^2 = \mu\eta^1\eta^3 = \mu\eta^2\eta^3 = 0 \end{cases}. \quad (11)$$

Since μ is nowhere zero and $\eta^i = f$, (11) is equivalent to

$$\begin{cases} f_1 \frac{\partial c}{\partial x^1} + f_2 \frac{\partial a}{\partial x^2} - a^2 - c^2 + \lambda + \mu f^2 = 0 \\ \lambda = -\mu f^2 \\ f^2 = 0 \end{cases}.$$

It follows that $\eta = 0$, $\lambda = 0$, and, from the first equation of the previous system, we get the relation between f_1 and f_2 . Since f_1 depends only on x^2 and f_2 depends only on x^1 , we deduce that the obtained ratio must be a constant. \square

PROPOSITION 3.8. *Let $(\mathbb{R}^3, \tilde{g}, V, \lambda, \mu)$ be an almost η -Ricci soliton with $V = \frac{\partial}{\partial x^3}$. If $f_1 = f_1(x^2)$, $f_2 = f_2(x^3)$, $\eta^i =: f$ for $i \in \{1, 2, 3\}$, and μ is nowhere zero, then $\eta = 0$. Moreover, $\lambda = 0$, V is a Killing vector field, and one of the functions f_1 and f_2 is constant.*

(i) *If f_1 is constant, then f_2 is constant, too.*

(ii) *If f_2 is constant, then f_1 is constant, or $f_1(x^2) = \frac{c_1}{x^2 - c_2}$ on an open interval $I \subset \mathbb{R}$ for $c_1 \in \mathbb{R} \setminus \{0\}$, $c_2 \in \mathbb{R} \setminus I$.*

PROOF. In this case, we have $a = f_2 \frac{f'_1}{f_1}$, $b = c = 0$, $d = \frac{f'_2}{f_2}$, and

$$\text{Ric}(E_1, E_1) = E_2(a) - a^2, \quad \text{Ric}(E_2, E_2) = E_2(a) - a^2 + E_3(d) - d^2, \quad \text{Ric}(E_3, E_3) = E_3(d) - d^2,$$

$$\text{Ric}(E_1, E_2) = \text{Ric}(E_1, E_3) = 0, \quad \text{Ric}(E_2, E_3) = E_3(a)$$

from Proposition 2.10, and the system (3) becomes

$$\left\{ \begin{array}{l} f_2 \frac{\partial a}{\partial x^2} - a^2 + \lambda + \mu(\eta^1)^2 = 0 \\ -d + f_2 \frac{\partial a}{\partial x^2} - a^2 + d' - d^2 + \lambda + \mu(\eta^2)^2 = 0 \\ d' - d^2 + \lambda + \mu(\eta^3)^2 = 0 \\ \mu\eta^1\eta^2 = \mu\eta^1\eta^3 = 0 \\ \frac{\partial a}{\partial x^3} + \mu\eta^2\eta^3 = 0 \end{array} \right. . \quad (12)$$

Since μ is nowhere zero and $\eta^i = f$, (12) is equivalent to

$$\left\{ \begin{array}{l} f_2 \frac{\partial a}{\partial x^2} - a^2 - d = 0 \\ d = d' - d^2 \\ \lambda = -d \\ f^2 = 0 \\ \frac{\partial a}{\partial x^3} = 0 \end{array} \right. .$$

It follows that $\eta = 0$ and $f_1'f_2' = 0$, which, together with the second equation of the system, implies that one of the functions f_1 and f_2 must be a constant. If $f_1 = k_1 \in \mathbb{R} \setminus \{0\}$, then $a = 0$, $d = 0$ (hence, f_2 is constant), and $\lambda = 0$. If $f_2 = k_2 \in \mathbb{R} \setminus \{0\}$, then $d = 0$, $\lambda = 0$, and

$$\left(\frac{f_1'}{f_1} \right)' = \left(\frac{f_1'}{f_1} \right)^2 .$$

Then, either f_1 is a constant, too, or $f_1' \neq 0$ everywhere, in which case, by integration, we obtain $-\frac{1}{\frac{f_1'(x^2)}{f_1(x^2)}} = x^2 + c_0$, i.e., $f_1(x^2) = \frac{c_1}{x^2 - c_2}$ on an open interval $I \subset \mathbb{R}$ with $c_1 \in \mathbb{R} \setminus \{0\}$, $c_2 \in \mathbb{R} \setminus I$. Since $b = d = 0$, we deduce that $\mathcal{L}_V \tilde{g} = 0$, i.e., V is a Killing vector field. \square

3.2 Almost η -Ricci solitons with $\eta = dx^3$

If $\eta^1 = \eta^2 = 0$ and $\eta^3 = 1$, then the system (2) becomes

$$\begin{cases} E_1(V^1) - aV^2 - bV^3 + \text{Ric}(E_1, E_1) + \lambda = 0 \\ E_2(V^2) - cV^1 - dV^3 + \text{Ric}(E_2, E_2) + \lambda = 0 \\ E_3(V^3) + \text{Ric}(E_3, E_3) + \lambda + \mu = 0 \\ E_1(V^2) + E_2(V^1) + aV^1 + cV^2 = 0 \\ \frac{1}{2}[E_1(V^3) + E_3(V^1) + bV^1] + \text{Ric}(E_1, E_3) = 0 \\ \frac{1}{2}[E_2(V^3) + E_3(V^2) + dV^2] + \text{Ric}(E_2, E_3) = 0 \end{cases} \quad (13)$$

THEOREM 3.9. *Let $(\mathbb{R}^3, \tilde{g}, V, \lambda, \mu)$ be an η -Ricci soliton with $\eta = dx^3$. If $f_i = f_i(x^i)$ for $i \in \{1, 2\}$, then*

$$\begin{cases} V^1(x^1, x^2, x^3) = -\lambda F_1(x^1) + c_1 F_2(x^2) + c_2 x^3 + c_3 \\ V^2(x^1, x^2, x^3) = -c_1 F_1(x^1) - \lambda F_2(x^2) + c_4 x^3 + c_5 \\ V^3(x^1, x^2, x^3) = -(\lambda + \mu)x^3 - c_2 F_1(x^1) - c_4 F_2(x^2) + c_6 \end{cases} \quad (14)$$

where F_i is an antiderivative of $\frac{1}{f_i}$ for $i \in \{1, 2\}$ and $c_i \in \mathbb{R}$ for $i \in \{1, 2, 3, 4, 5, 6\}$.

PROOF. In this case, we have $a = b = c = d = 0$ and $\text{Ric}(E_i, E_j) = 0$ for any $i, j \in \{1, 2, 3\}$ from Proposition 2.1, and the system (13) becomes

$$\begin{cases} \frac{\partial V^1}{\partial x^1} = -\frac{\lambda}{f_1} \\ \frac{\partial V^2}{\partial x^2} = -\frac{\lambda}{f_2} \\ \frac{\partial V^3}{\partial x^3} = -(\lambda + \mu) \\ f_1 \frac{\partial V^2}{\partial x^1} = -f_2 \frac{\partial V^1}{\partial x^2} \\ f_1 \frac{\partial V^3}{\partial x^1} = -\frac{\partial V^1}{\partial x^3} \\ f_2 \frac{\partial V^3}{\partial x^2} = -\frac{\partial V^2}{\partial x^3} \end{cases} \quad (15)$$

Since f_i depends only on x^i , from the first three equations of (15), we find that

$$\begin{cases} V^1(x^1, x^2, x^3) = -\lambda F_1(x^1) + h_1(x^2, x^3) \\ V^2(x^1, x^2, x^3) = -\lambda F_2(x^2) + h_2(x^1, x^3) \\ V^3(x^1, x^2, x^3) = -(\lambda + \mu)x^3 + h_3(x^1, x^2) \end{cases},$$

where $F'_i = \frac{1}{f_i}$ for $i \in \{1, 2\}$. From the last three equations of (15), we infer:

$$f_1(x^1) \frac{\partial h_2}{\partial x^1}(x^1, x^3) = -f_2(x^2) \frac{\partial h_1}{\partial x^2}(x^2, x^3), \quad (16)$$

$$f_1(x^1) \frac{\partial h_3}{\partial x^1}(x^1, x^2) = -\frac{\partial h_1}{\partial x^3}(x^2, x^3), \quad (17)$$

$$f_2(x^2) \frac{\partial h_3}{\partial x^2}(x^1, x^2) = -\frac{\partial h_2}{\partial x^3}(x^1, x^3). \quad (18)$$

Denoting $f_2(x^2) \frac{\partial h_1}{\partial x^2}(x^2, x^3) =: \bar{h}_1(x^3)$, we have

$$\frac{\partial h_1}{\partial x^2}(x^2, x^3) = \frac{\bar{h}_1(x^3)}{f_2(x^2)},$$

which, by integration, implies

$$h_1(x^2, x^3) = \bar{h}_1(x^3) F_2(x^2) + \hat{h}_1(x^3),$$

and, by replacing it in (17), it gives

$$-\bar{h}'_1(x^3) F_2(x^2) - f_1(x^1) \frac{\partial h_3}{\partial x^1}(x^1, x^2) = \hat{h}'_1(x^3).$$

Differentiating the previous relation with respect to x^2 , it implies

$$\bar{h}'_1(x^3) = -f_1(x^1) f_2(x^2) \frac{\partial^2 h_3}{\partial x^2 \partial x^1}(x^1, x^2),$$

which must be a constant, let's say, c_1 . Therefore,

$$\bar{h}_1(x^3) = c_1 x^3 + c_2$$

and

$$\frac{\partial^2 h_3}{\partial x^2 \partial x^1}(x^1, x^2) = -\frac{c_1}{f_1(x^1) f_2(x^2)},$$

which, by double integration, implies

$$h_3(x^1, x^2) = -c_1 F_1(x^1) F_2(x^2) + l_1(x^1) + l_2(x^2).$$

Also, from (16), we have $\frac{\partial h_2}{\partial x^1}(x^1, x^3) = -\frac{\bar{h}_1(x^3)}{f_1(x^1)}$, which gives

$$h_2(x^1, x^3) = -\bar{h}_1(x^3) F_1(x^1) + \hat{h}_2(x^3).$$

From (17), we find

$$\hat{h}_1'(x^3) = -f_1(x^1) l_1'(x^1),$$

which must be a constant, let's say, c_3 , which gives

$$\hat{h}_1(x^3) = c_3 x^3 + c_4.$$

Since $f_1(x^1) l_1'(x^1) = -c_3$, we get

$$l_1(x^1) = -c_3 F_1(x^1) + c_5.$$

From (18), we find

$$\hat{h}_2'(x^3) = 2c_1 F_1(x^1) - f_2(x^2) l_2'(x^2),$$

which must be a constant, let's say, c_6 , which gives

$$\hat{h}_2(x^3) = c_6 x^3 + c_7.$$

Since $2c_1 F_1(x^1) = f_2(x^2) l_2'(x^2) + c_6$ and F_1 is not constant, we get

$$c_1 = 0, \quad l_2(x^2) = -c_6 F_2(x^2) + c_8.$$

We have obtained

$$\begin{cases} V^1(x^1, x^2, x^3) = -\lambda F_1(x^1) + c_2 F_2(x^2) + c_3 x^3 + c_4 \\ V^2(x^1, x^2, x^3) = -\lambda F_2(x^2) - c_2 F_1(x^1) + c_6 x^3 + c_7 \\ V^3(x^1, x^2, x^3) = -(\lambda + \mu) x^3 - c_3 F_1(x^1) + c_5 - c_6 F_2(x^2) + c_8 \end{cases},$$

which, by changing the indices of some constants, gives (14). □

EXAMPLE 3.10. For $f_i = k_i \in \mathbb{R} \setminus \{0\}$, $i \in \{1, 2\}$, the vector field V with V^1 , V^2 and V^3 given by:

$$\begin{cases} V^1(x^1, x^2, x^3) = -\frac{\lambda}{k_1}x^1 + \frac{c_1}{k_2}x^2 + c_2x^3 + c_3 \\ V^2(x^1, x^2, x^3) = -\frac{c_1}{k_1}x^1 - \frac{\lambda}{k_2}x^2 + c_4x^3 + c_5 \\ V^3(x^1, x^2, x^3) = -\frac{c_2}{k_1}x^1 - \frac{c_4}{k_2}x^2 - (\lambda + \mu)x^3 + c_6 \end{cases},$$

where $c_i \in \mathbb{R}$ for $i \in \{1, 2, 3, 4, 5, 6\}$, is the potential vector field of the η -Ricci soliton $(\mathbb{R}^3, \tilde{g}, \lambda, \mu)$ for $\eta = dx^3$.

REMARK 3.11. If $f_i = f_i(x^i)$ for $i \in \{1, 2\}$, then V with V^1 , V^2 and V^3 given by:

$$\begin{cases} V^1(x^2, x^3) = c_1F_2(x^2) + c_2x^3 + c_3 \\ V^2(x^1, x^3) = -c_1F_1(x^1) + c_4x^3 + c_5 \\ V^3(x^1, x^2) = -c_2F_1(x^1) - c_4F_2(x^2) + c_6 \end{cases}$$

is a Killing vector field on $(\mathbb{R}^3, \tilde{g})$, where F_i is an antiderivative of $\frac{1}{f_i}$ for $i \in \{1, 2\}$ and $c_i \in \mathbb{R}$ for $i \in \{1, 2, 3, 4, 5, 6\}$.

THEOREM 3.12. Let $(\mathbb{R}^3, \tilde{g}, V, \lambda, \mu)$ be an almost η -Ricci soliton with $\eta = dx^3$. If $f_i = f_i(x^i)$ for $i \in \{1, 2\}$, and $V^i = V^i(x^3)$ for $i \in \{1, 2, 3\}$, then:

$$\begin{cases} V^1 = \frac{c_1}{f_1} \\ V^2 = \frac{c_2}{f_2} \\ (b-d)V^3 = (b-d)' - (b^2 - d^2) \\ (b-d)\lambda = b'd - bd' \\ (b-d)^2\mu = -(b-d)[(b-d)'' + b'd - bd'] + [(b-d)']^2 + (b-d)^2(b^2 + d^2) \end{cases},$$

with $c_1, c_2 \in \mathbb{R}$.

We have the following cases.

(i) On any open interval $I \subset \mathbb{R}$ on which $\left(\frac{f_1}{f_2}\right)' \neq 0$ everywhere (equivalent to $b \neq d$ on I), we have:

$$V^1 = \frac{c_1}{f_1}, \quad V^2 = \frac{c_2}{f_2}, \quad V^3 = \frac{(b-d)'}{b-d} - (b+d),$$

$$\lambda = \frac{b'd - bd'}{b-d}, \quad \mu = -\frac{(b-d)'' + b'd - bd'}{b-d} + \left(\frac{(b-d)'}{b-d}\right)^2 + b^2 + d^2,$$

with $c_1, c_2 \in \mathbb{R}$.

In particular:

(1) if f_1 is constant on an open interval $J \subset I$ (from which $f_2' \neq 0$ everywhere on J), we have on J :

$$V^1 = c_1, \quad V^2 = \frac{c_2}{f_2}, \quad V^3 = \frac{d' - d^2}{d},$$

$$\lambda = 0, \quad \mu = -\frac{d''}{d} + \left(\frac{d'}{d}\right)^2 + d^2,$$

with $c_1, c_2 \in \mathbb{R}$;

(2) if f_2 is constant on an open interval $J \subset I$ (from which $f_1' \neq 0$ everywhere on J), we have on J :

$$V^1 = \frac{c_1}{f_1}, \quad V^2 = c_2, \quad V^3 = \frac{b' - b^2}{b},$$

$$\lambda = 0, \quad \mu = -\frac{b''}{b} + \left(\frac{b'}{b}\right)^2 + b^2,$$

with $c_1, c_2 \in \mathbb{R}$.

(ii) On any open interval $I \subset \mathbb{R}$ on which $\frac{f_1}{f_2}$ is constant (equivalent to $b = d$ on I), we have:

$$V^1 = \frac{c_1}{f_1}, \quad V^2 = \frac{c_2}{f_2}, \quad bV^3 = b' - 2b^2 + \lambda,$$

with $c_1, c_2 \in \mathbb{R}$.

In addition:

(1) if $J \subset I$ is an open interval with $f_1' \neq 0$ (and $f_2' \neq 0$) everywhere on J , we have on J :

$$V^1 = \frac{c_1}{f_1}, \quad V^2 = \frac{c_2}{f_2}, \quad V^3 = \frac{b' - 2b^2 + \lambda}{b},$$

$$\mu = -\left(\frac{b' + \lambda}{b}\right)' + 2b^2 - \lambda,$$

with $c_1, c_2 \in \mathbb{R}$;

(2) if f_1 and f_2 are constant on an open interval $K \subset I$, we have on K :

$$V^1 = c_1, \quad V^2 = c_2, \quad V^3 = F,$$

$$\lambda = 0,$$

where $F' = -\mu$ and $c_1, c_2 \in \mathbb{R}$.

PROOF. In this case, we have $a = 0$, $b = \frac{f_1'}{f_1}$, $c = 0$, $d = \frac{f_2'}{f_2}$, and

$$\text{Ric}(E_1, E_1) = b' - b^2 - bd, \quad \text{Ric}(E_2, E_2) = d' - d^2 - bd, \quad \text{Ric}(E_3, E_3) = b' + d' - b^2 - d^2,$$

$$\text{Ric}(E_1, E_2) = \text{Ric}(E_1, E_3) = \text{Ric}(E_2, E_3) = 0$$

from Proposition 2.2, and the system (13) becomes

$$\left\{ \begin{array}{l} f_1 \frac{\partial V^1}{\partial x^1} - bV^3 + b' - b^2 - bd + \lambda = 0 \\ f_2 \frac{\partial V^2}{\partial x^2} - dV^3 + d' - d^2 - bd + \lambda = 0 \\ \frac{\partial V^3}{\partial x^3} + b' - b^2 + d' - d^2 + \lambda + \mu = 0 \\ f_1 \frac{\partial V^2}{\partial x^1} + f_2 \frac{\partial V^1}{\partial x^2} = 0 \\ f_1 \frac{\partial V^3}{\partial x^1} + \frac{\partial V^1}{\partial x^3} + bV^1 = 0 \\ f_2 \frac{\partial V^3}{\partial x^2} + \frac{\partial V^2}{\partial x^3} + dV^2 = 0 \end{array} \right. , \quad (19)$$

which is equivalent to

$$\left\{ \begin{array}{l} (V^1)' = -bV^1 \\ (V^2)' = -dV^2 \\ (V^3)' = -(b' - b^2 + d' - d^2) - (\lambda + \mu) . \\ bV^3 = b' - b^2 - bd + \lambda \\ dV^3 = d' - d^2 - bd + \lambda \end{array} \right.$$

From the first two equations we get either $V^i = 0$ or $\frac{(V^i)'}{V^i} = -\frac{f_i'}{f_i}$, i.e., $V^i = \frac{c_i}{f_i}$ with $c_i \in \mathbb{R} \setminus \{0\}$, for $i \in \{1, 2\}$. From the last two equations, we obtain

$$(b - d)V^3 = (b - d)' - (b^2 - d^2)$$

and

$$(b - d)\lambda = b'd - bd'$$

Differentiating the second-last equality, multiplying the result with $b - d$ and substituting in it the expressions of $(b - d)V^3$ and $(V^3)'$, we get

$$(b - d)^2\mu = -(b - d)[(b - d)'' + b'd - bd'] + [(b - d)']^2 + (b - d)^2(b^2 + d^2).$$

(i) For $x \in \mathbb{R}$, $\left(\frac{f_1}{f_2}\right)'(x) \neq 0$ if and only if $b(x) \neq d(x)$.

From the last three equalities from above, on any open interval $I \subset \mathbb{R}$ such that $\left(\frac{f_1}{f_2}\right)'(x) \neq 0$ for any $x \in I$ (equivalent to $b \neq d$ everywhere on I), we get the expressions of V^3 , λ and μ .

(ii) $\frac{f_1}{f_2}$ is constant on I if and only if $\left(\frac{f_1}{f_2}\right)' = 0$ on I , equivalent to $b = d$ on I .

In this case, (23) becomes

$$\begin{cases} (V^1)' = -bV^1 \\ (V^2)' = -bV^2 \\ (V^3)' = -2(b' - b^2) - (\lambda + \mu) \\ bV^3 = b' - 2b^2 + \lambda \end{cases}.$$

On any open interval $J \subset I$ where $f_1' \neq 0$ everywhere, we have $b \neq 0$ and

$$V^3 = \frac{b' - 2b^2 + \lambda}{b},$$

which, by differentiation and using the third relation, gives the expression of μ .

If f_1 and f_2 are constant on $K \subset I$, then $b = d = 0$ on K and (23) becomes

$$\begin{cases} \lambda = 0 \\ (V^1)' = 0 \\ (V^2)' = 0 \\ (V^3)' = -\mu \quad \square \end{cases}.$$

EXAMPLE 3.13. The Riemannian Sol_3 Lie group $(\mathbb{R}^3, \tilde{g})$, where $f_1(x^3) = e^{-x^3}$ and $f_2(x^3) = e^{x^3}$, is an η -Ricci soliton for $\eta = dx^3$, with

$$V = c_1 \frac{\partial}{\partial x^1} + c_2 \frac{\partial}{\partial x^2} \quad (c_1, c_2 \in \mathbb{R}), \quad \lambda = 0, \quad \mu = 2.$$

REMARK 3.14. If $f_i = f_i(x^3)$ for $i \in \{1, 2\}$, then V with V^1 , V^2 and V^3 given by:

$$\begin{cases} V^1(x^3) = \frac{c_1}{f_1(x^3)} \\ V^2(x^3) = \frac{c_2}{f_2(x^3)} \\ V^3 = 0 \end{cases}$$

is a Killing vector field on $(\mathbb{R}^3, \tilde{g})$, where $c_i \in \mathbb{R}$ for $i \in \{1, 2\}$.

THEOREM 3.15. *Let $(\mathbb{R}^3, \tilde{g}, V, \lambda, \mu)$ be an almost η -Ricci soliton with $\eta = dx^3$. If $f_1 = f_1(x^1)$, $f_2 = f_2(x^3)$, and $V^i = V^i(x^3)$ for $i \in \{1, 2, 3\}$, then:*

$$\begin{cases} V^1 = c_1 \\ V^2 = \frac{c_2}{f_2} \\ dV^3 = d' - d^2 \\ \lambda = 0 \\ d^2\mu = (d')^2 + d^4 - dd'' \end{cases},$$

with $c_1, c_2 \in \mathbb{R}$.

We have the following cases.

(i) On any open interval $I \subset \mathbb{R}$ on which f_2 is constant, we have:

$$V^1 = c_1, \quad V^2 = c_2, \quad V^3 = F,$$

$$\lambda = 0,$$

where $F' = -\mu$ and $c_1, c_2 \in \mathbb{R}$.

(ii) On any open interval $I \subset \mathbb{R}$ on which $f_2' \neq 0$ everywhere (equivalent to $d \neq 0$ on I), we have:

$$V^1 = c_1, \quad V^2 = \frac{c_2}{f_2}, \quad V^3 = \frac{d' - d^2}{d},$$

$$\lambda = 0, \quad \mu = -\frac{d''}{d} + \left(\frac{d'}{d}\right)^2 + d^2,$$

with $c_1, c_2 \in \mathbb{R}$.

PROOF. In this case, we have $a = b = c = 0$, $d = \frac{f_2'}{f_2}$, and

$$\text{Ric}(E_2, E_2) = \text{Ric}(E_3, E_3) = d' - d^2,$$

$$\text{Ric}(E_1, E_1) = \text{Ric}(E_1, E_2) = \text{Ric}(E_1, E_3) = \text{Ric}(E_2, E_3) = 0$$

from Proposition 2.4, and the system (13) becomes

$$\left\{ \begin{array}{l} f_1 \frac{\partial V^1}{\partial x^1} + \lambda = 0 \\ f_2 \frac{\partial V^2}{\partial x^2} - dV^3 + d' - d^2 + \lambda = 0 \\ \frac{\partial V^3}{\partial x^3} + d' - d^2 + \lambda + \mu = 0 \\ f_1 \frac{\partial V^2}{\partial x^1} + f_2 \frac{\partial V^1}{\partial x^2} = 0 \\ f_1 \frac{\partial V^3}{\partial x^1} + \frac{\partial V^1}{\partial x^3} = 0 \\ f_2 \frac{\partial V^3}{\partial x^2} + \frac{\partial V^2}{\partial x^3} + dV^2 = 0 \end{array} \right. , \quad (20)$$

equivalent to

$$\left\{ \begin{array}{l} \lambda = 0 \\ (V^1)' = 0 \\ (V^2)' = -dV^2 \\ (V^3)' = -d' + d^2 - \mu \\ dV^3 = d' - d^2 \end{array} \right. ,$$

from which we get

$$V^1 = c_1, \quad V^2 = \frac{c_2}{f_2},$$

with $c_1, c_2 \in \mathbb{R}$. Differentiating the last equality of the previous system, multiplying the result with d and substituting in it the expressions of dV^3 and $(V^3)'$, we get

$$d^2\mu = (d')^2 + d^4 - dd''.$$

(i) f_2 is constant on I if and only if $d = 0$ on I , and (20) becomes

$$\left\{ \begin{array}{l} \lambda = 0 \\ (V^1)' = 0 \\ (V^2)' = 0 \\ (V^3)' = -\mu \end{array} \right. .$$

(ii) $f_2' \neq 0$ on I if and only if $d \neq 0$ on I . Dividing the relations for dV^3 and $d^2\mu$ by d and d^2 , respectively, we obtain the expressions of V^3 and μ . \square

EXAMPLE 3.16. For $f_1(x^1) = e^{x^1}$, $f_2(x^3) = e^{x^3}$, the vector field

$$V = c_1 e^{x^1} \frac{\partial}{\partial x^1} + c_2 \frac{\partial}{\partial x^2} - \frac{\partial}{\partial x^3} \quad (c_1, c_2 \in \mathbb{R})$$

is the potential vector field of the η -Ricci soliton $(\mathbb{R}^3, \tilde{g}, 0, 1)$ for $\eta = dx^3$.

THEOREM 3.17. *Let $(\mathbb{R}^3, \tilde{g}, V, \lambda, \mu)$ be an almost η -Ricci soliton with $\eta = dx^3$. If $f_i = f_i(x^2)$ for $i \in \{1, 2\}$ and $V^i = V^i(x^3)$ for $i \in \{1, 2, 3\}$, then:*

$$\begin{cases} V^1 = c_1 \\ V^2 = c_2 \\ V^3 = F \\ \lambda = 2f_2^2 \left(\frac{f_1'}{f_1} \right)^2 - f_2 f_2' \frac{f_1'}{f_1} - f_2^2 \frac{f_1''}{f_1} \end{cases},$$

$$c_1 f_1' = 0 = c_2 f_1',$$

where $c_1, c_2 \in \mathbb{R}$, $F' = -(\lambda + \mu)$, that is, $\mu(x^2, x^3) = -\lambda(x^2) - G(x^3)$, with $G = G(x^3)$ a smooth function on \mathbb{R} , and $F' = G$.

We have the following cases.

(i) *If $I \subset \mathbb{R}$ is a nontrivial interval on which f_1 is constant, then:*

$$\lambda = 0, \quad F'(x^3) = -\mu(x^2, x^3) \quad \text{for } x^2 \in I, x^3 \in \mathbb{R}.$$

(ii) *If f_1 is constant, then:*

$$\lambda = 0, \quad \mu = \mu(x^3), \quad F'(x^3) = -\mu(x^3) \quad \text{for } x^3 \in \mathbb{R}.$$

(iii) *If f_1 is not constant, then $V^1 = V^2 = 0$.*

PROOF. In this case, we have $a = f_2 \frac{f_1'}{f_1}$, $b = c = d = 0$, and

$$\text{Ric}(E_1, E_1) = \text{Ric}(E_2, E_2) = E_2(a) - a^2,$$

$$\text{Ric}(E_1, E_2) = \text{Ric}(E_1, E_3) = \text{Ric}(E_2, E_3) = \text{Ric}(E_3, E_3) = 0$$

from Proposition 2.10, and the system (13) becomes

$$\left\{ \begin{array}{l} f_1 \frac{\partial V^1}{\partial x^1} - aV^2 + f_2 a' - a^2 + \lambda = 0 \\ f_2 \frac{\partial V^2}{\partial x^2} + f_2 a' - a^2 + \lambda = 0 \\ \frac{\partial V^3}{\partial x^3} + \lambda + \mu = 0 \\ f_1 \frac{\partial V^2}{\partial x^1} + f_2 \frac{\partial V^1}{\partial x^2} + aV^1 = 0 \\ f_1 \frac{\partial V^3}{\partial x^1} + \frac{\partial V^1}{\partial x^3} = 0 \\ f_2 \frac{\partial V^3}{\partial x^2} + \frac{\partial V^2}{\partial x^3} = 0 \end{array} \right. ,$$

that is,

$$\left\{ \begin{array}{l} aV^1 = 0 \\ aV^2 = 0 \\ (V^1)' = 0 \\ (V^2)' = 0 \\ (V^3)' = -(\lambda + \mu) \\ f_2 a' - a^2 + \lambda = 0 \end{array} \right. .$$

From the second-last equation, we deduce that $\lambda + \mu$ depends only on x^3 ; therefore,

$$0 = \frac{\partial(\lambda + \mu)}{\partial x^2} = \lambda' + \frac{\partial \mu}{\partial x^2} ,$$

which, by integration, gives

$$\mu(x^2, x^3) = -\lambda(x^2) - G(x^3)$$

with $G = G(x^3)$ a smooth function on \mathbb{R} and we will denote by F an antiderivative of G , i.e., $F' = G$. We obtain

$$\left\{ \begin{array}{l} V^1 = c_1 \\ V^2 = c_2 \\ V^3 = F \\ aV^1 = 0 \\ aV^2 = 0 \\ \lambda = -f_2 a' + a^2 \end{array} \right. , \tag{21}$$

where $F' = -(\lambda + \mu)$ and $c_1, c_2 \in \mathbb{R}$.

(i) f_1 is constant on I if and only if $a = 0$ on I ; therefore, $a' = 0$ and $\lambda = 0$ on I , hence $F' = -\mu$.

(ii) It follows from (i) for $I = \mathbb{R}$.

(iii) For f_1 not constant, there exists $x_0^2 \in \mathbb{R}$ such that $f_1'(x_0^2) \neq 0$, i.e., $a(x_0^2) \neq 0$ and, from the second- and third-last equations of (21), it follows that $V_1 = V_2 = 0$. \square

EXAMPLE 3.18. The manifold $\mathbb{H}^2 \times \mathbb{R} := \{(x^1, x^2, x^3) \in \mathbb{R}^3 : x^2 > 0\}$ with

$$\tilde{g} = \frac{1}{f^2} dx^1 \otimes dx^1 + \frac{1}{f^2} dx^2 \otimes dx^2 + dx^3 \otimes dx^3,$$

where $f(x^2) = x^2$, is an η -Ricci soliton for $\eta = dx^3$, with

$$V = \frac{\partial}{\partial x^3}, \quad \lambda = 1, \quad \mu = -1.$$

THEOREM 3.19. Let $(\mathbb{R}^3, \tilde{g}, V, \lambda, \mu)$ be an almost η -Ricci soliton with $\eta = dx^3$. If $f_1 = f_1(x^2)$, $f_2 = f_2(x^1)$, and $V^i = V^i(x^3)$ for $i \in \{1, 2, 3\}$, then:

$$\left\{ \begin{array}{l} V^1 = c_1 \\ V^2 = c_2 \\ V^3 = F \\ \lambda = c_1 f_1 \frac{f_2'}{f_2} + f_1^2 \left[\left(\frac{f_2'}{f_2} \right)^2 - \left(\frac{f_2'}{f_2} \right)' \right] + f_2^2 \left[\left(\frac{f_1'}{f_1} \right)^2 - \left(\frac{f_1'}{f_1} \right)' \right] \\ c_1 f_2 \frac{f_1'}{f_1} = -c_2 f_1 \frac{f_2'}{f_2}, \quad c_2 f_2 \frac{f_1'}{f_1} = c_1 f_1 \frac{f_2'}{f_2} \end{array} \right. ,$$

where $c_1, c_2 \in \mathbb{R}$ and $F' = -(\lambda + \mu)$.

For $c_1 \neq 0$ or $c_2 \neq 0$, we get $\lambda = 0$ and $F' = -\mu$.

PROOF. In this case, we have $a = f_2 \frac{f_1'}{f_1}$, $b = 0$, $c = f_1 \frac{f_2'}{f_2}$, $d = 0$, and

$$\text{Ric}(E_1, E_1) = \text{Ric}(E_2, E_2) = E_1(c) + E_2(a) - a^2 - c^2,$$

$$\text{Ric}(E_1, E_2) = \text{Ric}(E_1, E_3) = \text{Ric}(E_2, E_3) = \text{Ric}(E_3, E_3) = 0$$

from Proposition 2.8, and the system (13) becomes

$$\begin{cases} E_1(V^1) - aV^2 + E_1(c) + E_2(a) - a^2 - c^2 + \lambda = 0 \\ E_2(V^2) - cV^1 + E_1(c) + E_2(a) - a^2 - c^2 + \lambda = 0 \\ E_3(V^3) + \lambda + \mu = 0 \\ E_1(V^2) + E_2(V^1) + aV^1 + cV^2 = 0 \\ E_1(V^3) + E_3(V^1) = 0 \\ E_2(V^3) + E_3(V^2) = 0 \end{cases} \quad (22)$$

Since V^i for $i \in \{1, 2, 3\}$ depends only on x^3 , (22) becomes

$$\begin{cases} aV^2 = f_1 \frac{\partial c}{\partial x^1} + f_2 \frac{\partial a}{\partial x^2} - a^2 - c^2 + \lambda = cV^1 \\ aV^1 = -cV^2 \\ (V^1)' = 0 \\ (V^2)' = 0 \\ (V^3)' = -(\lambda + \mu) \end{cases}.$$

From the last three equations, we infer $V^1 = c_1$, $V^2 = c_2$ for $c_1, c_2 \in \mathbb{R}$, and $V^3 = F$, where $F' = -(\lambda + \mu)$, and the previous system is equivalent to

$$\begin{cases} V^1 = c_1 \\ V^2 = c_2 \\ \lambda = c_1 f_1 \frac{f_2'}{f_2} + f_1^2 \left[\left(\frac{f_2'}{f_2} \right)^2 - \left(\frac{f_2'}{f_2} \right)' \right] + f_2^2 \left[\left(\frac{f_1'}{f_1} \right)^2 - \left(\frac{f_1'}{f_1} \right)' \right] \\ c_2 a = c_1 c \\ c_1 a = -c_2 c \\ V^3 = F \end{cases}.$$

If $c_1 \neq 0$, then $(c_1^2 + c_2^2)a = 0$, hence $a = 0$, i.e., f_1 is constant. It follows that $c = 0$, i.e., f_2 is constant. In this case,

$$\lambda = 0, \quad V^3 = F,$$

where $F' = -\mu$. The same thing happens if $c_2 \neq 0$. So, we get the conclusion. \square

EXAMPLE 3.20. For $f_1(x^2) = e^{x^2}$ and $f_2(x^1) = e^{x^1}$, the vector field

$$V = e^{x^3} \frac{\partial}{\partial x^3}$$

is the potential vector field of the almost η -Ricci soliton

$$\left(\mathbb{R}^3, \tilde{g}, \lambda(x^1, x^2) = e^{2x^1} + e^{2x^2}, \mu(x^1, x^2, x^3) = -(e^{2x^1} + e^{2x^2} + e^{2x^3}) \right)$$

for $\eta = dx^3$.

THEOREM 3.21. *Let $(\mathbb{R}^3, \tilde{g}, V, \lambda, \mu)$ be an almost η -Ricci soliton with $\eta = dx^3$. If $f_1 = f_1(x^2)$, $f_2 = f_2(x^3)$, and $V^i = V^i(x^3)$ for $i \in \{1, 2, 3\}$, then:*

$$\left\{ \begin{array}{l} V^1 = c_1 \\ ac_1 = 0 \\ (V^2)' + dV^2 = -2\frac{f_1'f_2'}{f_1} \\ aV^2 = f_2^2 \left[\left(\frac{f_1'}{f_1} \right)' - \left(\frac{f_1'}{f_1} \right)^2 \right] + \lambda \\ (V^3)' = -(d' - d^2) - (\lambda + \mu) \\ dV^3 = f_2^2 \left[\left(\frac{f_1'}{f_1} \right)' - \left(\frac{f_1'}{f_1} \right)^2 \right] + \lambda + d' - d^2 \end{array} \right. ,$$

with $c_1 \in \mathbb{R}$.

In addition:

(i) if $V^1 \neq 0$, then f_1 is constant, and, in this case, we have:

$$\left\{ \begin{array}{l} V^1 = c_1 \\ V^2 = \frac{c_2}{f_2} \\ (V^3)' = -(d' - d^2) - \mu \\ dV^3 = d' - d^2 \\ \lambda = 0 \\ d^2\mu = d^4 + (d')^2 - dd'' \end{array} \right. ,$$

with $c_1 \in \mathbb{R} \setminus \{0\}$, $c_2 \in \mathbb{R}$; moreover,

(1) on any open interval $I \subset \mathbb{R}$ on which f_2 is constant and $V^1 \neq 0$, we have:

$$V^1 = c_1, \quad V^2 = c_2, \quad V^3 = F,$$

$$\lambda = 0,$$

where F is an antiderivative of $-\mu$ and $c_1 \in \mathbb{R} \setminus \{0\}$, $c_2 \in \mathbb{R}$;

(2) on any open interval $I \subset \mathbb{R}$ on which $f_2' \neq 0$ everywhere (equivalent to $d \neq 0$ on I) and $V^1 \neq 0$, we have:

$$V^1 = c_1, \quad V^2 = \frac{c_2}{f_2}, \quad V^3 = \frac{d' - d^2}{d},$$

$$\lambda = 0, \quad \mu = -\frac{d''}{d} + \left(\frac{d'}{d}\right)^2 + d^2,$$

with $c_1 \in \mathbb{R} \setminus \{0\}$, $c_2 \in \mathbb{R}$;

(ii) if $V^1 = 0$, we have:

$$\left\{ \begin{array}{l} V^1 = 0 \\ V^2 = \frac{F}{c_2|f_2|} \\ (V^3)' = -(d' - d^2) - (\lambda + \mu) \\ dV^3 = f_2^2 \left[\left(\frac{f_1'}{f_1}\right)' - \left(\frac{f_1'}{f_1}\right)^2 \right] + \lambda + d' - d^2 \\ \lambda = \frac{f_1' f_2}{c_2 f_1 |f_2|} F - f_2^2 \left[\left(\frac{f_1'}{f_1}\right)' - \left(\frac{f_1'}{f_1}\right)^2 \right] \\ d^2 \mu = d^4 + (d')^2 - dd'' + (d' - d^2)\lambda - d \frac{\partial \lambda}{\partial x^3} + [f_2 f_2'' - 3(f_2')^2] \left[\left(\frac{f_1'}{f_1}\right)' - \left(\frac{f_1'}{f_1}\right)^2 \right] \end{array} \right. ,$$

where F is an antiderivative of $-2c_2 \frac{f_1' f_2' |f_2|}{f_1}$ and $c_2 > 0$; moreover,

(1) on any open interval $I \subset \mathbb{R}$ on which f_2 is constant and $V^1 = 0$, we have:

$$V^1 = 0, \quad V^2 = c_3, \quad V^3 = F,$$

$$\lambda = -f_2^2 \left[\left(\frac{f_1'}{f_1}\right)' - \left(\frac{f_1'}{f_1}\right)^2 \right],$$

$$f_1 \text{ is constant or } c_3 = 0,$$

where F is an antiderivative of $-(\lambda + \mu)$ and $c_3 \in \mathbb{R}$;

(2) on any open interval $I \subset \mathbb{R}$ on which $f_2' \neq 0$ everywhere (equivalent to $d \neq 0$ on I) and $V^1 = 0$, we have:

$$V^1 = 0, \quad V^2 = -c_4 f_2 + \frac{c_6}{c_2 |f_2|}, \quad V^3 = -c_4^2 f_2^2 + \frac{c_4 c_6}{c_2} \frac{f_2}{|f_2|} + \left(\frac{f_2'}{f_2} \right)' \frac{f_2}{f_2'} - \frac{f_2'}{f_2},$$

$$\lambda = \frac{c_4 c_6}{c_2} \frac{f_2}{|f_2|}, \quad \mu = - \left[\left(\frac{f_2'}{f_2} \right)' - \left(\frac{f_2'}{f_2} \right)^2 \right] - \frac{c_4 c_6}{c_2} \frac{f_2}{|f_2|} - (V^3)',$$

with $c_2 > 0$, $c_4, c_6 \in \mathbb{R}$.

PROOF. In this case, we have $a = f_2 \frac{f_1'}{f_1}$, $b = c = 0$, $d = \frac{f_2'}{f_2}$, and

$$\text{Ric}(E_1, E_1) = E_2(a) - a^2, \quad \text{Ric}(E_2, E_2) = E_2(a) - a^2 + E_3(d) - d^2, \quad \text{Ric}(E_3, E_3) = E_3(d) - d^2,$$

$$\text{Ric}(E_1, E_2) = \text{Ric}(E_1, E_3) = 0, \quad \text{Ric}(E_2, E_3) = E_3(a)$$

from Proposition 2.10, and the system (13) becomes

$$\begin{cases} f_1 \frac{\partial V^1}{\partial x^1} - aV^2 + f_2 \frac{\partial a}{\partial x^2} - a^2 + \lambda = 0 \\ f_2 \frac{\partial V^2}{\partial x^2} - dV^3 + d' - d^2 + f_2 \frac{\partial a}{\partial x^2} - a^2 + \lambda = 0 \\ \frac{\partial V^3}{\partial x^3} + d' - d^2 + \lambda + \mu = 0 \\ f_1 \frac{\partial V^2}{\partial x^1} + f_2 \frac{\partial V^1}{\partial x^2} + aV^1 = 0 \\ f_1 \frac{\partial V^3}{\partial x^1} + \frac{\partial V^1}{\partial x^3} = 0 \\ \frac{1}{2} \left[f_2 \frac{\partial V^3}{\partial x^2} + \frac{\partial V^2}{\partial x^3} + dV^2 \right] + \frac{\partial a}{\partial x^3} = 0 \end{cases}, \quad (23)$$

which is equivalent to

$$\begin{cases} (V^1)' = 0 \\ aV^1 = 0 \\ (V^2)' + dV^2 = -2 \frac{\partial a}{\partial x^3} \\ aV^2 = f_2 \frac{\partial a}{\partial x^2} - a^2 + \lambda \\ (V^3)' = -(d' - d^2) - (\lambda + \mu) \\ dV^3 = f_2 \frac{\partial a}{\partial x^2} - a^2 + d' - d^2 + \lambda \end{cases}. \quad (24)$$

From the first two equations, it follows that either $V^1 = 0$, or $V^1 = c_1 \in \mathbb{R} \setminus \{0\}$ and $a = 0$, i.e., f_1 is constant. Let $V^1 = 0$. We get $V^2 = Fe^{-G}$, where F is an antiderivative of $-2\frac{f'_1 f'_2}{f_1} e^G$ and G is an antiderivative of d , i.e., $G = \ln(c_2|f_2|)$ with $c_2 > 0$, from which $F' = -2c_2\frac{f'_1 f'_2 |f_2|}{f_1}$, hence, $V^2 = \frac{F}{c_2|f_2|}$. From the fourth equation of the same system, we get the expression of λ . Differentiating the last equality of the previous system, multiplying the result with d and substituting in it the expressions of dV^3 and $(V^3)'$, we get

$$d^2\mu = d^4 + (d')^2 - dd'' + (d' - d^2)\lambda - d\frac{\partial\lambda}{\partial x^3} + [f_2 f_2'' - 3(f_2')^2] \left[\left(\frac{f'_1}{f_1}\right)' - \left(\frac{f'_1}{f_1}\right)^2 \right].$$

Also,

$$\begin{aligned} \mu &= -\lambda - (d' - d^2) - (V^3)', \\ dV^3 &= \frac{f'_1 f_2 F}{c_2 f_1 |f_2|} + \left(\frac{f'_2}{f_2}\right)' - \left(\frac{f'_2}{f_2}\right)^2. \end{aligned}$$

For $f'_2 \neq 0$ everywhere on an open interval $I \subset \mathbb{R}$ (equivalent to $d' \neq 0$ everywhere on I), we have on I :

$$V^3 = \frac{f'_1 |f_2| F}{c_2 f_1 f'_2} + \frac{d'}{d} - d.$$

From

$$(V^2)' + dV^2 = -2\frac{f'_1 f'_2}{f_1},$$

we deduce that $\frac{f'_1}{f_1}$ depends on x^3 , but since it depends only on x^2 , it must be a constant.

Denoting $\frac{f'_1}{f_1} =: c_4$, we get $f_1(x^2) = c_5 e^{c_4 x^2}$ with $c_4 \in \mathbb{R}$, $c_5 \in \mathbb{R} \setminus \{0\}$, and also that F depends only on x^3 (on I). We obtain

$$\begin{aligned} F &= -c_2 c_4 f_2 |f_2| + c_6 \quad (c_6 \in \mathbb{R}), \\ V^2 &= -c_4 f_2 + \frac{c_6}{c_2 |f_2|}, \\ V^3 &= -c_4^2 f_2^2 + \frac{c_4 c_6}{c_2} \frac{f_2}{|f_2|} + \left(\frac{f'_2}{f_2}\right)' \frac{f_2}{f'_2} - \frac{f'_2}{f_2}, \\ \lambda &= \frac{c_4 c_6}{c_2} \frac{f_2}{|f_2|}, \\ \mu &= - \left[\left(\frac{f'_2}{f_2}\right)' - \left(\frac{f'_2}{f_2}\right)^2 \right] - \frac{c_4 c_6}{c_2} \frac{f_2}{|f_2|} - (V^3)'. \end{aligned}$$

Let now $V^1 = c_1 \in \mathbb{R} \setminus \{0\}$, which implies that f_1 is constant. System (24) becomes

$$\begin{cases} V^1 = c_1 \\ (V^2)' = -dV^2 \\ (V^3)' = -(d' - d^2) - \mu, \\ dV^3 = d' - d^2 \\ \lambda = 0 \end{cases}$$

with $c_1 \in \mathbb{R} \setminus \{0\}$. From the second equation, it follows that either $V^2 = 0$ or $V^2 \neq 0$ everywhere, in which case $\frac{(V^2)'}{V^2} = -\frac{f_2'}{f_2}$, i.e., $V^2 = \frac{c_2}{f_2}$ with $c_2 \in \mathbb{R} \setminus \{0\}$. Differentiating the second-last equation, multiplying the result with d and substituting in it the expressions of dV^3 and $(V^3)'$, we get

$$d^2\mu = (d')^2 + d^4 - dd'' \quad \square$$

EXAMPLE 3.22. For $f_1(x^2) = e^{x^2}$ and $f_2(x^3) = e^{x^3}$, the vector field

$$V = (1 - e^{2x^3}) \frac{\partial}{\partial x^2} - e^{2x^3} \frac{\partial}{\partial x^3}$$

is a potential vector field of the almost η -Ricci soliton

$$\left(\mathbb{R}^3, \tilde{g}, \lambda = 1, \mu(x^3) = 2e^{2x^3} \right)$$

for $\eta = dx^3$.

3.3 Almost η -Ricci solitons with $\eta = dx^3$ and $V = \frac{\partial}{\partial x^3}$

If $V^1 = V^2 = \eta^1 = \eta^2 = 0$ and $V^3 = \eta^3 = 1$, then the system (2) becomes

$$\begin{cases} -b + \text{Ric}(E_1, E_1) + \lambda = 0 \\ -d + \text{Ric}(E_2, E_2) + \lambda = 0 \\ \text{Ric}(E_3, E_3) + \lambda + \mu = 0 \\ \text{Ric}(E_1, E_3) = 0 \\ \text{Ric}(E_2, E_3) = 0 \end{cases} \quad (25)$$

THEOREM 3.23. Let $V = \frac{\partial}{\partial x^3}$ and $\eta = dx^3$. If $f_i = f_i(x^3)$ for $i \in \{1, 2\}$, then $(\mathbb{R}^3, \tilde{g}, V, \lambda, \mu)$ is an almost η -Ricci soliton if and only if

$$\lambda = \frac{f_1'}{f_1} \left(\frac{f_2'}{f_2} + 1 \right) - \frac{f_1''}{f_1} + 2 \left(\frac{f_1'}{f_1} \right)^2, \quad \mu = -\frac{f_1'}{f_1} \left(\frac{f_2'}{f_2} + 1 \right) - \frac{f_2''}{f_2} + 2 \left(\frac{f_2'}{f_2} \right)^2,$$

and f_1 and f_2 satisfy

$$\frac{f_1'' - f_1'}{f_1} - 2 \left(\frac{f_1'}{f_1} \right)^2 = \frac{f_2'' - f_2'}{f_2} - 2 \left(\frac{f_2'}{f_2} \right)^2. \quad (26)$$

PROOF. In this case, we have $a = 0$, $b = \frac{f_1'}{f_1}$, $c = 0$, $d = \frac{f_2'}{f_2}$, and (25) becomes

$$\begin{cases} -b + \text{Ric}(E_1, E_1) + \lambda = 0 \\ -d + \text{Ric}(E_2, E_2) + \lambda = 0 \\ \text{Ric}(E_3, E_3) + \lambda + \mu = 0 \end{cases} \quad (27)$$

with:

$$\begin{aligned} \text{Ric}(E_1, E_1) &= \frac{f_1''}{f_1} - 2 \left(\frac{f_1'}{f_1} \right)^2 - \frac{f_1' f_2'}{f_1 f_2}, \\ \text{Ric}(E_2, E_2) &= \frac{f_2''}{f_2} - 2 \left(\frac{f_2'}{f_2} \right)^2 - \frac{f_1' f_2'}{f_1 f_2}, \\ \text{Ric}(E_3, E_3) &= \frac{f_1''}{f_1} - 2 \left(\frac{f_1'}{f_1} \right)^2 + \frac{f_2''}{f_2} - 2 \left(\frac{f_2'}{f_2} \right)^2. \end{aligned}$$

The two functions f_1 and f_2 must satisfy (26) due to the first two equations of (27), and we obtain the expressions of λ and μ from the first and the last equation of the same system. \square

EXAMPLE 3.24. For $f_i(x^3) = k_i e^{kx^3}$, $i \in \{1, 2\}$, $k, k_1, k_2 \in \mathbb{R} \setminus \{0\}$, $V = \frac{\partial}{\partial x^3}$, $\eta = dx^3$, $(\mathbb{R}^3, \tilde{g}, V, \lambda, \mu)$ is an η -Ricci soliton if and only if

$$\lambda = 2k^2 + k, \quad \mu = -k.$$

COROLLARY 3.25. Under the hypotheses of Theorem 3.32, if $f_1 = f_2 =: f(x^3)$, then $(\mathbb{R}^3, \tilde{g}, V)$ is an almost η -Ricci soliton with λ and μ as scalar functions if and only if they are given by

$$\lambda = 3 \left(\frac{f'}{f} \right)^2 + \frac{f' - f''}{f}, \quad \mu = \left(\frac{f'}{f} \right)^2 - \frac{f' + f''}{f}.$$

PROOF. It follows immediately from Theorem 3.32. \square

THEOREM 3.26. Let $V = \frac{\partial}{\partial x^3}$ and $\eta = dx^3$. If $f_1 = f_1(x^1)$, $f_2 = f_2(x^3)$, then $(\mathbb{R}^2 \times I, \tilde{g}|_{\mathbb{R}^2 \times I}, V, \lambda, \mu)$ is an almost η -Ricci soliton if and only if

$$f_2 : \mathbb{R}^2 \times I \rightarrow \mathbb{R}, \quad f_2(x^3) = c_0 e^{-x^3}, \quad \lambda = 0, \quad \mu = 1$$

for $c_0 \in \mathbb{R} \setminus \{0\}$, or

$$f_2 : \mathbb{R}^2 \times I \rightarrow \mathbb{R}, \quad f_2(x^3) = \frac{c_1}{c_2 e^{x^3} + 1}, \quad \lambda = 0, \quad \mu(x^3) = \frac{c_2 e^{x^3}}{c_2 e^{x^3} + 1}$$

for $c_1, c_2 \in \mathbb{R} \setminus \{0\}$ such that $e^{x^3} \neq -\frac{1}{c_2}$ for all $x^3 \in I$.

In particular, for $I = \mathbb{R}$, we additionally have the condition $c_2 > 0$.

PROOF. In this case, we have $a = b = c = 0$, $d = \frac{f_2'}{f_2}$, and (25) becomes

$$\begin{cases} \lambda = 0 \\ -d + d' - d^2 = 0. \\ d' - d^2 + \mu = 0 \end{cases}$$

If $d = 0$, then f_2 is constant and $\lambda = \mu = 0$ (which contradicts $\mu \neq 0$). If $d = -1$, then $\ln |f_2(x^3)| = -x^3 + k_0$ with $k_0 \in \mathbb{R}$, and we get $f_2(x^3) = \pm e^{-x^3+k_0}$ and $\mu = 1$. If $d \neq -1$ and $d \neq 0$ (hence, d is nowhere -1 neither 0), from the second equation of the above system we get

$$1 = \frac{d'}{d^2 + d} = \frac{d'}{d} - \frac{d'}{d+1},$$

which, by integration, gives

$$\ln \left(\left| \frac{d(x^3)}{d(x^3) + 1} \right| \right) = x^3 + k_1$$

with $k_1 \in \mathbb{R}$, from which

$$(1 - k_2 e^{x^3})d(x^3) = k_2 e^{x^3}$$

with $k_2 \in \mathbb{R} \setminus \{0\}$ and for all x^3 such that $e^{x^3} \neq \frac{1}{k_2}$. This infers

$$\frac{f_2'(x^3)}{f_2(x^3)} = d(x^3) = \frac{k_2 e^{x^3}}{1 - k_2 e^{x^3}},$$

and, by integrating, we get

$$f_2(x^3) = \frac{k_3}{1 - k_2 e^{x^3}}$$

with $k_3 \in \mathbb{R} \setminus \{0\}$ and

$$\mu(x^3) = d^2(x^3) - d'(x^3) = -d(x^3) = \frac{-k_2 e^{x^3}}{1 - k_2 e^{x^3}}. \quad \square$$

THEOREM 3.27. *Let $V = \frac{\partial}{\partial x^3}$ and $\eta = dx^3$. If $f_i = f_i(x^2)$ for $i \in \{1, 2\}$, then $(\mathbb{R}^3, \tilde{g}, V, \lambda, \mu)$ is an almost η -Ricci soliton if and only if*

$$\lambda = \frac{f_2}{f_1} \left[2 \frac{f_2}{f_1} (f_1')^2 - f_1' f_2' - f_1'' f_2 \right], \quad \mu = -\frac{f_2}{f_1} \left[2 \frac{f_2}{f_1} (f_1')^2 - f_1' f_2' - f_1'' f_2 \right].$$

PROOF. In this case, we have $a = f_2 \frac{f_1'}{f_1}$, $b = c = d = 0$, and (25) becomes

$$\begin{cases} \text{Ric}(E_1, E_1) + \lambda = 0 \\ \text{Ric}(E_2, E_2) + \lambda = 0 \\ \lambda + \mu = 0 \end{cases}$$

with:

$$\text{Ric}(E_1, E_1) = \text{Ric}(E_2, E_2) = f_2 f_2' \frac{f_1'}{f_1} + f_2^2 \frac{f_1''}{f_1} - 2 f_2^2 \left(\frac{f_1'}{f_1} \right)^2. \quad \square$$

EXAMPLE 3.28. For $f_1(x^2) = k_1 e^{-x^2}$, $f_2(x^2) = k_2 e^{x^2}$, $k_1, k_2 \in \mathbb{R} \setminus \{0\}$, $V = \frac{\partial}{\partial x^3}$, $\eta = dx^3$, $(\mathbb{R}^3, \tilde{g}, V, \lambda, \mu)$ is an almost η -Ricci soliton if and only if

$$\lambda(x^2) = 2k_2^2 e^{2x^2}, \quad \mu(x^2) = -2k_2^2 e^{2x^2}.$$

COROLLARY 3.29. *Under the hypotheses of Theorem 3.27, if $f_1 = f_2 =: f(x^2)$, then $(\mathbb{R}^3, \tilde{g}, V)$ is an almost η -Ricci soliton with λ and μ as scalar functions if and only if they are given by*

$$\lambda = (f')^2 - f'' f, \quad \mu = -(f')^2 + f'' f.$$

PROOF. It follows immediately from Theorem 3.27. \square

THEOREM 3.30. *Let $V = \frac{\partial}{\partial x^3}$ and $\eta = dx^3$. If $f_1 = f_1(x^2)$ and $f_2 = f_2(x^1)$, then $(\mathbb{R}^3, \tilde{g}, V, \lambda, \mu)$ is an almost η -Ricci soliton if and only if*

$$\lambda = 2 \left[\frac{f_2^2}{f_1^2} (f_1')^2 + \frac{f_1^2}{f_2^2} (f_2')^2 \right] - f_1^2 \frac{f_2''}{f_2} - f_2^2 \frac{f_1''}{f_1}, \quad \mu = -2 \left[\frac{f_2^2}{f_1^2} (f_1')^2 + \frac{f_1^2}{f_2^2} (f_2')^2 \right] + f_1^2 \frac{f_2''}{f_2} + f_2^2 \frac{f_1''}{f_1}.$$

PROOF. In this case, we have $a = f_2 \frac{f_1'}{f_1}$, $b = 0$, $c = f_1 \frac{f_2'}{f_2}$, $d = 0$, and (25) becomes

$$\begin{cases} \text{Ric}(E_1, E_1) + \lambda = 0 \\ \text{Ric}(E_2, E_2) + \lambda = 0 \\ \lambda + \mu = 0 \end{cases}$$

with:

$$\text{Ric}(E_1, E_1) = \text{Ric}(E_2, E_2) = f_2^2 \frac{f_1''}{f_1} + f_1^2 \frac{f_2''}{f_2} - 2 \left[f_2^2 \left(\frac{f_1'}{f_1} \right)^2 + f_1^2 \left(\frac{f_2'}{f_2} \right)^2 \right]. \quad \square$$

EXAMPLE 3.31. For $f_1(x^2) = e^{x^2}$, $f_2(x^1) = e^{x^1}$, $V = \frac{\partial}{\partial x^3}$, $\eta = dx^3$, $(\mathbb{R}^3, \tilde{g}, V, \lambda, \mu)$ is an almost η -Ricci soliton if and only if

$$\lambda(x^1, x^2) = e^{2x^1} + e^{2x^2}, \quad \mu(x^1, x^2) = -e^{2x^1} - e^{2x^2}.$$

THEOREM 3.32. Let $V = \frac{\partial}{\partial x^3}$, $\eta = dx^3$, and let $I \subset \mathbb{R}$ be an open interval. If $f_1 = f_1(x^2)$ for $x^2 \in \mathbb{R}$, $f_2 = f_2(x^3)$ for $x^3 \in I$, then $(\mathbb{R}^2 \times I, \tilde{g}, V, \lambda, \mu)$ is an almost η -Ricci soliton if and only if one of the following cases is satisfied:

(i)

$$\lambda = -f_2^2 \left[\left(\frac{f_1'}{f_1} \right)' - \left(\frac{f_1'}{f_1} \right)^2 \right], \quad \mu = f_2^2 \left[\left(\frac{f_1'}{f_1} \right)' - \left(\frac{f_1'}{f_1} \right)^2 \right]$$

for f_2 constant;

(ii)

$$\lambda = 0, \quad \mu = -\frac{f_2'}{f_2}$$

for f_1 constant and $f_2(x^3) = \frac{c_1}{c_2 e^{x^3} + 1}$ with $c_1, c_2 \in \mathbb{R} \setminus \{0\}$ and $e^{x^3} \neq -\frac{1}{c_2}$ on I ;

(iii)

$$\lambda = 0, \quad \mu = 1$$

for f_1 constant and $f_2(x^3) = c_0 e^{-x^3}$ with $c_0 \in \mathbb{R} \setminus \{0\}$.

PROOF. In this case, we have $a = f_2 \frac{f_1'}{f_1}$, $b = c = 0$, $d = \frac{f_2'}{f_2}$, and (25) becomes

$$\begin{cases} \text{Ric}(E_1, E_1) + \lambda = 0 \\ -d + \text{Ric}(E_2, E_2) + \lambda = 0 \\ \text{Ric}(E_3, E_3) + \lambda + \mu = 0 \\ \text{Ric}(E_2, E_3) = 0 \end{cases} \quad (28)$$

with:

$$\begin{aligned} \text{Ric}(E_1, E_1) &= f_2^2 \left[\left(\frac{f_1'}{f_1} \right)' - \left(\frac{f_1'}{f_1} \right)^2 \right], \\ \text{Ric}(E_2, E_2) &= f_2^2 \left[\left(\frac{f_1'}{f_1} \right)' - \left(\frac{f_1'}{f_1} \right)^2 \right] + d' - d^2, \\ \text{Ric}(E_3, E_3) &= d' - d^2, \\ \text{Ric}(E_2, E_3) &= \frac{f_1' f_2'}{f_1}. \end{aligned}$$

f_2 must satisfy $d' = d(d + 1)$ due to the first two equations of (28), and we obtain the cases:

- (i) f_2 constant;
- (ii) f_1 constant and $f_2(x^3) = \frac{c_1}{c_2 e^{x^3} + 1}$ on I with $c_1, c_2 \in \mathbb{R} \setminus \{0\}$ and $e^{x^3} \neq -\frac{1}{c_2}$ on I ;
- (iii) f_1 constant and $f_2(x^3) = c_0 e^{-x^3}$ with $c_0 \in \mathbb{R} \setminus \{0\}$. □

EXAMPLE 3.33. For $f_1 = 1$, $f_2(x^3) = \frac{1}{e^{x^3} + 1}$, $V = \frac{\partial}{\partial x^3}$, $\eta = dx^3$, $(\mathbb{R}^3, \tilde{g}, V, \lambda, \mu)$ is an η -Ricci soliton if and only if

$$\lambda = 0, \quad \mu(x^3) = \frac{e^{x^3}}{e^{x^3} + 1}.$$

References

- [1] Atashpeykara, P., Haji-Badali, A. The algebraic Ricci solitons of Lie groups $\mathbb{H}^2 \times \mathbb{R}$ and Sol_3 . *Journal of Finsler Geometry and its Applications* **2020**, 1(2), 105–114. <https://doi.org/10.22098/jfga.2020.1244>
- [2] Belarbi, L. Ricci solitons of the $\mathbb{H}^2 \times \mathbb{R}$ Lie group. *Electronic Research Archive* **2020**, 28(1), 157–163. <https://doi.org/10.3934/era.2020010>
- [3] Belarbi, L. Ricci solitons of the Sol_3 Lie group. *Authorea*. July 26, 2020. <https://doi.org/10.22541/au.159576131.13724173>
- [4] Blaga, A.M. Almost η -Ricci solitons in $(LCS)_n$ -manifolds. *Bull. Belg. Math. Soc. Simon Stevin* **2018**, 25(5), 641–653. <https://doi.org/10.36045/bbms/1547780426>

- [5] Blaga, A.M. η -Ricci solitons on para-Kenmotsu manifolds. *Balkan J. Geom. Appl.* **2015**, *20(1)*, 1–13.
- [6] Blaga, A.M. Solutions of some types of soliton equations in \mathbb{R}^3 . *Filomat* **2019**, *33(4)*, 1159–1162. <https://doi.org/10.2298/FIL1904159B>
- [7] Cao, H.-D., Cui, X. Curvature estimates for four-dimensional gradient steady Ricci solitons. *J. Geom. Anal.* **2020**, *30(1)*, 511–525. <https://doi.org/10.1007/s12220-019-00152-z>
- [8] Cao, H.-D., Liu, T. Curvature estimates for 4-dimensional complete gradient expanding Ricci solitons. *Journal für die reine und angewandte Mathematik* **2022**, *790*, 115–135. <https://doi.org/10.1515/crelle-2022-0039>
- [9] Cao, H.-D., Zhu, M. Linear stability of compact shrinking Ricci solitons. *Math. Ann.* **2024**, 1–17. <https://doi.org/10.1007/s00208-024-02824-w>
- [10] Cho, J.T., Kimura, M. Ricci solitons and real hypersurfaces in a complex space form. *Tohoku Math. J.* **2019**, *61(2)*, 205–212. <https://doi.org/10.2748/tmj/1245849443>
- [11] Chow, B., Lu, P., Ni, L. *Hamilton's Ricci Flow*, Graduate Studies in Mathematics 77, AMS, Providence, RI, USA, bf 2006.
- [12] Hamilton, R.S. The Ricci flow on surfaces, Math. and general relativity (Santa Cruz, CA, 1986). *Contemp. Math.* **1988**, *71*, AMS, 237–262. <https://doi.org/10.1090/conm/071/954419>
- [13] Pigola, S., Rigoli, M., Rimoldi, M., Setti, A.G. Ricci almost solitons. *Ann. Scuola Norm. Sup. Pisa Cl. Sci.* **2011**, *10(4)*, 757–799.

Department of Mathematics

West University of Timișoara

Faculty of Mathematics and Computer Science

Bld. V. Pârvan no. 4, 300223, Timișoara, Romania

adarablaga@yahoo.com