THE FORM-TYPE CALABI-YAU EQUATION ON A CLASS OF COMPLEX MANIFOLDS

LIDING HUANG

ABSTRACT. In this paper, we study the form type Calabi-Yau equation. We define the astheno-Ricci curvature and prove that there exists a solution for the form type Calabi-Yau equation if the astheno-Ricci curvature is non-positive.

1. INTRODUCTION

Let M be a compact Hermitian manifolds of complex dimensiona n with a Hermitian metric α . We use

(1.1)
$$H^{1,1}_{BC}(M,\mathbb{R}) = \frac{\{\text{d-closed real } (1,1)\text{-form}\}}{\{\sqrt{-1}\partial\overline{\partial}\psi : \psi \in C^{\infty}(M,\mathbb{R})\}}$$

to denotes the Bott-Chern cohomology group and $\operatorname{Ric}(\alpha)$ is the Chern Ricci form of α . The first Bott-Chern class is defined by $c_1^{BC} = [\operatorname{Ric}(\alpha)] \in$ $H_{BC}^{1,1}(M,\mathbb{R})$. Yau [38] proved Calabi conjecture [9] on Kähler manifolds that have many applications in geometry and physics. We define a Calabi-Yau manifold to be a compact complex manifold M with $c_1^{BC} = 0$ in $H_{BC}^{1,1}(M,\mathbb{R})$ [34]. Motivated by mathematics and mathematical physics, the non-Kähler Calabi-Yau manifolds has been extensively researched [33, 34, 39]. It is also related to Reid fantasy [29]. An important problem is to find an analog of Ricci-flat Kähler metrics on non-Kähler Calabi-Yau manifolds [34, 39].

Tosatti-Weinkove [36] proved that the existence of solutions for complex Monge-Ampère equations on Hermitian manifolds, which implies a version of Calabi-Yau type theorem for Hermitian metric (pluriclosed metrics). Székelyhidi-Tosatti-Weinkove [32] proved that every compact Calabi-Yau manifold admits a Gauduchon metric whose Chern-Ricci curvature is zero, by solving the Monge-Ampère type equation for (n-1)-plurisubharmonic functions.

A Hermitian metric ω_0 is balanced if $d(\omega_0^{n-1}) = 0$. There are a large class of balanced manifolds [21, 13, 18]. The balanced condition exhibits favorable mathematical properties. A sharp characterization for balanced metrics was gave by [24], and the balanced metrics can be utilized to explore

²⁰²⁰ Mathematics Subject Classification. Primary: 32W20; Secondary: 35J60,53C55, 58J05.

Key words and phrases. The form-type Calabi-Yau equation, balanced metric, the Chern-Ricci curvature, the Bismut Ricci curvature.

birational geometry since the existence of a balanced metric is invariant under birational transformations [1]. Fu-Li-Yau [13] proved that suppose $Y \to \underline{Y} \rightsquigarrow X_t$ be a conifold transition, then X_t is a balanced manifolds for tsmall enough, where Y is a Kähler Calabi-Yau manifolds and \underline{Y} is a singular variety with ODP singularities.

It is natural to seek a balanced metric which is Chern-Ricci flat on Calabi-Yau manifolds with a balanced metric. This problem is also related to the Strominger system. Let Ω be an no-where vanishing holomorphic (n, 0) form. Fu-Wang-Wu [14] want to seek a balanced metric $\hat{\omega}$ satisfying $\|\Omega\|_{\hat{\omega}} = 1$. It is equal to find a balanced metric which is Chern-Ricci flat. Popovici [28] also considered this problem. Fu-Wang-Wu [14] proposed the following approach to study this problem. Let ω_0 be a balanced metric, then it is suffice to find a metric $\hat{\omega}^{n-1} = \omega_0^{n-1} + \sqrt{-1}\partial\overline{\partial}(\varphi \alpha^{n-2})$ satisfying the equation

(1.2)
$$\hat{\omega}^n = e^{h+b} \alpha^n.$$

where α is a Hermitian metric. This equation called a "form-type Calabi-Yau equation" in [14]. It can be reduced to the Monge-Amère type equation for (n-1)-plurisubharmonic functions. If there exists a Hermitian metric has non-negative orthogonal bisectional curvature, Fu-Wang-Wu [15] solved this equation on Kähler manifolds. In Kähler case, this equation was solved by Tosatti-Weinkove [36]. If α is astheno-Kähler, there exists a solution for (1.2) using Székelyhidi-Tosatti-Weinkove's results [32].

For the balanced metric which is Chern-Ricci flat and the equation (1.2), the following two conjectures were stated in [34].

Conjecture 1.1. Let M be a compact complex manifold with $c_1^{BC} = 0$ and with a balanced metric ω_0 . Then there is a balanced metric $\hat{\omega}$ with $[\hat{\omega}^{n-1}] = [\omega_0^{n-1}]$ in $H^{2n-2}(M, R)$ with $Ric(\hat{\omega}) = 0$.

Conjecture 1.2. Let M be a compact complex manifold with a balanced metric ω_0 and a Hermitian metric α . Let h be a smooth function. Then there exists a constant b and a function φ satisfying

(1.3)
$$\begin{cases} \hat{\omega}^n = e^{h+b}\alpha^n;\\ \hat{\omega}^{n-1} = \omega_0^{n-1} + \sqrt{-1}\partial\overline{\partial}(\varphi\alpha^{n-2}) > 0. \end{cases}$$

Jost-Yau [23] defined the astheno-Kähler metric $\sqrt{-1}\partial\overline{\partial}\omega^{n-2} = 0$. Following Jost-Yau's definition, we introduce the following definition:

Definition 1.1. Let α be a Hermitian metric and * be the Hodge-star operator of α . We define the astheno-Ricci curvature of α by

$$*\sqrt{-1}\partial\overline{\partial}(\alpha^{n-2})$$

We say the astheno-Ricci curvature of α is non-positive if

$$*\sqrt{-1}\partial\overline{\partial}(\alpha^{n-2}) \le 0.$$

Remark 1.3. If the astheno-Ricci curvature of α is equal to 0, it is just an astheno-Kähler metric. The astheno-Ricci curvature is related to the Ricci curvature (see Proposition 2.4). Recently, George [17] study the equation (1.3) under the condition $\sqrt{-1}\partial\overline{\partial}\alpha^{n-2} \leq 0$. It is equal to the astheno-Ricci curvature is non-positive.

We confirm the conjecture 1.2 under this condition. More precisely, we have the following theorem

Theorem 1.4. Let M be a compact complex manifold with dimension n, equipped with Hermitian metrics α and ω_0 . Suppose the astheno-Ricci curvature of α is non-positive. Given smooth functions h on M, there exists a small constant A_0 such that for any $0 < A \leq A_0$, there exists a unique pair (φ, b) where $\varphi \in C^{\infty}(M)$ and $b \in \mathbb{R}$ solving

(1.4)
$$\begin{cases} \hat{\omega}^n = e^{h+b}\alpha^n; \\ \hat{\omega}^{n-1} = \omega_0^{n-1} + \sqrt{-1}\partial\overline{\partial}(\varphi\alpha^{n-2}) > 0, \|e^{\varphi}\|_{L^1(M)} = A, \end{cases}$$

where A_0 is a constant depending on $(M, \alpha), \omega_0$ and h. Here $||e^{\varphi}||_{L^1(M)} = \int_M |e^{\varphi}| \alpha^n$. Moreover, we have

(1.5)
$$\sup \varphi \le CA,$$

where C is a uniform constant depending only on $(M, \alpha), \omega_0$ and h.

Using Theorem 1.4, we can prove a Calabi-Yau type theorem. The Bismut connection are researched extensively (see [5, 12, 41] and the references therein). We consider the following Ricci curvature for Bismut connection:

$$\operatorname{Ric}^{B}(\omega) = \operatorname{Ric}(\omega) + dd^{*}\omega.$$

When the metirc is balanced, the Bismut connection and the Chern connection agree. Tosatti-Weinkove [37] proved a Calabi-Yau theorem on Ric^B in balanced class on Kähler manifolds. We have the following corollary:

Corollary 1.5. Let M be a compact complex manifold with a balanced metric ω_0 . Suppose there exists a Hermitian metric that the astheno-Ricci curvature is non-positive. Then for any $\Phi \in c_1^{BC} \in H^{1,1}_{BC}(M,\mathbb{R})$, there is a balanced metric $\hat{\omega}$ with $[\hat{\omega}^{n-1}] = [\omega_0^{n-1}]$ in $H^{2n-2}(M, \mathbb{R})$ with

$$Ric(\hat{\omega}) = Ric^B(\hat{\omega}) = \Phi.$$

In [11], Fernandez-Jordan-Streets studied the Busmut Hermitian-Einstein metric (a kind of Ricci curvature of Bismut connection) and proved that it is equal to a Hermitian-Einstein metric on exact holomorphic Courant algebroids. They also gave some examples of compact manifolds X with the first Chern class $c_1 = 0 \in H^2(X, Z)$ which do not admit a Bismut Hermitian-Einstein metric. For the Ricci curvature Ric^B , we conclude the following corollary:

Corollary 1.6. Let M be a compact complex manifold with $c_1^{BC} = 0$. Suppose there exists a balanced metric ω_0 and a Hermitian metric that the astheno-Ricci curvature is non-positive. Then there is a balanced metric $\hat{\omega}$ with $[\hat{\omega}^{n-1}] = [\omega_0^{n-1}]$ in $H^{2n-2}(M, R)$ with $Ric(\hat{\omega}) = Ric^B(\hat{\omega}) = 0$.

Since the equation involves the zero order terms, it is very difficult to solve this equation. To overcome the difficulty, we consider the condition that the astheno-Ricci curvature of the background metric is non-positive. We prove a new C^0 estimates for the equation (1.3). The normalized condition $\int_M e^{\varphi} \alpha^n = A$ ensures that the positive part φ_+ is integrable. Using Moser iteration, we can prove the solutions are "nearly non-positive". To be more specific, assuming that $\varphi \leq 1$, motivated by Fu-Yau [16], for a small constant A, we can prove

$$\varphi \le CA^{\frac{1}{n+1}}.$$

Then we obtain the above bound for the solutions via the continuity method (see Proposition 3.1 and Proposition 5.1).

The estimates plays an important role in the proof of the second order estimates. The assumption of the astheno-Ricci curvature can be used in the proof of C^0 -estimates and the openness. However, for the second order estimates, there are new difficulties come from $\xi(\varphi)$ in the quantity Q for the terms φB . We overcome these difficulties by our supremum estimates (see Lemma 4.10). Adapting Székelyhidi's argument [31] based on Blocki's work [3, 4], we established the infimum estimates. We expect that the analogous argument can be extended to study the equation (1.3) on general Hermitian manifolds.

For the second order estimate, following the idea of [22, 31, 32], we prove the following second order estimates

(1.6)
$$\sup_{M} |\sqrt{-1}\partial\overline{\partial}\varphi|_{\alpha} \le C(1+\sup_{M} |\nabla\varphi|_{\alpha}^{2}).$$

Since there exists a zero order term in the equation, some essential modifications of the arguments in [32] are needed. In this paper, we prove a special C^0 estimates such that we can use the argument in [32] to prove (1.6). Then the second order estimate follows from the blowup argument and Liouville type theorem [31].

Recently, George [17] also proved that there exists a unique solution for (1.3) under the non-positive of astheno-Ricci curvature. Compared to his results, our method is quite different. We assume the normalized condition and use a new method to prove the C^0 estimates. The proof of C^0 -estimates, the openness and the uniqueness of solutions heavily depends on the normalized condition. On the other hand, for the second order estimates, the supremum estimates are used to deal with the bad term which comes form the zero order terms in (2.7).

The organization of paper is as follows. In Section 2, we will introduce some notations and recall some important properties of the equation (1.3). The zero order estimate will be established in Section 3. We will derive the second order estimates and prove the higher order estimates in Section 4, and complete the proof of Theorem 1.4 in Section 5.

Acknowledgements. The author would also like to thank Professor Jixiang Fu, Jianchun Chu and Rirong Yuan for many useful discussions.

2. Preliminaries and notation

Let * be the Hodge star operator with respect to α . Set

$$\begin{split} \tilde{\chi} &= \frac{1}{(n-1)!} * (\omega_0^{n-1}), Z(\partial \varphi) = \frac{2}{(n-1)!} * \operatorname{Re}(\sqrt{-1}\partial \varphi \wedge \bar{\partial}(\alpha^{n-2})), \\ \tilde{B}(\varphi) &= \frac{1}{(n-1)!} * \sqrt{-1}\partial \overline{\partial}(\alpha^{n-2}). \end{split}$$

For convenience, we use Z and \tilde{B} to denote $Z(\partial \varphi)$ and $\tilde{B}(\varphi)$, respectively. Note

$$\frac{1}{(n-1)!} * (\sqrt{-1}\partial\overline{\partial}\varphi \wedge \alpha^{n-2}) = \frac{1}{n-1}(\Delta\varphi\alpha - \sqrt{-1}\partial\overline{\partial}\varphi).$$

Using the argument in [37, 32], the equation can be rewritten as

(2.1)
$$\begin{cases} (\tilde{\chi} + \frac{1}{n-1}((\Delta\varphi)\alpha - \sqrt{-1}\partial\overline{\partial}\varphi) + Z + \varphi\tilde{B})^n = e^{(n-1)(h+b)}\alpha^n, \\ \tilde{\omega} = \tilde{\chi} + \frac{1}{n-1}((\Delta\varphi)\alpha - \sqrt{-1}\partial\overline{\partial}\varphi) + Z + \varphi\tilde{B} > 0. \end{cases}$$

Let T be the linear map given by

(2.2)
$$T(\hat{\omega}) = \operatorname{tr}_{\alpha} \hat{\omega} \alpha - (n-1)\hat{\omega},$$

where $\hat{\omega}$ is a real smooth (1, 1) form. We also have

(2.3)
$$\hat{\omega} = \frac{1}{n-1} ((\operatorname{tr}_{\alpha} T(\hat{\omega}))\alpha - T(\hat{\omega})).$$

Denote

(2.4)
$$\omega = T(\tilde{\omega}), \chi = T(\tilde{\chi}), W = T(Z), B = T(\tilde{B}).$$

Then

(2.5)
$$\omega = \chi + \sqrt{-1}\partial\overline{\partial}\varphi + W(\partial\varphi) + \varphi B.$$

Suppose

$$\begin{aligned} \alpha &= \sqrt{-1}\alpha_{i\bar{j}}dz_i \wedge dz_{\bar{j}}, \, \omega = \sqrt{-1}g_{i\bar{j}}dz_i \wedge dz_{\bar{j}}, \\ \chi &= \sqrt{-1}\chi_{i\bar{j}}dz_i \wedge dz_{\bar{j}}, \, \tilde{\chi} = \sqrt{-1}\tilde{\chi}_{i\bar{j}}dz_i \wedge dz_{\bar{j}}, \\ \tilde{\omega} &= \sqrt{-1}\tilde{g}_{i\bar{j}}dz_i \wedge dz_{\bar{j}}, \, B = \sqrt{-1}B_{i\bar{j}}dz_i \wedge dz_{\bar{j}}, \end{aligned}$$

in local holomorphic coordinate system (z_1, \cdots, z_n) .

In a neighborhood of x_0 , we choose normal coordinates (z_1, \dots, z_n) such that at x_0 , we have $\alpha_{i\bar{j}} = \delta_{ij}$, $g_{i\bar{j}} = \lambda_i \delta_{i\bar{j}}$ and $\lambda_1 \ge \dots \ge \lambda_n$. Assume that $\tilde{\lambda}_1, \dots, \tilde{\lambda}_n$ are the eigenvalue of $\tilde{\omega}$ with respect to α . Then we have

(2.6)
$$\tilde{\lambda}_i = \frac{1}{n-1} \sum_{k \neq i} \lambda_i$$

and the equation can be rewritten as

(2.7)
$$\begin{cases} F(\omega) = f(\lambda_1, \cdots, \lambda_n) = \log \pi_k(\sum_{k \neq i} \lambda_i) = (n-1)(h+b), \\ \omega \in T(\Gamma_n), \end{cases}$$

where Γ_n is the positive orthant.

Define

$$F^{i\overline{j}} = \frac{\partial F}{\partial g_{i\overline{j}}}, \quad F^{i\overline{j},k\overline{l}} = \frac{\partial^2 F}{\partial g_{i\overline{j}}\partial g_{k\overline{l}}}$$

In addition, the equation (2.7) also can be viewed as a function on \tilde{g} . We denote

(2.8)
$$\tilde{F}^{i\bar{i}} = \frac{\partial F}{\partial \tilde{g}_{i\bar{j}}}$$

Set

(2.9)
$$f_k = \frac{\partial f}{\partial \lambda_i} = \frac{1}{n-1} \sum_{k \neq i} \frac{1}{\tilde{\lambda}_i} \text{ and } \tilde{f}_k = \frac{1}{\tilde{\lambda}_k}.$$

Then

(2.10)
$$\tilde{F}^{i\bar{j}} = \tilde{f}_i \delta_{ij} \text{ and } F^{i\bar{i}} = f_i \delta_{ij}$$

2.1. The properties for the form type Calabi-Yau equation. The following properties has been used in [32].

Lemma 2.1. (1)
$$\lambda_1 \leq \cdots \leq \lambda_n$$
.
(2) $\tilde{f}_n \leq \tilde{f}_{n-1} \leq \cdots \leq \tilde{f}_1$.
(3) $f_k \leq \tilde{f}_1 \leq (n-1)f_k, \ k \geq 2$.
(4) $\tilde{f}_k \leq (n-1)f_1, \ k \geq 2$.
(5) $\frac{1}{n(n-1)}\mathcal{F} \leq F^{k\bar{k}}, \ k \geq 2$,
where $\mathcal{F} = \sum_i F^{i\bar{i}}$.

Proof. By the definition of $\tilde{\lambda}_i$, the (1) and (2) follows. It is easy to see $f_k \leq \frac{1}{\tilde{\lambda}_1} = \tilde{f}_1$. Since $\tilde{\lambda}_i > 0$, we have $\tilde{f}_1 \leq (n-1)f_k$ and $\tilde{f}_k \leq (n-1)f_1$, $k \geq 2$. Note $\mathcal{F} = \sum_i \frac{1}{\tilde{\lambda}_i} \leq n \frac{1}{\tilde{\lambda}_1} \leq n(n-1)F^{k\bar{k}}$.

Székelyhidi [31] used the C-subsolution to prove the following properties which is a refinement of Guan [19, Theorem 2.18] and the properties are improved in [6, 40]. For the equation (2.7), it can be proved directly. Using the argument in [32], we prove

Lemma 2.2. There exist κ depending only on χ , n and h, such that

(1) we either have

$$f_k(\lambda)(\chi_{k\bar{k}} - \lambda_k) \ge \kappa \sum_k f_k(\lambda)$$

(2) or $f_k \ge \kappa \sum_i f_i$. In addition, $\sum_k f_k(\lambda) > \kappa$.

Proof. By directly calculation,

$$\sum_{k} f_k \chi_{k\bar{k}} = \sum_{i} \frac{1}{\tilde{\lambda}_i} \tilde{\chi}_{i\bar{i}} > \tau \sum_{i} \frac{1}{\tilde{\lambda}_i} = \tau \sum_{k} f_k(\lambda),$$

for some $\tau > 0$ depending on a lower bound for $\tilde{\chi}$. We also have

$$\sum_{k} \lambda_k f_k(\lambda) = n$$

Note that if λ_1 is large enough, then λ_n is also large which implies that λ_1 is small. Then we have the alternative (1) in Proposition (2.2). If for any λ_i is bounded, then (2) is true. In addition,

$$\sum_{k} f_k(\lambda) = \sum_{i=1}^n \frac{1}{\tilde{\lambda_i}} \ge n(\tilde{\lambda}_1 \cdots \tilde{\lambda}_n)^{-\frac{1}{n}} = ne^{-\frac{(n-1)(h+b)}{n}},$$

so the final claim in Proposition 2.2 also holds.

Assume that locally $Z = 2 \operatorname{Re}(Z_{i\bar{j}}^k \varphi_k)$ and $W = 2 \operatorname{Re}(W_{i\bar{j}}^k \varphi_k)$. We need the following properties about Z. It was proved in [32, p. 208].

Lemma 2.3. Choose normal coordinate system such that $\alpha_{i\bar{j}} = \delta_{ij}$ at x_0 . Then at x_0 , we have

- (1) $Z_{i\bar{j}}$ is independent of φ_i and $\varphi_{\bar{i}}$, i.e., $Z^i_{i\bar{i}}(x_0) = \overline{Z^i_{i\bar{i}}}(x_0) = 0$.
- (2) $\nabla_i Z_{i\bar{i}}(x_0)$ is independent of φ_i .

On the other hand, the linear operator of the equation (1.3) is

(2.11)
$$L(\psi) = F^{i\bar{j}}(\psi_{i\bar{j}} + 2\operatorname{Re}(W^k_{i\bar{j}}\psi_k) + B_{i\bar{j}}\psi)$$
$$= F^{i\bar{j}}(\psi_{i\bar{j}} + 2\operatorname{Re}(W^k_{i\bar{j}}\psi_k)) + \tilde{F}^{i\bar{j}}\tilde{B}_{i\bar{j}}\psi$$

2.2. The astheno-Ricci curvature. In this subsection, we will calculate the astheno-Ricci curvature $*\sqrt{-1}\partial\overline{\partial}(\omega^{n-2})$. Recall the curvature tensor are defined by

$$R_{i\bar{j}k\bar{l}} = -\partial_k \partial_{\bar{l}} \alpha_{i\bar{j}} + \alpha^{p\bar{q}} \partial_{\bar{l}} \alpha_{p\bar{j}} \partial_k \alpha_{i\bar{q}}.$$

Define the following Ricci curvature and scalar curvature

$$\operatorname{Ric}_{i\bar{j}} = \alpha^{k\bar{l}} R_{k\bar{l}i\bar{j}}, \operatorname{Ric}_{i\bar{j}}^{(2)} = \alpha^{k\bar{l}} R_{i\bar{j}k\bar{l}}, \operatorname{Ric}_{i\bar{j}}^{(3)} = \alpha^{k\bar{l}} R_{k\bar{j}i\bar{l}}, \operatorname{Ric}_{i\bar{j}}^{(4)} = \alpha^{k\bar{l}} R_{i\bar{l}k\bar{j}}$$

and

$$R = \alpha^{i\bar{j}} \operatorname{Ric}_{i\bar{j}}, R^{(2)} = \alpha^{i\bar{j}} \operatorname{Ric}_{i\bar{j}}^{(2)}, R^{(3)} = \alpha^{i\bar{j}} \operatorname{Ric}_{i\bar{j}}^{(3)}, R^{(4)} = \alpha^{i\bar{j}} \operatorname{Ric}_{i\bar{j}}^{(4)}.$$

Denote $T = \partial \alpha$ and assume $T = \frac{1}{2}T_{sj\bar{k}}dz_s \wedge dz_j \wedge d\bar{z}_k$ in local holomorphic coordinates system. Then we define $T_s = \alpha^{j\bar{k}}T_{sj\bar{k}}$ and $\tau = T_s dz_s$.

Phong-Picard-Zhang [26] calculated the $*(\sqrt{-1}\partial\overline{\partial}\alpha^{n-2})$ for conformal balanced metric. Now we give a formula for $*(\sqrt{-1}\partial\overline{\partial}\alpha^{n-2})$ for general Hermitian metric. From the following results, we can see that $*(\sqrt{-1}\partial\overline{\partial}\alpha^{n-2})$ is closed related to the Ricci curvature.

Proposition 2.4. (1) If n=3,

$$* (\sqrt{-1}\partial\overline{\partial}(\alpha^{n-2}))_{m\bar{l}}$$

$$= \sqrt{-1}(n-2)!(Ric_{m\bar{l}}^{(2)} - Ric_{m\bar{l}}^{(3)} + Ric_{m\bar{l}} - Ric_{m\bar{l}}^{(4)}$$

$$- \alpha^{j\bar{k}}\alpha^{s\bar{r}}T_{mj\bar{r}}\bar{T}_{\bar{l}\bar{k}s}) + \sqrt{-1}\frac{(n-2)!}{2}(2R - 2R^{(3)} - |T|^2)\alpha_{m\bar{l}}$$

$$\begin{array}{l} (2) \ If \ n \geq 4, \\ &* (\sqrt{-1}\partial\overline{\partial}(\alpha^{n-2}))_{m\bar{l}} \\ = &\sqrt{-1}(n-2)!(Ric_{m\bar{l}}^{(2)} - Ric_{m\bar{l}}^{(3)} + Ric_{m\bar{l}} - Ric_{m\bar{l}}^{(4)} \\ &- \alpha^{j\bar{k}}\alpha^{s\bar{r}}T_{mj\bar{r}}\bar{T}_{\bar{l}\bar{k}s}) + \sqrt{-1}\frac{(n-2)!}{2}(2R-2R^{(3)} - |T|^2)\alpha_{m\bar{l}} \\ &+ \frac{(n-2)!}{2}\alpha^{p\bar{q}}\alpha^{s\bar{r}}(T\wedge\bar{T})_{s\bar{r}q\bar{p}m\bar{l}} + \sqrt{-1}\frac{(n-2)!}{6}(3|T|^2 - 2|\tau|^2)\alpha_{m\bar{l}}. \end{array}$$

Proof. We have the following formula in [26, Lemma 2]

(2.12)

$$(*(\Phi \wedge \alpha^{n-3}))_{k\bar{l}} = \sqrt{-1}(n-3)! \alpha^{s\bar{r}} \Phi_{s\bar{r}k\bar{l}} + \sqrt{-1} \frac{(n-3)!}{2} (\operatorname{tr}\Phi) \alpha_{k\bar{l}}, n \ge 3;$$

$$(*(\Psi \wedge \alpha^{n-4}))_{k\bar{l}} = \frac{\sqrt{-1}(n-4)!}{2} \alpha^{s\bar{r}} \alpha^{p\bar{q}} \Phi_{s\bar{r}p\bar{q}k\bar{l}} + \sqrt{-1} \frac{(n-4)!}{6} (\operatorname{tr}\Phi) \alpha_{k\bar{l}}, n \ge 4,$$

where Φ is a (2,2) form and Ψ is a (3,3) form. Note

$$\sqrt{-1}\partial\overline{\partial}\alpha^{n-2} = (n-2)\sqrt{-1}\partial\overline{\partial}\alpha \wedge \alpha^{n-3} + (n-2)(n-3)\sqrt{-1}\partial\alpha \wedge \bar{\partial}\alpha \wedge \alpha^{n-4}$$

Now we calculate $\alpha^{j\bar{k}}(\sqrt{-1}\partial\overline{\partial}\alpha)_{j\bar{k}m\bar{l}}$, $\operatorname{tr}(\sqrt{-1}\partial\overline{\partial}\alpha)$, $\alpha^{p\bar{q}}\alpha^{s\bar{r}}(T\wedge\bar{T})_{s\bar{r}q\bar{p}j\bar{k}}$ and $\operatorname{tr}T \wedge T$ (see [26]): $\alpha^{j\bar{k}}(\sqrt{-1}\partial\overline{\partial}\alpha)_{j\bar{k}m\bar{l}} = \operatorname{Ric}_{m\bar{l}}^{(2)} - \operatorname{Ric}_{m\bar{l}}^{(3)} + \operatorname{Ric}_{m\bar{l}} - \operatorname{Ric}_{m\bar{l}}^{(4)} - \alpha^{j\bar{k}}\alpha^{s\bar{r}}T_{mj\bar{r}}\bar{T}_{\bar{l}\bar{k}s},$ $\alpha^{p\bar{q}}\alpha^{s\bar{r}}(T\wedge\bar{T})_{s\bar{r}q\bar{p}j\bar{k}} = \alpha^{p\bar{q}}\alpha^{s\bar{r}}(2T_{sj\bar{p}}\bar{T}_{\bar{r}\bar{k}q} + T_{qs\bar{k}}T_{\bar{p}\bar{r}j}) - 2\alpha^{s\bar{r}}(T_{js\bar{k}}T_{\bar{r}} + T_s\bar{T}_{\bar{k}\bar{r}j})$

$$-2T_j\bar{T}_{\bar{k}}$$

and

$$\mathrm{tr}(\sqrt{-1}\partial\overline{\partial}\alpha)=2R-2R^{(3)}-|T|^2,\ \ \mathrm{tr}T\wedge T=(3|T|^2-2|\tau|^2).$$
 Using (2.12), we have

$$(*(\sqrt{-1}\partial\overline{\partial}\alpha \wedge \alpha^{n-2}))_{m\bar{l}} = \sqrt{-1}(n-3)!(\operatorname{Ric}_{m\bar{l}}^{(2)} - \operatorname{Ric}_{m\bar{l}}^{(3)} + \operatorname{Ric}_{m\bar{l}} - \operatorname{Ric}_{m\bar{l}}^{(4)} - \alpha^{j\bar{k}}\alpha^{s\bar{r}}T_{mj\bar{r}}\bar{T}_{\bar{l}\bar{k}s}) + \sqrt{-1}\frac{(n-3)!}{2}(2R-2R^{(3)} - |T|^2)\alpha_{m\bar{l}}.$$

and

$$(*\sqrt{-1}\partial\alpha \wedge \bar{\partial}\alpha \wedge \alpha^{n-4})_{m\bar{l}} = \frac{(n-4)!}{2} \alpha^{p\bar{q}} \alpha^{s\bar{r}} (T \wedge \bar{T})_{s\bar{r}q\bar{p}m\bar{l}} + \sqrt{-1} \frac{(n-4)!}{6} (3|T|^2 - 2|\tau|^2) \alpha_{m\bar{l}}.$$

Then if n=3,

$$(*(\sqrt{-1}\partial\overline{\partial}(\alpha^{n-2})))_{m\bar{l}}$$

= $\sqrt{-1}(n-2)!(\operatorname{Ric}_{m\bar{l}}^{(2)} - \operatorname{Ric}_{m\bar{l}}^{(3)} + \operatorname{Ric}_{m\bar{l}} - \operatorname{Ric}_{m\bar{l}}^{(4)}$
- $\alpha^{j\bar{k}}\alpha^{s\bar{r}}T_{mj\bar{r}}\bar{T}_{\bar{l}\bar{k}s}) + \sqrt{-1}\frac{(n-2)!}{2}(2R-2R^{(3)}-|T|^2)\alpha_{m\bar{l}}$

and if
$$n = 4$$

 $(*(\sqrt{-1}\partial\overline{\partial}(\alpha^{n-2})))_{m\bar{l}}$
 $=\sqrt{-1}(n-2)!(\operatorname{Ric}_{m\bar{l}}^{(2)} - \operatorname{Ric}_{m\bar{l}}^{(3)} + \operatorname{Ric}_{m\bar{l}} - \operatorname{Ric}_{m\bar{l}}^{(4)}$
 $-\alpha^{j\bar{k}}\alpha^{s\bar{r}}T_{mj\bar{r}}\bar{T}_{\bar{l}\bar{k}s}) + \sqrt{-1}\frac{(n-2)!}{2}(2R - 2R^{(3)} - |T|^2)\alpha_{m\bar{l}}$
 $+\frac{(n-2)!}{2}\alpha^{p\bar{q}}\alpha^{s\bar{r}}(T \wedge \bar{T})_{s\bar{r}q\bar{p}m\bar{l}} + \sqrt{-1}\frac{(n-2)!}{6}(3|T|^2 - 2|\tau|^2)\alpha_{m\bar{l}}.$

3. The C^0 estimates

In this section, we will prove the C^0 estimates.

3.1. The upper bound for φ . In this subsection, we will prove the supremum estimates. Fu-Yau [16] proved there exists a large positive solution for Fu-Yau equation in complex surface. In higher dimensions, it was proved in [7, 8, 25, 27]. For the form type Calabi-yau equation, we prove that the solution are "nearly non-positive" if the astheno-Ricci curvature is non-positive.

Proposition 3.1. Suppose that

$$\|e^{\varphi}\|_{L^1(M,\alpha)} = A$$

and

$$\sup_{M} \varphi \le 1,$$

then we have

(3.1)
$$\sup_{M} \varphi \le M_0 A^{\frac{1}{n+1}}$$

where M_0 is a uniform constant depending on α, ω_0 and n.

Proof. Set $\varphi_+ = \max\{\varphi, 0\}$. It is easy to see that

(3.2)
$$\int_{M} \varphi_{+} \alpha^{n} \leq \int_{M} e^{\varphi} \alpha^{n} = A.$$

By the elliptic condition, we have

$$\int_{M} \varphi_{+}^{k} \left(\omega_{0}^{n-1} + \sqrt{-1} \partial \overline{\partial} (\varphi \alpha^{n-2}) \right) \wedge \alpha \geq 0.$$

Using integration by parts and the Cauchy-Schwarz inequality,

$$\begin{split} &\int_{M} \varphi_{+}^{k} \sqrt{-1} \partial \overline{\partial} (\varphi \alpha^{n-2}) \wedge \alpha \\ = &- \frac{4k}{n(k+1)^{2}} \int_{M} |\partial \varphi_{+}^{k+1}|_{\alpha}^{2} \alpha^{n} - k \int_{M} \varphi_{+}^{k} \sqrt{-1} \partial (\varphi) \wedge \overline{\partial} (\alpha^{n-2}) \wedge \alpha \\ &- \int_{M} \varphi_{+}^{k} \sqrt{-1} \partial \alpha \wedge \overline{\partial} \varphi \wedge \alpha^{n-2} - \int_{M} \varphi_{+}^{k+1} \sqrt{-1} \partial \alpha \wedge \overline{\partial} (\alpha^{n-2}) \\ \leq &- \frac{2k}{n(k+1)^{2}} \int_{M} |\partial \varphi_{+}^{k+1}|_{\alpha}^{2} \alpha^{n} + Ck \int_{M} \varphi_{+}^{k+1} \alpha^{n} \\ \leq &- \frac{2k}{n(k+1)^{2}} \int_{M} |\partial \varphi_{+}^{k+1}|_{\alpha}^{2} \alpha^{n} + Ck \int_{M} \varphi_{+}^{k} \alpha^{n}, \end{split}$$

where in the last inequality we used $\varphi \leq 1$. Then

$$\frac{2k}{n(k+1)^2} \int_M |\partial \varphi_+^{\frac{k+1}{2}}|_\alpha^2 \alpha^n \le Ck \int_M \varphi_+^k \alpha^n.$$

It follows, by Sobolev inequality,

$$\|\varphi_{+}^{\frac{(k+1)}{2}}\|_{L^{\frac{2n}{n-1}}} \leq \|\partial\varphi_{+}^{\frac{k+1}{2}}\|_{L^{2}} \leq (C(k+1)^{2})^{\frac{1}{2}} (\int_{M} \varphi_{+}^{k} \alpha^{n})^{\frac{1}{2}}.$$

Therefore,

$$|\varphi_+\|_{L^{\frac{n(k+1)}{n-1}}} \le (C(k+1)^2)^{\frac{1}{k+1}} \|\varphi_+\|_{L^k}^{\frac{k}{k+1}}, k \ge 1.$$

Choose $k_1 = 1, k_i = \frac{n}{n-1}(k_{i-1}+1)$. It is easy to see that

(3.3)
$$k_i = \left(\frac{n}{n-1}\right)^{i-1} (k_1 + n) - n.$$

Then we have (3.4)

$$\begin{split} \|\varphi_{+}\|_{L^{k_{i+1}}} &\leq \left(\left(C(k_{i}+1)\right)^{\frac{1}{k_{i}+1}} \|\varphi_{+}\|_{L^{k_{i}}}^{\frac{k_{i}}{k_{i}+1}} \\ &\leq \left(\left(C(k_{i}+1)\right)^{\frac{1}{k_{i}+1}} \left(\left(C(k_{i-1}+1)\right)^{\frac{k_{i}}{k_{i}+1}\frac{1}{k_{i-1}+1}} \|\varphi_{+}\|_{L^{k_{i-2}}}^{\frac{k_{i}}{k_{i}+1}\frac{k_{i-1}}{k_{i-1}+1}} \\ &\leq \left(\left(C(k_{i}+1)\right)^{\frac{1}{k_{i}+1}} \left(\left(C(k_{i-1}+1)\right)^{\frac{k_{i}}{k_{i}+1}\frac{1}{k_{i-1}+1}} \cdots \left(C(k_{1}+1)\right)^{\frac{k_{i}}{k_{i}+1}\frac{k_{i-1}}{k_{i-1}+1}\cdots\frac{1}{k_{1}+1}} \\ &\|\varphi_{+}\|_{L^{k_{1}}}^{\frac{k_{i}}{k_{i+1}}\frac{k_{i-1}}{k_{i-1}+1}\cdots\frac{k_{1}}{k_{1}+1}} \\ &= C^{\frac{1}{k_{i+1}}\left(\frac{n}{n-1}+\left(\frac{n}{n-1}\right)^{2}+\cdots+\left(\frac{n}{n-1}\right)^{i}\right)} \left(k_{i}+1\right)^{\frac{1}{k_{i}+1}} \left(k_{i-1}+1\right)^{\frac{1}{k_{i}+1}\frac{n}{n-1}} \cdots \\ &\left(k_{1}+1\right)^{\frac{1}{k_{i}+1}\left(\frac{n}{n-1}\right)^{i-1}} \|\varphi_{+}\|_{L^{1}}^{\frac{k_{i}}{k_{i-1}}+1\cdots\cdots\frac{k_{1}}{k_{1}+1}} \\ \end{aligned}$$
Ising $k_{i} = \frac{n}{n} \left(k_{i-1}+1\right)$ and (3.3) we have

Using $k_i = \frac{n}{n-1}(k_{i-1}+1)$ and (3.3), we have (3.5) $\frac{k_i}{k_i+1}\frac{k_{i-1}}{k_{i-1}+1}\cdots\frac{k_1}{k_1+1} = \frac{k_1}{(\frac{n-1}{n})^{i-1}k_{i+1}} \to \frac{k_1}{k_1+n}$, when $i \to \infty$.

In addition, we have

(3.6)
$$C^{\frac{1}{k_{i+1}} + \frac{k_{i}}{k_{i+1}} \frac{1}{k_{i-1}+1} + \dots + \frac{k_{i}}{k_{i+1}} \frac{k_{i-1}}{k_{i-1}+1} \dots \frac{1}{k_{1}+1}}{= C^{\frac{1}{k_{i+1}} (\frac{n}{n-1} + (\frac{n}{n-1})^{2} + \dots + (\frac{n}{n-1})^{i})}}{= C^{\frac{n((\frac{n}{n-1})^{i}-1)}{(\frac{n}{n-1})^{i}(k_{1}+n)-n}} \to C^{\frac{n}{(k_{1}+n)}}, \text{ when } i \to \infty$$

Using (3.3), we have

$$(k_{i}+1)^{\frac{1}{k_{i}+1}}(k_{i-1}+1)^{\frac{k_{i}}{k_{i}+1}\frac{1}{k_{i-1}+1}}\cdots(k_{1}+1)^{\frac{k_{i}}{k_{i}+1}\frac{k_{i-1}}{k_{i-1}+1}\cdots\frac{1}{k_{1}+1}}$$
$$(3.7) \qquad =(k_{i}+1)^{\frac{1}{k_{i}+1}}(k_{i-1}+1)^{\frac{1}{k_{i}+1}\frac{n}{n-1}}\cdots(k_{1}+1)^{\frac{1}{k_{i}+1}(\frac{n}{n-1})^{i-1}}$$
$$=e^{\sum_{a=0}^{i-1}\frac{(\frac{n}{n-1})^{a}\ln(k_{i-a}+1)}{k_{i}+1}} < C, \text{ when } i \to \infty.$$

By (3.2), letting $i \to \infty$ in (3.4), we have

(3.8)
$$\|\varphi_+\|_{L^{\infty}} \le C \|\varphi_+\|_{L^1}^{\frac{1}{n+1}} \le CA^{\frac{1}{n+1}}.$$

3.2. The lower bound for φ . First we use Székelyhidi 's argument [31], to prove the L^1 estimates. Consider the operator

(3.9)
$$\Delta \psi := \Delta_{\alpha} \psi + \operatorname{tr}_{\alpha} W(\psi) + (\operatorname{tr}_{\alpha} B) \varphi,$$

where $\Delta_{\alpha}\psi = \alpha^{i\bar{j}}\psi_{i\bar{j}}$ is the Chern Laplace operator, $\operatorname{tr}_{\alpha}W(\psi), \operatorname{tr}_{\alpha}B$ are the trace of $W(\psi)$, B, respectively, and $(\alpha^{i\bar{j}})$ is the inverse matrix of $(\alpha_{i\bar{j}})$.

Proposition 3.2. Let (M, α) be a compact Hermitian manifold with $\int_M \alpha^n = 1$. Suppose that φ satisfies

(3.10)
$$\int_{M} e^{\varphi} \alpha^{n} = A, \quad \Delta \varphi \ge -C_{0}$$

and $\sup \varphi \leq 1$ for some constant C_0 . Then there exists a constant C depending on A, C_0, ω_0, h and (M, α) such that

$$\int_M |\varphi| \, \alpha^n \le C.$$

Proof. By (3.10), $\sup_M e^{\varphi} \ge A$, which implies

$$(3.11) \qquad \qquad \sup \varphi \ge \ln A$$

Using Proposition 3.1, $|\sup_M \varphi| \leq C$. Note $\alpha^{ij} B_{i\bar{j}} \leq 0$. By the argument in [31], using weakly Harnack inequality, we can prove a bound on the integral of $|\sup \varphi - \varphi|$. This implies Proposition 3.2.

In the following, the infimum estimates will be proved. We follow the arguments of [31], which are generalizations of the arguments in [3, 4]. In [17], George also proved a similar result.

Proposition 3.3. Let φ be a smooth solution of (1.3) and $\sup \varphi \leq 1$. Suppose

$$*(\sqrt{-1}\partial\overline{\partial}\alpha)^{n-2} \le 0.$$

Then there exists a constant C depending on A, $\|h\|_{C^0}$, $\|\omega_0\|_{C^0}$ and (M, α) such that

(3.12)
$$\inf_{M} \varphi \ge -C.$$

Proof. If $\inf_M \varphi \geq 0$, the proof is complete. Now we assume $\inf_M \varphi < 0$ and denote $I = \inf_M \varphi$. Assume $I = \varphi(x_0)$ and choose a local coordinate chart (z_1, \ldots, z_n) near x_0 containing the unit ball $B_1(0) \subset \mathbb{R}^{2n}$ such that the point x_0 corresponds to the origin $0 \in \mathbb{R}^{2n}$. Define $v = \varphi + \varepsilon \sum_{i=1}^{2n} |z_i|^2$ for a small $\varepsilon > 0$. Then

$$v(0) = I, \quad v \ge I + \varepsilon \text{ on } \partial B_1(0).$$

Denote

(3.13)
$$P = \left\{ x \in B_1(0) : |Dv(x)| \le \frac{\varepsilon}{2}, v(y) \ge v(x) + Dv(x) \cdot (y - x), \\ \forall \ y \in B_1(0) \right\}.$$

Choose ϵ small enough such that $I + \epsilon \leq 0$. Then for any $p \in P$,

(3.14)
$$v(p) \le v(0) + |Dv||p| \le I + \epsilon \le 0.$$

Using the modified Alexandroff- Bakelman-Pucci maximum principle [31, Proposition 10], we have

(3.15)
$$c_0 \varepsilon^{2n} \le \int_P \det(D^2 v),$$

where c_0 is a constant depending only on n.

Claim 1. On P, $det(D^2v) \leq C$.

Proof of Claim 1. Note that v is a convex function on P and so $D^2 v \ge 0$ on P. Then

(3.16)
$$\sqrt{-1}\partial\overline{\partial}\varphi \ge \sqrt{-1}\partial\overline{\partial}v - C\varepsilon \operatorname{Id} \ge -C\varepsilon \operatorname{Id}$$
 on P

Combing with (3.14), $\varphi \tilde{B} \ge 0$ on P. If we choose ϵ small enough, then

$$\tilde{\chi} + \frac{1}{n-1}((\Delta \varphi)\alpha - \sqrt{-1}\partial\overline{\partial}\varphi) + Z + \varphi \tilde{B} \ge \frac{1}{2}\tilde{\chi},$$

which implies $\tilde{\lambda}_i \geq \frac{1}{C}$. By (2.1), we have

$$\tilde{\chi} + \frac{1}{n-1}((\Delta \varphi)\alpha - \sqrt{-1}\partial\overline{\partial}\varphi) + Z + \varphi \tilde{B} \le C\tilde{\chi},$$

and so $|\sqrt{-1}\partial\overline{\partial}\varphi| \leq C$ on *P*. Then

$$(3.17) \qquad \qquad |\sqrt{-1}\partial\overline{\partial}v| \le C \text{ on } P.$$

Recall that for Hermitian matrices $A \ge 0$ and $N \ge 0$, $\det(A+N) \ge \det(A) + \det(N)$. On P, we have $D^2v \ge 0$ and so

$$\det(\sqrt{-1}\partial\overline{\partial}v) = 2^{-2n}\det(D^2v + J^T \cdot D^2v \cdot J) \ge 2^{-2n+1}\det(D^2v).$$

Combining this with (3.17),

$$\det(D^2 v) \le 2^{2n-1} \det(\sqrt{-1}\partial\overline{\partial}v) \le C.$$

Using Claim 1 and (3.15),

 $(3.18) c_0 \varepsilon^{2n} \le C|P|.$

For each $x \in P$, choosing y = 0 in (3.13), then Since $|Dv| \leq \frac{\epsilon}{2}$,

$$I = v(0) \ge v(x) - \epsilon.$$

Since $I + \varepsilon \leq 0$, it follows that $|I + \varepsilon| \leq -v$ on P. Using (3.18), it follows

$$c_0 \varepsilon^{2n} \le |P| \le \frac{\int_P (-v)\chi^n}{|I + \varepsilon|}.$$

 $|I + \varepsilon| \leq \frac{1}{|P|} \int_{P} (-v) \chi^{n} \leq C$. This gives the required estimate of I.

4. Second order estimates

In this section, we prove the second order estimates.

Theorem 4.1. Suppose that

$$(4.1) -C_0 \le \varphi \le M_0 A^{\frac{1}{n+1}}$$

There exists a constant A_0 and C such that if $A \leq A_0$

(4.2)
$$\sup_{M} |\sqrt{-1}\partial\overline{\partial}\varphi|_{\alpha} \le C(\sup_{M} |\partial\varphi|_{\alpha}^{2} + 1),$$

where C depends on (M, α) , ω_0 , h, A and A_0 is a constant such that

$$M_0 A_0^{\frac{1}{n+1}} \|\tilde{B}\|_{\alpha} \le \frac{\kappa}{4}.$$

In this paper, we compute using covariant derivatives with respect to the Chern connection of α . Denote $g_{i\overline{j},k} = \nabla_k g_{i\overline{j}}$ and $g_{i\overline{j},k\overline{l}} = \nabla_{\overline{l}} \nabla_k g_{i\overline{j}}$. Let us recall the commutation formulas for covariant derivatives:

$$(4.3) \quad \begin{array}{l} \varphi_{i\overline{j}k} = \varphi_{k\overline{j}i} - T^{p}_{ki}\varphi_{p\overline{j}}, \\ \varphi_{i\overline{j}k\overline{l}} = \varphi_{k\overline{l}i\overline{j}} - T^{p}_{ki}\varphi_{p\overline{l}\overline{j}} - \overline{T^{q}_{lj}}\varphi_{k\overline{q}i} + \varphi_{p\overline{j}}R_{k\overline{l}i} \ ^{p} - \varphi_{p\overline{l}}R_{i\overline{j}k} \ ^{p} - T^{p}_{ik}\overline{T^{q}_{lj}}\varphi_{p\overline{q}}, \end{array}$$

where T_{ij}^k and $R_{i\overline{j}k}^{\ \ l}$ are components of torsion tensor and curvature tensor induced by the Chern connection of α .

Suppose that $\lambda_1 \geq \lambda_2 \geq \cdots \geq \lambda_n$ are the eigenvalues of $\Phi = (\Phi_k^i) = (\alpha^{i\bar{j}}\omega_{k\bar{j}})$ with respect to α . For convenience, we write $|\cdot| = |\cdot|_{\alpha}$ and

$$K = \sup_{M} |\partial \varphi|^2 + 1.$$

On $\Omega = \{\lambda_1 > 0\} \subset M$, consider

$$Q = \log \lambda_1 + \eta(|\partial \varphi|^2) + \xi(\varphi)$$

where

$$\xi(\varphi) = D_1 e^{-D_2(\varphi - \sup_M \varphi - 1)}, \quad \eta(s) = -\frac{1}{2}\log(2K - s),$$

for constants D_1 and D_2 to be determined later. By directly calculation,

(4.4)
$$\eta'' = 2(\eta')^2, \quad \frac{1}{8K} \le \eta' \le \frac{1}{4K}.$$

We may assume $\Omega \neq \emptyset$, otherwise we are done. Note $Q(z) \rightarrow -\infty$ as z approaches to $\partial\Omega$. Let $Q(x_0) = \max_M Q(x_0)$, where $x_0 \in \Omega$. Using the standard argument [31, 32, 37], to prove Theorem 4.1, it suffices to show

(4.5)
$$\lambda_1(x_0) \le CK$$

Near x_0 , choose a local normal coordinates (z_1, \dots, z_n) such that at x_0 ,

(4.6)
$$\alpha_{i\overline{j}} = \delta_{ij}, \quad g_{i\overline{j}} = \delta_{ij}g_{i\overline{i}}, \quad g_{1\overline{1}} \ge g_{2\overline{2}} \ge \cdots \ge g_{n\overline{n}}.$$

and

$$F^{1\overline{1}} \le F^{2\overline{2}} \le \dots \le F^{n\overline{n}}.$$

We assume that λ_1 is smooth and $\lambda_1 > \lambda_2$ at x_0 , using a viscosity argument (see [31, 32, 40])

Applying the maximum principle at x_0 , we see that

(4.7)
$$0 \ge L(Q) = \frac{L(\lambda_1)}{\lambda_1} - F^{i\bar{i}} \frac{|(\lambda_1)_i|^2}{\lambda_1^2} + \eta' L(|\partial \varphi|^2) + \eta'' F^{i\bar{i}} |(|\partial \varphi|^2)_i|^2 + \xi' L(\varphi) + \xi'' F^{i\bar{i}} |\varphi_i|^2$$

In the sequel, we will use the Einstein summation, and use C to denote a constant depending on $\|\varphi\|_{C^0}$, h, ω_0 , (M, α) , and C_D to denote a constant further depending on D_1 and D_2 ,

4.0.1. Lower bound for L(Q).

Proposition 4.2. For $\varepsilon \in (0, \frac{1}{3}]$, at x_0 , we have

(4.8)
$$0 \ge G - \sum_{k=1}^{n} \frac{C}{\lambda_{1}} F^{i\bar{i}} |g_{1\bar{1},i}| - N + \frac{3\eta'}{4} \sum_{j} F^{i\bar{i}} (|\varphi_{ij}|^{2} + |\varphi_{i\bar{j}}|^{2}) + \eta'' F^{i\bar{i}} |(|\partial\varphi|^{2})_{i}|^{2} + \xi' L(\varphi) + \xi'' F^{i\bar{i}} |\varphi_{i}|^{2} - C\mathcal{F},$$

where

$$G = -\frac{1}{\lambda_1} F^{i\overline{j},k\overline{l}} \nabla_1 g_{i\overline{j}} \nabla_{\overline{1}} g_{k\overline{l}}, N = F^{i\overline{i}} \frac{|(\lambda_1)_i|^2}{\lambda_1^2}.$$

The third term is the bad term that we need to control. To prove Proposition 4.2, we shall estimate the lower bounds of $L(\lambda_1)$ and $L(|\partial u|^2)$, respectively. First we deal with the term $F^{i\bar{i}}g_{1\bar{1},i\bar{i}}$ at x_0 . We have the following inequality.

Proposition 4.3. Assume $\lambda_1 \geq K$. At x_0 , we have

$$F^{i\bar{i}}g_{1\bar{1},i\bar{i}} \ge -F^{i\bar{j},k\bar{l}}\nabla_{1}g_{i\bar{j}}\nabla_{\bar{1}}g_{k\bar{l}} - \sum_{k=1}^{n} CF^{i\bar{i}}|g_{1\bar{k},i}| - \frac{1}{32K}F^{i\bar{i}}|\varphi_{pi}|^{2} - C\mathcal{F}\lambda_{1}.$$

Proof. By differentiating (2.7) along $\nabla_{\overline{1}} \nabla_1$,

_

(4.9)
$$F^{i\bar{i}}g_{i\bar{i},1\bar{1}} = F^{i\bar{i}}\nabla_{\bar{1}}\nabla_{1}(\chi_{i\bar{i}} + \varphi_{i\bar{i}} + W_{i\bar{i}} + \varphi B_{i\bar{i}}) = (n-1)h_{1\bar{1}} - F^{i\bar{j},k\bar{l}}\nabla_{1}g_{i\bar{j}}\nabla_{\bar{1}}g_{k\bar{l}}.$$

On the other hand, by (4.3),

(4.10)
$$F^{i\bar{i}}\varphi_{1\bar{1}i\bar{i}} = F^{i\bar{i}}\varphi_{i\bar{i}1\bar{1}} + \sum_{k=1}^{n} F^{i\bar{i}}O(|\varphi_{1\bar{k}i}|) + O(\lambda_1)\mathcal{F},$$

where $O(|\varphi_{1\bar{k}i}|)$, $O(\lambda_1)$ mean the terms can controlled by $|\varphi_{1\bar{k}i}|$, λ_1 respectively. Using (4.10) and (4.9), it follows

$$F^{ii}\varphi_{1\bar{1}i\bar{i}} + F^{ii}W_{i\bar{i},1\bar{1}}$$

$$\geq -C - F^{i\bar{j},k\bar{l}}\nabla_{1}g_{i\bar{j}}\nabla_{\bar{1}}g_{k\bar{l}} - C\mathcal{F}\lambda_{1} - \sum_{k=1}^{n} CF^{i\bar{i}}|\varphi_{1\bar{k}i}|.$$

Therefore,

$$\begin{split} F^{i\bar{i}}g_{1\bar{1},i\bar{i}} = & F^{i\bar{i}}\chi_{1\bar{1},i\bar{i}} + F^{i\bar{i}}\varphi_{1\bar{1}i\bar{i}} + F^{i\bar{i}}W_{1\bar{1},i\bar{i}} + F^{i\bar{i}}(\varphi B)_{1\bar{1},i\bar{i}} \\ \geq & -F^{i\bar{j},k\bar{l}}\nabla_{1}g_{i\bar{j}}\nabla_{\bar{1}}g_{k\bar{l}} - \sum_{k=1}^{n} CF^{i\bar{i}}|\varphi_{1\bar{k}i}| - C\mathcal{F}\lambda_{1} \\ & +F^{i\bar{i}}W_{1\bar{1},i\bar{i}} - F^{i\bar{i}}W_{i\bar{i},1\bar{1}} - C. \end{split}$$

Note $\varphi_{1\bar{k}i}=g_{1\bar{k},\bar{1}}-\chi_{1\bar{k},i}-W_{1\bar{k},i}-B_{1\bar{k},i}.$ We have

(4.11)
$$\sum_{k=1}^{n} F^{i\bar{i}} |\varphi_{1\bar{k}i}| \leq C(\sum_{p} F^{i\bar{i}} |\varphi_{p}| + \lambda_{1}\mathcal{F}) \\ \leq \frac{1}{96K} \sum_{p} F^{i\bar{i}} |\varphi_{pi}|^{2} + C\lambda_{1}\mathcal{F} + \sum_{k=1}^{n} CF^{i\bar{i}} |g_{1\bar{k},i}|.$$

It follows that

(4.12)
$$F^{i\bar{i}}g_{1\bar{1},i\bar{i}} \ge -F^{i\bar{j},k\bar{l}}\nabla_{1}g_{i\bar{j}}\nabla_{\overline{1}}g_{k\bar{l}} - \sum_{k=1}^{n} F^{i\bar{i}}|g_{1\bar{k},i}| - C\mathcal{F}\lambda_{1} + F^{i\bar{i}}W_{1\bar{1},i\bar{i}} - F^{i\bar{i}}W_{i\bar{i},1\bar{1}} - \frac{1}{96K}\sum_{p} F^{i\bar{i}}|\varphi_{pi}|^{2}.$$

Now we deal the terms involving W. First we deal with $F^{i\bar{i}}W_{i\bar{i},1\bar{1}}$.

Lemma 4.4. If $\lambda_1 \geq K$, we have

$$F^{i\bar{i}}W_{i\bar{i},1\bar{1}} \leq \frac{1}{96K} \sum_{p} F^{i\bar{i}} |\varphi_{pi}|^2 + C(F^{i\bar{i}}|g_{1\bar{1},i}| + \lambda_1 \mathcal{F}).$$

Proof. By (2.9) and (2.2), we have

$$F^{i\bar{i}}W_{i\bar{i},1\bar{1}} = \frac{1}{n-1}\sum_{k}\sum_{i\neq k}\tilde{F}^{i\bar{i}}W_{k\bar{k},1\bar{1}}$$
$$= \frac{1}{n-1}\sum_{i}\tilde{F}^{i\bar{i}}\sum_{k\neq i}W_{k\bar{k},1\bar{1}} = \tilde{F}^{i\bar{i}}Z_{i\bar{i},1\bar{1}}.$$

By Lemma 2.3, $Z_{1\bar{1},1\bar{1}}$ does not contain the terms $\varphi_{11\bar{1}}$ and $\varphi_{1\bar{1}}$ or their complex conjugates. By Lemma 2.1(3) (4), it follows that

$$\begin{aligned} F^{i\bar{i}}W_{i\bar{i},1\bar{1}} &\leq C(\tilde{F}^{1\bar{1}}\sum_{k>1}(|\varphi_{k1}|+|\varphi_{k\bar{1}1}|)+\sum_{i>1,k}\tilde{F}^{i\bar{i}}(|\varphi_{k1}|+|\varphi_{k\bar{1}1}|)+\lambda_{1}\mathcal{F}) \\ &\leq C(\sum_{i>1}F^{i\bar{i}}(|\varphi_{i1}|+|\varphi_{k\bar{1}1}|)+F^{1\bar{1}}\sum_{k}(|\varphi_{k1}|+|\varphi_{k\bar{1}1})+\lambda_{1}\mathcal{F}) \\ &\leq C(F^{1\bar{1}}(|\varphi_{11}|+|\varphi_{1\bar{1}1}|)+\sum_{i>1}F^{i\bar{i}}(|\varphi_{i1}|+|\varphi_{i\bar{1}1}|)+\lambda_{1}\mathcal{F}) \\ &\leq C(\sum_{p}F^{i\bar{i}}|\varphi_{pi}|+F^{i\bar{i}}|\varphi_{1\bar{1}i}|+\lambda_{1}\mathcal{F}). \end{aligned}$$

Here in the third inequality, we used $F^{1\overline{1}} \leq F^{i\overline{i}}$. Note that

$$\varphi_{1\bar{1}i} = g_{1\bar{1},\bar{i}} - \chi_{1\bar{1},i} - W_{1\bar{1},i} - (\varphi B_{1\bar{1}})_i,$$

which implies

$$F^{i\overline{i}}|\varphi_{1\overline{1}k}| \leq F^{i\overline{i}}|g_{1\overline{1},i}| + C(\sum_{p} F^{i\overline{i}}|\varphi_{pi}| + \lambda_1 \mathcal{F}).$$

Using the Cauchy-Schwarz inequality and $\lambda_1 \geq K$, we have

$$F^{i\bar{i}}W_{i\bar{i},1\bar{1}} \le \frac{1}{96K} \sum_{p} F^{i\bar{i}} |\varphi_{pi}|^2 + C(F^{i\bar{i}}|g_{1\bar{1},i}| + \lambda_1 \mathcal{F}).$$

Next we deal with the $F^{i\bar{i}}W_{1\bar{1},i\bar{i}}$. We have the following inequality.

Lemma 4.5. At x_0 , we have

(4.13)
$$F^{i\bar{i}}W_{1\bar{1},i\bar{i}} \ge -\sum_{p} \frac{1}{96K} F^{i\bar{i}} |\varphi_{pi}|^2 - C\lambda_1 \mathcal{F}.$$

Proof. By directly calculation,

(4.14)
$$F^{i\bar{i}}W_{1\bar{1},i\bar{i}} = F^{i\bar{i}}2\operatorname{Re}(\nabla_{\bar{i}}\nabla_{i}(W_{1\bar{1}}^{p}\varphi_{p}))$$
$$= 2\operatorname{Re}(F^{i\bar{i}}W_{1\bar{1}}^{p}\varphi_{pi\bar{i}}) + O(F^{i\bar{i}}|\varphi_{pi}|) + O(\lambda_{1})\mathcal{F}$$

Differentiating (2.7) along ∇_p , we have

$$F^{i\bar{i}}g_{i\bar{i},p} = (n-1)h_p,$$

which implies

(4.15)
$$|F^{ii}\varphi_{i\bar{i}p}| = |F^{ii}(g_{i\bar{i},p} - \chi_{i\bar{i},p} - W_{i\bar{i},p} - (\varphi B_{i\bar{i}})_p)| \le CK^{\frac{1}{2}}\mathcal{F} + |F^{i\bar{i}}W_{i\bar{i},p}|.$$

Here we used $K \ge 1$. Now we deal with $|F^{i\bar{i}}W_{i\bar{i},p}|$. By (2.2), we have

$$F^{ii}W_{i\bar{i},p} = \tilde{F}^{ii}Z_{i\bar{i},p}.$$

Then, using Lemma 2.3 and (2.9),

$$F^{i\bar{i}}W_{i\bar{i},p} \leq C(\tilde{F}^{1\bar{1}}\sum_{k>1}(|\varphi_{kp}|+|\varphi_{\bar{k}p}|)+\tilde{F}^{i\bar{i}}(|\varphi_{ip}|+|\varphi_{\bar{i}p}|)+K^{\frac{1}{2}}\mathcal{F})$$

$$\leq C(\sum_{k>1}F^{i\bar{i}}((|\varphi_{ip}|+|\varphi_{\bar{i}p}|)+F^{1\bar{1}}\sum_{k}(|\varphi_{kp}|+|\varphi_{\bar{k}p}|)+K^{\frac{1}{2}}\mathcal{F})$$

$$\leq C(\sum_{p}F^{i\bar{i}}(|\varphi_{pi}|+|\varphi_{\bar{i}p}|)+K^{\frac{1}{2}}\mathcal{F}).$$

Combing this inequality with (4.15),

(4.16)
$$|F^{i\overline{i}}\varphi_{i\overline{i}p}| \leq \sum_{p} F^{i\overline{i}}(|\varphi_{pi}| + |\varphi_{\overline{i}p}|) + K^{\frac{1}{2}}\mathcal{F}).$$

From (4.14), we have

$$F^{i\bar{i}}W_{1\bar{1},i\bar{i}} \ge -C(\sum_{p}F^{i\bar{i}}|\varphi_{pi}| + \lambda_{1}\mathcal{F})$$
$$\ge -\sum_{p}\frac{1}{96K}F^{i\bar{i}}|\varphi_{pi}|^{2} - C\lambda_{1}\mathcal{F}.$$

с		
L		
L		
L		

Combing (4.12), Lemma 4.4 with Lemma 4.5,

$$\begin{split} F^{i\overline{i}}g_{1\overline{1},i\overline{i}} &\geq -F^{i\overline{j},k\overline{l}}\nabla_{1}g_{i\overline{j}}\nabla_{\overline{1}}g_{k\overline{l}} - \sum_{k=1}^{n} CF^{i\overline{i}}|g_{1\overline{k},i}| \\ &- \frac{1}{32K}F^{i\overline{i}}|\varphi_{pi}|^{2} - C\lambda_{1}\mathcal{F}. \end{split}$$

Using the inequality about $L(g_{1\bar{1}})$ at x_0 , we immediately obtain the following inequality for $L(\lambda_1)$.

Proposition 4.6. Assume $\lambda_1 \geq CK$. For each $\varepsilon \in (0, \frac{1}{3}]$, at x_0 , we have

(4.17)
$$L(\lambda_{1}) \geq \sum_{p>1} F^{i\bar{i}} \frac{|g_{1\bar{p},i}|^{2}}{8n\lambda_{1}} - F^{i\bar{j},k\bar{l}} \nabla_{1}g_{i\bar{j}} \nabla_{\bar{1}}g_{k\bar{l}} - \sum_{k=1}^{n} CF^{i\bar{i}}|g_{1\bar{1},i}| - \frac{1}{32K} F^{k\bar{k}}|\varphi_{pk}|^{2} - C\lambda_{1}\mathcal{F}.$$

Proof. We need the following formulas (see e.g. [10, 30, 31]):

$$\frac{\partial \lambda_1}{\partial \Phi_p^q} = \delta_{1p} \delta_{1q},$$
$$\frac{\partial^2 \lambda_1}{\partial \Phi_p^q \partial \Phi_r^s} = (1 - \delta_{1p}) \frac{\delta_{1q} \delta_{1r} \delta_{ps}}{\lambda_1 - \lambda_p} + (1 - \delta_{1r}) \frac{\delta_{1s} \delta_{1p} \delta_{rq}}{\lambda_1 - \lambda_r}.$$

By directly calculation,

$$L(\lambda_{1}) = F^{i\overline{i}} \frac{\partial^{2}\lambda_{1}}{\partial \Phi_{p}^{q} \partial \Phi_{r}^{s}} \nabla_{i}(\Phi_{r}^{s}) \nabla_{\overline{i}}(\Phi_{p}^{q}) + F^{i\overline{i}} \frac{\partial\lambda_{1}}{\partial \Phi_{p}^{q}} \nabla_{i} \nabla_{\overline{i}}(\Phi_{p}^{q})$$

$$= F^{i\overline{i}} g_{1\overline{1},i\overline{i}} + \sum_{p>1} \frac{F^{i\overline{i}}(|g_{1\overline{p},i}|^{2} + |g_{p\overline{1},i}|^{2})}{\lambda_{1} - \lambda_{p}}$$

$$\geq F^{i\overline{i}} g_{1\overline{1},i\overline{i}} + \sum_{p>1} F^{i\overline{i}} \frac{|g_{1\overline{p},i}|^{2}}{4n\lambda_{1}}.$$

Using the Cauchy-Schwarz inequality, we have for $k \neq 1$,

(4.19)
$$|g_{1\bar{k},i}| \le \frac{1}{8n\lambda_1} |g_{1\bar{k},i}|^2 + C\lambda_1.$$

Combining this inequality with (4.18) and Proposition 4.3, we obtain Lemma 4.6. $\hfill \Box$

Now, we give the lower bound of $L(|\partial \varphi|^2).$

Lemma 4.7. At x_0 , we have

(4.20)
$$L(|\partial \varphi|^2) \ge \frac{3}{4} \sum_{j} F^{i\bar{i}}(|\varphi_{ij}|^2 + |\varphi_{i\bar{j}}^2) - CK\mathcal{F}.$$

Proof. By (4.3), (4.16) and the Cauchy-Schwarz inequality, we have

$$F^{ii}((|\partial\varphi|_{g}^{2})_{i\overline{i}})$$

$$= \sum_{k} F^{i\overline{i}}(|\varphi_{ki}|^{2} + |\varphi_{k\overline{i}}|^{2}) + 2\operatorname{Re}(\sum_{k} F^{i\overline{i}}\varphi_{k}\varphi_{\overline{k}i\overline{i}})$$

$$(4.21) \geq \sum_{k} F^{i\overline{i}}(|\varphi_{ik}|^{2} + |\varphi_{i\overline{k}}|^{2}) - CK^{\frac{1}{2}}\sum_{k} F^{i\overline{i}}(|\varphi_{ki}| + |\varphi_{k\overline{i}}|) - CK\mathcal{F}$$

$$\geq \frac{3}{4}\sum_{k} F^{i\overline{i}}(|\varphi_{ki}|^{2} + |\varphi_{k\overline{i}}|^{2}) - CK\mathcal{F}.$$

We will use the above computations to prove Proposition 4.2.

Proof of Proposition 4.2. Combining (4.7), (4.4) and Lemma 4.6 with Lemma 4.7, we obtain

$$(4.22) \qquad 0 \ge -\frac{1}{\lambda_{1}} F^{i\overline{j},k\overline{l}} \nabla_{1} g_{i\overline{j}} \nabla_{\overline{1}} g_{k\overline{l}} - \sum_{i=1}^{n} \frac{C}{\lambda_{1}} F^{i\overline{i}} |g_{1\overline{1},i}| - F^{i\overline{i}} \frac{|(\lambda_{1})_{i}|^{2}}{\lambda_{1}^{2}} + \frac{3\eta'}{4} \sum_{j} F^{i\overline{i}} (|\varphi_{ij}|^{2} + |\varphi_{i\overline{j}}|^{2}) + \eta'' F^{i\overline{i}} |(|\partial\varphi|^{2})_{i}|^{2} + \xi' L(\varphi) + \xi'' F^{i\overline{i}} |\varphi_{i}|^{2} - C\mathcal{F}.$$

Note that the first term is G and the third term is N.

Now we complete the proof of Theorem 4.1.

Proof. We now deal with two cases separately.

Case 1. Assume $\lambda_1 \geq -\delta \lambda_n$. Define the set

(4.23) $I = \{i: F^{i\bar{i}} > \delta^{-1} F^{1\bar{1}}\}.$

Now we decompose the term N into three terms based on I.

$$(4.24) \qquad N = F^{i\bar{i}} \frac{|(\lambda_1)_i|^2}{\lambda_1^2} \\ = \sum_{i \notin I} \frac{F^{i\bar{i}}|g_{1\bar{1},i}|}{\lambda_1^2} + 2\delta \sum_{i \in I} \frac{F^{i\bar{i}}|g_{1\bar{1},i}|}{\lambda_1^2} + (1 - 2\delta) \sum_{i \in I} \frac{F^{i\bar{i}}|g_{1\bar{1},i}|}{\lambda_1^2} \\ =: N_1 + N_2 + N_3.$$

Using $dQ(x_0) = 0$, we can prove a lower bound for the terms $-N_1$ and $-N_2$. Lemma 4.8. At x_0 , we have

$$\begin{split} -N_1 - N_2 &\geq -\eta'' \sum_{k \notin I} F^{i\bar{i}} ||\partial \varphi|_i^2|^2 - 2(\xi')^2 \delta^{-1} F^{1\bar{1}} K - 2\delta \eta'' \sum_{i \in I} F^{i\bar{i}} ||\partial \varphi|_i^2| \\ &- \frac{\xi''}{2} \sum_{k \in I} F^{i\bar{i}} |\varphi_i|^2. \end{split}$$

Proof. Since Q attains its maximum at x_0 , we have $dQ(x_0) = 0$. Then

(4.25)
$$\frac{g_{1\bar{1},i}}{\lambda_1} = -\eta'(|\partial\varphi|^2)_i - \xi'(\varphi)\varphi_i$$

for each $1 \le i \le n$. Using (4.25),

$$-\sum_{i \notin I} \frac{F^{ii} |g_{1\bar{1},i}|^2}{\lambda_1^2} = -\sum_{i \notin I} F^{i\bar{i}} |\eta'| \partial \varphi|_i^2 + \xi' \varphi_i|^2$$

$$\geq -2(\eta')^2 \sum_{i \notin I} F^{i\bar{i}} ||\partial \varphi|_i^2|^2 - 2(\xi')^2 \sum_{i \notin I} F^{i\bar{i}} |\varphi_k|^2$$

$$\geq -\eta'' \sum_{i \notin I} F^{i\bar{i}} ||\partial \varphi|_i^2|^2 - 2(\xi')^2 \delta^{-1} F^{1\bar{1}} K.$$

Now we choose δ such that

(4.26)
$$4\delta(\xi')^2 \le \frac{1}{2}\xi''.$$

Then, for $k \in I$, using the similar argument, we have

$$\begin{split} -2\delta \sum_{i\in I} \frac{F^{i\bar{i}} |g_{1\bar{1},i}|^2}{\lambda_1^2} &\geq -2\delta\eta'' \sum_{i\in I} F^{i\bar{i}} ||\partial\varphi|_i^2|^2 - 4\delta(\xi')^2 \sum_{i\in I} F^{i\bar{i}} |\varphi_i|^2\\ &\geq -2\delta\eta'' \sum_{i\in I} F^{i\bar{i}} ||\partial\varphi|_i^2|^2 - \frac{\xi''}{2} \sum_{i\in I} F^{i\bar{i}} |\varphi_i|^2. \end{split}$$

In the following, we will use G to control the term N_3 .

Lemma 4.9. For any $\epsilon > 0$, at x_0 , we have

$$G \ge N_3 - \sum_p \frac{F^{ii}}{12K} (|\varphi_{p\bar{i}}|^2 + |\varphi_{pi}|^2) + C\xi' F^{i\bar{i}} |\varphi_k|^2 + \epsilon C\xi' \mathcal{F} - CF.$$

Proof. By the concavity of the operator F (see [31, equation (67)]) and the definition of I,

$$G \ge \sum_{i \in I} \frac{F^{ii} - F^{1\overline{1}}}{\lambda_1 - \lambda_i} |\nabla_1 g_{i\overline{1}}|^2$$
$$\ge \sum_{i \in I} \frac{(1 - \delta)F^{i\overline{i}}}{\lambda_1 - \lambda_i} |\nabla_1 g_{i\overline{1}}|^2$$
$$\ge \sum_{i \in I} \frac{(1 - 2\delta)F^{i\overline{i}}}{\lambda_1} |\nabla_1 g_{i\overline{1}}|^2.$$

Here we used $\delta \lambda_1 \geq -\lambda_n$ in the last inequality. Next, we compare $g_{i\bar{1},1}$ to $g_{1\bar{1},i}$. By (4.3), we have

$$\begin{split} g_{i\bar{1},1} &= \nabla_1 (\chi_{i\bar{1}} + \varphi_{i\bar{1}} + W_{i\bar{1},1} + \varphi B_{i\bar{1}}) \\ &= \varphi_{i\bar{1}1} + W_{i\bar{1},1} + O(K^{\frac{1}{2}}) \\ &= \varphi_{1\bar{1}i} + W_{i\bar{1},1} + O(\lambda_1) \\ &= g_{1\bar{1},i} - W_{1\bar{1},i} + W_{i\bar{1},1} + O(\lambda_1). \end{split}$$

It follows that for any i, without summing

(4.27)
$$|g_{i\bar{1},1}|^2 \ge |g_{1\bar{1},i}|^2 - C(|W_{1\bar{1},i}|^2 + |W_{1\bar{i},1}|^2 + \lambda_1^2 + \lambda_1|g_{1\bar{1},i}| + |g_{1\bar{1},i}|(|W_{i\bar{1},1}| + |W_{1\bar{1},i}|)).$$

Now we deal with the terms involving W. We have the following inequalities

Claim 2. At x_0 , we have

$$\sum_{i \in I} \frac{1}{\lambda_1^2} F^{i\bar{i}} |W_{i\bar{1},1}|^2 + |W_{1\bar{1},i}|^2 \le C\mathcal{F} + \frac{C}{\lambda_1^2} \sum_p F^{i\bar{i}} |\varphi_{ip}|^2.$$

and

$$\begin{split} \sum_{i \in I} \frac{1}{\lambda_1^2} F^{i\bar{i}} |g_{1\bar{1},i}| (|W_{i\bar{1},1}| + |W_{1\bar{1},i}|) &\leq \sum_p \frac{F^{i\bar{i}}}{50K} (|\varphi_{p\bar{i}}|^2 + |\varphi_{pi}|^2) \\ &- C_{\epsilon} \xi' F^{i\bar{i}} |\varphi_i|^2 - \epsilon C \xi' \mathcal{F} + C \mathcal{F}. \end{split}$$

Proof. Assume $i \in I$. Then $i \neq 1$ and

$$\nabla_1 W_{i\bar{1}} = \operatorname{tr}_{\alpha} Z \nabla_1 \alpha_{i\bar{1}} - (n-1) \nabla_1 Z_{i\bar{1}}$$

Set $U = \sum_{p>1,q\geq 1} |u_{pq}|$. By Lemma 2.3, it follows that

(4.28)
$$|W_{i\bar{1},1}| + |W_{1\bar{1},i}| \le C(\lambda_1 + U).$$

Therefore, by Lemma 2.1(5),

(4.29)
$$\frac{1}{\lambda_1^2} F^{i\bar{i}}(|W_{i\bar{1},1}|^2 + |W_{1\bar{1},i}|^2) \le C\mathcal{F}(1 + \frac{U^2}{\lambda_1^2}) \le C\mathcal{F} + \frac{C}{\lambda_1^2} \sum_p F^{i\bar{i}} |\varphi_{ip}|^2.$$

Using (4.25) and (4.4), we have

(4.30)
$$|g_{1\bar{1},i}| \le \frac{C\lambda_1}{K^{\frac{1}{2}}} (\sum_r |\varphi_{\bar{r}i}| + \sum_r |\varphi_{ri}|) + C\lambda_1 |\xi'| |\varphi_i|.$$

By (4.28),

$$\begin{aligned} \frac{1}{\lambda_1^2} F^{i\bar{i}} |g_{1\bar{1},i}| (|W_{i\bar{1},1}| + |W_{1\bar{1},i}|) &\leq \frac{CF^{i\bar{i}}}{\lambda_1 K^{\frac{1}{2}}} (\sum_r |\varphi_{\bar{r}i}| + \sum_r |\varphi_{ri}|) (\lambda_1 + U) \\ &+ \frac{CF^{i\bar{i}}}{\lambda_1} |\xi'| |\varphi_i| (\lambda_1 + U). \end{aligned}$$

Now we deal with the first term. By directly calculation and Lemma 2.1(5),

(4.31)
$$\sum_{i \in I} \frac{CF^{ii}}{\lambda_1 K^{\frac{1}{2}}} (\sum_r |\varphi_{\bar{r}i}| + \sum_r |\varphi_{ri}|) (\lambda_1 + U)$$
$$\leq \frac{CF^{i\bar{i}}}{K^{\frac{1}{2}}} \sum_r |\varphi_{\bar{r}i}| + \frac{CF}{K^{\frac{1}{2}}} U + \frac{CF}{\lambda_1 K^{\frac{1}{2}}} U^2$$
$$\leq \frac{F^{i\bar{i}}}{100K} \sum_r (|\varphi_{\bar{r}i}|^2 + |\varphi_{pi}|^2) + CF.$$

Next, by the Cauchy-Schwarz inequality, for any $\epsilon>0,$ there exists a constant C_ϵ such that

(4.32)
$$\sum_{i \in I} \frac{CF^{i\bar{i}}}{\lambda_1} |\xi'| |\varphi_i| (\lambda_1 + U)$$
$$\leq -\epsilon \xi' \mathcal{F} - C_{\epsilon} \xi' F^{i\bar{i}} |\varphi_i|^2 - \frac{1}{\lambda_1} \mathcal{F} \xi' U^2 - \frac{1}{\lambda_1} \xi' F^{i\bar{i}} |\varphi_i|^2$$
$$\leq -\epsilon \xi' \mathcal{F} - C_{\epsilon} \xi' F^{i\bar{i}} |\varphi_i|^2 + \frac{1}{100K} \sum_p F^{i\bar{i}} |\varphi_{pi}|^2,$$

where we used the fact that $\xi' < 0$.

Combining the above three inequalities, we obtain

$$\sum_{i \in I} \frac{1}{\lambda_1^2} F^{i\bar{i}} |g_{1\bar{1},i}| (|W_{i\bar{1},1}| + |W_{1\bar{1},i}|) \le \sum_p \frac{F^{i\bar{i}}}{50K} (|\varphi_{p\bar{k}}|^2 + |\varphi_{pk}|^2) - C_{\epsilon} \xi' F^{i\bar{i}} |\varphi_i|^2 - \epsilon C \xi' \mathcal{F} + C \mathcal{F}.$$

In the following, we use Claim 2 to prove Lemma 4.9. Using (4.30), Lemma 2.1(5) and the Cauchy-Schwarz inequality, we have (4.33)

$$\begin{split} \sum_{i\in I}^{'} \frac{1}{\lambda_{1}} F^{i\bar{i}} |g_{1\bar{1},i}| \leq & \frac{1}{2K^{\frac{1}{2}}} \sum_{p} F^{i\bar{i}} |\varphi_{i\bar{p}}| + \frac{1}{2K^{\frac{1}{2}}} \mathcal{F}U - C_{\epsilon}\xi' F^{i\bar{i}} |\varphi_{i}|^{2} - \epsilon\xi' \mathcal{F} \\ \leq & \sum_{p} \frac{F^{i\bar{i}}}{100K} (|\varphi_{p\bar{i}}|^{2} + |\varphi_{pi}|^{2}) - C_{\epsilon}\xi' F^{i\bar{i}} |\varphi_{i}|^{2} - \epsilon\mathcal{F}\xi'. \end{split}$$

Combining this inequality with (4.27) and Claim 2, we finally obtain

$$\begin{split} \sum_{i \in I} \frac{F^{i\bar{i}} |\nabla_1 g_{i\bar{1}}|^2}{\lambda_1^2} \geq \sum_{i \in I} \frac{F^{i\bar{i}} |g_{1\bar{1},i}|^2}{\lambda_1^2} - \sum_p \frac{F^{i\bar{i}}}{30K} (|\varphi_{p\bar{i}}|^2 + |\varphi_{pi}|^2) \\ &+ C_{\epsilon} \xi' F^{i\bar{i}} |\varphi_i|^2 + \epsilon C \xi' \mathcal{F} - C \mathcal{F}, \end{split}$$

if we choose $\lambda_1 \geq CK$. This complete the proof of the Lemma.

Combining (4.8) with Lemma 4.8, Lemma 4.9 and (4.33), it follows

$$0 \ge \sum_{k} \frac{F^{ii}}{100K} (|\varphi_{pi}|^{2} + |\varphi_{p\bar{i}}|^{2}) + \frac{1}{2} \xi'' F^{i\bar{i}} |\varphi_{i}|^{2} + \xi' L(\varphi) - 2(\xi')^{2} \delta^{-1} F^{1\bar{1}} K - C\mathcal{F} + C_{\epsilon} \xi' F^{i\bar{i}} |\varphi_{i}|^{2} + \epsilon C \xi' \mathcal{F}.$$

In the following we deal with $\xi' L(\varphi)$.

Lemma 4.10.

$$\xi' F^{i\bar{i}} \varphi_{i\bar{i}} \ge \xi' F^{i\bar{i}} (g_{i\bar{i}} - \chi_{i\bar{i}}) + C_{\epsilon} \xi' F^{i\bar{i}} |\varphi_i|^2 + (\epsilon + M_0 A^{\frac{1}{n+1}}) \xi' \mathcal{F}.$$

Proof. Note

$$\xi' L(\varphi) = \xi' F^{i\overline{i}} (g_{i\overline{i}} - \chi_{i\overline{i}} - W_{i\overline{i}} - \varphi B_{i\overline{i}}).$$

By (2.2), we have

$$F^{i\bar{i}}W_{k\bar{k}} = \sum_{i} \tilde{F}^{i\bar{i}}Z_{i\bar{i}}$$

By Lemma 2.3 and Lemma 2.1(5), it follows that

$$|F^{i\overline{i}}W_{i\overline{i}}| \le C \sum_{i \ne k} \tilde{F}^{i\overline{i}} |\varphi_k| \le C_{\epsilon} F^{i\overline{i}} |\varphi_i|^2 + \epsilon \mathcal{F}.$$

Using (4.1), $\tilde{B} \leq 0$ and Proposition 3.1,

(4.34)
$$-\xi'\varphi F^{i\bar{i}}B_{i\bar{i}} = -\xi'\varphi \tilde{F}^{i\bar{i}}\tilde{B}_{i\bar{i}} \ge -M_0A^{\frac{1}{n+1}}|\xi'|\mathcal{F}$$

In conclusion,

$$\xi' F^{i\overline{i}} \varphi_{i\overline{i}} \ge \xi' F^{i\overline{i}} (g_{i\overline{i}} - \chi_{i\overline{i}}) + C_{\epsilon} \xi' F^{i\overline{i}} |\varphi_i|^2 + (\epsilon + M_0 A^{\frac{1}{n+1}}) \xi' \mathcal{F}.$$

Therefore, we have

(4.35)
$$0 \ge F^{1\bar{1}}(\frac{\lambda_1^2}{40K} - 2(\xi')^2 \delta^{-1}K) + (\frac{1}{2}\xi'' + C_{\epsilon}\xi')F^{i\bar{i}}|\varphi_i|^2 - C_0\mathcal{F} - \xi'F^{i\bar{i}}(g_{i\bar{i}} - \chi_{i\bar{i}}) + (\epsilon + M_0A^{\frac{1}{n+1}})\xi'\mathcal{F}.$$

By Proposition 2.2, there is a uniform positive number $\kappa > 0$ such that one of two possibilities occurs:

(a) We have
$$F^{ii}(\chi_{i\bar{i}} - g_{i\bar{i}}) > \kappa \mathcal{F}$$
, then

$$0 \ge F^{1\bar{1}}(\frac{\lambda_1^2}{40K} - 2(\xi')^2 \delta^{-1}K) + (\frac{1}{2}\xi'' + C_{\epsilon}\xi')F^{i\bar{i}}|\varphi_i|^2 - C_0\mathcal{F} + (-\kappa + \epsilon + M_0A^{\frac{1}{n+1}})\xi'\mathcal{F}.$$

We first choose $\epsilon > 0$ and A small enough such that $-\kappa + \epsilon + M_0 A^{\frac{1}{n+1}} \leq -\frac{\kappa}{2}$ and choose D_2 so large that $\frac{1}{2}\xi'' + C_{\epsilon}\xi' \geq 0$ which implies

$$0 \ge F^{1\bar{1}}(\frac{\lambda_1^2}{40K} - 2(\xi')^2 \delta^{-1}K) - C_0 \mathcal{F} - \frac{1}{2}\kappa \xi' \mathcal{F}.$$

By choosing D_2 large enough such that $-C_0 \mathcal{F} - \frac{1}{2}\kappa \xi' \mathcal{F} \ge 0$, we obtain the required upper bound for λ_1 .

(b) We have $F^{1\bar{1}} > \kappa \mathcal{F}$. Choose the constant that is similar to case(a). By (4.35),

$$0 \ge \kappa \mathcal{F}(\frac{\lambda_1^2}{40K} - 2(\xi')^2 \delta^{-1}K) - C\mathcal{F} + \epsilon C\xi' \mathcal{F} + C\xi' \mathcal{F} + \xi' F^{i\bar{i}} g_{i\bar{i}}.$$

Since $F^{i\bar{i}}g_{i\bar{i}} \leq \mathcal{F}\lambda_1$, we obtain

$$0 \ge \frac{\kappa \lambda_1^2}{40K^2} - C(1 + K^{-1} + \lambda_1 K^{-1}),$$

which implies

$$\lambda_1 \leq CK.$$

for a uniform constant C.

Case 2. We assume that $\delta \lambda_1 < -\lambda_n$ where δ is a fixed constant. Since $F^{n\bar{n}} \geq \frac{\mathcal{F}}{n}$ and $\lambda_n^2 > \delta^2 \lambda_1^2$,

$$\sum_{p} \frac{F^{i\bar{i}}}{6K} (|\varphi_{pi}|^2 + |\varphi_{p\bar{k}}|^2) \ge \frac{F^{n\bar{n}}}{6K} |\varphi_{n\bar{n}}|^2 \ge \frac{\delta^2 \mathcal{F}}{6nK} \lambda_1^2 - C\mathcal{F}.$$

Combining with (4.8),

$$0 \geq \frac{-F^{i\bar{i}}|g_{1\bar{1},i}|^2}{\lambda_1^2} + \frac{\delta^2}{6nK}\mathcal{F}\lambda_1^2 + \eta''F^{i\bar{i}}|\partial_i|\partial\varphi|^2|^2 + \xi'L(\varphi) - C(F^{i\bar{i}}\lambda_1^{-1}|g_{1\bar{1},i}| + \mathcal{F}).$$

By directly calculation,

$$F^{i\bar{i}}|\varphi_{i\bar{i}}| \le C\mathcal{F}\lambda_1$$

and

$$CF^{i\bar{i}}\lambda_1^{-1}|g_{1\bar{1},i}| \le \frac{1}{2}\frac{F^{i\bar{i}}|g_{1\bar{1},i}|^2}{\lambda_1^2} + C\mathcal{F}.$$

Then we obtain

(4.36)
$$0 \ge -\frac{3}{2} \frac{F^{ii} |g_{1\bar{1},i}|^2}{\lambda_1^2} + \frac{\delta^2}{6nK} \mathcal{F} \lambda_1^2 + F^{i\bar{i}} \eta'' |\partial_i |\partial\varphi|^2 |^2 - C \mathcal{F} \lambda_1.$$

By (4.25), it follows

$$\frac{3}{2} \frac{F^{ii}|g_{1\bar{1},i}|^2}{\lambda_1^2} = \frac{3}{2} F^{i\bar{i}} |\eta'| \partial \varphi|_i + \xi' \varphi_i|^2$$
$$\leq 2F^{i\bar{i}} \eta'^2 |\partial_i| \partial \varphi|^2|^2 + CF^{i\bar{i}} (\xi')^2 |\varphi_i|^2$$
$$\leq F^{i\bar{i}} \eta''| \partial_i| \partial \varphi|^2|^2 + C\mathcal{F}K.$$

Assume $\lambda_1 \geq K$. Returning to (4.36), we obtain,

$$0 \ge \frac{\delta^2 \lambda_1^2}{6nK} \mathcal{F} - C\lambda_1 \mathcal{F}.$$

Then we immediately deduce $\lambda_1 \leq CK$.

Using the blow-up argument [31], we have $\sup_M |\sqrt{-1}\partial\overline{\partial}\varphi| \leq C$. Using Evans-Krylov's theorem [35, 42], we have $\|\varphi\|_{C^{2,\alpha}} \leq C$. By a standard argument, we can prove the higher order estimates. Then we have the following Theorem:

Theorem 4.11. There exists a constant C such that

$$(4.37) \|\varphi\|_{C^k} \le C,$$

where C depends on $(M, \alpha), h, \omega_0, A$ and k.

5. Proofs of Theorem 1.4

In this section, we prove Theorem 1.4. If B = 0, using [32], we are done. Now we assume $B \neq 0$.

We consider the family of equations $(t \in [0, 1]),$ (5.1) $\log \frac{(\tilde{\chi} + \frac{1}{n-1}((\Delta \varphi)\alpha - \sqrt{-1}\partial \overline{\partial} \varphi) + Z + t\varphi \tilde{B})^n}{\alpha^n} = (n-1)(th + (1-t)h_0 + b_t),$

where $h_0 = \log \frac{\tilde{\chi}^n}{\alpha^n}$ and b_t are constant. Suppose φ satisfies the elliptic condition,

(5.2)
$$\tilde{\omega} = \tilde{\chi} + \frac{1}{n-1} ((\Delta \varphi)\alpha - \sqrt{-1}\partial \overline{\partial} \varphi) + Z + t\varphi \tilde{B} > 0$$

and the normalization condition

(5.3)
$$||e^{\varphi}||_{L^1} = A.$$

We define the space by

$$\mathcal{U}^{2,\alpha} := \Big\{ \phi \in C^{2,\alpha}(M) : \tilde{\omega} > 0, \ \int_M e^{\phi} \alpha^n = A \Big\}.$$

We shall prove that (5.1) is solvable for any $t \in [0, 1]$. Let I be the set

$$\{t \in [0,1] \mid \text{there exists } (\varphi,t) \in B_1 \text{ such that } \Phi(\varphi,t) = 0\}.$$

Thus, to prove Theorem 1.4, it suffices to prove that I = [0, 1]. Note that $\varphi_0 = \frac{1}{2} \ln A$ is a solution of (5.1) at t = 0. Hence, we have $0 \in I$. In the following, we prove that the set I is both open and closed.

5.1. **Openness.** Suppose that $(\varphi_{\hat{t}}, b_{\hat{t}})$ satisfies $(*)_{\hat{t}}$. In the following, we will show that when t is close to \hat{t} , there exists a pair $(\varphi_t, b_t) \in C^{\infty}(M) \times \mathbb{R}$ solving $(*)_t$.

We denote the linearized operator of (5.1) at $\varphi_{\hat{t}}$ by:

$$L_{\varphi_{\hat{i}}}(\psi) := F^{i\bar{j}}(\psi)_{i\bar{j}} + 2\operatorname{Re}(F^{i\bar{j}}W^k_{i\bar{j}}(\psi)_i) + t\tilde{F}^{i\bar{j}}\tilde{B}_{i\bar{j}}\psi.$$

Using maximum principle and $\ddot{B} \neq 0$,

(5.4)
$$\operatorname{Ker}(L_{\varphi_{\hat{x}}}) = \{0\}.$$

By the Fredholm alternative, $L_{\varphi_{\hat{t}}}$ is a bijective map from $C^{2,\alpha}(M)$ to $C^{\alpha}(M)$. Here $1 > \alpha > 0$. Note that the tangent space of $\mathcal{U}^{2,\alpha}$ at $\varphi_{\hat{t}}$ is given by

(5.5)
$$T_{\varphi_{\hat{i}}} \mathcal{U}^{2,\alpha} := \left\{ \psi \in C^{2,\alpha}(M) : \int_{M} e^{\varphi_{\hat{i}}} \psi \, \alpha^{n} = 0 \right\}$$

Let us consider the map

$$\Phi(\varphi, b) = \log \frac{(\tilde{\chi} + \frac{1}{n-1}((\Delta \varphi)\alpha - \sqrt{-1}\partial \overline{\partial}\varphi) + Z + t\varphi \tilde{B})^n}{\alpha^n} - b$$

which maps $\mathcal{U}^{2,\alpha} \times \mathbb{R}$ to $C^{\alpha}(M)$. It is clear that the linearized operator of Φ at $(\varphi_{\hat{t}}, \hat{t})$ is given by

(5.6)
$$(L_{\varphi_{\hat{t}}} - b) : T_{\varphi_{\hat{t}}} \mathcal{U}^{2,\alpha} \times \mathbb{R} \longrightarrow C^{\alpha}(M).$$

Note that for any $h \in C^{\alpha}(M)$, there exists a real function ψ on M such that

$$L_{\varphi_{\hat{t}}}(\psi) = h$$

and a function u_0 satisfying

$$(5.7) L_{\varphi_{\hat{x}}}(u_0) = 1.$$

Using maximum principle, we have $u_0 \leq 0$ and $u_0 \neq 0$. Then $\int_M u_0 e^{\varphi_{\hat{t}}} \alpha^n < 0$. We can choose b_0 such that $\psi + b_0 u_0 \in T_{\varphi_{\hat{t}}} \mathcal{U}^{2,\alpha}$, which implies

$$L_{\varphi_{\hat{i}}}(\psi + b_0 u_0) - b_0 = h$$

Hence, the map $L_{u_{\hat{t}}} - b$ is surjective.

On the other hand, suppose that there are two pairs $(\psi_1, b_1), (\psi_2, b_2) \in T_{u_i} \mathcal{U}^{2,\alpha} \times \mathbb{R}$ such that

$$L_{\varphi_{\hat{t}}}(\psi_1) - b_1 = L_{\varphi_{\hat{t}}}(\psi_2) - b_2.$$

It then follows that

$$L_{\varphi_{\hat{t}}}(\psi_1 - \psi_2) = b_1 - b_2$$

Since $L_{\varphi_{\hat{t}}}$ is bijective, by (5.7), we have $\psi_1 - \psi_2 = (b_1 - b_2)u_0$. Since

$$\int_{M} (\psi_1 - \psi_2) e^{\varphi_{\hat{t}}} \alpha^n = 0.$$

We have $b_1 = b_2$ which implies that $\psi_1 = \psi_2$. Then $L_{u_i} - b$ is injective.

Now we conclude that $L_{u_{\hat{t}}} - c$ is bijective. By the inverse function theorem, when t is close to \hat{t} , there exists a pair $(\varphi_t, b_t) \in \mathcal{U}^{2,\alpha} \times \mathbb{R}$ satisfying

$$\Phi(\varphi_t, b_t) = th + (1-t)h_0.$$

The standard elliptic theory shows that $\varphi_t \in C^{\infty}(M)$. Then I is open.

5.2. Closeness. Since $0 \in I$ and I is open, there exists $t_0 \in (0, 1]$ such that $[0, t_0) \subset I$. We need to prove $t_0 \in I$. First we show that $\{b_t\}$ is uniformly bounded.

Claim 3. There exists a constant C such that

$$(5.8) |b_t| \le C.$$

Proof. Suppose that $\varphi_t(p_t) = \max_M \varphi_t$ which implies $\sqrt{-1}\partial \overline{\partial} \varphi_t \leq 0$ at p_t . Using the equation (2.1), we have

(5.9)
$$th(p_t) + (1-t)h_0(p_t) + b_t \leq \frac{(\tilde{\chi} + \varphi_t(p_t)B)^n}{(n-1)\alpha^n} \leq \frac{(\tilde{\chi} + \min\{0, \varphi_t(p_t)\}\tilde{B})^n}{(n-1)\alpha^n}$$

Using the normalized condition $\int_M e^{\varphi} \alpha^n = A$, we have $\varphi_t(p_t) \ge \ln A$. Then we can obtain the upper bound of b_t :

(5.10)
$$b_{t} \leq \frac{(\tilde{\chi} + \varphi_{t}(p_{t})B)^{n}}{(n-1)\alpha^{n}} \leq \frac{(\tilde{\chi} + \min\{0, \ln A\}\tilde{B})^{n}}{(n-1)\alpha^{n}} - th(p_{t}) - (1-t)h_{0}(p_{t}).$$

The lower bound of b_t can be proved similarly by considering the minimum point. \Box

Now we prove the zero order estimate. In fact, we have

Proposition 5.1. There exists a constant A_0 such that if $A \leq A_0$,

(5.11)
$$\sup \varphi \le 2M_0 A^{\frac{1}{n+1}}, \ t \in [0, t_0).$$

where M_0 is the constant in Proposition 3.1.

Proof. Note that $\varphi_0 = \ln A$. Then $\sup_M \varphi_0 \leq 2M_0 A^{\frac{1}{n+1}}$, which satisfies (5.11). Thus, if (5.11) is false, there will exist $\tilde{t} \in (0, t_0)$ such that

(5.12)
$$\sup_{M} \varphi_{\tilde{t}} = 2M_0 A^{\frac{1}{n+1}}.$$

We may assume that $2M_0 A^{\frac{1}{n+1}} \leq 1$. Hence, we can apply Proposition 3.1 to $\varphi_{\tilde{t}}$ and we obtain

$$\sup_{M} \varphi_{\tilde{t}} \le M_0 A,$$

which contradicts to (5.12). This proves (5.11).

Combining Proposition 5.1 with Proposition 3.3, we have the zero order estimates and the second order estimates Theorem 4.1. Then C^{∞} a priori estimates of φ_t follows from Theorem 4.11. Combining this with the Arzelà-Ascoli theorem, I is closed.

Hence we prove the existence of solutions for the equation (1.4).

Now we prove uniqueness. Assume that we have two solutions (φ, b) and (φ', b') . Without loss of generality, we can assume there exists a point such that $\varphi(p) > \varphi'(p)$. Then we have $\inf_M(\varphi - \varphi') \leq 0$. If not, then $\varphi \geq \varphi'$. Using $\varphi(p) > \varphi'(p)$ and $\int_M e^{\varphi} \alpha^n = \int_M e^{\varphi'} \alpha^n$, it is a contraction. Note

(5.13)
$$\frac{(\tilde{\omega}_{\varphi'} + \frac{1}{n-1}((\Delta\theta)\alpha - \sqrt{-1}\partial\overline{\partial}\theta) + Z(\partial\theta) + \theta\overline{B})^n}{(\tilde{\omega}_{\varphi'})^n} = e^{(n-1)(b-b')}.$$

where $\tilde{\omega}_{\varphi'} = \tilde{\chi} + \frac{1}{n-1}((\Delta \varphi')\alpha - \sqrt{-1}\partial \overline{\partial} \varphi') + Z(\partial \varphi') + \varphi'\tilde{B}$ and $\theta = \varphi - \varphi'$. Consider the maximum point of $\varphi - \varphi'$, we can prove $b \geq b'$. Similarly, it follows that $b' \geq b$ and so b = b'. Then $F(\omega_{\varphi}) - F(\omega_{\varphi'}) = 0$, where $\omega_{\varphi} = T(\tilde{\omega}_{\varphi})$. Using the maximum principle, we can prove $\varphi = \varphi'$.

References

- Alessandrini, L., Bassanelli, G., Metric properties of manifolds bimeromorphic to compact Kähler spaces, J. Differential Geom. 37 (1993), 95–121.
- [2] Alesker, S., Shelukhin, E. On a uniform estimate for the quaternionic Calabi problem, Israel J. Math. 197 (2013), no. 1, 309–327.
- [3] Błocki, Z. On uniform estimate in Calabi-Yau theorem, Sci. China Ser. A 48 (2005), suppl., 244–247.
- Błocki, Z. On the uniform estimate in the Calabi-Yau theorem, II, Sci. China Math. 54 (2011), no. 7, 1375–1377.
- [5] Bismut, J.-M., A local index theorem for non-Kähler manifolds, Math. Ann. 284(1989), no. 4, 681–699.
- [6] Chu, J., McCleerey, N., Fully non-linear degenerate elliptic equations in complex geometry, J. Funct. Anal. 281 (2021), no. 9, Paper No. 109176, 45 pp.

- [7] Chu, J., Huang, L., Zhu, X., The Fu-Yau Equation in Higher Dimensions, Peking Math. J. 2 (2019), no. 1, 71–97.
- [8] Chu, J., Huang, L., Zhu, X., The Fu-Yau equation on compact astheno-Kähler manifolds, Adv. Math. 346 (2019), 908–945.
- Calabi, E. On Kähler manifolds with vanishing canonical class, Algebraic geometry and topology. A symposium in honor of S. Lefschetz, pp. 78–89. Princeton University Press, Princeton, N. J., 1957.
- [10] Chu, J., Tosatti, V., Weinkove, B. The Monge-Ampère equation for non-integrable almost complex structures, J. Eur. Math. Soc. (JEMS) 21 (2019), no. 7, 1949–1984.
- [11] Fernandez, M, Jordan, J., Streets, J., Non-Kähler Calabi-Yau geometry and pluriclosed flow, J. Math. Pures Appl. (9) 177 (2023), 329–367.
- [12] Fino,A., and Grantcharov,G., Properties of manifolds with skew-symmetric torsion and special holonomy, Adv. Math. 189 (2004), no. 2, 439–450.
- [13] Fu,J., Li,J, Yau, S.-T., Balanced metrics on non-Kahler Calabi-Yau threefolds, J. Differential Geom. 90 (2012) 81–130.
- [14] Fu, J., Wang, Z., Wu, D. Form-type Calabi-Yau equations, Math. Res. Lett. 17 (2010), no. 5, 887–903.
- [15] Fu, J., Wang, Z., Wu, D. Form-type Calabi-Yau equations on Kähler manifolds of nonnegative orthogonal bisectional curvature, Calc. Var. Partial Differential Equations 52 (2015), no. 1-2, 327–344.
- [16] Fu, J., Yau, S.-T., The theory of superstring with flux on non-Kähler manifolds and the complex Monge-Ampère equation, J. Differential Geom. 78 (2008), no. 3, 369–428.
- [17] George, M., Volume forms on balanced manifolds and the Calabi-Yau equation, preprint, arXiv:2406.00995.
- [18] Gray, A., Some examples of almost Hermitian manifolds, Illinois J. Math. 10 (1966), 353-366
- [19] Guan, B., Second-order estimates and regularity for fully nonlinear elliptic equations on Riemannian manifolds, Duke Math. J. 163 (2014), no. 8, 1491–1524.
- [20] Guan,B., and Nie, X., Fully nonlinear elliptic equations with gradient terms on Hermitian manifolds, Int. Math. Res. Not. IMRN 2023, no. 16, 14006–14042.
- [21] Hitchin, N., Kählerian twistor spaces, Proc. London Math. Soc. (3) 43 (1981), no. 1, 133–150.
- [22] Hou, Z., Ma, X.-N., Wu, D. A second order estimate for complex Hessian equations on a compact Kähler manifold, Math. Res. Lett. 17 (2010), no. 3, 547–561.
- [23] Jost, J., Yau, S-T, A nonlinear elliptic system for maps from Hermitian to Riemannian manifolds and rigidity theorems in Hermitian geometry. Acta Math. 170 (1993), no. 2, 221–254.
- [24] Michelsohn, M. L., On the existence of special metrics in complex geometry, Acta Math. 149, (1982), 261–295.
- [25] Phong, D. H.;, Picard, S., Zhang, X., The Fu-Yau equation with negative slope parameter, Invent. Math. 209 (2) (2017) 541–576.
- [26] Phong, D. H.;,Picard, S., Zhang, X., A flow of conformally balanced metrics with Kähler fixed points, Math. Ann. 374 (2019), no. 3–4, 2005–2040.
- [27] Phong, D. H.;, Picard, S., Zhang, X., Fu-Yau Hessian equations, J. Differential Geom. 118 (2021), no. 1, 147–187.
- [28] Popovici, D, Holomorphic deformations of balanced Calabi-Yau ∂∂-manifolds. Ann. Inst. Fourier (Grenoble) 69 (2019), no. 2, 673–728
- [29] Reid, M., The moduli space of 3-folds with K = 0 may nevertheless be irreducible, Math. Ann. 278(1987), 329-334, MR0909231, Zbl 0649.14021.
- [30] Spruck, J. Geometric aspects of the theory of fully nonlinear elliptic equations, Global theory of minimal surfaces, Amer. Math. Soc., Providence, RI, 2005, 283–309.
- [31] Székelyhidi, G. Fully non-linear elliptic equations on compact Hermitian manifolds, J. Differential Geom. 109 (2018), no. 2, 337–378.

- [32] Székelyhidi, G., Tosatti, V., Weinkove, B. Gauduchon metrics with prescribed volume form. Acta Math. 219(2017) no 1, 181–211.
- [33] Tseng, L-S., Yau, S-T., Non-Kähler Calabi-Yau manifolds, String-Math 2011, 241– 254, Proc. Sympos. Pure Math., 85, Amer. Math. Soc., Providence, RI, 2012.
- [34] Tosatti, V. Non-Kähler Calabi-Yau manifolds, Analysis, complex geometry, and mathematical physics: in honor of Duong H. Phong, 261–277, Contemp. Math., 644, Amer. Math. Soc., Providence, RI, 2015.
- [35] Tosatti, V., Wang, Y., Weinkove, B., Yang, X. C^{2,α} estimates for nonlinear elliptic equations in complex and almost complex geometry, Calc. Var. Partial Differential Equations 54 (2015), no. 1, 431–453.
- [36] Tosatti, V., Weinkove, B. The complex Monge-Ampère equation on compact Hermitian manifolds, J. Amer. Math. Soc. 23 (2010), no. 4, 1187–1195.
- [37] Tosatti, V. Weinkove, B., The Monge-Ampère equation for (n 1)-plurisubharmonic functions on a compact Kähler manifold, J. Amer. Math. Soc. 30 (2017), no. 2, 311– 346.
- [38] Yau, S.-T. On the Ricci curvature of a compact Kähler manifold and the complex Monge-Ampère equation, I, Comm. Pure Appl. Math. 31 (1978), no. 3, 339–411.
- [39] Yau, S.-T. Existence of canonical metrics in non-Kähler geometry, ICCM Not. 9 (2021), no. 1, 1–10.
- [40] Yuan, R., Fully non-linear elliptic equations on complex manifolds, preprint, arXiv:2404.12350.
- [41] Zhao, Q., Zheng, F., Strominger connection and pluriclosed metrics J. Reine Angew. Math. 796 (2023), 245–267.
- [42] Zhang, X., Zhang, X. Regularity estimates of solutions to complex Monge-Ampère equations on Hermitian manifolds, J. Funct. Anal. 260 (2011), no. 7, 2004–2026.

School of Mathematical Sciences, Xiamen University, Xiamen 361005, P. R. China

Email address: huangliding@xmu.edu.cn