

THE COHERENT STATE TRANSFORM AND AN APPLICATION TO THE HYPERBOLIC LAPLACIAN

TIMON RUBEN WEINMANN

ABSTRACT. We review some basic ideas from the theory of overfilling families and use the special case of coherent states to prove Weyl asymptotics for the euclidean Laplacian and the hyperbolic Laplacian on a domain with Dirichlet boundary conditions.

CONTENTS

1. Introduction	1
2. Coherent States	2
2.1. Overfilling Families	2
2.2. Examples	3
2.3. Definition of Coherent States	3
2.4. Weyl's Law for the Dirichlet Laplacian	4
3. The Hyperbolic Laplacian	6
3.1. Preliminaries	6
3.2. Lower Trance Bound	6
3.3. Upper Trace Bound	8
4. Asymptotic Formula	11
Acknowledgments	14
References	14

1. INTRODUCTION

Suppose $\Omega \subseteq \mathbf{R}^d$ be an open domain of finite measure. Denote by $\lambda_k, k \in \mathbf{N}$ the eigenvalues of the Laplacian on Ω with Dirichlet boundary conditions. One version of the Weyl law for the Dirichlet Laplacian reads

$$\lim_{\lambda \rightarrow \infty} \lambda^{-1-d/2} \sum_k (\lambda - \lambda_k)_+ = |\Omega| \int_{\mathbf{R}^d} (1 - \xi^2)_+ \frac{d\xi}{(2\pi)^d}.$$

There are countless proofs of this fact, see [FLW23] for instance. In this paper, we seek to establish a similar asymptotic formula for the hyperbolic Laplacian. If we consider the operator

$$H = -\partial_{x_1}^2 - e^{2x_1} \Delta_{\tilde{x}}$$

where $\Delta_{\tilde{x}} = \sum_{j=2}^d \partial_{x_j}^2$, on a bounded domain in $\Omega \subseteq \mathbf{R}^d$ and denote by λ_k its Dirichlet eigenvalues, we find

$$\lim_{\lambda \rightarrow \infty} \lambda^{-1-d/2} \sum_k (\lambda - \lambda_k)_+ = \int_{\mathbf{R}^d \times \Omega} \left(1 - \xi_1^2 - e^{2y_1} |\tilde{\xi}|^2\right)_+ \frac{d\xi dy}{(2\pi)^d}.$$

Date: June 25, 2024.

We prove this formula using the theory of coherent states. In section 2 we recall some definitions and theorems from the theory of overfilling families of functions as presented in [Ber73]. The discussion culminates in a proof of the Weyl asymptotic formula for the euclidean Laplacian. This is in fact a special case of the treatment of Weyl asymptotics for fractional Laplacians using coherent states considered in [Gei14]. Section 3 constitutes the principal part of this paper and is dedicated to proving the Weyl asymptotic formula for the hyperbolic Laplacian H using coherent states. Finally, in section 4, we discuss the asymptotic behavior of the error term in the Weyl law for the hyperbolic Laplacian and obtain the more precise formula

$$\sum_k (\lambda - \lambda_k)_+ = \lambda^{1+d/2} \int_{\mathbf{R}^d \times \Omega} (1 - \xi_1^2 - e^{2y_1} \tilde{\xi}^2)_+ \frac{d\xi dy}{(2\pi)^d} + O(\lambda^{d/2+2/3}).$$

2. COHERENT STATES

2.1. Overfilling Families. We introduce overfilling families and prove how they can be used to compute traces of operators following [Ber73].

Definition 2.1. *Let \mathcal{H} be a separable Hilbert space and let (A, μ) be a measure space. A family $e_\alpha \in \mathcal{H}$, $\alpha \in A$ of elements of \mathcal{H} indexed by A is said to be overfilling if the map $A \ni \alpha \mapsto e_\alpha \in \mathcal{H}$ is measurable and if for all $f \in \mathcal{H}$ one has*

$$\|f\|^2 = \int_A |\langle e_\alpha, f \rangle|^2 d\mu(\alpha).$$

Given an overfilling family, we obtain an isometric embedding $\mathcal{H} \rightarrow L^2(A)$ by setting

$$f(\alpha) = \langle e_\alpha, f \rangle$$

for $f \in \mathcal{H}$. Note in particular that $e_\alpha(\beta) = \langle e_\beta, e_\alpha \rangle = \overline{e_\beta(\alpha)}$. We can therefore use the same notation for the inner products in \mathcal{H} and in $L^2(A)$. Note that

$$\langle g, f \rangle = \int_A \langle g, e_\alpha \rangle \langle e_\alpha, f \rangle d\mu(\alpha)$$

for any $g, f \in \mathcal{H}$.

Theorem 2.2. *Let T be a non-negative self-adjoint trace-class operator on \mathcal{H} . Then*

$$\mathrm{tr} T = \int_A \langle e_\alpha, T e_\alpha \rangle d\mu(\alpha)$$

Proof. Indeed let ψ_k be an orthonormal basis of \mathcal{H} consisting of eigenfunctions of T ,

$$\begin{aligned} \int_A \langle e_\alpha, T e_\alpha \rangle d\mu(\alpha) &= \int_A \sum_k \langle e_\alpha, \psi_k \rangle \langle \psi_k, T e_\alpha \rangle d\mu(\alpha) \\ &= \sum_k \int_A \langle e_\alpha, \psi_k \rangle \langle \psi_k, T e_\alpha \rangle d\mu(\alpha) \\ &= \sum_k \int_A \langle T^* \psi_k, e_\alpha \rangle \langle e_\alpha, \psi_k \rangle d\mu(\alpha) \\ &= \sum_k \langle \psi_k, T \psi_k \rangle. \end{aligned}$$

Note that interchanging the sum and the integral is justified since

$$\begin{aligned} \sum_k \int_A |\langle e_\alpha, \psi_k \rangle \langle \psi_k, T e_\alpha \rangle| d\mu(\alpha) &\leq \sum_k \left(\int_A |\langle e_\alpha, \psi_k \rangle|^2 d\mu(\alpha) \right)^{1/2} \left(\int_A |\langle T \psi_k, e_\alpha \rangle|^2 d\mu(\alpha) \right)^{1/2} \\ &= \sum_k \|\psi_k\| \|T \psi_k\| = \sum_k \|T \psi_k\| < \infty \end{aligned}$$

by virtue of T being trace-class and the ψ_k being eigenfunctions. \square

2.2. Examples. Let us discuss two simple examples of overfilling families.

Example 2.3. Consider the case when $A = \mathbf{N}$ and $(e_n)_{n \in \mathbf{N}}$ is an orthonormal basis of \mathcal{H} . In this case the above procedure yields the well-known isometric isomorphism $\mathcal{H} \rightarrow \ell^2(\mathbf{N})$ and $f(n)$ is simply the n -th coefficient c_n in the Fourier expansions $f = \sum_n c_n e_n$. Given a trace-class operator T on \mathcal{H} we obtain $\text{tr} T = \sum_k T_{k,k}$, where $T_{k,l} = \langle e_k, T e_l \rangle$, as expected.

Example 2.4. Let $\Omega \subseteq \mathbf{R}^d$ be open and bounded. Let $e_\xi(x) = e^{i\xi x}$. By extending $f \in L^2(\Omega)$ by zero outside Ω , we may compute its Fourier transform. Plancherel's theorem yields

$$\int |f(x)|^2 dx = \int |\hat{f}(\xi)|^2 d\xi = \int |\langle e_\xi, f \rangle|^2 \frac{d\xi}{(2\pi)^d}.$$

One thus obtains an overfilling family with $A = \mathbf{R}^d$ and $d\mu(\xi) = (2\pi)^{-d} d\xi$.

2.3. Definition of Coherent States. The rest of this paper will deal with a class of overfilling families in L^2 -spaces constructed as follows. Let $g \in C_c^\infty(\mathbf{R}^d)$ be real-valued and such that $g(-x) = g(x)$ and $\int g(x)^2 dx = 1$. For $\xi, y \in \mathbf{R}^d$ define $e_{\xi,y}(x) = e^{i\xi x} g(x-y)$. Clearly $\|e_{\xi,y}\|_{L^2(\mathbf{R}^d)} = 1$. The space \mathbf{R}^{2d} indexing the family $(e_{\xi,y})$ shall be equipped with the measure $(2\pi)^{-d} d\xi dy$. The functions $e_{\xi,y} \in L^2(\mathbf{R}^d)$ are called coherent states.

Proposition 2.5. The family $e_{\xi,y}$, $(\xi, y) \in \mathbf{R}^{2d}$ is overfilling in $L^2(\mathbf{R}^d)$.

Proof. We need to show that

$$\int_{\mathbf{R}^{2d}} |\langle e_{\xi,y}, f \rangle|^2 \frac{d\xi dy}{(2\pi)^d} = \int_{\mathbf{R}^d} |f(x)|^2 dx.$$

Note that $\langle e_{\xi,y}, f \rangle = \int e^{-i\xi x} g(x-y) f(x) dx$ is the Fourier transform of $x \mapsto (2\pi)^{d/2} g(x-y) f(x)$. Therefore, Plancherel's theorem yields

$$\int_{\mathbf{R}^d} |\langle e_{\xi,y}, f \rangle|^2 d\xi = (2\pi)^d \int_{\mathbf{R}^d} g(x-y)^2 |f(x)|^2 dx$$

Integration with respect to y concludes the proof. \square

In the following section we will need to make use of the following technical

Lemma 2.6.

$$\langle e_{\xi,y}, -\partial_{x_j}^2 e_{\xi,y} \rangle = \xi_j^2 + \int_{\mathbf{R}^d} |\partial_{z_j} g(z)|^2 dz$$

Proof. Direct computation yields

$$\partial_{x_j} e_{\xi,y}(x) = i\xi_j e^{i\xi x} g(x-y) + e^{i\xi x} \partial_{x_j} g(x-y)$$

and

$$(1) \quad \partial_{x_j}^2 e_{\xi,y}(x) = -\xi_j^2 e^{i\xi x} g(x-y) + 2i\xi_j e^{i\xi x} \partial_{x_j} g(x-y) + e^{i\xi x} \partial_{x_j}^2 g(x-y).$$

Therefore

$$\langle e_{\xi,y}, -\partial_{x_j}^2 e_{\xi,y} \rangle = \int_{\mathbf{R}^d} \xi_j^2 (g(x-y))^2 dx + 2i\xi_j \int_{\mathbf{R}^d} \partial_{x_j} g(x-y) g(x-y) dx - \int_{\mathbf{R}^d} \partial_{x_j}^2 g(x-y) g(x-y) dx.$$

By symmetry of g , the second integral on the right vanishes and by integration by parts, the third integral is equal to $\int_{\mathbf{R}^d} (\partial_{z_j} g(z))^2 dz$. \square

Lemma 2.7. *The Laplacian acts on the functions $e_{\xi,y}$ in the following way:*

$$\langle e_{\xi,y}, -\Delta e_{\xi,y} \rangle_{L^2(\mathbf{R}^d)} = \xi^2 + \int_{\mathbf{R}} |\nabla g(z)|^2 dz.$$

Proof. Follows immediately from Lemma 2.6. \square

Instead of the entire space \mathbf{R}^d , consider now an open set $\Omega \subseteq \mathbf{R}^d$ of finite measure. For $\varepsilon > 0$ let $\Omega_\varepsilon = \{x \in \Omega: \text{dist}(x, \partial\Omega) > \varepsilon\}$. We impose on g the additional condition that $\text{supp } g \subseteq \{z \in \mathbf{R}^d: |z| < 1\}$. Note that $g^\varepsilon(z) = \varepsilon^{-d/2} g(z/\varepsilon)$ still satisfies the symmetry and normalization conditions demanded of g . We may therefore define $e_{\xi,y}^\varepsilon(x) = e^{i\xi x} g^\varepsilon(x-y)$.

Any function $f \in L^2(\Omega_\varepsilon)$ can be extended by zero to a function in $L^2(\mathbf{R}^d)$. Thus

$$\langle e_{\xi,y}^\varepsilon, f \rangle_{L^2(\mathbf{R}^d)} = \langle e_{\xi,y}^\varepsilon, f \rangle_{L^2(\Omega_\varepsilon)}.$$

The map $y \mapsto \langle e_{\xi,y}^\varepsilon, u \rangle$ vanished outside $\Omega_\varepsilon + \{z \in \mathbf{R}^d: |z| < \varepsilon\} \subseteq \Omega$, since $\text{supp } g^\varepsilon \subseteq \{z \in \mathbf{R}^d: |z| < \varepsilon\}$. Therefore

$$\int_{\mathbf{R}^d \times \mathbf{R}^d} |\langle e_{\xi,y}^\varepsilon, f \rangle_{L^2(\mathbf{R}^d)}|^2 \frac{d\xi dy}{(2\pi)^d} = \int_{\mathbf{R}^d \times \Omega} |\langle e_{\xi,y}^\varepsilon, f \rangle_{L^2(\Omega_\varepsilon)}|^2 \frac{d\xi dy}{(2\pi)^d}$$

Finally,

$$\int_{\mathbf{R}^d \times \Omega} |\langle e_{\xi,y}^\varepsilon, f \rangle_{L^2(\Omega_\varepsilon)}|^2 \frac{d\xi dy}{(2\pi)^d} = \|f\|_{L^2(\mathbf{R}^d)}^2 = \|f\|_{L^2(\Omega_\varepsilon)}^2.$$

This proves the following

Proposition 2.8. *Let $\Omega \subseteq \mathbf{R}^d$ be an open set of finite measure. Assume that g is supported in the open ball $\{z \in \mathbf{R}^d: |z| < 1\}$. Then the family $(e_{\xi,y}^\varepsilon)$, $y \in \Omega$, $\xi \in \mathbf{R}^d$ of coherent states constructed using g^ε is overfilling in $L^2(\Omega_\varepsilon)$.*

2.4. Weyl's Law for the Dirichlet Laplacian. To illustrate the utility of coherent states, we will prove the celebrated Weyl Law for the Dirichlet Laplacian on a domain in euclidean space.

Proposition 2.9. *Let λ_k , $k \in \mathbf{N}$ be the eigenvalues of the Dirichlet Laplacian in Ω counted with multiplicity. Then*

$$\liminf_{\lambda \rightarrow \infty} \lambda^{-1-d/2} \sum_k (\lambda - \lambda_k)_+ \geq |\Omega| \int_{\mathbf{R}^d} (1 - \xi^2)_+ \frac{d\xi}{(2\pi)^d}.$$

Proof. Denote by λ_k and $\tilde{\lambda}_k$ the k -th Dirichlet eigenvalue of the Laplacian with Dirichlet conditions in Ω , respectively Ω_ε . By domain monotonicity we have $\lambda_k \leq \tilde{\lambda}_k$ and consequently $(\lambda - \lambda_k)_+ \geq (\lambda - \tilde{\lambda}_k)_+$. By Theorem 2.2 and Proposition 2.8 therefore

$$\sum_k (\lambda - \lambda_k)_+ \geq \sum_k (\lambda - \tilde{\lambda}_k)_+ = \int_{\mathbf{R}^d \times \Omega} \langle e_{\xi,y}^\varepsilon, (\lambda + \Delta)_+ e_{\xi,y}^\varepsilon \rangle_{L^2(\Omega_\varepsilon)} \frac{d\xi dy}{(2\pi)^d}.$$

Clearly

$$\int_{\mathbf{R}^d \times \Omega} \langle e_{\xi,y}^\varepsilon, (\lambda + \Delta)_+ e_{\xi,y}^\varepsilon \rangle_{L^2(\Omega_\varepsilon)} \frac{d\xi dy}{(2\pi)^d} \geq \int_{\mathbf{R}^d \times \Omega_{2\varepsilon}} \langle e_{\xi,y}^\varepsilon, (\lambda + \Delta)_+ e_{\xi,y}^\varepsilon \rangle_{L^2(\Omega_\varepsilon)} \frac{d\xi dy}{(2\pi)^d}.$$

For $y \in \Omega_{2\varepsilon}$ the function $e_{\xi,y}^\varepsilon$ is supported in Ω_ε and consequently

$$\langle e_{\xi,y}^\varepsilon, (\lambda + \Delta)_+ e_{\xi,y}^\varepsilon \rangle_{L^2(\Omega_\varepsilon)} = \langle e_{\xi,y}^\varepsilon, (\lambda + \Delta)_+ e_{\xi,y}^\varepsilon \rangle_{L^2(\mathbf{R}^d)}.$$

Since $\mu \mapsto \mu_+$ is convex, applying Jensen's inequality to the spectral measure of $-\Delta$, we obtain

$$\langle e_{\xi,y}^\varepsilon, (\lambda + \Delta)_+ e_{\xi,y}^\varepsilon \rangle_{L^2(\mathbf{R}^d)} \geq \left(\langle e_{\xi,y}^\varepsilon, (\lambda + \Delta) e_{\xi,y}^\varepsilon \rangle_{L^2(\mathbf{R}^d)} \right)_+ = \left(\lambda - \xi^2 - \int_{\mathbf{R}^d} |\nabla g^\varepsilon(z)|^2 dz \right)_+.$$

Consequently,

$$\begin{aligned} \sum_k (\lambda - \lambda_k)_+ &\geq \int_{\mathbf{R}^d \times \Omega_{2\varepsilon}} \left(\lambda - \xi^2 - \int_{\mathbf{R}^d} |\nabla g^\varepsilon(z)|^2 dz \right)_+ \frac{d\xi dy}{(2\pi)^d} \\ &= \lambda^{1+d/2} \int_{\mathbf{R}^d \times \Omega_{2\varepsilon}} \left(1 - \xi^2 - \frac{1}{\lambda} \int_{\mathbf{R}^d} |\nabla g^\varepsilon(z)|^2 dz \right)_+ \frac{d\xi dy}{(2\pi)^d} \\ (2) \quad &\geq \lambda^{1+d/2} |\Omega_{2\varepsilon}| \int_{\mathbf{R}^d} \left(1 - \xi^2 - \frac{1}{\lambda} \int_{\mathbf{R}^d} |\nabla g^\varepsilon(z)|^2 dz \right)_+ \frac{d\xi}{(2\pi)^d}. \end{aligned}$$

Finally we obtain

$$\liminf_{\lambda \rightarrow \infty} \lambda^{-1-d/2} \sum_k (\lambda - \lambda_k)_+ \geq |\Omega_{2\varepsilon}| \int_{\mathbf{R}^d} (1 - \xi^2)_+ \frac{d\xi}{(2\pi)^d}$$

by dominated convergence. Letting $\varepsilon \rightarrow 0$ concludes the proof. \square

We now seek to prove an upper bound on $\sum_k (\lambda - \lambda_k)_+$. This result is due to Li and Yau [LY83] and our proof follows [FLW23].

Proposition 2.10.

$$\sum_k (\lambda - \lambda_k)_+ \leq |\Omega| \int_{\mathbf{R}^d} (\lambda - \xi^2)_+ \frac{d\xi}{(2\pi)^d}$$

Proof. Let $(\psi_k)_k \subseteq L^2(\Omega)$ be an orthonormal system of eigenfunctions corresponding to $(\lambda_k)_k$. Assume that $\psi_k \in H_0^1(\Omega)$. Then, since every ψ_k can be extended by zero to a function $H^1(\mathbf{R}^d)$, we have

$$\lambda_k = \int_{\Omega} |\nabla \psi_k(x)|^2 dx = \int_{\mathbf{R}^d} \xi^2 |\hat{\psi}_k(\xi)|^2 d\xi.$$

Therefore

$$(\lambda - \lambda_k)_+ = \left(\int_{\mathbf{R}^d} (\lambda - \xi^2) |\hat{\psi}_k(\xi)|^2 d\xi \right)_+ \leq \int_{\mathbf{R}^d} (\lambda - \xi^2)_+ |\hat{\psi}_k(\xi)|^2 d\xi,$$

by Jensen's inequality, since $|\hat{\psi}_k(\xi)|^2 d\xi$ is a probability measure.

$$\begin{aligned} \sum_k (\lambda - \lambda_k)_+ &\leq \sum_{\lambda_k < \lambda} \int_{\mathbf{R}^d} (\lambda - \xi^2)_+ |\hat{\psi}_k(\xi)|^2 d\xi \\ &= \int_{\mathbf{R}^d} (\lambda - \xi^2)_+ \sum_{\lambda_k < \lambda} |\hat{\psi}_k(\xi)|^2 d\xi \\ &\leq \int_{\mathbf{R}^d} (\lambda - \xi^2)_+ \sum_k |\hat{\psi}_k(\xi)|^2 d\xi \end{aligned}$$

Note that

$$\hat{\psi}_k(\xi) = \frac{1}{(2\pi)^{d/2}} \int_{\mathbf{R}^d} e^{-ix\xi} \psi_k(x) dx = \frac{1}{(2\pi)^{d/2}} \int_{\Omega} e^{-ix\xi} \psi_k(x) dx = \langle (2\pi)^{-d/2} e^{i\xi \cdot}, \psi_k \rangle_{L^2(\Omega)}.$$

Since the ψ_k form an orthonormal basis of $L^2(\Omega)$, we obtain

$$\sum_k |\hat{\psi}_k(\xi)|^2 = \sum_k | \langle (2\pi)^{-d/2} e^{i\xi \cdot}, \psi_k \rangle_{L^2(\Omega)} |^2 = \left\| (2\pi)^{-d/2} e^{i\xi \cdot} \right\|_{L^2(\Omega)}^2 = (2\pi)^{-d} |\Omega|.$$

This concludes the argument. \square

Corollary 2.11. *For the eigenvalues λ_k of the Dirichlet Laplacian on a domain Ω of finite measure, the following asymptotic formula holds*

$$\lim_{\lambda \rightarrow \infty} \lambda^{-1-d/2} \sum_k (\lambda - \lambda_k)_+ = |\Omega| \int_{\mathbf{R}^d} (1 - \xi^2)_+ \frac{d\xi}{(2\pi)^d}.$$

3. THE HYPERBOLIC LAPLACIAN

3.1. Preliminaries. On a Riemannian manifold (M, g) , the Laplace–Beltrami operator is locally given by

$$-\Delta_g = -\frac{1}{\sqrt{g}} \partial_{x^i} (\sqrt{g} g^{ij} \partial_{x^j}),$$

whereas the canonical Riemannian volume is given by

$$dV_g = \sqrt{g} dx^1 dx^2 \dots dx^d.$$

Consider now the hyperbolic space in the half-plane model

$$M = \mathbf{H}^d = \{(y, x) \in \mathbf{R} \times \mathbf{R}^{d-1} : y > 0\}, \quad g = \frac{1}{y^2} (dy^2 + dx^2).$$

The Laplace–Beltrami operator in this case is

$$-\Delta_g = -y^d \partial_y (y^{2-d} \partial_y) - y^2 \Delta_x,$$

where $\Delta_x = \sum_{j=1}^{d-1} \partial_{x_j}^2$. Performing the change of coordinates $y = e^t$, one obtains a unitary operator $L^2(\mathbf{H}^d, y^{-2} dx dy) \rightarrow L^2(\mathbf{R}^d, dx dt)$ and $-\Delta_g$ acts on $L^2(\mathbf{R}^d, dx dt)$ as $-\partial_t^2 - e^{2t} \Delta_x + (d-1)^2/4$. In [ILW24] this correspondence was used to extend a class of Lieb–Thirring inequalities from \mathbf{R}^d to \mathbf{H}^d . In the rest of this section we adopt the notation $\tilde{x} = (x_2, \dots, x_d)$ for $x \in \mathbf{R}^d$.

3.2. Lower Trance Bound. Consider the operator

$$H = -\partial_{x_1}^2 - e^{2x_1} \Delta_{\tilde{x}}$$

where $\Delta_{\tilde{x}} = \sum_{j=2}^d \partial_{x_j}^2$ on the space $L^2(\mathbf{R}^d)$. We are interested in its action on a coherent state of the form $e_{\xi, y}(x) = e^{i\xi x} g(x - y)$.

Lemma 3.1.

$$\langle e_{\xi, y}, H e_{\xi, y} \rangle_{L^2(\mathbf{R}^d)} = \xi_1^2 + e^{2y_1} |\tilde{\xi}|^2 \int_{\mathbf{R}^d} e^{2z_1} g(z)^2 dz + e^{2y_1} \int_{\mathbf{R}^d} e^{2z_1} |\nabla_{\tilde{z}} g(z)|^2 dz + \int_{\mathbf{R}^d} (\partial_{z_1} g(z))^2 dz$$

Proof. The contribution of $-\partial_{x_1}^2$ was already determined in 2.6. We compute $\langle e_{\xi, y}, -e^{2x_1} \partial_{x_j}^2 e_{\xi, y} \rangle_{L^2(\mathbf{R}^d)}$ for $j > 1$. Equation 1 in the proof of Lemma 2.6 implies

$$\begin{aligned} \langle e_{\xi, y}, -e^{2x_1} \partial_{x_j}^2 e_{\xi, y} \rangle &= \xi_j^2 \int_{\mathbf{R}^d} e^{2x_1} (g(x - y))^2 dx + 2i\xi_j \int_{\mathbf{R}^d} e^{2x_1} \partial_{x_j} g(x - y) g(x - y) dx \\ &\quad - \int_{\mathbf{R}^d} e^{2x_1} \partial_{x_j}^2 g(x - y) g(x - y) dx. \end{aligned}$$

The first term on the right equals

$$e^{2y_1} \xi_j^2 \int_{\mathbf{R}^d} e^{2(x_1-y_1)} (g(x-y))^2 dx = e^{2y_1} \xi_j^2 \int_{\mathbf{R}^d} e^{2z_1} (g(z))^2 dz.$$

For the second integral we find

$$\int_{\mathbf{R}^d} e^{2x_1} \partial_{x_j} g(x-y) g(x-y) dx = \int_{\mathbf{R}} e^{2x_1} \int_{\mathbf{R}^{d-1}} \partial_{x_j} g(x-y) g(x-y) d\tilde{x} dx_1,$$

where the inner integral vanishes due to symmetry of g . Finally, for the third integral we find, again by splitting the integral, that

$$\begin{aligned} \int_{\mathbf{R}^d} e^{2x_1} \partial_{x_j}^2 g(x-y) g(x-y) dx &= -e^{2y_1} \int_{\mathbf{R}} e^{2(x_1-y_1)} \int_{\mathbf{R}^{d-1}} \partial_{x_j}^2 g(x-y) g(x-y) d\tilde{x} dx_1 \\ &= e^{2y_1} \int_{\mathbf{R}} e^{2(x_1-y_1)} \int_{\mathbf{R}^{d-1}} (\partial_{x_j} g(x-y))^2 d\tilde{x} dx_1 \\ &= e^{2y_1} \int_{\mathbf{R}^d} e^{2z_1} (\partial_{z_j} g(z))^2 dz \end{aligned}$$

Summing over $j = 2, \dots, d$ completes the argument. \square

Theorem 3.2. *Let $\Omega \subseteq \mathbf{R}^d$ be a bounded open set and let λ_k , $k \in \mathbf{N}$ be the eigenvalues of the operator H on Ω with Dirichlet boundary conditions. Then*

$$\liminf_{\lambda \rightarrow \infty} \lambda^{-1-d/2} \sum_k (\lambda - \lambda_k)_+ \geq \int_{\mathbf{R}^d \times \Omega} \left(1 - \xi_1^2 - e^{2y_1} |\tilde{\xi}|^2\right)_+ \frac{d\xi dy}{(2\pi)^d}.$$

Proof. Let us again consider a g supported in the unit ball and construct $e_{\xi,y}^\varepsilon$ accordingly. By the same argument as for the Dirichlet Laplacian in the proof of Proposition 2.9, we have

$$\sum_k (\lambda - \lambda_k)_+ \geq \int_{\mathbf{R}^d \times \Omega_{2\varepsilon}} \left(\langle e_{\xi,y}^\varepsilon, (\lambda - H)e_{\xi,y}^\varepsilon \rangle_{L^2(\mathbf{R}^d)}\right)_+ \frac{d\xi dy}{(2\pi)^d}.$$

Using the expression obtained for the symbol of H in 3.1 then gives

$$(3) \quad \sum_k (\lambda - \lambda_k)_+ \geq \int_{\mathbf{R}^d \times \Omega_{2\varepsilon}} \left(\lambda - \xi_1^2 - e^{2y_1} |\tilde{\xi}|^2 c_3^\varepsilon - e^{2y_1} c_2^\varepsilon - c_1^\varepsilon\right)_+ \frac{d\xi dy}{(2\pi)^d},$$

where

$$\begin{aligned} c_1^\varepsilon &= \int_{\mathbf{R}^d} (\partial_{z_1} g^\varepsilon(z))^2 dz \\ c_2^\varepsilon &= \int_{\mathbf{R}^d} e^{2z_1} |\nabla_{\tilde{z}} g^\varepsilon(z)|^2 dz \\ c_3^\varepsilon &= \int_{\mathbf{R}^d} e^{2z_1} g^\varepsilon(z)^2 dz. \end{aligned}$$

Note that since $g(z)^2$ is a mollifier, we have $c_3^\varepsilon \rightarrow 1$ as $\varepsilon \rightarrow 0$. After a change of coordinates, we obtain

$$(4) \quad \sum_k (\lambda - \lambda_k)_+ \geq \lambda^{1+d/2} \int_{\mathbf{R}^d \times \Omega_{2\varepsilon}} \left(1 - \xi_1^2 - e^{2y_1} |\tilde{\xi}|^2 c_3^\varepsilon - \frac{1}{\lambda} e^{2y_1} c_2^\varepsilon - \frac{1}{\lambda} c_1^\varepsilon\right)_+ \frac{d\xi dy}{(2\pi)^d}.$$

Dominated convergence yields

$$\liminf_{\lambda \rightarrow \infty} \lambda^{-1-d/2} \sum_k (\lambda - \lambda_k)_+ \geq \int_{\mathbf{R}^d \times \Omega_{2\varepsilon}} \left(1 - \xi_1^2 - e^{2y_1} |\tilde{\xi}|^2 c_3^\varepsilon\right)_+ \frac{d\xi dy}{(2\pi)^d}.$$

Letting $\varepsilon \rightarrow 0$ and using dominated convergence concludes the argument. \square

3.3. Upper Trace Bound. Before proving an upper bound on the trace of the hyperbolic Laplacian, we need to consider some further technicalities concerning coherent states. Consider again the general situation where $g \in C_c^\infty(\mathbf{R}^d)$ is real-valued and such that $g(-x) = g(x)$ and $\int g(x)^2 dx = 1$. As before, define $e_{\xi,y}(x) = e^{i\xi x} g(x-y)$. The operator $\Phi: L^2(\mathbf{R}^d, dx) \rightarrow L^2(\mathbf{R}^{2d}, (2\pi)^{-d} dy d\xi)$ given by $(\Phi\psi)(y, \xi) = \langle e_{\xi,y}, \psi \rangle_{L^2(\mathbf{R}^d)}$ is called the coherent state transform. Since $(e_{\xi,y})$ is overfilling, Φ is an isometry, i.e. $\Phi^* \Phi = I_{L^2(\mathbf{R}^d)}$. Given $\psi \in L^2(\mathbf{R}^d, dx)$ and $\Psi \in \text{Im } \Phi$, we have

$$\begin{aligned} \langle \psi, \Phi^* \Psi \rangle &= \langle \Phi\psi, \Psi \rangle = \int_{\mathbf{R}^{2d}} \int_{\mathbf{R}^d} \overline{e_{\xi,y}(x)} \psi(x) dx \Psi(\xi, y) \frac{d\xi dy}{(2\pi)^d} \\ &= \int_{\mathbf{R}^d} \int_{\mathbf{R}^{2d}} e_{\xi,y}(x) \Psi(\xi, y) \frac{d\xi dy}{(2\pi)^d} \overline{\psi(x)} dx \end{aligned}$$

Consequently $\Phi^* \Psi(x) = \int_{\mathbf{R}^{2d}} e_{\xi,y}(x) \Psi(\xi, y) \frac{d\xi dy}{(2\pi)^d}$.

In this section's theorem we shall make use of the two following lemmas

Lemma 3.3. *Let $\psi \in C_c^\infty(\mathbf{R}^d)$. Then*

$$\langle \psi, \Phi^* \xi_1^2 \Phi \psi \rangle = \|\partial_{x_1} \psi\|^2 + \int_{\mathbf{R}^d} (\partial_{z_1} g(z))^2 dz \|\psi\|^2.$$

Proof. Indeed

$$\begin{aligned} \langle \psi, \Phi^* \xi_1^2 \Phi \psi \rangle &= \int_{\mathbf{R}^{4d}} e^{i\xi x} g(x-y) \xi_1^2 e^{-i\xi w} g(w-y) \psi(w) \overline{\psi(x)} \frac{dw d\xi dy dx}{(2\pi)^d} \\ &= \int_{\mathbf{R}^{4d}} \partial_{x_1} (e^{i\xi x}) \partial_{w_1} (e^{-i\xi w}) g(x-y) g(w-y) \psi(w) \overline{\psi(x)} \frac{dw d\xi dy dx}{(2\pi)^d} \\ (\text{Int. by parts}) &= - \int_{\mathbf{R}^{4d}} e^{i\xi x} \partial_{x_1} (g(x-y) \overline{\psi(x)}) \partial_{w_1} (e^{-i\xi w}) g(w-y) \psi(w) \frac{dw d\xi dy dx}{(2\pi)^d} \\ (\text{Int. by parts}) &= \int_{\mathbf{R}^{4d}} e^{i\xi x} \partial_{x_1} (g(x-y) \overline{\psi(x)}) e^{-i\xi w} \partial_{w_1} (g(w-y) \psi(w)) \frac{dw d\xi dy dx}{(2\pi)^d} \\ (\star) &= \int_{\mathbf{R}^{2d}} |\partial_{x_1} (g(x-y) \psi(x))|^2 dy dx \end{aligned}$$

In (\star) we used that $\int_{\mathbf{R}^d} e^{i\xi(x-w)} d\xi = (2\pi)^d \delta(x-w)$ in the weak sense as distributions. Note that $|\partial_{x_1} (g(x-y) \psi(x))|^2 = |\partial_{x_1} g(x-y) \psi(x)|^2 + |g(x-y) \partial_{x_1} \psi(x)|^2 + 2 \text{Re } \overline{\partial_{x_1} g(x-y) \psi(x)} g(x-y) \partial_{x_1} \psi(x)$. Since g is even, $\partial_1 g$ is odd and therefore

$$\int_{\mathbf{R}^d} \partial_{x_1} g(x-y) g(x-y) dy = \int_{\mathbf{R}^d} \partial_{z_1} g(z) g(z) dz = 0.$$

Consequently,

$$\begin{aligned} \langle \psi, \Phi^* \xi_1^2 \Phi \psi \rangle &= \int_{\mathbf{R}^{2d}} (|\partial_{x_1} g(x-y) \psi(x)|^2 + |g(x-y) \partial_{x_1} \psi(x)|^2) dy dx \\ &= \int_{\mathbf{R}^d} (\partial_{z_1} g(z))^2 dz \|\psi\|^2 + \int_{\mathbf{R}^d} g(z)^2 dz \|\partial_{x_1} \psi\|^2 \\ &= \int_{\mathbf{R}^d} (\partial_{z_1} g(z))^2 dz \|\psi\|^2 + \|\partial_{x_1} \psi\|^2 \end{aligned}$$

□

Lemma 3.4. *Assume that the function in the definition of coherent states can be decomposed as $g(z) = g_1(z_1)\tilde{g}(\tilde{z})$ where g_1 and \tilde{g} are non-negative-valued, symmetric, compactly supported and such that $\int_{\mathbf{R}} g_1(z_1)^2 dz_1 = \int_{\mathbf{R}^{d-1}} \tilde{g}(\tilde{z})^2 d\tilde{z} = 1$. Let $\psi \in C_c^\infty(\mathbf{R}^d)$. Then*

$$\langle \psi, \Phi^* e^{2y_1} |\tilde{\xi}|^2 \Phi \psi \rangle = \int_{\mathbf{R}^d} e^{2z_1} |\tilde{\nabla} g(z)|^2 dz \|e^{x_1} \psi\|^2 + \int_{\mathbf{R}^d} e^{2z_1} g(z)^2 dz \|e^{x_1} \nabla_{\tilde{x}} \psi\|^2$$

Proof. Similar to the proof above we find

$$\begin{aligned} \langle \psi, \Phi^* e^{2y_1} |\tilde{\xi}|^2 \Phi \psi \rangle &= \int_{\mathbf{R}^{4d}} e^{i\xi x} g(x-y) e^{2y_1} e^{2y_1} |\tilde{\xi}|^2 e^{-i\xi w} g(w-y) \psi(w) \overline{\psi(x)} \frac{dw d\xi dy dx}{(2\pi)^d} \\ &= \int_{\mathbf{R}^{4d}} e^{2y_1} \nabla_{\tilde{x}}(e^{i\xi x}) \nabla_{\tilde{w}}(e^{-i\xi w}) g(x-y) g(w-y) \psi(w) \overline{\psi(x)} \frac{dw d\xi dy dx}{(2\pi)^d} \\ &= - \int_{\mathbf{R}^{4d}} e^{2y_1} e^{i\xi x} \nabla_{\tilde{x}}(g(x-y) \overline{\psi(x)}) \nabla_{\tilde{w}}(e^{-i\xi w}) g(w-y) \psi(w) \frac{dw d\xi dy dx}{(2\pi)^d} \\ &= \int_{\mathbf{R}^{4d}} e^{2y_1} e^{i\xi x} \nabla_{\tilde{x}}(g(x-y) \overline{\psi(x)}) e^{-i\xi w} \nabla_{\tilde{w}}(g(w-y) \psi(w)) \frac{dw d\xi dy dx}{(2\pi)^d} \\ &= \int_{\mathbf{R}^{2d}} e^{2y_1} |\nabla_{\tilde{x}}(g(x-y) \psi(x))|^2 dy dx. \end{aligned}$$

Note that

$$\begin{aligned} |\nabla_{\tilde{x}}(g(x-y) \psi(x))|^2 &= |\nabla_{\tilde{x}} g(x-y)|^2 |\psi(x)|^2 + |g(x-y)|^2 |\nabla_{\tilde{x}} \psi(x)|^2 \\ &\quad + \sum_{j=2}^d 2 \operatorname{Re} \overline{\partial_{x_j} g(x-y) \psi(x)} g(x-y) \partial_{x_j} \psi(x), \end{aligned}$$

and that

$$\int_{\mathbf{R}^d} e^{2y_1} \partial_{x_j} g(x-y) g(x-y) dy = \int_{\mathbf{R}} e^{2y_1} g_1(x_1 - y_1)^2 dy_1 \int_{\mathbf{R}^{d-1}} \partial_{x_j} \tilde{g}(\tilde{x} - \tilde{y}) \tilde{g}(\tilde{x} - \tilde{y}) d\tilde{y},$$

where the second integral on the right vanishes due to the symmetry of \tilde{g} . Finally

$$\begin{aligned} \int_{\mathbf{R}^{2d}} e^{2y_1} |\nabla_{\tilde{x}} g(x-y)|^2 |\psi(x)|^2 dx dy &= \int_{\mathbf{R}^{2d}} e^{2(y_1-x_1)} |\nabla_{\tilde{x}} g(x-y)|^2 e^{2x_1} |\psi(x)|^2 dx dy \\ &= \int_{\mathbf{R}^d} e^{2z} |\nabla_{\tilde{z}} g(z)|^2 dz \int_{\mathbf{R}^d} e^{2x_1} |\psi(x)|^2 dx \end{aligned}$$

and

$$\begin{aligned} \int_{\mathbf{R}^{2d}} e^{2y_1} |g(x-y)|^2 |\nabla_{\tilde{x}} \psi(x)|^2 dx dy &= \int_{\mathbf{R}^{2d}} e^{2(y_1-x_1)} |g(x-y)|^2 e^{2x_1} |\nabla_{\tilde{x}} \psi(x)|^2 dx dy \\ &= \int_{\mathbf{R}^d} e^{2z_1} |g(z)|^2 dz \int_{\mathbf{R}^d} e^{2x_1} |\nabla_{\tilde{x}} \psi(x)|^2 dx. \end{aligned}$$

This concludes the proof. \square

We are now ready to state the main theorem of this subsection.

Theorem 3.5. *Let $\Omega \subseteq \mathbf{R}^d$ be a bounded open set and let λ_k , $k \in \mathbf{N}$ be the eigenvalues of the operator H on Ω with Dirichlet boundary conditions. Then*

$$\limsup_{\lambda \rightarrow \infty} \lambda^{-1-d/2} \sum_k (\lambda - \lambda_k)_+ \leq \int_{\mathbf{R}^d \times \Omega} \left(1 - \xi_1^2 - e^{2y_1} |\tilde{\xi}|^2 \right)_+ \frac{d\xi dy}{(2\pi)^d}$$

Proof. Assume that g is supported in the unit ball and that it can be decomposed as $g(z) = g_1(z_1)\tilde{g}(\tilde{z})$, where $g_1, \tilde{g} \geq 0$ are symmetric, compactly supported and satisfy $\int_{\mathbf{R}} g_1(z_1)^2 dz_1 = \int_{\mathbf{R}^{d-1}} \tilde{g}(\tilde{z})^2 d\tilde{z} = 1$. Define $g^\varepsilon(z) = \varepsilon^{-d/2}g(z/\varepsilon)$. Denote by Ω^ε the set $\{x \in \mathbf{R}^d: \text{dist}(x, \Omega) < \varepsilon\}$. As before, let $\lambda_k, k \in \mathbf{N}$ be the eigenvalues of H . Since H is clearly elliptic, we can find according eigenfunctions $\psi_k \in C_c^\infty(\Omega)$ such that $(\psi_k)_k$ is an orthonormal basis of $L^2(\Omega)$. It is

$$\lambda_k = \langle \psi_k, H\psi_k \rangle = \|\partial_{x_1}\psi_k\|^2 + \|e^{x_1}\nabla_{\tilde{x}}\psi_k\|^2.$$

For brevity we denote, again,

$$\begin{aligned} c_1^\varepsilon &= \int_{\mathbf{R}^d} (\partial_{z_1}g^\varepsilon(z))^2 dz \\ c_2^\varepsilon &= \int_{\mathbf{R}^d} e^{2z_1}|\nabla_{\tilde{z}}g^\varepsilon(z)|^2 dz \\ c_3^\varepsilon &= \int_{\mathbf{R}^d} e^{2z_1}g^\varepsilon(z)^2 dz. \end{aligned}$$

Denoting $R = \sup\{y_1 \mid y \in \Omega\}$, we find

$$\begin{aligned} (\lambda - \lambda_k)_+ &= \left(\lambda - \langle \psi_k, \Phi^* \xi_1^2 \Phi \psi_k \rangle + c_1^\varepsilon - (c_3^\varepsilon)^{-1} \langle \psi_k, \Phi^* e^{2y_1} \tilde{\xi}^2 \Phi \psi_k \rangle + (c_3^\varepsilon)^{-1} c_2^\varepsilon \|e^{x_1}\psi_k\|^2 \right)_+ \\ &\leq \left(\lambda - \langle \psi_k, \Phi^* \xi_1^2 \Phi \psi_k \rangle + c_1^\varepsilon - (c_3^\varepsilon)^{-1} \langle \psi_k, \Phi^* e^{2y_1} \tilde{\xi}^2 \Phi \psi_k \rangle + (c_3^\varepsilon)^{-1} c_2^\varepsilon e^{2R} \right)_+ \\ &= \left(\langle \Phi \psi_k, (\lambda - \xi_1^2 - (c_3^\varepsilon)^{-1} e^{2y_1} \tilde{\xi}^2 + c_1^\varepsilon + (c_3^\varepsilon)^{-1} c_2^\varepsilon e^{2R}) \Phi \psi_k \rangle \right)_+ \\ &\leq \langle \Phi \psi_k, (\lambda - \xi_1^2 - (c_3^\varepsilon)^{-1} e^{2y_1} \tilde{\xi}^2 + c_1^\varepsilon + (c_3^\varepsilon)^{-1} c_2^\varepsilon e^{2R})_+ \Phi \psi_k \rangle \end{aligned}$$

where the last inequality is due to Jensen's inequality. Summing over k then yields

$$\sum_k (\lambda - \lambda_k)_+ \leq \int_{\mathbf{R}^{2d}} (\lambda - \xi_1^2 - (c_3^\varepsilon)^{-1} e^{2y_1} \tilde{\xi}^2 + c_1^\varepsilon + (c_3^\varepsilon)^{-1} c_2^\varepsilon e^{2R})_+ \sum_k |\Phi \psi_k(\xi, y)|^2 \frac{d\xi dy}{(2\pi)^d}.$$

Since (ψ_k) is an orthonormal basis, we obtain

$$\sum_k |\Phi \psi_k(\xi, y)|^2 = \sum_k |\langle e_{\xi, y}, \psi_k \rangle|^2 = \|e_{\xi, y}\|_{L^2(\Omega)}^2 \leq \|e_{\xi, y}\|_{L^2(\mathbf{R}^d)}^2 = 1$$

Since the map $y \mapsto \Phi \psi(y, \xi)$ is supported in Ω^ε , we find

$$\sum_k |\Phi \psi_k(\xi, y)|^2 \leq \chi_{\Omega^\varepsilon}(y).$$

Hence

$$\begin{aligned} \sum_k (\lambda - \lambda_k)_+ &\leq \int_{\mathbf{R}^d \times \Omega^\varepsilon} (\lambda - \xi_1^2 - (c_3^\varepsilon)^{-1} e^{2y_1} \tilde{\xi}^2 + c_1^\varepsilon + (c_3^\varepsilon)^{-1} c_2^\varepsilon e^{2R})_+ \frac{d\xi dy}{(2\pi)^d} \\ &= \lambda^{1+d/2} \int_{\mathbf{R}^d \times \Omega^\varepsilon} (1 - \xi_1^2 - (c_3^\varepsilon)^{-1} e^{2y_1} \tilde{\xi}^2 + c_1^\varepsilon \lambda^{-1} + (c_3^\varepsilon)^{-1} c_2^\varepsilon e^{2R} \lambda^{-1})_+ \frac{d\xi dy}{(2\pi)^d}. \end{aligned}$$

Consequently

$$\limsup_{\lambda \rightarrow \infty} \lambda^{-1-d/2} \sum_k (\lambda - \lambda_k)_+ \leq \int_{\mathbf{R}^d \times \Omega^\varepsilon} (1 - \xi_1^2 - (c_3^\varepsilon)^{-1} e^{2y_1} \tilde{\xi}^2)_+ \frac{d\xi dy}{(2\pi)^d}.$$

Letting first $\varepsilon \rightarrow 0$ finishes the proof. \square

Corollary 3.6. *Let $\Omega \subseteq \mathbf{R}^d$ be a bounded open set and let λ_k , $k \in \mathbf{N}$ be the eigenvalues of the operator H on Ω with Dirichlet boundary conditions. Then*

$$\lim_{\lambda \rightarrow \infty} \lambda^{-1-d/2} \sum_k (\lambda - \lambda_k)_+ = \int_{\mathbf{R}^d \times \Omega} \left(1 - \xi_1^2 - e^{2y_1} |\tilde{\xi}|^2\right)_+ \frac{d\xi dy}{(2\pi)^d}$$

4. ASYMPTOTIC FORMULA

Recall that for the euclidean Laplacian with Dirichlet boundary conditions we obtained inequality 2 reading:

$$\sum_k (\lambda - \lambda_k)_+ \geq \lambda^{1+d/2} |\Omega_{2\varepsilon}| \int_{\mathbf{R}^d} \left(1 - \xi^2 - \frac{1}{\lambda} \int_{\mathbf{R}^d} |\nabla g^\varepsilon(z)|^2 dz\right)_+ \frac{d\xi}{(2\pi)^d}.$$

Denoting $B = \{\xi \in \mathbf{R}^d \mid \xi^2 \leq 1\}$, we can clearly make the estimate

$$\begin{aligned} \sum_k (\lambda - \lambda_k)_+ &\geq \lambda^{1+d/2} |\Omega_{2\varepsilon}| \int_B \left(1 - \xi^2 - \frac{1}{\lambda} \int_{\mathbf{R}^d} |\nabla g^\varepsilon(z)|^2 dz\right)_+ \frac{d\xi}{(2\pi)^d} \\ &\geq \lambda^{1+d/2} |\Omega_{2\varepsilon}| \int_B (1 - \xi^2)_+ \frac{d\xi}{(2\pi)^d} - \lambda^{d/2} \frac{|\Omega_{2\varepsilon}| |B|}{(2\pi)^d} \int_{\mathbf{R}^d} |\nabla g^\varepsilon(z)|^2 dz \\ &\geq \lambda^{1+d/2} |\Omega_{2\varepsilon}| \int_B (1 - \xi^2)_+ \frac{d\xi}{(2\pi)^d} - \lambda^{d/2} \frac{|\Omega| |B|}{(2\pi)^d} \int_{\mathbf{R}^d} |\nabla g^\varepsilon(z)|^2 dz \\ &\geq \lambda^{1+d/2} |\Omega| \int_B (1 - \xi^2)_+ \frac{d\xi}{(2\pi)^d} - R(\lambda, \varepsilon) \end{aligned}$$

where

$$R(\lambda, \varepsilon) = \lambda^{1+d/2} (|\Omega| - |\Omega_{2\varepsilon}|) \int_B (1 - \xi^2)_+ \frac{d\xi}{(2\pi)^d} + \lambda^{d/2} \frac{|\Omega| |B|}{(2\pi)^d} \int_{\mathbf{R}^d} |\nabla g^\varepsilon(z)|^2 dz$$

Note that

$$\int_{\mathbf{R}^d} |\nabla g^\varepsilon(z)|^2 dz = \varepsilon^{-2} \int_{\mathbf{R}^d} |\nabla g(z)|^2 dz.$$

Let $\varepsilon = \lambda^{-\alpha}$ with $\alpha > 0$. Then there are constants $L, C \geq 0$ such that

$$R(\lambda, \varepsilon) = L\lambda^{1+d/2} (|\Omega| - |\Omega_{2\varepsilon}|) + C\lambda^{d/2+2\alpha}.$$

Suppose that $|\Omega| - |\Omega_{2\varepsilon}| = O(\varepsilon)$. This is in particular the case when the inner Minkowski content of Ω exists which was proven to be the case when the boundary of Ω is Lipschitz-regular and Ω itself is bounded in [ACV08]. Then,

$$R(\lambda, \lambda^{-\alpha}) = C\lambda^{d/2+2\alpha} + O(\lambda^{1+d/2-\alpha}).$$

Minimizing $\max\{d/2 + 2\alpha, 1 + d/2 - \alpha\}$, we find $\alpha = 1/3$. Consequently

$$\sum_k (\lambda - \lambda_k)_+ \geq \lambda^{1+d/2} |\Omega| \int_B (1 - \xi^2)_+ \frac{d\xi}{(2\pi)^d} + O(\lambda^{d/2+2/3}).$$

Combining this with the upper estimate 2.10 we obtain the following

Theorem 4.1. *Let Ω be a bounded domain whose boundary is Lipschitz-regular. Then the eigenvalues λ_k of the Dirichlet Laplacian on Ω satisfy*

$$\sum_k (\lambda - \lambda_k)_+ = \lambda^{1+d/2} |\Omega| \int_B (1 - \xi^2)_+ \frac{d\xi}{(2\pi)^d} + O(\lambda^{d/2+2/3}).$$

This is a significantly weaker result than the one obtained in [FL20] where the following two-term asymptotic formula was proven for bounded Lipschitz domains:

$$\sum_k (\lambda - \lambda_k)_+ = L_d |\Omega| \lambda^{d/2+1} - \frac{L_{d-1}}{4} \mathcal{H}^{d-1}(\partial\Omega) \lambda^{(d-1)/2+1} + o(\lambda^{(d-1)/2+1}),$$

for constants $L_d, L_{d-1} \geq 0$.

We intend to prove a similar asymptotic formula for the hyperbolic Laplacian H .

Lemma 4.2. *Assume that $\Omega \subseteq \mathbf{R}^d$ is open and bounded and that $\partial\Omega$ is Lipschitz-regular. Then the trace of the hyperbolic Laplacian on Ω with Dirichlet boundary conditions satisfies*

$$\sum_k (\lambda - \lambda_k)_+ \geq \lambda^{1+d/2} \int_{\mathbf{R}^d \times \Omega} \left(\lambda - \xi_1^2 - e^{2y_1} |\tilde{\xi}|^2 \right)_+ \frac{d\xi dy}{(2\pi)^d} + O(\lambda^{d/2+2/3}).$$

Proof. Recall that in 3 we found

$$\sum_k (\lambda - \lambda_k)_+ \geq \int_{\mathbf{R}^d \times \Omega_{2\varepsilon}} \left(\lambda - \xi_1^2 - e^{2y_1} |\tilde{\xi}|^2 c_3^\varepsilon - e^{2y_1} c_2^\varepsilon - c_1^\varepsilon \right)_+ \frac{d\xi dy}{(2\pi)^d}.$$

Defining $B_\varepsilon = \{(\xi, y) \in \mathbf{R}^d \times \Omega_{2\varepsilon} : \xi_1^2 + e^{2y_1} \tilde{\xi}^2 \leq 1\}$, one evidently has

$$\begin{aligned} \sum_k (\lambda - \lambda_k)_+ &\geq \int_{B_\varepsilon} \left(\lambda - \xi_1^2 - e^{2y_1} |\tilde{\xi}|^2 c_3^\varepsilon - e^{2y_1} c_2^\varepsilon - c_1^\varepsilon \right)_+ \frac{d\xi dy}{(2\pi)^d} \\ &= \lambda^{1+d/2} \int_{B_\varepsilon} \left(1 - \xi_1^2 - e^{2y_1} |\tilde{\xi}|^2 c_3^\varepsilon - \frac{1}{\lambda} e^{2y_1} c_2^\varepsilon - \frac{1}{\lambda} c_1^\varepsilon \right)_+ \frac{d\xi dy}{(2\pi)^d} \end{aligned}$$

For the integrand we find

$$\left(1 - \xi_1^2 - e^{2y_1} |\tilde{\xi}|^2 c_3^\varepsilon - \frac{1}{\lambda} e^{2y_1} c_2^\varepsilon - \frac{1}{\lambda} c_1^\varepsilon \right)_+ \geq \left(1 - \xi_1^2 - e^{2y_1} |\tilde{\xi}|^2 \right)_+ - e^{2y_1} |\tilde{\xi}|^2 (c_3^\varepsilon - 1)_+ - \frac{1}{\lambda} e^{2y_1} c_2^\varepsilon - \frac{1}{\lambda} c_1^\varepsilon.$$

Consequently,

$$\begin{aligned} \sum_k (\lambda - \lambda_k)_+ &\geq \lambda^{1+d/2} \int_B \left(1 - \xi_1^2 - e^{2y_1} |\tilde{\xi}|^2 \right)_+ \frac{d\xi dy}{(2\pi)^d} - \lambda^{1+d/2} \int_{B \setminus B_\varepsilon} \left(1 - \xi_1^2 - e^{2y_1} |\tilde{\xi}|^2 \right)_+ \frac{d\xi dy}{(2\pi)^d} \\ &\quad - \lambda^{1+d/2} (c_3^\varepsilon - 1)_+ \frac{|B|}{(2\pi)^d} - \lambda^{d/2} c_2^\varepsilon e^{2R} \frac{|B|}{(2\pi)^d} - \lambda^{d/2} c_1^\varepsilon \frac{|B|}{(2\pi)^d}, \end{aligned}$$

where $B = \{(\xi, y) \in \mathbf{R}^d \times \Omega : \xi_1^2 + e^{2y_1} \tilde{\xi}^2 \leq 1\}$ and $R = \sup\{y_1 : y \in \Omega\}$. Note that

$$\int_{B \setminus B_\varepsilon} \left(1 - \xi_1^2 - e^{2y_1} |\tilde{\xi}|^2 \right)_+ \frac{d\xi dy}{(2\pi)^d} \leq \frac{|B \setminus B_\varepsilon|}{(2\pi)^d},$$

and that

$$\begin{aligned} B \setminus B_\varepsilon &= \{(\xi, y) \in \mathbf{R}^d \times (\Omega \setminus \Omega_{2\varepsilon}) : \xi_1^2 + e^{2y_1} \tilde{\xi}^2 \leq 1\} \\ &\subseteq \{(\xi, y) \in \mathbf{R}^d \times (\Omega \setminus \Omega_{2\varepsilon}) : \xi_1^2 + e^{2r} \tilde{\xi}^2 \leq 1\} \\ &= \{\xi \in \mathbf{R}^d : \xi_1^2 + e^{2r} \tilde{\xi}^2 \leq 1\} \times (\Omega \setminus \Omega_{2\varepsilon}), \end{aligned}$$

where $r = \inf\{y_1 : y \in \Omega\}$. Clearly, the set $\{\xi \in \mathbf{R}^d : \xi_1^2 + e^{2r} \tilde{\xi}^2 \leq 1\}$ is bounded and thus of finite measure. Therefore, assuming that $\partial\Omega$ is Lipschitz-regular, we can infer that

$$\frac{|B \setminus B_\varepsilon|}{(2\pi)^d} = O(\varepsilon), \quad \varepsilon \rightarrow 0.$$

Consider now the term $c_3^\varepsilon - 1$. We know that

$$c_3^\varepsilon = \int_{\mathbf{R}^d} e^{2z_1} g^\varepsilon(z)^2 dz = \int_{\mathbf{R}} e^{2z_1} g_1^\varepsilon(z_1)^2 dz_1 = \int_{\mathbf{R}} e^{2z_1} g_1(z_1)^2 dz_1.$$

Hence

$$c_3^\varepsilon - 1 = \int_{\mathbf{R}} (e^{2z_1} - 1) g_1(z_1)^2 dz_1 = \int_{\mathbf{R}} \sum_{k=1}^{\infty} \frac{(2z_1)^k}{k!} \varepsilon^k g_1(z_1)^2 dz_1 = O(\varepsilon), \quad \varepsilon \rightarrow 0.$$

For c_2^ε we find

$$c_2^\varepsilon = \int_{\mathbf{R}^d} e^{2z_1} |\nabla_{\tilde{z}} g^\varepsilon(z)|^2 dz = \varepsilon^{-2} \int_{\mathbf{R}^d} e^{2z_1} g(z_1)^2 dz_1 \int_{\mathbf{R}^d} e^{2z_1} |\nabla_{\tilde{z}} \tilde{g}(\tilde{z})|^2 d\tilde{z} = O(\varepsilon^{-2}), \quad \varepsilon \rightarrow 0,$$

and finally

$$c_1^\varepsilon = \int_{\mathbf{R}^d} (\partial_{z_1} g^\varepsilon(z))^2 dz = \int_{\mathbf{R}} (\partial_{z_1} g_1^\varepsilon(z_1))^2 dz_1 \int_{\mathbf{R}^{d-1}} \tilde{g}(\tilde{z})^2 d\tilde{z} = \varepsilon^{-2} \int_{\mathbf{R}} (\partial_{z_1} g_1(z_1))^2 dz_1 = O(\varepsilon^{-2}).$$

Linking ε to λ , again, via $\varepsilon = \lambda^{-\alpha}$ we obtain

$$\sum_k (\lambda - \lambda_k)_+ \geq \lambda^{1+d/2} \int_{\mathbf{R}^d \times \Omega} \left(\lambda - \xi_1^2 - e^{2y_1} |\tilde{\xi}|^2 \right)_+ \frac{d\xi dy}{(2\pi)^d} - R(\lambda),$$

where $R(\lambda)$ is a linear combination of terms that are in $O(\lambda^{d/2+1-\alpha})$ and $O(\lambda^{d/2-2\alpha})$ respectively. Again, minimizing $\max\{d/2 + 2\alpha, 1 + d/2 - \alpha\}$ yields $\alpha = 1/3$ and thus

$$\sum_k (\lambda - \lambda_k)_+ \geq \lambda^{1+d/2} \int_{\mathbf{R}^d \times \Omega} \left(\lambda - \xi_1^2 - e^{2y_1} |\tilde{\xi}|^2 \right)_+ \frac{d\xi dy}{(2\pi)^d} + O(\lambda^{d/2+2/3}).$$

□

In contrast to the case of the euclidean Laplacian, we need to prove an asymptotic upper bound for the hyperbolic Laplacian as well.

Lemma 4.3. *Let the same conditions as in the preceding lemma be satisfied. Then*

$$\sum_k (\lambda - \lambda_k)_+ \leq \lambda^{1+d/2} \int_{\mathbf{R}^d \times \Omega} (1 - \xi_1^2 - e^{2y_1} \tilde{\xi}^2)_+ \frac{d\xi dy}{(2\pi)^d} + O(\lambda^{d/2+2/3})$$

Proof. Recall that

$$\sum_k (\lambda - \lambda_k)_+ \leq \lambda^{1+d/2} \int_{\mathbf{R}^d \times \Omega^\varepsilon} (1 - \xi_1^2 - (c_3^\varepsilon)^{-1} e^{2y_1} \tilde{\xi}^2 + \lambda^{-1} c_1^\varepsilon + \lambda^{-1} (c_3^\varepsilon)^{-1} c_2^\varepsilon e^{2R})_+ \frac{d\xi dy}{(2\pi)^d}.$$

Let $\varepsilon = \lambda^{-\alpha}$. Then $((1 - (c_3^\varepsilon)^{-1}) = O(\lambda^{-\alpha})$, $\lambda^{-1} c_1^\varepsilon = O(\lambda^{-1+2\alpha})$ and $\lambda^{-1} c_2^\varepsilon = O(\lambda^{-1+2\alpha})$. If $\alpha < 1/2$, then all these terms converge to 0. In this case, for sufficiently large λ , the integrand is supported in $B^\varepsilon = \{(\xi, y) \in \mathbf{R}^d \times \Omega^\varepsilon : \xi_1^2 + e^{2y_1} \tilde{\xi}^2 \leq 2\}$. Furthermore, the integrand can be estimated from above in the following way

$$(1 - \xi_1^2 - (c_3^\varepsilon)^{-1} e^{2y_1} \tilde{\xi}^2 + c_1^\varepsilon \lambda^{-1} + (c_3^\varepsilon)^{-1} c_2^\varepsilon e^{2R} \lambda^{-1})_+ \leq (1 - \xi_1^2 - e^{2y_1} \tilde{\xi}^2)_+ + (1 - (c_3^\varepsilon)^{-1})_+ e^{2R} \tilde{\xi}^2 + \lambda^{-1} c_1^\varepsilon + \lambda^{-1} (c_3^\varepsilon)^{-1} c_2^\varepsilon e^{2R}.$$

Define $B = \{(\xi, y) \in \mathbf{R}^d \times \Omega: \xi_1^2 + e^{2y_1} \tilde{\xi}^2 \leq 2\}$. Then

$$\begin{aligned} \sum_k (\lambda - \lambda_k)_+ &\leq \lambda^{1+d/2} \int_B (1 - \xi_1^2 - e^{2y_1} \tilde{\xi}^2)_+ \frac{d\xi dy}{(2\pi)^d} \\ &\quad + \lambda^{1+d/2} \int_{B^\varepsilon \setminus B} (1 - \xi_1^2 - e^{2y_1} \tilde{\xi}^2)_+ \frac{d\xi dy}{(2\pi)^d} \\ &\quad + 2e^R \lambda^{d/2+1} ((1 - (c_3^\varepsilon)^{-1})_+ \frac{|B^\varepsilon|}{(2\pi)^d} + \lambda^{d/2} c_1^\varepsilon \frac{|B^\varepsilon|}{(2\pi)^d} + \lambda^{d/2} (c_3^\varepsilon)^{-1} c_2^\varepsilon e^{2R} \frac{|B^\varepsilon|}{(2\pi)^d}). \end{aligned}$$

Note that

$$\int_{B^\varepsilon \setminus B} (1 - \xi_1^2 - e^{2y_1} \tilde{\xi}^2)_+ \frac{d\xi dy}{(2\pi)^d} \leq \frac{|B^\varepsilon \setminus B|}{(2\pi)^d} = O(\varepsilon) = O(\lambda^{-\alpha}).$$

Therefore, the error term is of the order $O(\lambda^{\max\{d/2+2\alpha, 1+d/2-\alpha\}})$. As for the lower bound, the exponent is minimized for $\alpha = 1/3$. This concludes the argument. \square

Combining the two preceding lemmas, we obtain

Theorem 4.4. *Let $\Omega \subseteq \mathbf{R}^d$ be a bounded open set with Lipschitz boundary and let λ_k , $k \in \mathbf{N}$ be the eigenvalues of the operator H on Ω with Dirichlet boundary conditions. Then*

$$\sum_k (\lambda - \lambda_k)_+ = \lambda^{1+d/2} \int_{\mathbf{R}^d \times \Omega} (1 - \xi_1^2 - e^{2y_1} \tilde{\xi}^2)_+ \frac{d\xi dy}{(2\pi)^d} + O(\lambda^{d/2+2/3}).$$

ACKNOWLEDGMENTS

I would like to thank Ari Laptev for his help in preparing this paper.

REFERENCES

- [ACV08] Luigi Ambrosio, Andrea Colesanti, and Elena Villa. Outer Minkowski content for some classes of closed sets. *Math. Ann.*, 342(4):727–748, 2008.
- [Ber73] F. A. Berezin. Covariant and contravariant symbols of operators. *Math. USSR, Izv.*, 6:1117–1151, 1973.
- [FL20] Rupert L. Frank and Simon Larson. Two-term spectral asymptotics for the Dirichlet Laplacian in a Lipschitz domain. *J. Reine Angew. Math.*, 766:195–228, 2020.
- [FLW23] Rupert L. Frank, Ari Laptev, and Timo Weidl. *Schrödinger operators: eigenvalues and Lieb-Thirring inequalities*, volume 200 of *Camb. Stud. Adv. Math.* Cambridge: Cambridge University Press, 2023.
- [Gei14] Leander Geisinger. A short proof of Weyl’s law for fractional differential operators. *J. Math. Phys.*, 55(1):011504, 7, 2014.
- [ILW24] Alexei Ilyin, Ari Laptev, and Timon Weinmann. Lieb-Thirring inequalities on manifolds with constant negative curvature. *J. Geom. Anal.*, 34(2):17, 2024. Id/No 62.
- [LY83] Peter Li and Shing-Tung Yau. On the Schrödinger equation and the eigenvalue problem. *Commun. Math. Phys.*, 88:309–318, 1983.

IMPERIAL COLLEGE LONDON

Email address: t.weinmann21@imperial.ac.uk