

DOMAIN THEORY IN UNIVALENT FOUNDATIONS I: DIRECTED COMPLETE POSETS AND SCOTT'S D_∞

TOM DE JONG

ABSTRACT. We develop domain theory in constructive and predicative univalent foundations (also known as homotopy type theory). That we work predicatively means that we do not assume Voevodsky's propositional resizing axioms. Our work is constructive in the sense that we do not rely on excluded middle or the axiom of (countable) choice. Domain theory studies so-called directed complete posets (dcpos) and Scott continuous maps between them and has applications in a variety of fields, such as programming language semantics, higher-type computability and topology. A common approach to deal with size issues in a predicative foundation is to work with information systems, abstract bases or formal topologies rather than dcpos, and approximable relations rather than Scott continuous functions. In our type-theoretic approach, we instead accept that dcpos may be large and work with type universes to account for this. A priori one might expect that iterative constructions of dcpos may result in a need for ever-increasing universes and are predicatively impossible. We show, through a careful tracking of type universe parameters, that such constructions can be carried out in a predicative setting. In particular, we give a predicative reconstruction of Scott's D_∞ model of the untyped λ -calculus. Our work is formalised in the AGDA proof assistant and its ability to infer universe levels has been invaluable for our purposes.

1. INTRODUCTION

Domain theory [AJ94] is a well-established subject in mathematics and theoretical computer science with applications to programming language semantics [Sco72; Sco93; Plo77], higher-type computability [LN15], topology, and more [Gie+03].

We explore the development of domain theory from the univalent point of view [Voe15; Uni13]. This means that we work with the stratification of types as singletons, propositions, sets, 1-groupoids, etc. Our work does not require any higher inductive types other than the propositional truncation, and the only consequences of univalence needed here are function extensionality and propositional extensionality. Additionally, we work constructively and predicatively, as described below.

This paper develops the general theory of the central objects of study in domain theory: *directed complete posets (dcpos)* and the *Scott continuous* functions between them. We show how we can predicatively construct products, exponentials, bilimits and free pointed dcpos, culminating in the construction of Scott's famous D_∞ which provides a mathematical model of the untyped λ -calculus.

1.1. Constructivity. That we work constructively means that we do not assume excluded middle, or weaker variants, such as Bishop's LPO [Bis67], or the axiom of choice (which implies excluded middle), or its weaker variants, such as the axiom of countable choice. An advantage of working constructively and not relying on these additional logical axioms is that our development is valid in every $(\infty, 1)$ -topos [Shu19] and not just those in which the logic is classical.

Key words and phrases. constructivity, predicativity, univalent foundations, homotopy type theory, HoTT/UF, propositional resizing, domain theory, dcpo, directed complete poset.

Our commitment to constructivity has the particular consequence that we cannot simply add a least element to a set to obtain the free pointed dcpo. Instead of adding a single least element representing an undefined value, we must work with a more complex type of partial elements (Section 5). Similarly, the booleans under the natural ordering fail to be a dcpo, so we use the type of (small) propositions, ordered by implication, instead.

1.2. Predicativity. Our work is predicative in the sense that we do not assume Voevodsky’s *resizing* rules [Voe11; Voe15] or axioms. In particular, powersets of small types are large.

There are several (philosophical, model-theoretic, proof-theoretic, etc.) arguments for keeping the type theory predicative, see for instance [Uem19; Swa19b; Swa19a], and [dJE23, Section 1.1] for a brief overview, but here we only mention one that we consider to be amongst the most interesting. Namely, the existence of a computational interpretation of propositional impredicativity axioms for univalent foundations is an open problem.

A common approach to deal with domain-theoretic size issues in a predicative foundation is to work with information systems [Sco82a; Sco82b], abstract bases [AJ94] or formal topologies [Sam87; Sam03; Coq+03] rather than dcpos, and approximable relations rather than *Scott continuous functions*.

Instead, we work directly with dcpos and Scott continuous functions. In dealing with size issues, we draw inspiration from category theory and make crucial use of type universes and type equivalences to capture *smallness*. For example, in our development of the Scott model of PCF [dJon21; Har20], the dcpos have carriers in the second universe \mathcal{U}_1 and least upper bounds for directed families indexed by types in the first universe \mathcal{U}_0 . Moreover, up to equivalence of types, the order relation of the dcpos takes values in the lowest universe \mathcal{U}_0 . Seeing a poset as a category in the usual way, we can say that these dcpos are large, but locally small, and have small filtered colimits. The fact that the dcpos have large carriers is in fact unavoidable and characteristic of predicative settings, as proved in [dJE23].

Because the dcpos have large carriers it is a priori not clear that complex constructions of dcpos, involving countably infinite iterations of exponentials for example, do not result in a need for ever-increasing universes and are predicatively possible. We show that they are possible through a careful tracking of type universe parameters, and this is also illustrated by the construction of Scott’s D_∞ .

Since keeping track of these universes is prone to mistakes, we have formalised our work in AGDA (see Section 1.5); its ability to infer and keep track of universe levels has been invaluable.

1.3. Related work. In short, the distinguishing features of our work are: (i) the adoption of homotopy type theory as a foundation, (ii) a commitment to predicatively and constructively valid reasoning, (iii) the use of type universes to avoid size issues concerning large posets.

The standard works on domain theory, e.g. [AJ94; Gie+03], are based on traditional impredicative set theory with classical logic. A constructive, topos valid, and hence impredicative, treatment of some domain theory can be found in [Tay99, Chapter III].

Domain theory has been studied predicatively in the setting of formal topology [Sam87; Sam03; Coq+03] in [MV04; Neg02; SVV96] and the more recent categorical papers [Kaw17; Kaw21]. In this predicative setting, one avoids size issues by working with information systems [Sco82a; Sco82b], abstract bases [AJ94] or formal topologies, rather than dcpos, and approximable relations rather than Scott continuous functions. Hedberg [Hed96] presented some of these ideas in Martin-Löf

Type Theory and formalised them in the proof assistant ALF [Mag94], a precursor to AGDA. A modern formalisation in AGDA based on Hedberg's work was recently carried out in Lidell's master thesis [Lid20].

Our development differs from the above line of work in that it studies dcpos directly and uses type universes to account for the fact that dcpos may be large. An advantage of this approach is that we can work with (Scott continuous) functions rather than the arguably more involved (approximable) relations.

Another approach to formalising domain theory in type theory can be found in [BKV09; Doc14]. Both formalisations study ω -chain complete preorders, work with setoids, and make use of COQ's impredicative sort `Prop`. A setoid is a type equipped with an equivalence relation that must be respected by all functions. The particular equivalence relation given by equality is automatically respected of course, but for general equivalence relations this must be proved explicitly. The aforementioned formalisations work with preorders, rather than posets, because they are setoids where two elements x and y are related if $x \leq y$ and $y \leq x$. Our development avoids the use of setoids thanks to the adoption of the univalent point of view. Moreover, we work predicatively and we work with the more general directed families rather than ω -chains, as we intend the theory to also be applicable to topology and algebra [Gie+03].

There are also constructive accounts of domain theory aimed at program extraction [BK09; PM21]. Both these works study ω -chain complete posets (ω -cpos) and define notions of ω -continuity for them. The former [BK09] is notably predicative, but makes use of additional logical axioms: countable choice, dependent choice and Markov's Principle, which are validated by a realisability interpretation. The latter [PM21] uses constructive logic to extract witnesses but employs classical logic in the proofs of correctness by phrasing them in the double negation fragment of constructive logic. By contrast, we study (continuous) dcpos rather than (ω -continuous) ω -cpos and is fully constructive without relying on additional principles such as countable choice or Markov's Principle.

Yet another approach is the field of *synthetic domain theory* [Ros86; Ros87; Hy191; Reu99; RS99]. Although the work in this area is constructive, it is still impredicative, as it is based on topos logic; but more importantly it has a focus different from that of regular domain theory. The aim is to isolate a few basic axioms and find models in (realisability) toposes where every object is a domain and every morphism is continuous. These models often validate additional axioms, such as Markov's Principle and countable choice, and moreover (necessarily) falsify excluded middle. We have a different goal, namely to develop regular domain theory constructively and predicatively, but in a foundation compatible with excluded middle and choice, while not relying on them or on Markov's Principle or countable choice.

1.4. Relation to our other work. This paper and its follow-up [dJE24] (referred to as Part II) present a revised and expanded treatment of the results in our conference paper [dJE21]. In [dJE21] (and also [dJon21]) the definition of a poset included the requirement that the carrier is a set, because we only realised later that this was redundant (Lemma 3.3). Products of dcpos were not discussed in these works and are included in this paper, after having been in AGDA by Brendan Hart [Har20] for a final year MSci project supervised by Martín Escardó and myself.

Besides the construction of Scott's D_∞ in constructive and predicative univalent foundations, this paper lays the foundations for Part II [dJE24] which presents the predicative theory of continuous and algebraic domains. These results are also included in the author's PhD thesis [dJon22].

1.5. Formalisation. All of our results are formalised in AGDA, building on Escardó’s TYPE TOPOLOGY development [Esc+]. Hart’s previously cited work [Har20] was also ported to the current TYPE TOPOLOGY development by Escardó [Har23]. The reference [dJon24] precisely links each numbered environment (including definitions, examples and remarks) in this paper to its implementation. The HTML rendering has clickable links and so is particularly suitable for exploring the development: <https://www.cs.bham.ac.uk/~mhe/TypeTopology/DomainTheory.Part-I.html>.

1.6. Organisation. The paper is organised as follows:

- Section 2: A brief introduction to univalent foundations with a particular focus on type universes and the propositional truncation, as well as a discussion of impredicativity in the form of Voevodsky’s propositional resizing axioms.
- Section 3: Directed complete posets (dcpos), constructively and predicatively.
- Section 4: Scott continuous maps: morphism between dcpos.
- Section 5: The constructive lifting of a set and of a dcpo.
- Section 6: Products and exponentials of dcpos.
- Section 7: Bilimits of dcpos.
- Section 8: A predicative reconstruction of Scott’s D_∞ which models the untyped λ -calculus.

2. FOUNDATIONS

We work within intensional Martin-Löf Type Theory and we include $+$ (binary sum), Π (dependent product), Σ (dependent sum), Id (identity type), and inductive types, including $\mathbf{0}$ (empty type), $\mathbf{1}$ (type with exactly one element $\star : \mathbf{1}$) and \mathbb{N} (natural numbers). In general we adopt the same conventions of [Uni13]. In particular, we simply write $x = y$ for the identity type $\text{Id}_X(x, y)$ and use \equiv for the judgemental equality, and for dependent functions $f, g : \Pi_{x:X} A(x)$, we write $f \sim g$ for the pointwise equality $\Pi_{x:X} f(x) = g(x)$.

2.1. Universes. We assume a universe \mathcal{U}_0 and two operations: for every universe \mathcal{U} , a successor universe \mathcal{U}^+ with $\mathcal{U} : \mathcal{U}^+$, and for every two universes \mathcal{U} and \mathcal{V} another universe $\mathcal{U} \sqcup \mathcal{V}$ such that for any universe \mathcal{U} , we have $\mathcal{U}_0 \sqcup \mathcal{U} \equiv \mathcal{U}$ and $\mathcal{U} \sqcup \mathcal{U}^+ \equiv \mathcal{U}^+$. Moreover, $(-) \sqcup (-)$ is idempotent, commutative, associative, and $(-)^+$ distributes over $(-) \sqcup (-)$. We write $\mathcal{U}_1 \equiv \mathcal{U}_0^+$, $\mathcal{U}_2 \equiv \mathcal{U}_1^+$, \dots and so on. If $X : \mathcal{U}$ and $Y : \mathcal{V}$, then $X + Y : \mathcal{U} \sqcup \mathcal{V}$ and if $X : \mathcal{U}$ and $Y : X \rightarrow \mathcal{V}$, then the types $\Sigma_{x:X} Y(x)$ and $\Pi_{x:X} Y(x)$ live in the universe $\mathcal{U} \sqcup \mathcal{V}$; finally, if $X : \mathcal{U}$ and $x, y : X$, then $\text{Id}_X(x, y) : \mathcal{U}$. The type of natural numbers \mathbb{N} is assumed to be in \mathcal{U}_0 and we postulate that we have copies $\mathbf{0}_{\mathcal{U}}$ and $\mathbf{1}_{\mathcal{U}}$ in every universe \mathcal{U} . This has the useful consequence that while we do not assume cumulativity of universes, embeddings that lift types to higher universes are definable. For example, the map $(-) \times \mathbf{1}_{\mathcal{V}}$ takes a type in any universe \mathcal{U} to an equivalent type in the higher universe $\mathcal{U} \sqcup \mathcal{V}$. All our examples go through with just two universes \mathcal{U}_0 and \mathcal{U}_1 , but the theory is more easily developed in a general setting.

2.2. The univalent point of view. Within this type theory, we adopt the univalent point of view [Uni13]. A type X is a *proposition* (or *truth value* or *sub-singleton*) if it has at most one element, i.e. we have an element of the type $\text{is-prop}(X) \equiv \prod_{x,y:X} x = y$. A major difference between univalent foundations and other foundational systems is that we *prove* that types are propositions or properties. For instance, we can show (using function extensionality) that the axioms of directed complete poset form a proposition. A type X is a *set* if any two elements can be identified in at most one way, i.e. we have an element of the type $\prod_{x,y:X} \text{is-prop}(x = y)$.

2.3. Extensionality axioms. The univalence axiom [Uni13] is not needed for our development, although we do pause to point out its consequences in two places, namely in Section 2.5 and Remark 5.9.

We assume function extensionality and propositional extensionality, often tacitly:

- (i) *Propositional extensionality*: if P and Q are two propositions, then we postulate that $P = Q$ holds exactly when we have both $P \rightarrow Q$ and $Q \rightarrow P$.
- (ii) *Function extensionality*: if $f, g : \prod_{x:X} A(x)$ are two (dependent) functions, then we postulate that $f = g$ holds exactly when $f \sim g$.

Function extensionality has the important consequence that the propositions form an exponential ideal, i.e. if X is a type and $Y : X \rightarrow \mathcal{U}$ is such that every $Y(x)$ is a proposition, then so is $\prod_{x:X} Y(x)$ [Uni13, Example 3.6.2]. In light of this, universal quantification is given by Π -types in our type theory.

2.4. The propositional truncation. In Martin-Löf Type Theory, an element of $\prod_{x:X} \sum_{y:Y} \phi(x, y)$, by definition, gives us a function $f : X \rightarrow Y$ such that $\prod_{x:X} \phi(x, f(x))$. In some cases, we wish to express the weaker “for every $x : X$, there exists some $y : Y$ such that $\phi(x, y)$ ” without necessarily having an assignment of x 's to y 's. A good example of this is when we define directed families later (see Definition 3.4). This is achieved through the propositional truncation.

Given a type $X : \mathcal{U}$, we postulate that we have a proposition $\|X\| : \mathcal{U}$ with a function $|-| : X \rightarrow \|X\|$ such that for every proposition $P : \mathcal{V}$ in any universe \mathcal{V} , every function $f : X \rightarrow P$ factors (necessarily uniquely, by function extensionality) through $|-|$. Diagrammatically,

$$\begin{array}{ccc} X & \xrightarrow{f} & P \\ & \searrow & \nearrow \\ & |-| & \|X\| \end{array}$$

Notice that the induction and recursion principles automatically hold up to an identification: writing \bar{f} for the dashed map above, we have an identification $\bar{f}(|x|) = f(x)$ for every $x : X$ because P is assumed to be a proposition. This is sufficient for our purposes and we do not require these equalities to hold judgementally.

Existential quantification $\exists_{x:X} Y(x)$ is given by $\|\Sigma_{x:X} Y(x)\|$. One should note that if we have $\exists_{x:X} Y(x)$ and we are trying to prove some proposition P , then we may assume that we have $x : X$ and $y : Y(x)$ when constructing our element of P . Similarly, we can define disjunction as $P \vee Q \equiv \|P + Q\|$.

We assume throughout that every universe is closed under propositional truncations, meaning that if $X : \mathcal{U}$ then $\|X\| : \mathcal{U}$ as well. We also stress that the propositional truncation is the only higher inductive type used in our work.

2.5. Size and impredicativity. We introduce the notion of smallness and use it to define propositional resizing axioms, which we take to be the definition of impredicativity in univalent foundations.

Definition 2.1 (Smallness). A type X in any universe is said to be \mathcal{U} -small if it is equivalent to a type in the universe \mathcal{U} . That is, X is \mathcal{U} -small $\equiv \Sigma_{Y:\mathcal{U}} (Y \simeq X)$.

Here, the symbol \simeq refers to Voevodsky's notion of equivalence [Uni13]. Notice that the type $(X \text{ is } \mathcal{U}\text{-small})$ is a proposition if and only if the univalence axiom holds, see [Esc19, Sections 3.14 and 3.36.3].

Definition 2.2 (Type of propositions $\Omega_{\mathcal{U}}$). The type of propositions in a universe \mathcal{U} is $\Omega_{\mathcal{U}} \equiv \sum_{P:\mathcal{U}} \text{is-prop}(P) : \mathcal{U}^+$.

Observe that $\Omega_{\mathcal{U}}$ itself lives in the successor universe \mathcal{U}^+ . We often think of the types in some fixed universe \mathcal{U} as *small* and accordingly we say that $\Omega_{\mathcal{U}}$ is *large*. Similarly, the powerset of a type $X : \mathcal{U}$ is large. Given our predicative setup, we must pay attention to universes when considering powersets:

Definition 2.3 (\mathcal{V} -powerset $\mathcal{P}_{\mathcal{V}}(X)$, \mathcal{V} -subsets). Let \mathcal{V} be a universe and $X : \mathcal{U}$ type. We define the \mathcal{V} -powerset $\mathcal{P}_{\mathcal{V}}(X)$ as $X \rightarrow \Omega_{\mathcal{V}} : \mathcal{V}^+ \sqcup \mathcal{U}$. Its elements are called \mathcal{V} -subsets of X .

Definition 2.4 (\in, \subseteq). Let x be an element of a type X and let A be an element of the powerset $\mathcal{P}_{\mathcal{V}}(X)$. We write $x \in A$ for the type $\text{pr}_1(A(x))$. The first projection pr_1 is needed because $A(x)$, being of type $\Omega_{\mathcal{V}}$, is a pair. Given two \mathcal{V} -subsets A and B of X , we write $A \subseteq B$ for $\prod_{x:X}(x \in A \rightarrow x \in B)$.

Function extensionality and propositional extensionality imply that $A = B$ if and only if $A \subseteq B$ and $B \subseteq A$.

One could ask for a *resizing axiom* asserting that $\Omega_{\mathcal{U}}$ has size \mathcal{U} , which we call *the propositional impredicativity of \mathcal{U}* . A closely related axiom is *propositional resizing*, which asserts that every proposition $P : \mathcal{U}^+$ has size \mathcal{U} . Without the addition of such resizing axioms, the type theory is said to be *predicative*. As an example of the use of impredicativity in mathematics, we mention that the powerset has unions of arbitrary subsets if and only if propositional resizing holds [Esc19, Section 3.36.6].

We note that the resizing axioms are actually theorems when classical logic is assumed. This is because if $P \vee \neg P$ holds for every proposition in $P : \mathcal{U}$, then the only propositions (up to equivalence) are $\mathbf{0}_{\mathcal{U}}$ and $\mathbf{1}_{\mathcal{U}}$, which have equivalent copies in \mathcal{U}_0 , and $\Omega_{\mathcal{U}}$ is equivalent to a type $\mathbf{2}_{\mathcal{U}} : \mathcal{U}$ with exactly two elements.

3. DIRECTED COMPLETE POSETS

We offer the following overture in preparation of our development, especially if the reader is familiar with domain theory in a classical, set-theoretic setting.

The basic object of study in domain theory is that of a *directed complete poset* (dcpo). In (impredicative) set-theoretic foundations, a dcpo can be defined to be a poset that has least upper bounds of all directed subsets. A naive translation of this to our foundation would be to proceed as follows. Define a poset in a universe \mathcal{U} to be a type $P : \mathcal{U}$ with a reflexive, transitive and antisymmetric relation $\sqsubseteq - : P \times P \rightarrow \mathcal{U}$. Since we wish to consider posets and not categories we require that the values $p \sqsubseteq q$ of the order relation are *subsingletons*. Then we could say that the poset (P, \sqsubseteq) is *directed complete* if every directed family $I \rightarrow P$ with indexing type $I : \mathcal{U}$ has a least upper bound (supremum). The problem with this definition is that there are no interesting examples in our constructive and predicative setting. For instance, assume that the poset $\mathbf{2}$ with two elements $0 \sqsubseteq 1$ is directed complete, and consider a proposition $A : \mathcal{U}$ and the directed family $A + \mathbf{1} \rightarrow \mathbf{2}$ that maps the left component to 0 and the right component to 1. By case analysis on its hypothetical supremum (Definition 3.6), we conclude that the negation of A is decidable. This amounts to weak excluded middle (which is equivalent to De Morgan's Law) and is constructively unacceptable.

To try to get an example, we may move to the poset $\Omega_{\mathcal{U}_0}$ of propositions in the universe \mathcal{U}_0 , ordered by implication. This poset does have all suprema of families $I \rightarrow \Omega_{\mathcal{U}_0}$ indexed by types I in the *first universe* \mathcal{U}_0 , given by existential quantification. But if we consider a directed family $I \rightarrow \Omega_{\mathcal{U}_0}$ with I in the *same universe* as $\Omega_{\mathcal{U}_0}$ lives, namely the *second universe* \mathcal{U}_1 , existential quantification gives a proposition in the *second universe* \mathcal{U}_1 and so doesn't give its supremum. In this example, we get a poset such that

- (i) the carrier lives in the universe \mathcal{U}_1 ,

- (ii) the order has truth values in the universe \mathcal{U}_0 , and
- (iii) suprema of directed families indexed by types in \mathcal{U}_0 exist.

Regarding a poset as a category in the usual way, we have a large, but locally small, category with small filtered colimits (directed suprema). This is typical of all the concrete examples that we consider, such as the dcpos in the Scott model of PCF [dJon21] and Scott's D_∞ model of the untyped λ -calculus (Section 8). We may say that the predicativity restriction increases the universe usage by one. However, for the sake of generality, we formulate our definition of dcpo with the following universe conventions:

- (i) the carrier lives in a universe \mathcal{U} ,
- (ii) the order has truth values in a universe \mathcal{T} , and
- (iii) suprema of directed families indexed by types in a universe \mathcal{V} exist.

So our notion of dcpo has three universe parameters \mathcal{U} , \mathcal{V} and \mathcal{T} . We will say that the dcpo is *locally small* when \mathcal{T} is not necessarily the same as \mathcal{V} , but the order has \mathcal{V} -small truth values. Most of the time we mention \mathcal{V} explicitly and leave \mathcal{U} and \mathcal{T} to be understood from the context.

We now define directed complete poset in constructive and predicative univalent foundations. We carefully explain our use of the propositional truncation in our definitions and, as mentioned above, the type universes involved.

Definition 3.1 (Preorder, reflexivity, transitivity). A *preorder* (P, \sqsubseteq) is a type $P : \mathcal{U}$ together with a proposition-valued binary relation $\sqsubseteq : P \rightarrow P \rightarrow \Omega_{\mathcal{T}}$ satisfying

- (i) *reflexivity*: for every $p : P$, we have $p \sqsubseteq p$, and
- (ii) *transitivity*: for every $p, q, r : P$, if $p \sqsubseteq q$ and $q \sqsubseteq r$, then $p \sqsubseteq r$.

Definition 3.2 (Poset, antisymmetry). A *poset* is a preorder (P, \sqsubseteq) that is *antisymmetric*: if $p \sqsubseteq q$ and $q \sqsubseteq p$, then $p = q$ for every $p, q : P$.

Lemma 3.3. *If (P, \sqsubseteq) is a poset, then P is a set.*

Proof. For every $p, q : P$, the composite

$$(p = q) \xrightarrow{\text{by reflexivity}} (p \sqsubseteq q) \times (q \sqsubseteq p) \xrightarrow{\text{by antisymmetry}} (p = q)$$

is constant since $(p \sqsubseteq q) \times (q \sqsubseteq p)$ is a proposition. By [Kra+17, Lemma 3.11] it therefore follows that P must be a set. \square

From now on, we will simply write “let P be a poset” leaving the partial order \sqsubseteq implicit. We will often use the symbol \sqsubseteq for partial orders on different carriers when it is clear from the context which one it refers to.

Definition 3.4 ((Semi)directed family). A family $\alpha : I \rightarrow P$ of elements of a poset P is *semidirected* if whenever we have $i, j : I$, there exists some $k : I$ such that $\alpha_i \sqsubseteq \alpha_k$ and $\alpha_j \sqsubseteq \alpha_k$. We frequently use the shorthand $\alpha_i, \alpha_j \sqsubseteq \alpha_k$ to denote the latter requirement. Such a family is *directed* if it is semidirected and its domain I is inhabited.

The name “semidirected” matches Taylor’s terminology [Tay99, Definition 3.4.1].

Remark 3.5. Note our use of the propositional truncation in defining when a family is *directed*. To make this explicit, we write out the definition in type-theoretic syntax: a family $\alpha : I \rightarrow P$ is directed if

- (i) we have an element of $\|I\|$, and
- (ii) $\prod_{i,j:I} \|\Sigma_{k:I} (\alpha_i \sqsubseteq \alpha_k) \times (\alpha_j \sqsubseteq \alpha_k)\|$.

The use of the propositional truncation ensures that the types (i) and (ii) are propositions and hence that being (semi)directed is a property of a family. The type

(ii) without truncation would instead express an assignment of a chosen $k : I$ for every $i, j : I$ instead.

Definition 3.6 ((Least) upper bound, supremum). An element x of a poset P is an *upper bound* of a family $\alpha : I \rightarrow P$ if $\alpha_i \sqsubseteq x$ for every $i : I$. It is a *least upper bound* of α if it is an upper bound, and whenever $y : P$ is an upper bound of α , then $x \sqsubseteq y$. By antisymmetry, a least upper bound is unique if it exists, so in this case we will speak of *the* least upper bound of α , or sometimes the *supremum* of α .

Definition 3.7 (\mathcal{V} -directed complete poset, \mathcal{V} -dcpo, $\bigsqcup \alpha$, $\bigsqcup_{i:I} \alpha_i$). For a universe \mathcal{V} , a *\mathcal{V} -directed complete poset* (or *\mathcal{V} -dcpo*, for short) is a poset D such that every directed family $\alpha : I \rightarrow D$ with $I : \mathcal{V}$ has a supremum in D that we denote by $\bigsqcup \alpha$ or $\bigsqcup_{i:I} \alpha_i$.

Remark 3.8. Explicitly, we ask for an element of the type

$$\prod_{I:\mathcal{V}} \prod_{\alpha:I \rightarrow D} (\text{is-directed } \alpha \rightarrow \Sigma_{x:D} (x \text{ is-sup-of } \alpha)),$$

where $(x \text{ is-sup-of } \alpha)$ is the type expressing that x is the supremum of α . Even though we used Σ and not \exists in this expression, this type is still a proposition: By [Uni13, Example 3.6.2], it suffices to prove that the type $\Sigma_{x:D} (x \text{ is-sup-of } \alpha)$ is a proposition. So suppose that we have $x, y : D$ with $p : x \text{ is-sup-of } \alpha$ and $q : y \text{ is-sup-of } \alpha$. Being the supremum of a family is a property because the partial order is proposition-valued. Hence, by [Uni13, Lemma 3.5.1], to prove that $(x, p) = (y, q)$, it suffices to prove that $x = y$. But this follows from antisymmetry and the fact that x and y are both suprema of α .

We will sometimes leave the universe \mathcal{V} implicit, and simply speak of a dcpo. On other occasions, we need to carefully keep track of universe levels. To this end, we make the following definition.

Definition 3.9 (\mathcal{V} -DCPO $_{\mathcal{U}, \mathcal{T}}$). Let \mathcal{V}, \mathcal{U} and \mathcal{T} be universes. We write \mathcal{V} -DCPO $_{\mathcal{U}, \mathcal{T}}$ for the type of \mathcal{V} -dcpos with carrier in \mathcal{U} and order taking values in \mathcal{T} .

Remark 3.10. In particular, it is very important to keep track of the universe parameters of the lifting (Section 5) and of exponentials (Section 6) in order to ensure that it is possible to construct Scott's D_∞ (and the Scott model of PCF [dJon21]) in our predicative setting, as we do in Section 8.

In many examples and applications, we require our dcpos to have a least element.

Definition 3.11 (Pointed dcpo). A dcpo D is *pointed* if it has a least element which we will denote by \perp_D , or simply \perp .

Definition 3.12 (Local smallness). A \mathcal{V} -dcpo D is *locally small* if $x \sqsubseteq y$ is \mathcal{V} -small for every $x, y : D$.

Lemma 3.13. A \mathcal{V} -dcpo D is locally small if and only if we have $\sqsubseteq_{\mathcal{V}} : D \rightarrow D \rightarrow \mathcal{V}$ such that $x \sqsubseteq y$ holds precisely when $x \sqsubseteq_{\mathcal{V}} y$ does.

Proof. The \mathcal{V} -dcpo D is locally small exactly when we have an element of

$$\prod_{x,y:D} \Sigma_{T:\mathcal{V}} (T \simeq x \sqsubseteq y).$$

But this type is equivalent to

$$\Sigma_{R:D \rightarrow D \rightarrow \mathcal{V}} \prod_{x,y:D} (R(x, y) \simeq x \sqsubseteq y)$$

by distributivity of Π over Σ [Uni13, Theorem 2.5.17]. \square

Nearly all examples of \mathcal{V} -dcpos in this thesis will be locally small. We now introduce two fundamental examples of dcpos: the type of subsingletons and powersets.

Example 3.14 (The type of subsingletons as a pointed dcpo). For any type universe \mathcal{V} , the type $\Omega_{\mathcal{V}}$ of subsingletons in \mathcal{V} is a poset if we order the propositions by implication. Note that antisymmetry holds precisely because of propositional extensionality. Moreover, $\Omega_{\mathcal{V}}$ has a least element, namely $\mathbf{0}_{\mathcal{V}}$, the empty type in \mathcal{V} . We also claim that $\Omega_{\mathcal{V}}$ has suprema for all (not necessarily directed) families $\alpha : I \rightarrow \Omega_{\mathcal{V}}$ with $I : \mathcal{V}$. Given such a family α , its least upper bound is given by $\exists_{i:I} \alpha_i$. It is clear that this is indeed an upper bound for α . And if P is a subsingleton such that $\alpha_i \sqsubseteq P$ for every $i : I$, then to show that $(\exists_{i:I} \alpha_i) \rightarrow P$ it suffices to construct a map $(\Sigma_{i:I} \alpha_i) \rightarrow P$ as P is a subsingleton. But this is easy because we assumed that $\alpha_i \sqsubseteq P$ for every $i : I$. Finally, paying attention to the universe levels we observe that $\Omega_{\mathcal{V}} : \mathcal{V}\text{-DCPO}_{\mathcal{V}^+, \mathcal{V}}$.

Example 3.15 (Powersets as pointed dcpos). Recalling our treatment of subset and powersets from Section 2.5, we show that powersets give examples of pointed dcpos. Specifically, for every type $X : \mathcal{U}$ and every type universe \mathcal{V} , the subset inclusion \subseteq makes $\mathcal{P}_{\mathcal{V}}(X)$ into a poset, where antisymmetry holds by function extensionality and propositional extensionality. Moreover, $\mathcal{P}_{\mathcal{V}}(X)$ has a least element of course: the empty set \emptyset . We also claim that $\mathcal{P}_{\mathcal{V}}(X)$ has suprema for all (not necessarily directed) families $\alpha : I \rightarrow \mathcal{P}_{\mathcal{V}}(X)$ with $I : \mathcal{V}$. Given such a family α , its least upper bound is given by $\bigcup \alpha \equiv \lambda x . \exists_{i:I} x \in \alpha_i$, the set-theoretic union, which is well-defined as $(\exists_{i:I} x \in \alpha_i) : \mathcal{V}$. It is clear that this is indeed an upper bound for α . And if A is a \mathcal{V} -subset of X such that $\alpha_i \subseteq A$ for every $i : I$, then to show that $\bigcup \alpha \subseteq A$ it suffices to construct for every $x : X$, a map $(\Sigma_{i:I} \alpha_i) \rightarrow (x \in A)$ as $x \in A$ is a subsingleton. But this is easy because we assumed that $\alpha_i \subseteq A$ for every $i : I$. Finally, paying attention to the universe levels we observe that $\mathcal{P}_{\mathcal{V}}(X) : \mathcal{V}\text{-DCPO}_{\mathcal{V}^+ \sqcup \mathcal{U}, \mathcal{V} \sqcup \mathcal{U}}$. In the case that $X : \mathcal{U} \equiv \mathcal{V}$, we obtain the simpler, locally small $\mathcal{P}_{\mathcal{V}}(X) : \mathcal{V}\text{-DCPO}_{\mathcal{V}^+, \mathcal{V}}$.

Of course, $\Omega_{\mathcal{V}}$ is easily seen to be equivalent to $\mathcal{P}_{\mathcal{V}}(\mathbf{1}_{\mathcal{V}})$, so Example 3.15 subsumes Example 3.14, but it is instructive to understand Example 3.14 first.

Proposition 3.16 (ω -completeness). *Every \mathcal{V} -dcpo D is ω -complete, viz. if we have elements $x_0 \sqsubseteq x_1 \sqsubseteq x_2 \sqsubseteq \dots$ of D , then the supremum of $(x_n)_{n:\mathbb{N}}$ exists in D .*

Proof. Recalling from Section 2.1 that we can lift types to higher universes and using the fact that \mathbb{N} is in the base universe \mathcal{U}_0 , we obtain a type $\text{lift}_{\mathcal{U}_0, \mathcal{V}}(\mathbb{N})$ in the universe \mathcal{V} that is equivalent to \mathbb{N} . Now $\text{lift}_{\mathcal{U}_0, \mathcal{V}}(\mathbb{N}) \simeq \mathbb{N} \xrightarrow{x(-)} D$ is a directed family as $x_n \sqsubseteq x_{n+1}$ for every natural number n , and it is indexed by a type in \mathcal{V} . Hence, it has a least upper bound in D which is the supremum of $(x_n)_{n:\mathbb{N}}$. \square

4. SCOTT CONTINUOUS MAPS

We discuss an appropriate notion of morphism between \mathcal{V} -dcpos, namely one that requires preservation of directed suprema and the order (Lemma 4.4).

Definition 4.1 (Scott continuity). A function $f : D \rightarrow E$ between two \mathcal{V} -dcpos is (Scott) continuous if it preserves directed suprema, i.e. if $I : \mathcal{V}$ and $\alpha : I \rightarrow D$ is directed, then $f(\bigsqcup \alpha)$ is the supremum in E of the family $f \circ \alpha$.

Remark 4.2. When we speak of a Scott continuous function between D and E , then we will always assume that D and E are both \mathcal{V} -dcpos for some arbitrary but fixed type universe \mathcal{V} .

Lemma 4.3. *Being Scott continuous is a property. In particular, two Scott continuous maps are equal if and only if they are equal as functions.*

Proof. By [Uni13, Example 3.6.2] and the fact that being the supremum of a family is a property, cf. Remark 3.8. \square

Lemma 4.4. *If $f : D \rightarrow E$ is Scott continuous, then it is monotone, i.e. $x \sqsubseteq_D y$ implies $f(x) \sqsubseteq_E f(y)$.*

Proof. Given $x, y : D$ with $x \sqsubseteq y$, consider the directed family $\mathbf{2}_{\mathcal{V}} \xrightarrow{\alpha} D$ defined by $\alpha(0) \equiv x$ and $\alpha(1) \equiv y$. Its supremum is y and f must preserve it. Hence, $f(y)$ is an upper bound of $f(\alpha(0)) \equiv f(x)$, so $f(x) \sqsubseteq f(y)$, as we wished to show. \square

Lemma 4.5. *If $f : D \rightarrow E$ is monotone and $\alpha : I \rightarrow D$ is directed, then so is $f \circ \alpha$.*

Proof. Since α is directed, I is inhabited, so it remains to prove that $f \circ \alpha$ is semidirected. If we have $i, j : I$, then by directedness of α , there exists $k : I$ such that $\alpha_i, \alpha_j \sqsubseteq \alpha_k$. By monotonicity, we obtain $f(\alpha_i), f(\alpha_j) \sqsubseteq f(\alpha_k)$ as desired. \square

Lemma 4.6. *A monotone map $f : D \rightarrow E$ between \mathcal{V} -dcpos is Scott continuous if and only if $f(\bigsqcup \alpha) \sqsubseteq \bigsqcup f \circ \alpha$.*

Note that we are justified in writing $\bigsqcup f \circ \alpha$ because Lemma 4.5 tells us that $f \circ \alpha$ is directed by the assumed monotonicity of f .

Proof. The left-to-right implication is immediate. For the converse, note that it only remains to show that $f(\bigsqcup \alpha) \sqsupseteq \bigsqcup f \circ \alpha$. But for this it suffices that $f(\alpha_i) \sqsubseteq f(\bigsqcup \alpha)$ for every $i : I$, which holds as $\bigsqcup \alpha$ is an upper bound of α and f is monotone. \square

Remark 4.7. In constructive mathematics it is not possible to exhibit a discontinuous function from $\mathbb{N}^{\mathbb{N}}$ to \mathbb{N} , because sheaf [TvD88, Chapter 15] and realizability models [vOos08, e.g. Proposition 3.1.6] imply that it is consistent to assume that all such functions are continuous. This does not mean, however, that we cannot exhibit a discontinuous function between dcpos. In fact, the negation map $\neg : \Omega \rightarrow \Omega$ is not monotone and hence not continuous. If we were to preclude such examples, then we can no longer work with the full type Ω of all propositions, but instead we must restrict to a subtype of propositions, for example by using dominances [Ros86]. Indeed, this approach is investigated in the context of topos theory in [Pha91; Lon95] and for computability instead of continuity in univalent foundations in [EK17].

Definition 4.8 (Strictness). A Scott continuous function $f : D \rightarrow E$ between pointed dcpos is *strict* if $f(\perp_D) = \perp_E$.

Lemma 4.9. *A poset D is a pointed \mathcal{V} -dcpo if and only if it has suprema for all semidirected families indexed by types in \mathcal{V} that we will denote using the \bigvee symbol. In particular, a pointed \mathcal{V} -dcpo has suprema of all families indexed by propositions in \mathcal{V} .*

Moreover, if f is a Scott continuous and strict map between pointed \mathcal{V} -dcpos, then f preserves suprema of semidirected families.

Proof. If D is complete with respect to semidirected families indexed by types in \mathcal{V} , then it is clearly a \mathcal{V} -dcpo and it is pointed because the supremum of the family indexed by the empty type is the least element. Conversely, if D is a pointed \mathcal{V} -dcpo and $\alpha : I \rightarrow D$ is a semidirected family with $I : \mathcal{V}$, then the family

$$\begin{aligned} \hat{\alpha} : I + \mathbf{1}_{\mathcal{V}} &\rightarrow D \\ \text{inl}(i) &\mapsto \alpha_i \\ \text{inr}(\star) &\mapsto \perp \end{aligned}$$

is directed and hence has a supremum in D which is also the least upper bound of α .

A pointed \mathcal{V} -dcpo must have suprema for all families indexed by propositions in \mathcal{V} , because any such family is semidirected. Finally, suppose that $\alpha : I \rightarrow D$

is semidirected and that $f : D \rightarrow E$ is Scott continuous and strict. Using the $(-)$ -construction from above, we see that

$$\begin{aligned} f(\bigvee \alpha) &\equiv f(\bigsqcup \hat{\alpha}) \\ &= \bigsqcup f \circ \hat{\alpha} && \text{(by Scott continuity of } f\text{)} \\ &= \bigsqcup \widehat{f \circ \alpha} && \text{(since } f \text{ is strict)} \\ &\equiv \bigvee f \circ \alpha, \end{aligned}$$

finishing the proof. \square

Proposition 4.10.

- (i) *The identity on any dcpo is Scott continuous.*
 - (ii) *For dcpos D and E and $y : E$, the constant map $x \mapsto y : D \rightarrow E$ is Scott continuous.*
 - (iii) *If $f : D \rightarrow E$ and $g : E \rightarrow E'$ are Scott continuous, then so is $g \circ f$.*
- Moreover, if D is pointed, then the identity on D is strict, and if f and g are strict in (iii), then so is $g \circ f$.*

Proof. The proofs of (i) and (ii) are obvious. For (iii), let $\alpha : I \rightarrow D$ be directed and notice that $g(f(\bigsqcup \alpha)) = g(\bigsqcup f \circ \alpha) = \bigsqcup g \circ f \circ \alpha$ by respectively continuity of f and g . The claims about strictness are also clear. \square

Definition 4.11 (Isomorphism). A Scott continuous map $f : D \rightarrow E$ is an *isomorphism* if we have a Scott continuous inverse $g : E \rightarrow D$.

Lemma 4.12. *Every $f : D \rightarrow E$ isomorphism between pointed dcpos is strict.*

Proof. Let $y : E$ be arbitrary and notice that $\perp_D \sqsubseteq g(y)$ because \perp_D is the least element of D . By monotonicity of f , we get $f(\perp_D) \sqsubseteq f(g(y)) = y$ which shows that $f(\perp_D)$ is the least element of E . \square

Definition 4.13 (Scott continuous retract). A dcpo D is a (*Scott*) *continuous retract* of E if we have Scott continuous maps $s : D \rightarrow E$ and $r : E \rightarrow D$ such that s is a section of r . We denote this situation by $D \xrightleftharpoons[r]{s} E$.

Lemma 4.14. *If D is a continuous retract of E and E is locally small, then so is D .*

Proof. We claim that $x \sqsubseteq_D y$ and $s(x) \sqsubseteq_E s(y)$ are equivalent, which proves the lemma as E is assumed to be locally small. One direction of the equivalence is given by the fact that s is monotone. In the other direction, assume that $s(x) \sqsubseteq s(y)$ and note that $x = r(s(x)) \sqsubseteq r(s(y)) = y$, as r is monotone and s is a section of r . \square


5. LIFTING

We now turn to constructing pointed \mathcal{V} -dcpos from sets. First of all, every discretely ordered set is a \mathcal{V} -dcpo, where discretely ordered means that we have $x \sqsubseteq y$ exactly when $x = y$. This is because if $\alpha : I \rightarrow X$ is a directed family into a discretely ordered set X , then α has to be constant (by semidirectedness), so α_i is its supremum for any $i : I$. And since directedness includes the condition that the domain is inhabited, it follows that α must have a supremum in X . In fact, ordering X discretely yields the free \mathcal{V} -dcpo on the set X in the categorical sense.

With excluded middle, the situation for *pointed* \mathcal{V} -dcpos is also very straightforward. Simply order the set X discretely and add a least element, as depicted for $X \equiv \mathbb{N}$ in the Hasse diagram of Proposition 5.1. The point of that proposition is to show, by a reduction to the constructive taboo LPO [Bis67], that this approach is

constructively unsatisfactory. In fact, in [dJE23] we proved a general constructive no-go theorem showing that there is a nontrivial dcpo with decidable equality if and only if weak excluded middle holds.

Proposition 5.1. *If the poset $\mathbb{N}_\perp = (\mathbb{N} + \mathbf{1}, \sqsubseteq)$ with order depicted by the Hasse diagram*



is ω -complete, then LPO holds. In particular, by Proposition 3.16, if it is \mathcal{U}_0 -directed complete, then LPO holds.

Proof. Let $\alpha : \mathbb{N} \rightarrow \mathbf{2}$ be an arbitrary binary sequence. We show that $\exists_{n:\mathbb{N}} \alpha_n = 1$ is decidable. Define the family $\beta : \mathbb{N} \rightarrow \mathbb{N}_\perp$ by

$$\beta_n \equiv \begin{cases} \text{inl}(k) & \text{if } k \text{ is the least integer below } n \text{ for which } \alpha_k = 1, \text{ and} \\ \text{inr}(\star) & \text{else.} \end{cases}$$

Then β is a chain, so by assumption it has a supremum s in \mathbb{N}_\perp . By the induction principle of coproducts, we have $s = \text{inr}(\star)$ or we have $k : \mathbb{N}$ such that $s = \text{inl}(k)$. If the latter holds, then $\alpha_k = 1$, so $\exists_{n:\mathbb{N}} \alpha_n = 1$ is decidable. We claim that $s = \text{inr}(\star)$ implies that $\neg(\exists_{n:\mathbb{N}} \alpha_n = 1)$. Indeed, assume for a contradiction that $\exists_{n:\mathbb{N}} \alpha_n = 1$. Since we are proving a proposition, we may assume to have $n : \mathbb{N}$ with $\alpha_n = 1$. Then, $\beta_n = \text{inl}(k)$ for a natural number $k \leq n$. Since s is the supremum of β we have $\text{inl}(k) = \beta_n \sqsubseteq s = \text{inr}(\star)$, but $\text{inr}(\star)$ is the least element of \mathbb{N}_\perp , so by antisymmetry $\text{inl}(k) = \text{inr}(\star)$, which is impossible. \square

Our solution to the above will be to work with the lifting monad, sometimes known as the partial map classifier monad from topos theory [Joh77; Ros86; Ros87; Koc91], which has been extended to constructive type theory by Reus and Streicher [RS99] and recently to univalent foundations by Escardó and Knapp [EK17; Kna18].

Remark 5.2. Escardó and Knapp's work [EK17; Kna18] yields an approach to partiality in univalent foundations and aim to avoid (weak) countable choice, which is not provable in constructive univalent foundations [CMR17; Coq18; Swa19b; Swa19a]. This is to be contrasted to other approaches to partiality in Martin-Löf Type Theory. The first is Capretta's delay monad [Cap05], which uses coinduction. Arguably, the correct notion of equality of Capretta's delay monad is that of weak bisimilarity where two partial elements are considered equal when they are both undefined, or, when one of them is, so is the other and they have equal values in this case. This prompted the authors of [CUV19] to consider its quotient by weak bisimilarity, but they use countable choice to show that quotient is again a monad. Again using countable choice, they show that their quotient yields free pointed ω -complete posets (ω -cpo). In [ADK17] the authors use a so-called quotient inductive-inductive type (QIIT) to construct the free pointed ω -cpo, essentially by definition of the QIIT. It was shown in [CUV19] that a simpler higher inductive type actually suffices. Regardless, we stress that our approach yields free dcpo's as opposed to ω -cpo's and does not use countable choice or higher inductive types other than the propositional truncation. For a discussion on ω -completeness, countable choice and their relations to the Scott model of PCF, see [dJon21, Section 8] and [dJon22, Section 5.2.4].

Definition 5.3 (Lifting, partial element, $\mathcal{L}_\mathcal{V}(X)$; [EK17, Section 2.2]). We define the type of *partial elements* of a type $X : \mathcal{U}$ with respect to a universe \mathcal{V} as

$$\mathcal{L}_\mathcal{V}(X) \equiv \Sigma_{P:\Omega_\mathcal{V}}(P \rightarrow X)$$

and we also call it the *lifting* of X with respect to \mathcal{V} .

Every (total) element of X gives rise to a partial element of X through the following map, which will be shown to be the unit of the monad later.

Definition 5.4 (η_X). The map $\eta_X : X \rightarrow \mathcal{L}_{\mathcal{V}}(X)$ is defined by mapping x to the tuple $(\mathbf{1}_{\mathcal{V}}, \lambda u . x)$, where we have omitted the witness that $\mathbf{1}_{\mathcal{V}}$ is a subsingleton. We sometimes omit the subscript in η_X .

Besides these total elements, the lifting has another distinguished element that will be the least with respect to the order with which we shall equip the lifting.

Definition 5.5 (\perp). For every type $X : \mathcal{U}$ and universe \mathcal{V} , we denote the element $(\mathbf{0}_{\mathcal{V}}, \varphi) : \mathcal{L}_{\mathcal{V}}(X)$ by \perp . (Here φ is the unique map from $\mathbf{0}_{\mathcal{V}}$ to X .)

Next we introduce appropriate names for the projections from the type of partial elements.

Definition 5.6 (is-defined and value). We write $\text{is-defined} : \mathcal{L}_{\mathcal{V}}(X) \rightarrow \Omega_{\mathcal{V}}$ for the first projection and $\text{value} : \Pi_{l : \mathcal{L}_{\mathcal{V}}(X)} (\text{is-defined}(l) \rightarrow X)$ for the second projection.

Thus, with this terminology, the element \star witnesses that $\eta(x)$ is defined with value x for every $x : X$, while \perp is not defined because $\text{is-defined}(\perp)$ is the empty type.

Excluded middle says exactly that such elements are the only elements of the lifting of a type X , as the following proposition shows. Thus, the lifting generalises the classical construction of adding a new element.

Proposition 5.7 ([EK17, Section 2.2]). *The map $X + \mathbf{1} \xrightarrow{[\eta, \text{const}_{\perp}]} \mathcal{L}_{\mathcal{V}}(X)$ is an equivalence for every type $X : \mathcal{U}$ if and only if excluded middle in \mathcal{V} holds.*

Proof. Excluded middle in \mathcal{V} is equivalent to the map $[\text{const}_{\mathbf{0}}, \text{const}_{\mathbf{1}}] : \mathbf{2}_{\mathcal{V}} \rightarrow \Omega_{\mathcal{V}}$ being an equivalence. But if that map is an equivalence, then it follows that the map $[\eta, \text{const}_{\perp}] : X + \mathbf{1} \rightarrow \mathcal{L}_{\mathcal{V}}(X)$ is also an equivalence for every type X . Conversely, we can take $X \equiv \mathbf{1}_{\mathcal{V}}$ to see that $[\text{const}_{\mathbf{0}}, \text{const}_{\mathbf{1}}] : \mathbf{2}_{\mathcal{V}} \rightarrow \Omega_{\mathcal{V}}$ must be an equivalence. \square

Lemma 5.8. *Two partial elements $l, m : \mathcal{L}_{\mathcal{V}}(X)$ of a type X are equal if and only if we have $\text{is-defined}(l) \iff \text{is-defined}(m)$ and the diagram*

$$\begin{array}{ccc}
 \text{is-defined}(m) & \xrightarrow{\text{value}(m)} & X \\
 & \searrow & \nearrow \\
 & & \text{is-defined}(l) \\
 & & \text{value}(l)
 \end{array}$$

commutes.

Proof. By the characterisation of the identity type of Σ -types [Uni13, Theorem 2.7.2], we have

$$(l = m) \simeq (\Sigma_{e : \text{is-defined}(l) = \text{is-defined}(m)} \text{transport}^{\lambda P.P \rightarrow X} (e, \text{value}(l)) = \text{value}(m))$$

By path induction on e we can prove that

$$\text{transport}^{\lambda P.P \rightarrow X} (e, \text{value}(l)) = \text{value}(l) \circ \tilde{e}^{-1},$$

where \tilde{e} is the equivalence $\text{is-defined}(l) \simeq \text{is-defined}(m)$ induced by e . Hence, using function extensionality and propositional extensionality, the right hand side of the equivalence given above is logically equivalent to

$$\Sigma_{(e_1, e_2) : \text{is-defined}(l) \leftrightarrow \text{is-defined}(m)} \text{value}(l) \circ e_2 \sim \text{value}(m),$$

as desired. \square

Remark 5.9. It is possible to promote the logical equivalence of Lemma 5.8 to an equivalence of types using univalence and a generalised *structure identity principle* [Esc19, Section 3.33], as done in [Esc21, Lemma 44] and formalised in [Esc+, Lifting.IdentityViaSIP]. But the above logical equivalence will suffice for our purposes.

Theorem 5.10 (Lifting monad, Kleisli extension, $f^\#$; [EK17, Section 2.2]). *The lifting is a monad with unit η . That is, for every map $f : X \rightarrow \mathcal{L}_V(Y)$ we have a map $f^\# : \mathcal{L}_V(X) \rightarrow \mathcal{L}_V(Y)$, the Kleisli extension of f , such that*

- (i) $\eta_X^\# \sim \text{id}_{\mathcal{L}_V(X)}$ for every type X ,
- (ii) $f^\# \circ \eta_X \sim f$ for every map $f : X \rightarrow \mathcal{L}_V(Y)$, and
- (iii) $(g^\# \circ f)^\# \sim g^\# \circ f^\#$ for every two maps $f : X \rightarrow \mathcal{L}_V(Y)$ and $g : Y \rightarrow \mathcal{L}_V(Z)$.

Proof. Given $f : X \rightarrow \mathcal{L}_V(Y)$, we define

$$\begin{aligned} f^\# : \mathcal{L}_V(X) &\rightarrow \mathcal{L}_V(Y) \\ (P, \varphi) &\mapsto (\Sigma_{p:P} \text{is-defined}(f(\varphi(p))), \psi), \end{aligned}$$

where $\psi(p, q) \equiv \text{value}(f(\varphi(p)), q)$.

Now for the proof of (i): Let $(P, \varphi) : \mathcal{L}_V(X)$ be arbitrary and we calculate that

$$\begin{aligned} \eta^\#(P, \varphi) &\equiv (\Sigma_{p:P} \text{is-defined}(\eta(\varphi(p))), \lambda(p, q) \cdot \text{value}(\eta(\varphi(p)), q)) \\ &\equiv (P \times \mathbf{1}, \lambda(p, q) \cdot \varphi(p)) \\ &= (P, \varphi), \end{aligned}$$

where the final equality is seen to hold using Lemma 5.8. For (ii), let $x : X$ and $f : X \rightarrow \mathcal{L}_V(Y)$ be arbitrary and observe that

$$\begin{aligned} f^\#(\eta(x)) &\equiv f^\#(\mathbf{1}, \lambda u \cdot x) \\ &\equiv (\mathbf{1} \times \text{is-defined}(f(x)), \lambda(u, p) \cdot \text{value}(f(x), p)) \\ &= (\text{is-defined}(f(x)), \lambda p \cdot \text{value}(f(x), p)) \\ &\equiv f(x) \end{aligned}$$

where the penultimate equality is another easy application of Lemma 5.8. We see that these proofs amount to the fact that $\mathbf{1}$ is the unit for taking the product of types. For (iii) the proof amounts to the associativity of Σ . \square

Remark 5.11. It should be noted that if $X : \mathcal{U}$, then $\mathcal{L}_V(X) : \mathcal{V}^+ \sqcup \mathcal{U}$, so in general the lifting is a monad “across universes”. But this increase in universes does not hinder us in stating and proving the monad laws and using them in later proofs.

Remark 5.12. The equalities of Theorem 5.10 do not include any coherence conditions which may be needed when X is not a set but a higher type. We will restrict to the lifting of sets, but the more general case is considered in [Esc21] where the coherence conditions are not needed for its goals either.

Definition 5.13 (Lifting functor, $\mathcal{L}_V(f)$). The functorial action of the lifting could be defined from the unit and Kleisli extension as $\mathcal{L}_V(f) \equiv (\eta_Y \circ f)^\#$ for $f : X \rightarrow Y$. But it is equivalent and easier to define $\mathcal{L}_V(f)$ directly by post-composition:

$$\mathcal{L}_V(f)(P, \varphi) \equiv (P, f \circ \varphi).$$

We now work towards showing that $\mathcal{L}_V(X)$ is the free pointed \mathcal{V} -dcpo on a set X .

Proposition 5.14. *The relation $\sqsubseteq : \mathcal{L}_V(X) \rightarrow \mathcal{L}_V(X) \rightarrow \mathcal{V}^+ \sqcup \mathcal{U}$ given by*

$$l \sqsubseteq m \equiv \text{is-defined}(l) \rightarrow l = m$$

is a partial order on $\mathcal{L}_\mathcal{V}(X)$ for every set $X : \mathcal{U}$. Moreover, it is equivalent to the more verbose relation

$$(P, \varphi) \sqsubseteq' (Q, \psi) \equiv \Sigma_{f:P \rightarrow Q} (\varphi \sim \psi \circ f)$$

that is valued in $\mathcal{V} \sqcup \mathcal{U}$.

Proof. Note that \sqsubseteq is subsingleton-valued because X is assumed to be a set. The other properties follow using Lemma 5.8. \square

In light of Remark 3.10, we carefully keep track of the universe parameters of the lifting in the following proposition.

Proposition 5.15 (cf. [EK17, Theorem 1]). *For a set $X : \mathcal{U}$, the lifting $\mathcal{L}_\mathcal{V}(X)$ ordered as in Proposition 5.14 is a pointed \mathcal{V} -dcpo. In full generality we have $\mathcal{L}_\mathcal{V}(X) : \mathcal{V}\text{-DCPO}_{\mathcal{V}+\perp\mathcal{U}, \mathcal{V}+\perp\mathcal{U}}$, but if $X : \mathcal{V}$, then $\mathcal{L}_\mathcal{V}(X)$ is locally small.*

Proof. By Proposition 5.14 we have a poset and it is clear that \perp from Definition 5.5 is its least element. Now let $(Q_{(-)}, \varphi_{(-)}) : I \rightarrow \mathcal{L}_\mathcal{V}(X)$ be a directed family with $I : \mathcal{V}$. We claim that the map $\Sigma_{i:I} Q_i \xrightarrow{(i,q) \mapsto \varphi_i(q)} X$ is constant. Indeed, given $i, j : I$ with $p : Q_i$ and $q : Q_j$, there exists $k : I$ such that $(Q_i, \varphi_i), (Q_j, \varphi_j) \sqsubseteq (Q_k, \varphi_k)$ by directedness of the family. But by definition of the order and the elements $p : Q_i$ and $q : Q_j$, this implies that $(Q_i, \varphi_i) = (Q_j, \varphi_j) = (Q_k, \varphi_k)$ which in particular tells us that $\varphi_i(p) = \varphi_j(q)$. Hence, by [Kra+17, Theorem 5.4], we have a (dashed) map ψ making the diagram

$$\begin{array}{ccc} \Sigma_{i:I} Q_i & \xrightarrow{(i,q) \mapsto \varphi_i(q)} & X \\ \downarrow |-| & & \uparrow \psi \\ \exists_{i:I} Q_i & & \end{array}$$

commute. We claim that $(\exists_{i:I} Q_i, \psi)$ is the least upper bound of the family. To see that it is an upper bound, let $j : I$ be arbitrary. By the commutative diagram and Proposition 5.14 we see that $(Q_j, \varphi_j) \sqsubseteq (\exists_{i:I} Q_i, \psi)$, as desired. Moreover, if (P, ρ) is an upper bound for the family, then $(Q_i, \varphi_i) = (P, \rho)$ for all $i : I$ such that Q_i holds. Hence, $(\exists_{i:I} Q_i, \psi) \sqsubseteq (P, \rho)$, as desired. Finally, local smallness in the case that X is a type in \mathcal{V} follows from Proposition 5.14. \square

Proposition 5.16. *The Kleisli extension $f^\# : \mathcal{L}_\mathcal{V}(X) \rightarrow \mathcal{L}_\mathcal{V}(Y)$ is Scott continuous for any map $f : X \rightarrow \mathcal{L}_\mathcal{V}(Y)$.*

Proof. It is straightforward to prove that $f^\#$ is monotone. Hence, it remains to prove that $f^\#(\bigsqcup \alpha) \sqsubseteq \bigsqcup f^\# \circ \alpha$ for every directed family $\alpha : I \rightarrow \mathcal{L}_\mathcal{V}(X)$. So suppose that $f^\#(\bigsqcup \alpha)$ is defined. Then we have to show that it equals $\bigsqcup f^\# \circ \alpha$. By our assumption and definition of $f^\#$ we get that $\bigsqcup \alpha$ is defined too. By the definition of suprema in the lifting and because we are proving a proposition, we may assume to have $i : I$ such that α_i is defined. But since $\alpha_i \sqsubseteq \bigsqcup \alpha$, we get $\alpha_i = \bigsqcup \alpha$ and hence, $f^\#(\alpha_i) = f^\#(\bigsqcup \alpha)$. Finally, $f^\#(\alpha_i) \sqsubseteq \bigsqcup f^\# \circ \alpha$, but by assumption $f^\#(\bigsqcup \alpha)$ is defined and hence so is $f^\#(\alpha_i)$ which implies $f^\#(\bigsqcup \alpha) = f^\#(\alpha_i) = \bigsqcup f^\# \circ \alpha$, as desired. \square

Recall from Lemma 4.9 that pointed \mathcal{V} -dcpos have suprema of families indexed by propositions in \mathcal{V} . We make use of this fact in the following lemma.

Lemma 5.17. *For a set X , every partial element $(P, \varphi) : \mathcal{L}_\mathcal{V}(X)$ is equal to supremum $\bigvee_{p:P} \eta_X(\varphi(p))$.*

Proof. Note that if $p : P$, then $(P, \varphi) = \eta_X(\varphi(p))$, so that the lemma follows from antisymmetry. \square

The lifting $\mathcal{L}_{\mathcal{V}}(X)$ gives the free pointed \mathcal{V} -dcpo on a set X . Keeping track of universes, it holds in the following generality:

Theorem 5.18. *If $X : \mathcal{U}$ is a set, then for every pointed \mathcal{V} -dcpo $D : \mathcal{V}\text{-DCPO}_{\mathcal{U}', \mathcal{T}'}$ and function $f : X \rightarrow D$, there is a unique strict continuous function $\bar{f} : \mathcal{L}_{\mathcal{V}}(X) \rightarrow D$ making the diagram*

$$\begin{array}{ccc} X & \xrightarrow{f} & D \\ & \searrow \eta_X & \nearrow \bar{f} \\ & & \mathcal{L}_{\mathcal{V}}(X) \end{array}$$

commute.

Proof. We define $\bar{f} : \mathcal{L}_{\mathcal{V}}(X) \rightarrow D$ by $(P, \varphi) \mapsto \bigvee_{p:P} f(\varphi(p))$, which is well-defined by Lemma 4.9 and easily seen to be strict and continuous. For uniqueness, suppose that we have $g : \mathcal{L}_{\mathcal{V}}(X) \rightarrow D$ strict and continuous such that $g \circ \eta_X = f$ and let (P, φ) be an arbitrary element of $\mathcal{L}_{\mathcal{V}}(X)$. Then,

$$\begin{aligned} g(P, \varphi) &= g\left(\bigvee_{p:P} \eta_X(\varphi(p))\right) && \text{(by Lemma 5.17)} \\ &= \bigvee_{p:P} g(\eta_X(\varphi(p))) && \text{(by Lemma 4.9 and strictness and continuity of } g) \\ &= \bigvee_{p:P} f(\varphi(p)) && \text{(by assumption on } g) \\ &\equiv \bar{f}(P, \varphi), \end{aligned}$$

as desired. \square

Finally, we consider a variation of Proposition 5.14 which allows us to freely add a least element to a \mathcal{V} -dcpo instead of just a set.

Proposition 5.19. *For a poset D whose order takes values in \mathcal{T} , the binary relation $\sqsubseteq : \mathcal{L}_{\mathcal{V}}(D) \rightarrow \mathcal{L}_{\mathcal{V}}(D) \rightarrow \mathcal{V} \sqcup \mathcal{T}$ given by*

$$(P, \varphi) \sqsubseteq (Q, \psi) \equiv \Sigma_{f:P \rightarrow Q} (\Pi_{p:P} (\varphi(p) \sqsubseteq_D \psi(f(p))))$$

is a partial order on $\mathcal{L}_{\mathcal{V}}(D)$.

Proof. Similar to Proposition 5.14, but using that \sqsubseteq_D is reflexive, transitive and antisymmetric. \square

Proposition 5.20. *For a dcpo $D : \mathcal{V}\text{-DCPO}_{\mathcal{U}, \mathcal{T}}$, the lifting $\mathcal{L}_{\mathcal{V}}(D)$ ordered as in Proposition 5.19 is a pointed \mathcal{V} -dcpo. In general, $\mathcal{L}_{\mathcal{V}}(D) : \mathcal{V}\text{-DCPO}_{\mathcal{V} \sqcup \mathcal{U}, \mathcal{V} \sqcup \mathcal{T}}$, but if D is locally small, then so is $\mathcal{L}_{\mathcal{V}}(D)$.*

Proof. The element \perp from Definition 5.5 is still the least element with respect to the new order. If $\alpha : I \rightarrow \mathcal{L}_{\mathcal{V}}(D)$ is directed, then we consider $\Phi : (\Sigma_{i:I} Q_i) \rightarrow D$ given by $(i, q) \mapsto \varphi_i(q)$. This family is semidirected, for if we have $i, j : I$ with $p : Q_i$ and $q : Q_j$, then there exists $k : I$ such that $\alpha_i, \alpha_j \sqsubseteq \alpha_k$ in $\mathcal{L}_{\mathcal{V}}(D)$ by directedness of α , which implies that $\Phi(i, p) \sqsubseteq \Phi(j, q)$ in D . Thus, if we know that $\exists_{i:I} Q_i$, then the family Φ is directed and must have a supremum in D . Hence we have a partial element $(\exists_{i:I} Q_i, \psi) : \mathcal{L}_{\mathcal{V}}(D)$ where ψ takes the witness that the domain of Φ is inhabited to the directed supremum $\bigsqcup \Phi$ in D . It is not hard to verify that this partial element is the least upper bound of α in $\mathcal{L}_{\mathcal{V}}(D)$, completing the proof. \square

The lifting $\mathcal{L}_{\mathcal{V}}(D)$ with the partial order of Proposition 5.19 gives the free *pointed* \mathcal{V} -dcpo on a \mathcal{V} -dcpo D . Keeping track of universes, it holds in the following generality:

Theorem 5.21. *If $D : \mathcal{V}\text{-DCPO}_{\mathcal{U},\mathcal{T}}$ is a \mathcal{V} -dcpo, then for every pointed \mathcal{V} -dcpo $E : \mathcal{V}\text{-DCPO}_{\mathcal{U}',\mathcal{T}'}$ and continuous function $f : D \rightarrow E$, there is a unique strict continuous function $\bar{f} : \mathcal{L}_{\mathcal{V}}(D) \rightarrow E$ making the diagram*

$$\begin{array}{ccc} D & \xrightarrow{f} & E \\ \eta_D \searrow & & \nearrow \bar{f} \\ & \mathcal{L}_{\mathcal{V}}(D) & \end{array}$$

commute.

Proof. Similar to the proof of Theorem 5.18. \square

Notice how Theorem 5.21 generalises Theorem 5.18 as any set can be viewed as a discretely ordered \mathcal{V} -dcpo.

6. PRODUCTS AND EXPONENTIALS

We describe two constructions of \mathcal{V} -dcpos, namely products and exponentials. Exponentials will be crucial in Scott's D_∞ construction, as discussed in Section 8. Products are not needed for this purpose, but they allow us to state the universal property of the exponential (Proposition 6.7).

Definition 6.1 (Product of (pointed) dcpos, $D_1 \times D_2$). The *product* of two \mathcal{V} -dcpos D_1 and D_2 is given by the \mathcal{V} -dcpo $D_1 \times D_2$ defined as follows. Its carrier is the cartesian product of the carriers of D_1 and D_2 . The order is given componentwise, i.e. $(x, y) \sqsubseteq_{D_1 \times D_2} (x', y')$ if $x \sqsubseteq_{D_1} x'$ and $y \sqsubseteq_{D_2} y'$. Accordingly directed suprema are also given componentwise. That is, given a directed family $\alpha : I \rightarrow D_1 \times D_2$, one quickly verifies that the families $\text{pr}_1 \circ \alpha$ and $\text{pr}_2 \circ \alpha$ are also directed. We then define the supremum of α as $(\bigsqcup \text{pr}_1 \circ \alpha, \bigsqcup \text{pr}_2 \circ \alpha)$. Moreover, if D and E are pointed, then so is $D \times E$ by taking the least elements in both components.

Remark 6.2. Notice that if $D_1 : \mathcal{V}\text{-DCPO}_{\mathcal{U},\mathcal{T}}$ and $D_2 : \mathcal{V}\text{-DCPO}_{\mathcal{U}',\mathcal{T}'}$, then for their product we have $D_1 \times D_2 : \mathcal{V}\text{-DCPO}_{\mathcal{U} \sqcup \mathcal{U}', \mathcal{T} \sqcup \mathcal{T}'}$, which simplifies to $\mathcal{V}\text{-DCPO}_{\mathcal{U},\mathcal{T}}$ when $\mathcal{U}' \equiv \mathcal{U}$ and $\mathcal{T}' \equiv \mathcal{T}$.

Proposition 6.3. *The product defined above satisfies the appropriate universal property: the projections $\text{pr}_1 : D_1 \times D_2 \rightarrow D_1$ and $\text{pr}_2 : D_1 \times D_2 \rightarrow D_2$ are Scott continuous and if $f : E \rightarrow D_1$ and $g : E \rightarrow D_2$ are Scott continuous functions from a \mathcal{V} -dcpo E , then there is a unique Scott continuous map $k : E \rightarrow D_1 \times D_2$ such that the diagram*

$$\begin{array}{ccc} & D_1 \times D_2 & \\ \text{pr}_1 \swarrow & \uparrow & \searrow \text{pr}_2 \\ D_1 & k & D_2 \\ f \swarrow & \uparrow & \nearrow g \\ & E & \end{array}$$

commutes.

Proof. The projections are Scott continuous by definition of directed suprema in $D_1 \times D_2$. Moreover, if $f : E \rightarrow D_1$ and $g : E \rightarrow D_2$ are Scott continuous maps, then we see that we have no choice but to define $k : E \rightarrow D_1 \times D_2$ by $e \mapsto (f(e), g(e))$. Moreover, this assignment is Scott continuous, because for a directed family $\alpha : I \rightarrow E$, we have $k(\bigsqcup \alpha) \equiv (f(\bigsqcup \alpha), g(\bigsqcup \alpha)) = (\bigsqcup f \circ \alpha, \bigsqcup g \circ \alpha) \equiv \bigsqcup k \circ \alpha$ by Scott continuity of f and g and the definition of directed suprema in $D_1 \times D_2$. \square

Lemma 6.4. *A map $f : D_1 \times D_2 \rightarrow E$ is Scott continuous if and only if the maps $f(x, -) : D_2 \rightarrow E$ and $f(-, y) : D_1 \rightarrow E$ are Scott continuous for every $x : D_1$ and $y : D_2$.*

Proof. Straightforward as in the classical case. \square

Definition 6.5 (Exponential of (pointed) dcpos, E^D). The *exponential* of two \mathcal{V} -dcpos D and E is given by the poset E^D defined as follows. Its carrier is the type of Scott continuous functions from D to E . The order is given pointwise, i.e. $f \sqsubseteq_{E^D} g$ holds if $f(x) \sqsubseteq_E g(x)$ for every $x : D$. Notice that if E is pointed, then so is E^D with least element given the constant function $\lambda x : D . \perp_E$ which is Scott continuous by Proposition 4.10(ii). Finally, it is straightforward to show that E^D is \mathcal{V} -directed complete, so that E^D is another \mathcal{V} -dcpo.

Note that the exponential E^D is a priori not locally small even if E is because the partial order quantifies over all elements of D . But if D is continuous (a notion that we study in detail in the Part II paper [dJE24]) then E^D will be locally small when E is.

Remark 6.6. Recall from Remark 3.10 that it is necessary to carefully keep track of the universe parameters of the exponential. In general, the universe levels of E^D can be quite large and complicated. For if $D : \mathcal{V}\text{-DCPO}_{\mathcal{U}, \mathcal{T}}$ and $E : \mathcal{V}\text{-DCPO}_{\mathcal{U}', \mathcal{T}'}$, then the exponential E^D has a carrier in the universe

$$\mathcal{V}^+ \sqcup \mathcal{U} \sqcup \mathcal{T} \sqcup \mathcal{U}' \sqcup \mathcal{T}'$$

and an order relation that takes values in $\mathcal{U} \sqcup \mathcal{T}'$.

Even if $\mathcal{V} = \mathcal{U} \equiv \mathcal{T} \equiv \mathcal{U}' \equiv \mathcal{T}'$, the carrier of E^D still lives in the larger universe \mathcal{V}^+ , because the type expressing Scott continuity for \mathcal{V} -dcpos quantifies over all types in \mathcal{V} . Actually, the scenario where $\mathcal{U} = \mathcal{U}' = \mathcal{V}$ cannot happen in a predicative setting unless D and E are trivial, in a sense made precise in [dJE23].

Even so, in many applications such as those in [dJon21] or Section 8, if we take $\mathcal{V} \equiv \mathcal{U}_0$ and all other parameters to be $\mathcal{U} \equiv \mathcal{T} \equiv \mathcal{U}' \equiv \mathcal{T}' \equiv \mathcal{U}_1$, then the situation is much simpler and D , E and the exponential E^D are all elements of $\mathcal{U}_0\text{-DCPO}_{\mathcal{U}_1, \mathcal{U}_1}$ with all of them being locally small (remember that this is defined up to equivalence). This turns out to be a very favourable situation for both the Scott model of PCF [dJon21] and Scott's D_∞ model of the untyped λ -calculus (Section 8).

In the proposition below we can have $D : \mathcal{V}\text{-DCPO}_{\mathcal{U}, \mathcal{T}}$ and $E : \mathcal{V}\text{-DCPO}_{\mathcal{U}', \mathcal{T}'}$ for arbitrary universes \mathcal{U} , \mathcal{T} , \mathcal{U}' and \mathcal{T}' . In particular, the universe parameters of D and E , apart from the universe of indexing types, need not be the same.

Proposition 6.7. *The exponential defined above satisfies the appropriate universal property: the evaluation map $\text{ev} : E^D \times D \rightarrow E$, $(g, x) \mapsto g(x)$ is Scott continuous and if $f : D' \times D \rightarrow E$ is a Scott continuous function, then there is a unique Scott continuous map $\bar{f} : D' \rightarrow E^D$ such that the diagram*

$$\begin{array}{ccc} D' \times D & & \\ \bar{f} \times \text{id}_D \downarrow & \searrow f & \\ E^D \times D & \xrightarrow{\text{ev}} & E \end{array}$$

commutes.

Proof. Straightforward application of Lemma 6.4, as in the classical case. \square

For completeness, we recall the following theorem which lies at the heart of the Scott model of PCF studied in [dJon21]:

Theorem 6.8 (Least fixed point, μ). *Every Scott continuous endomap f on a pointed \mathcal{V} -dcpo D has a least fixed point given by*

$$\mu(f) := \bigsqcup_{n \in \mathbb{N}} f^n(\perp).$$

Specifically, the following two conditions hold:

- (i) $f(\mu(f)) = \mu(f)$, and
- (ii) for every $x : D$, if $f(x) \sqsubseteq x$, then $\mu(f) \sqsubseteq x$.

Moreover, the assignment $f \mapsto \mu(f)$ defines a Scott continuous map $D^D \rightarrow D$.

Proof. Following the proof [AJ94, Theorem 2.1.19]; see [dJon21, Theorem 16] for details. \square

7. BILIMITS

Recall that in a \mathcal{V} -dcpo D we can take suprema of directed families $\alpha : I \rightarrow D$. It is a striking feature of directed complete posets that this act is reflected in the *category* of dcpos, although it does require us to specify an appropriate notion of one dcpo being “below” another one. This notion will be exactly that of an embedding-projection pair. The technical results developed in this section will find application in the construction of Scott’s D_∞ , a model of the untyped λ -calculus, as discussed in Section 8. We refer the reader to the conclusion (Section 9) for a discussion on how our predicative reconstruction of D_∞ compares Scott’s original work [Sco72].

Definition 7.1 (Deflation). An endofunction $f : D \rightarrow D$ on a poset D is a *deflation* if $f(x) \sqsubseteq x$ for all $x : D$.

Definition 7.2 (Embedding-projection pair). An *embedding-projection pair* from a \mathcal{V} -dcpo D to a \mathcal{V} -dcpo E consists of two Scott continuous functions $\varepsilon : D \rightarrow E$ (the *embedding*) and $\pi : E \rightarrow D$ (the *projection*) such that:

- (i) ε is a section of π , and
- (ii) $\varepsilon \circ \pi$ is a deflation.

For the remainder of this section, fix the following setup, where we try to be as general regarding universe levels as we can be. We fix a directed preorder (I, \sqsubseteq) with $I : \mathcal{V}$ and \sqsubseteq takes values in some universe \mathcal{W} . Now suppose that (I, \sqsubseteq) indexes a family of \mathcal{V} -dcpos with embedding-projection pairs between them, i.e. we have

- for every $i : I$, a \mathcal{V} -dcpo $D_i : \mathcal{V}\text{-DCPO}_{\mathcal{U}, \mathcal{T}}$, and
- for every $i, j : I$ with $i \sqsubseteq j$, an embedding-projection pair $(\varepsilon_{i,j}, \pi_{i,j})$ from D_i to D_j .

Moreover, we require that the following compatibility conditions hold:

$$\text{for every } i : I, \text{ we have } \varepsilon_{i,i} = \pi_{i,i} = \text{id}; \tag{1}$$

$$\text{for every } i \sqsubseteq j \sqsubseteq k \text{ in } I, \text{ we have } \varepsilon_{i,k} \sim \varepsilon_{j,k} \circ \varepsilon_{i,j} \text{ and } \pi_{i,k} \sim \pi_{i,j} \circ \pi_{j,k}. \tag{2}$$

Example 7.3. If $I \equiv \mathbb{N}$ with the usual ordering, then we are looking at a diagram of \mathcal{V} -dcpos like this

$$D_0 \begin{array}{c} \xrightarrow{\varepsilon_{0,1}} \\ \xleftarrow{\pi_{0,1}} \end{array} D_1 \begin{array}{c} \xrightarrow{\varepsilon_{1,2}} \\ \xleftarrow{\pi_{1,2}} \end{array} D_2 \begin{array}{c} \xrightarrow{\varepsilon_{2,3}} \\ \xleftarrow{\pi_{2,3}} \end{array} D_3 \begin{array}{c} \xrightarrow{\varepsilon_{3,4}} \\ \xleftarrow{\pi_{3,4}} \end{array} \cdots$$

where, for example, we have not pictured $\varepsilon_{1,1} : D_1 \hookrightarrow D_1$ and $\varepsilon_{0,2} : D_0 \hookrightarrow D_2$ explicitly, as they are equal to $\text{id}_{D_1} : D_1 \rightarrow D_1$ and the composition of $D_0 \xrightarrow{\varepsilon_{0,1}} D_1$ and $D_1 \xrightarrow{\varepsilon_{1,2}} D_2$, respectively.

The goal is now to construct another \mathcal{V} -dcpo D_∞ with embedding-projections pairs $(\varepsilon_{i,\infty} : D_1 \hookrightarrow D_\infty, \pi_{i,\infty} : D_\infty \rightarrow D_i)$ for every $i : I$, such that $(D_\infty, (\varepsilon_{i,\infty})_{i:I})$ is the colimit of the diagram given by $(\varepsilon_{i,j})_{i \sqsubseteq j \text{ in } I}$ and $(D_\infty, (\pi_{i,\infty})_{i:I})$ is the limit of the diagram given by $(\pi_{i,j})_{i \sqsubseteq j \text{ in } I}$. In other words, $(D_\infty, (\varepsilon_{i,\infty})_{i:I}, (\pi_{i,\infty})_{i:I})$ is both the colimit and the limit in the category of \mathcal{V} -dcpos with embedding-projections pairs between them. We say that it is the *bilimit*.

Definition 7.4 (D_∞). We define a poset D_∞ as follows. Its carrier is given by dependent functions $\sigma : \prod_{i:I} D_i$ satisfying $\pi_{i,j}(\sigma_j) = \sigma_i$ whenever $i \sqsubseteq j$. That is, the carrier is the type

$$\sum_{\sigma : \prod_{i:I} D_i} \prod_{i,j:I, i \sqsubseteq j} \pi_{i,j}(\sigma_j) = \sigma_i.$$

Note that this defines a subtype of $\prod_{i:I} D_i$ as the condition $\prod_{i,j:I, i \sqsubseteq j} \pi_{i,j}(\sigma_j) = \sigma_i$ is a property by [Uni13, Example 3.6.2] and the fact that each D_i is a set.

These functions are ordered pointwise, i.e. if $\sigma, \tau : \prod_{i:I} D_i$, then $\sigma \sqsubseteq_{D_\infty} \tau$ exactly when $\sigma_i \sqsubseteq_{D_i} \tau_i$ for every $i : I$.

Lemma 7.5. *The poset D_∞ is \mathcal{V} -directed complete with suprema calculated pointwise. Paying attention to the universe levels involved, we have $D_\infty : \mathcal{V}\text{-DCPO}_{\mathcal{U} \sqcup \mathcal{V} \sqcup \mathcal{W}, \mathcal{U} \sqcup \mathcal{T}}$.*

Proof. If $\alpha : A \rightarrow D_\infty$ is a directed family, then the family $\alpha_i : A \rightarrow D_i$ given by $\alpha_i(a) \equiv (\alpha(a))_i$ is directed again, and we define the supremum of α in D_∞ as the function $i \mapsto \bigsqcup \alpha_i$. To see that this indeed defines an element of D_∞ , observe that for every $i, j : I$ with $i \sqsubseteq j$ we have

$$\begin{aligned} \pi_{i,j}(\bigsqcup \alpha_j) &\equiv \pi_{i,j}(\bigsqcup \alpha_j) \\ &= \bigsqcup \pi_{i,j} \circ \alpha_j && \text{(by Scott continuity of } \pi_{i,j}\text{)} \\ &\equiv \bigsqcup_{a:A} (\pi_{i,j}((\alpha(a))_j)) \\ &= \bigsqcup \alpha_i && \text{(as } \alpha(a) \text{ is an element of } D_\infty\text{),} \end{aligned}$$

as desired. \square

Remark 7.6. We allow for general universe levels here, which is why D_∞ lives in the relatively complicated universe $\mathcal{U} \sqcup \mathcal{V} \sqcup \mathcal{W}$. In concrete examples, the situation often simplifies. E.g., in Section 8 we find ourselves in the favourable situation described in Remark 6.6 where $\mathcal{V} \equiv \mathcal{W} \equiv \mathcal{U}_0$ and $\mathcal{U} \equiv \mathcal{T} \equiv \mathcal{U}_1$, so that we get $D_\infty : \mathcal{U}_0\text{-DCPO}_{\mathcal{U}_1, \mathcal{U}_1}$, as the bilimit of a diagram of dcpos $D_n : \mathcal{U}_0\text{-DCPO}_{\mathcal{U}_1, \mathcal{U}_1}$ indexed by natural numbers.

Definition 7.7 ($\pi_{i,\infty}$). For every $i : I$, we define the Scott continuous function $\pi_{i,\infty} : D_\infty \rightarrow D_i$ by $\sigma \mapsto \sigma_i$.

Lemma 7.8. *The map $\pi_{i,\infty} : D_\infty \rightarrow D_i$ is Scott continuous for every $i : I$.*

Proof. This holds because suprema in D_∞ are calculated pointwise and $\pi_{i,\infty}$ selects the i -th component. \square

While we could closely follow [Sco72] up until this point, we will now need a new idea to proceed. Our goal is to define maps $\varepsilon_{i,\infty} : D_i \rightarrow D_\infty$ for every $i : I$ so that $\varepsilon_{i,\infty}$ and $\pi_{i,\infty}$ form an embedding-projection pair. We give an outline of the idea for defining this map $\varepsilon_{i,\infty}$. For an arbitrary element $x : D_i$, we need to construct $\sigma : D_\infty$ at component $j : I$, say. If we had $k : I$ such that $i, j \sqsubseteq k$, then we could define $\sigma_j : D_j$ by $\pi_{j,k}(\varepsilon_{i,k}(x))$. Now semidirectedness of I tells us that there exists such a $k : I$, so the point is to somehow make use of this propositionally truncated fact. This is where [Kra+17, Theorem 5.4] comes in. We define a map $\kappa_{i,j}^x : (\sum_{k:I} (i \sqsubseteq k) \times (j \sqsubseteq k)) \rightarrow D_j$ by sending k to $\pi_{j,k}(\varepsilon_{i,k}(x))$ and show it to be

constant, so that it factors through the truncation of its domain. In the special case that $I \equiv \mathbb{N}$, as in [Sco72], we could simply take k to be the sum of the natural numbers i and j , but this does not work in the more general directed case, of course.

Definition 7.9 ($\kappa_{i,j}^x$). For every $i, j : I$ and $x : D_i$ we define the function

$$\kappa_{i,j}^x : (\Sigma_{k:I} (i \sqsubseteq k) \times (j \sqsubseteq k)) \rightarrow D_j$$

by mapping $k : I$ with $i, j \sqsubseteq k$ to $\pi_{j,k}(\varepsilon_{i,k}(x))$.

Lemma 7.10. *The function $\kappa_{i,j}^x$ is constant for every $i, j : I$ and $x : D_i$. Hence, $\kappa_{i,j}^x$ factors through $\exists_{k:I} (i \sqsubseteq k) \times (j \sqsubseteq k)$ by [Kra+17, Theorem 5.4].*

Proof. If we have $k_1, k_2 : I$ with $i \sqsubseteq k_1, k_2$ and $j \sqsubseteq k_1, k_2$, then by semidirectedness of I , there exists some $k : K$ with $k_1, k_2 \sqsubseteq k$ and hence,

$$\begin{aligned} (\pi_{j,k_1} \circ \varepsilon_{i,k_1})(x) &= (\pi_{j,k_1} \circ \pi_{k_1,k} \circ \varepsilon_{k_1,k} \circ \varepsilon_{i,k_1})(x) && \text{(since } \varepsilon_{k_1,k} \text{ is a section of } \pi_{k_1,k}\text{)} \\ &= (\pi_{j,k} \circ \varepsilon_{i,k})(x) && \text{(by Equation (2))} \\ &= (\pi_{j,k} \circ \pi_{k_2,k} \circ \varepsilon_{k_2,k} \circ \varepsilon_{i,k_2})(x) && \text{(since } \varepsilon_{k_2,k} \text{ is a section of } \pi_{k_2,k}\text{)} \\ &= (\pi_{j,k_2} \circ \varepsilon_{i,k_2})(x) && \text{(by Equation (2)),} \end{aligned}$$

proving that $\kappa_{i,j}^x$ is constant. \square

Definition 7.11 ($\rho_{i,j}$). For every $i, j : I$, the type $\exists_{k:I} (i \sqsubseteq k) \times (j \sqsubseteq k)$ has an element since (I, \sqsubseteq) is directed. Thus, Lemma 7.10 tells us that we have a function $\rho_{i,j} : D_i \rightarrow D_j$ such that if $i, j \sqsubseteq k$, then the equation

$$\rho_{i,j}(x) = \kappa_{i,j}^x(k) \equiv \pi_{j,k}(\varepsilon_{i,k}(x)) \quad (3)$$

holds for every $x : D_i$.

Definition 7.12 ($\varepsilon_{i,\infty}$). The map ρ induces a map $\varepsilon_{i,\infty} : D_i \rightarrow D_\infty$ by sending $x : D_i$ to the function $\lambda j : I. \rho_{i,j}(x)$. To see that this is well-defined, assume that we have $j_1 \sqsubseteq j_2$ in J and $x : D_i$. We have to show that $\pi_{j_1,j_2}((\varepsilon_{i,\infty}(x))_{j_2}) = (\varepsilon_{i,\infty}(x))_{j_1}$. By semidirectedness of I and the fact that we are looking to prove a proposition, we may assume to have $k : I$ with $i \sqsubseteq k$ and $j_1 \sqsubseteq j_2 \sqsubseteq k$. Then,

$$\begin{aligned} \pi_{j_1,j_2}((\varepsilon_{i,\infty}(x))_{j_2}) &\equiv \pi_{j_1,j_2}(\rho_{i,j_2}(x)) \\ &= \pi_{j_1,j_2}(\pi_{j_2,k}(\varepsilon_{i,k}(x))) && \text{(by Equation (3))} \\ &= \pi_{j_1,k}(\varepsilon_{i,k}(x)) && \text{(by Equation (2))} \\ &= \rho_{i,j_1}(x) && \text{(by Equation (3))} \\ &\equiv (\varepsilon_{i,\infty}(x))_{j_1}, \end{aligned}$$

as desired.

This completes the definition of $\varepsilon_{i,\infty}$. From this point on, we can typically work with it by using Equation (3) and the fact that $(\varepsilon_{i,\infty}(x))_j$ is defined as $\rho_{i,j}(x)$.

Lemma 7.13. *The map $\rho_{i,j} : D_i \rightarrow D_j$ is Scott continuous for every $i, j : I$.*

Proof. Since we are proving a property, we may use semidirectedness of I to get $k : I$ with $i, j \sqsubseteq k$. Then, $\rho_{i,j} \sim \pi_{j,k} \circ \varepsilon_{i,k}$ by Equation (3). But the functions $\pi_{j,k}$ and $\varepsilon_{i,k}$ are continuous and continuity is preserved by function composition, so $\rho_{i,j}$ is continuous, as we wished to show. \square

Lemma 7.14. *The map $\varepsilon_{i,\infty} : D_i \rightarrow D_\infty$ is Scott continuous for every $i : I$.*

Proof. If $\alpha : A \rightarrow D_i$ is directed, then for every $j : I$ we have

$$\begin{aligned} (\varepsilon_{i,\infty}(\bigsqcup \alpha))_j &\equiv \rho_{i,j}(\bigsqcup \alpha) \\ &= \bigsqcup \rho_{i,j} \circ \alpha && \text{(by Lemma 7.13)} \\ &\equiv \bigsqcup_{a:A} (\varepsilon_{i,\infty}(\alpha(a)))_j \\ &\equiv (\bigsqcup (\varepsilon_{i,\infty} \circ \alpha))_j && \text{(as suprema in } D_\infty \text{ are calculated pointwise).} \end{aligned}$$

Hence, $\varepsilon_{i,\infty}(\bigsqcup \alpha) = \bigsqcup (\varepsilon_{i,\infty} \circ \alpha)$ and $\varepsilon_{i,\infty}$ is seen to be Scott continuous. \square

Theorem 7.15. *For every $i : I$, the pair $(\varepsilon_{i,\infty}, \pi_{i,\infty})$ is an embedding-projection pair from D_i to D_∞ .*

Proof. Scott continuity of both maps is given by Lemmas 7.8 and 7.14. To see that $\varepsilon_{i,\infty}$ is a section of $\pi_{i,\infty}$, observe that for every $x : D_i$, we have

$$\begin{aligned} \pi_{i,\infty}(\varepsilon_{i,\infty}(x)) &\equiv (\varepsilon_{i,\infty}(x))_i \\ &\equiv \rho_{i,i}(x) \\ &= \pi_{i,i}(\varepsilon_{i,i}(x)) && \text{(by Equation (3))} \\ &\equiv x && \text{(by Equation (1)),} \end{aligned}$$

so that $\varepsilon_{i,\infty}$ is indeed a section of $\pi_{i,\infty}$. It remains to prove that $\varepsilon_{i,\infty}(\pi_{i,\infty}(\sigma)) \sqsubseteq \sigma$ for every $\sigma : D_\infty$. The order is given pointwise, so let $j : I$ be arbitrary and since we are proving a proposition, assume that we have $k : I$ with $i, j \sqsubseteq k$. Then,

$$\begin{aligned} (\varepsilon_{i,\infty}(\pi_{i,\infty}(\sigma)))_j &\equiv (\varepsilon_{i,\infty}(\sigma_i))_j \\ &\equiv \rho_{i,j}(\sigma_i) \\ &= \pi_{j,k}(\varepsilon_{i,k}(\sigma_i)) && \text{(by Equation (3))} \\ &= \pi_{j,k}(\varepsilon_{i,k}(\pi_{i,k}(\sigma_k))) && \text{(since } \sigma \text{ is an element of } D_\infty) \end{aligned}$$

But $\pi_{i,k} \circ \varepsilon_{i,k}$ is deflationary and $\pi_{j,k}$ is monotone, so

$$\begin{aligned} &\sqsubseteq \pi_{j,k}(\sigma_k) \\ &= \sigma_j && \text{(since } \sigma \text{ is an element of } D_\infty), \end{aligned}$$

finishing the proof. \square

Lemma 7.16. *The maps $\pi_{i,\infty}$ and $\varepsilon_{i,\infty}$ respectively commute with $\pi_{i,j}$ and $\varepsilon_{i,j}$ whenever $i \sqsubseteq j$, viz. the diagrams*

$$\begin{array}{ccc} D_\infty & \xrightarrow{\pi_{i,\infty}} & D_i \\ & \searrow \pi_{j,\infty} & \nearrow \pi_{i,j} \\ & D_j & \end{array} \qquad \begin{array}{ccc} D_i & \xrightarrow{\varepsilon_{i,\infty}} & D_\infty \\ & \searrow \varepsilon_{i,j} & \nearrow \varepsilon_{j,\infty} \\ & D_j & \end{array}$$

commute for all $i, j : I$ with $i \sqsubseteq j$.

Proof. If $i \sqsubseteq j$ and $\sigma : D_\infty$ is arbitrary, then

$$\pi_{i,j}(\pi_{j,\infty}(\sigma)) \equiv \pi_{i,j}(\sigma_j) = \sigma_i$$

precisely because σ is an element of D_∞ , which proves the commutativity of the first diagram. For the second, let $x : D_i$ be arbitrary and we compare $(\varepsilon_{j,\infty}(\varepsilon_{i,j}(x)))$ and $\varepsilon_{i,\infty}(x)$ componentwise. So let $j' : I$ be arbitrary. Since we are proving a proposition, we may assume to have $k : I$ with $j, j' \sqsubseteq k$ by semidirectedness of I . We now calculate that

$$\begin{aligned} (\varepsilon_{j,\infty}(\varepsilon_{i,j}(x)))_{j'} &\equiv \rho_{j,j'}(\varepsilon_{i,j}(x)) \\ &= \pi_{j',k}(\varepsilon_{j,k}(\varepsilon_{i,j}(x))) && \text{(by Equation (3))} \\ &= \pi_{j',k}(\varepsilon_{i,k}(x)) && \text{(by Equation (2))} \end{aligned}$$

$$\begin{aligned}
 &= \rho_{i,j'}(x) && \text{(by Equation (3))} \\
 &\equiv (\varepsilon_{i,\infty}(x))_{j'}
 \end{aligned}$$

as desired. \square

Theorem 7.17. *The \mathcal{V} -dcpo D_∞ with the maps $(\pi_{i,\infty})_{i:I}$ is the limit of the diagram $((D_i)_{i:I}, (\pi_{i,j})_{i \sqsubseteq j})$. That is, given a \mathcal{V} -dcpo $E : \mathcal{V}\text{-DCPO}_{\mathcal{U}', \mathcal{T}'}$ and Scott continuous functions $f_i : E \rightarrow D_i$ for every $i : I$ such that the diagram*

$$\begin{array}{ccc}
 E & \xrightarrow{f_i} & D_i \\
 & \searrow f_j & \nearrow \pi_{i,j} \\
 & & D_j
 \end{array} \tag{4}$$

commutes for every $i \sqsubseteq j$, we have a unique Scott continuous function $f_\infty : E \rightarrow D_\infty$ making the diagram

$$\begin{array}{ccc}
 E & \xrightarrow{f_i} & D_i \\
 & \searrow f_\infty & \nearrow \pi_{i,\infty} \\
 & & D_\infty
 \end{array} \tag{5}$$

commute for every $i : I$.

Proof. Note that Equation (5) dictates that we must have $(f_\infty(y))_i = f_i(y)$ for every $i : I$. Hence, we define $f_\infty : E \rightarrow D_\infty$ as $f_\infty(y) \equiv \lambda i : I. f_i(y)$, which is Scott continuous because each f_i is and suprema are calculated pointwise in D_∞ . To see that f_∞ is well-defined, i.e. that $f_\infty(y)$ is indeed an element of D_∞ , observe that for every $i \sqsubseteq j$, the equation $\pi_{i,j}((f_\infty(y))_j) \equiv \pi_{i,j}(f_j(y)) = f_i(y)$ holds because of Equation (4). \square

It should be noted that in the above universal property E can have its carrier in any universe \mathcal{U}' and its order taking values in any universe \mathcal{T}' , even though we required all D_i to have their carriers and orders in two fixed universes \mathcal{U} and \mathcal{T} , respectively.

Lemma 7.18. *If $i \sqsubseteq j$ in I , then $\varepsilon_{i,\infty}(\sigma_i) \sqsubseteq \varepsilon_{j,\infty}(\sigma_j)$ for every $\sigma : D_\infty$.*

Proof. The order of D_∞ is pointwise, so we compare $\varepsilon_{i,\infty}(\sigma_i)$ and $\varepsilon_{j,\infty}(\sigma_j)$ at an arbitrary component $k : I$. We may assume to have $m : I$ such that $j, k \sqsubseteq m$ by semidirectedness of I . We then calculate that

$$\begin{aligned}
 (\varepsilon_{i,\infty}(\sigma_i))_k &\equiv \rho_{i,k}(\sigma_i) \\
 &= (\pi_{k,m} \circ \varepsilon_{i,m})(\sigma_i) && \text{(by Equation (3))} \\
 &= (\pi_{k,m} \circ \varepsilon_{i,m} \circ \pi_{i,j})(\sigma_j) && \text{(since } \sigma \text{ is an element of } D_\infty) \\
 &= (\pi_{k,m} \circ \varepsilon_{j,m} \circ \varepsilon_{i,j} \circ \pi_{i,j})(\sigma_j) && \text{(by Equation (2))}
 \end{aligned}$$

But $\varepsilon_{i,j} \circ \pi_{i,j}$ is deflationary and $\pi_{k,m} \circ \varepsilon_{j,m}$ is monotone, so

$$\begin{aligned}
 &\sqsubseteq (\pi_{k,m} \circ \varepsilon_{j,m})(\sigma_j) \\
 &= \rho_{j,k}(\sigma_j) && \text{(by Equation (3))} \\
 &\equiv (\varepsilon_{j,\infty}(\sigma_j))_k,
 \end{aligned}$$

as we wished to show. \square

Lemma 7.19. *Every element $\sigma : D_\infty$ is equal to the directed supremum $\bigsqcup_{i:I} \varepsilon_{i,\infty}(\sigma_i)$.*

Proof. The domain of the family is inhabited, because (I, \sqsubseteq) is assumed to be directed. Moreover, if we have $i, j : I$, then there exists $k : I$ with $i, j \sqsubseteq k$, which implies $\varepsilon_{i,\infty}(\sigma_i), \varepsilon_{j,\infty}(\sigma_j) \sqsubseteq \varepsilon_{k,\infty}(\sigma_k)$ by Lemma 7.18. Thus, the family $i \mapsto \varepsilon_{i,\infty}(\sigma_i)$ is indeed directed. To see that its supremum is indeed σ we use antisymmetry at an arbitrary component $j : I$. Firstly, observe that

$$\begin{aligned} \sigma_j &= \pi_{j,j}(\varepsilon_{j,j}(\sigma_j)) && \text{(by Equation (1))} \\ &= \rho_{j,j}(\sigma_j) && \text{(by Equation (3))} \\ &\equiv (\varepsilon_{j,\infty}(\sigma_j))_j \\ &\sqsubseteq (\bigsqcup_{i:I} \varepsilon_{i,\infty}(\sigma_i))_j && \text{(since suprema are computed pointwise in } D_\infty\text{).} \end{aligned}$$

Secondly, to prove that $(\bigsqcup_{i:I} \varepsilon_{i,\infty}(\sigma_i))_j \sqsubseteq \sigma_j$ it suffices to show that $(\varepsilon_{i,\infty}(\sigma_i))_j \sqsubseteq \sigma_j$ for every $i : I$. But this just says that $\varepsilon_{i,\infty} \circ \pi_{i,\infty}$ is a deflation, which was proved in Theorem 7.15. \square

Although the composites $\varepsilon_{i,\infty} \circ \pi_{i,\infty}$ are deflations for each $i : I$, the supremum of all of them is the identity. This fact will come in useful in Section 8.

Lemma 7.20. *The family $i \mapsto \varepsilon_{i,\infty} \circ \pi_{i,\infty}$ is directed in the exponential $D_\infty^{D_\infty}$ and its supremum is the identity on D_∞ .*

Proof. The order and suprema are given pointwise in exponentials, so this follows from Lemma 7.19. \square

Theorem 7.21. *The \mathcal{V} -dcpo D_∞ with the maps $(\varepsilon_{i,\infty})_{i:I}$ is the colimit of the diagram $((D_i)_{i:I}, (\varepsilon_{i,j})_{i \sqsubseteq j})$. That is, given a \mathcal{V} -dcpo $E : \mathcal{V}\text{-DCPO}_{\mathcal{U}', \mathcal{T}'}$ and Scott continuous functions $g_i : D_i \rightarrow E$ for every $i : I$ such that the diagram*

$$\begin{array}{ccc} D_i & \xrightarrow{g_i} & E \\ & \searrow \varepsilon_{i,j} & \nearrow g_j \\ & & D_j \end{array} \quad (6)$$

commutes for every $i \sqsubseteq j$, we have a unique Scott continuous function $g_\infty : D_\infty \rightarrow E$ making the diagram

$$\begin{array}{ccc} D_i & \xrightarrow{g_i} & E \\ & \searrow \varepsilon_{i,\infty} & \nearrow g_\infty \\ & & D_\infty \end{array} \quad (7)$$

commute for every $i : I$.

Proof. Note that any such Scott continuous function g_∞ must satisfy

$$\begin{aligned} g_\infty(\sigma) &= g_\infty(\bigsqcup_{i:I} \varepsilon_{i,\infty}(\sigma_i)) && \text{(by Lemma 7.19)} \\ &= \bigsqcup_{i:I} g_\infty(\varepsilon_{i,\infty}(\sigma_i)) && \text{(as } g_\infty \text{ is assumed to be Scott continuous)} \\ &= \bigsqcup_{i:I} g_i(\sigma_i) && \text{(by Equation (7))} \end{aligned}$$

for every $\sigma : D_\infty$. Accordingly, we define g_∞ by $g_\infty(\sigma) := \bigsqcup_{i:I} g_i(\sigma_i)$, where we verify that the family is indeed directed: If we have $i, j : I$, then there exists $k : I$ with $i, j \sqsubseteq k$, and we have

$$\begin{aligned} g_i(\sigma_i) &= g_i(\pi_{i,k}(\sigma_k)) && \text{(since } \sigma \text{ is an element of } D_\infty\text{)} \\ &= g_k(\varepsilon_{i,k}(\pi_{i,k}(\sigma_k))) && \text{(by Equation (6))} \\ &\sqsubseteq g_k(\sigma_k) && \text{(since } \varepsilon_{i,k} \circ \pi_{i,k} \text{ is deflationary and } g_k \text{ is monotone),} \end{aligned}$$

and similarly, $g_j(\sigma_j) \sqsubseteq g_k(\sigma_k)$. To see that g_∞ satisfies Equation (7), let $x : D_i$ be arbitrary and first observe that

$$g_\infty(\varepsilon_{i,\infty}(x)) \equiv \bigsqcup_{j:I} g_j \left((\varepsilon_{i,\infty}(x))_j \right) \equiv \bigsqcup_{j:I} g_j(\rho_{i,j}(x)).$$

We now use antisymmetry to prove that this is equal to $g_i(x)$. In one direction this is easy as $g_i(x) = (g_i \circ \pi_{i,i} \circ \varepsilon_{i,i})(x) \equiv g_i(\rho_{i,i}(x)) \sqsubseteq \bigsqcup_{j:I} g_j(\rho_{i,j}(x))$. In the other direction, it suffices to prove that $g_j(\rho_{i,j}(x)) \sqsubseteq g_i(x)$ for every $j : I$. By directedness of I there exists $k : I$ with $i, j \sqsubseteq k$ so that

$$g_j(\rho_{i,j}(x)) = (g_j \circ \pi_{j,k} \circ \varepsilon_{i,k})(x) \quad (\text{by Equation (3)})$$

$$= (g_k \circ \varepsilon_{j,k} \circ \pi_{j,k} \circ \varepsilon_{i,k})(x) \quad (\text{by Equation (6)})$$

But $\varepsilon_{j,k} \circ \pi_{j,k}$ is deflationary and g_k is monotone, so

$$\begin{aligned} & \sqsubseteq (g_k \circ \varepsilon_{i,k})(x) \\ & = g_i(x) \end{aligned} \quad (\text{by Equation (6)}),$$

as we wished to show.

Finally, we verify that g_∞ is Scott continuous. We first check that g_∞ is monotone. If $\sigma \sqsubseteq \tau$ in D_∞ , then $g_\infty(\sigma) \equiv \bigsqcup_{i:I} g_i(\sigma_i) \sqsubseteq \bigsqcup_{i:I} g_i(\tau_i) \equiv g_\infty(\tau)$, as each g_i is monotone. It remains to show that $g_\infty(\bigsqcup \alpha) \sqsubseteq \bigsqcup (g_\infty \circ \alpha)$ for every directed family $\alpha : A \rightarrow D_\infty$. By definition of g_∞ , it suffices to show that $g_i((\bigsqcup \alpha)_i) \sqsubseteq \bigsqcup (g_\infty \circ \alpha)$ for every $i : I$. By continuity of g_i it is enough to establish that $g_i((\alpha(a))_i) \sqsubseteq \bigsqcup (g_\infty \circ \alpha)$ for every $a : A$. But this holds as $g_i((\alpha(a))_i) \sqsubseteq g_\infty(\alpha(a)) \sqsubseteq \bigsqcup (g_\infty \circ \alpha)$, completing our proof. \square

Proposition 7.22. *The bilimit of locally small dcpos is locally small, i.e. if every \mathcal{V} -dcpo D_i is locally small for all $i : I$, then so is D_∞ .*

Proof. If every D_i is locally small, then for every $i : I$, we have a specified \mathcal{V} -valued partial order $\sqsubseteq_{\mathcal{V}}^i$ on D_i such that for every $i : I$ and every $x, y : D_i$, we have an equivalence $(x \sqsubseteq_{D_i} y) \simeq (x \sqsubseteq_{\mathcal{V}}^i y)$. Hence, $(\sigma \sqsubseteq_{D_\infty} \tau) \equiv (\prod_{i:I} (\sigma_i \sqsubseteq_{D_i} \tau_i)) \simeq (\prod_{i:I} (\sigma_i \sqsubseteq_{\mathcal{V}}^i \tau_i))$, but the latter is small, because $I : \mathcal{V}$ and $\sqsubseteq_{\mathcal{V}}^i$ is \mathcal{V} -valued. \square

8. SCOTT'S D_∞ MODEL OF THE UNTYPED λ -CALCULUS

We construct Scott's D_∞ [Sco72] predicatively. Formulated precisely, we construct a pointed $D_\infty : \mathcal{U}_0\text{-DCPO}_{\mathcal{U}_1, \mathcal{U}_1}$ such that D_∞ is isomorphic to its self-exponential $D_\infty^{D_\infty}$, employing the machinery from Section 7.

Definition 8.1 (D_n). We inductively define pointed dcpos $D_n : \mathcal{U}_0\text{-DCPO}_{\mathcal{U}_1, \mathcal{U}_1}$ for every natural number n by setting $D_0 \equiv \mathcal{L}_{\mathcal{U}_0}(\mathbf{1}_{\mathcal{U}_0})$ and $D_{n+1} \equiv D_n^{D_n}$.

In light of Remark 3.10 we highlight the fact that every D_n is a \mathcal{U}_0 -dcpo with carrier in \mathcal{U}_1 by the discussion of universe parameters of exponentials in Remark 6.6.

Definition 8.2 (ε_n, π_n). We inductively define for every natural number n , two Scott continuous maps $\varepsilon_n : D_n \rightarrow D_{n+1}$ and $\pi_n : D_{n+1} \rightarrow D_n$:

- (i)
 - $\varepsilon_0 : D_0 \rightarrow D_1$ is given by mapping $x : D_0$ to the continuous function that is constantly x ,
 - $\pi_0 : D_1 \rightarrow D_0$ is given by evaluating a continuous function $f : D_0 \rightarrow D_0$ at \perp which is itself continuous by Proposition 6.7,
- (ii)
 - $\varepsilon_{n+1} : D_{n+1} \rightarrow D_{n+2}$ takes a continuous function $f : D_n \rightarrow D_n$ to the continuous composite $D_{n+1} \xrightarrow{\pi_n} D_n \xrightarrow{f} D_n \xrightarrow{\varepsilon_n} D_{n+1}$, and
 - $\pi_{n+1} : D_{n+2} \rightarrow D_{n+1}$ takes a continuous function $f : D_{n+1} \rightarrow D_{n+1}$ to the continuous composite $D_n \xrightarrow{\varepsilon_n} D_{n+1} \xrightarrow{f} D_{n+1} \xrightarrow{\pi_n} D_n$.

Lemma 8.3. *The maps (ε_n, π_n) form an embedding-projection pair for every natural number n .*

Proof. We prove this by induction on n . For $n \equiv 0$ and arbitrary $x : D_0$, we have

$$\pi_0(\varepsilon_0(x)) \equiv \pi_0(\text{const}_x) \equiv \text{const}_x(\perp) \equiv x,$$

so ε_0 is indeed a section of π_0 . Moreover, for arbitrary $f : D_1$, we have

$$\varepsilon_0(\pi_0(f)) \equiv \varepsilon_0(f(\perp)) \equiv \text{const}_{f(\perp)},$$

so that for arbitrary $x : D_0$ we get $(\varepsilon_0(\pi_0(f)))(x) \equiv f(\perp) \sqsubseteq f(x)$ by monotonicity of f , proving that $\varepsilon_0 \circ \pi_0$ is deflationary.

Now suppose that the result holds for a natural number n ; we prove it for $n + 1$. For arbitrary $f : D_n \rightarrow D_n$, we calculate that

$$\pi_{n+1}(\varepsilon_{n+1}(f)) \equiv \pi_n \circ \varepsilon_{n+1}(f) \circ \varepsilon_n \equiv \pi_n \circ \varepsilon_n \circ f \circ \pi_n \circ \varepsilon_n = f,$$

as ε_n is a section of π_n by induction hypothesis. The proof that $\varepsilon_{n+1} \circ \pi_{n+1}$ is a deflation is similar. \square

In order to apply the tools from Section 7, we will need embedding-projection pairs $(\varepsilon_{n,m}, \pi_{n,m})$ from D_n to D_m whenever $n \leq m$.

Definition 8.4 $(\varepsilon_{n,m}, \pi_{n,m})$. We define Scott continuous maps $\varepsilon_{n,m} : D_n \rightarrow D_m$ and $\pi_{n,m} : D_m \rightarrow D_n$ for every two natural numbers $n \leq m$ as follows:

- (i) $\varepsilon_{n,m}$ and $\pi_{n,m}$ are both defined to be the identity if $n = m$;
- (ii) if $n < m$, then we define $\varepsilon_{n,m}$ as the composite

$$D_n \xrightarrow{\varepsilon_n} D_{n+1} \rightarrow \cdots \rightarrow D_{m-1} \xrightarrow{\varepsilon_{m-1}} D_m$$

and $\pi_{n,m}$ as the composite

$$D_m \xrightarrow{\pi_m} D_{m-1} \rightarrow \cdots \rightarrow D_{n+1} \xrightarrow{\pi_{n+1}} D_n$$

which yields embedding-projection pairs as they are compositions of them.

Instantiating the framework of Section 7 with the above diagram of objects $D_n : \mathcal{U}_0\text{-DCPO}_{\mathcal{U}_1, \mathcal{U}_1}$, we arrive at the construction of D_∞ and appropriate embedding-projection pairs. Observe that D_∞ is a \mathcal{U}_0 -dcpo with carrier and order taking values in \mathcal{U}_1 , just like each D_n , as was also mentioned in Remark 7.6.

Definition 8.5 (D_∞) . Applying Definitions 7.4, 7.7 and 7.12 to the above diagram yields $D_\infty : \mathcal{U}_0\text{-DCPO}_{\mathcal{U}_1, \mathcal{U}_1}$ with embedding-projection pairs $(\varepsilon_{n,\infty}, \pi_{n,\infty})$ from D_n to D_∞ for every natural number n .

Lemma 8.6. *The function $\pi_n : D_{n+1} \rightarrow D_n$ is strict for every natural number n . Hence, so is $\pi_{n,m}$ whenever $n \leq m$.*

Proof. Both statements are proved by induction. \square

Proposition 8.7. *The dcpo D_∞ is pointed.*

Proof. Since every D_n is pointed, we can consider the function $\sigma : \prod_{n:\mathbb{N}} D_n$ given by $\sigma(n) \equiv \perp_{D_n}$. Then σ is an element of D_∞ by Lemma 8.6 and it is the least, so D_∞ is indeed pointed. \square

We now work towards showing that D_∞ is isomorphic to the exponential $D_\infty^{D_\infty}$. Note that this exponential is again an element of $\mathcal{U}_0\text{-DCPO}_{\mathcal{U}_1, \mathcal{U}_1}$ by Remark 6.6, so the universe parameters do not increase.

Definition 8.8 (Φ_n). For every natural number n , we define the continuous maps

$$\begin{aligned} \Phi_{n+1} : D_{n+1} &\rightarrow D_\infty^{D_\infty} \\ f &\mapsto (D_\infty \xrightarrow{\pi_{n,\infty}} D_n \xrightarrow{f} D_n \xrightarrow{\varepsilon_{n,\infty}} D_\infty) \end{aligned}$$

and $\Phi_0 : D_0 \rightarrow D_\infty^{D_\infty}$ as $\Phi_1 \circ \varepsilon_0$.

Lemma 8.9. For every two natural numbers $n \leq m$, the diagram

$$\begin{array}{ccc} D_n & \xrightarrow{\Phi_n} & D_\infty^{D_\infty} \\ & \searrow \varepsilon_{n,m} & \nearrow \Phi_m \\ & & D_m \end{array}$$

commutes.

Proof. By induction on the difference of the two natural numbers, it suffices to prove that for every natural number n , the diagram

$$\begin{array}{ccc} D_n & \xrightarrow{\Phi_n} & D_\infty^{D_\infty} \\ & \searrow \varepsilon_n & \nearrow \Phi_{n+1} \\ & & D_{n+1} \end{array}$$

commutes. But this follows from Lemma 7.16 and unfolding the definition of Φ_n . \square

Definition 8.10 (Φ). The map $\Phi : D_\infty \rightarrow D_\infty^{D_\infty}$ is defined as the unique Scott continuous map induced by the Φ_n via Theorem 7.21.

Lemma 8.11. For $\sigma : D_\infty$ we have $\Phi(\sigma) = \bigsqcup_{n:\mathbb{N}} \Phi_{n+1}(\sigma_{n+1})$.

Proof. Recalling the proof of Theorem 7.21 we have $\Phi(\sigma) \equiv \bigsqcup_{n:\mathbb{N}} \Phi_n(\sigma_n)$, from which the claim follows easily. \square

We now define a map in the other direction using that D_∞ is also the limit.

Definition 8.12 (Ψ_n). For every natural number n , we define the continuous maps

$$\begin{aligned} \Psi_{n+1} : D_\infty^{D_\infty} &\rightarrow D_{n+1} \\ f &\mapsto (D_n \xrightarrow{\varepsilon_{n,\infty}} D_\infty \xrightarrow{f} D_\infty \xrightarrow{\pi_{n,\infty}} D_n) \end{aligned}$$

and $\Psi_0 : D_\infty^{D_\infty} \rightarrow D_0$ as $\pi_0 \circ \Psi_1$.

Lemma 8.13. For every two natural numbers $n \leq m$, the diagram

$$\begin{array}{ccc} D_\infty^{D_\infty} & \xrightarrow{\Psi_n} & D_n \\ & \searrow \Psi_m & \nearrow \pi_{n,m} \\ & & D_m \end{array}$$

commutes.

Proof. Similar to Lemma 8.9. \square

Definition 8.14 (Ψ). The map $\Psi : D_\infty^{D_\infty} \rightarrow D_\infty$ is defined as the unique Scott continuous map induced by the Ψ_n via Theorem 7.17.

Lemma 8.15. For $f : D_\infty^{D_\infty}$ we have $\Psi(f) = \bigsqcup_{n:\mathbb{N}} \varepsilon_{n+1,\infty}(\Psi_{n+1}(f))$.

Proof. Notice that

$$\begin{aligned}\Psi(f) &= \bigsqcup_{n:\mathbb{N}} \varepsilon_{n,\infty}(\pi_{n,\infty}(\Psi(f))) && \text{(by Lemma 7.19)} \\ &= \bigsqcup_{n:\mathbb{N}} \varepsilon_{n,\infty}(\Psi_n(f)) && \text{(by Equation (5)),}\end{aligned}$$

from which the claim follows easily. \square

Theorem 8.16. *The maps Φ and Ψ are inverses and hence, D_∞ is isomorphic to $D_\infty^{D_\infty}$.*

Proof. We first recall that

$$\varepsilon_{n,\infty} \text{ is a section of } \pi_{n,\infty} \text{ for every } n \in \mathbb{N}. \quad (*)$$

For arbitrary $\sigma : D_\infty$ we then calculate that

$$\begin{aligned}\Psi(\Phi(\sigma)) &= \Psi(\bigsqcup_{n:\mathbb{N}} \Phi_{n+1}(\sigma_{n+1})) && \text{(by Lemma 8.11)} \\ &= \bigsqcup_{n:\mathbb{N}} \Psi(\Phi_{n+1}(\sigma_{n+1})) && \text{(by continuity of } \Psi) \\ &= \bigsqcup_{n:\mathbb{N}} \bigsqcup_{m:\mathbb{N}} \varepsilon_{m+1,\infty}(\Psi_{m+1}(\Phi_{n+1}(\sigma_{n+1}))) && \text{(by Lemma 8.15)} \\ &= \bigsqcup_{n:\mathbb{N}} \varepsilon_{n+1,\infty}(\Psi_{n+1}(\Phi_{n+1}(\sigma_{n+1}))) \\ &\equiv \bigsqcup_{n:\mathbb{N}} \varepsilon_{n+1,\infty}(\pi_{n,\infty} \circ \varepsilon_{n,\infty} \circ \sigma_{n+1} \circ \pi_{n,\infty} \circ \varepsilon_{n,\infty}) && \text{(by definition)} \\ &= \bigsqcup_{n:\mathbb{N}} \varepsilon_{n+1,\infty}(\sigma_{n+1}) && \text{(by } (*)) \\ &= \sigma && \text{(by Lemma 7.19),}\end{aligned}$$

so Φ is indeed a section of Ψ . Moreover, for arbitrary $f : D_\infty^{D_\infty}$ we calculate that

$$\begin{aligned}\Phi(\Psi(f)) &= \Phi(\bigsqcup_{n:\mathbb{N}} \varepsilon_{n+1,\infty}(\Psi_{n+1}(f))) && \text{(by Lemma 8.15)} \\ &= \bigsqcup_{n:\mathbb{N}} \Phi(\varepsilon_{n+1,\infty}(\Psi_{n+1}(f))) && \text{(by continuity of } \Phi) \\ &= \bigsqcup_{n:\mathbb{N}} \bigsqcup_{m:\mathbb{N}} \Phi_{m+1}(\pi_{m+1,\infty}(\varepsilon_{n+1,\infty}(\Psi_{n+1}(f)))) && \text{(by Lemma 8.11)} \\ &= \bigsqcup_{n:\mathbb{N}} \Phi_{n+1}(\pi_{n+1,\infty}(\varepsilon_{n+1,\infty}(\Psi_{n+1}(f)))) \\ &= \bigsqcup_{n:\mathbb{N}} \Phi_{n+1}(\Psi_{n+1}(f)) && \text{(by } (*)) \\ &\equiv \bigsqcup_{n:\mathbb{N}} (\varepsilon_{n,\infty} \circ \pi_{n,\infty} \circ f \circ \varepsilon_{n,\infty} \circ \pi_{n,\infty}) && \text{(by definition)} \\ &= (\bigsqcup_{n:\mathbb{N}} \varepsilon_{n,\infty} \circ \pi_{n,\infty}) \circ f \circ (\bigsqcup_{m:\mathbb{N}} \varepsilon_{m,\infty} \circ \pi_{m,\infty}) \\ &= f && \text{(by Lemma 7.20),}\end{aligned}$$

finishing the proof. \square

Remark 8.17. Of course, Theorem 8.16 is only interesting when D_∞ is not the trivial one-element dcpo. Fortunately, D_∞ has (infinitely) many elements besides the least element \perp_{D_∞} . For instance, we can consider $x := \eta(\star) : D_0$ and observe that $\varepsilon_{0,\infty}(x)$ is an element of D_∞ not equal to \perp_{D_∞} , since $x \neq \perp_{D_0}$.

9. CONCLUDING REMARKS

We have shown how to develop basic domain theory in constructive and predicative univalent foundations with type universes playing a fundamental role in keeping track of relative sizes. We highlighted the use of the propositional truncation and illustrated our development with a predicative reconstruction of Scott's D_∞ .

A priori it is not clear that D_∞ should exist in predicative univalent foundations and it is one of the contributions of this work that this is indeed possible. Our construction largely follows the classical development of Scott's original paper [Sco72], but with some crucial differences. First of all, we carefully keep track of the universe parameters and try to be as general as possible. In the particular case of Scott's D_∞ model of the untyped λ -calculus, we obtain a \mathcal{U}_0 -dcpo whose carrier lives in the second universe \mathcal{U}_1 . The ability of the proof assistant AGDA to infer universe levels has been invaluable in this regard. Secondly, difference arises from proof relevance and these

complications are tackled with techniques in univalent foundations and [Kra+17, Theorem 5.4] in particular, as discussed right before Lemma 7.10, for example. Finally, we generalised Scott’s treatment from sequential bilimits to directed bilimits.

In the Part II paper [dJE24], we further show that D_∞ is an algebraic dcpo with a small compact basis. More generally, we take inspiration from category theory to present a predicative theory of continuous and algebraic domains.

10. ACKNOWLEDGEMENTS

I am grateful to Andrej Bauer and Dana Scott as they provided the spark for this paper by (independently) asking me whether we could have D_∞ in univalent foundations. I also thank Andrej and Vincent Rahli for valuable feedback on my thesis. Finally, I am very thankful to Martín Escardó with whom I had many fruitful discussions on the broader topic of developing domain theory constructively and predicatively in univalent foundations.

This work was supported by Cambridge Quantum Computing and Ilyas Khan [Dissertation Fellowship in Homotopy Type Theory]; and The Royal Society [grant reference URF\R1\191055].

REFERENCES

- [ADK17] Thorsten Altenkirch, Nils Anders Danielsson, and Nicolai Kraus. “Partiality, Revisited. The Partiality Monad as a Quotient Inductive-Inductive Type”. In: *Foundations of Software Science and Computation Structures. 20th International Conference, FOSSACS 2017, Held as Part of the European Joint Conferences on Theory and Practice of Software, ETAPS 2017, Uppsala, Sweden, April 22–29, 2017, Proceedings*. Ed. by Javier Esparza and Andrzej S. Murawski. Vol. 10203. Lecture Notes in Computer Science. Springer, 2017, pp. 534–549. DOI: [10.1007/978-3-662-54458-7_31](https://doi.org/10.1007/978-3-662-54458-7_31).
- [AJ94] Samson Abramsky and Achim Jung. “Domain theory”. In: *Handbook of Logic in Computer Science*. Ed. by S. Abramsky, Dov M. Gabray, and T. S. E. Maibaum. Vol. 3. Clarendon Press, 1994, pp. 1–168. Updated online version available at <https://www.cs.bham.ac.uk/~axj/pub/papers/handy1.pdf>.
- [Bis67] Erret Bishop. *Foundations of Constructive Analysis*. McGraw-Hill Book Company, 1967.
- [BK09] Andrej Bauer and Iztok Kavkler. “A constructive theory of continuous domains suitable for implementation”. In: *Annals of Pure and Applied Logic* 159.3 (2009): *Joint Workshop Domains VIII — Computability over Continuous Data Types, Novosibirsk, September 11–15, 2007*. Ed. by Yuri L. Ershov et al., pp. 251–267. DOI: [10.1016/j.apal.2008.09.025](https://doi.org/10.1016/j.apal.2008.09.025).
- [BKV09] Nick Benton, Andrew Kennedy, and Carsten Varming. “Some Domain Theory and Denotational Semantics in Coq”. In: *Theorem Proving in Higher Order Logics. 22nd International Conference, TPHOLs 2009, Munich, Germany, August 17–20, 2009, Proceedings*. Ed. by Stefan Berghofer et al. Vol. 5674. Lecture Notes in Computer Science. Springer, 2009, pp. 115–130. DOI: [10.1007/978-3-642-03359-9_10](https://doi.org/10.1007/978-3-642-03359-9_10).
- [Cap05] Venanzio Capretta. “General Recursion via Coinductive Types”. In: *Logical Methods in Computer Science* 1.2 (2005). DOI: [10.2168/LMCS-1\(2:1\)2005](https://doi.org/10.2168/LMCS-1(2:1)2005).

- [CMR17] Thierry Coquand, Bassel Manna, and Fabian Ruch. “Stack semantics of type theory”. In: *2017 32nd Annual ACM/IEEE Symposium on Logic in Computer Science (LICS)*. IEEE, 2017, pp. 1–12. DOI: [10.1109/LICS.2017.8005130](https://doi.org/10.1109/LICS.2017.8005130).
- [Coq+03] Thierry Coquand et al. “Inductively generated formal topologies”. In: *Annals of Pure and Applied Logic* 124.1–3 (2003), pp. 71–106. DOI: [10.1016/s0168-0072\(03\)00052-6](https://doi.org/10.1016/s0168-0072(03)00052-6).
- [Coq18] Thierry Coquand. “A survey of constructive presheaf models of univalence”. In: *ACM SISLOG News* 5.3 (2018), pp. 54–65. DOI: [10.1145/3242953.3242962](https://doi.org/10.1145/3242953.3242962).
- [CUV19] James Chapman, Tarmo Uustalu, and Niccolò Veltri. “Quotienting the delay monad by weak bisimilarity”. In: *Mathematical Structures in Computer Science* 29.1 (2019): *Best Papers Presented at ICTAC 2015*. Published online: 17th Oct. 2017, pp. 67–92. DOI: [10.1017/S0960129517000184](https://doi.org/10.1017/S0960129517000184).
- [dJE21] Tom de Jong and Martín Hötzel Escardó. “Domain Theory in Constructive and Predicative Univalent Foundations”. In: *29th EACSL Annual Conference on Computer Science Logic (CSL 2021)*. Ed. by Christel Baier and Jean Goubault-Larrecq. Vol. 183. Leibniz International Proceedings in Informatics (LIPIcs). Schloss Dagstuhl–Leibniz-Zentrum für Informatik, 2021, 28:1–28:18. DOI: [10.4230/LIPIcs.CSL.2021.28](https://doi.org/10.4230/LIPIcs.CSL.2021.28). Expanded version with full proofs available on arXiv: [2008.01422](https://arxiv.org/abs/2008.01422) [math.LO].
- [dJE23] Tom de Jong and Martín Hötzel Escardó. “On Small Types in Univalent Foundations”. In: *Logical Methods in Computer Science* 19.2 (2023), 8:1–8:33. DOI: [10.46298/lmcs-19\(2:8\)2023](https://doi.org/10.46298/lmcs-19(2:8)2023).
- [dJE24] Tom de Jong and Martín Hötzel Escardó. “Domain theory in univalent foundations II: Continuous and algebraic domains”. 2024. arXiv: [2407.06956](https://arxiv.org/abs/2407.06956) [cs.LO].
- [dJon21] Tom de Jong. “The Scott model of PCF in univalent type theory”. In: *Mathematical Structures in Computer Science* 31.10 (2021): *Homotopy Type Theory 2019*, pp. 1270–1300. DOI: [10.1017/S0960129521000153](https://doi.org/10.1017/S0960129521000153).
- [dJon22] Tom de Jong. “Domain Theory in Constructive and Predicative Univalent Foundations”. PhD thesis. University of Birmingham, 2022. arXiv: [2301.12405](https://arxiv.org/abs/2301.12405) [cs.LO]. URL: <https://etheses.bham.ac.uk/id/eprint/13401/>.
- [dJon24] Tom de Jong. *Index for “Domain Theory in Univalent Foundations I: Directed complete posets and Scott’s D_∞ ”*. <https://www.cs.bham.ac.uk/~mhe/TypeTopology/DomainTheory.Part-I.html>. 2024. <https://github.com/martinescardo/TypeTopology/tree/master/source/DomainTheory/Part-I.lagda>, commit [1d83144](https://github.com/martinescardo/TypeTopology/commit/1d83144).
- [Doc14] Robert Dockins. “Formalized, Effective Domain Theory in Coq”. In: *Interactive Theorem Proving. 5th International Conference, ITP 2014, Held as Part of the Vienna Summer of Logic, VSL 2014, Vienna, Austria, July 14–17, 2014, Proceedings*. Ed. by Gerwin Klein and Ruben Gamboa. Vol. 8558. Lecture Notes in Computer Science. Springer, 2014, pp. 209–225. DOI: [10.1007/978-3-319-08970-6_14](https://doi.org/10.1007/978-3-319-08970-6_14).
- [EK17] Martín H. Escardó and Cory M. Knapp. “Partial Elements and Recursion via Dominances in Univalent Type Theory”. In: *26th EACSL Annual Conference on Computer Science Logic (CSL 2017)*. Ed. by Valentin Goranko and Mads Dam. Vol. 82. Leibniz International Proceedings in Informatics (LIPIcs). Schloss Dagstuhl–Leibniz-Zentrum für Informatik, 2017, 21:1–21:16. DOI: [10.4230/LIPIcs.CSL.2017.21](https://doi.org/10.4230/LIPIcs.CSL.2017.21).

- [Esc+] Martín Hötzel Escardó et al. *TypeTopology*. <https://www.cs.bham.ac.uk/~mhe/TypeTopology/index.html>. AGDA development. <https://github.com/martinescardo/TypeTopology>, commit 1d83144.
- [Esc19] Martín Hötzel Escardó. “Introduction to Univalent Foundations of Mathematics with Agda”. Nov. 2019. arXiv: 1911.00580 [cs.LO].
- [Esc21] Martín Hötzel Escardó. “Injective types in univalent mathematics”. In: *Mathematical Structures in Computer Science* 31.1 (2021), pp. 89–111. DOI: 10.1017/s0960129520000225.
- [Gie+03] G. Gierz et al. *Continuous Lattices and Domains*. Vol. 93. Encyclopedia of Mathematics and its Applications. Cambridge University Press, 2003. DOI: 10.1017/CB09780511542725.
- [Har20] Brendan Hart. “Investigating Properties of PCF in Agda”. Final year MSci project. School of Computer Science, University of Birmingham, 2020. URL: <https://raw.githubusercontent.com/BrendanHart/Investigating-Properties-of-PCF/master/InvestigatingPropertiesOfPCFInAgda.pdf>. AGDA code available at <https://github.com/BrendanHart/Investigating-Properties-of-PCF>.
- [Har23] Brendan Hart. *Formalisation of the Scott model of PCF in Agda*. <https://www.cs.bham.ac.uk/~mhe/TypeTopology/Published.PCF.Lambda.index.html>. 2023. <https://github.com/martinescardo/TypeTopology/tree/master/source/PCF/Lambda/index.lagda>, commit 1d83144.
- [Hed96] Michael Hedberg. “A type-theoretic interpretation of constructive domain theory”. In: *Journal of Automated Reasoning* 16 (1996), pp. 369–425. DOI: 10.1007/BF00252182.
- [Hyl91] J. M. E. Hyland. “First steps in synthetic domain theory”. In: *Category Theory. Proceedings of the International Conference held in Como, Italy, July 22–28, 1990*. Ed. by Aurelio Carboni, Maria Cristina Pedicchio, and Giuseppe Rosolini. Vol. 1488. Lecture Notes in Mathematics. Springer, 1991, pp. 131–156. DOI: 10.1007/BFB0084217.
- [Joh77] P. T. Johnstone. *Topos Theory*. Academic Press, 1977. Reprinted by Dover Publications in 2014.
- [Kaw17] Tatsuji Kawai. “Geometric theories of patch and Lawson topologies”. 2017. arXiv: 1709.06403 [math.CT].
- [Kaw21] Tatsuji Kawai. “Predicative theories of continuous lattices”. In: *Logical Methods in Computer Science* 17.2 (2021). DOI: 10.23638/LMCS-17(2:22)2021.
- [Kna18] Cory Knapp. “Partial Functions and Recursion in Univalent Type Theory”. PhD thesis. School of Computer Science, University of Birmingham, June 2018. URL: <https://etheses.bham.ac.uk/id/eprint/8448/>.
- [Koc91] Anders Kock. “Algebras for the partial map classifier monad”. In: *Category Theory. Proceedings of the International Conference held in Como, Italy, July 22–28, 1990*. Ed. by Aurelio Carboni, Maria Cristina Pedicchio, and Giuseppe Rosolini. Vol. 1488. Lecture Notes in Mathematics. Springer, 1991, pp. 262–278. DOI: 10.1007/BFB0084225.
- [Kra+17] Nicolai Kraus et al. “Notions of Anonymous Existence in Martin-Löf Type Theory”. In: *Logical Methods in Computer Science* 13.1 (2017). DOI: 10.23638/LMCS-13(1:15)2017.
- [Lid20] David Lidell. “Formalizing domain models of the typed and the untyped lambda calculus in Agda”. Master’s thesis. Department of Computer Science and Engineering, Chalmers University of Technology and University of Gothenburg, 2020. URL: <http://hdl.handle.net/2077/67193>.

- [LN15] John Longley and Dag Normann. *Higher-Order Computability. Theory and Applications of Computability*. Springer, 2015. DOI: [10.1007/978-3-662-47992-6](https://doi.org/10.1007/978-3-662-47992-6).
- [Lon95] John Longley. “Realizability Toposes and Language Semantics”. PhD thesis. Department of Computer Science, University of Edinburgh, 1995. URL: <https://www.lfcs.inf.ed.ac.uk/reports/95/ECS-LFCS-95-332>.
- [Mag94] Lena Magnusson. “The Implementation of ALF. a Proof Editor Based on Martin-Löf’s Monomorphic Type Theory with Explicit Substitution”. PhD thesis. Department of Computing Science, Chalmers University of Technology and University of Gothenburg, 1994. URL: <http://hdl.handle.net/2077/12916>.
- [MV04] Maria Emilia Maietti and Silvio Valentini. “Exponentiation of Scott Formal Topologies”. In: *Electronic Notes in Theoretical Computer Science* 73 (2004): *Proceedings of the Workshop on Domains VI*. Ed. by Martín Escardó and Achim Jung, pp. 111–131. DOI: [10.1016/j.entcs.2004.08.005](https://doi.org/10.1016/j.entcs.2004.08.005).
- [Neg02] Sara Negri. “Continuous domains as formal spaces”. In: *Mathematical Structures in Computer Science* 12.1 (2002), pp. 19–52. DOI: [10.1017/S0960129501003450](https://doi.org/10.1017/S0960129501003450).
- [Pha91] Wesley Phao. “Domain Theory in Realizability Toposes”. PhD thesis. Department of Computer Science, University of Edinburgh, 1991. URL: <https://www.lfcs.inf.ed.ac.uk/reports/91/ECS-LFCS-91-171>.
- [Plo77] G.D. Plotkin. “LCF considered as a programming language”. In: *Theoretical Computer Science* 5.3 (1977), pp. 223–255. DOI: [10.1016/0304-3975\(77\)90044-5](https://doi.org/10.1016/0304-3975(77)90044-5).
- [PM21] Dirk Pattinson and Mina Mohammadian. “Constructive Domains with Classical Witnesses”. In: *Logical Methods in Computer Science* 17.1 (2021). DOI: [10.23638/LMCS-17\(1:19\)2021](https://doi.org/10.23638/LMCS-17(1:19)2021).
- [Reu99] Bernhard Reus. “Formalizing Synthetic Domain Theory”. In: *Journal of Automated Reasoning* 23 (1999), pp. 411–444. DOI: [10.1023/A:1006258506401](https://doi.org/10.1023/A:1006258506401).
- [Ros86] G. Rosolini. “Continuity and effectiveness in topoi”. PhD thesis. University of Oxford, 1986. URL: https://www.researchgate.net/publication/35103849_Continuity_and_effectiveness_in_topoi.
- [Ros87] G. Rosolini. “Categories and effective computations”. In: *Category Theory and Computer Science. Edinburgh, U.K., September 7–9, 1987. Proceedings*. Ed. by David H. Pitt, Axel Poigné, and David E. Rydeheard. Vol. 283. Lecture Notes in Computer Science. Springer, 1987. DOI: [10.1007/3-540-18508-9_17](https://doi.org/10.1007/3-540-18508-9_17).
- [RS99] Bernhard Reus and Thomas Streicher. “General synthetic domain theory — a logical approach”. In: *Mathematical Structures in Computer Science* 9.2 (1999), pp. 177–223. DOI: [10.1017/S096012959900273X](https://doi.org/10.1017/S096012959900273X).
- [Sam03] Giovanni Sambin. “Some points in formal topology”. In: *Theoretical Computer Science* 305.1–3 (2003): *Topology in Computer Science*, pp. 347–408. DOI: [10.1016/S0304-3975\(02\)00704-1](https://doi.org/10.1016/S0304-3975(02)00704-1).
- [Sam87] Giovanni Sambin. “Intuitionistic formal spaces — a first communication”. In: *Mathematical logic and its applications*. Ed. by Dimiter G. Skordev. Springer, 1987, pp. 187–204. DOI: [10.1007/978-1-4613-0897-3_12](https://doi.org/10.1007/978-1-4613-0897-3_12).
- [Sco72] Dana S. Scott. “Continuous lattices”. In: *Toposes, Algebraic Geometry and Logic. Dalhousie University, Halifax, January 16–19, 1971*. Ed. by

- F. W. Lawvere. Vol. 274. Lecture Notes in Mathematics. Springer, 1972, pp. 97–136. DOI: [10.1007/BFB0073967](https://doi.org/10.1007/BFB0073967).
- [Sco82a] Dana S. Scott. “Domains for denotational semantics”. In: *Automata, Languages and Programming. Ninth Colloquium. Aarhus, Denmark, July 1982*. Ed. by Mogens Nielsen and Erik Meineche Schmidt. Vol. 140. Lecture Notes in Computer Science. Springer-Verlag, 1982, pp. 577–610. DOI: [10.1007/BFB0012801](https://doi.org/10.1007/BFB0012801).
- [Sco82b] Dana S. Scott. “Lectures on a Mathematical Theory of Computation”. In: *Theoretical Foundations of Programming Methodology. Lecture Notes of an International Summer School, directed by F. L. Bauer, E. W. Dijkstra and C. A. R. Hoare*. Ed. by Manfred Broy and Gunther Schmidt. Vol. 91. NATO Advanced Study Institutes Series. Springer, 1982, pp. 145–292. DOI: [10.1007/978-94-009-7893-5_9](https://doi.org/10.1007/978-94-009-7893-5_9).
- [Sco93] Dana S. Scott. “A type-theoretical alternative to ISWIM, CUCH, OWHY”. In: *Theoretical Computer Science* 121.1–2 (1993), pp. 411–440. DOI: [10.1016/0304-3975\(93\)90095-B](https://doi.org/10.1016/0304-3975(93)90095-B).
- [Shu19] Michael Shulman. “All $(\infty, 1)$ -toposes have strict univalent universes”. Apr. 2019. arXiv: [1904.07004](https://arxiv.org/abs/1904.07004) [math.CT].
- [SVV96] Giovanni Sambin, Silvio Valentini, and Paolo Virgili. “Constructive domain theory as a branch of intuitionistic pointfree topology”. In: *Theoretical Computer Science* 159.2 (1996), pp. 319–341. DOI: [10.1016/0304-3975\(95\)00169-7](https://doi.org/10.1016/0304-3975(95)00169-7).
- [Swa19a] Andrew W Swan. “Choice, Collection and Covering in Cubical Sets”. <https://www.youtube.com/watch?v=r9KbE0zyr1g>. Talk at *Homotopy Type Theory Electronic Seminar Talks (HoTTEST)*, online. Nov. 6, 2019. URL: <https://www.math.uwo.ca/faculty/kapulkin/seminars/hotttestfiles/Swan-2019-11-06-HoTTEST.pdf>.
- [Swa19b] Andrew W Swan. “Counterexamples in Cubical Sets”. Talk at *Mathematical Logic and Constructivity: The Scope and Limits of Neutral Constructivism*, Stockholm, Sweden. Aug. 20, 2019. URL: <https://logic.math.su.se/mloc-2019/slides/Swan-mloc-2019-slides.pdf>.
- [Tay99] Paul Taylor. *Practical Foundations of Mathematics*. Vol. 59. Cambridge Studies in Advanced Mathematics. Cambridge University Press, 1999.
- [TvD88] A. S. Troelstra and D. van Dalen. *Constructivism in Mathematics. An Introduction*. Volume II. Ed. by J. Barwise et al. Vol. 123. Studies in Logic and the Foundations of Mathematics. Elsevier, 1988.
- [Uem19] Taichi Uemura. “Cubical Assemblies, a Univalent and Impredicative Universe and a Failure of Propositional Resizing”. In: *24th International Conference on Types for Proofs and Programs (TYPES 2018)*. Ed. by Peter Dybjer, José Espírito Santo, and Luís Pinto. Vol. 130. Leibniz International Proceedings in Informatics (LIPIcs). Schloss Dagstuhl–Leibniz-Zentrum für Informatik, 2019, 7:1–7:20. DOI: [10.4230/LIPIcs.TYPES.2018.7](https://doi.org/10.4230/LIPIcs.TYPES.2018.7).
- [Uni13] The Univalent Foundations Program. *Homotopy Type Theory: Univalent Foundations of Mathematics*. Institute for Advanced Study: <https://homotopytypetheory.org/book>, 2013.
- [Voe11] Vladimir Voevodsky. “Resizing Rules — their use and semantic justification”. Talk at *18th International Conference on Types for Proofs and Programs (TYPES)*, Bergen, Norway. Sept. 11, 2011. URL: https://www.math.ias.edu/vladimir/sites/math.ias.edu.vladimir/files/2011_Bergen.pdf.

- [Voe15] Vladimir Voevodsky. “An experimental library of formalized mathematics based on the univalent foundations”. In: *Mathematical Structures in Computer Science* 25.5 (2015): *From type theory and homotopy theory to Univalent Foundations of Mathematics*, pp. 1278–1294. DOI: [10.1017/S0960129514000577](https://doi.org/10.1017/S0960129514000577).
- [vOos08] Jaap van Oosten. *Realizability: An Introduction to its Categorical Side*. Ed. by S. Abramsky et al. Vol. 152. Studies in Logic and the Foundations of Mathematics. Elsevier, 2008. DOI: [10.1016/s0049-237X\(08\)X8001-2](https://doi.org/10.1016/s0049-237X(08)X8001-2).

SCHOOL OF COMPUTER SCIENCE, UNIVERSITY OF BIRMINGHAM, BIRMINGHAM, UK
Email address: tom.dejong@nottingham.ac.uk
URL: <https://www.tdejong.com>