# Scaling limit of the KPZ equation with non-integrable spatial correlations

July 19, 2024

# Luca Gerolla<sup>1</sup>, Martin Hairer<sup>1,2</sup>, Xue-Mei Li<sup>1,2</sup>

<sup>1</sup> Imperial College London, SW7 2AZ London, UK.

<sup>2</sup> EPFL, 1015 Lausanne, Switzerland.

Email: luca.gerolla16@imperial.ac.uk, martin.hairer@epfl.ch, xue-mei.li@epfl.ch

## Abstract

We study the large scale fluctuations of the KPZ equation in dimensions  $d \ge 3$  driven by Gaussian noise that is white in time Gaussian but features nonintegrable spatial correlation with decay rate  $\kappa \in (2, d)$  and a suitable limiting profile. We show that its scaling limit is described by the corresponding additive stochastic heat equation. In contrast to the case of compactly supported covariance, the noise in the stochastic heat equation retains spatial correlation with covariance  $|x|^{-\kappa}$ . Surprisingly, the noise driving the limiting equation turns out to be the scaling limit of the noise driving the KPZ equation so that, under a suitable coupling, one has convergence in probability, unlike in the case of integrable correlations where fluctuations are enhanced in the limit and convergence is necessarily weak.

Mathematics Subject Classification: 60H15, 60F17

*Keywords:* KPZ equation, stochastic heat equation, fluctuations, long range correlations

#### Contents

1	Introduction Analysis of the stationary solution			
2				
	2.1	Notations and conventions	6	
	2.2	Forward and backward stationary solutions	7	
		A priori bounds		
3	The	main homogenisation theorem	16	
4	Proof of the main results			
	4.1	Main convergence result	26	
	4.2	Convergence in Hölder spaces	32	
	4.3	Proof of the main results	35	

#### **1** Introduction

In this article, we investigate the large-scale dynamics of the Kardar–Parisi–Zhang (KPZ) equation on  $\mathbb{R}^d$  in dimension  $d \ge 3$ . The KPZ equation

$$\partial_t h = \frac{1}{2}\Delta h + \frac{1}{2}|\nabla h|^2 + \beta\xi , \qquad (1.1)$$

was originally introduced in [KPZ86] to model random interface growth. Here,  $\beta$  is a coupling constant, and  $\xi$  is a mean zero Gaussian noise. Over time, it has evolved into one of the most extensively studied stochastic PDEs due to its ability to capture the universal behaviour of many probabilistic models on one hand and the mathematical challenges it presents for a well-posed solution theory on the other. Very recently, significant achievements have been made, demonstrating that, in dimension 1, the rescaled solution  $\varepsilon^{\frac{1}{2}}h(\varepsilon^{-\frac{3}{2}}t,\varepsilon^{-1}x) - C_{\varepsilon}t$  converges to the KPZ fixed point [QS23, Vir20]. This accomplishment built upon the remarkable progress made in the last couple of decades [BG97, BDJ99, ACQ11, BQS11, AKQ14, MQR21]. While the KPZ equation offers insights into universal phenomena, it poses substantial mathematical challenges when it comes to defining a robust and mathematically rigorous solution that exhibits the aforementioned physical relevance [Hai13, Hai14, GIP15, GP17, Kup16].

In the original model, the mean zero Gaussian noise  $\xi$  has short range correlations. In that case, it was shown recently in a number of works [MU18, GRZ18, CCM19, DGRZ20, CNN22, LZ22] that the large-scale behaviour of the KPZ equation in dimensions  $d \ge 3$  in the weak coupling regime is described by the Edwards– Wilkinson model, namely simply the additive stochastic heat equation driven by space-time white noise.

In the present article, we consider instead spatially coloured noise with long range correlations, and we are interested to explore how these correlations affect the KPZ scaling limit in the supercritical regime  $d \ge 3$ . In such settings, where the correlation is not integrable, so that the form of the effective variance in the above references is infinite, an effective fluctuation theory for the stochastic heat equation (SHE) has been developed [GHL23]. Like there, we consider large scales correlations of polynomial decay  $|x|^{-\kappa}$ . Our result on the KPZ scaling limit introduced below aligns with the physics literature [KS99, KCDW14] which predicts a Gaussian limit when  $\kappa > 2$ .

We begin by describing the noise  $\xi$  considered:

Assumption 1.1. The Gaussian noise  $\xi$  is white in time with smooth spatial covariance R, formally  $\mathbb{E}[\xi(t, x)\xi(s, y)] = \delta(t-s)R(x-y)$ . There exists  $\kappa \in (2, d)$  such that  $R(x) \leq (1+|x|)^{-\kappa}$  and, for any  $x \in \mathbb{R}^d \setminus \{0\}$ ,  $\lim_{\varepsilon \to 0} \varepsilon^{-\kappa} R(\varepsilon^{-1}x) = |x|^{-\kappa}$ .

For example, the noise obtained by convolving (in space) a space-time white noise with a smooth function  $\phi : \mathbb{R}^d \to \mathbb{R}_+$  behaving at infinity like  $\phi(x) \sim |x|^{-\frac{d+\kappa}{2}}$  leads to  $R(x) = \int_{\mathbb{R}^d} \phi(x+y)\phi(y)dy$ , which satisfies Assumption 1.1.

Writing h for the solution to (1.1), but with a slowly varying initial condition of the form  $\varepsilon^{\frac{\kappa}{2}-1}h_0(\varepsilon x)$ , we rescale it by setting, for  $\varepsilon \in (0, 1]$ ,

$$h_{\varepsilon}(t,x) = \varepsilon^{1-\frac{\kappa}{2}} h(t/\varepsilon^2, x/\varepsilon) - \frac{t}{2}\beta^2 R(0)\varepsilon^{-1-\frac{\kappa}{2}} .$$

A simple calculation shows that  $h_{\varepsilon}$  then solves the rescaled KPZ equation

$$\partial_t h_{\varepsilon} = \frac{1}{2} \Delta h_{\varepsilon} + \frac{1}{2} \varepsilon^{\frac{\kappa}{2} - 1} |\nabla h_{\varepsilon}|^2 - \frac{1}{2} \beta^2 R(0) \varepsilon^{-1 - \frac{\kappa}{2}} + \beta \xi^{\varepsilon} , \qquad (1.2)$$

with initial condition  $h_0$ . Here  $\xi^{\varepsilon}$  is the rescaling of  $\xi$  such that

$$\mathbb{E}[\xi^{\varepsilon}(t,x)\xi^{\varepsilon}(s,y)] = \delta(t-s)\varepsilon^{-\kappa}R(\frac{x-y}{\varepsilon}).$$
(1.3)

Under this scaling,  $\xi^{\varepsilon}$  converges in law, as  $\varepsilon \to 0$ , to a non-trivial Gaussian limit  $\xi^0$  with covariance

$$\mathbb{E}[\xi^0(t,x)\xi^0(s,y)] = \delta(t-s)|x-y|^{-\kappa} .$$
(1.4)

It will be convenient to upgrade this convergence to convergence in probability (which can always be achieved by Skorokhod's representation theorem):

**Definition 1.2.** A coupling between the  $\xi^{\varepsilon}$  is *good* if, for every  $\psi \in C_c^{\infty}(\mathbb{R}^{d+1})$ , there exists a random variable  $\xi^0(\psi)$  such that  $\xi^{\varepsilon}(\psi) \Rightarrow \xi^0(\psi)$  in probability.

If one can write  $\xi(t, x) = \int_{\mathbb{R}^d} \phi(x - y)\eta(t, y) \, dy$ , for  $\eta$  a space-time white noise and for  $\phi$  as described just above, then a natural good coupling is given by

$$\xi^{\varepsilon}(t,x) = \varepsilon^{-(d+\kappa)/2} \int_{\mathbb{R}^d} \phi((x-y)/\varepsilon) \eta(t,y) \, dy \;. \tag{1.5}$$

Given the limiting noise  $\xi^0$ , we write  $\mathcal{U}$  for the solution to the corresponding additive stochastic heat equation:

$$\partial_t \mathcal{U} = \frac{1}{2} \Delta \mathcal{U} + \beta \xi^0 , \qquad \mathcal{U}(0, \cdot) = h_0 .$$
 (1.6)

Our main result is then the following.

**Theorem 1.3.** Let  $d \ge 3$  and let  $\xi$  satisfy Assumption 1.1. For any  $p \ge 1$ ,  $\alpha < 1 - \frac{\kappa}{2}$ , and  $\sigma < -1 - \frac{\kappa}{2}$ , there exists a constant c < 0 and a positive value  $\beta_0 = \beta_0(p, \alpha)$  such that for any  $\beta < \beta_0$ , for any deterministic initial condition  $h_0 \in C_b(\mathbb{R}^d)$ , and any good coupling of the noise, one has, for any  $T > T_0 > 0$ 

$$\lim_{\varepsilon \to 0} (h_{\varepsilon}(t, x) - \varepsilon^{1 - \frac{\kappa}{2}} c) = \mathcal{U} ,$$

in probability in  $L^p([T_0, T], \mathcal{C}^{\alpha}(E))$ , for any compact  $E \subset \mathbb{R}^d$ .

We provide the definition and comment on the constant c in Remark 1.6 below, whilst the proof of this result will be given in Section 4.3, where we also provide the definition of the spaces  $C^{\alpha}(E)$  and their semi-norms. In fact, we will show a limit theorem for general functions  $\Phi(u)$  of solutions to the multiplicative stochastic heat equation

$$\partial_t u(t,x) = \frac{1}{2} \Delta u(t,x) + \beta u(t,x) \xi(t,x) , \qquad u(0,\cdot) = 1 , \qquad (1.7)$$

from which Theorem 1.3 essentially follows as a special case with  $\Phi = \log$ . (We still need a little bit more work since we allow for general initial conditions in Theorem 1.3, while our general fluctuation result, Theorem 1.4, holds for  $u_0 \equiv 1$ . It is this additional work that is performed at the end of Section 4.3.) Under the hypotheses stated earlier, (1.7) admits a unique solution [PZ00, Thm 0.2]. Since we are interested in its large-scale fluctuations, we write  $u_{\varepsilon}(t, x) = u(t/\varepsilon^2, x/\varepsilon)$  for the diffusively rescaled solution, which solves

$$\partial_t u_{\varepsilon}(t,x) = \frac{1}{2} \Delta u_{\varepsilon}(t,x) + \beta \varepsilon^{\frac{\kappa}{2}-1} u_{\varepsilon}(t,x) \xi^{\varepsilon}(t,x) , \qquad u_{\varepsilon}(0,\cdot) = 1, \qquad (1.8)$$

with  $\xi^{\varepsilon}$  as before. We then have the following functional central limit theorem, which will be restated as Theorem 4.2 below and proven in Section 4.3.

**Theorem 1.4.** Consider a map  $\Phi : \mathbb{R}_+ \to \mathbb{R}$  satisfying Assumption 4.1 and let

$$u_{\varepsilon}^{\Phi}(t,x) = \varepsilon^{1-\frac{\kappa}{2}} \left( \Phi(u_{\varepsilon}(t,x)) - \mathbb{E}[\Phi(u_{\varepsilon}(t,x))] \right)$$

With assumptions on d, R, p,  $\alpha$  and  $\sigma$  as in Theorem 1.3, there exists  $\beta_0^{\Phi} = \beta_0^{\Phi}(\alpha, p) > 0$  and  $\nu_{\Phi} \ge 0$  such that, for all  $\beta < \beta_0^{\Phi}$ , for any T > 0, and any good coupling,  $u_{\varepsilon}^{\Phi}$  converges in probability to  $\nu_{\Phi}\mathcal{U}^0$  in the space  $L^p([0, T], \mathcal{C}^{\alpha}(E))$  for any compact  $E \subset \mathbb{R}^d$ . Here,  $\mathcal{U}^0$  denotes the solution to the additive stochastic heat equation

$$\partial_t \mathcal{U}^0 = \frac{1}{2} \Delta \mathcal{U}^0 + \beta \xi^0 , \qquad \mathcal{U}^0(0, \cdot) = 0 .$$
(1.9)

Our class of possible transformations includes the Cole–Hopf transform  $\Phi(u) = \log u$ , for which one has  $\nu_{\Phi} = 1$ . In this case we recover the fluctuations of KPZ since  $h_{\varepsilon}(t,x) = \varepsilon^{1-\frac{\kappa}{2}} \log u_{\varepsilon}(t,x)$  solves (1.2) with flat initial conditions  $h_0 \equiv 0$ . We discuss and provide the explicit expression of  $\nu_{\Phi}$  in the general case in Remark 1.7 below.

The main tool (and this is also one of the main novelties of this article) to prove the above is a homogenisation result for the fundamental solution of (1.7). In order to formulate it, write  $w_{s,y}(t,x)$  for the solution to (1.7) at times  $t \ge s$  with initial condition at time s given by  $w_{s,y}(s, \cdot) = \delta_y$ :

**Theorem 1.5.** Let  $d \ge 3$  and  $\xi$  satisfies Assumption 1.1. Then for any  $p \ge 1$ , for sufficiently small  $\beta$ , there exist stationary random fields  $\vec{Z}$  and  $\vec{Z}$  such that, with  $\mu = \frac{1}{2} - \frac{1}{\kappa}$ ,

$$\left\|\frac{w_{s,y}(t,x)}{P_{t-s}(y-x)} - \vec{Z}(t,x)\vec{Z}(s,y)\right\|_p \lesssim (1+t-s)^{-\mu}(1+(1+t-s)^{-\frac{1}{2}}|x-y|) .$$
(1.10)

A slightly more detailed version of this statement will be formulated as Proposition 3.1 below. The field  $\vec{Z}$  is the analogue of the pullback field already appearing in [GHL23, DGRZ21, CCM19, CCM20] where it was denoted by Z. It is obtained as the limit  $\vec{Z}(t, x) = \lim_{s \to -\infty} u^{(s)}(t, x)$ , where  $u^{(s)}$  denotes the solution to (1.7) with initial condition 1 at time s. It turns out that the field  $\vec{Z}$  is given by " $\vec{Z}$  running backwards in time" (which explains our notations) in a sense that will be made precise below, see Theorem 2.1 and equation (2.8). A shadow of the field  $\vec{Z}$  can be seen in the constant  $\bar{c}$  appearing in [DGRZ21, Eq. 1.6], whose inverse would correspond to the expectation of  $\vec{Z}$  (which in our case equals 1 as a consequence of the fact that we consider noise that is white in time). More explicitly, these fields satisfy the equations

$$\vec{Z}(t,x) = 1 + \beta \int_{-\infty}^{t} \int_{\mathbb{R}^d} P_{t-r}(x-z)\vec{Z}(r,z)\xi(dr,dz) , \qquad (1.11)$$

$$\bar{Z}(s,y) = 1 + \beta \int_s^\infty \int_{\mathbb{R}^d} P_{r-s}(y-z)\bar{Z}(r,z)\,\xi(dr,dz) , \qquad (1.12)$$

with the stochastic integral appearing in the second expression being of backwards Itô type.

Using the Clark–Ocone formula we recover a divergence representation for the fluctuations of  $\Phi(u)$ , which can be expressed in terms of the fundamental solution of (1.7). Relying on the large time homogenisation of  $w_{s,y}$  from Theorem 1.5 and mixing of the stationary fields, we are able to approximate the fluctuations of  $\Phi(u)$ by the fluctuations of the linear SHE (Proposition 4.5) and then prove Theorem 1.4. We shall explain more in details the argument and carry out the proof in Section 4. *Remark* 1.6. Unlike what happens in the case of short-range correlations, the constant c in Theorem 1.3 is all that is left from the KPZ nonlinearity since neither the covariance of the noise nor the coefficient in front of the Laplacian end up being affected by the scaling limit. The constant is given by  $c = \mathbb{E}[\log Z]$  with  $Z \stackrel{\text{law}}{=} \vec{Z}(0,0)$ . Since  $\mathbb{E}Z = 1$  and Z has non-zero variance by (1.11), c is strictly negative. The way one can intuitively understand the presence of c is the following: since we start with a fixed  $\varepsilon$ -independent bounded continuous initial condition  $h_0$  which looks quite different from the stationary solutions (which exhibit oscillations of size about  $\varepsilon^{1-\frac{\kappa}{2}}$ ), the KPZ nonlinearity and the renormalisation constant appearing in (1.2) don't quite balance out initially. By the time the process has become locally sufficiently close to stationarity, this imbalance has already caused it to shoot down by a substantial amount.

This strongly suggests that one can find a collection  $Y_{\varepsilon}$  of stationary processes such that  $\lim_{\varepsilon \to 0} Y_{\varepsilon} = 0$  in distribution and such that a statement analogous to Theorem 1.3, but without the constant c, holds provided that one starts  $h_{\varepsilon}$  with initial condition  $Y_{\varepsilon} + h_0$ . In fact, we believe that one should be able to take

$$Y_{\varepsilon}(x) \stackrel{\text{\tiny haw}}{=} \varepsilon^{1-\frac{\alpha}{2}} (\log(u_{\varepsilon}(t_{\varepsilon}, x)) - c) ,$$

(but independent of the noise  $\xi$ ) for a suitable sequence of times  $t_{\varepsilon} \to 0$ , and show that  $\lim_{\varepsilon \to 0} h_{\varepsilon} = \mathcal{U}$ .

Remark 1.7. With the notations above at hand, one has  $\nu_{\Phi} = \mathbb{E}[Z \Phi'(Z)]$  in Theorem 1.4 where  $Z = \vec{Z}(0,0)$ . We will see in Lemma 2.6 that our assumptions on  $\Phi$  guarantee that this quantity is finite. Given the dependence of the pullback stationary solution  $\vec{Z}$  on  $\beta$ , we note that  $\nu_{\Phi}$  actually depends on  $\beta$  in general and that, as  $\beta \to 0$ ,  $\nu_{\Phi}$  converges to  $\Phi'(1)$ . In the special case  $\Phi(x) = x$ , one has  $\nu_{\Phi} = 1$  and we recover the result of [GHL23], at least in the case  $\sigma(u) = u$ .

Note that the effective variance differs from that obtained for the case of integrable R in [GRZ18, MU18, DGRZ20, CNN22, LZ22, GL20, DGRZ21, CCM19]. There, the effective variance  $\nu_{\text{eff}}$  is given by the expression

$$\nu_{\text{eff}}^2 = \nu_{\Phi}^2 \int_{\mathbb{R}^d} R(x) \mathbb{E}[\vec{Z}(0,x)\vec{Z}(0,0)] dx$$

In their works, the fluctuation scale depends on the dimension, while in our setting the fluctuations scale  $\varepsilon^{\kappa-2}$  depends on the decay coefficient  $\kappa$  of the noise covariance.

#### Structure of the article

The remainder of the article is structured as follows. In Section 2, we introduce the random fields  $\vec{Z}$  and  $\vec{Z}$  and we collect some useful a priori bounds on these fields as well as on the solutions to the multiplicative stochastic heat equation (1.7) and their Malliavin derivatives. Section 3 forms the bulk of the article and contains the proof of the homogenisation result. Finally, in Section 4, we use all of these ingredients to provide a proof of Theorems 1.3 and 1.4.

#### Acknowledgements

The authors are grateful to Alex Dunlap and Yu Gu for discussions on this topic and in particular for pointing out that our proof may yield convergence in probability in this setting. This research was partially supported by the Royal Society through MH's research professorship. XML acknowledges support from EPSRC grant EP/V026100/1 and EP/S023925/1. LG is supported by the EPSRC Centre for Doctoral Training in Mathematics of Random Systems: Analysis, Modelling and Simulation (EP/S023925/1).

#### **2** Analysis of the stationary solution

#### 2.1 Notations and conventions

The background probability space is  $(\Omega, \mathcal{F}, \mathbb{P})$ , and we write  $\mathcal{F}_t$  for the filtration generated by  $\xi$ , i.e.  $\mathcal{F}_t$  is the smallest  $\mathbb{P}$ -complete  $\sigma$ -algebra such that the random variables  $\{\xi(\phi) : \operatorname{supp} \phi \subset (-\infty, t] \times \mathbb{R}^d, \phi \in \mathcal{C}_c^{\infty}(\mathbb{R}_+ \times \mathbb{R}^d)\}$  are all  $\mathcal{F}_t$ -measurable. We generally interpret an integrable function  $\xi$  as a distribution by the following expression:

$$\xi(\phi) = \int_{\mathbb{R} \times \mathbb{R}^d} \xi(s, y) \phi(s, y) ds dy.$$

The notation  $\|\cdot\|_p$  stands for the  $L^p(\Omega)$  norm. Moreover,

- $\triangleright$  With  $a \leq b$  we mean  $a \leq Cb$  for some constant C > 0, independent of  $\varepsilon$ .
- $\triangleright \ \mathcal{C}^{\infty}_{c}(\mathbb{R}^{d})$  denotes the set of smooth functions on  $\mathbb{R}^{d}$  with compact support.
- $\triangleright$  For  $E \subset \mathbb{R}^d$  open and  $k \in \mathbb{N}$ , we denote with  $\mathcal{C}^k(E)$  the space of k times continuously differentiable functions on E and  $\mathcal{C}^k_b(E)$  the subset of bounded  $\mathcal{C}^k(E)$  functions with all partial derivatives up to order k bounded.

#### 2.2 Forward and backward stationary solutions

For the purpose of this section, it will be convenient to assume that the underlying probability space  $\Omega$  is the space of space-time distributions and that the noise  $\xi$  is simply the canonical process.

With  $u^{(s)}(t, x)$  denoting the solution to (1.7) with initial condition  $u^{(s)}(s, \cdot) \equiv 1$  at time s, we recall [GHL23, Thm 2.10]. To avoid confusion, we remark that in [GHL23],  $u^{(K)}(t, x)$  denotes the solution that starts from time -K. A result of similar nature can be found in [GL20, Thm 1.1].

**Theorem 2.1.** Let  $d \ge 3$  and  $\xi$  satisfies Assumption 1.1. Given any  $p \ge 1$ , there exists a positive constant  $\beta_1(p) = \beta_1(p, d, R)$  and a space-time stationary random field  $\vec{Z}$  such that, for  $\beta \le \beta_1(p)$ , one has

$$\|u^{(s)}(t,x) - \vec{Z}(t,x)\|_{L_p(\Omega)} \lesssim 1 \wedge (t-s)^{\frac{2-\kappa}{4}},$$
 (2.1)

uniformly over all  $s \leq t$  and  $x \in \mathbb{R}^d$ . Recall that  $\kappa \in (2, d)$ . Furthermore, for any compact region  $\mathfrak{K} \subset \mathbb{R}^d$  and every  $t \in \mathbb{R}$ , one has

$$\lim_{s \to -\infty} \mathbb{E} \sup_{x \in \mathfrak{K}} |u^{(s)}(t,x) - \vec{Z}(t,x)|^p \to 0.$$

It follows from (2.1) that  $\vec{Z}$  is a random fixed point for the random dynamical system generated by the stochastic heat equation (1.7). In fact, we have even more structure as we will describe now. Our probability space admits a natural "time reversal" involution  $\mathcal{R}: \Omega \to \Omega$  and "space-time translations"  $\mathcal{T}_{t,x}: \Omega \to \Omega$  such that, formally

$$(\mathcal{R}\xi)(t,x) = \xi(-t,x), \qquad (\mathcal{T}_{t,x}\xi)(s,y) = \xi(s+t,x+y),$$

with the obvious rigorous interpretations in terms of test functions. These operations naturally yield linear operators acting on the space of random variables  $A: \Omega \to \mathbb{R}$  by

$$(\mathcal{R}A)(\xi) \stackrel{\text{def}}{=} A(\mathcal{R}\xi) , \qquad (\mathcal{T}_{t,x}A)(\xi) \stackrel{\text{def}}{=} A(\mathcal{T}_{t,x}\xi) .$$
(2.2)

Since the law of  $\xi$  is invariant under  $\mathcal{R}$  and  $\mathcal{T}_{t,x}$ , the operations in (2.2) are isometries for every  $L^p(\Omega, \mathbb{P})$ -space.

We will repeatedly use the fact that, by uniqueness of solutions to (1.7), one has the following.

**Lemma 2.2.** For any  $t \in \mathbb{R}$ ,  $x \in \mathbb{R}^d$ , one has  $\mathcal{T}_{t,x}\mathcal{R} = \mathcal{R}\mathcal{T}_{-t,x}$  and the following holds almost surely:

$$u^{(s)}(t,x)(\mathcal{T}_{0,y}\xi) = u^{(s)}(t,x+y)(\xi) ,$$
  
$$u^{(s)}(t,x)(\mathcal{T}_{r,0}\xi) = u^{(s+r)}(t+r,x)(\xi) .$$

Given any random variable A as above, we then define the stationary random fields

$$\overline{A}(t,x) = \mathcal{T}_{t,x}A$$
,  $\overline{A}(t,x) = \mathcal{T}_{t,x}\mathcal{R}A$ . (2.3)

If we now define

$$Z = \lim_{s \to -\infty} Z_s , \qquad Z_s = u^{(s)}(0,0) , \qquad (2.4)$$

then we see that this definition is consistent with our notation since by Lemma 2.2  $\vec{Z}(t, x) := \mathcal{T}_{t,x}Z$ , defined as in (2.3), satisfies

$$\vec{Z}(t,x) = \lim_{s \to -\infty} u^{(s)}(0,0)(\mathcal{T}_{t,x}\xi) = \lim_{s \to -\infty} u^{(t+s)}(t,x)(\xi)$$
(2.5)  
$$= \lim_{s \to -\infty} u^{(s)}(t,x) ,$$

which is indeed the same quantity appearing in (2.1), as well as in (1.10), by virtue of (1.11). With the notation above

$$\bar{Z}(s,y) = \mathcal{T}_{s,y}\mathcal{R}Z = \mathcal{R}\mathcal{T}_{-s,y}Z = \mathcal{R}\,\bar{Z}(-s,y)\,,\qquad(2.6)$$

so that

$$\overline{Z}(s,y)(\xi) = \lim_{r \to -\infty} u^{(r)}(-s,y)(\mathcal{R}\xi) ,$$

which is indeed consistent with (1.12) via the change of variables  $r \mapsto -r$  (and noting that such a change of variables does indeed turn Itô integrals into backwards Itô integrals by simple Riemann sum approximations).

Let us now explain why  $\overline{Z}$  appears in (1.10). If it were the case that

$$w_{s,y}(t,x) = P_{t-s}(y-x)\vec{Z}(t,x)\vec{Z}(s,y) ,$$

then, as a consequence of the fact that  $\mathbb{E}\vec{Z} = 1$  and the law of large number we could recover  $\bar{Z}$  by

$$\bar{Z}(s,y) = \lim_{t \to \infty} \int_{\mathbb{R}^d} w_{s,y}(t,x) \, dx$$

It is therefore natural to consider the stochastic process

$$C_{s,y}(t) = \int_{\mathbb{R}^d} w_{s,y}(t,x) \, dx \; . \tag{2.7}$$

As we will see shortly, it is then indeed the case that one has the identity  $\overline{Z}(s, y) = \lim_{t\to\infty} C_{s,y}(t)$ . In fact we will show more, namely that with  $Z_s$  as in (2.4), one has the exact identity  $C_{s,y}(t) = \overline{Z}_{s-t}(s, y)$ .

**Proposition 2.3.** Under the conditions of Theorem 2.1, one has  $C_{s,y}(t) = \overline{Z}_{s-t}(s, y)$ , and

$$||C_{s,y}(t) - \overleftarrow{Z}(s,y)||_p \lesssim 1 \wedge (t-s)^{\frac{2-\kappa}{4}}.$$

*Proof.* By the definition (2.3),  $\overline{Z}_{s-t}(s, y) := \mathcal{T}_{s,y}\mathcal{R}Z_{s-t} = \mathcal{R}\mathcal{T}_{-s,y}u^{(s-t)}(0, 0)$ . By Lemma 2.2,

$$\bar{Z}_{s-t}(s,y) = \mathcal{R}u^{(-t)}(-s,y), \qquad \lim_{t \to \infty} \bar{Z}_{s-t}(s,y) = \bar{Z}(s,y).$$
(2.8)

We shall use the method of duality. The basic observation is that if u denotes the SHE running forward in time with initial condition  $u_s$  at some time s and vdenotes the one running backwards in time with terminal condition  $v_t$  at some time t > s then  $\langle u_r, v_r \rangle$ , where the bracket denotes the inner product on  $L^2(\mathbb{R}^d)$ , is independent of  $r \in [s, t]$ . Formally, this is because

$$\partial_r \langle u_r, v_r \rangle = \langle \Delta u_r + \xi u_r, v_r \rangle - \langle u_r, \Delta v_r + \xi v_r \rangle = 0.$$
 (2.9)

One can make this precise in the following way. Fix a mollifier  $\rho \in C_c^{\infty}(\mathbb{R} \times \mathbb{R}^d)$ such that  $\int \rho = 1$  and  $\rho(t, x) = \rho(-t, x) = \rho(t, -x)$ . For any  $\delta > 0$ , we write  $\rho_{\delta}(t, x) = \delta^{-2-d}\rho(t/\delta^2, x/\delta)$  and we set  $\xi_{\delta}(t, x) = (\mathcal{T}_{t,x}\xi)(\rho_{\delta})$ .

Fix then some initial time s and constant  $K_{\delta}$  and consider the solution  $u^{\delta}(r, x)$  to the random PDE

$$\partial_r u^{\delta} = \frac{1}{2} \Delta u^{\delta} + \beta u^{\delta} \xi_{\delta} - K_{\delta} u^{\delta} , \quad u^{\delta}(s, \cdot) = \delta_y .$$
 (2.10)

We also fix some final time t and consider the solution  $v^{\delta}$  to the random *backwards* PDE

$$\partial_r v^{\delta} = -\frac{1}{2} \Delta v^{\delta} - \beta v^{\delta} \xi_{\delta} + K_{\delta} v^{\delta} , \quad v^{\delta}(t, \cdot) = 1 .$$
 (2.11)

At this stage we note that if we set  $\tilde{v}^{\delta}(r, x) = \mathcal{R}v^{\delta}(-r, x)$ , then  $\tilde{v}^{\delta}$  satisfies the *forwards* PDE

$$\partial_r \tilde{v}^{\delta} = \frac{1}{2} \Delta \tilde{v}^{\delta} + \beta \tilde{v}^{\delta} \xi_{\delta} - K_{\delta} \tilde{v}^{\delta} , \quad \tilde{v}^{\delta}(-t, \cdot) = 1 .$$
 (2.12)

At this point we note that (2.10) fits into the setting of [HL18, Thm 5.2] (in fact, the regularity of our limiting noise  $\xi$  is better than what is considered there so some of the arguments could be simplified). Combining this with [HS23] which shows that the BPHZ lift of  $\xi^{\delta}$  to the regularity structure of [HL18, Sec. 2.1] converges as  $\delta \to 0$ , we conclude that, provided that the constants  $K_{\delta}$  are chosen appropriately,  $u^{\delta}$  converges as  $\delta \to 0$  to the Green's function  $w_{s,y}$ . This convergence takes place in an exponentially weighted space that enforces rapid decay of the solution.

By the exact same argument, one finds that  $\tilde{v}^{\delta}$  converges to  $u^{(-t)}$  as  $\delta \to 0$ , this time in a weighted space that allows for some slow growth of the solutions. In particular, this shows that  $v_r^{\delta}$  converges to

$$v_r(x) = \mathcal{R}u^{(-t)}(-r, x) = \bar{Z}_{r-t}(r, x)$$
.

We conclude that, for any  $r \in [s, t]$ , one has

$$\lim_{\delta \to 0} \langle u_r^{\delta}, v_r^{\delta} \rangle = \int_{\mathbb{R}^d} w_{s,y}(r, x) \bar{Z}_{r-t}(r, x) \, dx \; .$$

On the other hand, it follows immediately from (2.10) and (2.11) that  $\langle u_r^{\delta}, v_r^{\delta} \rangle$  is independent of r, as in the formal calculation (2.9). Since  $Z_0 = 1$  and  $\tilde{Z}_0(t, \cdot) = 1$ , it follows that

$$\dot{Z}_{s-t}(s,y) = \int_{\mathbb{R}^d} w_{s,y}(t,x) \dot{Z}_0(t,x) \, dx = C_{s,y}(t) ,$$

which is precisely as claimed. The rate of convergence follows from (2.8) and Theorem 2.1.

#### 2.3 A priori bounds

Our main goal in this subsection is to establish  $L^p$  bounds on the Malliavin derivative of the random variable u as well as on  $\Phi(u)$ . We write  $\mathcal{H}$  for the reproducing kernel space of the noise  $\xi$ , which is formed by completing the space  $\mathcal{C}_c^{\infty}(\mathbb{R} \times \mathbb{R}^d)$ under the inner product

$$\langle f,g \rangle_{\mathcal{H}} = \int_{-\infty}^{\infty} \int_{\mathbb{R}^{2d}} f(s,y_1)g(s,y_2)R(y_1 - y_2) \, dy_1 \, dy_2 \, ds \,,$$
 (2.13)

after quotienting out zero norm elements. The family of random variables

$$\xi(f) = \int f(s, y) \,\xi(ds, dy) \,, \qquad f \in \mathcal{H} \,,$$

defines an isonormal Gaussian process on  $\mathcal{H}$ :  $\mathbb{E}[\xi(f)\xi(g)] = \langle f,g \rangle_{\mathcal{H}}$ . Denote by Dthe Malliavin derivative D of a random variable,  $D(\xi(f)) = f$ . Let  $\mathcal{C}_p^{\infty}(\mathbb{R}^n)$  denote the space of smooth functions  $F \in \mathcal{C}^{\infty}(\mathbb{R}^n)$  with F and its partial derivatives growing at most polynomially. Denoting by S the class of smooth random variables of the form  $F(\xi(f_1), \ldots, \xi(f_n))$ , we set

$$DF(\xi(f_1),\ldots,\xi(f_n)) = \sum_{i=1}^n \partial_i F(\xi(f_1),\ldots,\xi(f_n)) f_i.$$

The Malliavin derivative operator  $D: L^p(\Omega) \to L^p(\Omega; \mathcal{H})$ , with initial domain S is closable for any p > 1. We define  $\mathbb{D}^{1,p}$  as the completion of smooth random variables S under the norm

$$\|X\|_{\mathbb{D}^{1,p}}^p \stackrel{\text{\tiny def}}{=} \mathbb{E}(|X|^p) + \mathbb{E}(\|DX\|_{\mathcal{H}}^p).$$

In all the cases we consider, our random variables X will be such that their Malliavin derivatives DX have natural function-valued representatives, and we denote their pointwise evaluations by  $D_{t,x}X$ . For example, for a deterministic function h,  $D_{r,z}\xi(h) = h(r, z)$ . The  $L^2$  adjoint of D is denoted by  $\delta$  and is known as the divergence operator or Skorokhod integral. It is shown in [Nua06, Prop. 1.5.4] that, for every  $p \ge 2$ ,  $\delta$  extends to a bounded operator  $\delta : \mathbb{D}^{1,p}(\mathcal{H}) \to L^p(\Omega)$ . For a deterministic  $f \in \mathcal{H}$ , we have  $\delta(f) = \xi(f)$  and  $D(\xi(f)) = f$ . If  $w \in \text{Dom } \delta$  is  $\mathcal{F}_t$ -adapted,  $\delta(w)$  coincides with the Itô integral. We refer to e.g. [Nua06] for more details on Malliavin calculus.

Before stating the lemma, we recall some covariance estimates – used for the a priori bounds on the moments below and Theorems 1.3-1.5 in next sections.

**Lemma 2.4.** If  $R(x) \leq (1 + |x|^{\kappa})^{-1}$ , where  $\kappa \in (2, d)$ , then uniformly in t > 0 and in  $x \in \mathbb{R}^d$ ,

$$\int_{\mathbb{R}^d} P_t(x-z) R(z) \, dz \lesssim (1+|x|+\sqrt{t})^{-\kappa}, \quad (2.14)$$

$$\int_0^\infty \int_{\mathbb{R}^d} P_r(z-x) R(z) \, dr \, dz \lesssim 1 \wedge |x|^{2-\kappa}, \tag{2.15}$$

$$\int_0^t \int_{\mathbb{R}^{2d}} P_r(z_1 - x) P_{t-r}(z_2) R(z_1 - z_2) \, dr \, dz_1 \, dz_2 \lesssim t(1 + |x| + \sqrt{t})^{-\kappa}.$$
(2.16)

*Proof.* The first two estimates follow from the proof of [GHL23, Lem. 2.3]. By a change of variables, the left hand side of (2.16) equals

$$\int_0^t \int_{\mathbb{R}^d} P_t(y-x) R(y) \, dr \, dy \lesssim t(1+|x|+\sqrt{t})^{-\kappa} \; ,$$

so that (2.16) follows from (2.14).

In the following we gather moment estimates on u,  $\vec{Z}$  and the Green's function  $w_{s,y}$  appearing in Theorem 1.5. Crucially, for sufficiently small  $\beta$ , the negative moments of u are also uniformly bounded.

**Lemma 2.5.** Assume that R satisfies Assumption 1.1. For any  $p \ge 2$ , there exist constants  $\beta_0(p) > 0$  and  $C_p$ , such that if  $\beta \le \beta_0$ , the following holds for the solution of (1.7).

(i) The solution of (1.7) has positive and negative moments of all orders:

$$\sup_{s \ge 0, x \in \mathbb{R}^d} \|u(s, x)\|_p \leqslant C_p , \quad \sup_{s \ge 0, x \in \mathbb{R}^d} \|u(s, x)^{-1}\|_p \leqslant C_p .$$
(2.17)

(ii) For all  $x \in \mathbb{R}^d$  and t > 0,  $u(t, x) \in \mathbb{D}^{1,p}$ . The Malliavin derivative of u is given by

$$D_{s,y}u(t,x) = \beta u(s,y)w_{s,y}(t,x)\mathbf{1}_{[0,t]}(s), \qquad (2.18)$$

and the process w satisfies the pointwise estimate

$$||w_{s,y}(t,x)||_p \lesssim P_{t-s}(x-y)$$
. (2.19)

(iii) The process  $C_{s,y}(t)$  defined in (2.7) satisfies  $\sup_y \sup_{t>s \ge 0} ||C_{s,y}(t)||_p \lesssim 1$ and

$$\|C_{s,y}(t) - C_{s,y}(\tau)\|_p \lesssim \beta (1 + (t \wedge \tau) - s)^{\frac{2-\kappa}{4}} .$$
(2.20)

Assertion (i) also holds if we replace u by  $\vec{Z}$ , and one has

$$D_{r,z}\vec{Z}(t,x) = \beta \vec{Z}(r,z)w_{r,z}(t,x)\mathbf{1}_{(-\infty,t]}(r) .$$
(2.21)

*Proof.* The positive moment bounds in (2.17) follow from [GHL23, Lem. 2.8]. Concerning the Malliavin derivative, we note that by the mild solution formulation we have  $u(t, x) = 1 + \beta \delta(v_{t,x})$ , where

$$v_{t,x}(r,z) = P_{t-r}(x-z)u(r,z)\mathbf{1}_{[0,t]}(r)$$

Differentiating on both sides and using the commutation relation [Nua06, Eq. 1.46], it follows that for  $s \in [0, t]$  one has

$$D_{s,y}u(t,x) = \beta P_{t-s}(x-y)u(s,y) + \beta \int_{s}^{t} \int_{\mathbb{R}^{d}} P_{t-r}(x-z)D_{s,y}u(r,z)\,\xi(dr,dz) \;.$$

This shows that  $D_{s,y}u$  solves the same equation as u itself, but with initial condition  $\beta u(s, \cdot)\delta_y$  at time s, so that (2.18) follows by linearity.

The moment bound on  $w_{s,y}$  follows by the same steps as in the proof of [GHL23, Lem. 2.8] (we omit tracking the same constants here). Consider  $w_{s,y}^{(0)}(t,x) := P_{t-s}(x-y)$ , and

$$w_{s,y}^{(n+1)}(t,x) = P_{t-s}(x-y) + \beta \int_s^t \int_{\mathbb{R}^d} P_{t-r}(x-z) w_{s,y}^{(n)}(r,z) \,\xi(dr,dz) \;.$$

Letting  $\tilde{w}_{s,y}^{(n)}(t,x) = w_{s,y}^{(n+1)}(t,x) - w_{s,y}^{(n)}(t,x)$ , by (3.10) one has

$$\begin{split} \|\tilde{w}_{s,y}^{(n)}(t,x)\|_{p} \\ &\leqslant \beta C_{p} \bigg( \int_{s}^{t} \int_{\mathbb{R}^{2d}} \Big( \prod_{i=1}^{2} P_{t-r}(x-z_{i}) \|\tilde{w}_{s,y}^{(n-1)}(r,z_{i})\|_{p} \Big) |R(z_{1}-z_{2})| \, dz \, dr \bigg)^{\frac{1}{2}} \\ &\lesssim \beta C_{p} \sup_{r>s} \sup_{z} \|w_{s,y}^{(n-1)}(r,z)\|_{p} \, . \end{split}$$

Thus, for sufficiently small  $\beta \leq \beta_0(p)$ ,  $w_{s,y}^{(n)}$  converges to  $w_{s,y}$  as  $n \to \infty$ . Moreover, again by (3.10), we have

$$\|w_{s,y}^{(n+1)}(t,x)\|_{p}^{2} \lesssim P_{t-s}(x-y) + \beta^{2} \int_{s}^{t} \int_{\mathbb{R}^{2d}} R(z_{1}-z_{2}) \Big(\prod_{i=1}^{2} P_{t-r}(x-z_{i}) \|w_{s,y}^{(n)}(s,z_{i})\|_{p} \Big) dz dr .$$

Given these bounds and Lemma 2.4, (2.19) follows by [GHL23, Lem. 2.7].

In view of Proposition 2.3, the bounds on the mass process C are an immediate consequence of (2.1) in the uniform moment bounds on  $\vec{Z}$ .

The uniform control on the negative moments of u in (2.17) is a consequence of the fact that  $\log u(t, x)$  has sub-Gaussian left tails, namely for  $\theta > 0$ ,

$$\mathbb{P}(\log u(t,x) \leqslant -\theta) \leqslant C e^{-\frac{\theta^2}{2}}.$$
(2.22)

The proof of (2.22) follows by [CCM20, Thm. 1.3], given that u is uniformly  $L^2$  bounded for any  $\beta \leq \beta_0(2)$ . Their strategy to discretize and cut-off of the mollifier  $\phi$  does not require  $\phi$  to be compactly supported. Then, all steps carry through once noted that even with long range covariance R, uniformly in t > 0,

$$\mathbf{E}_{B^{(1)},B^{(2)}} \Big[ \int_0^t R(B_s^{(1)} - B^{(2)}) \, ds \, e^{\beta^2 \int_0^t R(B_s^{(1)} - B^{(2)}) \, ds} \Big] \lesssim \sup_{\substack{\beta \leqslant \beta_0(2), \\ t > 0}} \mathbb{E}[u(t,0)^2] \lesssim 1,$$

where  $\mathbf{E}_{B^{(1)},B^{(2)}}$  denotes the expectation with respect to Brownian motions  $B^{(i)}$  independent of  $\xi$ . The first inequality above follows by noting that, via Feynman-Kac formula, one has an explicit expression for the second moment of u (cf. [MSZ16, Lem. 3.1]):

$$\mathbb{E}[u(t,0)^{2}] = \mathbf{E}_{B^{(1)},B^{(2)}} \left[ e^{\beta^{2} \int_{0}^{t} R(B_{s}^{(1)} - B^{(2)}) \, ds} \right]$$

Regarding  $\vec{Z}$ , we note that it solves the fixed point problem

$$\vec{Z}(t,x) = 1 + \beta \int_{-\infty}^{t} \int_{\mathbb{R}^d} P_{t-s}(x-y)\vec{Z}(s,y)\,\xi(ds,dy)\,,\qquad(2.23)$$

whence we conclude that

$$D_{r,z}\vec{Z}(t,x) = \beta P_{t-r}(x-z)\vec{Z}(r,z) + \beta \int_{-\infty}^{t} \int_{\mathbb{R}^d} P_{t-s}(x-y)D_{r,z}\vec{Z}(s,y)\,\xi(ds,dy).$$

Since  $D_{r,z}\vec{Z}(t,x) = 0$  for r > t, assertion (2.21) holds. Since the bounds in (i) are uniform in time, their extension to  $\vec{Z}$  is immediate. It remains to show (2.20), which follows from Proposition 2.3, the fact that  $\bar{Z}(s,y)(\xi) = \lim_{r \to -\infty} u^{(r)}(-s,y)(\mathcal{R}\xi)$ , and Theorem 2.1.

With positive and negative moment bounds in place, we can Malliavin differentiate transformations of u and then approximate fluctuations through the homogenisation result.

**Lemma 2.6.** Assume  $\Psi \in C^1(\mathbb{R}_+)$  and that, for some  $M, q \ge 1$ , satisfies for all  $z \in \mathbb{R}_+$ ,

$$|\Psi(z)| + |\Psi'(z)| \leqslant M(z^{-q} + z^q).$$
(2.24)

Then, for any p > 1 there exists  $\beta_0^{M,q}(p) = \beta_0(p, d, R, q, M)$  such that for any  $\beta \leq \beta_0^{M,q}(p)$ , we have  $\Psi(u(t, x)) \in \mathbb{D}^{1,p}$  with

$$D_{s,y}[\Psi(u(t,x))] = \beta \Psi'(u(t,x))u(s,y)w_{s,y}(t,x)\mathbf{1}_{[0,t]}(s), \qquad (2.25)$$

and the following bounds are satisfied:

$$\|\Psi(u(t,x))\|_p \lesssim M$$
,  $\|\Psi'(u(t,x))\|_p \lesssim M$ , (2.26)

$$\|D_{s,y}\Psi(u(t,x))\|_p \lesssim MP_{t-s}(x-y) , \quad \|\Psi(u(t,x))\|_{\mathbb{D}^{1,p}} \lesssim M , \qquad (2.27)$$

uniformly over s, t, x, y. Moreover, uniformly over  $x \in \mathbb{R}^d$  we have

$$\sup_{t>0} \left| \operatorname{cov} \left( \Psi(u(t,x)), \Psi(u(t,0)) \right) \right| \lesssim M \left( 1 \wedge |x|^{2-\kappa} \right) . \tag{2.28}$$

All the estimates above also hold with  $\vec{Z}(t, x)$  in place of u(t, x). Finally, we also have the following convergence rate

$$\|\Psi(u^{(s)}(t,x)) - \Psi(\vec{Z}(t,x))\|_p \lesssim 1 \wedge (t-s)^{\frac{2-\kappa}{4}}$$
. (2.29)

*Proof.* For  $\beta \leq \beta_0(pq)$ , bounds (2.26) follow immediately from (2.17) and the assumptions on  $\Psi$ . Observe that  $\beta_0(p)$  is non-increasing in p.

Regarding the Malliavin derivative, let us introduce an approximation by bounded functions  $\Psi_n \in \mathcal{C}_b^1(\mathbb{R}_+)$  such that  $\Psi_n = \Psi$  on  $[\frac{1}{n}, n]$ . Consider  $\varrho_n : \mathbb{R}_+ \to \mathbb{R}_+$  defined by

$$\varrho_n(x) := \begin{cases}
\frac{1}{2n} + \frac{n}{2}x^2 & \text{for } x \in [0, \frac{1}{n}], \\
x & \text{for } x \in [\frac{1}{n}, n], \\
\frac{3}{2}n - \frac{1}{2n}(x - 2n)^2 & \text{for } x \in [n, 2n], \\
\frac{3}{2}n & \text{for } x \in [2n, +\infty).
\end{cases}$$

Then, we let  $\Psi_n := \Psi \circ \varrho_n$ . Note that  $\varrho_n \in \mathcal{C}_b^1(\mathbb{R}_+)$  with  $\sup_x |\varrho'_n(x)| \leq 1$  and range  $\varrho_n(x) \in [\frac{1}{2n}, \frac{3}{2}n]$ . Thus,  $\Psi_n \in \mathcal{C}_b^1(\mathbb{R}_+)$  for all n. Moreover, we note that uniformly in n:

$$|\varrho_n(x)| \leq |x|+1, \qquad |\varrho_n(x)|^{-1} \leq \frac{1}{|x|}+1.$$

Then, given growth control (2.24) of  $\Psi$ , for  $\beta \leq \beta_0(pq)$  we can also infer

$$\sup_{n>0,s\geqslant 0,x\in\mathbb{R}^d} \|\Psi_n(u(s,x))\|_p \lesssim 1, \qquad \sup_{n>0,s\geqslant 0,x\in\mathbb{R}^d} \|\Psi'_n(u(s,x))\|_p \lesssim 1.$$

Since  $u(t, x) \in \mathbb{D}^{1,p}$  by Lemma 2.5, we have  $\Psi_n(u(t, x)) \in \mathbb{D}^{1,p}$  with  $D\Psi_n(u(t, x)) = \Psi'_n(u(t, x))Du(t, x)$  by [Nua06, Prop. 1.2.3]. We let

$$F_{r,z}^{(n)}(t,x) = D_{r,z}(\Psi_n(u(t,x))) - \Psi'(u(t,x))D_{r,z}u(t,z)$$
  
=  $\mathbf{1}_{\{\frac{1}{n} \leq u(t,x) \leq n\}^c}(\Psi'_n(u(t,x)) - \Psi'(u(t,x)))D_{r,z}u(t,x).$ 

Then, for sufficiently small  $\beta$  we use the moments estimates on  $u, \Psi', \Psi'_n$  and Du,

$$\|F_{r,z}^{(n)}(t,x)\|_{p} \leq \mathbb{P}(\{u < \frac{1}{n}\} \cup \{u > n\})^{\frac{1}{3p}} \|\Psi_{n}'(u(t,x)) - \Psi'(u(t,x))\|_{3p} \|D_{r,z}u(t,x)\|_{3p}$$

$$\lesssim n^{-\frac{1}{p}} M P_{t-r}(x-z).$$
 (2.30)

By Minkowski, Hölder's inequalities and (2.15),

$$\mathbb{E}[\|F^{(n)}(t,x)\|_{\mathcal{H}}^{p}] = \mathbb{E}\Big[\Big(\int_{0}^{t}\int_{\mathbb{R}^{2d}}\prod_{i=1}^{2}F^{(n)}_{r,z_{i}}(t,x)R(z_{1}-z_{2})\,dr\,dz\Big)^{\frac{p}{2}}\Big]$$

$$\leqslant \Big(\int_{0}^{t}\int_{\mathbb{R}^{2d}}\|\prod_{i=1}^{2}F^{(n)}_{r,z_{i}}(t,x)\|_{\frac{p}{2}}R(z_{1}-z_{2})\,dr\,dz\Big)^{\frac{p}{2}}$$

$$\leqslant n^{-1}M^{p}\Big(\int_{0}^{t}\int_{\mathbb{R}^{2d}}\prod_{i=1}^{2}P_{t-r}(x-z_{i})R(z_{1}-z_{2})\,dr\,dz\Big)^{\frac{p}{2}}$$

$$\leqslant CM^{p}n^{-1}.$$

Thus,  $F^{(n)}(t,x) \to 0$  in  $L^p(\Omega, \mathcal{H})$  as  $n \to \infty$ . On the other hand,  $\Psi_n(u(t,x)) \to \Psi(u(t,x))$  in  $L^p(\Omega)$ . Since D is closed, we conclude  $\Psi(u(t,x)) \in \text{Dom}(D)$  and  $D_{s,y}\Psi(u(t,x)) = \Psi'(u(t,x))D_{s,y}u(t,x)$ . Then, given (2.18), representation (2.25) holds. Moreover, for sufficiently small  $\beta$ , given (2.19) and the uniform moments bounds on  $\Psi'(u)$  and u, we have  $\|D_{r,z}\Psi(u(t,x))\|_p \leq MP_{t-r}(x-z)$ . Since the inequality constant is independent of (t,x), we estimate similarly to above

$$\mathbb{E}[\|D\Psi(u(t,x))(t,x)\|_{\mathcal{H}}^{p}] \leq \left(\int_{0}^{t} \int_{\mathbb{R}^{2d}} \prod_{i=1}^{2} \|D_{r,z_{i}}\Psi(u(t,x))\|_{p} R(z_{1}-z_{2}) dr dz\right)^{\frac{p}{2}} \\ \leq \left(\int_{0}^{\infty} \int_{\mathbb{R}^{2d}} \prod_{i=1}^{2} P_{t-r}(x-z_{i})R(z_{1}-z_{2}) dr dz\right)^{\frac{p}{2}} \lesssim 1,$$

where we used (2.15). This concludes the proof of (2.27).

Using the Clark–Ocone formula in the same way as in [GHL23, Lem. 2.12], we obtain the bound

$$|\operatorname{cov}(\Psi(u(s,x)),\Psi(u(s,0)))| \lesssim M \int_0^t \int_{\mathbb{R}^{2d}} P_{t-r}(x-z_1)P_{t-r}(z_2)R(z_1-z_2)\,dz\,dr,$$

so that (2.28) follows by (2.15).

It remains to show (2.29), for which we make use of the identity

$$\Psi(u^{(s)}(t,x)) - \Psi(\vec{Z}(t,x)) = \left(u^{(s)}(t,x) - \vec{Z}(t,x)\right) \int_0^1 \Psi'(\theta u^{(s)}(t,x) + (1-\theta)\vec{Z}(t,x)) \, d\theta$$

Using the fact that u is non-negative and the uniform a priori bounds on positive and negative moments of u and  $\vec{Z}$ , one obtains a uniform  $L^{2p}$  bound on  $\int_0^1 \Psi'(\theta u(t, x) + (1 - \theta)Z(t, x))d\theta$  for sufficiently small  $\beta$ . Thus, (2.1) implies

$$\|\Psi(u^{(s)}(t,x)) - \Psi(\vec{Z}(t,x))\|_p \lesssim M \|u^{(s)}(t,x) - \vec{Z}(t,x)\|_{2p} \lesssim 1 \wedge (t-s)^{\frac{2-\kappa}{4}},$$

which is precisely (2.29), concluding the proof.

## **3** The main homogenisation theorem

In view of Theorem 1.5, recalling that  $\vec{Z}(t,x) = \lim_{s \to -\infty} u^{(s)}(t,x)$ , we define

$$\rho_{s,y}(t,x) = \frac{w_{s,y}(t,x)}{P_{t-s}(x-y)} - C_{s,y}(t)\vec{Z}(t,x) .$$
(3.1)

By Proposition 2.3 and (2.20), we know that

$$\|C_{s,y}(t) - \bar{Z}(s,y)\|_p \lesssim \beta (1+t-s)^{\frac{2-\kappa}{4}}$$

Since  $\kappa > 2$ , our main homogenisation result then follows from the following proposition. The proof relies on a kernel estimate and moment bounds on Skorokhod correction terms proved below.

**Proposition 3.1.** Let  $\mu = \frac{1}{2} - \frac{1}{\kappa}$ . Then, given  $p \ge 1$ , there exists  $\tilde{\beta}(p) > 0$ , such that for  $\beta \le \tilde{\beta}$ , the following holds

$$\sup_{x,y} \sup_{t>s\geq 0} \frac{\|\rho_{s,y}(t,x)\|_p}{(1+t-s)^{-\mu}(1+(1+t-s)^{-\frac{1}{2}}|x-y|)} \lesssim \beta .$$
(3.2)

*Proof.* Let us consider  $\tilde{w}_{s,y}(t,x) = w_{s,y}(t,x)/P_{t-s}(x-y)$ . Since  $w_{s,y}(t,x)$  solves

$$w_{s,y}(t,x) = P_{t-s}(x-y) + \beta \int_{s}^{t} \int_{\mathbb{R}^{d}} P_{t-r}(x-z) w_{s,y}(r,z) \,\xi(dr,dz) \,, \quad (3.3)$$

one has the identity

$$\tilde{w}_{s,y}(t,x) = 1 + \beta \int_{s}^{t} \int_{\mathbb{R}^{d}} q_{s,t}^{x,y}(r,z) \tilde{w}_{s,y}(r,z) \xi(dr,dz) , \qquad (3.4)$$

where  $q_{s,t}^{x,y}$  is kernel of the Brownian bridge starting from s at x and ending at y at time t,

$$q_{s,t}^{x,y}(r,z) = \frac{P_{r-s}(z-y)P_{t-r}(x-z)}{P_{t-s}(x-y)} = P_{(r-s)(t-r)/(t-s)}(z-y-\frac{r-s}{t-s}(x-y)) .$$
(3.5)

On the other hand, integrating (3.3) over x, we obtain

$$C_{s,y}(t) = 1 + \beta \int_{s}^{t} \int_{\mathbb{R}^{d}} P_{r-s}(z-y)\tilde{w}_{s,y}(r,z)\xi(dr,dz) .$$
(3.6)

As a consequence, we can wrte  $\rho_{s,y}$  as

$$\rho_{s,y}(t,x) = \tilde{w}_{s,y}(t,x) - C_{s,y}(t)\vec{Z}(t,x) 
= 1 + \beta \int_{s}^{t} \int_{\mathbb{R}^{d}} q_{s,t}^{x,y}(r,z)\tilde{w}_{s,y}(r,z)\xi(dr,dz) 
- \left(1 + \beta \int_{-\infty}^{t} \int_{\mathbb{R}^{d}} P_{t-r}(x-z)\vec{Z}(r,z)\xi(dr,dz)\right)C_{s,y}(t) 
= \beta \int_{s}^{t} \int_{\mathbb{R}^{d}} (q_{s,t}^{x,y}(r,z) - P_{r-s}(y-z))\tilde{w}_{s,y}(r,z)\xi(dr,dz) 
- \beta \int_{-\infty}^{t} \int_{\mathbb{R}^{d}} P_{t-r}(x-z)\vec{Z}(r,z)\xi(dr,dz)C_{s,y}(t) .$$
(3.7)

To treat the term  $C_{s,y}(t)$ , we split the last integral into two parts, integrating over  $r \in (-\infty, s)$  leads to a contribution proportional to

$$E_1 := \int_{-\infty}^s \int_{\mathbb{R}^d} P_{t-r}(x-z) \vec{Z}(r,z) \,\xi(dr,dz) \,C_{s,y}(t) \,. \tag{3.8}$$

For the other region  $r \in [s,t]$ , we rewrite it in terms of a Skorokhod integral using the fact (cf. [Nua06, Prop. 1.3.3]) that  $F \int G d\xi = \delta(FG) + \langle G, DF \rangle_{\mathcal{H}}$ with  $F = C_{s,y}(t)$  and  $G(r, z) = P_{t-r}(x - z)\vec{Z}(r, z)\mathbf{1}_{[s,t]}(r)$ . Moreover,  $C_{s,y}(r)$  is  $\mathcal{F}_r$ -adapted and we obtain, writing  $\delta\xi$  for the Skorokhod integration:

$$\begin{split} &\int_{s}^{t} \int_{\mathbb{R}^{d}} P_{t-r}(x-z) \vec{Z}(r,z) \,\xi(dr,dz) \,C_{s,y}(t) \\ &= \int_{s}^{t} \int_{\mathbb{R}^{d}} P_{t-r}(x-z) \vec{Z}(r,z) C_{s,y}(t) \,\delta\xi(dr,dz) \\ &+ \int_{s}^{t} \int_{\mathbb{R}^{2d}} P_{t-r}(x-z_{1}) \vec{Z}(r,z_{1}) \,D_{r,z_{2}} C_{s,y}(t) R(z_{1}-z_{2}) \,dr \,dz \\ &= \int_{s}^{t} \int_{\mathbb{R}^{d}} P_{t-r}(x-z) \vec{Z}(r,z) C_{s,y}(r) \,\xi(dr,dz) \\ &+ \int_{s}^{t} \int_{\mathbb{R}^{d}} P_{t-r}(x-z) \vec{Z}(r,z) (C_{s,y}(t)-C_{s,y}(r)) \,\delta\xi(dr,dz) \\ &+ \int_{s}^{t} \int_{\mathbb{R}^{2d}} P_{t-r}(x-z_{1}) \vec{Z}(r,z_{1}) \,D_{r,z_{2}} C_{s,y}(t) R(z_{1}-z_{2}) \,dr \,dz \;. \end{split}$$

Bringing all these back to (3.7) and given the definition (3.1) of  $\rho_{s,y}$ , we obtain

$$\rho_{s,y}(t,x) = \beta \int_s^t \int_{\mathbb{R}^d} P_{t-r}(x-z)\rho_{s,y}(r,z)\xi(dr,dz) + \beta E_0 - \beta \sum_{i=1}^3 E_i , \ (3.9)$$

where  $E_1$  was defined in (3.8) and the other terms are given by

$$\begin{split} E_0 &:= \int_s^t \int_{\mathbb{R}^d} \left( q_{s,t}^{x,y}(r,z) - P_{r-s}(y-z) - P_{t-r}(x-z) \right) \tilde{w}_{s,y}(r,z) \,\xi(dr,dz) \;, \\ E_2 &:= \int_s^t \int_{\mathbb{R}^d} P_{t-r}(x-z) \vec{Z}(r,z) (C_{s,y}(t) - C_{s,y}(r)) \,\delta\xi(dr,dz) \;, \\ E_3 &:= \int_s^t \int_{\mathbb{R}^{2d}} P_{t-r}(x-z_1) \vec{Z}(r,z_1) \, D_{r,z_2} C_{s,y}(t) R(z_1-z_2) \, dr \, dz \;. \end{split}$$

We can now estimate their  $L^p$  norms. Given a time interval I = [a, b] and an adapted random process f(s, y), by Burkholder–Davis–Gundy (BDG) inequality, followed by Minkowski and Hölder inequalities, we have the following estimate

$$\left\|\int_{I\times\mathbb{R}^d} f(s,y)\xi(ds,dy)\right\|_p \leqslant C_p \mathbb{E}\left[\left(\int_I \int_{\mathbb{R}^{2d}} f(s,y_1)f(s,y_2)R(y_1-y_2)\,dy\,ds\right)^{\frac{p}{2}}\right]^{\frac{1}{p}}$$

$$\leq C_p \Big( \int_I \int_{\mathbb{R}^{2d}} \|f(s, y_1)\|_p \|f(s, y_2)\|_p R(y_1 - y_2) \, dy \, ds \Big)^{\frac{1}{2}} \,. \tag{3.10}$$

Given the uniform moments bounds, on  $C_{s,y}$  and  $\vec{Z}$ , from Lemma 2.5 and the kernel estimate (2.14), for  $\beta \leq \beta_0(2p)$  we have

$$\begin{split} \|E_1\|_p &\lesssim \left\| \int_{-\infty}^s P_{t-r}(x-z)\vec{Z}(r,z)\,\xi(dr,dz) \right\|_{2p} \|C_{s,y}(t)\|_{2p} \\ &\lesssim \Big( \int_{-\infty}^s \int_{\mathbb{R}^{2d}} \prod_{j=1}^2 \Big( P_{t-r}(x-z_j) \|\vec{Z}(r,z_j)\|_{2p} \Big) R(z_1-z_2)\,dr\,dz \Big)^{\frac{1}{2}} \\ &\lesssim \Big( \int_{-\infty}^s (1+t-r)^{-\frac{\kappa}{2}} dr \Big)^{\frac{1}{2}} \lesssim (1+t-s)^{\frac{2-\kappa}{4}} \,. \end{split}$$

For  $E_0$ , we note that by (2.19),

$$\sup_{x,y} \sup_{t>s \ge 0} \|\tilde{w}_{s,y}(t,x)\|_p \lesssim 1 .$$

Then, estimating moments like above, followed by change of variables  $z_i \mapsto z_i - y$ ,  $r \mapsto r - s$ , and Lemma 3.3 below, we can bound  $E_0$  as follows for  $\beta \leq \beta_0(p)$ :

$$\begin{split} \|E_0\|_p &\leqslant C_p \Big( \int_s^t \int_{\mathbb{R}^{2d}} \prod_{i=1}^2 \Big( q_{s,t}^{x,y}(r,z_i) - P_{r-s}(y-z_i) - P_{t-r}(x-z_i) \Big) R(z_1-z_2) \, dr \, dz \Big)^{\frac{1}{2}} \\ &= C_p \Big( \int_0^{t-s} \int_{\mathbb{R}^{2d}} \prod_{i=1}^2 \Big( q_{0,t-s}^{x-y,0}(r,z_i) - P_{r-s}(z_i) - P_{t-r}(x-y-z_i) \Big) R(z_1-z_2) \, dr \, dz \Big)^{\frac{1}{2}} \\ &\lesssim (1+t-s)^{-\mu} (1+(1+t-s)^{-\frac{1}{2}} |x-y|) \, . \end{split}$$

By Lemma 3.2 below, for  $\beta \leq \beta_1(3p) \wedge \beta_0(3p)$  also the terms  $E_2$  and  $E_3$  satisfy

$$||E_i||_p \lesssim (1+t-s)^{-\mu}, \quad i=2,3.$$

Among the estimates, the worst decay rate is the bound on  $E_0$ , and we obtain

$$\sum_{i=0}^{3} \|E_i\|_p \lesssim m(t-s, x-y) ,$$

where we denote the space-time weight

$$m(r,z) = (1+r)^{-\mu}(1+(1+r)^{-\frac{1}{2}}|z|).$$

Finally, let us consider the weighted norm

$$M := \sup_{x,y} \sup_{t>s \ge 0} \frac{\|\rho_{s,y}(t,x)\|_p}{m(t-s,x-y)} \,.$$

We estimate the first term in (3.9) as follows:

$$\begin{split} \left| \int_{s}^{t} \int_{\mathbb{R}^{d}} P_{t-r}(x-z)\rho_{s,y}(r,z)\,\xi(dr,dz) \right\|_{p} \\ &\leqslant C_{p} \Big( \int_{s}^{t} \int_{\mathbb{R}^{2d}} \prod_{i=1}^{2} \left( P_{t-r}(x-z_{i}) \| \rho_{s,y}(r,z_{i}) \|_{p} \right) R(z_{1}-z_{2})\,dr\,dz \Big)^{\frac{1}{2}} \\ &\lesssim M \Big( \int_{s}^{t} (1+r-s)^{-2\mu} \int_{\mathbb{R}^{2d}} \Big( \prod_{i=1}^{2} P_{t-r}(x-z_{i}) \Big) R(z_{1}-z_{2})\,dr\,dz \Big)^{\frac{1}{2}} \\ &+ M \Big( \sum_{j=1}^{2} \int_{s}^{t} (1+r-s)^{-1-2\mu} \int_{\mathbb{R}^{2d}} \Big( \prod_{i=1}^{2} |z_{j}-y|^{2} P_{t-r}(x-z_{i}) \Big) R(z_{1}-z_{2})\,dr\,dz \Big)^{\frac{1}{2}} \\ &\lesssim M \Big( \int_{s}^{t} (1+r-s)^{-2\mu} (1+t-r)^{-\frac{\kappa}{2}} \,dr \Big)^{\frac{1}{2}} \\ &+ M \Big( \int_{s}^{t} (1+r-s)^{-1-2\mu} (1+t-r)^{-\frac{\kappa}{2}} \,dr \Big)^{\frac{1}{2}} \end{split}$$
(3.11)

where we used (2.14) and the fact that  $\int_{\mathbb{R}^d} |z|^2 P_r(z-x) dz \leq r+|x|^2$  for the second integral. For  $\alpha, \beta > 0$ , we note

$$\int_0^T (1+\tau)^{-\alpha} (1+T-\tau)^{-1-\beta} d\tau \lesssim (1+T)^{-\min(\alpha,\beta)} .$$
 (3.12)

This follows by splitting the integration region. On the interval [0,T/2], we have  $T-\tau \geqslant T/2$  and

$$\int_0^{\frac{T}{2}} (1+\tau)^{-\alpha} (1+T-\tau)^{-1-\beta} d\tau \lesssim (1+T)^{-1-\beta} T \lesssim (1+T)^{-\beta} ,$$

while on [T/2, T], we have

$$\int_{\frac{T}{2}}^{T} (1+\tau)^{-\alpha} (1+T-\tau)^{-1-\beta} d\tau \lesssim (1+T)^{-\alpha} \int_{0}^{\infty} (1+\tau)^{-1-\beta} d\tau \lesssim (1+T)^{-\alpha} .$$

Then, going back to (3.11), we apply (3.12) to both terms and obtain

$$\left\|\int_{s}^{t}\int_{\mathbb{R}^{d}}P_{t-r}(x-z)\rho_{s,y}(r,z)\,\xi(dr,dz)\right\|_{p} \lesssim M\ m(t-s,x-y)\ .$$

Hence, by the estimates above, for some C > 0 we have

$$\|\rho_{s,y}(t,x)\|_p \leqslant C\beta M m(t-s,x-y) + C\beta m(t-s,x-y) .$$

Taking  $\tilde{\beta} > 0$  such that  $C\tilde{\beta} < 1$  and  $\tilde{\beta} \leq \beta_1(3p) \wedge \beta_0(3p)$ , we conclude (3.2).  $\Box$ 

**Lemma 3.2.** Let  $p \ge 1$ . With notation as in the proof of Proposition 3.1, for any  $\beta \le \beta_1(3p) \land \beta_0(3p)$ , for i = 2, 3, we have the following uniformly in  $x, y \in \mathbb{R}^d$  and  $t > s \ge 0$ :

$$\begin{aligned} \|E_2(s,t,y,x)\|_p &\lesssim \beta \left(1+t-s\right)^{\frac{2-\kappa}{4}},\\ \|E_3(s,t,y,x)\|_p &\lesssim \beta \left(1+t-s\right)^{1-\frac{\kappa}{2}}. \end{aligned}$$

*Proof.* Let us start with the Skorokhod correction term  $E_3$ . Similarly to Lemma 2.5, where we obtained  $D_{s,y}u(t,x) = \beta u(s,y)w_{s,y}(t,x)\mathbf{1}_{[0,t]}(s)$ , one has

$$D_{r,z}w_{s,y}(t,x) = \beta w_{s,y}(r,z)w_{r,z}(t,x)\mathbf{1}_{[s,t]}(r) ,$$

so that, integrating over x,

$$D_{r,z}C_{s,y}(t) = \beta w_{s,y}(r,z)C_{r,z}(t)\mathbf{1}_{[s,t]}(r) .$$
(3.13)

Therefore, the term  $E_3$  is given by

$$E_{3} = \int_{s}^{t} \int_{\mathbb{R}^{2d}} P_{t-r}(x-z_{1})\vec{Z}(r,z_{1}) D_{r,z_{2}}C_{s,y}(t)R(z_{1}-z_{2}) dr dz$$
$$= \beta \int_{s}^{t} \int_{\mathbb{R}^{2d}} P_{t-r}(x-z_{1})\vec{Z}(r,z_{1}) w_{s,y}(r,z_{2})C_{r,z_{2}}(t)R(z_{1}-z_{2}) dr dz$$

Using the uniform moments estimates on  $\vec{Z}$ , w and C from Lemma 2.5, by Minkowski and Hölder's inequalities, followed by (2.16) we obtain for  $\beta \leq \beta_0(3p)$ :

$$\begin{split} \|E_3\|_p &\leq \beta \int_s^t \int_{\mathbb{R}^{2d}} P_{t-r}(x-z_1) \|\vec{Z}(r,z_1)w_{s,y}(r,z_2)C_{r,z_2}(t)\|_p R(z_1-z_2) \, dr \, dz \\ &\lesssim \beta \int_s^t \int_{\mathbb{R}^{2d}} P_{t-r}(x-z_1)P_{r-s}(z_2-y)R(z_1-z_2) \, dr \, dz \\ &\lesssim \beta (1+t-s)^{1-\frac{\kappa}{2}} \, . \end{split}$$

To estimate the  $L^p$  norm of  $E_2$ , we use the continuity of the Skorokhod integral  $\delta : \mathbb{D}^{1,p}(\mathcal{H}) \to L^p$  [Nua06, Prop. 1.5.4]. Letting

$$F_{s,t,x,y}(r,z) = P_{t-r}(x-z)\vec{Z}(r,z)(C_{s,y}(t) - C_{s,y}(r))\mathbf{1}_{[s,t]}(r) ,$$

we have  $E_2 = \delta(F_{s,t,x,y})$ . Therefore, dropping subscripts of F going forward, we have

$$||E_2||_p \lesssim ||F||_{L^p(\Omega,\mathcal{H})} + ||DF||_{L^p(\Omega,\mathcal{H}\otimes\mathcal{H})}.$$
(3.14)

We see that F is a pointwise product of a Malliavin differentiable random variable M and a deterministic element  $h \in \mathcal{H}$ . Namely, F(r, z) = M(r, z)h(r, z), where

$$M(r,z) = Z(r,z)(C_{s,y}(t) - C_{s,y}(r)), \qquad h(r,z) = P_{t-r}(x-z)\mathbf{1}_{[s,t]}(r).$$

Given (2.20), we note that

$$||M(r,z)||_p \lesssim ||\vec{Z}(r,z)||_{2p} ||C_{s,y}(t) - C_{s,y}(r)||_{2p} \lesssim \beta (1+r-s)^{\frac{2-\kappa}{4}}.$$

With this, using Minkowski inequality, (2.14) and (3.12), we bound the first term:

$$\begin{split} \|F\|_{L^{p}(\Omega,\mathcal{H})} &= \mathbb{E}\Big[\big(\int_{\mathbb{R}}\int_{\mathbb{R}^{2d}}F(r,z_{1})F(r,z_{2})R(z_{1}-z_{2})\,dr\,dz\big)^{\frac{p}{2}}\Big]^{\frac{1}{p}} \\ &\leqslant \Big(\int_{s}^{t}\int_{\mathbb{R}^{2d}}\|M(r,z_{1})M(r,z_{2})\|_{\frac{p}{2}}\,h(r,z_{1})h(r,z_{2})R(z_{1}-z_{2})\,dr\,dz\Big)^{\frac{1}{2}} \\ &\lesssim \Big(\beta^{2}\int_{s}^{t}(1+r-s)^{1-\frac{\kappa}{2}}\int_{\mathbb{R}^{2d}}P_{t-r}(x-z_{1})P_{t-r}(x-z_{2})R(z_{1}-z_{2})\,dr\,dz\Big)^{\frac{1}{2}} \\ &\lesssim \beta\Big(\int_{s}^{t}(1+r-s)^{1-\frac{\kappa}{2}}(1+t-r)^{-\frac{\kappa}{2}}\,dr\Big)^{\frac{1}{2}} \lesssim \beta(1+t-s)^{\frac{2-\kappa}{4}} \,. \end{split}$$

For the derivative term, by Minkowski and Hölder's inequalities we have

$$\begin{split} \|DF\|_{L^{p}(\Omega,\mathcal{H}\otimes\mathcal{H})} &= \mathbb{E}\Big[\big(\int_{\mathbb{R}}\int_{\mathbb{R}^{2d}} \langle DM(r,z_{1}), DM(r,z_{2})\rangle_{\mathcal{H}} \ h(r,z_{1})h(r,z_{2})R(z_{1}-z_{2}) \, dz \, dr\big)^{\frac{p}{2}}\Big]^{\frac{1}{p}} \\ &\leqslant \Big(\int_{s}^{t}\int_{\mathbb{R}^{2d}} \|\langle DM(r,z_{1}), DM(r,z_{2})\rangle_{\mathcal{H}}\|_{\frac{p}{2}} h(r,z_{1})h(r,z_{2})R(z_{1}-z_{2}) \, dz \, dr\Big)^{\frac{1}{2}} \\ &\leqslant \Big(\int_{s}^{t}\int_{\mathbb{R}^{2d}} \Big(\prod_{i=1}^{2} \|DM(r,z_{i})\|_{L^{p}(\Omega,\mathcal{H})} h(r,z_{i})\Big)R(z_{1}-z_{2}) \, dz \, dr\Big)^{\frac{1}{2}} \,. \end{split}$$

We note that  $\vec{Z}$  and C's are adapted, satisfying identities (2.21) and (3.13) respectively. Then,

$$\begin{split} D_{u,\bar{z}}M(r,z) &= D_{u,\bar{z}}\vec{Z}(r,z)\left(C_{s,y}(t) - C_{s,y}(r)\right) + \vec{Z}(r,z) D_{u,\bar{z}}(C_{s,y}(t) - C_{s,y}(r)) \\ &= \beta w_{u,\bar{z}}(r,z)\vec{Z}(u,\bar{z})(C_{s,y}(t) - C_{s,y}(r))\mathbf{1}_{[s,r]}(u) \\ &+ \beta \vec{Z}(r,z)w_{s,y}(u,\bar{z})(C_{u,\bar{z}}(t) - C_{u,\bar{z}}(r))\mathbf{1}_{[s,r]}(u) \\ &+ \beta \vec{Z}(r,z)w_{s,y}(u,\bar{z})C_{u,\bar{z}}(t)\mathbf{1}_{[r,t]}(u) =: \beta \sum_{j=1}^{3} m_{r,z}^{(j)}(u,\bar{z}) \,. \end{split}$$

By Minkowski and Hölder inequalities

$$\|DM(r,z)\|_{L^{p}(\Omega,\mathcal{H})} \lesssim \sum_{j,k=1}^{3} \left( \int_{s}^{t} \int_{\mathbb{R}^{2d}} \|m_{r,z}^{(j)}(u,\bar{z}_{1})\|_{p} \|m_{r,z}^{(k)}(u,\bar{z}_{2})\|_{p} R(\bar{z}_{1}-\bar{z}_{2}) \, d\bar{z} \, du \right)^{\frac{1}{2}}$$
(3.15)

Given the a priori estimates from Lemma 2.5, for  $\beta \leq \beta_0(3p)$  we have:

$$||m_{r,z}^{(1)}(u,\bar{z})||_p \lesssim P_{r-u}(z-\bar{z})(1+r-s)^{\frac{2-\kappa}{4}}\mathbf{1}_{[s,r]}(u)$$

$$\begin{split} & \|m_{r,z}^{(2)}(u,\bar{z})\|_p \lesssim P_{u-s}(\bar{z}-y)(1+r-u)^{\frac{2-\kappa}{4}} \mathbf{1}_{[s,r]}(u) , \\ & \|m_{r,z}^{(3)}(u,\bar{z})\|_p \lesssim P_{u-s}(\bar{z}-y) \mathbf{1}_{[r,t]}(u) . \end{split}$$

Hence, by Lemma 2.4, for j = k = 1 we obtain

$$\begin{split} &\int_{s}^{t} \int_{\mathbb{R}^{2d}} \left( \prod_{i=1}^{2} \| m_{r,z}^{(1)}(u,\bar{z}_{i}) \|_{p} \right) R(\bar{z}_{1}-\bar{z}_{2}) \, d\bar{z} \, du \\ &\lesssim (1+r-s)^{1-\frac{\kappa}{2}} \int_{s}^{r} \int_{\mathbb{R}^{2d}} \left( \prod_{i=1}^{2} P_{r-u}(z-\bar{z}_{i}) \right) R(\bar{z}_{1}-\bar{z}_{2}) \, d\bar{z} \, du \lesssim (1+r-s)^{1-\frac{\kappa}{2}} \, . \end{split}$$

On the other hand, by (2.14) and (3.12),

$$\begin{split} &\int_{s}^{t} \int_{\mathbb{R}^{2d}} \|m_{r,z}^{(1)}(u,\bar{z}_{1})\|_{p} \|m_{r,z}^{(2)}(u,\bar{z}_{2})\|_{p} R(\bar{z}_{1}-\bar{z}_{2}) \, d\bar{z} \, du \\ &\lesssim (1+r-s)^{\frac{2-\kappa}{4}} \int_{s}^{r} (1+r-u)^{\frac{2-\kappa}{4}} \int_{\mathbb{R}^{2d}} P_{r-u}(z-\bar{z}_{1}) P_{u-s}(\bar{z}_{2}-y) \, R(\bar{z}_{1}-\bar{z}_{2}) \, d\bar{z} \, du \\ &\lesssim (1+r-s)^{\frac{2-\kappa}{4}} \int_{s}^{r} (1+r-u)^{\frac{2-\kappa}{4}} (1+u-s)^{-\frac{\kappa}{2}} \, du \lesssim (1+r-s)^{1-\frac{\kappa}{2}} \, . \end{split}$$

Estimating in a similar way the other terms in (3.15) involving  $m_{r,z}^{(2)}$  and  $m_{r,z}^{(3)}$ , we obtain

$$\|DM(r,z)\|_{L^p(\Omega,\mathcal{H})} \lesssim \beta(1+r-s)^{\frac{2-\kappa}{4}}.$$

With this, using again (2.14) and (3.12), we return to estimate DF:

$$||DF||_{L^p(\Omega,\mathcal{H}\otimes\mathcal{H})}$$

$$\lesssim \beta \Big( \int_{s}^{t} (1+r-s)^{1-\frac{\kappa}{2}} \int_{\mathbb{R}^{2d}} P_{t-r}(x-z_1) P_{t-r}(x-z_2) R(z_1-z_2) \, dr \, dz \, dz \, dr \Big)^{\frac{1}{2}} \\ \lesssim \beta \Big( \int_{s}^{t} (1+r-s)^{1-\frac{\kappa}{2}} (1+t-r)^{-\frac{\kappa}{2}} \, dr \Big)^{\frac{1}{2}} \lesssim \beta (1+t-s)^{\frac{2-\kappa}{4}} \, .$$

Gathering bounds back to (3.14), we obtain  $||E_2(s,t,y,x)||_p \lesssim \beta (1+t-s)^{\frac{2-\kappa}{4}}$ .

The following kernel estimate leads to the  $L^p$  homogenisation rate.

**Lemma 3.3.** Let  $x \in \mathbb{R}^d$ , t > 0,  $P_r(x)$  the heat kernel, and let

$$H_{t,x}(r,z) = P_{r(t-r)/t}(z + \frac{r}{t}x) - P_r(z) - P_{t-r}(z+x) .$$

Let  $\tilde{\mu} = 1 - \frac{2}{\kappa}$ , then the following holds:

$$J_t(x) := \int_0^t \int_{\mathbb{R}^{2d}} \left( \prod_{i=1}^2 |H_{t,x}(r,z_i)| \right) R(z_1 - z_2) \, dz \, dr \lesssim (1+t)^{-\tilde{\mu}} (1 + (1+t)^{-1} |x|^2) \, .$$

*Remark* 3.4. Since  $P_{r(t-r)/t}(z + \frac{r}{t}x) = q_{0,t-s}^{x,0}(r,z)$ , this is really just a quantitative way of stating that, when considered over a very large time interval, a Brownian bridge is close to a Brownian motion near either end of the interval. Furthermore, the integral  $J_t$  is such that the contribution coming from the "bulk" is small.

*Proof.* Let  $\alpha = \frac{2}{\kappa}$  such that  $\tilde{\mu} = \alpha(\frac{\kappa}{2} - 1)$ . By Lemma 2.4, we have  $J_{t,x} \leq 1$  uniformly in t. Then, in the following we can assume  $t > 2^{\frac{1}{1-\alpha}}$ . We consider three regions of integration,

$$J_t(x) = J_{0,t^{\alpha}}(x) + J_{t^{\alpha},t-t^{\alpha}}(x) + J_{t-t^{\alpha},t}(x) ,$$

where

$$J_{t_1,t_2}(x) = \int_{t_1}^{t_2} \int_{\mathbb{R}^{2d}} \left( \prod_{i=1}^2 |H_{t,x}(r,z_i)| \right) R(z_1 - z_2) \, dz \, dr \, .$$

By changing variables  $z \mapsto z + x$  and  $r \mapsto t - r$ , we have  $J_{0,t^{\alpha}}(x) = J_{t-t^{\alpha},t}(-x)$ . Hence, it suffices to estimate  $J_{0,t^{\alpha}}(x)$  and  $J_{t^{\alpha},t-t^{\alpha}}(x)$ .

For  $J_{t^{\alpha},t-t^{\alpha}}(x)$ , since  $r \in [t^{\alpha},t-t^{\alpha}]$ , using (2.14) for each Gaussian kernel:

$$\begin{aligned} J_{t^{\alpha},t-t^{\alpha}}(x) &= \int_{t^{\alpha}}^{t-t^{\alpha}} \int_{\mathbb{R}^{d}} |H_{t,x}(r,z_{1})| \Big( \int_{\mathbb{R}^{d}} |H_{t,x}(r,z_{2})| R(z_{1}-z_{2}) dz_{2} \Big) dz_{1} dr \\ &\lesssim \int_{t^{\alpha}}^{t-t^{\alpha}} r^{-\frac{\kappa}{2}} \int_{\mathbb{R}^{d}} |H_{t,x}(r,z_{1})| dz_{1} dr \lesssim t^{\alpha(1-\frac{\kappa}{2})} . \end{aligned}$$

In the remainder we estimate  $J_{0,t^{\alpha}}(x)$ . For this, we split  $H_{t,x}$  into two terms:

$$|H_{t,x}(r,z)| = H^1_{t,x}(r,z) + H^2_{t,x}(r,z) ,$$

where

$$H_{t,x}^{(1)}(r,z) = |P_{r(t-r)/t}(z + \frac{r}{t}x) - P_r(z)|, \qquad H_{t,x}^{(2)}(r,z) = P_{t-r}(z+x).$$

So that,

$$J_{0,t^{\alpha}}(x) \leqslant \sum_{i,j=1}^{2} \int_{0}^{t^{\alpha}} \int_{\mathbb{R}^{2d}} H_{t,x}^{(i)}(r,z_1) H_{t,x}^{(j)}(r,z_2) R(z_1-z_2) \, dz \, dr = \sum_{i,j=1}^{2} J_{0,t^{\alpha}}^{(i,j)}(x) \, .$$

By (2.14) from Lemma 2.4, we have uniformly in  $z_1$ :

$$\int_{\mathbb{R}^d} H_{t,x}^{(2)}(r,z_2) R(z_1-z_2) \, dz_2 \lesssim (1+t-r)^{-\frac{\kappa}{2}}$$

Hence, recalling  $t > 2^{\frac{1}{1-\alpha}}$ , for any i = 1, 2, we have

$$J_{0,t^{\alpha}}^{(i,2)}(x) \lesssim \int_{0}^{t^{\alpha}} (t-r)^{-\frac{\kappa}{2}} \int_{\mathbb{R}^{d}} H_{t,x}^{(i)}(r,z_{1}) \, dz_{1} \, dr \lesssim (t-t^{\alpha})^{1-\frac{\kappa}{2}} \lesssim t^{1-\frac{\kappa}{2}} \, .$$

We now need to estimate  $J_{0,t^{\alpha}}^{(1,1)}(x)$ . We rewrite

$$H_{t,x}^{(1)}(r,z) = |P_{r(t-r)/t}(z+\frac{r}{t}x) - P_r(z)| \leq \int_0^1 |\partial_\theta P_{r-\theta\frac{r^2}{t}}(z+\theta\frac{r}{t}x)| \, d\theta \, .$$

Letting  $v = r^2/t$  and w = rx/t, with  $\theta \in [0, 1]$  we can compute

$$\begin{aligned} |\partial_{\theta} P_{r-\theta v}(z+\theta w)| &= \Big| \frac{dv}{2(r-\theta v)} + \frac{v|z+\theta w|^2}{2(r-\theta v)^2} - \frac{\langle z, w \rangle + \theta |w|^2}{r-\theta v} \Big| P_{r-\theta v}(z+\theta w) \\ &\lesssim \Big( \frac{v}{r-v} + \frac{v(|z|^2+|w|^2)}{(r-v)^2} + \frac{|z||w|+|w|^2}{r-v} \Big) P_{r-\theta v}(z+\theta w) \,. \end{aligned}$$

Given  $r < t^{\alpha}$  and  $t > 2^{\frac{1}{1-\alpha}}$ , we have

$$\begin{aligned} \frac{v}{r-v} &= \frac{r/t}{1-r/t} \lesssim t^{\alpha-1} , \qquad |w| \leqslant t^{\alpha-1} |x| ,\\ \frac{v}{(r-v)^2} &\leqslant \frac{v}{r^2 - 2rv} \lesssim t^{-1} , \qquad \frac{|w|}{r-v} &= \frac{r|x|/t}{r-r^2/t} \lesssim t^{-1} |x| \end{aligned}$$

Therefore, letting  $\mu = \alpha(\kappa/2 - 1)/2 = \tilde{\mu}/2$ , we have the following bound

$$\begin{split} |\partial_{\theta} P_{r-\theta r^{2}/t}(z+\theta \frac{r}{t}x)|/P_{r-\theta r^{2}/t}(z+\theta \frac{r}{t}x) \\ &\lesssim t^{\alpha-1}+t^{-1}(|z|^{2}+t^{2(\alpha-1)}|x|^{2})+t^{-1}(|x||z|+t^{\alpha-1}|x|^{2}) \\ &\lesssim t^{\alpha-1}+t^{-1}(|z|^{2}+|x||z|+t^{\alpha-1}|x|^{2}) \\ &\lesssim t^{\alpha-1}+(t^{-1-2\mu}+t^{\alpha-2})|x|^{2}+(t^{-1}+t^{-1+2\mu})|z|^{2} \\ &\lesssim t^{\alpha-1}+t^{-1-2\mu}|x|^{2}+t^{-1+2\mu}|z|^{2} \;, \end{split}$$

where we used the fact that  $|x||z| \lesssim t^{-2\mu}|x|^2 + t^{2\mu}|z|^2$  and  $\alpha - 1 \leqslant -2\mu$  when  $\alpha \leqslant 2/\kappa$ . Thus,

$$H_{t,x}^{(1)}(r,z) \lesssim \left(t^{\alpha-1} + t^{-1-2\mu}|x|^2 + t^{-1+2\mu}|z|^2\right) \int_0^1 P_{r-\theta r^2/t}(z+\theta\frac{r}{t}x) \, d\theta \,. \tag{3.16}$$

Uniformly in  $\theta$ , by Lemma 2.4 and since  $\int_{\mathbb{R}^d} |z|^2 P_r(z+x) dz \lesssim r+|x|^2$ , we have

$$\begin{split} \int_{0}^{t^{\alpha}} \int_{\mathbb{R}^{2d}} |z_{1}|^{2} P_{r-\theta r^{2}/t}(z_{1}+\theta\frac{r}{t}x) H_{t,x}^{(1)}(r,z_{2}) R(z_{1}-z_{2}) dr dz \\ &= \int_{0}^{t^{\alpha}} \int_{\mathbb{R}^{d}} |z_{1}|^{2} P_{r-\theta r^{2}/t}(z_{1}+\theta\frac{r}{t}x) \Big( \int_{\mathbb{R}^{d}} H_{t,x}^{(1)}(r,z_{2}) R(z_{1}-z_{2}) dz_{2} \Big) dz_{1} dr \\ &\lesssim \int_{0}^{t^{\alpha}} \left( (r-\theta\frac{r^{2}}{t}) + \theta^{2} \frac{r^{2}}{t^{2}} |x|^{2} \right) ((r(t-r)/t)^{-\frac{\kappa}{2}} + r^{-\frac{\kappa}{2}})) dr \\ &\lesssim \int_{0}^{t^{\alpha}} r^{1-\frac{\kappa}{2}} + r^{2-\frac{\kappa}{2}} t^{-2} |x|^{2} dr \lesssim t^{\alpha(2-\frac{\kappa}{2})} + t^{\alpha(3-\frac{\kappa}{2})-2} |x|^{2} = t^{\alpha-2\mu} + t^{2\alpha-2-2\mu} |x|^{2} \end{split}$$

,

and

$$\int_{0}^{t^{\alpha}} \int_{\mathbb{R}^{2d}} P_{r-\theta r^{2}/t}(z_{1}+\theta \frac{r}{t}x) H_{t,x}^{(1)}(r,z_{2}) R(z_{1}-z_{2}) dr dz \lesssim 1$$

With (3.16) and these estimates, we obtain

$$J^{(1,1)}_{0,t^\alpha}(x) \lesssim t^{\alpha-1} + t^{-1-2\mu} |x|^2 + t^{2\alpha-3} |x|^2 \lesssim t^{\alpha-1} + t^{-1-2\mu} |x|^2 \; .$$

Therefore, since  $\max(1 - \frac{\kappa}{2}, \alpha - 1) \leq -2\mu$ , we obtain

$$J_{0,t^{\alpha}}(x) \lesssim t^{1-\frac{\kappa}{2}} + t^{\alpha-1} + t^{-1-2\mu}|x|^2 \lesssim t^{-2\mu} + t^{-1-2\mu}|x|^2$$

Combining the estimates of  $J_{0,t^{\alpha}}(x)$  and  $J_{t^{\alpha},t-t^{\alpha}}(x)$ , we conclude the proof.

## 4 Proof of the main results

In order to prove Theorem 1.3, we are going to consider general transformations of u (still as in (1.8), i.e. with initial condition equal to 1) and prove the following for the class of functions  $\Phi$  below.

Assumption 4.1. Let  $\Phi \in C^2(\mathbb{R}_+;\mathbb{R})$  be a function satisfying for some q > 0,

$$|\Phi(z)| + |\Phi'(z)| + |\Phi''(z)| \lesssim z^{-q} + z^q , \qquad \forall z \in \mathbb{R}_+ .$$
 (4.1)

**Theorem 4.2.** Let  $d \ge 3$ ,  $\xi$  and  $\Phi$  satisfy Assumptions 1.1 and 4.1 respectively and set

$$u_{\varepsilon}^{\Phi}(t,x) = \varepsilon^{1-\frac{\kappa}{2}} \left( \Phi(u_{\varepsilon}(t,x)) - \mathbb{E}[\Phi(u_{\varepsilon}(t,x))] \right) \, .$$

Then, given any  $p \ge 1$ ,  $\alpha < 1 - \frac{\kappa}{2}$  and  $\sigma < -1 - \frac{\kappa}{2}$ , there exists  $\beta_0^{\Phi}(\alpha, p) > 0$  and  $\nu_{\Phi} \ge 0$  such that, for all  $\beta < \beta_0^{\Phi}$  and any good coupling of the noise, the random processes  $u_{\varepsilon}^{\Phi}$  converge in probability to  $\nu_{\Phi}\mathcal{U}^0$  in  $L^p([0,T], \mathcal{C}^{\alpha}(E))$  as  $\varepsilon \to 0$ , for any T > 0 and compact  $E \subset \mathbb{R}^d$ . Here,  $\mathcal{U}^0$  denotes the solution to the additive stochastic heat equation

$$\partial_t \mathcal{U}^0 = \frac{1}{2} \Delta \mathcal{U}^0 + \beta \xi^0 , \qquad \mathcal{U}^0(0, \cdot) \equiv 0 .$$

The effective variance is given by  $\nu_{\Phi} = \mathbb{E}[Z \Phi'(Z)]$ , where Z denotes a real-valued random variable with the same law as  $\vec{Z}(t, x)$ .

The proof of this theorem will be given in Section 4.3 below. Given a test function  $g : \mathbb{R}^d \to \mathbb{R}$  and  $\Phi : \mathbb{R}_+ \to \mathbb{R}$ , in the following we write

$$Y_t^{\varepsilon,g} = \varepsilon^{1-\frac{\kappa}{2}} \int_{\mathbb{R}^d} (\Phi(u_\varepsilon(t,x)) - \mathbb{E}[\Phi(u_\varepsilon(t,x))])g(x)dx.$$

In order to prove Theorem 4.2, we first derive a divergence representation for  $Y_t^{\varepsilon,g}$  by using the Clark–Ocone formula.

**Lemma 4.3.** Let  $\Phi \in C^2(\mathbb{R}_+)$  satisfy (4.1), then for  $\beta$  sufficiently small,

$$Y_t^{\varepsilon,g} = \varepsilon^{1-\frac{\kappa}{2}} \int_0^{\frac{t}{\varepsilon^2}} \int_{\mathbb{R}^d} \int_{\mathbb{R}^d} \mathbb{E}[\Phi'(u_\varepsilon(t,x))D_{s,y}u_\varepsilon(t,x)|\mathcal{F}_s] g(x)dx\,\xi(ds,dy).$$
(4.2)

Proof. By Lemma 2.6 and Clark–Ocone formula

$$Y_t^{\varepsilon,g} = \int_{\mathbb{R}\times\mathbb{R}^d} \mathbb{E}[D_{s,y}Y_t^{\varepsilon,g}|\mathcal{F}_s]\xi(ds,dy)$$
$$= \varepsilon^{1-\frac{\kappa}{2}} \int_0^{\frac{t}{\varepsilon^2}} \int_{\mathbb{R}^d} \int_{\mathbb{R}^d} \mathbb{E}[D_{s,y}\Phi(u_\varepsilon(t,x))|\mathcal{F}_s]g(x)\,dx\,\xi(ds,dy)\;,$$

and the claim follows from the chain rule.

The strategy of proof of Theorem 4.2 is then as follows. We use the expression for the Malliavin derivative of the solution combined with the homogenisation result to argue that, for  $\tau \gg s$ ,

$$\mathbb{E}[D_{s,y}\Phi(u(\tau,x))|\mathcal{F}_s] = \beta \mathbb{E}[\Phi'(u(\tau,x))u(s,y)w_{s,y}(\tau,x)|\mathcal{F}_s]$$
  
$$\approx \beta \mathbb{E}[\Phi'(\vec{Z}(\tau,x))\vec{Z}(s,y)u(s,y)P_{\tau-s}(x-y)\vec{Z}(\tau,x)|\mathcal{F}_s].$$

At that point, we note that the process  $\vec{Z}$  is expected to mix relatively well and that furthermore for  $|\tau - s| \gg 1$ ,  $\tilde{Z}(s, y)$  is  $\mathcal{F}_s$ -independent, of expectation 1, and roughly independent of  $\vec{Z}(\tau, x)$  so that one gets

$$\mathbb{E}[D_{s,y}\Phi(u(\tau,x))|\mathcal{F}_s] \approx \beta u(s,y)P_{\tau-s}(x-y)\mathbb{E}[\Phi'(\vec{Z}(\tau,x))\vec{Z}(\tau,x)] \\ \stackrel{\text{def}}{=} \beta u(s,y)P_{\tau-s}(x-y)\nu_{\Phi} .$$
(4.3)

The approximations are proved in the next section. With these in place, in Section 4.2 we can show  $Y_t^{\varepsilon,g}$  converges strongly towards  $\mathcal{U}^0$  and conclude the proof of Theorem 4.2. In the final section, we consider the KPZ equation with non-flat initial conditions and prove our main result Theorem 1.3.

## 4.1 Main convergence result

The main result of this section is Proposition 4.5 below, which shows that we can approximate  $Y_t^{\varepsilon,g}$  by  $\nu_{\Phi}$  times the large-scale fluctuations of the linear SHE which, by the mild formulation, are given by

$$X_t^{\varepsilon,g} = \beta \varepsilon^{1-\frac{\kappa}{2}} \int_0^{\frac{t}{\varepsilon^2}} \int_{\mathbb{R}^d} \int_{\mathbb{R}^d} P_{t/\varepsilon^2 - s}(x - y) u(s, y) \varepsilon^d g(\varepsilon x) \, dx \, \xi(ds, dy) \, . \tag{4.4}$$

Then, in Proposition 4.6, we show that as  $\varepsilon \to 0$  these are close to the solution of the stochastic heat equation (1.9) driven by noise  $\xi^{\varepsilon}$ , which can be written as

$$\mathcal{U}_{t}^{\varepsilon}(g) := \beta \varepsilon^{1-\frac{\kappa}{2}} \int_{0}^{\frac{t}{\varepsilon^{2}}} \int_{\mathbb{R}^{d}} \int_{\mathbb{R}^{d}} P_{t/\varepsilon^{2}-s}(x-y) \varepsilon^{d} g(\varepsilon x) \, dx \, \xi(ds, dy) \,. \tag{4.5}$$

Since  $\mathcal{U}^{\varepsilon}$  converges to the limit  $\mathcal{U}^0$ , we conclude that  $Y_t^{\varepsilon,g}$  converges to  $\nu_{\Phi}\mathcal{U}^0$  as in Theorem 4.2. We first write

$$J_{\kappa,\delta}(f) \stackrel{\text{def}}{=} \int_{\mathbb{R}^{4d}} \left( \prod_{i=1}^{2} |f(x_i)| |x_i - y_i|^{\delta - d} \right) |y_1 - y_2|^{-\kappa} \, dy \, dx \,, \qquad (4.6)$$

and we recall the following [GHL23, Lem. 4.10].

**Lemma 4.4.** For any compactly supported bounded function  $f : \mathbb{R}^d \to \mathbb{R}$  and any  $\delta \in (0, \frac{\kappa}{2})$ , one has  $J_{\kappa,\delta}(f) \leq M^{2d+2\delta-\kappa} ||f||_{\infty}^2$ , where M denotes the diameter of the support of f. Furthermore, setting  $f_x^{\lambda}(y) = \lambda^{-d} f(\frac{y-x}{\lambda})$ , one has the scaling relation

$$J_{\kappa,\delta}(f_x^{\lambda}) = \lambda^{2\delta - \kappa} J_{\kappa,\delta}(f) .$$
(4.7)

**Proposition 4.5.** Let  $p \ge 1$ . For  $\beta$  sufficiently small, for any test function  $g \in C_c(\mathbb{R}^d)$  and t > 0, the following holds for any  $\delta_0 \in (0, 1)$ :

$$\|Y_t^{\varepsilon,g} - \nu_{\Phi} X_t^{\varepsilon,g}\|_p \lesssim \varepsilon^{\frac{1}{2}\min(1-\delta_0, 1-\frac{2}{\kappa})}, \qquad (4.8)$$

with precise rate given by (4.13).

*Proof.* Recall we denote with  $u^{(s)}(t, x)$  the solution of SHE for  $t \ge s$  with initial condition 1 at time s. Without loss of generality, assume  $\varepsilon < t$  and let  $\tau = t/\varepsilon^2$ . By Lemma 4.3 and the expression for the Malliavin derivative (2.25),

$$Y_t^{\varepsilon,g} = \beta \varepsilon^{1-\frac{\kappa}{2}} \int_0^\tau \int_{\mathbb{R}^d} \int_{\mathbb{R}^d} \mathbb{E}[\Phi'(u(\tau,x))u(s,y)w_{s,y}(\tau,x)|\mathcal{F}_s] \varepsilon^d g(\varepsilon x) \, dx \, \xi(ds,dy)$$
(4.9)

Given the large times asymptotics, the homogenisation of  $w_{s,y}(\tau, x)$  and mixing of the stationary process, the proof consists in approximating this fluctuations expression through separation of timescales. Namely, given moments control, the contribution coming from  $\tau - s < \varepsilon^{-1}$  is negligeable. On the other hand, we let  $\Psi(u) = \Phi'(u)u$  and for  $\tau - s > \varepsilon^{-1}$  we show

$$\begin{split} \mathbb{E}[\Phi'(u(\tau,x))u(s,y)w_{s,y}(\tau,x)|\mathcal{F}_s] \\ &\approx \mathbb{E}[\Phi'(\vec{Z}(\tau,x))u(s,y)w_{s,y}(\tau,x)|\mathcal{F}_s] \\ &\approx \mathbb{E}[\Phi'(\vec{Z}(\tau,x))u(s,y)P_{\tau-s}(x-y)C_{s,y}(\tau)\vec{Z}(\tau,x)|\mathcal{F}_s] \\ &\approx \mathbb{E}[u(s,y)P_{\tau-s}(x-y)C_{s,y}(\frac{s+\tau}{2})\Psi(u^{(\frac{s+\tau}{2})}(\tau,x))|\mathcal{F}_s] \\ &= P_{\tau-s}(x-y)u(s,y)\mathbb{E}[\Psi(u^{(\frac{s+\tau}{2})}(\tau,x))] \approx P_{\tau-s}(x-y)u(s,y)\nu_{\Phi} . \end{split}$$

Define the indicator functions  $\overline{\chi}_{\tau}^{\varepsilon}(s) = \mathbf{1}_{[\tau-\varepsilon^{-1},\tau]}(s)$  and  $\underline{\chi}_{\tau}^{\varepsilon}(s) = \mathbf{1}_{[0,\tau-\varepsilon^{-1}]}(s)$ . Then, to show (4.8), we write the conditional projection in (4.9) as

$$\mathbb{E}[\Phi'(u(\tau, x))u(s, y)w_{s, y}(\tau, x)|\mathcal{F}_s] - \nu_{\Phi}P_{\tau-s}(x-y)u(s, y) = \sum_{i=1}^5 S_i(\tau, x, s, y) ,$$

where

$$S_1 = \left( \mathbb{E}[\Phi'(u(\tau, x))u(s, y)w_{s,y}(\tau, x)|\mathcal{F}_s] - \nu_{\Phi}P_{\tau-s}(x-y)u(s, y) \right) \underline{\chi}_{\tau}^{\varepsilon}(s)$$

denotes the integrand on  $[0,\tau-\varepsilon^{-1}]$  and, on the remaining interval, we have 4 terms:

$$\begin{split} S_{2} &= \mathbb{E}[(\Phi'(u(\tau, x)) - \Phi'(\vec{Z}(\tau, x))u(s, y)w_{s,y}(\tau, x) | \mathcal{F}_{s}] \,\overline{\chi}_{\tau}^{\varepsilon}(s) ,\\ S_{3} &= \mathbb{E}[\Phi'(\vec{Z}(\tau, x))u(s, y)(w_{s,y}(\tau, x) - P_{\tau-s}(x - y)C_{s,y}(\tau)\vec{Z}(\tau, x)) | \mathcal{F}_{s}] \,\overline{\chi}_{\tau}^{\varepsilon}(s) ,\\ S_{4} &= P_{\tau-s}(x - y)u(s, y) \,\mathbb{E}[C_{s,y}(\tau)\Psi(\vec{Z}(\tau, x)) - C_{s,y}(\frac{s+\tau}{2})\Psi(u^{(\frac{s+\tau}{2})}(\tau, x)) | \mathcal{F}_{s}] \,\overline{\chi}_{\tau}^{\varepsilon}(s) ,\\ S_{5} &= P_{\tau-s}(x - y)u(s, y)(\mathbb{E}[C_{s,y}(\frac{s+\tau}{2})\Psi(u^{(\frac{s+\tau}{2})}(\tau, x)) | \mathcal{F}_{s}] - \nu_{\Phi}) \,\overline{\chi}_{\tau}^{\varepsilon}(s) . \end{split}$$

With this decomposition, we can express the difference as follows:

$$Y_t^{\varepsilon,g} - \nu_{\Phi} X_t^{\varepsilon,g} = \beta \varepsilon^{1-\frac{\kappa}{2}} \sum_{k=1}^5 \mathcal{I}_{k,\varepsilon} ,$$

where

$$\mathcal{I}_{k,\varepsilon} := \int_0^\tau \int_{\mathbb{R}^d} \int_{\mathbb{R}^d} S_k(\tau, x, s, y) \, \varepsilon^d g(\varepsilon x) \, dx \, \xi(ds, dy) \; .$$

Estimating the  $L^p$  norms of  $S_j$ , we show that each term  $\mathcal{I}_{j,\varepsilon}$  in the decomposition converges to 0. Since, by BDG inequality as in (3.10), we have for  $k = 1, \ldots, 5$ ,

$$\|\mathcal{I}_{k,\varepsilon}\|_{p}^{2} \leqslant \int_{0}^{\tau} \int_{\mathbb{R}^{4d}} \left(\prod_{i=1}^{2} \|S_{k}(\tau, x_{i}, s, y_{i})\|_{p} |g_{\varepsilon}(x_{i})|\right) R(y_{1} - y_{2}) \, ds \, dx \, dy \,, \quad (4.10)$$

where  $g_{\varepsilon}(x) = \varepsilon^d g(\varepsilon x)$ .

By the moment estimates from Lemmas 2.5–2.6 (in particular (2.19) to bound  $w_{s,y}$ ), we have  $||S_1(\tau, x, s, y)||_p \leq P_{\tau-s}(x-y)$  provided that  $\beta$  is sufficiently small. For  $\delta \in (0, d)$ , we have (cf. [GHL23, Lem. 2.1])

$$P_s(x) \lesssim s^{-\frac{\delta}{2}} |x|^{\delta - d}$$
 (4.11)

Inserting this bound into (4.10) yields

$$\|\mathcal{I}_{1,\varepsilon}\|_p^2 \lesssim \int_{\tau-\varepsilon^{-1}}^{\tau} \int_{\mathbb{R}^{4d}} \Big(\prod_{i=1}^2 P_{\tau-s}(x_i-y_i)|g_{\varepsilon}(x_i)|\Big) R(y_1-y_2) \, ds \, dx \, dy$$

Recall that  $t = \varepsilon^2 \tau$ . By the change of variables  $x_i \mapsto \varepsilon x_i, y_i \mapsto \varepsilon y_i, s \mapsto \varepsilon^2 s$ , and the fact that  $P_{r/\varepsilon^2}(z/\varepsilon) = \varepsilon^d P_r(z)$ , we have

$$\|\mathcal{I}_{1,\varepsilon}\|_p^2 \lesssim \varepsilon^{\kappa-2} \int_{t-\varepsilon}^t \int_{\mathbb{R}^{4d}} \prod_{i=1}^2 \Big( P_{t-s}(x_i - y_i) |g(x_i)| \Big) \varepsilon^{-\kappa} R(\frac{y_1 - y_2}{\varepsilon}) \, ds \, dx \, dy \, .$$

Since  $\varepsilon^{-\kappa} R(\frac{y_1-y_2}{\varepsilon}) \lesssim |y_1-y_2|^{-\kappa}$ , using (4.11) with  $\delta_0 < 1$ , we estimate

$$\begin{aligned} \|\mathcal{I}_{1,\varepsilon}\|_{p}^{2} &\lesssim \varepsilon^{\kappa-2} \int_{t-\varepsilon}^{t} \int_{\mathbb{R}^{4d}} \left(\prod_{i=1}^{2} P_{t-s}(x_{i}-y_{i}) \left|g(x_{i})\right|\right) |y_{1}-y_{2}|^{-\kappa} \, ds \, dx \, dy \\ &\lesssim \varepsilon^{\kappa-2} \int_{0}^{\varepsilon} s^{-\delta_{0}} \, ds \, \int_{\mathbb{R}^{4d}} \left(\prod_{i=1}^{2} |x_{i}-y_{i}|^{\delta_{0}-d} \left|g(x_{i})\right|\right) |y_{1}-y_{2}|^{-\kappa} \, dx \, dy \\ &\lesssim \varepsilon^{\kappa-2} \varepsilon^{1-\delta_{0}} J_{\kappa,\delta_{0}}(g) \;, \end{aligned}$$

where  $J_{\kappa,\delta}$  is defined as in (4.6).

For  $\mathcal{I}_{2,\varepsilon}$ , first apply Cauchy-Schwartz inequality, then the moment bounds (2.19) and (2.29), we have

$$\begin{split} \|S_2(\tau, x, s, y)\|_p &\lesssim \|\Phi'(u(\tau, x)) - \Phi'(\vec{Z}(\tau, x)\|_{3p} \|w_{s,y}(\tau, x)\|_{3p} \\ &\lesssim (1+\tau)^{\frac{2-\kappa}{4}} P_{\tau-s}(x-y) \lesssim \varepsilon^{\frac{\kappa-2}{4}} P_{\tau-s}(x-y) \,. \end{split}$$

By a change of variables and (4.11) with  $\delta_1 < 1$  as above, from (4.10) we obtain

$$\begin{aligned} \|\mathcal{I}_{2,\varepsilon}\|_{p}^{2} &\lesssim \varepsilon^{\frac{\kappa}{2}-1} \int_{0}^{\tau-\varepsilon^{-1}} \int_{\mathbb{R}^{4d}} \left(\prod_{i=1}^{2} P_{\tau-s}(x_{i}-y_{i}) \left|g_{\varepsilon}(x_{i})\right|\right) R(y_{1}-y_{2}) \, ds \, dx \, dy \\ &\lesssim \varepsilon^{\kappa-2} \, \varepsilon^{\frac{\kappa}{2}-1} \int_{0}^{t-\varepsilon} \int_{\mathbb{R}^{4d}} \left(\prod_{i=1}^{2} P_{t-s}(x_{i}-y_{i}) \left|g(x_{i})\right|\right) |y_{1}-y_{2}|^{-\kappa} \, ds \, dx \, dy \\ &\lesssim \varepsilon^{\kappa-2} \, \varepsilon^{\frac{\kappa}{2}-1} \, t^{1-\delta_{1}} \, J_{\kappa,\delta_{1}}(g) \,. \end{aligned}$$

$$(4.12)$$

For the remaining  $\mathcal{I}_{j,\varepsilon}$ , we shall make use of the fact that  $s < \tau - \varepsilon^{-1}$ . Using the uniform moment bounds on  $\Psi(u) = \Phi'(u)u$  and on u, and the homogenisation result Proposition 3.1, we have

$$\begin{split} \|S_3\|_p &\lesssim P_{\tau-s}(x-y) \, \|\rho_{s,y}(\tau,x)\|_{2p} \\ &\lesssim P_{\tau-s}(x-y) \, (1+\tau-s)^{-\mu} (1+(1+\tau-s)^{-\frac{1}{2}}|x-y|) \\ &\lesssim \varepsilon^{\mu} P_{\tau-s}(x-y) + \varepsilon^{\mu} |x-y| P_{\tau-s}(x-y) \, . \end{split}$$

Here  $\mu = \frac{1}{2} - \frac{1}{\kappa}$  and  $\rho_{s,y}$  is given by (3.1). As for  $\mathcal{I}_{2,\varepsilon}$ , the contribution from  $\varepsilon^{\mu}P_{\tau-s}(x-y)$  can be bounded by  $\varepsilon^{\kappa-2} \varepsilon^{2\mu} t^{1-\delta_1} J_{\kappa,\delta_1}(g)$ . The contribution from  $\varepsilon^{\mu}|x-y|P_{\tau-s}(x-y)$  can be estimated as follows:

$$\begin{split} \varepsilon^{2\mu} \int_0^{\tau-\varepsilon^{-1}} \int_{\mathbb{R}^{4d}} \Big( \prod_{i=1}^2 |x_i - y_i| P_{\tau-s}(x_i - y_i)| g_{\varepsilon}(x_i)| \Big) R(y_1 - y_2) \, ds \, dx \, dy \\ &\lesssim \varepsilon^{2\mu} \varepsilon^{\kappa-2} \int_0^{t-\varepsilon} (t-s)^{-\delta_2} \, ds \, \int_{\mathbb{R}^{4d}} \Big( \prod_{i=1}^2 |x_i - y_i|^{1+\delta_2 - d} \, |g(x_i)| \Big) |y_1 - y_2|^{-\kappa} \, dx \, dy \\ &\lesssim \varepsilon^{2\mu} \varepsilon^{\kappa-2} \, t^{1-\delta_2} \, J_{\kappa, 1+\delta_2}(g) \, . \end{split}$$

We have employed again the heat kernel bound (4.11), with  $\delta_2 < \frac{\kappa}{2} - 1$ . Summing these up, we have:

$$\|\mathcal{I}_{3,\varepsilon}\|_p^2 \lesssim \varepsilon^{2\mu} \varepsilon^{\kappa-2} (t^{1-\delta_1} J_{\kappa,\delta_1}(g) + t^{1-\delta_2} J_{\kappa,1+\delta_2}(g)) .$$

For  $\mathcal{I}_{4,\varepsilon}$  and  $\mathcal{I}_{5,\varepsilon}$ , we make use of mixing, separating  $C_{s,y}$  and  $\vec{Z}$  in independent parts. We first apply (2.29). For  $s < \tau - \varepsilon^{-1}$  and q > 1 we have

$$\|\Psi(\vec{Z}(\tau,x)) - \Psi(u^{(\frac{s+\tau}{2})}(\tau,x))\|_q \lesssim \beta(1 + \frac{1}{2}(\tau-s))^{\frac{2-\kappa}{4}} \lesssim \varepsilon^{\frac{\kappa-2}{4}},$$

On the other hand, by (2.20) we also have

$$\|C_{s,y}(\tau) - C_{s,y}(\frac{s+\tau}{2})\|_q \lesssim \beta (1 + \frac{1}{2}(\tau-s))^{\frac{2-\kappa}{4}} \lesssim \varepsilon^{\frac{\kappa-2}{4}}.$$

Therefore, for sufficiently small  $\beta$ ,

$$\|C_{s,y}(\tau)\Psi(\vec{Z}(\tau,x)) - C_{s,y}(\frac{s+t/\varepsilon^2}{2})\Psi(u^{(\frac{s+\tau}{2})}(\tau,x))\|_{2p} \lesssim \varepsilon^{\frac{\kappa-2}{4}},$$

We have used trinagle inequality and Lemma 2.5. Consequently,

$$||S_4||_p \lesssim \varepsilon^{\frac{\kappa-2}{4}} P_{\tau-s}(x-y) .$$

This is the same estimates as for  $\mathcal{I}_{2,\varepsilon}$ , therefore,  $\|\mathcal{I}_{4,\varepsilon}\|_p^2 \lesssim \varepsilon^{\kappa-2}\varepsilon^{\frac{\kappa}{2}-1}t^{1-\delta_1}J_{\kappa,\delta_1}(g)$ . By the integral expression for  $C_{s,y}$  in (3.6),  $C_{s,y}(\frac{s+\tau}{2})$  is of expectation 1 and independent of  $u^{(\frac{s+\tau}{2})}(\tau, x)$ . In addition, both terms are  $\mathcal{F}_s$ -independent, therefore we have

$$\mathbb{E}[C_{s,y}(\frac{s+\tau}{2})\Psi(u^{(\frac{s+\tau}{2})}(\tau,x))|\mathcal{F}_s] = \mathbb{E}[C_{s,y}(\frac{s+\tau}{2})]\mathbb{E}[\Psi(u^{(\frac{s+\tau}{2})}(\tau,x))]$$
$$= \mathbb{E}[\Psi(u^{(\frac{s+\tau}{2})}(\tau,x)) - \Psi(\vec{Z}(\tau,x))] + \nu_{\Phi}.$$

Using the bounds in (2.29), we obtain

$$||S_5||_p \lesssim P_{\tau-s}(x-y) ||\Psi(u^{(\frac{s+\tau}{2})}(\tau,x)) - \Psi(\vec{Z}(\tau,x))||_{2p} \lesssim P_{\tau-s}(x-y)(1+\frac{1}{2}(\tau-s))^{\frac{2-\kappa}{4}} \lesssim \varepsilon^{\frac{\kappa-2}{4}} P_{\tau-s}(x-y) .$$

This is the same bound for  $S_4$ , hence  $\|\mathcal{I}_{5,\varepsilon}\|_p^2 \lesssim \varepsilon^{\kappa-2}\varepsilon^{\frac{\kappa}{2}-1}t^{1-\delta_1}J_{\kappa,\delta_1}(g)$ . Putting all estimates together, observing that  $\mu = \frac{1}{2} - \frac{1}{\kappa} < \frac{\kappa-2}{4}$ , we obtain

$$\|Y_t^{\varepsilon,g} - \nu_{\Phi} X_t^{\varepsilon,g}\|_p \lesssim \varepsilon^{\frac{1-\delta_0}{2}} J_{\kappa,\delta_0}(g)^{\frac{1}{2}} + \varepsilon^{\mu} \left( t^{\frac{1-\delta_1}{2}} J_{\kappa,\delta_1}(g)^{\frac{1}{2}} + t^{\frac{1-\delta_2}{2}} J_{\kappa,1+\delta_2}(g)^{\frac{1}{2}} \right),$$
(4.13)

where  $\delta_0, \delta_1 \in (0, 1)$  and  $\delta_2 \in (0, \frac{\kappa}{2} - 1)$ . The implicit constants are uniform in  $t, \varepsilon > 0$ . We conclude the proof of (4.8). 

**Proposition 4.6.** Let  $p \ge 1$  and  $\beta$  sufficiently small. The following holds for any  $g \in \mathcal{C}_c(\mathbb{R}^d)$ , t > 0, and  $\theta \in (2, \kappa \land (d+2-\kappa))$ :

$$\|X_t^{\varepsilon,g} - \mathcal{U}_t^{\varepsilon}(g)\|_p \lesssim \varepsilon^{\frac{\theta-2}{2p}} t^{\frac{1-\delta}{2}(1+\frac{1}{p})} J_{\kappa+\theta-2\delta,\delta}(g)^{\frac{1}{2p}} J_{\kappa,\delta}(g)^{\frac{1}{2}(1-\frac{1}{p})} .$$
(4.14)

*Proof.* Recall that  $X_t^{\varepsilon,g}$  is defined in (4.4) and  $\mathcal{U}_t^{\varepsilon}$  is defined in (4.5). In the former, we substitute u(s, y) by  $1 + \int_0^s \int_{\mathbb{R}^d} P_{s-r}(y-z)u(r,z)\xi(dr,dz)$  allowing to recover  $\mathcal{U}_t^{\varepsilon}(g)$  and an additional term:  $X_t^{\varepsilon,g} = \mathcal{U}_t^{\varepsilon}(g) + B_{\varepsilon}(g)$ , where

$$B_{\varepsilon} = \beta^{2} \varepsilon^{1-\frac{\kappa}{2}} \int_{0}^{\tau} \int_{\mathbb{R}^{d}} \int_{0}^{s} \int_{\mathbb{R}^{d}} \int_{\mathbb{R}^{d}} P_{\tau-s}(x-y) \cdot P_{s-r}(y-z)u(r,z) \varepsilon^{d} g(\varepsilon x) dx \,\xi(dr,dz) \,\xi(ds,dy)$$

$$(4.15)$$

and  $\tau = t/\varepsilon^2$ . We first show that  $B_{\varepsilon}$  converges to 0 in  $L^2$ . By Itô's isometry,

$$||B_{\varepsilon}||_{2}^{2} = \beta^{4} \varepsilon^{2-\kappa} \int_{0}^{\tau} \int_{0}^{s} \int_{\mathbb{R}^{6d}} \left( \prod_{i=1}^{2} P_{\tau-s}(x_{i}-y_{i}) P_{s-r}(y_{i}-z_{i}) \right) R(y_{1}-y_{2})$$
$$R(z_{1}-z_{2}) \mathbb{E}[u(r,z_{1})u(r,z_{2})] \varepsilon^{2d} g(\varepsilon x_{1}) g(\varepsilon x_{2}) \, ds \, dr \, dx \, dy \, dz$$

Using the uniform  $L^2$  bound on u and a change of variables as before<sup>(1)</sup>, we obtain

$$||B_{\varepsilon}||_{2}^{2} \lesssim \int_{0}^{t} \int_{0}^{s} \int_{\mathbb{R}^{6d}} \left( \prod_{i=1}^{2} P_{t-s}(x_{i}-y_{i})P_{s-r}(y_{i}-z_{i}) \right) \varepsilon^{-2} R(\frac{y_{1}-y_{2}}{\varepsilon})$$
$$\varepsilon^{-\kappa} R(\frac{z_{1}-z_{2}}{\varepsilon}) |g(x_{1})g(x_{2})| \, ds \, dr \, dx \, dy \, dz \, .$$

Using  $\varepsilon^{-\kappa} R(z/\varepsilon) \lesssim |z|^{-\kappa}$  and the kernel bound (4.11) with  $\delta \in (0, 1)$ , we obtain

$$||B_{\varepsilon}||_{2}^{2} \lesssim \int_{0}^{t} \int_{0}^{s} (t-s)^{-\delta} (s-r)^{-\delta} dr \, ds \, I_{\delta,\kappa}^{\varepsilon}(g) \lesssim t^{2-2\delta} \, I_{\delta,\kappa}^{\varepsilon}(g)$$

where, with  $G(x) = g(x_1)g(x_2)$ ,

$$I_{\delta,\kappa}^{\varepsilon}(g) = \int_{\mathbb{R}^{6d}} \left( \prod_{i=1}^{2} |x_i - y_i|^{\delta-d} |y_i - z_i|^{\delta-d} \right) \varepsilon^{-2} R(\frac{y_1 - y_2}{\varepsilon}) |z_1 - z_2|^{-\kappa} |G(x)| \, dx \, dy \, dz$$

Using the fact that, for  $\beta_1 + \beta_2 + d < 0$  and  $\beta_i < -d$ ,  $\int_{\mathbb{R}^d} |y - z|^{\beta_1} |z - w|^{\beta_2} dz = c_{\beta_1,\beta_2} |y - w|^{\beta_1 + \beta_2 + d}$ , we can integrate out the  $z_i$  variables and obtain

$$I^{\varepsilon}_{\delta,\kappa}(g) = c \int_{\mathbb{R}^{4d}} \left( \prod_{i=1}^{2} |x_i - y_i|^{\delta-d} \right) \varepsilon^{-2} R(\frac{y_1 - y_2}{\varepsilon}) |y_1 - y_2|^{2\delta - \kappa} |G(x)| \, dx \, dy \; .$$

Let us denote

$$K_{\varepsilon}(x) = \varepsilon^{-2} R(x/\varepsilon) |x|^{2\delta - \kappa}$$

We know that  $R(x) \leq |x|^{-\theta}$ , for any  $\theta \in [0, \kappa]$ . Let us pick  $\theta \in (2, \kappa \wedge (d+2-\kappa))$ , so that  $\theta + \kappa - d < 2$  and we fix  $\delta \in (\frac{1}{2}(\theta + \kappa - d), 1)$ . Then, uniformly in  $\varepsilon \in (0, 1)$ ,

$$K_{\varepsilon}(x) \lesssim \varepsilon^{\theta-2} |x|^{2\delta-\kappa-\theta}$$
 (4.16)

<sup>&</sup>lt;sup>(1)</sup>Namely,  $x_i \mapsto \varepsilon x_i, y_i \mapsto \varepsilon y_i, z_i \mapsto \varepsilon z_i, s \mapsto \varepsilon^2 s, r \mapsto \varepsilon^2 r$ , combined with  $P_{r/\varepsilon^2}(z/\varepsilon) = \varepsilon^d P_r(z)$ .

Note that, by the choice of  $\theta$  and  $\delta$ ,  $K_{\varepsilon}(x)$  is integrable at 0 and decays fast enough at infinity to integrate it over the other kernels. In fact, by (4.16) we obtain

$$I_{\delta,\kappa}^{\varepsilon}(g) \lesssim \varepsilon^{\theta-2} \int_{\mathbb{R}^{4d}} \left( \prod_{i=1}^{2} |x_i - y_i|^{\delta-d} \right) |y_1 - y_2|^{2\delta - \kappa - \theta} |G(x)| \, dx \, dy$$
$$= \varepsilon^{\theta-2} J_{\kappa+\theta-2\delta,\delta}(g) \,,$$

where J is defined as in (4.6) and bounded by Lemma 4.4, as long as  $\delta < \frac{1}{2}(\kappa + \theta - 2\delta)$ . This is the case, given that  $\delta < 1$  (thus  $\delta < \frac{1}{4}(\kappa + \theta)$  as required). Then,

$$|B_{\varepsilon}||_2 \lesssim \varepsilon^{\frac{\theta-2}{2}} t^{1-\delta} J_{\kappa+\theta-2\delta,\delta}(g)^{\frac{1}{2}}$$
.

On the other hand, estimating the  $L^p$  norms of  $X^{\varepsilon}$  and  $\mathcal{U}^{\varepsilon}$  by BDG and a change of variables as in (4.12) in the previous proof, we have

$$\begin{split} \|B_{\varepsilon}\|_{p} &\lesssim \|X_{t}^{\varepsilon,g}\|_{p} + \|\mathcal{U}_{t}^{\varepsilon}(g)\|_{p} \\ &\lesssim \varepsilon^{1-\frac{\kappa}{2}} \Big(\int_{0}^{\tau} \int_{\mathbb{R}^{4d}} \Big(\prod_{i=1}^{t} P_{\tau-s}(x_{i}-y_{i})\varepsilon^{d}g(\varepsilon x_{i})\Big) R(y_{1}-y_{2}) \, dx \, dy \, ds\Big)^{\frac{1}{2}} \\ &\lesssim \Big(\int_{0}^{t} \int_{\mathbb{R}^{4d}} \Big(\prod_{i=1}^{2} P_{t-s}(x_{i}-y_{i})|g(x_{i})|\Big)|y_{1}-y_{2}|^{-\kappa} \, ds \, dx \, dy\Big)^{\frac{1}{2}} \\ &\lesssim t^{\frac{1-\delta}{2}} J_{\kappa,\delta}(g)^{\frac{1}{2}} \,, \end{split}$$

By interpolation, for any  $\theta \in (2, \kappa \land (d+2-\kappa))$  and  $\delta \in (\frac{1}{2}(\theta + \kappa - d), 1)$ ,

$$\|X_{t}^{\varepsilon,g} - \mathcal{U}_{t}^{\varepsilon}(g)\|_{p} = \|B_{\varepsilon}\|_{p} \leqslant \|B_{\varepsilon}\|_{2}^{\frac{1}{p}} \|B_{\varepsilon}\|_{2(p-1)}^{1-\frac{1}{p}} \lesssim \varepsilon^{\frac{\theta-2}{2p}} t^{\frac{1-\delta}{2}(1+\frac{1}{p})} J_{\kappa+\theta-2\delta,\delta}(g)^{\frac{1}{2p}} J_{\kappa,\delta}(g)^{\frac{1}{2}(1-\frac{1}{p})},$$
(4.17)

with implicit constants uniform in  $t, \varepsilon > 0$ , concluding the proof.

# 4.2 Convergence in Hölder spaces

Before going into the proof of Theorem 4.2, we introduce the spaces of locally Hölder continuous distributions  $C^{\alpha}(\mathbb{R}^d)$ , of negative exponent, and prove moment estimates in these spaces. For a function  $g \in C(\mathbb{R}^d)$ ,  $z \in \mathbb{R}^d$  and  $\lambda > 0$ , we denote

$$g_x^{\lambda}(y) := \lambda^{-d} g(\lambda^{-1}(y-x)) ,$$

where we drop the corresponding index if  $\lambda = 1$  or x = 0. We denote with  $\mathcal{D}$  the space of smooth functions with compact support, and  $\mathcal{D}'$  the dual space of distributions. Given  $\alpha < 0$ , let  $r = \lceil -\alpha \rceil$  be the smallest positive integer bigger than  $-\alpha$  and consider the following subset of  $\mathcal{C}^r(\mathbb{R}^d)$ :

$$\mathcal{B}_r := \{g \in \mathcal{D} : \operatorname{supp}(g) \subset B_1, \|g\|_{\mathcal{C}^r} \leq 1\},\$$

where  $B_a$  denotes the open ball of radius *a* centred at 0.

We say that a distribution  $\zeta$  belongs to  $\mathcal{C}^{\alpha}(\mathbb{R}^d)$  if for every compact set  $E \subset \mathbb{R}^d$ ,

$$\|\zeta\|_{\alpha;E} \stackrel{\text{def}}{=} \sup_{g \in \mathcal{B}_r} \sup_{\lambda \in (0,1)} \sup_{x \in E} |\lambda^{-\alpha}| \langle \zeta, g_x^{\lambda} \rangle| < \infty .$$
(4.18)

See [Hai14, Defn. 3.7]. The space  $C^{\alpha}$  is metrisable, with

$$d_{\alpha}(\zeta,\zeta') \stackrel{\text{def}}{=} \sum_{m \ge 1} 2^{-m} (1 \land \|\zeta - \zeta'\|_{\alpha;B_m}) , \qquad (4.19)$$

making it a Fréchet space. For  $\alpha' > \alpha$ ,  $C^{\alpha'}$  is compactly embedded in  $C^{\alpha}$ .

The spaces of  $C^{\alpha}$  can be characterised in multiple ways, here we use the characterisation from [CZ20] in terms of a single test function. We fix an arbitrary reference test function  $\varphi \in D$  such that  $\operatorname{supp} \varphi \subset B_1$  and  $\int \varphi = 1$  for the remainder of the article. We shall use the following notation for its rescaled version:

$$\varphi^{(n)} = 2^{nd}\varphi(2^n \cdot) , \qquad \varphi^{(n)}_x = 2^{nd}\varphi(2^n(\cdot - x))$$

Although  $\varphi$  is fixed, the following shows that control on  $\langle \zeta, \varphi_x^{(k)} \rangle$  allows to get a similar bound for all test functions and control (4.18).

**Theorem 4.7** ([CZ20, Theorem 12.4]). Let  $\zeta \in D'$  be a distribution and let  $\alpha \in (-\infty, 0]$ . Then, one has  $\zeta \in C^{\alpha}$  if and only if, for any compact set E and uniformly over  $k \in \mathbb{N}$ ,

$$\sup_{x \in E} |\langle \zeta, \varphi_x^{(k)} \rangle| \lesssim 2^{-kc}$$

Furthermore, the semi-norms of  $\zeta$  can be estimated by

$$\|\zeta\|_{\alpha;E} \leqslant C \sup_{x \in E_2} \sup_{k \in \mathbb{N}} 2^{k\alpha} |\langle \zeta, \varphi_x^{(k)} \rangle|$$
(4.20)

where  $C = C(\varphi, \alpha, d)$  is an explicit constant and  $E_2 = \{z : d(z, E) \leq 2\}$ .

This characterisation of  $C^{\alpha}$  spaces allows for a criterion to estimate norms in Hölder spaces. We recall the following definition from [GHL23, Defn. 4.6].

**Definition 4.8.** A random distribution  $\zeta$  is said to belong to  $C_p^{\alpha_0}$  if there exists C > 0 such that the following estimates hold for any  $n \ge 1$ :

$$\sup_{x \in \mathbb{R}^d} \|\langle \zeta, \varphi_x^{(n)} \rangle\|_p \leqslant C 2^{-n\alpha_0}, \tag{4.21}$$

$$\sup_{\substack{x,y \in \mathbb{R}^d, \\ 0 < |x-y| \le 2^{-n}}} \|\langle \zeta, \varphi_x^{(n)} - \varphi_y^{(n)} \rangle\|_p \leqslant C 2^{-n(\alpha_0 - 1)} |x-y| .$$
(4.22)

The smallest such constant C is denoted by  $\|\zeta\|_{\mathcal{C}_p^{\alpha_0}}$ . We write  $\mathcal{C}_{p,T}^{\gamma_0,\alpha_0}$ , as a shorthand, for the space  $\mathcal{C}^{\gamma_0}([0,T], \mathcal{C}_p^{\alpha_0})$ .

Given any compact  $E \subset \mathbb{R}^d$ , we denote by  $\mathcal{C}^{\alpha}(E)$  the space of distributions such that the seminorm in (4.18) is finite. Then, by [GHL23, Prop. 4.8] we have the following:

**Proposition 4.9.** Let  $\alpha_0 < 0$ , p > d, and  $\gamma_0 > \frac{1}{p}$ . Then, for any  $\gamma \in (0, \gamma_0 - \frac{1}{p})$ ,  $\alpha < \alpha_0 - \frac{d}{p}$  and any compact  $E \subset \mathbb{R}^d$ , one has the continuous embeddings

$$\mathcal{C}_p^{\alpha_0} \subset L^p(\Omega, \mathcal{C}^{\alpha}(E)), \qquad \mathcal{C}_{p,T}^{\gamma_0, \alpha_0} \subset L^p(\Omega, \mathcal{C}^{\gamma}([0,T], \mathcal{C}^{\alpha}(E))).$$

*Remark* 4.10. In [GHL23, Prop. 4.8], these inclusions are stated for  $E = \mathbb{R}^d$ , but the proof given there allows to replace  $\mathbb{R}^d$  by E.

We have the following:

**Lemma 4.11.** Given any  $\alpha_0 < 1 - \frac{\kappa}{2}$ ,  $p \ge 1$ , for sufficiently small  $\beta$ , there exist  $a = a(\alpha_0, p, \kappa) > 0$ , such that the following holds for any T > 0:

$$\sup_{0 < t \leq T} \|Y_t^{\varepsilon} - \nu_{\Phi} \mathcal{U}_t^{\varepsilon}\|_{\mathcal{C}_p^{\alpha_0}} \lesssim \varepsilon^a .$$
(4.23)

In particular, for every  $\alpha < \alpha_0 - \frac{d}{p}$ , there exists a > 0 such that  $\|\sup_{0 < t \leq T} \|Y_t^{\varepsilon} - \nu_{\Phi} \mathcal{U}_t^{\varepsilon}\|_{\alpha, E}\|_p \lesssim \varepsilon^a$ , for any compact  $E \subset \mathbb{R}^d$ . Furthermore, for any  $\gamma_0 \in (0, \frac{1}{2})$ ,  $\alpha_0 < 1 - \frac{\kappa}{2} - 2\gamma_0$ , and  $p \ge 1$ , one has  $\sup_{\varepsilon \in (0,1]} \|\mathcal{U}^{\varepsilon}\|_{\mathcal{C}_p^{\gamma_0, \alpha_0}} < \infty$ .

*Proof.* By (4.13), for  $\delta_0, \delta_1 \in (0, 1), \delta_2 \in (0, \frac{\kappa}{2} - 1)$ , and  $\mu = \frac{1}{2} - \frac{1}{\kappa}$ , one has

$$\sup_{t \in [0,T]} \|Y_t^{\varepsilon,g} - \nu_{\Phi} X_t^{\varepsilon,g}\|_p \lesssim \varepsilon^{\frac{1-\delta_0}{2}} J_{\kappa,\delta_0}(g)^{\frac{1}{2}} + \varepsilon^{\mu} \Big( J_{\kappa,\delta_1}(g)^{\frac{1}{2}} + J_{\kappa,1+\delta_2}(g)^{\frac{1}{2}} \Big).$$

Recall, Lemma 4.4,  $J_{\kappa,\delta}(g) \leq M^{2d+2\delta-\kappa} ||g||_{\infty}^2$ , where *M* is the diameter of the support of *g*, and the scaling property (4.7).

Let  $g \in \overline{\mathcal{D}}$  and  $\delta \in (0, 1)$ . Then the following holds uniformly in  $\lambda \in (0, 1)$ :

$$\sup_{t>0} \sup_{x\in\mathbb{R}^d} \|Y_t^{\varepsilon,g_x^{\lambda}} - \nu_{\Phi} X_t^{\varepsilon,g_x^{\lambda}}\|_p \lesssim \varepsilon^{(\frac{1-\delta}{2})\wedge\mu} \lambda^{\delta-\frac{\kappa}{2}} (J_{\kappa,\delta}(g)^{\frac{1}{2}} + J_{\kappa,1+\delta_2}(g)^{\frac{1}{2}})$$
$$\lesssim \varepsilon^{(\frac{1-\delta}{2})\wedge\mu} \lambda^{\delta-\frac{\kappa}{2}} \|g\|_{\infty} .$$

Hence, for any  $\alpha_0 < 1 - \frac{\kappa}{2}$ , by picking<sup>(2)</sup>  $\delta = \delta(\alpha_0)$  sufficiently close to 1 and applying this to the test function  $\varphi_x^{(n)}$ , we have

$$\sup_{t \in (0,T]} \sup_{n \ge 1} \sup_{x \in \mathbb{R}^d} 2^{n\alpha_0} \| \langle Y_t^{\varepsilon} - \nu_{\Phi} X_t^{\varepsilon}, \varphi_x^{(n)} \rangle \|_p \lesssim \varepsilon^{(\frac{1-\sigma}{2}) \wedge \mu}$$

which implies (4.21). Since the bound holds for any  $g \in \overline{D}$ , (4.22) is also satisfied (cf. [GHL23, Rem. 4.7]). We obtain

$$\sup_{\varepsilon \in (0,T]} \|Y_t^{\varepsilon} - \nu_{\Phi} X_t^{\varepsilon}\|_{\mathcal{C}_p^{\alpha_0}} \lesssim \varepsilon^{(\frac{1-\sigma}{2}) \wedge \mu}$$

It remains to control  $||X_t^{\varepsilon} - U_t^{\varepsilon}||_{\mathcal{C}_p^{\alpha_0}}$ . Following the same steps as above, for any  $\theta \in (2, \kappa \land (d+2-\kappa))$  and  $\delta \in (\frac{1}{2}(\theta + \kappa - d), 1)$ , by Proposition 4.6 and the scaling relation (4.7), we have uniformly in  $\lambda \in (0, 1)$ :

<sup>&</sup>lt;sup>(2)</sup>If  $\alpha_0 > -\frac{\kappa}{2}$ , can set  $\delta = \alpha_0 + \frac{\kappa}{2}$ .

 $\sup_{x \in \mathbb{R}^d} \sup_{t \in (0,T]} \|X_t^{\varepsilon,g_x^{\lambda}} - \mathcal{U}_t^{\varepsilon}(g_x^{\lambda})\|_p \lesssim \varepsilon^{\frac{\theta-2}{2p}} T^{\frac{1-\delta}{2}(1+\frac{1}{p})} \lambda^{(4\delta-\kappa-\theta)\frac{1}{2p}} \lambda^{(2\delta-\kappa)\frac{p-1}{2p}} \|g\|_{\infty}$  $\lesssim \varepsilon^{\frac{\theta-2}{2p}} \lambda^{(\delta-\frac{\kappa}{2})(1+\frac{1}{p})} \|g\|_{\infty} ,$ 

where we used the fact that  $\theta < \kappa$ .

For any  $\alpha_0 < 1 - \frac{\kappa}{2}$ , picking sufficiently high  $p(\alpha_0) \ge 1$  and  $\delta(\alpha_0) < 1$ , we have  $(\delta - \frac{\kappa}{2})(1 + \frac{1}{p}) > \alpha_0$ . Then, by the same reasoning as above, we obtain

$$\sup_{t \in (0,T]} \|X_t^{\varepsilon} - \mathcal{U}_t^{\varepsilon}\|_{\mathcal{C}_p^{\alpha_0}} \lesssim \varepsilon^{\frac{\theta-2}{2p}}$$

Letting  $a = \frac{1}{2}\min(1-\delta, 1-\frac{2}{\kappa}, \frac{\theta-2}{p})$ , by triangle inequality we conclude that (4.23) holds. The final claim follows by Proposition 4.9, as long as  $\alpha < \alpha_0 - \frac{d}{p}$ .

Regarding the bound on  $\mathcal{U}_t^{\varepsilon}$ , we note that the definition of  $\xi^{\varepsilon}$  implies the scaling relation

$$\mathcal{U}_t^{\varepsilon}(g_x^{\lambda}) = \varepsilon^{1-\frac{\kappa}{2}} \mathcal{U}_{t/\varepsilon^2}(g_{x/\varepsilon}^{\lambda/\varepsilon}) \,,$$

so that the required bound follows immediately from [GHL23, Lem. 4.14].

#### **4.3 Proof of the main results**

**Lemma 4.12.** Consider a good coupling for the  $\xi^{\varepsilon}$  satisfying Assumption 1.1. Given a test function  $\psi \in C_c^{\infty}(\mathbb{R}^{d+1})$  and  $t \in \mathbb{R}$ , write  $\tilde{\psi}_t(s, x) = \psi(s, x)\mathbf{1}_{s \geq t}$ . Then, one has  $\lim_{\varepsilon \to 0} \xi^{\varepsilon}(\tilde{\psi}_t) = \xi^0(\tilde{\psi}_t)$  in probability.

*Proof.* Let  $\chi: \mathbb{R} \to \mathbb{R}_+$  be a smooth increasing function such that  $\chi(s) = 0$  for  $s \leq 0$  and  $\chi(s) = 1$  for  $s \geq 1$ , and set  $\chi_{\delta}(t) = \chi(t/\delta)$ . Setting  $\tilde{\psi}_t^{\delta}(s, x) = \chi_{\delta}(s-t)\psi(s, x)$ , we then note that, as a consequence of Itô's isometry, one can find a constant C (depending on  $\psi$ ) such that

$$\mathbb{E}\xi^{\varepsilon}(\tilde{\psi}_t^{\delta} - \tilde{\psi}_t)^2 \leqslant C\delta , \qquad (4.24)$$

uniformly over all  $\varepsilon, \delta \leq 1$ . Since  $\tilde{\psi}_t^{\delta}$  is a smooth function, one furthermore has  $\xi^{\varepsilon}(\tilde{\psi}_t^{\delta}) \to \xi^0(\tilde{\psi}_t^{\delta})$  in probability as  $\varepsilon \to 0$  for any fixed  $\delta$ . Combining this with (4.24), the claim follows at once.

With the Lemmas above we are now in the position to prove Theorem 4.2. Write  $L^p([0,T], C^{\alpha}(E))$  for the space with seminorms,

$$\|\zeta\|_{\alpha,E,p,T} := \left(\int_0^T \|\zeta_t\|_{\alpha,E}^p \, dt\right)^{\frac{1}{p}},$$

where the semi-norms of  $C^{\alpha}(E)$  is as in Sec. 4.2. We see that for any  $\gamma > 0$ , the space  $C^{\gamma}([0,T], C^{\alpha}(E))$  is continuously embedded in  $L^{p}([0,T], C^{\alpha}(E))$ .

*Proof of Theorem 4.2.* From Lemma 4.11, given  $\alpha < 1 - \frac{\kappa}{2}$ , we pick  $p = p(\alpha)$  sufficiently high such that, as  $\varepsilon \to 0$ ,

$$\left\|\sup_{t\in[0,T]} \|Y_t^{\varepsilon} - \nu_{\Phi}\mathcal{U}_t^{\varepsilon}\|_{\alpha,E}\right\|_p \longrightarrow 0.$$

This implies that  $Y^{\varepsilon} - \nu_{\Phi} \mathcal{U}^{\varepsilon}$  converges in probability to zero in  $L^{p}([0,T], \mathcal{C}^{\alpha}(E))$ .

It therefore remains to show that, for any good coupling, one has  $\mathcal{U}^{\varepsilon} \Rightarrow \mathcal{U}^{0}$ in probability in  $L^{p}([0,T], \mathcal{C}^{\alpha}(E))$ . For this, we make use of [HL22, Prop. 3.12] which implies that it suffices to show tightness of the laws of  $\mathcal{U}^{\varepsilon}$  and convergence in probability of  $\mathcal{U}^{\varepsilon}(\psi)$  for any test function  $\psi \in \mathcal{C}^{\infty}_{c}(\mathbb{R}^{d+1})$ .

By Lemma 4.11 and Proposition 4.9, we obtain for any  $\gamma' \in (0, \frac{1}{2})$  and any  $\alpha' < 1 - \frac{\kappa}{2} - 2\gamma'$ ,

$$\sup_{\varepsilon \in (0,1]} \left\| \left\| \mathcal{U}^{\varepsilon} \right\|_{\mathcal{C}^{\gamma'}([0,T],\mathcal{C}^{\alpha'}(E))} \right\|_p \lesssim 1 \; .$$

Since  $C^{\gamma'}([0,T], C^{\alpha'}(E))$  is compactly embedded in  $C^{\gamma}([0,T], C^{\alpha}(E))$  for any  $\gamma \in (0, \gamma')$  and  $\alpha < \alpha'$ , we conclude that the required tightness holds.

To get convergence of  $\mathcal{U}^{\varepsilon}(\psi)$ , we note that  $\mathcal{U}^{\varepsilon}(\psi) = \xi^{\varepsilon}(P\psi)$  with

$$(P\psi)(s,y) = \beta \mathbf{1}_{s\geq 0} \int_0^\infty \int_{\mathbb{R}^d} \psi(t,x) P_{t-s}(x-y) \, dx \, dt \; ,$$

so that the desired convergence follows from Lemma 4.12.

We now have all the ingredients in place to prove the main result of this article, namely the identification of the large-scale limit of the KPZ equation given in Theorem 1.3.

*Proof of Theorem 1.3.* Given the initial condition  $h_0$ , let

$$\psi_{\varepsilon}(y) = \exp(\varepsilon^{\frac{\kappa}{2}-1}h_0(y)) - 1$$

and denote by  $w_{0,z}^{\varepsilon}(t,x) = \varepsilon^{-d} w_{0,z/\varepsilon}(t/\varepsilon^2, x/\varepsilon)$  the rescaled fundamental solution of the linear stochastic heat equation. Since the initial condition for  $u_{\varepsilon}$  is 1, it follows from the Cole–Hopf transform that

$$h_{\varepsilon}(t,x) = \varepsilon^{1-\frac{\kappa}{2}} \log \left( u_{\varepsilon}(t,x) + \int_{\mathbb{R}^d} w_{0,y}^{\varepsilon}(t,x) \psi_{\varepsilon}(y) \, dy \right)$$

Since  $2|\log(x+y) - \log(x) - y/x| \leq (y/x)^2$  for x, y > 0, it follows that

$$\begin{aligned} \left| h_{\varepsilon}(t,x) - \varepsilon^{1-\frac{\kappa}{2}} \log \left( u_{\varepsilon}(t,x) \right) - \varepsilon^{1-\frac{\kappa}{2}} (u_{\varepsilon}(t,x))^{-1} \int_{\mathbb{R}^d} w_{0,y}^{\varepsilon}(t,x) \psi_{\varepsilon}(y) \, dy \right| \\ &\leqslant \frac{1}{2} \varepsilon^{1-\frac{\kappa}{2}} \Big( (u_{\varepsilon}(t,x))^{-1} \int_{\mathbb{R}^d} w_{0,y}^{\varepsilon}(t,x) \psi_{\varepsilon}(y) \, dy \Big)^2 \,. \end{aligned}$$

By the a priori bounds for  $u_{\varepsilon}(t, x)$ )<sup>-1</sup> from Lemma 2.5 and (2.19) and since  $\|\psi_{\varepsilon}\|_{\infty} \lesssim \varepsilon^{\frac{\kappa}{2}-1}$ , we obtain for any  $p \ge 1$ 

$$\left\| (u_{\varepsilon}(t,x))^{-1} \int_{\mathbb{R}^d} w_{0,y}^{\varepsilon}(t,x) \psi_{\varepsilon}(y) \, dy \right\|_p \lesssim \varepsilon^{\frac{\kappa}{2}-1} \int_{\mathbb{R}^d} P_t(x-y) \, dy = \varepsilon^{\frac{\kappa}{2}-1} \, .$$

On the other hand, by Taylor approximating  $\psi_{\varepsilon}(y)$ , we have  $\|\psi_{\varepsilon}(y) - \varepsilon^{\frac{\kappa}{2}-1}h_0\|_{\infty} \lesssim \varepsilon^{\kappa-2}$  and consequently,

$$\left|\varepsilon^{1-\frac{\kappa}{2}}(u_{\varepsilon}(t,x))^{-1}\left(\int_{\mathbb{R}^d} w_{0,y}^{\varepsilon}(t,x)\psi_{\varepsilon}(y)\,dy - \int_{\mathbb{R}^d} w_{0,y}^{\varepsilon}(t,x)h_0(y)\,dy\right)\right| \lesssim \varepsilon^{\frac{\kappa}{2}-1},$$

Combining these bounds, we conclude that

$$\left\|h_{\varepsilon}(t,x) - \varepsilon^{1-\frac{\kappa}{2}} \log\left(u_{\varepsilon}(t,x)\right) - (u_{\varepsilon}(t,x))^{-1} \int_{\mathbb{R}^d} w_{0,y}^{\varepsilon}(t,x) h_0(y) \, dy\right\|_p \lesssim \varepsilon^{\frac{\kappa}{2}-1}$$

As a consequence of Theorem 1.5, with  $\mu = \frac{1}{2} - \frac{1}{\kappa}$ , for any  $p \ge 1$  we have

$$\begin{split} \|w_{0,y}^{\varepsilon}(t,x) - \vec{Z}(t/\varepsilon^{2},x/\varepsilon)\vec{Z}(0,y/\varepsilon)\varepsilon^{-d}P_{t/\varepsilon^{2}}(\frac{x-y}{\varepsilon})\|_{p} \\ &\lesssim (t/\varepsilon^{2})^{-\mu}(1+(t/\varepsilon^{2})^{-\frac{1}{2}}\varepsilon^{-1}|x-y|)\varepsilon^{-d}P_{t/\varepsilon^{2}}(\frac{x-y}{\varepsilon}) \\ &\lesssim \varepsilon^{2\mu}t^{-\mu}(1+t^{-\frac{1}{2}}|x-y|)P_{t}(x-y) \;, \end{split}$$

where we used the fact that  $\varepsilon^{-d}P_{t/\varepsilon^2}(\frac{x}{\varepsilon}) = P_t(x)$ .

Given that  $h_0$  is bounded and the uniform control of negative moments of  $u_{\varepsilon}$  from Lemma 2.5(i), we obtain

$$\begin{split} \left\| \int_{\mathbb{R}^d} (u_{\varepsilon}(t,x))^{-1} \Big( w_{0,y}^{\varepsilon}(t,x) - \vec{Z}(t/\varepsilon^2, x/\varepsilon) \vec{Z}(0,y/\varepsilon) \varepsilon^{-d} P_{t/\varepsilon^2}(\frac{x-y}{\varepsilon}) \Big) h_0(y) \, dy \right\|_p \\ \lesssim \varepsilon^{2\mu} t^{-\mu} \, \|u_{\varepsilon}^{-1}(t,x)\|_{2p} \int_{\mathbb{R}^d} (1 + t^{-\frac{1}{2}} |x-y|) P_t(x-y) \, dy \lesssim \varepsilon^{2\mu} t^{-\mu} \, . \end{split}$$

On the other hand, by Theorem 2.1 we also have

$$\left\|\frac{\vec{Z}(t/\varepsilon^2, x/\varepsilon)}{u_{\varepsilon}(t, x)} - 1\right\|_p \lesssim \|u_{\varepsilon}^{-1}(t, x)\|_{2p} \|\vec{Z}(t/\varepsilon^2, x/\varepsilon) - u_{\varepsilon}(t, x)\|_{2p} \lesssim t^{\frac{2-\kappa}{4}} \varepsilon^{\frac{\kappa}{2}-1}.$$

Combining the estimates, we obtain

$$\left\|\int_{\mathbb{R}^d} \left(\frac{w_{0,y}^{\varepsilon}(t,x)}{u_{\varepsilon}(t,x)} - \check{Z}(0,y/\varepsilon)P_t(x-y)\right)h_0(y)\,dy\right\|_p \lesssim \varepsilon^{\frac{\kappa}{2}-1}t^{\frac{2-\kappa}{4}} + \varepsilon^{2\mu}t^{-\mu}\,.$$

Hence, noting that  $2\mu < \frac{\kappa}{2} - 1$  and using the fact that  $t \ge T_0 > 0$ , we have

$$\left\|h_{\varepsilon}(t,x) - \varepsilon^{1-\frac{\kappa}{2}} \log\left(u_{\varepsilon}(t,x)\right) - \int_{\mathbb{R}^d} \bar{Z}(0,y/\varepsilon) P_t(x-y) h_0(y) \, dy\right\|_p \lesssim \varepsilon^{2\mu} \, . \tag{4.25}$$

Let  $c = \mathbb{E}[\log \vec{Z}(t, x)]$ ,  $F(t) = \mathbb{E}[\log (u(t, x))]$  and set  $F_{\varepsilon}(t) = F(t/\varepsilon^2)$ . Then, we decompose  $h_{\varepsilon}$  as follows:

$$h_{\varepsilon}(t,x) - \varepsilon^{1-\frac{\kappa}{2}}c = u_{\varepsilon}^{\log}(t,x) + \int_{\mathbb{R}^d} P_t(x-y)h_0(y)\,dy + R_{0,\varepsilon}(t) + \sum_{i=1}^2 R_{i,\varepsilon}(t,x)\,,$$
(4.26)

where

$$\begin{split} u_{\varepsilon}^{\log}(t,x) &= \varepsilon^{1-\frac{\kappa}{2}} (\log \left( u_{\varepsilon}(t,x) \right) - \mathbb{E}[\log \left( u_{\varepsilon}(t,x) \right)]) , \quad R_{0,\varepsilon}(t) = \varepsilon^{1-\frac{\kappa}{2}} (F_{\varepsilon}(t) - c) ,\\ R_{1,\varepsilon}(t,x) &= h_{\varepsilon}(t,x) - \varepsilon^{1-\frac{\kappa}{2}} \log \left( u_{\varepsilon}(t,x) \right) - \int_{\mathbb{R}^d} \breve{Z}(0,y/\varepsilon) P_t(x-y) h_0(y) \, dy ,\\ R_{2,\varepsilon}(t,x) &= \int_{\mathbb{R}^d} (\breve{Z}(0,y/\varepsilon) - 1) P_t(x-y) h_0(y) \, dy . \end{split}$$

By Theorem 4.2 with  $\Phi = \log$ , we know that  $\nu_{\Phi} = 1$  and  $u_{\varepsilon}^{\log}$  converge in probability to  $\mathcal{U}^0$  in  $L^p([0,T], \mathcal{C}^{\alpha}(E))$  for a good coupling of  $\xi^{\varepsilon}$ , where  $\mathcal{U}^0$  is the solution of the stochastic heat equation (1.9) with zero initial conditions. Hence, the first two terms in (4.26) converge in law to the claimed limit  $\mathcal{U}$ , the solution of (1.6) with initial conditions  $h_0$ . It remains to show that the remainder terms  $R_{i,\varepsilon}$  vanish in  $L^p([T_0,T], \mathcal{C}^{\alpha}(E))$ .

Firstly, we show that for t > 0:

$$|F(t) - c| \lesssim 1 \wedge |t|^{(2-\kappa)/2}$$

Then, this implies  $|R_{0,\varepsilon}(t)| \leq \varepsilon^{\frac{\kappa}{2}-1}t^{1-\frac{\kappa}{2}}$ , which vanishes uniformly in  $t \geq T_0$  as  $\varepsilon \to 0$ . Recall that

$$F(t) - c = \mathbb{E}(\log u(t, x) - \log \vec{Z}(t, x)) .$$

It follows that, by splitting into the u < Z and  $u \ge Z$  case,

$$\left|F(t) - c - \mathbb{E}\left(\frac{u(t,x) - \vec{Z}(t,x)}{u(t,x)}\right)\right| \lesssim \mathbb{E}\left(\frac{u(t,x) - \vec{Z}(t,x)}{u(t,x) \wedge \vec{Z}(t,x)}\right)^2,$$

which in turn is bounded by  $1 \wedge |t|^{(2-\kappa)/2}$  by combining Theorem 2.1 and Lemma 2.5(i). The trick now is to write

$$\frac{u(t,x) - \vec{Z}(t,x)}{u(t,x)} = \int \frac{1 - \vec{Z}(0,x)}{u(t,x)} w_{0,y}(t,x) \, dy$$

and to note that  $\vec{Z}(0, x)$  is independent of both u(t, x) and  $w_{0,y}(t, x)$ , so that the expectation of this expression vanishes, and concluding the claim on |F(t) - c|.

From (4.25), we see that  $||R_{1,\varepsilon}(t,x)||_p \leq \varepsilon^{2\mu}$  uniformly in x and  $t \geq T_0$ . Hence, given that the bounds are uniform in space, integrating against any spatial test function  $g \in \overline{D}$  implies that  $R_{0,\varepsilon}$  and  $R_{1,\varepsilon}$  converge to zero in  $L^p([T_0,T], \mathcal{C}^{\alpha}(E))$ .

It remains to estimate  $R_{2,\varepsilon}$ . We claim, for any  $\delta \in (0, 1)$ , uniformly in  $\lambda \in (0, 1)$ ,

$$\left\| \int_{\mathbb{R}^d} R_{2,\varepsilon}(t,x) g_z^{\lambda}(x) \, dx \right\|_2 \lesssim \varepsilon^{\frac{\kappa}{2} - 1} t^{-\frac{\delta}{2}} (J_{\kappa - 2,\delta}(g_z^{\lambda}))^{\frac{1}{2}} \,. \tag{4.27}$$

By Lemma 4.4, for any  $g \in \overline{\mathcal{D}}$  we obtain

$$\sup_{t \ge T_0} \left\| \int_{\mathbb{R}^d} R_{2,\varepsilon}(t,x) g_z^{\lambda}(x) \, dx \right\|_2 \lesssim \varepsilon^{\frac{\kappa}{2} - 1} \lambda^{\delta + 1 - \frac{\kappa}{2}} \lesssim \varepsilon^{\frac{\kappa}{2} - 1} \lambda^{1 - \frac{\kappa}{2}}$$

As in the proof of Lemma 4.11, this implies that  $||R_{2,\varepsilon}(t,\cdot)||_{\mathcal{C}_2^{\alpha_0}} \lesssim \varepsilon^{\frac{\kappa}{2}-1}$  for any  $\alpha_0 \leqslant 1 - \frac{\kappa}{2}$ , uniformly in  $t \ge T_0$ . On the other hand, by the uniform moments bounds, we know

$$\sup_{t \ge T_0} \left\| \int_{\mathbb{R}^d} R_{2,\varepsilon}(t,x) g_z^{\lambda}(x) \, dx \right\|_p \lesssim \sup_{t,x} \|R_{2,\varepsilon}(t,x)\|_p \lesssim 1$$

Then, given  $\alpha < 1 - \frac{\kappa}{2}$ , picking  $p \ge 1$  sufficiently large such that  $\alpha < \alpha_0 - \frac{d}{p}$ , by Proposition 4.9 and interpolation we conclude that  $\sup_{t \ge T_0} || ||R_{2,\varepsilon}(t,\cdot)||_{\alpha,E}||_p \to 0$  as  $\varepsilon \to 0$  for  $p \in [2,\infty]$ . Therefore,  $R_{2,\varepsilon}$  vanishes in  $L^p([T_0,T], \mathcal{C}^{\alpha}(E))$ .

It remains to show (4.27), namely that  $\tilde{Z}(0, y/\varepsilon)$  satisfies a law of large numbers. We know that  $\tilde{Z}$  has constant expectation 1 and decorrelates in space by (2.28):

$$\Lambda(0, y/\varepsilon) := \operatorname{cov}(\bar{Z}(0, 0), \bar{Z}(0, y/\varepsilon)) \lesssim 1 \wedge \varepsilon^{\kappa - 2} |y|^{2-\kappa}$$

Then, we compute the second moments and use (4.11) with  $\delta \in (0, 1)$ ,

$$\begin{split} \mathbb{E}\Big[\Big(\int_{\mathbb{R}^d} R_{2,\varepsilon}(t,x)g_z^{\lambda}(x)\,dx\Big)^2\Big] \\ &= \int_{\mathbb{R}^{4d}} \Lambda(y_1/\varepsilon,y_2/\varepsilon)\prod_{i=1}^2 \left(P_t(x_i-y_i)h_0(y_i)g_z^{\lambda}(x_i)\right)\,dx\,dy \\ &\lesssim \varepsilon^{\kappa-2}t^{-\delta}\int_{\mathbb{R}^{4d}} |y_1-y_2|^{2-\kappa}\prod_{i=1}^2 \left(|x_i-y_i|^{\delta-d}|g_z^{\lambda}(x_i)|\right)\,dx\,dy \;, \end{split}$$

which implies (4.27), given definition (4.6) of  $J_{\kappa-2,\delta}$ . This concludes the proof of Theorem 1.3.

## References

- [ACQ11] G. AMIR, I. CORWIN, and J. QUASTEL. Probability distribution of the free energy of the continuum directed random polymer in 1+1 dimensions. *Comm. Pure Appl. Math.* 64, no. 4, (2011), 466–537. doi:10.1002/cpa.20347.
- [AKQ14] T. ALBERTS, K. KHANIN, and J. QUASTEL. The intermediate disorder regime for directed polymers in dimension 1 + 1. Ann. Probab. 42, no. 3, (2014), 1212–1256. doi:10.1214/13-A0P858.

- [BDJ99] J. BAIK, P. DEIFT, and K. JOHANSSON. On the distribution of the length of the longest increasing subsequence of random permutations. J. Amer. Math. Soc. 12, no. 4, (1999), 1119–1178. doi:10.1090/S0894-0347-99-00307-0.
- [BG97] L. BERTINI and G. GIACOMIN. Stochastic Burgers and KPZ equations from particle systems. *Comm. Math. Phys.* 183, no. 3, (1997), 571–607. doi:10.1007/s002200050044.
- [BQS11] M. BALÁZS, J. QUASTEL, and T. SEPPÄLÄINEN. Fluctuation exponent of the KPZ / stochastic Burgers equation. J. Amer. Math. Soc. 24, no. 3, (2011), 683–708. doi:10.1090/S0894-0347-2011-00692-9.
- [CCM19] F. COMETS, C. COSCO, and C. MUKHERJEE. Space-time fluctuation of the Kardar–Parisi–Zhang equation in  $d \ge 3$  and the Gaussian Free Field. *arXiv* preprints (2019). arXiv:1905.03200.
- [CCM20] F. COMETS, C. COSCO, and C. MUKHERJEE. Renormalizing the Kardar-Parisi-Zhang equation in  $d \ge 3$  in weak disorder. J. Stat. Phys. **179**, no. 3, (2020), 713–728. arXiv:1902.04104. doi:10.1007/s10955-020-02539-7.
- [CNN22] C. COSCO, S. NAKAJIMA, and M. NAKASHIMA. Law of large numbers and fluctuations in the sub-critical and  $L^2$  regions for SHE and KPZ equation in dimension  $d \ge 3$ . *Stochastic Process. Appl.* **151**, (2022), 127–173. doi:10.1016/j.spa.2022.05.010.
- [CZ20] F. CARAVENNA and L. ZAMBOTTI. Hairer's reconstruction theorem without regularity structures. *EMS Surv. Math. Sci.* 7, no. 2, (2020), 207–251. doi:10.4171/emss/39.
- [DGRZ20] A. DUNLAP, Y. GU, L. RYZHIK, and O. ZEITOUNI. Fluctuations of the solutions to the KPZ equation in dimensions three and higher. *Probab. Theory Related Fields* **176**, no. 3-4, (2020), 1217–1258. doi:10.1007/s00440-019-00938-w.
- [DGRZ21] A. DUNLAP, Y. GU, L. RYZHIK, and O. ZEITOUNI. The random heat equation in dimensions three and higher: the homogenization viewpoint. Arch. Ration. Mech. Anal. 242, no. 2, (2021), 827–873. doi:10.1007/s00205-021-01694-9.
- [GHL23] L. GEROLLA, M. HAIRER, and X.-M. LI. Fluctuations of stochastic PDEs with long-range correlations. *arXiv preprint* (2023). arXiv:2303.09811.
- [GIP15] M. GUBINELLI, P. IMKELLER, and N. PERKOWSKI. Paracontrolled distributions and singular PDEs. *Forum Math. Pi* **3**, (2015), e6, 75. doi:10.1017/fmp.2015.2.
- [GL20] Y. GU and J. LI. Fluctuations of a nonlinear stochastic heat equation in dimensions three and higher. SIAM J. Math. Anal. 52, no. 6, (2020), 5422– 5440. doi:10.1137/19M1296380.
- [GP17] M. GUBINELLI and N. PERKOWSKI. KPZ reloaded. *Comm. Math. Phys.* **349**, no. 1, (2017), 165–269. doi:10.1007/s00220-016-2788-3.
- [GRZ18] Y. GU, L. RYZHIK, and O. ZEITOUNI. The Edwards-Wilkinson limit of the random heat equation in dimensions three and higher. *Comm. Math. Phys.* 363, no. 2, (2018), 351–388. doi:10.1007/s00220-018-3202-0.

- [Hai13] M. HAIRER. Solving the KPZ equation. Ann. of Math. (2) **178**, no. 2, (2013), 559–664. doi:10.4007/annals.2013.178.2.4.
- [Hai14] M. HAIRER. A theory of regularity structures. *Invent. Math.* 198, no. 2, (2014), 269–504. doi:10.1007/s00222-014-0505-4.
- [HL18] M. HAIRER and C. LABBÉ. Multiplicative stochastic heat equations on the whole space. J. Eur. Math. Soc. (JEMS) 20, no. 4, (2018), 1005–1054. arXiv:1504.07162. doi:10.4171/JEMS/781.
- [HL22] M. HAIRER and X.-M. LI. Generating diffusions with fractional Brownian motion. Comm. Math. Phys. 396, no. 1, (2022), 91–141. arXiv:2109.06948. doi:10.1007/s00220-022-04462-2.
- [HS23] M. HAIRER and R. STEELE. The BPHZ theorem for regularity structures via the spectral gap inequality. *arXiv preprint* (2023). arXiv:2301.10081.
- [KCDW14] T. KLOSS, L. CANET, B. DELAMOTTE, and N. WSCHEBOR. Kardar– Parisi–Zhang equation with spatially correlated noise: A unified picture from nonperturbative renormalization group. *Phys. Rev. E* 89, no. 2(2014). doi:10.1103/PhysRevE.89.022108.
- [KPZ86] M. KARDAR, G. PARISI, and Y.-C. ZHANG. Dynamic scaling of growing interfaces. *Phys. Rev. Lett.* 56, (1986), 889–892. doi:10.1103/PhysRevLett.56.889.
- [KS99] E. KATZAV and M. SCHWARTZ. Self-consistent expansion for the Kardar– Parisi–Zhang equation with correlated noise. *Phys. Rev. E* 60, (1999), 5677– 5680. doi:10.1103/PhysRevE.60.5677.
- [Kup16] A. KUPIAINEN. Renormalization group and stochastic PDEs. Ann. Henri Poincaré 17, no. 3, (2016), 497–535. doi:10.1007/s00023-015-0408-y.
- [LZ22] D. LYGKONIS and N. ZYGOURAS. Edwards-Wilkinson fluctuations for the directed polymer in the full  $L^2$ -regime for dimensions  $d \ge$ 3. Ann. Inst. Henri Poincaré Probab. Stat. 58, no. 1, (2022), 65–104. doi:10.1214/21-aihp1173.
- [MQR21] K. MATETSKI, J. QUASTEL, and D. REMENIK. The KPZ fixed point. *Acta Math.* 227, no. 1, (2021), 115–203. doi:10.4310/acta.2021.v227.n1.a3.
- [MSZ16] C. MUKHERJEE, A. SHAMOV, and O. ZEITOUNI. Weak and strong disorder for the stochastic heat equation and continuous directed polymers in  $d \ge 3$ . *Electron. Commun. Probab.* **21**, (2016), Paper No. 61, 12. doi:10.1214/16-ECP18.
- [MU18] J. MAGNEN and J. UNTERBERGER. The scaling limit of the KPZ equation in space dimension 3 and higher. J. Stat. Phys. **171**, no. 4, (2018), 543–598. doi:10.1007/s10955-018-2014-0.
- [Nua06] D. NUALART. *The Malliavin calculus and related topics*. Probability and its Applications (New York). Springer-Verlag, Berlin, second ed., 2006, xiv+382. doi:10.1007/3-540-28329-3.
- [PZ00] S. PESZAT and J. ZABCZYK. Nonlinear stochastic wave and heat equations. *Probab. Theory Related Fields* 116, no. 3, (2000), 421–443. doi:10.1007/s004400050257.

- [QS23] J. QUASTEL and S. SARKAR. Convergence of exclusion processes and the KPZ equation to the KPZ fixed point. J. Amer. Math. Soc. **36**, no. 1, (2023), 251–289. doi:10.1090/jams/999.
- [Vir20] B. VIRÁG. The heat and the landscape I. *arXiv preprint* (2020). arXiv:2008.07241.