

A novel translationally invariant supersymmetric chain with inverse-square interactions: partition function, thermodynamics and criticality

Bireswar Basu-Mallick, Federico Finkel, Artemio González-López

Depto. de Física Teórica, Facultad de Ciencias Físicas, Plaza de las Ciencias 1, Universidad Complutense de Madrid, 28040 Madrid, SPAIN

E-mail: bireswar.basumallick@saha.ac.in, ffinkel@ucm.es, artemio@ucm.es

September 16, 2024

Abstract. We introduce a novel family of translationally-invariant $\text{su}(m|n)$ supersymmetric spin chains with long-range interaction not directly associated to a root system. We study the symmetries of this model, establishing in particular the existence of a boson-fermion duality characteristic of this type of systems. Taking advantage of the relation of the new chains with an associated many-body supersymmetric spin dynamical model, we are able to compute their partition function in closed form for all values of m and n and for an arbitrary number of spins. When both m and n are even, we show that the partition function factorizes as the product of the partition functions of two supersymmetric Haldane–Shastry spin chains, which in turn leads to a simple expression for the thermodynamic free energy per spin in terms of the Perron eigenvalue of a suitable transfer matrix. We use this expression to study the thermodynamics of a large class of these chains, showing in particular that the specific heat presents a single Schottky peak at approximately the same temperature as a suitable k -level model. We also analyze the critical behavior of the new chains, and in particular the ground state degeneracy and the existence of low energy excitations with a linear energy-momentum dispersion relation. In this way we show that the only possible critical chains are the ones with $m = 0, 1, 2$. In addition, using the explicit formula for the partition function we are able to establish the criticality of the $\text{su}(0|n)$ and $\text{su}(2|n)$ chains with even n , and to evaluate the central charge of their associated conformal field theory.

Keywords: integrable spin chains and vertex models; solvable lattice models; quantum criticality.

1. Introduction

In our previous paper [1] we introduced a novel translationally invariant spin chain with inverse-square interactions depending on both spin permutation and spin reversal operators, which reduces to the celebrated Haldane–Shastry (HS) chain [2, 3] when the latter operators are replaced by plus or minus the identity. The new model, which consists of N sites each of which is occupied by particles of a single species transforming under the fundamental representation of the $\mathfrak{su}(m)$ Lie algebra, is exactly solvable, in the sense that its partition function can be evaluated in closed form for arbitrary N . The ultimate reason for this is that the spin chain can be obtained as the strong coupling limit of a one-dimensional many-body spin dynamical model, whose partition function can be computed in closed form in this limit. The chain’s partition function can then be evaluated applying Polychronakos’s freezing trick [4, 5], which basically amounts to modding out the dynamical degrees of freedom of the spin dynamical model.

A remarkable property of the new solvable chain introduced in Ref. [1] is the fact that, unlike what is the case with practically all integrable spin chains with long-range interactions, it is not directly associated to a single classical (extended) root system (see, e.g., [6–8]). More precisely, since the latter chain is translationally invariant the interaction between two sites depends only on their distance, as is the case for spin chains of HS type constructed from the A_{N-1} root system. On the other hand, the chain’s Hamiltonian includes not only spin permutation but also spin reversal operators, in a combination characteristic of spin chains related to the D_N root system.

The structure of the new chain’s partition function turns out to be particularly simple in the case of even m . Indeed, in this case the model’s Hamiltonian is unitarily equivalent to the sum of two independent HS chain Hamiltonians, an ordinary one of type $\mathfrak{su}(\frac{m}{2})$ and a supersymmetric one of $\mathfrak{su}(1|1)$ type. This yields an elegant description of the spectrum in terms of Haldane’s (supersymmetric) motifs [9–12] of $\mathfrak{su}(\frac{m}{2})$ and $\mathfrak{su}(1|1)$ types, and also establishes the invariance of the model under the direct sum of the (super-)Yangians $Y(\mathfrak{gl}(0|m/2))$ and $Y(\mathfrak{gl}(1|1))$. In fact, the general $\mathfrak{su}(m|n)$ supersymmetric version of the original HS chain was introduced early on by Haldane himself [10]. This model, which consists of two species of particles behaving as bosons and fermions with m and n internal degrees of freedom, respectively, was thoroughly studied in Ref. [13]. In particular, a closed-form expression for its partition function was derived in the latter reference, which was used in turn in Ref. [14] to obtain a complete description of the spectrum in terms of $\mathfrak{su}(m|n)$ -supersymmetric bond vectors and their associated motifs.

In view of the above, it is natural to consider the $\mathfrak{su}(m|n)$ -supersymmetric generalization of the translationally invariant spin chain of Ref. [1]. This is, indeed, the first aim of this paper. More precisely, we shall construct the latter chain from its associated spin dynamical model and evaluate its partition function in closed form for all values of m and n . We shall show that when both m and n are even the model’s Hamiltonian is unitarily equivalent to the sum of the Hamiltonians of two

supersymmetric HS chains of $\text{su}(1|1)$ and $\text{su}(\frac{m}{2}|\frac{n}{2})$ types. As in the non-supersymmetric case, this automatically entails a simple description of the chain's spectrum in terms of $\text{su}(1|1)$ and $\text{su}(\frac{m}{2}|\frac{n}{2})$ supersymmetric motifs.

The thermodynamics of the original HS chain and its rational and hyperbolic variants has been extensively studied in the literature since its very inception [15–17]. More recently, the description of the spectrum in terms of Haldane motifs was systematically used in Ref. [18] to derive a closed-form expression of the thermodynamic functions of all (non-supersymmetric) spin chains of HS type related to the A_{N-1} root system. This method was later extended to $\text{su}(m|n)$ supersymmetric chains of HS type in Ref. [19]. Our second aim is to take advantage of the motif-based description of the spectrum of the $\text{su}(m|n)$ supersymmetric chain introduced in this paper when both m and n are even to compute the free energy of this model in the thermodynamic limit, deriving closed-form expressions for its main thermodynamic functions. In particular, using these expressions we shall show that the specific heat per spin exhibits a single Schottky peak, whose temperature is close to the temperature of the Schottky peak of an appropriate k -level system. When either m or n is odd, we shall use the exact expression of the partition function derived in this work to study the thermodynamic functions for N finite but as large as possible. We shall show that, barring finite-size effects at very low temperatures, their behavior is qualitatively analogous to that of their counterparts when both m and n are even.

One-dimensional spin chains are one of the simplest systems exhibiting quantum phase transitions [20]. This typically occurs when the spectrum is gapless in the thermodynamic limit, and the model's low-energy sector can be effectively described by a suitable $(1+1)$ -dimensional CFT (or, if the ground state is degenerate, by several copies thereof). For this to be possible the system's ground state must have finite degeneracy in the thermodynamic limit, and it must possess low-energy excitations above the ground state exhibiting a linear energy-momentum relation with a characteristic Fermi velocity v_F . This is the case, for example, for the original (antiferromagnetic) $\text{su}(m)$ HS chain [9, 21, 22], whose low-energy excitations are known to be governed by the $\text{su}(m)_1$ Wess–Zumino–Novikov–Witten model [23–25]. The critical behavior of $\text{su}(m|n)$ -supersymmetric HS spin chains was also analyzed in Ref. [26], where it was established that only the $\text{su}(1|m)$ chain (i.e., with one bosonic and n fermionic degrees of freedom) is also critical. Another hallmark of critical quantum systems is the low temperature behavior of their free energy, which should behave as the free energy of a $(1+1)$ -dimensional CFT. In other words, at low temperatures we should have [27, 28]

$$f(T) = f(0) - \frac{\pi c T^2}{6v_F} + o(T^2), \quad (1.1)$$

where c is the central charge of the associated CFT. In particular, the analysis of the low-temperature behavior of the free energy of a critical quantum system provides an efficient way of computing the central charge of its associated CFT, which in turn determines its universality class. Our third objective is to analyze the critical behavior of the $\text{su}(m|n)$ supersymmetric chains introduced in this paper, and in particular to determine for what

values of m and n they are critical and what is their central charge. By studying the ground state degeneracy of these models, we shall show that they can only be critical for $m = 0, 1, 2$. We shall then use the motif-based description of the spectrum to prove that the $\text{su}(0|n)$ and $\text{su}(2|n)$ chains are critical for all even values of n . This will be confirmed by the analysis of the low-temperature behavior of the thermodynamic free energy per spin, which also yields the central charge of the associated CFTs.

The rest of the paper is organized as follows. In Section 2 we introduce the novel $\text{su}(m|n)$ spin chains under consideration, and explain how they can be obtained from a suitable (supersymmetric) spin dynamical model in the strong coupling limit. Exploiting this connection, in Section 3 we evaluate in closed form the chain's partition function for arbitrary m , n and N , and use the explicit expression thus obtained to study the ground state degeneracy. In Section 4 we analyze the the new chain's main symmetries, showing that they possess a remarkable "twisted" translation invariance as well as a boson-fermion duality characteristic of this type of models. Section 5 is devoted to the study of the chain's thermodynamics. Taking advantage of the simple structure of the partition function, we are able to find analytic expressions for the thermodynamic free energy per spin and the main thermodynamic functions when m and n are both even. We also study the behavior of these functions, showing in particular that the specific heat features a single Schottky peak. In Section 6 we examine the critical behavior of the supersymmetric chains under study, deriving the partial results explained above. We present our conclusions and outline several paths for future research in Section 7. The paper ends with two technical appendixes, in which we discuss the precise connection of the $\text{su}(m|n)$ Lie superalgebra with our model and derive an asymptotic approximation for an integral used to ascertain the low-temperature behavior of the free energy per spin of the critical chains.

2. The model

The model we shall deal with in this paper is the supersymmetric version of the spin chain introduced in Ref. [1]. It describes a one-dimensional array of N spins, each of which can be either a boson or a fermion, lying on the upper unit half-circle at uniformly spaced positions $\zeta_k = e^{2i\theta_k}$, with

$$\theta_k := \frac{k\pi}{2N}, \quad 1 \leq k \leq N. \quad (2.1)$$

More precisely, we shall suppose that there are m bosonic and n fermionic degrees of freedom, so that the Hilbert space of the system is $\mathcal{S}^{(m|n)} = \otimes_{i=1}^N \mathcal{S}_i^{(m|n)}$ with $\mathcal{S}_i^{(m|n)} = \mathbb{C}^{m+n}$. The canonical basis in this space shall be denoted by

$$|s_1 \cdots s_N\rangle \equiv |\mathbf{s}\rangle := |s_1\rangle \otimes \cdots \otimes |s_N\rangle, \quad 1 \leq s_i \leq m+n. \quad (2.2)$$

We shall regard the basis states $|s_i\rangle$ with $s_i \in B := \{1, \dots, m\}$ as bosonic, and those with $s_i \in F := \{m+1, \dots, m+n\}$ as fermionic. The model's Hamiltonian admits a simple expression in terms of the $\text{su}(m|n)$ supersymmetric spin permutation and spin flip operators, whose definition we shall next recall.

The supersymmetric spin permutation operators $S_{ij}^{(m|n)} = S_{ji}^{(m|n)}$ (with $i < j$) are defined by

$$S_{ij}^{(m|n)} |\dots s_i \dots s_j \dots\rangle := (-1)^{\nu(s_i, \dots, s_j)} |\dots s_j \dots s_i \dots\rangle, \quad (2.3)$$

where $\nu(s_i, \dots, s_j)$ is 0 (respectively 1) if $s_i, s_j \in B$ (respectively $s_i, s_j \in F$), and is otherwise equal to the number of fermionic spins s_k with $i + 1 \leq k \leq j - 1$. Note that $S_{ij}^{(m|0)}$ is a standard permutation operator P_{ij} , while $S_{ij}^{(0|n)} = -P_{ij}$. Likewise, the action of the spin flip operators $S_i^{(m\varepsilon_B|n\varepsilon_F)}$ on the canonical basis vectors is given by

$$S_i^{(m\varepsilon_B|n\varepsilon_F)} |\dots s_i \dots\rangle := \sigma(s_i) |\dots s'_i \dots\rangle, \quad (2.4)$$

with

$$\sigma(s_i) = \begin{cases} \varepsilon_B, & s_i \in B, \\ \varepsilon_F, & s_i \in F, \end{cases} \quad (2.5)$$

where $\varepsilon_B, \varepsilon_F \in \{\pm 1\}$ are two fixed signs and $s_i \mapsto s'_i$ is the ‘‘spin flip’’ (involution) defined by

$$s'_i = \begin{cases} m + 1 - s_i, & s_i \in B, \\ 2m + n + 1 - s_i, & s_i \in F. \end{cases} \quad (2.6)$$

The precise connection between the spin permutation and reversal operators just defined with the Lie superalgebra $\mathfrak{su}(m|n)$ is explained out in [Appendix A](#). In terms of these operators, the model’s Hamiltonian is defined as

$$\mathcal{H}^{(m\varepsilon_B|n\varepsilon_F)} = \frac{1}{4} \sum_{1 \leq i < j \leq N} \left(\frac{1 - S_{ij}^{(m|n)}}{\sin^2(\theta_i - \theta_j)} + \frac{1 - \tilde{S}_{ij}^{(m\varepsilon_B|n\varepsilon_F)}}{\cos^2(\theta_i - \theta_j)} \right), \quad (2.7)$$

where we have set

$$\tilde{S}_{ij}^{(m\varepsilon_B|n\varepsilon_F)} := S_{ij}^{(m|n)} S_i^{(m\varepsilon_B|n\varepsilon_F)} S_j^{(m\varepsilon_B|n\varepsilon_F)}.$$

In what follows we shall also usually suppress the superindices m, n, ε_B , and ε_F from all operators, writing simply $S_{ij}, S_i, \tilde{S}_{ij}$ and \mathcal{H} .

Remark 1. Since the Hamiltonian (2.7) only contains products of two spin flip operators, it is clear that \mathcal{H} is invariant under the replacement $\varepsilon_B \rightarrow -\varepsilon_B, \varepsilon_F \rightarrow -\varepsilon_F$. For this reason, from now on we shall suppose without loss of generality that

$$\varepsilon_B = 1, \quad \varepsilon_F = \pm 1. \quad \square$$

Remark 2. When $m = 0$ and $\varepsilon_F = 1$ the Hamiltonian (2.7) reduces to the Hamiltonian of the non-supersymmetric spin chain introduced in Ref. [1]. On the other hand, if the spin flip operators $S_i^{(m|n)}$ are replaced by ± 1 then $\tilde{S}_{ij}^{(m|n)} = S_{ij}^{(m|n)}$ and (2.7) becomes

$$\begin{aligned} & \frac{1}{4} \sum_{1 \leq i < j \leq N} \left(\sin^{-2}(\theta_i - \theta_j) + \cos^{-2}(\theta_i - \theta_j) \right) \left(1 - S_{ij}^{(m|n)} \right) \\ & = \sum_{1 \leq i < j \leq N} \sin^{-2} \left(\frac{(i-j)\pi}{N} \right) \left(1 - S_{ij}^{(m|n)} \right) \equiv 2\mathcal{H}_{\text{HS}}^{(m|n)}, \end{aligned}$$

where $\mathcal{H}_{\text{HS}}^{(m|n)}$ denotes the Hamiltonian of the $\mathfrak{su}(m|n)$ Haldane–Shastry spin chain with the standard normalization (see, e.g., Ref. [13]). \square

It shall be useful for the sequel to define the sets

$$\begin{aligned} B_+ &= \{1, \dots, \lfloor m/2 \rfloor\}, & F_+ &= \{m+1, \dots, m + \lfloor n/2 \rfloor\}, \\ B_0 &= \{1, \dots, \lfloor (m+1)/2 \rfloor\}, & F_0 &= \{m+1, \dots, m + \lfloor (n+1)/2 \rfloor\}, \end{aligned}$$

where $\lfloor x \rfloor$ denotes the integer part of the real number x . Note that $B_+ = B_0$ for m even, while $B_0 = B_+ \cup \{(m+1)/2\}$ for m odd, and similarly for F_+ and F_0 . We shall colloquially say that a spin s_i is *positive* (respectively *non-negative*), and write $s_i \succ 0$ (respectively $s_i \succeq 0$) if $s_i \in B_+ \cup F_+$ (respectively $s_i \in B_0 \cup F_0$).

Remark 3. The operators S_{ij} and $S_i S_j$ generate the Weyl algebra of the D_N root system through the non-trivial relations

$$S_{ij}^2 = (S_i S_j)^2 = 1, \quad S_{ij} S_{jk} = S_{ik} S_{ij} = S_{jk} S_{ik}, \quad S_{ij} S_i S_k = S_j S_k S_{ij},$$

where the indices i, j, k are all distinct. Note, however, that the Hamiltonian (2.7) does *not* have the standard form[‡]

$$\mathcal{H}_D = \sum_{i < j} \left[f(\xi_i - \xi_j)(1 - S_{ij}) + f(\xi_i + \xi_j)(1 - \tilde{S}_{ij}) \right] \quad (2.8)$$

for a spin chain with *real* sites ξ_k associated to the D_N root system [29, 30]. Nor is it purely of A_N type like, e.g., the supersymmetric version of the original HS chain [13], due to the presence of the operators S_i in the Hamiltonian. In fact, the model (2.7) (as its non-supersymmetric version) is not directly associated to an extended root system, unlike most chains of HS type considered so far in the literature. \square

Remark 4. The Hamiltonian (2.7) can be written as

$$\mathcal{H} = \sum_{1 \leq i < j \leq N} \left(\frac{1 - S_{ij}}{|\zeta_i - \zeta_j|^2} + \frac{1 - \tilde{S}_{ij}}{|\zeta_i + \zeta_j|^2} \right) \quad (2.9)$$

in terms of the chain site coordinates $\zeta_k = e^{2i\theta_k}$. The first term in the Hamiltonian is the usual spin-spin interaction between the spins at sites ζ_i and ζ_j . On the other hand, the second term describes a non-standard interaction of the spin at site ζ_i with the reflection with respect to the origin of the spin at site ζ_j . Note, finally, that the fact that the chain sites lie on the *upper* unit half-circle means that, in spite of appearances, the chain (2.7) should be regarded as *open*. \square

An essential ingredient in the solvability of the chain (2.7) is its close connection with the supersymmetric dynamical spin model with Hamiltonian

$$\begin{aligned} H &= -\Delta + 8a \sum_{1 \leq i < j \leq N} \left(\frac{a - S_{ij}}{|z_i - z_j|^2} + \frac{a - \tilde{S}_{ij}}{|z_i + z_j|^2} \right) \\ &= -\Delta + 2a \sum_{i < j} \left[\frac{a - S_{ij}}{\sin^2(x_i - x_j)} + \frac{a - \tilde{S}_{ij}}{\cos^2(x_i - x_j)} \right] \end{aligned} \quad (2.10)$$

[‡] From now on, unless otherwise stated all summations and products will range over the set $\{1, \dots, N\}$.

and its scalar counterpart

$$\begin{aligned} H_{\text{sc}} &= -\Delta + 8a(a-1) \sum_{1 \leq i < j \leq N} \left(\frac{1}{|z_i - z_j|^2} + \frac{1}{|z_i + z_j|^2} \right) \\ &= -\Delta + 8a(a-1) \sum_{i < j} \sin^{-2}(2(x_i - x_j)), \end{aligned} \quad (2.11)$$

where

$$z_k = e^{2ix_k}, \quad (2.12)$$

$\Delta := \sum_i \frac{\partial^2}{\partial x_i^2} = -4 \sum_i \left(z_i \frac{\partial}{\partial z_i} \right)^2$ and $a > 1/2$. Indeed, it can be readily checked that the chain sites (2.1) are the coordinates of the minimum of the scalar potential

$$U(\mathbf{x}) = \sum_{1 \leq i < j \leq N} \left(\frac{1}{|z_i - z_j|^2} + \frac{1}{|z_i + z_j|^2} \right) = \sum_{i < j} \sin^{-2}(2(x_i - x_j))$$

of H_{sc} in the configuration space

$$A = \left\{ \mathbf{x} := (x_1, \dots, x_N) \in \mathbb{R}^N : x_1 < x_2 < \dots < x_N < x_1 + \frac{\pi}{2} \right\}, \quad (2.13)$$

of H_{sc} and H (this minimum is unique up to an inessential rigid translation [31]). We obviously can write

$$H = H_{\text{sc}} + 8aH_{\text{spin}}(\mathbf{x}),$$

where

$$H_{\text{spin}}(\mathbf{x}) := \frac{1}{4} \sum_{i < j} \left[\frac{1 - S_{ij}}{\sin^2(x_i - x_j)} + \frac{1 - \tilde{S}_{ij}}{\cos^2(x_i - x_j)} \right] \quad (2.14)$$

and

$$H_{\text{spin}}(\theta_1, \dots, \theta_N) = \mathcal{H}.$$

Hence in the large coupling limit $a \rightarrow \infty$ the energies of the spin dynamical model (2.10) are approximately given by

$$E_{ij} \simeq E_i + 8a\mathcal{E}_j,$$

where E_i and \mathcal{E}_j are two arbitrary eigenvalues of the Hamiltonians H_{sc} and \mathcal{H} , respectively. The above relation, although not suited for computing the spectrum of \mathcal{H} in terms of those of H and H_{sc} , yields however the following *exact* formula for the partition function \mathcal{Z} of the chain (2.7):

$$\mathcal{Z}(T) = \lim_{a \rightarrow \infty} \frac{Z(8aT)}{Z_{\text{sc}}(8aT)}. \quad (2.15)$$

Remark 5. In view of Eq. (2.12), it is natural to interpret the variables x_k appearing in the Hamiltonians (2.10) and (2.11) as half the *angular* coordinates of the particles, and $z_k = e^{2ix_k}$ as their actual (or *physical*) coordinates in the unit circle. Thus the latter Hamiltonians describe the motion of a system of particles (with or without $\text{su}(m|n)$ “spin”) in the unit circle. \square

Remark 6. The Hamiltonians H and H_{sc} in Eqs. (2.10)–(2.11) are closely related to the Hamiltonians of the Sutherland spin and scalar dynamical models, respectively given by

$$H_{\text{S}} = -\Delta + 2a \sum_{i < j} \sin^{-2}(x_i - x_j)(a - S_{ij}), \quad H_{\text{S,sc}} = -\Delta + 2a(a - 1) \sum_{i < j} \sin^{-2}(x_i - x_j).$$

Indeed, if the supersymmetric spin reversal operators S_i are replaced by plus or minus the identity then H and H_{sc} respectively reduce to $4H_{\text{S}}(2\mathbf{x})$ and $4H_{\text{S,sc}}(2\mathbf{x})$. \square

3. Evaluation of the partition function

In this section we shall use the freezing trick formula (2.15) to derive an exact expression for the partition function of the supersymmetric chain (2.7). To begin with, we quote the formula derived in Ref. [1] for the spectrum of the scalar Hamiltonian (2.11) in the center of mass (CM) frame:

$$E_{\mathbf{n}} = 4 \sum_i \left(2p_i + a(N - 2i + 1) \right)^2, \quad p_i := n_i - \frac{|\mathbf{n}|}{N}. \quad (3.1)$$

Here the multiindex $\mathbf{n} := (n_1, \dots, n_N)$ labeling the spectrum has non-negative integer components n_i satisfying

$$n_1 \geq n_2 \geq \dots \geq n_{N-1} \geq n_N = 0,$$

and we are using the notation

$$|\mathbf{n}| := \sum_i n_i.$$

Expanding $E_{\mathbf{n}}$ in powers of a we find that

$$E_{\mathbf{n}} = E_0 + 16a \sum_i p_i(N - 2i + 1) + O(1), \quad (3.2)$$

where

$$E_0 = 4a^2 \sum_i (N - 2i + 1)^2 = \frac{4}{3} N(N^2 - 1)a^2 \quad (3.3)$$

is the ground-state energy of H_{sc} in the CM frame. Hence

$$\lim_{a \rightarrow \infty} q^{-\frac{E_0}{8a}} Z_{\text{sc}}(8aT) = \sum_{n_1 \geq \dots \geq n_{N-1} \geq 0} q^{2 \sum_i p_i(N+1-2i)} = \prod_{i=1}^{N-1} (1 - q^{2i(N-i)})^{-1}, \quad (3.4)$$

where[§]

$$q := e^{-1/T}$$

(see Ref. [32] for details on the evaluation of the sum).

The computation of the spectrum of the dynamical spin Hamiltonian H in Eq. (2.10) proceeds along same lines as for its non-supersymmetric version in Ref. [1]. We shall therefore omit unnecessary details, referring to the latter reference where needed. To begin with, let us denote by Λ_s the total supersymmetric symmetrizer with respect to

[§] For convenience, we are setting the Boltzmann constant k_B equal to one.

simultaneous permutations of the particle's spatial coordinates and spins, determined by the relations

$$\Pi_{ij}\Lambda_s = \Lambda_s\Pi_{ij} = \Lambda_s, \quad 1 \leq i < j \leq N.$$

Here

$$\Pi_{ij} = K_{ij}S_{ij} = S_{ij}K_{ij},$$

where K_{ij} is the coordinate permutation operator defined by

$$K_{ij}f(z_1, \dots, z_i, \dots, z_j, \dots, z_N) = f(z_1, \dots, z_j, \dots, z_i, \dots, z_N),$$

and S_{ij} is the supersymmetric spin permutation operator acting on the internal (spin) $\mathcal{S}^{(m|n)}$ defined above. Note that Λ_s is an ordinary symmetrizer in the purely bosonic case $n = 0$, and an antisymmetrizer in the purely fermionic one $m = 0$. For instance, for $N = 2, 3$ we respectively have

$$\Lambda_s = \frac{1}{2}(1 + \Pi_{12}), \quad \Lambda_s = \frac{1}{6}(1 + \Pi_{12} + \Pi_{13} + \Pi_{23} + \Pi_{12}\Pi_{23} + \Pi_{12}\Pi_{13}).$$

Likewise, we define the total (i.e., with respect to coordinates and spin variables) supersymmetric flip operators Π_i by

$$\Pi_i = K_i S_i = S_i K_i,$$

where K_i is the operator flipping the *physical* coordinate of the i -th particle with respect to the origin:

$$K_i f(z_1, \dots, z_i, \dots, z_N) = f(z_1, \dots, -z_i, \dots, z_N).$$

The projectors Λ_0^\pm onto the spaces of states even (“+”) or odd (“−”) under the action of the supersymmetric flip operators Π_i are then given by

$$\Lambda_0^\varepsilon = \frac{1}{N!} \left(1 + \sum_{n=1}^N \sum_{i_1 < \dots < i_n} \varepsilon^n \Pi_{i_1} \dots \Pi_{i_n} \right), \quad \varepsilon = \pm 1.$$

Following the procedure in Ref. [1], we next construct a Schauder (i.e., non-orthonormal) basis on which the dynamical spin Hamiltonian H acts triangularly. To this end, we define the scalar functions

$$\varphi_{\mathbf{n}}(\mathbf{z}) = \mu(\mathbf{z}) \prod_i z_i^{n_i}, \quad n_1 \geq \dots \geq n_N,$$

where n_i is an integer and

$$\mu(\mathbf{z}) := \prod_{i < j} |z_i - z_j|^a$$

is the ground state wave function of the scalar Hamiltonian H_{sc} . It can then be shown that the states

$$\Phi_{\mathbf{n}, \mathbf{s}}^\varepsilon(\mathbf{z}) = \left(\prod_i z_i \right)^{-|\mathbf{n}|/N} \Lambda_s \Lambda_0^\varepsilon(\varphi_{\mathbf{n}}(\mathbf{z})|\mathbf{s}\rangle), \quad \varepsilon = \pm 1, \quad (3.5)$$

will in fact constitute a Schauder basis of the Hilbert space of H provided that the quantum numbers \mathbf{n} and \mathbf{s} are chosen so that the set of all such states is linearly

independent. Note that the prefactor in the RHS of Eq. (3.5) ensures that all the states $\Phi_{\mathbf{n},\mathbf{s}}^\varepsilon$ have vanishing linear momentum, i.e., that we are working in the CM frame. Moreover, since $\Phi_{\mathbf{n},\mathbf{s}}^\varepsilon$ changes at most by a sign if we add to each n_i an arbitrary integer, we shall from now normalize the multiindex \mathbf{n} by taking $n_N = 0$. Following Ref. [1], we shall impose the following conditions on the quantum numbers $\mathbf{n} \in (\mathbb{N} \cup \{0\})^N$ and $\mathbf{s} \in (B \cup F)^N$ to ensure the linear independence of the states (3.5):

$$\text{B1) } n_1 \geq \dots \geq n_N = 0.$$

$$\text{B2) } n_i = n_j \implies s_i \geq s_j, \text{ and } s_i > s_j \text{ if } s_i, s_j \in F.$$

$$\text{B3) } s_i \in B_0 \cup F_0 \text{ if } (-1)^{n_i} \sigma(s_i) = \varepsilon, \text{ and } s_i \in B_+ \cup F_+ \text{ if } (-1)^{n_i} \sigma(s_i) = -\varepsilon.$$

To understand these conditions, note that the states (3.5) verify

$$\Phi_{P_{ij}\mathbf{n}, P_{ij}\mathbf{s}}^\varepsilon = (-1)^{\nu(s_i, \dots, s_j)} \Phi_{\mathbf{n}, \mathbf{s}}^\varepsilon, \quad (3.6)$$

$$\Phi_{\mathbf{n}, P_i\mathbf{s}}^\varepsilon = \varepsilon (-1)^{n_i} \sigma(s_i) \Phi_{\mathbf{n}, \mathbf{s}}^\varepsilon, \quad (3.7)$$

where P_{ij} acts on the vectors \mathbf{n} and \mathbf{s} by permuting their i -th and j -th components and P_i acts on the multiindex \mathbf{s} by flipping its i -th component:

$$P_i(s_1, \dots, s_i, \dots, s_N) = (s_1, \dots, s'_i, \dots, s_N).$$

In particular, the wave functions $\Phi_{P_{ij}\mathbf{n}, P_{ij}\mathbf{s}}^\varepsilon$ and $\Phi_{\mathbf{n}, P_i\mathbf{s}}^\varepsilon$ define the same quantum state as $\Phi_{\mathbf{n}, \mathbf{s}}^\varepsilon$. Thus, taking advantage of (3.6), we can order the spin quantum numbers occupying the same positions as a sequence of consecutive equal components of the multiindex \mathbf{n} by applying a suitable permutation to \mathbf{n} and \mathbf{s} . This is the content of the second condition. As to the third one, note first of all that if $s_i \notin B_0 \cup F_0$ then $s'_i \in B_0 \cup F_0$, and by Eq. (3.7) the spin flip $s_i \mapsto s'_i$ does not change the state. Thus we can take $s_i \in B_0 \cup F_0$ by flipping the i -th spin if necessary. Whether s_i can be equal to $(m+1)/2$ (when m is odd) or $m+1+(n+1)/2$ (when n is odd) depends on the value of ε . Indeed, in either case $P_i\mathbf{s} = \mathbf{s}$, and hence (3.7) implies that

$$\Phi_{\mathbf{n}, \mathbf{s}}^\varepsilon = \Phi_{\mathbf{n}, P_i\mathbf{s}}^\varepsilon = \varepsilon (-1)^{n_i} \sigma(s_i) \Phi_{\mathbf{n}, \mathbf{s}}^\varepsilon \implies (-1)^{n_i} \sigma(s_i) = \varepsilon.$$

Thus s_i can take the value $(m+1)/2$ (when m is odd) or $m+1+(n+1)/2$ (when n is odd) if and only if $(-1)^{n_i} \sigma(s_i) = \varepsilon$. The last two remarks thus account for condition iii).

Proceeding in much the same way as for the non-supersymmetric case studied in Ref. [1], one can show that the states (3.5) with quantum numbers \mathbf{n} and \mathbf{s} obeying conditions B1)–B3) above can be ordered in such a way that the action of the spin Hamiltonian H on them is triangular. It follows that the eigenvalues of H (in the CM frame) coincide with its diagonal matrix elements in the latter basis, given by

$$E_{\mathbf{n}, \mathbf{s}}^\varepsilon = 4 \sum_i \left(p_i + a(N+1-2i) \right)^2 = E_{\mathbf{n}}, \quad p_i = n_i - \frac{|\mathbf{n}|}{N}, \quad (3.8)$$

where the quantum numbers $\varepsilon = \pm 1$, \mathbf{n} , \mathbf{s} satisfy conditions B1)–B3) (see Ref [1] for the details). Since the RHS of this equation depends only on the multiindex \mathbf{n} , the partition function of the dynamical spin Hamiltonian (2.10) is given by

$$Z(q) = \sum_{n_1 \geq \dots \geq n_{N-1} \geq 0} D(\mathbf{n}) q^{E_{\mathbf{n}}}, \quad (3.9)$$

where the *spin degeneracy* $D(\mathbf{n})$ is equal to the number of choices of pairs $(\varepsilon, \mathbf{s})$ satisfying conditions B1)–B3) for a given multiindex \mathbf{n} (with $n_1 \geq \dots \geq n_N = 0$). Using the expansion (3.2)–(3.3) of $E_{\mathbf{n}}$ in powers of a we obtain

$$\lim_{a \rightarrow \infty} q^{-\frac{E_0}{8a}} Z(8aT) = \sum_{n_1 \geq \dots \geq n_{N-1} \geq 0} D(\mathbf{n}) q^{\sum_i p_i (N+1-2i)}. \quad (3.10)$$

From Eqs. (2.15), (3.4), and (3.10) we then deduce the following formula for the partition function of the supersymmetric chain (2.7):

$$\mathcal{Z}(q) = \prod_{i=1}^{N-1} (1 - q^{2i(N-i)}) \cdot \sum_{n_1 \geq \dots \geq n_{N-1} \geq 0} D(\mathbf{n}) q^{\sum_i p_i (N+1-2i)}. \quad (3.11)$$

The first step in the evaluation of the partition function $\mathcal{Z}(q)$ is thus a combinatorial problem, namely the evaluation of the spin degeneracy $D(\mathbf{n})$ for an arbitrary integer multiindex $\mathbf{n} = (n_1, \dots, n_N)$ satisfying $n_1 \geq \dots \geq n_N = 0$. Due to condition B3), it is clear that this combinatorial problem depends crucially on the parity of m and n . Before starting a detailed analysis, it is convenient to parametrize the multiindex \mathbf{n} as follows:

$$\mathbf{n} = (\underbrace{\nu_1, \dots, \nu_1}_{\ell_1}, \dots, \underbrace{\nu_{r-1}, \dots, \nu_{r-1}}_{\ell_{r-1}}, \underbrace{0, \dots, 0}_{\ell_r}), \quad (3.12)$$

where $\nu_1 > \dots > \nu_{r-1} > \nu_r = 0$ (with $\nu_k \in \mathbb{Z}$), $\ell_1 + \dots + \ell_r = N$ (with $\ell_i > 0$ for all i) and $r = 1, \dots, N$. Conditions B1)–B3) imply that

$$D(\mathbf{n}) = \prod_{i=1}^r d_+(\ell_i, \nu_i) + \prod_{i=1}^r d_-(\ell_i, \nu_i) =: d(\boldsymbol{\ell}, \boldsymbol{\nu}), \quad (3.13)$$

where $\boldsymbol{\ell} = (\ell_1, \dots, \ell_r)$, $\boldsymbol{\nu} = (\nu_1, \dots, \nu_{r-1}, \nu_r = 0)$, and $d_{\varepsilon}(\ell, \nu)$ is the number of choices of ℓ spins $\{s_1, \dots, s_{\ell}\}$ satisfying

C1) $s_i \geq s_j$, and $s_i > s_j$ if $s_i, s_j \in F$.

C2) $s_i \in B_0 \cup F_0$ if $(-1)^{\nu} \sigma(s_i) = \varepsilon$, and $s_i \in B_+ \cup F_+$ if $(-1)^{\nu} \sigma(s_i) = -\varepsilon$.

Thus $d_{\pm}(\ell, \nu)$ is the contribution to the spin degeneracy of a constant “sector” (ν, \dots, ν) of length ℓ in the multiindex \mathbf{n} when ε takes the fixed value ± 1 . It is also clear that

$$d_{\pm}(\ell, \nu) = \sum_{k=0}^{\ell} d_{\pm}^B(k, \nu) d_{\pm}^F(\ell - k, \nu), \quad (3.14)$$

where $d_{\pm}^B(k, \nu)$ denotes the number of choices of k bosonic spin satisfying conditions C1)–C2) when $\varepsilon = \pm 1$, and similarly for $d_{\pm}^F(\ell - k, \nu)$. From the parametrization (3.12) of the multiindex \mathbf{n} and Eqs. (3.11)–(3.13) we obtain the following formula for the partition function of the supersymmetric chain (2.7)

$$\mathcal{Z}(q) = \prod_{i=1}^{N-1} (1 - q^{2i(N-i)}) \sum_{r=1}^N \sum_{\boldsymbol{\ell} \in \mathcal{P}_N(r)} \sum_{\nu_1 > \dots > \nu_{r-1} > 0} d(\boldsymbol{\ell}, \boldsymbol{\nu}) q^{\sum_i p_i (N+1-2i)}, \quad (3.15)$$

where $\mathcal{P}_N(r)$ denotes the set of compositions (i.e., ordered partitions) of the integer N into r parts.

3.1. m and n even

In this case condition C2) simplifies to

$$\text{C2')} \quad s_i \in B_+ \cup F_+,$$

independently of the value of ε , ν and ε_F . In particular, in this case the partition function is the same for $\varepsilon_F = 1$ and $\varepsilon_F = -1$. Since the cardinals of the sets B_+ and F_+ are respectively $m/2$ and $n/2$, by condition C1) we have||

$$d_{\pm}^B(k, \nu) = \binom{\frac{m}{2} + k - 1}{k}, \quad d_{\pm}^F(\ell - k, \nu) = \binom{\frac{n}{2}}{\ell - k}, \quad (3.16)$$

and therefore

$$d_{\pm}(\ell, \nu) = \sum_{k=0}^{\ell} \binom{\frac{m}{2} + k - 1}{k} \binom{\frac{n}{2}}{\ell - k}. \quad (3.17)$$

Thus $d_{\varepsilon}(\ell, \nu) \equiv d(\ell)$ is in this case independent of ν and ε . It follows that

$$d(\boldsymbol{\ell}, \boldsymbol{\nu}) = 2 \prod_{k=1}^r d(\ell_k) \equiv d(\boldsymbol{\ell}),$$

where the factor of 2 takes care of the two possible choices of $\varepsilon = \pm 1$, and Eq. (3.15) simplifies to

$$\mathcal{Z}(q) = 2 \prod_{i=1}^{N-1} \left(1 - q^{2i(N-i)}\right) \sum_{r=1}^N \sum_{\boldsymbol{\ell} \in \mathcal{P}_N(r)} \prod_{k=1}^r d(\ell_k) \sum_{\nu_1 > \dots > \nu_{r-1} > 0} q^{\sum_i p_i(N+1-2i)}. \quad (3.18)$$

To evaluate the last sum we define

$$\tilde{\nu}_i = \nu_i - \nu_{i+1}, \quad i = 1, \dots, r-1,$$

with $\nu_r = 0$, so that the sum over $\nu_1 > \dots > \nu_{r-1} > 0$ turns into an unrestricted sum over the positive integers $\tilde{\nu}_1, \dots, \tilde{\nu}_{r-1}$. Taking into account that

$$\sum_i p_i(N+1-2i) = \sum_{i=1}^{r-1} \tilde{\nu}_i \mathcal{E}(L_i), \quad L_i := \sum_{j=1}^i \ell_j,$$

where the *dispersion relation* \mathcal{E} is given by

$$\mathcal{E}(i) = i(N-i)$$

(cf. [1, 32]), we readily obtain

$$\sum_{\nu_1 > \dots > \nu_{r-1} > 0} q^{\sum_i p_i(N+1-2i)} = \sum_{\tilde{\nu}_1, \dots, \tilde{\nu}_{r-1} = 1}^{\infty} \prod_{i=1}^{r-1} q^{\tilde{\nu}_i \mathcal{E}(L_i)} = \prod_{i=1}^{r-1} \sum_{\tilde{\nu}_i = 1}^{\infty} q^{\tilde{\nu}_i \mathcal{E}(L_i)} = \prod_{i=1}^{r-1} \frac{q^{\mathcal{E}(L_i)}}{1 - q^{\mathcal{E}(L_i)}}.$$

Substituting into Eq. (3.18) we arrive at the following explicit function for the partition function of the supersymmetric chain (2.7) in this case:

$$\mathcal{Z}(q) = 2 \prod_{i=1}^{N-1} \left(1 + q^{\mathcal{E}(i)}\right) \cdot \sum_{r=1}^N \sum_{\boldsymbol{\ell} \in \mathcal{P}_N(r)} \prod_{i=1}^r d(\ell_i) \cdot q^{\sum_{i=1}^{r-1} \mathcal{E}(L_i)} \prod_{i=1}^{N-r} \left(1 - q^{\mathcal{E}(L'_i)}\right), \quad (3.19)$$

|| Here and in what follows we shall set $\binom{n}{k} = 0$ for $k > n \geq 0$ and $\binom{n}{0} = 1$ for $n \leq 0$.

with $d(\ell)$ defined in Eq. (3.17)

$$\{L'_1, \dots, L'_{N-r}\} = \{1, \dots, N-1\} \setminus \{L_1, \dots, L_{r-1}\}.$$

Thus when both m and n are even the partition function of the $\text{su}(m|n)$ supersymmetric chain (2.7) can be factored as the product

$$\mathcal{Z}^{(m|n)}(q) = \mathcal{Z}_{\text{HS}}^{(1|1)}(q) \mathcal{Z}_{\text{HS}}^{\left(\frac{m}{2}|\frac{n}{2}\right)}(q), \quad (3.20)$$

of the partition functions of the $\text{su}(1|1)$ and $\text{su}\left(\frac{m}{2}|\frac{n}{2}\right)$ HS spin chains [13]. In other words, in this case the supersymmetric chain Hamiltonian (2.7) is unitarily equivalent to the sum

$$\mathcal{H}_{\text{HS}}^{(1|1)} \otimes \mathbb{1} + \mathbb{1} \otimes \mathcal{H}_{\text{HS}}^{\left(\frac{m}{2}|\frac{n}{2}\right)}, \quad (3.21)$$

where $\mathcal{H}_{\text{HS}}^{(p|q)}$ denotes the Hamiltonian of the $\text{su}(p|q)$ HS chain. This property, which is not evident from Eq. (2.7), is the essential ingredient used to arrive at the description of the model's spectrum in terms of Haldane motifs and their corresponding Young tableaux akin to the one developed in Ref. [1] for the non-supersymmetric case.

The combinatorial formula (3.19) becomes particularly simple when $m, n \leq 2$. Indeed, when $m = 2$ and $n = 0$ by Eq. (3.17) we have

$$\prod_{i=1}^r d(\ell_i) = \prod_{i=1}^r \binom{i}{i} = 1,$$

so that

$$\mathcal{Z}_{\text{HS}}^{(1|0)}(q) = \sum_{r=1}^N \sum_{\ell \in \mathcal{P}_N(r)} q^{\sum_{i=1}^{r-1} \mathcal{E}(L_i)} \prod_{i=1}^{N-r} (1 - q^{\mathcal{E}(L'_i)}) = \prod_{i=1}^N (1 - q^{\mathcal{E}(i)} + q^{\mathcal{E}(i)}) = 1,$$

and therefore

$$\mathcal{Z}^{(2|0)}(q) = \mathcal{Z}_{\text{HS}}^{(1|1)}(q) = 2 \prod_{i=1}^{N-1} (1 + q^{\mathcal{E}(i)}). \quad (3.22)$$

Similarly, when $m = 0$ and $n = 2$ the spin degeneracy is given by

$$\prod_{i=1}^r d(\ell_i) = \begin{cases} 1, & \ell_i = 1 \quad \forall i, \\ 0, & \text{otherwise,} \end{cases}$$

so that

$$\mathcal{Z}_{\text{HS}}^{(0|1)}(q) = q^{\sum_{i=1}^{N-1} \mathcal{E}(i)} = q^{\frac{1}{6}N(N^2-1)}$$

and

$$\mathcal{Z}^{(0|2)}(q) = q^{\frac{1}{6}N(N^2-1)} \mathcal{Z}_{\text{HS}}^{(1|1)}(q) = 2q^{\frac{1}{6}N(N^2-1)} \prod_{i=1}^{N-1} (1 + q^{\mathcal{E}(i)}) = q^{\frac{1}{6}N(N^2-1)} \mathcal{Z}^{(2|0)}(q). \quad (3.23)$$

Finally, when $m = n = 2$ we have

$$\mathcal{Z}^{(2|2)}(q) = \left[\mathcal{Z}_{\text{HS}}^{(1|1)}(q) \right]^2 = 4 \prod_{i=1}^{N-1} (1 + q^{\mathcal{E}(i)})^2;$$

in particular, in this case the energy levels are at least four times degenerate.

In general, from Eq. (3.19) it follows that in this case all the energy levels have even degeneracy. This is an immediate consequence of the independence of the spin degeneracy factor from the value of $\varepsilon = \pm 1$. Another direct consequence of Eq. (3.19) is the fact that, with the normalization chosen for the Hamiltonian (2.7), all energies are nonnegative integers. Furthermore, the degeneracy of the model's zero mode is given by

$$\mathcal{Z}(0) = 2 \sum_{r=1}^N \sum_{\ell \in \mathcal{P}_N(r)} \prod_{i=1}^r d(\ell_i) \cdot q^{\sum_{i=1}^{r-1} \varepsilon(L_i)} \Big|_{q=0}.$$

A cursory inspection of the latter formula shows that the only partition of N contributing to the zero mode is $\ell = (N)$, so that

$$\mathcal{Z}(0) = d((N)) = 2 \sum_{k=0}^N \binom{\frac{m}{2} + k - 1}{k} \binom{\frac{n}{2}}{N - k}. \quad (3.24)$$

Since the RHS does not vanish unless $m = 0$, we conclude that the ground state degeneracy of the supersymmetric chain (2.7) with $m \neq 0$ and n both even coincides with the RHS of Eq. (3.24), which is in turn twice the ground state degeneracy of the $\text{su}(\frac{m}{2}, \frac{n}{2})$ HS chain.

3.2. m even and n odd

In this case condition C2) simply states that $s_i \in B_+$ if s_i is bosonic, and therefore $d_{\pm}^B(k, \nu)$ is still given by Eq. (3.16). On the other hand, for fermionic spins condition C2) reads

$$\text{C2F)} \quad s_i \in F_0 \text{ if } (-1)^\nu \varepsilon_F = \varepsilon, \text{ and } s_i \in F_+ \text{ if } (-1)^\nu \varepsilon_F = -\varepsilon.$$

Since

$$|F_0| = \frac{n+1}{2}, \quad |F_+| = \frac{n-1}{2},$$

a moment's thought reveals that

$$d_{\pm}^F(\ell - k, \nu) = \binom{\frac{1}{2}(n \pm \varepsilon_F (-1)^\nu)}{\ell - k}, \quad (3.25)$$

and therefore

$$d_{\pm}(\ell, \nu) = \sum_{k=0}^{\ell} \binom{\frac{m}{2} + k - 1}{k} \binom{\frac{1}{2}(n \pm \varepsilon_F (-1)^\nu)}{\ell - k}. \quad (3.26)$$

Hence

$$d_{\pm}(\ell, \nu; \varepsilon_F = 1) = d_{\mp}(\ell, \nu; \varepsilon_F = -1),$$

from which it follows that the spin degeneracy $d(\ell, \nu)$ given by Eq. (3.13) —and hence the partition function (3.15)— is again independent of ε_F . As in the previous case, it is convenient to rewrite $\mathcal{Z}(q)$ in terms of the independent variables $\tilde{\nu}_i = \nu_i - \nu_{i+1}$, namely

$$\mathcal{Z}(q) = \prod_{i=1}^{N-1} (1 - q^{2\varepsilon(i)}) \sum_{r=1}^N \sum_{\ell \in \mathcal{P}_N(r)} \sum_{\tilde{\nu}_1, \dots, \tilde{\nu}_{r-1}=1}^{\infty} d(\ell, \nu) q^{\sum_{i=1}^{r-1} \tilde{\nu}_i \varepsilon(L_i)}, \quad (3.27)$$

with

$$\nu_i = \sum_{j=i}^{r-1} \tilde{\nu}_j.$$

Since in this case $d_{\pm}(\ell, \nu)$ depends only on the parity of the integer ν , it is convenient to define

$$\tilde{\nu}_i = 2\tilde{n}_i - \delta_i,$$

where $\tilde{n}_i \in \mathbb{N}$ and $\delta_i \in \{0, 1\}$. We then have

$$(-1)^{\nu_i} = (-1)^{\Delta_i}, \quad \text{with} \quad \Delta_i := \sum_{k=i}^{r-1} \delta_k, \quad (3.28)$$

and therefore

$$\begin{aligned} \sum_{\tilde{\nu}_1, \dots, \tilde{\nu}_{r-1}=1}^{\infty} d(\ell, \boldsymbol{\nu}) q^{\sum_{i=1}^{r-1} \tilde{\nu}_i \mathcal{E}(L_i)} &= \sum_{\delta_1, \dots, \delta_{r-1}=0}^1 d(\ell, \boldsymbol{\Delta}) \sum_{\tilde{n}_1, \dots, \tilde{n}_{r-1}=1}^{\infty} q^{\sum_{i=1}^{r-1} (2\tilde{n}_i - \delta_i) \mathcal{E}(L_i)} \\ &= \sum_{\delta_1, \dots, \delta_{r-1}=0}^1 d(\ell, \boldsymbol{\Delta}) \prod_{i=1}^{r-1} \frac{q^{(2-\delta_i)\mathcal{E}(L_i)}}{1 - q^{2\mathcal{E}(L_i)}}, \end{aligned} \quad (3.29)$$

where $\boldsymbol{\Delta} = (\Delta_1, \dots, \Delta_{r-1}, \Delta_r = 0)$. Combining this equation with Eq. (3.27) we finally obtain the following explicit formula for the partition function $\mathcal{Z}(q)$:

$$\mathcal{Z}(q) = \sum_{r=1}^N \sum_{\ell \in \mathcal{P}_N(r)} \prod_{i=1}^{N-r} (1 - q^{2\mathcal{E}(L'_i)}) \sum_{\delta_1, \dots, \delta_{r-1}=0}^1 d(\ell, \boldsymbol{\Delta}) q^{\sum_{i=1}^{r-1} (2-\delta_i)\mathcal{E}(L_i)}. \quad (3.30)$$

The spin degeneracy factor can be somewhat simplified when $m = 2$ and $n = 1$. Indeed, for $n = 1$ the factor

$$\binom{\frac{1}{2}(n \pm \varepsilon_F(-1)^\nu)}{\ell - k} = \binom{\frac{1}{2}(1 \pm \varepsilon_F(-1)^\nu)}{\ell - k}$$

in Eq. (3.26) is equal to 1 if $(-1)^\nu = \mp \varepsilon_F$ and $k = \ell$, or $(-1)^\nu = \pm \varepsilon_F$ and $k = \ell - 1, \ell$, and is zero otherwise. Thus for $m = 2$ and $n = 1$ Eq. (3.26) reduces to

$$d_{\pm}(\ell, \nu) = 1 + \frac{1}{2}(1 \pm \varepsilon_F(-1)^\nu),$$

whence it follows that

$$\begin{aligned} d_{\pm}(\ell, \boldsymbol{\nu}) &= 2^{|\{i: (-1)^{\nu_i} = \pm \varepsilon_F, 1 \leq i \leq r\}|} \\ \implies d(\ell, \boldsymbol{\nu}) &= 2^{|\{i: (-1)^{\nu_i} = -1, 1 \leq i \leq r\}|} + 2^{|\{i: (-1)^{\nu_i} = 1, 1 \leq i \leq r\}|}, \end{aligned}$$

where $|A|$ denotes the cardinal of the set A .

As in the previous case, from Eq. (3.30) it follows that the energies are nonnegative integers, and by Eqs. (3.13) and (3.26) the degeneracy of the zero mode is given by

$$\mathcal{Z}(0) = d((N), (0)) = \sum_{k=0}^N \binom{\frac{m}{2} + k - 1}{k} \left[\binom{\frac{1}{2}(n-1)}{N-k} + \binom{\frac{1}{2}(n+1)}{N-k} \right]. \quad (3.31)$$

Since the right hand side does not vanish for $m \neq 0$, it follows that when $m \neq 0$ the ground state is the zero mode, with degeneracy equal to the RHS of the latter equation. In particular, for $n = 1$ and arbitrary (even) m the ground state degeneracy is simply

$$2 \binom{\frac{m}{2} + N - 1}{N} + \binom{\frac{m}{2} + N - 2}{N - 1}.$$

Thus for $m = 2$ and $n = 1$ the ground state of the supersymmetric chain (2.7) has zero energy and is three times degenerate, regardless of the value of N .

3.3. m odd and n even

This case is similar to the previous one, with the roles of m and n reversed. In other words,

$$d_{\pm}^B(k, \nu) = \binom{\frac{1}{2}(m \pm (-1)^\nu) + k - 1}{k}, \quad d_{\pm}^F(\ell - k, \nu) = \binom{\frac{n}{2}}{\ell - k} \quad (3.32)$$

for both $\varepsilon_F = 1$ and $\varepsilon_F = -1$, since as n is even condition C3) for fermions simply states that $s_i \in F_+$. We thus have

$$d_{\pm}(\ell, \nu) = \sum_{k=0}^{\ell} \binom{\frac{n}{2}}{\ell - k} \binom{\frac{1}{2}(m \pm (-1)^\nu) + k - 1}{k}. \quad (3.33)$$

Equation (3.30) for the partition function is still valid in this case —with $d_{\pm}(\ell, \nu)$ defined by Eq. (3.33) instead of (3.26)—, since in its derivation we only used the fact that $d(\ell, \nu)$ in Eq. (3.26) depended on ν through $(-1)^\nu$.

As in the previous case, the spin degeneracy factor simplifies to some extent when $m = 1$. Indeed, in this case the second binomial coefficient in Eq. (3.33) is equal to 1 for all values of $k = 0, \dots, \ell$ when $(-1)^\nu = \varepsilon$, so that

$$d_{\varepsilon}(\ell, \nu) = \sum_{k=0}^{\ell} \binom{\frac{n}{2}}{\ell - k}, \quad (-1)^\nu = \varepsilon.$$

In particular,

$$d_{\varepsilon}(\ell, \nu) = 2^{n/2}, \quad (-1)^\nu = \varepsilon \quad \text{and} \quad \ell \geq \frac{n}{2}.$$

On the other hand, when $(-1)^\nu = -\varepsilon$ the latter binomial coefficient reduces to

$$\binom{k-1}{k} = \begin{cases} 1, & k = 0 \\ 0, & 1 \leq k \leq \ell, \end{cases}$$

and therefore

$$d_{-\varepsilon}(\ell, \nu) = \binom{\frac{n}{2}}{\ell}, \quad (-1)^\nu = -\varepsilon.$$

As in the previous cases, the energies are nonnegative integers, and the degeneracy of the zero mode can be easily computed from the formula

$$\mathcal{Z}(0) = d((N), (0)) = \sum_{k=0}^N \binom{\frac{n}{2}}{N-k} \left[\binom{\frac{1}{2}(m-1) + k - 1}{k} + \binom{\frac{1}{2}(m+1) + k - 1}{k} \right]. \quad (3.34)$$

Since this number does not vanish, it follows that the ground state has again energy zero and degeneracy given by the previous equation. In particular, for $m = 1$ the ground state degeneracy is

$$\binom{\frac{n}{2}}{N} + \sum_{k=0}^N \binom{\frac{n}{2}}{k},$$

which reduces to $2^{n/2}$ (independent of N) when $N > n/2$.

3.4. m and n odd

In this case the fermionic spins satisfy condition C2F) above, so that $d_{\pm}^F(\ell - k, \nu)$ is given by Eq. (3.25). The bosonic spins verify the analogous condition

$$\text{C2B) } s_i \in B_0 \text{ if } (-1)^\nu = \varepsilon, \text{ and } s_i \in B_+ \text{ if } (-1)^\nu = -\varepsilon,$$

so that $d_{\pm}^B(k, \nu)$ is given by Eq. (3.33). From Eq. (3.14) it then follows that

$$d_{\pm}(\ell, \nu; \varepsilon_F) = \sum_{k=0}^{\ell} \binom{\frac{1}{2}(m \pm (-1)^\nu) + k - 1}{k} \binom{\frac{1}{2}(n \pm \varepsilon_F(-1)^\nu)}{\ell - k}. \quad (3.35)$$

Note that in this case $d_{\pm}(\ell, \nu; \varepsilon_F = 1) \neq d_{\mp}(\ell, \nu; \varepsilon_F = -1)$, and hence the degeneracy factor $d(\boldsymbol{\ell}, \boldsymbol{\nu})$ in Eq. (3.13) is *not* the same for both values of ε_F . More precisely,

$$\begin{aligned} d(\boldsymbol{\ell}, \boldsymbol{\nu}; \varepsilon_F = 1) &= \prod_{i=1}^r \sum_{k=0}^{\ell_i} \binom{\frac{1}{2}(m + (-1)^{\nu_i}) + k - 1}{k} \binom{\frac{1}{2}(n + (-1)^{\nu_i})}{\ell_i - k} \\ &\quad + \prod_{i=1}^r \sum_{k=0}^{\ell_i} \binom{\frac{1}{2}(m - (-1)^{\nu_i}) + k - 1}{k} \binom{\frac{1}{2}(n - (-1)^{\nu_i})}{\ell_i - k}, \end{aligned} \quad (3.36)$$

$$\begin{aligned} d(\boldsymbol{\ell}, \boldsymbol{\nu}; \varepsilon_F = -1) &= \prod_{i=1}^r \sum_{k=0}^{\ell_i} \binom{\frac{1}{2}(m + (-1)^{\nu_i}) + k - 1}{k} \binom{\frac{1}{2}(n - (-1)^{\nu_i})}{\ell_i - k} \\ &\quad + \prod_{i=1}^r \sum_{k=0}^{\ell_i} \binom{\frac{1}{2}(m - (-1)^{\nu_i}) + k - 1}{k} \binom{\frac{1}{2}(n + (-1)^{\nu_i})}{\ell_i - k}. \end{aligned} \quad (3.37)$$

The partition function in this case is again given by Eq. (3.30), with $d(\boldsymbol{\ell}, \boldsymbol{\nu})$ replaced by $d(\boldsymbol{\ell}, \boldsymbol{\nu}; \varepsilon_F)$ in Eqs. (3.36)-(3.37). In particular,

$$\mathcal{Z}(q; \varepsilon_F = 1) \neq \mathcal{Z}(q; \varepsilon_F = -1).$$

As an example, consider the simplest case $m = n = 1$. To begin with, if $\varepsilon_F = 1$ the spin flip operator S_i reduces to the identity. Hence $\tilde{S}_{ij} = S_{ij} \equiv S_{ij}^{(1|1)}$ and

$$\mathcal{H} = \frac{1}{4} \sum_{i < j} (1 - S_{ij}^{(1|1)}) \left(\sin^{-2}(\theta_i - \theta_j) + \cos^{-2}(\theta_i - \theta_j) \right) = \sum_{i < j} \frac{1 - S_{ij}^{(1|1)}}{\sin^2(2(\theta_i - \theta_j))},$$

which (apart from a conventional factor of $1/2$) is the Hamiltonian of the $\text{su}(1|1)$ Haldane–Shastry model [13].

Consider next the case $m = n = 1$ and $\varepsilon_F = -1$. Much as in the previous cases, the chain's energies in this case are nonnegative integers, and the zero mode degeneracy can

be readily computed from Eq. (3.30) for the partition function from the usual formula $\mathcal{Z}(0) = d((N), (0))$. In general, when $m = n = 1$ we have

$$\binom{\frac{1}{2}(1 + \varepsilon(-1)^\nu) + k - 1}{k} = \begin{cases} 1, & (-1)^\nu = \varepsilon \\ 1, & (-1)^\nu = -\varepsilon \text{ and } k = 0 \\ 0, & (-1)^\nu = -\varepsilon \text{ and } k > 0 \end{cases}$$

$$\binom{\frac{1}{2}(1 - \varepsilon(-1)^\nu)}{\ell - k} = \begin{cases} 1, & (-1)^\nu = \varepsilon \text{ and } k = \ell \\ 1, & (-1)^\nu = -\varepsilon \text{ and } k = \ell - 1, \ell \\ 0, & \text{otherwise,} \end{cases}$$

and therefore (assuming, again, that $\varepsilon_F = -1$)

$$d_\varepsilon(\ell, \nu) = \begin{cases} 1, & (-1)^\nu = \varepsilon \\ 1, & (-1)^\nu = -\varepsilon \text{ and } \ell = 1 \\ 0, & (-1)^\nu = -\varepsilon \text{ and } \ell > 1. \end{cases}$$

It follows that $\prod_{i=1}^r d_\varepsilon(\ell_i, \nu_i)$ is equal to 1 if $\ell_i = 1$ for all i such that $(-1)^{\nu_i} = -\varepsilon$, and vanishes otherwise. We thus see that in this case the spin degeneracy depends crucially on the number of components of ℓ greater than 1, or equivalently of sectors of length greater than one in the multiindex \mathbf{n} . More precisely, calling

$$\lambda(\ell) = |\{i : \ell_i > 1, 1 \leq i \leq r\}|,$$

the spin degeneracy of the $\text{su}(1, 1)$ supersymmetric chain (2.7) is given by

$$d(\ell, \nu) = \begin{cases} 2, & \ell = (1, \dots, 1) \\ 1, & \lambda(\ell) = 1 \\ 1, & \lambda(\ell) > 1 \text{ and } (-1)^{\nu_i} = (-1)^{\nu_j} \text{ if } \ell_i, \ell_j > 1 \\ 0, & \text{otherwise.} \end{cases} \quad (3.38)$$

In particular, the degeneracy of the zero mode is in this case

$$\mathcal{Z}(0) = d((N), (0)) = 1.$$

In other words, when $m = n = 1$ and $\varepsilon_F = -1$ the ground state is non-degenerate for all values of N . Combining Eq. (3.38) for the spin degeneracy with Eqs. (3.27) and (3.29) we also obtain the following more explicit formula for the partition function of the $\text{su}(1|1)$ chain:

$$\begin{aligned} \mathcal{Z}^{(1|1)}(q; \varepsilon_F = -1) &= q^{e(N)} \mathcal{Z}_{\text{HS}}^{(1|1)}(q) \\ &+ \prod_{i=1}^{N-1} (1 + q^{\mathcal{E}(i)}) \sum_{1 \leq k < j \leq N} q^{e(k) + e(N-j+1)} \prod_{i=k}^{j-1} (1 - q^{\mathcal{E}(i)}) \\ &+ \sum_{r=2}^{N-2} \sum_{\substack{\ell \in \mathcal{P}_N(r) \\ \lambda(\ell) > 1}} \prod_{i=1}^{N-r} (1 - q^{2\mathcal{E}(L'_i)}) \sum_{\substack{\delta_1, \dots, \delta_{r-1} \in \{0,1\} \\ \ell_i, \ell_j > 1 \Rightarrow (-1)^{\Delta_i} = (-1)^{\Delta_j}}} q^{\sum_{i=1}^{r-1} (2-\delta_i)\mathcal{E}(L_i)}, \end{aligned}$$

where

$$e(k) = \sum_{i=1}^{k-1} \mathcal{E}(i) = \frac{1}{6}k(k-1)(3N-2k+1).$$

Proceeding as above it is straightforward to show that for both $\varepsilon_F = 1$ and $\varepsilon_F = -1$ the ground state is the zero mode, and its degeneracy is given by

$$\mathcal{Z}(0) = d((N), (0)) = \sum_{k=0}^N \left[\binom{\frac{1}{2}(m+1) + k - 1}{k} \binom{\frac{1}{2}(n + \varepsilon_F)}{N - k} + \binom{\frac{1}{2}(m-1) + k - 1}{k} \binom{\frac{1}{2}(n - \varepsilon_F)}{N - k} \right] \quad (3.39)$$

In particular,

$$\mathcal{Z}(0) = \begin{cases} \binom{\frac{1}{2}(m-1) + N}{N} + \binom{\frac{1}{2}(m-1) + N - 1}{N} + \binom{\frac{1}{2}(m + \varepsilon_F) + N - 2}{N - 1}, & n = 1 \\ 2^{\frac{1}{2}(n + \varepsilon_F)} + \binom{\frac{1}{2}(n - \varepsilon_F)}{N}, & m = 1. \end{cases}$$

4. Symmetries

In this Section we shall examine two basic symmetries of the supersymmetric chain (2.7), namely invariance under “twisted” translations (defined below) and boson-fermion duality.

4.1. Twisted translations

Although the interaction strengths in Eq. (2.7) depend on the differences $\theta_i - \theta_j$, the model clearly lacks translation invariance due to its fundamentally *open* nature. Indeed, while a formal “translation” $k \mapsto k + 1$ along the chain maps the k -th site located at $\zeta_k = e^{2i\theta_k}$ into the $(k + 1)$ -th for $1 \leq k < N$, the N -th site at $\zeta_N = e^{2i\theta_N} = -1$ is mapped to the point $\zeta_{N+1} = -e^{i\pi/N} \neq \zeta_1 = e^{i\pi/N}$. The lack of translation invariance of the Hamiltonian (2.7) can be more formally shown by computing the action on the \mathcal{H} of the (left) translation operator T , defined by the relation

$$T|s_1, \dots, s_N\rangle := |s_2, \dots, s_N, s_1\rangle.$$

Clearly $T^\dagger = T^{-1}$ and

$$T^\dagger S_{ij} T = S_{i+1, j+1}, \quad T^\dagger \tilde{S}_{ij} T = \tilde{S}_{i+1, j+1}$$

with

$$S_{k, N+1} \equiv S_{1k}, \quad \tilde{S}_{k, N+1} \equiv \tilde{S}_{1k}. \quad (4.1)$$

A straightforward calculation then shows that

$$\begin{aligned} T^\dagger \mathcal{H} T &= \frac{1}{4} \sum_{2 \leq i < j \leq N} \left[\frac{1 + S_{ij}}{\sin^2(\theta_i - \theta_j)} + \frac{1 + \tilde{S}_{ij}}{\cos^2(\theta_i - \theta_j)} \right] + \frac{1}{4} \sum_{j=2}^N \left[\frac{1 + \tilde{S}_{1j}}{\sin^2(\theta_j - \theta_1)} + \frac{1 + S_{1j}}{\cos^2(\theta_j - \theta_1)} \right] \\ &= \mathcal{H} - \sum_{j=2}^N \frac{\cos(2(\theta_j - \theta_1))}{\sin^2(2(\theta_j - \theta_1))} S_{1j} (1 - S_1 S_j) \end{aligned}$$

(cf. Ref. [1] for more details). Remarkably, a form of translation invariance can be recovered by combining an ordinary translation with a spin flip in one of the chain's ends. More precisely, let us define the *twisted translation* operator \mathcal{T} by

$$\mathcal{T} := TS_1,$$

whose action on the canonical spin basis (2.2) is given by

$$\mathcal{T}|s_1, \dots, s_N\rangle = |s_2, \dots, s_N, s'_1\rangle.$$

Note that \mathcal{T} is unitary, since both T and S_1 are; indeed, S_1 is idempotent ($S_1^2 = S_1$) and self-adjoint. It is straightforward to show that

$$\mathcal{T}^\dagger S_{ij} \mathcal{T} = S_{i+1, j+1}, \quad \mathcal{T}^\dagger \tilde{S}_{ij} \mathcal{T} = \tilde{S}_{i+1, j+1},$$

provided that we set

$$S_{k, N+1} \equiv \tilde{S}_{1k}, \quad \tilde{S}_{k, N+1} \equiv S_{1k}.$$

From these relations it readily follows that

$$\mathcal{T}^\dagger \mathcal{H} \mathcal{T} = \mathcal{H},$$

so that the chain Hamiltonian (2.7) commutes with the elements of the group of twisted translations generated by \mathcal{T} . Note that the identity $S_i T = T S_{i+1}$ implies that $\mathcal{T}^k = T^k S_k \cdots S_1$ for $k = 1, \dots, N$. In particular,

$$\mathcal{T}^N = S_N \cdots S_1 \neq \mathbb{1}, \quad \text{but} \quad \mathcal{T}^{2N} = \mathbb{1}.$$

Thus the twisted translation group generated by the operator \mathcal{T} is a cyclic group of order $2N$, i.e., *twice* the order of the standard translation group generated by the ordinary translation operator T .

4.2. Boson-fermion duality

We shall next analyze the relation between the Hamiltonians $\mathcal{H}^{(m \in B | n \in F)}$ and $\mathcal{H}^{(n \in F | m \in B)}$ (which, by Remark 1, coincides with $\mathcal{H}^{(n \in B | m \in F)}$), differing by the exchange of the bosonic and fermionic degrees of freedom. To this end, we start by defining the operator $\chi^{(m|n)} : \mathcal{S}^{(m|n)} \rightarrow \mathcal{S}^{(n|m)}$ by

$$\chi^{(m|n)}|s_1, \dots, s_N\rangle = |\hat{s}_1, \dots, \hat{s}_N\rangle,$$

where

$$\hat{s}_i = \begin{cases} s_i + n, & m \in B, \\ s_i - m, & s_i \in F. \end{cases}$$

In other words, the state $|\hat{s}_1, \dots, \hat{s}_N\rangle$ is obtained replacing the k -th bosonic (resp. fermionic) spin in $|s_1, \dots, s_N\rangle$ by the k -th fermionic (resp. bosonic) spin. Following Refs. [12, 33], we next define $\rho : \mathcal{S}^{(m|n)} \rightarrow \mathcal{S}^{(m|n)}$ by

$$\rho|s_1, \dots, s_N\rangle = (-1)^{\sum_k k\pi(s_k)}|s_1, \dots, s_N\rangle,$$

where

$$\pi(s_i) = \begin{cases} 0, & s_i \in B, \\ 1, & s_i \in F, \end{cases}$$

and consider the operator

$$U := \chi^{(m|n)} \rho : \mathcal{S}^{(m|n)} \rightarrow \mathcal{S}^{(n|m)}.$$

The operator U is clearly unitary, since it maps one orthonormal basis into another one. Moreover, since obviously $\rho^{-1} = \rho$ and $(\chi^{(m|n)})^{-1} = \chi^{(n|m)}$ it follows that $U^{-1} = \rho \chi^{(n|m)}$. We then have

$$\begin{aligned} U^{-1} S_{ij}^{(n|m)} U |s_1, \dots, s_N\rangle &= (-1)^{\sum_k k\pi(s_k)} \rho \chi^{(n|m)} S_{ij}^{(n|m)} |\hat{s}_1, \dots, \hat{s}_N\rangle \\ &= (-1)^{\sum_k k\pi(s_k)} (-1)^{\nu(\hat{s}_i, \dots, \hat{s}_j)} \rho \chi^{(n|m)} |\hat{s}_1, \dots, \hat{s}_j, \dots, \hat{s}_i, \dots, \hat{s}_N\rangle \\ &= \varepsilon(s_i, \dots, s_j) |s_1, \dots, s_j, \dots, s_i, \dots, s_N\rangle \end{aligned}$$

with

$$\begin{aligned} \varepsilon(s_i, \dots, s_j) &= (-1)^{\sum_k k\pi(s_k)} (-1)^{\sum_{k \neq i, j} k\pi(s_k)} (-1)^{i\pi(s_j) + j\pi(s_i)} (-1)^{\nu(\hat{s}_i, \dots, \hat{s}_j)} \\ &= (-1)^{(j-i)(\pi(s_i) - \pi(s_j))} (-1)^{\nu(\hat{s}_i, \dots, \hat{s}_j)}. \end{aligned}$$

Clearly, when $\pi(s_i) = \pi(s_j)$ we have

$$\varepsilon(s_i, \dots, s_j) = (-1)^{\nu(\hat{s}_i, \dots, \hat{s}_j)} = (-1)^{\pi(\hat{s}_i)} = -(-1)^{\pi(s_i)} = -(-1)^{\nu(s_i, \dots, s_j)}.$$

On the other hand, when $\pi(s_i) \neq \pi(s_j)$ the identity

$$\nu(s_i, \dots, s_j) + \nu(\hat{s}_i, \dots, \hat{s}_j) = j - i - 1$$

implies that also in this case $\varepsilon(s_i, \dots, s_j) = -(-1)^{\nu(s_i, \dots, s_j)}$. Hence

$$U^{-1} S_{ij}^{(n|m)} U |s_1, \dots, s_N\rangle = -(-1)^{\nu(s_i, \dots, s_j)} |s_1, \dots, s_j, \dots, s_i, \dots, s_N\rangle = -S_{ij}^{(m|n)} |s_1, \dots, s_N\rangle,$$

and therefore

$$U^{-1} S_{ij}^{(n|m)} U = -S_{ij}^{(m|n)}. \quad (4.2)$$

Similarly,

$$\begin{aligned} U^{-1} S_i^{(n|m)} U |s_1, \dots, s_N\rangle &= (-1)^{\sum_k k\pi(s_k)} \rho \chi^{(n|m)} S_i^{(n|m)} |\hat{s}_1, \dots, \hat{s}_N\rangle \\ &= (-1)^{\sum_k k\pi(s_k)} \sigma(\hat{s}_i) \rho \chi^{(n|m)} |\hat{s}_1, \dots, (\hat{s}_i)', \dots, \hat{s}_N\rangle \\ &= \sigma(\hat{s}_i) |s_1, \dots, s_i', \dots, s_N\rangle = \sigma(s_i) |s_1, \dots, s_i', \dots, s_N\rangle \\ &= S_i^{(m|n)} |s_1, \dots, s_i, \dots, s_N\rangle, \end{aligned}$$

and thus

$$U^{-1} S_i^{(n|m)} U = S_i^{(m|n)}. \quad (4.3)$$

Combining Eqs. (4.2) and (4.3) we obtain

$$U^{-1} \tilde{S}_{ij}^{(n|m)} U = -\tilde{S}_{ij}^{(m|n)},$$

whence

$$U^{-1} \mathcal{H}^{(n|m)} U = \frac{1}{4} \sum_{i < j} \left(\frac{1 + S_{ij}^{(m|n)}}{\sin^2(\theta_i - \theta_j)} + \frac{1 + \tilde{S}_{ij}^{(m|n)}}{\cos^2(\theta_i - \theta_j)} \right) = \mathcal{E}_0 - \mathcal{H}^{(m|n)}, \quad (4.4)$$

where

$$\mathcal{E}_0 = \frac{1}{2} \sum_{i < j} \left[\sin^{-2}(\theta_i - \theta_j) + \cos^{-2}(\theta_i - \theta_j) \right] = 2 \sum_{i < j} \sin^{-2} \left(\frac{\pi(j-i)}{N} \right) = \frac{1}{3} N(N^2 - 1) \quad (4.5)$$

(see Ref. [32] for the evaluation of the sum). In other words, the partition functions of $\mathcal{H}^{(n|m)}$ and $\mathcal{H}^{(m|n)}$ are related by

$$\mathcal{Z}^{(n|m)}(q) = q^{\mathcal{E}_0} \mathcal{Z}^{(m|n)}(q^{-1}). \quad (4.6)$$

In particular, for $m = n$ we have

$$\mathcal{Z}^{(m|m)}(q) = q^{\mathcal{E}_0} \mathcal{Z}^{(m|m)}(q^{-1}), \quad (4.7)$$

implying that the spectrum of $\mathcal{H}^{(m|m)}$ is symmetric about $\mathcal{E}_0/2$. Note that both equations (4.6) and (4.7) can be explicitly checked in the cases $(m, n) = (0, 2), (2, 0), (2, 2)$ studied in Section 3.1.

5. Thermodynamics

With the help of the partition function $\mathcal{Z}_N(q)$ of the $\text{su}(m|n)$ chain (2.7) with N spins computed in Section 3 one can in principle obtain the thermodynamic free energy per particle

$$f(T) = -T \lim_{N \rightarrow \infty} N^{-1} \log \mathcal{Z}_N(q), \quad q = e^{-1/T} \equiv e^{-\beta}, \quad (5.1)$$

from which all the other thermodynamic functions (internal energy, specific heat at constant volume, entropy per particle, etc.) can be derived through the usual formulas

$$u = \frac{\partial}{\partial \beta}(\beta f), \quad c_V = -\beta^2 \frac{\partial u}{\partial \beta}, \quad s = \beta^2 \frac{\partial f}{\partial \beta} = \beta(u - f).$$

In fact, for the previous formulas to make sense we must first normalize the Hamiltonian (2.7) so that the average energy per spin tends to a finite constant in the thermodynamic limit $N \rightarrow \infty$. The average energy $\langle \mathcal{H} \rangle = (m+n)^{-N} \text{tr} \mathcal{H}$ of the Hamiltonian (2.7) is easily computed by taking into account the identities

$$\text{tr} S_{ij} = \text{tr}(\tilde{S}_{ij}) = (m+n)^{N-2} (m-n),$$

whence

$$\langle \mathcal{H} \rangle = \left(1 + \frac{n-m}{(m+n)^2} \right) \sum_{i < j} \sin^{-2}(\pi(i-j)/N) = \frac{1}{6} N(N^2 - 1) \left(1 + \frac{n-m}{(m+n)^2} \right)$$

(see Ref. [32] for the evaluation of the sum). Note that this is twice the value of the average energy of the $\text{su}(m|n)$ HS chain [13]. Thus in this section we shall take

$$\mathcal{H} = \frac{J}{4N^2} \sum_{1 \leq i < j \leq N} \left(\frac{1 - S_{ij}}{\sin^2(\theta_i - \theta_j)} + \frac{1 - \tilde{S}_{ij}}{\cos^2(\theta_i - \theta_j)} \right), \quad (5.2)$$

whose average energy per spin $\langle \mathcal{H} \rangle / N$ tends to a constant in the thermodynamic limit. Note that the constant J , which sets the energy scale, could be of either sign.

In practice, except in the case of even m and n that we shall discuss below, the complexity of the expressions for \mathcal{Z}_N makes it unfeasible to compute the thermodynamic




Figure 1. Thermodynamic functions of the $\text{su}(m|n)$ chains (5.2) with $m+n=4$ (the $\text{su}(0|4)$ chain has been omitted since, as we shall show below, it is thermodynamically equivalent to the $\text{su}(4|0)$ chain). The color code is as indicated in the plot of f (top left), and in all plots the temperature is measured in units of J . The thermodynamic functions of the $\text{su}(3|1)$ and $\text{su}(1|3)$ chains, for whose $N \rightarrow \infty$ limit there is no known closed-form expression, have been computed using the exact partition function for $N=14$ spins.

free energy in closed form. However, even in these cases it is possible to obtain an approximation of the thermodynamic free energy through the formula

$$f(T) \simeq f_N(T) = -\frac{T}{N} \log \mathcal{Z}_N(q) \quad (5.3)$$

with a sufficiently large N . For instance, in Fig. 1 we present the plots of the thermodynamic functions of the $\text{su}(3|1)$ and $\text{su}(1|3)$ chains computed from Eq. (5.3) with $N=14$, compared to their analogues for the $\text{su}(2|2)$ and $\text{su}(4|0)$ chains in the thermodynamic limit (cf. Eqs. (5.14) and (5.11)–(5.13)). In particular, this and similar plots suggest that in the thermodynamic limit the thermodynamics of the $\text{su}(3|1)$ and $\text{su}(1|3)$ chains are independent of ε_F .

Remark 7. From the duality relation (4.6) it is immediate to obtain a relation between the thermodynamic functions of the $\text{su}(m|n)$ and $\text{su}(n|m)$ chains with opposite values of the parameter J . Indeed, we can rewrite Eq. (4.7) more explicitly as

$$\mathcal{Z}_N^{(m|n)}(T; -J) = e^{J\beta\varepsilon_0/N^2} \mathcal{Z}_N^{(n|m)}(T; J),$$

where the factor of J/N^2 in the exponential is due to the new normalization (5.2) of the

chain's Hamiltonian. From Eqs. (4.5) and (5.1) we then obtain

$$f^{(m|n)}(T; -J) = f^{(n|m)}(T; J) - \frac{J}{3} \lim_{N \rightarrow \infty} \left(1 - \frac{1}{N^2} \right) = f^{(n|m)}(T; J) - \frac{J}{3}. \quad (5.4)$$

For this reason, in what follows we shall restrict ourselves without loss of generality to the case $J > 0$. In particular, for $m = n$ the previous formula becomes

$$f^{(m|m)}(T; -J) = f^{(m|m)}(T; J) - \frac{J}{3}, \quad (5.5)$$

from which it easily follows that

$$u^{(m|m)}(T; -J) = u^{(m|m)}(T; J) - \frac{J}{3}, \quad (5.6)$$

$$c_V^{(m|m)}(T; -J) = c_V^{(m|m)}(T; J), \quad (5.7)$$

$$s^{(m|m)}(T; -J) = s^{(m|m)}(T; J). \quad (5.8)$$

Hence in this case the $J > 0$ and $J < 0$ cases are actually equivalent. \square

As mentioned above, when m and n are both even the factorization (3.20) of the partition function makes it possible to express the thermodynamic free energy per spin $f^{(m|n)}$ of the $\text{su}(m|n)$ chain (5.2) in terms of the free energies per spin of the $\text{su}(1|1)$ and $\text{su}(\frac{m}{2}|\frac{n}{2})$ Haldane–Shastry chains as

$$f^{(m|n)}(T) = f_{\text{HS}}^{(1|1)}(T) + f_{\text{HS}}^{(\frac{m}{2}|\frac{n}{2})}(T). \quad (5.9)$$

In fact, a closed form expression for the free energy per spin $f_{\text{HS}}^{(p|q)}(T)$ of the $\text{su}(p|q)$ HS chain was developed in Ref. [19], namely

$$f_{\text{HS}}^{(p|q)}(T) = -T \int_0^1 \log \lambda^{(p|q)}(x) dx,$$

where $\lambda^{(p|q)}(x)$ is the Perron–Frobenius (i.e., largest in modulus) eigenvalue of the transfer matrix $A^{(p|q)}(x)$ of order $m + n$ introduced in Ref. [19]. More precisely, the matrix elements of $A^{(p|q)}(x)$ are defined by

$$A_{\alpha\gamma}^{(p|q)}(x) = e^{-\beta J \rho(x) \delta(\alpha, \gamma)}, \quad 1 \leq \alpha, \gamma \leq m + n,$$

where

$$\rho(x) = x(1 - x)$$

is the continuous version of the dispersion relation $\mathcal{E}(i)/N^2$ (with $i/N \rightarrow x$) and

$$\delta(\alpha, \gamma) = \begin{cases} 0, & \alpha < \gamma \text{ or } \alpha = \gamma \in B, \\ 1, & \alpha > \gamma \text{ or } \alpha = \gamma \in F. \end{cases}$$

For instance, for $p = q = 1$ we have

$$A^{(1|1)}(x) = \begin{pmatrix} 1 & 1 \\ e^{-\beta J \rho(x)} & e^{-\beta J \rho(x)} \end{pmatrix}$$

so that

$$\lambda^{(1|1)}(x) = 1 + e^{-\beta J \rho(x)}.$$

On the other hand, the Perron–Frobenius eigenvalue for the case $q = 0$ was found in Ref. [34] to be

$$\lambda^{(p|0)}(x) = \frac{1 - e^{-\beta J \rho(x)}}{1 - e^{-\beta J \rho(x)/p}} = e^{(1-p)\beta J \rho(x)/(2p)} \frac{\sinh(\beta J \rho(x)/2)}{\sinh(\beta J \rho(x)/(2p))}.$$

Thus when m is even the free energy of the chain (5.2) in the non-supersymmetric case $n = 0$ is explicitly given by

$$f^{(m|0)}(T) = \frac{J}{6} \left(1 - \frac{1}{m}\right) - T \int_0^1 \log \left(\frac{\sinh(\beta J \rho(x))}{\sinh(\beta J \rho(x)/m)} \right) dx. \quad (5.10)$$

Note also that, since the integral in Eq. (5.10) does not depend on the sign of J , for even m we have

$$\begin{aligned} f^{(m|0)}(T; -J) &= -\frac{J}{3} \left(1 - \frac{1}{m}\right) + f^{(m|0)}(T; J), \\ u^{(m|0)}(T; -J) &= -\frac{J}{3} \left(1 - \frac{1}{m}\right) + u^{(m|0)}(T; J), \\ c_V^{(m|0)}(T; -J) &= c_V^{(m|0)}(T; J), \quad s^{(m|0)}(T; -J) = s^{(m|0)}(T; J). \end{aligned}$$

In particular combining the previous equation for $f^{(m|0)}(T; -J)$ with the duality relation (5.4) we deduce that

$$f^{(0|m)}(T; J) = f^{(m|0)}(T; -J) + \frac{J}{3} = f^{(m|0)}(T; J) + \frac{J}{3m}.$$

Hence the $\text{su}(m|0)$ and $\text{su}(0|m)$ cases are thermodynamically equivalent.

Interestingly, the free energy (5.10) of the $\text{su}(m|0)$ chain with even m coincides with the free energy of the ordinary (bosonic) $\text{su}(m|0)$ HS chain with J replaced by $2J$. In other words, both models are equivalent in the thermodynamic limit up to a suitable rescaling of the coupling constant J . By the remark preceding Eq. (2.9) in Section 2, we thus obtain in this case the same thermodynamics as if we replaced the spin flip operators S_i by plus or minus the identity. For the sake of completeness, we list below the explicit formulas for the internal energy, specific heat at constant volume and entropy per particle of the $\text{su}(m|0)$ chain with even m (the corresponding formulas for the $\text{su}(0|n)$ chain with even n are similar and shall therefore be omitted):

$$u^{(m|0)}(T) = \frac{J}{6} \left(1 - \frac{1}{m}\right) - J \int_0^1 \rho(x) \left[\coth(\beta J \rho(x)) - \frac{1}{m} \coth(\beta J \rho(x)/m) \right] dx, \quad (5.11)$$

$$c_V^{(m|0)}(T) = \beta^2 J^2 \int_0^1 \rho^2(x) \left[\frac{1}{m^2} \text{csch}^2(\beta J \rho(x)/m) - \text{csch}^2(\beta J \rho(x)) \right] dx, \quad (5.12)$$

$$\begin{aligned} s^{(m|0)}(T) &= \int_0^1 \left[\log \left(\frac{\sinh(\beta J \rho(x))}{\sinh(\beta J \rho(x)/m)} \right) \right. \\ &\quad \left. - \beta J \rho(x) \left(\coth(\beta J \rho(x)) - \frac{1}{m} \coth(\beta J \rho(x)/m) \right) \right] dx. \end{aligned} \quad (5.13)$$




Figure 2. Thermodynamic functions of the $\text{su}(2|0)$ chain (5.2) with $N = 20$ spins (dashed red line) and its thermodynamic limit (continuous blue line). In all plots the temperature is measured in units of J , and the insets show the low temperature behavior of each thermodynamic function. The significance of the continuous green lines in the insets of the plots of the free energy (top left) and the entropy (bottom left) is explained in the body of the article.

In Fig. 2 we plot the thermodynamic functions of the $\text{su}(2|0)$ chain for $N = 20$ spins, compared to their thermodynamic limits given by Eqs. (5.10)-(5.13). We see that the agreement between both plots is excellent even for this relatively low number of spins, except at very low temperatures (which in our units corresponds to $T \lesssim 2 \cdot 10^{-2}$). This low temperature discrepancy is, in fact, an inevitable finite size effect. Indeed, at very low temperatures we have

$$\mathcal{Z}_N(T) \simeq d_{\text{GS}} e^{-\beta E_{\text{GS}}} + d_1 e^{-\beta E_1},$$

where E_{GS} and $E_1 = E_{\text{GS}} + \Delta E$ denote respectively the energies of the ground state and the first excited state, and d_{GS} and d_1 are their degeneracies. Thus for finite N the chain's thermodynamic functions behave at low temperatures as

$$\begin{aligned} f_N(T) &\simeq \frac{E_{\text{GS}}}{N} - \frac{\log d_{\text{GS}}}{N} T - \frac{d_1}{d_{\text{GS}}} \frac{T}{N} e^{-\beta \Delta E}, \\ u_N(T) &\simeq \frac{E_{\text{GS}}}{N} + \frac{\Delta E}{N} \frac{d_1}{d_{\text{GS}}} e^{-\beta \Delta E}, \\ c_{V,N}(T) &\simeq \frac{d_1}{d_{\text{GS}}} \frac{(\Delta E)^2}{NT^2} e^{-\beta \Delta E}, \end{aligned}$$

$$s_N(T) \simeq \frac{\log d_{\text{GS}}}{N} + \frac{d_1}{d_{\text{GS}}} \frac{\Delta E}{NT} e^{-\beta \Delta E}.$$

In our case ($m = 2, n = 0$) we have $E_{\text{GS}} = 0$ and $d_{\text{GS}} = 2$, so that at low temperatures the free energy for finite N is approximately a linear function of the temperature with slope equal to $-(\log 2)/N$. This is corroborated by the inset in the upper left corner of Fig. 2, where the line $f = -(\log 2)T/20$ has been plotted in green. It is also clear from the previous equations that the remaining thermodynamic functions differ from their zero temperature values by exponentially small terms of the form $T^{-k}e^{-\beta \Delta E}$ for suitable $k \in \{0, 1, 2\}$; this behavior is again apparent from the corresponding insets in Fig. 2 (note that $\Delta E = 19$ and $d_1 = 4$ for $N = 20$). In particular, the zero temperature value of the entropy for a finite number of spins N is

$$s_N(0) = \frac{\log 2}{N} \neq 0;$$

this can be seen from the bottom left inset of Fig. 2, where we have plotted in green the horizontal line $s = (\log 2)/20 \simeq 0.0346574$.

Apart from the previous non-supersymmetric cases, it is also possible to evaluate the thermodynamic functions of the chain (5.2) in a few genuinely supersymmetric cases with m and n even using the results of Ref. [19]. To begin with, for $m = n = 2$ we have

$$f^{(2|2)}(T; J) = 2f_{\text{HS}}^{(1|1)}(T; J) = -2T \int_0^1 \log(1 + e^{-\beta J \rho(x)}) dx = f_{\text{HS}}^{(2|2)}(T; 2J), \quad (5.14)$$

where we have used the explicit expression for $f_{\text{HS}}^{(2|2)}$ in Ref. [19]. Thus the thermodynamics of the $\text{su}(2|2)$ chain is equivalent to that of its HS counterpart with J replaced by $2J$, obtained by setting $S_i = 1$ in Eq. (5.2). Likewise, for $m = 4, n = 2$, using the expression of $f_{\text{HS}}^{(2|1)}$ in Ref. [19] we obtain

$$\begin{aligned} f^{(4|2)}(T) &= f_{\text{HS}}^{(1|1)}(T) + f_{\text{HS}}^{(2|1)}(T) \\ &= -T \int_0^1 \log(1 + e^{-\beta J \rho(x)}) dx \\ &\quad - T \int_0^1 \log \left[1 + \frac{1}{2} e^{-\beta J \rho(x)} + \frac{1}{2} e^{-\beta J \rho(x)} \sqrt{1 + 8e^{\beta J \rho(x)}} \right] dx. \end{aligned}$$

Although it is straightforward to obtain the energy, specific heat at constant volume and entropy from the previous formula, the resulting expressions are somewhat lengthy and shall therefore be omitted. The free energy of the $\text{su}(2|4)$ case is easily obtained from the latter equation for $f^{(4|2)}$ using the duality relation (5.4):

$$\begin{aligned} f^{(2|4)}(T) &= -T \int_0^1 \log(1 + e^{-\beta J \rho(x)}) dx \\ &\quad - T \int_0^1 \log \left[\frac{1}{2} + e^{-\beta J \rho(x)} + \frac{1}{2} \sqrt{1 + 8e^{-\beta J \rho(x)}} \right] dx. \end{aligned} \quad (5.15)$$

Finally, the thermodynamic free energy per particle for the case $m = n = 4$ can also be computed in closed form using the results of Ref. [19], namely

$$\begin{aligned} f^{(4|4)}(T) &= f_{\text{HS}}^{(1|1)}(T) + f_{\text{HS}}^{(2|2)}(T) \\ &= -T \int_0^1 \log(1 + e^{-\beta J \rho(x)}) dx - 2T \int_0^1 \log(1 + e^{-\beta J \rho(x)/2}) dx \end{aligned}$$




Figure 3. Thermodynamic functions of the $\text{su}(2|2)$, $\text{su}(2|4)$, $\text{su}(4|2)$ and $\text{su}(4|4)$ chains (5.2), whose thermodynamic functions have closed-form expressions in the thermodynamic limit (color code in top left figure). In all plots the temperature is measured in units of J . The thermodynamic functions of the non-supersymmetric $\text{su}(2|0)$ chain are also shown for comparison purposes.

$$\begin{aligned}
 &= \frac{J}{6} - T \int_0^1 \log \left(8 \cosh \left(\frac{\beta J \rho(x)}{2} \right) \cosh^2 \left(\frac{\beta J \rho(x)}{4} \right) \right) dx \\
 &\equiv f_{\text{HS}}^{(1|1)}(T) + f_{\text{HS}}^{(1|1)}(2T).
 \end{aligned}$$

In this case the expressions for the remaining thermodynamic functions are relatively simple, to wit

$$\begin{aligned}
 u^{(4|4)}(T) &= J \int_0^1 \left(\frac{1}{1 + e^{\beta J \rho(x)}} + \frac{1}{1 + e^{\beta J \rho(x)/2}} \right) \rho(x) dx, \\
 &= \frac{J}{6} - \frac{J}{2} \int_0^1 \left[\tanh \left(\frac{\beta J \rho(x)}{2} \right) + \tanh \left(\frac{\beta J \rho(x)}{4} \right) \right] \rho(x) dx, \\
 c_V^{(4|4)}(T) &= \frac{\beta^2 J^2}{8} \int_0^1 \left[2 \text{sech}^2 \left(\frac{\beta J \rho(x)}{2} \right) + \text{sech}^2 \left(\frac{\beta J \rho(x)}{4} \right) \right] \rho^2(x) dx, \\
 s^{(4|4)}(T) &= \int_0^1 \left[\log \left(8 \cosh \left(\frac{\beta J \rho(x)}{2} \right) \cosh^2 \left(\frac{\beta J \rho(x)}{4} \right) \right) \right. \\
 &\quad \left. - \frac{\beta J \rho(x)}{2} \left(\tanh \left(\frac{\beta J \rho(x)}{2} \right) + \tanh \left(\frac{\beta J \rho(x)}{4} \right) \right) \right] dx.
 \end{aligned}$$

In particular, note that the previous formulas are obviously consistent with the general relations (5.5)–(5.8). See Fig. 3 for a plot of the main thermodynamic functions for the truly supersymmetric cases in which these functions have the closed-form expressions




Figure 4. Free energy per spin of the $\text{su}(1|1)$ (with $\varepsilon = -1$), $\text{su}(2|1)$, $\text{su}(1|2)$, $\text{su}(3|0)$, $\text{su}(3|1)$ and $\text{su}(1|3)$ chains (5.2) (with $\varepsilon_F = 1$ in the last two cases) for $N = 14$ spins (solid lines) vs. the temperature T (in units of J). The dashed lines represent the thermodynamic free energy per spin of the $\text{su}(1|1)$, $\text{su}(2|1)$, $\text{su}(1|2)$, and $\text{su}(3|0)$ HS chains, as well as the free energy per spin of the $\text{su}(3|1)$ and $\text{su}(1|3)$ HS chains (whose thermodynamic limit is not explicitly known) for $N = 14$ spins.

listed above.

Remark 8. We have noted above that the thermodynamic free energy per spin of the $\text{su}(2|2)$, $\text{su}(m|0)$, and $\text{su}(0|m)$ (with even m) chains (5.2) coincides with the free energy of their HS counterparts with the coupling J rescaled to $2J$. Equivalently, by Remark 3, the free energy per spin of the latter chains does not change if we replace the spin flip operators S_i by (plus or minus) the identity. It is also apparent from Fig. 1 that the thermodynamic free energies per spin of the $\text{su}(3|1)$ and $\text{su}(1|3)$ chains with $N = 14$ spins are practically independent of ε_F . These facts suggest that, in general, in the thermodynamic limit the free energy per spin of the $\text{su}(m|n)$ chains (5.2) with arbitrary m and n is independent of the representation of the spin reversal operators chosen. In particular, if this conjecture holds the free energy of the chain (5.2) should coincide with the free energy of its HS counterpart with J replaced by $2J$, obtained when S_i is set to (plus or minus) the identity in Eq. (5.2). Although, as remarked above, the thermodynamic free energy per spin of the $\text{su}(m|n)$ supersymmetric HS chain is not known exactly except for relatively low values of m and n , we have numerically verified the latter conjecture in a few more cases by comparing the free energy per spin of both the chain (5.2) and its HS counterpart (with J replaced to $2J$) for a finite number of spins; see, e.g., Fig. 4. \square

It is apparent from Figs. 1-3 that the qualitative behavior of the thermodynamic functions of all the chains discussed in this work is very similar. In particular, the specific heat per spin of all of these models features a single Schottky peak, typical of many k -level systems. In the $\text{su}(m|0)$ —or, equivalently, $\text{su}(n|0)$ — case this fact can be explained by noting that the free energy (5.10) of the $\text{su}(m|0)$ chain can be written as

$$f^{(m|0)}(T) = -T \int_0^1 \log \left(\sum_{k=0}^{m-1} e^{-\frac{2kJ\beta\rho(x)}{m}} \right) dx.$$

As noted in Ref [34], replacing $\rho(x)$ by its mean value over the interval $[0, 1]$, $1/6$,

this partition function reduces to that of an m -level system with equally spaced, non-degenerate levels with energies

$$E_i = \frac{iJ}{3m}, \quad i = 0, \dots, m-1.$$

This suggests that the specific heat of the latter m -level system should qualitatively behave as the specific heat per spin of the $\text{su}(m|0)$ chain (5.2). That this is indeed the case can be seen from Fig. 5. This result also follows from the discussion of Ref. [34], since as noted above the thermodynamic functions of the $\text{su}(m|0)$ chain (5.2) coincide with those of the ferromagnetic $\text{su}(m)$ HS chain up to a trivial rescaling of the coupling J . More interestingly, a similar argument can be applied to the $\text{su}(2|2)$ and $\text{su}(4|4)$ chains. Indeed, in the former case the thermodynamic free energy per spin can be expressed as

$$f^{(2|2)}(T) = -T \int_0^1 \log(1 + 2e^{-\beta J \rho(x)} + e^{-2\beta J \rho(x)}) dx,$$

which is expected to behave qualitatively as the free energy of a 3-level system with energies $E_k = kJ/6$ (with $k = 0, 1, 2$) and degeneracies $g_0 = g_2 = 1, g_1 = 2$. The specific heat of the this three-level system, namely

$$c_V = \frac{\beta^2 J^2}{72} \text{sech}^2(\beta J/12),$$

is again in reasonable agreement with that of the $\text{su}(2|2)$ chain (5.2) (cf. Fig. 5). In particular, the temperature of the Schottky peak of the three-level system, $T_m \simeq 0.0695 J$, is of the same order of magnitude as the corresponding temperature for the $\text{su}^{(2|2)}$ chain (5.2), $T_m^{(2|2)} \simeq 0.0809 J$. Similarly, expressing the free energy per spin of the $\text{su}(4|4)$ chain as

$$\begin{aligned} f^{(4|4)}(T) &= -T \int_0^1 \log \left[(1 + e^{-\beta J \rho(x)}) (1 + e^{-\beta J \rho(x)/2})^2 \right] dx \\ &= -T \int_0^1 \log \left(1 + 2e^{-\beta J \rho(x)/2} + 2e^{-\beta J \rho(x)} + 2e^{-3\beta J \rho(x)/2} + e^{-2\beta J \rho(x)} \right) dx \end{aligned}$$

and replacing again $\rho(x)$ by its average over the interval $[0, 1]$ we obtain a 5-level system with energies $E_k = kJ/12$ (with $k = 0, \dots, 4$) and degeneracies $g_0 = g_4 = 1, g_1 = g_2 = g_3 = 2$, whose specific heat is given by

$$c_V = \frac{\beta^2}{72} e^{-\beta J/12} \frac{1 + 2e^{-\beta J/12} + 2e^{-\beta J/6} + 2e^{-\beta J/4} + e^{-\beta J/3}}{(1 + e^{-\beta J/12})^2 (1 + e^{-\beta J/6})^2}.$$

This specific heat provides again a reasonable approximation to the specific heat per spin $c_V^{(4|4)}$, particularly at high temperatures (cf. Fig. 5). Again, the temperature of the Schottky peak of the 5-level system, given by $T_m \simeq 0.0443 J$, is in excellent agreement with the analogous temperature $T_m^{(4|4)} \simeq 0.0491 J$ for the $\text{su}(4|4)$ chain.

The situation is somewhat murkier for the $\text{su}(2|4)$ and $\text{su}(4|2)$ chains, whose thermodynamic free energy does has a more complicated structure than its $\text{su}(m|0)$, $\text{su}(1|1)$, $\text{su}(2|2)$ and $\text{su}(4|4)$ counterparts. In the former case, at low temperatures we




Figure 5. Specific heat per spin of the $\text{su}(2|2)$, $\text{su}(4|4)$, and $\text{su}(2|4)$ chains (5.2) (solid curves), compared to their k -level counterparts (dashed curves). In all plots the temperature is measured in units of J .

can approximate the thermodynamic free energy as

$$\begin{aligned} f^{(2|4)}(T) &\sim -T \int_0^1 \log \left[(1 + e^{-\beta J \rho(x)})(1 + 3e^{-\beta J \rho(x)}) \right] dx \\ &= -T \int_0^1 \log \left(1 + 4e^{-\beta J \rho(x)} + 3e^{-2\beta J \rho(x)} \right) dx, \end{aligned}$$

whose associated 3-level system has energies $iJ/6$ (with $i = 0, 1, 2$) and degeneracies $g_0 = 1$, $g_1 = 4$, $g_2 = 3$. The specific heat of this 3-level system,

$$c_V = \frac{\beta^2}{9} e^{-2\beta J/3} \frac{1 + 3e^{-\beta J/6} + 3e^{-\beta J/3}}{(1 + 4e^{-\beta J/6} + 3e^{-\beta J/3})^2},$$

should therefore provide a rough approximation to the specific heat per spin $c_V^{(2|4)}$. Although in this case the agreement between both specific heats is noticeably poorer than in the previous cases, the temperature of the Schottky peak of the three-level model, $T_m \simeq 0.0611J$, is remarkably close to its counterpart $T_m^{(2|4)} \simeq 0.0684J$.

6. Critical behavior

In this section we shall briefly analyze the critical behavior of the supersymmetric chains (5.2). As is well known, one of the characteristic features of a conformal field theory (CFT) is that its ground state (i.e., the vacuum) is non-degenerate, and that its spectrum contains a subset of low-lying gapless excitations above the ground state with a linear energy-momentum dispersion relation. These two fundamental properties are thus a necessary condition for a finite-dimensional quantum system to be critical, i.e., for its low-energy sector to be described by a CFT in the thermodynamic limit. More precisely, the ground state of such a system must have at most a finite degeneracy¶, and its low-energy excitations must be gapless and feature a linear energy-momentum

¶ By “finite” we mean independent of the number of particles in the thermodynamic limit. If the ground state of a critical system is g times degenerate, its low-energy excitations will be described by g identical copies of a single CFT.

relation. We shall therefore start by identifying in what cases the chain (5.2) satisfies these necessary conditions for criticality.

Remark 9. Since (as noted in the previous section) we can assume without loss of generality that the coupling J is positive, and the criticality properties of the chain (5.2) are obviously independent of the size of J , for the sake of conciseness we shall set $J = 1$ throughout this section. \square

To begin with, we shall analyze for what values of m and n the ground state of the $\text{su}(m|n)$ chain (5.2) has a finite degeneracy. For $m \neq 0$, this problem can be solved using the explicit formulas for the ground state degeneracy g of the ground state. From these formulas it is straightforward to show that, as is the case with the ordinary supersymmetric HS chain, this degeneracy cannot remain finite as $N \rightarrow \infty$ unless $m = 1, 2$. Indeed, for $m > 2$ and n both even by Eq. (3.24) we have

$$g > 2 \sum_{k=0}^N (k+1) \binom{\frac{n}{2}}{N-k} \geq 2(N+1).$$

Similarly, for $m > 2$ even and n odd Eq. (3.31) implies that

$$g > \sum_{k=0}^N (k+1) \left[\binom{\frac{1}{2}(n-1)}{N-k} + \binom{\frac{1}{2}(n+1)}{N-k} \right] \geq 2(N+1).$$

The case $m > 2$ odd and n even is analogous, namely (by Eq. (3.34))

$$g > \sum_{k=0}^N (k+2) \binom{\frac{n}{2}}{N-k} \geq N+2.$$

Finally, when $m > 2$ and n are both odd Eq. (3.39) yields

$$g > \sum_{k=0}^N \left[(k+1) \binom{\frac{1}{2}(n+\varepsilon_F)}{N-k} + \binom{\frac{1}{2}(n-\varepsilon_F)}{N-k} \right] \geq N+2.$$

Let us next study the ground state degeneracy for the cases $m = 1, 2$. For $m = 2$, this degeneracy is given by

$$g = \begin{cases} 2 \sum_{k=0}^N \binom{\frac{n}{2}}{k} \leq 2^{\frac{n}{2}+1}, & n \text{ even} \\ \sum_{k=0}^N \left[\binom{\frac{1}{2}(n-1)}{k} + \binom{\frac{1}{2}(n+1)}{k} \right] \leq 3 \cdot 2^{\frac{1}{2}(n-1)}, & n \text{ odd,} \end{cases} \quad (6.1)$$

which is indeed finite (note that the equality holds for $N \geq n/2$ if n is even or $N \geq (n-1)/2$ if n is odd). The case $m = 1$ is dealt with analogously. Indeed,

$$g = \begin{cases} \binom{\frac{n}{2}}{N} + \sum_{k=0}^N \binom{\frac{n}{2}}{k} \leq \binom{\frac{n}{2}}{N} + 2^{\frac{n}{2}}, & n \text{ even} \\ \binom{\frac{1}{2}(n-\varepsilon_F)}{N} + \sum_{k=0}^N \binom{\frac{1}{2}(n+\varepsilon_F)}{k} \leq \binom{\frac{1}{2}(n-\varepsilon_F)}{N} + 2^{\frac{1}{2}(n+\varepsilon_F)}, & n \text{ odd,} \end{cases}$$

which is finite (note that in this case the ground state degeneracy is simply $2^{\frac{n}{2}}$ for n even and $2^{\frac{1}{2}(n+\varepsilon_F)}$ for n odd, if N is respectively greater than $n/2$ or $(n-\varepsilon_F)/2$).

From the previous analysis it follows that the chain (5.2) can only be critical for $m = 0, 1, 2$. The determination of the ground state degeneracy of the $\text{su}(0|n)$ chain, or of the existence of low-lying levels with a linear energy-momentum relation in all of the latter cases, can be carried out with the results of the previous sections only when m and n are both even, since only in this case the partition function can be expressed in terms of the partition functions of the $\text{su}(1|1)$ and $\text{su}(\frac{m}{2}|\frac{n}{2})$ HS chains. In particular, the criticality of the $m = 1$ case —more precisely, the existence of low-lying levels with a linear energy-momentum dispersion relation— cannot be ascertained with the techniques of this paper, and thus remains an open problem.

When m and n are both even, the equivalence of the chain Hamiltonian (5.2) to the sum of two non-interacting $\text{su}(1|1)$ and $\text{su}(\frac{m}{2}|\frac{n}{2})$ HS chains (cf. Eq.(3.21)) entails that the energy spectrum of our model can be expressed in terms of $\text{su}(1|1)$ and $\text{su}(\frac{m}{2}|\frac{n}{2})$ bond vectors and their corresponding (supersymmetric) motifs [14]. More precisely, the spectrum of the chain (5.2) with m and n even can be generated from the formula

$$E(\boldsymbol{\sigma}, \boldsymbol{\sigma}') = \frac{1}{N^2} \sum_{i=1}^{N-1} [\delta(\sigma_i, \sigma_{i+1}) + \delta'(\sigma'_i, \sigma'_{i+1})] i(N-i), \quad (6.2)$$

where $\boldsymbol{\sigma} \in \{1, 2\}^N$, $\boldsymbol{\sigma}' \in \{1, \dots, (m+n)/2\}^N$ are respectively $\text{su}(1|1)$ and $\text{su}(\frac{m}{2}|\frac{n}{2})$ bond vectors, and the functions δ and δ' are defined by

$$\delta(j, k) = \begin{cases} 0, & j < k \text{ or } j = k = 1 \\ 1, & j > k \text{ or } j = k = 2 \end{cases}$$

and

$$\delta'(j, k) = \begin{cases} 0, & j < k \text{ or } j = k \in \{1, \dots, \frac{m}{2}\} \\ 1, & j > k \text{ or } j = k \in \{\frac{m}{2} + 1, \dots, \frac{m+n}{2}\}. \end{cases}$$

Equation (6.2) implies that the ground state of the supersymmetric chain (5.2) with even m and n is obtained from the bond vectors $\boldsymbol{\sigma}$ and $\boldsymbol{\sigma}'$ yielding respectively the ground states of the $\text{su}(1|1)$ and $\text{su}(\frac{m}{2}|\frac{n}{2})$ HS chains. In particular, the degeneracy of this ground state is the product of the degeneracies of the ground states of the $\text{su}(1|1)$ and $\text{su}(\frac{m}{2}|\frac{n}{2})$ chains. Since the ground state of the $\text{su}(1|1)$ HS chain is obviously obtained from the two bond vectors $(1, \dots, 1, s)$ with $s = 1, 2$, the previous observation entails that the ground state degeneracy of the $\text{su}(m|n)$ chain (5.2) with even m and n is twice the degeneracy of the ground state of the $\text{su}(\frac{m}{2}|\frac{n}{2})$ HS chain. For $m = 2$ and $N \geq n/2$, the ground state of the latter chain is the zero mode obtained from an $\text{su}(1|\frac{n}{2})$ bond vector of the form

$$\underbrace{(1, \dots, 1)}_{N-k}, s_1, \dots, s_k,$$

with $2 \leq s_1 < \dots < s_k \leq n/2$ and $k \leq n/2$. Since the number of such bond vectors is $2^{n/2}$, the ground state degeneracy of the $\text{su}(2|n)$ chain (5.2) with even n and $N \geq n/2$ is $2^{\frac{n}{2}+1}$, in agreement with the remark after Eq. (6.1). Moreover, it is known [19, 26] that both the $\text{su}(1|1)$ and the $\text{su}(1|\frac{n}{2})$ chains have low energy excitations obtained by slightly

varying the $\text{su}(1|1)$ and $\text{su}(1|\frac{n}{2})$ bond vectors near their ends. The typical energy of one of these excitations is thus

$$\Delta E = \frac{i}{N^2}(N - i),$$

with $i \ll N$, and its momentum is also known to be $\Delta p = \pm 2\pi i/N$ [26]. Thus the Fermi velocity of the low energy excitations of both the $\text{su}(1|1)$ and $\text{su}(1|\frac{n}{2})$ HS chains is

$$v_F = \lim_{N \rightarrow \infty} \left| \frac{\Delta E}{\Delta p} \right| = \frac{1}{2\pi}.$$

Combining these excitations we obviously obtain low energy excitations above the ground state of the $\text{su}(2|n)$ chain (5.2) with a finite Fermi velocity $1/(2\pi)$, so that this chain does fulfill the two necessary conditions for criticality mentioned at the beginning of this section. In fact, it is shown in Ref. [26] that in the thermodynamic limit the low energy excitations of the $\text{su}(1|p)$ HS chains are described by a CFT of p non-interacting massless Majorana fermion fields. It follows that the $\text{su}(2|n)$ chain (5.2) is indeed critical, with central charge

$$c^{(2|n)} = \frac{1}{2} + \frac{n}{4} = \frac{1}{4}(n + 2). \quad (6.3)$$

A similar analysis can be performed for the $\text{su}(0|n)$ chain (5.2) with even n . The main difference is that in this case the $\text{su}(0|\frac{n}{2})$ HS chain contains no bosons, and thus (for $N > n/2$) its ground state has positive energy. As explained in Ref. [34], the ground state in this case is obtained from $\text{su}(0|\frac{n}{2})$ bond vectors of the form

$$(1, 2, \dots, \frac{n}{2}, \dots, 1, 2, \dots, \frac{n}{2}, s_1, \dots, s_{N-rn/2})$$

or their reflected analogues

$$(s_1, \dots, s_{N-rn/2}, 1, 2, \dots, \frac{n}{2}, \dots, 1, 2, \dots, \frac{n}{2}),$$

where $r = \lfloor 2N/n \rfloor$ and

$$1 \leq s_1 < \dots < s_{N-rn/2} \leq \frac{n}{2}.$$

Hence the ground state degeneracy of the $\text{su}(0|\frac{n}{2})$ HS chain is equal to

$$\begin{cases} 1, & 2N = nr \\ 2 \binom{\frac{n}{2}}{N - \frac{rn}{2}}, & 2N > nr, \end{cases}$$

and the degeneracy of the ground state of the $\text{su}(0|\frac{n}{2})$ chain (5.2) is twice the above number. In particular, the ground state degeneracy of the latter model remains finite as $N \rightarrow \infty$. Furthermore, the $\text{su}(0|\frac{n}{2})$ HS chain is also known to possess low energy excitations with finite Fermi velocity $v_F = 1/(2\pi)$, described by the $\text{su}(n/2)_1$ WZNW model with central charge $(n/2) - 1$ [26]. Reasoning as above we deduce that the $\text{su}(0|n)$ chain (5.2) is critical, with central charge

$$c^{(0|n)} = \frac{1}{2} + \frac{n}{2} - 1 = \frac{1}{2}(n - 1). \quad (6.4)$$

Summarizing, we have just shown that the $\text{su}(0|n)$ and $\text{su}(2|n)$ chains (5.2) with even n are critical, while the $\text{su}(m|n)$ chains with $m > 2$ are not. Moreover, the $\text{su}(1|n)$ chains with arbitrary n and the $\text{su}(2|n)$ chains with odd n have been shown to have a finite ground state degeneracy, and could thus be critical. On the other hand, with the results of this paper nothing can be concluded about the $\text{su}(0|n)$ chains with odd n , although our numerical calculations suggest that these chains have a non-degenerate ground state and therefore could be critical. We briefly sum up the above results in Table 1.

(m, n)	Finite GS degeneracy	c	Associated CFT
$(0, 2p)$	✓	$p - \frac{1}{2}$	1 MMF + $\text{su}(p)_1$ WNZW
$(0, 2p - 1)$	*	?	?
$(1, n)$	✓	?	?
$(2, 2p)$	✓	$\frac{1}{2}(p + 1)$	$p + 1$ MMFs
$(2, 2p - 1)$	✓	?	?

Table 1. Summary of results on the critical character of the $\text{su}(m|n)$ supersymmetric chain (2.7). Here p and n are positive integers, and the abbreviations GS and MMF stand respectively for “ground state” and “massless Majorana fermion”. The asterisk (*) indicates that the finite degeneracy of the ground state of the $\text{su}(0|2p - 1)$ chain has been verified only for particular values of p and the number of spins N , while the interrogation mark (?) stands for “not known”. As shown above, the chains with $m \geq 3$ have infinite ground state degeneracy in the thermodynamic limit, and cannot thus be critical.

As remarked in the Introduction, at low temperature the free energy of a critical quantum system should have the same asymptotic behavior as the free energy of the CFT describing its low energy sector, given by Eq. (1.1). In particular, from the growth of the free energy of a quantum critical system at low temperatures one can infer the central charge of its associated CFT, and thus identify the system’s universality class. Since the free energy of the $\text{su}(m|n)$ chains (5.2) with $m = 0, 2$ and n even discussed above obeys Eq. (5.9), it should verify Eq. (1.1) with $v_F = 1/(2\pi)$ (the Fermi velocity of the critical supersymmetric HS chains) and central charge

$$c^{(m|n)} = c_{\text{HS}}^{(1|1)} + c_{\text{HS}}^{(\frac{m}{2}|\frac{n}{2})} = \frac{1}{2} + c_{\text{HS}}^{(\frac{m}{2}|\frac{n}{2})}. \quad (6.5)$$

This formula is in fact in full agreement with the result obtained above studying the behavior of the low-lying energy excitations (cf. Eqs. (6.3) and (6.4)).

The low temperature behavior (1.1) of the free energy per spin of the $\text{su}(0|n)$ chain with even n , the $\text{su}(2|2)$ and the $\text{su}(2|4)$ chains (and, in particular, Eq. (6.5) for the central charge) can be explicitly checked using the formulas for their thermodynamic free energy per spin derived in the previous section. Indeed, in all of these cases the free energy can be expressed in terms of integrals of the form

$$I_\beta[\varphi] := \int_0^{1/2} dx \log(\varphi(e^{-\beta\rho(x)})), \quad (6.6)$$

where $\varphi(z)$ is a smooth function such that $\varphi(z) > 0$ for $z > 0$, $\varphi(0) = 1$ and⁺

$$\varphi(z) = 1 + O(z) \quad (6.7)$$

as $z \rightarrow 0+$. Using Eq. (B.2) in Appendix B to approximate this integral, it is straightforward to derive the asymptotic behavior at low temperatures of the thermodynamic free energy per spin of the $\text{su}(0|n)$ (with even n), $\text{su}(2|2)$ and $\text{su}(2|4)$ chains. To begin with, using Eqs. (5.4) and (5.10) (with $J = 1$) the free energy of the $\text{su}(0|n)$ chain (5.2) with n even can be expressed as

$$\begin{aligned} f^{(0|n)}(T) &= \frac{1}{6} \left(1 + \frac{1}{n}\right) - T \int_0^1 dx \log \left(\frac{\sinh(\beta\rho(x))}{\sinh(\beta\rho(x)/n)} \right) \\ &= \frac{1}{3n} - T \int_0^1 dx \log \left(\frac{1 - e^{-2\beta\rho(x)}}{1 - e^{-2\beta\rho(x)/n}} \right). \end{aligned}$$

Taking into account that $\rho(x)$ is symmetric about $x = 1/2$ we can rewrite the previous formula as

$$f^{(0|n)}(T) = f^{(0|n)}(0) - 2T \left(I_\beta[1 - z^2] - I_{\beta/n}[1 - z^2] \right).$$

From Eq. (B.2) we then have

$$\begin{aligned} I_\beta[1 - z^2] &= T \int_0^\infty dy \log(1 - e^{-2y}) + O(T^2) = \frac{T}{2} \int_0^\infty dy \log(1 - e^{-y}) + O(T^2) \\ &= -\frac{T}{2} \zeta(2) + O(T^2) = -\frac{\pi^2 T}{12} + O(T^2), \end{aligned}$$

and therefore

$$f^{(0|n)}(T) - f^{(0|n)}(0) = -(n-1) \frac{\pi^2 T^2}{6} + O(T^3) = -\frac{1}{2}(n-1) \frac{\pi T^2}{6v_F} + O(T^3).$$

Comparing with Eq. (1.1) we conclude that in this case the central charge is given by Eq. (6.4). Likewise, by Eq. (5.14) with $J = 1$ the thermodynamic free energy of the $\text{su}(2|2)$ chain is given by

$$f^{(2|2)}(T) = 2f_{\text{HS}}^{(1|1)}(T) = -4TI_\beta[1 + z],$$

and thus

$$\begin{aligned} f^{(2|2)}(T) &= -4T^2 \int_0^\infty dy \log(1 + e^{-y}) + O(T^3) = -2T^2 \zeta(2) + O(T^3) \\ &= -\frac{\pi^2 T^2}{3} + O(T^3) = -\frac{\pi T^2}{6v_F} + O(T^3). \end{aligned}$$

Hence the central charge of the $\text{su}(2|2)$ chain is $c^{(2|2)} = 1$, in agreement with Eq. (6.3).

Finally, by Eq. (5.15) with $J = 1$ we have

$$f^{(2|4)}(T) = -2TI_\beta[1 + z] - 2TI_\beta\left[\frac{1}{2} + z + \frac{1}{2}\sqrt{1 + 8z}\right],$$

and therefore

$$\begin{aligned} f^{(2|4)}(T) &= -\frac{\pi^2 T^2}{12} - 2T^2 \int_0^\infty dy \log\left(\frac{1}{2} + e^{-y} + \frac{1}{2}\sqrt{1 + 8e^{-y}}\right) + O(T^3) \\ &= -\frac{\pi^2 T^2}{6} - \frac{\pi^2 T^2}{3} + O(T^3) = -\frac{\pi^2 T^2}{2} + O(T^3) = -\frac{3}{2} \frac{\pi T^2}{6v_F} + O(T^3), \end{aligned}$$

⁺ Note that that $z^\alpha = O(z)$ as $z \rightarrow 0+$ for $\alpha \geq 1$.




Figure 6. Thermodynamic free energy per spin of the $\text{su}(0|2)$, $\text{su}(2|2)$ and $\text{su}(2|4)$ critical chains (solid line) compared to their low temperature approximations (dashed curves). As usual, the temperature is measured in units of J .

so that the central charge of the $\text{su}(2|4)$ chain is again given by Eq. (6.3) with $n = 4$. This concludes the verification of Eq. (6.5) for the $\text{su}(0|n)$, $\text{su}(2|2)$ and $\text{su}(2|4)$ chains (see Fig. 6 for a plot comparing $f(T) - f(0)$ for the latter chains with its low temperature approximations derived above).

7. Conclusions and outlook

In this paper we have introduced a novel family of $\text{su}(m|n)$ supersymmetric, translationally invariant spin chains with long-range interactions. The new chain’s spin-spin interaction term, which depends on supersymmetric permutation and spin reversal operators, reduces to the interaction term of the celebrated (supersymmetric) Haldane–Shastry chain when the spin reversal operators are replaced by (plus or minus) the identity. On the other hand, by contrast with all previously known spin chains of Haldane–Shastry type, the new model is not directly associated with an extended root system. We show that the spin chain under study can be obtained from a suitable many-body spin dynamical model in the strong coupling limit, and take advantage of this fact to evaluate the chain’s partition function in closed form. The structure of the partition function turns out to depend crucially on the parity of the integers m and n . In particular, we show that when both m and n are even the partition function exactly factorizes as the product of the partition functions of an $\text{su}(1|1)$ and an $\text{su}(\frac{m}{2}|\frac{n}{2})$ Haldane–Shastry spin chains. We also study in detail the chain’s symmetries, showing that it features a remarkable invariance under translations combined spin “twists” (i.e., spin flips at one end of the chain), as well as a general boson-fermion duality as is the case with supersymmetric spin chains of Haldane–Shastry type. In fact, the structure of the partition mentioned above when both m and n are even strongly suggests that in this case the model admits the direct sum of the Yangians $Y(\mathfrak{gl}(1|1))$ and $Y(\mathfrak{gl}(\frac{m}{2}|\frac{n}{2}))$ as an exact symmetry for an arbitrary number of sites.

The explicit knowledge of the new chain’s partition function makes it possible to study its thermodynamics in a systematic way. More precisely, when both m and n are

even we are able to find a closed-form expression for the thermodynamic free energy per spin for low values of m and n , and for arbitrary even m or n in the non-supersymmetric case $mn = 0$. For other values of m and n , we compute the free energy per spin for a finite number of spins from the partition function and show that the thermodynamic functions behave similarly as in the above mentioned cases. In particular, the specific heat at constant volume exhibits a single marked Schottky peak, whose appearance can be understood by noting the qualitative similarity of the chain's free energy with the free energy of a suitable k -level system. Our results also have led us to conjecture that in the thermodynamic limit the free energy per spin is independent of the particular representation of the spin reversal operators S_i used. In particular, this result applied to the trivial representation $S_i = \pm 1$ would imply that the thermodynamic free energy per spin of the new chain exactly coincides with its analogue for the corresponding HS chain with the coupling J doubled. This equality has been in fact verified, either exactly or numerically, in several cases.

We have also studied in detail the critical behavior of the new chain introduced in this paper. To begin with, from the ground state degeneracy we have been able to show that the model cannot be critical when $m > 2$. With the help of the closed-form expression for the partition function, we have established the criticality of the chains with $m = 0, 2$ and even n . We have also checked that at low temperatures the free energy per spin of the $\text{su}(0|n)$ (with even n), $\text{su}(2|2)$, and $\text{su}(2|4)$ chains behaves as expected for a critical system, and have computed the central charge of their associated CFTs.

The above results suggest several open problems and related lines for future work. In the first place, it would be desirable to find a simple description of the spectrum when either m or n (or both) are odd in terms of some variant of the supersymmetric motifs introduced in Section 6. A potential candidate for such a variant could be the branched motifs introduced in Ref. [35] to generate the spectrum of the supersymmetric Polychronakos–Frahm (PF) spin chain of BC_N type. Note that a motif-based description of the spectrum could shed light on the existence of low energy excitations with a linear energy-momentum relation, which is a necessary condition for criticality, for the $\text{su}(0|n)$ and $\text{su}(2|n)$ chains with odd n and the $\text{su}(1|n)$ chain. Another problem worth investigating is the existence of a factorization of the partition function of the $\text{su}(m|n)$ chains with odd m or n in terms of the partition functions of HS spin chains of A_{N-1} type, akin to Eq. (3.20) for the case of even m and n . This would automatically yield a motif-based description of the spectrum, and imply the existence of a Yangian symmetry for arbitrary values of m and n . More generally, it would certainly be desirable to find a closed-form expression for the thermodynamic free energy per spin valid for all values of m and n . In particular, the low-temperature behavior of this expression would shed light on the critical character of the $\text{su}(0|n)$ and $\text{su}(2|n)$ chains with odd n , as well as the $\text{su}(1|n)$ chain, since it could be used to compute the central charge of the hypothetical CFTs associated to these models. In any case, rigorously determining the criticality of the latter chains, and identifying their associated CFTs, are natural open

problems suggested by the present work. Finally, it would also be worthwhile to probe our conjecture on the independence of the free thermodynamic energy per spin on the particular representation of the spin reversal operators chosen, not only for the chains introduced in this work but also for similar models (for instance, HS chains of BC_N type).

Acknowledgments

This work was partially supported by grant GRFN24/24 from Universidad Complutense de Madrid. The authors would like to thank Professors Diptiman Sen and Germán Sierra for useful discussions.

Appendix A. The $\text{su}(m|n)$ superalgebra

In this appendix we shall spell out the precise connection between the $\text{su}(m|n)$ Lie superalgebra and the supersymmetric chain (2.7). To begin with, we note that the basis states (2.2) can be obtained from a Fock vacuum $|\emptyset\rangle$ as

$$|s_1 \cdots s_N\rangle = c_{1,s_1}^\dagger \cdots c_{N,s_N}^\dagger |\emptyset\rangle,$$

where $c_{i\alpha}^\dagger$ creates a particle of type α (i.e., a boson for $\alpha = 1, \dots, m$ and a fermion for $\alpha = m+1, \dots, m+n$) at site i . The operators $c_{i\alpha}, c_{i\alpha}^\dagger$ satisfy the usual canonical (anti)commutation relations

$$[c_{i\alpha}, c_{j\beta}^\dagger]_\pm := c_{i\alpha} c_{j\beta}^\dagger - (-1)^{p(\alpha)p(\beta)} = \delta_{ij} \delta_{\alpha\beta},$$

where the parity $p(\alpha)$ is defined as 0 for bosons and 1 for fermions. The chain's Hilbert space coincides with the subspace of the Fock space defined by the conditions

$$\sum_{\alpha=1}^{m+n} c_{k\alpha}^\dagger c_{k\alpha} = \mathbb{1}_k, \quad k = 1, \dots, N, \quad (\text{A.1})$$

enforcing restriction that there be exactly one particle per site. It is straightforward to check that the $\text{su}(m|n)$ supersymmetric permutation and spin flip operators S_{ij} and S_i appearing in the Hamiltonian (2.7) can be expressed in terms of the operators

$$E_k^{\alpha\beta} := c_{k\alpha}^\dagger c_{k\beta}, \quad k = 1, \dots, N, \quad \alpha, \beta = 1, \dots, m+n, \quad (\text{A.2})$$

as

$$S_{ij} = \sum_{\alpha,\beta=1}^{m+n} (-1)^{p(\beta)} E_i^{\alpha\beta} E_j^{\beta\alpha}, \quad S_i = \sum_{\alpha=1}^{m+n} \sigma(\alpha) E_i^{\alpha'\alpha}, \quad (\text{A.3})$$

where the σ and prime ($'$) notation was defined in Eqs. (2.5)-(2.6). It is also easily verified that the operators (A.2) at each site $k = 1, \dots, N$ realize the commutation relations of the $\text{gl}(m|n)$ Lie superalgebra, namely [36, 37]

$$[E^{\alpha\beta}, E^{\gamma\delta}]_\pm := E^{\alpha\beta} E^{\gamma\delta} - (-1)^{p(\alpha,\beta)p(\gamma,\delta)} E^{\gamma\delta} E^{\alpha\beta} = \delta_{\beta\gamma} E^{\alpha\delta} - (-1)^{p(\alpha,\beta)p(\gamma,\delta)} \delta_{\alpha\delta} E^{\gamma\beta},$$

where $p(\mu, \nu) = p(\mu) + p(\nu)$ is the parity of the generator $E^{\mu\nu}$. Note that the linear combination $\sum_{\alpha=1}^{m+n} E^{\alpha\alpha}$ is a linear Casimir of the $\mathfrak{gl}(n|m)$ superalgebra, since

$$\left[E^{\alpha\beta}, \sum_{\gamma=1}^{m+n} E^{\gamma\gamma} \right]_{\pm} = \left[E^{\alpha\beta}, \sum_{\gamma=1}^{m+n} E^{\gamma\gamma} \right] = \sum_{\gamma=1}^{m+n} (\delta_{\beta\gamma} E^{\alpha\gamma} - \delta_{\alpha\gamma} E^{\gamma\beta}) = E^{\alpha\beta} - E^{\alpha\beta} = 0.$$

For the representations (A.2) of $\mathfrak{gl}(m|n)$ in terms of creation and annihilation operators at site k this Casimir takes the value 1 on account of the constraint (A.1). Let us next define the supertrace of an operator A_k acting on the k -th site as

$$\text{str } A_k = \sum_{\alpha=1}^{m+n} (-1)^{p(\alpha)} {}_k \langle \alpha | A | \alpha \rangle_k,$$

where $|\alpha\rangle_k = c_{k\alpha}^\dagger |\emptyset\rangle$. Clearly

$$\text{str } E_k^{\alpha\beta} = \sum_{\gamma=1}^{m+n} (-1)^{p(\gamma)} \langle \emptyset | c_{k\gamma}^\dagger c_{k\alpha}^\dagger c_{k\beta} c_{k\gamma} | \emptyset \rangle = \sum_{\gamma=1}^{m+n} (-1)^{p(\gamma)} \delta_{\gamma\alpha} \delta_{\beta\gamma} = (-1)^{p(\alpha)} \delta_{\alpha\beta}.$$

In order to obtain a representation of the Lie superalgebra $\mathfrak{sl}(m|n)$, we must impose the additional condition that the supertrace of its generators vanish. When $m \neq n$, this is can be done in a symmetric way by defining the new generators

$$J_k^{\alpha\beta} = E_k^{\alpha\beta} - \frac{(-1)^{p(\alpha)}}{m-n} \delta_{\alpha\beta} \mathbb{1}_k, \quad (\text{A.4})$$

since $\text{str } \mathbb{1}_k = m - n$. Note that the dimension of $\mathfrak{sl}(m|n)$ is $(m+n)^2 - 1$, since we have the constraint

$$\sum_{\alpha=1}^{m+n} J_k^{\alpha\alpha} = \sum_{\alpha=1}^{m+n} E_k^{\alpha\alpha} - \mathbb{1}_k = 0 \quad (\text{A.5})$$

by Eq. (A.1). When $m = n$ it is still possible to construct the $\mathfrak{sl}(m|m)$ generators out of the $\mathfrak{gl}(m|n)$ generators (A.2) in a less symmetric way by defining, for instance, $J_k^{\alpha\beta} = E_k^{\alpha\beta}$ for $\alpha \neq \beta$ and

$$J_k^{\alpha\alpha} = (-1)^{p(\alpha)} E_k^{\alpha\alpha} - (-1)^{p(\alpha+1)} E_k^{\alpha+1, \alpha+1}, \quad \alpha = 1, \dots, N-1.$$

In fact, while the Lie superalgebra $\mathfrak{sl}(m|n)$ is simple when $m \neq n$ [36, 38], $\mathfrak{sl}(m|m)$ is not. Indeed, when $m = n$ the one-dimensional linear space spanned by the Casimir $\sum_{\alpha=1}^{2m} E_k^{\alpha\alpha}$ —which belongs to $\mathfrak{sl}(m|n)$ for $m = n$ —is obviously an ideal of $\mathfrak{sl}(m|m)$. For the sake of simplicity (as is usually done in the literature) we shall therefore restrict ourselves in what follows to the case $m \neq n$.

When $m \neq n$, an elementary calculation shows that the $\mathfrak{sl}(m|n)$ generators (A.4) at site k satisfy exactly the same commutation relation as the $\mathfrak{gl}(m|n)$ generators (A.2), namely

$$[J_k^{\alpha\beta}, J_k^{\gamma\delta}]_{\pm} = \delta_{\beta\gamma} J_k^{\alpha\delta} - (-1)^{p(\alpha, \beta)p(\gamma, \delta)} \delta_{\alpha\delta} J_k^{\gamma\beta}, \quad \alpha, \beta = 1, \dots, m+n.$$

Formally these are also the $\mathfrak{su}(m|n)$ commutation relations, since $\mathfrak{sl}(m|n)$ is the complexification of $\mathfrak{su}(m|n)$.

It is straightforward to express the operators S_{ij} and S_i in terms of the $\text{su}(m|n)$ — or, strictly speaking, $\text{sl}(m|n)$ — generators $J_k^{\alpha\beta}$ at sites i and j . Indeed, using Eq. (A.5) we obtain

$$\begin{aligned}
 \sum_{\alpha,\beta} (-1)^{p(\beta)} J_i^{\alpha\beta} J_j^{\beta\alpha} &\equiv \sum_{\alpha,\beta} (-1)^{p(\alpha)} (J_i^{\alpha\beta})^\dagger J_j^{\alpha\beta} \\
 &= \sum_{\alpha} \left(c_{i\alpha}^\dagger c_{i\alpha} - \frac{(-1)^{p(\alpha)}}{m-n} \mathbb{1} \right) (-1)^{p(\alpha)} J_j^{\alpha\alpha} + \sum_{\alpha \neq \beta} (-1)^{p(\beta)} c_{i\alpha}^\dagger c_{i\beta} c_{j\beta}^\dagger c_{j\alpha} \\
 &= \sum_{\alpha} (-1)^{p(\alpha)} c_{i\alpha}^\dagger c_{i\alpha} J_j^{\alpha\alpha} + \sum_{\alpha \neq \beta} c_{i\alpha}^\dagger c_{j\beta}^\dagger c_{i\beta} c_{j\alpha} \\
 &= \sum_{\alpha} (-1)^{p(\alpha)} c_{i\alpha}^\dagger c_{i\alpha} \left(c_{j\alpha}^\dagger c_{j\alpha} - \frac{(-1)^{p(\alpha)}}{m-n} \mathbb{1} \right) + \sum_{\alpha \neq \beta} c_{i\alpha}^\dagger c_{j\beta}^\dagger c_{i\beta} c_{j\alpha} \\
 &= \sum_{\alpha,\beta} c_{i\alpha}^\dagger c_{j\beta}^\dagger c_{i\beta} c_{j\alpha} - \frac{\mathbb{1}}{m-n} \sum_{\alpha} c_{i\alpha}^\dagger c_{i\alpha} = S_{ij} - \frac{\mathbb{1}}{m-n},
 \end{aligned}$$

and therefore

$$S_{ij} = \sum_{\alpha,\beta} (-1)^{p(\beta)} J_i^{\alpha\beta} J_j^{\beta\alpha} + \frac{\mathbb{1}}{m-n}. \quad (\text{A.6})$$

Similarly,

$$\begin{aligned}
 S_i &= \sum_{\alpha} \sigma(\alpha) E_i^{\alpha'\alpha} = \sum_{\alpha} \sigma(\alpha) \left(J_i^{\alpha'\alpha} - \frac{(-1)^{p(\alpha)}}{m-n} \delta_{\alpha\alpha'} \mathbb{1} \right) \\
 &= \sum_{\alpha} \sigma(\alpha) J_i^{\alpha'\alpha} + \frac{\pi(n)\varepsilon_F - \pi(m)\varepsilon_B}{m-n} \mathbb{1},
 \end{aligned} \quad (\text{A.7})$$

where $\pi(\alpha) = (1 - (-1)^\alpha)/2$ is the parity of the integer α . Since $\tilde{S}_{ij} = S_{ij} S_i S_j$, it follows that the operators \tilde{S}_{ij} appearing in the chain Hamiltonian (2.7) can be expressed as a fourth degree polynomial in the $\text{su}(m|n)$ generators (A.4) at sites i and j . Obviously the same is then true for the full Hamiltonian (2.7), and it is precisely in this sense that the latter model can be regarded as an $\text{su}(m|n)$ supersymmetric chain.

Although not strictly necessary, we can replace the non-Hermitian generators $J_k^{\alpha\beta}$ in Eq. (A.4) by the Hermitian linear combinations

$$\begin{aligned}
 A_k^{\alpha\beta} &= \frac{1}{2} (J_k^{\alpha\beta} + J_k^{\beta\alpha}), \quad 1 \leq \alpha \leq \beta \leq m+n; \\
 B_k^{\alpha\beta} &= \frac{i}{2} (J_k^{\beta\alpha} - J_k^{\alpha\beta}), \quad 1 \leq \alpha < \beta \leq m+n,
 \end{aligned}$$

which obey the constraint

$$\sum_{\alpha=1}^{m+n} A_k^{\alpha\alpha} \equiv \sum_{\alpha=1}^{m+n} J_k^{\alpha\alpha} = 0.$$

After a straightforward calculation we arrive at the relations

$$\begin{aligned}
 S_{ij} &= \sum_{\alpha=1}^{m+n} (-1)^{p(\alpha)} A_i^{\alpha\alpha} A_j^{\alpha\alpha} + 2 \sum_{r<s} (A_i^{rs} A_j^{rs} + B_i^{rs} B_j^{rs}) - 2 \sum_{\rho<\sigma} (A_i^{\rho\sigma} A_j^{\rho\sigma} + B_i^{\rho\sigma} B_j^{\rho\sigma}) \\
 &\quad + 2i \sum_{r,\rho} (A_i^{r\rho} B_j^{r\rho} - B_i^{r\rho} A_j^{r\rho}) + \frac{\mathbb{1}}{m-n},
 \end{aligned}$$

$$\begin{aligned}
 S_i &= 2\varepsilon_B \sum_{r \leq m/2} A_i^{rr'} + 2\varepsilon_F \sum_{\rho \leq m+n/2} A_i^{\rho\rho'} + \pi(m)\varepsilon_B \left(A_i^{\frac{m+1}{2} \frac{m+1}{2}} - \frac{\mathbb{1}}{m-n} \right) \\
 &\quad + \pi(n)\varepsilon_F \left(A_i^{m+\frac{n+1}{2} m+\frac{n+1}{2}} + \frac{\mathbb{1}}{m-n} \right),
 \end{aligned}$$

where r and s (respectively ρ and σ) run over the bosonic indices $1, \dots, m$ (respectively the fermionic indices $m+1, \dots, m+n$).

Appendix B. Asymptotic approximation of the integral (6.6)

The asymptotic behavior of the integral $I_\beta[\varphi]$ in Eq. (6.6) as $\beta \rightarrow \infty$ can be ascertained performing the change of variable $\beta\rho(x) = y$, or equivalently

$$x = \rho^{-1}(Ty) = \frac{1}{2} \left(1 - \sqrt{1 - 4Ty} \right), \quad 0 \leq y \leq \beta/4,$$

whence

$$\begin{aligned}
 I_\beta[\varphi] &= T \int_0^{\beta/4} dy \frac{\log(\varphi(e^{-y}))}{\sqrt{1-4Ty}} \\
 &= TI[\varphi] - T \int_{\beta/4}^\infty dy \log(\varphi(e^{-y})) + T \int_0^{\beta/4} dy \left(\frac{1}{\sqrt{1-4Ty}} - 1 \right) \log(\varphi(e^{-y}))
 \end{aligned}$$

with

$$I[\varphi] := \int_0^\infty dy \log(\varphi(e^{-y})).$$

Note that the assumption (6.7) on $\varphi(z)$ implies that

$$\log(\varphi(z)) = \log(1 + O(z)) = O(z) \tag{B.1}$$

as $z \rightarrow 0+$, from which it follows that $I[\varphi]$ is convergent and

$$\int_{\beta/4}^\infty dy \log(\varphi(e^{-y})) = O(e^{-\beta/4}).$$

To cope with the (integrable) singularity of the integrand at the upper integration limit in the integral

$$\int_0^{\beta/4} dy \left(\frac{1}{\sqrt{1-4Ty}} - 1 \right) \log(\varphi(e^{-y})),$$

we note that

$$\frac{1}{\sqrt{1-4z}} - 1 = h'(z), \quad \text{with } h(z) = \frac{1}{2} (1 - 2z - \sqrt{1-4z}).$$

Taking into account that $h(0) = 0$ and integrating by parts we obtain

$$\begin{aligned}
 \int_0^{\beta/4} dy \left(\frac{1}{\sqrt{1-4Ty}} - 1 \right) \log(\varphi(e^{-y})) &= \int_0^{\beta/4} dy h'(Ty) \log(\varphi(e^{-y})) \\
 &= \frac{1}{4T} \log(\varphi(e^{-\beta/4})) + \frac{1}{T} \int_0^{\beta/4} dy h(Ty) e^{-y} \frac{\varphi'(e^{-y})}{\varphi(e^{-y})}.
 \end{aligned}$$

The first term is clearly $O(\beta e^{-\beta/4})$ as $T \rightarrow 0+$ by Eq. (B.1). On the other hand, from the elementary inequalities

$$0 \leq h(z) \leq 4z^2$$

it follows that

$$\frac{1}{T} \left| \int_0^{\beta/4} dy h(Ty) e^{-y} \frac{\varphi'(e^{-y})}{\varphi(e^{-y})} \right| \leq 4T \int_0^\infty dy y^2 e^{-y} \left| \frac{\varphi'(e^{-y})}{\varphi(e^{-y})} \right| = O(T),$$

since the integral is convergent. Indeed, by Eq. (B.1) we have

$$e^{-y} \left| \frac{\varphi'(e^{-y})}{\varphi(e^{-y})} \right| = O(e^{-y})$$

as $y \rightarrow \infty$. Putting everything together we conclude that

$$I_\beta[\varphi] = TI[\varphi] + O(T^2) \tag{B.2}$$

as $T \rightarrow 0+$.

References

- [1] Basu-Mallick B, Finkel F and González-López A, *A novel class of translationally invariant spin chains with long-range interactions*, 2020 *J. High Energy Phys.* **08** 099(43)
- [2] Haldane F D M, *Exact Jastrow–Gutzwiller resonating-valence-bond ground state of the spin-1/2 antiferromagnetic Heisenberg chain with $1/r^2$ exchange*, 1988 *Phys. Rev. Lett.* **60** 635
- [3] Shastry B S, *Exact solution of an $S = 1/2$ Heisenberg antiferromagnetic chain with long-ranged interactions*, 1988 *Phys. Rev. Lett.* **60** 639
- [4] Polychronakos A P, *Lattice integrable systems of Haldane–Shastry type*, 1993 *Phys. Rev. Lett.* **70** 2329
- [5] Polychronakos A P, *Exact spectrum of $SU(n)$ spin chain with inverse-square exchange*, 1994 *Nucl. Phys. B* **419** 553
- [6] Olshanetsky M A and Perelomov A M, *Quantum integrable systems related to Lie algebras*, 1983 *Phys. Rep.* **94** 313
- [7] Corrigan E and Sasaki R, *Quantum versus classical integrability in Calogero–Moser systems*, 2002 *J. Phys. A: Math. Gen.* **35** 7017
- [8] Odake S and Sasaki R, *Equilibria of ‘discrete’ integrable systems and deformation of classical orthogonal polynomials*, 2004 *J. Phys. A: Math. Gen.* **37** 11841
- [9] Haldane F D M, Ha Z N C, Talstra J C, Bernard D and Pasquier V, *Yangian symmetry of integrable quantum chains with long-range interactions and a new description of states in conformal field theory*, 1992 *Phys. Rev. Lett.* **69** 2021
- [10] Haldane F D M, *Physics of the ideal semion gas: spinons and quantum symmetries of the integrable Haldane–Shastry spin chain*, in A Okiji and N Kawakami, eds., *Correlation Effects in Low-dimensional Electron Systems*, Springer Series in Solid-state Sciences, volume 118 (Springer Berlin, Heidelberg), pp. 3–20
- [11] Hikami K and Basu-Mallick B, *Supersymmetric Polychronakos spin chain: motif, distribution function, and character*, 2000 *Nucl. Phys. B* **566** 511
- [12] Basu-Mallick B, Bondyopadhyaya N, Hikami K and Sen D, *Boson-fermion duality in $SU(m|n)$ supersymmetric Haldane–Shastry spin chain*, 2007 *Nucl. Phys. B* **782** 276
- [13] Basu-Mallick B and Bondyopadhyaya N, *Exact partition functions of $SU(m|n)$ supersymmetric Haldane–Shastry spin chain*, 2006 *Nucl. Phys. B* **757** 280
- [14] Basu-Mallick B, Bondyopadhyaya N and Hikami K, *One-dimensional vertex models associated with a class of Yangian invariant Haldane–Shastry like spin chains*, 2010 *Symmetry Integr. Geom.* **6** 091(13)
- [15] Haldane F D M, *“Spinon gas” description of the $S = \frac{1}{2}$ Heisenberg chain with inverse-square exchange: exact spectrum and thermodynamics*, 1991 *Phys. Rev. Lett.* **66** 1529
- [16] Frahm H, *Spectrum of a spin chain with inverse-square exchange*, 1993 *J. Phys. A: Math. Gen.* **26** L473

- [17] Frahm H and Inozemtsev V I, *New family of solvable 1D Heisenberg models*, 1994 *J. Phys. A: Math. Gen.* **27** L801
- [18] Enciso A, Finkel F and González-López A, *Thermodynamics of spin chains of Haldane–Shastry type and one-dimensional vertex models*, 2012 *Ann. Phys.-New York* **327** 2627
- [19] Finkel F, González-López A, León I and Rodríguez M A, *Thermodynamics and criticality of supersymmetric spin chains with long-range interactions*, 2018 *J. Stat. Mech.-Theory E.* **2018** 043101(47)
- [20] Sachdev S, *Quantum Phase Transitions* (Cambridge University Press), second edition 2011
- [21] Schoutens K, *Yangian symmetry in conformal field theory*, 1994 *Phys. Lett. B* **331** 335
- [22] Bouwknegt P and Schoutens K, *The $\widehat{SU(n)}_1$ WZW models. Spinon decomposition and Yangian structure*, 1996 *Nucl. Phys. B* **482** 345
- [23] Polyakov A and Wiegmann P B, *Theory of non-abelian Goldstone bosons in two dimensions*, 1983 *Phys. Lett. B* **131** 121
- [24] Witten E, *Non-abelian bosonization in two dimensions*, 1984 *Commun. Math. Phys.* **92** 455
- [25] Knizhnik V G and Zamolodchikov A B, *Current algebra and Wess–Zumino model in two dimensions*, 1984 *Nucl. Phys. B* **247** 83
- [26] Basu-Mallick B, Bondyopadhyaya N and Sen D, *Low energy properties of the $SU(m|n)$ supersymmetric Haldane–Shastry spin chain*, 2008 *Nucl. Phys. B* **795** 596
- [27] Blöte H W J, Cardy J L and Nightingale M P, *Conformal invariance, the central charge, and universal finite-size amplitudes at criticality*, 1986 *Phys. Rev. Lett.* **56** 742
- [28] Affleck I, *Universal term in the free energy at a critical point and the conformal anomaly*, 1986 *Phys. Rev. Lett.* **56** 746
- [29] Basu-Mallick B, Finkel F and González-López A, *Exactly solvable D_N -type quantum spin models with long-range interaction*, 2009 *Nucl. Phys. B* **812** 402
- [30] Basu-Mallick B, Finkel F and González-López A, *The spin Sutherland model of D_N type and its associated spin chain*, 2011 *Nucl. Phys. B* **843** 505
- [31] Finkel F and González-López A, *A new perspective on the integrability of Inozemtsev’s elliptic spin chain*, 2014 *Ann. Phys.-New York* **351** 797
- [32] Finkel F and González-López A, *Global properties of the spectrum of the Haldane–Shastry spin chain*, 2005 *Phys. Rev. B* **72** 174411(6)
- [33] Barba J C, Finkel F, González-López A and Rodríguez M A, *An exactly solvable supersymmetric spin chain of BC_N type*, 2009 *Nucl. Phys. B* **806** 684
- [34] Finkel F and González-López A, *Thermodynamics and criticality of $su(m)$ spin chains of Haldane–Shastry type*, 2022 *Phys. Rev. E* **106** 054120(11)
- [35] Basu-Mallick B and Sinha M, *Appearance of branched motifs in the spectra of BC_N type Polychronakos spin chains*, 2020 *Nucl. Phys. B* **952** 114914(37)
- [36] Rittenberg V, *A guide to Lie superalgebras*, in P Kramer and A Rieckers, eds., *Group Theoretical Methods in Physics. Sixth International Colloquium, Tübingen 1977, Lecture Notes in Physics*, volume 79 (Springer-Verlag), pp. 3–21
- [37] Jarvis P D and Green H S, *Casimir invariants and characteristic identities for generators of the general linear, special linear and orthosymplectic graded Lie algebras*, 1979 *J. Math. Phys.* **20** 2115
- [38] Kac V G, *Lie superalgebras*, 1977 *Adv. Math.* **26** 8