

Set risk measures

Marcelo Righi

Eduardo Horta

Marlon Moresco

Abstract

We introduce the concept of set risk measures (SRMs), which are real-valued maps defined on the space of all non-empty, closed, and bounded sets of almost surely bounded random variables. Traditional risk measures typically operate on spaces of random variables, but SRMs extend this framework to sets of random variables. We establish an axiom scheme for SRMs, similar to classical risk measures but adapted for set operations. The main technical contribution is an axiomatic dual representation of convex SRMs by using regular, finitely additive measures on the unit ball of the dual space of essentially bounded random variables. We explore worst-case SRMs, which evaluate risk as the supremum of individual risks within a set, and provide a collection of examples illustrating the applicability of our framework to systemic risk, portfolio optimization, and decision-making under uncertainty. This work extends the theory of risk measures to a more general and flexible setup, accommodating a broader range of financial and mathematical applications.

Keywords: Risk measures; Set analysis; Robustness; Uncertainty; Convex analysis.

Index of Notation

\mathbf{ba}	The space of all finitely additive signed measures on (Ω, \mathcal{F}) that are absolutely continuous with respect to p and with finite total variation
\mathbf{ba}_+	The set of elements $q \in \mathbf{ba}$ s.t. $q(A) \geq 0$ for all $A \in \mathcal{F}$
$\mathbf{ba}_{1,+}$	The set of elements $q \in \mathbf{ba}_+$ s.t. $q(\Omega) = 1$
$\text{CB}(M)$	Space of all non-empty, closed and bounded subsets of a metric space M
$\text{CBC}(M)$	Space of all non-empty, closed, bounded and convex subsets of a metric vector space M
$\langle q, x \rangle$	Integral of the random variable x w.r.t. the measure q
$\langle Q, X \rangle$	$\sup_{(q,x) \in Q \times X} \langle q, x \rangle$
p, q, \dots	Probability measures on (Ω, \mathcal{F})
P, Q, \dots	Sets of probability measures
$\overline{\mathbb{R}}$	The extended real line, i.e., $\overline{\mathbb{R}} = \mathbb{R} \cup \{-\infty, \infty\}$
x, y, z, \dots	Random variables
X, Y, Z, \dots	Sets of random variables
$\mathcal{X}, \mathcal{Y}, \mathcal{Z}, \dots$	Collections of sets of random variables
$\alpha, \gamma, \epsilon, \lambda$	Real scalars
\dot{x}	For any object x , the singleton containing x , that is, $\dot{x} = \{x\}$
U	The unit ball in $(\mathbf{ba}, \ \cdot\ _{TV})$
U_+	The non-negative elements in U , i.e., $U_+ = U \cap \mathbf{ba}_+$

1 Introduction

Risk measures are a prominent topic in the mainstream literature on Mathematical Finance, especially since the landmark paper of [Artzner et al. \(1999\)](#)—see [Pflug and Römisch \(2007\)](#), [Delbaen \(2012\)](#), [Rüschendorf \(2013\)](#), [Follmer and Schied \(2016\)](#) and [Shapiro et al. \(2021\)](#) for a comprehensive review. The canonical theory, as summarized for example in [Delbaen \(2002\)](#), considered risk measures as real-valued functionals defined on the domain L^∞ of bounded random variables. From there, more general domains have been considered in order to both generalize and adapt the approach, for instance, L^λ spaces ($\lambda \geq 1$) in [Kaina and Rüschendorf \(2009\)](#), non-integrable random variables in [Delbaen \(2009\)](#), finite spaces in [Kou et al. \(2013\)](#), Orlicz spaces in [Cheridito and Li \(2009\)](#) and [Gao and Xanthos \(2018\)](#), spaces generated by risk measures in [Pichler \(2013\)](#), spaces of probability distributions as in [Frittelli et al. \(2014\)](#), continuous and bounded random variables over a Polish space as in [Bion-Nadal and Kervarec \(2012\)](#), [Delbaen \(2022\)](#), [Nendel \(2024\)](#).

Beyond scalar random variables, [Burgert and Rüschendorf \(2006, 2008\)](#), [Rüschendorf \(2006\)](#), [Ivar and Walter \(2011\)](#), [Ekeland et al. \(2012\)](#), [Wei and Hu \(2014\)](#) and [Feinstein and Rudloff \(2022\)](#)—to name a few—, study real valued risk measures whose domain is a space of random vectors, thus partially tackling the problem of measuring the risk of a portfolio. The works of [Chen et al. \(2013\)](#), [Kromer et al. \(2016a\)](#), and [Biagini and Frittelli \(2010\)](#), among many others, investigate a similar setup but with the interpretation for systemic risk measures. Beyond finite index sets, we may cite works of risk measures over spaces of stochastic processes, as addressed in [Cheridito et al. \(2004\)](#), [Cheridito et al. \(2005\)](#), [Frittelli and Scandolo \(2006\)](#), [Arai \(2014\)](#) and [Assa and Morales \(2010\)](#), among others. Additionally, there is a stream of literature—see [Jouini et al. \(2004\)](#), [Hamel and Heyde \(2010\)](#), [Hamel et al. \(2011\)](#), [Ararat et al. \(2017\)](#), to name a few—focusing on set-valued risk measures, i.e., functionals whose codomain is a vector space, or, more generally, a space of closed convex sets.

In this paper, we introduce the concept of a *set risk measure* (SRM, for short), by contemplating real valued maps defined on the domain $\text{CBC}(L^\infty)$ comprised of all non-empty, closed and bounded sets of almost surely bounded random variables. The challenge of working in this setup is elevated, since this domain is not a linear space, and thus many conventional tools available for topological vector spaces, are not directly applicable. To circumvent such difficulties, we rely on topological, algebraic and order structures. The key technical insight is using Hörmander’s Theorem ([Beer, 1993](#), Theorem 3.2.9), which allows one to embed $\text{CBC}(L^\infty)$ as a closed, convex cone into a Banach

space of continuous functions. To the best of our knowledge, this framework is entirely new in the literature.

We begin in Section 2 by introducing an *axiom scheme* for our proposed set risk measures. Some of these axioms are direct generalizations of standard axioms for traditional risk measures; others are specific to our framework, for instance those dealing with set operations. In our setup, similarly to the classical case, a set risk measure induces an acceptance set that is a collection of sets. We introduce an analogous axiom scheme for these acceptance sets, and establish a connection between them, vis-à-vis the axioms for the set risk measure.

Next, in the main technical step of this paper, we provide in Section 3 an axiomatic dual representation for convex set risk measures in terms of regular, finitely additive measures over the unit ball of \mathbf{ba} , the topological dual of L^∞ . The representation is obtained by relying on the important fact that the set of non-empty, closed, bounded and convex sets in L^∞ can be embedded as a cone into the space of continuous and bounded real valued functions on the unit ball of \mathbf{ba} . Furthermore, under continuity properties, we show that this dual representation can be expressed in terms of *bona fide* probability measures.

In Section 4, we also study in detail the framework of *worst-case risk measures*, an important class of set risk measures which appraise the risk of a collection of financial positions as the pointwise supremum of the individual risks (as measured by some traditional risk measure) of the positions in the set. This has become a prominent stream of studies such as Föllmer and Schied (2002), Laeven and Stajje (2013), Ang et al. (2018), Wang and Ziegel (2018), Bartl et al. (2019), Bellini et al. (2018), Guo and Xu (2019), Qian et al. (2019), Righi (2019), Bernard et al. (2023), Fadina et al. (2021), Righi (2024). We show necessary and sufficient conditions for a SRM to be a worst case risk measure, and refine some results regarding its continuity and dual representation.

Section 5 provides a collection of illustrative examples, showcasing the practical applicability of our theoretical framework to systemic risk, portfolio optimization, and decision-making under uncertainty. These examples demonstrate the flexibility and broad relevance of SRMs, highlighting their potential to address complex challenges in financial risk management.

2 Set risk measures

We consider a probability space (Ω, \mathcal{F}, p) . By letting $L^0 := L^0(\Omega, \mathcal{F}, p)$ and $L^\infty := L^\infty(\Omega, \mathcal{F}, p)$ denote the spaces of (equivalence classes under p -a.s. equality of) finite

and essentially bounded random variables, respectively. All equalities and inequalities are taken in the p -almost sure sense, in particular, constant random variables *are* real numbers. When not explicitly mentioned, we consider L^∞ equipped with its strong topology. We write 1_A for the indicator function of an event $A \in \mathcal{F}$. Convergences will be denoted by an arrow (\rightarrow) with an affixed subscript indicating the type of convergence, for instance, $x_n \rightarrow_{L^\infty} x$ means convergence in essential supremum norm; $x_n \rightarrow_{q\text{-a.s.}} x$ indicates q -a.s. convergence, for any probability measure q . Similarly, norms are denoted with an affixed subscript. In what follows, for any object x , the singleton containing x will be denoted by \dot{x} , that is, $\dot{x} = \{x\}$. We denote by $\text{cl}(X)$ the **closure** of X , and by $\text{conv}(X)$ the **closed convex hull** of $X \subseteq L^\infty$. In both cases, the closure is taken with respect to the supremum norm in L^∞ .

Consider the power set $\text{pwr}(L^\infty)$, i.e., $X \in \text{pwr}(L^\infty)$ if and only if $X \subseteq L^\infty$. We start off by endowing $\text{pwr}(L^\infty)$ with a metric structure, as follows: For any pair $X, Y \subseteq L^\infty$, define the **distance** between the two sets as

$$d(X, Y) = \inf_{x \in X, y \in Y} \|x - y\|_\infty.$$

The **Hausdorff distance** on $\text{pwr}(L^\infty)$ is then defined through

$$d_H(X, Y) = \max \left\{ \sup_{x \in X} d(\dot{x}, Y), \sup_{y \in Y} d(\dot{y}, X) \right\}.$$

In what follows, the notation $X_n \rightarrow X$ means $d_H(X_n, X) \rightarrow 0$. We now recall some properties of d_H that will be used without further mention (see [Beer, 1993](#), for details).

1. $d_H(X, Y) = \sup_{x \in L^\infty} |d(\dot{x}, X) - d(\dot{x}, Y)|$.
2. If $Y = \dot{y}$ is a singleton, then $d_H(X, Y) = \sup_{x \in X} \|x - y\|_\infty$.
3. $d_H(X, Y) \in [0, \infty]$. If both sets are bounded, then d_H is guaranteed to be finite. Further, $d_H(X, Y) = 0$ if and only if $\text{cl}(X) = \text{cl}(Y)$.
4. $d(\dot{x}, X) \leq d(\dot{x}, Y) + d_H(X, Y)$ for any $x \in L^\infty$.
5. If $\text{int}(X \cap Y) \neq \emptyset$, then there is a $\lambda > 0$ such that, for any $Z \subseteq L^\infty$ satisfying $d_H(X, Z) < \lambda$, it holds that $Y \cap Z \neq \emptyset$.
6. d_H is an extended pseudometric on $\text{pwr}(L^\infty)$.

We write $\text{CB}(L^\infty)$ to denote the collection of non-empty, closed and bounded subsets of L^∞ , while $\text{CBC}(L^\infty)$ denotes the collection of subsets in $\text{CB}(L^\infty)$ that are convex in L^∞ . Importantly, d_H yields a complete metric on $\text{CB}(L^\infty)$.

In view of the above, our “domain of choice” to work with here will be the complete

metric space $(\text{CB}(L^\infty), d_H)$. Clearly, the map $(X, Y) \mapsto X \cup Y$ is continuous on $\text{CB}(L^\infty)$, and $(X, Y) \mapsto \text{conv}(X \cup Y)$ is continuous on $\text{CBC}(L^\infty)$. In addition, notice that the collection of finite sets in L^∞ is dense in the collection of non-empty compact sets in L^∞ .

Next, we focus on algebraic structures on $\text{pwr}(L^\infty)$. It is important to notice that, while these properties *do* depend on the topology of L^∞ , in $\text{pwr}(L^\infty)$ they are indeed purely algebraic. We define the **Minkowski sum** of two sets X and Y as the operation $X + Y = \text{cl}(\{x + y : x \in X, y \in Y\})$, and we shall write $X + \dot{x}$ and $X + x$ indistinctly when no confusion may arise. This operation is commutative and associative, with $\dot{0}$ as the identity element. Likewise, we introduce multiplication by scalars through $\lambda X = \{\lambda x : x \in X\}$ for $\lambda \in \mathbb{R}$, and $X - Y = X + (-1Y)$. Contrary to first-intuition, such operations do not yield a linear space, since the Minkowski sum does not fulfil the cancellation law.

Now, as $d_H(X + \dot{\lambda}, Y + \dot{\lambda}) = d_H(X, Y)$ and $d_H(\lambda X, \lambda Y) = |\lambda|d_H(X, Y)$ for any $\lambda \in \mathbb{R}$, we have that both the Minkowski sum and the multiplication by scalar are continuous on respective topologies. Also, notice that the distributive property $\lambda(X + Y) = \lambda X + \dot{\lambda}Y$ holds for any $\lambda \in \mathbb{R}$ and any $X, Y \in \text{CB}(L^\infty)$, and that $\lambda_1(\lambda_2 X) = (\lambda_1 \lambda_2)X$ as well. However, in general $\lambda_1 X + \lambda_2 X \neq (\lambda_1 + \lambda_2)X$. In summary, $\text{pwr}(L^\infty)$ is a **conlinear** space, and $\text{CB}(L^\infty)$ is a conlinear subspace of $\text{pwr}(L^\infty)$ (for a review on conlinear spaces, see [Schrage, 2009](#); [Hamel, 2005](#)). With a slight abuse of language, we employ in $\text{pwr}(L^\infty)$ some vector space concepts such as convex set, convex hull, linear function etc., whenever those operations are well-defined. We call $X \mapsto \|X\| = \sup_{x \in X} \|x\|_\infty = \inf\{\alpha \in \mathbb{R} : p(|x| > \alpha) = 0, \forall x \in X\}$ a norm in $\text{pwr}(L^\infty)$. From properties of both supremum and $\|\cdot\|_\infty$, we get that $X \mapsto \|X\|$ is sub-additive, positive homogeneous, and finite for bounded X . In yet another abuse of notation, we say X_n **converges** to X in this essential supremum (uniform) norm when $\|X_n - X\| \rightarrow 0$. Furthermore, since $d_H(X, Y) \leq \|X - Y\|$, it follows that $\|X_n - X\| \rightarrow 0$ implies in $d_H(X_n, X) \rightarrow 0$.

Last but not least, we introduce the following pre-order in $\text{pwr}(L^\infty)$: we say that Y **is greater than** X , denoted by $X \leq Y$, if for any $x \in X$ there is a $y \in Y$ such that $x \leq y$, and we write $Y \geq X$ to mean $X \leq Y$. Reflexivity and transitivity clearly hold for “ \leq ”, while anti-symmetry does not hold. Additionally, this pre-order is compatible with the algebraic operations in $\text{CB}(L^\infty)$, in the sense that if $X \leq Y$, then $X + Z \leq Y + Z$, for any $Z \in \text{CB}(L^\infty)$, $\lambda X \leq \lambda Y$ for any $\lambda \geq 0$, and $-Y \leq -X$. Indeed, we have $X \leq Y$ if and only if $X \subseteq Y - L_+^\infty$. The restriction of “ \leq ” to the monotone subsets of L^∞ , i.e., the sets such that $X + L_+^\infty = X$, coincides with the order complete lattice set inclusion \subseteq .

Let the **absolute value** of X be defined through $|X| = \{|x|: x \in X\}$. Additionally, for any $X \in \text{pwr}(L^\infty)$, its essential supremum, denoted $\text{ess sup } X$, is the (almost surely) unique $y \in L^\infty$ that satisfies $y \geq x$ for any $x \in X$, and $y \leq z$ for any $z \in L^\infty$ such that $z \geq x$ for any $x \in X$. The essential infimum of X is then defined as $\text{ess inf } X = -\text{ess sup } -X$. It is possible to show (see Theorem A.37 in [Follmer and Schied, 2016](#)) that $\text{ess sup } X$ is well-defined, by allowing the extended real line to be its codomain, and can be taken as the pointwise supremum $\text{ess sup } X(\omega) = \sup_{x \in X} x(\omega)$ when X is countable. The essential supremum of a set is: finite if and only if there exists a finite upper bound for the almost sure supremum of those variables; integrable if and only if there exists an integrable random variable that dominates the set; bounded if and only if there exists a finite constant that serves as an upper bound for almost all values of those random variables. Note that $\|X\| \geq \text{ess sup } |X|$.

To sum up, we have that, based on the topological, algebraic and order structures introduced above, the space $\text{CB}(L^\infty)$, endowed with the topology induced by d_H , is a preordered Abelian topological semigroup—see [Beer \(1993\)](#), Chapter 3, for instance.

Definition 2.1. Any functional $R: \text{CB}(L^\infty) \rightarrow \mathbb{R}$ is called a **set risk measure** (SRM henceforth), and may have the following properties, for any $X, Y \in \text{CB}(L^\infty)$, any $\alpha \in \mathbb{R}$, any $0 \leq \lambda \leq 1$ and any $\gamma \geq 0$:

- (i) **Monotonicity:** If $X \leq Y$, then $R(X) \geq R(Y)$.
- (ii) **Translation invariance:** $R(X + \alpha) = R(X) - \alpha$.
- (iii) **Set-monotonicity:** If $X \subseteq Y$, then $R(X) \leq R(Y)$.
- (iv) **Union-boundedness:** $R(X \cup Y) \leq \max\{R(X), R(Y)\}$.
- (v) **WC-boundedness:** $R(X) \leq \sup_{x \in X} R(x)$.
- (vi) **Convexity:** $R(\lambda X + (1 - \lambda)Y) \leq \lambda R(X) + (1 - \lambda)R(Y)$.
- (vii) **Positive homogeneity:** $R(\gamma X) = \gamma R(X)$.
- (viii) **Set-convexity:** $R(\text{conv}(X)) \leq R(X)$.

An SRM is called **monetary** if it satisfies (i) and (ii). It is called **convex** if it is monetary and satisfies (vi), **coherent** if it is convex and satisfies (vii). In what follows, we assume that SRMs are always **normalized**, i.e. $R(\dot{0}) = 0$.

Assuming normalization does not affect our main results. For instance, this property is always present under positive homogeneity. Additionally, for a monetary SRM, say R , we have that R' defined as $R'(X) = R(X) - R(\dot{0})$ is monetary and normalized, and also inherits from R any other property from [Definition 2.1](#). If R is monetary, then clearly it

is finite since $-\|X\| \leq R(X) \leq \|X\|$ for any bounded X . Further, translation invariance implies that $R(X + \lambda)$ is non-increasing in $\lambda \in \mathbb{R}$.

Set-monotonicity means that adding positions to an ensemble increases its risk, that is, a set with larger cardinality will be riskier. Of course, one could also consider the property of “set anti-monotonicity”, but we leave this for future work in other contexts. This appears naturally when X arises from uncertainty—see Example 5.6. It implies $R(\dot{0}) \leq R(X - X)$ for any $X \in \text{CB}(L^\infty)$. It further implies

$$R(X \cup Y) \geq \max\{R(X), R(Y)\} \geq \min\{R(X), R(Y)\} \geq R(X \cap Y),$$

which can be thought of as quasi-concavity and quasi-convexity. The last inequality makes sense when $X \cap Y \neq \emptyset$. WC-boundedness in turn has an interpretation that, to make some set of positions acceptable, one needs less than the larger amount of money among all positions in this set, as this value also makes acceptable all the other positions in the set (the acronym WC stands for *worst-case*, see Section 4). Set-convexity means that diversification alongside positions within the set does not increase risk.

Spaces such as $\text{CB}(L^\infty)$ are subsets of *hyperspaces* (or: a space equipped with a hypertopology), i.e., a topological space comprised of all non-empty, closed subsets of a underlying topological space, so that the canonical map $x \mapsto \dot{x}$ is a homeomorphism onto its image (Lucchetti and Pasquale, 1994). This is the case for the Hausdorff distance we consider here. Since the canonical mapping is injective, clearly there is no trivial surjection from L^∞ to $\text{CB}(L^\infty)$. Thus, the approach of SRMs can not be reduced to the traditional setup for risk measures, which contemplates mappings of the form $\rho: L^\infty \rightarrow \mathbb{R}$. On the other hand, the map defined as $\rho(x) = R(\dot{x})$ yields a *bona fide* risk measure. We call ρ monetary, convex and coherent analogously as for R .

Remark 2.2. If $X \in \text{CB}(L^\infty)$, then obviously $X^c \notin \text{CB}(L^\infty)$. Thus, properties for other set operations such as $R(X^c)$, $R(X \setminus Y)$ and $R(X \Delta Y)$ are outside our proposed scope. While this kind of operation may be interesting, particularly for risk budgeting and risk sharing problems, its exploration is left for future works. Other algebraic and order properties from risk measures on random variables, such as quasi-convexity, cash-subadditivity, relevance, surplus invariance and star-shapedness, on the other hand, can be directly adapted to our framework. Besides, properties that depend on distributions and quantiles, such as law invariance and comonotonic additivity, require a more in-depth discussion on the proper probability distribution of a set of random variables—possibly beyond the collection of finite-dimensional distributions. We leave this stream of inquiry for future research.

Definition 2.3. The **acceptance set** of a SRM R is defined as

$$\mathcal{A}_R = \{X \in \text{CB}(L^\infty) : R(X) \leq 0\}. \quad (2.1)$$

Proposition 2.4. Let R be a monetary SRM. Then,

(i) \mathcal{A}_R is non-empty and satisfies the following:

(a) if $X \in \mathcal{A}_R$ and $Y \geq X$, then $Y \in \mathcal{A}_R$.

(b) $\inf\{\alpha \in \mathbb{R} : \alpha \in \mathcal{A}_R\} = 0$.

(ii) for all $X \in \text{CB}(L^\infty)$,

$$R(X) = \inf\{\alpha \in \mathbb{R} : X + \alpha \in \mathcal{A}_R\}. \quad (2.2)$$

(iii) R is lower semi-continuous if and only if \mathcal{A}_R is closed.

(iv) R is set-monotone if and only if, for each $X \in \text{CB}(L^\infty)$, the conditions $X \subseteq Y$ and $Y \in \mathcal{A}_R$ imply $X \in \mathcal{A}_R$.

(v) R is WC-bounded if and only if, for each $X \in \text{CB}(L^\infty)$, the following implication holds: if $x \in \mathcal{A}_R$ for all $x \in X$, then $X \in \mathcal{A}_R$.

(vi) R is union-bounded if and only if \mathcal{A}_R is stable under a finite number of unions of its elements.

(vii) R is convex if and only if \mathcal{A}_R is a convex set in $\text{CB}(L^\infty)$.

(viii) R is positive homogeneous if and only if \mathcal{A}_R is a cone in $\text{CB}(L^\infty)$.

(ix) R is set-convex if and only if \mathcal{A}_R is stable under the *convex hull* operation.

Proof. For item (i), the claim directly follows from the fact that R is monetary and normalized.

For item (ii), note that if $X \in \text{CB}(L^\infty)$ then $X + \alpha \in \text{CB}(L^\infty)$ for any real α . A direct calculation leads to

$$\inf\{\alpha \in \mathbb{R} : X + \alpha \in \mathcal{A}_R\} = \inf\{\alpha \in \mathbb{R} : R(X) \leq \alpha\} = R(X),$$

for any $X \in \text{CB}(L^\infty)$.

For item (iii), if R is lower semi-continuous, then $\mathcal{A}_R \equiv R^{-1}((-\infty, 0])$ is closed. Conversely, assume \mathcal{A}_R is closed. For any $\alpha \in \mathbb{R}$, we have by translation invariance that

$$\begin{aligned} & \{X \in \text{CB}(L^\infty) : R(X) \leq \alpha\} \\ &= \{X \in \text{CB}(L^\infty) : R(X + \alpha) \leq 0\} = \{X - \alpha : X \in \mathcal{A}_R\}. \end{aligned}$$

Thus, all lower sublevel sets of R are closed, which implies that it is lower semi-continuous.

For item (iv), let R be set-monotone, $X \subseteq Y \in \mathcal{A}_R$, and $X \in \text{CB}(L^\infty)$. Then, $R(X) \leq R(Y) \leq 0$, which implies $X \in \mathcal{A}_R$. For the converse, for any $\alpha \in \mathbb{R}$, if $Y + \alpha \in \mathcal{A}_R$, then $X + \alpha \in \mathcal{A}_R$. Hence, $R(X) = \inf\{\alpha \in \mathbb{R} : X + \alpha \in \mathcal{A}_R\} \leq \inf\{\alpha \in \mathbb{R} : Y + \alpha \in \mathcal{A}_R\} = R(Y)$.

For item (v), first let R be WC-bounded and let $X \in \text{CB}(L^\infty)$ be such that $\dot{x} \in \mathcal{A}_R$ for all $x \in X$. Then $R(X) \leq \sup_{x \in X} R(\dot{x}) \leq 0$ guarantees that $X \in \mathcal{A}_R$. For the “if” part of the statement, fix $\alpha \in \mathbb{R}$ and $X \in \text{CB}(L^\infty)$ such that $\{x + \alpha\} \in \mathcal{A}_R$ for all $x \in X$. Then $X + \alpha = \bigcup_{x \in X} \{x + \alpha\} \in \mathcal{A}_R$. Thus, we get

$$\begin{aligned} R(X) &= \inf\{\alpha \in \mathbb{R} : X + \alpha \in \mathcal{A}_R\} \\ &\leq \inf\{\alpha \in \mathbb{R} : \{x + \alpha\} \in \mathcal{A}_R \text{ for all } x \in X\} \\ &= \inf\{\alpha \in \mathbb{R} : R(\dot{x}) \leq \alpha\} \text{ for all } x \in X \\ &\leq \sup_{x \in X} R(\dot{x}). \end{aligned}$$

For item (vi), let $X, Y \in \mathcal{A}_R$. If $R(X \cup Y) \leq \max\{R(X), R(Y)\} \equiv 0$, then $X \cup Y \in \mathcal{A}_R$. For the converse, note that translation invariance ensures both $X + R(X)$ and $Y + R(Y)$ lie in \mathcal{A}_R . Then, by monotonicity, we have $X + \max\{R(X), R(Y)\} \in \mathcal{A}_R$ and $Y + \max\{R(X), R(Y)\} \in \mathcal{A}_R$. By hypothesis, \mathcal{A}_R is stable under finitely many unions, which implies

$$\begin{aligned} X \cup Y + \max\{R(X), R(Y)\} \\ = (X + \max\{R(X), R(Y)\}) \cup (Y + \max\{R(X), R(Y)\}) \in \mathcal{A}_R. \end{aligned}$$

Hence, $R(X \cup Y + \max\{R(X), R(Y)\}) \leq 0$ and, by translation invariance again, $R(X \cup Y) \leq \max\{R(X), R(Y)\}$.

The “only if” part of items (vii) and (viii) are both straightforward. The “if” part of both statements comes from items (vii) and (viii) in Proposition 2.6, respectively.

For item (ix), set-convexity of R implies that, for any $X \in \mathcal{A}_R$, one has $R(\text{conv}(X)) \leq R(X) \leq 0$. Hence, $\text{conv}(X) \in \mathcal{A}_R$. For the converse, it holds for any $X \in \text{CB}(L^\infty)$ and any $\alpha \in \mathbb{R}$ such that $X + \alpha \in \mathcal{A}_R$, that $\text{conv}(X + \alpha) = \text{conv}(X) + \alpha \in \mathcal{A}_R$. Thus, $R(\text{conv}(X)) \leq R(X)$. ■

We now consider the converse problem of generating a SRM from a given collection of acceptable ensembles of positions.

Definition 2.5. If $\mathcal{A} \subseteq \text{CB}(L^\infty)$ is non-empty, then its induced SRM is defined as

$$R_{\mathcal{A}}(X) = \inf\{\alpha \in \mathbb{R} : X + \dot{\alpha} \in \mathcal{A}\}, \quad X \in \text{CB}(L^\infty). \quad (2.3)$$

Proposition 2.6. Let $\mathcal{A} \subseteq \text{CB}(L^\infty)$ satisfy the following

1. if $X \in \mathcal{A}$ and $Y \geq X$, then $Y \in \mathcal{A}$.
2. $\inf\{\alpha \in \mathbb{R} : \dot{\alpha} \in \mathcal{A}\} = 0$.

Then:

- (i) $R_{\mathcal{A}}$ is a monetary SRM.
- (ii) if R is an arbitrary monetary SRM, then $R_{\mathcal{A}_R} = R$.
- (iii) If \mathcal{A} is closed, then $R_{\mathcal{A}}$ is lower semi-continuous and $\mathcal{A}_{R_{\mathcal{A}}} = \mathcal{A}$. Conversely, if $R_{\mathcal{A}}$ is lower semi-continuous, then $\mathcal{A}_{R_{\mathcal{A}}} = \text{cl}(\mathcal{A})$.
- (iv) if $X \in \mathcal{A}$ holds whenever $X \in \text{CB}(L^\infty)$ is a subset of some $Y \in \mathcal{A}$, then $R_{\mathcal{A}}$ is set-monotone.
- (v) if, for every $X \in \text{CB}(L^\infty)$, the implication holds that $X \in \mathcal{A}$ whenever $\dot{x} \in \mathcal{A}$ for all $x \in X$, then R is WC-bounded.
- (vi) if \mathcal{A} is stable under pairwise unions of its elements, then $R_{\mathcal{A}}$ is union-bounded.
- (vii) if \mathcal{A} is a convex set, then $R_{\mathcal{A}}$ is a convex SRM.
- (viii) if \mathcal{A} is a cone, then $R_{\mathcal{A}}$ is a positive homogeneous SRM.
- (ix) if \mathcal{A} is stable under the convex hull operation, then $R_{\mathcal{A}}$ is set-convex.

Proof. Item (i) is straightforward from the properties of \mathcal{A} .

Item (ii) follows from Proposition 2.4, item (ii).

For item (iii), in view of Proposition 2.4, item (iii), it suffices to show that $\mathcal{A}_{R_{\mathcal{A}}}$ is closed. Since \mathcal{A} is closed, we have that $X + R_{\mathcal{A}}(X) \in \mathcal{A}$, for any $X \in \text{CB}(L^\infty)$. We then see that

$$\mathcal{A}_{R_{\mathcal{A}}} = \{X \in \text{CB}(L^\infty) : X + \dot{\alpha} \in \mathcal{A} \text{ for some } \alpha \leq 0\} = \mathcal{A}.$$

Now, let $R_{\mathcal{A}}$ be lower semi-continuous. It follows that $\mathcal{A}_{R_{\mathcal{A}}}$, being a lower sublevel set, is closed. Take $X \in \text{cl}(\mathcal{A})$. Then, there is a sequence $\{X_n, n \geq 1\} \subseteq \mathcal{A}$ such that $X_n \rightarrow_{d_H} X$. Thus, $R_{\mathcal{A}}(X) \leq \liminf_n R_{\mathcal{A}}(X_n) \leq 0$, which implies $X \in \mathcal{A}_{R_{\mathcal{A}}}$. Hence, $\text{cl}(\mathcal{A}) \subseteq \mathcal{A}_{R_{\mathcal{A}}}$.

To see that $\mathcal{A}_{R_{\mathcal{A}}} \subseteq \text{cl}(\mathcal{A})$, let $X \in \mathcal{A}_{R_{\mathcal{A}}}$. Then $R_{\mathcal{A}}(X) \leq 0$, and thus, there is a scalar sequence $\{\alpha_n, n \geq 1\} \subseteq \mathbb{R}$ such that $\alpha_n \downarrow R_{\mathcal{A}}(X)$ and $X + \dot{\alpha}_n \in \mathcal{A}$. By hypothesis on

\mathcal{A} , we also have that $X_n = X + \dot{\alpha}_n - R_{\mathcal{A}}(X) \in \mathcal{A}$ for any $n \in \mathbb{N}$. Therefore,

$$\begin{aligned} d_H(X_n, X) &= \sup_{z \in L^\infty} \left| \inf_{x \in X} \|z - (x + \alpha_n - R_{\mathcal{A}}(X))\|_\infty - \inf_{x \in X} \|z - x\|_\infty \right| \\ &\leq \sup_{z \in L^\infty} \left| \inf_{x \in X} \|z - x\|_\infty - \inf_{x \in X} \|z - x\|_\infty \right| + |\alpha_n - R_{\mathcal{A}}(X)| \\ &= |\alpha_n - R_{\mathcal{A}}(X)| \rightarrow 0. \end{aligned}$$

Thus, we get $X_n \rightarrow_{d_H} X$, which implies $X \in \text{cl}(\mathcal{A})$, and we get the desired inclusion and, by consequence, $\mathcal{A}_{R_{\mathcal{A}}} = \text{cl}(\mathcal{A})$.

Items (iv) to (vi) follow the same steps as items (iv) to (vi) in Proposition 2.4.

For item (vii) we have, given any $\alpha_1, \alpha_2 \in \mathbb{R}$ satisfying $X + \dot{\alpha}_1, Y + \dot{\alpha}_2 \in \mathcal{A}$, that $\lambda(X + \dot{\alpha}_1) + (1 - \lambda)(Y + \dot{\alpha}_2) \in \mathcal{A}$, for all $\lambda \in [0, 1]$. Thus, $R_{\mathcal{A}}(\lambda X + (1 - \lambda)Y) \leq \lambda \alpha_1 + (1 - \lambda)\alpha_2$. Hence, by taking the infimum over those α_1 and α_2 , we get the claim.

For item (viii), let $X \in \text{CB}(L^\infty)$ and $\lambda \geq 0$. In fact, due to normalization, without loss of generality we can take $\lambda > 0$. A direct calculation yields

$$\begin{aligned} R_{\mathcal{A}}(\lambda X) &= \inf\{\alpha \in \mathbb{R} : \lambda X + \dot{\alpha} \in \mathcal{A}\} \\ &= \inf\{\alpha \in \mathbb{R} : \lambda(X + \lambda^{-1}\alpha) \in \mathcal{A}\} \\ &= \inf\{\alpha \in \mathbb{R} : X + \lambda^{-1}\alpha \in \mathcal{A}\} \\ &= \inf\{\lambda\alpha \in \mathbb{R} : X + \dot{\alpha} \in \mathcal{A}\} \\ &= \lambda R_{\mathcal{A}}(X) \end{aligned}$$

For item (ix), the claim follows the same steps as item (ix) in Proposition 2.4. \blacksquare

Remark 2.7. By Proposition 2.4 and item (iii), if \mathcal{A} is closed, then the converse implications in items (iv) to (ix) also hold. In this case, the infimum in eq. (2.3) is attained, which is always the case for the infimum in eq. (2.2).

3 Dual representations

In what follows, we shall use the term *measure* in the broad sense, allowing it to refer to both finitely- and countably additive set functions, indicating the type of additivity explicitly in each case. Nevertheless, a *probability measure* is always understood in the σ -additive sense. We denote the topological dual $(L^\infty)'$ of $L^\infty \equiv L^\infty(p)$ by $\mathbf{ba} \equiv \mathbf{ba}(\mathcal{F})$, which is the space of all finitely additive signed measures on (Ω, \mathcal{F}) that are absolutely continuous with respect to p and with finite total variation norm $\|\cdot\|_{TV}$. The canonical

bilinear form on $\mathbf{ba} \times L^\infty$ is denoted simply as $\langle q, x \rangle = \int_\Omega x dq$. Moreover, we let $\mathbf{ba}_{1,+}$ denote the subset of \mathbf{ba} comprised of finitely additive, non-negative measures q on \mathcal{F} with unit total variation, and $P^{\text{ac}} \subseteq \mathbf{ba}_{1,+}$ is the set of *probability measures* q that are absolutely continuous with respect to p , with Radon-Nikodym derivative dq/dp . With no risk of confusion, we write, for any collection $Q \subseteq \mathbf{ba}$, the **norm of** Q as $\|Q\| := \sup_{q \in Q} \|q\|_{TV}$.

We also let U denote the $\|\cdot\|_{TV}$ -unit ball on \mathbf{ba} , and $U_+ = U \cap \mathbf{ba}_+$, where \mathbf{ba}_+ are the non-negative elements in \mathbf{ba} . In addition, we let $\mathbf{ba}^r := \mathbf{ba}^r(\mathcal{A}_U)$ be the set of regular, signed finitely additive measures of bounded variation on $\mathcal{A}_U \subseteq \text{pwr}(\mathbf{ba})$, the *algebra* generated by the open subsets of U . We follow [Aliprantis and Border \(2006\)](#) and say that a finitely additive measure μ is **regular** if (i) it is finite-valued on compact sets; (ii) $\mu(A) = \sup\{\mu(K) : K \subseteq A \text{ compact}\}$, and; (iii) $\mu(A) = \inf\{\mu(V) : V \supseteq A \text{ open}\}$. Notice that an element $\mu \in \mathbf{ba}^r$ is a measure on a class of subsets of \mathbf{ba} , in particular $\mathbf{ba}^r \not\subseteq \mathbf{ba}$. We define $\mathbf{ba}_{1,+}^r$ similarly to $\mathbf{ba}_{1,+}$ and so on.

Let $\overline{\mathbb{R}} := \mathbb{R} \cup \{-\infty, \infty\}$. For each $Q \subseteq \mathbf{ba}$ we have that the functional $f_Q : \text{pwr}(L^\infty) \rightarrow \overline{\mathbb{R}}$, defined through

$$f_Q(X) := \langle Q, X \rangle := \sup_{x \in X, q \in Q} \langle q, x \rangle, \quad X \subseteq L^\infty,$$

is sub-additive, positive homogeneous, additive for constants, and satisfies the Cauchy-Schwarz type inequality $f_Q(X) \leq \|X\| \cdot \|Q\|$. Thus, it is finite when both X and Q are bounded, and monotone for any $Q \subseteq \mathbf{ba}_+$. Notice that by taking closures and convex hulls one does not alter the supremum due to the continuity and bi-linear properties of $\langle q, x \rangle$; thus, for example $f_{\text{int}(Q)}(\text{int}(X)) = f_Q(X)$. Furthermore, when $Q = \dot{q}$ is a singleton, the map is additive. Additionally, when $(\text{ess sup } X) \in X$, we have that $f_Q(X) = f_Q(\text{ess sup } X)$. We also have that $d_H(X, Y) = \sup_{q \in U} |\langle \dot{q}, X \rangle - \langle \dot{q}, Y \rangle|$ for any $X, Y \in \text{CBC}(L^\infty)$.

Remark 3.1. Most concepts and results of the paper can be adjusted for a more general base domain $\text{CB}(L^\lambda(p))$, $\lambda \in [1, \infty)$, under the replacement of \mathbf{ba} by $L^\gamma(p)$, where $1/\lambda + 1/\gamma = 1$. In this case, duality is given by $f_Y(X) = \langle X, Y \rangle = \sup_{x \in X, y \in Y} \langle p, xy \rangle$, where $X \subseteq L^\lambda$, $Y \subseteq L^\gamma$.

Before stating our first dual representation theorem below, we need the following facts and definitions. Let $C_b(U)$ be the Banach lattice comprised of all the real valued functions f defined on $(U, \|\cdot\|_{TV})$ such that f is continuous and bounded. Furthermore,

for each $X \in \text{CBC}(L^\infty)$, define $\widehat{X}: U \rightarrow \mathbb{R}$ through

$$\widehat{X}(q) := \sup_{x \in X} \langle q, x \rangle \equiv \langle \dot{q}, X \rangle \equiv f_{\dot{q}}(X), \quad q \in U. \quad (3.1)$$

A key step in the proof of Theorem 3.2 below is *Hörmander's Theorem* (see Theorem 3.2.9 in Beer, 1993), according to which $\text{CBC}(L^\infty)$ is embedded as a closed convex cone $\mathbb{C} \subseteq C_b(U)$ under the algebraic and isometric embedding $X \mapsto \widehat{X}$.

Theorem 3.2. R is a convex, lower semi-continuous, set-monotone and set-convex SRM if and only if

$$R(X) = \max_{\mu \in \mathbf{ba}_{1,+}^r} \left\{ \int_{U_+} \langle \dot{q}, -X \rangle d\mu(q) - \tau_R(\mu) \right\}, \quad X \in \text{CB}(L^\infty), \quad (3.2)$$

where $\tau_R: \mathbf{ba}^r \rightarrow [0, \infty]$ is the non-negative, convex and weak* lower semi-continuous functional defined through

$$\tau_R(\mu) = \sup_{X \in \mathcal{A}_R} \int_{U_+} \langle \dot{q}, X \rangle d\mu(q), \quad \mu \in \mathbf{ba}^r. \quad (3.3)$$

Proof. The “if” part of the claim can be directly checked, so we focus on the “only if” part. By set-monotonicity and set-convexity, we have that $R(X) = R(\text{conv}(X))$ for any $X \in \text{CB}(L^\infty)$. Thus, it is enough to understand the properties of R in $\text{CBC}(L^\infty)$, the space of non-empty, closed and convex sets in L^∞ . Note that sum and multiplication by non-negative scalar are continuous operations. Furthermore, by the Banach-Alaoglu Theorem, elements of $\text{CBC}(L^\infty)$ are weak* compact.

Now, take $C_b(U)'$ as the topological dual of $C_b(U)$, and f and f' for elements of $C_b(U)$ and $C_b(U)'$, respectively. By Theorem 14.10 in Aliprantis and Border (2006), we can identify $C_b(U)'$ with \mathbf{ba}^r . In this case, for $\mu \approx f'$, the bi-linear form is $\langle f, f' \rangle = f'(f) = \langle \mu, f \rangle = \int_U f d\mu$.

From the Fenchel-Moreau Conjugation Theorem, a functional $\Pi: C_b(U) \rightarrow \mathbb{R} \cup \{\infty\}$ is proper, convex and lower semi-continuous if and only if

$$\Pi(f) = \sup_{f' \in C_b(U)'} \{f'(f) - \Pi'(f')\},$$

where $\Pi'(f') = \sup_{f \in C_b(U)} \{f'(f) - \Pi(f)\}$. Let then

$$\Pi(f) = \begin{cases} R(X), & \text{if } f = \widehat{X} \text{ for some } X \in \text{CBC}(L^\infty) \\ \infty, & \text{otherwise.} \end{cases}$$

Thus, $R|_{\text{CBC}(L^\infty)}(X) = \Pi|_{\mathbb{C}}(\widehat{X})$, where we recall that \mathbb{C} denotes the range of $X \mapsto \widehat{X}$. We then get, for any $X \in \text{CB}(L^\infty)$, that

$$\begin{aligned} R(X) &= R(\text{conv}(X)) \\ &= \sup_{f' \in C_b(U)'} \{f'(\text{conv}(X)) - \Pi'(f')\} \\ &= \sup_{\mu \in \mathbf{ba}_{1,+}^r} \left\{ \int_{U_+} \langle \dot{q}, \text{conv}(X) \rangle d\mu(q) - \Pi'(\mu) \right\} \\ &= \sup_{\mu \in \mathbf{ba}_{1,+}^r} \left\{ \int_{U_+} \langle \dot{q}, X \rangle d\mu(q) - \tau_R(\mu) \right\}, \end{aligned}$$

where

$$\begin{aligned} \Pi'(\mu) &= \sup_{f \in C_b(U)} \{f'(f) - \Pi(f)\} \\ &= \sup_{X \in \text{CB}(L^\infty)} \left\{ \int_{U_+} \langle \dot{q}, \text{conv}(X) \rangle d\mu(q) - R(\text{conv}(X)) \right\} \\ &= \sup_{X \in \mathcal{A}_R} \int_{U_+} \langle \dot{q}, \text{conv}(X) \rangle d\mu(q) \\ &= \sup_{X \in \mathcal{A}_R} \int_{U_+} \langle \dot{q}, X \rangle d\mu(q). \end{aligned}$$

The last two equalities in both deductions, as well as the integral being over U_+ , arise from monotonicity and translation invariance of R . Concerning properties of τ_R , convexity is straightforward. Non-negativity follows since $\dot{0} \in \mathcal{A}_R$, and weak* lower semi-continuity is due the fact that τ_R is the pointwise supremum over \mathbb{C} of the weak* continuous linear maps $f' \mapsto f'(f)$. Since $\mathbf{ba}_{1,+}^r$ is weak* compact by Alaoglu's Theorem, and $\mu \mapsto \int_{U_+} \langle \dot{q}, -X \rangle d\mu(q) - \tau_R(\mu)$ is weak* upper semi-continuous, the supremum is attained. \blacksquare

Remark 3.3. By the Extended Namioka Theorem (see [Biagini and Frittelli, 2010](#); [Kaina and Rüschemdorf, 2009](#)) we get that, if a functional $\Pi: C_b(U) \rightarrow \mathbb{R} \cup \{\infty\}$ is proper, convex, and monotone, then Π is continuous on $\text{int}(\text{dom } \Pi)$. Thus, if R is a convex, set-monotone and

set-convex SRM, we have that it is continuous on $A = T^{-1}(\text{int}(\mathbb{C}))$, where T is the embedding from $\text{CBC}(L^\infty)$ into $C_b(U)$ appearing in the proof of the theorem. In particular, a dual representation as that in eq. (3.2) holds for any $X \in A$.

Remark 3.4. Given any $\mu \in \mathbf{ba}^r$, that is, $\mu: \mathcal{A}_U \rightarrow \overline{\mathbb{R}}$ is regular, finitely additive and of bounded variation, it can be extended to a finitely additive measure μ^{ext} defined on $\text{pwr}(U)$. The existence of such extensions follows immediately from the Hahn-Banach Extension Theorem applied to the subspace $C_b(U)$ of $B(U)$, the space of all bounded real functions on U —see [Luxemburg \(1991\)](#) and Corollary 14.11 of [Aliprantis and Border \(2006\)](#), for instance. Furthermore, positivity and regularity are preserved in the extension (Theorem 2.1 in [Bachman and Sultan, 1980](#)). Let then $\mathbf{ba}^{\text{ext}} \subseteq \mathbf{ba}(\text{pwr}(U))$ be the set of such extensions. It is convex and weak* compact (i.e., in the $\sigma(\mathbf{ba}(\text{pwr}(U)), B(U))$ topology). Since any $f \in C_b(U)$ also belongs to $B(U)$, a convex duality argument similar to that in Theorem 3.2 leads to a representation of R over \mathbf{ba}^{ext} as

$$R(X) = \max_{\mu \in \mathbf{ba}_{1,+}^{\text{ext}}} \left\{ \int_{U_+} \langle \dot{q}, -X \rangle d\mu(q) - \tau_R(\mu) \right\}, \quad X \in \text{CB}(L^\infty).$$

Definition 3.5. Let X, X_1, X_2, \dots be elements in $\text{CB}(L^\infty)$, and let R be a SRM. We say that R is:

- (i) **continuous from above** if it holds that $R(X_n) \rightarrow R(X)$ whenever $\widehat{X}_n|_{U_+} \downarrow \widehat{X}|_{U_+}$ pointwise.
- (ii) **continuous from below** if it holds that $R(X_n) \rightarrow R(X)$ whenever $\widehat{X}_n|_{U_+} \uparrow \widehat{X}|_{U_+}$ pointwise.
- (iii) **Fatou continuous** if it holds that $R(X) \leq \liminf_n R(X_n)$ whenever $\{\widehat{X}_n|_{U_+}, n \geq 1\}$ is a bounded sequence in $C_b(U_+)$ such that $\widehat{X}_n|_{U_+} \rightarrow \widehat{X}|_{U_+}$ pointwise.
- (iv) **Lebesgue continuous** if it holds that $R(X_n) \rightarrow R(X)$ whenever $\{\widehat{X}_n|_{U_+}, n \geq 1\}$ is a bounded sequence in $C_b(U_+)$ such that $\widehat{X}_n|_{U_+} \rightarrow \widehat{X}|_{U_+}$ pointwise.
- (v) **represented over probabilities** if, for every $X \in \text{CB}(L^\infty)$,

$$R(X) = \max_{\mu \in \mathcal{P}^{\text{ext}}} \left\{ \int_{U_+} \langle \dot{q}, -X \rangle d\mu(q) - \tau_R(\mu) \right\}, \quad (3.4)$$

where $\mathcal{P}^{\text{ext}} := \mathcal{P}^{\text{ext}}(\text{pwr}(U))$ are the probability measures over $(U, \text{pwr } U)$ such that $\mu|_{\mathcal{A}_U} \in \mathbf{ba}^r$.

Remark 3.6. By Ulam's Theorem (see Theorem 1.12.40 and Corollary 1.12.41 in [Bogachev, 2006](#)), under the Continuum Hypothesis, \mathcal{P}^{ext} contains only discrete probability mea-

tures. Thus, for example, the integrals appearing in eq. (3.4) are actually sums over q , with q varying in the support of μ .

Theorem 3.7. Let R be a convex, lower semi-continuous, set-monotone and set-convex SRM. Then the following hold:

- (i) R is Fatou continuous if and only if it is continuous from above.
- (ii) R is Lebesgue continuous if and only if it is continuous from above and continuous from below.
- (iii) if R is represented over probabilities, then R is Lebesgue continuous.

Proof. For the “only if” part of item (i), let R be Fatou continuous and assume $\widehat{X}_n|_{U_+} \downarrow \widehat{X}|_{U_+}$ pointwise. Notice that $\{\widehat{X}_n|_{U_+}, n \geq 1\}$ is bounded since $\widehat{X}_1 \geq \widehat{X}_n \geq \widehat{X}$ on U_+ . Monotonicity and Fatou continuity then imply, by Theorem 3.2, that

$$\limsup_n R(X_n) \leq R(X) \leq \liminf_n R(X_n).$$

For the “if” part, let R be continuous from above and a bounded sequence $\{\widehat{X}_n|_{U_+}, n \geq 1\}$ in $C_b(U_+)$ such that $\widehat{X}_n|_{U_+} \rightarrow \widehat{X}|_{U_+}$ pointwise. Notice that, letting $Z := \bigcup_{n \in \mathbb{N}} X_n$, one has $\widehat{Z} \in C_b(U_+)$. Define, for $\alpha \in \mathbb{N}$, $y_\alpha = \text{ess sup } \bigcup_{n \geq \alpha} X_n$. Then, we get for any $q \in U_+$ that $\langle q, y_1 \rangle, \langle q, y_2 \rangle, \dots$ is non-increasing and $\langle q, y_n \rangle \geq \langle q, X_n \rangle$. Thus, $\langle q, y_n \rangle \downarrow \langle q, X \rangle$. Continuity from above and monotonicity then imply

$$R(X) = \liminf_n R(y_n) \leq \liminf_n R(X_n).$$

Regarding item (ii), let R be Lebesgue continuous. Take $\widehat{X}_n|_{U_+} \uparrow \widehat{X}|_{U_+}$ pointwise and notice that this implies that $\{\widehat{X}_n|_{U_+}, n \geq 1\}$ is a bounded sequence in $C_b(U_+)$. Lebesgue continuity then implies $R(X_n) \rightarrow R(X)$. A similar reasoning with $\widehat{X}_n|_{U_+} \uparrow \widehat{X}|_{U_+}$ leads to continuity from above. For the converse implication, let R be continuous from below, and let $\widehat{X}_n|_{U_+} \uparrow \widehat{X}|_{U_+}$ pointwise. Again, this yields that $\{\widehat{X}_n|_{U_+}, n \geq 1\}$ is a bounded sequence in $C_b(U_+)$. Similarly to item (ii), we can show that $R(X) \geq \limsup_n R(X_n)$. Further, by item (i) we have that R is Fatou continuous, which grants $R(X) \leq \liminf_n R(X_n)$. Hence, R is Lebesgue continuous.

For item (iii), assume R is represented over probabilities. Take $\widehat{X}_n|_{U_+} \rightarrow \widehat{X}|_{U_+}$ pointwise such that $\{\widehat{X}_n, n \geq 1\}$ is bounded in $C_b(U_+)$. By the Dominated Convergence

Theorem, we have that

$$\begin{aligned}
R(X) &= \max_{\mu \in \mathcal{P}^{\text{ext}}} \left\{ \lim_n \int_{U_+} \langle \dot{q}, -X_n \rangle d\mu(q) - \tau_R(\mu) \right\} \\
&\leq \liminf_n \max_{\mu \in \mathcal{P}^{\text{ext}}} \left\{ \int_{U_+} \langle \dot{q}, -X_n \rangle d\mu(q) - \tau_R(\mu) \right\} \\
&= \liminf_n R(X_n).
\end{aligned}$$

Hence, R is Fatou continuous. By item (i), it is continuous from above. Now, take $\{X_n, n \geq 1\}$ such that $\widehat{X}_n \uparrow \widehat{X}$. Assume, without loss of generality (due to translation invariance), that $\widehat{X}_n|_{U_+} \leq 0$. Let for each $k \in \mathbb{R}$ the set $L_k = \{\mu \in \mathbf{ba}_{1,+}^{\text{ext}} : \tau(\mu) \leq k\}$, and notice that such sets are weak* compact as they are bounded, closed and convex. Moreover, the mappings $\Lambda_n(\mu) := \int_{U_+} \langle \dot{q}, -X_n \rangle d\mu(q)$ are weak* continuous over each L_k . Since, $\Lambda_n(\mu) \uparrow \Lambda(\mu) := \int_{U_+} \langle \dot{q}, -X \rangle d\mu(q)$, we have by Dini's Lemma that Λ_n converges uniformly to Λ . We then obtain for any $k > 0$ that

$$\max_{\mu \in L_k} \left\{ \int_{U_+} \langle \dot{q}, -X_n \rangle d\mu(q) - \tau_R(\mu) \right\} \rightarrow \max_{\mu \in L_k} \left\{ \int_{U_+} \langle \dot{q}, -X \rangle d\mu(q) - \tau_R(\mu) \right\}.$$

Now, note that for each $n \in \mathbb{N}$ and each $X \in \text{CB}(L^\infty)$ we have

$$R(X) \leq R(X_n) = \max_{\mu \in \mathcal{P}^{\text{ext}}} \left\{ \int_{U_+} \langle \dot{q}, -X_n \rangle d\mu(q) - \tau_R(\mu) \right\}.$$

Since the linear part $\int_{U_+} \langle \dot{q}, -X_n \rangle d\mu(q)$ is non-positive for any $\mu \in \mathcal{P}^{\text{ext}}$, we have that the supremum in the representation is not altered if taken only over those μ such that $\tau_R(\mu) \leq \|X\| - R(X)$, i.e., $\mu \in L_{\|X\| - R(X)}$. Then, by the uniform convergence of Λ_n to Λ we obtain that $\lim_n R(X_n) = R(X)$. Hence, R is continuous from below. Hence, from item (ii) it is Lebesgue continuous. \blacksquare

Remark 3.8. While Fatou continuity is necessary for a dual representation under probabilities, it is, in general, not sufficient. Nonetheless, when U_+ is Polish, the recent papers of [Delbaen \(2022\)](#) and [Nendel \(2024\)](#) show that, in the setting of traditional risk measures, under some technical conditions we have that Fatou continuity is equivalent to a representation over the supremum of probabilities in the Borel sigma-algebra of U_+ .

Corollary 3.9. R is a coherent, lower semi-continuous, set-monotone and set-convex

SRM if and only if

$$R(X) = \max_{\mu \in \mathbf{ba}_{1,+R}^r} \int_{U_+} \langle \dot{q}, -X \rangle d\mu(q), \quad X \in \text{CB}(L^\infty), \quad (3.5)$$

where $\mathbf{ba}_{1,+R}^r$ is non-empty, convex and weak* closed, defined as

$$\begin{aligned} \mathbf{ba}_{1,+R}^r &= \left\{ \mu \in \mathbf{ba}_{1,+}^r : \int_{U_+} \langle \dot{q}, -X \rangle d\mu(q) \leq R(X) \text{ for all } X \in \text{CB}(L^\infty) \right\} \\ &= \{ \mu \in \mathbf{ba}_{1,+}^r : \tau_R(\mu) = 0. \} \end{aligned}$$

Furthermore, R is Lebesgue continuous if $\mathbf{ba}_{1,+R}^r \subseteq \mathcal{P}^{\text{ext}}$.

Proof. From Theorem 3.2, positive homogeneity for R implies that τ_R can take values only in $\{0, \infty\}$. Finiteness of R then assures the claim. The properties of $\mathbf{ba}_{1,+R}^r$ are directly inherited from those of τ_R . The last claim is directly obtained from Theorem 3.7. \blacksquare

4 Worst-case SRMs

In this section, we will focus on set risk measures R that are of the *worst-case* type, that is, when $R(X) = \sup_{x \in X} R(x)$ for all $X \in \text{CB}(L^\infty)$. Before proceeding, we begin recalling the notion of *Painlevé-Kuratowski convergence* (Aubin and Frankowska, 2009; Beer, 1993). Let X, X_1, X_2, \dots be subsets of L^∞ . We say that X_n **set-converges** to X , written $X_n \xrightarrow{s} X$ or $\lim_n X_n = X$, if X is closed and $d(\dot{x}, X_n) \rightarrow d(\dot{x}, X)$ for any $x \in L^\infty$. For instance, a sequence of closed balls converge: if $x_n \rightarrow x$ and $\epsilon_n \rightarrow \epsilon$, then $\lim_n \{y \in L^\infty : \|x_n - y\|_\infty \leq \epsilon_n\} = \{y \in L^\infty : \|x - y\|_\infty \leq \epsilon\}$. Notice that if $d_H(X_n, X) \rightarrow 0$, then $\lim_n X_n = \text{cl}(X)$, as seen from Corollary 5.1.11 and Theorem 5.2.10 in Beer (1993). However, the converse implication may not be true. Both convergences coincide if and only if the space is compact, and since L^∞ is not compact as a Banach space, these two notions of convergence do not agree in general (Beer, 1993). Additionally, the **inner limit** and the **outer limit** of $\{X_n, n \geq 1\}$ are defined, respectively, as

$$\lim \text{inn}_n \{X_n\} = \{x \in L^\infty : \limsup_n d(\dot{x}, X_n) = 0\}$$

and

$$\lim \text{out}_n \{X_n\} = \{x \in L^\infty : \liminf_n d(\dot{x}, X_n) = 0\}.$$

We shall use the following properties without further mention, see [Aubin and Frankowska \(2009\)](#) for details:

1. $\lim \text{inn}_n \{X_n\} \subseteq \lim \text{out}_n \{X_n\}$.
2. $x \in \lim \text{inn}_n \{X_n\}$ if and only if, for each n there is an $x_n \in X_n$ such that $x_n \rightarrow x$.
3. $x \in \lim \text{out}_n \{X_n\}$ if and only if there exist integers $k_1 < k_2 < \dots$ and random variables $x_{k_n} \in X_{k_n}$ such that $x_{k_n} \rightarrow x$.
4. $\lim \text{inn}_n \{X_n\}$ and $\lim \text{out}_n \{X_n\}$ are closed sets.
5. $\lim_n X_n = X$ if and only if $\lim \text{inn}_n \{X_n\} = \lim \text{out}_n \{X_n\} = X$.
6. if $\{X_n\}$ is increasing (i.e., $X_n \subseteq X_{n+1}$ for all n), then $\lim_n X_n = \text{cl}(\cup_{n=1}^{\infty} X_n)$
7. if $\{X_n\}$ is decreasing (i.e., $X_n \supseteq X_{n+1}$ for all n), then $\lim_n X_n = \cap_{n=1}^{\infty} \text{cl}(X_n)$.

Now let, for any $f: L^\infty \rightarrow \overline{\mathbb{R}}$, its **proper domain**, **epi-graph** and **hypo-graph** be defined, respectively, as $\text{dom } f = \{x \in L^\infty : f(x) > -\infty\}$, $\text{epi } f = \{(x, r) \in L^\infty \times \mathbb{R} : f(x) \leq r\}$ and $\text{hyp } f = \{(x, r) \in L^\infty \times \mathbb{R} : f(x) \geq r\}$. When we have $\text{epi } f_n \xrightarrow{s} \text{epi } f$ we say that $\{f_n, n \geq 1\}$ **epi-converges** to f , and similarly for **hyp-convergence**. In order to link epi-convergence to convergence of the infimum, we need the following definition. We say that a sequence of functions $\{f_n: L^\infty \rightarrow \overline{\mathbb{R}}, n \geq 1\}$ is **tight** if, for each $\epsilon > 0$, there is a compact set $X_\epsilon \subseteq L^\infty$ and $N_\epsilon \in \mathbb{N}$ such that, for all $n \geq N_\epsilon$, it holds that

$$\sup_{x \in X_\epsilon} f_n(x) \geq \sup_{x \in L^\infty} f_n(x) - \epsilon.$$

The functions **epi-converge tightly** if, in addition to being tight, they also epi-converge to some function. It is straightforward to verify that when $\{f_n: L^\infty \rightarrow \overline{\mathbb{R}}, n \geq 1\}$ is contained in a compact set, then it is tight. Such property implies that hypo-convergence implies convergence and finiteness of the supremum. This is formalized in the lemma below.

Lemma 4.1 (Proposition 7.3.5 of [Aubin and Frankowska \(2009\)](#)). Let $f: L^\infty \rightarrow \overline{\mathbb{R}}$ and $\{f_n: L^\infty \rightarrow \overline{\mathbb{R}}, n \geq 1\}$ be proper. If $\text{epi } f_n \xrightarrow{s} \text{epi } f$, then we have the following:

- (i) $\limsup_n \inf_{\{x \in L^\infty\}} f_n(x) \leq \inf_{\{x \in L^\infty\}} f(x)$.
- (ii) if some sub-sequence $\{x_{k_n} \in \arg \min_{\{x \in L^\infty\}} f_{k_n}, n \geq 1\}$ converges, then we have $\inf_{\{x \in L^\infty\}} f_{k_n}(x) \rightarrow \inf_{\{x \in L^\infty\}} f(x)$.
- (iii) $\{f_n: L^\infty \rightarrow \overline{\mathbb{R}}, n \geq 1\}$ is tight if and only if $\inf_{\{x \in L^\infty\}} f_n(x) \rightarrow \inf_{\{x \in L^\infty\}} f(x)$.

Likewise, if $\text{hyp } f_n \xrightarrow{s} \text{hyp } f$, then we have the following:

- (i') $\liminf_n \sup_{\{x \in L^\infty\}} f_n(x) \geq \sup_{\{x \in L^\infty\}} f(x)$.

- (ii') if some sub-sequence $\{x_{k_n} \in \arg \max_{x \in L^\infty} f_{k_n}, n \geq 1\}$ converges, then we have $\sup_{\{x \in L^\infty\}} f_{k_n}(x) \rightarrow \sup_{\{x \in L^\infty\}} f(x)$.
- (iii') $\{f_n: L^\infty \rightarrow \overline{\mathbb{R}}, n \geq 1\}$ is tight if and only if $\sup_{\{x \in L^\infty\}} f_n(x) \rightarrow \sup_{\{x \in L^\infty\}} f(x)$.

Theorem 4.2. Let R be a SRM. Then R is set-monotone and WC-bounded if and only if the **worst-case representation**

$$R(X) = \sup_{x \in X} R(\dot{x}) \tag{4.1}$$

holds for all $X \in \text{CB}(L^\infty)$. In this case, R also fulfills union-boundedness. Further, under these equivalent conditions, R is monetary if and only if the restriction of R to singletons is monetary. In the latter scenario, the following holds:

- (i) R is lower semi-continuous.
- (ii) The acceptance set of R is given by the condition $X \in \mathcal{A}_R$ if and only if $R(\dot{x}) \leq 0$ for all $x \in X$.
- (iii) The bound $|R(X) - R(Y)| \leq \|X - Y\|$ holds for all pairs $X, Y \in \text{CB}(L^\infty)$.

Proof. The claim follows from Proposition 2.4 because R being set-monotone and WC-bounded is equivalent to $X \in \mathcal{A}_R$ if and only if $\dot{x} \in \mathcal{A}_R$ for any $x \in X$ and any $X \in \text{CB}(L^\infty)$. For union-boundedness, we have that

$$R(X \cup Y) = \sup_{x \in X \cup Y} R(\dot{x}) = \max \left\{ \sup_{x \in X} R(\dot{x}), \sup_{x \in Y} R(\dot{x}) \right\} = \max\{R(X), R(Y)\}.$$

Moreover, the equivalence for monotonicity and translation invariance is directly obtained.

To prove R fulfills lower semi-continuity, it is sufficient to show that it is true for set convergence, as this implies the property w.r.t. d_H . Let then $X_n \xrightarrow{s} X$. Define, for each $n \in \mathbb{N}$, the mappings $f, f_n: L^\infty \rightarrow \overline{\mathbb{R}}$ through

$$f_n(x) = \begin{cases} R(\dot{x}), & x \in X_n \\ -\infty, & \text{otherwise} \end{cases} \quad \text{and} \quad f(x) = \begin{cases} R(\dot{x}), & x \in X \\ -\infty, & \text{otherwise.} \end{cases}$$

We show that $\text{hyp } f_n \xrightarrow{s} \text{hyp } f$. Let $(x, r) \in \text{hyp } f$. Then, $f(x) = R(\dot{x}) \geq r$. Since $X_n \xrightarrow{s} X$, there is both $x_n \in X_n$ and $x_{k_n} \in X_{k_n}$ for all $n \in \mathbb{N}$ such that $x_n \rightarrow x$ and $x_{k_n} \rightarrow x$. We have that $(x_n, R(\dot{x}_n)) \in \text{hyp } f_n$. Thus, $(x_n, R(\dot{x}_n) - (R(\dot{x}) - r)) \in \text{hyp } f_n$. Continuity of $x \mapsto R(\dot{x})$, which holds since it is monetary (see Lemma 4.3 in Follmer and Schied, 2016) then implies $R(\dot{x}_n) \rightarrow R(\dot{x}) \geq r$. Therefore, $(x_n, R(\dot{x}_n) - (R(\dot{x}) - r)) \rightarrow (x, r)$. A

similar reasoning holds for to $\{x_{k_n}\}$. Thus, $\lim \text{inn}_n\{\text{hyp } f_n\} = \lim \text{out}_n\{\text{hyp } f_n\} = \text{hyp } f$. Such equality implies $\lim_n \text{hyp } f_n = \text{hyp } f$. Hence, by item 1 in Lemma 4.1 we have the following

$$\begin{aligned} \liminf_n R(X_n) &= \liminf_n \sup_{x \in X_n} R(\dot{x}) = \liminf_n \sup_{x \in X} f_n(x) \\ &\geq \sup_{x \in X} f(x) = \sup_{x \in X} R(\dot{x}) = R(X). \end{aligned}$$

Hence, we have the desired lower semi-continuity. The deduction for \mathcal{A}_R is straightforward. Regarding the remaining inequality, let $\|X - Y\| < \infty$. We then have, by Lipschitz continuity, that

$$\begin{aligned} |R(X) - R(Y)| &= \left| \sup_{x \in X} R(\dot{x}) - \sup_{y \in Y} R(\dot{y}) \right| \\ &\leq \sup_{x \in X, y \in Y} |R(\dot{x}) - R(\dot{y})| \\ &\leq \sup_{x \in X, y \in Y} \|x - y\|_\infty \\ &= \sup_{z \in X - Y} \|z\|_\infty = \|X - Y\|. \end{aligned}$$

This completes the proof. ■

Remark 4.3. Under finiteness, WC-boundedness is equivalent to the following implication: if, for some real scalar α , $R(\dot{x}) \leq \alpha$ for all $x \in X$ then $R(X) \leq \alpha$. Both these properties are implied by the stronger one, which we call **element monotonicity**: for all $X, Y \in \text{CB}(L^\infty)$, if $R(\dot{x}) \leq R(\dot{y})$ for any $x \in X$ and any $y \in Y$, then $R(X) \leq R(Y)$. That this is indeed a stronger condition can be checked by taking $Y = \{-\sup_{x \in X} R(\dot{x})\}$. In addition, under set-monotonicity, Theorem 4.2 implies all these properties are equivalent, since the worst-case representation (4.1) implies element monotonicity.

Remark 4.4. In view of the last theorem, under the worst-case representation there is a link between SRMs and usual, univariate risk measures on L^∞ . Indeed, let ρ be a monetary risk measure on L^∞ , and suppose R is defined on $\text{CB}(L^\infty)$ through $R(X) = \sup_{x \in X} \rho(x)$. By recalling that the acceptance set of ρ is $A_\rho = \{x \in L^\infty : \rho(x) \leq 0\}$, we have by Theorem 4.2 that $\mathcal{A}_R = \{X \in \text{CB}(L^\infty) : X \subseteq A_\rho\}$, that is, $\mathcal{A}_R = \text{pwr}(A_\rho) \cap \text{CB}(L^\infty)$ or yet, for $X \in \text{CB}(L^\infty)$ one has $X \in \mathcal{A}_R$ if and only if $X \subseteq A_\rho$.

Theorem 4.5. Let R be a monetary, set-monotone, and WC-bounded SRM. Then, the following are equivalent:

- (i) R is convex.
- (ii) $x \mapsto R(\hat{x})$ is convex.
- (iii) R is set-convex.
- (iv) it holds, for every $X \in \text{CB}(L^\infty)$, that

$$R(X) = \max_{q \in \mathbf{ba}_{1,+}} \{ \langle \dot{q}, -X \rangle - \tau_R(\delta_q) \}, \quad (4.2)$$

where τ_R is as in (3.3) and δ_q is the Dirac measure concentrated at q .

In particular, R is coherent if and only if $x \mapsto R(\hat{x})$ is coherent, and this is equivalent to the representation

$$R(X) = \max_{q \in \mathbf{ba}_{1,+R}} \langle \dot{q}, -X \rangle, \quad X \in \text{CB}(L^\infty), \quad (4.3)$$

where

$$\begin{aligned} \mathbf{ba}_{1,+R} &= \{q \in \mathbf{ba}_{1,+} : \langle \dot{q}, -X \rangle \leq R(X) \text{ for all } X \in \text{CB}(L^\infty)\} \\ &= \{q \in \mathbf{ba}_{1,+} : \tau_R(\delta_q) = 0\}. \end{aligned}$$

Proof. Regarding (i) \Leftrightarrow (ii), the equivalence is straightforward. For (ii) \Leftrightarrow (iii), notice that taking convex combinations does not alter the supremum that represents R , which implies $R(X) = R(\text{conv}(X))$ for any $X \in \text{CB}(L^\infty)$. For the converse, let $x, y \in L^\infty$ and $\lambda \in [0, 1]$. Then,

$$R(\{\lambda x + (1 - \lambda)y\}) \leq R(\text{conv}(\{x, y\})) \leq R(\{x, y\}) = \max\{R(\hat{x}), R(\hat{y})\}.$$

Thus, $x \mapsto R(\hat{x})$ is quasi-convex. By translation invariance, it is a convex risk measure in L^∞ . Additionally, we have (i) \Leftrightarrow (iii) by noticing that

$$R(\lambda X + (1 - \lambda)Y) \leq R(\text{conv}(X \cup Y)) \leq R(X \cup Y) = \max\{R(X), R(Y)\}.$$

Before proving the representation, we need the following definitions: given a risk-measure ρ in L^∞ with acceptance set $A_\rho = \rho^{-1}(-\infty, 0]$, the corresponding **penalty term** is the mapping $\tau_\rho: \mathbf{ba}_{1,+} \rightarrow \mathbb{R}_+ \cup \{\infty\}$, defined through $\tau_\rho(q) = \sup_{x \in A_\rho} \langle q, -x \rangle$. As is well known in the literature, the penalty term is convex and lower semi-continuous in the total variation norm. Theorem 4.33 in Follmer and Schied (2016)] states that ρ is

a convex risk measure if and only if it can be represented as

$$\rho(x) = \max_{q \in \mathbf{ba}_{1,+}} \{ \langle q, -x \rangle - \tau_\rho(q) \}, \quad x \in L^\infty.$$

In this setup, assume (iv) holds. Then it is clear that (4.2) defines a set-monotone and WC-bounded, convex and set-convex SRM. Thus, the equivalences in items (i) to (iii) hold. For the converse implication, let $x \mapsto R(\dot{x}) =: \rho(x)$. By Theorem 4.2 we have that R is lower semi-continuous. We then get that $R(X) = \sup_{x \in X} R(\dot{x})$ holds if and only if

$$\begin{aligned} R(X) &= \sup_{x \in X} \sup_{q \in \mathbf{ba}_{1,+}} \{ \langle q, -x \rangle - \tau_\rho(q) \} \\ &= \sup_{q \in \mathbf{ba}_{1,+}} \left\{ \sup_{x \in X} \langle q, -x \rangle - \tau_\rho(q) \right\} \\ &= \sup_{q \in \mathbf{ba}_{1,+}} \{ \langle \dot{q}, -X \rangle - \tau_\rho(q) \}. \end{aligned}$$

Regarding the penalty term, recall from Theorem 4.2 that, for $X \in \text{CB}(L^\infty)$, one has $X \in \mathcal{A}_R$ if and only if $\dot{x} \in \mathcal{A}_R$ for all $x \in X$. Thus, the penalty term is given by

$$\begin{aligned} \tau_\rho(q) &= \sup_{\dot{x} \in \mathcal{A}_R} \langle q, -x \rangle \\ &= \sup_{X \in \mathcal{A}_R} \sup_{x \in X} \langle q, -x \rangle \\ &= \sup_{X \in \mathcal{A}_R} \langle \dot{q}, -X \rangle = \tau_R(\delta_q). \end{aligned}$$

Finally, for the coherent case, the equivalence for positive homogeneity is clear. From (4.2) we get that τ_R only takes 0 and ∞ as possible values, and then

$$\begin{aligned} \mathbf{ba}_{1,+R} &= \{ q \in \mathbf{ba}_{1,+} : \tau_R(q) = 0 \} \\ &= \{ q \in \mathbf{ba}_{1,+} : \langle \dot{q}, -X \rangle \leq R(X) \text{ for all } X \in \text{CB}(L^\infty) \}. \end{aligned}$$

It is straightforward, from linearity and continuity of $q \mapsto \langle \dot{q}, -X \rangle$, that $\mathbf{ba}_{1,+R}$ is non-empty, closed and convex. This concludes the proof. \blacksquare

Remark 4.6. The connection between eqs. (3.2) and (4.2) is that the latter is equivalent to the former by taking the supremum over $\{ \mu \in \mathbf{ba}_{1,+}^r : \mu = \delta_q, q \in \mathbf{ba}_{1,+} \}$. Since $\langle Q, X \rangle = \sup_{q \in Q} \langle \dot{q}, X \rangle$, the representation (4.2) can be taken over $\text{CB}(\mathbf{ba})$ as

$$R(X) = \max_{Q \in \text{CB}(\mathbf{ba}_{1,+})} \{ \langle Q, -X \rangle - \tau_R(Q) \}, \quad X \in \text{CB}(L^\infty).$$

When $x \mapsto R(\dot{x})$ is Fatou continuous, the representations in the last theorem are taken over probability measures. In this case, one has to replace the maximum for a supremum. If in addition $x \mapsto R(\dot{x})$ is Lebesgue continuous, then the supremum over probability measures is attained.

Remark 4.7. A question of interest here is whether the supremum in the worst-case formulation is attained. Under monotonicity, we have $\sup_{x \in X} R(\dot{x}) \leq R(\text{ess inf } X)$ for any $X \in \text{CB}(L^\infty)$. The converse inequality holds when $\text{ess inf } X \in X$. Thus, under the additional assumption of set-monotonicity, we get

$$R(X) = \max_{x \in X} R(\dot{x}) = R(\text{ess inf } X).$$

Alternatively, if $x \mapsto R(\dot{x})$ is convex, then we have that

$$R(X) = R(\text{conv}(X)) = \sup_{x \in \text{conv}(X)} R(\dot{x}), \quad X \in \text{CB}(L^\infty).$$

Noticing that $\text{conv}(X) \in \text{CB}(L^\infty)$ for any $X \in \text{CB}(L^\infty)$, it is clear that X is weak* compact. Then, when $x \mapsto R(\dot{x})$ is weak* upper semi-continuous, which is equivalent to Lebesgue continuity, we have that the supremum is then attained for any X .

5 Examples

In the examples below, $R: \text{CB}(L^\infty) \rightarrow \mathbb{R}$ stands for a SRM, while $\rho: L^\infty \rightarrow \mathbb{R}$ is a traditional risk measure. Additionally, for any metric space S , we write $\text{Fin}(S)$ and $\text{K}(S)$ as the collection of finite (respectively, compact) subsets of S .

We know from Theorem 4.2 and Remark 4.4 that it is possible to construct a SRM R from a given risk measure ρ via $R(X) := \sup_{x \in X} \rho(x)$. Our first set of examples explores this construction for a few well-known risk measures in the literature.

Example 5.1 (Expected loss). The **expected loss** (EL) is a coherent risk measure defined through $\text{EL}_q(x) = -\langle q, x \rangle$, for some fixed probability $q \in P^{\text{ac}}$ and allowable x . The corresponding worst-case SRM is

$$R_{\text{EL}}(X) = \sup_{x \in X} \text{EL}_q(x), \quad X \in \text{CB}(L^\infty).$$

Clearly, this SRM satisfy all properties on Definition 2.1, additionally, as $R_{\text{EL}}(-X) = \widehat{X}(q)$, the continuities on Definition 3.5 are also satisfied. This SRM can also serve as basis to define any coherent traditional risk measure ρ , as a SRM thought the choice of

X , as $\rho(x) = \sup_{q \in Q} \langle q, -x \rangle = \sup_{q \in Q} \langle p, -x \frac{dq}{dp} \rangle$ for some set $Q \subseteq P^{\text{ac}}$, we can write $\rho(x) = R_{\text{EL}}(X_x)$, where $X_x = \{x \frac{dq}{dp} : q \in Q\}$.

Example 5.2 (Expected shortfall). Two *tail risk measures* that are both the most studied in the literature and the most widely applied in financial practice are the **Value-at-Risk** (VaR) and the **Expected shortfall** (ES). They are defined, for $\alpha \in (0, 1)$ and allowable x , as $\text{VaR}^\alpha(x) = -F_x^{-1}(\alpha)$ and $\text{ES}^\alpha(x) = \frac{1}{\alpha} \int_0^\alpha \text{VaR}^s(x) ds$, respectively. ES is coherent, while VaR is not convex. Now, for α as above, let

$$Q_\alpha = \left\{ q \in P^{\text{ac}} : \left\| \frac{dq}{dp} \right\|_\infty \leq 1/\alpha \right\}.$$

It is well known that ES^α admits the representation $\text{ES}^\alpha(x) = \sup_{q \in Q_\alpha} \langle q, -x \rangle$. Drawing from this representation, we can define an Expected Shortfall-like SRM as

$$R_{\text{ES}}(X) = \langle Q_\alpha, -X \rangle, \quad X \in \text{CB}(L^\infty).$$

Notice that $R_{\text{ES}}(-X) = \langle Q_\alpha, X \rangle = \sup_{x \in X} \sup_{q \in Q_\alpha} \langle q, x \rangle = \sup_{x \in X} \text{ES}^\alpha(-x)$, so R_{ES} is of the worst-case type.

Example 5.3 (Shortfall risk measures). Let $\ell: \mathbb{R} \rightarrow \mathbb{R}$ be a strictly convex and increasing function. With some abuse of notation, we write $\ell(X) = \{\ell \circ x : x \in X\}$. It is clear that $\ell(X) \in \text{CB}(L^\infty)$ for any $X \in \text{CB}(L^\infty)$. For some fixed $Q \in \text{CB}(P^{\text{ac}})$ and λ_0 in the interior of $\ell(\mathbb{R})$ we define $\mathcal{A}_\ell = \{X \in \text{CB}(L^\infty) : \langle Q, \ell(-X) \rangle \leq \lambda_0\}$. The induced risk measure

$$R_{\mathcal{A}_\ell}(X) = \inf\{\alpha \in \mathbb{R} : \langle Q, \ell(-(X + \alpha)) \rangle \leq \lambda_0\}$$

is called a **shortfall SRM**. Since \mathcal{A}_ℓ is a convex acceptance set, in view of Proposition 2.4, $R_{\mathcal{A}_\ell}$ is a convex SRM. For an exponential loss function $\ell(\alpha) = e^{\lambda\alpha}$, $\lambda > 0$, and $\lambda_0 = 1$, the resulting SRM is linked to the entropic risk measure, which is convex and defined as $S^\gamma(x) = \frac{1}{\gamma} \log(\langle p, e^{-\gamma x} \rangle)$, $\gamma \geq 0$. In this case, we get

$$R_{\text{ent}}(X) = \inf\{\alpha \in \mathbb{R} : \langle Q, e^{\lambda(-X-\alpha)} \rangle \leq 1\} = \frac{1}{\lambda} \log \langle Q, e^{-\lambda X} \rangle.$$

Further, by taking R_{EL} as a basis, we can recover any coherent risk measure ρ simply by a suitable choice of X , that is, for each $x \in L^\infty$, let $X_x = \{x \frac{dq}{dp} : q \in Q_\rho\}$, where Q_ρ is the dual set associated with ρ . Then $\rho(x) = R_{\text{EL}}(X_x)$.

Example 5.4 (New SRMs from old SRMs). Given a collection R_1, \dots, R_n of *base* SRMs,

it is possible to obtain new SRMs by composition with a multivariate transformation $g: \mathbb{R}^n \rightarrow \mathbb{R}$. Below, we provide a few examples. First, consider the **risk averse** R_{\max} defined through

$$R_{\max}(X) = \max\{R_1(X), \dots, R_n(X)\}, \quad X \in \text{CB}(L^\infty).$$

This map inherits the properties from R_1, \dots, R_n , and, in this case, we get $\mathcal{A}_{R_{\max}} = \bigcap_{i=1}^n \mathcal{A}_{R_i}$. In view of Theorem 4.2, under convexity (respectively, coherence) of the base SRMs, we get $\tau_{R_{\max}} = \text{conv}(\min_{i \in \{1, \dots, n\}} \tau_{R_i})$ (respectively, dual set $\mathbf{ba}_{1,+}^{R_{\max}} = \text{conv}(\bigcup_{i=1}^n \mathbf{ba}_{1,+}^{R_i})$). Alternatively, consider the **permissive** SRM, R_{\min} , given by

$$R_{\min}(X) = \min\{R_1(X), \dots, R_n(X)\}, \quad X \in \text{CB}(L^\infty).$$

In this case, we get $\mathcal{A}_{R_{\min}} = \bigcup_{i=1}^n \mathcal{A}_{R_i}$, but notice that R_{\min} does not inherit convexity from R_1, \dots, R_n . In particular, this implies R_{\min} lacks a representation under a penalty function or a dual set. It also fails to inherit WC-boundedness, since we can easily take some $X \in \text{CB}(L^\infty)$ such that

$$\min_{i=1, \dots, n} \sup_{x \in X} R_i(x) > \sup_{x \in X} \min_{i=1, \dots, n} R_i(x) = \sup_{x \in X} R_{\min}(x).$$

Last but not least, one could consider an averaging of the form

$$\bar{R}(X) = \sum_{i=1}^n \lambda_i R_i(X), \quad X \in \text{CB}(L^\infty),$$

where $\lambda_1, \dots, \lambda_n \in [0, 1]$ are convex weights such that $\sum_{i=1}^n \lambda_i = 1$. This map does inherit from the base SRMs all the properties of Definition 2.1, with exception to WC-boundedness, since we can take some $X \in \text{CB}(L^\infty)$ such that

$$\sum_{i=1}^n \lambda_i \sup_{x \in X} R_i(x) > \sup_{x \in X} \sum_{i=1}^n \lambda_i R_i(x) = \sup_{x \in X} \bar{R}(x).$$

The three SRMs above can be considered as a way to tackle uncertainty about the adequacy/quality of the risk assessments offered by each individual base SRM R_1, \dots, R_n .

Example 5.5 (Maximum of finite, convex combinations). Assume ρ is a traditional risk measure. Let $\text{Fin}_n \subseteq \text{CB}(L^\infty)$ be the class of subsets of $\text{CB}(L^\infty)$ having exactly $n < \infty$

elements, and put $\Delta_n = \{\boldsymbol{\lambda} \in [0, 1]^n : \sum_{i=1}^n \lambda_i = 1\}$. Define

$$R(X) = \sup_{n \in \mathbb{N}} \sup_{\boldsymbol{\lambda} \in \Delta_n} \sup_{x \in X \cap \text{Fin}_n} \rho\left(\sum_{i=1}^n \lambda_i x_i\right), \quad X \in \text{CB}(L^\infty).$$

Notice that, since we are taking the suprema above in the above expression, the choice of indexing $X \cap \text{Fin}_n = \{x_1, \dots, x_n\}$ is irrelevant. If the supremum is attained, we can interpret the SRM R as appraising the risk of a set X of financial positions by considering the ρ -riskiest among all finite convex combination of its elements.

Example 5.6 (Knightian uncertainty). This examples offers an alternative approach to tackle uncertainty. Two topics broadly discussed in the literature, *model misspecification* and *Knightian uncertainty in a dynamic decision-making framework* are theoretical frameworks that try to overcome the difficulties arising from an environment of imperfect information, where decision-makers are confronted with the consequences of their risk assessments. To give an example, [Moresco et al. \(2023\)](#) consider a worst-case SRM similar to (4.1). In the context of financial losses represented by x , uncertainty may arise as the agent faces ambiguity about whether x accurately models the true payoff. In such instances, it becomes prudent for the agent to contemplate a set of alternative random variables. These alternatives are payoffs that are considered “close” to x or share common attributes, such as similar distributional features. The uncertainty set of x is then written as $u(x)$, where $u: L^\infty \rightarrow \text{CB}(L^\infty)$ is translation invariant, positive homogeneous, convex, and monotone, i.e. if $x \leq y$, then $u(x) =: X \leq Y := u(y)$. There is a plethora of uncertainty sets considered in the literature, including uncertainty sets defined via, e.g., mixtures of distributions ([Zhu and Fukushima, 2009](#)), moment constraints ([Natarajan et al., 2009](#)), divergence constraints ([Wang et al., 2016](#)), and combinations of moment and divergence constraints ([Bernard et al., 2023](#)). Those are prime examples of uncertainty sets that are constructed as neighborhoods around a reference distribution or random variable, with radius given by a tolerance distance, that is

$$u_D(x) := \{y \in L^\infty : D(x, y) \leq \varepsilon_X\},$$

where D can be, for example, a semi-norm ([Gotoh et al., 2013](#)), the Wasserstein distance ([Pflug and Wozabal, 2007](#)), Kullback–Leibler (KL) divergence ([Calafiore, 2007](#)), general f -divergences, and expected scores ([Moresco et al., 2023](#)).

Given a traditional coherent risk measure $\rho: L^\infty \rightarrow \mathbb{R}$ and an uncertainty set u , the

corresponding **worst-case robust** risk measure ρ_u is given by

$$\rho_u(x) = \sup_{y \in u(x)} \rho(y) = R_\rho(u(x)) = \langle Q^\rho, -u(x) \rangle, \quad x \in L^\infty.$$

where R_ρ is the worst-case SRM associated with ρ , and the set $Q^\rho = \{Q \in \mathbf{ba}_{1,+} : \sup_{x \in A_\rho} \langle q, -x \rangle < \infty\}$.

Example 5.7 (Systemic risk). A SRM can be understood as the risk of a system of financial institutions, stocks in a financial market, or even a finite-time discrete stochastic process under a proper filtration on (Ω, \mathcal{F}, p) . In the literature, a **systemic risk measure** is a functional $\rho_S: (L^\infty)^n \rightarrow \mathbb{R}$ which satisfies monotonicity, convexity, positive homogeneity and, in addition,

Risk Convexity: if $R(z(\omega)) = \lambda R(x(\omega)) + (1 - \lambda)R(y(\omega))$ for a given scalar $\lambda \in [0, 1]$ and almost all $\omega \in \Omega$, then $R(z) \leq \lambda R(x) + (1 - \lambda)R(y)$.

Normalization: $\rho(1) = n$.

An aggregation function is surjective map $\Lambda: \mathbb{R}^n \rightarrow \mathbb{R}$ that is monotone, concave, and satisfies $\Lambda((L^\infty)^n) = L^\infty$. We have that R is a systemic risk measure if and only if there is a coherent risk measure ρ on L^∞ and an aggregation function Λ such that $R = \rho \circ \Lambda$, see [Chen et al. \(2013\)](#) and [Kromer et al. \(2016b\)](#) for details. In this setup we get that $R(x) = \inf\{\alpha \in \mathbb{R} : \Lambda(x) + \alpha \in A_\rho\}$. Thus, systemic risk is the minimal capital added to secure the system after aggregating individual risks. Nonetheless, it might be more relevant to measure systemic risk as the minimal cash amount that secures the aggregated system by adding the capital into the single institutions before aggregating their individual risks. This way of measuring systemic risk can be expressed by

$$R(x) = \inf \left\{ \sum_{i=1}^n \alpha_i : \alpha = (\alpha_1, \dots, \alpha_n) \in \mathbb{R}^n, \Lambda(x + \alpha) \in A \right\},$$

where A is the acceptance set of a coherent risk measure. Then, restricted to such space, a systemic risk measure is similar to a coherent SRM, with an adapted translation invariance, $R(x + \lambda) = R(x) - n\lambda$, and some technical properties to avoid pathological patterns.

We now show how the two approaches above can be adapted to construct SRMs. Let an aggregation function be a surjective map $\Lambda: \mathbf{CB}(L^\infty) \rightarrow L^\infty$ that is monotone, translation invariant, normalized, concave and continuous. Prominent examples are $\Lambda(X) = \text{ess inf } X$ or $\Lambda(X) = \text{ess sup } X$. For some convex risk measure ρ we have a

SRM defined as

$$R(X) := R_{\rho, \Lambda}(X) = \inf \{ \alpha \in \mathbb{R} : \Lambda(X) + \alpha \in A_\rho \}, \quad X \in \text{CB}(L^\infty).$$

It is straightforward to verify that such map is a convex SRM. It can be monetary or coherent, by dropping or demanding the necessary properties for both Λ and ρ . Now, by translation invariance,

$$R(X) := R_{A, \Lambda}(X) = \inf \{ \alpha \in \mathbb{R} : \Lambda(X + \dot{\alpha}) \in A \}, \quad X \in \text{CB}(L^\infty),$$

where A is a convex acceptance set. Both maps coincide over $A = A_\rho$ or $\rho = \rho_A$. By demanding Λ to be set-monotone, union-bounded, or WC-bounded, it is a direct check that R also satisfies such properties. This is the case for $\Lambda(X) = \text{ess inf } X$. In this case we get that $R(X) = -\inf_{x \in X} \text{ess inf } x = \sup_{x \in X} \text{ML}(x)$, where ML stands for the **maximum loss** risk measure.

A particularly interesting case for practical applications is when the SRM is defined on the subspace $\text{Fin}(L^\infty)$ of $\text{CB}(L^\infty)$ comprised of the non-empty, finite subsets of L^∞ . Under continuity, a risk measure $R: \text{Fin}(L^\infty) \rightarrow \mathbb{R}$ can be extended to $\text{K}(L^\infty)$, since $\text{Fin}(L^\infty)$ is dense in $\text{K}(L^\infty)$: for any $X \in \text{K}(L^\infty)$, we can find a bounded sequence $X_n \in \text{Fin}(L^\infty)$ such that $X_n \rightarrow X$. Hence, R can be extended through

$$R(X) = \lim_{X_n \rightarrow X} R(X_n),$$

which is well-defined for any $X \in \text{K}(L^\infty)$. This SRM does not satisfy our definition of translation invariance in general, so it is not a monetary SRM. We leave the discussion of how to properly adapt this concept to non-finite sets to future work.

Example 5.8. A usual risk-neutral stochastic problem has the form

$$\min_{a \in A} \langle p, \ell(a, x) \rangle,$$

where A is a set of possible/allowable *actions*, $x \in (L^\infty)^n$, and $\ell: A \times \mathbb{R}^n \rightarrow \mathbb{R}$ is a suitable loss function. A possible risk-averse counterpart to the above is

$$\min_{a \in A} \rho(\ell(a, x)).$$

which, in the conditions of Theorem 4.2, can be extended to

$$\min_{a \in A} R(\ell(a, X)) = \min_{a \in A} \sup_{x \in X} \rho(\ell(a, x)), \quad X \in \text{CB}(L^\infty)^n.$$

A natural question that arises in this venue is whether this problem is *robust*, in the sense of how it responds to small perturbations in X . We have the following result.

Proposition 5.9. Assume R is a set-monotone and WC-bounded monetary SRM. If $X_n \xrightarrow{s} X$, and $\bigcup_{n \in \mathbb{N}} X_n$ is precompact, then

$$\min_{a \in A} R(\ell(a, X_n)) \rightarrow \min_{a \in A} R(\ell(a, X)).$$

Proof. In this case we have that the sequence $\{f_n, n \geq 1\}$ as defined in the proof of Theorem 4.2 is tight. By item 3 in Lemma 4.1, hypo-convergence then implies convergence and finiteness of the supremum. Hence, we have that

$$\sup_{x \in X_n} R(\dot{x}) = \sup_{x \in X} f_n(x) \rightarrow \sup_{x \in X} f(x) = \sup_{x \in X} R(\dot{x}).$$

Thus, we have $R(X_n) \rightarrow R(X)$. For ease of notation, let $g_n, g: \mathbb{R} \rightarrow \mathbb{R}$ be defined as $g_n(a) = R(\ell(a, X_n))$ for each $n \in \mathbb{N}$, and $g(a) = R(\ell(a, X))$. Then, we have that $g_n \rightarrow g$ pointwise. That is, if $a_n \rightarrow a$, then $g_n(a_n) = R(\ell(a_n, X_n)) \rightarrow R(\ell(a, X)) = g(a)$, which implies $\text{epi } g_n \xrightarrow{s} \text{epi } g$. Now, take

$$a_n^* \in \arg \min_{a \in A} R(\ell(a, X_n))$$

for each $n \in \mathbb{N}$. Since, translation invariance and monotonicity imply $|R(X)| \leq \|X\|$ for any $X \in \text{CB}(L^\infty)$, we get that $\{a_n^*, n \geq 1\}$ is bounded since $\bigcup_{n \in \mathbb{N}} X_n$ is precompact, and hence bounded. Then, by the Bolzano-Weierstrass Theorem, a_n^* converges (by taking a sub-sequence if needed). Thus, from item 2 in Lemma 4.1 we get that $\inf_{a \in A} R(\ell(a, X_n)) \rightarrow \inf_{a \in A} R(\ell(a, X))$. The claim follows, since the infimum is attained. ■

Example 5.10 (Financial problems). SRMs appear naturally in the context of portfolio optimization. Let X be a set of n available assets; then, the risk of the optimum portfolio,

given a return μ and a traditional risk measure ρ , is given by

$$R^\mu(X) = \inf \left\{ \rho \left(\sum_{i=1}^n w_i x_i \right) : \sum_{i=1}^n w_i = 1, \sum_{i=1}^n \langle p, w_i x_i \rangle \geq \mu, x_i \in X \forall i \leq n \right\}.$$

In this case, $R^\mu(X)$ is the risk of the optimum portfolio given a required return μ . With greater generality, given a set of decisions D , where each $d \in D$ leads to an outcome $x_d \in L^\infty$, and $X = \{x_d \in L^\infty : d \in D\}$ the risk of the optimal decision is then $R(X) = \inf_{x \in X} \rho(x)$.

Risk contribution is another field where SRMs can be employed. The risk of a position x changes when it is merged with another position y . For instance, if a monetary risk measure ρ deems a position x as unacceptable, but it becomes a sub-portfolio of y , then the marginal contribution $\rho_y(x) := \rho(y+x) - \rho(x)$ may alter the risk profile of x , making it acceptable according to the measure $\rho_y(\cdot)$ without contributing to the risk of y , and perhaps even reducing, the risk of y (Canna et al., 2020). This can be written as a SRM when $X = \{x, y\}$, as

$$R(\{x, y\}) = \rho(y+x) - \min\{\rho(x), \rho(y)\}.$$

If X is given by a set of finitely many positions, the SRM is given by

$$R(X) = \rho \left(\sum_{x \in X} x \right) - \min_{x \in X} \{\rho(x)\}.$$

The domain of this risk measure can be expanded to $K(L^\infty)$ by the same argument as in Example 5.7. Similar in nature is the *diversification benefit* (Embrechts et al., 2009; Koumou and Dionne, 2022; Han et al., 2022), for a finite X , the diversification benefit within X can be cast as a SRM through

$$R(X) = \sum_{x \in X} \rho(x) - \rho \left(\sum_{x \in X} x \right)$$

Similarly, considering now the problem of risk sharing, assume X to be finite, and let Λ be a set of coherent risk measures. Then the risk of each position in X may be shared among the risk measures in Λ . The *optimal sharing* is then given by the SRM

$$R(X) = \inf_{\rho \in \Lambda} \left\{ \sum_{x \in X} \rho(x) \right\}.$$

If this is taken over all X having the same aggregated risk, we arrive at the formulation of the **inf-convolution**,

$$\inf \left\{ R(X) : \sum_{x \in X} = x \right\} = \square_{\rho \in \Lambda} \rho(x), \quad x \in L^\infty.$$

For a generalization of inf-convolution to countable sets, see [Righi and Moresco \(2022\)](#).

Example 5.11. A prominent example is based on the concept of Bochner integral. See the chapter 11.8 in [Aliprantis and Border \(2006\)](#) for specific details and results. For any $X \in \text{CB}(L^\infty)$, we have that the set $X(\omega) = \{x(\omega) : x \in X\}$ is both closed and bounded in \mathbb{R} (this is true p -a.s.). Boundedness is directly inherited from that of X , while closedness follows since $|x_n(\omega) - x(\omega)| \leq \|x_n - x\|_{L^\infty}$. Thus, $X(\omega) \in K(\mathbb{R})$. Notice that $K(\mathbb{R})$ is in fact a Banach space, since sum and multiplication by scalar for compact spaces are closed operations. A function $\varphi : \Omega \rightarrow K(\mathbb{R})$ is **simple** if $\varphi(\omega) = \sum_{i=1}^n K_i 1_{A_i}(\omega)$, for some collection K_1, \dots, K_n of compact sets in \mathbb{R} and some events $\{A_i\}_{i=1}^n \subseteq \mathcal{F}$. The **Bochner integral** (w.r.t p) of a simple function is the monotone, linear operator from the linear space of step functions to $K(\mathbb{R})$ defined as $\int_\Omega \varphi \, dp = \sum_{i=1}^n K_i p(A_i)$. We say that $\omega \mapsto X(\omega)$ is **strongly measurable** if there is a sequence of simple functions $\{\varphi_n : \Omega \rightarrow K(\mathbb{R}), n \geq 1\}$, such that

$$\|X(\omega) - \varphi_n(\omega)\|_{K(\mathbb{R})} \rightarrow 0, \quad p\text{-a.s.},$$

where $\|\cdot\|_{K(\mathbb{R})}$ is the norm in $K(\mathbb{R})$. In addition, we say that X is **Bochner integrable** if there is a sequence of simple functions (which can be taken to be the same one as above) such that $\omega \mapsto \|X(\omega) - \varphi_n(\omega)\|_{K(\mathbb{R})}$ is Lebesgue integrable with respect to p for each $n \in \mathbb{N}$ and

$$\lim_{n \rightarrow \infty} \int_\Omega \|X(\omega) - \varphi_n(\omega)\|_{K(\mathbb{R})} \, dp(\omega) = 0.$$

In this case, we define the **Bochner integral** of X with respect to p as

$$\int_\Omega X \, dp = \lim_{n \rightarrow \infty} \int_\Omega \varphi_n \, dp,$$

where the limit is taken on the norm of $K(\mathbb{R})$. Thus, examples of SRMs over the subspace

of Bochner integrable sets in $\text{CB}(L^\infty)$ are, but of course not limited to:

$$R_1(X) = -\min \int_{\Omega} X \, dp,$$

$$R_2(X) = -\min \bigcup_{q \subseteq Q_{R_2}} \int_{\Omega} X \, dq,$$

where $Q_{R_2} \subseteq Q$ is non-empty, closed and convex. R_2 is well-defined since $L^\infty(p) \subseteq L^\infty(q)$ for any $q \in Q$. Both R_1 and R_2 are coherent SRMs. Of course, many other real valued operators over $K(\mathbb{R})$ can be considered beyond the minimum.

6 Concluding remarks

In this paper, we have introduced the concept of set risk measures (SRMs), broadening the traditional framework of risk measures to encompass sets of random variables. This innovative approach addresses the limitations of classical risk measures by offering a more general and flexible setup, suitable for a wide range of financial and mathematical applications. Our main contributions include the establishment of an axiomatic scheme for SRMs, the derivation of a dual representation for convex SRMs, and the exploration of worst-case SRMs.

The axiomatic framework presented in Section 2 provides a solid foundation for understanding and working with SRMs, highlighting essential properties such as monotonicity, translation invariance, and convexity. In Section 3, we developed a dual representation for convex SRMs using regular, finitely additive measures, offering new insights into the structure and behavior of these measures. This dual representation is a significant theoretical advancement, extending the classical duality theory to the realm of set risk measures.

Section 4 focused on worst-case SRMs, a critical subclass of SRMs that appraise risk based on the supremum of individual risks within a set. We provided necessary and sufficient conditions for SRMs to be considered worst-case measures and refined some results regarding their continuity and dual representation. These findings underscore the practical importance of worst-case SRMs in financial risk management, where they offer robust tools for evaluating and mitigating risk.

Our work extends the theory of risk measures to a more inclusive and adaptable framework, accommodating various scenarios in systemic risk, portfolio optimization, and decision-making under uncertainty. The examples presented in Section 5 illustrate the practical applicability of our framework, demonstrating its potential to address com-

plex challenges in these areas.

Looking ahead, there are several promising avenues for future research. One potential direction is the exploration of SRMs in more specific financial contexts, such as credit risk, market risk, and operational risk. Additionally, further investigation into the connections between SRMs and other advanced risk measures, such as coherent and spectral risk measures, could yield valuable insights. Another interesting area for future research is the application of SRMs to machine learning and artificial intelligence, where they could play a crucial role in developing robust models for risk assessment and decision-making under uncertainty.

In conclusion, the introduction of SRMs represents a significant advancement in the field of risk measurement, offering a versatile and powerful tool for analyzing and managing risk in diverse settings. We hope that our work will inspire further research and development in this area, leading to more effective and comprehensive approaches to risk management.

References

- Aliprantis, C. D. and Border, K. C. (2006). *Infinite Dimensional Analysis*. Springer, Berlin. [13](#), [14](#), [16](#), [33](#)
- Ang, M., Sun, J., and Yao, Q. (2018). On the dual representation of coherent risk measures. *Annals of Operations Research*, 262(1):29–46. [4](#)
- Arai, T. (2014). Convex risk measures for cadlag processes on orlicz hearts. *SIAM Journal on Financial Mathematics*, 5(1):609–625. [3](#)
- Ararat, Ç., Hamel, A. H., and Rudloff, B. (2017). Set-valued shortfall and divergence risk measures. *International Journal of Theoretical and Applied Finance*, 20(05):1750026. [3](#)
- Artzner, P., Delbaen, F., Eber, J.-M., and Heath, D. (1999). Coherent measures of risk. *Mathematical Finance*, 9(3):203–228. [3](#)
- Assa, H. and Morales, M. (2010). Risk measures on the space of infinite sequences. *Mathematics and Financial Economics*, 2:253–275. [3](#)
- Aubin, J.-P. and Frankowska, H. (2009). *Set-valued analysis*. Springer Science & Business Media. [19](#), [20](#)

- Bachman, G. and Sultan, A. (1980). On regular extensions of measures. *Pacific Journal of Mathematics*, 86(2):389–395. [16](#)
- Bartl, D., Drapeau, S., and Tangpi, L. (2019). Computational aspects of robust optimized certainty equivalents and option pricing. *Mathematical Finance*, in press. [4](#)
- Beer, G. (1993). *Topologies on closed and closed convex sets*, volume 268. Springer Science & Business Media. [3](#), [5](#), [7](#), [14](#), [19](#)
- Bellini, F., Laeven, R. J. A., and Rosazza Gianin, E. (2018). Robust return risk measures. *Mathematics and Financial Economics*, 12(1):5–32. [4](#)
- Bernard, C., Pesenti, S. M., and Vanduffel, S. (2023). Robust distortion risk measures. *Mathematical Finance*. (forthcoming). [4](#), [28](#)
- Biagini, S. and Frittelli, M. (2010). *On the Extension of the Namioka-Klee Theorem and on the Fatou Property for Risk Measures*, pages 1–28. Springer Berlin Heidelberg, Berlin, Heidelberg. [3](#), [15](#)
- Bion-Nadal, J. and Kervarec, M. (2012). Risk measuring under model uncertainty. *Ann. Appl. Probab.*, 22(1):213–238. [3](#)
- Bogachev, V. I. (2006). *Measure Theory*, volume 1. Springer Berlin, Heidelberg, 1st edition. [16](#)
- Burgert, C. and Rüschendorf, L. (2006). On the optimal risk allocation problem. *Statistics & Decisions*, 24:153 – 171. [3](#)
- Burgert, C. and Rüschendorf, L. (2008). Allocation of risks and equilibrium in markets with finitely many traders. *Insurance: Mathematics and Economics*, 42(1):177 – 188. [3](#)
- Calafiore, G. C. (2007). Ambiguous risk measures and optimal robust portfolios. *SIAM Journal on Optimization*, 18(3):853–877. [28](#)
- Canna, G., Centrone, F., and Rosazza Gianin, E. (2020). Capital allocation rules and acceptance sets. *Mathematics and Financial Economics*, 14:759–781. [32](#)
- Chen, C., Iyengar, G., and Moallemi, C. C. (2013). An axiomatic approach to systemic risk. *Management Science*, 59(6):1373–1388. [3](#), [29](#)

- Cheridito, P., Delbaen, F., and Kupper, M. (2004). Coherent and convex monetary risk measures for bounded cadlag processes. *Stochastic Processes and their Applications*, 112(1):1–22. [3](#)
- Cheridito, P., Delbaen, F., and Kupper, M. (2005). Coherent and convex monetary risk measures for unbounded cadlag processes. *Finance and Stochastics*, 9(3):369–387. [3](#)
- Cheridito, P. and Li, T. (2009). Risk measures on Orlicz Hearts. *Mathematical Finance*, 19(2):189–214. [3](#)
- Delbaen, F. (2002). Coherent risk measures on general probability spaces. *Advances in Finance and Stochastics*, pages 1–37. [3](#)
- Delbaen, F. (2009). Risk measures for non-integrable random variables. *Mathematical Finance: An International Journal of Mathematics, Statistics and Financial Economics*, 19(2):329–333. [3](#)
- Delbaen, F. (2012). *Monetary utility functions*. Osaka University Press. [3](#)
- Delbaen, F. (2022). Monetary utility functions on $c_b(x)$ spaces. *arXiv preprint arXiv:2209.08137*. [3](#), [18](#)
- Ekeland, I., Galichon, A., and Henry, M. (2012). Comonotonic measures of multivariate risks. *Mathematical Finance: An International Journal of Mathematics, Statistics and Financial Economics*, 22(1):109–132. [3](#)
- Embrechts, P., Furrer, H., and Kaufmann, R. (2009). Different kinds of risk. In *Handbook of financial time series*, pages 729–751. Springer. [32](#)
- Fadina, T., Liu, Y., and Wang, R. (2021). A framework for measures of risk under uncertainty. *arXiv preprint arXiv:2110.10792*. [4](#)
- Feinstein, Z. and Rudloff, B. (2022). Scalar multivariate risk measures with a single eligible asset. *Mathematics of Operations Research*, 47(2):899–922. [3](#)
- Föllmer, H. and Schied, A. (2002). Convex measures of risk and trading constraints. *Finance and stochastics*, 6(4):429–447. [4](#)
- Follmer, H. and Schied, A. (2016). *Stochastic finance: an introduction in discrete time*. Walter de Gruyter GmbH. [3](#), [7](#), [21](#), [23](#)
- Frittelli, M., Maggis, M., and Peri, I. (2014). Risk measures on $p(r)$ and value at risk with probability/loss function. *Mathematical Finance*, 24(3):442–463. [3](#)

- Frittelli, M. and Scandolo, G. (2006). Risk measures and capital requirements for processes. *Mathematical finance*, 16(4):589–612. [3](#)
- Gao, N. and Xanthos, F. (2018). On the c-property and-representations of risk measures. *Mathematical Finance*, 28(2):748–754. [3](#)
- Gotoh, J.-Y., Shinozaki, K., and Takeda, A. (2013). Robust portfolio techniques for mitigating the fragility of CVaR minimization and generalization to coherent risk measures. *Quantitative Finance*, 13(10):1621–1635. [28](#)
- Guo, S. and Xu, H. (2019). Distributionally robust shortfall risk optimization model and its approximation. *Mathematical Programming*, 174:473–498. [4](#)
- Hamel, A. (2005). *Variational principles on metric and uniform spaces*. PhD thesis, Halle (Saale), Univ., Habil.-Schr., 2005. [6](#)
- Hamel, A. H. and Heyde, F. (2010). Duality for set-valued measures of risk. *SIAM Journal on Financial Mathematics*, 1(1):66–95. [3](#)
- Hamel, A. H., Heyde, F., and Rudloff, B. (2011). Set-valued risk measures for conical market models. *Mathematics and financial economics*, 5(1):1–28. [3](#)
- Han, X., Lin, L., and Wang, R. (2022). Diversification quotients: Quantifying diversification via risk measures. *arXiv preprint arXiv:2206.13679*. [32](#)
- Ivar, E. and Walter, S. (2011). Law invariant risk measures on $l^\infty(\mathbb{R}^d)$. *Statistics & Risk Modeling*, 28(3):195–225. [3](#)
- Jouini, E., Meddeb, M., and Touzi, N. (2004). Vector-valued coherent risk measures. *Finance and stochastics*, 8(4):531–552. [3](#)
- Kaina, M. and Rüschendorf, L. (2009). On convex risk measures on lp-spaces. *Mathematical Methods of Operations Research*, 69(3):475–495. [3](#), [15](#)
- Kou, S., Peng, X., and Heyde, C. C. (2013). External risk measures and basel accords. *Mathematics of Operations Research*, 38(3):393–417. [3](#)
- Koumou, G. B. and Dionne, G. (2022). Coherent diversification measures in portfolio theory: An axiomatic foundation. *Risks*, 10(11):205. [32](#)
- Kromer, E., Overbeck, L., and Zilch, K. (2016a). Systemic risk measures on general measurable spaces. *Mathematical Methods of Operations Research*, 84:323–357. [3](#)

- Kromer, E., Overbeck, L., and Zilch, K. (2016b). Systemic risk measures on general measurable spaces. *Mathematical Methods of Operations Research*, 84:323–357. 29
- Laeven, R. J. A. and Stajje, M. (2013). Entropy coherent and entropy convex measures of risk. *Mathematics of Operations Research*, 38(2):265–293. 4
- Lucchetti, R. and Pasquale, A. (1994). A new approach to a hyperspace theory. *Journal of Convex Analysis*, 1(2). 8
- Luxemburg, W. A. J. (1991). Integration with respect to finitely additive measures. In *Positive Operators, Riesz Spaces, and Economics*, pages 109–150, Berlin, Heidelberg. Springer Berlin Heidelberg. 16
- Moresco, M., Mailhot, M., and Pesenti, S. (2023). Uncertainty propagation and dynamic robust risk measures. *arXiv preprint arXiv:2308.12856*. 28
- Natarajan, K., Pachamanova, D., and Sim, M. (2009). Constructing risk measures from uncertainty sets. *Operations Research*, 57(5):1129–1141. 28
- Nendel, M. (2024). Lower semicontinuity of monotone functionals in the mixed topology on c_b . *arXiv preprint arXiv:2210.09133*. 3, 18
- Pflug, G. and Römisch, W. (2007). *Modeling, Measuring and Managing Risk*. World Scientific, 1 edition. 3
- Pflug, G. and Wozabal, D. (2007). Ambiguity in portfolio selection. *Quantitative Finance*, 7(4):435–442. 28
- Pichler, A. (2013). The natural banach space for version independent risk measures. *Insurance: Mathematics and Economics*, 53(2):405 – 415. 3
- Qian, P.-Y., Wang, Z.-Z., and Wen, Z.-W. (2019). A composite risk measure framework for decision making under uncertainty. *Journal of the Operations Research Society of China*, 7(1):43–68. 4
- Righi, M. (2019). A theory for combinations of risk measures. *Working Paper*. 4
- Righi, M. (2024). Robust convex risk measures. *arXiv preprint arXiv:2406.12999*. 4
- Righi, M. B. and Moresco, M. R. (2022). Inf-convolution and optimal risk sharing with countable sets of risk measures. *Annals of Operations Research*, pages 1–32. 33

- Rüschendorf, L. (2006). Law invariant convex risk measures for portfolio vectors. *Statistics & Decisions*, 24(1):97–108. [3](#)
- Rüschendorf, L. (2013). *Mathematical Risk Analysis*. Springer. [3](#)
- Schrage, C. (2009). *Set-valued convex analysis*. PhD thesis, Martin-Luther-Universität Halle-Wittenberg. [6](#)
- Shapiro, A., Dentcheva, D., and Ruszczyński, A. (2021). *Lectures on stochastic programming: modeling and theory*. SIAM. [3](#)
- Wang, R. and Ziegel, J. (2018). Scenario-based risk evaluation. *Working Paper*. [4](#)
- Wang, Z., Glynn, P. W., and Ye, Y. (2016). Likelihood robust optimization for data-driven problems. *Computational Management Science*, 13:241–261. [28](#)
- Wei, L. and Hu, Y. (2014). Coherent and convex risk measures for portfolios with applications. *Statistics & Probability Letters*, 90:114–120. [3](#)
- Zhu, S. and Fukushima, M. (2009). Worst-case conditional Value-at-Risk with application to robust portfolio management. *Operations Research*, 57(5):1155–1168. [28](#)