

POSITIVE SCALAR CURVATURE WITH POINT SINGULARITIES

SIMONE CECCHINI, GEORG FRENCK, AND RUDOLF ZEIDLER

ABSTRACT. We show that in every dimension $n \geq 8$, there exists a smooth closed manifold M^n which does not admit a smooth positive scalar curvature (“psc”) metric, but M admits an L^∞ -metric which is smooth and has psc outside a singular set of codimension ≥ 8 . This provides counterexamples to a conjecture of Schoen. In fact, there are such examples of arbitrarily high dimension with only single point singularities. In addition, we provide examples of L^∞ -metrics on \mathbb{R}^n for certain $n \geq 8$ which are smooth and have psc outside the origin, but cannot be smoothly approximated away from the origin by everywhere smooth Riemannian metrics of non-negative scalar curvature. This stands in precise contrast to established smoothing results via Ricci–DeTurck flow for singular metrics with stronger regularity assumptions. Finally, as a positive result, we describe a KO-theoretic condition which obstructs the existence of L^∞ -metrics that are smooth and of psc outside a finite subset. This shows that closed enlargeable spin manifolds do not carry such metrics.

1. INTRODUCTION

Studying weak lower curvature bounds in various singular settings has been a core aspect of Riemannian geometry since a long time. While for spaces with *scalar curvature* bounded below such a theory is not yet fully developed, various approaches with results in different directions have been proposed recently (for instance [2; 4; 7; 15; 18; 25; 26; 28; 32; 39; 41]). In this paper, we investigate scalar curvature lower bounds for Riemannian L^∞ -metrics in the sense of the following definition.

Definition 1.1 (cf. [28, Definition 1.3]). Let M be a smooth manifold. A *uniformly Euclidean metric* (or for short “ L^∞ -metric”) is a measurable symmetric $(0, 2)$ -tensor g on M such that there exists a smooth Riemannian metric g_0 on M and $\Lambda > 0$ with

$$\Lambda^{-1}g_0(\xi, \xi) \leq g(\xi, \xi) \leq \Lambda g_0(\xi, \xi)$$

for almost all $x \in M$ and all tangent vectors $\xi \in T_x M$.

For uniformly Euclidean metrics which are smooth outside a singular subset of sufficiently high codimension, it makes sense to study the behavior of lower scalar curvature bounds on the regular subset of the metric. The first general results in this direction have been obtained by Li–Mantoulidis [28], where the following conjecture of Schoen was popularized.

Conjecture 1.2 (Schoen, cf. [28, Conjecture 1.5]). Let M be a closed manifold that does not support any smooth Riemannian metric of positive scalar curvature (i.e. M is Yamabe non-positive). Let $S \subset M$ be a closed submanifold of codimension ≥ 3

2020 *Mathematics Subject Classification.* 53C21 (Primary) 57R65; 58J22 (Secondary).

Co-funded by the Deutsche Forschungsgemeinschaft (SPP2026 “Geometry at Infinity”).

Co-funded by the European Union (ERC Starting Grant 101116001 – COMSCAL). Views and opinions expressed are however those of the author(s) only and do not necessarily reflect those of the European Union or the European Research Council. Neither the European Union nor the granting authority can be held responsible for them.

Co-funded by the Deutsche Forschungsgemeinschaft (DFG, German Research Foundation) – Project numbers 523079177; 427320536; 390685587.

and let g be an L^∞ -metric on M which is smooth on $M \setminus S$ and satisfies $\text{scal}_g \geq 0$ on $M \setminus S$. Then g is smooth on all of M and $\text{Ric}_g = 0$.

In other words, this conjecture predicts that subsets of codimension at least three are invisible for the purposes of scalar curvature in a quite strong sense. This differs from the setting of codimension one and two, where it is well known that such statements do not hold without additional regularity assumptions or boundary conditions, compare [28, §8.1–8.2]. Likewise, the condition of being uniformly Euclidean cannot be dropped from Conjecture 1.2, otherwise suitable Schwarzschild metrics would yield counterexamples with point singularities, compare [28, §8.3].

Conjecture 1.2 has attracted considerable interest and it has been confirmed in many specific cases. For instance, if $\dim(M) = 3$, it holds in general by Li–Mantoulidis [28, Corollary 1.6]. In the 4-dimensional case, far-reaching results have been established by Kazaras [23]. In a closely related direction, a version of Llarull’s theorem for metrics with L^∞ point singularities was recently proved by Chu–Lee–Zhu [12]. Fill-in problems with L^∞ point singularities in dimension three were studied by Mantoulidis–Miao–Tam [31, §4]. We also note that related results for conical singularities and the positive mass theorem have been proved by Dai–Sun–Wang [13; 14]. In sharp contrast to all of this, our main result shows that L^∞ -regularity is in general not enough to retain positive scalar curvature (“psc”) obstructions on closed manifolds, even assuming smoothness outside a single point.

Theorem A. *For every $n \geq 8$, there exists a closed smooth manifold M of dimension n which does not admit a smooth psc metric, but which admits an L^∞ -metric which is smooth and of uniformly positive scalar curvature on the complement of a connected submanifold $S \subseteq M$ of codimension ≥ 8 . Moreover, the metric we obtain is of regularity $W^{2,p/2}$, for every $p < \text{codim}(S \subset M)$. If $8 \leq n \equiv 0, 1, 2$, or $4 \pmod{8}$, then there are examples where $S = \{*\}$ is a single point.*

In particular, Conjecture 1.2 is false in every dimension $n \geq 8$ and there are counterexamples where the singular set has arbitrarily high codimension.

Remark 1.3. Due to the Sobolev $W^{2,p/2}$ -regularity, in our examples the scalar curvature even exists in the weak sense on all of M as a uniformly positive $L^{p/2}$ -function for all $p < \text{codim}(S \subset M)$. In particular, our metrics have positive distributional scalar curvature in the sense of Lee–LeFloch [25, Definition 2.1].

The counterexamples in Theorem A are obtained via classification results based on surgery theory which allow to construct psc metrics with conical singularities on all high-dimensional simply connected manifolds (see Section 2 for details). To contrast this with the existing positive results concerning Conjecture 1.2, note that the proofs usually proceed by first blowing up the singularity to a complete psc metric on the complement of the singular set. Next, appropriate topological obstructions are used to exclude the existence of such complete psc metrics, leading to a contradiction. The last step fails precisely because this is unobstructed on punctured simply connected manifolds in high dimensions.

A more analytic approach to study Conjecture 1.2, and lower (scalar) curvature bounds on metrics with low regularity in general, is to regularize them using geometric flows, for a selection of works in this direction see [2; 4; 5; 9; 11; 24; 27; 40]. However, all existing results following this line need additional assumptions on the metric beyond being uniformly Euclidean—for instance, additional Sobolev or C^0 regularity near the singular set or being sufficiently L^∞ -close to a smooth background metric. It stands to reason that Theorem A must impose definite restrictions for such a procedure to work for the scalar curvature of L^∞ -metrics.

Remark 1.4. Indeed, Chu–Lee [11, Theorem 6.2] showed that uniformly Euclidean metrics of $W^{1,n}$ -regularity can be smoothed via Ricci–DeTurck flow (see also Lamm–Simon [24] who treated the $W^{2,2}$ case in dimension four). Since our counterexamples from Theorem A in the point singularity case are in $W^{2,p/2} \subset W^{1,p}$ for all $p < n$, this means that $W^{1,n}$ is the sharp threshold on the Sobolev scale for such a smoothing result in general.

In another direction, Burkhardt–Guim [5] recently showed that one may smooth away certain L^∞ singularities on \mathbb{R}^n using Ricci–DeTurck flow while preserving non-negativity of the scalar curvature, provided that the singular metric is sufficiently L^∞ -close to the Euclidean metric. Since the counterexamples in Theorem A are based on high-dimensional simply connected manifolds which do not admit psc, at first glance this may look like a subtle global phenomenon of the underlying closed manifold. However, as we note in the following theorem, they actually lead to examples of L^∞ -metrics on \mathbb{R}^n with a single point singularity which are non-smoothable. In particular, this answers [5, Question 3].

Theorem B. *For $n \geq 8$, $n \equiv 0, 1, 2$, or $4 \pmod{8}$, there exists a metric g on \mathbb{R}^n which has regularity $W_{\text{loc}}^{2,p/2}$ for each $p < n$ such that*

- $\Lambda^{-1}g_0 \leq g \leq \Lambda g_0$ a.e. for some fixed $\Lambda > 0$ and g_0 the Euclidean metric,
- g is smooth on $\mathbb{R}^n \setminus \{0\}$,
- $\text{scal}_g \geq \lambda|x|^{-2} > 0$ on $\mathbb{R}^n \setminus \{0\}$ for some $\lambda > 0$,

but there exists no family $(g_t)_{t \in (0, \varepsilon)}$ of smooth metrics on \mathbb{R}^n satisfying both

- (i) $\text{scal}_{g_t} \geq 0$ for all $t \in (0, \varepsilon)$,
- (ii) $g_t \xrightarrow{C_{\text{loc}}^\infty(\mathbb{R}^n \setminus \{0\})} g$ for $t \searrow 0$.

Theorems A and B suggest to us that the setting of L^∞ -metrics with small singular sets is in general insufficient for studying lower scalar curvature bounds in a local sense. Instead, the fact that Conjecture 1.2 does hold true in certain cases appears to be more of a global topological phenomenon. Thus, as a companion to our counterexamples which are based on simply connected spin obstructions, we note a stronger global spin obstruction to psc which does enable a positive result in the direction of Conjecture 1.2 for point singularities. To formulate it, let M be a closed n -dimensional spin manifold with fundamental group Γ . Denote by $\alpha(M; 1) \in \text{KO}_n(\mathbb{R})$ the Hitchin genus [21] of M , that is, the KO-theoretic index of the Cl_n -linear Dirac operator on M . Let $C^*\Gamma$ be the maximal real group C^* -algebra of Γ , and let $\alpha(M; \Gamma) \in \text{KO}_n(C^*\Gamma)$ be the Rosenberg index of M [34–36], see also [43, §2] for a brief survey. The inclusion $\mathbb{R} \hookrightarrow C^*\Gamma$, sending $\lambda \in \mathbb{R}$ into $\lambda \cdot 1_{C^*\Gamma}$, induces a map $j: \text{KO}_n(\mathbb{R}) \rightarrow \text{KO}_n(C^*\Gamma)$. We will use the condition that $j(\alpha(M; 1)) \neq \alpha(M; \Gamma)$. In other words, this means that the Rosenberg index of M does not come from the trivial group.

Theorem C. *Let M be a closed spin n -dimensional manifold with fundamental group Γ . Suppose that $j(\alpha(M; 1)) \neq \alpha(M; \Gamma)$. Let $S \subset M$ be a finite set, and let g be an L^∞ -metric on M which is smooth outside of S . If $\text{scal}_g \geq 0$ on $M \setminus S$, then $\text{Ric}_g \equiv 0$ on $M \setminus S$.*

Remark 1.5. The condition $j(\alpha(M; 1)) \neq \alpha(M; \Gamma)$ is satisfied for enlargeable spin manifolds [19; 20]. This includes spin manifolds that admit a non-zero degree map to the torus, in particular manifolds of the form $M = T^n \# N$, where N is an arbitrary spin manifold. See [6, Remarks 1.10–1.12] for further discussions of examples. Moreover, all 3-manifolds not carrying psc metrics are enlargeable and thus satisfy this condition, which is the case treated by Li–Mantoulidis [28, Theorem 1.6].

Conversely, by design this is never satisfied for simply connected manifolds and so avoids the counterexamples of Theorem A. It is essentially the most general such condition currently conceivable in the spin setting (compare [37, Conjecture 1.5]).

We note that the proof of Theorem C follows the same lines as in the works of Kazaras [23] and Li–Mantoulidis [28]. Our contribution is identifying the correct KO-theoretic obstruction to complete psc metrics which already appeared in work of Cecchini [6], see also Cecchini–Zeidler [8, §3.2]. Moreover, as in the classical setting, the positive mass theorem with L^∞ point singularities can be reduced to a corresponding psc obstruction statement for closed manifolds of the form $T^n \# N$. Indeed, in [23, Proof of Corollary B], Kazaras extended Lohkamp’s argument [30, Theorem 6.1] to uniformly Euclidean metrics. Therefore, Theorem C directly implies the following.

Corollary D. *Let (M, g) be a complete asymptotically flat n -dimensional spin Riemannian manifold, where g is an L^∞ -metric which is smooth outside a finite set $S \subset M$. If $\text{scal}_g \geq 0$ on $M \setminus S$, then the ADM mass of each asymptotically flat end is non-negative.*

The paper is organized as follows. In Section 2 we present the counterexamples to Schoen’s conjecture and prove Theorem A. Section 3 is devoted to the proof of Theorem B. Finally, in Section 4 we prove Theorem C.

Acknowledgements. We thank Otis Chodosh, Johannes Ebert and Sven Hirsch for insightful conversations. SC gratefully acknowledges the hospitality of Mathematics Münster during his visit as a Young Research Fellow, which significantly contributed to the development of this paper.

2. COUNTEREXAMPLES TO SCHOEN’S CONJECTURE

In this section, we prove Theorem A. The proof is based on the following classification result which stands in contrast to [42, Theorem A] and complements [17, Corollary C].

Proposition 2.1. *Every closed simply connected manifold M of dimension $n \geq 5$ admits an L^∞ -metric g which lies in $W^{2,p/2}$ for every $p < n$, is smooth away from a point $p \in M$, and whose scalar curvature is uniformly positive on $M \setminus \{p\}$.*

The idea of the proof is the following: First, we remove an open disk. On this new manifold with boundary, there is a *collared* psc metric g , that is, $g = h + dt^2$ near the boundary. Then we interpolate this to a conical metric $(t/\lambda)^2 h + dt^2$ which has an L^∞ -singularity at $t = 0$. We start with the following lemma.

Lemma 2.2. *Let M be a closed simply connected spin manifold of dimension $n \geq 5$. Let $D^n \subset M$ be an embedded disk and $M' = M \setminus D^n$. Then there exists a collared psc metric g on M' .*

Proof. First, we perform 2-surgeries on M away from D^n to obtain M_2 which is 2-connected. This is possible as every embedded 2-sphere has trivial normal bundle because of the spin condition.

We write $M'_2 := M_2 \setminus D^n$ and we observe that the inclusion $\partial M'_2 = S^{n-1} \hookrightarrow M'_2$ is 2-connected by construction. If $n \geq 6$, the surgery theorem of Gromov–Lawson [17] and Schoen–Yau [38] in its cobordism formulation [16, Theorem 1.5] implies that there exists a psc metric g_2 on M'_2 which is collared, that is, $g_2 = h + dt^2$ on a collar $\partial M'_2 \times [0, \varepsilon) \subset M'_2$. Since the reverse surgery of a 2-surgery has codimension 3, we can reverse the surgeries from the beginning to obtain a collared psc metric g on $M' = M \setminus D^n$ which agrees with g_2 near $\partial M' = \partial M'_2$.

If on the other hand $n = 5$, M itself already admits a psc metric by [42, Theorem A]¹ which can be deformed to equal a torpedo-metric on an embedded $D^5 \subset M$ ([10, Theorem 1.1], see also [16, Theorem 1.2]). Removing D^5 will yield a psc metric on M' which is collared. \square

Remark 2.3. The statement of Lemma 2.2 is also true if M is simply connected and non-spin: In this case, [17, Corollary C] yields a psc metric on M which—as in the proof above (using [10, Theorem 1.1; 16, Theorem 1.2])—can be deformed to be a torpedo-metric on an embedded disk $D^n \subset M$. Deleting this disk with the torpedo-metric yields the desired collared metric on the complement.

Proof of Proposition 2.1. If M does not admit a spin structure then by [17, Corollary C] there even exists a smooth positive scalar curvature metric on M . Therefore, we may assume that M is spin.

In this case let g be a collared psc metric on $M' = M \setminus D^n$ provided by Lemma 2.2, where D^n is some embedded disk around the point $p \in M$. Denote the restriction of g to the boundary $\partial M'$ by h . Then h has uniformly positive scalar curvature due to compactness. Let $C := \inf_{x \in \partial M'} \text{scal}_h(x) > 0$. Now, let $R > 1$ be such that there exists a smooth, monotonically decreasing function $\alpha: \mathbb{R} \rightarrow [0, 1]$ (see Figure 1) such that $\alpha|_{(-\infty, -R+\varepsilon]} = 1$ and $\alpha|_{[-\varepsilon, \infty)} = 0$ with

$$-\frac{1}{2} \sqrt{\frac{C}{4(n-1)(n-2)}} \leq \alpha'(t) \leq 0 \quad \text{and} \quad |\alpha''(t)| \leq \frac{C}{8(n-1)}.$$

Furthermore, let $\lambda \in \mathbb{R}$ be such that

$$\lambda \geq \max \left\{ R, 2 \sqrt{\frac{4(n-1)(n-2)}{C}} \right\} > 1.$$




FIGURE 1. The auxiliary function $\alpha: \mathbb{R} \rightarrow [0, 1]$ and the warping function $f: \mathbb{R} \rightarrow (-\infty, 1]$.

We define $f(t) := \alpha(t) \frac{t+R}{\lambda} + (1 - \alpha(t))$ and consider the conical metric

$$g_{\text{cone}} = f(t)^2 h + dt^2$$

on $S^{n-1} \times (-R, 0]$. Its scalar curvature is given by

$$(2.1) \quad \text{scal}_{g_{\text{cone}}} = \frac{1}{f(t)^2} \left(\text{scal}_h - 2(n-1)f(t)f''(t) - (n-1)(n-2)f'(t)^2 \right).$$

¹This also follows directly from [17, Theorem B] and the fact that $\Omega_5^{\text{spin}} = 0$.




FIGURE 2. The metric \hat{g} on M : On $M \setminus D^n$ we have a psc metric g which is collared, then there is a transition region $S^{n-1} \times [-R + \varepsilon, 0]$ and on $S^{n-1} \times (-R, R + \varepsilon]$ the metric g_{cone} is the linear scaling of h .

We note that, by our choices, we have for $t > -R$

$$\begin{aligned}
 f(t) &= -\underbrace{\alpha(t)\left(1 - \frac{t+R}{\lambda}\right)}_{\in[0,1]} + 1 \in (0, 1] \\
 |f'(t)| &= |\alpha'(t)\left(\frac{t+R}{\lambda} - 1\right) + \alpha(t)\frac{1}{\lambda}| \leq |\alpha'(t)| \cdot \underbrace{\left|\frac{t+R}{\lambda} - 1\right|}_{\leq 1} + \underbrace{|\alpha(t)|}_{\leq 1} \cdot \frac{1}{\lambda} \\
 &\leq -\alpha'(t) + \frac{1}{\lambda} \leq \sqrt{\frac{C}{4(n-1)(n-2)}} \\
 (2.2) \quad &\Rightarrow (n-1)(n-2)f'(t)^2 \leq \frac{C}{4}
 \end{aligned}$$

$$\begin{aligned}
 f''(t) &= \alpha''(t)\left(\frac{t+R}{\lambda} - 1\right) + \underbrace{\frac{2\alpha'(t)}{\lambda}}_{\leq 0} \leq |\alpha''(t)| \left(1 - \frac{t+R}{\lambda}\right) \\
 &\leq |\alpha''(t)| \leq \frac{C}{8(n-1)} \\
 (2.3) \quad &\Rightarrow 2(n-1)f(t)f''(t) \leq 2(n-1)f''(t) \leq \frac{C}{4}.
 \end{aligned}$$

Inserting Equation (2.2) and Equation (2.3) into Equation (2.1), we obtain:

$$\text{scal}_{g_{\text{cone}}} = \frac{1}{f(t)^2} \left(\text{scal}_h - 2(n-1)f(t)f''(t) - (n-1)(n-2)f'(t)^2 \right) \geq \frac{C}{2} > 0.$$

Note that $f(-R) = 0$ and, since $g_{\text{cone}} = h + dt^2$ for $t > -\varepsilon$, we can extend the metric g on M' by g_{cone} to a metric \hat{g} on M which in general has a true singularity at the point $p \in M$ corresponding to $t = -R$. A visualization of \hat{g} can be found in Figure 2. Near the singularity we have $\text{scal}_{\hat{g}} > \frac{C}{2}$, so \hat{g} has uniformly positive scalar curvature on all of $M \setminus \{p\}$.

It remains to verify that \hat{g} is uniformly Euclidean and satisfies the desired Sobolev regularity on all of M . Since the metric is smooth away from the singularity, it suffices to show that the metric has these properties on some neighborhood including the singularity, where the metric is precisely conical. This follows from a direct local computation which we record in Lemma 2.4 below. \square

Lemma 2.4. *Let h be a Riemannian metric on S^{n-1} . Then the metric $g = t^2h + dt^2$ on $D^n \setminus \{0\} \cong S^{n-1} \times (0, 1]$, identified via $x \mapsto (x/|x|, |x|)$, is uniformly Euclidean on the entire disk D^n and its coefficients lie in $W^{2,p/2}(D^n) \cap L^\infty(D^n)$ for every $p < n$.*

Proof. Since S^{n-1} is compact, the metric h is “uniformly round”, that is, there exists $\Lambda_\circ > 1$ such that $\Lambda_\circ^{-1}h_\circ \leq h \leq \Lambda_\circ h_\circ$ for h_\circ the round metric on S^{n-1} . Then $g_\circ = t^2h_\circ + dt^2$ corresponds to the standard Euclidean metric on D^n and we have

$$\Lambda_\circ^{-1}g_\circ \leq \Lambda_\circ^{-1}t^2h_\circ + dt^2 \leq t^2h + dt^2 = g \leq \Lambda_\circ t^2h_\circ + dt^2 \leq \Lambda_\circ g_\circ,$$

so g is uniformly Euclidean on D^n .

To verify the Sobolev regularity, let $\Phi: D^n \setminus \{0\} \rightarrow S^{n-1} \times (0, 1]$ denote the diffeomorphism sending $x = (x^1, \dots, x^n)$ to $(x/|x|, |x|)$. Then

$$\frac{\partial \Phi}{\partial x^i} = \frac{1}{|x|} \left(\frac{\partial}{\partial x^i} - x^i \frac{\mathbf{x}}{|x|^2}, x^i \right),$$

where $\mathbf{x} = \sum_{k=1}^n x^k \frac{\partial}{\partial x^k}$ is the identity viewed as vector field. Thus

$$g_{ij}(x) = g_x \left(\frac{\partial \Phi}{\partial x^i}, \frac{\partial \Phi}{\partial x^j} \right) = h_{x/|x|} \left(\frac{\partial}{\partial x^i} - x^i \frac{\mathbf{x}}{|x|^2}, \frac{\partial}{\partial x^j} - x^j \frac{\mathbf{x}}{|x|^2} \right) + \frac{x^i x^j}{|x|^2}$$

where we used the identification $t = |x|$ under the diffeomorphism Φ . We deduce that we can write $g_{ij}(x) = f_{ij}(x/|x|)$ for some smooth functions $f_{ij}: \mathbb{R}^n \rightarrow \mathbb{R}$. Hence, $g_{ij} \in \mathcal{O}(1)$, $\partial_k g_{ij} \in \mathcal{O}(t^{-1})$ and $\partial_l \partial_k g_{ij} \in \mathcal{O}(t^{-2})$ as $t \rightarrow 0$. This shows that each g_{ij} is in $W^{2,p/2}(D^n) \cap L^\infty(D^n)$, for every $p < n$. \square

To prove Theorem A, we need closed simply connected manifolds which do not support positive scalar curvature. In dimensions ≥ 5 such examples must be spin and they exist whenever the receptacle for the Hitchin α -invariant [21] does not vanish. The latter is the group KO_n which is \mathbb{Z} if $n \equiv 0, 4 \pmod{8}$, $\mathbb{Z}/2\mathbb{Z}$ if $n \equiv 1, 2 \pmod{8}$ and vanishes otherwise.

Remark 2.5. If $4 \leq n \equiv 0, 1, 2$, or $4 \pmod{8}$, then there exists a simply connected closed spin n -manifold B which does not admit a smooth positive scalar curvature metric because $\alpha(B; 1) \neq 0 \in \mathrm{KO}_n$. Indeed, the Atiyah–Bott–Shapiro orientation yields a surjective homomorphism from the spin cobordism group $\Omega_n^{\mathrm{spin}} \rightarrow \mathrm{KO}_n$ for $n \geq 0$ and for $n \geq 4$ we can use surgery to find simply connected representatives. More concretely, for $n \equiv 0$ or $4 \pmod{8}$ we can just take B to be a product of K3 surfaces to obtain a simply connected spin manifold with non-vanishing \widehat{A} -genus. In the cases $8 \leq n \equiv 1, 2 \pmod{8}$ one can find exotic spheres with non-vanishing Hitchin genus in $\mathbb{Z}/2\mathbb{Z}$. This follows from classical computations of Adams [1] and Milnor [33], see also the discussion in [22, §4].

Proof of Theorem A. In the cases $8 \leq n \equiv 0, 1, 2$, or $4 \pmod{8}$ there exists a simply connected spin n -manifold B which does not admit a smooth positive scalar curvature as explained in Remark 2.5 above. By Proposition 2.1 however, there exists an L^∞ -metric of regularity $W^{2,p/2}$ on B which is smooth away from a point and has uniformly positive scalar curvature. This metric is already the desired counterexample.

The cases of other dimension parities follow from the fact that the product of a smooth metric on another manifold N and an L^∞ -metric of regularity $W^{k,p}$ on B which is smooth on $B \setminus \{*\}$ yields an L^∞ -metric on $B \times N$ which is of regularity $W^{k,p}$ and smooth on $B \times N \setminus (\{*\} \times N)$. Therefore, on the one hand, $M := B \times \mathbb{T}^k$ admits an L^∞ -metric which is smooth with uniformly positive scalar curvature outside $S = \{*\} \times \mathbb{T}^k \subset M$. Moreover, such metric has regularity $W^{2,p/2}$ for every $p < n$. Note that the codimension of S in M is equal to $\dim(B) = n \geq 8$. On the other hand, $\alpha(B \times \mathbb{T}^k; \mathbb{Z}^k) \neq 0$ (see for instance [44, Appendix B]) and so M does not support a smooth metric of positive scalar curvature. \square

3. NON-SMOOTHABILITY OF L^∞ -METRICS ON \mathbb{R}^n WITH POSITIVE SCALAR CURVATURE OUTSIDE A POINT

Here we prove Theorem B by a similar construction as in the previous section.

Proof of Theorem B. Let n be as specified in the statement of the theorem. Then by Remark 2.5 and Lemma 2.2 there exists a closed n -manifold M which does not admit psc but there exists a psc metric on g_M on $M \setminus D^n$ that is collared near the boundary. In particular, we can arrange that $g_M = h + dt^2$ near $\partial(M \setminus D^n) = S^{n-1}$ for some metric h on S^{n-1} with $\text{scal}_h \geq 1$. We define an L^∞ -metric on \mathbb{R}^n by

$$g := \left(\frac{t}{\lambda}\right)^2 h + dt^2$$

where $\lambda := \sqrt{2(n-1)(n-2)}$. As in the proof of Lemma 2.4, since any metric on S^{n-1} is “uniformly round” and the Euclidean metric is given by $g_o = t^2 h_o + dt^2$, this means that there exists a constant $\Lambda > 1$ such that $\Lambda^{-1} g_o \leq g \leq \Lambda g_o$. The metric g is smooth on $\mathbb{R}^n \setminus \{0\}$ and the desired $W_{\text{loc}}^{2,p/2}$ -regularity near 0 also follows from Lemma 2.4. Its scalar curvature on $\mathbb{R}^n \setminus \{0\}$ is given by

$$\text{scal}_g = \left(\frac{\lambda}{t}\right)^2 \left(\text{scal}_h - \frac{(n-1)(n-2)}{\lambda^2}\right) \geq \frac{1}{2} \lambda^2 t^{-2} > 0.$$

Furthermore, the induced metric on $S_\lambda^{n-1} = \{t = \lambda\} \subset \mathbb{R}^n$ is precisely given by h . The mean curvature² H_λ of S_λ^{n-1} satisfies

$$H_\lambda = (n-1) \frac{1}{\lambda} > 0.$$

Let us now proceed by contradiction and assume the existence of the family of smooth metrics g_t with non-negative scalar curvature and converging to g in $C_{\text{loc}}^\infty(M \setminus \{0\})$. Then there exists an $\varepsilon > 0$ such that $\text{scal}_{g_\varepsilon}(p) \geq \frac{1}{4} \lambda^2$ for all $\lambda/2 \leq |p| \leq \lambda$, the metric h_ε induced by g_ε on S_λ^n satisfies $\text{scal}_{h_\varepsilon} \geq \frac{1}{2} > 0$ and the mean curvature of S_λ^n is $H_\lambda \geq (n-1)/(2\lambda)$. Consider now the restriction of g_ε to D_λ^n . It has non-negative scalar curvature overall, positive scalar curvature near the mean convex boundary ∂D_λ^n and the induced metric on the boundary equals h_ε and has positive scalar curvature. Thus we can assume that g_ε is collared while retaining non-negative scalar curvature and without changing the restriction to the boundary S_λ^{n-1} , again by [3, Corollary 4.9]³.

Furthermore, if ε is small enough, h_ε and h are isotopic through psc metrics, since the space of psc metrics is an open subspace of the space of all metrics in the (weak) C^∞ -topology. Hence, there is a concordance from h to h_ε by [17, Lemma 3] (see also [16, Lemma 2.5]), that is, there is a psc metric G on the cylinder $S^{n-1} \times [0, 1]$ which is collared near both boundaries and restricting to $h_\varepsilon + dt^2$ and $h + dt^2$ on the respective boundary components.

Gluing G onto $g_\varepsilon|_{D_\lambda^n}$, we obtain a metric on D^n of non-negative scalar curvature with collared boundary that restricts to h near the boundary. Then we can glue this metric onto g_M to obtain a metric on the closed manifold M which has non-negative and non-vanishing scalar curvature. But this would mean that M also has a metric of positive scalar curvature (for instance, run the Ricci flow for a short time), a contradiction to the choice of M . \square

²Using the convention that the mean curvature of the unit sphere with respect to the Euclidean metric is equal to $n-1 > 0$.

³Even though it is not stated in loc.cit. the deformation principle they prove leaves the metric on the boundary fixed ([3, Theorem 3.7 (b)]). In fact the proof of [3, Theorem 4.8] also works for a fixed psc metric h on the boundary and then [3, Theorem 4.5] with $\mathcal{X} = \{h\}$.

Theorem B affirmatively answers [5, Question 3]. In *loc. cit.* it is shown that an L^∞ -metric g admits a “smoothing family” $(g_t)_{t>0}$ such as in the statement of Theorem B, provided the metric g is L^∞ -close to the Euclidean one. Burkhardt-Guim [5, Question 3] then asks if the assumption of L^∞ -closeness is necessary. Indeed, our Theorem B shows that this condition cannot be replaced by merely requiring the metric to be uniformly bilipschitz to the Euclidean metric.

4. RELATIVE DIRAC OBSTRUCTIONS

The proof of Theorem C is based on the following proposition.

Proposition 4.1. *Let M be a closed n -dimensional spin manifold with fundamental group Γ . Let S be a finite subset of M , and let g be a uniformly Euclidean metric on M which is smooth on $M \setminus S$. If $\text{scal}_g > 0$ on $M \setminus S$, then $j(\alpha(M; 1)) = \alpha(M; \Gamma)$.*

Proof. To simplify the notation, set $\hat{M} = M \setminus S$ and $\hat{g} = g|_{\hat{M}}$. We will first construct a complete psc metric on \hat{M} following [28, Proof of Proposition 6.2] and [23, Proof of Theorem A]. Consider the modified conformal Laplacian

$$\mathcal{L}_\sigma := -4 \frac{n-1}{n-2} \Delta + \sigma,$$

where Δ is the Laplace-Beltrami operator for (\hat{M}, \hat{g}) , and σ is a positive function on $M \setminus S$ such that $0 < \sigma \leq \min(1, \text{scal}_{\hat{g}})$. Let G be the positive distributional solution of the elliptic PDE

$$\mathcal{L}_\sigma G = \delta_S \quad \text{on } M,$$

where δ_S is the Dirac delta measure on S . By [29, Theorem 7.1] (see also [9, Appendix A] for more details), there exists $c > 0$ such that

$$(4.1) \quad c^{-1} \text{dist}_g(S, x)^{2-n} \leq G(x) \leq c \text{dist}_g(S, x)^{2-n} \quad \text{on } M \setminus S.$$

For $\varepsilon > 0$, consider the metric $g_\varepsilon := (1 + \varepsilon G)^{\frac{4}{n-2}} \hat{g}$ on \hat{M} . By (4.1), g_ε is complete. By direct computation,

$$\text{scal}_{g_\varepsilon} = (1 + \varepsilon G)^{-\frac{n+2}{n-2}} (\text{scal}_{\hat{g}} + \varepsilon (\text{scal}_{\hat{g}} - \sigma) G) \geq \text{scal}_{\hat{g}} (1 + \varepsilon G)^{-\frac{n+2}{n-2}} > 0.$$

In conclusion, we constructed a complete metric of positive scalar curvature on \hat{M} . By a direct application of [6, Theorem C], we conclude that $j(\alpha(M; 1)) = \alpha(M; \Gamma)$. \square

Proof of Theorem C. We proceed by contraposition. Suppose that $\text{Ric}_g \not\equiv 0$ on $M \setminus S$. By [28, Section 7, Proof of Theorem 1.7], there exists a uniformly Euclidean metric g_1 on M which is smooth on $M \setminus S$ and such that $\text{scal}_{g_1} > 0$ on $M \setminus S$. Then Proposition 4.1 shows that $j(\alpha(M; 1)) = \alpha(M; \Gamma)$. \square

REFERENCES

- [1] J. F. Adams. “On the groups $J(X)$. IV”. In: *Topology* 5 (1966), pp. 21–71. DOI: [10.1016/0040-9383\(66\)90004-8](https://doi.org/10.1016/0040-9383(66)90004-8).
- [2] R. H. Bamler. “A Ricci flow proof of a result by Gromov on lower bounds for scalar curvature”. In: *Math. Res. Lett.* 23.2 (2016), pp. 325–337. DOI: [10.4310/MRL.2016.v23.n2.a2](https://doi.org/10.4310/MRL.2016.v23.n2.a2).
- [3] C. Bär and B. Hanke. “Boundary conditions for scalar curvature”. In: *Perspectives in scalar curvature. Vol. 2*. Singapore: World Scientific, 2023, pp. 325–377. DOI: [10.1142/9789811273230_0010](https://doi.org/10.1142/9789811273230_0010).
- [4] P. Burkhardt-Guim. “Pointwise lower scalar curvature bounds for C^0 metrics via regularizing Ricci flow”. In: *Geom. Funct. Anal.* 29.6 (2019), pp. 1703–1772. DOI: [10.1007/s00039-019-00514-3](https://doi.org/10.1007/s00039-019-00514-3).
- [5] P. Burkhardt-Guim. *Smoothing L^∞ Riemannian metrics with nonnegative scalar curvature outside of a singular set*. 2024. arXiv: [2406.04564v1](https://arxiv.org/abs/2406.04564v1) [math.DG].

- [6] S. Cecchini. “A long neck principle for Riemannian spin manifolds with positive scalar curvature”. In: *Geom. Funct. Anal.* 30.5 (2020), pp. 1183–1223. DOI: [10.1007/s00039-020-00545-1](https://doi.org/10.1007/s00039-020-00545-1).
- [7] S. Cecchini, B. Hanke, and T. Schick. *Lipschitz rigidity for scalar curvature*. To appear in *J. Eur. Math. Soc. (JEMS)*. 2023. arXiv: [2206.11796](https://arxiv.org/abs/2206.11796) [math.DG].
- [8] S. Cecchini and R. Zeidler. “Scalar curvature and generalized Callias operators”. In: *Perspectives in scalar curvature. Vol. 1*. World Sci. Publ., Hackensack, NJ, 2023, pp. 515–542.
- [9] M.-C. Cheng, M.-C. Lee, and L.-F. Tam. “Singular metrics with negative scalar curvature”. In: *Int. J. Math.* 33.7 (2022). Id/No 2250047, p. 27. DOI: [10.1142/S0129167X22500471](https://doi.org/10.1142/S0129167X22500471).
- [10] V. Chernysh. *On the homotopy type of the space $\mathcal{R}^+(M)$* . 2004. arXiv: [math/0405235](https://arxiv.org/abs/math/0405235) [math.GT].
- [11] J. Chu and M.-C. Lee. *Ricci-DeTurck flow from rough metrics and applications*. 2022. arXiv: [2204.05843](https://arxiv.org/abs/2204.05843) [math.DG].
- [12] J. Chu, M.-C. Lee, and J. Zhu. *Llarull’s theorem on punctured sphere with L^∞ metric*. 2024. arXiv: [2405.19724](https://arxiv.org/abs/2405.19724) [math.DG].
- [13] X. Dai, Y. Sun, and C. Wang. *Positive mass theorem for asymptotically flat spin manifolds with isolated conical singularities*. 2023. arXiv: [2310.13285](https://arxiv.org/abs/2310.13285) [math.DG].
- [14] X. Dai, Y. Sun, and C. Wang. *Positive mass theorem for asymptotically flat manifolds with isolated conical singularities*. 2024. arXiv: [2401.07186](https://arxiv.org/abs/2401.07186) [math.DG].
- [15] C. Dong and A. Song. *Stability of Euclidean 3-space for the positive mass theorem*. 2023. arXiv: [2302.07414](https://arxiv.org/abs/2302.07414) [math.DG].
- [16] J. Ebert and G. Frenck. “The Gromov-Lawson-Chernysh surgery theorem”. In: *Bol. Soc. Mat. Mex. (3)* 27.2 (2021), Paper No. 37, 43. DOI: [10.1007/s40590-021-00310-w](https://doi.org/10.1007/s40590-021-00310-w).
- [17] M. Gromov and H. B. Lawson Jr. “The classification of simply connected manifolds of positive scalar curvature”. In: *Ann. of Math. (2)* 111.3 (1980), pp. 423–434. DOI: [10.2307/1971103](https://doi.org/10.2307/1971103).
- [18] M. Gromov. “Dirac and Plateau billiards in domains with corners”. In: *Cent. Eur. J. Math.* 12.8 (2014), pp. 1109–1156. DOI: [10.2478/s11533-013-0399-1](https://doi.org/10.2478/s11533-013-0399-1).
- [19] B. Hanke and T. Schick. “Enlargeability and index theory”. In: *J. Differential Geom.* 74.2 (2006), pp. 293–320. URL: <http://projecteuclid.org/euclid.jdg/1175266206>.
- [20] B. Hanke and T. Schick. “Enlargeability and index theory: infinite covers”. In: *K-Theory* 38.1 (2007), pp. 23–33. DOI: [10.1007/s10977-007-9004-3](https://doi.org/10.1007/s10977-007-9004-3).
- [21] N. J. Hitchin. “Harmonic spinors”. In: *Adv. Math.* 14 (1974), pp. 1–55. DOI: [10.1016/0001-8708\(74\)90021-8](https://doi.org/10.1016/0001-8708(74)90021-8).
- [22] M. Joachim and D. J. Wraith. “Exotic spheres and curvature”. In: *Bull. Amer. Math. Soc. (N.S.)* 45.4 (2008), pp. 595–616. DOI: [10.1090/S0273-0979-08-01213-5](https://doi.org/10.1090/S0273-0979-08-01213-5).
- [23] D. Kazaras. “Desingularizing positive scalar curvature 4-manifolds”. In: *Math. Ann.* (2024). DOI: [10.1007/s00208-024-02829-5](https://doi.org/10.1007/s00208-024-02829-5).
- [24] T. Lamm and M. Simon. “Ricci flow of $W^{2,2}$ -metrics in four dimensions”. In: *Comment. Math. Helv.* 98.2 (2023), pp. 261–364. DOI: [10.4171/CMH/553](https://doi.org/10.4171/CMH/553).
- [25] D. A. Lee and P. G. LeFloch. “The positive mass theorem for manifolds with distributional curvature”. In: *Commun. Math. Phys.* 339.1 (2015), pp. 99–120. DOI: [10.1007/s00220-015-2414-9](https://doi.org/10.1007/s00220-015-2414-9).
- [26] M.-C. Lee, A. Naber, and R. Neumayer. “ d_p -convergence and ϵ -regularity theorems for entropy and scalar curvature lower bounds”. In: *Geom. Topol.* 27.1 (2023), pp. 227–350. DOI: [10.2140/gt.2023.27.227](https://doi.org/10.2140/gt.2023.27.227).
- [27] M.-C. Lee and L.-F. Tam. *Continuous metrics and a conjecture of Schoen*. 2021. arXiv: [2111.05582](https://arxiv.org/abs/2111.05582) [math.DG].
- [28] C. Li and C. Mantoulidis. “Positive scalar curvature with skeleton singularities”. In: *Math. Ann.* 374.1-2 (2019), pp. 99–131. DOI: [10.1007/s00208-018-1753-1](https://doi.org/10.1007/s00208-018-1753-1).
- [29] W. Littman, G. Stampacchia, and H. F. Weinberger. “Regular points for elliptic equations with discontinuous coefficients”. In: *Ann. Scuola Norm. Sup. Pisa Cl. Sci.*

- (3) 17 (1963), pp. 43–77. URL: http://www.numdam.org/item/ASNSP_1963_3_17_1-2_43_0/.
- [30] J. Lohkamp. “Scalar curvature and hammocks”. In: *Math. Ann.* 313.3 (1999), pp. 385–407. DOI: [10.1007/s002080050266](https://doi.org/10.1007/s002080050266).
- [31] C. Mantoulidis, P. Miao, and L.-F. Tam. “Capacity, quasi-local mass, and singular fill-ins”. In: *J. Reine Angew. Math.* 768 (2020), pp. 55–92. DOI: [10.1515/crelle-2019-0040](https://doi.org/10.1515/crelle-2019-0040).
- [32] P. Miao. “Positive mass theorem on manifolds admitting corners along a hypersurface”. In: *Adv. Theor. Math. Phys.* 6.6 (2002), pp. 1163–1182. DOI: [10.4310/ATMP.2002.v6.n6.a4](https://doi.org/10.4310/ATMP.2002.v6.n6.a4).
- [33] J. W. Milnor. *Remarks concerning spin manifolds*. Differ. and Combinat. Topology, Sympos. Marston Morse, Princeton, 55–62 (1965). 1965.
- [34] J. Rosenberg. “ C^* -algebras, positive scalar curvature and the Novikov conjecture. II”. In: *Geometric methods in operator algebras (Kyoto, 1983)*. Vol. 123. Pitman Res. Notes Math. Ser. Longman Sci. Tech., Harlow, 1986, pp. 341–374.
- [35] J. Rosenberg. “ C^* -algebras, positive scalar curvature, and the Novikov conjecture”. In: *Inst. Hautes Études Sci. Publ. Math.* 58 (1983), 197–212 (1984). URL: http://www.numdam.org/item?id=PMIHES_1983__58__197_0.
- [36] J. Rosenberg. “ C^* -algebras, positive scalar curvature, and the Novikov conjecture. III”. In: *Topology* 25.3 (1986), pp. 319–336. DOI: [10.1016/0040-9383\(86\)90047-9](https://doi.org/10.1016/0040-9383(86)90047-9).
- [37] T. Schick. “The topology of positive scalar curvature”. In: *Proceedings of the International Congress of Mathematicians—Seoul 2014. Vol. II*. Kyung Moon Sa, Seoul, 2014, pp. 1285–1307.
- [38] R. Schoen and S. T. Yau. “On the structure of manifolds with positive scalar curvature”. In: *Manuscripta Math.* 28.1-3 (1979), pp. 159–183. DOI: [10.1007/BF01647970](https://doi.org/10.1007/BF01647970).
- [39] Y. Shi and L.-F. Tam. “Positive mass theorem and the boundary behaviors of compact manifolds with nonnegative scalar curvature”. In: *J. Differential Geom.* 62.1 (2002), pp. 79–125. URL: <http://projecteuclid.org/euclid.jdg/1090425530>.
- [40] M. Simon. “Deformation of C^0 Riemannian metrics in the direction of their Ricci curvature”. In: *Commun. Anal. Geom.* 10.5 (2002), pp. 1033–1074. DOI: [10.4310/CAG.2002.v10.n5.a7](https://doi.org/10.4310/CAG.2002.v10.n5.a7).
- [41] C. Sormani. “Scalar curvature and intrinsic flat convergence”. In: *Measure theory in non-smooth spaces*. Warsaw: De Gruyter Open, 2017, pp. 288–338.
- [42] S. Stolz. “Simply connected manifolds of positive scalar curvature”. In: *Ann. Math. (2)* 136.3 (1992), pp. 511–540. DOI: [10.2307/2946598](https://doi.org/10.2307/2946598).
- [43] R. Zeidler. “Width, largeness and index theory”. In: *SIGMA, Symmetry Integrability Geom. Methods Appl.* 16 (2020), paper 127, 15. DOI: [10.3842/SIGMA.2020.127](https://doi.org/10.3842/SIGMA.2020.127).
- [44] R. Zeidler. “Band width estimates via the Dirac operator”. In: *J. Differ. Geom.* 122.1 (2022), pp. 155–183. DOI: [10.4310/jdg/1668186790](https://doi.org/10.4310/jdg/1668186790).

(Simone Cecchini) DEPARTMENT OF MATHEMATICS, TEXAS A&M UNIVERSITY, COLLEGE STATION, TX 77843, USA

Email address: cecchini@tamu.edu

URL: simonececchini.org

(Georg Frenck) INSTITUT FÜR MATHEMATIK, UNIVERSITÄT AUGSBURG, UNIVERSITÄTSSTRASSE 14, 86159 AUGSBURG, BAVARIA, GERMANY

Email address: math@frenck.net

URL: frenck.net

(Rudolf Zeidler) UNIVERSITY OF MÜNSTER, MATHEMATISCHES INSTITUT, EINSTEINSTR. 62, 48149 MÜNSTER, GERMANY

Email address: math@rzejidler.eu

URL: www.rzejidler.eu