

Self-consistent equilibrium of a force-free magnetic flux rope

Oleg K. Cheremnykh

Space Research Institute, Pr. Glushkova 40 k.4/1, Kyiv 03187, Ukraine

Volodymyr M. Lashkin*

¹*Institute for Nuclear Research, Pr. Nauki 47, Kyiv 03028, Ukraine and*

²*Space Research Institute, Pr. Glushkova 40 k.4/1, Kyiv 03187, Ukraine*

We present an exact solution to the problem of a self-consistent equilibrium force-free magnetic flux rope. Unlike other approaches, we use magnetostatic equations and assume only a relatively rapid decrease in the axial magnetic field at infinity. For the first time we obtain a new nonlinear equation for the axial current density, the derivation of which does not require any phenomenological assumptions. From the resulting nonlinear equation, we analytically find the radial profiles of the components of the magnetic field strength and current density.

The concept of a magnetic flux rope is widely used in solar physics and has been an area of active research over the years. In particular, it is assumed that such structures play a dominant role in the release of flare energy and in the formation of coronal mass ejections [1–5]. Magnetic clouds, which are large-scale magnetic ropes, are also often observed in interplanetary space [6, 7]. As a theoretical model, the magnetic ropes are usually considered as plasma tubes with a helical magnetic field. Among these models, much attention has been paid to the so-called force-free plasma configurations, in which the current is parallel to the magnetic field. It is believed [8] that in closed plasma systems a force-free magnetic field is a field with minimal energy and therefore the force-free magnetic ropes are stable entities widespread in space plasma.

The simplest force-free configurations usually correspond to cylindrical symmetry, and then the force-free plasma tube is considered in a cylindrical coordinate system (r, φ, z) as a plasma cylinder with two components of magnetic field strength B_φ and B_z . Since within the framework of such a model there is only one equation for the magnetic field components (see, e.g., [4]), there is an arbitrariness in the choice of configurations. The most well-known and frequently used force-free cylindrical magnetic configurations are those obtained by Lundquist [9–11] and Gold-Hoyle [12–14]. The force-free magnetic fields of Lundquist and Gold-Hoyle are exact solutions to the equations of plasma hydrostatics, however, they are not the only solutions within the framework of the force-free plasma cylinder model and the problem of finding other solutions is still ongoing [15–18]. It should be noted that the results obtained by Lundquist [10] and Gold-Hoyle [12] were obtained under quite significant additional restrictions. In [10] it was assumed that the magnetic field strength and current density are related by a constant value, and in [12] a constant angle of twist of the magnetic field lines was assumed. That is, the forceless

equilibrium in those works was "set by hand".

In this paper, we do not make any additional assumptions about the behaviour of equilibrium quantities in force-free configurations and use the approach of [19], which allows us to analytically find a self-consistent equilibrium. Following this paper, we find and add an integral equilibrium equation to the three commonly used local equilibrium equations. We obtain a new nonlinear equation for the axial current density and this makes it possible to determine uniquely the self-consistent equilibrium state of a force-free magnetic rope.

We consider an inhomogeneous equilibrium state of a force-free plasma configuration, which is established self-consistently when a helical current flows in a low-pressure plasma. Since in the configuration under consideration the Lorentz force significantly exceeds the pressure gradient, the standard equations of magnetic hydrostatics, in which we neglect the gas-kinetic pressure,

$$\mathbf{j} \times \mathbf{B} = 0, \quad (1)$$

$$\nabla \times \mathbf{B} = \frac{4\pi}{c} \mathbf{j}, \quad (2)$$

$$\nabla \cdot \mathbf{B} = 0, \quad (3)$$

are a good approximation for describing the equilibrium of a force-free magnetic rope [4]. Throughout the paper, cgs units are used. Here \mathbf{j} is the electric current density, \mathbf{B} is the magnetic field strength, and c is the speed of light. From equation (1) it follows that the electric current flows along the magnetic field lines, and therefore we can write

$$\mathbf{j} = \mu \mathbf{B}, \quad (4)$$

where unlike [10] we do not assume the coefficient μ to be a constant.

Let us assume that in a cylindrical coordinate system the components of the magnetic field strength \mathbf{B} and electric current density \mathbf{j} depend only on the radial coordinate r , that is, we consider an axially symmetric configuration with $\mathbf{B} = (0, B_\varphi(r), B_z(r))$ and

*Electronic address: vlashkin62@gmail.com

$\mathbf{j} = (0, j_\varphi(r), j_z(r))$. It can be seen that in this case eq. (3) is satisfied automatically. Magnetic field lines lie on cylindrical surfaces with radius r and form spirals with a twist of field lines per unit length of the plasma cylinder k equal to

$$k = \frac{d\varphi}{dz} = \frac{B_\varphi}{rB_z}. \quad (5)$$

Equations (1) and (2) in a cylindrical coordinate system have the form

$$j_z B_\varphi - j_\varphi B_z = 0, \quad (6)$$

and

$$j_\varphi = -\frac{c}{4\pi} \frac{dB_z}{dr}, \quad j_z = \frac{c}{4\pi r} \frac{d}{dr}(rB_\varphi), \quad (7)$$

respectively. Eliminating the current density from eqs. (6) and (7), we have equation

$$\frac{1}{2} \frac{dB_z^2}{dr} + \frac{B_\varphi}{r} \frac{d}{dr}(rB_\varphi) = 0, \quad (8)$$

containing two independent radial profiles $B_z(r)$ and $B_\varphi(r)$. A unique solution to the equilibrium equations of a magnetic flux rope can be obtained if one more equation for equilibrium quantities is added to them. In a recent work [19] on the study of the equilibrium of a magnetic flux rope with a finite plasma pressure with a constant B_z , it was shown that such an equation can be found from the integral equilibrium equation. For a force-free plasma configuration, we similarly obtain an additional equilibrium local equation using the integral equilibrium equation.

Rewriting eq. (8) in the form

$$r^2 \frac{dB_z^2}{dr} + \frac{d}{dr}(rB_\varphi)^2 = 0, \quad (9)$$

and integrating it over r from 0 to ∞ , we have

$$\int_0^\infty \frac{dB_z^2}{dr} r^2 dr = - (rB_\varphi)^2 \Big|_0^\infty. \quad (10)$$

Integrating eq. (7) one can obtain

$$rB_\varphi \Big|_{r \rightarrow \infty} = \frac{2I_z}{c}, \quad (11)$$

where I_z is the total current flowing through the plasma in the z -direction:

$$I_z = 2\pi \int_0^\infty j_z r dr. \quad (12)$$

Note that the regularity condition on B_φ ensures that $rB_\varphi \rightarrow 0$ at $r = 0$. Equation (10) then becomes

$$\int_0^\infty \frac{dB_z^2}{dr} r^2 dr = -\frac{4I_z^2}{c^2}. \quad (13)$$

Integrating by parts the left side of this equation gives

$$(rB_z)^2 \Big|_0^\infty - 2 \int_0^\infty B_z^2 r dr = -\frac{4I_z^2}{c^2}. \quad (14)$$

Assuming that B_z^2 is bounded at $r = 0$ and vanishes at infinity faster than $\sim 1/r^2$, we obtain the following integral equilibrium equation for a force-free magnetic flux rope:

$$\int_0^\infty B_z^2 r dr = \frac{2I_z^2}{c^2}. \quad (15)$$

Taking into account eq. (12), one can rewrite eq. (15) in the form

$$\int_0^\infty \left(B_z^2 - \frac{4\pi j_z I_z}{c^2} \right) r dr = 0. \quad (16)$$

It can be seen that equation (16) is satisfied if the z -components of the magnetic field strength B_z and current density j_z satisfy the condition:

$$B_z^2 = \alpha j_z, \quad (17)$$

where $\alpha = 4\pi I_z/c^2$ is a constant determined by the total current. Setting $r = 0$ in eq. (17), the constant α can also be written as $\alpha = B_0^2/j_0$, where we have introduced notations for the values of the axial magnetic field and current density on the axis:

$$B_0 = B_z \Big|_{r=0}, \quad j_0 = j_z \Big|_{r=0}. \quad (18)$$

It is assumed that the axial current density j_z is non-zero. This assumption is consistent with the subsequent explicit expression for j_z . Then, the total current can be written as

$$I_z = \frac{c^2 B_0^2}{4\pi j_0}. \quad (19)$$

Equation (17) is the desired additional equilibrium local equation and, along with eqs. (1)-(4), allows us to self-consistently find all equilibrium quantities. From eqs. (7), (8) and (17) we can obtain a nonlinear equation for the axial current density j_z :

$$\frac{1}{r} \frac{d}{dr} \left(\frac{r}{j_z} \frac{dj_z}{dr} \right) + \frac{8\pi}{I_z} j_z = 0. \quad (20)$$

The solution of this equation with boundary conditions

$$j_z = j_0 \quad \text{at } r = 0, \quad j_z \rightarrow 0 \quad \text{at } r \rightarrow \infty, \quad (21)$$

along with eqs. (7), (8) and (17) allows us to self-consistently find all equilibrium quantities. As will be shown below, this solution can be extended to the problem where the current is mainly contained in a plasma cylinder with a certain radius a . Such a cylinder can be considered as a force-free magnetic flux rope. Let us represent the axial current density in the form $j_z = j_0 \exp \psi$,

where the function $\psi(r)$, according to (21), satisfies the boundary conditions $\psi = 0$ at $r = 0$ and $\psi \rightarrow -\infty$ at $r \rightarrow \infty$. Then from eq. (20) we obtain a nonlinear equation for the function ψ :

$$\frac{1}{r} \frac{d}{dr} \left(r \frac{d\psi}{dr} \right) + \beta \exp \psi = 0, \quad (22)$$

where $\beta = 32\pi^2 j_0 / (\alpha c^2)$. The solution to equation (22) was found in [19] and has the form:

$$\psi = \ln \frac{1}{(1 + r^2/r_*^2)^2}, \quad (23)$$

where

$$r_*^2 = \frac{8}{\beta} = \frac{c^2 B_0^2}{4\pi^2 j_0^2} = \frac{I_z}{\pi j_0}. \quad (24)$$

Note that $r_* = 2b$, where b is the unit of length introduced in the numerical simulation of force-free configurations in [20]. The radial dependence of the axial current density j_z is readily found from (23):

$$j_z = \frac{j_0}{(1 + r^2/r_*^2)^2}. \quad (25)$$

The value r_* can be treated as the radius of the cross section of the cylinder, inside of which the total axial current $I_z = \pi j_0 r_*^2$ flows with a uniform current density j_0 . From eqs. (7), (8) and (17) we find for the azimuthal magnetic field:

$$B_\varphi = -\frac{\alpha c}{8\pi j_z} \frac{dj_z}{dr}, \quad (26)$$

and substituting (25) into (26) we have:

$$B_\varphi = B_0 \frac{(r/r_*)}{(1 + r^2/r_*^2)}, \quad (27)$$

where

$$B_0 = \frac{2\pi r_* j_0}{c} = \frac{2\sqrt{\pi I_z j_0}}{c}. \quad (28)$$

The axial magnetic field B_z can be found from eq. (17) and (25):

$$B_z = \frac{B_0}{(1 + r^2/r_*^2)}. \quad (29)$$

Note that the B_0 defined by eq. (28) is consistent, as can be seen from eq. (29), with the boundary condition (18). In turn, from eqs. (7), (28), and (29) we have for the azimuthal current density:

$$j_\varphi = j_0 \frac{(r/r_*)}{(1 + r^2/r_*^2)^2}. \quad (30)$$

Equations (25), (27), (29), and (30) give expressions for the components of magnetic field strength and current density for an equilibrium force-free magnetic flux

rope and present the main result of this Letter. This force-free plasma configuration is analytically obtained from the magnetostatic equations only under the assumption that the axial magnetic field decrease at infinity faster than $\sim 1/r$. It can be seen that all equilibrium quantities depend on two free parameters j_0 and B_0 (or j_0 and I_z), which determine the self-consistent equilibrium of the force-free configuration of the magnetic flux rope. From the obtained expressions it also follows that $j_\varphi/j_z = B_\varphi/B_z = r/r_*$, so that the magnetic field and current density at distances small compared to r_* have mainly z -components, and at large distances, respectively, φ -components. Thus, the configurations of magnetic field lines and streamlines resemble the helical ropes.

From eqs. (5), (27) and (29) it follows that the twisting of the magnetic field lines for the resulting equilibrium magnetic field is equal to $k = 1/r_*$. Up to the change $1/r_* \rightarrow k$, the expressions for B_φ and B_z coincide with the expressions for components of the Gold-Hoyle magnetic field strength [12]. However, unlike the Gold-Hoyle solution, the twist of the field lines for the resulting solution is not an arbitrary constant value, but is expressed in terms of j_0 and I_z , and has the form:

$$k = \sqrt{\frac{\pi j_0}{I_z}} = \frac{2\pi j_0}{c B_0}, \quad (31)$$

where k is a constant, but depends on the values of j_0 and I_z . As follows from eqs. (25), (27), (29), and (30), the coefficient μ in equation (4) is not a constant but depends on r and is equal to

$$\mu = \frac{j_0}{B_0} \frac{1}{(1 + r^2/r_*^2)} = \frac{c}{2} \sqrt{\frac{j_0}{\pi I_z}} \frac{1}{(1 + \pi j_0 r^2 / I_z)}. \quad (32)$$

Radial profiles of equilibrium quantities B_φ , B_z , j_φ , and j_z are shown in Fig. 1. Note that they qualitatively coincide with the profiles obtained from the numerical simulation in [20]. The axial magnetic field B_z and current density j_z decrease monotonically with increasing r and exceed the azimuthal magnetic field B_φ and current density j_φ up to $r = r_*$, and then become smaller compared to the azimuthal magnetic field and current density. The azimuthal magnetic field and current density increase from zero at $r = 0$ to maximum values at $r = r_*$ and $r \approx 0.577r$ respectively. Then at $r = r_*$, they are equalized with the axial magnetic field and current density and decrease monotonically.

Since the radial profiles of the found equilibrium quantities do not vanish anywhere and disappear only at infinity one can introduce the concept of cross section radius a of a force-free magnetic flux rope. It can be defined as the distance at which the current flowing in the cylinder can be neglected with sufficient accuracy. The axial $I_z(r)$ and azimuthal $I_\varphi(r)$ currents flowing inside a cylinder of radius r and height $2\pi R$ are:

$$I_z(r) = 2\pi \int_0^r j_z r dr = I_z \frac{(r/r_*)^2}{(1 + r^2/r_*^2)}, \quad (33)$$




FIG. 1: Radial dependences of the normalized equilibrium quantities f : axial magnetic field B_z , azimuthal magnetic field B_φ , axial current density j_z , and azimuthal current density j_φ .

$$I_\varphi(r) = 2\pi \int_0^r j_\varphi R dr = I_\varphi \frac{(r/r_*)^2}{(1 + r^2/r_*^2)}, \quad (34)$$

where $I_\varphi = I_z R/r_*$ is the total azimuthal current. It can be seen that although the currents are distributed to infinity, the main part of them is concentrated near the z -axis. For example, assuming that the magnetic rope is bounded by a cylindrical surface of radius a , from eqs. (33) and (34) it can be seen that if

$$a = 3r_* = 3\sqrt{\frac{I_z}{\pi j_0}} = \frac{3cB_0}{2\pi j_0}, \quad (35)$$

then about 90% of the total current flows inside this cylinder. At distances greater than a , the total current and the axial magnetic field with such accuracy are equal to zero, and the azimuthal magnetic field decreases as $\sim 1/r$.

From eqs. (27) and (29) we find the energy stored

inside the force-free magnetic flux rope:

$$E = 2\pi \int_0^a \left(\frac{B_\varphi^2}{8\pi} + \frac{B_z^2}{8\pi} \right) r dr = \frac{I_z^2}{2c^2} \ln \left(1 + \frac{a^2}{r_*^2} \right). \quad (36)$$

It can be seen that the energy is determined by the total axial current and weakly depends on a/r_* .

In conclusion, we have obtained the integral equilibrium equation for a cylindrically symmetric force-free magnetic flux rope. Using this equation, a new local relationship between the axial magnetic field strength and electric current density was found. This differs significantly from the Gold-Hoyle work [12], as well as from subsequent works [15–18] on equilibrium of a force-free magnetic flux rope, which used some phenomenological assumptions. As a result, we have derived a new nonlinear equation (which apparently has never been encountered before in other physical problems either) for the axial current density from the equations of magnetostatics without any phenomenological assumptions, except for the physically natural assumption that the axial magnetic field decreases at infinity faster than $1/r$. We have found an analytical solution to this equation, which made it possible to completely solve the problem of finding a self-consistent equilibrium of the force-free magnetic flux rope. We have obtained exact analytical expressions for the self-consistent components of the electric current density and magnetic field. It is shown that all equilibrium quantities are determined by the total axial current (or the amplitude of the axial magnetic field) and the maximum density of this current. It was established that the found magnetic field qualitatively coincides with the Gold-Hoyle magnetic field. However, unlike the Gold-Hoyle solution, in the solution we obtained, the twist of the magnetic field lines is not an arbitrary value, but is determined by the axial current.

The results obtained may be of interest for studies in the solar atmosphere, solar wind, as well as in other astrophysical and geophysical problems with low-pressure plasma.

-
- [1] Aschwanden M. J., *New Millenium Solar Physics* (Springer, Berlin) 2019.
- [2] Chen J., *Phys. Plasmas*, **24** (2017) 090501.
- [3] Daughton W., Scudder J. and Karimabadi H., *Phys. Plasmas*, **13** (2006) 072101.
- [4] Priest E. R., *Solar Magnetohydrodynamics* (Kluwer Academic Publishers, Dordrecht) 2000.
- [5] Ryutova M., *Physics of Magnetic Flux Tubes* (Springer-Verlag, Berlin) 2015.
- [6] Klein L. W. and Burlaga L. F., *J. Geophys. Res.*, **87** (1982) 613.
- [7] Krimigis S., Sarris E. and Armstrong T., *Trans. AGU*, **57** (1976) 304.
- [8] Woltjer L., *Proc. Nat. Acad. Sci.*, **44** (1958) 489.
- [9] Bothmer V. and R. Schwenn, *J. Geophys. Res.*, **104** (1999) 1.
- [10] Lundquist S., *Phys. Rev.*, **83** (1951) 307.
- [11] Miyamoto K., *Fundamentals of Plasma Physics and Controlled Fusion* (Springer, Berlin) 2005.
- [12] Gold T. and Hoyle F., *MNRAS*, **120** (1960) 89.
- [13] Farrugia C. J., Janoo L. A., Tobert R. B., et al. 1999, in *Solar Wind Nine* eds. S. R. Habbal, R. Esser, J. V. Hollweg, P. A. Isenberg (Woodbury, New York: AIP), *AIP Conf. Proc.*, **471** (1999) 745.
- [14] Allanson O., Wilson F. and Neukirch T., *Phys. Plasmas*, **23** (2016) 092106.

- [15] Marubashi K. and Lepping R. P., *Ann. Geophys.*, **25** (2007) 2453.
- [16] Vandas M. and Romashets E. P., *Astronomy & Astrophysics*, **608** (2017) A118.
- [17] Vandas M and, Romashets E. P., *Astronomy & Astrophysics*, **627** (2019) A90.
- [18] Wang Y., Shen C., Liu R. et al., *J. Geophys. Res.*, **123** (2018) 3238.
- [19] Cheremnykh O. K., Fedun V. and Lashkin V. M., *Phys. Plasmas*, **30** (2023) 124501.
- [20] Alfvén H. and Fälthammar C.-G., *Cosmical Electrodynamics: Fundamental Principles* (Clarendon Press, Oxford) 1965.