

Nonlinear excitations in multi-dimensional nonlocal lattices

Brian Choi*

August 22, 2024

Abstract

We study the formation of breathers in multi-dimensional lattices with nonlocal coupling that decays algebraically. By variational methods, the exact relationship between various parameters (dimension, nonlinearity, nonlocal parameter α) that defines positive excitation thresholds is characterized. At the anti-continuum regime, there exists a family of unique ground states that determines excitation thresholds. We not only characterize the sharp spatial decay of ground states, which varies continuously in α , but also identify the time decay of dispersive waves, which undergoes a discontinuous transition in α .

Keywords. nonlocal dynamics, excitation threshold, localization, breather, dispersive decay

MSC 2020 34A08, 34A12, 37K58, 37K60, 37K40

1 Introduction.

Discrete breathers on nonlinear lattices, solitary waves that are localized in space and periodic in time, have been an active area of research in both experiments and mathematical studies for decades [1, 2]. Our focus is on discrete Hamiltonian systems with long-range interactions (LRI) whose equations of motion provide a variety of dynamics ranging from breathers with slowly decaying tails to nonlinear dispersive decay due to an interplay between linear dispersion (or lattice diffraction) characterized by the nonlocal lattice coupling and nonlinearity. LRI rises naturally in the numerical discretization of fractional differential equations, among many others. Relevant to physical applications and numerical studies are the following non-exhaustive references: [3] (energy transport in biomolecules), [4, 5] (optical waveguide arrays), [6] (nonlinear optical realization of the fractional Schrödinger equation), [7, 8, 9] (quantum systems), [10, 11, 12] (existence and stability of breathers), and [13, 14] (breathers in the Klein-Gordon-type coupled oscillators with global coupling). Among many sub-topics related to breathers and LRI, our interest is in extending the theory of excitation thresholds beyond the nearest-neighbor interaction (NNI) [15, 16] to LRI with a nonlocal generalization of the discrete nonlinear Schrödinger equation (DNLS) as the main model; see also [17, 18, 19, 20, 21, 22] for excitation thresholds of DNLS featuring nonlinear hopping, saturable nonlinearities, and periodic/gap solitons.

To illustrate the existence of excitation thresholds (see (2.9) for a precise definition), consider the focusing DNLS

$$i\dot{u}_n = -\Delta_d u_n - |u_n|^{p-1} u_n \quad (1.1)$$

where $u : \mathbb{Z}^d \times \mathbb{R} \rightarrow \mathbb{C}$, $p > 1$ and $\Delta_d u_n := -2du_n + \sum_{|m-n|=1} u_m$ with $|\cdot|$ being the l^1 metric. Comparing

(1.1) with the NLS $i\partial_t u = -\Delta u - |u|^{p-1} u$, both (1.1) and NLS are invariant under the time-translation and gauge symmetry, which yield energy and mass conservation, respectively. Although the solutions of NLS are invariant under $u(x, t) \mapsto \lambda^{-\frac{2}{p-1}} u(\frac{x}{\lambda}, \frac{t}{\lambda^2})$ for $\lambda > 0$, there is no such scaling symmetry for (1.1) due to the underlying lattice structure.

For $\omega > 0$, $Q(x) \in \mathbb{R}$, substituting $u(x, t) = e^{i\omega t} Q(x)$ into NLS, we obtain a nonlinear eigenvalue equation $\omega Q = \Delta Q + |Q|^{p-1} Q$. Since $\|\lambda^{-\frac{2}{p-1}} Q(\frac{x}{\lambda})\|_{L^2} = \lambda^{\frac{d}{2} - \frac{2}{p-1}} \|Q\|_{L^2}$, if Q is a non-trivial solution for some $\omega > 0$ and $p \neq 1 + \frac{4}{d}$, another non-trivial solution to the eigenvalue equation with $\omega' > 0$ with an arbitrarily small L^2 -norm can be constructed by the scaling symmetry. In other words, the NLS allows a family of breather solutions at any L^2 -norm (zero excitation threshold, or in short, $\nu_0 = 0$) unless $p = 1 + \frac{4}{d}$; if $p = 1 + \frac{4}{d}$, then any initial data with L^2 -norm less than $\|Q\|_{L^2}$, where $Q \in H^1(\mathbb{R}^d)$ is the ground state of NLS, disperse

*Corresponding author. United States Military Academy, choigh@bu.edu

and decay to zero in $L^\infty(\mathbb{R}^d)$ as $t \rightarrow \infty$ [23] and therefore some positive L^2 -norm ($\nu_0 > 0$) is required for localization to persist globally in time.

With the lack of scaling symmetry, the dynamics of lattice systems exhibits distinct behavior relative to their continuum analogue. By variational methods, the existence of excitation thresholds for DNLS was shown and classified in [15], thereby proving the heuristics based on scaling arguments laid out in [16]. More precisely, it was shown that $\nu_0 > 0$ if and only if $p \geq 1 + \frac{4}{d}$. Motivated by the work above, our goal is to investigate the role of LRI in excitation thresholds (Propositions 3.1 and 3.2). For the coupling strength decaying algebraically as $|n|^{-(d+\alpha)}$, it is shown that any power-type nonlinearity is mass-supercritical for $0 < \alpha \ll 1$, and hence $\nu_0 > 0$. Our definition of LRI, adopted from [24], requires the coupling coefficients to be summable, and therefore $\alpha = 0$ is not considered. However Proposition 3.3 considers this singular case under a logarithmic weight where it is shown that $\nu_0 > 0$ for any $p > 1$. On the other hand, when $\alpha \geq 2$, the effect of LRI is replaced by that of NNI, which recovers the result of [15].

Once the criteria of existence/non-existence of ground states have been identified, it is of interest to study the spatial profile of ground states, their stability properties, and a generic dynamical behavior of initial data near ground states as $t \rightarrow \infty$. We revisit the topic of the slowly-decaying tail behavior of ground states [7, 13, 14] induced by LRI and show the fully algebraic nature of the decay of (1.2) via rigorous error analysis including the sharp order of decay. At the anti-continuum regime (Proposition 3.6), the ground states are unique, modulo symmetries, and are minimizers of the Gagliardo-Nirenberg type inequalities (2.8) defined similar to the Weinstein functional [23]. This determines an explicit formula for the excitation thresholds. In Section 4, we numerically confirm the excitation of breather-like solutions for localized initial data with sufficiently large mass. The solutions decompose into a hump of peak intensity and radiation where the radiation component, for any time, decays algebraically, consistent with Proposition 2.2. For sufficiently small mass, the solutions exhibit global scattering and nonlinear dispersive decay. It is shown, by a detailed study on the polylogarithms (see also [25, Appendix]), that $\alpha = 1$ is the threshold nonlocal parameter at which the order of dispersive decay exhibits a discontinuous transition (see Proposition 4.1); interestingly, this dichotomy persists in the continuum limit as suggested in [26, Remark 2]. In general, the non-analyticity of polylogarithms at the origin affects the dispersive properties of (1.2), and thus have consequences on the global dynamical behavior.

An outline of this paper is as follows: basic definitions are given in Section 1.1; Section 2 revises the method of concentration compactness in obtaining breather solutions and confirms its applicability to LRI; the sharp decay of ground states is given in Section 2.2; assuming $J_{|n|} \sim |n|^{-(d+\alpha)}$, a precise relationship between d, p, α and the excitation threshold is given in Section 3; that the ground states near the anti-continuum regime are unique minimizers of the excitation threshold functional is given in Section 3.3; in Section 4, the initial value problem in 1D lattice is given as an application of previous ideas.

1.1 Formulation

The behavior of breathers depends crucially on the nonlocal coupling. Let $J : (\mathbb{N} \cup \{0\})^n \rightarrow [0, \infty)$ where $n \in \{1, \dots, d\}$ and $J(0, \dots, 0) = 0$. For the sake of setup, assume J is radial ($n = 1$) where an example when $n > 1$ is studied in Section 3.2; the non-radial coupling, which could potentially rise from direction-dependent LRI, could have promising applications in photonics and optoelectronics [27], and moreover see [28, 29] for the analyses of NLS-type models with anisotropic diffraction whose numerical discretization gives rise to nonlocal non-radial coupling. Suppose throughout the summability condition and define a non-negative operator

$$\bar{J} := \sum_{m \in \mathbb{Z}^d} J_{|m|} < \infty; \quad Lu_n = \sum_{m \in \mathbb{Z}^d \setminus \{n\}} J_{|n-m|} (u_n - u_m),$$

from which our main model is given by

$$i\dot{u}_n = \kappa Lu_n - |u_n|^{p-1} u_n; \quad -\omega q_n = \kappa Lq_n - |q_n|^{p-1} q_n, \quad (1.2)$$

where the time-independent model is obtained by taking $u_n = e^{i\omega t} q_n$; unless the anti-continuum limit is discussed, assume $\kappa = 1$. The two conserved quantities

$$M[u] := \|u\|_{l^2}^2 = \sum_n |u_n|^2; \quad H[u] := \frac{1}{2} \langle Lu, u \rangle - \frac{\|u\|_{l^{p+1}}^{p+1}}{p+1}$$

are used throughout. By direct estimation, it can be seen that L is bounded on $l^2(\mathbb{Z}^d)$ with an upper bound in operator norm $2\bar{J}$. Equivalently let $\xi \in \mathbb{T}^d = \mathbb{R}^d/(2\pi\mathbb{Z}^d) = (-\pi, \pi]^d$ and define $\sigma(\xi) = 2 \sum_{m \neq 0} J_{|m|} \sin^2\left(\frac{m \cdot \xi}{2}\right)$ where $\widehat{Lu}(\xi) = \sigma(\xi)\widehat{u}(\xi)$ and $\widehat{u}(\xi) = \mathcal{F}[u](\xi) = \sum_{m \in \mathbb{Z}^d} u_m e^{-im \cdot \xi}$.

2 Constrained optimization and ground states.

Define

$$I_\nu = \inf_{q \in l^2(\mathbb{Z}^d)} \{H[q] : M[q] = \nu\}.$$

If $\nu = 0$, then $q = 0$, and therefore assume $\nu > 0$. A ground state at mass ν is defined as a minimizer of I_ν . The purpose of this section is twofold: determine a sufficient condition under which a ground state exists (Proposition 2.1) and the sharp order of decay of ground states (Proposition 2.2).

2.1 Existence of ground states.

Lemma 2.1. *If J is radial, then $-\frac{\nu^{\frac{p+1}{2}}}{p+1} \leq I_\nu \leq 0$.*

Proof. By $l^2 \hookrightarrow l^{p+1}$,

$$H[q] \geq -\frac{\nu^{\frac{p+1}{2}}}{p+1} > -\infty.$$

Let $R > 0$. Define $q_n = A := (\frac{\nu}{R^d})^{\frac{1}{2}}$ if $|n| \leq R$ and $q_n = 0$, otherwise; note that $M[q] \simeq \nu$. The second term in the Hamiltonian vanishes for large R since

$$\sum_n |q_n|^{p+1} \simeq \nu^{\frac{p+1}{2}} R^{-\frac{d(p-1)}{2}} \xrightarrow{R \rightarrow \infty} 0. \quad (2.1)$$

In estimating $\langle Lq, q \rangle$, assume $|n| \leq R$, $|m| > R$ since otherwise, $|q_n - q_m|^2 = 0$. Take $R > 0$, $k_0 \in \mathbb{N}$ sufficiently large such that $k_0 \ll R$ and $(R + k_0)^d - R^d \simeq d k_0 R^{d-1}$.

Consider $\mathcal{N} = \{|m'| > R : \min_{|n'| \leq R} |m' - n'| < k_0\}$. If $m \in \mathcal{N}$, then $J_{|n-m|} |q_n - q_m|^2 \leq \bar{J} \nu R^{-d}$. Since $|B(m; k_0) \cap B(0; R)| \leq |B(m; k_0)| \lesssim k_0^d$ and $|\mathcal{N}| \lesssim k_0 R^{d-1}$, we have

$$\sum_{m \in \mathcal{N}, n \in B(m; k_0)} J_{|n-m|} |q_n - q_m|^2 \lesssim \nu \bar{J} k_0^{d+1} R^{-1}. \quad (2.2)$$

For $n \in B(0; R) \setminus B(m; k_0)$, set $k = |n - m|$. Then,

$$\sum_{m \in \mathcal{N}, n \in B(0; R) \setminus B(m; k_0)} J_{|n-m|} |q_n - q_m|^2 \lesssim |\mathcal{N}| A^2 \sum_{k_0 \leq k \lesssim R} k^{d-1} J_k \lesssim k_0 \nu R^{-1} \sum_{k \geq k_0} k^{d-1} J_k, \quad (2.3)$$

and therefore the RHS of (2.3) is bounded above by any $\epsilon > 0$ by taking k_0, R sufficiently large, provided that $k_0 \ll R$.

Consider $m \in S := B(0; 2R) \setminus (\mathcal{N} \cup B(0; R))$. A similar analysis yields

$$\sum_{m \in S, n \in B(0; R)} J_{|n-m|} |q_n - q_m|^2 \lesssim A^2 \sum_{|m| \leq 2R} \sum_{k \geq k_0} k^{d-1} J_k \lesssim \nu \sum_{k \geq k_0} k^{d-1} J_k = o(1), \quad (2.4)$$

as $k_0 \rightarrow \infty$. Lastly let $|m| > 2R$, $|n| \leq R$. Setting $j = n - m$, the lower bound $|j| > R$ holds. Then the contribution to $\langle Lu, u \rangle$ is at most

$$\nu R^{-d} \sum_{|n| \leq R} \sum_{|m| > 2R} J_{|n-m|} \leq \nu R^{-d} \sum_{|n| \leq R} \sum_{|j| > R} J_{|j|} \lesssim \nu \sum_{|j| > R} J_{|j|} = o(1), \quad (2.5)$$

and hence the claim by (2.1), (2.2), (2.3), (2.4), (2.5). \square

The case when $I_\nu = 0$ can be characterized by the Gagliardo-Nirenberg-type inequality (2.6).

Lemma 2.2. $I_\nu = 0$ if and only if

$$\|q\|_{l^{p+1}}^{p+1} \leq \frac{p+1}{2} \nu^{-\frac{p-1}{2}} \|q\|_{l^2}^{p-1} \langle Lq, q \rangle, \quad \forall q \in l^2(\mathbb{Z}^d). \quad (2.6)$$

Proof. (2.6) is equivalent to

$$M[q] = \nu \Rightarrow H[q] \geq 0. \quad (2.7)$$

If $I_\nu = 0$, then $H[q] \geq I_\nu = 0$ whenever $M[q] = \nu$. Conversely, (2.7) implies $I_\nu \geq 0$, which implies $I_\nu = 0$ by Lemma 2.1. \square

If there exists $\nu > 0$ such that (2.6) holds, then the inequality holds with ν replaced by $0 < \nu' < \nu$. Motivated by this observation, define

$$\nu_0 := \sup\{\nu : (2.6) \text{ holds}\} \geq 0.$$

By continuity, $I_{\nu_0} = 0$. Equivalently define

$$K = \inf_{u \in l^2(\mathbb{Z}^d) \setminus \{0\}} \frac{\|u\|_{l^2}^{p-1} \langle Lu, u \rangle}{\|u\|_{l^{p+1}}^{p+1}} \geq 0. \quad (2.8)$$

Comparing K with the coefficients of (2.6), we have

$$\nu_0 = \left(\frac{p+1}{2} \kappa K \right)^{\frac{2}{p-1}}, \quad (2.9)$$

where $\kappa > 0$ is the coupling coefficient if L were replaced by κL . In particular, (2.9) correctly predicts that there is no excitation threshold in the anti-continuum limit $\kappa \rightarrow 0$ with a unique ground state of mass $\nu > 0$ given by $q_* = \nu^{\frac{1}{2}} \delta$ where δ is the Kronecker delta supported at the origin (modulo translation). For ν sufficiently large, ground states are found from the strict negativity of I_ν as [15, Theorem 2.1].

Proposition 2.1. *Let $I_\nu < 0$ and $\{q^{(k)}\} \subseteq l^2(\mathbb{Z}^d)$ be a minimizing sequence. Then there exist a subsequence $\{q^{(n_k)}\}$, $m_k \in \mathbb{Z}^d$, $\phi_k \in \mathbb{R}$, and $q_* \in l^2(\mathbb{Z}^d)$ with non-negative entries such that $e^{i\phi_k} q^{(n_k)}(\cdot - m_k) \xrightarrow[k \rightarrow \infty]{} q_*$ in $l^2(\mathbb{Z}^d)$, and hence $I_\nu = H[q_*]$. There exists $\omega(\nu) > 0$ such that*

$$-wq_* = Lq_* - q_*^p. \quad (2.10)$$

Proof. Although a straightforward application of the method of concentration compactness is applied, we give a sketch of the proof to address a technical issue rising from LRI; for a detailed proof applied to DNLS, see [30] or the appendix of [15].

There exists $\{q^{(n_k)}\}$ whose qualitative behavior is either of vanishing, dichotomy, or compactness. We rule out the first two from which the existence of strong limit $q_* \in l^2(\mathbb{Z}^d)$ follows. Then the non-negativity follows by observing $\langle Lq, q \rangle \geq \langle L|q|, |q| \rangle$ since $|q_n - q_m| \geq ||q_n| - |q_m||$. Then the Euler-Lagrange equation with $\omega(\nu)$ as the Lagrange multiplier yields (2.10).

By definition of minimizing sequence and I_ν , there exists $\epsilon_k > 0$ such that $\epsilon_k \xrightarrow[k \rightarrow \infty]{} 0$ and, without re-labeling the subsequence indices, $I_\nu = H[q^{(k)}] + \epsilon_k$. By Lemma 2.1 and the interpolation of Lebesgue spaces,

$$0 \geq I_\nu \geq -\frac{\|q^{(k)}\|_{l^{p+1}}^{p+1}}{p+1} + \epsilon_k \geq -\frac{\nu \|q^{(k)}\|_{l^\infty}^{p-1}}{p+1} + \epsilon_k.$$

Vanishing subsequence implies $\|q^{(k)}\|_{l^\infty} \xrightarrow[k \rightarrow \infty]{} 0$, which implies $I_\nu = 0$.

The dichotomy of concentration of mass cannot occur due to the strict subadditivity, $I_\nu < I_{\nu_1} + I_{\nu - \nu_1}$ for $\nu_1 \in (0, \nu)$, which follows straightforwardly from $I_{\theta\nu_1} < \theta I_{\nu_1}$ for $\theta > 1$, $\nu_1 > 0$. Indeed,

$$I_{\theta\nu_1} \leq \inf_{M[q]=\nu_1} \{H[\sqrt{\theta}q]\} = \theta \cdot \inf_{M[q]=\nu_1} \left\{ \langle Lq, q \rangle - \frac{\theta^{\frac{p-1}{2}}}{p+1} \|q\|_{l^{p+1}}^{p+1} \right\} < \theta I_{\nu_1},$$

where the last strict inequality follows from $p, \theta > 1$.

The claim is by contradiction. Let $\nu_1 \in (0, \nu)$. For any $\epsilon > 0$, suppose there exist $\{a^{(k)}\}, \{b^{(k)}\} \subseteq l^2(\mathbb{Z}^d)$ with compact support and $\text{dist}(a^{(k)}, b^{(k)}) \xrightarrow[k \rightarrow \infty]{} 0$ satisfying

$$|M[a^{(k)}] - \nu_1| < \epsilon, |M[b^{(k)}] - (\nu - \nu_1)| < \epsilon, \|q^{(k)} - (a^{(k)} + b^{(k)})\|_{l^2} < \epsilon,$$

for sufficiently large $k \in \mathbb{N}$. We claim $H[q^{(k)}] = H[a^{(k)}] + H[b^{(k)}] + o(1)$ as $k \rightarrow \infty, \epsilon \rightarrow 0$. It suffices to show $\langle La^{(k)}, b^{(k)} \rangle \xrightarrow[k \rightarrow \infty]{} 0$, which follows from

$$\begin{aligned} \sum_n \sum_{m \neq n} J_{|n-m|} (a_n^{(k)} - a_m^{(k)}) b_n^{(k)} &= \sum_n \sum_{m \neq 0} J_{|m|} (a_n^{(k)} - a_{n+m}^{(k)}) b_n^{(k)} \\ &= \sum_{m \neq 0} J_{|m|} \sum_n (a_n^{(k)} - a_{n+m}^{(k)}) b_n^{(k)} \\ &\lesssim \sum_{|m| \geq R_k} J_{|m|} (\nu_1 (\nu - \nu_1))^{\frac{1}{2}} \xrightarrow[k \rightarrow \infty]{} 0, \end{aligned}$$

where $R_k = \text{dist}(a^{(k)}, b^{(k)})$. Then $I_\nu = H[a^{(k)}] + H[b^{(k)}] + o(1)$, which implies $I_\nu \geq I_{\nu_1} + I_{\nu - \nu_1}$ as $k \rightarrow \infty, \epsilon \rightarrow 0$. \square

Remark 2.1. *An immediate consequence of any minimizing sequence being precompact modulo symmetries (translations and gauge) is that the set of ground states at any mass is orbitally stable. Spectral stability in the case of LRI is expected (see [31] for the spectral stability of DNLS), which we leave for further studies.*

Remark 2.2. *Taking the contrapositive of (2.6), if the inequality fails for some $q \in l^2(\mathbb{Z}^d)$ for some $\nu > 0$, then $I_\nu < 0$ by Lemma 2.1, and there exists a minimizer of I_ν by Proposition 2.1. Therefore for all $\nu > \nu_0$, we have $I_\nu < 0$, and conversely, for all $\nu \leq \nu_0, I_\nu = 0$ where $I_{\nu_0} = 0$ is by continuity. The existence of minimizer at ν_0 is unclear.*

2.2 Decay of ground states.

It was claimed in [7, Section 3] that when $d = 1$ and $J_k = k^{-(1+\alpha)}$, the tail behavior of the localized states is exponential if $\alpha > 2$ and algebraic if $1 < \alpha < 2$. Soon after in [13], an analysis on a similar model (Klein-Gordon type with cubic potential and LRI) was reported with numerical evidence that the decay near the origin undergoes an exponential-to-algebraic crossover for $|n| > n_c$ for some transitional lattice site $n_c \in \mathbb{N}$. Here we provide a rigorous error analysis with the sharp order of decay. If $q \in l^2(\mathbb{Z}^d)$ is a ground state satisfying (2.10), then there exists $\omega > 0$ such that

$$q = (\omega + L)^{-1} q^p = K * q^p \quad (2.11)$$

where

$$K_n = [\mathcal{F}^{-1}(\omega + \sigma(\xi))^{-1}]_n = (2\pi)^{-d} \int_{\mathbb{T}^d} \frac{e^{in \cdot \xi}}{\omega + \sigma(\xi)} d\xi. \quad (2.12)$$

For $d = 1, J_k = k^{-(1+\alpha)}$, (2.12) was approximated in [7] for $|n| \gg 1$ by

$$\frac{1}{2\pi} \int_{-\infty}^{\infty} \frac{\cos(n\xi)}{\omega + a\xi^2} d\xi = \frac{e^{-\sqrt{\frac{\omega}{a}}|n|}}{2\sqrt{a\omega}}$$

when $\alpha > 2$, for some $a > 0$, using the properties of the polylogarithm, which yields an exponential decay of q . Note that $\sigma(\xi) \simeq |\xi|^2$ is valid for $|\xi| \ll 1$ by Lemma 3.1, but not for $|\xi| \gtrsim 1$, and therefore an extension of the integral from \mathbb{T}^d to \mathbb{R}^d should be dealt with care.

Proposition 2.2. *Let $q \in l^2(\mathbb{Z}^d)$ satisfy (2.10) with $M[q] = \nu$. For any $0 < \alpha < \infty$, the decay estimate*

$$|q_n| \lesssim \nu^{\frac{\alpha}{2}} \langle n \rangle^{-(d+\alpha)} \quad (2.13)$$

holds, and for $\alpha = \infty, q = O(|n|^{-\beta})$ for any $\beta > 0$ where the implicit constant is independent of β . Furthermore if $\epsilon > 0, \alpha < \infty$, then

$$q \notin O(|n|^{-(d+\alpha+\epsilon)}). \quad (2.14)$$

Proof. By an explicit computation of $\mathcal{F}^{-1}[\sin^2\left(\frac{m \cdot \xi}{2}\right)]_n$, we have

$$s_n := \mathcal{F}^{-1}\sigma_n = \sum_{m \neq 0} J_{|m|} \mathcal{F}^{-1}[\sin^2\left(\frac{m \cdot \xi}{2}\right)]_n = \begin{cases} \frac{1}{2} \sum_{m \neq 0} J_{|m|} & , \text{ if } n = 0 \\ -\frac{1}{2} J_{|n|} & , \text{ if } n \neq 0. \end{cases} \quad (2.15)$$

Assume $\alpha \in (0, \infty)$. To obtain a decay estimate of q from (2.11), an argument similar to the Picard iteration is implemented. Note that the decay of K corresponds to the regularity of $(\omega + \sigma(\xi))^{-1}$, and therefore of $\sigma(\xi)$, which in turn corresponds to the decay of s_n ; the symbols $(\omega + \sigma(\xi))^{-1}$ and $\sigma(\xi)$ are in the same Hölder space since $\sigma \geq 0$ and $\omega > 0$. In short, $K = O(|n|^{-(d+\alpha)})$.

By $l^2(\mathbb{Z}^d) \hookrightarrow l^\infty(\mathbb{Z}^d)$, for any $\epsilon > 0$, there exists $R > 0$ such that $|q_n| \leq \epsilon$ whenever $|n| > R$. For $|n| > 2R$, the convolution in (2.11) is split into $|m| \leq R$ and $|m| > R$ to obtain

$$\begin{aligned} q_n &\leq \|q\|_{l^2}^p \sum_{|m| \leq R} K_{n-m} + \epsilon^{p-1} \sum_{|m| > R} K_{n-m} q_m \\ &\lesssim \|q\|_{l^2}^p \sum_{|m| \leq R} \langle n-m \rangle^{-(d+\alpha)} + \epsilon^{p-1} \sum_{|m| > R} K_{n-m} q_m =: S_1 + S_2. \end{aligned} \quad (2.16)$$

Since $|n-m| \geq \frac{|n|}{2}$, we have

$$S_1 \leq C \langle n \rangle^{-(d+\alpha)} \simeq \|q\|_{l^2}^p R^d \langle n \rangle^{-(d+\alpha)}.$$

For $\delta > 0$, let $C_0(\delta) > 0$ be the best constant that satisfies

$$q_n \leq C_0(\delta + \langle n \rangle^{-(d+\alpha)}). \quad (2.17)$$

Indeed C_0 exists since $\frac{q_n}{\delta + \langle n \rangle^{-(d+\alpha)}} \leq \frac{\|q\|_{l^2}^p}{\delta}$. Define $C_1 > 0$ by

$$\sum_m K_{n-m} \langle m \rangle^{-(d+\alpha)} \leq C_1 \langle n \rangle^{-(d+\alpha)}, \quad (2.18)$$

which can be shown by comparing the series with the corresponding integral. Observe that the α -dependence of C_1 is uniform on $\alpha \in [\alpha_0, \infty)$ for a fixed $\alpha_0 > 0$. By (2.16), (2.17),

$$\begin{aligned} q_n &\leq C \langle n \rangle^{-(d+\alpha)} + C_0 \epsilon^{p-1} \sum_{|m| > R} K_{n-m} (\delta + \langle m \rangle^{-(d+\alpha)}) \\ &\leq (\epsilon^{p-1} \sum_m |K_m|) C_0 \delta + (C + C_0(\epsilon^{p-1} C_1)) \langle n \rangle^{-(d+\alpha)} \\ &\leq (C + \frac{C_0}{2})(\delta + \langle n \rangle^{-(d+\alpha)}), \end{aligned}$$

where $0 < \epsilon \ll 1$ was chosen at the last inequality, implying $C_0 \leq C + \frac{C_0}{2}$, and therefore $C_0 \leq 2C$. By (2.17) and $C > 0$, the estimate (2.13) follows by taking $\delta \rightarrow 0$ in (2.17). If $\alpha = \infty$, a similar analysis as above shows $O(|n|^{-\beta})$ decay where the implicit constant can be chosen independent of β by the remark following (2.18).

Suppose there exists $\epsilon > 0$ satisfying $d + \alpha + \epsilon \notin \mathbb{Z}$ and $\epsilon < (p-1)(d+\alpha)$ such that $q = O(|n|^{-(d+\alpha+\epsilon)})$. By (2.11), this implies $\mathcal{F}^{-1}[\sigma \hat{q}] = s * q = O(|n|^{-(d+\alpha+\epsilon)})$. Let $R \gg 1$ such that $R^{-\frac{\epsilon}{2}} \sum_m |s_m| \ll 1$ and $|q_{m_0}| > \frac{\|q\|_{l^\infty}}{2}$ for some $|m_0| < R$. Let $2R \leq |n| \leq 2^{-\frac{1}{d+\alpha}} R^{\frac{d+\alpha+\epsilon/2}{d+\alpha}}$. Then,

$$\begin{aligned} |n|^{-(d+\alpha+\epsilon)} &\gtrsim |(s * q)_n| \geq \left| \sum_{|m| \leq R} s_{n-m} q_m \right| - \left| \sum_{|m| > R} s_{n-m} q_m \right| \\ &\gtrsim \sum_{|m| \leq R} J_{|n-m|} q_m - \sum_{|m| > R} \frac{|s_{n-m}|}{|m|^{d+\alpha+\epsilon}} \\ &\gtrsim \|q\|_{l^\infty} |n|^{-(d+\alpha)} - (R^{-\frac{\epsilon}{2}} \sum_m |s_m|) R^{-(d+\alpha+\epsilon/2)} \gtrsim |n|^{-(d+\alpha)}, \end{aligned} \quad (2.19)$$

where an explicit computation (2.15) and $q_n \geq 0$ were used in the second inequality. \square

3 Excitation thresholds for LRI.

3.1 Radial algebraic decay.

J is defined to be of order $\alpha \in (0, \infty)$ if $\lim_{k \rightarrow \infty} k^{d+\alpha} J_k = A(d, \alpha) \in (0, \infty)$; J is of order ∞ if $\lim_{k \rightarrow \infty} k^{d+\alpha} J_k = 0$ for any $\alpha < \infty$. Suppose $J_1 > 0$ so that $\sigma(\xi) = 0$ if and only if $\xi = 0$; this assumes that the nearest-neighbor coupling is nontrivial. Observe that if $J_1 = 1$ and $J_k = 0$ whenever $k > 1$, then the center-difference discrete Laplacian is recovered.

The existence of excitation thresholds hinges upon the following lemma, which is a higher-dimensional extension of [32, Appendix A.1]. Let $\langle z \rangle = (|z|^2 + 1)^{\frac{1}{2}}$. In what follows, assume for convenience the leading coefficient of the series given by σ to be 1.

Lemma 3.1. *Given a radial J of order $\alpha \in (0, \infty]$, define*

$$\alpha' = \begin{cases} \frac{\alpha}{2} & , \text{ if } \alpha \in (0, 2) \\ 1 & , \text{ if } \alpha \in [2, \infty]. \end{cases} \quad (3.1)$$

Then,

$$\sigma(\xi) \simeq_{d, \alpha} \delta(\xi) := \begin{cases} |\xi|^\alpha & , \text{ if } \alpha \in (0, 2) \\ |\xi|^2 \langle \log |\xi| \rangle & , \text{ if } \alpha = 2 \\ |\xi|^2 & , \text{ if } \alpha \in (2, \infty]. \end{cases} \quad (3.2)$$

Proof. Assume $\alpha \in (0, \infty)$; the case $\alpha = \infty$ follows similarly. Let $k_0 \in \mathbb{N}$ such that $J_k \simeq k^{-(d+\alpha)}$ whenever $k \geq k_0$. Since \mathbb{T}^d is compact, it suffices to show (3.2) for $|\xi| < \frac{\pi}{k_0}$. For $k = |m|$ and $\cos \psi = \frac{m \cdot \xi}{|m||\xi|}$,

$$\sigma(\xi) = \left(\sum_{1 \leq k \leq \frac{\pi}{|\xi|}} + \sum_{k > \frac{\pi}{|\xi|}} \right) \sum_{|m|=k} J_k \sin^2 \left(\frac{k|\xi| \cos \psi}{2} \right) =: I_1 + I_2.$$

By $J_k \simeq k^{-(d+\alpha)}$, we have

$$I_2 \lesssim \sum_{k > \frac{\pi}{|\xi|}} k^{d-1} J_k \lesssim |\xi|^\alpha \lesssim \delta(\xi).$$

Recalling the one-dimensional result in [32, Appendix A.1],

$$I_1 \lesssim \sum_{k=1}^{\infty} k^{d-1} J_k \sin^2 \left(\frac{k|\xi|}{2} \right) \lesssim \delta(\xi),$$

and hence the upper bound.

Let $e_d = (0, \dots, 0, 1) \in \mathbb{Z}^d$. To show the lower bound, we show

$$\delta(\xi) \lesssim \sigma(|\xi|e_d) \lesssim \sigma(\xi). \quad (3.3)$$

Given $\xi \in \mathbb{T}^d$, $m \in \mathbb{Z}^d \setminus \{0\}$, let $\theta, \psi \in [0, \pi]$ be the angles between m and $|\xi|e_d, \xi$, respectively. Then,

$$\sigma(|\xi|e_d) = \left(\sum_{1 \leq k \leq \frac{\pi}{|\xi|}} + \sum_{k > \frac{\pi}{|\xi|}} \right) \sum_{|m|=k} J_k \sin^2 \left(\frac{k|\xi| \cos \theta}{2} \right) =: A_1 + A_2. \quad (3.4)$$

Observing that $\sin^2(z)$ is an increasing function on $z \in [0, \frac{\pi}{2}]$,

$$A_1 \lesssim \sum_{1 \leq k \leq \frac{\pi}{|\xi|}} k^{d-1} J_k \sin^2 \left(\frac{k|\xi|}{2} \right). \quad (3.5)$$

Since $\sin^2 z \lesssim \sin^2 \frac{z}{2}$ for $z \in [0, \frac{\pi}{2}]$ and $|\{m \in \mathbb{Z}^d : |m| = k, |\cos \psi| \geq \frac{1}{2}\}| \gtrsim k^{d-1}$, we have

$$\begin{aligned} \sigma(\xi) &\geq \sum_{1 \leq k \leq \frac{\pi}{|\xi|}} \sum_{|m|=k} J_k \sin^2 \left(\frac{k|\xi| \cos \psi}{2} \right) \geq \sum_{1 \leq k \leq \frac{\pi}{|\xi|}} \sum_{\{|m|=k, |\cos \psi| \geq \frac{1}{2}\}} J_k \sin^2 \left(\frac{k|\xi|}{4} \right) \\ &\gtrsim \sum_{1 \leq k \leq \frac{\pi}{|\xi|}} k^{d-1} J_k \sin^2 \left(\frac{k|\xi|}{4} \right) \gtrsim \sum_{1 \leq k \leq \frac{\pi}{|\xi|}} k^{d-1} J_k \sin^2 \left(\frac{k|\xi|}{2} \right), \end{aligned} \quad (3.6)$$

and therefore $A_1 \lesssim \sigma(\xi)$ by (3.5).

Since $A_2 \lesssim \sum_{k > \frac{\pi}{|\xi|}} k^{d-1} J_k \lesssim |\xi|^\alpha$, it suffices to show $|\xi|^\alpha \lesssim \text{RHS of (3.5)}$, which follows from

$$\text{RHS of (3.5)} \gtrsim \sum_{k_0 \leq k \leq \frac{\pi}{|\xi|}} \frac{\sin^2 \left(\frac{k|\xi|}{2} \right)}{k^{1+\alpha}} = \sum_{k \geq k_0} \frac{\sin^2 \left(\frac{k|\xi|}{2} \right)}{k^{1+\alpha}} - \sum_{k > \frac{\pi}{|\xi|}} \frac{\sin^2 \left(\frac{k|\xi|}{2} \right)}{k^{1+\alpha}},$$

and

$$|\xi|^\alpha + \sum_{k > \frac{\pi}{|\xi|}} \frac{\sin^2 \left(\frac{k|\xi|}{2} \right)}{k^{1+\alpha}} \lesssim |\xi|^\alpha \lesssim \delta(\xi) \simeq \sum_{k \geq k_0} \frac{\sin^2 \left(\frac{k|\xi|}{2} \right)}{k^{1+\alpha}}, \quad (3.7)$$

where the last step is by [32, Appendix A.1].

To show the other lower bound of (3.3), observe that

$$\sigma(|\xi|e_d) \simeq \sum_{m_d=1}^{\infty} \sum_{m_1, \dots, m_{d-1}} J_{|m|} \sin^2 \left(\frac{m_d \xi_d}{2} \right) \gtrsim \sum_{m_d > k_0} \left(\sum_{m_1, \dots, m_{d-1}} \frac{1}{|m|^{d+\alpha}} \right) \sin^2 \left(\frac{m_d \xi_d}{2} \right). \quad (3.8)$$

A $(d-1)$ -repeated application of an elementary estimate

$$\int_1^{\infty} \frac{dx}{x^p + y} \simeq_p y^{-\frac{p-1}{p}}, \quad y \geq 1, \quad p > 1,$$

yields

$$\begin{aligned} \sum_{m_1, \dots, m_{d-1}} \frac{1}{|m|^{d+\alpha}} &\gtrsim \sum_{m_2, \dots, m_{d-1}} \sum_{m_1 \geq 1} \frac{1}{|m_1|^{d+\alpha} + \dots + |m_d|^{d+\alpha}} \\ &\gtrsim \sum_{m_2, \dots, m_{d-1}} \frac{1}{|m_2|^{d+\alpha-1} + \dots + |m_d|^{d+\alpha-1}} \gtrsim \dots \gtrsim \frac{1}{|m_d|^{1+\alpha}}, \end{aligned}$$

and therefore by [32, Appendix A.1] again,

$$\text{RHS of (3.8)} \gtrsim \sum_{m_d > k_0} \frac{\sin^2 \left(\frac{m_d \xi_d}{2} \right)}{|m_d|^{1+\alpha}} \simeq \delta(\xi).$$

□

Proposition 3.1. *If $p < 1 + \frac{4\alpha'}{d}$, then $K = 0$. Consequently $\nu_0 = 0$, and I_ν has a minimizer for any $\nu > 0$.*

The proposition is shown by an explicit construction of test functions similar to [15, Proposition 4.1]. An extension of this idea to LRI requires more technicality.

Lemma 3.2. *Let $u_n = e^{-\rho|n|^\gamma}$ for $n \in \mathbb{Z}$, $\rho > 0$, $\gamma \in (0, 2)$. Then as $\rho \rightarrow 0$, $\sum_{n \in \mathbb{Z}} |u_n|^q \simeq (q\rho)^{-\frac{1}{\gamma}}$ holds for any $q \in (0, \infty)$. For $\rho \ll 1$,*

$$\int_{-\pi}^{\pi} |\xi|^\beta |\widehat{u}(\xi)|^2 d\xi \simeq \begin{cases} \rho^{\frac{\beta-1}{\gamma}} & , \text{ if } \beta \in (-1, 1 + 2\gamma) \\ \rho^2 |\log \rho| & , \text{ if } \beta = 1 + 2\gamma \\ \rho^2 & , \text{ if } \beta > 1 + 2\gamma. \end{cases} \quad (3.9)$$

Proof. Observe that u_n is the fundamental solution of the fractional heat equation

$$\partial_\rho u = -(-\Delta)^{\frac{\gamma}{2}} u, \quad (\xi, \rho) \in \mathbb{T} \times \mathbb{R} \quad (3.10)$$

on the Fourier side where ρ is the time variable. Let $p_\rho(x)$ be the fundamental solution of (3.10) posed on \mathbb{R} . From the semigroup theory of $\exp(-(-\Delta)^{\frac{\gamma}{2}})$, we have (see [33, Section 2.6], for example)

$$p_\rho(x) \simeq \min(\rho^{-\frac{1}{\gamma}}, \frac{\rho}{|x|^{1+\gamma}}), \quad x \in \mathbb{R}. \quad (3.11)$$

On \mathbb{T} , the fundamental solution \tilde{p}_ρ is periodized from p_ρ , i.e.,

$$\tilde{p}_\rho(\xi) = \sum_{m \in \mathbb{Z}} p_\rho(\xi + 2m\pi) = \sum_{n \in \mathbb{Z}} e^{-\rho|n|^\gamma} \cos n\xi,$$

where the last equality is by the Poisson summation formula. By the relationship above, \tilde{p}_ρ obeys (3.11) for $|\xi| \leq \pi$, and therefore

$$\widehat{u}(\xi) = \tilde{p}_\rho(\xi) \simeq \frac{\rho}{\rho^{1+\frac{1}{\gamma}} + |\xi|^{1+\gamma}}.$$

Setting $\xi = 0$ and replacing ρ by $q\rho$, the desired l^q -norm follows.

Let I denote the integral in (3.9) where the domain is restricted to $[0, \pi]$ by the evenness of the integrand. Let $k_0 \geq 1$ be the least integer satisfying $\frac{\pi}{2^{k_0-1}} \leq \rho^{\frac{1}{\gamma}}$. Decomposing $I = I_1 + I_2 + I_3$ on $[0, \rho^{\frac{1}{\gamma}}] \cup [\rho^{\frac{1}{\gamma}}, \frac{\pi}{2}] \cup [\frac{\pi}{2}, \pi]$, respectively, where

$$\begin{aligned} I_1 &\simeq \rho^{-\frac{2}{\gamma}} \int_0^{\rho^{\frac{1}{\gamma}}} |\xi|^\beta d\xi \simeq \rho^{\frac{\beta-1}{\gamma}}, \\ I_2 &\simeq \rho^2 \sum_{k=1}^{k_0} \int_{\frac{\pi}{2^{k+1}}}^{\frac{\pi}{2^{k-1}}} \frac{2^{-\beta k} d\xi}{(\rho^{1+\frac{1}{\gamma}} + |\xi|^{1+\gamma})^2} \simeq \rho^2 \sum_{k=1}^{k_0} 2^{(1+2\gamma-\beta)k}, \\ I_3 &\simeq \rho^2 \int_{\frac{\pi}{2}}^\pi |\xi|^\beta d\xi \simeq \rho^2, \end{aligned} \quad (3.12)$$

(3.9) follows from the sign consideration of the geometric series in (3.12). \square

Proof of Proposition 3.1. Once $K = 0$ is shown, the rest of the statement follows from Remark 2.2. Let

$$u_n = e^{-\rho|n|} = \prod_{j=1}^d e^{-\rho|n_j|}.$$

For $\rho \ll 1$,

$$\|u\|_{l^2}^{p-1} \simeq \rho^{-\frac{d(p-1)}{2}}, \quad \|u\|_{l^{p+1}}^{p+1} \simeq \rho^{-d}.$$

Let $\alpha \in (0, \infty] \setminus \{2\}$. By Lemma 3.1 and the Plancherel Theorem,

$$\langle Lu, u \rangle \simeq \sum_{i=1}^d \int_{-\pi}^\pi |\xi_i|^{2\alpha'} |f(\xi_i)|^2 d\xi_i \cdot \prod_{j \neq i} \int_{-\pi}^\pi |f(\xi_j)|^2 d\xi_j,$$

where $f(\xi_j)$ is the discrete Fourier transform of $e^{-\rho|n_j|}$. By Lemma 3.2,

$$\int_{-\pi}^\pi |\xi_i|^{2\alpha'} |f(\xi_i)|^2 d\xi_i = O(\rho^{2\alpha'-1}), \quad \rho \rightarrow 0. \quad (3.13)$$

Then by the condition on p ,

$$\frac{\|u\|_{l^2}^{p-1} \langle Lu, u \rangle}{\|u\|_{l^{p+1}}^{p+1}} \lesssim \rho^{-\frac{d(p-1)}{2} + 2\alpha'} \xrightarrow{\rho \rightarrow 0} 0.$$

If $\alpha = 2$, then $\rho(\xi) \lesssim |\xi|^{2-\epsilon}$ for any $\epsilon > 0$. Choose $\epsilon > 0$ such that $p < 1 + \frac{2}{d}(2-\epsilon)$. Then a similar analysis as above yields $K = 0$. \square

Proposition 3.2. *If $p \geq 1 + \frac{4\alpha'}{d}$, then $\nu_0 > 0$. If $\nu > \nu_0$, then $I_\nu < 0$ and has a minimizer. Conversely if $\nu < \nu_0$, then $I_\nu = 0$ and no minimizer exists.*

Proof. Given $\nu_0 > 0$, Remark 2.2 implies the existence of minimizer for any $\nu > \nu_0$ and, conversely, $I_\nu = 0$ for any $\nu < \nu_0$. If $q_* \in l^2(\mathbb{Z}^d)$ were a minimizer of I_ν for $\nu < \nu_0$, then (2.6) yields

$$\|q_*\|_{l^{p+1}}^{p+1} = \frac{p+1}{2} \langle Lq_*, q_* \rangle \leq \frac{p+1}{2} \nu_0^{-\frac{p-1}{2}} \nu^{\frac{p-1}{2}} \langle Lq_*, q_* \rangle, \quad (3.14)$$

implying $\nu_0 \leq \nu$, a contradiction.

To show estimates of the form (2.6) hold, let $\theta \in (0, 1)$ satisfy $\frac{\theta\alpha'}{d} = \frac{1}{2} - \frac{1}{p+1}$. Note by standard algebra that the condition on θ holds if and only if

$$\begin{aligned} & \{\alpha \in [2, \infty], d \in \{1, 2\}\} \text{ or } \{\alpha \in [2, \infty], d = 3, p < \frac{d+2}{d-2}\} \\ & \text{ or } \{\alpha \in [1, 2), d = 1\} \text{ or } \{\alpha \in (0, 2), d > \alpha, p < \frac{d+\alpha}{d-\alpha}\}. \end{aligned} \quad (3.15)$$

By Lemma 3.1,

$$\|\nabla_d |\alpha' u|\|_{l^2}^2 \simeq \int |\xi|^{2\alpha'} |\widehat{u}(\xi)|^2 d\xi \lesssim \int \sigma(\xi) |\widehat{u}(\xi)|^2 d\xi = \langle Lu, u \rangle,$$

where $|\nabla_d| = (-\Delta_d)^{\frac{1}{2}}$. Then by the discrete Gagliardo-Nirenberg inequality, we have

$$\|u\|_{l^{p+1}} \lesssim \|u\|_{l^2}^{1-\theta} \|\nabla_d |\alpha' u|\|_{l^2}^\theta \lesssim \|u\|_{l^2}^{1-\theta} \langle Lu, u \rangle^{\frac{\theta}{2}}. \quad (3.16)$$

Let the parameters satisfy (3.15). Then (3.16) and $p \geq 1 + \frac{4\alpha'}{d}$ imply

$$\|u\|_{l^{p+1}} \lesssim \|u\|_{l^2}^{\frac{p-1}{p+1}} \langle Lu, u \rangle^{\frac{1}{p+1}} \left(\frac{\langle Lu, u \rangle}{\|u\|_{l^2}^2} \right)^{\frac{d(p-1)-4\alpha'}{4\alpha'(p+1)}} \lesssim \|u\|_{l^2}^{\frac{p-1}{p+1}} \langle Lu, u \rangle^{\frac{1}{p+1}},$$

and hence $\nu_0 > 0$. If $p \gg 1$ does not satisfy (3.15), then we can bootstrap from the previous argument by choosing $p_1 < p$ satisfying (3.15), and hence

$$\sum_n |u_n|^{p_1+1} = \|u\|_{l^{p_1+1}}^{p_1+1} \lesssim \|u\|_{l^2}^{p_1-1} \langle Lu, u \rangle. \quad (3.17)$$

Assume $\|u\|_{l^2} = 1$. By $l^2 \hookrightarrow l^\infty$, it follows that $\|u\|_{l^{p_1+1}}^{p_1+1} \lesssim \langle Lu, u \rangle$, and the general case follows from replacing u by $\frac{u}{\|u\|_{l^2}}$. \square

Remark 3.1. *The relationship between ν_0 and the existence of ground states, as stated in Proposition 3.2, is general, and applies to Propositions 3.3 and 3.5.*

At a formal level when $\alpha = 0$, J behaves as $|n|^{-d}$ as $n \rightarrow \infty$, which is not summable. However with a sufficient logarithmic decay at the infinity, the limiting regime of Proposition 3.2 yields positive excitation thresholds for any power nonlinearity. The proof is similar to that in Lemma 3.1 and Proposition 3.2, and thus is presented in Appendix A.

Proposition 3.3. *Suppose there exists $\lim_{k \rightarrow \infty} k^d \log^q(k) J_k \in (0, \infty)$ for $q > 1$. For any $p > 1$, then, $\nu_0 > 0$.*

Proof. Let $k_0 \in \mathbb{N}$ such that $J_k \simeq_{k_0} \frac{1}{k^d \log^q k}$ for any $k \geq k_0$. We claim

$$\sigma(\xi) \simeq_{d,q} \langle \log |\xi| \rangle^{-(q-1)}. \quad (3.18)$$

Let $\alpha \ll 1$ such that $1 + \frac{2\alpha}{d} < p < \frac{d+\alpha}{d-\alpha}$ and $\frac{\theta\alpha}{2d} = \frac{1}{2} - \frac{1}{p+1}$ for some $\theta \in (0, 1)$; the general case when $p \geq \frac{d+\alpha}{d-\alpha}$ follows as the bootstrapping argument (3.17). By (3.16), (3.18), it follows that $\|u\|_{l^{p+1}}^{p+1} \lesssim \|u\|_{l^2}^{p-1} \langle Lu, u \rangle$ for any $u \in l^2(\mathbb{Z}^d)$, and hence $\nu_0 > 0$. It suffices to show (3.18) for $|\xi| \ll 1$.

To obtain the upper bound, let $M \simeq (|\xi|^{\frac{2}{d+2}} |\log |\xi||^{\frac{q-1}{d+2}})^{-1}$. Then,

$$\begin{aligned} \sum_{1 \leq |m| \leq M} J_{|m|} \sin^2 \left(\frac{m \cdot \xi}{2} \right) &\lesssim \bar{J} |\xi|^2 \sum_{k=1}^M k^{d+1} \lesssim |\xi|^2 M^{d+2} \simeq \langle \log |\xi| \rangle^{-(q-1)}, \\ \sum_{|m| > M} J_{|m|} \sin^2 \left(\frac{m \cdot \xi}{2} \right) &\lesssim \sum_{k > M} \frac{1}{k \log^q k} \simeq \frac{1}{(\log M)^{q-1}} \lesssim \langle \log |\xi| \rangle^{-(q-1)}. \end{aligned}$$

To obtain the lower bound, we show

$$\langle \log |\xi| \rangle^{-(q-1)} \lesssim \sigma(|\xi| e_d) \lesssim \sigma(\xi).$$

Define A_1, A_2 as (3.4) and argue as (3.5), (3.6) to obtain $A_1 \lesssim \sigma(\xi)$. Since

$$A_2 = \sum_{k > \frac{\pi}{|\xi|}} \sum_{|m|=k} J_k \sin^2 \left(\frac{k|\xi| \cos \theta}{2} \right) \lesssim \sum_{k > \frac{\pi}{|\xi|}} \frac{1}{k \log^q k} \simeq \langle \log |\xi| \rangle^{-(q-1)},$$

it suffices to show

$$\langle \log |\xi| \rangle^{-(q-1)} \lesssim \sum_{k \geq k_0} \frac{\sin^2 \left(\frac{k|\xi|}{2} \right)}{k \log^q k} \quad (3.19)$$

by an argument similar to (3.7). Since $\sin^2 \left(\frac{k|\xi|}{2} \right) \geq \frac{1}{2}$ whenever $k \in E_j := [\frac{(2j+1)\pi}{|\xi|}, \frac{(2j+3/2)\pi}{|\xi|}]$ for $j \in \mathbb{N} \cup \{0\}$, it follows that

$$\begin{aligned} \text{RHS of (3.19)} &\gtrsim \sum_{j=0}^{\infty} \sum_{k \in E_j} \frac{1}{k \log^q k} \simeq \sum_{j=0}^{\infty} \left(\log \frac{(2j+1)\pi}{|\xi|} \right)^{-(q-1)} - \left(\log \frac{(2j+\frac{3}{2})\pi}{|\xi|} \right)^{-(q-1)} \\ &\gtrsim \sum_{j=0}^{\infty} \frac{1}{(2j+3/2)\pi \log^q \left(\frac{(2j+3/2)\pi}{|\xi|} \right)} \\ &\gtrsim \sum_{j \gg |\xi|^{-1}} \frac{1}{j \log^q j} \simeq \langle \log |\xi| \rangle^{-(q-1)}, \end{aligned} \quad (3.20)$$

where the difference quotient was estimated below by the derivative of $(\log(\cdot))^{-(q-1)}$ in (3.20). To obtain the lower bound, assume $d \geq 2$; $d = 1$ case follows straightforwardly. It can be shown that for any $r, q > 1$, there exists $y_0(r, q) > 1$ such that for any $y \geq y_0$, the estimate

$$\int_y^{\infty} \frac{dx}{x^r \log^q x} \gtrsim_{r,q} \frac{1}{y^{r-1} \log^q y} \quad (3.21)$$

holds. Let $k_1 = \max(k_0, y_0(2, q), \dots, y_0(d, q))$. Then,

$$\begin{aligned} \sigma(|\xi| e_d) &\gtrsim \sum_{m_d > k_1} \left(\sum_{m_1, \dots, m_{d-1}} \frac{1}{|m|^d \log^q |m|} \right) \sin^2 \left(\frac{m_d \xi_d}{2} \right) \\ &\gtrsim \sum_{m_d > k_1} \sum_{m_2, \dots, m_{d-1}} \sum_{|m_1| \gtrsim |m_2| + \dots + |m_d|} \frac{1}{|m_1|^d \log^q |m_1|} \sin^2 \left(\frac{m_d \xi_d}{2} \right) \\ &\gtrsim \sum_{m_d > k_1} \sum_{m_2, \dots, m_{d-1}} \frac{1}{(|m_2|^{d-1} + \dots + |m_d|^{d-1}) \log^q (|m_2| + \dots + |m_d|)} \sin^2 \left(\frac{m_d \xi_d}{2} \right), \end{aligned}$$

where (3.21) was used in the last inequality. A $(d-1)$ repeated applications of (3.21) yields

$$\sigma(|\xi| e_d) \gtrsim \sum_{m_d > k_1} \frac{\sin^2 \left(\frac{m_d \xi_d}{2} \right)}{|m_d| \log^q |m_d|} \gtrsim \langle \log |\xi| \rangle^{-(q-1)},$$

where the last inequality is by an approach similar to (3.20). \square

3.2 Non-radial algebraic decay.

Consider the following specific form of J when $n = d = 2$. For $i = 1, 2$, let $J^{(i)} : \mathbb{N} \cup \{0\} \rightarrow [0, \infty)$ be an interaction kernel of order $\alpha_i \in (0, \infty]$ and let $\alpha_1 \geq \alpha_2$ by symmetry. Suppose $J_0^{(i)} = 0$, $\sum_{k=1}^{\infty} J_k^{(i)} < \infty$. Define

$$J_{k_1, k_2} = J_{k_1}^{(1)} \delta_{k_2} + \delta_{k_1} J_{k_2}^{(2)}.$$

Our motivation is to extend the ideas of Section 3.1 to characterize the ground states of the two-dimensional model given by (1.2) where

$$\begin{aligned} (Lu)_{n,m} &= \sum_{(n', m') \neq (n, m)} J_{|n-n'|, |m-m'|} (u_{n,m} - u_{n', m'}) \\ &= \sum_{n' \neq n} J_{|n-n'|}^{(1)} (u_{n,m} - u_{n', m}) + \sum_{m' \neq m} J_{|m-m'|}^{(2)} (u_{n,m} - u_{n, m'}) =: (L_1 u)_{n,m} + (L_2 u)_{n,m}, \end{aligned}$$

and $L = \mathcal{F}^{-1}(\sigma(\xi_1, \xi_2))\mathcal{F}$ where

$$\sigma(\xi_1, \xi_2) = \sigma_1(\xi_1) + \sigma_2(\xi_2) := \sum_{m \in \mathbb{Z}} J_{|m|}^{(1)} \sin^2\left(\frac{m\xi_1}{2}\right) + J_{|m|}^{(2)} \sin^2\left(\frac{m\xi_2}{2}\right). \quad (3.22)$$

Proposition 3.4. *Let α'_i be defined as (3.1). If $p < 1 + \frac{4\alpha'_1\alpha'_2}{\alpha'_1 + \alpha'_2}$, then I_ν has a minimizer for any $\nu > 0$, or equivalently, $\nu_0 = 0$.*

Proof. As Proposition 3.1, it suffices to show the claim for $\alpha_i \neq 2$ for $i = 1, 2$. Let $\gamma = \frac{\alpha'_2}{\alpha'_1} \in (0, 1]$ and define

$$u_{n,m} = v_n w_m := e^{-\rho|n|} e^{-\rho|m|^\gamma}.$$

A routine computation using Lemmas 3.1 and 3.2 yields

$$\|u\|_{l^2}^{p-1} \simeq \rho^{-\frac{p-1}{2}(1+\frac{1}{\gamma})}, \quad \|u\|_{l^{p+1}}^{p+1} \simeq \rho^{-(1+\frac{1}{\gamma})}, \quad \langle Lu, u \rangle \simeq \rho^{2\alpha'_1 - 1 - \frac{1}{\gamma}},$$

and the claim follows as Proposition 3.1. \square

To show $\nu_0 > 0$ for sufficiently large nonlinearity, we observe that the variational characterization of ground states extends straightforwardly from radial coupling to our consideration in this section. In particular, Lemmas 2.1 and 2.2 and Proposition 2.1 hold. We identify the range of parameters that ensures (2.6) holds for all l^2 -functions.

Proposition 3.5. *If $p \geq 1 + \frac{4\alpha'_1\alpha'_2}{\alpha'_1 + \alpha'_2}$, then $\nu_0 > 0$.*

Since the dispersion relation depends on directions, the multiplier (3.22) is not radial on \mathbb{T}^2 , and to address this issue, the anisotropic modification of the Littlewood-Paley decomposition is reviewed. For $\alpha > 0$, define $D = \mathcal{F}^{-1}(\xi_1^2 + |\xi_2|^\alpha)^{1/2}\mathcal{F}$. Define an even function $\phi \in C_c^\infty((-2, 2); [0, 1])$ where $\phi(z) = 1$ for $z \in [-1, 1]$. Let $\psi(z) := \phi(z) - \phi(2z)$. For $N \in 2^{\mathbb{Z}}$, let

$$\psi_N(\xi_1, \xi_2) = \psi\left(\frac{\sqrt{\xi_1^2 + |\xi_2|^\alpha}}{\pi N}\right); \quad P_N := \mathcal{F}^{-1}\psi_N\mathcal{F}.$$

On lattices, the dyadic decomposition is restricted to $N \leq 1$. The mapping properties of D and P_N were studied in \mathbb{R}^2 in [29, Lemma 3.1, 3.3]. These results can be transferred to \mathbb{Z}^2 with ease.

Lemma 3.3. *Let $1 \leq q_1 \leq q_2 \leq \infty$, $N \leq 1$, and $\alpha > 0$. Then,*

$$\|P_N f\|_{l^{q_2}(\mathbb{Z}^2)} \lesssim N^{(1+\frac{2}{\alpha})(\frac{1}{q_1} - \frac{1}{q_2})} \|P_N f\|_{l^{q_1}(\mathbb{Z}^2)}.$$

Lemma 3.4. *Let $s \in \mathbb{R}$, $r \in [1, \infty]$, $N \leq 1$. Then,*

$$\|D^s P_N f\|_{l^r(\mathbb{Z}^2)} \simeq N^s \|P_N f\|_{l^r(\mathbb{Z}^2)}.$$

As Proposition 3.2, the key idea is to show the Gagliardo-Nirenberg inequality involving the operator D .

Proof of Proposition 3.5. Adopting the method in [34, Proposition 4] along with Lemmas 3.3 and 3.4, the Gagliardo-Nirenberg inequality

$$\|u\|_{l^{p+1}} \lesssim \|u\|_{l^2}^{1-\theta} \|D^s u\|_{l^2}^\theta \quad (3.23)$$

can be shown for $s > 0$, $\theta \in (0, 1)$, $\alpha > 0$ satisfying $s\theta = (1 + \frac{2}{\alpha})(\frac{1}{2} - \frac{1}{p+1})$. Let $\alpha = \frac{2\alpha'_2}{\alpha'_1}$, $s = \alpha'_1$. Then,

$$\|D^s u\|_{l^2}^2 \simeq \iint (|\xi_1|^{2s} + |\xi_2|^{\alpha s}) |\widehat{u}(\xi)|^2 d\xi_1 d\xi_2 \lesssim \langle Lu, u \rangle \quad (3.24)$$

by Lemma 3.1. By (3.23), (3.24),

$$\|u\|_{l^{p+1}}^{p+1} \lesssim \|u\|_{l^2}^{p-1} \langle Lu, u \rangle \left(\frac{\langle Lu, u \rangle}{\|u\|_{l^2}^2} \right)^{\frac{\alpha'_1 + \alpha'_2}{4\alpha'_1 \alpha'_2} \left(p - \left(1 + \frac{4\alpha'_1 \alpha'_2}{\alpha'_1 + \alpha'_2} \right) \right)} \lesssim \|u\|_{l^2}^{p-1} \langle Lu, u \rangle,$$

and hence $\nu_0 > 0$. If $p > 1$ does not satisfy the scaling condition for (3.23), then the claim follows from the bootstrapping argument (3.17) by applying (3.23) to $p_1 \ll p$. \square

3.3 Estimation of excitation thresholds.

An estimation of (2.9) is given at the anti-continuum regime. An explicit computation of the derivatives in κ is needed to estimate thresholds.




Figure 1: Peak intensity versus time for various α (left) and ν (right) with $\kappa = 0.1$.

Proposition 3.6. *Let J be radial and $\nu > 0$. Then there exists $\kappa_0 > 0$ and a smooth map $[0, \kappa_0) \rightarrow l^2(\mathbb{Z}^d)$ given by $\kappa \mapsto q(\kappa)$ such that $\{q(\kappa)\}$ defines a family of unique ground states satisfying $-\omega(\kappa)q = \kappa Lq - q^p$ and $M[q(\kappa)] = \nu$ where*

$$\omega(\kappa; \nu) = \nu^{\frac{p-1}{2}} - \kappa \sum_{m \neq 0} J_{|m|} - \kappa^2 \frac{(p-3)\nu^{-\frac{p-1}{2}}}{2} \sum_{m \neq 0} J_{|m|}^2 + O(\kappa^3). \quad (3.25)$$

Furthermore $q(\kappa)$ is a unique minimizer of (2.8) for any $\kappa \in (0, \kappa_0)$, and therefore

$$\nu_0 = \left(\frac{p+1}{2} J \kappa \right)^{\frac{2}{p-1}} + o(\kappa^{\frac{2}{p-1}}), \quad \kappa \rightarrow 0. \quad (3.26)$$

Proof. Let δ_{n_0} be the Kronecker delta supported at $n_0 \in \mathbb{Z}^d$. Let $\Phi : X \times \mathbb{R}_\kappa \rightarrow X := l^2(\mathbb{Z}^d) \times \mathbb{R}_\omega$ where $(q, \omega, \kappa) \mapsto ((\omega + \kappa L)q - q^p, \|q\|_{l^2}^2 - \nu) =: (\Phi_1, \Phi_2)$. By Proposition 2.1, a ground state has non-negative entries and hence it suffices to consider $l^2(\mathbb{Z}^d)$ as a Hilbert space over the reals. If $\kappa = 0$, the nonlinear eigenvalue equation reduces to $\omega q = q^p$, and therefore $q_n \in \{\omega^{\frac{1}{p-1}}, 0\}$. Since any ground state is necessarily supported at a single site, let $z = (q_*, \omega_*, \kappa_*) := (\nu^{\frac{1}{2}} \delta_{n_0}, \nu^{\frac{p-1}{2}}, 0)$ where we may assume $n_0 = 0$ by the translation symmetry. It can be shown that the Fréchet derivative in X at z , $D\Phi(z)$, is invertible, and therefore the Implicit Function Theorem (IFT) yields the desired (unique) smooth map on $[0, \kappa_0)$. Taking the κ -derivatives of $\Phi(q(\kappa), \omega(\kappa)) = 0$ gives

$$\begin{aligned} (\omega + \kappa L - pq^{p-1})\dot{q} + (\dot{\omega} + L)q &= 0; \quad \langle q, \dot{q} \rangle = 0, \\ (\omega + \kappa L - pq^{p-1})\ddot{q} + (2\dot{\omega} + 2L - p(p-1)[q^{p-2}\dot{q}])\dot{q} + \ddot{\omega}q &= 0; \quad \langle \dot{q}, \dot{q} \rangle + \langle q, \ddot{q} \rangle = 0. \end{aligned}$$

An explicit evaluation at $\kappa = 0$ yields

$$\begin{aligned} \dot{q}(0)_n &= \nu^{\frac{2-p}{2}} J_{|n|}(1 - \delta); \quad \dot{\omega}(0) = - \sum_{m \neq 0} J_{|m|} \\ \ddot{q}(0)_n &= \begin{cases} -\nu^{\frac{3-2p}{2}} \sum_{m \neq 0} J_{|m|}^2 & , n = 0 \\ 2\nu^{\frac{3-2p}{2}} \sum_{m \notin \{0, n\}} J_{|m|} J_{|n-m|} & , n \neq 0 \end{cases}; \quad \ddot{\omega}(0) = (3-p)\nu^{-\frac{p-1}{2}} \sum_{m \neq 0} J_{|m|}^2, \end{aligned}$$

and therefore

$$q(\kappa)_n = \begin{cases} \nu^{\frac{1}{2}} - \frac{\nu^{\frac{3-2p}{2}}}{2} \kappa^2 \sum_{m \neq 0} J_{|m|}^2 + o(\kappa^2) & , n = 0 \\ \nu^{\frac{2-p}{2}} \kappa J_{|n|} + o(\kappa) & , n \neq 0, \end{cases} \quad (3.27)$$

and likewise for ω , which shows (3.25). Note that the error terms of $q(\kappa)$ can be uniformly bounded in n since the convolution terms of $\ddot{q}(0)$ can be uniformly bounded by the Young's inequality. Better approximations could be obtained by taking the higher order Taylor expansions.

To show $q(\kappa)$ is a unique minimizer for $\kappa > 0$, assume $M[v] = \nu$; otherwise consider $\tilde{v} = \frac{\nu^{1/2}}{\|v\|_{l^2}} v$. Since $H[v] \geq H[q(\kappa)]$, we have

$$\frac{\langle \kappa L v, v \rangle}{\|v\|_{l^{p+1}}^{p+1}} \geq \frac{2H[q(\kappa)]}{\|v\|_{l^{p+1}}^{p+1}} + \frac{2}{p+1}. \quad (3.28)$$

Then the RHS of (3.28) $> \frac{\langle \kappa L q(\kappa), q(\kappa) \rangle}{\|q(\kappa)\|_{l^{p+1}}^{p+1}}$, and hence the claim, if and only if $\|v\|_{l^{p+1}} < \|q(\kappa)\|_{l^{p+1}}$ for any $\kappa \in (0, \kappa_0)$ by shrinking κ_0 if necessary.

By contradiction, suppose there exists $\{\kappa_j\} \subseteq (0, \kappa_0)$ such that $\kappa_j \xrightarrow{j \rightarrow \infty} 0$ and $\|v\|_{l^{p+1}} \geq \|q(\kappa_j)\|_{l^{p+1}}$. By (3.27), $\|q(\kappa_j)\|_{l^{p+1}} \xrightarrow{j \rightarrow \infty} \|q(0)\|_{l^{p+1}} = \nu^{\frac{1}{2}}$ strictly from the right, and therefore for any j sufficiently large, we have

$$\nu^{\frac{1}{2}} < \|q(\kappa_j)\|_{l^{p+1}} \leq \|v\|_{l^{p+1}} \leq \|v\|_{l^2} = \nu^{\frac{1}{2}}. \quad \square$$

Remark 3.2. In Figure 1, the dynamics under (1.2) with the coupling $\kappa = 0.1$, $d = 1$, $J_{|n|} = |n|^{-(1+\alpha)}$, $p = 3$ is generated with the initial condition $q(\kappa) = \nu^{\frac{1}{2}} \delta + \frac{\nu^{-\frac{1}{2}} \kappa}{|n|^{1+\alpha}} (1 - \delta)$, i.e., the first-order approximation of $q(\kappa)$ in (3.27). The vanishing Dirichlet boundary condition, $u_n = 0$ for $|n| > 50$, is imposed. Observe in (3.26) that if the zeroth-order approximation of $q(\kappa)$, i.e., $q(\kappa) \approx \nu^{\frac{1}{2}} \delta$, were used, then $\nu_0 \approx 4\kappa\zeta(1 + \alpha)$. Hence if $\alpha \ll 1$, then $\nu_0 \gg 1$ and no ground state can form by Proposition 3.2, which can be seen in the left plot. Similarly the right plot suggests the solution to decay for $\nu \ll \nu_0$ for fixed α .

Remark 3.3. Variational approaches do not ensure uniqueness in general. Uniqueness in Proposition 3.6 is a consequence of IFT, and therefore cannot be generalized to an arbitrary κ . For PDEs, uniqueness of ground states to nonlocal equations have been established for certain models [35, 36]. It is of interest whether similar results could be obtained for nonlocal models in lattices or networks.

4 Applications to the 1D algebraic lattice

For concreteness, let $d = 1$, $J_k = k^{-(1+\alpha)}$ for $\alpha < \infty$. For initial data of small mass below the energy threshold, ground states do not exist by Proposition 3.2. In fact, it was shown in [37] that the solutions to DNLS decay to zero in l^r for $r > 2$. Analogously, nonlinear decay estimates and global scattering are shown in this section by developing linear dispersive estimates and realizing the nonlinear evolution as perturbation. On the other hand, we provide numerical evidence that localized initial data with sufficiently large mass evolve into a localized breather-like component and an algebraically-decaying radiation.




Figure 2: The evolution of a localized initial condition $u(0) = 5\delta_0$ under $\kappa = 0.1$, $p = 3$, and the Dirichlet boundary condition $u_n = 0$, $|n| > 100$. Left: $\log |u_n(t)|$ on $t \in [0, 30]$ for $\alpha = 0.5$. Right: for $\alpha = 1.5$, the linear fit is $y = -2.499x - 1.779$, and for $\alpha = 0.5$, $y = -1.501x - 1.756$. The left plot is a visual illustration of the decomposition into a localized state and radiation. Note that the intensity is not stationary since $u(0)$ is not an exact solution to the nonlinear eigenvalue problem (2.10). The decay of radiation, whose order is $d + \alpha$, is in excellent agreement with Proposition 2.2.

To show linear dispersive estimates, the basics of the polylogarithm functions are revised. The linear flow of (1.2) with $(u_n(0))_n = (\phi_n)_n \in l^2$ is given by

$$u(t) = U(t)\phi = K_t * \phi; (K_t)_n := \frac{1}{2\pi} \int_{-\pi}^{\pi} e^{i(n\xi - \sigma(\xi)t)} d\xi. \quad (4.1)$$

By scaling, let

$$\sigma(\xi) = \sum_{k=1}^{\infty} \frac{\sin^2\left(\frac{k\xi}{2}\right)}{k^{1+\alpha}} = \frac{1}{2}\zeta(1+\alpha) - \frac{1}{4}(\text{Li}_{1+\alpha}(e^{i\xi}) + \text{Li}_{1+\alpha}(e^{-i\xi})),$$

where the polylogarithm $\text{Li}_s(z)$ is a holomorphic function in $s, z \in \mathbb{C}$; see [38] and references therein for more details regarding this special function. The radius of convergence (in z) is 1, inside which

$$\text{Li}_s(z) := \sum_{k=1}^{\infty} \frac{z^k}{k^s}.$$

Outside the radius of convergence, there exists an analytic continuation in $\mathbb{C} \setminus [1, \infty)$ where for $\Re s > 0$, the polylogarithm can be identified with the Bose-Einstein integral

$$\text{Li}_s(z) = \frac{1}{\Gamma(s)} \int_0^{\infty} \frac{t^{s-1}}{e^t/z - 1} dt. \quad (4.2)$$

An important property of polylogarithms pertinent to our interest is

$$\frac{d}{d\xi} \text{Li}_s(e^{\pm i\xi}) = \pm i \text{Li}_{s-1}(e^{\pm i\xi}), \quad s \in \mathbb{C}, \quad \xi \notin 2\pi\mathbb{Z}. \quad (4.3)$$

Although $\sigma \in C(\mathbb{T})$ by the summability of J , it is not analytic at the origin. Hence the derivatives of σ are locally identified with those of the polylogarithms, by uniqueness of analytic continuation, which furnishes the following dispersive estimates.

Proposition 4.1 (Linear dispersive decay). *For any $\alpha \in (0, \infty)$, there exists $C = C(\alpha) > 0$ such that*

$$\|U(t)\phi\|_{l^\infty} \leq C\langle t \rangle^{-\rho} \|\phi\|_{l^1}, \quad (4.4)$$

where $\rho = \rho_+ := \frac{1}{3}$ if $\alpha \in (1, \infty)$ and $\rho = \rho_- := \frac{1}{2}$ if $\alpha \in (0, 1]$.

Remark 4.1. *Our proof, based on the Van der Corput Lemma, which was also used in [37], splits into three cases: i) $\alpha \in (0, 1)$; ii) $\alpha = 1$; iii) $\alpha \in (1, \infty)$. The case $\alpha = 1$ reduces to the well-known dilogarithm. The other cases are based on detailed estimates using the Bose-Einstein integral. For the sake of the overall flow of presentation, the proof is given in Appendix A.*

Let $2 \leq q, r \leq \infty$. Say (q, r) is ρ_+ -admissible if $\frac{3}{q} + \frac{1}{r} \leq \frac{1}{2}$ and ρ_- -admissible if $\frac{2}{q} + \frac{1}{r} \leq \frac{1}{2}$. By applying [39, Theorem 1.2], we obtain

Corollary 4.1 (Strichartz estimate). *For any ρ_+ -admissible pairs (or ρ_- -admissible pairs) $(q, r), (\tilde{q}, \tilde{r})$, we have*

$$\begin{aligned} \|U(t)\phi\|_{L_t^q l^r} &\lesssim \|\phi\|_{l^2}, \\ \left\| \int_0^t U(t-\tau)F(\tau)d\tau \right\|_{L_t^q l^r} &\lesssim \|F\|_{L_t^{\tilde{q}} l^{\tilde{r}}}. \end{aligned} \quad (4.5)$$




Figure 3: Peak intensity versus time with $(\kappa, p) = (1, 3)$. For $\alpha = 1.5$, the linear fit is $y = -0.30555x - 0.67053$, and for $\alpha = 0.5$, $y = -0.45583x - 0.96974$ computed in Matlab. The vanishing Dirichlet boundary condition, $u_n = 0, |n| > 100$ is used. As predicted in Proposition 4.1, the rates of dispersive decay ρ_{\pm} is illustrated by the slopes of the log-log plot. This numerically verifies Propositions 4.2 and 4.3 where small data disperse over the lattice without self-collapse.

Assuming small data, the linear dispersive estimates extend to the nonlinear regime under certain technical hypotheses. The conditions below are most likely not sharp.

Proposition 4.2 (Nonlinear dispersive decay). *Suppose that*

$$p > \begin{cases} 3 & , \alpha \in (0, 1] \\ \max(4, 1 + 4\alpha') & , \alpha \in (1, \infty) \end{cases} ; \quad q \in \begin{cases} \left(\frac{2(p+1)}{p-1}, p+1 \right] & , \alpha \in (0, 1] \\ \left(\frac{2(p+1)}{p-2}, p+1 \right] & , \alpha \in (1, \infty). \end{cases}$$

Then,

$$\|u(t)\|_{l^q} \leq C\langle t \rangle^{-(\rho_{\pm})\frac{(q-2)}{q}} \|\phi\|_{l^{q'}}, \quad (4.6)$$

whenever $\|\phi\|_{l^{q'}} \ll_{\alpha, p, q} 1$ where $C(\alpha, p, q) > 0$ depends on the constants of the Strichartz estimates (4.5).

Proof. The proof is in the spirit of [37, Theorem 7] where an a priori bound is obtained to run a fixed point argument. We only give a sketch of proof for $\alpha \in (0, 1]$ to illustrate the source of our technical hypotheses.

Let $X = \{(u_n(t)) : \|u(t)\|_{l^q} \lesssim \langle t \rangle^{-(\rho_-)\frac{(q-2)}{q}} \|\phi\|_{l^{q'}}\}$ and

$$\Gamma u := U(t)\phi + i \int_0^t U(t-t') (|u(t')|^{p-1}u(t')) dt'. \quad (4.7)$$

To show the existence of a fixed point in X , and therefore $u = \Gamma u$, observe that the interpolation of $U(t)$ for $l^1 \rightarrow l^\infty$ (4.4) and $l^2 \rightarrow l^2$ (unitary action) yields (4.6) with $u(t)$ replaced by $U(t)\phi$ for all $q \in [2, \infty]$. The condition $q \leq p+1$ ensures $\|u(t')\|_{l^{pq'}} \leq \|u(t')\|_{l^q}$, and the lower bound on q yields the estimate $\int_0^t \langle t-t' \rangle^{-\frac{q-2}{2q}} \langle t' \rangle^{-\frac{p(q-2)}{2q}} \lesssim \langle t \rangle^{-\frac{q-2}{2q}}$. Taking $\|\phi\|_{l^{q'}} \ll_{\alpha, p, q} 1$, an a priori estimate

$$\begin{aligned} \|\Gamma u\|_{l^q} &\lesssim \langle t \rangle^{-(\rho_-)\frac{(q-2)}{q}} \|\phi\|_{l^{q'}} + \int_0^t \langle t-t' \rangle^{-(\rho_-)\frac{(q-2)}{q}} \|u(t')\|_{l^{pq'}}^p dt' \\ &\lesssim \langle t \rangle^{-(\rho_-)\frac{(q-2)}{q}} (\|\phi\|_{l^{q'}} + \|\phi\|_{l^{q'}}^p) \lesssim \langle t \rangle^{-(\rho_-)\frac{(q-2)}{q}} \|\phi\|_{l^{q'}}, \end{aligned}$$

holds and the standard contraction argument yields the desired nonlinear decay. \square

A stronger assumption on the power of nonlinearity yields a well-defined wave operator and global scattering for small data.

Proposition 4.3 (Scattering). *Suppose $p \geq 5$ if $\alpha \in (0, 1]$ and $p \geq 7$ if $\alpha \in (1, \infty)$. There exists $\delta > 0$ such that whenever $\|\phi\|_{l^2} < \delta$, there exists a unique asymptotic state $u_\pm \in l^2(\mathbb{Z})$ with $\|u_\pm\|_{l^2} = \|\phi\|_{l^2}$ such that*

$$u(t) - U(t)u_\pm \xrightarrow[t \rightarrow \pm\infty]{} 0 \text{ in } l^2(\mathbb{Z}). \quad (4.8)$$

Furthermore the map $\phi \mapsto u_\pm$ defines a homeomorphism on the δ -neighborhood of the origin.

Proof. The proof is standard in the literature, having developed the linear dispersive estimates. However we illustrate the main ideas for $\alpha \in (0, 1]$ for the sake of completeness. Assume $t \in [0, \infty)$ without loss of generality.

A fixed point argument using $\Gamma : X \rightarrow X$ where Γ is by (4.7) and $X = C(\mathbb{R}; l^2) \cap L_t^{\frac{4p}{3}} l^{\frac{2p}{p-3}}$ yields a global solution $u \in X$ with $\|\phi\|_{l^2} \ll 1$ where the implicit constant depends on α, p and the Strichartz constants. Since

$$\left\| \int_{t_1}^{t_2} U(-t') (|u(t')|^{p-1}u(t')) dt' \right\|_{l^2} \lesssim \|u\|_{L_{t \in [t_1, t_2]}^{4p} l^p}^p \leq \|u\|_{L_{t \in [t_1, t_2]}^{\frac{4p}{3}} l^{\frac{2p}{p-3}}}^p \xrightarrow[t_1, t_2 \rightarrow \infty]{} 0,$$

there exists $u_+ = \lim_{t \rightarrow \infty} U(-t)u(t) \in l^2$ such that (4.8) holds by reasoning as above. Injectivity follows from the uniqueness of limit.

To show surjectivity, let $u_+ \in l^2(\mathbb{Z})$ and define

$$\tilde{\Gamma} u(t) := U(t)u_+ - i \int_t^\infty U(t-t') (|u(t')|^{p-1}u(t')) dt',$$

and let $T \gg 1$ such that $\|U(t)u_+\|_{L_{t \in [T, \infty)}^{\frac{4p}{3}} l^{\frac{2p}{p-3}}} \ll 1$. Another fixed point argument using $\tilde{\Gamma}$ yields $u \in$

$C([T, \infty); l^2) \cap L_t^{\frac{4p}{3}} l^{\frac{2p}{p-3}}$ such that its asymptotic state is u_+ . Then flow $u(T)$ backwards in time to obtain $\phi = u(0)$. Continuity follows from the contractivity of $\Gamma, \tilde{\Gamma}$ and the Strichartz estimates. \square

Acknowledgement

This work was partially supported by the United States Military Academy via the Faculty Research Funds. The author thanks A Aceves for constructive comments.

A Appendix

Proof of Proposition 4.1. Our proof is based on the Van der Corput Lemma (VCL) applied to K_t in (4.1), and hence the higher-order derivatives of σ are estimated from below. By unitarity, we have $\|U(t)\phi\|_{l^\infty} \leq \|\phi\|_{l^1}$, and therefore assume $|t| \geq 1$. By (4.3),

$$\sigma''(\xi) = \frac{1}{4} (\text{Li}_{\alpha-1}(e^{i\xi}) + \text{Li}_{\alpha-1}(e^{-i\xi})); \quad \sigma'''(\xi) = \frac{i}{4} (\text{Li}_{\alpha-2}(e^{i\xi}) - \text{Li}_{\alpha-2}(e^{-i\xi})).$$

For special cases, $\alpha \in \{1, 2\}$, we show by direct computation. Let $\alpha = 1$. Since $\text{Li}_0(z) = \frac{z}{1-z}$ for $z \neq 1$, it follows that $|\sigma''(\xi)| = \frac{1}{4}$ for $\xi \neq 0$. By VCL,

$$2\pi|(K_t)_n| = \left| \int_{-\pi}^{\pi} e^{it(\frac{n}{t}\xi - \sigma(\xi)t)} d\xi \right| \leq 2\delta + \left| \int_{[-\pi, \pi] \setminus (-\delta, \delta)} e^{it(\frac{n}{t}\xi - \sigma(\xi)t)} d\xi \right| \leq 2\delta + C|t|^{-\frac{1}{2}} \leq 2C|t|^{-\frac{1}{2}},$$

where $\delta \leq \frac{C|t|^{-1/2}}{2}$. For $\alpha = 2$ and $z \in \mathbb{C} \setminus [1, \infty)$, it can be shown that $\text{Li}_1(z) = -\log(1-z)$ and

$$\sigma''(\xi) = -\frac{1}{4} (\log(1 - e^{i\xi}) + \log(1 - e^{-i\xi})), \quad \sigma'''(\xi) = -\frac{1}{4} \cot\left(\frac{\xi}{2}\right).$$

Near the origin, $|\sigma'''(\xi)| \gtrsim 1$, and near $\xi = \pm\pi$, we have $\sigma''(\xi) = -\frac{\log 2}{2} + \frac{|k \mp \pi|^2}{16} + O(|k \mp \pi|^3)$ by the Taylor expansion, and hence the decay of $O(|t|^{-\frac{1}{3}})$ by VCL.

For $\alpha > 2$, the integral representation (4.2) is applied. If $|\xi| \in (\frac{\pi}{2}, \pi]$, then

$$|\text{Li}_{\alpha-1}(e^{i\xi}) + \text{Li}_{\alpha-1}(e^{-i\xi})| = \frac{2}{\Gamma(\alpha-1)} \int_0^\infty \frac{t^{\alpha-2}(1 - e^t \cos \xi)}{e^{2t} - 2e^t \cos \xi + 1} dt \geq \frac{1}{2\Gamma(\alpha-1)} \int_0^\infty t^{\alpha-2} e^{-2t} dt = 2^{-\alpha},$$

since $e^{2t} - 2e^t \cos \xi + 1 \leq 4e^{2t}$ and $1 - e^t \cos \xi \geq 1$.

Let $\xi_0 \ll 1$. For $|\xi| \in (0, \xi_0]$, let $t_0 = -\log \cos \xi > 0$. For $t \in [0, t_0]$, the first-order Taylor expansion yields

$$\frac{1 - e^t \cos \xi}{e^{2t} - 2e^t \cos \xi + 1} \leq \frac{(-\cos \xi)t + 1 - \cos \xi}{2 - 2 \cos \xi}$$

to obtain

$$\begin{aligned} \frac{\Gamma(\alpha-1)}{2} |\text{Li}_{\alpha-1}(e^{i\xi}) + \text{Li}_{\alpha-1}(e^{-i\xi})| &\geq \int_{t_0}^\infty \frac{t^{\alpha-2}(e^t \cos \xi - 1)}{e^{2t} - 2e^t \cos \xi + 1} dt - \int_0^{t_0} \frac{t^{\alpha-2}(1 - e^t \cos \xi)}{e^{2t} - 2e^t \cos \xi + 1} dt \\ &\geq \int_{t_0}^\infty \frac{t^{\alpha-2}(e^t \cos \xi - 1)}{2e^{2t}} dt - \int_0^{t_0} \frac{t^{\alpha-2}((- \cos \xi)t + 1 - \cos \xi)}{2 - 2 \cos \xi} dt \\ &= \frac{1}{2} (-\log \cos \xi)^{\alpha-1} \left(\frac{-\cos \xi \cdot \log \cos \xi}{\alpha(1 - \cos \xi)} - \frac{1}{\alpha-1} + \int_1^\infty t^{\alpha-2} ((\cos \xi)^{t+1} - (\cos \xi)^{2t}) dt \right), \end{aligned} \quad (\text{A.1})$$

where (A.1) from its previous step could be assisted with softwares such as Maple or Mathematica. For $t < \epsilon_0 \xi^{-2}$ for some $0 < \epsilon_0 \ll 1$, we have

$$\int_1^\infty t^{\alpha-2} ((\cos \xi)^{t+1} - (\cos \xi)^{2t}) dt \geq \frac{\xi^2}{4} \int_1^{\epsilon_0 \xi^{-2}} t^{\alpha-2} (t-1) dt \gtrsim \frac{\xi^{2-2\alpha}}{\alpha(\alpha-1)},$$

and therefore

$$(\text{A.1}) \gtrsim (-\log \cos \xi)^{\alpha-1} \xi^{2-2\alpha} \gtrsim 1,$$

where it can be shown by direct computation that $\frac{-\cos \xi \cdot \log \cos \xi}{\alpha(1 - \cos \xi)} - \frac{1}{\alpha-1} \geq 0$ for any $0 \leq \xi < \frac{\pi}{2}$.

For $|\xi| \in (\xi_0, \frac{\pi}{2}]$,

$$|\text{Li}_{\alpha-2}(e^{i\xi}) - \text{Li}_{\alpha-2}(e^{-i\xi})| = \frac{2|\sin \xi|}{\Gamma(\alpha-2)} \int_0^\infty \frac{t^{\alpha-3} e^t}{e^{2t} - 2e^t \cos \xi + 1} dt \geq |\sin \xi_0| > 0,$$

by $e^{2t} - 2e^t \cos \xi + 1 \leq 2e^{2t}$. Hence $\max_{\xi \neq 0} (|\sigma''(\xi)|, |\sigma'''(\xi)|) \gtrsim_{\alpha} 1$.

For $\alpha \in (1, 2)$, although (4.2) does not yield a convergent integral for $\sigma'''(\xi)$, the derivative relation (4.3) gives

$$4\sigma'''(\xi) = \frac{d}{d\xi}(\text{Li}_{\alpha-1}(e^{i\xi}) + \text{Li}_{\alpha-1}(e^{-i\xi})) = \frac{1}{\Gamma(\alpha-1)} \int_0^{\infty} t^{\alpha-2} \frac{d}{d\xi} \frac{2(e^t \cos \xi - 1)}{e^{2t} - 2e^t \cos \xi + 1} dt,$$

as long as $\xi \neq 0$. Similarly for $\alpha \in (0, 1)$, $|\sigma''(\xi)|$ can be estimated from below by $\frac{d}{d\xi}(\text{Li}_{\alpha}(e^{i\xi}) - \text{Li}_{\alpha}(e^{-i\xi}))$. Then estimating the integrals as above ($\alpha > 2$), the proposition follows. \square

References

- [1] Y. V. Kartashov, B. A. Malomed, and L. Torner, “Solitons in nonlinear lattices,” *Reviews of Modern Physics*, vol. 83, no. 1, pp. 247–305, 2011.
- [2] P. G. Kevrekidis, “Non-linear waves in lattices: past, present, future,” *IMA Journal of Applied Mathematics*, vol. 76, no. 3, pp. 389–423, 2011.
- [3] S. F. Mingaleev, P. L. Christiansen, Y. B. Gaididei, M. Johansson, and K. Rasmussen, “Models for energy and charge transport and storage in biomolecules,” *Journal of Biological Physics*, vol. 25, pp. 41–63, 1999.
- [4] N. K. Efremidis and D. N. Christodoulides, “Discrete solitons in nonlinear zigzag optical waveguide arrays with tailored diffraction properties,” *Physical Review E*, vol. 65, no. 5, p. 056607, 2002.
- [5] A. Szameit, T. Pertsch, S. Nolte, A. Tünnermann, and F. Lederer, “Long-range interaction in waveguide lattices,” *Physical Review A—Atomic, Molecular, and Optical Physics*, vol. 77, no. 4, p. 043804, 2008.
- [6] S. Liu, Y. Zhang, B. A. Malomed, and E. Karimi, “Experimental realisations of the fractional Schrödinger equation in the temporal domain,” *Nature Communications*, vol. 14, no. 1, p. 222, 2023.
- [7] Y. B. Gaididei, S. Mingaleev, P. L. Christiansen, and K. Rasmussen, “Effects of nonlocal dispersive interactions on self-trapping excitations,” *Physical Review E*, vol. 55, no. 5, p. 6141, 1997.
- [8] A. G. Choudhury and A. R. Chowdhury, “On a long range variant of the discrete NLS equation and algebraic Bethe ansatz,” *Physica Scripta*, vol. 53, no. 2, p. 129, 1996.
- [9] N. Korabel and G. M. Zaslavsky, “Transition to chaos in discrete nonlinear Schrödinger equation with long-range interaction,” *Physica A: Statistical Mechanics and its Applications*, vol. 378, no. 2, pp. 223–237, 2007.
- [10] P. Kevrekidis, “Non-nearest-neighbor interactions in nonlinear dynamical lattices,” *Physics Letters A*, vol. 373, no. 40, pp. 3688–3693, 2009.
- [11] P. Kevrekidis, “The drastic role of beyond-nearest-neighbor interactions on discrete two-dimensional dynamical lattices: a case example,” *Journal of Optics*, vol. 15, no. 4, p. 044013, 2013.
- [12] P. L. Christiansen, Y. B. Gaididei, M. Johansson, K. Rasmussen, V. Mezentsev, and J. J. Rasmussen, “Solitary excitations in discrete two-dimensional nonlinear Schrödinger models with dispersive dipole-dipole interactions,” *Physical Review B*, vol. 57, no. 18, p. 11303, 1998.
- [13] S. Flach, “Breathers on lattices with long range interaction,” *Physical Review E*, vol. 58, no. 4, p. R4116, 1998.
- [14] M. Molerón, C. Chong, A. J. Martínez, M. A. Porter, P. G. Kevrekidis, and C. Daraio, “Nonlinear excitations in magnetic lattices with long-range interactions,” *New Journal of Physics*, vol. 21, no. 6, p. 063032, 2019.
- [15] M. I. Weinstein, “Excitation thresholds for nonlinear localized modes on lattices,” *Nonlinearity*, vol. 12, no. 3, p. 673, 1999.
- [16] S. Flach, K. Kladko, and R. MacKay, “Energy thresholds for discrete breathers in one-, two-, and three-dimensional lattices,” *Physical Review Letters*, vol. 78, no. 7, p. 1207, 1997.
- [17] M. Kastner, “Energy thresholds for discrete breathers,” *Physical Review Letters*, vol. 92, no. 10, p. 104301, 2004.
- [18] J. Cuevas, J. Eilbeck, and N. Karachalios, “Thresholds for breather solutions of the discrete nonlinear Schrödinger equation with saturable and power nonlinearity,” *Discrete and Continuous Dynamical Systems*, vol. 21, no. 2, pp. 445–475, 2008.
- [19] J. Cuevas, N. Karachalios, and F. Palmero, “Energy thresholds for the existence of breather solutions and travelling waves on lattices,” *Applicable Analysis*, vol. 89, no. 9, pp. 1351–1385, 2010.
- [20] N. I. Karachalios, B. Sánchez-Rey, P. G. Kevrekidis, and J. Cuevas, “Breathers for the discrete nonlinear Schrödinger equation with nonlinear hopping,” *Journal of Nonlinear Science*, vol. 23, pp. 205–239, 2013.

- [21] M. Cheng, “Solitary waves of the Schrödinger lattice system with nonlinear hopping,” *Journal of Function Spaces*, no. 1, p. 385649, 2015.
- [22] M. Cheng and A. Pankov, “Gap solitons in periodic Schrödinger lattice system with nonlinear hopping,” *Electronic Journal of Differential Equations*, no. 287, 2016.
- [23] M. I. Weinstein, “Nonlinear Schrödinger equations and sharp interpolation estimates,” *Communications in Mathematical Physics*, vol. 87, no. 4, pp. 567 – 576, 1982.
- [24] V. E. Tarasov, “Continuous limit of discrete systems with long-range interaction,” *Journal of Physics A: Mathematical and General*, vol. 39, no. 48, p. 14895, 2006.
- [25] G. Gori, T. Macrì, and A. Trombettoni, “Modulational instabilities in lattices with power-law hoppings and interactions,” *Physical Review E—Statistical, Nonlinear, and Soft Matter Physics*, vol. 87, no. 3, p. 032905, 2013.
- [26] Y. Hong and C. Yang, “Strong convergence for discrete nonlinear Schrödinger equations in the continuum limit,” *SIAM Journal on Mathematical Analysis*, vol. 51, no. 2, pp. 1297–1320, 2019.
- [27] X. Li, H. Liu, C. Ke, W. Tang, M. Liu, F. Huang, Y. Wu, Z. Wu, and J. Kang, “Review of anisotropic 2d materials: controlled growth, optical anisotropy modulation, and photonic applications,” *Laser & Photonics Reviews*, vol. 15, no. 12, p. 2100322, 2021.
- [28] T. Gou, H. Hajaiej, and A. G. Stefanov, “On the solitary waves for anisotropic nonlinear Schrödinger models on the plane,” *European Journal of Mathematics*, vol. 9, no. 3, p. 55, 2023.
- [29] B. Choi and A. Aceves, “Well-posedness of the mixed-fractional nonlinear Schrödinger equation on \mathbb{R}^2 ,” *Partial Differential Equations in Applied Mathematics*, vol. 6, p. 100406, 2022.
- [30] P.-L. Lions, “The concentration-compactness principle in the calculus of variations. The limit case, Part 1,” *Revista Matemática Iberoamericana*, vol. 1, no. 1, pp. 145–201, 1985.
- [31] A. G. Stefanov, R. M. Ross, and P. G. Kevrekidis, “Ground states in spatially discrete non-linear Schrödinger models,” *Nonlinearity*, vol. 36, no. 8, p. 4053, 2023.
- [32] K. Kirkpatrick, E. Lenzmann, and G. Staffilani, “On the continuum limit for discrete NLS with long-range lattice interactions,” *Communications in Mathematical Physics*, vol. 317, no. 3, pp. 563–591, 2013.
- [33] M. Kwaśnicki, “Ten equivalent definitions of the fractional Laplace operator,” *Fractional Calculus and Applied Analysis*, vol. 20, no. 1, pp. 7–51, 2017.
- [34] Y. Hong and C. Yang, “Uniform Strichartz estimates on the lattice,” *Discrete and Continuous Dynamical Systems*, vol. 39, no. 6, pp. 3239–3264, 2019.
- [35] C. J. Amick and J. F. Toland, “Uniqueness and related analytic properties for the Benjamin-Ono equation—a nonlinear Neumann problem in the plane,” *Acta Math*, vol. 167, no. 1-2, p. 107–126, 1991.
- [36] R. L. Frank, E. Lenzmann, and L. Silvestre, “Uniqueness of radial solutions for the fractional Laplacian,” *Communications on Pure and Applied Mathematics*, vol. 69, no. 9, pp. 1671–1726, 2016.
- [37] A. Stefanov and P. G. Kevrekidis, “Asymptotic behaviour of small solutions for the discrete nonlinear Schrödinger and Klein–Gordon equations,” *Nonlinearity*, vol. 18, no. 4, p. 1841, 2005.
- [38] “NIST Digital Library of Mathematical Functions.” <https://dlmf.nist.gov/25.12>, Release 1.2.1 of 2024-06-15. F. W. J. Olver, A. B. Olde Daalhuis, D. W. Lozier, B. I. Schneider, R. F. Boisvert, C. W. Clark, B. R. Miller, B. V. Saunders, H. S. Cohl, and M. A. McClain, eds.
- [39] M. Keel and T. Tao, “Endpoint Strichartz estimates,” *American Journal of Mathematics*, vol. 120, no. 5, pp. 955–980, 1998.