

Notes on scalar curvature lower bounds of steady gradient Ricci solitons

Shota Hamanaka

September 4, 2024

Abstract

We provide new type of decay estimate for scalar curvatures of steady gradient Ricci solitons. We also give certain upper bound for the diameter of a Riemannian manifold whose ∞ -Bakry–Émery Ricci tensor is bounded by some positive constant from below. For the proofs, we use μ -bubbles introduced by Gromov.

1 Introduction

A triple (M, g, f) consisting of a Riemannian manifold (M, g) and a function f on M is called *steady gradient Ricci soliton* if

$$\text{Ric}_g + \text{Hess}_g f = 0,$$

where Ric_g and $\text{Hess}_g f$ denote respectively the Ricci tensor of g and the Hessian of f with respect to g . Hamilton have proven [9] that

$$R_g + |\nabla f|_g^2 = C_0 \tag{1}$$

for some constant C_0 . Here, R_g denotes the scalar curvature of g . Moreover, by [27, Theorem 1.3], $R_g \geq 0$. Steady Ricci solitons appear as Type II singularity models of Ricci flows with nonnegative curvature operator and positive Ricci curvature (see [8, 9] and [2, Theorem 3.4]). First of all, from [18, Theorem 1.7] and [10, Theorem 2.2], the following holds.

Fact 1.1. *Let (M, g) be a complete steady gradient Ricci soliton. Then $\inf_M R_g = 0$.*

Proof. From (1),

$$\limsup_{d_g(p,x) \rightarrow \infty} \frac{-f(x)}{d_g(p,x)} < +\infty$$

Keywords: Scalar curvature, Ricci flow, Surfaces with prescribed mean curvature
2020 Mathematics Subject Classification: 53C20, 53E20, 53A10.

for some fixed point $p \in M$. Moreover, from [18, Theorem 1.7],

$$\limsup_{r \rightarrow +\infty} \frac{\log \text{Vol}_g(B_g(p, r))}{r} = 0,$$

where $\text{Vol}(B_g(p, r))$ denotes the volume of the geodesic ball $B_g(p, r)$ centered at $p \in M$ of radius $r > 0$ with respect to g . Hence from the traced soliton identity: $R_g + \Delta_g f = 0$ and [10, Theorem 2.2],

$$\inf_M R_g = \inf_M \Delta_g(-f) \leq 0.$$

See [6, 7, 25] for another proofs. \square

Remark 1.1. This has been proven by Fernández-López-García-Río [7] using Omori-Yau maximum principle. Later, Wu [25] has proven this fact as a corollary of certain decay estimate of the potential function. Moreover Chow-Lu [6] have shown that $\inf_M |\text{Ric}_g| = 0$.

Fact 1.1 has some similarities in mean curvature flows. In [17], Ma proved that if M is a complete convex translating soliton in \mathbb{R}^{n+1} , then its infimum of the scalar mean curvature is zero (see [17, Theorem 1 (3)]). For convexity of translating solitons, see also the recent work by Xie-Yu [26]. And, Sun [22] proved that for a (real) two-dimensional complete symplectic (resp. almost-calibrated Lagrangian) translating soliton Σ in \mathbb{C}^2 with quadratic area growth, if its Kähler angle is not too large, then its norm of the mean curvature vector must be zero.

In this note, we provide another proof (using μ -bubbles) of Fact 1.1 under certain much stronger condition. Beyond Fact 1.1, we will also investigate the decay of the scalar curvature by using μ -bubbles. Wu [25, Corollary 1.3] has investigated the order of decay of $\inf_M R_g$. We will give another type of decay estimate of $\inf_M R_g$ in the following our main theorem.

The main result of this note is the following theorem.

Theorem 1.1. *Let $n \leq 7$. Suppose that (M^n, g, f) is an n -dimensional complete non-compact nonparabolic steady gradient Ricci soliton. Assume that*

$$\lim_{d_g(p, x) \rightarrow \infty} G(x) \cdot d_g(p, x) = 0 \tag{2}$$

where $G(\cdot)$ is the minimal positive Green's function with the pole at $p \in M$. Moreover we assume that there is a constant $C \geq 0$ such that

$$\text{Ric}_g(x) \geq -C d_g(p, x)^{-2} \cdot G(x) \tag{3}$$

for all $x \in M$ with $d_g(p, x) \gg 1$. Then there is a positive constant $C = C(n) > 0$ such that

$$\liminf_{x \rightarrow \infty} R_g(x) d_g(x, p) \leq C C_0^{1/2}.$$

Here C_0 denotes the constant in the equation (1).

Recall that a complete Riemannian manifold (M, g) is *nonparabolic* if it admits a positive symmetric Green's function. From Propositions 3.1 and 3.2 below, we can immediately obtain the following corollary.

Corollary 1.1. *Let $n \leq 7$. Suppose that (M^n, g, f) is an n -dimensional complete non-compact nonparabolic steady gradient Ricci soliton with $\text{Ric}_g \geq 0$. Assume that*

$$\lim_{d_g(p,x) \rightarrow \infty} G(x) \cdot d_g(p, x) = 0$$

where $G(\cdot)$ is the minimal positive Green's function with the pole at $p \in M$. Then there is a positive constant $C = C(n) > 0$ such that

$$\liminf_{x \rightarrow \infty} R_g(x) d_g(x, p) \leq CC_0^{1/2}.$$

Here C_0 denotes the constant in the equation (1).

Remark 1.2. Suppose that (M^n, g, f) is an n -dimensional complete non-compact steady gradient Ricci soliton with $\text{Ric}_g \geq 0$. If there is a point $p \in M$ such that $\text{Ric}_g(p) = 0$, then (M, g) is Ricci flat (see [5, Remark 6.58]).

Remark 1.3. Munteanu–Wang have recently proven [19, Lemma 2.4] that C can be taken to be zero in dimension three without assuming that (M, g, f) is a steady gradient Ricci soliton and the assumption (2).

This paper is organized as follows. We prove our main theorem (Theorem 1.1) in Subsection 4.3. Before proving Theorem 1.1, we will provide some propositions in Section 2. The proofs of Propositions 2.1, 2.2 are given respectively in subsections 4.1, 4.2 as a prelude to proving our main theorem. In Section 3, we provide some preliminaries that will be key tools to prove our Propositions 2.1, 2.2 and Theorem 1.1. In Appendix (= Section 5), we will give a (partially) new upper bound of the diameter of a Riemannian manifold whose ∞ -Bakry–Emery Ricci tensor is bounded by some positive constant from below.

Acknowledgements

This work was supported by JSPS KAKENHI Grant Number 24KJ0153. Part of this paper was written during a short stay at Mathematics Münster. The author would be grateful to Prof. Rudolf Zeidler and Ms. Claudia–Maria Rüdiger their hospitality and kind support. The author would like to thank Prof. Keita Kunikawa for notifying him the paper [22].

2 Propositions

In this section and Section 4, we will provide some propositions which are weaker than Fact 1.1 and prove them as a prelude to proving our main theorem (Theorem 1.1).

For a steady gradient Ricci soliton (M, g, f) , from (1) and $R_g \geq 0$ ([27, Theorem 1.3]), it turns out that $C_0^{-1} \inf_M R_g$ (C_0 is the constant in (1)) is a

scale invariant non-negative constant. Here, “scale invariant” means that this quantity is invariant under scaling: $g \mapsto c \cdot g$, where $c > 0$ is some positive constant.

Proposition 2.1. *Let $n \leq 7$. Suppose that (M^n, g, f) is an n -dimensional complete non-compact steady gradient Ricci soliton. Assume that there is a constant $C \geq 0$ and a point $p \in M$ such that*

$$\text{Ric}_g(x) \geq -C d_g(p, x)^{-\alpha} \quad (4)$$

for all $x \in M$ with $d_g(p, x) \gg 1$ for some positive constant $\alpha > 0$. Then

$$C_0^{-1} \inf_M R_g \leq A(n),$$

where

$$A(n) := \frac{\frac{4}{3} + \frac{(n-3)(n+1)}{4(n-1)}}{\frac{7}{3} + \frac{(n-3)(n+1)}{4(n-1)}}.$$

Proposition 2.2. *Let $n \leq 7$. Suppose that (M^n, g, f) is an n -dimensional complete non-compact nonparabolic steady gradient Ricci soliton. Assume that $G(x) \rightarrow 0$ as $d_g(p, x) \rightarrow \infty$ (i.e., for any $\varepsilon > 0$, there is $\delta = \delta(\varepsilon) > 0$ such that $G(x) < \varepsilon$ for all $x \in M$ with $d_g(p, x) \geq \delta$), and there is a constant $C \geq 0$ such that*

$$\text{Ric}_g(x) \geq -C d_g(p, x)^{-2} \cdot G(x) \quad (5)$$

for all $x \in M$ with $d_g(p, x) \gg 1$. Here, $G(\cdot)$ is the minimal positive Green’s function with the pole at $p \in M$. Then $\inf_M R_g = 0$.

Remark 2.1. From Propositions 3.1 and 3.2, the assumption of Proposition 2.2 is satisfied if $\text{Ric}_g \geq 0$.

Remark 2.2. Munteanu–Wang have recently proven the same statement for dimension 3 without assuming that (M, g, f) is a steady gradient Ricci soliton [19, Theorem 3.5]. Their proof is based on their analysis of certain harmonic functions and it relies on the fact that the manifold is three dimensional. Our proof is instead relies on μ -bubbles introduced by Gromov.

A complete non-compact Riemannian manifold is nonparabolic if and only if it has at least one nonparabolic end. (See immediately after [13, Definition 20.5].) Munteanu–Sesum have proven [18, Theorem 1.5] that any steady gradient Ricci soliton has at most one nonparabolic end.

From the recent result of Bamler–Chan–Ma–Zhang [1, Theorem 1.1] and the criterion (6) below, if (M^n, g, f) is a complete steady gradient Ricci soliton with $n \geq 4$, $\text{Ric}_g \geq 0$ and the corresponding Ricci flow $(M, g_t)_{t \in \mathbb{R}}$ has a uniformly bounded Nash entropy (see the condition (1.4) in [1]), which is either

- $(M, g_t)_{t \in \mathbb{R}}$ arises as a singularity model, or
- (M, g) has bounded curvature,

then (M, g) is nonparabolic.

Example 2.1. The Cigar soliton is the unique two dimensional complete steady gradient Ricci soliton with positive Gaussian curvature. And its infimum of the scalar curvature is zero. Moreover, since the Cigar soliton has linear volume growth, it is parabolic (i.e., it is not nonparabolic).

Example 2.2. $(\text{Cigar} \times \mathbb{R}, g_{\text{Cigar}} + g_{\text{eucl}})$ is a three dimensional steady gradient Ricci soliton with nonnegative curvature and its infimum of the scalar curvature is zero. Since the Cigar soliton has linear volume growth, it turns out that $\text{Cigar} \times \mathbb{R}$ has quadratic volume growth (by the coarea formula). Hence $\text{Cigar} \times \mathbb{R}$ is parabolic.

Example 2.3. The n -dimensional Bryant solitons on \mathbb{R}^n ($n \geq 3$) are rotationally symmetric and positive sectional curvature. The volume of geodesic balls $B_r(o)$ grow on the order $r^{\frac{n+1}{2}}$, and the curvature decay is of $O(r^{-1})$. In particular, the n -dimensional Bryant soliton satisfies $\lim_{d(p,x) \rightarrow +\infty} R(x) = 0$, and it is parabolic if $n = 3$ and nonparabolic if $n \geq 4$.

Example 2.4. Lai [11] recently found a family of n -dimensional ($n \geq 3$) steady gradient Ricci soliton which is $\mathbb{Z}_2 \times O(n-1)$ -symmetric but rotationally symmetric with positive curvature operator. Moreover, she [12] has also proven that there exists a $\mathbb{Z}_2 \times O(2)$ -symmetric three dimensional flying wing which is asymptotic to a sector with angle θ for all $\theta \in (0, \pi)$. (Note that $\theta = 0, \pi$ respectively corresponds to the Bryant soliton and $\text{Cigar} \times \mathbb{R}$.) From [11, Corollary 1.5], for such a three-dimensional flying wing (M, g, f) , $\inf_M R_g = 0$.

Question 2.1. Are such three-dimensional flying wings nonparabolic?

3 Preliminaries

3.1 Green's functions on a Riemannian manifold with Ricci lower bound

Let (M^n, g) be a smooth complete non-compact Riemannian manifold. Recall that it is called *nonparabolic* if it admits a positive symmetric Green's function. It is well known that in this case the minimal positive Green's function $G(x, y)$ may be obtained as the limit of the Dirichlet Green's function of a sequence of compact exhaustive domains of the manifold. Then,

$$\Delta_x G(x, y) = -\delta_x(y), \quad G(x, y) = G(y, x) > 0.$$

The first key tool used to prove our main theorems is the following gradient estimate for positive harmonic functions by Li-Wang [14] (see also [13]).

Proposition 3.1 ([14, Lemma 2.1] or [13]). *Let (M^n, g) be a complete Riemannian manifold. Suppose that h is a positive harmonic function defined on the geodesic ball $B_g(p, 2R) \subset M$ of radius $2R$ centered at p and $B_g(p, 2R) \cap \partial M = \emptyset$, and*

$$\text{Ric}_g \geq -(n-1)\rho^2$$

for some constant $\rho \in \mathbb{R}$. Then there is a positive constant $C = C(n) > 0$ such that

$$\frac{|\nabla h|_g^2(x)}{h(x)^2} \leq C(1 + \varepsilon^{-1})R^{-2} + \frac{(4(n-1)^2 + 2\varepsilon)\rho^2}{4 - 2\varepsilon}$$

for all $x \in B_g(p, R)$ and for any $\varepsilon < 2$.

By [24, Corollary of Theorem 2], a complete non-compact Riemannian manifold (M^n, g) with $\text{Ric}_g \geq 0$ is nonparabolic if and only if

$$\int_1^\infty \frac{t}{\text{Vol}_g(B_g(p, t))} dt < +\infty, \quad (6)$$

where $\text{Vol}_g(B_g(p, t))$ is the volume of geodesic ball $B_g(p, t)$ of radius t centered at p with respect to g . The second key tool is the following estimate of the minimal positive Green's function $G(x, y)$ by Li-Yau [15, Theorem 5.2].

Proposition 3.2 ([15, Theorem 5.2]). *Suppose that (M^n, g) is a complete nonparabolic Riemannian manifold with $\text{Ric}_g \geq 0$. Then the minimal positive Green's function $G(x, y)$ satisfies that there is a positive constant $C = C(n) > 0$ such that*

$$C^{-1} \int_{d_g(x, y)}^\infty \frac{t}{\text{Vol}_g(B_g(x, t))} dt \leq G(x, y) \leq C \int_{d_g(x, y)}^\infty \frac{t}{\text{Vol}_g(B_g(x, t))} dt$$

for all $x \neq y$. In particular, $G(x, y) \rightarrow 0$ as $d_g(x, y) \rightarrow \infty$.

3.2 Warped μ -bubbles

Let (X, g) be an oriented connected Riemannian manifold together with a decomposition $\partial X = \partial_- X \sqcup \partial_+ X$, where $\partial_\pm X$ are (non-empty) unions of boundary components. Fix a smooth function $u > 0$ on X and a smooth function h on the interior $\overset{\circ}{X}$. Choose a Caccioppoli set Ω_0 with smooth boundary, which contains an open neighborhood of $\partial_- X$ and is disjoint from $\partial_+ X$. Consider the following functional

$$\mathcal{A}_{u, h}(\Omega) = \int_{\partial^* \Omega} u d\mathcal{H}^{n-1} - \int_X (\chi_\Omega - \chi_{\Omega_0}) h u d\mathcal{H}^n$$

for all Caccioppoli sets Ω with $\Omega \Delta \Omega_0 \Subset \overset{\circ}{X}$. Here, \mathcal{H}^{n-1} denotes the $(n-1)$ -Hausdorff measure with respect to the distance d_g induced from g . This implies that Ω contains an open neighborhood of $\partial_- X$ and is disjoint from $\partial_+ X$. A Caccioppoli set minimizing $\mathcal{A}_{u, h}$ in this class $\mathcal{C}(X)$ is called a *warped μ -bubble*. The existence and regularity of a minimizer of $\mathcal{A}_{u, h}$ was given in [28, Proposition 2.1] and [3, Proposition 12].

Proposition 3.3 ([28, Proposition 2.1] or [3, Proposition 12]). *Let $n \leq 7$. Suppose that $h(x) \rightarrow \pm\infty$ as $x \rightarrow \partial_\mp X$. Then there is a smooth minimizer of $\mathcal{A}_{u, h}$ such that $\Omega \Delta \Omega_0 \Subset \overset{\circ}{X}$.*

The first and second variation formula are given in [3, Lemma 13] and [3, Lemma 14] respectively (see also [21, 4.1, 4.3]). We use the the second variation formula in the form of [4, Theorem 4.3].

Proposition 3.4 (First variation [21, Lemma 4.10]). *Suppose $\Omega \in \mathcal{C}(X)$ is a smooth and let Σ be a connected component of $\partial\Omega \setminus \partial_- X$. For any smooth function ϕ on Σ let V_ϕ be a vector field on X , which vanishes outside a small neighborhood of Σ and agree with $\phi\nu$ on Σ . Here, ν is the outwards pointing unit normal of Σ . Let Φ_t be the flow generated by V_ϕ with $\Phi_0 = \text{id}$. Then*

$$\left. \frac{d}{dt} \right|_{t=0} \mathcal{A}_{u,h}(\Phi_t(\Omega)) = \int_{\Sigma} (Hu + g(\nabla^X u, \nu) - hu)\phi \, d\mathcal{H}^{n-1}.$$

Here, $\nabla^X u \in \mathfrak{X}(X)$ denotes the gradient vector field of u with respect to g and H is the scalar mean curvature of Σ . In particular, a smooth μ -bubble Ω satisfies

$$H = -u^{-1}g(\nabla^X u, \nu) + h \quad (7)$$

along $\partial\Omega$.

Proposition 3.5 (Second variation [4, In the proof of Theorem 4.3]). *Suppose $\Omega \in \mathcal{C}(X)$ is a smooth μ -bubble and let Σ be a connected component of $\partial\Omega \setminus \partial_- X$. For any smooth function $\phi \in C_0^\infty(\Sigma)$ on Σ let V_ϕ be a vector field on X , which vanishes outside a small neighborhood of Σ and agree with $\phi\nu$ on Σ . Here, ν is the outwards pointing unit normal of Σ . Let Φ_t be the flow generated by V_ϕ with $\Phi_0 = \text{id}$. Then*

$$\begin{aligned} \left. \frac{d^2}{dt^2} \right|_{t=0} \mathcal{A}_{u,h}(\Phi_t(\Omega)) &= \int_{\Sigma} \phi^2 (\Delta^X u - \Delta^\Sigma u) - 2\phi^2 u^{-1} g(\nabla^X u, \nu)^2 \, d\mathcal{H}^{n-1} \\ &\quad + \int_{\Sigma} u (|\nabla^\Sigma \phi|_g^2 - (|A_\Sigma|^2 + \text{Ric}_g(\nu, \nu))\phi^2) \, d\mathcal{H}^{n-1} \quad (8) \\ &\quad + \int_{\Sigma} \phi^2 g(\nabla^X u, \nu)h - \phi^2 u g(\nabla^X h, \nu) \, d\mathcal{H}^{n-1}. \end{aligned}$$

Here, $\Delta^X u, \nabla^\Sigma \phi$ and A_Σ denote respectively the Laplacian of u with respect to g , the gradient of ϕ with respect to the induced metric $g|_\Sigma$ and the second fundamental form of Σ with respect to g .

4 Proofs of Propositions 2.1, 2.2 and Theorem 1.1

In this section we prove Propositions 2.1, 2.2 and Theorem 1.1.

We will consider the smooth connected compact Riemannian manifold $(X, g|_X)$ for some $X \subset M$, and warped μ -bubbles on it. Using Proposition 3.3 we can find a warped μ -bubble Ω minimizing

$$\mathcal{A}_{u,h}(\Omega) = \int_{\partial^* \Omega} u \, d\mathcal{H}^{n-1} - \int_X (\chi_\Omega - \chi_{\Omega_0}) hu \, d\mathcal{H}^n$$

for all Caccioppoli sets Ω with $\Omega \Delta \Omega_0 \in \hat{X}$ for some reference Caccioppoli set Ω_0 . And set Σ be a component of $\partial\Omega$ contained in \hat{X} . Then, from Proposition 3.3, Σ is compact and smooth. Moreover, since Ω is a minimizer of $\mathcal{A}_{u,h}$, its second derivative at Ω (i.e., the right hand side of (8)) is nonnegative.

Lemma 4.1 ([4, p. 13]).

$$\begin{aligned} \int_{\Sigma} u |\nabla^{\Sigma} \phi|^2 - \phi^2 \Delta^{\Sigma} u &\leq \frac{4}{3} \int_{\Sigma} |\nabla_{\Sigma}(u^{1/2} \phi)|_g^2 \\ &= \frac{1}{3} \int_{\Sigma} u^{-1} \phi^2 |\nabla_{\Sigma} u|_g^2 + \frac{4}{3} \int_{\Sigma} \phi g(\nabla_{\Sigma} u, \nabla_{\Sigma} \phi) \quad (9) \\ &\quad + \frac{4}{3} \int_{\Sigma} u |\nabla_{\Sigma} \phi|_g^2 \end{aligned}$$

for all $\phi \in C_0^{0,1}(\Sigma)$.

Take $u := e^{\psi f}$ where ψ is some smooth function on X and f is the potential function of the Ricci soliton. Then

$$\begin{aligned} \nabla^X u &= u \nabla^X(\psi f) = u \psi \nabla^X f + u f \nabla^X \psi, \\ \Delta^X u &= u \psi \Delta^X f + 2ug(\nabla^X \psi, \nabla^X f) + u f \Delta^X \psi + u \psi^2 |\nabla^X f|_g^2 + f |\nabla^X \psi|_g^2 \\ &\quad - u \psi R_g + 2ug(\nabla^X \psi, \nabla^X f) + u f \Delta^X \psi + u \psi^2 |\nabla^X f|_g^2 \\ &\quad + u f^2 |\nabla^X \psi|_g^2 + 2u \psi f g(\nabla^X \psi, \nabla^X f). \end{aligned}$$

We have used the traced soliton equation: $R_g + \Delta^X f = 0$ in the last equality. Moreover, from (7),

$$|A_{\Sigma}|^2 \geq \frac{1}{n-1} H^2 = \frac{1}{n-1} |h - u^{-1} g(\nabla^X u, \nu)|^2.$$

Putting these together into (8) and using (9), we obtain that

$$\begin{aligned} 0 &\leq \int_{\Sigma} -\phi^2 u \psi R_g + \phi^2 u \psi^2 |\nabla^X f|_g^2 + \phi^2 u f^2 |\nabla^X \psi|_g^2 + 2\phi^2 u \psi f g(\nabla^X \psi, \nabla^X f) \\ &\quad + \int_{\Sigma} -2\phi^2 u \psi^2 g(\nabla^X f, \nu)^2 - 2\phi^2 u f^2 g(\nabla^X \psi, \nu)^2 \\ &\quad + \int_{\Sigma} -4\phi^2 u \psi f g(\nabla^X f, \nu) \cdot g(\nabla^X \psi, \nu) + \int_{\Sigma} \phi^2 u f \Delta^X \psi + 2\phi^2 u g(\nabla^X \psi, \nabla^X f) \\ &\quad - \int_{\Sigma} \frac{\phi^2 u}{n-1} |h - \psi g(\nabla^X f, \nu) - f g(\nabla^X \psi, \nu)|^2 - \text{Ric}_g(\nu, \nu) \phi^2 u \\ &\quad + \int_{\Sigma} \phi^2 u \psi g(\nabla^X f, \nu) h + \phi^2 u f g(\nabla^X \psi, \nu) h - \int_{\Sigma} \phi^2 u g(\nabla^X h, \nu) \\ &\quad + \frac{1}{3} \int_{\Sigma} \phi^2 u \psi^2 |\nabla^{\Sigma} f|_g^2 + \phi^2 u f^2 |\nabla^{\Sigma} \psi|_g^2 + \frac{2}{3} \int_{\Sigma} \phi^2 u \psi f g(\nabla^{\Sigma} \psi, \nabla^{\Sigma} f) \\ &\quad + \frac{4}{3} \int_{\Sigma} \phi u \psi g(\nabla^{\Sigma} f, \nabla^{\Sigma} \phi) + \phi u f g(\nabla^{\Sigma} \psi, \nabla^{\Sigma} \phi) + \frac{4}{3} \int_{\Sigma} u |\nabla^{\Sigma} \phi|_g^2. \end{aligned}$$

4.1 Proof of Proposition 2.1

Fix a point $p \in M$. Let ρ_0 be a smoothing defined on $\overline{B_g(p, L')} \setminus B(p, L)$ ($0 < L < L'$) of the distance function $d_g(p, \cdot)|_{\overline{B_g(p, L')} \setminus B(p, L)}$ from a fixed point $p \in M$ such that $\rho_0(x) = d(p, x)$ for all $x \in \partial(\overline{B_g(p, L')} \setminus B(p, L))$ and $|\nabla \rho_0|_g \leq 2$.

From the Sard's theorem, we can assume that $\partial(\overline{B_g(p, L')} \setminus B(p, L))$ is smooth compact $(n-1)$ -dimensional submanifold of M . Let X be one of connected components of $\overline{\{x \in M \mid \rho(x) \leq L'\}} \setminus \{x \in M \mid \rho(x) < L\}$. We set the component of $\partial\{x \in M \mid \rho(x) \leq L'\}$ as $\partial_- X$. Take some reference Caccioppoli set Ω_0 with $\partial_- X \subset \Omega_0 \subset X \setminus \partial_+ X$, and $\psi = \phi \equiv 1$ in the above calculation. Then we get

$$\begin{aligned} 0 \leq & - \int_{\Sigma} u R_g + CL^{-\alpha} + \frac{4}{3} \int_{\Sigma} u |\nabla^X f|_g^2 - \frac{n+1}{n-1} u g(\nabla^X f, \nu)^2 \\ & - \frac{1}{n-1} \int_{\Sigma} u (h^2 - (n+1)hg(\nabla^X f, \nu)) + \int_{\Sigma} u |\nabla^X h|_g. \end{aligned}$$

(We have also used the assumption (4) here.)

Using Young's inequality,

$$(n+1)g(\nabla^X f, \nu)h \leq \frac{n+1}{2\varepsilon} g(\nabla^X f, \nu)^2 + \frac{\varepsilon(n+1)}{2} h^2.$$

Taking $\varepsilon = \frac{2}{n+1}\delta$ ($\delta \in (0, 1)$), we get

$$(n+1)g(\nabla^X f, \nu)h \leq \left(\frac{2}{n+1}\right)^2 \delta^{-1} g(\nabla^X f, \nu)^2 + \delta h^2,$$

and the $g(\nabla^X f, \nu)^2$ -terms can be estimated as

$$\begin{aligned} \left(\frac{1}{n-1} \left(\frac{n+1}{2}\right)^2 \delta^{-1} - \frac{n+1}{n-1}\right) g(\nabla^X f, \nu)^2 &= \frac{(n+1-4\delta)(n+1)}{4\delta(n-1)} g(\nabla^X f, \nu)^2 \\ &\leq \frac{(n+1-4\delta)(n+1)}{4\delta(n-1)} |\nabla^X f|_g^2. \end{aligned}$$

Putting them together into the above stability inequality, we obtain that

$$\begin{aligned} 0 \leq & - \int_{\Sigma} u R_g + \left(\frac{4}{3} + \frac{(n+1-4\delta)(n+1)}{4\delta(n-1)}\right) \int_{\Sigma} u |\nabla^X f|_g^2 \\ & - \frac{1-\delta}{n-1} \int_{\Sigma} u h^2 + \int_{\Sigma} u |\nabla^X h|_g. \end{aligned}$$

If $\inf_M R(g) = 0$, then the desired estimate is trivial, hence we assume that $\inf_M R(g) > 0$. In order to obtain a contradiction, we suppose that

$$- \left(\frac{7}{3} + \frac{(n+1-4\delta)(n+1)}{4\delta(n-1)}\right) \inf_M R(g) + \left(\frac{4}{3} + \frac{(n+1-4\delta)(n+1)}{4\delta(n-1)}\right) C_0 < 0. \quad (10)$$

Let L_0 be an arbitrary positive real constant. Since the diameter of (M, g) is infinite, we can take L' so that $L_0 < L' - L$ and neither of the sets $\partial_- X, \partial_+ X$ are non-empty. Let $\rho : X \rightarrow \left(-\frac{L'-L}{2}, \frac{L'-L}{2}\right)$ be a smoothing of the signed distance function of the set $\{x \in M \mid d_g(\partial_- X, x) = d_g(\partial_+ X, x)\}$ with $|\text{Lip } \rho| \leq 2$ and $\rho \equiv \pm \frac{L'-L}{2}$ on $\partial_\pm X$. Hence, in particular,

$$\rho \rightarrow \begin{cases} -\frac{L'-L}{2} & \text{at } \partial_- X, \\ \frac{L'-L}{2} & \text{at } \partial_+ X. \end{cases}$$

Take h as

$$h(x) := -\frac{2\pi A}{L'-L} \tan\left(\frac{\pi}{L'-L}\rho(x)\right)$$

for any real positive constant A (we will take $A = \frac{n-1}{1-\delta}$). Then it satisfies

$$-\frac{1}{A}h^2(x) + |\nabla^X h|(x) \leq 4A \left(\frac{\pi}{L'-L}\right)^2.$$

Thus, taking L and L_0 large enough, we finally obtain that

$$0 \leq \frac{d^2}{dt^2} \Big|_{t=0} \mathcal{A}_{u,h}(\Phi_t(\Omega)) < 0.$$

This is impossible. Therefore our supposition (10) was not correct and hence

$$-\left(\frac{7}{3} + \frac{(n+1-4\delta)(n+1)}{4\delta(n-1)}\right) \inf_M R(g) + \left(\frac{4}{3} + \frac{(n+1-4\delta)(n+1)}{4\delta(n-1)}\right) C_0 \geq 0.$$

Sorting this out, we finally obtain that

$$\frac{\inf_M R(g)}{C_0} \leq A(\delta, n) \quad \text{for all } \delta \in (0, 1),$$

where

$$A(\delta, n) := \frac{\frac{4}{3} + \frac{(n+1-4\delta)(n+1)}{4\delta(n-1)}}{\frac{7}{3} + \frac{(n+1-4\delta)(n+1)}{4\delta(n-1)}} \quad (< 1).$$

And it easily turns out that

- $A(\delta, n)$ is non-increasing with respect to δ , and
- $A(1, n)$ is increasing with respect to n .

Hence $\frac{\inf_M R(g)}{C_0} \leq A(1, n) = A(n)$ and $A(\delta, n) > A(1, n) \geq A(1, 2) = \frac{7}{19}$. \square

4.2 Proof of Proposition 2.2

Take $\phi \equiv 1$ and $\psi(\cdot) := G(p, \cdot)$, where G is the minimal positive Green's function with the pole at p . Take X as a component of the set $\{x \in M \mid a \leq \psi(x) \leq 2a\}$ ($0 < a \leq 1$). By Sard's theorem, we can take a so that X is smooth compact connected manifold with manifold. From the assumption, we can assume that $X \subset M \setminus \overline{B_g(p, L)}$ for sufficiently large $L > 0$, which is determined later. Take some reference Caccioppoli set Ω_0 with $\partial_- X \subset \Omega_0 \subset X \setminus \partial_+ X$. From the identity (1), $|\nabla^X f| \leq C_0^{1/2}$ and

$$|f(x)| \leq C_0^{1/2} d_g(p, x) + |f(p)|. \quad (11)$$

Using these, Proposition 3.1 and (5), we obtain that

$$0 \leq \int_{\Sigma} u\psi (-R_g + Ca\psi) + B(h, |\nabla h|_g)$$

for some positive constant $C > 0$. Here $B(h, |\nabla h|_g)$ is the terms containing h or $|\nabla h|$ and this is explicitly expressed as

$$B(h, |\nabla h|_g) = -\frac{1}{n-1} u h^2 + \frac{n+1}{n-1} (u\psi g(\nabla^X f, \nu)h + ufg(\nabla^X \psi, \nu)h) + u|\nabla^X h|.$$

Now suppose that $\inf_M R_g > 0$. Then, from Proposition 3.1, we can take L large enough so that $C\psi < \frac{1}{4} \inf_M R_g$ on Σ . Then, using Young's inequality, we can estimate the term $B(h, |\nabla h|_g)$ as

$$B(h, |\nabla h|_g) \leq (-A\psi^{-1}h^2 + \psi^{-1}|\nabla h|_g + C_1\psi g(\nabla^X f, \nu)^2 + C_2f^2g(\nabla^X \psi, \nu)^2) u\psi$$

for some positive constants A, C_1 and C_2 . The third and fourth term of the right hand side of the previous inequality are estimated by using Proposition 3.1 and (11) so that these are $< \frac{1}{4} \inf_M R_g u\psi$ on Σ by taking L large enough. Take h as

$$h(x) := \frac{A^{-1}}{1 - a^{-1}\psi(x)} - \frac{A^{-1}}{1 - (2a)^{-1}\psi(x)}.$$

Then

$$\begin{aligned} |\nabla^X h|_g &\leq \frac{(a^{-1} - (2a)^{-1})|\nabla^X \psi|\psi^2 + (a^{-1} - (2a)^{-1})|\nabla^X \psi|(a^{-1}\psi^2 + (2a)^{-1}\psi^2)}{A(1 - a^{-1}\psi)^2(1 - (2a)^{-1}\psi)^2} \\ &\leq \frac{(2a)^{-1} \frac{C}{L} \psi^2}{A(1 - a^{-1}\psi)^2(1 - (2a)^{-1}\psi)^2}. \end{aligned}$$

for some positive constant $C = C(n) > 0$. Here, we have used Proposition 3.1 in the last inequality. Thus, taking L large enough so that $C/L \leq 1$, we finally obtain that

$$|\nabla^X h|_g \leq \frac{(2a)^{-1}\psi^2}{A(1 - a^{-1}\psi)^2(1 - (2a)^{-1}\psi)^2}.$$

On the other hand, since $a \leq 1$,

$$-Ah^2 = -\frac{(2a)^{-2}\psi^2}{A(1-a^{-1}\psi)^2(1-(2a)^{-1}\psi)^2} \leq -\frac{(2a)^{-1}\psi^2}{A(1-a^{-1}\psi)^2(1-(2a)^{-1}\psi)^2}.$$

Therefore $-A\psi^{-1}h^2 + \psi^{-1}|\nabla^X h|_g \leq 0$. As a result, we come up with the conclusion that

$$0 \leq \frac{d^2}{dt^2} \Big|_{t=0} \mathcal{A}_{u,h}(\Phi_t(\Omega)) < 0,$$

and conclude the proof by contradiction. \square

4.3 Proof of Theorem 1.1

Finally, we give a proof of Theorem 1.1.

Take Ω_0, ϕ and ψ in the same way as in the proof of Proposition 2.2. Then we obtain that

$$\begin{aligned} 0 \leq & \int_{\Sigma} u\psi d_g(p, x)^{-1} (-R_g d_g(p, x) + 2\psi^{-1} d_g(p, x) \cdot g(\nabla^X \psi, \nabla^X f)) \\ & + \int_{\Sigma} u\psi d_g(p, x)^{-1} d_g(p, x) (\text{other terms}). \end{aligned}$$

Here, by taking L large enough and using (2) and (3), $d_g(p, x)$ (other terms) can be arbitrary small as in the proof of Proposition 2.2. From Proposition 3.1 and (1), the second term can be estimated as

$$2\psi^{-1} d_g(p, x) \cdot g(\nabla^X \psi, \nabla^X f) \leq CC_0^{1/2}$$

for some positive constant $0 < C = C(n)$. Therefore, if we assume that

$$\liminf_{x \rightarrow \infty} R_g d_g(p, x) > CC_0^{1/2},$$

then we obtain that

$$0 \leq \frac{d^2}{dt^2} \Big|_{t=0} \mathcal{A}_{u,h}(\Phi_t(\Omega)) < 0.$$

This is a contradiction. \square

5 Appendix

Using warped μ -bubbles, we can also prove the following diameter upper bounds.

Proposition 5.1. *Let $n \leq 7$ and (M^n, g, f) be a tuple that consists of an n -dimensional complete Riemannian manifold (M, g) and a function on M satisfying*

$$\text{Ric}_g + \text{Hess}_g f \geq \frac{\lambda}{2} g \tag{12}$$

for some positive constant λ . Assume that $\sup_M |\nabla f|_g < \infty$. Then M is compact and

$$\text{diam}(M, g) \leq \pi \sqrt{\frac{2(n-1)}{\lambda A}} \cdot \frac{A(1+A)^{1/4}}{((2+A)\sqrt{1+A} - 2(A+1))^{1/2}}, \quad (13)$$

where

$$A := \frac{(n-1)\lambda}{2 \sup_M |\nabla f|_g^2}.$$

Remark 5.1. The right hand side of the estimate diverges to $+\infty$ as $A \rightarrow 0$ and converges to $\pi \sqrt{\frac{2(n-1)}{\lambda}}$ as $A \rightarrow +\infty$ respectively. In particular, the case of $A = +\infty$ (which corresponds to the case of $f \equiv \text{const}$) recovers Myers's theorem [20].

Proof. Firstly, note that the 2nd variation of \mathcal{A} can also be written as the following alternative form (see [4, Proof of Theorem 4.3]):

$$\begin{aligned} 0 \leq \frac{d^2}{dt^2} \mathcal{A}(\Omega_t) \Big|_{t=0} &= \int_{\Sigma} \psi^2 \text{Hess}_g u(\nu, \nu) - \psi^2 g(\nabla_{\Sigma} u, \nabla_{\Sigma} \psi) \\ &\quad + \int_{\Sigma} \psi^2 g(\nabla_M u, \nu) H \\ &\quad - \int_{\Sigma} u (\psi \Delta_{\Sigma} \psi + (|A_{\Sigma}|^2 + \text{Ric}(\nu, \nu)) \psi^2) \\ &\quad - \int_{\Sigma} \psi^2 g(\nabla_M u, \nu) h + \psi^2 u g(\nabla_M h, \nu). \end{aligned}$$

We will prove the proposition by contradiction. Suppose that

$$\text{diam}(M, g) > (1 + \tilde{\delta}) \sqrt{\frac{2(n-1)(1+a)^2 \pi^2}{\lambda \varepsilon_{\delta} (1-\delta)}} =: L$$

for fixed arbitrary constants $\tilde{\delta}, a > 0$. Here, we set

$$\varepsilon_{\delta} := \left(1 + \frac{(n-1)\lambda \tilde{\delta}}{2 \sup_M |\nabla f|_g^2} \right)^{-1} \frac{(n-1)\lambda \tilde{\delta}}{2 \sup_M |\nabla f|_g^2}.$$

Then there exist two points $p, q \in M$ such that $d_g(p, q) = L$. Consider the set $M_{p,q} := \{x \in M \mid d_g(x, p) = d_g(x, q)\}$, and take $X \subset M$ as one of the components of the set $\{x \in M \mid d_g(M_{p,q}, x) \leq L/2\}$. (More precisely, in order to construct such an X , we need to take a smoothing of the distance function and perturb the boundaries by using Sard's theorem (see the proof of Proposition 2.1). But, since we can perform such smoothing with arbitrary small error, we can still discuss without loss of generality.) By Proposition 3.3, we can find a warped μ -bubble Ω of the functional $\mathcal{A}_{u,h}$ with the choice $u = e^{-f}$ and

$\phi = \text{const}$ and some reference Caccioppoli set Ω_0 . Set $\Sigma := \partial\Omega \setminus \partial_- X$. Then, by Proposition 3.5, (8) and our assumption (12), we obtain that

$$\begin{aligned} 0 &\leq \int_{\Sigma} u \left[\psi^2 |\nabla^M h| - \frac{\psi^2}{n-1} |A_{\Sigma}|^2 - \frac{\lambda}{2} \psi^2 \right] \\ &\leq \int_{\Sigma} u \left[\psi^2 |\nabla^M h| - \frac{\psi^2}{n-1} (h^2 + 2g(\nabla^M f, \nu)h + g(\nabla^M f, \nu)^2) - \frac{\lambda}{2} \psi^2 \right]. \end{aligned}$$

Using Young's inequality, we obtain that

$$2g(\nabla^M f, \nu)h \leq \frac{1}{1-\varepsilon} g(\nabla^M f, \nu)^2 + (1-\varepsilon)h^2$$

for all $\varepsilon \in (0, 1)$. So,

$$0 \leq \int_{\Sigma} u \left[\psi^2 \left(|\nabla^M h| - \frac{\varepsilon}{n-1} h^2 \right) + \frac{\psi^2}{n-1} \cdot \frac{\varepsilon}{1-\varepsilon} g(\nabla^M f, \nu)^2 - \frac{\lambda}{2} \psi^2 \right].$$

Thus we take h as

$$h(x) := -\frac{(n-1)(1+a)\pi}{\varepsilon L} \tan\left(\frac{\pi}{L}\rho(x)\right),$$

where $\rho(x)$ is a smoothing of the signed distance function from $M_{p,q}$ with $|\text{Lip } \rho| \leq 1+a$. Then it holds that

$$|\nabla^M h| - \frac{\varepsilon}{n-1} h^2 \leq \frac{(n-1)(1+a)^2 \pi^2}{\varepsilon L^2}.$$

Then, we get

$$0 \leq \int_{\Sigma} u \psi^2 \left[\frac{(n-1)(1+a)^2 \pi^2}{\varepsilon L^2} + \frac{1}{n-1} \cdot \frac{\varepsilon}{1-\varepsilon} \sup_M |\nabla^M f|^2 - \frac{\lambda}{2} \right].$$

For $\delta \in (0, 1)$, take ε_{δ} so that

$$\frac{1}{n-1} \cdot \frac{\varepsilon_{\delta}}{1-\varepsilon_{\delta}} \sup_M |\nabla^M f|^2 = \frac{\lambda}{2} \delta,$$

equivalently,

$$\varepsilon_{\delta} = \left(1 + \frac{(n-1)\lambda\delta}{2 \sup_M |\nabla^M f|^2} \right)^{-1} \frac{(n-1)\lambda\delta}{2 \sup_M |\nabla^M f|^2}.$$

Then it satisfies that

$$\frac{1}{n-1} \cdot \frac{\varepsilon_{\delta}}{1-\varepsilon_{\delta}} \sup_M |\nabla^M f|^2 - \frac{\lambda}{2} = -\frac{\lambda}{2}(1-\delta).$$

Summing up with this and the definition of L , we get a contradiction:

$$0 \leq \int_{\Sigma} u \psi^2 \left[\frac{(n-1)(1+a)^2 \pi^2}{\varepsilon} \cdot \frac{1}{L^2} + \frac{1}{n-1} \cdot \frac{\varepsilon_{\delta}}{1-\varepsilon_{\delta}} \sup_M |\nabla^M f|^2 - \frac{\lambda}{2} \right] < 0.$$

Hence we obtain that

$$\text{diam}(M, g) \leq (1 + \tilde{\delta}) \sqrt{\frac{2(n-1)(1+a)^2\pi^2}{\lambda\varepsilon_\delta(1-\delta)}}$$

for all $\tilde{\delta} > 0$ and $a > 0$ by contradiction. Therefore we finally reach the conclusion that

$$\text{diam}(M, g) \leq \sqrt{\frac{(n-1)\pi^2}{\lambda\varepsilon_\delta(1-\delta)}} \quad \text{for all } \delta \in (0, 1).$$

In particular, M is compact.

The right hand side of the previous estimate can be written as

$$\pi \sqrt{\frac{2(n-1)}{\lambda A}} \sqrt{\frac{1+A\delta}{\delta(1-\delta)}} =: \pi \sqrt{\frac{2(n-1)}{\lambda A}} \cdot F(\delta).$$

An easy calculation implies that $F(\delta)$ attains its minimum

$$\frac{A(1+A)^{1/4}}{((2+A)\sqrt{1+A} - 2(A+1))^{1/2}}$$

at $\delta = -A^{-1} + A^{-1}\sqrt{1+A} \in (0, 1)$. Hence we obtain the desired estimate. \square

Remark 5.2. This estimate (13) is not better than that of [16, (1.4)] and [23, Theorem 2.1]:

$$\text{diam}(M, g) \leq 4\lambda^{-1} \max_M |\nabla f|_g + \lambda^{-1} \sqrt{16 \max_M |\nabla f|_g^2 + 2(n-1)\lambda\pi^2} \quad (14)$$

when $\max_M |\nabla f|_g$ is small enough. Indeed,

$$\frac{\text{(RHS) of (13)}}{\text{(RHS) of (14)}} = \frac{(n-1)\pi\lambda}{\left((n-1)^2\lambda^2\pi^2 - 2(n-1)\lambda\pi^2\sqrt{2(n-1)\lambda B} + O(B^2)\right)^{1/2}}$$

for all sufficiently small $B := \max_M |\nabla f|_g \ll 1$. Hence, as $B \rightarrow 0$,

$$\frac{\text{(RHS) of (13)}}{\text{(RHS) of (14)}} \searrow 1.$$

On the other hand,

$$\frac{\text{(RHS) of (13)}}{\text{(RHS) of (14)}} \rightarrow 0$$

as $B \rightarrow +\infty$.

Moreover they have proven this type of estimate (14) under more general setting (i.e., not only for the case of (M^n, g) is a steady gradient Ricci soliton with $n \leq 7$ but also for the case of (M^n, g, X) is a tuple that consists of a Riemannian n -manifold (M^n, g) for all $n \geq 2$ and a vector field X on M satisfying $\text{Ric}_g + \frac{1}{2}\mathcal{L}_X g \geq \frac{\lambda}{2}g$ for some positive constant λ with $\sup_M |X|_g < \infty$).

Remark 5.3. As observed in Remark 5.2, our upper bound (13) is better than (14) when $\sup_M |\nabla f|_g$ is large enough. Hence we can also obtain certain sufficient condition for four dimensional compact connected shrinking gradient Ricci solitons to satisfy the Hitchin–Thorpe inequality, which is better than [23, (1.10)] when $\sup_M |\nabla f|_g$ is large enough (see [23, Proof of Corollary 1.9]).

References

- [1] R. H. Bamler, P.-Y. Chan, Z. Ma and Y. Zhang, An optimal volume growth estimate for noncollapsed steady gradient Ricci solitons, *Peking Math. J.* **6** (2023), 353–364.
- [2] H.-D. Cao, Recent progress on Ricci solitons, arXiv preprint arXiv:0908.2006 (2009).
- [3] O. Chodosh and C. Li, Generalized soap bubbles and the topology of manifolds with positive scalar curvature, *Ann. Math.* **199** (2024), 707–740.
- [4] O. Chodosh, C. Li, P. Minter and D. Stryker, Stable minimal hypersurfaces in \mathbb{R}^5 , arXiv preprint arXiv:2401.01492v1 (2024).
- [5] B. Chow, P. Lu and L. Ni, Hamilton’s Ricci flow, *Gradient Studies in Mathematics* **77**, AMS, Providence (2006).
- [6] B. Chow and P. Lu, Mild Ricci curvature restrictions for steady gradient Ricci solitons, arXiv preprint arXiv:1103.5505 (2011).
- [7] M. Fernández-López and E. García-Río, Maximum principles and gradient Ricci solitons, *J. Differ. Equ.* **251** (2011), 73–81.
- [8] R. S. Hamilton, Eternal solutions to the Ricci flow, *J. Diff Geom.* **38** (1993), 1–11.
- [9] R. S. Hamilton, The formations of singularities in the Ricci Flow, *Surveys in Differential Geometry* **2** (1993), 7–136.
- [10] L. Karp, Differential inequalities on complete Riemannian manifolds and applications, *Math. Ann.* **272** (1985), 449–459.
- [11] Y. Lai, A family of 3d steady gradient solitons that are flying wings, arXiv preprint arXiv:2010.07272 (2020).
- [12] Y. Lai, 3D flying wings for any asymptotic cones, arXiv preprint arXiv:2207.02714 (2022).
- [13] P. Li, *Geometric analysis*, Cambridge Studies in Advanced Mathematics, **134**, Cambridge: Cambridge University Press (2012).
- [14] P. Li and J. Wang, Complete manifolds with positive spectrum, II *J. Differential Geom.* **62** (2002), 143–162.

- [15] P. Li and S.-T. Yau, On the parabolic kernel of the Schrödinger operator, *Acta Math.* **156** (1986), 153–201.
- [16] M. Limoncu, Modifications of the Ricci tensor and applications, *Arch. Math.* **96** (2010), 191–199.
- [17] L. Ma, Volume growth and Bernstein theorems for translating solitons, *J. Math. Anal. Appl.* **473** (2019), 1244–1252.
- [18] O. Munteanu and N. Sesum, On gradient Ricci solitons, *J. Geom. Anal.* **23** (2013), 539–561.
- [19] O. Munteanu and J. Wang, Geometry of three-dimensional manifolds with positive scalar curvature, arXiv preprint arXiv:2201.05595v3 (2024).
- [20] S. B. Myers, Riemannian manifolds with positive mean curvature, *Duke Math. J.* **8** (1941), 401–404.
- [21] D. Råde, Scalar and mean curvature comparison via μ -bubbles, *Calc. Var.* **62** (2023), pp. 187.
- [22] J. Sun, Mean curvature decay in symplectic and lagrangian translating solitons, *Geom. Dedicata* **172** (2014), 207–215.
- [23] H. Tadano, An upper diameter bound for compact Ricci solitons with application to the Hitchin–Thorpe inequality. II, *J. Math. Phys.* **59** (2018).
- [24] N. Th Varopoulos, The Poisson kernel on positively curved manifolds, *J. Funct. Anal.* **44** (1981), 359–380.
- [25] P. Wu, On the potential function of gradient steady Ricci solitons, *J. Geom. Anal.* **23** (2013), 221–228.
- [26] J. Xie and J. Yu, Convexity of 2-Convex Translating and Expanding Solitons to the Mean Curvature Flow in \mathbb{R}^{n+1} , *J. Geom. Anal.* **33** (2023), pp. 252.
- [27] Z. H. Zhang, On the completeness of gradient Ricci solitons, *Proc. Am. Math. Soc.* **137** (2009), 2755–2759.
- [28] J. Zhu, Width estimate and doubly warped product, *Trans. Am. Math. Soc.* **374** (2021), 1497–1511.

E-mail adress: hamanaka.shota.sci@osaka-u.ac.jp

DEPARTMENT OF MATHEMATICS, GRADUATE SCHOOL OF SCIENCE, OSAKA UNIVERSITY, TOYONAKA, OSAKA 560-0043, JAPAN