# SKEW-BRAUER GRAPH ALGEBRAS

<span id="page-0-0"></span>ANA GARCÍA-ELSENER, VICTORIA GUAZZELLI, AND YADIRA VALDIVIESO

Abstract. In this work, we introduce a new class of algebras called skew-Brauer graph algebras, which generalize the well-known Brauer graph algebras. We establish that skew-Brauer graph algebras are symmetric and can be defined using a Brauer graph with additional information. We show that the class of trivial extensions of skew-gentle algebras coincides with a subclass of skew-Brauer graph algebras, where the associated skew-Brauer graph has multiplicity function identically equal to one, generalizing a result over gentle algebras. Furthermore, we characterize skew-Brauer algebras of finite representation type. Finally, we provide a geometric interpretation of cut-sets and reflections of algebras using orbifold dissections.

### 1. INTRODUCTION

Brauer graph algebras are defined combinatorially through a graph with some additional data on their vertices, given by a multiplicity function and an order over the edges attached to each vertex. They appeared first in representation theory of groups, and were defined in [DF78] to classify indecomposable modules over quasi-Frobenious algebras.

Brauer graph algebras have been recently related to cluster algebras of surfaces, gentle algebras and mirror symmetry, see for example [MS14, Sch15, OPS18], moreover they have played an important role in the representation theory of tame symmetric algebras. Most of the relevant questions about their homological properties have been studied, for example, it is well understood their derived equivalence classes [OZ22] [AZ22], the classification of their two-term tilting complexes [AAC18] and their first Hochschild cohomology group [CSS20]. A natural relation between trivial extension of gentle algebras and a subclass of Brauer graph algebras has been described in [Sch15], more precisely, the paper shows that any trivial extension of a gentle algebra is actually a Brauer graph algebra, in which the multiplicity function is identically equal to one.

Several generalizations of Brauer graph algebras have been defined using slightly modified graphs or definitions, in which important properties of the algebra, and its modules, can be directly understood from its graph. See for example [GS17] where the so-called Brauer configuration algebras are defined, or [Sko06] where the modified Brauer algebras are constructed.

In this paper, we define what we call a skew-Brauer graph, which is by definition a Brauer graph Γ, that has a set of distinguished vertices  $\Gamma_0^{\times}$  with valency one and multiplicity function equal to one. Using this extra data we define a path algebra called skew-Brauer graph algebra, in order to link this new family of algebras and the trivial extensions of skew-gentle algebras, which are skew-group algebras of gentle algebras with a  $\mathbb{Z}_2$  action.

Skew-gentle algebras were defined in [GdlP99] to study a subclass of matrix-problems, and describe explicitly their Auslander-Reiten sequences. This family of tame algebras is also related to several

AGE, VG and YV would like to thank the Isaac Newton Institute for Mathematical Sciences, Cambridge, for support and hospitality during the programme "Cluster Algebras and Representation Theory" where part of the work on this paper was undertaken, this work was supported by EPSRC grant EP/R014604/1. YV was supported by EPSRC grant no EP/K032208/1 and the Simons Foundation. YV is also supported by the European Union's Horizon 2020 research and innovation programme under the Marie Sklodowska-Curie grant agreement No H2020-MSCA-IF-2018-838316. AGE was supported by PICT 2021-I-A-01154 Agencia Nacional de Promoción Científica y Tecnológica and by the EPSRC grant EP/R009325/1.

subjects, for example, they were used in [\[GLFS16\]](#page-0-0) to prove that a large class of Jacobian algebras arising from closed surfaces are also tame. Skew-gentle algebras are also a generalization of gentle algebras, and as they are, by definition, skew-group algebras of gentle, both families, gentle and skew-gentle algebras share several properties, for example, both of them are Gorenstein, see [\[GR05\]](#page-0-0).

A skew-gentle algebra can be also defined as a gentle algebra with some extra data, that is a set of special loops attached to a set of special vertices, and a set of special relations, some of them non-admissible. Using this definition, we show that the original skew-gentle algebra and the underline gentle algebra also share some properties.

In this work, we study trivial extensions of skew-gentle algebras, and we show that there is a simple way to define such trivial extension using the trivial extension of the underline gentle algebra. Even more, we show that any trivial extension of a skew-gentle algebra is a skew-Brauer graph algebra. More precisely, we have the following result.

Theorem A (Theorem 4.5, Theorem 4.9 and Theorem 4.11 )

- i) Any skew-Brauer graph algebra is a symmetric finite dimensional algebra.
- ii) The class of trivial extension of skew-gentle algebras coincides with the subclass of skew-Brauer graph algebras with multiplicity function identically equal to one.

Deciding whether a trivial extension of an algebra is of finite representation type is a question that has been addressed by several authors. In this work, we consider two different tools that were used to solve that problem. The first one is the admissible cuts or cut sets, defined in [\[Fer99, FP06\]](#page-0-0), to provide a complete characterization of trivial extension of finite representation type. The second one is reflections of quivers and relations, defined in [\[HW83\]](#page-0-0), to show that any trivial extension is of finite representation type if and only if it is related to an algebra of Dynkin type through sequences of reflections. In this paper, we study reflections and cut sets that preserve the class of skew-gentle algebras and their trivial extensions, in this context, we have the following results.

**Theorem B** (Theorem 5.4 and Theorem 5.7) Let  $A$  be a skew-gentle algebra,  $A'$  be its underline gentle algebra and i be a source in A. Then the reflection  $S_i$  of A' gives rise a sequence of reflections  $\overline{S_i}$  on A such that  $\overline{S_i}A$  is skew-gentle. Futhermore there exists a cut set  $\mathcal D$  of  $T(A)$  such that  $\overline{S_i} = T(A)/\langle \mathcal D \rangle$ 

Following a natural question, we also show the following characterisation of skew-Brauer graph algebras of finite representation type, see definition of skew-Brauer tree graph in Section 6.

**Theorem C** (Theorem 6.10 and Theorem 6.11) Let  $B_{\Gamma} \times$  be a skew-Brauer graph algebra. Then  $B_{\Gamma} \times$ is of finite representation type if and only if  $\Gamma^\times$  is a skew-Brauer tree graph. Moreover, in this case,  $B_{\Gamma} \times$  is isomorphic to the trivial extension of an iterated tilted algebra of type  $\mathbb{D}$ .

Additionally, we present a geometric interpretation of cut sets and reflections using orbifold dissections, a geometric model inspired on the one for the derived category of skew-gentle algebras and the Koszul dual algebra, see [\[AB19\]](#page-0-0) and [\[LFSV20\]](#page-0-0). More specifically, we establish the following theorems.

**Theorem D** (Theorem 7.7) Let A be a skew-gentle algebra with Sp the set of special vertices,  $A^{sg}$  be its admissible presentation and  $(S, M, P, O, M)$  be its orbifold  $\times$ -dissection. Then, any good cut set of the trivial extension  $T(A^{sg})$  of  $A^{sg}$  is induced by a sequence of contracting and adding boundary segments in  $(S, M, P, O, M)$ .

As a consequence of Theorem B and Theorem D, one can prove that the sequences of reflections described in Theorem B are also induced by contracting and adding boundary segments.

### Remark:

- (1) Generalizations of Brauer graphs have been studied by several authors. In particular, generalizations related to trivial extension of iterated tilted of type  $\mathbb{D}_n$  appeared in [\[Nor21\]](#page-0-0), where skew-Brauer tree graphs are named as modified Brauer tree (and double edge algebras).
- (2) This work started in November 2021 in the Newton Institute, since then, the authors presented preliminary results in several conferences and workshops  $1, 2, 3$  $1, 2, 3$  $1, 2, 3$  $1, 2, 3$  $1, 2, 3$ .
- (3) Recently, we found a similar definition of skew-Brauer graph algebras in [\[Sot24\]](#page-0-0).

**Acknowledgment:** We would like to thank Elsa Ferández for an insightful discussion on trivial extension of iterated tilted algebras of Dynkin type.

# 2. Background

Let K be an algebraically closed field of characteristic  $\neq 2$ . Let  $Q = (Q_0, Q_1)$  be a finite quiver, where  $Q_0$  denotes the set of vertices and  $Q_1$  denotes the set of arrows. For an arrow  $\alpha$  denote  $s(\alpha)$  the starting vertex and  $t(\alpha)$  the ending vertex. A path p of length  $l > 0$  from the vertex x to the vertex y in Q is an ordered set of arrows  $\{\alpha_1, \ldots, \alpha_l\}$  such that  $s(\alpha_1) = x$ ,  $t(\alpha_l) = y$  and  $t(\alpha_i) = s(\alpha_{i+1})$ for all  $1 \leq i \leq l-1$ , and we define  $s(p) = s(\alpha_1)$  and  $t(p) = t(\alpha_l)$ , and by abuse of notation, we write  $p = \alpha_1 \alpha_2 \ldots \alpha_l$ . A path p is a cycle if  $s(p) = t(p)$ . In addition, for each vertex i we denote by  $e_i$  the stationary path of length 0.

Consider the algebra  $KQ$  generated by paths of any length, and denote by  $J$  the two-sided ideal generated by arrows. A two-sided ideal I of  $KQ$  is called *admissible* if there exists  $m \geq 2$  such that  $J^m \subset I \subset J^2$ , where  $J^m$  is the ideal generated by paths of length m and  $J^2$  is the ideal generated by paths of length 2. The set of generators in  $I$  is called  $a$  set of relations.

The pair  $(Q, I)$  is said to be a *bound quiver*, and the algebra  $KQ/I$  is the bound quiver algebra. Every basic and connected finite dimensional K-algebra A can be defined by a bound quiver with an admissible ideal  $(Q_A, I_A)$  [\[ASS](#page-0-0)+06, Theorem 3.7], in this case  $Q_A$  is called the ordinary quiver of A.

Finally a path p in Q is non-zero in  $(Q, I)$  if  $p \notin I$ , and a non-zero path p is maximal if for any arrow  $\alpha \in Q$ , the paths  $\alpha p$  and  $p\alpha$  are elements of I.

2.1. Gentle and skew-gentle algebras. Skew-gentle algebras were defined in [\[GdlP99\]](#page-0-0) as certain skew-group algebras of gentle algebras. We first recall the definition of gentle algebra.

**Definition 2.1.** A K-algebra A is gentle if it is Morita equivalent to a path algebra  $KQ/I$ , where

- (1)  $Q$  is a quiver such that there is at most two arrows ending and starting at each vertex  $x$ ;
- (2) for every arrow  $\alpha$  in  $Q_1$  there is at most one arrow  $\beta$  such that  $t(\alpha) = s(\beta)$  and  $\alpha\beta \in I$ , and there is at most one arrow  $\beta'$  such that  $t(\alpha) = s(\beta')$  and  $\alpha\beta' \notin I$ ;
- (3) for every arrow  $\alpha$  in  $Q_1$  there is at most one arrow  $\gamma$  such that  $s(\alpha) = t(\gamma)$  and  $\gamma \alpha \in I$ , and there is at most one arrow  $\gamma'$  such that  $s(\alpha) = t(\gamma')$  and  $\gamma' \alpha \notin I$ ;
- (4)  $I$  is a 2-sided ideal of  $KQ$  generated by some paths of length 2;
- (5)  $I$  is an admissible ideal of  $KQ$ .

If the bound quiver  $(Q, I)$  satisfies  $(1), (2), (3)$  and  $(4)$ , then we say that the quotient  $KQ/I$  is locally gentle.

<span id="page-2-0"></span><sup>1</sup> II Encuentro conjunto RSME-UMA, December 12-16, 2022. Website [https://www.rsme.es/congresos-y](https://www.rsme.es/congresos-y-escuelas/congresos-conjuntos-con-otras-sociedades-cientificas/ii-encuentro-conjunto-rsme-uma/)[escuelas/congresos-conjuntos-con-otras-sociedades-cientificas/ii-encuentro-conjunto-rsme-uma/](https://www.rsme.es/congresos-y-escuelas/congresos-conjuntos-con-otras-sociedades-cientificas/ii-encuentro-conjunto-rsme-uma/)

<span id="page-2-1"></span><sup>2</sup>Combinatorial aspects of Representation Theory, March 28-31, 2023. Website [https://wiki.math.ntnu.no/combrep/program/](https://wiki.math.ntnu.no/combrep/program)

<span id="page-2-2"></span><sup>3</sup>FD seminar, Talk 133. September 14th, 2023. Website<https://www.fd-seminar.xyz/talks/2023-09-14/>

**Definition 2.2.** A K-algebra  $\Lambda$  is *(locally) skew-gentle* if it is Morita equivalent to an algebra  $KQ/I$ where

- (1)  $Q_1 = Q'_1 \cup S$  where S is a set of special loops,
- (2)  $I = \langle I' \cup \{ f^2 f | f \in S \} \rangle;$
- (3)  $KQ'/I'$  is a (locally) gentle algebra where  $Q' = (Q_0, Q'_1);$
- (4) if  $f \in S$  then  $x = s(f) = t(f)$  is the start or the end of exactly one arrow in  $Q'_1$  and, if there is an arrow  $\alpha \in Q'_1$  with  $t(\alpha) = x$  and an arrow  $\beta \in Q'_1$  with  $s(\beta) = x$ , then  $\alpha\beta \in I'$ . Moreover, there is no other loop at the vertex  $x$ .

We say that a vertex  $x \in Q_0$  is special if there exists a special loop at x and we denote by Sp the set of special vertices.

*Remark* 2.3. Given a skew-gentle algebra  $A = KQ/I$ , we observe that:

- the algebra  $\Lambda$ , obtained from  $A$  by deleting all the special loops, is gentle;
- if A has at least one special vertex, then the ideal  $I$  is not admissible. However, in [\[GdlP99\]](#page-0-0), the authors present the quiver with admissible relations for a skew-gentle algebra. The reader can see this construction in Example 3.5.

2.2. Trivial extensions. Trivial extensions of artin algebras play an important role in representation theory. In this section we describe a construction for the quiver and relations of trivial extension of a bound quiver algebra. We mainly follow the articles  $[FP02]$ ,  $[FFT+22]$  and  $[Sch15]$ .

The trivial extension of an algebra A is the algebra  $T(A) = A \ltimes DA$  with underline vector space  $A \oplus D A$  and with the multiplication defined by  $(a, f)(b, g) = (ab, ag + fb)$ , for  $a, b \in A$  and  $f, g \in DA$ , where  $D = \text{Hom}_K(-, K)$ .

For every finite dimensional algebra  $A = KQ_A/I_A$ , where  $I_A$  is an admissible ideal, we can construct its trivial extension  $T(A)$  based on the quiver and relations of A and a basis of the two-side ideal soc<sub> $A^e$ </sub> A, where  $A^e$  is the enveloping algebra.

Recall that as K-vector space  $\operatorname{soc}_{A^e} A$  can be defined as the intersection of

 $\text{soc}(A A) = \{a \in A : Ja = 0\}$  and  $\text{soc}(A_A) = \{a \in A : aJ = 0\},\$ 

where J denotes the Jacobson radical, i.e. the ideal generated by arrows. See [\[Lam91,](#page-0-0) Exercise 4.20] and  $[ASS<sup>+</sup>06, Lemma 2.10]$ .

Let  $\mathcal{M} = \{p_1, \ldots, p_r\}$  be K-basis of soc<sub>A</sub><sup>e</sup> A. The ordinary quiver of  $T(A)$  is given by:

- (1)  $(Q_{T(A)})_0 = (Q_A)_0$
- $(2)$   $(Q_{T(A)})_1 = (Q_A)_1 \cup \{\beta_{p_1}, \ldots, \beta_{p_r}\},$  where  $\beta_{p_i}$  is an arrow from  $t(p_i)$  to  $s(p_i)$  for each  $p_i \in \mathcal{M},$ with  $1 \leq i \leq r$ .

To give an explicit description of the ideal of relations  $I_{T(A)}$  we need to recall some definitions from [\[FP02,](#page-0-0) Section 3].

**Definition 2.4.** Let  $M = \{p_1, \ldots, p_r\}$  be a basis of soc<sub>A<sup>e</sub></sup> A. An oriented cycle  $C = \alpha_1 \ldots \alpha_l \beta_{p_j}$  in</sub>  $KQ_{T(A)}$  is said to be *elementary* if  $p_j^*(\alpha_1 \dots \alpha_l) \neq 0$  where  $\alpha_1 \dots \alpha_l$  is a path in  $KQ_A$ ,  $p_j \in \mathcal{M}$ , and  $p_j^*$  is the dual of  $p_j$ . In this case, we say that  $\omega(C) = p_j^*(\alpha_1 \dots \alpha_l) \in K$  is the *weight* of C.

Let C be a cycle of a quiver  $Q$  and  $q$  be a path contained in C. The supplement of  $q$  in C is the (possibly stationary) path p such that  $C = pq$  up to cyclic equivalence.

For each vertex  $x \in Q_{T(A)}$  we define the two-sided ideal  $I'_x$  in  $Q_{T(A)}$  generated by

- (1) Oriented cycles from  $x$  to  $x$  which are not contained in an elementary cycle.
- (2) The elements of the form  $\omega(C')C \omega(C)C'$ , where C and C' are elementary cycles starting and ending at x.

The ideal of relations for the trivial extension of a finite dimensional algebra was described in  $[FT^+22]$ .

**Theorem 2.5.** [\[FST](#page-0-0)<sup>+</sup>22, Theorem A.1] Let  $A = KQ_A/I_A$  a finite dimensional algebra and let  $T(A) =$  $KQ_{T(A)}/I_{T(A)}$  be its trivial extension. Then, the quiver  $Q_{T(A)}$  is as above and the ideal  $I_{T(A)}$  is generated by the union of the following sets.

- (1) A generating set of the ideal of relations  $I_A$  of  $A$ .
- (2) The paths that are not contained in any elementary cycle.
- (3) Linear combination of paths  $\rho \in e_x K Q_{T(A)} e_y$  such that  $\rho q \in I'_x$  or  $q \rho \in I'_y$  for any supplement path q in an elementary cycle C, where  $I'_x$  is the ideal defined above.

2.3. Repetitive algebra and reflections. In this section, we recall the definition of the repetitive algebra and reflections, both of them linked to an alternative construction for a trivial extension of an algebra. We mainly follow [\[Sch99, FP02, HW83\]](#page-0-0).

Let  $A = KQ/I$  be a finite dimensional algebra with I be an ideal generated by a set of relations  $\rho$  for  $Q$  which are either zero relations or commutativity relations, and let  $M$  be a set of maximal paths in  $(Q, I)$ . The repetitive algebra  $\widehat{A}$  is defined as the path algebra  $K\widehat{Q}/\langle\widehat{\rho}\rangle$ , where  $\widehat{Q}$  and  $\widehat{\rho}$  are defined as follows.

- (1) The set of vertices of  $\widehat{Q}$  is the set  $\{x[n] : x \in Q_0 \text{ and } n \in \mathbb{Z}\}.$
- (2) In the same way, every arrow  $\alpha : x \to y$  in  $Q_1$  induces arrows  $\alpha[n] : x[n] \to y[n]$  in  $\widehat{Q}$  for all  $n \in \mathbb{Z}$ .
- (3) For each maximal path  $p \in \mathcal{M}$  in  $(Q, I)$ , with  $x = s(p)$  and  $y = t(p)$ , there is an arrow  $\overline{p}[n] : y[n] \to x[n+1]$  in  $\widehat{Q}$ , for all  $n \in \mathbb{Z}$ . These arrows are called *connecting arrows*.

To define the set of relations  $\hat{\rho}$ , we need to introduce some definitions for paths in  $(Q, I)$ . Let p be a path in  $(Q, I)$ , we denote by  $p[n]$  a the corresponding path in  $\hat{Q}$ . If  $p = p_1p_2$  is a maximal path in  $(Q, I)$ , then  $p_2[n]\overline{p}[n]p_1[n+1]$  is called full path in  $\widetilde{Q}$ .

The set of relations  $\hat{\rho}$  of  $\hat{Q}$  are given by the following list.

- For each monomial relation  $p \in \rho$ , the path  $p[n]$  is in  $\widehat{\rho}$ , for all  $n \in \mathbb{Z}$ .
- For each commutative relation  $p_1 p_2 \in \rho$ , then  $p_1[n] p_2[n]$  is in  $\hat{\rho}$ , for all  $n \in \mathbb{Z}$ .
- Let p be a path in  $\tilde{Q}$  which contains a connecting arrow. If p is not a sub-path of a full path, then  $p \in \widehat{\rho}$ .
- If  $p = p_1p_2p_3$  and  $q = q_1q_2q_3$  are maximal paths in  $(Q, I)$  such that  $p_2 = q_2$ , the  $p_3[n]\overline{p}[n]p_1[n+1]$  $1] - q_3[n]\overline{q}[n]q_1[n+1]$  is in  $\widehat{\rho}$ , for all  $n \in \mathbb{Z}$ .

To define reflections over an algebra, we need to consider  $\nu$  the Nakayama automorphism of  $\widehat{A}$ , such that  $\nu(x[n]) = x[n+1]$  for every  $x \in Q_0$  and every  $n \in \mathbb{Z}$ .

Let x be a source of  $Q_A$ , we denote by  $\sigma_x^- Q_A$  the full sub-quiver of  $\widehat{Q}_A$  with vertex set

$$
S = \{y[0] \in (\widehat{Q_A})_0 \mid y \in (Q_A)_0, y \neq x\} \cup \{x[1]\}.
$$

We note that  $\sigma_x^- Q_A$  is a *complete v-slice*, which is defined as a connected full subquiver of  $\widehat{Q}_A$  which contains exactly one vertex from each  $\nu$ -orbit of vertices of  $Q_{\hat{A}}$ .

The **negative reflection**  $S_x^-(A)$  of A is the algebra  $\text{End}_{\widehat{A}}$  $\sqrt{2}$  $\oplus$  $\bigoplus_{x[z]\in S}e_{x[z]}A$  $\setminus$ , where  $\{e_{x[z]}\}_{x[z]\in S}$  is a set of primitive orthogonal idempotents of  $\widehat{A}$ .

A negative reflection sequence of  $Q_A$  is a sequence of vertices  $x_1, \ldots, x_r$  of  $Q_A$  such that  $x_j$  is a source of  $\sigma_{x_{j-1}}^- \dots \sigma_{x_1}^- Q_A$  for  $1 \leq j \leq r$ .

Similarly, given a sink x of  $Q_A$ , a positive reflection of an algebra is defined, considering  $\sigma_x^+ Q_A$  as the full-subquiver of  $\widehat{Q}_A$  with vertex set

$$
S = \{y[0] \in (\widehat{Q_A})_0 \mid y \in (Q_A)_0, y \neq x\} \cup \{x[-1]\}.
$$

Remark 2.6. The quiver of the negative (positive) reflection  $S_x(A)^-$  (resp.  $S_x(A)^+$ ) coincides with the quiver  $\sigma_x^- Q_A$  (resp.  $\sigma_x^+ Q_A$ ).

# 3. Trivial extension of a skew-gentle algebra

In this section, we describe the quiver and relations for the trivial extension of the admissible version of a skew-gentle algebra. To do this, we introduce a construction of quivers and relations that depends on a set of distinguished vertices from a given quiver, which, by abuse of notation and convenience, we denote by Sp. This construction is similar to the one presented in [\[GdlP99\]](#page-0-0), and revisited in [\[AB19, LFSV20\]](#page-0-0), implemented to define the admissible presentation of any skew-gentle algebra.

**Definition 3.1.** (sg-quiver  $Q^{sg}$ ) Let Q be a quiver with a set of *distinguished vertices*  $Sp$ , such that there is no loop incident at any distinguished vertex. We define the  $sg\text{-}quiver Q^{sg}$  as follows. We start by duplicating the distinguished vertices in  $Q$ , that is, for each vertex  $x$  in  $Q_0$  we define

$$
Q_0^{sg}(x) = \begin{cases} \{x^+, x^-\} & \text{if } x \in Sp\\ \{x\} & \text{otherwise,} \end{cases}
$$

obtaining the set of vertices  $Q_0^{sg} := \bigcup_{x \in Q_0} Q_0^{sg}(x)$ . For each arrow  $\alpha$  in  $Q_1$  we define the set of arrows

$$
Q^{sg}_1(\alpha) := \{(x,\alpha,y) \mid x \in Q^{sg}_0(s(\alpha)), y \in Q^{sg}_0(t(\alpha))\}
$$

obtaining the set of arrows  $Q_1^{sg} := \bigcup_{\alpha \in Q_1} Q_1^{sg}(\alpha)$ .

In other words, while arrows incident to non-distinguished vertices remain the same, for an arrow in Q of the form  $x \xrightarrow{\alpha} y$ , if one of the vertices, say y, belongs to Sp then we will have two arrows in  $Q^{sg}$ . If both vertices x and y are distinguished there will be four arrows in  $Q^{sg}$ . See these cases below.



To shorten notation, we use upper scripts for new arrows in  $Q^{sg}$ , for instance,  $(x, \alpha, y^+)$  can be denoted  $x \xrightarrow{\alpha^+} y^+,$  and  $(x^-, \alpha, y^+)$  can be denoted  $x^- \xrightarrow{-\alpha^+} y^+,$  see Example 3.5. In our proofs we denote arrows after distinguished vertex duplication by  $\epsilon \alpha^{\epsilon'}$ , where the indices  $\epsilon, \epsilon'$  will be always considered in  $\{+,-,\emptyset\}$ .

The next definition generalizes the construction of the admissible ideal of a skew-gentle algebra.

**Definition 3.2.** (sg-ideal  $I^{sg}$ ) Let  $(Q, I, Sp, C)$  be a tuple, where  $(Q, I)$  is a bound quiver, with  $Sp \subset Q_0$  a subset of distinguished vertices such that there are no loops at any  $x \in Sp$ . Let C be a (possibly empty) set of distinguished cycles such that there is at most one distinguished cycle starting at each vertex in  $Sp.$  Moreover, the ideal I is generated by

- A (possibly empty) set of monomial relations such that if  $\alpha\beta$  is a monomial relation of length two, then  $t(\alpha) = s(\beta) \notin Sp$ ,
- the difference  $C_1 C_2$  of any two distinguished cycles  $C_1$  and  $C_2$  starting at the same nondistinguished vertex.

For the quiver Q in the tuple  $(Q, I, Sp, \mathcal{C})$ , consider the quiver  $Q^{sg}$ . We define the sg-ideal  $I^{sg}$  of  $KQ^{sg}$  as the ideal generated by

- **Type a:** Commutative relations  $({}^{\varepsilon}\alpha^{+})({}^{\varepsilon}\beta^{\varepsilon'}) ({}^{\varepsilon}\alpha^{-})({}^{\varepsilon}\beta^{\varepsilon'})$  such that  $\alpha\beta$  is a path of length 2 in Q with  $t(\alpha) \in Sp$ , for all possible  $\varepsilon, \varepsilon' \in \{+, -, \emptyset\}.$
- **Type b:** Commutative relations  $(\alpha_1^{\varepsilon_2})(\varepsilon_2 \alpha_2^{\varepsilon_3}) \dots (\varepsilon_k \alpha_k) (\beta_1^{\varepsilon_2})(\varepsilon_2' \beta_2^{\varepsilon_3'}) \dots (\varepsilon_l' \beta_l)$ , for each difference of two distinguished cycles  $C_1 - C_2 = \alpha_1 \dots \alpha_k - \beta_1 \dots \beta_l$  starting at the same non-distinguished vertex, for all possible  $\varepsilon_i$ ,  $\varepsilon'_j = \{+, -, \emptyset\}$  with  $i = 1, ..., k$  and  $j = 1, ..., l$ .
- **Type c**: Monomial relations  $({}^{\varepsilon_1}\alpha_1^{\varepsilon_2})({}^{\varepsilon_2}\alpha_2^{\varepsilon_3})\dots({}^{\varepsilon_k}\alpha_k^{\varepsilon_{k+1}})$  where  $\alpha_1 \dots \alpha_k$  is a monomial relation of *I*, for all possible  $\varepsilon_i = \{+, -, \emptyset\}$  and  $i = 1, ..., k + 1$ .
- **Type d:** A monomial relation  $({}^{\varepsilon_1}\alpha_1^{\varepsilon_2})({}^{\varepsilon_2}\alpha_2^{\varepsilon_3})\dots({}^{\varepsilon_k}\alpha_k^{\varepsilon_{k+1}}),$  for each distinguished cycle  $C = \alpha_1 \dots \alpha_k$ starting at a distinguished vertex, such that  $\varepsilon_1 \neq \varepsilon_{k+1}$ , for all possible  $\varepsilon_i = \{+, -\emptyset\}$  and  $i = 1, \ldots, k + 1.$

Given  $A = KQ/I$  a bound-quiver algebra with  $(Q, I, Sp, C)$  as in Definition 3.2, for abuse of notation we denote by  $A^{sg}$  the bound quiver algebra  $KQ^{sg}/I^{sg}$ . We say that  $A^{sg}$  is the sq-bound quiver algebra of A.

To show that the previous definitions generalize the construction of the admissible presentation of a skew-gentle algebra, we need to introduce the following definition.

**Definition 3.3.** Let  $A = KQ/I$  be skew-gentle algebra with Sp a set of special vertices and S a set of special loops. The *auxiliary gentle algebra A'* is the algebra  $KQ'/I'$  where  $Q'$  is the quiver obtained from Q by deleting the special loops and I' is the ideal generated by the relations of length two  $\alpha\beta$  of I such that  $t(\alpha) \notin Sp$ . We say that Q' is the *auxiliary quiver* of A and I' is the *auxiliary ideal* of A.

Remark 3.4. With the notation above, given a skew-gentle  $A = KQ/I$ , the quiver of the admissible presentation of A is the quiver  $(Q')^{sg}$  and its ideal is the ideal  $(I')^{sg}$ , with  $Sp$  as the set of distinguished vertices and an empty set of distinguished cycles, see [\[GdlP99\]](#page-0-0) and [\[LFSV20,](#page-0-0) Section 2.1] . In this context, by abuse of notation, we denote  $(Q')^{sg}$  by  $Q^{sg}$ ,  $(I')^{sg}$  by  $I^{sg}$ , and we say that  $A^{sg} = KQ^{sg}/I^{sg}$ is the admissible presentation for the skew gentle algebra A.

**Example 3.5.** Let  $KQ/I$  be the skew-gentle algebra with quiver  $Q$ 



ideal  $I = \langle f_i^2 - f_i, \alpha \beta, \gamma \delta, \lambda \gamma \rangle$  and  $Sp = \{1, 2\}$  the set of special vertices. Following Remark3.4 to compute the quiver of the admissible presentation of  $A$ , we consider  $Q'$  be the auxiliary quiver of  $Q$ and  $Sp = \{1, 2\}$  the set of distinguished vertices and  $I' = \langle \gamma \delta, \lambda \gamma \rangle$ .

Following the notation, the sg-bound quiver algebra  $A^{sg}$  of  $A' = KQ'/I'$  is the algebra  $KQ^{sg}/I^{sg}$ , where  $Q^{sg}$  is the quiver



and  $(I')^{sg} = \langle (a^+ \alpha^+)(a^+ \beta) - (a^+ \alpha^-)(a^- \beta), (a^+ \beta^+)(a^- \beta^+)(a^- \beta), \gamma \delta, \lambda \gamma \rangle$ , which coincides with the admissible presentation of A as defined in [\[GdlP99\]](#page-0-0).

Remark 3.6. Let  $p = \alpha_1 \ldots \alpha_r$  be a non-zero path from x to y in a bound quiver algebra  $KQ/I$ . Then, p induces a set of non-zero paths  $({}^{\varepsilon_1}\alpha_1^{\varepsilon_2})\cdots({}^{\varepsilon_r}\alpha_r^{\varepsilon_{r+1}})$  in the sg-bound quiver algebra  $KQ^{sg}/I^{sg}$ , for all possible  $\varepsilon_i = \{+,-,\emptyset\}$ , with  $i = 1,\ldots,r+1$ . Notice that by the commutativity relations of Type (a) of the sg-ideal  $I^{sg}$  given in Definition 3.2, any two paths  $({}^{\varepsilon_1}\alpha_1^{\varepsilon'_2}) \cdots ({}^{\varepsilon'_r}\alpha_r^{\varepsilon_{r+1}})$  and  $({}^{\varepsilon_1}\alpha_1^{\varepsilon_2}) \cdots ({}^{\varepsilon_r}\alpha_r^{\varepsilon_{r+1}})$ from  $x^{\varepsilon_1}$  to  $y^{\varepsilon_{r+1}}$ 

are equivalent modulo  $I^{sg}$ . Then, we denote the unique representative path in  $KQ^{sg}/I^{sg}$  by  $\epsilon_1 p^{\epsilon_{r+1}}$ .

In order to compute the quiver of the trivial extension of a skew-gentle algebra  $A$ , we need to consider a basis of the socle of the admissible presentation  $A^{sg}$  of A (see section 2.2). For our next purpose, we recall that a path p (in an admissible presentation  $A = KQ/I$ ) is called maximal, if  $p\alpha = 0$  and  $\alpha p = 0$  in A for any arrow  $\alpha$  in Q.

**Lemma 3.7.** Let  $A = KQ/I$  be a skew-gentle algebra and let  $A' = KQ'/I'$  be its auxiliary gentle algebra. Consider  $A^{sg} = KQ^{sg}/I^{sg}$  to be the admissible presentation of A. The following statements hold.

- (a) A non-zero path in A' induces at most four linearly independent non-zero paths in  $A^{sg}$ . Conversely, any non-zero path in  $A^{sg}$  is induced by a path in  $A'$ .
- (b) A path in A' is maximal if and only if any induced path in  $A^{sg}$  is a maximal path.

*Proof.* Let A be a skew-gentle algebra with special vertices  $Sp$ . We consider  $A^{sg}$  the admissible presentation of  $A$  and  $A'$  its auxiliary gentle algebra. We consider  $Sp$  the set of distinguished vertices of  $A'$ .

(a) Let  $p = \alpha_1 \dots \alpha_r$  be a non-zero path in A'. Following Remark 3.6, p induces set of paths  $\epsilon_p \epsilon'$  for all possible  $\varepsilon, \varepsilon' = \{+, -, \emptyset\}.$ 

In particular, a path  $p$  in  $A'$  induces exactly:

- four different equivalence classes of paths in  $A^{sg}$  if and only if  $s(p)$  and  $t(p)$  are special vertices, each one with different starting and ending vertex;
- two different equivalence classes of paths in  $A^{sg}$  if and only if one and only one of the vertices  $s(p)$  or  $t(p)$  is a special vertex, coinciding exclusively either at the starting or a the ending vertex;
- exactly one equivalence class of paths in  $A^{sg}$  if and only if none of the vertices  $s(p)$  and  $t(p)$ are special.

It is also clear that any non-zero path  $({}^{\varepsilon_1}\alpha_1^{\varepsilon_2})\cdots({}^{\varepsilon_r}\alpha_r^{\varepsilon_{r+1}})$  in  $A^{sg}$  is induced by the non-zero path  $p = \alpha_1 \dots \alpha_r$  in A'.

(b) Let  $p = \alpha_1 \dots \alpha_r$  be a maximal path in A', and let  $\epsilon_p^{\epsilon}$  be an induced path in  $A^{sg}$ , with  $\epsilon, \epsilon' \in$  $\{+, -,\emptyset\}$ . Suppose that  ${}^{\varepsilon}p^{\varepsilon'}$  is not maximal in  $A^{sg}$ , then there exist an arrow  ${}^{\varepsilon_1}\beta^{\varepsilon}$  (or  ${}^{\varepsilon'}\gamma^{\varepsilon_2}$ ) such that the path  $({}^{\varepsilon_1}\beta^{\varepsilon})(\varepsilon'p^{\varepsilon'})\neq 0$  (or  $({}^{\varepsilon}p^{\varepsilon'})({}^{\varepsilon'}\gamma^{\varepsilon_2})\neq 0$ , respectively). Therefore, such a path in  $A^{sg}$  is induced by a non-zero path  $\beta p$  (or  $p\gamma$ , respectively) in A', contradicting the fact that p is a maximal path. With similar argument, we claim that any maximal path  ${}^{\varepsilon}p^{\varepsilon'}$  in  $A^{sg}$  with all possible  $\varepsilon, \varepsilon' \in \{+, -, \emptyset\}$ is induced by a maximal path  $p$  in  $A'$ . . □

Recall that by [\[Sch15,](#page-0-0) Section 2.1], when A is gentle the set of maximal paths forms a basis for  $\operatorname{soc}_{A^e}(A)$ .

**Lemma 3.8.** Let  $A = KQ/I$  be a skew-gentle algebra with Sp a set of special vertices and  $A^{sg} =$  $KQ^{sg}/I^{sg}$  be its admissible presentation. Then the set of maximal paths M in  $A^{sg}$  forms a basis of  $\operatorname{soc}_{(A^{sg})^e}(A^{sg}).$ 

*Proof.* Let A' be the auxiliary gentle algebra of A. Denote  $\operatorname{soc}_{(A^{sg})^e}(A^{sg})$  by R. As a K-vector space, R is the direct sum

$$
R = \bigoplus_{x,y \in Q_0^{sg}} e_x R e_y
$$

where  $e_x$  and  $e_y$  are the trivial paths at a vertex x and y in  $Q_0^{sg}$ .

Let x and y be two vertices in  $Q^{sg}$ , and let x' and y' be their associated vertices in A'. Since A' is gentle, there are at most two linearly independent classes of paths  $p' = \alpha_1 \dots \alpha_n$  and  $q' = \beta_1 \dots \beta_m$ from  $x'$  to  $y'$  (see [\[BMM03,](#page-0-0) Lemma 3]). By Lemma 3.7 and Remark 3.6, there are at most two linearly independent classes of paths  $p$  and  $q$  from  $x$  to  $y$ .

Note that there are at most four arrows ending at x and at most four arrows starting at  $y$ , which correspond to arrows ending at  $x'$  and arrows starting at  $y'$ . We will show that the paths p and q induce a basis for  $e_xRe_y$ . We need to analyze each case based on the number of arrows ending at x and starting at y, respectively, and if there is exactly one independent class of paths from x and y or two, but we will describe two of them.

Suppose there are two independent class of path from  $x$  to  $y$  and suppose that there is exactly four arrows ending at x and four arrows staring at y, then there is exactly two arrows  $\gamma_1$  and  $\gamma_2$  ending at x' and two arrows  $\delta_1$  and  $\delta_2$  staring at y', and x' and y' are not special vertices of A'.

Since A' is gentle, either  $\gamma_1\alpha_1 \notin I$  or  $\gamma_2\alpha_1 \notin I$ , and either  $\gamma_1\beta_1 \notin I$  or  $\gamma_2\beta_1 \notin I$ . Therefore, p' and  $q'$  are not maximal in A', and by Lemma 3.7, neither are p and q in  $A^{sg}$ , and as a consequence  $e_x Re_y = \{0\}.$ 

Suppose now that there are exactly two arrows  $\gamma_1$  and  $\gamma_2$  ending at x and no arrows starting at y. In this case, we need to consider two cases,  $x'$  is a special vertex or not. In the first case, we have that  $\gamma_1$  and  $\gamma_2$  come from an arrow  $\gamma$  in A', and there is exactly one independent class of path p' from x' to y', since there is exactly one arrow staring at x'. Hence  $\gamma \alpha_1 \notin I$ , and p' is not maximal in A', neither p in  $A^{sg}$ , then  $e_x Re_y = \{0\}.$ 

If x' is not a special vertex, we have that either  $\gamma_1$  and  $\gamma_2$  are arrows of A', or they come from an arrow  $\gamma$  in A'. In the first case, by definition of gentle and a similar analysis as in the first case, the path or paths from  $x'$  to  $y'$  are not maximal, and neither the associated path or paths in  $A^{sg}$ .

In the second case, if there  $p \neq q$ , then either  $\gamma \alpha_1 \in I$  or  $\gamma \beta_1 \in I$ ; hence, p is maximal or q is maximal respectively. As a consequence  $\{p\}$  or  $\{q\}$  is a basis of  $e_x Re_y$ .

Finally, if there is exactly one independent class of paths, namely p, then either  $\gamma \alpha_1 \in I$  or  $\gamma \alpha_1 \notin I$ . In the first situation, p is maximal and is a basis of  $e_xRe_y$ , while in the second case, p is not maximal and therefore  $e_x R e_y = \{0\}$ .

**Lemma 3.9.** Let  $A = KQ/I$  be a skew-gentle algebra with special vertices  $Sp$ ,  $A^{sg} = KQ^{sg}/I^{sg}$  its admissible presentation and let  $A' = KQ'/I'$  be its auxiliary gentle algebra. Then the quiver  $Q_{T(A^{sg})}$ of the trivial extension of  $A^{sg}$  coincides with the sg-quiver  $Q^{sg}_{T(A')}$  of the trivial extension of  $A'$  with distinguished vertices Sp.

*Proof.* It follows by Lemma 3.8 and by Lemma 3.7 (b).

$$
\qquad \qquad \Box
$$

**Lemma 3.10.** Let  $A = KQ/I$  be a skew-gentle algebra with special vertices Sp with  $A^{sg} = KQ^{sg}/I^{sg}$ its admissible presentation and let  $A' = KQ'/I'$  be its auxiliary gentle algebra. We denote by  $T(A^{sg})$ and  $T(A')$  the trivial extension of  $A^{sg}$  and  $A'$ , respectively. Then a cycle C in  $T(A')$  is elementary if and only if any induced cycle  ${}^{\epsilon}C^{\epsilon}$  in  $Q_{T(A')}^{sg}$  is an elementary cycle in  $T(A^{sg})$ , for all possible  $\varepsilon \in \{+, -, \emptyset\}.$ 

*Proof.* Let  $T(A')$  be the trivial extension of A', and let C be an elementary cycle in  $T(A')$ . Then,  $C = \rho_1 \beta_\rho \rho_2$ , where  $\rho = \rho_2 \rho_1$  is a maximal path in A' and  $\beta_\rho$  is the added arrow in  $Q_{T(A')}$  with respect to the path  $\rho$ , see Section 2.2. Following Lemma 3.7, the induced paths  $\epsilon \rho^{\epsilon'}$ , for all possible  $\varepsilon, \varepsilon' \in \{+, -, \emptyset\}$ , are maximal paths in  $A^{sg}$ . By Lemma 3.8 we can give a basis M of the socle of  $A^{sg}$  such that  $\epsilon \rho^{\epsilon'} \in \mathcal{M}$ . Then, by Definition 2.4 there exist arrows  $\epsilon' \beta \epsilon$  in  $Q_{T(A^{sg})}$  such that  $({^{\varepsilon}\rho}^{\varepsilon'})({^{\varepsilon'}}\beta^{\varepsilon})$  are elementary cycles in  $T(A^{sg})$ . Notice that the arrows  $^{\varepsilon'}\beta^{\varepsilon}$  in  $Q^{sg}_{T(A')}$ , for all possible

 $\varepsilon, \varepsilon' \in \{+, -, \emptyset\}$ , are induced by the arrow  $\beta_{\rho}$  in  $T(A')$ . Therefore, the cycles  ${}^{\varepsilon}C^{\varepsilon} = ({}^{\varepsilon}\rho^{\varepsilon'})({}^{\varepsilon'}\beta^{\varepsilon})$  in  $T(A^{sg})$  are elementary cycles in  $T(A^{sg})$  induced by C.

With similar arguments one can show that any elementary cycle  ${}^{\varepsilon}C^{\varepsilon}$  in  $T(A^{sg})$  is induced by an elementary cycle C in  $T(A')$ . ).  $\Box$ 

**Theorem 3.11.** Let  $A = KQ/I$  be a skew-gentle algebra with special vertices Sp, let  $A^{sg}$  be its admissible presentation, and  $A' = KQ'/I'$  be its auxiliary gentle algebra. Then the trivial extension  $T(A^{sg})$  is isomorphic to the sg-bound quiver algebra of the trivial extension  $T(A')$  with Sp as the set of distinguished vertices and  $C$  the set of elementary cycles of  $T(A')$  as the set of distinguished cycles.

*Proof.* Let  $A = KQ/I$  be a skew-gentle algebra with special vertices  $Sp$  and let  $A^{sg} = KQ^{sg}/I^{sg}$  and  $A' = KQ'/I'$  be the algebras as described in the statement. By Lemma 3.9, the quivers  $Q_{T(A^{sg})}$  and  $Q_{T(A')}$  of the trivial extensions  $T(A^{sg})$  and  $T(A')$  coincide, then it is enough to verify that the ideal  $I_{T(A^{sg})}$  of the trivial extension coincides with the sg-ideal  $I_{T(A')}^{sg}$  of  $T(A')$ .

• Let x be a vertex in Sp, and let  $\alpha, \beta$  be arrows in  $Q_{T(A')}$  such that  $\cdot \stackrel{\alpha}{\to} x \stackrel{\beta}{\to} \cdot$ . The commutative relations  $({\epsilon \alpha^+})({^+ \beta^{\epsilon'}}) - ({\epsilon \alpha^-})({^- \beta^{\epsilon'}})$  in  $I^{sg}_{T(A')}$  are of type (a). For the ideal  $I_{T(A^{sg})}$  we have to consider two cases:

- (1) If  $\alpha$  and  $\beta$  are arrows of  $Q'$ , then such commutative relations are generators of  $I_{A^{sg}}$ , and therefore also of  $I_{T(A^{sg})}$ .
- (2) We assume that  $\beta$  (or  $\alpha$ ) is not an arrow of  $Q'$ .

Then, the arrows  $\epsilon''\beta^{\epsilon'}$  in  $Q_{T(A^{sg})}$  do not belong to  $Q_{A^{sg}}$ , with  $\epsilon'' \in \{+, -\}$ , and  $\epsilon' \in$  $\{+,-,\emptyset\}$ . That it to say that the mentioned arrows are new arrows in the quiver  $Q_{T(A^{sg})}$ which are in correspondence with maximal paths  $({\epsilon}' \rho^{\varepsilon})({\epsilon} {\alpha}^{\varepsilon} ...)$  in  $A^{sg}$ , with  $\varepsilon \in \{+, -\varnothing\}.$ Therefore, the commutative relations  $({\epsilon_{\alpha}}^{+})(+{\beta\epsilon^{\prime}}) - ({\epsilon_{\alpha}}^{-})(-{\beta\epsilon^{\prime}})$  in  $I_{T(A^{sg})}$  arise from the third condition of [\[FST](#page-0-0)+22, Theorem A.1], since the paths  $({\epsilon \alpha^+})({^+ \beta^{\epsilon'}})$  and  $({\epsilon \alpha^-})({^-\beta^{\epsilon'}})$  have the same complement  $\epsilon' \rho^{\epsilon}$  in an elementary cycle.

• Let  $C_1$  and  $C_2$  be two elementary cycles of  $T(A')$  starting at the same non-distinguished vertex. Then, by Lemma 3.10, we have that  ${}^{\varepsilon}C_1^{\varepsilon}$  and  ${}^{\varepsilon}C_2^{\varepsilon}$  are elementary cycles of  $Q(A^{sg})$ , with  $\varepsilon = \emptyset$ . Therefore, it follows from the third condition of  $[\text{FST}^+22, \text{Theorem A.1}]$  that  ${}^{\varepsilon}C_1^{\varepsilon} - {}^{\varepsilon}C_2^{\varepsilon}$  are generators of  $I_T(A^{sg})$ , and for the sg-ideal  $I_{T(A')}^{sg}$  of  $T(A')$ , such relations arise from relations of Type (b) in Definition 3.1.

• Finally, monomial relations of  $I_{T(A^{sg})}$  arise from monomial relations of  $I^{sg}$  or from paths which are not contained in any elementary cycle of  $T(A^{sg})$ . On the other hand, monomial relations of the sg-ideal  $I_T^{sg}(A')$  of  $T(A')$  arise from relations of Type (c) and (d) of Definition 3.1. It follows from Lemma 3.10 that these two mentioned sets are equal.  $\Box$ 

In the following example, we illustrate how to use Theorem 3.11 to compute the trivial extension of a skew-gentle algebra starting from its auxiliary gentle algebra.

**Example 3.12.** Let  $A = KQ/I$  be the skew gentle algebra given in Example 3.5 and let A' be its auxiliary gentle algebra. The trivial extension  $T(A')$  of A' is given by the quiver  $Q_{T(A')}$ 



and the ideal  $I_{T(A')}$  is generated by

(1)  $\gamma \delta, \lambda \gamma$ ; (2)  $\beta\beta_{p_2}, \beta_{p_2}\beta_{p_1}, \alpha\beta\gamma\beta_{p_1}\alpha, \beta\gamma\beta_{p_1}\alpha\beta, \gamma\beta_{p_1}\alpha\beta\gamma, \beta_{p_1}\alpha\beta\gamma\beta_{p_1}, \delta\lambda\beta_{p_2}\delta, \lambda\beta_{p_2}\delta\lambda, \beta_{p_2}\delta\lambda\beta;$ (3)  $\beta_{p_2}\delta\lambda - \gamma\beta_{p_1}\alpha\beta, \delta\lambda\beta_{p_2} - \beta_{p_1}\alpha\beta\gamma,$ 

where  $\beta_{p_1}$  and  $\beta_{p_2}$  correspond to the maximal paths  $p_1 = \alpha \beta \gamma$  and  $p_2 = \delta \lambda$ , respectively. It is follows from Theorem 3.11, that the trivial extension  $T(A^{sg})$  of  $A^{sg}$  is isomorphic to the sg-bound quiver algebra of  $T(A')$  with  $Sp = \{1,2\}$  as the set of distinguished vertices and  $\mathcal{C} = \{\alpha\beta\gamma\beta_{p_1}, \delta\lambda\beta_{p_2}\}\$ as the set of distinguished cycles. Then  $T(A^{sg})$  is the algebra with quiver



FIGURE 1. Quiver of the trivial extension  $T(A^{sg})$ 

and the ideal  $I_{T(A^{sg})}$  is generated by:

Type a:  $({}^+\alpha^+)({}^+\beta)-({}^+\alpha^-)({}^-\beta), ({}^-\alpha^+)({}^+\beta)-({}^-\alpha^-)({}^-\beta),$ Type b:  $\beta_{p_2}\delta\lambda - \gamma(\beta_{p_1})(-\alpha^{-})(-\beta), \beta_{p_2}\delta\lambda - \gamma(\beta_{p_1}^{+})(+\alpha^{+})(+\beta), \delta\lambda\beta_{p_2} - (\beta_{p_1}^{+})(+\alpha^{+})(+\beta)\gamma$  and  $\delta \lambda \beta_{p_2} - (\beta_{p_1}^-)(-\alpha^-)(-\beta)\gamma,$ Type c:  $({^+\beta})\beta_{p_2}, (\bar{\phantom{x}}\beta)\beta_{p_2}, \beta_{p_2}(\beta_{p_1}^+), \beta_{p_2}(\beta_{p_1}^-), ({^+\alpha^+})(^+\beta)\gamma(\beta_{p_1}^+)(^+\alpha^+), ({^+\beta})\gamma(\beta_{p_1}^+)(^+\alpha^+)(^+\beta),$  $\gamma(\beta^+_{p_1})({}^+\alpha^+)({}^+\beta)\gamma,\ (\beta^+_{p_1})({}^+\alpha^+)({}^{\dot +}\beta)\gamma(\beta^+_{p_1}),\ \delta\lambda\beta_{p_2}\delta,\lambda\beta_{p_2}\delta\lambda,\beta_{p_2}\delta\lambda\beta,$ Type d:  $({}^+\alpha^+)({}^+\beta)\gamma(\beta_{p_1}^+)({}^+\alpha^-),({}^-\alpha^-)({}^-\beta)\gamma(\beta_{p_1}^-)({}^-\alpha^+).$ 

### 4. Skew-Brauer graph algebras and trivial extension of skew-gentle algebras

In this section we define a generalization of Brauer graph algebras, that we call skew-Brauer graph algebras, and we show that the trivial extension of any skew-gentle algebra is isomorphic to a skew-Brauer graph algebra.

Given an undirected graph Γ, we denote by  $val(v)$  the *valency* of a vertex v, i.e. the amount of edges incident to  $v$  in  $\Gamma$ .

We recall that a *Brauer graph*  $\Gamma$  is a tuple  $(\Gamma_0, \Gamma_1, \mathfrak{m}, o)$  where  $(\Gamma_0, \Gamma_1)$  is an unoriented graph with  $\Gamma_0$  the set of vertices and  $\Gamma_1$  the set of edges and

- (1) **m** is a multiplicity function  $\mathfrak{m}: \Gamma_0 \to \mathbb{Z}^+$ , assigning a positive number to each  $v \in \Gamma_0$ .
- (2) **o** is an *orientation* of the edges attached to each  $v \in \Gamma_0$ . For a vertex v such that val(v) = t we denote  $\mathfrak{o}(v) = (x_1, \ldots, x_t)$ . In the particular case that val $(v) = 1$  and the only edge attached to v is x, if  $m(v) = 1$  the cyclic ordering is given by the single element x, but when  $m(v) > 1$ we set  $\mathfrak{o}(v) = (x, x)$  so that in the cyclic order the successor (and the predecessor) of x is x.

**Definition 4.1.** A skew-Brauer graph  $\Gamma^{\times}$  is tuple  $\Gamma^{\times} = (\Gamma_0', \Gamma_0^{\times}, \Gamma_1, \mathfrak{m}, \mathfrak{o})$  such that  $\Gamma = (\Gamma_0' \sqcup$  $\Gamma_0^{\times}, \Gamma_1, \mathfrak{m}, \mathfrak{o}$  is a Brauer graph with  $\Gamma_0' \sqcup \Gamma_0^{\times}$  a disjoint union of the set of vertices of  $\Gamma$ , and such that any vertex  $v \in \Gamma_0^{\times}$  satisfies  $\mathfrak{m}(v) = 1$ ,  $\mathfrak{o}(v) = (x_v)$ , and v is adjacent to a single vertex  $w \in \Gamma_0'$ .

We call the vertices of  $\Gamma_0^{\times}$  distinguished vertices and the edges attached to an distinguished vertex will be called *distinguished edges*. To emphasise that a vertex v in  $\Gamma$  is distinguished, we replace v by a decorated vertex  $v^{\times}$ .

**Example 4.2.** Consider the graph in Figure 2 where  $\mathfrak{m}(v_i) = 1$  for  $i \in \{0, 1, 4\}$  and  $\mathfrak{m}(v_i) \in \mathbb{N}$  for  $j \in \{2, 3\}.$ 



FIGURE 2. Skew-Brauer graph with two distinguished vertices.

This is a skew-Brauer graph with two distinguished vertices  $v_1^{\times}$  and  $v_0^{\times}$ . We have  $\mathfrak{o}(v_0) = (1)$ ,  $\mathfrak{o}(v_1) = (2), \mathfrak{o}(v_2) = (1, 2, 3, 4), \mathfrak{o}(v_3) = (3, 4, 5) \text{ and } \mathfrak{o}(v_4) = (5).$ 

Following [\[Sch18\]](#page-0-0), given a Brauer graph Γ, we define the Brauer quiver  $Q_{\Gamma} = (Q_0, Q_1)$  as follows:

- (1) The set of vertices  $Q_0$  is in bijection with the edges of  $\Gamma$ .
- (2) There is an arrow  $x \to y$  each time x and y are edges incident to a vertex  $v \in \Gamma_0$  and y is the immediate successor of x in the cyclic order  $\mathfrak{o}(v)$ .
- (3) For each non-distinguised vertex v of Γ with  $\mathfrak{m}(v)$ val $(v) \geq 2$  gives rise an oriented cycle  $C_v: x_1 \xrightarrow{\alpha_1} x_2 \to \cdots x_t \xrightarrow{\alpha_t} x_1$  in Q, up to cycle equivalence, established by the orientation  $\mathfrak{o}$ , i. e. if  $\mathfrak{o}(v) = (x_1, \ldots, x_t)$  (that we will set as clockwise in our examples when  $t > 1$ ). We say that  $C_v$  is a special cycle at v.

We define the skew-Brauer quiver  $Q_{\Gamma^{\times}}$  from a graph  $\Gamma^{\times}$  as a pair  $Q_{\Gamma^{\times}} = (Q_{\Gamma}, Sp)$  where  $Q_{\Gamma}$  is the Brauer quiver of  $\Gamma$  and  $Sp$  is the set of distinguished vertices that is in bijection with the set  $\Gamma_0^{\times}$ . As before, to emphasise that a vertex v is distinguished in  $Q_{\Gamma^{\times}}$ , we replace v by a decorated vertex  $v^{\times}$ .

Finally, given a skew-Brauer quiver  $Q_{\Gamma^{\times}}$ , we define its *admissible skew-Brauer quiver*  $Q_{\Gamma^{\times}}^{sg}$  as the one constructed from  $Q_{\Gamma}$ × following the rules described in Definition 3.1.

Following the previous definitions, the skew-Brauer graph in Figure 2 defines the skew-Brauer quiver  $Q_{\Gamma^{\times}}$  depicted in Figure 3, and its admissible skew-Brauer quiver coincides with the quiver given in Figure 1.



FIGURE 3. Skew-Brauer quiver with two distinguished vertices

Note that given a skew-Brauer graph  $\Gamma^{\times}$ , there is exactly one special cycle  $C_v$  in  $Q_{\Gamma^{\times}}$  for every  $v \in \Gamma_0$  with  $\mathfrak{m}(v)$ val $(v) \geq 2$ , up to cycle permutation, and each special cycle  $C_v$  induces  $2^r$  cycles  $\{C_{v,1}, C_{v,2}, \ldots, C_{v,2^r}\}\$  in  $Q_{\Gamma^{\times}}^{sg}$ , where r is the number of special edges attached to v, which we call sg-special cycles at v. Each one of the cycles  $C_{v,l}$  and  $C_v$  are considered up to cyclic permutation.

4.1. Skew-Brauer algebras and a set of relations. Let  $\Gamma^{\times}$  be a skew-Brauer graph, where  $Q_{\Gamma^{\times}}$  is its skew-Brauer quiver with distinguished set of vertices  $Sp$ , and let  $Q_{\Gamma^{\times}}^{sg}$  be its admissible skew-Brauer quiver. We define the ideal  $I_{\Gamma^{\times}}$  in  $KQ_{\Gamma^{\times}}^{sg}$  as the one generated by the following four types of relations.

Relations of type 0: Difference of two paths of the form

$$
({}^\varepsilon\alpha^+)({}^+\beta^{\varepsilon'})-({}^\varepsilon\alpha^-)({}^-\beta^{\varepsilon'})
$$

for all possible  $\varepsilon = \{+, -, \emptyset\}$  and  $\varepsilon' = \{+, -, \emptyset\}.$ 

Relations of type I: Difference of powers of two sg-special cycles  $C_{v,l}$  and  $C_{v',l'}$  at vertices v and v' of Γ

$$
C_{v,l}^{\mathfrak{m}(v)}-C_{v',l'}^{\mathfrak{m}(v')}
$$

where both cycles start at a vertex  $y \in Q_{\Gamma^\times}^{sg}$  with  $\mathfrak{m}(v)$  val $(v) \neq 1$  and  $\mathfrak{m}(v')$  val $(v') \neq 1$ .

Relations of type IIa: Paths of the form

$$
C_{v,l}^{\mathfrak{m}(v)} \alpha
$$

where  $\alpha$  is the first arrow of the sg-special cycle  $C_{v,l}$ .

Relations of type IIb: Paths of the form

$$
C_{v,l}^{\mathfrak{m}(v)-1}(\epsilon_1 \alpha_1^{\epsilon_2}) \cdots (\epsilon_r \alpha_r^{\epsilon_{r+1}})
$$

for each special cycle  $C_v = \alpha_1 \cdots \alpha_r$  in  $Q_{\Gamma} \times$  such that  $s(\alpha_1)$  is a distinguished vertex of  $Q_{\Gamma} \times$ , for all possibles  $\varepsilon_k = \{+, -, \emptyset\}$  where  $k = 1, ..., r + 1$  with  $\varepsilon_1 \neq \varepsilon_{r+1}$ .

Relations of type III: Paths of length two of the form

αβ

where  $\alpha\beta$  is not a sub-path of any sg-special cycle  $C_{v,l}$ , except when  $\alpha = \beta$  and  $\alpha$  is a loop induced by a vertex v in  $\Gamma^{\times}$ , i.e. where val $(v) = 1$  and  $\mathfrak{m}(v) > 1$ .<sup>[4](#page-12-0)</sup>

**Definition 4.3.** Let  $\Gamma^{\times} = (\Gamma'_0, \Gamma_0^{\times}, \Gamma_1, \mathfrak{m}, \mathfrak{o})$  be a skew-Brauer graph. The algebra  $B_{\Gamma^{\times}} = KQ_{\Gamma^{\times}}^{sg}/I_{\Gamma^{\times}}$ is called the *skew-Brauer algebra* associated to the skew-Brauer graph  $\Gamma^{\times}$ .

Remark 4.4. We note that any skew-Brauer graph  $\Gamma^\times = (\Gamma_0', \Gamma_0^{\times}, \Gamma_1, \mathfrak{m}, \mathfrak{o})$  with  $\Gamma_0^\times = \emptyset$  is a Brauer graph. Therefore, in that case, the definition of skew-Brauer graph algebras coincides with the definition of Brauer graph algebras given in [\[Sch15\]](#page-0-0).

**Theorem 4.5.** Skew-Brauer graph algebras (with arbitrary multiplicity function  $\mathfrak{m}$ ) are finite dimensional and symmetric.

*Proof.* To show that  $B_{\Gamma^{\times}}$  is a finite dimensional algebra, it is enough to show that the ideal  $I_{\Gamma^{\times}}$  is admissible. Since any sg-special cycle  $C_{v,l}$  is either a cycle of length at least 2 or a loop surroundings a vertex v with multiplicity function  $\mathfrak{m}(v) > 1$ , we have that  $I_{\Gamma^{\times}} \subset J^2$ , where  $J^2$  denotes the ideal generated by paths of length 2.

Moreover, it follows from the relations of type I, IIa, IIb, and III, that any path of length at least  $N+1$ , where

$$
N = \max\left\{\text{length } C_{v,l}^{\mathfrak{m}(v)}\right\},\
$$

is an element of the ideal  $I_{\Gamma^{\times}}$ , therefore  $I_{\Gamma^{\times}}$  is admissible, as we claimed.

<span id="page-12-0"></span> $^{4}$ See Examples 2.2 and 2.3 (1) in [\[Sch18\]](#page-0-0).

By [\[Yam96\]](#page-0-0), to prove that  $B_{\Gamma^{\times}}$  is symmetric it is enough to prove that the K-lineal form  $\phi: B_{\Gamma^{\times}} \to K$ given by

$$
\phi(x) = \begin{cases} 1 & \text{if } x = C_{v,l}^{\mathfrak{m}(v)} \text{ is the } \mathfrak{m}(v)\text{-power of a sg-special cycle for some } v \in \Gamma^{\times} \\ 0 & \text{otherwise} \end{cases}
$$

satisfies  $\phi(ab) = \phi(ba)$  for any paths a, b in  $B_{\Gamma^{\times}}$ .

Let a and b be any two paths in  $B_{\Gamma}$ × such that ab is not zero. By relations of type IIa, IIb, and III, we notice that a, b and ab is a sub-path of the  $\mathfrak{m}(v)$ -power of a sg-cycle  $C_{v,l}^{\mathfrak{m}(v)}$  for some vertex  $v \in \Gamma^{\times}$ . If  $ab = C_{v,l}^{\mathfrak{m}(v)}$ , then  $t(ab) = t(b) = s(a)$ , and as a consequence ba is a cyclic permutation of  $C_{v,l}^{\mathfrak{m}(v)}$ , then  $\phi(ba) = 1 = \phi(ab)$ .

It is easy to see that if ab is not the  $\mathfrak{m}(v)$ -th power of any sg-special cycle  $C_{v,l}$ , then  $\phi(ab) = 0$  $\phi(ba)$ .

Remark 4.6. The following example shows a simple way to compute the indecomposable projective modules of skew-Brauer graph algebras corresponding to vertices of  $Q_{\Gamma^{\times}}^{sg}$  using its associated Brauer graph. It also shows that skew-Brauer graph algebras are not special biserial.

**Example 4.7.** Let  $\Gamma^{\times} = (\Gamma_0', \Gamma_0^{\times}, \Gamma_1, \mathfrak{m}, \mathfrak{o})$  be the skew-Brauer graph of the Example 4.2, and  $\Gamma = (\Gamma_0' \sqcup \Gamma_0^{\times}, \Gamma_1, \mathfrak{m}, \mathfrak{o})$  its associated Brauer graph. For simplicity we set  $\mathfrak{m}(v_i) = 1$  for every  $i = 0, 1, 2, 3, 4, 5$ . Following [\[Sch18\]](#page-0-0), one can compute the indecomposable projective  $B_{\Gamma}$ -module  $P_i$  of each vertex  $i \in Q_{\Gamma}$  from  $\Gamma$ , for example, the indecomposable modules  $P_2$  and  $P_3$  are as follows



Since the vertex 2 corresponds to a distinguished vertex in  $Q_{\Gamma^{\times}}$ , then there are two indecomposable projectives  $B_{\Gamma}$  ×-modules  $P_2^+$  and  $P_2^-$ , both of them are obtained from  $P_2$  by replacing every distinguished vertex *i* in their composition series of  $rad(P_2)/soc(P_2)$  by  $i^+$  and  $i^-$  and adding the respective linear maps induced by the arrows in  $Q_{\Gamma}^{sg}$ , and then replacing the socle and the top by 2<sup>+</sup> and 2<sup>−</sup> respectively. While, for  $P_3$  there is only one projective  $B_{\Gamma} \times$ -module  $\widehat{P_3}$  which is obtained from  $P_3$  by replacing every distinguished vertex i in their composition series of  $rad(P_2)/soc(P_2)$  by  $i^+$  and  $i^-$  and considering their respective linear maps induced by the arrows in  $Q_{\Gamma}^{sg}$ . Then the projectives  $P_2^+, P_2^$ and  $P_3$  are as follows.



4.2. The trivial extension of a skew-gentle algebra. In this section we state a relation between trivial extension of skew gentle algebras and a subfamily of skew-Brauer graph algebras. To do that, we follow the construction of the generalized ribbon graph given in [\[LFSV20,](#page-0-0) Remark 3.4].

Given a  $A = KQ/I$  skew-gentle algebra with a set of special vertices  $Sp$ , we say that a path p in Q is sp-maximal if  $p\alpha = \alpha p = 0$  in A for every arrow  $\alpha$  in Q that is not a special loop and if  $s(p)$  (or  $t(p)$ ) is a special vertex then the first arrow ( resp. or the last arrow) is the special loop attached to it.

We define a skew-Brauer graph  $\Gamma_A^\times = \Gamma^\times$  as follows.

- The set of vertices  $\Gamma'_0$  of  $\Gamma^{\times}_A$  is in bijection with the union of
	- i) all sp-maximal paths,
	- ii) trivial paths  $e_x$  such that x is either the source or the target of only one arrow or x is the target of exactly one arrow  $\alpha$  and the source of exactly one arrow  $\beta$ , and  $\alpha\beta \neq 0$  in A.
- The set of distinguished vertices  $\Gamma_0^{\times}$  is in bijection with trivial paths  $e_x$  such that x is a special vertex of Q.
- The set of edges of  $\Gamma_A^{\times}$  is in bijection with the vertices of  $Q_0$  (note that this includes the special vertices).

The multiplicity function  $\mathfrak{m}$  is the constant function equal to 1, and the cyclic ordering of the edges attached to each vertex  $v_0 \in \Gamma_0$  is given either by the order of the sp-maximal associated to  $v_0$  or  $\mathfrak{o}(v) = (e_x)$  otherwise.

**Lemma 4.8.** Let A be a skew-gentle algebra with Sp the set of special vertices and  $A'$  be its auxiliary gentle algebra. Then the sp-maximal paths of the skew-gentle algebra A are in bijection with the maximal paths of the auxiliary gentle algebra  $A'$ .

*Proof.* Let  $\rho = \alpha_1 \dots \alpha_k$  be a sp-maximal path in A and  $\rho' = \beta_1 \dots \beta_k$  be the path in A' where  $\beta_i = \alpha_i$ if  $\alpha_i$  is not a special loop and  $\beta_i = e_{t(\alpha_{i-1})}$  otherwise. We claim that  $\rho'$  is a maximal path in A'.

Notice that  $\rho' \neq 0$ , otherwise  $\beta_i \beta_{i+1}$  is a zero relation in A' for some i, which implies that  $\beta_i = \alpha_i$ and  $\beta_{i+1} = \alpha_{i+1}$  and  $\alpha_i \alpha_{i+1}$  is a zero relation in A, as a consequence  $\rho$  is not a sp-maximal path.

Suppose  $\rho'$  is not a maximal path in A'. Then there exist at least one arrow in  $Q_{A'}$  such that  $\alpha \rho' \neq 0$ or  $\rho' \alpha \neq 0$  in A'. Without loss of generality, suppose that  $\alpha \rho' \neq 0$  in A'. We notice that if the start  $s(\rho) = x$  of  $\rho$  is not a special vertex, then  $\alpha$  is not a special loop and  $\alpha$  is an arrow of  $Q_A$ , then  $\alpha \rho$ is not zero in  $A$ , a contradiction with the hypothesis. Now, if  $x$  is a special vertex, then the first arrow  $\alpha_1$  of  $\rho$  is the special loop attached to x and  $\alpha \rho$  is not zero in A, again a contradiction with the hypothesis. □

**Theorem 4.9.** Let A be a skew-gentle algebra and  $\Gamma_A^{\times} = (\Gamma_0', \Gamma_0^{\times}, \Gamma_1, \mathfrak{m} = 1, \mathfrak{o})$  be its skew-Brauer graph. Then  $T(A^{sg}) \simeq B_{\Gamma_A^{\times}}$ .

*Proof.* Let  $A'$  be the auxiliary gentle algebra of  $A$ . As a consequence of Lemma 4.8, is it easy to check that  $\Gamma_A^{\times}$  as Brauer graph, nameley, the Brauer graph  $\Gamma = (\Gamma_0' \cup \Gamma_0^{\times}, \Gamma_1, \mathfrak{m}, \mathfrak{o})$  coincides with the Brauer graph  $\Gamma_{A'}$ .

By [\[Sch15,](#page-0-0) Theorem 1.2], the trivial extension  $T(A')$  of  $A'$  is isomorphic to the Brauer graph algebra  $B_{\Gamma_{A'}}$  and moreover, by Theorem 3.11 we have that  $T(A)$  is isomorphic to  $KQ_{T(A')}^{sg} / I_{T(A')}^{sg}$ . It is easy to check that the last presentation is isomorphic to the skew-Brauer graph algebra  $B_{\Gamma_A^{\times}}$  described in Section 4.1. Therefore, the statement follows.

To prove the second theorem of this section, we need to recall more definitions. Following [\[FP02\]](#page-0-0), let A be an algebra and let  $T(A)$  be the trivial extension of A. Recall that a cycle C in  $Q_{T(A)}$  is called *elementary cycle* if  $C = \alpha_j \dots \alpha_1 \beta_p \alpha_n \dots \alpha_{j+1}$ , with  $\alpha_1, \dots, \alpha_n \in (Q_A)_1$  and  $\beta_p$  the new arrow of  $T(A)$ , corresponding to the element p of soc<sub>Ae</sub> A.

Let D be a set of arrows of  $Q_{T(A)}$ . We say that D is an *admissible cut* of  $Q_{T(A)}$  if it consists of exactly one arrow of each elementary cycle in  $Q_{T(A)}$ . Moreover, we say that the algebra B is an *admissible* cut of  $T(A)$  if it is the quotient algebra  $T(A)/\langle \mathcal{D} \rangle$ , where  $\mathcal D$  is an admissible cut of  $Q_{T(A)}$ . We say that  $T(A)/\langle \mathcal{D} \rangle$  is the quotient algebra of  $T(A)$  induced by  $\mathcal{D}$ .

Notice that the definition of admissible cut is given for trivial extensions of path algebras. However, it is possible to generalize this idea to give a recognition theorem of trivial extension, see  $[FT^+22]$ for details. To do this, given an algebra A we consider a finite set  $\mathcal E$  of non-zero cycles of length at least two, then a *cut set*  $\mathcal D$  of A with respect to  $\mathcal E$  is a subset of arrows  $\{\alpha_1,\ldots,\alpha_k\}$  of  $Q_A$  containing exactly one arrow in each cycle of  $\mathcal{E}$ , and such that  $\alpha_i$  appears at most once in each cycle C of  $\mathcal{E}$ . This definition corresponds to definition of *allowable cut* given in  $[FT^+22]$ .

Remark 4.10. If A is a trivial extension of an algebra  $B$ , then any admissible cut is indeed a cut set with respect to the set of the elementary cycles.

**Theorem 4.11.** Let K be an algebraically closed field and let  $\Gamma^{\times} = (\Gamma_0, \Gamma_0^{\times}, \Gamma_1, \mathfrak{m}, \mathfrak{o})$  be a skew-Brauer graph with  $\mathfrak{m}(v) = 1$  for every vertex v in  $\Gamma_0$  and  $B_{\Gamma^\times} = KQ_{\Gamma^\times}/I_{\Gamma^\times}$  be its associated skew-Brauer graph algebra. Then there exists (not necessarily unique) skew-gentle algebra A, such that  $T(A^{sg}) \simeq B_{\Gamma^{\times}}$ .

*Proof.* Let  $\Gamma^{\times} = (\Gamma_0, \Gamma_0^{\times}, \Gamma_1, \mathfrak{m}, \mathfrak{o})$  be the skew-Brauer graph with  $\mathfrak{m}(v) = 1$  for every vertex v in  $\Gamma_0$ and  $B_{\Gamma^{\times}}$  be a skew-Brauer algebra with  $\mathcal E$  the set of sg-special cycles and  $Sp$  the set of distinguished vertices.

Consider  $B_{\Gamma}$  be the Brauer graph algebra with  $\Gamma = (\Gamma_0 \cup \Gamma_0^{\times}, \Gamma_1, \mathfrak{m}, \mathfrak{o})$  the Brauer graph obtained from  $\Gamma^{\times}$  by forgetting the extra information of distinguished vertices. We notice that  $Q_{\Gamma}$  is also obtained from  $Q_{\Gamma^{\times}} = (Q_{\Gamma}, Sp)$  by forgetting the extra information of Sp. We claim that any admissible cut of  $B_{\Gamma}$  induces a cut set of  $B_{\Gamma^{\times}}$ .

By [\[Sch15\]](#page-0-0), there exists an admissible cut  $\mathcal D$  on  $B_\Gamma$  such that  $B_\Gamma/\langle \mathcal D \rangle$  is gentle. Let  $\mathcal D'$  be the set of arrows

$$
\mathcal{D}' = \{^{\varepsilon} \alpha^{\varepsilon'} \mid \alpha \in \mathcal{D} \quad \text{for all possible} \quad \varepsilon, \varepsilon' \in \{+, -, \emptyset\} \}
$$

in  $B_{\Gamma^{\times}}$ .

Recall that by definition of  $B_{\Gamma^{\times}}$ , the set of sg-special cycles E are induced by the set of special cycles of  $B_{\Gamma_A}$ . To fix notation, for each special cycle  $C_v$  of  $Q_{\Gamma}$ , denote by  $\{C_{v,1}, \ldots C_{v,2^r}\}$  the set of  $2^r$ sg-special cycles in  $Q_{\Gamma^\times}^{sg}$  (up to cycle equivalence).

Suppose that there is a sg-elementary cycle  $C_{v,l} = \epsilon_1 \alpha_1^{\epsilon'_1} \dots \epsilon_m \alpha_m^{\epsilon'_m}$  that contains two different arrows  $\epsilon_i \alpha_i^{\epsilon'_i}$  and  $\epsilon_j \alpha_j^{\epsilon'_j}$  of  $\mathcal{D}'$ . Then by construction the cycle  $C_v = \alpha_1 \dots \alpha_m$  is special and  $\alpha_i$  and  $\alpha_j$  are arrows of D. Since D is an admissible cut, then  $\alpha = \alpha_i = \alpha_j$ , and because  $B_{\Gamma}/\langle D \rangle$  is gentle  $\alpha$  appears exactly once in  $C_v$ , this implies that  ${}^{\varepsilon_i} \alpha_i^{\varepsilon'_i} = {}^{\varepsilon_j} \alpha_i^{\varepsilon'_j}$ .

To finish the proof, it is enough to observe that the algebra  $B_{\Gamma^{\times}}/\langle \mathcal{D}'\rangle$  is the admissible version of  $B_{\Gamma}/\langle \mathcal{D} \rangle$  with Sp the set of distinguished vertices of  $B_{\Gamma} \times$ .

Motivated by the cut sets of a skew-Brauer algebra  $B_{\Gamma^{\times}}$  used in Theorem 4.11, which are induced by cut sets of its auxiliary Brauer algebra  $B_{\Gamma}$ , we introduce the following definition.

**Definition 4.12.** Let  $\Gamma^{\times} = (\Gamma_0, \Gamma_0^{\times}, \Gamma_1, \mathfrak{m}, \mathfrak{o})$  be a skew-Brauer graph with  $\mathfrak{m}(v) = 1$  for every v in  $\Gamma_0$  and  $B_{\Gamma^{\times}} = KQ_{\Gamma^{\times}}/I_{\Gamma^{\times}}$  be its associated skew-Brauer graph algebra. A cut set  $\mathcal{D}$  is good if it is induced by a cut set  $\mathcal{D}'$  of its auxiliary Brauer-graph algebra  $B_{\Gamma}$ , namely

$$
\mathcal{D} = \{^{\varepsilon} \alpha^{\varepsilon'} \mid \alpha \in \mathcal{D}' \text{ for all possible } \varepsilon, \varepsilon' \in \{+, -, \emptyset\} \}
$$

We observe that any good cut set of a skew-Brauer graph algebra induces a skew-gentle algebra. We show that converse is not true in the following example.

**Example 4.13.** In this example, we show a cut set  $D$  of skew-Brauer algebra that is not good, but the quotient algebra induced by  $D$  is skew-gentle.

Let  $B = kQ_{\Gamma^{\times}}/I_{\Gamma^{\times}}$  be a skew-Brauer algebra with quiver  $Q_{\Gamma^{\times}}$  and skew-Brauer graph  $\Gamma^{\times}$  that are depicted below

$$
Q_{\Gamma^\times}: \qquad \xrightarrow{\begin{array}{c} f_1 \\ \downarrow \end{array}} 1^\times \xrightarrow{\alpha \qquad \qquad \beta \qquad \qquad } 2 \xrightarrow{\beta \qquad \qquad } 3^\times \qquad \qquad \Gamma^\times: \quad v_1^\times \longrightarrow v_2 \longrightarrow v_3 \longrightarrow v_4^\times
$$

To compute the cut set, see that the admissible presentation  $B^{sg} = K Q_{\Gamma^{\times}}^{sg} / I_{\Gamma^{\times}}^{sg}$  of the algebra B is given by quiver  $Q_{\Gamma^\times}^{sg}$ 



with sg-special cycles  $({}^+\alpha)(\gamma^+),({}^-\alpha)(\gamma^-),(\beta^+)({}^+\delta),(\beta^-)({}^-\delta)$  and their permutations, and the ideal  $I_{\Gamma^{\times}}^{sg}$  is generated by all relations

- (1)  $C C'$ , where C and C' are sg-special cycles that start in the same vertex, for example  $(\gamma^+)(^+\alpha) - (\beta^-)(^-\delta);$
- (2) the paths of length 2 that start and end in two different vertices of the set  $\{1^+, 1^-, 3^+, 3^-\}$ , for example  $({}^{\dagger}\alpha)(\beta^-);$
- (3) the paths of length 3 of the form  $C\eta$ , where C is a sg-special cycle, and  $\eta$  is the first arrow of C, for example,  $(\gamma^+)(^{\dagger}\alpha)(\gamma^+);$
- (4) and finally, the paths of length 3 of the form  $(\eta_1^{\varepsilon})(\eta_2)(\eta_1^{\varepsilon'})$  where  $(\eta_1^{\varepsilon})(\varepsilon\eta_2)$  is a sg-special cycle that starts at 2,  $\varepsilon$ ,  $\varepsilon'$  are either  $+$ ,  $-$  and  $\varepsilon \neq \varepsilon'$ , for example  $(\gamma^+)(^+\alpha)(\gamma^-)$ .

Consider  $\mathcal{D} = \{^+ \alpha, \gamma^-, ^+ \delta, \beta^- \}$  a cut set of  $B^{sg}$ . We notice that the quotient algebra of  $B^{sg}$  induced by D is the algebra  $A = KQ/I$  where the quiver Q is and I is the ideal generated by the relations:



$$
(^{-\alpha})(\gamma^{+}), (-\alpha)(\beta^{+}), (-\delta)(\gamma^{+}), (-\delta)(\beta^{+}).
$$

Finally, A is isomorphic to the skew-gentle algebra  $A' = KQ'/I'$  where Q' is the quiver

$$
\begin{array}{c}\n f_1 \\
f_2 \\
1 \xrightarrow{\qquad \qquad } 1 \\
\hline\n a \rightarrow 2 \xrightarrow{\qquad \qquad } 3\n\end{array}
$$

and  $I' = \langle \alpha \beta, f_1^2 - f_1, f_2^2 - f_2 \rangle$ .

### 5. Admissible cuts, the repetitive algebra and reflections

In this section we consider the non-admissible presentation of a skew-gentle algebra, and we describe the quiver of its repetitive algebra. We also describe the positive and negative reflections on a skewgentle algebra. Moreover, we analyze the good cuts over a skew-gentle algebra and the relationship between these two notions.

The following definition of repetitive algebra for skew-gentle algebras, in their non-admissible presentation, is an extension of the one given for basic algebras by Hughes and Waschbüsch.

**Definition 5.1.** Let  $A = KQ/I$  be a skew-gentle algebra with Sp the set of special vertices, S be the set of special loops and M be the set of sp-maximal paths. The repetitive algebra  $\hat{A}$  of A is  $\widehat{A} = K\widehat{Q}/\widehat{I}$  where  $\widehat{Q}$  and  $\widehat{I}$  are defined as follows

- The set of vertices of  $\widehat{Q}$  is the set  $\{x[n] : x \in Q_0 \text{ and } n \in \mathbb{Z}\}\)$ . We say that  $x[n]$  is a special vertex in  $\widehat{Q}$  if an only if x is a special vertex in  $Q$  and  $n \in \mathbb{Z}$ .
- The set of arrows  $\widehat{Q}_1$  is the set  $\{\alpha[n] : x[n] \to y[n] : \alpha \in Q_1 \text{ and } n \in \mathbb{Z}\}.$
- Each sp-maximal path  $p: x \to y$  induces a set of *connecting arrow*  $\overline{p}[n]: y[n] \to x[n+1]$  for all  $n \in \mathbb{Z}$ .

To define the set of relation  $\hat{\rho}$  of  $\hat{Q}$ , we need to introduce the following definition. If  $p = p_q p_2$  is a sg-maximal path in  $(Q, I)$ , then  $p_2[n]\overline{p}[n]p_1[n + 1]$  is a sg-full path in  $\widehat{Q}$ .

The set of relations  $\widehat{\rho}$  of  $\widehat{Q}$  are given by the following list.

- (1) Each monomial relation  $p \in \rho$  induces a path  $p[n]$  in  $\hat{\rho}$ , for all  $n \in \mathbb{Z}$ .
- (2) Each relation  $f^2 f$  in  $\rho$  induces a relation  $f^2[n] f[n]$  in  $\hat{\rho}$ , for all  $n \in \mathbb{Z}$ .<br>(2) Are nothing  $\hat{\rho}$  which contains a connection arrange is an almost of  $\hat{\gamma}$  if w
- (3) Any path p in  $\hat{Q}$  which contains a connecting arrow is an element of  $\hat{\rho}$  if p is nor a sub-path of a sg-full subpath.

**Lemma 5.2.** Let  $A = KQ/I$  be a skew-gentle algebra and denote by  $A^{sg}$  its admissible presentation. Let A' be the auxiliary gentle algebra of A. Then, the quiver  $Q_{\widehat{A}sg}$  of the repetitive algebra of  $A^{sg}$ coincides with the sg-quiver  $Q_{\widetilde{\mathcal{M}}}^{sg}$  $\frac{sq}{A'}$  of the repetitive algebra of A', considering the set of distinguished vertices as the set of special vertices of A.

*Proof.* It follows from Lemma 3.9 and that the sq-maximal paths in  $(Q, I)$  are in correspondence with maximal paths in  $(Q', I')$ , where  $KQ'/I'$  is the auxiliary gentle algebra A' . □

**Example 5.3.** Let  $A = KQ/I$  be the skew-gentle algebra with quiver

$$
\begin{array}{ccc}\n f_1 & & f_2 \\
\searrow & & \\
1 & \xrightarrow{\alpha} & 2 & \xrightarrow{\beta} & 3\n \end{array}
$$

and ideal  $I = \langle \alpha \beta, f_1^2 - f_1, f_2^2 - f_2 \rangle$ .

To compute the repetitive algebra of A, we need to find the  $sg$ -maximal paths in  $(Q, I)$ . In this case, there are exactly two sg-maximal path, namely  $p_1 = f_1 \alpha$  and  $p_2 = \beta f_2$ . Then by the definition, the quiver  $Q_{\widehat{A}}$  of repetitive algebra of the algebra A is the following.







One can check that the set of relations coincides with the relations induced by the sg-ideal of the repetitive algebra  $A$ .

Next result shows when reflections on a skew-gentle algebra returns a skew-gentle algebra.

**Theorem 5.4.** Let  $A = KQ/I$  be a skew-gentle algebra, with Sp the set of special vertices, and x be a source in its auxiliary gentle algebra  $A'$ . If x is a special vertex (resp. non special vertex), then the negative reflection sequence  $S_{x+}^{-}S_{x-}^{-}(A^{sg})$  (resp. the negative reflection  $S_{x}^{-}(A^{sg})$ ) is skew-gentle.

A similar result follows when x is a sink in A′ if we consider positive reflections.

*Proof.* Let x be a source of  $Q_0$ . If  $x \in Sp$  or  $x \notin Sp$ , in both cases it follows from Lemma 5.2 that the sg-quiver  $(\sigma_x Q_{A'})^{sg}$  coincides with the quivers  $\sigma_{x+} \sigma_{x-} Q_{A^{sg}}$  and  $\sigma_x Q_{A^{sg}}$ , respectively. It is easy to see that the algebra  $S_x^{-}(A')$  is a gentle algebra, where x is a sink vertex for its ordinary quiver. Therefore the sg-algebra  $(S_x^-(A'))^{sg}$  is skew-gentle, which is isomorphic to  $S_{x^+}S_{x^-}(A)$  and  $S_x(A)$ , respectively.  $\Box$ 

The following example shows that certain sequences of reflections over a skew-gentle algebra, not defined as in the previous theorem, can also result on a skew-gentle algebra.

**Example 5.5.** Let  $A = KQ/I$  be the skew-gentle algebra from Example 5.3.

By the Theorem 5.4, the algebras  $S_{1+}^- S_{1-}^- (A^{sg})$  and  $S_{3+}^+ S_{3-}^+ (A^{sg})$  are skew-gentle. We note that the algebra  $S_3^+$ ,  $S_1^-$  ( $A^{sg}$ ) is also skew-gentle, but it is not obtained by a sequence of reflections as described in Theorem 5.4.

To show the link between reflections and admissible cuts, we need to use the following result, which is part of a work in progress  $\text{[ABF]}$ . For the benefit of the reader, we include the proof of this result.

**Lemma 5.6** ( $\widehat{ABF^+}$ ). Let  $A = KQ/I$  be a gentle algebra,  $T(A)$  its trivial extension, and x a source vertex of A. Then, there exists an admissible cut  $D$  of  $B$  such that:

- (1)  $S_x^-(A)$  is isomorphic to the quotient  $T(A)/\langle \mathcal{D} \rangle$ .
- (2) the admissible cut  $\mathcal D$  contains the arrows in Q that start at x; in other words,  $\mathcal D \cap Q_1 = {\alpha \in \mathbb N}$  $Q_1 | s(\alpha) = x$ .

*Proof.* Let x be a source vertex in A. We denote by  $B = S_x^{-}(A)$  the negative reflection of A. By [\[HW83\]](#page-0-0), we have that the trivial extension algebra  $T(A)$  of A coincides with the trivial extension algebra  $T(B)$  of B, and moreover, by [\[FST](#page-0-0)+22], there exists an admissible cut D of  $T(A)$  such that  $B=T(A)/<\mathcal{D}>.$ 

To prove the second statement, let  $\gamma : y \to z$  be an element of  $\mathcal{D} \cap (Q)_1$ . Suppose  $s(\gamma) \neq x$ . By construction,  $(\gamma, 0) : (y, 0) \to (z, 0)$  is an arrow of  $\sigma_x^{-}(Q_A) = Q_B$ , contradicting the fact that  $\gamma$  is in D. Therefore  $s(\gamma) = x$ .

Conversely, if  $\gamma$  is an arrow of  $Q_1$  such that  $s(\gamma) = x$ , since the vertex x is a sink of Q we have that  $\gamma \notin (Q_B)_1$  and consequently  $\gamma \in \mathcal{C}$ .

**Theorem 5.7.** Let  $A = KQ/I$  be a skew-gentle algebra with Sp as the set of special vertices and let x be a source vertex in the auxiliary gentle algebra  $A'$  of  $A$ . If x is a special vertex (resp. x is not a special vertex), then,  $S_{x+}^{-}S_{x-}^{-}(A^{sg}) = T(A^{sg})/\langle \mathcal{D} \rangle$  (resp.  $S_{x}^{-}(A^{sg}) = T(A^{sg})/\langle \mathcal{D} \rangle$ ) where  $\mathcal D$  is the good cut of  $T(A^{sg})$  induced by the cut set  $\mathcal{D}'$  such that

$$
\mathcal{D}' \cap (Q_{A'})_1 = \{ \alpha \in (Q_{A'})_1 \mid s(\alpha) = x \}.
$$

*Proof.* Let x be a source vertex in  $A'$  and suppose that x is a special vertex of A. We denote by  $B = S_{x+}^{-} S_{x-}^{-} (A^{sg})$  the negative reflection sequence of  $A^{sg}$ . By Theorem 5.4 we know that B is a skew-gentle algebra and  $B' = S_x^{-1}(A')$  is the auxiliary gentle algebra of B.

By Lemma 5.6, there exists an admissible cut  $\mathcal D$  of A' such that  $\mathcal D' \cap (Q_{A'})_1 = \{ \alpha \in (Q_{A'})_1 \mid s(\alpha) = x \},$ which induces a good cut set  $\mathcal D$  in  $T(A^{sg})$ . The result follows from Theorem 5.4.

#### 6. Skew-Brauer graph algebras of finite representation type

In this section, we show that the class of skew-Brauer graph algebras of finite representation type are characterized by the structure of the skew-Brauer graph as a graph and the number of distinguished vertices on it. In the remainder of this section, we consider non-trivial skew-Brauer graphs  $\Gamma^{\times}$ , that is with at least one distinguished vertex.

**Definition 6.1.** A skew-Brauer graph  $\Gamma^{\times} = (\Gamma_0, \Gamma_0^{\times}, \Gamma_1, \mathfrak{m}, \mathfrak{o})$  is a *skew-Brauer tree* if the underline graph  $(\Gamma_0 \cup \Gamma_0^{\times}, \Gamma_1)$  is a tree graph,  $\mathfrak{m}(v) = 1$  for all  $v \in \Gamma_0 \cup \Gamma_0^{\times}$ , and there is exactly one distinguished vertex in  $\Gamma^{\times}$ . We say that  $B_{\Gamma^{\times}}$  is a skew-Brauer three algebra if  $\Gamma^{\times}$  is a skew-Brauer tree graph.

Recall that a *Brauer tree algebra* is a Brauer algebra defined by a graph  $\Gamma$  that is a tree, with  $\mathfrak{m}(v) = 1$ for all but at most one  $v \in \Gamma_0$ . Since skew-Brauer graph algebras are symmetric, we also recall the classification of symmetric finite representation type algebras.

**Lemma 6.2.** ( $\text{ACH22}$ , Theorem 2.11) A non-simple basic symmetric algebra is of finite representation type if and only if it is isomorphic to an algebra in the following mutually exclusive classes.

- (1) Trivial extensions of iterated tilted algebras of Dynkin type.
- (2) Brauer tree algebras with exactly one vertex v with multiplicity  $\mathfrak{m}(v) > 1$ .
- (3) Modified Brauer tree algebras with exactly one vertex v with multiplicity  $\mathfrak{m}(v) > 1$ .

In the previous lemma, the family of Brauer tree algebras with  $\mathfrak{m}(v) = 1$  for all v coincides with the trivial extensions of iterated tilted algebras of Dynkin type A, see [\[WZ22,](#page-0-0) Lemma 3.1]. The definition of Brauer graph algebras, Brauer tree algebras and modified Brauer tree algebras can be found in [\[Sch18,](#page-0-0) Section 2.3] and [\[Sko06,](#page-0-0) Section 2.8 and Section 3.6]. We notice that modified Brauer tree algebras are not necessarily biserial, see for example the projective-injective  $P_1$  in  $|$ Sko06, Example 3.7].

**Lemma 6.3.** Let  $B_{\Gamma^{\times}} = KQ^{sg}/I^{sg}$  be a skew-Brauer graph algebra of finite representation type, where  $\Gamma^{\times}$  is not a skew-Brauer graph with two edges. Then  $\Gamma^{\times}$  has at most one distinguished vertex.

*Proof.* Let  $B = B_{\Gamma^{\times}}$  be a skew-Brauer graph algebra of finite representation type. Suppose that  $\Gamma^{\times}$ has at least two distinguished vertices  $v^{\times}$  and  $w^{\times}$ . By definition  $\Gamma^{\times}$  is connected, then there exists a minimal path  $\rho$  of edges in  $\Gamma^{\times}$  joining  $v^{\times}$  and  $w^{\times}$  as the one depicted in the following figure.

 $v^{\times}$   $\frac{a_1}{\cdots}$   $x_1$   $\frac{a_2}{\cdots}$   $x_2$   $\frac{a_3}{\cdots}$   $\cdots$   $\frac{a_{k-1}}{k}$   $x_k$   $\frac{a_k}{\cdots}$   $w^{\times}$ 

We denote by  $Q_{\Gamma}$  be the quiver associated to the underline Brauer graph  $\Gamma$  of  $\Gamma^{\times}$ . Recall that there is a unique special cycle  $C_{x_i}$  in  $Q_{\Gamma}$ , up to cycle equivalence, induced by the edges attached to the vertex  $x_i \in \Gamma$ , for  $i = \{1, \ldots, k\}.$ 

For each  $i \in \{1, ..., k\}$ , we define a sub-path  $\rho_i$  of the special cycle  $C_{x_i}$  from the vertex  $a_{i+1}$  to  $a_i$  if i is odd or the sub-path from  $a_i$  to  $a_{i+1}$  if i is even. Then the walk  $\rho_1^{-1}\rho_2 \ldots \rho_k^{(-1)^k}$  $k<sup>(-1)</sup>$  defines a non-linearly oriented sub-quiver  $Q'$  of type  $A_n$  without relations in  $Q_{\Gamma}$  such that the first and the last vertex are distinguished. As a consequence B contains an algebra of Dynkin type  $\tilde{D}_n$  as a sub-algebra, which is a contradiction because B is an algebra of finite representation type.  $\Box$ 

**Lemma 6.4.** ( $[ASS^+06, II.3$  $[ASS^+06, II.3$  Lemma 3.6], [\[Sch18,](#page-0-0) Section 2.3] and [\[Sko06,](#page-0-0) Section 3.6])

- (1) If two basic connected isomorphic algebras are defined by a bound quiver with an admissible ideal, then their quivers coincide.
- (2) Let  $KQ/I$  be a Brauer graph algebra defined by an admissible ideal, then for all  $x \in Q_0$  we have that x is the start (end) of exactly two arrows or x is the start (end) of exactly one arrow.
- (3) Let  $KQ/I$  be a modified Brauer tree algebra defined by an admissible ideal, then for all  $x \in Q_0$ we have that x is the start (end) of exactly two arrows or x the start (end) of exactly one arrow and Q must contain exactly one loop.

It follows from the previous lemma that any bound quiver algebra A cannot be isomorphic to a Brauer graph algebra or a modified Brauer graph algebra if its quiver  $Q_A$  has three or more arrows ending at (or starting from) any of its vertices.

In the following lemma, we describe some properties of a subclass of skew-Brauer graph algebras that are relevant to this section.

**Lemma 6.5.** Let  $B = B_{\Gamma_i^{\times}}$  be a skew-Brauer graph algebra associated with one of the following skew-Brauer graphs.

- $\Gamma_1^{\times}$ :  $\times$   $\longrightarrow$   $v \longrightarrow w$  $\frac{x}{2}$ :  $\times$   $\frac{y}{2}$   $\times$
- (1) If B is either with  $B_{\Gamma_1^{\times}}$  where  $\mathfrak{m}(w) = 1$  or  $B_{\Gamma_2^{\times}}$ , then B is isomorphic to a Brauer graph algebra.
- (2) If  $B = B_{\Gamma_1^{\times}}$  with  $\mathfrak{m}(w) \neq 1$ , then B is not of finite representation type.

*Proof.* By the definition of skew-Brauer graph algebras, the quiver  $Q_B$  of B is one of the quivers depicted in Figure 4: the first two quivers correspond to  $\Gamma_1^{\times}$  with  $\mathfrak{m}(w) = 1$  and  $\mathfrak{m}(m) \neq 1$ , respectively, while the last one corresponds to  $\Gamma_2^{\times}$  with arbitrary  $\mathfrak{m}(v)$ .

$$
1^{+} \xleftarrow{\alpha^{+}} \frac{\alpha^{+}}{\alpha \beta} 2 \xleftarrow{\alpha^{-}} 1^{-}
$$
\n
$$
1^{+} \xleftarrow{\alpha^{+}} \frac{\gamma}{\gamma \beta} 2 \xleftarrow{\alpha^{-}} \frac{\gamma}{\gamma \beta} 1^{-}
$$
\n
$$
1^{+} \xleftarrow{\alpha^{+}} \frac{\gamma}{\gamma \beta} 2 \xleftarrow{\alpha^{-}} \frac{\gamma}{\gamma \beta} 1^{-}
$$
\n
$$
1^{+} \xleftarrow{\alpha^{+}} \frac{\gamma}{\gamma \beta^{+}} 2^{-}
$$
\n
$$
2^{+} \xleftarrow{\alpha^{-}} \frac{\gamma}{\gamma \beta^{+}} 1^{-}
$$

Figure 4. Quivers of skew-Brauer graphs with underline tree graph and 2 edges

To prove the first statement, suppose that B is either  $B_{\Gamma_1^{\times}}$  with  $\mathfrak{m}(w) = 1$  or  $B_{\Gamma_2^{\times}}$ . Let C be any cyclic permutation of the elementary cycle  $({}^+\beta^{\epsilon})({}^{\epsilon}\alpha^+)$  or  $({}^-\beta^{\epsilon})({}^{\epsilon}\alpha^-)$ , for any possible  $\epsilon \in \{+,-,\emptyset\}$ , and let m be a natural number greater than 1. It follows from relations of type 0 and relations of

type III, that any m-power  $C<sup>m</sup>$  of the cycle C is zero in B, and as a consequence one can assume that  $\mathfrak{m}(v) = 1$ . Then we have that B is isomorphic to a Brauer graph algebra  $B_{\Gamma}$  with  $\Gamma$  one of the following two Brauer graphs, respectively.



with  $m(v_i) = 1$  for  $i = 1, 2, 3, 4$ .

Now, suppose that  $B = B_{\Gamma_1^{\times}}$  where  $\mathfrak{m}(w) \neq 1$ . Then the algebra is given by the second quiver in Figure 4 and the ideal

$$
I_{B_{\Gamma_1^{\times}}} = \langle (\alpha^+)(^+\beta) - (\alpha^-)(^-\beta), \gamma^{\mathfrak{m}(w)+1}, \gamma^{\mathfrak{m}(w)} - (\alpha^-)(^-\beta), \gamma^{\mathfrak{m}(w)} - (\alpha^+)(^+\beta),
$$
  

$$
(^+\beta)\gamma, (^+\beta)(\alpha^-), (^-\beta)(\alpha^+), (^-\beta)\gamma, \gamma(\alpha^+), \gamma(\alpha^-)\rangle.
$$

We observe that  $B$  is an algebra of infinite representation type, since there is band module associated to the word  $\gamma^{-1}(\alpha)$  $^{+})(^{+}\beta).$ 

**Lemma 6.6.** Let B be a skew-Brauer graph algebra, and suppose that B is neither  $B_{\Gamma_1^{\times}}$  where  $\mathfrak{m}(w)$  = 1 or  $B_{\Gamma_2^{\times}}$ , where  $\Gamma_i^{\times}$  is one of the following two skew-Brauer graphs.

$$
\Gamma_1^\times \colon \times \text{---} \bullet \text{---} w \qquad \qquad \Gamma_2^\times \colon \times \text{---} \bullet \text{---} \times
$$

Then B is not a Brauer tree algebra.

Proof. Let  $\Gamma^{\times}$  be a skew-Brauer graph with a distinguished vertex  $v^{\times}$ . By definition of skew-Brauer graph, there exists a non-distinguished vertex  $v \in \Gamma^\times$  such that v and  $v^\times$  have exactly one edge x in common. We claim that if v has valency at least three,  $B_{\Gamma^{\times}}$  is not a Brauer graph algebra.

Since v has at least valency three, consider x and  $y = \mathfrak{o}^{-1}(x)$  two different edges attached to v, where y might be a loop. Then by Remark 4.6, there are at most two projectives in  $B_{\Gamma^{\times}}$  associated to y, namely  $P_y$  or  $P_{y+}$  and  $P_{y-}$ , in both cases the projectives are not biserial, see Example 4.7. As a consequence  $B_{\Gamma}$  is not a Brauer graph algebra, and in particular is not a Brauer tree algebra.

Now, we suppose that v has valency two. In this case, x and  $y = \mathfrak{o}^{-1}(x)$  are two different edges attached to v and y is not a loop, otherwise, the valency of v would be grater than 2. We denote by w the vertex in  $\Gamma^{\times}$  such that y is the edge connecting v and w.

By hypothesis, the valency of w is at least two, or if the valency of w is one, then  $\mathfrak{m}(w) \neq 1$ . In the second case, B is the skew-Brauer graph  $B_{\Gamma_1^{\times}}$  with  $\mathfrak{m}(w) \neq 1$ , and by Lemma 6.5 is not of finite representation type, therefore can not be isomorphic to a Brauer tree algebra.

Now we suppose that w has valency at least two. By the definition of the quiver  $Q = Q_{\Gamma^{\times}}$ , the local configuration for the vertex  $y \in Q$  associated to the edge y in  $\Gamma^{\times}$  (here y receives the same name by abuse of notation) is as follows.



where  $\mathfrak{o}(y)$  and  $\mathfrak{o}^{-}(y)$  might be equal. Then there are tree arrows ending at y and, by Lemma 6.4, B is not a Brauer graph algebra as we claim.

**Lemma 6.7.** Let  $\Gamma^\times$  be a skew-Brauer graph with at least two edges. If  $B_{\Gamma^\times}$  is of finite representation type, then it is not a modified Brauer tree algebra.

*Proof.* Since  $B_{\Gamma^{\times}}$  is of finite representation type, by Lemma 6.3,  $\Gamma^{\times}$  has exactly one distinguished vertex  $v^{\times}$ . By definition of skew-Brauer graph, there is exactly one edge x incident to  $v^{\times}$ . We denote by v the other vertex of  $x$ .

Consider the edge  $y = \mathfrak{o}(x)$  having v and u as vertices. This is the minimal configuration for a skew-Brauer graph with a distinguished vertex. We analyse cases for the rest of the graph.

$$
v^\times \xrightarrow{~~x~~} v \xrightarrow{~~y~~} u
$$

Case 1: suppose  $\mathfrak{m}(u) > 1$  or  $\text{val}(u) > 1$ . Then the vertex  $y \in Q_{\Gamma^{\times}}$  is the end of three arrows.

Case 2: the following three  $m(u) = 1$ , val $(u) = 1$  and val $(v) > 1$  hold. Then there are exactly two arrows ending at  $y \in Q_{\Gamma}$  and one arrow starting at y.

Case 3: the graph  $\Gamma^{\times}$  is the one depicted above. In that case either there is no loop in  $Q_{\Gamma^{\times}}$  or the vertex  $y$  is the start and end of three arrows, one of them being a loop.

Given the three cases, by Lemma 6.4 (3) the algebra  $B_{\Gamma^{\times}}$  is not a modified Brauer tree algebra.  $\square$ 

So far we have established that, except for small pathological cases (see Lemma 6.5), skew-Brauer graph algebras of finite representation type are not Brauer graph algebras nor modified Brauer tree algebras. So we should explore the existence of representation finite skew-Brauer graph algebras in the family of trivial extensions of iterated tilted algebras of type D or E. The following result shows that any skew-Brauer graph algebra of finite representation type is not the trivial extension of a iterated tilted algebra of Dynkin type E.

**Lemma 6.8.** Let  $B_{\Gamma^{\times}}$  be a skew-Brauer graph algebra of finite representation type. Then  $B_{\Gamma^{\times}}$  is not isomorphic to a trivial extension of an iterated tilted algebra of Dynkin type E.

*Proof.* By Lemma 6.3, the skew-Brauer graph  $\Gamma^{\times}$  has exactly one distinguished vertex, let  $v^{\times}$  be such vertex and let u be the non-distinguished vertex of  $\Gamma^{\times}$  that shares an edge with  $v^{\times}$ . Since  $\Gamma^{\times}$  has exactly one distinguished vertex, there are exactly two distinguished cycles in  $Q_{\Gamma}$ × induced by the edges attached to u. These cycles share all arrows except two, having the following diagram.



We notice that vertices  $a_1^+$  and  $a_1^-$  have exactly one arrow ending and one arrow starting at them, a property that is not allowed in the quivers of trivial extension of iterated tilted algebras of Dynkin type E, see [\[Fer99,](#page-0-0) Section 3] for a complete list of quivers of trivial extension of iterate tilted algebras of Dynkin type.  $\Box$ 

Remark 6.9. In general, computing cut sets (or admissible cuts) is not an easy task, as the computations depend on the set of elementary cycles of the trivial extension of an algebra. However, in the context of trivial extension of gentle algebras, if we consider its Brauer graph, any set of arrows  $D$  in  $Q_{\Gamma}$  containing exactly one arrow of each special cycle is a cut set, and furthermore  $T(B/\langle D \rangle)$  is isomorphic to B, see [\[Sch15,](#page-0-0) Theorem 1.3].

**Theorem 6.10.** Let  $B_{\Gamma^{\times}}$  be a skew-Brauer graph algebra of finite representation. Then  $B_{\Gamma^{\times}}$  is isomorphic to a trivial extension of iterated tilted algebra of Dynkin type  $\mathbb{D}_n$ . Moreover,  $\Gamma^\times$  is a skew-Brauer three graph.

*Proof.* Let  $\Gamma^{\times} = (\Gamma_0, \Gamma_0^{\times}, \Gamma_1, \mathfrak{m}, \mathfrak{o})$  be a skew-Brauer graph, let  $\Gamma = (\Gamma_0 \cup \Gamma_0^{\times}, \Gamma_1, \mathfrak{m}, \mathfrak{o})$  be its underline Brauer graph, and  $\hat{\Gamma} = (\Gamma_0 \cup \Gamma_0^{\times}, \Gamma_1)$  be its underline graph.

It follows from Lemma 6.2, Lemma 6.6, Lemma 6.7, and Lemma 6.8 that  $B = B_{\Gamma} \times$  is isomorphic to the trivial extension of an iterated tilted algebra A of Dynkin type D.

To prove that  $\Gamma^{\times}$  is a skew-Brauer tree, we observe first that since B is of finite representation type, by Lemma 6.3 there is exactly one vertex  $v^{\times}$ .

As B is isomorphic to the trivial extension of an iterated tilted algebra A of finite representation type, by [\[Yam06,](#page-0-0) Theorem 2], we have that A is triangular, and as a consequence, the multiplicity function m is identically equal to one.

Finally to prove our claim, we need to show that  $\widehat{\Gamma}$  is a tree.

Suppose that  $\hat{\Gamma} = (\Gamma_0 \cup \Gamma_0^{\times}, \Gamma_1)$  is not a tree, then there exists a cycle  $\mathcal{C}$  in  $\hat{\Gamma}$ 

$$
x_1 \stackrel{a_1}{\longrightarrow} x_2 \stackrel{a_2}{\longrightarrow} x_3 \stackrel{a_3}{\longrightarrow} \cdots \stackrel{a_{k-1}}{\longrightarrow} x_k \stackrel{a_k}{\longrightarrow} x_{k+1} = x_1
$$

such that each  $x_i \neq x_j$  if  $i \neq j$  and  $a_k \neq a_1$ . We denote by  $V = \{x_1, \ldots, x_k\}$  the set of vertices of C.

We notice that for each vertex  $x_i$  there exactly one special cycle  $C_{x_i}$ , up to cycle equivalence, in  $Q_{\Gamma}$ , given by the edges attached to each  $x_i$ , for  $i = 0, \ldots, k$ . Since there are no repetition of vertices in C and each arrow of  $Q_{\Gamma}$  belongs to exactly one especial cycle, then by Remark 6.9 the set

$$
\mathcal{D} = \bigcup_{i=1}^k \{ \alpha_{x_i} \in Q_{\Gamma} \mid \alpha_{x_i} : a_i \to \mathfrak{o}(a_i) \} \bigcup \mathcal{D}_2
$$

is a cut set of  $Q_{\Gamma}$ , where  $\mathcal{D}_2$  is any set of arrows containing exactly one arrow  $\alpha_x$  that is not a loop of each special cycle  $C_x$  with  $x \notin V$ . By Theorem 4.11 and Theorem [\[FST](#page-0-0)+22, Theorem 3.6] there exists a good cut  $\mathcal{D}'$  of  $Q_B$  induced by  $\mathcal D$  such that  $T(B/\langle \mathcal{D}' \rangle) \simeq B$ .

Finally, to prove the claim, observe that the quiver  $Q_{\Gamma}$  has an oriented cycle induced by the cycle  $\mathcal{C}$ in  $\Gamma$  given by the sub-paths of  $C_{x_i}$  from the vertex  $a_{i+1}$  to the vertex  $a_i$  in  $Q_A$  for  $i = 2, \ldots, k$  and the sub-path of  $C_{x_1}$  from the vertex  $a_1$  to  $a_k$ , and therefore  $B/\langle \mathcal{D}' \rangle$  has also an oriented cycle, which is a contradiction with the hypothesis of B being the trivial extension of an iterated tilted algebra of Dynkin type  $\mathbb{D}_n$ .

**Theorem 6.11.** Let  $B_{\Gamma}$ <sup>\*</sup> be a skew-Brauer tree algebra. If  $\Gamma$ <sup>\*</sup> is not one of the following skew-Brauer graphs, then  $B_{\Gamma}^{\times}$  is of finite representation type.

 $\times$  w w  $\times$   $\times$   $\times$   $\times$   $\times$ 

*Proof.* By Theorem 4.9,  $B = B_{\Gamma^{\times}}$  is a trivial extension of a skew-gentle algebra. We will show that B is actually a trivial extension of a iterated tilted algebra of Dynkin type  $\mathbb{D}_n$ .

Let  $v^{\times}$  be the unique distinguished vertex in  $\Gamma^{\times}$ , u be the unique non-distinguished vertex in  $\Gamma^{\times}$  that shares an edge with  $v^{\times}$ , and  $u_2, \ldots, u_k$  be the non-distinguished vertices in  $\Gamma^{\times}$  that share an edge with u. Denote these last edges by  $a_2, \ldots, a_k$  respectively, see the picture below.



Since the underlying graph of  $\Gamma^{\times}$  is a tree, the subgraph obtained by deleting the vertex u is disconnected, consisting of connected components  $\Gamma_1, \Gamma_2, \ldots, \Gamma_k$ . These components are in bijection with the vertices  $v^{\times}, u_2, \ldots, u_k$ , where each  $u_i$  is a vertex of the connected component  $\Gamma_i$  and  $\Gamma_i$  is a tree for  $i = 2, ..., k$ . Furthermore,  $\Gamma_1$  is associated with the distinguished vertex  $v^{\times}$ .

By [\[WZ22,](#page-0-0) Lemma 3.1], each connected component  $\Gamma_i$  for  $i = 2, \ldots, k$  induces a trivial extension of an iterated tilted algebra of Dynkin type  $\mathbb{A}_n$ . Then any cut set  $\mathcal{C}_i$  of  $\Gamma_i$  defines an iterated tilted algebra of Dynkin type  $\mathbb{A}_n$ 

Even more, by construction of  $Q_{\Gamma}$ ×, there are exactly two distinguished cycles in  $Q_{\Gamma}$ × induced by the edges attached to  $u$  that share all arrows except two, having the following diagram.



It is clear that the set  $\mathcal{C} = \bigcup_{k=1}^{k}$  $\bigcup_{i=2}$   $C_i \cup \{\alpha^+, \alpha^-\}\}$  is a cut set of  $B_{\Gamma^\times}$ . Then the induced quotient algebra  $A = B/\langle C \rangle$  of B has the following quiver

 $a_1^+ \xrightarrow{+_{\beta}} a_2^* \longrightarrow a_3^* \longrightarrow \dots \longrightarrow a_{k-1}^* \longrightarrow a_k^* \xrightarrow{\alpha^-} a_1^-$ 

where each  $a_i^*$  has attached an iterated tilted algebra of Dynkin type  $\mathbb{A}_n$ . Then by [\[AS06,](#page-0-0) Section 7, Main theorem, case D3 and [\[Kel91\]](#page-0-0) A is an iterated tilted algebra of Dynkin type  $\mathbb{D}$ .  $\Box$ 

## 7. A geometric interpretation using dissections

7.1. Computing on dissections. In this section we use generalized dissections of orbifold surfaces, that were originally considered in [\[LFSV20, AB19\]](#page-0-0) to study the derived categories of skew-gentle algebras, to give a geometric version of the good cut set. We follow [\[OPS18, LFSV20, AB19\]](#page-0-0) for the next definitions and results.

A marked surface is a tuple  $(S, M, P)$  where S is a compact and oriented Riemann surface with boundary, M is a finite set of marked points in the boundary  $\partial S$ , and P is a finite set of marked points in the interior of S. We say that a marked surface  $(S, M, P)$  is unpunctured if  $P = \emptyset$ .

A boundary segment is a curve in  $\partial S$  that is delimited by its endpoints in M and does not contain other points in M. An arc is a curve  $\gamma: [0,1] \to S$  that is injective on  $(0,1)$ , it is considered up to injective homotopy over that interval, and such that  $\gamma(0)$  and  $\gamma(1)$  belong to M or to P. We say that an arc  $\gamma$  is pendant if it is incident to  $p \in P$  and p is the endpoint of exactly one arc.

A •-dissection of  $(S, M, P)$  is a set  $D = \{\gamma_1, \ldots, \gamma_s\}$  of arcs splitting the surface into (degenerate) polygons whose edges are arcs except for exactly one boundary segment. In this context, a degenerate polygon is one that contains pendant arcs as one of its edges, see for example Figure 5 (left). The tuple  $(S, M, P, D)$  is called *surface dissection*.

An orbifold marked surface is a tuple  $(S, M, P, O)$  where  $(S, M, P)$  is a marked surface and O is a finite set of marked points in the interior called *orbifold points*. A special arc  $\gamma^{\times}$  is an arc connecting a point in M and an orbifold point.

A  $\times$ -dissection of an orbifold marked surface  $(S, M, P, O)$  is a set  $D = \{ \gamma_1, \ldots, \gamma_s, \gamma_{s+1}^{\times}, \ldots, \gamma_r^{\times} \}$  of arcs such that  $\{\gamma_1,\ldots,\gamma_s\}$  is a •-dissection of  $(S, M, P)$  and each  $x \in \mathcal{O}$  is an endpoint of exactly one arc in  $\{\gamma_{s+1}^{\times}, \ldots, \gamma_r^{\times}\}\.$  The tuple  $(S, M, P, O, D)$  is called *orbifold dissection*.

By  $[LFSV20, Lemma 5.3]$ , the orbifold  $\times$ -dissection splits the surface into (degenerate) polygons. Such polygons have all edges in  $\{\gamma_1, \ldots, \gamma_s\}$  except for one edge that is a boundary segment. In this case, a degenerate polygon may also contain special arcs or pendant arcs as one of its edges. The list of such polygons might contain what we call trivial polygons, i.e. bigons enclosed by an edge in the dissection and one boundary segment.

The  $\times$ -dissection  $(S, M, P, O, D)$  is defined in [\[LFSV20,](#page-0-0) Sect. 4] as the dual dissection of another orbifold dissection. In this construction, the orbifold dissection  $(S, M, P, O, D)$  has two different kinds of points: the red points and the black points, in our definition, we are omitting the latter. The authors also show that  $\times$ -dissection is related to the Koszul dual algebra A! of A. In this section, we establish a different convention to construct a quiver Q and an ideal I that recovers, in our case, a skew-gentle algebra A and also data of the trivial extension  $T(A^{sg})$  of  $A^{sg}$ .

We define the quiver of the skew gentle algebra  $A = KQ/I$  from a  $\times$ -dissection  $(S, M, P, O, D)$ following the rules:

- (r1) Each arc  $\gamma_i$  of D gives rise to a vertex i of Q. In particular, special arcs correspond to special vertices of Q.
- (r2) Each internal angle between arcs in a non-trivial polygon gives rise to an arrow oriented in counterclockwise direction, taking the marked point as pivot. Additionally, each pendant arc (resp. special arc)  $\gamma_i$  gives rise to a loop (resp. special loop)  $f_i$  in  $Q$  such that  $s(f_i) = t(f_i) = i$ .
- $(r3)$  The set of relations of I is given as follows:
	- (i) Any pair of consecutive arrows  $\alpha\beta$  from two different polygons define a relation  $\alpha\beta \in I$ .
	- (ii) A non-special loop  $f_i$  defines a relation  $f_i^2 = 0$ .
	- (iii) A special loop  $f_i$  defines the relation  $f_i^2 = f_i$ .
	- (iv) If  $\alpha$  and  $\beta$  are two consecutive arrows in the same polygon, but  $t(\alpha) = s(\beta)$  corresponds to a special or a pendant arc, then  $\alpha\beta$  is a relation.

**Example 7.1.** In Figure 5 we show a **•**-dissection and a  $\times$ -dissection, each one with three polygons. One dissection has an interior point, the other one has an orbifold point.



FIGURE 5. Two dissections for the disk that define the same Q.

On the terms of Section 3, each  $\times$ -dissection defines a tuple  $(Q, I, Sp, \emptyset)$  where the vertices of Q correspond to arcs, the arrows of Q arise from angles and pendant arcs (i.e. we must not consider special loops) and the ideal I is defined as above but without considering the relations  $\alpha\beta$  when  $t(\alpha) = s(\beta)$  corresponds to a special arc. The special arcs define the vertices in Sp and C is empty. The admissible presentation for the skew gentle algebra  $KQ/I$  from the  $\times$ -dissection is defined as  $KQ^{sg}/I^{sg}.$ 

**Example 7.2.** The ×-dissection in Figure 6 defines the tuple  $(Q, I, Sp, \emptyset)$  where  $I = \langle \beta \gamma, \gamma \beta, \alpha \delta, \varepsilon \lambda \rangle$ ,  $Sp = \{4\}$  and  $C = \emptyset$ .



FIGURE 6. A  $\times$ -dissection of the torus with one boundary component.

As we mentioned before, a  $\times$ -dissection defines a skew-gentle algebra  $A = KQ/I$ , and we can translate this information to a tuple  $(Q, I, Sp, \emptyset)$ , but it also gives us information of the trivial extension  $T(A^{sg})$ . This information is obtained as follows.

By construction each non-trivial (degenerate) polygon  $P_i$  defines a sp-maximal path  $p_i$  in A. Moreover, by Lemma 4.8, the set of  $sp$ -maximal paths are in bijection with the set of maximal paths in  $A'$ . Hence, each  $p_i$  gives rise a maximal path  $p'_i$  in  $A'$  and a new arrow  $\beta_{p'_i}$  such that  $p'_i \beta_{p'_i}$  is an elementary cycle of A′ . As a consequence there is a bijection between non-trivial (degenerate) polygons and elementary cycles of A'. Then, the x-dissection defines a tuple  $(Q^*, I^*, Sp, \mathcal{C})$ , where  $Q^*$  is obtained from Q by adding the arrow  $\beta_{p_i}$  for each non-trivial polygon  $P_i$ . The family C is formed by all possible permutations of the cycles  $C_{P_i} = p_i \beta_{p_i}$  arising from non-trivial (degenerate) polygons, while  $Sp$ remains the same. The new ideal  $I^*$  contains:

- (1) all the quadratic relations in I
- (2) quadratic relations of the form  $\beta_{p_i} \alpha$  where  $\alpha$  is an arrow such that  $t(\beta_{p_i}) = s(\alpha)$  and it is not the first arrow of  $p_i$ , and of the form  $\omega \beta_{p_i}$  where  $\omega$  is an arrow such that  $t(\omega) = s(\beta_{p_i})$  and  $\omega$ is not the last arrow of  $p_i$ .
- (3) monomial relations of the form  $C\alpha$  where C is a cycle from the family C and  $\alpha$  is the first arrow of  $C$ ,
- (4) relations of the form  $C_1 C_2$  where  $C_1$  and  $C_2$  are distinguished cycles starting at the same vertex  $x \notin Sp$ . These relations arise when a red arc is an edge of two non trivial polygons or a repeated edge in a single non trivial polygon. See the next example.

**Example 7.3.** The  $\times$ -dissection in Figure 6 is formed by two polygons. The polygon  $P_1$  is delimited by edges  $1, 2, 3, 2, 1, 5$  and boundary, and the polygon  $P_2$  is delimited by edges 3,5 and boundary. The quiver  $Q^*$  for the ×-dissection is the one below. In this case  $P_1$  contains the maximal path  $p_1 = \gamma \delta \varepsilon \alpha \beta \omega$  and generates a new arrow  $\beta_{p_1}$  (in red) and  $P_2$  contains the maximal path  $p_2 = \lambda$  and generates the new arrow  $\beta_{p_2}$  (in blue).



There are two non trivial polygons and there are four edges shared by polygons, indeed:

- a) the arc 3 sits between  $P_1$  and  $P_2$ ,
- b) the arc 5, between  $P_1$  and  $P_2$ ,
- c) the arc 1, between  $P_1$  and  $P_1$ ,
- d) the arc 2, between  $P_1$  and  $P_1$ .

The relations of the form  $C_1 - C_2$  arise for each one of these cases. The ideal  $I^*$  is generated by the relations:

- $βγ, γβ, αδ, ελ$  (quadratic relations in I)
- $\alpha\beta\omega\beta_{p_1}\gamma\delta\varepsilon \lambda\beta_{p_2}$  from a)
- $\beta_{p_2} \lambda \beta_{p_1} \gamma \delta \varepsilon \alpha \beta \omega$  from b)
- $\gamma \delta \varepsilon \alpha \beta \omega \beta_{p_1} \omega \beta_{p_1} \gamma \delta \varepsilon \alpha \beta$  from c)
- $\delta \varepsilon \alpha \beta \omega \beta_{p_1} \gamma \beta \omega \beta_{p_1} \gamma \delta \varepsilon \alpha$  from d)
- Relations of the form  $C\alpha$ , with  $\alpha$  the first arrow of the cycle for each cyclic orientation of the cycles  $C_{P_1} = p_1 \beta_{p_1}$  and  $C_{P_2} = p_2 \beta_{p_2}$ .
- $\beta_{p_1}\omega$ ,  $\beta_{p_2}\alpha$ ,  $\lambda\beta_{p_1}$ ,  $\omega\beta_{p_2}$  (new quadratic relations)

We have built a tuple  $(Q^*, I^*, Sp, \mathcal{C})$  from the ×-dissection. It follows from our definition that the algebra  $B = K(Q^*)^{sg}/(I^*)^{sg}$  is the trivial extension for the skew-gentle algebra A.

**Proposition 7.4.** Let A be a skew-gentle with Sp the set of special vertices, and let  $(S, M, P, \mathcal{O}, D)$ be its  $\times$ -dissection. The quiver of the trivial extension  $T(A^{sg})$  of the algebra  $A^{sg}$  coincides with the sg-bound quiver arising from the tuple  $(Q^*, I^*, Sp, C)$ .

*Proof.* This follows from the fact that the algebra A can be defined from the  $\times$ -dissection, and from the the construction proposed for the tuple, where we get a new arrow  $\beta_{p_i}$  for each maximal path as in Theorem 3.11.  $\Box$ 

Example 7.5. The skew-Brauer graph for Example 7.3 is depicted in Figure 7:



Figure 7. Skew-Brauer graph.

The sg-bound quiver from the tuple  $(Q^*, I^*, Sp, C)$  defines a skew-Brauer graph algebra (with  $\mathfrak{m} = 1$ ) for all vertices) arising from the graph in Figure 7 and this algebra is the trivial extension of the one defined by  $(Q, I, Sp, \emptyset)$  arising from the  $\times$ -dissection in Figure 6.

7.2. Contracting and adding boundary segments. Let  $\mathcal{X} = (S, M, P, \mathcal{O}, D)$  be a  $\times$ -dissection that splits the surface into (degenerate) polygons. To fix notation, let  $P_1, P_2, \ldots, P_m$  be the non-trivial polygons.

For a polygon  $P_i$  and an interior angle  $\theta$  of  $P_i$  (or a marked point  $p_i$  in the interior of  $P_i$  attached to a pendant arc), we define *contraction-addition at*  $(P_i, \theta)$  (resp.  $(P_i, p_i)$ ) as follows.

- (1) For the polygon  $P_i$ , there is exactly one edge, say b, that is a boundary segment. We note that  $b$  can be a boundary component with one marked point. If  $b$  is a boundary segment with two different endpoints, then we contract  $b$  by identifying the points. If  $b$  is a boundary component with one marked point, then we contract  $b$  to a single marked point. In both cases, we obtain a polygon  $\widetilde{P}_i$  with no boundary segment as an edge.
- (2) For the case in which we have an angle  $\theta$ , there is a red marked point p in  $P_i$ . If p is in a boundary component, we can add a new boundary segment, splitting the red marked point (see Figure 8 left). Otherwise, if  $p$  is interior, we add a new boundary component in the angle  $\theta$  (see Figure 8 center).
- $(2')$  For the case in which we have an interior marked point  $p_i$ , we add a boundary component, having  $p_i$  has a marked point (Figure 8 right)



FIGURE 8. Adding boundary segment at an angle  $\theta$  or a boundary component at  $p_i$ .

In both cases, (2) and (2'), we obtain a new polygon  $P'_i$  with exactly one edge that is a boundary segment. We denote by  $\tau_{(P_i,\theta)}\mathcal{X}$  (resp.  $\tau_{(P_i,p_i)}\mathcal{X}$ ) the new ×-dissection. The new dissection splits S into (degenerate) polygons, and the set of non-trivial polygons is  $\{P_1, \ldots, P_m\} \setminus P_i \cup \{P'_i\}.$ 

**Example 7.6.** A  $\times$ -dissection for the annulus and the new  $\times$ -dissection after a contraction-addition at  $\theta$ .



FIGURE 9. A  $\times$ -dissection X and the new  $\times$ -dissection  $\tau_{(P_i,\theta)}\mathcal{X}$ .

Observe that both  $\times$ -dissections in Figure 9, generate the same tuple  $(Q^*, I^*, Sp, C)$  and this tuple defines the skew-Brauer graph algebra B (with  $m = 1$ ) from Example 4.2. Each one of two  $\times$ dissections defines a skew gentle algebra that is a good cut (Definition 4.12) for B.

**Theorem 7.7.** Let A be a skew-gentle with Sp the set of special vertices, and  $\mathcal{X} = (S, M, P, \mathcal{O}, D)$  its  $\times$ -dissection and let D be a good cut set of  $T(A^{sg})$ . Then there is a sequence of contraction-addition moves taking X to  $\widetilde{\mathcal{X}}$ , where  $\widetilde{\mathcal{X}}$  is the  $\times$ -dissection of the quotient algebra of  $T(A^{sg})$  induced by  $\mathcal{D}$ .

*Proof.* Let D be a good cut of  $T(A^{sg})$ . By definition there exists an admissible cut D' of the trivial extension  $T(A')$  where A' is the associated gentle algebra of A. By [\[Sch15,](#page-0-0) Theorem 1.3],  $\mathcal{D}'$  is a set of arrows containing exactly one arrow from each one of the special cycles of  $T(A')$ , which by Lemma 4.8 are in bijection with the non-trivial (degenerate) polygons of  $\mathcal{X}$ . To fix notation, denote by  $\alpha_i$  the arrow of  $\mathcal{D}'$  associated with the non-trivial polygon  $P_i$ .

Each arrow  $\alpha_i$  of  $\mathcal{D}'$  is either opposite an internal angle  $\theta_i$  of the non-trivial polygon  $P_i$ , corresponds to a pendant arc with end point  $p_i$ , or corresponds to a boundary segment of  $P_i$ . Only in the first two cases, we need to apply an contraction-addition at  $(P_i, \theta_i)$  or  $(P_i, p_i)$  respectively.

Since  $\mathcal{D}'$  is an admissible cut of  $T(A')$ , there are no repeated polygons in the list of contractionaddition at  $(P_i, \theta_i)$  and  $(P_i, p_i)$  induced by the arrows of  $\mathcal{D}'$ . Then we can define a sequence  $\tau$  of contraction-addition, and it is well-defined. Finally, by construction  $T(A^{sg})/\langle \mathcal{D} \rangle$  is isomorphic to the skew-gentle algebra obtained from  $\tau \mathcal{X}$ .

7.3. On derived equivalent skew-gentle algebras. The following example shows that contractionaddition does not necessarily induces derived equivalent skew-gentle algebras.



FIGURE 10. ×-dissection  $\mathcal X$  (left) and the new  $\times$ -dissection  $\tau_{(P,\theta)}\mathcal X$  (right).

**Example 7.8.** Let A and B the skew-gentle algebras associated to the  $\times$ -dissections depicted in Figure 10. One can see that A and B are related by an contraction-addition at  $\theta$ , and even more the  $\times$ -dissections are induced by good cuts of a skew-Brauer graph algebra with multiplicity  $\mathfrak{m} = 1$ .

Let A be the skew-gentle algebra defined by X in Figure 10, and consider B defined by  $\tau_{(P,\theta)}\mathcal{X}$ .

$$
1 \xrightarrow{\alpha_1} 2 \xrightarrow{\alpha_2} 3 \bigcirc f_3 \qquad 1 \xrightarrow{\alpha_1} 2 \xrightarrow{\alpha_2} 3 \bigcirc f_3
$$

Hence A is defined by the quiver on the left and taking the ideal  $\langle \beta_1 \alpha_2, f_3^2 - f_3 \rangle$ , while B is defined by the quiver on the right considering the ideal  $\langle \beta_2 \alpha_1, \alpha_1 \beta_2, f_3^2 - f_3 \rangle$ .

Following [\[LFSV20,](#page-0-0) Theorem 7.3] and looking at the  $\times$ -dissections<sup>[5](#page-29-0)</sup>, it is easy to see that the determinants of their q-Cartan matrices are det  $(C_A)_q = 1$  and det  $(C_B)_q = 1 - q^2$ . The ordinary Cartan matrices are obtained by evaluating  $q = 1$ , hence we can find their determinants det  $(C_A) = 1$  and

<span id="page-29-0"></span><sup>&</sup>lt;sup>5</sup>Theorem 7.3 in [\[LFSV20\]](#page-0-0) can be re stated det  $(C_A)_q = \prod_{k \geq 1} (1 - (-q)^k)^{c_k}$  where  $c_k$  is the number of red punctures with  $k$  incident red arcs.

 $\det (C_B) = 0$ . Since these determinants are derived invariants [\[BS05,](#page-0-0) Proposition 1.5], we conclude that  $A$  and  $B$  are not derived equivalent.

7.4. Reflections as geometric moves on a ×-dissection. By Theorem 5.7 and Theorem 7.7, any sequence of reflections in a skew-gentle algebra A can be interpreted as a sequence of contractionaddition on its ×-dissection. In this section, we present another interpretation of reflections as local moves in  $\times$ -dissections. We will present examples of reflections on  $\times$ -dissections and show how these moves behave locally.

Let A be a skew-gentle algebra defined by a  $\times$ -dissection. Let y be a vertex that is a source or a target in  $Q_{A'}$ , where A' is the auxiliary gentle algebra of A. Such a vertex will be associated to a red arc y that can be a special arc, as is Figure 11 right, or a regular red arc. If the vertex y is a target, the arc y is such that if we pivot counterclockwise on one of its endpoints, we meet the boundary edge of a polygon. See Figure 11 left.

Let y be a non-special target vertex. Consider  $p_a$  and  $p_b$  the maximal paths in A' ending at y. See that one of these paths might be trivial, that is the case when one of the polygons having  $y$  as an edge is a trivial polygon. Let  $\beta_{p_1}$  and  $\beta_{p_2}$  the new arrows in  $Q_{\widehat{A}}$ , then  $S_y^+(A)$  is given by a quiver and the (non-admissible) ideal defined by a new  $\times$ -dissection, exchanging locally the vertex y and the vertex y' as in the next figure. If y is special, we see the local exchange  $S_y^-(A)$  in the right part of the figure. The negative and positive reflections for non-special and special vertices are analogous.



FIGURE 11. Positive reflection  $S_y^+$  with y a non-special target vertex (right). Negative reflection  $S_y^-$  with y a special source vertex (left)

**Example 7.9.** Let A be the skew gentle algebra defined by the  $\times$ -dissection on the left of Figure 9. Thus  $A = KQ/I$  where Q is the quiver on the left,



and  $I = \langle f_1^2 - f_1, f_2^2 - f_2, \alpha_1 \alpha_2, \alpha_3 \beta_3, \beta_1 \alpha_3 \rangle$ . Then  $\beta_{p_a}$  is the new arrow arising from the path  $p_a = \alpha_1 \alpha_2 \alpha_3$  and  $S_1^{-}(A)$  is the skew-gentle algebra defined by the quiver on the right and the ideal  $J = \langle f_{1'}^2 - f_{1'}, f_2^2 - f_2, \alpha_3 \beta_3, \beta_1 \alpha_3 \rangle$ , that is associated to the ×-dissection after applying a move as the one in Figure 11 right.

## **REFERENCES**

- [AAC18] Takahide Adachi, Takuma Aihara, and Aaron Chan. Classification of two-term tilting complexes over brauer graph algebras. Mathematische Zeitschrift, 290(1-2):1–36, 2018.
- [AB19] Claire Amiot and Thomas Brüstle. Derived equivalences between skew-gentle algebras using orbifolds.  $arXiv$ preprint arXiv:1912.04367, 2019.
- [ABF<sup>+</sup>] María Alejandra Álvarez, Diego Bravo, Elsa Fernández, Victoria Guazzelli, Monique Müller, Nadina Rojas, and Sonia Trepode. Trivial extension of admissible cuts and HW reflexions. Work in progress 2024.
- [ACH22] Takuma Aihara, Aaron Chan, and Takahiro Honma. On representation-finite gendo-symmetric algebras with only one non-injective projective module. Journal of Algebra, 603:61–88, 2022.
- [AS06] Ibrahim Assem and Andrzej Skowronski. Algèbres pré-inclinées et catégories dérivées. In Séminaire d'Algèbre Paul Dubreil et Marie-Paul Malliavin: Proceedings, Paris 1987–1988 (39ème Année), pages 1–34. Springer, 2006.
- [ASS<sup>+</sup>06] Ibrahim Assem, Daniel Simson, Andrzej Skowronski, et al. Elements of the Representation Theory of Associative Algebras: Volume 1: Techniques of Representation Theory. Cambridge University Press, 2006.
- [AZ22] Mikhail Antipov and Alexandra Zvonareva. Brauer graph algebras are closed under derived equivalence. Mathematische Zeitschrift, 301(2):1963–1981, 2022.
- [BMM03] Viktor Bekkert, Eduardo N. Marcos, and Héctor A. Merklen. Indecomposables in derived categories of skewedgentle algebras. Comm. Algebra, 31(6):2615–2654, 2003.
- [BS05] Rafał Bocian and Andrzej Skowroński. Weakly symmetric algebras of euclidean type. 2005.
- [CSS20] Cristian Chaparro, Sibylle Schroll, and Andrea Solotar. On the lie algebra structure of the first hochschild cohomology of gentle algebras and brauer graph algebras. Journal of Algebra, 558:293–326, 2020.
- [DF78] Peter W Donovan and M R Freislich. The indecomposable modular representations of certain groups with dihedral sylow subgroup. Mathematische Annalen, 238:207–216, 1978.
- [Fer99] Elsa Fernández. Extensiones triviales y álgebras inclinadas iteradas. PhD thesis, Ph. D. Thesis, Universidad Nacional del Sur, Argentina, 1999.
- [FP02] Elsa A Fern´andez and Marıa In´es Platzeck. Presentations of trivial extensions of finite dimensional algebras and a theorem of sheila brenner. Journal of Algebra, 249(2):326–344, 2002.
- [FP06] Elsa A Fernández and Mara Inés Platzeck. Isomorphic trivial extensions of finite dimensional algebras. Journal of Pure and Applied Algebra, 204(1):9–20, 2006.
- [FST+22] Elsa Fernandez, Sibylle Schroll, Hipolito Treffinger, Sonia Trepode, and Yadira Valdivieso. Characterisations of trivial extensions. arXiv preprint arXiv:2206.04581, 2022.
- [GdlP99] Christof Geiß and José Antonio de la Peña. Auslander-reiten components for clans. Boletín de la Sociedad Matemática Mexicana: Tercera Serie, 5(2):307-326, 1999.
- [GLFS16] Christof Geiß, Daniel Labardini-Fragoso, and Jan Schröer. The representation type of jacobian algebras. Advances in Mathematics, 290:364–452, 2016.
- [GR05] Christof Geiß and Idun Reiten. Gentle algebras are gorenstein. Representations of algebras and related topics, pages 129–133, 2005.
- [GS17] Edward L Green and Sibylle Schroll. Brauer configuration algebras: A generalization of brauer graph algebras. Bulletin des Sciences Mathématiques, 141(6):539–572, 2017.
- [HW83] David Hughes and Josef Washbüsch. Trivial extensions of tilted algebras. Proceedings of the London Mathematical Society, 46:347–364, 1983.
- [Kel91] Bernhard Keller. Algèbres héréditaires par morceaux de type  $D_n$ . C. R. Acad. Sci. Paris Sér. I Math., 312(7):483–486, 1991.
- [Lam91] Tsit-Yuen Lam. A first course in noncommutative rings, volume 131. Springer, 1991.
- [LFSV20] Daniel Labardini-Fragoso, Sibylle Schroll, and Yadira Valdivieso. Derived categories of skew-gentle algebras and orbifolds. arXiv preprint arXiv:2006.05836, 2020.
- [MS14] Bethany Rose Marsh and Sibylle Schroll. The geometry of brauer graph algebras and cluster mutations. Journal of Algebra, 419:141–166, 2014.
- [Nor21] Anya Nordskova. Derived picard groups of symmetric representation-finite algebras of type d. arXiv preprint arXiv:2112.09257, 2021.
- [OPS18] Sebastian Opper, Pierre-Guy Plamondon, and Sibylle Schroll. A geometric model for the derived category of gentle algebras. arXiv preprint arXiv:1801.09659, 2018.
- [OZ22] Sebastian Opper and Alexandra Zvonareva. Derived equivalence classification of brauer graph algebras. Advances in Mathematics, 402:108341, 2022.
- [Sch99] Jan Schröer. On the quiver with relations of a repetitive algebra. Archiv der Mathematik, 72(6):426–432, 1999.
- [Sch15] Sibylle Schroll. Trivial extensions of gentle algebras and brauer graph algebras. Journal of Algebra, 444:183– 200, 2015.
- [Sch18] Sibylle Schroll. Brauer graph algebras: a survey on Brauer graph algebras, associated gentle algebras and their connections to cluster theory. In Homological methods, representation theory, and cluster algebras, CRM Short Courses, pages 177–223. Springer, Cham, 2018.
- [Sko06] Andrzej Skowronski. Selfinjective algebras: finite and tame type. Trends in representation theory of algebras and related topics, 406:169–238, 2006.
- [Sot24] Valentine Soto. Tilting mutations as generalized kauer moves for (skew) brauer graph algebras with multiplicity. arXiv preprint arXiv:2406.10634, 2024.
- [WZ22] Qi Wang and Yingying Zhang. On τ−tilting modules over trivial extensions of gentle tree algebras. arXiv preprint arXiv:2204.06418, 2022.
- [Yam96] Kunio Yamagata. Frobenius algebras. In Handbook of algebra, Vol. 1, volume 1 of Handb. Algebr., pages 841–887. Elsevier/North-Holland, Amsterdam, 1996.
- [Yam06] Kunio Yamagata. On algebras whose trivial extensions are of finite representation type. In Representations of Algebras: Proceedings of the Third International Conference on Representations of Algebras Held in Puebla, Mexico, August 4–8 1980, pages 364–371. Springer, 2006.

[AGE] Facultad de Ciencias Exactas y Naturales (CEMIM), Universidad Nacional de Mar del Plata and CONICET. Dean Funes 3350. (7600) Mar del Plata, Argentina.

## Email address: elsener@mdp.edu.ar

[VG] Facultad de Ciencias Exactas y Naturales, Universidad Nacional de Mar del Plata. Dean Funes 3350. (7600) Mar del Plata, Argentina.

## $Email \;address\colon\; \mathtt{vguazzelli@mdp.edu.ar}$

[YV] DEPARTAMENTO DE MATEMÁTICAS, UNIVERSIDAD DE LAS AMÉRICAS PUEBLA. EX HACIENDA STA. CATARINA MÁRTIR S/N. (72810) SAN ANDRÉS CHOLULA, PUEBLA, MÉXICO.

#### Email address: yadira.valdivieso@udlap.mx