# REMARKS ON LIPSCHITZ GEOMETRY ON GLOBALLY CONIC SINGULAR MANIFOLDS

ANDRÉ COSTA, VINCENT GRANDJEAN, AND MARIA MICHALSKA

In honour of Professor Pham Tiến Sơn for his sixtieth birthday.

Abstract. We study metric properties of manifolds with conic singularities and present a natural interplay between metrically conic and metrically asymptotically conic behaviour. As a consequence, we prove that a singular sub-manifold is Lipschitz normally embedded, i.e. its inner and outer metric structures are equivalent, in an ambient singular manifold, whenever the singularities are conic and the ends of the manifold are asymptotically conic, which answers positively a question of [\[7\]](#page-20-0).

#### **CONTENTS**



#### <span id="page-0-0"></span>**INTRODUCTION**

The aim of this note is to explain within the context of conic singular manifolds how the results on Lipschitz geometry of algebraic sets of our recent works [\[9,](#page-20-2) [7\]](#page-20-0), compare [\[8\]](#page-20-3), are occurrences of a general simple principle.

A diffeomorphism between two smooth Riemannian manifolds is always a locally bi-Lipschitz mapping. Yet in presence of singularities a counterpart statement is not clear. Thus we investigate this question here for metrically conic singularities. We will showcase the relations between conic singular manifolds, manifolds with conic metric and their quotient spaces in Section [2.](#page-8-0)

The natural interplay between metrically conic and asymptotically metrically conic behaviour is presented in Corollary [2.12](#page-14-0) and Proposition [2.16.](#page-15-1) Thus Lipschitz geometry of globally conic singular manifolds becomes much clearer and results conjectured in [\[7\]](#page-20-0) follow in Theorem [3.9](#page-19-0) which states that every connected globally conic singular sub-manifold of a globally conic singular manifold is Lipschitz normally embedded, i.e. the inner and outer metric structures are equivalent.

Key words and phrases. conic metric, asymptotically conic metric, conic singularity, Lipschitz geometry, Lipschitz normally embedded set.

#### 1. Preliminaries

<span id="page-1-0"></span>Throughout the paper smooth means  $C^{\infty}$ -smooth, all manifolds without or with boundary are smooth, all Riemannian metrics are assumed smooth.

#### 1.1. Metric structures of subsets

For convenience we allow a distance function to attain value  $\infty$ , for instance for points in different connected components of a set. Thus isometries of metric spaces are connectedcomponent-wise isometries.

A pair  $(M, d)$  is a pseudo-metric space, if the function  $d : M \times M \to \mathbb{R}_{\geq 0} \cup \{\infty\}$  is a pseudodistance, i.e. for all  $\mathbf{x}, \mathbf{x}', \mathbf{x}'' \in M$  it satisfies the following conditions:

$$
d(\mathbf{x}, \mathbf{x}) = 0
$$
,  $d(\mathbf{x}, \mathbf{x}') = d(\mathbf{x}', \mathbf{x})$  and  $d(\mathbf{x}, \mathbf{x}'') \leq d(\mathbf{x}, \mathbf{x}') + d(\mathbf{x}', \mathbf{x}'')$ .

Let  $(M, d)$  be a pseudo-metric space. The length of an arc  $c : [a, b] \to M$  is

$$
\ell(\mathbf{c}) = \sup \left| \sum_{i=1}^k d(\mathbf{c}(t_{i-1}), \mathbf{c}(t_i)) \right|,
$$

where the supremum is taken over all partitions  $a = t_0 < \cdots < t_k = b, k \in \mathbb{N}$ , of the interval I.

**Definition 1.1.** Let  $(M, d)$  be a pseudo-metric space and X be a subset of M.

(i) The outer pseudo-metric structure is  $(X, d_X)$ , where  $d_X$  is the restriction of the pseudodistance in M to X.

(ii) The inner pseudo-metric structure is  $(X, d_{\text{in}}^X)$ , where  $d_{\text{in}}^X$  is the infimum of lengths of arcs within X connecting a given pair of points.

Whenever the inner (rep. outer) pseudo-distance is a distance on  $X$ , the inner (resp. outer) structure is naturally an inner (resp. outer) metric structure on X.

A mapping  $f : (M, d) \to (M', d')$  between pseudo-metric spaces is *pseudo-Lipschitz* if there exists a positive constant L such that the inequality  $d'(f(x), f(y)) \le Ld(x, y)$  is satisfied for all  $x, y \in M$ .

A special type of subsets of a (pseudo-)metric space are those for which their outer and inner (pseudo-)metric structure are indistinguishable from the (pseudo-)Lipschitz point of view:

**Definition 1.2.** Let  $(M, d)$  be a pseudo-metric space and X be a subset of M such that its inner and outer pseudo-distances in  $X$  are both distance functions.

(i) The subset  $X$  is Lipschitz normally embedded *(shortened to LNE)* in  $M$  if its outer and inner distances are equivalent, i.e. the identity map

$$
id : (X, d_X) \to (X, d_{\text{in}}^X)
$$

is bi-Lipschitz.

(ii) The subset X is locally LNE at  $x$  if there exists a neighbourhood U of  $x$  in M such that  $X \cap U$  is LNE in M.

(iii) The subset X is locally LNE if it is locally LNE at each point of its closure in  $M$ .

(iv) The space  $(X, d_X)$  is a length space when the identity map of point (i) is an isometry.

When writing that a subset of a metric space is LNE, it is always understood that it is LNE with respect to the metric structure of the given ambient space. By convention, the empty set is LNE. This notion is the same as Whitney's  $P$  condition of [\[22\]](#page-21-0) and Gromov's quasi-convexity of [\[14\]](#page-21-1). We use the least ambivalent name of Lipschitz normally embedded after the paper [\[1\]](#page-20-4).

### <span id="page-2-0"></span>1.2. Singular metrics and pseudo-distances on manifolds

Let  $(M, \partial M)$  be a smooth manifold with boundary (possibly empty). A *singular Riemannian* metric over a manifold M is a smooth 2-symmetric tensor over M everywhere positive semidefinite. The *singular locus*  $\text{sing}(g)$  of the singular Riemannian metric g over M is the set of points of M at which g is not positive-definite. It is a closed subset of M and  $g_{|M\setminus\text{sing}(g)}$  is a Riemannian metric.

Two singular Riemannian metrics q and h over  $(M, \partial M)$  are *equivalent* if there exist constants  $a, b > 0$  such that

$$
a g(\mathbf{x})(\xi, \xi) \leq h(\mathbf{x})(\xi, \xi) \leq b g(\mathbf{x})(\xi, \xi), \ \forall (\mathbf{x}, \xi) \in TM.
$$

Given a singular Riemannian metric q over a manifold  $M$ , the length of a smooth curve  $c$ :  $[a, b] \to M$  is

$$
\ell_g(\mathbf{c}) = \int_a^b \left( g(\mathbf{c}'(t), \mathbf{c}'(t)) \right)^{\frac{1}{2}} dt.
$$

Then M is equipped with the pseudo-distance  $d_g$  obtained by taking the infimum of the length of piecewise smooth curves connecting a given pair of points, this pseudo-distance is a distance function on  $M \setminus \text{sing}(g)$ .

**Remark 1.3.** By for instance  $[2, 21]$  $[2, 21]$ , given a length structure d on the manifold M, smooth outside of the diagonal  $M \times M$ , there exists a Riemannian metric g on M such that  $d = d_q$ . By Myers-Steenrod theorem a mapping between Riemannian manifolds is a smooth Riemannian isometry if and only if it is an isometry of the underlying metric spaces.

In particular, a diffeomorphism between smooth manifolds is locally bi-Lipschitz regardless of the choice of Riemannian metrics. Thus the Lipschitz structure on a manifold depends only on its smooth structure and is independent on the Riemannian structure (this is already true for  $\mathcal{C}^1$  regularity).

### 1.3. Collar neighbourhoods and p-sub-manifolds

Let  $(M, \partial M)$  be a smooth manifold with boundary  $\partial M$  (possibly empty).

A boundary defining function in M is a smooth function over M, positive in  $M \setminus \partial M$ , vanishing on  $\partial M$  and with no critical point in  $\partial M$ . Such smooth functions always exist. When  $\partial M$  is compact, given any boundary defining function x, there exists  $\eta > 0$  such that  $[0, \eta]$  contains no critical value of x.

An embedded smooth sub-manifold of  $(M, \partial M)$  is a subset N such that (i) it is a smooth manifold  $(N, \partial N)$  with boundary  $\partial N$  (possibly empty); (ii) its manifold topology coincides with the induced topology from M; (iii) the inclusion mapping  $N \hookrightarrow M$  is a smooth injective immersion.

**Definition 1.4.** ([\[17,](#page-21-3) Section I.7]) Let  $(M, \partial M)$  be a smooth manifold with boundary. A subset N of M is a p-sub-manifold if: (i) it is a smooth sub-manifold with boundary  $\partial N$  of  $M$ ; (ii)  $\partial N$  is contained in  $\partial M$ ; (iii) N is transverse to  $\partial M$ :

$$
\mathbf{x} \in \partial N \implies T_{\mathbf{x}}N + T_{\mathbf{x}}\partial M = T_{\mathbf{x}}M.
$$

The smooth cylinder over a manifold N of height  $\eta$  is

$$
Cyl(N, \eta) := N \times [0, \eta)
$$

and the open cylinder over  $N$  is

$$
\text{Cyl}(N,\eta)^o:=\text{Cyl}(N,\eta)\setminus (N\times\{0\})=N\times (0,\eta)
$$

We recall the collar neighbourhood result below, proof can be found for instance in [\[7\]](#page-20-0).

**Proposition-Definition 1.5.** Let  $(M, \partial M)$  be a manifold with smooth compact boundary. Let x be a boundary defining function. There exist  $\eta > 0$  such that the function x has no critical value in  $[0, \eta]$ , an open neighbourhood  $U_x^{\eta}$  of  $\partial M$  in M and a smooth diffeomorphism

(1.1) 
$$
\phi_x^{\eta}: U_x^{\eta} := M \cap \{x < \eta\} \to \mathrm{Cyl}(\partial M, \eta), \ \mathbf{z} \mapsto (\mathbf{y}(\mathbf{z}), x(\mathbf{z})),
$$

which we call the collar neighbourhood diffeomorphism.

## 1.4. Spherical blowing-up

Let  $(M, g)$  be a Riemannian manifold possibly with non-empty boundary  $\partial M$ . Let a be a point of  $M \setminus \partial M$ . There exists a positive radius  $\varepsilon$  such that the exponential map

$$
\exp_{\mathbf{a}}:B(\mathbf{a},\varepsilon)\to B_g(\mathbf{a},\varepsilon)
$$

centred at **a** maps diffeomorphically the open  $\varepsilon$ -ball  $B(\mathbf{a}, \varepsilon)$  of  $T_{\mathbf{a}}M$  onto the open  $\varepsilon$ -geodesic ball  $B_q(\mathbf{a}, \varepsilon)$  of M. Let  $\mathbf{S}_\mathbf{a}$  be the unit-sphere of  $T_\mathbf{a}M$ . The following mapping

 $\rho_{\mathbf{a}} : \mathbf{S}_{\mathbf{a}} \times [0, 1) \to B(\mathbf{a}, \varepsilon), \ \ (\mathbf{u}, t) \to \varepsilon t \mathbf{u}.$ 

is smooth.

The blowing-up of the Riemannian manifold  $(M, g)$  centred at **a** is the smooth surjective mapping

(1.2) 
$$
\mathbf{bl}_{\mathbf{a}}^M : [M, \mathbf{a}] \to M
$$

where [M, a] is a smooth manifold with boundary  $S_a \times 0 \cup \partial M$ . It is obtained as the gluing of  $\mathbf{S}_a \times [0, 1)$  with  $M \setminus \mathbf{a}$  identifying  $\mathbf{S}_a \times (0, 1)$  with the pointed open ball  $\mathbf{B}_q(\mathbf{a}, \varepsilon) \setminus \mathbf{a}$  via exp<sub>a</sub> ∘ $\rho_{\mathbf{a}}$ . The mapping  $bl_a$  is just the quotient mapping of the gluing.

Therefore, the spherical blow up  $\mathbf{bl}_a$  maps  $\mathbf{S}_a \times 0$  onto a and induces a diffeomorphism  $[M, a] \setminus$  $\mathbf{S}_a \times 0 \to M \setminus \mathbf{a}$ . The pull-backed metric  $(\mathbf{bl}_\mathbf{a}^M)^* g$  is a singular Riemannian metric over  $[M, \mathbf{a}]$ with singular locus  $S_a \times 0$ . In particular, given two Riemannian metrics g and h over M, the spherically blown-up manifolds  $[(M, g), \mathbf{a}]$  and  $[(M, h), \mathbf{a}]$  are diffeomorphic as manifolds with boundary.

### <span id="page-4-2"></span>1.5. Conic metrics

Let  $(M, \partial M)$  be a smooth manifold with compact boundary. We present below the classical material of [\[3,](#page-20-6) [4\]](#page-20-7) with some additional notions necessary for this paper.

**Definition 1.6.** A conic metric  $g^c$  on the smooth manifold  $(M, \partial M)$  is a singular Riemannian metric over M with  $\text{sing}(g^c) = \partial M$ , for which there exists a smooth collar neighbourhood diffeomorphism

<span id="page-4-1"></span>
$$
\varphi = \phi_r^{\eta} : M \cap \{r < \eta\} \to \text{Cyl}(\partial M, \eta)
$$

such that

(1.3) 
$$
\varphi_* g^c = dr^2 + r^2 G(r, \mathbf{y}; dr, dy)
$$

and  $y \to G(0, y)|_{0 \times \partial M}$  is a Riemannian metric over  $\partial M$ . When  $g^c$  is a conic metric, the triple  $(M, \partial M, g^c)$  is said to be a manifold with a conic metric.

Note that the singular locus of conic metrics lies on the boundary of the manifold and if  $(S, \partial S)$  is a p-sub-manifold of the manifold with conic metric  $(M, \partial M, g^c)$ , then  $(S, \partial S, g^c|_S)$  is a manifold with a conic metric.

**Definition 1.7.** The pseudo-distance  $d^c := d_{g^c}$  over M induced by the conic metric  $g^c$  is the conic distance. For a sub-manifold S of M, the pseudo-distance obtained by restricting to S the conic distance  $d^c$  is its outer conic distance whereas the pseudo-distance obtained from the conic metric tensor  $g^c$  restricted to  $S$  is its inner pseudo-distance.

The following special cases of conic metrics will be of great use:

**Definition 1.8.** (1) Let N be a compact manifold without boundary and let  $\eta > 0$ . A model conic metric *on the cylinder* Cyl $(N, \eta)$  is a singular Riemannian metric  $g_{\text{mod}}^c$  over Cyl $(N, \eta)$  of the following form

(1.4) 
$$
g_{\text{mod}}^c = dr^2 + r^2 g^{\partial}(r, \mathbf{y}; \mathrm{d}\mathbf{y})
$$

where  $r \to g^{\partial}(r)$  is a smooth family of Riemannian metrics over N.

(2) A simple conic metric over the manifold  $(M, \partial M)$  with compact boundary is any conic metric g<sup>c</sup> for which there exists a collar neighbourhood diffeomorphism  $\varphi: U_r^n \to Cyl(\partial M, \eta)$ and a Riemannian metric  $g_{\partial M}$  over  $\partial M$  such that

$$
\varphi_* g^c = \mathrm{d}r^2 + r^2 \, g_{\partial M}.
$$

Any model conic metric on a cylinder Cyl(N,  $\eta$ ) over  $N = \partial M$  extends as a conic metric onto  $(M, \partial M)$  and there is no other type of conic metrics. More precisely:

<span id="page-4-0"></span>**Theorem 1.9** (Theorem 1.2, [\[18\]](#page-21-4)). Let  $(M, \partial M, g^c)$  be a manifold with conic metric. There exists a collar neighbourhood diffeomorphism  $\varphi: U \to Cyl(\partial M, \eta)$  such that  $\varphi_* g^c$  is a model conic metric over  $Cyl(\partial M, \eta)$ .

**Example 1.10.** 1) Let  $\mathbf{bl}_0 : [\mathbb{R}^n, \mathbf{0}] = \mathbf{S}^{n-1} \times \mathbb{R}_{\geq 0} \to \mathbb{R}^n$  be the spherical blowing-up of  $\mathbb{R}^n$ . The pull-back  $\mathbf{bl}_0^*(eucl)$  of the Euclidean metric eucl over the blown-up space  $[\mathbb{R}^n, \mathbf{0}]$  is the simplest example of a model conic metric

$$
\mathbf{bl}_0^*(eucl) = dr^2 + r^2 du^2.
$$

2) Warped metrics over  $Cyl(N, \eta)$  are model conic metrics, these are singular metrics of the form

$$
\mathrm{d}r^2 + (rf(r))^2 g_N
$$

where f is a smooth positive function over  $[0, \eta)$  and  $g_N$  is a Riemannian metric over N, see also [\[13\]](#page-21-5).

<span id="page-5-4"></span>An essential by-product of the demonstration of Theorem [1.9](#page-4-0) is the following

**Corollary 1.11.** If  $d^c$  is a conic distance, then the function  $d^c(-, \partial M)$  is a boundary defining function. In particular there exists a positive  $r_0$  such that each level  $\{d^c(-, \partial M) = r\}$  is diffeomorphic to ∂M for any  $0 \le r \le r_0$ .

Moreover, for any two conic metrics g and h over  $(M, \partial M)$  there exists a collar neighbourhood U of  $\partial M$  such that g and h restricted to U are equivalent and thus the identity mapping id :  $(U, d_g) \rightarrow (U, d_h)$  is pseudo-bi-Lipschitz with respect to the conic distances  $d_g$  and  $d_h$ , see Section [1.2.](#page-2-0) In particular, we get

<span id="page-5-1"></span><span id="page-5-0"></span>Corollary 1.12. Any two conic metrics over a compact manifold are equivalent.

**Proposition-Definition 1.13.** Any conic metric  $g^c$  on the manifold M has (1) an associated Riemannian metric on the boundary, *i.e.* a Riemannian metric  $g^{\partial} := G_{|\partial M}$  on  $\partial M$  of representation [\(1.3\)](#page-4-1), and (2) an associated simple conic metric near the boundary  $\partial M$ , i.e. a simple conic metric h on a neighbourhood U of  $\partial M$  such that the singular metrics  $g^c$  and h are equivalent over U and have the same associated Riemannian metric on the boundary.

*Proof.* Let  $g^c$  be a conic metric over  $(M, \partial M)$ . By definition  $g^{\partial} := G_{|\partial M}$  is Riemannian over  $\partial M$ . Let  $\varphi: U_r^{\eta} \to \mathrm{Cyl}(\partial M, \eta)$  be a collar neighbourhood diffeomorphism of Theorem [1.9](#page-4-0) such that  $\varphi_* g^c$  is a model conic metric

$$
\varphi_* g^c = \mathrm{d}r^2 + r^2 g^{\partial}(r).
$$

Thus  $g^{\partial}(0)$  is a Riemannian metric over  $\partial M$ , and the model conic metric

$$
g_0^c = \mathrm{d}r^2 + r^2 g^{\partial}(0)
$$

is equivalent to  $g^c$  over a smaller cylinder Cyl $(\partial M, \eta')$  for some positive height  $\eta' < \eta$ . Since  $\varphi^* g_0^c$  extends as a conic metric  $g^s$  over  $(M, \partial M)$ , the proof ends by Corollary [1.12.](#page-5-0)

<span id="page-5-2"></span>The following, though elementary, is a useful variation of the law of cosines in the conic context:

**Proposition 1.14.** Let  $N$  be a compact smooth manifold without boundary. Let

$$
g^c = \mathrm{d}r^2 + r^2 g^\partial(r)
$$

be a model conic metric on  $Cyl(N, \eta)$ . Denote by  $d^c$  the conic distance on  $Cyl(N, \eta)$  and by  $d_N$ the distance on N induced by  $g^{\partial}(0)$ . Then there exist positive constants  $A^c, B^c$ , depending only on  $g^c$  and  $\eta$ , such that

<span id="page-5-3"></span>(1.5) 
$$
A^{c}(|r - r'| + \min(r, r') d_{N}(\mathbf{y}, \mathbf{y}')) \leq d^{c}(\mathbf{x}, \mathbf{x}') \leq B^{c}(|r - r'| + \min(r, r') d_{N}(\mathbf{y}, \mathbf{y}'))
$$
  
for all  $\mathbf{x} = (\mathbf{y}, r)$  and  $\mathbf{x}' = (\mathbf{y}, r')$  of  $\text{Cyl}(N, \eta)$ .

*Proof.* If  $r \to g^{\partial}(r)$  is constant and equal to a tensor  $g_N$  on  $N \times \{0\}$ , then by straightforward computation we get

$$
\frac{|r-r'|}{2} + \frac{\min(r,r')}{2} d_N(\mathbf{y},\mathbf{y}') \leq d^c(\mathbf{x},\mathbf{x}') \leq |r-r'| + \min(r,r') d_N(\mathbf{y},\mathbf{y}')
$$

for the associated simple conic metric  $dr^2 + r^2g_N$ . The general case is deduced from the constant case above. See also [\[10\]](#page-20-8).  $\Box$ 

### 1.6. Asymptotically conic metrics

Due to considerations similar to the previous Section [1.5](#page-4-2) but for scattering metrics, see [\[19,](#page-21-6) [15\]](#page-21-7), we will give here a simplified presentation of asymptotically conic metrics without taking away from generality, compare Proposition-Definition [1.13](#page-5-1) and Remark [1.16.](#page-6-0)

Let  $M$  be a smooth manifold with compact boundary,  $N$  a smooth manifold without boundary and  $\eta$  a positive number.

**Definition 1.15.** An asymptotically conic metric over the cylinder Cyl $(N, \eta)$  is a smooth Riemannian metric  $g^{\infty}$  on the open cylinder Cyl(N,  $\eta$ )<sup>o</sup> such that the Riemannian metric  $r^4 g^{\infty}$ extends as a smooth model conic metric  $g^0$  over  $Cyl(N, \eta)$ .

The conic metric  $g^0$  is the conic metric associated with the asymptotically conic metric  $g^{\infty}$  and over  $Cyl(N, \eta)^o = N_{\mathbf{y}} \times (0, \infty)_r$  we can write

$$
g^{\infty} = \frac{dr^2}{r^4} + \frac{g^{\partial}(r)}{r^2} = \frac{1}{r^4} g^0
$$

where  $\mathbb{R}_{\geq 0} \ni r \to g^{\partial}(r)$  is a smooth family of Riemannian metrics over N.

An asymptotically conic metric (ac-metric) on the manifold  $(M, \partial M)$  is a Riemannian metric  $g^{\infty}$  on  $M \setminus \partial M$  for which there exists a collar neighbourhood diffeomorphism  $\varphi : U_r^n \to$  $Cyl(\partial M, \eta)$  such that  $\varphi_* g^{\infty}$  is an asymptotically conic metric over  $Cyl(\partial M, \eta)$ .

<span id="page-6-0"></span>The Riemannian metric  $r^2 g^{\infty}|_{\partial M} = g^{\partial}(0)$  on  $\partial M$  is the Riemannian metric on the boundary associated to the ac-metric.

Remark 1.16. An ac-metric is also known as a scattering metric [\[19,](#page-21-6) [15\]](#page-21-7). The standard definition does not require that it be a conformal factor of a model conic metric, but just of a conic metric. Yet, a change of coordinates over  $Cyl(N, \eta)$  allows this in a neighbourhood of the boundary  $N \times 0$ , see [\[15\]](#page-21-7).

Let us illustrate the notion with a simple example.

Example 1.17. Let  $bl_{\infty} : S^{n-1} \times \mathbb{R}_{>0} \to \mathbb{R}^n \setminus 0$  be the spherical blowing-up of  $\mathbb{R}^n$  at infinity, defined by

$$
\mathbf{bl}_{\infty}(\mathbf{u},R) = \frac{\mathbf{u}}{R}.
$$

The pull-back  $\mathbf{bl}_{\infty}^*(eucl)$  over  $\mathbf{S}^{n-1} \times \mathbb{R}_{>0}$  is the simplest example of an asymptotically conic metric

$$
\mathbf{bl}_{\infty}^*(eucl) = \frac{1}{r^4} [\mathrm{d}r^2 + r^2 \mathrm{d}\mathbf{u}^2].
$$

Every ac-metric  $g^{\infty}$  introduces a distance function  $d^{\infty}$  on M such that  $d^{\infty}(\mathbf{x}, \mathbf{y}) = \infty$  only if **x** or y lie in ∂M or in different connected components of M.

**Proposition 1.18.** Let  $Cyl(N, \eta)$  be equipped with an ac-metric  $g^{\infty}$  and the induced distance  $d^{\infty}$ . Let  $g^0 = dr^2 + r^2 g^{\partial}(r)$  be the associated model conic metric and denote  $g_N := g^{\partial}(0)$ . For points  $\mathbf{x} = (\mathbf{y}, r)$  and  $\mathbf{x}' = (\mathbf{y}, r')$  of  $Cyl(N, \eta)$ <sup>o</sup> define

<span id="page-7-1"></span>
$$
e(\mathbf{x}, \mathbf{x}') = \left| \frac{1}{r} - \frac{1}{r'} \right| + \min\left( \frac{1}{r}, \frac{1}{r'} \right) d_N(\mathbf{y}, \mathbf{y}').
$$

Then there exist positive constants  $A^{\infty}, B^{\infty}$ , depending only on  $g^0$  and  $\eta$ , such that

(1.6) 
$$
A^{\infty} e(\mathbf{x}, \mathbf{x}') \leq d^{\infty}(\mathbf{x}, \mathbf{x}') \leq B^{\infty} e(\mathbf{x}, \mathbf{x}')
$$

for all  $\mathbf{x}, \mathbf{x}' \in \text{Cyl}(N, \eta)^o$ .

*Proof.* Proof as in Proposition [1.14,](#page-5-2) see [\[10\]](#page-20-8) and [\[11\]](#page-21-8) as well.

### 1.7. Manifolds with globally conic metrics

Let  $(M, \partial M)$  be a compact smooth manifold.

**Definition 1.19.** A globally conic metric on M is a conic metric over  $M \setminus \partial M_{\infty}$ , where  $\partial M_{\infty}$  a finite union of connected components of  $\partial M$ , which is an asymptotically conic metric on some neighbourhood of  $\partial M_{\infty}$  in M.

Then the triple  $(M, \partial M, g)$  is a manifold with a globally conic metric and the subset  $\partial M_{\infty}$  is the boundary at infinity of  $(M, \partial M, q)$ .

From the point of view of Lipschitz geometry, the following Proposition [1.20](#page-7-0) shows interchangeability of globally conic metrics, similarly to Riemannian structure, compare Section [1.2.](#page-2-0)

<span id="page-7-0"></span>**Proposition 1.20.** Let  $(M, \partial M, q)$  and  $(N, \partial N, h)$  be two smooth manifolds with globally conic metrics and  $\phi : M \to N$  be a diffeomorphism such that  $\phi(\partial M_{\infty}) = \partial N_{\infty}$ . Then the conic metrics  $\phi_*(g)$  and h are equivalent over  $N \setminus \partial N_{\infty}$ . In particular any two globally conic metrics on a compact manifold yielding the same boundary at infinity are equivalent.

*Proof.* By Corollary [1.12](#page-5-0) any two conic metrics over  $Cyl(N, \eta)$  are equivalent over  $Cyl(N, \varepsilon)$  for some positive  $\varepsilon \leq \eta$ . Thus by definition any two ac-metrics over Cyl(N,  $\eta$ ) are equivalent over Cyl $(N, \varepsilon)$ <sup>o</sup> for some positive  $\varepsilon \leq \eta$ . Since the singular locus and boundary at infinity of the globally conic metrics  $\phi_*(g)$  and h are thus equivalent.

### 1.8. Conic inversion

Let  $(N, g_N)$  be a connected compact Riemannian manifold without boundary. Denote by

$$
P = N_{\mathbf{y}} \times (\mathbb{R}_{\geq 0})_r
$$

the infinite (non-negative) cylinder over N. Let  $P^0 = N \times (0, \infty)$  be the open infinite cylinder over N.

The *conic inversion* of an open cylinder  $P^0$  is the mapping

$$
\iota^{0}: P^{0} \to P^{0}, \ \ (\mathbf{y}, r) \mapsto (\mathbf{y}, R) = \left(\mathbf{y}, \frac{1}{r}\right).
$$

Gluing two disjoint copies of P along their interiors  $P^0$  identified by the inversion  $\iota^0$  yields the smooth compact manifold with boundary

$$
\overline{P} = N \times [0, \infty].
$$

The inversion  $\iota^o$  extends onto  $\overline{P}$  as a diffeomorphism  $\iota$ , inverting  $\partial(N \times \mathbb{R}_{\geq 0})$  and  $\partial(N \times (0, \infty])$ . Consider the following singular Riemannian metrics

$$
g^0 = dr^2 + r^2 g^{\partial}(r)
$$
 over  $N_{\mathbf{y}} \times (\mathbb{R}_{\geq 0})_r$  and  $g^{\infty} = dR^2 + R^2 g^{\partial}(R)$  over  $N_{\mathbf{y}} \times (0, \infty]_R$ ;

Both are model conic metrics over their respective domains. On  $P^0$  we get

$$
r^{-4}g^{0} = (\iota^{0})^{*}g^{\infty},
$$

thus  $r^{-4}g^0$  is an asymptotically conic metric over  $P^0$ .

Let  $d^0$  be the conic distance over  $P^0$  obtained from  $g^0$ , and let  $d^{\infty}$  be the distance over  $P^0$ obtained from  $(\iota^0)^* g^{\infty}$ . A simple, though essential, metric property relating conic with asymptotically conic distances is summarized as follows.

<span id="page-8-2"></span>**Proposition 1.21.** There exist positive constants  $A', B'$  such that for all  $\mathbf{x}, \mathbf{x}' \in P^0$  the following holds

(1.7) 
$$
A' \cdot d^{0}(\mathbf{x}, \mathbf{x}') \leq r \cdot r' \cdot d^{\infty}(\mathbf{x}, \mathbf{x}') \leq B' \cdot d^{0}(\mathbf{x}, \mathbf{x}').
$$

Proof. From Estimates [\(1.5\)](#page-5-3) and [\(1.6\)](#page-7-1) we can assume that

 $d^0(\mathbf{x}, \mathbf{x}') = |r - r'| + \min(r, r')d_N(\mathbf{y}, \mathbf{y}'), \text{ and } d^\infty(\mathbf{x}, \mathbf{x}') = |R - R'| + \min(R, R')d_N(\mathbf{y}, \mathbf{y}'),$ with  $R = r^{-1}$  and  $R' = (r')^{-1}$ . We immediately obtain

$$
r \cdot r' \cdot d^{\infty} \ge d^0.
$$

<span id="page-8-0"></span>Distinguishing the cases  $r \geq 2r'$  and  $r' \leq r \leq 2r'$ , we get the other inequality.

#### 2. Globally conic singular manifolds

The interest in metrically conical singularities in Riemannian geometry dates back at least to the seminal works [\[3,](#page-20-6) [5,](#page-20-9) [6,](#page-20-10) [4\]](#page-20-7). Definition presented in this section is a variation on the standard notion of singular manifold with metrically conical points. For more on the local differential geometry of conic singular manifolds see [\[17,](#page-21-3) [12\]](#page-21-9).

We will give natural characterizations of (globally) conic singular manifolds: by an atlas of isometries in Definition [2.1,](#page-8-1) isometric resolution in Section [2.4,](#page-12-0) as a quotient space in Section [2.2](#page-10-0) and by a system of conic charts in Section [2.3.](#page-11-0)

### 2.1. Conic singular manifolds

Recall from Section [1.2](#page-2-0) that if  $M$  is a manifold with a conic metric or an ac-metric, then the metric induces a length distance  $d_M$  on  $M \setminus \partial M$ .

<span id="page-8-1"></span>**Definition 2.1** (Conic singular manifold). A complete length space  $(E, d)$  is an n-dimensional conic singular manifold if there exists a finite subset  $\Sigma_E$  of E, an open covering U of E and smooth n-dimensional manifolds  $M_U$  with compact boundary such that for any  $U \in \mathcal{U}$  one of the following is satisfied

(1) there exists an isometry

$$
\phi_U : (M_U, d_{M_U}) \to (U, d),
$$

where  $(M_U, g_U)$  is a Riemannian manifold with empty boundary

(2) there exists a point  $\mathbf{a} \in U \cap \Sigma_E$  and a continuous surjective mapping  $\phi_U : M_U \to U$  such that the restriction

 $(\phi_U)_{|M_U \setminus \partial M_U} : (M_U \setminus \partial M_U, d_{M_U}) \to (U \setminus \mathbf{a}, d)$ 

is an isometry, where  $(M_U, \partial M_U, g_U)$  is a manifold with a conic metric

The smallest among subsets  $\Sigma_E$  satisfying the definition is called the set  $E_{\text{sing}}$  of singular points of E. Its complement  $E^o$  is the set of smooth points of E.

Moreover, a conic singular manifold is a globally conic singular manifold if additionally

(3) there exist a compact subset K of E such that  $E \setminus K = U \in \mathcal{U}$ , and an isometry

 $\phi_U: (M_U \setminus \partial M_U, d_{M_U}) \to (U, d),$ 

where  $(M_U, \partial M_U, g_U)$  is a manifold with an asymptotically conic metric.



FIGURE 1. A globally conic singular manifold and its smooth model.

Note that the transition mappings  $\phi_U^{-1}$  $U^{-1} \circ \phi_V$ , appropriately restricted, are smooth Riemannian isometries (see [\[20,](#page-21-10) [21\]](#page-21-2)), in particular smooth diffeomorphisms. Thus we get the following Proposition [2.2.](#page-9-0)

<span id="page-9-0"></span>**Proposition 2.2.** If  $(E, d)$  is a (globally) conic singular manifold, then

(1) The set of smooth points  $E^o$  is a smooth manifold.

(2) The Riemannian metrics  $(\phi_U)_*(g_{U|M_U \setminus \partial M_U})$  glue to a Riemannian metric  $g^o$  over  $E^o$ .

(3) The metric spaces  $(E^o, d^o)$  and  $(E^o, d_{|E^o})$  are locally isometric, where  $d^o$  is the distance function induced by the Riemann tensor  $g^o$  on  $E^o$ .

### <span id="page-10-0"></span>2.2. Conic singular manifolds as quotient spaces

Let M be a manifold with conic metric g and denote  $M^o := M \setminus \partial M$ . Recall that the conic tensor induces a pseudo-distance  $d^c$  on M which is infinite between points in different connected components of M and is equal zero between points in the same boundary component.

Let  $\Sigma$  be a finite set and  $\sigma : M \to M^o \cup \Sigma$  a surjective map such that  $\sigma_{|M^o} = id_{M^o}$  and for each boundary component N of  $\partial M$  we have  $\sigma(N) \in \Sigma$ . Let  $M^{\sigma}$  denote the quotient space obtained from M by the mapping  $\sigma$ . With a slight abuse of notation we will say that the mapping  $\sigma : M \to M^{\sigma}$  is a boundary collapsing map.

Let us introduce distance on  $M^{\sigma}$  as

$$
d^{\sigma}(\mathbf{x}, \mathbf{y}) := \min \left( d^{c}(\mathbf{x}, \mathbf{y}), d^{c}(\mathbf{x}, \sigma^{-1}(p_0)) + \sum_{j=1}^{k} d^{c}(\sigma^{-1}(p_{j-1}), \sigma^{-1}(p_j)) + d^{c}(\sigma^{-1}(p_k), \mathbf{y}) \right),
$$

where minimum is taken with respect to all chains of points  $p_0, \ldots, p_k$  in  $\Sigma$ .



FIGURE 2. Distance between points in the quotient space.

<span id="page-10-1"></span>**Proposition 2.3** (Conic singular manifold as quotient space). Let  $\sigma : M \to M^{\sigma}$  be a boundary collapsing map on a manifold M with a conic metric. Then  $(M^{\sigma}, d^{\sigma})$  is a conic singular manifold. If additionally, the manifold M is densely embedded in a compact manifold with a globally conic metric, then  $(M^{\sigma}, d^{\sigma})$  is a globally conic singular manifold.

*Proof.* It suffices to note that the space  $(M^{\sigma}, d^{\sigma})$  is a length space, since  $d^{\sigma}$  is a distance function and the conic distance between a point of  $M$  and a component of its compact boundary is either realized along a smooth arc or infinite.

Note that a boundary collapsing map  $\sigma$  can be alternatively described as a composition of metric identification on each connected component of  $M$  and then glueing points between different connected components, the choice of which points to glue depends on  $\sigma$  and may produce topologically distinct spaces.

#### <span id="page-11-1"></span><span id="page-11-0"></span>2.3. Conic charts

**Proposition-Definition 2.4.** Let  $(E, d)$  be a complete length space such that  $E^o$ , the complement of a finite subset  $\Sigma$ , is a Riemannian manifold without boundary with metric tensor  $g^o$ .

The space  $(E, d)$  is a conic singular manifold of dimension n if and only if every point a of E admits a conic chart  $\varphi_a$  centred at **a**, *i.e.* a continuous surjective mapping

$$
\varphi_{\mathbf{a}} : \text{Cyl}(N_{\mathbf{a}}, \eta) \to U_{\mathbf{a}},
$$

such that

(1)  $U_a$  is an open neighbourhood of  $a$ ,

(2) Cyl(N<sub>a</sub>,  $\eta$ ) is an n-dimensional cylinder over a smooth compact manifold without boundary  $N_{\mathbf{a}}$ ,

(3)  $(\varphi_{\mathbf{a}})|_{\text{Cyl}(N_{\mathbf{a}},\eta)^o} : \text{Cyl}(N_{\mathbf{a}},\eta)^o \to U \setminus \mathbf{a}$  is a diffeomorphism,

(4)  $\varphi_{\mathbf{a}}^* g^o$  extends smoothly as a conic metric over  $Cyl(N_{\mathbf{a}}, \eta)$ .

Moreover,  $(E, d)$  is a globally conic singular manifold if and only if additionally there is an asymptotically conic chart

$$
\varphi_{\infty}: \text{Cyl}(N_{\infty}, \eta)^{o} \to U_{\infty},
$$

where

(1')  $U_{\infty}$  is a complement of a compact subset of E,

(2') Cyl( $N_{\infty}, \eta$ ) is an n-dimensional cylinder over a compact smooth manifold without boundary  $N_{\infty}$ 

 $(3') \varphi_{\infty}$  is a diffeomorphism,

(4')  $\varphi^*_{\infty} g^o$  is an ac-metric over Cyl $(N_{\infty}, \eta)^o$ .

A conic chart is a (asymptotically) model conic chart if the metric in point  $\left(4\right)$  (resp. in point  $(4')$ ) is model.

*Proof.* We check those conditions imply the (globally) conic singular nature of  $(E, d)$  using the pull-backs of  $g^o$ . For  $\mathbf{a} \in E_{sing}$ , let  $\varphi_{\mathbf{a}}$  be the mapping  $\phi_U$  of point (2) of Definition [2.1.](#page-8-1) For  $\mathbf{a} \in U \subset E^o$ , the mapping  $\phi_U \circ \mathbf{bl}_{\mathbf{a}_U}$ , where  $\mathbf{a}_U = \phi_U^{-1}$  $U^{-1}(\mathbf{a})$ , will satisfy the requirements.  $\Box$ 

<span id="page-11-2"></span>Using conic charts, Corollary [1.11](#page-5-4) can be rephrased as

**Lemma 2.5.** Under assumptions and notation of Proposition-Definition [2.4](#page-11-1) let  $d_{\bf a}: E \to \mathbb{R}$  be the distance function to a in the singular conic manifold  $(E, d)$ .

Possibly shrinking  $U_a$ , the function  $\varphi_a^*d_a$  extends as a boundary defining function in Cyl( $N_a$ ,  $\eta$ ). Moreover, if  $\mathbf{a} \in E \backslash U_{\infty}$ , then possibly shrinking  $U_{\infty}$ , the function  $\varphi_{\infty}^*(\frac{1}{d_{\epsilon}})$  $\frac{1}{d_{\mathbf{a}}}$ ) extends as a boundary defining function in  $Cyl(N_{\infty}, \eta)$ .

<span id="page-11-3"></span>**Proposition 2.6.** Let  $(E, d)$  be a (resp. globally) conic singular manifold with two (resp. asymptotically) conic charts  $\varphi_{\bf a} : Cyl(N_{\bf a}, \eta) \to U_{\bf a}$  and  $\varphi'_{\bf a} : Cyl(N'_{\bf a}, \eta) \to U'_{\bf a}$  at a point  ${\bf a} \in E$ (resp.  $\mathbf{a} = \infty$ ). Then  $N_{\mathbf{a}}$  is diffeomorphic to  $N'_{\mathbf{a}}$ .

*Proof.* Since an appropriate restriction of the mapping  $\varphi_a$  is a Riemannian isometry between  $Cyl(N_{\mathbf{a}}, \eta)^{\circ}$  and  $U_{\mathbf{a}} \setminus \mathbf{a}$ , for  $0 < \varepsilon < \eta$  with  $\eta$  small enough, we find that  $(\varphi_{\mathbf{a}}^* d_{\mathbf{a}})^{-1}(\varepsilon)$ ,

 $((\varphi_{\mathbf{a}}')^*d_{\mathbf{a}})^{-1}(\varepsilon)$  and  $(d_{\mathbf{a}})^{-1}(\varepsilon)$  are isometric manifolds by Lemma [2.5,](#page-11-2) using collar neighbourhood diffeomorphisms. Since the levels of a boundary defining function for values small enough are diffeomorphic to the boundary, we get the claim.

Therefore, by Proposition [2.6](#page-11-3) and Theorem [1.9](#page-4-0) our definition of conic charts is not restrictive when we ask the conic metric to be model.

<span id="page-12-0"></span>Corollary 2.7. A complete length space is a conic singular manifold if and only if every point admits model conic charts.

### 2.4. Conic resolution and completion

<span id="page-12-1"></span>From definition of conic singular manifolds and Proposition [2.2](#page-9-0) we obtain immediately the following which can be treated as a converse to Proposition [2.3.](#page-10-1)

**Proposition 2.8** (Isometric resolution of conic singularities). A complete length space  $(E, d)$ is a conic singular manifold if and only if there exists a manifold M with conic metric g and a continuous surjective mapping

$$
\phi: M \to E
$$

such that  $\phi(\partial M) = E_{sing}$  and the restriction  $\phi_{|M\setminus\partial M}$  is a Riemannian isometry over  $E \setminus E_{sing}$ .

A complete length space  $(E, d)$  is a globally conic singular manifold if and only if additionally there exists a compact manifold  $\overline{M}$  such that  $M = \overline{M} \setminus \partial M_{\infty}$  for some union of boundary components  $\partial M_{\infty}$  of  $\overline{M}$  and g is a globally conic metric on  $\overline{M}$ .

Given a (globally) conic singular manifold E, a pair  $(M, \phi)$  as in Proposition [2.8](#page-12-1) will be referred to as a *(global) conic resolution of E with boundary at infinity*  $\partial M_{\infty}$ *.* A (global) conic resolution s essentially unique as the following shows.

<span id="page-12-2"></span>**Proposition 2.9.** Let  $(M_p, \partial M_p, g_p^c)$  be two (resp. globally) conic manifolds for  $p = 1, 2$  such that there exist mappings  $\phi_p : M_p \to E$  satisfying Proposition [2.8.](#page-12-1) Then for  $p = 1, 2$  the manifolds  $(M_p, \partial M_p)$  are diffeomorphic and  $M_p \setminus \partial M_p$  are isometric. Moreover, the boundaries  $\partial M_p$  (resp. and boundaries at infinity  $\partial M_{p,\infty}$ ) are isometric with respect to the Riemannian metric on the boundary associated to the (resp. asymptotically) conic metric.

*Proof.* Let  $M_p^o = M_p \setminus \partial M_p$  and  $\phi_p^o = \phi_p|_{M_p^o}$ . Thus  $(\phi_2^o)^{-1} \circ \phi_1^o : M_1^o \to M_2^o$  is a Riemannian isometry.

Now, let  $N=\cup_{\mathbf{a}\in E_{\text{sing}}}N_{\mathbf{a}}$ . Set

$$
U_\eta:=\{d(-,E_\textup{sing})<\eta\}\ \ \text{and}\ \ \Sigma_\eta:=\{d(-,E_\textup{sing})=\eta\}.
$$

If  $\eta$  is small enough, we define the mapping  $\varphi_{\eta}: \mathrm{Cyl}(N, \eta) \to U_{\eta}$  as the union  $\Box_{\mathbf{a} \in E_{\text{sing}}} \varphi_{\mathbf{a}}$  of the conic charts centred at  $E_{\text{sing}}$ . Lemma [2.5](#page-11-2) implies that  $\Sigma_{\varepsilon}$  is diffeomorphic to N for  $0 < \varepsilon < \eta$ . Thus  $(\phi_p^o)^{-1} \circ \varphi_\eta : \text{Cyl}(N, \eta) \setminus N \to U_\eta^p \setminus \partial M_p$  is an isometry, where

$$
U_{\eta}^{p} = \phi_{p}^{-1}(U_{\eta}) = \{d_{p}^{c}(-, \partial M_{p}) < \eta\} \text{ and } \Sigma_{\eta}^{p} := \phi_{p}^{-1}(\Sigma_{\eta}) = \{d_{p}^{c}(-, \partial M_{p}) = \eta\}.
$$

Thus  $\phi_p^*(d(-, E_{sing}))$  extends as a boundary defining function of  $\partial M_p$  by Lemma [2.5,](#page-11-2) so that we deduce  $\partial M_p = N$ . This is enough to guarantee that  $U^p_\eta$  is isometric to Cyl $(N, \eta)$  via a collar neighbourhood diffeomorphism of  $\partial M_p$  and a smaller  $\eta$ . Then we glue isometrically. Isometry on the boundary follows from consideration as in Proposition [2.6.](#page-11-3) **Proposition-Definition 2.10.** For any globally conic singular manifold  $(E, d)$  there exists a compact conic singular manifold  $(\overline{E}, \overline{d})$  called a conic completion of  $(E, d)$  such that

(1) the space E is topologically embedded in  $\overline{E} = E \sqcup \infty_E$ , where the set of conic ends  $\infty_E$  of E is finite

(2) the distances d and  $\overline{d}$  are equivalent over any compact subset of E

(3) there exists a neighbourhood U of  $E_{\text{sing}}$  such that  $d_{|U} = \overline{d}_{|U}$ 

(4) the global conic resolution space  $\overline{M} = M \cup \partial M_{\infty}$  of  $(E, d)$  is diffeomorphic to the conic resolution space of  $(E, d)$ 

(5) the Riemannian tensor  $\bar{g}^o$  on  $\bar{E}^o$  has the same associated Riemannian metric on  $\partial M_{\infty}$  as the tensor  $g^o$  of  $E^o$ 

*Proof.* Let  $(E, d)$  be a globally conic singular manifold and  $\overline{M} = M \cup \partial M_{\infty}$  be its global conic resolution with the globally conic metric  $g$ . By Proposition [2.8](#page-12-1) the topological space  $E$ embedds isometrically as an open dense subset into  $\overline{E} := E \cup \partial M_{\infty}$ . By Proposition [2.9](#page-12-2) the set  $\partial M_{\infty}$  is unique up to diffeomorphism and has k connected components. If  $\sigma : M \to E$  is the boundary collapsing function of  $E$ , then let us extend it as a boundary collapsing function  $\sigma : \overline{M} \to E \sqcup \{p_1,\ldots,p_k\}$  so that each boundary component of  $\partial M_\infty$  maps to a different point  $p_j \in \infty_E := \{p_1, \ldots, p_k\}.$ 

Let  $\overline{q}$  be the conic metric on  $\overline{M}$  obtained as a smooth glueing of the conic metric q restricted to a neighbourhood of  $\partial M$  with a conic metric h over  $\overline{M}$ , having the same Riemannian metric associated on  $\partial M_{\infty}$  as the ac-metric g near  $\partial M_{\infty}$ .

Let now  $\overline{E}$  be equipped with the distance  $\overline{d} := d^{\overline{\sigma}}$  constructed from  $(\overline{M}, \overline{g})$  as in Section [2.2.](#page-10-0)

Claim (1) is obvious. Claim (2) follows from Corollary [1.12](#page-5-0) and claims (3) from choice of boundary collapsing function and the tensor  $\overline{q}$ . Point (4) is obvious since by construction M is the resolution space for both E and  $\overline{E}$  which is unique up to diffeomorphism by Proposition [2.9.](#page-12-2) Thus (5) follows from (4) and the construction of  $\overline{q}$ .

A conic completion can be considered as a compactification of each end of E by one point, and finding an appropriate bounded metric on this compactification. Note that a conic completion can be a smooth manifold, for instance the n-dimensional sphere with standard distance is a conic completion of  $\mathbb{R}^n$ , so although  $\overline{E}_{\text{sing}}$  contains  $E_{\text{sing}}$ , it may differ from  $E_{\text{sing}} \cup \infty_E$ .

The conic completion is unique up to bi-Lipschitz homeomorphism as the following shows.

**Proposition 2.11.** For any two conic completions  $(\overline{E}_p, \overline{d}_p)$ ,  $p = 1, 2$ , of a globally conic singular manifold  $(E, d)$  there exists a bi-Lipschitz homeomorphism  $(\overline{E}_1, \overline{d}_1) \rightarrow (\overline{E}_2, \overline{d}_2)$ , which is a diffeomorphism  $\overline{E}_1 \setminus \overline{E}_{1,\text{sing}}$  onto  $\overline{E}_2 \setminus \overline{E}_{2,\text{sing}}$ .

*Proof.* For  $p = 1, 2$  let  $(\overline{M}_p, \partial \overline{M}_p)$  be globally conic resolution spaces of  $(E, d)$  which yield the conic completions  $\overline{E}_p$  when equipped with conic metrics  $h_p$  by the boundary collapsing maps  $\sigma_p : \overline{M}_p \to \overline{E}_p$ . Let  $\psi : (\overline{M}_1, \partial \overline{M}_1) \to (\overline{M}_2, \partial \overline{M}_2)$  be the diffeomorphism of Proposition [2.9.](#page-12-2) Since  $\Psi$  is an isometry outside the boundary, we necessarily get  $\psi(\partial \overline{M}_{1,\infty}) = \partial \overline{M}_{2,\infty}$ and the mapping  $\Psi := \sigma_2 \circ \psi \circ \sigma_1$  yields a homeomorphism  $(\overline{E}_1, \overline{d}_1) \to (\overline{E}_2, \overline{d}_2)$ , which is a diffeomorphism outside the singular points. Since  $\psi_*(h_1)$  is a conic metric over  $(\overline{M}_2, \partial \overline{M}_2)$ , by Corollary [1.12](#page-5-0) it is bi-pseudo-Lipschitz with respect to conic distances introduced by the conic

metrics  $h_p$ . By the quotient construction of conic completions, the mapping  $\Psi$  is bi-Lipschitz. If a point of  $\overline{E}_p \setminus \infty_E$  is smooth, then  $\Psi$  extends as a diffeomorphism through this point, thus  $\Psi$  is a diffeomorphism  $\overline{E}_1 \setminus \overline{E}_{1,\text{sing}}$  onto  $\overline{E}_2 \setminus \overline{E}_{2,\text{sing}}$ .

In this context we obtain the global version of Proposition [1.21](#page-8-2) as follows, which is a central result of the paper.

<span id="page-14-0"></span>**Corollary 2.12.** Let  $(\overline{E}, \overline{d})$  be a conic completion of a globally conic manifold  $(E, d)$ . There exist positive constants A, B such that

$$
A\overline{d}(\mathbf{x}, \mathbf{x}') \leq r r' d(\mathbf{x}, \mathbf{x}') \leq B\overline{d}(\mathbf{x}, \mathbf{x}'),
$$

for  $\mathbf{x}, \mathbf{x}' \in E$  with  $r := \overline{d}(\mathbf{x}, \infty_E)$  and  $r' := \overline{d}(\mathbf{x}', \infty_E)$ .

*Proof.* Let  $\overline{\varphi}$ :  $\overline{P} = Cyl(\overline{N}, \overline{\eta}) \rightarrow \overline{U}$  be the union of conic charts of  $\overline{E}$  centred at all the points of  $\infty_E$ . Let  $\overline{g}^c$  be the conic metric over  $\overline{P}$  consistent with the pull-back  $\overline{\varphi}^*(\overline{d}|_{\overline{U}})$ . We can assume that  $\varphi$ , the restriction of  $\overline{\varphi}$  to  $\overline{P}^{\circ}$  is an asymptotic chart of E. Let  $g^{\infty}$  be the asymptotically conic metric with distance function equal to the pull-back  $\varphi^*(d|_{E \cap \overline{U}})$ . Therefore the desired estimates holds true in  $E \cap \overline{U}$  by Proposition [1.21](#page-8-2) used with  $\overline{d}^c$  and  $d^{\infty}$ , pseudo-distances induced respectively from  $\overline{g}^c$  and  $g^{\infty}$ , thus holds in  $V = \text{clos}_E(\overline{U})$ . Since  $K = \overline{E} \setminus \overline{U}$  is a compact subset of E containing a neighbourhood of  $E_{sing}$ , both  $d, \overline{d}$  are bounded over K and thus r, r,' are bounded above and below over K, therefore the estimates hold over K. Let  $\mathbf{x} \in K$  and  $\mathbf{x}' \in \overline{U}$ . Let  $\mathbf{y} \in K \cap V$  such that  $d(\mathbf{x}, \mathbf{x}') = d(\mathbf{x}, \mathbf{y}) + d(\mathbf{y}, \mathbf{x}')$ . Thus we deduce from the estimates in K and in V that  $A\overline{d}(\mathbf{x}, \mathbf{x}') \leq r \cdot r' \cdot d(\mathbf{x}, \mathbf{x}')$ . The remaining estimate is obtained similarly.  $\Box$ 

### 2.5. Globally conic singular sub-manifolds

The following definition coincides with the standard definition, for instance for spherical blowings-up, compare [\[7\]](#page-20-0). Let  $(E, d)$  be a (globally) conic singular manifold. The *strict transform of a* closed subset X of E by a conic chart  $\varphi_a$  for  $a \in E \cup \{\infty\}$  is the subset of Cyl( $N_a, \eta$ ) defined as

$$
\mathrm{st}_{\mathbf{a}}X := \mathrm{clos}_{\mathrm{Cyl}(N_{\mathbf{a}}, \eta)}(\varphi_{\mathbf{a}}^{-1}(X \setminus \mathbf{a})).
$$

**Definition 2.13.** Let X be a closed subset of a conic singular manifold E such that outside of a finite subset it is a sub-manifold of  $E^o$ . The subset X is:

(1) a conic singular sub-manifold of E if its strict transform by every conic chart is a p-submanifold nearby the boundary of the conic chart domain,

 $(2)$  an asymptotically conic singular sub-manifold of the globally conic singular manifold E if its strict transform by the asymptotic conic chart is a p-sub-manifold nearby the boundary of the conic chart domain,

 $(3)$  a globally conic singular sub-manifold of the globally conic singular manifold E if it is both conic singular and asymptotically conic singular sub-manifold.

<span id="page-14-1"></span>**Proposition 2.14.** A (globally) conic singular sub-manifold equipped with the inner metric is a (globally) conic singular manifold.

*Proof.* The restriction of a(n asymptotic) conic chart to a p-sub-manifold yields a(n asymptotic) conic chart onto the (globally) conic singular sub-manifold and the (globally) conic metric



Figure 3. A conic singular sub-manifold and its smooth model as a p-sub-manifold.

restricted to a p-sub-manifold is a (globally) conic metric on the p-sub-manifold. The (globally) conic singular sub-manifold with its inner metric is also a length space, where we can point out the length-minimizing curves using for instance the quotient construction of Section [2.2.](#page-10-0)  $\Box$ 

**Definition 2.15.** Let X be a closed subset of the globally conic singular manifold  $(E, d)$ . The conic completion in  $\overline{E}$  of the subset X is simply its closure in the conic completion  $\overline{E}$ .

Note that the conic completion of a globally singular sub-manifold  $X$  in  $E$  is due to Proposi-tion [2.14](#page-14-1) isometric to a conic completion of  $X$ , whenever  $X$  is considered with the inner metric. Therefore, conic completion gives a convenient way to characterize a globally conic singular manifold as follows.

<span id="page-15-1"></span><span id="page-15-0"></span>**Proposition 2.16.** Let  $(E, d)$  be a globally conic singular manifold and let  $(\overline{E}, \overline{d})$  be its conic completion. The closed subset  $X$  is a globally conic singular sub-manifold of  $E$  if and only if its conic completion  $\overline{X}$  in the conic completion  $\overline{E}$  is a conic singular sub-manifold.

## 3. Lipschitz geometry on globally conic singular manifolds

By definition, a manifold which is Riemannian, conic singular or globally conic singular is LNE in itself as a length space. A smooth sub-manifold of a Riemannian manifold is always locally LNE, yet it may fail to be LNE, see [\[7\]](#page-20-0). In this section we show that conic singular sub-manifolds are locally LNE and globally conic singular manifolds are LNE in an ambient globally conic singular manifold, generalizing results of [\[7\]](#page-20-0).

### 3.1. Preliminary remarks on LNE subsets of conic singular manifolds

The next result generalizes [\[1,](#page-20-4) Proposition 2.1],[\[16,](#page-21-11) Proposition 2.4] and [\[9,](#page-20-2) Lemma 2.6] in the context of conic singular manifolds.

<span id="page-15-2"></span>**Proposition 3.1.** A connected compact subset of the conic singular manifold  $(E, d)$  is LNE if and only if it is locally LNE. In particular, any connected compact  $C^1$  embedded sub-manifold of  $E^o$ , possibly with boundary, is LNE.

The LNE property is hereditary in the sense of Proposition [3.2](#page-16-0) below. Equivalence of induced metrics on an LNE subset shows that LNE subsets preserve ambient Lipschitz properties, for example Hölder exponents.

<span id="page-16-0"></span>**Proposition 3.2** (Proposition 1.9, [\[7\]](#page-20-0)). Let X be a subset of a conic singular manifold  $(E, d)$ which is locally LNE at a point **x**. A subset Y of X is locally LNE at **x** in  $(X, d_{\text{in}}^X)$  if and only if it is locally LNE at  $x$  in  $(E, d)$ .

<span id="page-16-1"></span>The next result is the following analogue of a cone type result in the Euclidean case.

**Proposition 3.3.** Let  $(E, d)$  be a conic singular manifold and  $\mathbf{a} \in E_{sing}$ . Let  $\phi : \text{Cyl}(N, \eta) \to U$ be a model conic chart on E centred at **a**, let  $g^c$  be the model conic metric on  $Cyl(N, \eta)$  and  $g_N$  be the Riemannian metric over N associated with  $g^c$ . If Y is a LNE subset of  $(N, g^{\partial}(0))$ then there exists a positive height  $\eta_Y < \eta$  such that each image  $\phi(Y \times [0, \varepsilon])$  is LNE in  $(E, d)$ for any  $0 < \varepsilon \leq \eta_Y$ .

*Proof.* The model conic metric  $g^c$  writes as

$$
g^c := \mathrm{d}r^2 + r^2 g^{\partial}(r)
$$

with  $g_N = g^{\partial}(0)$ . By Corollary [1.12,](#page-5-0) up to shrinking  $\eta$ , it is equivalent as a singular Riemannian metric to the simple conic metric

$$
h^c = \mathrm{d}r^2 + r^2 g_N
$$

In particular the inner conic distances induced respectively by  $g^c$  and  $h^c$  over any subset of  $Cyl(N, \eta)$  are equivalent.

Let  $d^N$  be the distance on N induced by  $g_N$ , and  $d_{h^c}$  be the conic distance induced by  $h^c$  on  $Q = \text{Cyl}(N, \eta)$ . Let X be a LNE subset of N. We recall Estimates [\(1.5\)](#page-5-3)

$$
\frac{|r-r'|}{2}+\min(r,r')\frac{d_N(\mathbf{y},\mathbf{y}')}{2} \leq d_{h^c}(\mathbf{x},\mathbf{x}') \leq |r-r'|+\min(r,r')d_N(\mathbf{y},\mathbf{y}').
$$

Let  $d_N^Y$  be the inner distance on Y obtained for  $g_N$ , and let  $d_{hc}^Y$  be the inner conic distance over Q obtained from  $h^c$ , that is the inner distance over  $Q \setminus \partial Q$ . Let L be a LNE constant for Y. For points  $\mathbf{x}, \mathbf{x}' \in Q$  with  $r' \geq r$ , then clearly

$$
d_{h^c}^Q(\mathbf{x}, \mathbf{x}') \ge |r - r'| \text{ and } d_{h^c}^Q(\mathbf{x}, \mathbf{x}') \ge r L d_N(\mathbf{y}, \mathbf{y}')
$$

Since the following estimates also holds true

$$
d_{h^c}^Q(\mathbf{x}, \mathbf{x}') \le |r - r'| + r L d^N(\mathbf{y}, \mathbf{y}')
$$

we conclude that any cylinder  $Cyl(X, \varepsilon)$  with  $\varepsilon \leq \eta$  is LNE w.r.t. to  $h^c$ .

Let  $d_{q^d}^Q$  $Q_{\sigma^c}$  be the inner (pseudo-)distance induced by  $g^c$  over Q and let  $X := \phi(Q)$ . By definition the spaces  $(Q, d_{g^c}^Q)$  and  $(X, d_{\text{in}}^X)$ , are locally (pseudo-)isometric. Since  $g^c$  and  $h^c$  are equivalent over  $Q$ , the result is proved.

### <span id="page-16-2"></span>3.2. Conic singular sub-manifolds are locally Lipschitz normally embedded

**Theorem 3.4.** Let X be a subset of the conic singular manifold  $(E, d)$  such that the germ  $(X, a)$  at the point  $a \in E$  is that of a closed conic singular sub-manifold. Then X is locally LNE at a.

*Proof.* Without loss of generality we can assume that  $X \setminus \mathbf{a}$  is a smooth sub-manifold of a smooth manifold  $E \setminus \mathbf{a}$ . There exists a conic chart  $\phi : P = \text{Cyl}(N, \eta) \to U$ , centred at  $\mathbf{a}$  in E such that  $\phi^* g^o$  extends smoothly as the model conic metric  $g^c$  over P.

By definition, the closure  $\mathcal X$  of  $\phi^{-1}(X \setminus \mathbf{a})$  in P is a p-sub-manifold. It suffices to show that the outer conic distance and inner conic distance over  $\mathcal X$  obtained from the conic metric  $g^c$  are equivalent.

Since  $(\mathbf{y}, r)$  are global coordinates over P, the model conic metric  $g^c$  writes

$$
g^c = dr^2 + r^2 g^{\partial}(r).
$$

Since the LNE property by Corollary [1.12](#page-5-0) stays true regardless of the conic metric we are working with, we can assume without loss of generality, up to shrinking  $\eta \leq 1$ , that  $g^c$  is a simple conic metric, i.e. the family  $r \to g^{\partial}(r)$  is constant and equal to the associated Riemmanian metric on the boundary  $g_N := g^{\partial}(0)$  and there exists a collar neighbourhood diffeomorphism

$$
\tau: \mathcal{X} \to \mathcal{C} = \partial \mathcal{X} \times [0, \eta), \quad (\mathbf{y}, r) \mapsto (\omega(\mathbf{y}, r), r)).
$$

<span id="page-17-0"></span>**Lemma 3.5.** The mapping  $\phi \circ \tau \circ \phi^{-1}$  :  $(\phi(\mathcal{X}), d) \to (\phi(\mathcal{C}), d)$  is bi-Lipschitz for the outer metric space structures.

Proof. Consider P equipped with the product metric

$$
cyl := dr^2 + g_N.
$$

Let  $| - |_{cyl}$  be the induced norm on TP and  $d^{cyl}$  be the induced distance on P. Let  $| - |_N$  be the norm on TN obtained from  $g_N$ , and  $d^N$  be the induced distance on N. Last, let  $|-|c|$  be the semi-norm induced by  $g^c$  on TP and  $d^c$  be the conic distance on P. We recall that, given  $\mathbf{v} = (\xi, \lambda) \in T_{(\mathbf{y}, r)}P = T_{\mathbf{y}}N \times \mathbb{R}$ , the norms of  $\mathbf{v}$  are as follows

$$
|\mathbf{v}|_{\text{cyl}}^2 = \lambda^2 + |\xi|_N^2
$$
 and  $|\mathbf{v}|_c^2 = \lambda^2 + r^2 |\xi|_N^2$ .

Observe that the identity mapping  $(P, d^{\text{cyl}}) \rightarrow (P, d^c)$  is (pseudo-)Lipschitz with Lipschitz constant 1 whenever  $\eta \leq 1$ .

Observe that the restriction of  $| - |_{cyl}$  to TN is simply  $| - |_{N}$ . Given  $z = (y, r) \in \mathcal{X}$ , the (restriction to TX of the) cyl-norms for the derivatives of smooth mappings  $\tau$  and  $\omega$  are

$$
||D_{\mathbf{z}}\tau||_{\text{cyl}} \text{ and } ||D_{\mathbf{z}}\omega||_{\text{cyl}},
$$

Analogously, over  $\mathcal{X} \setminus \partial \mathcal{X}$ , the (restriction to  $T\mathcal{X}$  of the)  $g^c$ -norms for  $\tau$  and  $\omega$  is denoted by

$$
||D_{\mathbf{z}}\tau||_c \text{ and } ||D_{\mathbf{z}}\omega||_c,
$$

Given a subset S of P and  $r \geq 0$ , let

$$
S_r := S \cap (N \times r).
$$

Since  $\omega|_{\partial \mathcal{X}\times 0}$  is a diffeomorphism and  $\tau(\mathcal{X}_r) = \mathcal{C}_r$ , the mapping  $\partial_N \omega(\mathbf{z}) = D_{\mathbf{z}} \omega|_{T_{\mathbf{z}}\mathcal{X}_r}$  is invertible, up to shrinking  $\eta$  there exists a positive constant K such that

$$
\frac{1}{K} \leq \|\partial_N \omega(\mathbf{z})\|_{\text{cyl}} \leq K, \ \forall \mathbf{z} \in \mathcal{X}.
$$

Observe that

$$
\|\partial_N \omega(\mathbf{z})\|_{\text{cyl}} = \|\partial_N \omega(\mathbf{z})\|_{\text{c}}.
$$

Let  $\mathbf{v} = (\xi, \lambda) \in T_{\mathbf{z}}\mathcal{C} = T_{\mathbf{y}}\partial \mathcal{X} \times \mathbb{R} \subset T_{\mathbf{z}}P = T_{\mathbf{y}}N \times \mathbb{R}$  with  $\mathbf{z} \in \mathcal{X}$ . We get

 $D_{\mathbf{z}}\omega \cdot \mathbf{v} = \partial_N \omega(\mathbf{z}) \cdot \mathbf{v} + \lambda \partial_r \omega.$ 

Thus there exists a positive constant A so that

$$
|(\partial_r \omega)(\mathbf{z})|_{\text{cyl}} \leq A \cdot ||\partial_N \omega(\mathbf{z})||_{\text{cyl}}, \quad \forall \mathbf{z} \in \mathcal{X},
$$

which yields

$$
|D_{\mathbf{z}}\omega \cdot \mathbf{v}|_c = r \cdot |D_{\mathbf{z}}\omega \cdot \mathbf{v}|_{\text{cyl}} \le rB \cdot ||\partial_N \omega(\mathbf{z})||_{\text{cyl}} \cdot |\mathbf{v}|_{\text{cyl}}, \text{ where } B = \sqrt{1 + A^2}.
$$

Thus

(3.1) 
$$
||D_{\mathbf{z}}\omega \cdot \mathbf{v}||_{c} \leq B \cdot ||\partial_{N}\omega(\mathbf{z})||_{cyl}, \quad \mathbf{z} \in \mathcal{X}.
$$

Let  $\mathbf{1}_{\mathbb{R}}$  be the identity mapping of  $\mathbb{R}$ , the latter being identified with  $T_{\mathbf{z}}(0, r_0)$  and  $T_{\tau(\mathbf{z})}(0, r_0)$ . Since  $D_{\mathbf{z}}\tau = D_{\mathbf{z}}\omega \oplus \mathbf{1}_{\mathbb{R}}$ , we deduce that

$$
||D_{\mathbf{z}}\tau||_{c} \leq 1 + r \cdot ||D_{\mathbf{z}}\omega|| \leq 1 + B \cdot |\partial_{N}\omega(\mathbf{z})||_{cyl}
$$

Thus  $\tau$  induces (pseudo-)Lipschitz homeomorphisms  $(\mathcal{X}_i, d^c) \to (\mathcal{C}_i, d^c)$ , where  $\mathcal{X}_1, \ldots, \mathcal{X}_s$ , are the connected components of X and  $\mathcal{C}_i = \tau(\mathcal{X}_i)$ . Since the smooth inverse  $\tau^{-1}$  is of the form  $(r, y) \rightarrow (\mu(r, y), r)$  the exact same arguments show that  $\tau^{-1}$  is also (pseudo-)Lipschitz with respect to the outer conic metrics  $d^c$  over each  $\mathcal{C}_i$ .

Since  $\partial X = \bigcup_{i=1}^s S_i$ , where  $S_i$  is a smooth compact connected sub-manifold of  $S^{n-1}$  if  $a \in E_{\text{sm}}^d$ or of N if  $\mathbf{a} \in E_{\text{sing}}^d$ , and  $\mathcal{S}_i \cap \mathcal{S}_j = \emptyset$  for  $1 \leq i < j \leq s$ , we deduce that

$$
\delta := \min\{d^{\text{cyl}}(\mathcal{S}_i, \mathcal{S}_j) : 1 \leq i < j \leq k\} > 0.
$$

From which follows that for any  $r \leq \eta$ , up to shrinking  $\eta$ , we get

$$
\min\{d^c(\mathcal{X}_i)_r,(\mathcal{X}_j)_r):1\leq i
$$

Fix a pair  $1 \leq i < j \leq s$  and let  $\mathbf{z}_k = (\mathbf{y}_k, r_k) \in \mathcal{X}_k$  with  $k = i, j$ . Then

$$
d^{c}(\mathbf{z}_{i}, \partial P) + d^{c}(\partial P, \mathbf{z}_{j}) \geq d^{c}(\mathbf{z}_{i}, \mathbf{z}_{j}) \geq d^{c}((\mathcal{X}_{i})_{r_{i}}, (\mathcal{X}_{j})_{r_{j}}).
$$

Since  $\tau|_{\partial X}$  is the identity mapping of  $\partial X$ , up to shrinking  $\eta$ , we deduce the following estimates

$$
r_i + r_j \ge d^c(\mathbf{z}_i, \mathbf{z}_j) \ge \frac{\delta}{4} [r_i + r_j]
$$

When  $\mathbf{w}_k = \tau(\mathbf{z}_k) = (\omega_k, r_k)$ , for  $k = i, j$ , we get

$$
d^{c}(\mathbf{w}_{i}, \mathbf{w}_{j}) \leq d^{c}(\mathbf{w}_{i}, \partial M) + d^{c}(\partial M, \mathbf{w}_{j}) = r_{i} + r_{j} \leq \frac{4}{\delta} d^{c}(\mathbf{z}_{i}, \mathbf{z}_{j})
$$

We conclude that  $\tau : (\mathcal{X} \setminus \partial \mathcal{X}, d^c) \to (\mathcal{C} \setminus \partial \mathcal{C}, d^c)$  is (pseudo-)Lipschitz. The same arguments will give that  $\tau^{-1}$  is (pseudo-)Lipschitz as well.

Since the image of the cylinder  $\phi(\mathcal{C})$  is LNE with respect to the metric d by Proposition [3.3,](#page-16-1) so is  $\phi(\mathcal{X})$ . Therefore X is locally LNE at **a**.

By Theorem [3.4,](#page-16-2) Lemma [3.5](#page-17-0) and Proposition [3.3](#page-16-1) we get a family of LNE representatives as follows.

**Corollary 3.6.** If X is a germ at a point a of a conic singular sub-manifold in a conic singular manifold  $(E, d)$ , then there exists a positive number  $\eta_X$  and a germ of a non-negative function  $r:(E, \mathbf{a}) \to \mathbb{R}$ , smooth outside of  $\mathbf{a}$  and vanishing only at the point  $\mathbf{a}$  such that  $X \cap r^{-1}([0, \eta])$ is locally LNE at **a** for every  $\eta < \eta_X$ .

Theorem [3.4](#page-16-2) and Proposition [3.1](#page-15-2) yield the following relevant implications.

<span id="page-19-1"></span>Corollary 3.7. A conic singular sub-manifold of a conic singular manifold is locally LNE.

Corollary 3.8. A connected compact conic singular sub-manifold of a conic singular manifold is LNE.

### <span id="page-19-0"></span>3.3. Globally conic singular sub-manifolds are Lipschitz normally embedded

**Theorem 3.9.** Let X be a globally conic singular sub-manifold of the globally conic singular manifold  $(E, d)$ . Then each connected component of X is LNE in E.

Proof. Without loss of generality we can assume that X is connected.

Let  $(\overline{E}, \overline{d})$  be a conic completion of  $(E, d)$  and let  $Y := \overline{X}$  be the closure of X in  $\overline{E}$ . Let  $\infty_X = \infty_E \cap Y$  be the conic ends of X. Since  $(Y, \overline{d}_{in}^Y)$  is a conic completion of the globally conic singular manifold  $(X, d_{\text{in}}^X)$ , we get

$$
\overline{d}_{\text{in}}^Y|_X = \overline{d}_{\text{in}}^X.
$$

Since Y is a conic singular sub-manifold of the compact conic singular manifold  $\overline{E}$ , by Corol-lary [3.8](#page-19-1) it is LNE, thus there exits a positive constant  $L'$  such that

$$
\overline{d}_Y \le \overline{d}_{\text{in}}^Y \le L' \overline{d}_Y.
$$

Corollary [2.12,](#page-14-0) restricting from  $\overline{E}$  to Y, implies there exist positive constants  $A', B'$  such that

$$
A'\overline{d}_Y(\mathbf{x}, \mathbf{x}') \leq r r' d_X(\mathbf{x}, \mathbf{x}') \leq B'\overline{d}_Y(\mathbf{x}, \mathbf{x}').
$$

Define  $\tilde{r} = \overline{d}_{\text{in}}^Y(\mathbf{x}, \infty_X)$  and  $\tilde{r}' = \overline{d}_{\text{in}}^Y(\mathbf{x}', \infty_X)$ , Corollary [2.12](#page-14-0) applied to Y with the inner distance  $\overline{d}_{\text{in}}^Y$  gives positive constants  $A'', B''$  such that

$$
A''\overline{d}_{\rm in}^Y(\mathbf{x},\mathbf{x}') \leq \tilde{r}\tilde{r}' d_{\rm in}^X(\mathbf{x},\mathbf{x}') \leq B''\overline{d}_{\rm in}^Y(\mathbf{x},\mathbf{x}').
$$

Therefore over  $X$  we get

$$
d_X \le d_{\text{in}}^X \le L d_X \quad \text{with} \quad L = \frac{B''}{A'} (L')^3,
$$

since  $\overline{d}_Y \leq \overline{d}_{in}^Y \leq L' \overline{d}_Y$ , yielding  $r \leq \tilde{r} \leq L'$ 

By Theorems [3.4](#page-16-2) and [3.9](#page-19-0) we easily get the following variation, of interest for applications.

**Corollary 3.10.** Let  $(E, d)$  be a globally conic singular manifold and let X be its closed connected subset which is asymptotically conic. Then X is LNE in  $(E, d)$  if and only if its conic completion  $\overline{X}$  is LNE in  $(\overline{E}, \overline{d})$ .

### 3.4. Examples

The Euclidean space  $(\mathbb{R}^n, |-|)$  is naturally a globally conic singular manifold and its conic completion is simply  $S<sup>n</sup>$  via the inverse of the stereographic projection. The main result of

 $r$ .

[\[8\]](#page-20-3) states that each connected component of a generic real algebraic set of  $\mathbb{R}^n$  is a globally conic-singular sub-manifold (usually non-singular) of  $\mathbb{R}^n$ , thus is LNE.

Consider the following singular Riemannian metric over  $\mathbb{R}^2$ :

$$
h = [2x^{2} - 2xy + y^{2}]dx^{2} + 2[x^{2} + xy - y^{2}]dxdy + [x^{2} + 2xy + 2y^{2}]dy^{2}.
$$

Denoting  $r = \sqrt{x^2 + y^2}$ , we check that h is equivalent to  $r^2$ eucl over  $\mathbb{R}^2$ , where eucl is the Euclidean metric tensor. Therefore  $g^o = r^{-2}h$  is a Riemannian metric over  $E^o = \mathbb{R}^2 \setminus \mathbf{0}$ . Let  $E = \mathbb{R}^2$  and d be the length distance over E induced from  $g^o$ . Since  $g^o$  is equivalent to  $eucl|_{E^o}$ , the space( $E, d$ ) is a complete length space. In polar coordinates, we find

$$
g^o = 2dr^2 + 2r dr d\theta + r^2 d\theta^2.
$$

The following mapping is continuous and surjective

$$
\phi: M = \mathbf{S}^1 \times \mathbb{R}_{\geq 0} \to E, \ \ (\theta, r) \mapsto \begin{cases} \ r e^{i(\theta - \ln r)} & \text{if } r > 0 \\ 0 & \text{if } r = 0 \end{cases},
$$

and over  $M \setminus \partial M$ , we verify that

$$
\phi^*(g^o) = dr^2 + r^2 d\theta^2,
$$

thus extends as a conic metric over M. Thus  $(E, d)$  is a conic singular manifold. Using the conic inversion and  $\phi$  we check that  $(E, d)$  is also asymptotically conic.

Observe that although  $\phi$  is not differentiable over  $S^1 \times 0$  as a mapping with values in  $\mathbb{R}^2$ , it is yet a globally conic resolution of  $(E, d)$  in the sense of Proposition [2.8.](#page-12-1)

Observe also that the (image of the unit speed)  $g^o$ -geodesic from  $\mathbf{x}_0 = r_0 e^{\theta_0}$  to **0** is the logarithmic spiral parametrized in polar coordinates as  $r \to (\theta_0 + \ln r - \ln r_0, r)$ . Obviously, a similar phenomenon holds true for any  $g^o$ -geodesic leaving any compact of E.

#### <span id="page-20-1"></span>**REFERENCES**

- <span id="page-20-5"></span><span id="page-20-4"></span>[1] Lev Birbrair and Tadeusz Mostowski. Normal embeddings of semialgebraic sets. Michigan Math. J., 47(1):125–132, 2000.
- <span id="page-20-6"></span>[2] Dmitri Burago, Yuri Burago, and Sergei Ivanov. A course in metric geometry, volume 33 of Graduate Studies in Mathematics. American Mathematical Society, Providence, RI, 2001.
- [3] Jeff Cheeger. On the spectral geometry of spaces with cone-like singularities. Proc. Nat. Acad. Sci. U.S.A., 76(5):2103–2106, 1979.
- <span id="page-20-9"></span><span id="page-20-7"></span>[4] Jeff Cheeger. Spectral geometry of singular Riemannian spaces. J. Differential Geom., 18(4):575–657, 1983.
- [5] Jeff Cheeger and Michael Taylor. On the diffraction of waves by conical singularities. I. Comm. Pure Appl. Math., 35(3):275–331, 1982.
- <span id="page-20-10"></span>[6] Jeff Cheeger and Michael Taylor. On the diffraction of waves by conical singularities. II. Comm. Pure Appl. Math., 35(4):487–529, 1982.
- <span id="page-20-3"></span><span id="page-20-0"></span>[7] André Costa, Vincent Grandjean, and Maria Michalska. Global Lipschitz geometry of conic singular submanifolds with applications to algebraic sets. To appear in Doc. Mathemat.
- [8] Andr´e Costa, Vincent Grandjean, and Maria Michalska. One point compactification and Lipschitz normally embedded definable subsets.  $arXiv$ , 2023, <https://arxiv.org/abs/2305.07469>.
- <span id="page-20-2"></span>[9] Andr´e Costa, Vincent Grandjean, and Maria Michalska. Characterization of Lipschitz normally embedded complex curves. Bull. Sci. Math., 190:Paper No. 103369, 35, 2024.
- <span id="page-20-8"></span>[10] André Costa, Vincent Grandjean, and Maria Michalska. On globally hbc-singular spaces and subspaces. In preparation, 2024.
- <span id="page-21-9"></span><span id="page-21-8"></span>[11] Vincent Grandjean and Roger Oliveira. Stereographic compactification and affine bi-lipschitz homeomorphisms. Glasgow Mathematical Journal, page 1–15, 2024.
- <span id="page-21-5"></span>[12] Daniel Grieser. A natural differential operator on conic spaces. Discrete Contin. Dyn. Syst., pages 568–577, 2011.
- <span id="page-21-1"></span>[13] Daniel Grieser and Jørgen Olsen Lye. Geodesics orbiting a singularity. J. Geom., 115(1):Paper No. 1, 31, 2024.
- <span id="page-21-7"></span>[14] Mikhael Gromov. Structures métriques pour les variétés riemanniennes, volume 1 of Textes Mathématiques. CEDIC, Paris, 1981.
- <span id="page-21-11"></span>[15] Andrew Hassell, Terence Tao, and Jared Wunsch. Sharp Strichartz estimates on nontrapping asymptotically conic manifolds. Amer. J. Math., 128(4):963–1024, 2006.
- <span id="page-21-3"></span>[16] Dmitry Kerner, Helge Møller Pedersen, and Maria A. S. Ruas. Lipschitz normal embeddings in the space of matrices. Math. Z., 290(1-2):485–507, 2018.
- <span id="page-21-4"></span>[17] Richard Melrose. *Differential Analysis on Manifolds with Corners*. unfinished book, <http://math.mit.edu/~rbm>, 1996.
- <span id="page-21-6"></span>[18] Richard Melrose and Jared Wunsch. Propagation of singularities for the wave equation on conic manifolds. Invent. Math., 156(2):235–299, 2004.
- <span id="page-21-10"></span>[19] Richard Melrose and Maciej Zworski. Scattering metrics and geodesic flow at infinity. Invent. Math., 124(1- 3):389–436, 1996.
- <span id="page-21-2"></span>[20] S. B. Myers and N. E. Steenrod. The group of isometries of a Riemannian manifold. Ann. of Math. (2), 40(2):400–416, 1939.
- <span id="page-21-0"></span>[21] Richard S. Palais. On the differentiability of isometries. Proc. Amer. Math. Soc., 8:805–807, 1957.
- [22] Hassler Whitney. Analytic extensions of differentiable functions defined in closed sets. Trans. Amer. Math. Soc., 36(1):63–89, 1934.

A. COSTA, CENTRO DE CIÊNCIAS E TECNOLOGIAS, UNIVERSIDADE ESTADUAL DO CEARÁ, CAMPUS DO Itaperi, 60.714-903 Fortaleza - CE, Brasil

V. GRANDJEAN, DEPARTAMENTO DE MATEMÁTICA, DEPARTAMENTO DE MATEMÁTICA, UNIVERSIDADE FEDeral de Santa Catarina, 88.040-900 Florianopolis - SC, Brasil, Brasil ´

M. MICHALSKA, WYDZIAŁ MATEMATYKI I INFORMATYKI, UNIWERSYTET ŁÓDZKI, BANACHA 22, 90-238 LÓDŹ, POLAND

Email address: andrecosta.math@gmail.com, vincent.grandjean@ufsc.br, maria.michalska@wmii.uni.lodz.pl