

Topological properties of black holes in five-dimensional gauged supergravity

Yucheng He ^{1a} Changxiang Lei ^{2a} Deyou Chen ^{3a}

^a*School of Science, Xihua University, Chengdu 610039, China*

ABSTRACT: In this paper, we study the topological properties of five-dimensional rotating charged black holes in different ensembles. The topological numbers for the black holes are gotten in the grand canonical and canonical ensembles, which are 1. When the charge and cosmological constant disappear, their topological numbers are 0. When the pressure is lower than the critical pressure, the phase transition exists in the canonical ensemble. However, the phase transition also exists in the grand canonical ensemble when the pressure is higher than the critical pressure and two independent rotational parameters are $a = 1$ and $b = -1$. This may be due to the fact that the values of the rotational parameters change the supersymmetry of the black hole and lead to the phase transition.

¹E-mail: heyucheng365@hotmail.com

²E-mail: leichangxiangqqp@outlook.com

³E-mail: deyouchen@hotmail.com

Contents

1	Introduction	1
2	Five-dimensional black holes in gauged supergravity	2
3	Topological numbers for five-dimensional black holes	4
3.1	Review of topological approach	4
3.2	Topological numbers for five-dimensional black holes in canonical ensemble	5
3.3	Topological numbers for five-dimensional black holes in grand canonical ensemble	8
4	Conclusion	12

1 Introduction

Recently, Wei, Liu and Mann proposed an elegant method to study the black holes. They treated the black hole solutions as defects in the thermodynamic parameter space [1], and used Duan’s topological current ϕ -mapping theory [2, 3] to study the black holes’ properties. In this work, the local properties of the black holes are characterized by their winding numbers, and their positive/negative values reflect the local stabilities/unstabilities of the black holes. The globe properties are characterized by their topological numbers, and each black hole is endowed with a topological number, therefore, the black holes can be classified by their topological numbers. Inspired by this work, the topological numbers of different black holes were calculated in different ensembles, and the influences of the black holes’ parameters such as the angular momentum, charge and cosmological constant on the topological numbers have been extensively discussed [4–41]. An interesting phenomenon was found, namely, the topological numbers for some black holes were affected by the ensembles and the values of the black holes’ parameters [42–47]. In view of this, it is natural for people to perform topological classification on black holes in various spacetimes, and to test whether the relevant parameters of black holes affect their classification.

In this paper, we study the topological numbers for the charged rotating black holes in five-dimensional minimal gauged supergravity in the different ensembles, and test whether these numbers are affected by the ensembles and parameters’ values for these black holes. These black holes’ solution are parameterized by four nontrivial parameters, namely, the mass, charge, and two independent rotation parameters in the two orthogonal 2–planes, which was obtained in [48]. An interest phenomenon for this gauged supergravity is in the context of the AdS/CFT correspondence, and their bulk properties in anti-de Sitter spacetimes are related to these of strongly coupled conformal field theories on the four-dimensional boundary of AdS₅ [49, 50]. Another interest for studying these black holes is that the unfixed cosmological constant can reveal deeper physics and produce many meaningful results. Therefore, it is necessary to study the topological properties of these black holes.

The rest is organized as follows. In the next section, we give a brief review of the solution of the charged rotating black hole in five-dimensional gauged supergravity and discuss its thermodynamic properties. In Sec. 3, we first review the topological approach, and then calculate the topological numbers for the charged rotating black holes and Myers-Perry black holes in canonical and grand canonical ensembles, respectively. Sec. 4 is devoted to our conclusion.

2 Five-dimensional black holes in gauged supergravity

In the five-dimensional gauged supergravity, the general solution of charged rotating black holes in Boyer-Lindquist type coordinates is given by [48]

$$ds^2 = -\frac{\Delta_\theta[(1+g^2r^2)\rho^2 dt + 2q\nu]dt}{\Xi_a\Xi_b\rho^2} + \frac{2q\nu\omega}{\rho^2} + \frac{f}{\rho^4} \left(\frac{\Delta_\theta dt}{\Xi_a\Xi_b} - \omega \right)^2 + \frac{\rho^2 dr^2}{\Delta_r} + \frac{\rho^2 d\theta^2}{\Delta_\theta} + \frac{r^2 + a^2}{\Xi_a} \sin^2 \theta d\varphi^2 + \frac{r^2 + b^2}{\Xi_b} \cos^2 \theta d\psi^2, \quad (2.1)$$

with an electromagnetic potential

$$A = \frac{\sqrt{3}q}{\rho^2} \left(\frac{\Delta_\theta dt}{\Xi_a\Xi_b} - \omega \right), \quad (2.2)$$

where

$$\begin{aligned} \nu &= b \sin^2 \theta d\varphi + a \cos^2 \theta d\psi, \\ \omega &= a \sin^2 \theta \frac{d\varphi}{\Xi_a} + b \cos^2 \theta \frac{d\psi}{\Xi_b}, \\ f &= 2m\rho^2 - q^2 + 2abqg^2\rho^2, \\ \rho^2 &= r^2 + a^2 \cos^2 \theta + b^2 \sin^2 \theta, \\ \Xi_a &= 1 - a^2g^2, \quad \Xi_b = 1 - b^2g^2, \\ \Delta_\theta &= 1 - a^2g^2 \cos^2 \theta - b^2g^2 \sin^2 \theta, \\ \Delta_r &= \frac{(r^2 + a^2)(r^2 + b^2)(1 + g^2r^2) + q^2 + abq}{r^2} - 2m, \end{aligned} \quad (2.3)$$

a and b are two independent rotation parameters, and their positive/negative values represent the rotational directions. g is a physical quantity expressed as the cosmological constant $g^2 = -\frac{\Lambda}{4}$. m and q are parameters related to the ADM mass M and charge Q as follows

$$\begin{aligned} M &= \frac{m\pi(2\Xi_a + 2\Xi_b - \Xi_a\Xi_b) + 2\pi qabg^2(\Xi_a + \Xi_b)}{4\Xi_a^2\Xi_b^2}, \\ Q &= \frac{\sqrt{3}\pi q}{4\Xi_a\Xi_b}, \end{aligned} \quad (2.4)$$

respectively. The metric (2.1) describes a Myers-Perry black hole when $g^2 = q = 0$, and describes a static AdS black hole when $a = b = 0$. The electromagnetic potential at the event horizon is

$$\Phi = l^\mu A_\mu = \frac{\sqrt{3}qr_h^2}{(r_h^2 + a^2)(r_h^2 + b^2) + abq}. \quad (2.5)$$

where $l^\mu = \frac{\partial}{\partial t} + \Omega_a \frac{\partial}{\partial \varphi} + \Omega_b \frac{\partial}{\partial \psi}$ is the Killing vector, and r_h is the event horizon determined by $\Delta_r = 0$. The Hawking temperature and Bekenstein-Hawking entropy are

$$\begin{aligned}
T &= \frac{r_h^4 [1 + g^2 (a^2 + b^2 + 2r_h^2)] - (ab + q)^2}{2\pi r_h [(r_h^2 + a^2)(r_h^2 + b^2) + abq]}, \\
S &= \frac{\pi^2 [(r_h^2 + a^2)(r_h^2 + b^2) + abq]}{2r_h \Xi_a \Xi_b}.
\end{aligned} \tag{2.6}$$

The angular momenta and angular velocities at the horizon are

$$\begin{aligned}
J_a &= \frac{\pi [2ma + qb (1 + g^2 a^2)]}{4\Xi_a^2 \Xi_b}, \\
J_b &= \frac{\pi [2mb + qa (1 + g^2 b^2)]}{4\Xi_a \Xi_b^2}, \\
\Omega_a &= \frac{a (r_h^2 + b^2) (1 + g^2 r_h^2) + bq}{(r_h^2 + a^2)(r_h^2 + b^2) + abq}, \\
\Omega_b &= \frac{b (r_h^2 + a^2) (1 + g^2 r_h^2) + aq}{(r_h^2 + a^2)(r_h^2 + b^2) + abq}.
\end{aligned} \tag{2.7}$$

The above thermodynamic quantities obey the first law of thermodynamics,

$$dM = TdS + \Omega_a dJ_a + \Omega_b dJ_b + \Phi dQ, \tag{2.8}$$

where the cosmological constant is fixed. Recently, the black holes' thermodynamics in the extended phase spaces have been researched, and some interesting phenomena, such as Van der Waals-like phase transition, microstructures of black holes, etc, were found. In these phase spaces, the cosmological constant is regarded as a variable related to pressure $P = -\frac{\Lambda}{8\pi} = \frac{(n-1)(n-2)}{16\pi l^2}$, where l is the bulk curvature radius and n is a dimension of spacetimes, its conjugate quantity is a thermodynamic volume V . Thus the mass M is interpreted as thermodynamic enthalpy instead of internal energy. Considering the varied constant, the first law of this black hole is

$$dM = TdS + \Omega_a dJ_a + \Omega_b dJ_b + \Phi dQ + VdP. \tag{2.9}$$

The phase transition occurs at $\frac{\partial_{r_h} T}{\partial_{r_h} S} = 0$. Using this condition, we get the critical pressure

$$P_c = -\frac{ab(ab+q)^3 + 3(a^2+b^2)(ab+q)^2 r_h^2 + (ab+q)(8ab+5q)r_h^4 + (a^2+b^2)r_h^6 - r_h^8}{3ab(a^2+b^2)(ab+q)r_h^4 + 2\pi(12a^2b^2+a^4+10abq+b^4)r_h^6 + 5(a^2+b^2)r_h^8 + 2r_h^{10}}. \tag{2.10}$$

Then the critical horizon radius for the phase transition obeys

$$\begin{aligned}
&3ab(a^2+b^2)^2(ab+q)^3 + 3(a^2+b^2)(ab+q)^2(a^4+9a^2b^2+b^4+7abq)r_h^2 \\
&+ 2(ab+q)^2(14a^4+43a^2b^2+14b^4+15abq)r_h^4 + (a^2+b^2)(ab+q)(74ab+59q)r_h^6 \\
&+ 3(a^4+22a^2b^2+b^4+30abq+10q^2)r_h^8 + 5(a^2+b^2)r_h^{10} - 2r_h^{12} = 0.
\end{aligned} \tag{2.11}$$

3 Topological numbers for five-dimensional black holes

3.1 Review of topological approach

According to Ref. [1], the generalized free energy is defined by

$$\mathcal{F} = E - \frac{S}{\tau}, \quad (3.1)$$

where E and S are the energy and entropy of a black hole system, respectively. τ is a variable and can be seen as the inverse temperature of the cavity enclosing the black hole. This free energy is off-shell except at $\tau = \frac{1}{T}$. To calculate the topological numbers, a vector is constructed as

$$\phi = \left(\frac{\partial \mathcal{F}}{\partial r_h}, -\cot \Theta \csc \Theta \right). \quad (3.2)$$

In this vector, $0 < r_h < +\infty$ and $0 \leq \Theta \leq \pi$. Zero points of the vector obtained at $\tau = 1/T$ and $\Theta = \pi/2$ correspond to the on-shell black hole solution. Other points are not the solutions of Einstein field equations, thus they are the off-shell states. ϕ^Θ diverges at $\Theta = 0$ and $\Theta = \pi$, which leads to that the direction of the vector is outward.

Using Duan's ϕ -mapping topological current theory [2, 3], we define a topological current

$$j^\mu = \frac{1}{2\pi} \varepsilon^{\mu\nu\rho} \varepsilon_{ab} \partial_\nu n^a \partial_\rho n^b, \quad (3.3)$$

where $\mu, \nu, \rho = 0, 1, 2$, $a, b = 1, 2$, $\partial_\nu = \frac{\partial}{\partial x^\nu}$ and $x^\nu = (\tau, r_h, \Theta)$. τ is seen as a time parameter of the topological defect. n^a is a unit vector defined by $\left(\frac{n^r}{\|n\|}, \frac{n^\Theta}{\|n\|} \right)$. It is easily to prove the current is conserved. Using the Jacobi tensor $\varepsilon^{ab} J^\mu \left(\frac{\phi}{x} \right) = \varepsilon^{\mu\nu\rho} \partial_\nu \phi^a \partial_\rho \phi^b$ and two-dimensional Laplacian Green function $\Delta_{\phi^a} \ln \|\phi\| = 2\pi \delta^2(\phi)$, the current is rewritten as

$$j^\mu = \delta^2(\phi) j^\mu \left(\frac{\phi}{x} \right), \quad (3.4)$$

which is nonzero only when $\phi^a(x^i) = 0$. Then a topological number in a parameter region Σ is calculated as

$$W = \int_{\Sigma} j^0 d^2 x = \sum_{i=1}^N \beta_i \eta_i = \sum_{i=1}^N w_i, \quad (3.5)$$

where $j^0 = \sum_{i=1}^N \beta_i \eta_i \delta^2(\vec{x} - \vec{z}_i)$ is the density of the current. β_i is Hopf index which counts the number of the loops that ϕ^a makes in the vector ϕ space when x^μ goes around the zero point z_i . Clearly, this index is always positive. $\eta_i = \text{sign} J^0(\phi/x)_{z_i} = \pm 1$ is the Brouwer degree. w_i is the winding number for the i -th zero point of the vector in the region and its values is independent on the shape of the region. In the researches, people found the this value is determined by the (un-)stable black holes. In the following, we will study the topological numbers for the five-dimensional black holes.

3.2 Topological numbers for five-dimensional black holes in canonical ensemble

Using Eq. (3.1), we get the generalized free energy

$$\mathcal{F} = \frac{\pi (2\Xi_a + 2\Xi_b - \Xi_a\Xi_b) [(r_h^2 + a^2) (r_h^2 + b^2) (1 + g^2 r_h^2) + 2abq + q^2]}{8\Xi_a^2\Xi_b^2 r_h^2} + \frac{\pi qabg^2 (\Xi_a + \Xi_b)}{2\Xi_a^2\Xi_b^2} - \frac{\pi^2 [(r_h^2 + a^2) (r_h^2 + b^2) + abq]}{2r_h\Xi_a\Xi_b\tau}. \quad (3.6)$$

Thus the components of the vector ϕ are gotten as follows

$$\phi^{r_h} = -\frac{\pi [\Xi_a\Xi_b - 3(\Xi_a + \Xi_b)] [2g^2 r_h^6 + (g^2 (a^2 + b^2) + 1) r_h^4 - (ab + q)^2]}{4r_h^3\Xi_a^2\Xi_b^2} + \frac{\pi^2 [a^2 b^2 + abq - (a^2 + b^2) r_h^2 - 3r_h^4]}{2r_h^2\Xi_a\Xi_b\tau}, \quad (3.7)$$

$$\phi^\Theta = -\cot \Theta \csc \Theta. \quad (3.8)$$

Zero points are determined by $\phi^{r_h} = 0$ and $\Theta = \pi/2$. We solve it and get

$$\tau = \frac{2\pi r_h \Xi_a \Xi_b [a^2 b^2 + abq - (a^2 + b^2) r_h^2 - 3r_h^4]}{[\Xi_a \Xi_b - 3(\Xi_a + \Xi_b)] [2g^2 r_h^6 + (g^2 (a^2 + b^2) + 1) r_h^4 - (ab + q)^2]}. \quad (3.9)$$




Figure 1: Topological properties of the five-dimensional charged rotating black hole, where $a/r_0 = 0$, $b/r_0 = 1$, $q/r_0^2 = 1$, $Pr_0^2 = 0.02$ and $P_c r_0^2 = 0.00631$. Zero points of the vector ϕ^{r_h} in the plane $r_h - \tau$ are plotted in the left picture. The unit vector field n on a portion of the plane $\Theta - r_h$ at $\tau/r_0 = 5$ is plotted in the right picture. The zero point is at $(r_h/r_0, \Theta) = (3.37, \pi/2)$.

The above equation shows the relation between the inverse temperature and horizon radius for the zero points. The critical pressure for different values of the parameters is calculated numerically. Topological properties of the black holes are shown in Figure (1) - (6).




Figure 2: Topological properties of the five-dimensional charged rotating black hole, where $a/r_0 = 0$, $b/r_0 = 1$, $q/r_0^2 = 1$, $P r_0^2 = 0.001$ and $P_c r_0^2 = 0.00631$. Zero points of the vector ϕ^{r_h} in the plane $r_h - \tau$ are plotted in the left picture. The unit vector field n on a portion of the plane $\Theta - r_h$ at $\tau/r_0 = 25$ is plotted in the right picture. The zero points are at $(r_h/r_0, \Theta) = (1.10, \pi/2)$, $(5.44, \pi/2)$ and $(14.37, \pi/2)$, respectively.




Figure 3: Topological properties of the five-dimensional charged rotating black hole, where $a/r_0 = 1$, $b/r_0 = 1$, $q/r_0^2 = 1$, $P r_0^2 = 0.01$ and $P_c r_0^2 = 0.00286$. Zero points of the vector ϕ^{r_h} in the plane $r_h - \tau$ are plotted in the left picture. The unit vector field n on a portion of the plane $\Theta - r_h$ at $\tau/r_0 = 5$ is plotted in the right picture. The zero point is at $(r_h/r_0, \Theta) = (8.17, \pi/2)$.




Figure 4: Topological properties of the five-dimensional charged rotating black hole, where $a/r_0 = 1$, $b/r_0 = 1$, $q/r_0^2 = 1$, $P r_0^2 = 0.001$ and $P_c r_0^2 = 0.00286$. Zero points of the vector ϕ^{r_h} in the plane $r_h - \tau$ are plotted in the left picture. The unit vector field n on a portion of the plane $\Theta - r_h$ at $\tau/r_0 = 25$ is plotted in the right picture. The zero points are at $(r_h/r_0, \Theta) = (1.62, \pi/2)$, $(5.38, \pi/2)$ and $(14.23, \pi/2)$, respectively.




Figure 5: Topological properties of the five-dimensional charged rotating black hole, where $a/r_0 = 1$, $b/r_0 = -1$, $q/r_0^2 = 1$, $Pr_0^2 = 0.05$ and $P_c r_0^2 = 0.00468$. Zero points of the vector ϕ^{r_h} in the plane $r_h - \tau$ are plotted in the left picture. The unit vector field n on a portion of the plane $\Theta - r_h$ at $\tau/r_0 = 2$ is plotted in the right picture. The zero point is at $(r_h/r_0, \Theta) = (2.38, \pi/2)$.




Figure 6: Topological properties of the five-dimensional charged rotating black hole, where $a/r_0 = 1$, $b/r_0 = -1$, $q/r_0^2 = 1$, $Pr_0^2 = 0.001$ and $P_c r_0^2 = 0.00468$. Zero points of the vector ϕ^{r_h} in the plane $r_h - \tau$ are plotted in the left picture. The unit vector field n on a portion of the plane $\Theta - r_h$ at $\tau/r_0 = 25$ is plotted in the right picture. The zero points are at $(r_h/r_0, \Theta) = (0.17, \pi/2)$, $(5.42, \pi/2)$ and $(14.24, \pi/2)$, respectively.

In Figure (1) and (2), we first order $a = 0$ and study the case where only one rotational parameter exists. When the pressure is larger than the critical pressure we plot in Figure (1). In this figure, the horizon radius decreases monotonically with the increase of the inverse temperature, which shows that the phase transition doesn't exist. Thus the black hole is stable for any temperature. We use the method developed in [1] and find that the winding number is 1. There is only one zero point in the right picture of the figure. The winding number can also be calculated by using the method in [51, 52], and it is independent of the loop that surround the zero point. Thus its topological number is 1. When the pressure is less than the critical pressure, a generation point and an annihilation point exist in Figure (2), and they divide the black hole into three branches: a large black hole (LBH), an intermediate black hole (IBH), and a small black hole (SBH). These three branches coexist in the range $\tau_a < \tau < \tau_b$, which implies that a LBH/SBH phase transition (PT) exists in this range. For the LBH and SBH branches, they are locally stable, and both have a winding number of 1. For the IBH branch, it is locally unstable, and its winding number is -1 . Thus the topological number for this black hole is also 1.

When two rotational parameters coexist in the black hole, we draw Figure (3) - (6). The positive

and negative values of a and b represent the direction of rotation for the black hole on φ and ψ , respectively. When the pressure is larger than the critical pressure, the radius decreases monotonically with τ 's value in Figure (3) and (5). When the pressure is less than the critical pressure, there is also a LBH/SBH phase transition in Figure (4) and (6), which is similar to the case in Figure (2). Comparing to Figure (2), we find some differences in Figure (4) and (6). Due to the presence of the parameter a , τ 's values corresponding to the generation point and annihilation point change, and its range for the PT decrease. All of them show that the topological number for this black hole is 1, which is fully confirms to the result obtained in [1] that the topological number is a universal number independent of the black hole parameters.

The metric (2.1) describes a Myers-Perry black hole when $g^2 = q = 0$. To study the topological number for this black hole, we draw Figure (7) and (8) when $a = 1, b = 0$ and $a = 1, b = 1$, respectively. Due to the vanish of the cosmological constant, we don't need consider the influence of the pressure in the figures. In Figure (7), a generation point divides the black hole into two regions. When τ 's value exceeds a certain value, there are two different black holes, where the black holes with a small radius being stable and with a large radius being unstable. Their winding numbers are 1 and -1, respectively. Thus the topological number is 0. When τ 's is less than a certain value, there is no black hole existed. Similarly, the topological number in (8) is also 0.




Figure 7: Topological properties of the Myers-Perry black hole, where $a/r_0 = 1$ and $b/r_0 = 0$. Zero points of the vector ϕ^{r_h} in the plane $r_h - \tau$ are plotted in the left picture. The unit vector field n on a portion of the plane $\Theta - r_h$ at $\tau/r_0 = 25$ is plotted in the right picture. The zero points are at $(r_h/r_0, \Theta) = (0.21, \pi/2)$ and $(1.49, \pi/2)$, respectively.

3.3 Topological numbers for five-dimensional black holes in grand canonical ensemble

In [41, 42, 45], the authors found that the topological class of the four-dimensional dyonic AdS black hole is ensemble dependent. We recalculate the topological numbers in the grand canonical ensemble to verify whether the numbers are consistent with these gotten in the canonical ensemble in this section. In a canonical ensemble, a black hole in a cavity has an energy exchange with the outside, and its temperature, volume and particle number remain be unchanged. In a grand canonical ensemble, the black hole in the cavity exchanges energy and charge with the outside, and its temperature, volume and chemical potential remain be unchanged.

In the grand canonical ensemble, the generalized free energy is




Figure 8: Topological properties of the Myers-Perry black hole, where $a/r_0 = 1, b/r_0 = 1$. Zero points of the vector ϕ^{r_h} in the plane $r_h - \tau$ are plotted in the left picture. The unit vector field n on a portion of the plane $\Theta - r_h$ at $\tau/r_0 = 25$ is plotted in the right picture. The zero points are at $(r_h/r_0, \Theta) = (1.17, \pi/2)$ and $(3.22, \pi/2)$, respectively.

$$\begin{aligned}
\mathcal{F} &= M - Q\Phi - \frac{S}{\tau} \\
&= \frac{\pi (2\Xi_a + 2\Xi_b - \Xi_a\Xi_b) [(r_h^2 + a^2) (r_h^2 + b^2) (1 + g^2 r_h^2) + 2abq + q^2]}{8\Xi_a^2\Xi_b^2 r_h^2} \\
&\quad + \frac{\pi qabg^2 (\Xi_a + \Xi_b)}{2\Xi_a^2\Xi_b^2} - \frac{3\pi q^2 r_h^2}{4\Xi_a\Xi_b [(r_h^2 + a^2) (r_h^2 + b^2) + abq]} \\
&\quad - \frac{\pi^2 [(r_h^2 + a^2) (r_h^2 + b^2) + abq]}{2r_h\Xi_a\Xi_b\tau}. \tag{3.10}
\end{aligned}$$

Using the definition $\phi^{r_h} = \frac{\partial \mathcal{F}}{\partial r_h}$, we get the component of the vector,

$$\begin{aligned}
\phi^{r_h} &= - \frac{\pi [\Xi_a\Xi_b - 3(\Xi_a + \Xi_b)] [2g^2 r_h^6 + r_h^4 (g^2 (a^2 + b^2) + 1) - (ab + q)^2]}{4r_h^3\Xi_a^2\Xi_b^2} \\
&\quad + \frac{3\pi q^2 r_h [r_h^4 - ab(ab + q)]}{2\Xi_a\Xi_b [r_h^4 + r_h^2 (a^2 + b^2) + ab(ab + q)]^2} \\
&\quad + \frac{\pi^2 [a^2 b^2 + abq - r_h^2 (a^2 + b^2) - 3r_h^4]}{2r_h^2\Xi_a\Xi_b\tau}. \tag{3.11}
\end{aligned}$$

We solve $\phi^{r_h} = 0$ and obtain the relation

$$\tau = \frac{2\pi r_h \Xi_a \Xi_b [a^2 b^2 + abq - r_h^2 (a^2 + b^2) - 3r_h^4]}{\frac{6q^2 r_h^4 \Xi_a \Xi_b [ab(ab+q) - r_h^4]}{[r_h^2 (a^2 + b^2) + ab(ab+q) + r_h^4]^2} + [\Xi_a \Xi_b - 3(\Xi_a + \Xi_b)] [2g^2 r_h^6 + r_h^4 (g^2 (a^2 + b^2) + 1) - (ab + q)^2]} \tag{3.12}$$

To study the topological properties in the grand canonical ensemble, we draw Figure (9) - (14). There is only one rotational parameter in Figure (9) and (10). When the pressure is larger than the critical pressure, the horizon radius monotonically decreases with the increase of the temperature, the black hole is local stable and there is no PT. When the pressure is less than the critical pressure, there is a generation point and an annihilation point which divide the black hole into LBH, IBH and

SBH branches. The LBH/SBH PT exists in the range $\tau_a < \tau < \tau_b$. These two figures show that the topological number is 1. When $a = b \neq 0$, the similar result is found in Figure (11) and (12).

We draw Figure (13) and (14) when $b/r_0 = -1$. It is clearly that the topological number is also 1. In the figures, no matter how the pressure is taken, the LBH/SBH PT always exists for this black hole. This strange phenomenon exists in the five-dimensional charged rotating black hole in the grand canonical ensemble. The reason for this phenomenon may be that the values of a and b change the supersymmetry. In [53, 54], the authors found that the fraction of supersymmetry is in general 1/4 for charged rotating black hole solutions in five dimensional gauged supergravity. However, when $a = -b$, the black hole can be viewed as a solution of the pure $N = 2$ gauged supergravity theory. In which case, the preserved supersymmetry becomes 1/2. When the pressure is higher than the critical pressure, although there is still a LBH/SBH PT, we find that the distance between the generation points and annihilation points in Figure (13) is smaller than that in Figure (14).




Figure 9: Topological properties of the five-dimensional charged rotating black hole in the grand canonical ensemble, where $a/r_0 = 0$, $b/r_0 = 1$, $q/r_0^2 = 1$, $Pr_0^2 = 0.03$ and $P_c r_0^2 = 0.00631$. Zero points of the vector ϕ^{r_h} in the plane $r_h - \tau$ are plotted in the left picture. The unit vector field n on a portion of the plane $\Theta - r_h$ at $\tau/r_0 = 5$ is plotted in the right picture. The zero point is at $(r_h/r_0, \Theta) = (1.27, \pi/2)$.




Figure 10: Topological properties of the five-dimensional charged rotating black hole in the grand canonical ensemble, where $a/r_0 = 0$, $b/r_0 = 1$, $q/r_0^2 = 1$, $Pr_0^2 = 0.001$ and $P_c r_0^2 = 0.00631$. Zero points of the vector ϕ^{r_h} in the plane $r_h - \tau$ are plotted in the left picture. The unit vector field n on a portion of the plane $\Theta - r_h$ at $\tau/r_0 = 20$ is plotted in the right picture. The zero points are at $(r_h/r_0, \Theta) = (0.96, \pi/2)$, $(3.68, \pi/2)$ and $(21.12, \pi/2)$, respectively.




Figure 11: Topological properties of the five-dimensional charged rotating black hole in the grand canonical ensemble, where $a/r_0 = 1$, $b/r_0 = 1$, $q/r_0^2 = 1$, $Pr_0^2 = 0.01$ and $P_c r_0^2 = 0.00361$. Zero points of the vector ϕ^{r_h} in the plane $r_h - \tau$ are plotted in the left picture. The unit vector field n on a portion of the plane $\Theta - r_h$ at $\tau/r_0 = 5$ is plotted in the right picture. The zero point is at $(r_h/r_0, \Theta) = (1.26, \pi/2)$.




Figure 12: Topological properties of the five-dimensional charged rotating black hole in the grand canonical ensemble, where $a/r_0 = 1$, $b/r_0 = 1$, $q/r_0^2 = 1$, $Pr_0^2 = 0.001$ and $P_c r_0^2 = 0.00361$. Zero points of the vector ϕ^{r_h} in the plane $r_h - \tau$ are plotted in the left picture. The unit vector field n on a portion of the plane $\Theta - r_h$ at $\tau/r_0 = 25$ is plotted in the right picture. The zero points are at $(r_h/r_0, \Theta) = (1.12, \pi/2)$, $(5.41, \pi/2)$ and $(14.22, \pi/2)$, respectively.




Figure 13: Topological properties of the five-dimensional charged rotating black hole in the grand canonical ensemble, where $a/r_0 = 1$, $b/r_0 = -1$, $q/r_0^2 = 1$, $Pr_0^2 = 0.01$ and $P_c r_0^2 = 0.00468$. Zero points of the vector ϕ^{r_h} in the plane $r_h - \tau$ are plotted in the left picture. The unit vector field n on a portion of the plane $\Theta - r_h$ at $\tau/r_0 = 8$ is plotted in the right picture. The zero points are at $(r_h/r_0, \Theta) = (0.38, \pi/2)$, $(1.97, \pi/2)$ and $(3.43, \pi/2)$, respectively.




Figure 14: Topological properties of the five-dimensional charged rotating black hole in the grand canonical ensemble, where $a/r_0 = 1$, $b/r_0 = -1$, $q/r_0^2 = 1$, $Pr_0^2 = 0.001$ and $P_c r_0^2 = 0.00468$. Zero points of the vector ϕ^{r_h} in the plane $r_h - \tau$ are plotted in the left picture. The unit vector field n on a portion of the plane $\Theta - r_h$ at $\tau/r_0 = 25$ is plotted in the right picture. The zero points are at $(r_h/r_0, \Theta) = (0.10, \pi/2)$, $(1.98, \pi/2)$ and $(3.42, \pi/2)$, respectively.

4 Conclusion

In this work, we studied the topological properties of the five-dimensional gauged supergravity black holes in the canonical and grand canonical ensembles. The result shows that their topological numbers are 1 in these ensembles. When the charge and cosmological constant disappear, their topological numbers are 0. This indicates that the cosmological constant and black holes' charges play important roles in the topological classification.

In the work, the cosmological constant was regarded as a variable related to the pressure, and the critical pressure was discussed. We calculated the topological numbers for the black holes when the pressure is greater and less than the critical pressure, respectively. The topological number for the charged rotating black hole is 1 for any parameters' values. In the canonical ensemble, when the pressure is greater than the critical pressure, there is no generation point and annihilation point. However, when the pressure is less than the critical pressure, there is a generation point and an annihilation point. For the Myers-Perry black hole, since there is no cosmological constant, its topological number is 0, and there is only one generation point. From the above analysis, we conclude that the topological number is always 1 when the black hole is charged and is 0 when the black hole is uncharged. The existence of the PT is related to the critical pressure. However, the LBH/SBH PT exists in the grand canonical ensemble when the pressure is higher than the critical pressure. The reason may be that the values of the rotational parameters change the supersymmetry of the black hole.

References

- [1] S.W. Wei, Y.X. Liu and R.B. Mann, *Black hole solutions as topological thermodynamic defects*, *Phys. Rev. Lett.* **129** (2022) 191101.
- [2] Y.S. Duan and M.L. Ge, *$SU(2)$ gauge theory and electrodynamics of N moving magnetic monopoles*, *Sci. Sin.* **9** (1979) 1072.
- [3] Y.S. Duan, *The structure of the topological currenty*, *SLAC-PUB-3301*, (1984).
- [4] P.K. Yerra, C. Bhamidipati and S. Mukherji, *Topology of critical points and Hawking-Page transition*, *Phys. Rev. D* **106** (2022) 064059.

- [5] M.S. Ali, H. El Moumni, J. Khalloufi and K. Masmar, *Topology of Born-Infeld-AdS black hole phase transition*, *Annals Phys.* **465** (2024) 169679.
- [6] P.K. Yerra and C. Bhamidipati, *Topology of black hole thermodynamics in Gauss-Bonnet gravity*, *Phys. Rev. D* **105** (2022) 104053.
- [7] S.W. Wei and Y.X. Liu, *Topology of equatorial timelike circular orbits around stationary black holes*, *Phys. Rev. D* **107** (2023) 064006.
- [8] P.K. Yerra and C. Bhamidipati, *Topology of Born-Infeld AdS black holes in 4D novel Einstein-Gauss-Bonnet gravity*, *Phys. Lett. B* **835** (2022) 137591.
- [9] N.C. Bai, L. Li and J. Tao, *Topology of black hole thermodynamics in Lovelock gravity*. *Phys. Rev. D* **107** (2023) 064015.
- [10] Z.Y. Fan, *Topological interpretation for phase transitions of black holes*, *Phys. Rev. D* **107** (2023) 044026.
- [11] C. Fang, J. Jiang and M. Zhang, *Revisiting thermodynamic topologies of black holes*, *JHEP* **2301** (2023) 102.
- [12] D. Wu, *Topological classes of rotating black holes*, *Phys. Rev. D* **107** (2023) 024024.
- [13] D. Wu and S.Q. Wu, *Topological classes of thermodynamics of rotating AdS black holes*, *Phys. Rev. D* **107** (2023) 084002.
- [14] D. Wu, *Classifying topology of consistent thermodynamics of the four-dimensional neutral Lorentzian NUT-charged spacetimes*, *Eur. Phys. J. C* **83** (2023) 365.
- [15] X. Ye and S.W. Wei, *Distinct topological configurations of equatorial timelike circular orbit for spherically symmetric (hairy) black holes*, *JCAP* **07** (2023) 049.
- [16] Y. Du and X.D. Zhang, *Topological classes of BTZ black holes*, [arXiv:2302.11189 [gr-qc]].
- [17] M. Zhang and J. Jiang, *Bulk-boundary thermodynamic equivalence: a topology viewpoint*, *JHEP* **2306** (2023) 115.
- [18] M.R. Alipour, M.A.S. Afshar, S. Noori Gashti and J. Sadeghi, *Topological classification and black hole thermodynamics*, *Phys. Dark Univ.* **42** (2023) 101361.
- [19] J. Sadeghi, S. Noori Gashti, M.R. Alipour and M.A.S. Afshar, *Bardeen black hole thermodynamics from topological perspective*, *Annals Phys.* **455** (2023) 169391.
- [20] D. Wu, *Topological classes of thermodynamics of the four-dimensional static accelerating black holes*, *Phys. Rev. D* **108** (2023) 084041.
- [21] M.Y. Zhang, H. Chen, H. Hassanabadi, Z.W. Long and H. Yang, *Topology of nonlinearly charged black hole chemistry via massive gravity*, *Eur. Phys. J. C* **83** (2023) 773.
- [22] R. Li, C. Liu, K. Zhang and J. Wang, *Topology of the landscape and dominant kinetic path for the thermodynamic phase transition of the charged Gauss-Bonnet-AdS black holes*, *Phys. Rev. D* **108** (2023) 044003.
- [23] S.W. Wei, Y.P. Zhang, Y.X. Liu and R.B. Mann, *Static spheres around spherically symmetric black hole spacetime*, *Phys. Rev. Res.* **5** (2023) 043050.
- [24] R. Li and J. Wang, *Generalized free energy landscapes of charged Gauss-Bonnet-AdS black holes in diverse dimensions*, *Phys. Rev. D* **108** (2023) 044057.
- [25] N.J. Gogoi and P. Phukon, *Topology of thermodynamics in R-charged black holes*, *Phys. Rev. D* **107** (2023) 106009.

- [26] D. Wu, *Consistent thermodynamics and topological classes for the four-dimensional Lorentzian charged Taub-NUT spacetimes*, *Eur. Phys. J. C* **83** (2023) 589.
- [27] T.N. Hung and C.H. Nam, *Topology in thermodynamics of regular black strings with Kaluza-Klein reduction*, *Eur. Phys. J. C* **83** (2023) 582.
- [28] M.S. Ali, H. El Moumni, J. Khalloufi and K. Masmar, *Topology of Born-Infeld-AdS black hole phase transitions: Bulk and CFT sides*, *Annals Phys.* **465** (2024) 169679.
- [29] J. Sadeghi, M.A.S. Afshar, S. Noori Gashti and M.R. Alipour, *Thermodynamic topology and photon spheres in the hyperscaling violating black holes*, *Astropart. Phys.* **156** (2024) 102920.
- [30] F. Barzi, H. El Moumni and K. Masmar, *Rényi topology of charged-flat black hole: Hawking-Page and Van-der-Waals phase transitions*, *JHEAp* **42** (2024) 63.
- [31] Z.Q. Chen and S.W. Wei, *Thermodynamics, Ruppeiner geometry, and topology of Born-Infeld black hole in asymptotic flat spacetime*, *Nucl. Phys. B* **996** (2023) 116369.
- [32] M. Rizwan and K. Jusufi, *Topological classes of thermodynamics of black holes in perfect fluid dark matter background*, *Eur. Phys. J. C* **83** (2023) 944
- [33] M.U. Shahzad, A. Mehmood, S. Sharif and A. Övgün, *Criticality and topological classes of neutral Gauss-Bonnet AdS black holes in 5D*, *Annals Phys.* **458** (2023) 169486.
- [34] K. Bhattacharya, K. Bamba and D. Singleton, *Topological interpretation of extremal and Davies-type phase transitions of black holes*, *Phys. Lett. B* **854** (2024) 138722.
- [35] M.U. Shahzad, A. Mehmood, A. Malik and A. Övgün, *Topological behavior of 3D regular black hole with zero point length*, *Phys. Dark Univ.* **44** (2024) 101437.
- [36] Y. Du and X. Zhang, *Topological classes of black holes in de-Sitter spacetime*, *Eur. Phys. J. C* **83** (2023) 927.
- [37] C. Liu, R. Li, K. Zhang and J. Wang, *Generalized free energy and dynamical state transition of the dyonic AdS black hole in the grand canonical ensemble*, *JHEP* **11** (2023) 068.
- [38] C. Fairoos and T. Sharqui, *Topological nature of black hole solutions in dRGT massive gravity*, *Int. J. Mod. Phys. A* **38** (2023) 2350133.
- [39] V.P. Cunha, C.A.R. Herdeiro and J.P.A. Novo, *Light rings on stationary axisymmetric spacetimes: Blind to the topology and able to coexist*, *Phys. Rev. D* **109** (2024) 064050.
- [40] B. Hazarika and P. Phukon, *Thermodynamic topology of black holes in $f(R)$ gravity*, *PTEP* **2024** (2024) 043E01.
- [41] P.K. Yerra, C. Bhamidipati and S. Mukherji, *Topology of critical points in boundary matrix duals*, *JHEP* **03** (2024) 138.
- [42] N.J. Gogoi and P. Phukon, *Thermodynamic topology of 4D Dyonically AdS black holes in different ensembles*, *Phys. Rev. D* **108** (2023) 066016.
- [43] C. Liu and J. Wang, *Topological natures of the Gauss-Bonnet black hole in AdS space*, *Phys. Rev. D* **107** (2023) 064023.
- [44] B. Hazarika and P. Phukon, *Thermodynamic topology of $D = 4, 5$ Horava Lifshitz black hole in two ensembles*, [arXiv:2312.06324 [hep-th]].
- [45] N.J. Gogoi and P. Phukon, *Thermodynamic topology of 4D Euler-Heisenberg-AdS black hole in different ensembles*, *Phys. Dark Univ.* **44** (2024) 101456.
- [46] D.Y. Chen, Y.C. He and J. Tao, *Topological classes of higher-dimensional black holes in massive gravity*, *Eur. Phys. J. C* **83** (2023) 872.

- [47] D.Y. Chen, Y.C. He, J. Tao and W. Yang, *Topology of Hořava-Lifshitz black holes in different ensembles*, *Eur. Phys. J. C* **84** (2024) 96.
- [48] Z.W. Chong, M. Cvetič, H. Lu and C.N. Pope, *General non-extremal rotating black holes in minimal five-dimensional gauged supergravity*, *Phys. Rev. Lett.* **95** (2005) 161301.
- [49] J.M. Maldacena, *The Large N limit of superconformal field theories and supergravity*, *Adv. Theor. Math. Phys.* **2** (1998) 231.
- [50] D.Y. Chen, X. Jiang and H.T. Yang, *Holographic $T\bar{T}$ deformed entanglement entropy in $dS_3 = CFT_2$* , *Phys. Rev. D* **109** (2024) 026011.
- [51] S.W. Wei, *Topological charge and black hole photon spheres*, *Phys. Rev. D* **102** (2020) 064039.
- [52] S.W. Wei and Y.X. Liu, *Topology of black hole thermodynamics*, *Phys. Rev. D* **105** (2022) 104003.
- [53] D. Klemm and W.A. Sabra, *Charged rotating black holes in 5D Einstein-Maxwell (A)dS gravity*, *Phys. Lett. B* **503** (2001) 147.
- [54] J.B. Gutowski and H.S. Reall, *Supersymmetric AdS(5) black holes*, *JHEP* **02** (2004) 006.