FROBENIUS-PERRON DIMENSIONS OF CONJUGACY CLASSES AND AN ITO-MICHLER-TYPE RESULT IN MODULAR FUSION CATEGORIES

SEBASTIAN BURCIU

ABSTRACT. The influence of certain arithmetic conditions on the sizes of conjugacy classes of a finite group on the group structure has been extensively studied in recent years. In this paper, we explore analogous properties for fusion categories. In particular, we establish an Ito-Michler-type result for modular fusion categories.

1. INTRODUCTION

Fusion categories have emerged as a powerful tool in both mathematics and theoretical physics, offering a categorical framework for understanding symmetry and structure in various contexts, such as quantum field theory, topology, and representation theory. These categories generalize many classical notions from group theory, especially in settings where group symmetries alone are insufficient to describe complex phenomena.

Over the past few decades, considerable attention has been given to understanding the structure of a finite group G based on information about the arithmetic properties of its irreducible character degrees or, dually, of the sizes of its conjugacy classes. One of the central questions that has been extensively studied is how the structure of a group G can be determined when certain arithmetic properties of its irreducible character set, $\text{Irr}(G)$, are known. Significant contributions to this area have been made recently. For a summary of the known results in this direction one may consult the expository papers of [\[Man87\]](#page-20-0) and $|Lew08|$.

Building on the work of Zhu [\[Zhu94\]](#page-20-2), Cohen and Westreich [\[CW00\]](#page-20-3) introduced the concept of conjugacy classes for semisimple Hopf algebras. This new framework allowed them to extend several results from

Date: November 6, 2024.

²⁰²⁰ *Mathematics Subject Classification.* 16T30; 18M20.

The author is supported by a Research Grant GAR 2023 (cod 114), supported from the Donors' Recurrent Fund of the Romanian Academy, managed by the "PATRIMONIU" Foundation.

the theory of finite group representations to the context of semisimple Hopf algebra representations. More recently, Shimizu [\[Shi17\]](#page-20-4) expanded on this idea by defining conjugacy classes for fusion categories, generalizing the notion introduced by Cohen and Westreich. In the same work, Shimizu associated a central element, termed the conjugacy class sum, to each conjugacy class, which parallels the role of the sum of elements in a group conjugacy class in group theory. Shimizu also posed the question of which results from [\[CW00,](#page-20-3) [CW14\]](#page-20-5) could be further extended from semisimple Hopf algebras to fusion categories.

The primary goal of this article is to establish analogous results for the Frobenius-Perron dimensions of conjugacy classes in weakly integral fusion categories and to compare these findings with the corresponding results for finite groups. It is noteworthy that there are currently very few results concerning the Frobenius-Perron dimensions of conjugacy classes in fusion categories. Cohen and Westreich proved in [\[CW14,](#page-20-5) Theorem 4.5] that a simple non-commutative quasitriangular semisimple Hopf algebra possesses no non-trivial conjugacy classes whose dimensions are powers of a prime number.

The key ingredient in our study is Harada's identity on the product of conjugacy class sums for weakly integral fusion categories, see [\[Bur24,](#page-19-0) Theorem 1.2]. Our first main results in this direction are the following:

Let C be a weakly integral fusion category and $\{\mathcal{C}^j\}_{j=0}^m$ the set of all its conjugacy classes and C_j their associated class sums.

Theorem 1.1. *In any weakly integral braided fusion category* C *with a commutative Grothendieck ring* K(C) *the following holds:*

(1.1)
$$
\frac{\mathrm{FPdim}(\mathcal{C}_{\mathrm{pt}})}{\mathrm{FPdim}(\mathcal{C})}\left[\prod_{j=0}^{m}\mathrm{FPdim}(\mathcal{C}^{j})\right]^{2\mathrm{rank}(\mathcal{C})+1} \in \mathbb{Z}.
$$

For the definition of conjugacy classes \mathcal{C}^j and their associated class sums C_j , refer to Section [2.](#page-3-0) Moreover, \mathcal{C}_{pt} denotes the maximal pointed fusion subcategory of C.

Theorem 1.2. *Let* C *be a weakly integral braided fusion category, and let* p *be a prime divisor of* $\text{FPdim}(\mathcal{C})$ *such that* $\text{FPdim}(\mathcal{C}) = p^{\alpha}N$ *with* $\alpha \geq 1$ and $(p, N) = 1$. If p does not divide $\text{FPdim}(\mathcal{C}_i)$ for any conjugacy *class* C^j *of* C *then* p^{α} *divides* $\text{FPdim}(\mathcal{C}_{\text{pt}})$ *.*

A well-known result, the Ito-Michler theorem, see [\[Mic86\]](#page-20-6), states that the degree of every irreducible character of a finite group G is coprime to a prime p if and only if a Sylow p -subgroup of G is both abelian and normal.

In a similar vein, for weakly-integral modular tensor categories, we obtain the following:

Proposition 1.3. *Let* C *be a weakly-integral modular fusion category, and let* p *be a prime divisor of* $\text{FPdim}(\mathcal{C})$ *such that* $\text{FPdim}(\mathcal{C}) = p^{\alpha}N$ *with* $\alpha \geq 1$ *and* $(p, N) = 1$ *. If* p *does not divides the Frobenius-Perron dimension of any of its simple objects then* p ^α *divides* |U(C)|*, the order of the universal grading group of* C*.*

Furthermore, in this case we have:

 $\mathcal{C} \simeq \mathcal{D} \boxtimes \mathcal{D}'$

where $\mathcal D$ *is a pointed modular tensor category of dimension* p^{α} and $\mathcal D'$ *is a modular tensor category of dimension* N*.*

As a direct consequence of the above result we obtain the following:

Theorem 1.4. *Every weakly integral modular category* C *decomposes as a Deligne product* $\mathcal{C} \simeq \mathcal{D} \boxtimes \mathcal{E}$ *of two weakly-integral modular categories where:*

- (1) D *is pointed.*
- (2) FPdim(D) and FPdim(E) are coprime integers.
- (3) E has the property that, for any prime divisor $p \mid \text{FPdim}(\mathcal{E})$, *there exists a simple object in* E *whose Frobenius-Perron dimension is divisible by* p*.*

For braided nilpotent fusion categories, we establish the following divisibility property:

Theorem 1.5. *In any braided nilpotent fusion category* C*, the following holds:*

$$
\mathrm{FPdim}(\mathcal{C}^j) \mid \frac{\mathrm{FPdim}(\mathcal{C})}{\mathrm{FPdim}(\mathcal{C}_{\mathrm{pt}})}.
$$

for any conjugacy class C ^j *of* C*. Moreover, the set of prime numbers dividing* $FPdim(\mathcal{C})/FPdim(\mathcal{C}_{pt})$ *coincides with the set of primes numbers dividing at least one element of the set* $\{FPdim(\mathcal{C}^j)\}_{j\in\mathcal{J}}$ *.*

The structure of this paper is as follows: In Section [2,](#page-3-0) we review the basic concepts of pivotal fusion categories from [\[Shi17\]](#page-20-4) that are essential for the rest of the paper. Section [3](#page-7-0) presents several results concerning the characters of Grothendieck rings in fusion categories, which are crucial for proving Theorem [1.5.](#page-2-0) These results may also hold independent theoretical interest. In Section [4,](#page-11-0) we establish Theorem [1.1](#page-1-0) and its related results, Proposition [1.3](#page-2-1) and Theorem [1.4.](#page-2-2) Finally, in Section [5,](#page-17-0) we prove Theorem [1.5](#page-2-0) along with some other consequences.

Acknowledgements. The author thanks Sebastien Palcoux for insightful discussions and for carefully reviewing a preliminary version of this manuscript.

2. Preliminaries

In this paper, we work over the base field $\mathbb C$. For a comprehensive overview of fusion categories, we direct the reader to the monograph [\[EGNO15\]](#page-20-7).

Let $\mathcal C$ be a fusion category. We denote the maximal pointed fusion subcategory of C by \mathcal{C}_{pt} , and let Irr(C) be a set of representatives for the isomorphism classes of simple objects in C. The size of $\text{Irr}(\mathcal{C})$ is referred to as the *rank of* C and is denoted by rank (C) .

A *pivotal structure* in a rigid monoidal category C is an isomorphism $j : id_{\mathcal{C}} \to (\mathcal{V}^*)^*$ between monoidal functors. A *pivotal monoidal category* refers to a rigid monoidal category equipped with such a pivotal structure. Recall from [Mü03] that a pivotal fusion category is *spherical* if and only if $\dim(V) = \dim(V^*)$ for all simple objects V, where dim denotes the *quantum dimension* of C. A fusion category C is called *pseudo-unitary* if its Frobenius–Perron dimension $FPdim(\mathcal{C})$ equals its categorical dimension $\dim(\mathcal{C})$. In this case, according to [\[ENO05,](#page-20-9) Proposition 8.23], the category C possesses a unique spherical structure, known as the *canonical spherical structure*, under which the categorical dimensions of all simple objects are positive. With this spherical structure, the categorical dimension of any object X in $\mathcal C$ matches its Frobenius–Perron dimension, i.e., $FPdim(X) = dim(X)$ for every object X .

Recall that a fusion category is called *weakly integral* if its Frobenius-Perron is an integer. Moreover, the fusion category is called *integral* if the Frobenius-Perron dimension of each object is an integer. Any weakly integral fusion category is inherently pseudo-unitary and possesses a canonical spherical structure, under which the categorical dimensions match the Frobenius–Perron dimensions [\[ENO05,](#page-20-9) Proposition 8.24].

For any fusion category \mathcal{C} , the monoidal center (or the Drinfeld center) of C is a braided fusion category $\mathcal{Z}(\mathcal{C})$ equipped with a forgetful functor $F : \mathcal{Z}(\mathcal{C}) \to \mathcal{C}$, see [\[EGNO15\]](#page-20-7).

The forgetful functor F also admits a right adjoint functor $R : \mathcal{C} \to$ $\mathcal{Z}(\mathcal{C})$ and $Z := FR : \mathcal{C} \to \mathcal{C}$ is a Hopf comonad defined as an end. Indeed, following [\[Shi17,](#page-20-4) Section 2.6] one has that

(2.1)
$$
Z(V) \simeq \int_{X \in \mathcal{C}} X \otimes V \otimes X^*
$$

It is known that $A := Z(1)$ (the image of the unit object) has the structure of a (central) commutative algebra in $\mathcal{Z}(\mathcal{C})$, and it is called *the adjoint algebra* of C.

The vector space $CE(\mathcal{C}) := Hom_{\mathcal{C}}(1, A)$ is referred to as the *the set of central elements*. The vector space $CF(\mathcal{C}) := Hom_{\mathcal{C}}(A, \mathbf{1})$ is referred to as the space of class functions of C. Both $CF(\mathcal{C})$ and $CE(\mathcal{C})$ have a semisimple C-algebra structure, as described in [\[Shi17\]](#page-20-4). Additionally, $CE(\mathcal{C})$ is a commutative algebra, which means it decomposes as a direct product of copies of the field C.

Since $R : \mathcal{C} \to \mathcal{Z}(\mathcal{C})$ is a right adjoint to the forgetful functor $F : \mathcal{Z}(\mathcal{C}) \to \mathcal{C}$, [\[Shi17,](#page-20-4) Theorem 3.8] establishes that this adjunction induces an isomorphism of C-algebras:

(2.2)
$$
\qquad \text{CF}(\mathcal{C}) \xrightarrow{\cong} \text{End}_{\mathcal{Z}(\mathcal{C})}(R(\mathbf{1})), \ \ \chi \mapsto Z(\chi) \circ \delta_{\mathbf{1}}.
$$

where δ_1 is the comultiplication of the Hopf comonad Z associated at the unit 1 of the fusion category \mathcal{C} .

Example 2.1. If H is a semisimple Hopf algebra, then, for $C =$ Rep(*H*)*, as shown in* [\[Shi17,](#page-20-4) Subsect. 3.7]*, it follows that* $CF(\mathcal{C}) \simeq$ C(H)*, the* character algebra *of* H*. Recall that* C(H) *consists of all linear trace maps* $f : H \to \mathbb{C}$.

2.1. The bijection between central primitive idempotents and conjugacy classes. Suppose that $\mathcal C$ is a pivotal fusion category with a commutative Grothendieck ring $K(\mathcal{C})$. Since the monoidal center $\mathcal{Z}(\mathcal{C})$ is also a fusion category, we can write $R(\mathbf{1}) = \bigoplus_{i=0}^{m} C_i$ as a direct sum of simple objects in $\mathcal{Z}(\mathcal{C})$. Note that any two distinct conjugacy classes are not isomorphic to each other, since from the isomorphism of Equation (2.2) the endomorphism ring of $R(1)$ is a commutative C-algebra. Thus $\mathcal{C}^0, \ldots, \mathcal{C}^m$ represent the conjugacy classes of \mathcal{C} . Since the unit object $\mathbf{1}_{\mathcal{Z}(C)}$ is always a sub-object of $R(1)$, we can assume $C^0 = \mathbf{1}_{\mathcal{Z}(C)}$. In [\[Shi17,](#page-20-4) Section 5], Shimizu also introduced a canonical pairing between $CF(\mathcal{C})$ and $CE(\mathcal{C})$ with values in \mathbb{C} . This pairing is defined by the condition:

$$
(2.3) \t\t f \circ a = \langle f, a \rangle = id_1
$$

for all $f \in CF(\mathcal{C})$ and $a \in CE(\mathcal{C})$.

Let $\tilde{F}_0, \tilde{F}_1, \ldots, \tilde{F}_m \in \text{End}_{\mathcal{Z}(\mathcal{C})}(R(1))$ be the canonical projections onto each of these conjugacy classes. Let also $F_0, F_1, \ldots F_m$ be also the corresponding primitive idempotents of $CF(\mathcal{C})$ under the canonical adjunction isomorphism $CF(\mathcal{C}) \simeq End_{\mathcal{Z}(\mathcal{C})}(R(1))$ from Equation [\(2.2\)](#page-4-0). This establishes a canonical bijection between the (central) primitive idempotents of $CF(\mathcal{C})$ and the conjugacy classes of \mathcal{C} .

Let $\mathcal{J} := \{0, 1, \ldots, m\}$ represent the set of indices corresponding to the central primitive idempotents of CF(C). To each F_i with $j \in \mathcal{J}$, the bijection described above assigns a conjugacy class \mathcal{C}^j .

2.2. On the definition of conjugacy classes. In a pivotal fusion category, Shimizu defines the elements $C_j := \mathcal{F}_{\lambda}^{-1}$ $\chi^{-1}(F_j) \in \mathrm{CE}(\mathcal{C})$ as the *conjugacy class sums* corresponding to the conjugacy class \mathcal{C}^j . Here, $\lambda \in \mathrm{CF}(\mathcal{C})$ is a cointegral of $\mathcal C$ such that $\langle \lambda, u \rangle = 1$ (see [\[Shi17,](#page-20-4) Section 5]). Here $u : \mathbf{1} \to A$ is the unit of the central algebra A. Additionally, the *Fourier transform* of C associated with λ is defined as the linear map:

(2.4)
$$
\mathcal{F}_{\lambda} : CE(\mathcal{C}) \to CF(\mathcal{C})
$$
 given by $a \mapsto \lambda \leftarrow \mathcal{S}(a)$

where S denotes the antipode.

Example 2.2. If $C = \text{Vec}_G^{\omega}$ is a pointed fusion category, then each *conjugacy class sum of* G *corresponds to a representation of the group algebra* $\mathbb{C}[G]$ *. Furthermore, the space of central elements* $CE(\mathcal{C})$ *is isomorphic to* $K(G)$ *, the Grothendieck group of* $\mathbb{C}[G]$ *. In particular, if* G *is abelian, the conjugacy classes of* C *can be identified with elements of* G *, since in this case, the dual group* \widehat{G} *is isomorphic to* G *.*

2.3. The integral of a fusion category. Let $\mathcal C$ be a fusion category, and let $A = Z(1)$ be its adjoint algebra as defined earlier. Following [\[Shi17,](#page-20-4) Definition 5.7], an *integral* in C is a morphism $\Lambda: 1 \to A$ in C that satisfies

$$
m\circ (\mathrm{id}_A\otimes\Lambda)=\epsilon_1\otimes\Lambda.
$$

where $m : A \otimes A \rightarrow 1$ is the multiplication of the adjoint algebra A. Here, ϵ_1 is counit $\epsilon_V : Z(V) \to V$ specialized at $V = 1$, the unit of the fusion category \mathcal{C} .

It is well known that the integral of a finite fusion category is unique up to a scalar. Let $\Lambda \in \text{CE}(\mathcal{C})$ be the idempotent integral of C and define $\widehat{\Lambda} := \dim(\mathcal{C})\Lambda$.

Let $\{V_0, \ldots, V_m\}$ be a complete set of representatives of the isomorphism classes of simple objects in \mathcal{C} . Shimizu, in [\[Shi17,](#page-20-4) Section 6], provided a construction of the (central) primitive elements $E_i \in \text{CE}(\mathcal{C}),$ corresponding to the irreducible characters $\chi_i := \chi_{V_i}$ associated to each simple object V_i of $\mathcal C$. This correspondence is defined by the condition:

$$
\langle \chi_i, E_j \rangle = \delta_{i,j},
$$

for all i, j .

Note that [\[Bur20,](#page-19-1) Lemma 4.1], together with Equation (4.7) from the same paper, provides an alternative pair of dual bases for the canonical

pairing. More precisely, it shows that $\{F_j\}$ and $\frac{C_j}{\dim(C_j)}$ form another pair of dual bases expressed as:

$$
(2.5) \t\t\t $F_j, \frac{C_{j'}}{\dim(\mathcal{C}^{j'})} >= \delta_{j,j'}$
$$

for all $j, j' \in \mathcal{J}$.

Without loss of generality, we can assume that $V_0 = 1$, the unit of the fusion category C. This gives $\chi_0 = \epsilon_1$, and the associated primitive idempotent is $E_0 = \Lambda_{\mathcal{C}}$, see [\[Shi17,](#page-20-4) Lemma 6.1].

The following lemma is straightforward, but for reader's convenience, a proof is provided.

Lemma 2.1. *Let* C *be a pivotal fusion category with a commutative Grothendieck ring. We have the following formula for the integral element of* C*:*

(2.6)
$$
\widehat{\Lambda}_{\mathcal{C}} = \sum_{j \in \mathcal{J}} C_j.
$$

Proof. We will show that both sides of the Equation (2.6) have the same image under \mathcal{F}_{λ} . For the right hand side, by the definition of conjugacy class sums C_j , we have $\mathcal{F}_\lambda(\sum_{j\in\mathcal{J}} C_j) = \sum_{j\in\mathcal{J}} F_j = \epsilon_1$. For the left hand side, according to [\[Shi17,](#page-20-4) Equation (6.10)], we have

(2.7)
$$
\mathcal{F}_{\lambda}(E_{i^*}) = \frac{d_i}{\dim(\mathcal{C})} \chi_i
$$

for any $0 \le i \le m$. In the case $i = 0$, one has $\chi_0 = \epsilon_1$ and $E_0 = \Lambda_{\mathcal{C}}$. Since $d_0 = 1$ it follows that $\mathcal{F}_{\lambda}(E_0) = \epsilon_1$.

2.4. Definition of the support $\mathcal{J}_\mathcal{D}$. Let C be a fusion category whose Grothendieck ring is commutative. For any fusion subcategory $\mathcal{D} \subseteq \mathcal{C}$, recall the notion of its support from [\[Bur20,](#page-19-1) Subsection 4.2]. It is well known that there is an inclusion of $\mathbb{C}\text{-algebras } \iota : \text{CF}(\mathcal{D}) \hookrightarrow \text{CF}(\mathcal{C})$. Consequently, there exists a subset $\mathcal{J}_{\mathcal{D}}^{\mathcal{C}} \subseteq \mathcal{J}^{\mathcal{C}} := \{0, \ldots, m\}$ such that

(2.8)
$$
\lambda_{\mathcal{D}} = \sum_{j \in \mathcal{J}_{\mathcal{D}}} F_j
$$

since $\lambda_{\mathcal{D}}$ is an idempotent element in CF(C). The set $\mathcal{J}_{\mathcal{D}}$ is referred to as the *support* of the fusion subcategory D.

For a fusion category \mathcal{C} , the fusion subcategory \mathcal{C}_{ad} represents the trivial component in the universal grading of \mathcal{C} (see [\[GN08\]](#page-20-10)). Alternatively, \mathcal{C}_{ad} is the smallest fusion subcategory of $\mathcal C$ which contains all the objects $X \otimes X^*$, with $X \in \text{Irr}(\mathcal{C})$.

A grading of a fusion category $\mathcal C$ by a group G is defined by a map $deg: Irr(\mathcal{C}) \to G$ with the property that for any simple objects X, Y, Z of C, if $X \otimes Y$ contains Z then $deg(Z) = deg(X) deg(Y)$. The term "grading" also refers to the decomposition $\mathcal{C} = \bigoplus_{g \in G} \mathcal{C}_g$, where \mathcal{C}_g is the full additive subcategory generated by the simple objects of degree $g \in G$.

The subcategory C_1 corresponding to $g = 1$, is called the *trivial component of the grading* and is a fusion subcategory. A grading is *trivial* if $C_1 = C$, and *faithful* if the map deg : $\text{Irr}(\mathcal{C}) \rightarrow G$ is surjective. The set of all possible gradings of $\mathcal C$ by G depends functorially on G , leading to the concept of a *universal grading*. As shown in [\[GN08,](#page-20-10) Section 3.2], such a grading always exists, and the associated group is denoted by $U(\mathcal{C})$. The following results are proven in [\[GN08,](#page-20-10) Section] 3.2]: (see [\[GN08,](#page-20-10) (Lemma 3.4, Theorem 3.5, and Proposition 3.9)].

- (1) Any faithful grading of $\mathcal C$ by a group G is determined by a surjective group homomorphism $\pi: U(\mathcal{C}) \to G$.
- (2) The universal grading deg : $\text{Irr}(\mathcal{C}) \to U(\mathcal{C})$ is faithful.
- (3) The trivial component of the universal grading equals \mathcal{C}_{ad} .
- (4) Any fusion subcategory $\mathcal{D} \subseteq \mathcal{C}$ that contains \mathcal{C}_{ad} can be written as form $\mathcal{D} = \bigoplus_{g \in G} C_g$ for some subgroup $G \leq U(\mathcal{C})$.

It follows that $\text{FPdim}(\mathcal{C}) = |U(\mathcal{C})|\text{FPdim}(\mathcal{C}_{ad})$.

Let C be a fusion category. Following [\[GN08\]](#page-20-10), we define a sequence of subcategories: $\mathcal{C}^{(0)} = \mathcal{C}, \, \mathcal{C}^{(1)} = \mathcal{C}_{ad}$, and for all $n \geq 1, \, \mathcal{C}^{(n)} = \mathcal{C}_{ad}^{(n-1)}$.

The descending sequence

$$
C = C^{(0)} \supseteq C^{(1)} \supseteq \cdots \supseteq C^{(n)} \supseteq \ldots
$$

is called the *upper central series*. A fusion category $\mathcal C$ is said to be *nilpotent* if its upper central eventually reaches Vec (the category of finite dimensional C-vector spaces), meaning that $\mathcal{C}^{(n)}$ = Vec for some n. The smallest such n is called the *nilpotency class* of C.

3. Restriction of characters of Grothendieck rings

Throughout this section, let $\mathcal C$ be a fusion category with a commutative Grothendieck ring, $K(\mathcal{C})$. Then $CF(\mathcal{C})$ is a commutative semisimple algebra, and $CF(\mathcal{C}) \simeq K(\mathcal{C})$, as shown in [\[Shi17,](#page-20-4) Example 4.4]. As above, Let $\mathcal{J}^{\mathcal{C}} := \{0, \ldots, m\}$ represent the set of indices corresponding to the conjugacy classes of \mathcal{C} . It is clear that the rank of \mathcal{C} is rank $(C) = |\mathcal{J}^c| = m + 1$.

Suppose $\mathcal D$ is a fusion subcategory of $\mathcal C$. Note that the Grothendieck ring $K(\mathcal{D})$ is also commutative, and, as in the case above, $K(\mathcal{D}) \simeq$ $CF(\mathcal{D})$ as algebras. We let $\mathcal{J}^{\mathcal{D}} := \{0, \ldots, m'\}$ represent the set of

indices corresponding to the conjugacy classes of \mathcal{D} . As above, it is clear that the rank of D is rank $(\mathcal{D}) = |\mathcal{J}^{\mathcal{D}}| = m + 1$.

For any object $V \in \mathcal{C}$, following [\[Shi17,](#page-20-4) Sect. 4.3], consider the end

$$
\bar{Z}(V): \mathcal{D}^{\mathrm{op}} \times \mathcal{D} \to \mathcal{C}, \quad \bar{Z}(V) := \int_{X \in \mathcal{D}} X \otimes V \otimes X^*.
$$

Let $\bar{\pi}_{V; X}: \mathbb{Z}(V) \to X \otimes V \otimes X^*$ denote the universal dinatural maps that define this end. By the universal property of \overline{Z} , for any $V \in \mathcal{C}$ there exists a unique canonical map in \mathcal{C} :

$$
(3.1)\quad Z(V) \xrightarrow{q_V} \bar{Z}(V)
$$

such that $\bar{\pi}_{V; X} \circ q_V = \pi_{V; X}$ for every object X in \mathcal{D} .

In [\[Bur20,](#page-19-1) Appendix] we demonstrated that q is a morphism of Hopf comonads. The map $q_1 : Z(1) \to Z(1)$ induces two associated maps:

$$
\pi_f = q_1^* : \text{CF}(\mathcal{D}) \to \text{CF}(\mathcal{C}), \ \chi \mapsto \chi \circ q_1,
$$

$$
\pi_e = q_{1*} : \text{CE}(\mathcal{C}) \to \text{CE}(\mathcal{D}), \ z \mapsto q_1 \circ z.
$$

Furthermore, by [\[Bur20,](#page-19-1) Lemma 7.7] it is known that q_1^* is a monomorphism and q¹[∗] is an epimorphism. Both maps are, in fact, C-algebra homomorphisms.

Example 3.1. If π : $H \rightarrow K$ *is a surjective Hopf algebra homomorphism, then clearly we have* $\text{Rep}(K) \subseteq \text{Rep}(H)$ *. In this case,* $\pi_e = \pi|_{Z(H)} : Z(H) \rightarrow Z(K)$ is the restriction of π to the center $Z(H)$ *of* H. Furthermore, π_f corresponds to the canonical inclusion of *the character algebra* $C(K)$ *of* K *into that* H *, i.e,* $C(K) \hookrightarrow C(H)$ *.*

Let, ${F_i}_{0 \leq i \leq m}$ be the (central) primitive idempotents of CF(C), and let $\{\mathcal{C}^j\}_{0\leq j\leq m}$ be their associated conjugacy classes. For each (central) primitive idempotent $F_j \in CF(\mathcal{C})$, let $\mu_j : CF(\mathcal{C}) \to \mathbb{C}$ denote its associated linear character. Similarly, let ${G_t}_{0\leq t\leq m'}$ be the (central) primitive idempotents of CF(D), with $\{\mathcal{D}^t\}_{0\leq t\leq m'}$ being their associated conjugacy classes. Additionally, let $\nu_t : CF(\mathcal{D}) \to \mathbb{C}$ denote the characters associated with these idempotents.

Using the map π_f we can identify $CF(\mathcal{D})$ as subring of $CF(\mathcal{C})$.

From the proof of [\[Bur23,](#page-19-2) Theorem 6.12], for all $\alpha \in \mathrm{CF}(\mathcal{C})$, we have

(3.2)
$$
\mu_j(\alpha) = \langle \alpha, \frac{C_j}{\dim(C^j)} \rangle
$$

We say that two characters $\mu_j, \mu_{j'} : \text{CF}(\mathcal{C}) \to \mathbb{C}$ are equivalent or *in the same class*, (denoted $\mu_j \simeq \mu_{j'}$) with respect to \mathcal{D} if their restrictions to $CF(D)$ are the same.

The set of equivalence classes is denoted by $Cl(C : D)$ and it is referred to as *the set of classes* of C with respect to D . This set is in bijection with the set of the characters of $\nu_t : CF(D) \to \mathbb{C}$ with $0 \leq t \leq m'.$

Moreover, an element $C \in \text{Cl}(\mathcal{C} : \mathcal{D})$ consists of all the indices j such that the restriction $\mu_j \downarrow_{CF(D)}$ corresponds to the same character in CF(D), denoted as $\rho_C := \nu_s$ for some $s \leq m'$. Thus, we use ρ_C to represent the restriction to $CF(D)$ of all characters μ_i belonging to some class $C \in \text{Cl}(\mathcal{C} : \mathcal{D}).$

As noted above, the set of character classes of C relative to $\mathcal D$ corresponds bijectively to the characters of $CF(D)$. Consequently, the central primitive idempotents of $CF(D)$ are also indexed by the classes $C \in \text{Cl}(\mathcal{C} : \mathcal{D})$. We denote the central primitive idempotent associated with a class $C \in \text{Cl}(\mathcal{C} : \mathcal{D})$ by $G_C \in \text{CF}(\mathcal{D})$. Thus, the elements G_C for $C \in \text{Cl}(\mathcal{C} : \mathcal{D})$ form a linear basis for $\text{CF}(\mathcal{D})$, consisting of the complete set of (central) primitive idempotents of the commutative Calgebra $CF(\mathcal{D})$. For the remainder of this section, we identify the set of indices $\mathcal{J}^{\mathcal{D}}$ of the central primitive idempotents of $CF(\mathcal{D})$ with the elements of $Cl(C : \mathcal{D})$. For any primitive idempotent G_C of $CF(\mathcal{D})$, as described in Subsection [2.1,](#page-4-1) let $D_C \in CE(\mathcal{D})$ denote the corresponding conjugacy class sum of $\mathcal D$ associated with G_C .

Proposition 3.1. Let C be a pivotal fusion category with a commuta*tive Grothendieck ring,* K(C)*, and let* D *be a fusion subcategory. With the above notation, we have*

(3.3)
$$
\pi_f(G_C) = \sum_{j \in C} F_j
$$

for any class C*.*

Proof. Since π_f is an algebra homomorphism and $\pi_f(G_s)$ is idempotent, it can be expressed as a sum of primitive idempotents in $CF(\mathcal{C})$. Suppose that $\pi_e(G_C) = \sum_{j \in \mathcal{B}_C} F_j$ for some subset $\mathcal{B}_C \subseteq \{0, 1, \ldots, m\}.$ This implies that $\mu_j(G_m) = \mu_{j'}(G_m) = 1$ for any $j, j' \in \mathcal{B}_C$, indicating that j and j' belong to the same character class, denoted as C . Indeed, we have $\mu_j(G_t) = \mu_{j'}(G_t) = 0$ for any $t \neq s$, which leads to $\mu_j|_{\text{CF}(\mathcal{D})} = \mu_{j'}|_{\text{CF}(\mathcal{D})}.$

Conversely, if $j' \in C$ then $\mu_{j'} \downarrow_{CF(D)} = \mu_j \downarrow_{CF(D)}$. This implies that $\mu_{j'}(G_C) = 1$, and therefore $j' \in \mathcal{B}_C$.

Recall the canonical pairing between $CF(\mathcal{C})$ and $CE(\mathcal{C})$, which we denote as $\langle, \rangle_{\mathcal{C}}$. For the fusion subcategory $\mathcal{D} \subseteq \mathcal{C}$, there is an analogous pairing, written here as $\langle, \rangle_{\mathcal{D}}$. The proof of the following lemma follows directly from the definitions of the two canonical pairings.

Lemma 3.2. *The two canonical pairings on the fusion categories* C *and* D *are compatible, satisfying the relation:*

$$
(3.4) $\pi_f(\chi), \ z >_c = <\alpha, \pi_e(z) >_D$
$$

for all $\chi \in \mathrm{CF}(\mathcal{D})$ *and* $z \in \mathrm{CE}(\mathcal{C})$ *.*

3.1. On the projection of conjugacy class sums.

Theorem 3.3. *Let* C *be a pivotal fusion category with a commutative Grothendieck ring, and* D *be a fusion subcategory of* C *. Let* $j \in \mathcal{J}$ *be any index and suppose that* $j \in C$ *for some* $C \in \text{Cl}(\mathcal{C} : \mathcal{D})$ *. Then:*

(3.5)
$$
\pi_e\left(\frac{C_j}{\dim(C^j)}\right) = \frac{D_C}{\dim(D^C)}
$$

Proof. We can write

$$
\pi_e(\frac{C_j}{\dim(\mathcal{C}^j)}) = \sum_{C_1 \in \text{Cl}(\mathcal{C}:\mathcal{D})} \alpha_{jC_1} \mathcal{D}_{C_1}
$$

for some scalars $\alpha_{i}C_1 \in \mathbb{C}$.

Let now $C_2 \in \text{Cl}(\mathcal{C} : \mathcal{D})$ be an arbitrary class, and let $\chi = G_{C_2}$ and $z = C_j$. By Lemma [3.2](#page-10-0) we have:

$$
<\sum_{s\in C_2} F_s, \frac{C_j}{\dim(C^j)}>_c=_{\mathcal{D}},
$$

which in turn can be written as

$$
<\sum_{s \in C_2} F_s, \frac{C_j}{\dim(C^j)} >_c = _p.
$$

The right hand side of the last above equation, using the dual bases form Equation [2.5](#page-6-1) for D , can be written as:

$$
\langle G_{C_2}, \sum_{C_1 \in \text{Cl}(\mathcal{C}:\mathcal{D})} \alpha_{jC_1} D_{C_1} >_{\mathcal{D}} = \alpha_{jC_2} \dim(\mathcal{D}^{C_2}).
$$

On the other hand, using also the dual bases form Equation [2.5,](#page-6-1) for C this time, the left hand side equals $\delta_{j\in C_2}$.

Thus, we find that α_{jC_2} dim $(\mathcal{D}^{C_2}) = \delta_{j \in C_2}$ for all character classes $C_2 \in \text{Cl}(\mathcal{C} : \mathcal{D})$. Since $j \in C$, this gives that $\alpha_{jC} = \frac{1}{\dim(\mathcal{D}^C)}$ and $\alpha_{jC_2} = 0$ for all $C_2 \neq C$. This establishes Equation [\(3.5\)](#page-10-1).

For a fusion category $\mathcal C$ we define the natural coend projection $\epsilon_{\mathcal C}$: $A \to \mathbf{1} \in \mathrm{CF}(\mathcal{C})$ as the map $A \xrightarrow{\pi_{1,1}} \mathbf{1} \otimes \mathbf{1}^* \simeq \mathbf{1}$. This projection is referred to as ϵ_1 in [\[Shi17\]](#page-20-4), but we use the notation ϵ_c to emphasize the category \mathcal{C} .

Recall that in the previous section, we defined $\chi_0 \in \mathrm{CF}(\mathcal{C})$ as the irreducible character corresponding to the tensor unit $\mathbf{1}_{\mathcal{C}}$ of \mathcal{C} . Without loss of generality, we can also assume that $\alpha_0 \in \mathrm{CF}(\mathcal{D})$ is the character associated to the unit object $1_{\mathcal{D}}$ of \mathcal{D} . It is clear that from its definition that $\pi_f(\alpha_0) = \chi_0$.

Lemma 3.4. Let $\mathcal{D} \subseteq \mathcal{C}$ be a fusion subcategory of a pivotal fusion *category* C*. Under the above notation, we have:*

$$
(3.6) \qquad \qquad \epsilon_{\mathcal{D}} \circ q_1 = \epsilon_{\mathcal{C}}
$$

and

$$
(3.7) \t\t \pi_e(\Lambda_{\mathcal{C}}) = \Lambda_{\mathcal{D}}
$$

Proof. From the naturality of q_1 , we can conclude that $\epsilon_{\mathcal{D}} \circ q_1 = \epsilon_{\mathcal{C}}$, since $\epsilon_{\mathcal{C}}: A \to \mathbf{1}$ serves as the universal end projection $A \xrightarrow{\pi_{1,1}} \mathbf{1} \otimes \mathbf{1}^* \simeq \mathbf{1}$.

For the second assertion, we first show that $\pi_e(\Lambda_{\mathcal{C}}) \neq 0$. According to [\[Shi17,](#page-20-4) Lemma 6.1], the corresponding idempotent of χ_0 is $E_0 = \Lambda_c$. By applying Lemma [3.2,](#page-10-0) we have:

$$
1 = \langle \pi_f(\alpha_0), E_0 \rangle = \langle \alpha_0, \pi_e(\Lambda_c) \rangle,
$$

which implies that $\pi_e(\Lambda_{\mathcal{C}}) \neq 0$. Moreover, for any $z \in \text{CE}(\mathcal{C})$, we find:

$$
\epsilon_{\mathcal{C}}(z)\pi_{e}(\Lambda_{\mathcal{C}})=\pi_{e}(z\Lambda_{\mathcal{C}})=\pi_{e}(z)\pi_{e}(\Lambda_{\mathcal{C}})
$$

Note that $\epsilon_{\mathcal{D}}(\pi(z)) = \epsilon_{\mathcal{C}}(z)$, and since π_e is surjective, it follows that:

$$
w\pi_e(\Lambda_{\mathcal{C}})=\epsilon_{\mathcal{D}}(w)\pi_e(\Lambda_{\mathcal{C}})
$$

for any $w \in \text{CE}(\mathcal{D})$. Given that $\pi_e(\Lambda_{\mathcal{C}}) \neq 0$ it follows $\pi_e(\Lambda_{\mathcal{C}}) = \Lambda_{\mathcal{D}}$. \Box

4. On the dimensions of conjugacy classes

For any weakly integral fusion category, whose Grothendieck group $K(\mathcal{C})$ is commutative, the following Harada type identity holds, as established in [\[Bur24,](#page-19-0) Theorem 1]:

(4.1)
$$
(\prod_{j=0}^{m} \frac{C_j}{\mathrm{FPdim}(\mathcal{C}^j)})^2 = \frac{\mathrm{FPdim}(\mathcal{C}_{\mathrm{pt}})}{\mathrm{FPdim}(\mathcal{C})} \left(\sum_{j \in \mathcal{J}_{\mathcal{C}_{\mathrm{pt}}}} C_j\right)
$$

Recall that the support of D in C is the subset of indices of $\mathcal{J}_\mathcal{D} \subseteq \mathcal{J}$ such that $\lambda_{\mathcal{D}} = \sum_{j \in \mathcal{J}_{\mathcal{D}}} F_j$.

Since the elements $\{C_i\}$ form a linear basis for $CE(\mathcal{C})$, we can express the product of two basis elements as:

(4.2)
$$
C_i C_j = \sum_{k=0}^{m} c_{ij}^k C_k.
$$

where $c_{ij}^k \in \mathbb{C}$ are *the structure constants* of C. It is known that these constants are in fact integers in the case $\mathcal{C} = \text{Rep}(G)$ for a finite group G. Burnside provided a formula relating these structure constants to the values of irreducible characters, see [\[Isa76,](#page-20-11) pp. 45]. A similar result for the structure constants of a pivotal fusion category with a commutative Grothendieck ring was presented in [\[Bur21,](#page-19-3) Theorem 1.1].

A braided fusion category with a spherical structure is known as a *premodular* fusion category. The structure constants for premodular categories were investigated in [\[Bur21\]](#page-19-3).

In the case of premodular categories, a specific formula for the structure constants c_{ij}^k was provided in the proof of [\[Bur21,](#page-19-3) Theorem 1.2]. It was demonstrated that:

(4.3)
$$
C_i C_j = \sum_{k=1}^m \left(\sum_{v \in \mathcal{A}_k} \frac{\widehat{N}_{\sigma(i)\sigma(j)}^v \widehat{d}_v}{\dim(\mathcal{C}^k)} \right) C_k
$$

which leads to the following expression for the structure constants:

(4.4)
$$
c_{ij}^k = \sum_{v \in \mathcal{A}_k} \frac{\widehat{N}_{\sigma(i)\sigma(j)}^v \widehat{d}_v}{\dim(\mathcal{C}^k)}.
$$

By [\[ENO05,](#page-20-9) Prop 8.23], since every weakly integral braided fusion category is premodular, the results of [\[Bur21,](#page-19-3) Theorem 1.2] apply in this context. Therefore from Equation [\(4.4\)](#page-12-0) we can conclude that:

$$
(4.5) \qquad \dim(\mathcal{C}^k)c_{ij}^k \in \mathbb{Z}_{\geq 0}
$$

for any structure constant c_{ij}^k in a weakly integral braided fusion category. Note the that the fusion rules $\hat{N}^v_{\sigma(i)\sigma(j)}$ are integers, and since C is weakly integral, the quantum dimensions d_v are also integers. For further details, see the proof of [\[Bur21,](#page-19-3) Theorem 1.2].

4.1. Proof of Theorem [1.1.](#page-1-0)

Proof. Let $\widehat{P} := \prod_{j=0}^m \text{FPdim}(\mathcal{C}^j)$. Since C is weakly integral, by [\[ENO05,](#page-20-9) Proposition 8.27] implies that the Frobenius–Perron dimensions of simple objects in C_{ad} are integers. Given that the conjugacy classes \mathcal{C}^j are direct sums of simple objects of the adjoint subcategory, it follows that $\mathrm{FPdim}(\mathcal{C}^j)$ are also integers and therefore $\widehat{P}\in \mathbb{Z}$.

From the proof of [\[Bur21,](#page-19-3) Theorem 1.2], we know that $\dim(\mathcal{C}^k)c_{ij}^k \in$ $\mathbb{Z}_{\geq 0}$ for any premodular fusion category. Therefore, multiplying equa-tion [\(4.2\)](#page-11-1) by \widehat{P} , we have:

$$
\widehat{P}C_iC_j = \sum_{k=0}^{m} (\widehat{P}c_{ij}^k)C_k
$$

This shows that the coefficients $\widehat{P}c_{ij}^k \in \mathbb{Z}$. By induction, for any rclass sums, the product $\widehat{P}^{r-1}C_{i_1}C_{i_2}\ldots C_{i_r}$ yields integral coefficients for each C_k when expressed as a linear combination of conjugacy class sums.

Next, by multiplying by \widehat{P}^{2m+1} in Equation [\(4.1\)](#page-11-2), we obtain:

(4.6)
$$
\widehat{P}^{2m+1}(\prod_{j=0}^{m} C_j)^2 = \frac{\widehat{P}^{2m+3} \text{FPdim}(\mathcal{C}_{\text{pt}})}{\text{FPdim}(\mathcal{C})} \big(\sum_{j \in \mathcal{J}_{\mathcal{C}_{\text{pt}}}} C_j\big)
$$

Thus, in the product $\widehat{P}^{2m+1}(\prod_{j=0}^m C_j)^2$, all C_j with $j \in \mathcal{J}_{\mathcal{C}_{\mathrm{pt}}}$ have a integral coefficient. Thus, $\frac{\hat{P}^{2m+3}\text{FPdim}(\mathcal{C}_{pt})}{\text{FPdim}(\mathcal{C})} \in \mathbb{Z}_{>0}$ and the proof is complete since $\text{rank}(\mathcal{C}) = m + 1$.

4.2. Fusion categories with rational structure constants.

Theorem 4.1. *Let* C *be a weakly integral fusion category, and suppose that all its structure constants* c_{ij}^k *are rational numbers. Let* $D \in \mathbb{Z}$ *be such that* $Dc_{ij}^k \in \mathbb{Z}$ *for all* i, j, k *. Then:*

 $(1) \frac{D^{2m+1}\hat{P}^2\text{FPdim}(\mathcal{C}_{\text{pt}})}{\text{FPdim}(\mathcal{C})} \in \mathbb{Z}.$ (2) $d_i | D$ FPdim (C) (3) $d_i | D^{2m+2} \widehat{P}^2 \text{FPdim}(\mathcal{C}_{\text{pt}})$.

Proof. The proof of the first item follows the same method as the previous theorem, but this time multiplying Harada's identity by D^{2m+1} . This ensures an integral coefficient for C_0 on the left hand side of the equation. The second item is a direct consequence of [\[BP24b,](#page-19-4) Theorem 4.7]. The third item follows directly from the first two divisibility r esults.

Lemma 4.2. *Let* C *be a weakly integral fusion category, and suppose that all its structure constants* c_{ij}^k *are rational numbers. Let* \mathcal{D} *be a fusion subcategory of* D*. Then* D *has also rational structure constants.*

Proof. Recall the map $\pi_e : CE(\mathcal{C}) \to CE(\mathcal{D})$ from Section [3.](#page-7-0) Let \mathcal{D}_E and \mathcal{D}_F be two conjugacy classes of D with $E, F \in \text{Cl}(\mathcal{C} : \mathcal{D})$. Since π_e

is a surjective map, as in Equation [\(3.5\)](#page-10-1), there are indices $s \in E$, $t \in F$ such that $\pi_e(C_t) = \frac{\dim(\mathcal{C}^t)}{\dim(\mathcal{D}^t)} D_E$ and $\pi_e(C_s) = \frac{\dim(\mathcal{C}^s)}{\dim(\mathcal{D}^s)} D_F$. Then

$$
D_E D_F = \frac{\dim(\mathcal{D}^E)}{\dim(\mathcal{C}^t)} \frac{\dim(\mathcal{D}^F)}{\dim(\mathcal{C}^s)} \pi_e(C_t) \pi_e(C_s)
$$

On the other hand, since π_e is an algebra homomorphism one has

$$
\pi_e(C_t)\pi_e(C_s) = \pi_e(C_tC_s) = \pi_e(\sum_{k=0}^m c_{ts}^k C_k)
$$

\n
$$
= \sum_{k=0}^m c_{ts}^k \frac{\dim(\mathcal{C}^k)}{\dim(\mathcal{D}^{\pi(k)})} D_{\pi(k)}
$$

\n
$$
= \sum_{C \in \text{Cl}(\mathcal{C}:\mathcal{D})} \left(\sum_{k \in C} c_{ts}^k \frac{\dim(\mathcal{C}^k)}{\dim(\mathcal{D}^C)}\right) D_C
$$

\n
$$
= \sum_{C \in \text{Cl}(\mathcal{C}:\mathcal{D})} \left(\sum_{k \in C} c_{ts}^k \dim(\mathcal{C}^k)\right) \frac{D_C}{\dim(\mathcal{D}^C)}
$$

Therefore

$$
D_t D_s = \frac{\dim(\mathcal{D}^t)}{\dim(\mathcal{C}^{t_1})}\frac{\dim(\mathcal{D}^s)}{\dim(\mathcal{C}^{s_1})}\sum_{C \in \text{Cl}(\mathcal{C}:\mathcal{D})} \big(\sum_{k \in C} c_{t_1 s_1}^k \dim(\mathcal{C}^k)\big) \frac{D_C}{\dim(\mathcal{D}^C)}
$$

which shows that the coefficient of each class sum D_C is a rational number. \Box

Proposition 4.3. *Let* C *be a weakly integral fusion category with a commutative Grothendieck ring and suppose that all its structure constants are integral, i.e.,* $c_{ij}^k \in \mathbb{Z}$ *. With above notation, one has:*

(4.7)
$$
\frac{\mathrm{FPdim}(\mathcal{C}_{\mathrm{pt}})}{\mathrm{FPdim}(\mathcal{C})}\left[\prod_{j=0}^{m}\mathrm{FPdim}(\mathcal{C}^{j})\right]^{2} \in \mathbb{Z}.
$$

Moreover, if U(C) *has odd order, then we have:*

(4.8)
$$
\frac{\mathrm{FPdim}(\mathcal{C}_{\mathrm{pt}})}{\mathrm{FPdim}(\mathcal{C})}\Big[\prod_{j=0}^{m}\mathrm{FPdim}(\mathcal{C}^{j})\Big] \in \mathbb{Z}.
$$

Proof. If C has integral structure constants then, we can take $D = 1$ since the coefficient of C_0 in the left hand side of Equation [\(4.1\)](#page-11-2) is already an integer. Therefore, in this case, there is no need to multiply the equation by \widehat{P}^{2m+1} , and we directly obtain:

$$
\frac{\mathrm{FPdim}(\mathcal{C}_{\mathrm{pt}})}{\mathrm{FPdim}(\mathcal{C})} \big[\prod_{j=0}^m \mathrm{FPdim}(\mathcal{C}^j) \big]^2 \in \mathbb{Z}.
$$

Using the dual version of [\[BP24a,](#page-19-5) Corollary 4.11], in the case in which $U(\mathcal{C})$ has odd order, we also have the second divisibility result. \square

A well-known result regarding the prime factors of the sizes of conjugacy classes in finite groups can be found, for example, in [\[CH90,](#page-19-6) Proposition 4. Let G be a finite group and p a prime divisor of G . Then, p does not divide the size of any conjugacy class of G if and only if the Sylow p-subgroup of G is contained in its center. In this case, Schur–Zassenhaus theorem, see [\[Rob96\]](#page-20-12), ensures that $G \simeq H \times K$ where H is its Sylow p-subgroup and K is another subgroup of G . In particular this shows that p^{α} divides the order of the abelian group G/G' , i.e. dimension of $\text{Rep}(G)_{pt}$

As a consequence of Theorem [1.1,](#page-1-0) a weaker version of this result can be proven for weakly integral braided fusion categories in Theorem [1.2.](#page-1-1)

4.3. Proof of Theorem [1.2.](#page-1-1)

Proof. Since $\frac{\widehat{P}^{2m+3}\mathrm{FPdim}(\mathcal{C}_{\mathrm{pt}})}{\mathrm{FPdim}(\mathcal{C})} \in \mathbb{Z}_{>0}$, and p does not divide \widehat{P} , we conclude that p^{α} \Box FPdim(\mathcal{C}_{pt}).

For finite groups we obtain the following:

Corollary 4.4. Let G be a finite group of order $|G| = p^{\alpha} N$ with $\alpha \geq 1$ and $(p, N) = 1$ *. If* p does not divide the size of any conjugacy class of G *, then* p^{α} *divides* $\frac{|G|}{|G'|}$ *.*

Proof. Corollary 1.2 implies that
$$
p^{\alpha} | \text{FPdim}(\mathcal{C}_{pt}) = \frac{|G|}{|G'|}
$$
.

4.4. Proof of Proposition [1.3.](#page-2-1)

Proof. Since C is non-degenerate, we have $\mathcal{Z}(\mathcal{C}) \simeq \mathcal{C} \boxtimes \mathcal{C}^{\text{rev}}$, as shown in [\[DGNO07,](#page-20-13) Proposition 3.7]. Thus, any simple object in $\mathcal{Z}(\mathcal{C})$ is of the form $V_i \boxtimes V_m$, and from the hypothesis, the prime p does not divide the Frobenius-Perron dimension of any such simple object. Given that the conjugacy classes of C are simple objects in $\mathcal{Z}(\mathcal{C})$, it follows that the conditions of Theorem [1.2](#page-1-1) are met. Consequently, p^{α} divides $FPdim(\mathcal{C}_{\text{pt}})$. Furthermore, in any modular tensor category, it is known from [\[GN08,](#page-20-10) Theorem 6.2] that $G(\mathcal{C}) \simeq U(\mathcal{C})$, which implies that p^{α} divides the order of $U(\mathcal{C})$.

Since p^{α} | FPdim (\mathcal{C}_{pt}) there exists a fusion subcategory a $\mathcal{D} \subseteq \mathcal{C}_{pt}$ with FPdim $(\mathcal{D}) = p^{\alpha}$. Let \mathcal{D}' be the Müger centralizer of $\mathcal D$ in $\mathcal C$. By [\[DGNO07,](#page-20-13) Theorem 3.10], we know that $\text{FPdim}(\mathcal{D}') = N$, and thus $\mathcal{D} \cap \mathcal{D}' = \text{Vec.}$ Furthermore, [\[DGNO07,](#page-20-13) Corollary 3.5] implies that $\mathcal D$ is braided non-degenerate and therefore $\mathcal C \simeq \mathcal D \boxtimes \mathcal D'$. According to [\[DGNO07,](#page-20-13) Theorem 3.10] \mathcal{D}' is also non-degenerate. \Box

Remark 4.5. *The fact that* $p | \text{FPdim}(\mathcal{C}_{\text{pt}})$ *also follows from* [\[BP24a,](#page-19-5) Equation (1.14)]*. It is worth noting that for finite groups, as shown in* [\[Mic86\]](#page-20-6)*, the proof of the Ito-Michler Theorem is significantly more complex and relies on the Classification of Finite Simple Groups.*

4.5. Proof of Theorem [1.4.](#page-2-2)

Proof. We apply Proposition [1.3](#page-2-1) repeatedly for each prime that does not divide the dimension of any simple object in C. Suppose $FPdim(\mathcal{C}) =$ $p_1^{\alpha_1} \dots p_n^{\alpha_n}$. Without loss of generality, assume that the primes p_i for $1 \leq i \leq r$ do not divide the dimension of any simple object in C.

Applying Proposition [1.3](#page-2-1) with $p = p_1$, we obtain a decomposition $\mathcal{C} = \mathcal{D}_1 \boxtimes \mathcal{E}_1$, where \mathcal{D}_1 is a pointed modular fusion category with $FPdim(\mathcal{D}_1) = p_1^{\alpha_1}$. Since $FPdim(\mathcal{E}_1) = p_2^{\alpha_2} \dots p_n^{\alpha_n}$, we can apply Propo-sition [1.3](#page-2-1) again with $p = p_2$, yielding $\mathcal{E}_1 = \mathcal{D}_2 \boxtimes \mathcal{E}_2$, where \mathcal{D}_2 is a pointed modular fusion category with $\text{FPdim}(\mathcal{D}_2) = p_2^{\alpha_2}$.

Continuing in this way for each prime p_i with $1 \leq i \leq r$, we obtain

$$
\mathcal{C} \simeq \mathcal{D}_1 \boxtimes \ldots cd_r \boxtimes \mathcal{E}_r.
$$

Thus we set $\mathcal{D} := \mathcal{D}_1 \boxtimes \ldots \mathcal{D}_r$ and $\mathcal{E} := \mathcal{E}_r$.

Corollary 4.6. *For any weakly integral perfect modular fusion category* $\mathcal{C},$ if p is a prime divisor of $\mathrm{FPdim}(\mathcal{C})$, then there exists a simple object *in* C *whose squared dimension is divisible by* p*.*

Example 4.1. Let $\mathcal{A} = \text{Rep}(D(S_3))$ be the fusion category of represen*tations of the Drinfeld double of the symmetric group* S_3 *. As described, for instance, in* [\[CW14\]](#page-20-5)*, this category contains* 2 *invertible objects,* 4 *simple objects of dimension* 2*, and* 2 *simple objects of dimension* 3*. Additionally, the* 2 *invertible objects together with the* 4 *simple objects of dimension* 2 *form a fusion subcategory* C *with Frobenius-Perron dimension* $\text{FPdim}(\mathcal{C}) = 2.1^2 + 4.2^2 = 18$ *. In fact, it can be verified that* $\mathcal{C} = \mathcal{A}_{ad}$.

For $p = 3$ *we have* $p | \text{FPdim}(\mathcal{C}) \text{ (with } \alpha = 2)$ *. Moreover,* p *does not divide the dimension of any simple object in* \mathcal{C} *, and* $p \nmid \text{FPdim}(\mathcal{C}_{pt}) = 2$ *. Therefore, in Proposition [1.3,](#page-2-1) the assumption that* C *is non-degenerate cannot be omitted. From Theorem [1.2,](#page-1-1) it also follows that* C *has at least one conjugacy class with a dimension divisible by* 3*.*

The example above, $\mathcal{C} = \text{Rep}(D(S_3))_{ad}$, also illustrates that, in general, the set of prime divisors of the Frobenius-Perron dimensions of simple objects does not always coincide with the set of prime divisors of the Frobenius-Perron dimensions of the conjugacy classes.

5. Proof of Theorem [1.5](#page-2-0)

We would like to thank A. Moreto for suggesting the proof of the following result:

Proposition 5.1. *For any weakly integral fusion category with a commutative Grothendieck ring* K(C)*, the following inequality holds:*

(5.1)
$$
\text{FPdim}(\mathcal{C}^j) \leq \frac{\text{FPdim}(\mathcal{C})}{\text{FPdim}(\mathcal{C}_{\text{pt}})},
$$

for any conjugacy class \mathcal{C}^j of \mathcal{C} .

Proof. Since $\mathcal{C}_{pt} \subseteq \mathcal{C}$, by [\[Bur20,](#page-19-1) Lemma 7.5, item 1)] there exists a surjective homomorphism of C-algebras:

$$
\pi_e : CE(\mathcal{C}) \to CE(\mathcal{C}_{pt}),
$$

as discussed in Section [3.](#page-7-0)

Let $\mathcal{C}_{\mathrm{pt}} := \mathrm{Vec}_G^{\omega}$ for some abelian group G and some $\omega \in H^3(G, \mathbb{C}^*)$. It follows from Example [2.2,](#page-5-0) we have $CE(\mathcal{C}_{pt}) = \mathbb{C}[G]$. Note that the conjugacy classes of Vec_G^ω correspond to the elements of $G.$

According to Theorem [3.3](#page-10-2) for any conjugacy class \mathcal{C}^j of \mathcal{C} , there exists a group element $g_i \in G$ such that:

$$
\pi_e(C_j) = \dim(\mathcal{C}^j) g_j.
$$

Since $\mathcal C$ is weakly integral, it is a pseudo-unitary fusion category implying that $\dim(\mathcal{C}^j) = \text{FPdim}(\mathcal{C}^j) > 0$ for any $j \in \mathcal{J}$.

Additionally, by Lemma [3.4,](#page-11-3) we have:

(5.2)
$$
\pi_e(\widehat{\Lambda}_{\mathcal{C}}) = \frac{\text{FPdim}(\mathcal{C})}{\text{FPdim}(\mathcal{C}_{\text{pt}})} \widehat{\Lambda}_{\mathcal{C}_{\text{pt}}}.
$$

Since $CE(\mathcal{C}_{pt}) = \mathbb{C}[G]$, it follows that $\widehat{\Lambda}_{\mathcal{C}_{pt}} = \sum_{g \in G} g$. Recall that, according to [\[Bur20,](#page-19-1) Proposition 4.1] (applied for $\mathcal{D} = \mathcal{C}$) we have:

$$
\widehat{\Lambda}_{\mathcal{C}} = \sum_{j \in \mathcal{J}} C_j.
$$

By the same [\[Bur20,](#page-19-1) Proposition 4.1] applied for $\mathcal{D} = \mathcal{C}_{pt}$, we obtain:

$$
\widehat{\Lambda}_{\mathcal{C}_{\mathrm{pt}}} = \sum_{j \in \mathcal{J}_{\mathcal{C}_{\mathrm{pt}}}} C_j.
$$

Let $\mathcal{B}_j := \{l \in \mathcal{J} \mid \pi_e(C_l) = \text{FPdim}(\mathcal{C}^l)g_j\}.$ Counting the multiplicity of g_j in the left hand side of Equation [\(5.2\)](#page-17-1) we find:

$$
\sum_{l \in \mathcal{B}_j} \mathrm{FPdim}(\mathcal{C}^l) = \frac{\mathrm{FPdim}(\mathcal{C})}{\mathrm{FPdim}(\mathcal{C}_{\mathrm{pt}})}.
$$

Since $\text{FPdim}(\mathcal{C}^l) > 0$ for any $l \in \mathcal{J}$, we conclude that:

$$
\mathrm{FPdim}(\mathcal{C}^l) \leq \frac{\mathrm{FPdim}(\mathcal{C})}{\mathrm{FPdim}(\mathcal{C}_{\mathrm{pt}})}.
$$

Remark 5.2. The above proof shows that the possible equality $\dim(C^j)$ = $FPdim(\mathcal{C})$ $\frac{\text{FPdim}(\mathcal{C})}{\text{FPdim}(\mathcal{C}_{pt})}$ holds for a particular conjugacy class \mathcal{C}^j if and only if the set \mathcal{B}_i *contains exactly one element. This indicates that the corresponding character class relative to* \mathcal{C}_{pt} *consists of a single element.*

Recall that C is called a p-fusion category if $\text{FPdim}(\mathcal{C}) = p^n$ for some $n \geq 1$.

Corollary 5.3. *For any* p*-fusion category one has*

$$
\mathrm{FPdim}(\mathcal{C}^j) \mid \frac{\mathrm{FPdim}(\mathcal{C})}{\mathrm{FPdim}(\mathcal{C}_{\mathrm{pt}})}.
$$

Proof. Note that both FPdim (C^j) and $\frac{\text{FPdim}(C)}{\text{FPdim}(C_{pt})}$ are non-negative integers that are powers of p-power. The previous inequality therefore implies the divisibility result.

5.1. Proof of Theorem [1.5.](#page-2-0)

Proof. Note that any nilpotent fusion category is weakly integral because it can be viewed as iterated G-extensions of Vect. Furthermore, any braided nilpotent fusion category can be uniquely decomposed into a Deligne product of p-categories, as established in [\[DGNO10,](#page-20-14) Theorem 1.1]. Consequently, the divisibility property follows from Corollary [5.3.](#page-18-0) Indeed, if $\mathcal{C} \simeq \prod_{p|F}$ Pdim (\mathcal{C}) , \mathcal{C}_p with each \mathcal{C}_p being a p-fusion category then it is straightforward to see that $\mathcal{Z}(\mathcal{C}) \simeq \prod_{p| \text{FPdim}(\mathcal{C})} \mathcal{Z}(\mathcal{C}_p)$. Thus, each conjugacy class of $\mathcal C$ is a Deligne product of a conjugacy class of each \mathcal{C}_p . Moreover, $\mathcal{C}_{pt} \simeq \prod_{p|F \in \text{Pdim}(\mathcal{C})} (\mathcal{C}_p)_{pt}$ and the proof of the first part of the Theorem follows from Corollary [5.3.](#page-18-0)

Since $\mathcal C$ is weakly integral Harada's identity applies and yields:

$$
\widehat{P}^{2m-2} \left(\prod_{j=0}^{m} C_j \right)^2 = \frac{\widehat{P}^{2m} \dim(\mathcal{C}_{\mathrm{pt}})}{\dim(\mathcal{C})} \left(\sum_{j \in \mathcal{J}_{\mathcal{C}_{\mathrm{pt}}}} C_j \right)
$$

where $\widehat{P} := \prod_{j=0}^m \dim(C^j)$. Given that the multiplicity of C_0 on the left hand side is an integer, it follows that $\dim(\mathcal{C})/\dim(\mathcal{C}_{pt})$ divides \widehat{P}^{2m} . Additionally, since each $\dim(\mathcal{C}^j)$ divides this ratio, the proof is \Box complete. \Box

 \Box

5.2. Nilpotent modular categories. Let $\mathcal C$ be a modular category of Frobenius-Perron dimension $\text{FPdim}(\mathcal{C}) = p^n$ for some prime number p. Since FPdim(C_{pt}) divides FPdim(C), it follows that FPdim(C_{pt}) = p^m for some $0 \leq m \leq n$.

Note that [\[CPS24,](#page-19-7) Lemma 2] establishes that in any modular tensor category of dimension p^n one has that

$$
2 \le m \le n - 3
$$

Using Theorem [1.1](#page-1-0) the upper bound can be improved as follows:

Lemma 5.4. *Let* C *be a an integral modular category of Frobenius-Perron dimension* $\text{FPdim}(\mathcal{C}) = p^n$ *for some prime number p. Suppose that* $FPdim(\mathcal{C}_{pt}) = p^m$ *for some* $0 \leq m \leq n$ *. Then either* $2 \leq m \leq n-4$ *or the Frobenius-Perron dimension of the simple objects of* C *are either* 1 *or* p*.*

Proof. Suppose that $m = n - 3$. Note that $\text{FPdim}(\mathcal{C}^j) = d_j^2$ and therefore, since $\text{FPdim}(\mathcal{C}^j) \mid \frac{\text{FPdim}(\mathcal{C})}{\text{FPdim}(\mathcal{C}_{\text{pt}})} = p^{n-m}$ this implies that $d_j^2 \mid p^3$ and therefore $d_i \in \{1, p\}.$

Additionally, consider the class equation:

$$
\mathrm{FPdim}(\mathcal{C}) = \sum_{j \in \mathcal{J}_{\mathcal{C}_{ad}}} \mathrm{FPdim}(\mathcal{C}^j) + \sum_{j \in \mathcal{J} \setminus \mathcal{J}_{\mathcal{C}_{ad}}} \mathrm{FPdim}(\mathcal{C}^j).
$$

Since p^2 | FPdim (C^j) for all $j \notin \mathcal{J}_{C_{ad}}$ it follows that p^2 | $|\mathcal{J}_{C_{ad}}|$ = $|U(\mathcal{C})|$. On the other hand, since C is modular, it follows that $|U(\mathcal{C})|$ = $|FPdim(\mathcal{C}_{pt})|$, and therefore $m \geq 2$.

REFERENCES

- [BP24a] S. Burciu and S. Palcoux. Burnside type results for fusion rings. *arXiv:2302.07604*, 2024.
- [BP24b] S. Burciu and S. Palcoux. Structure constants, Isaacs property and Extended-Haagerup fusion categories. *accepted to Comm. Alg.*, 2024.
- [Bur20] S. Burciu. Conjugacy classes and centralizers for pivotal fusion categories. *Monatshefte für Mathematik*, 193(2):13–46, 2020.
- [Bur21] S. Burciu. Structure constants for premodular categories. *Bull. Lond. Math. Soc.*, 53(3):777–791, 2021.
- [Bur23] S. Burciu. On the Galois symmetries for the character table of an integral fusion category. *J. Algebra Appl.*, 22(1):Paper No. 2350026, 26, 2023.
- [Bur24] S. Burciu. On Harada's identity and some other consequences. *Bull. Belgian Math. Soc. Simon Stevin*, 31(2):211 – 219, 2024.
- [CH90] D. Chillag and M. Herzog. On the length of the conjugacy classes of finite groups. *J. Alg.*, 131(1):110–125, 1990.
- [CPS24] A. Czenky, J. Plavnik, and A. Schopieray. On Frobenius-Schur exponent bounds. *arXiv:2404.06643*, 2024.
- [CW00] M. Cohen and S. Westreich. Conjugacy Classes, Class Sums and Character Tables for Hopf Algebras. *Comm. Algebra*, 39(12):4618–4633, 200.
- [CW14] M. Cohen and S. Westreich. Character tables and normal left coideal subalgebras. *J. Pure and Appl. Alg.*, 218(10):1845–1866, 2014.
- [DGNO07] V. Drinfeld, S. Gelaki, D. Nikshych, and V. Ostrik. Group-theoretical properties of nilpotent modular categories. *arXiv:0704.0195v2*, 2007.
- [DGNO10] V. Drinfeld, S. Gelaki, D. Nikshych, and V. Ostrik. On braided fusion categories, I. *Sel. Math. New Ser.*, 16:1–119, 2010.
- [EGNO15] P. Etingof, S. Gelaki, D. Nikshych, and V. Ostrik. *Tensor categories*, volume 205. Mathematical Surveys and Monographs, American Mathematical Society, Providence, RI, 2015.
- [ENO05] P. Etingof, D. Nikshych, and V. Ostrik. On fusion categories. *Ann. of Math. (2)*, 162(2):581–642, 2005.
- [GN08] S. Gelaki and D. Nikshych. Nilpotent fusion categories. *Adv. in Math.*, 217(3):1053–1071, 2008.
- [Isa76] M. Isaacs. *Character theory of finite Groups*. Academis Press, New York, San Francisco, London, 1976.
- [Lew08] M. L. Lewis. An overview of graphs associated with character degrees and conjugacy class sizes in finite groups. *The Rocky Mountain J. Math.*, 38(1):175–211, 2008.
- [Man87] O. Manz. Arithmetical conditions on character degrees and group structure. *Proc. Sympos. Pure Math.*, 41(2):65–69, 1987.
- [Mic86] G. O. Michler. Brauer's conjectures and the classification of finite simple groups. In Vlastimil Dlab, Peter Gabriel, and Gerhard Michler, editors, *Representation Theory II Groups and Orders*, pages 129–142, Berlin, Heidelberg, 1986. Springer Berlin Heidelberg.
- [Mü03] M. Müger. From subfactors to categories and topology i: Frobenius algebras in and morita equivalence of tensor categories. *Journal of Pure and Applied Algebra*, 180(1):81–157, 2003.
- [Rob96] D. J. S. Robinson. *A Course in the Theory of Groups*. *Lecture Notes in Mathematics*, Springer-Verlag, New York, 1996.
- [Shi17] K. Shimizu. The monoidal center and the character algebra. *J. Pure Appl. Alg.*, 221(9):2338–2371, 2017.
- [Zhu94] Y. Zhu. Hopf algebras of prime dimension. *Internat. Math. Res. Not.*, 1:53–59, 1994.

Inst. of Math. "Simion Stoilow" of the Romanian Academy P.O. Box 1-764, RO-014700, Bucharest, Romania

Email address: sebastian.burciu@imar.ro