Reverse stealth construction and its thermodynamic **IMPRINTS**

Cristián Erices^{∗a,b}, Luis Guajardo^{†c}, and Kristiansen Lara^{‡d}

^aVicerrectoría Académica, Universidad Central de Chile, Toesca 1783, Santiago 8320000, Chile

 b Departamento de Matemática, Física y Estadística, Universidad Católica del Maule, Av. San Miguel 3605, Talca 3480094, Chile

^cInstituto de Investigación Interdisciplinario, Vicerrectoría Académica, Universidad de Talca, 3465548 Talca, Chile. d Centro de Estudios Científicos (CECs), Av. Arturo Prat 514, Valdivia, Chile.

Abstract

We study a class of solutions within the context of modified gravity theories, characterized by a non-trivial field that does not generate any back-reaction on the metric. These stealth configurations are effectively defined by the stealth conditions, which correspond to a vanishing stress-energy tensor. In this work, we introduce a novel approach to constructing this class of solutions. In contrast to the standard procedure, the starting point requires satisfying the stealth conditions for a given ansatz independently of the gravitational dynamics. This approach simultaneously determines the nontrivial field and the geometries capable of supporting it as a stealth configuration. Consequently, a gravity model can accommodate a stealth field only if its vacuum solution falls within the geometries permissible under stealth conditions. By applying this reverse procedure in the non-minimal $R\phi^2$ coupling, we recover all previously known stealth configurations and present new solutions. Although it seems intuitive to assume that this "gravitationally undetectable" scalar field leaves no physical traces, it remarkably reveals thermodynamic imprints, as its presence screens the black hole mass and modifies the entropy according to the first law.

[∗]cristian.erices@ucentral.cl

[†] luis.guajardo.r@gmail.com

[‡]klara@cecs.cl

Contents

1 Introduction

The no-hair conjecture states that black holes cannot be described by any quantity apart from their mass, electric charge, and angular momentum [\[1](#page-25-0)]. This means that in an appropriately chosen reference frame, isolated black holes can be described by the well-known Kerr-Newman solution [\[2](#page-25-1)[–4\]](#page-25-2). According to this conjecture, in the final stage of gravitational collapse, black holes are defined solely by these quantities, which adhere to a Gauss law and are measurable at infinity. This implies that any other information about the matter that falls into a black hole is irretrievably lost. This can be understood from a mathematical perspective; in General Relativity (GR), a set of assumptions renders black hole solutions incompatible with non-trivial matter fields, apart from the electromagnetic one. Specifically, it has been shown that there are no non-trivial regular solutions in GR when minimally coupled scalar fields are considered [\[5\]](#page-25-3). Consequently, under particular conditions that violate these assumptions of the no-hair conjecture, hairy black hole solutions are expected to emerge. One of these assumptions is violated if there is a non-minimal coupling between the scalar field and gravity. This provided one of the earliest examples of a black hole solution with a non-trivial scalar field in an asymptotically flat spacetime more than five decades ago [\[6\]](#page-25-4).

Exploring hairy black holes in scalar-tensor theories has been especially interesting, considering the great interest resurfaced in the community when it was proved that the generalized Galileon theory [\[7,](#page-25-5) [8\]](#page-25-6) is equivalent to the Horndeski theory [\[9\]](#page-25-7). Horndeski theory was proposed fifty years ago as the most general scalar-tensor theory, whose action principle leads to second-order field equations, and consequently, it is free of Ostrogradski instabilities [\[10\]](#page-25-8). Subsequently, the first hairy black hole solution in Horndeski theory [\[11\]](#page-25-9) was found, and a plethora of hairy black hole solutions were investigated in scalar-tensor theories, motivated by no-hair extensions specially adapted to these theories [\[12](#page-25-10)[–14\]](#page-25-11).

However, during the last decade, it was proved that more general scalar-tensor theories can generate higher-order field equations and still avoid Ostrogradski instabilities by adding suitable higher-order modifications. This generalization of Horndeski theory was presented first in Beyond Horndeski and then in Degenerate Higher Order Scalar Tensor (DHOST) theories, providing a rich class of effective field theories [\[15](#page-25-12)[–20\]](#page-26-0).

Some particularly interesting configurations in modified theories of gravity possess a non-trivial field and do not generate any back-reaction, resulting in a metric that coincides with GR solutions. In scalartensor theories, this translates to a vanishing effective stress-energy tensor originating from all the scalartensor terms in the action except for the purely metric sector. These solutions, initially introduced in [\[21](#page-26-1)], are the so-called stealth black holes and represent the simplest example of hairy black holes that are regular at the horizon.

The search for stealth solutions in scalar-tensor theories has a long story. They were introduced almost two decades ago, dubbing this phenomenon a "Gravitational Cheshire effect" [\[22](#page-26-2)]. Over the years, this became a seminal work in the field, showing that it was possible to dress a flat spacetime with a real scalar field that satisfies a nonlinear Klein-Gordon equation without curving spacetime. This configuration is possible due to the non-minimal coupling of the scalar field with the curvature and a self-interacting potential. A footprint of the coupling remains in the stress-energy tensor even when gravity is switched off, and the condition for a vanishing stress-energy tensor fixes the self-interaction potential as a local function of the scalar field depending on two coupling constants. The solutions can describe shock waves and, in the Euclidean continuation, D-dimensional instanton configurations.

Stealth black holes with planar topology were obtained in the bi-scalar extension of Horndeski [\[23\]](#page-26-3). For the shift-symmetric Horndeski theory, it was found that Schwarzschild and a partially self-tuned de-Sitter Schwarzschild black hole are supported by a space and time-dependent scalar field [\[24](#page-26-4)], and a Kerr background can be supported by a static scalar field [\[25\]](#page-26-5). Schwarzschild and Kerr backgrounds with nontrivial scalar field were also found in shift-symmetric breaking Horndeski theories [\[26,](#page-26-6) [27\]](#page-26-7). Stealth scalar fields were described in shift-symmetric scalar-tensor theories of gravity in the family of Degenerate Higher Order Scalar-Tensor (DHOST) theories for Schwarzschild-(A)dS black holes [\[28\]](#page-26-8), rotating backgrounds such as Kerr black holes [\[29\]](#page-26-9) and in cosmological setups [\[30](#page-27-0)[–32\]](#page-27-1). The existence and geometric properties of stealth solutions have been explored in different scenarios other than scalar-tensor theories, such as 3D New Massive Gravity [\[33\]](#page-27-2), hypergravity [\[34\]](#page-27-3), and interacting gauge fields [\[35](#page-27-4)[–37\]](#page-27-5).

To find stealth black holes in those scenarios, one must circumvent the underlying no-hair theorems and additionally ensure that the stress-energy tensor vanishes. In general terms, a scalar-tensor action can be decomposed as

$$
S[\phi, g] = S_g[g] + S_{st}[\phi, g] \,, \tag{1.1}
$$

where S_g is a purely geometric part and consequently depends only on the metric tensor $g_{\mu\nu}$. The last term $S_{\rm st}$ denotes the scalar-tensor part and is the only one that contains the scalar field ϕ .

A nontrivial^{[1](#page-3-0)} scalar field ϕ defines a stealth solution if the following conditions are satisfied

$$
\mathcal{E}_{\mu\nu} = 0 \;, \tag{1.2a}
$$

$$
T_{\mu\nu} = 0. \tag{1.2b}
$$

Here, we denote by

$$
\mathcal{E}_{\mu\nu} = \frac{\delta S_g}{\delta g^{\mu\nu}} \,, \tag{1.3}
$$

to the Euler-Lagrange equations generated by the purely metric sector S_q . It is worth emphasizing that $\mathcal{E}_{\mu\nu}$ is not necessarily the Einstein tensor $G_{\mu\nu}$, as S_g can contain purely geometric terms other than the Einstein-Hilbert one. On the other hand, the stress-energy tensor associated with the scalar-tensor part is

$$
T_{\mu\nu} = -\frac{2}{\sqrt{-g}} \frac{\delta S_{\rm st}}{\delta g^{\mu\nu}} \,. \tag{1.4}
$$

Equations [\(1.2\)](#page-3-1) defy Wheeler's paradigm, which states that "spacetime tells matter how to move; matter tells spacetime how to curve" [\[38\]](#page-27-6). In a broader sense that fits with scalar-tensor theories, the scalar field ϕ is generating no backreaction in the spacetime, yielding an effective equation $\mathcal{E}_{\mu\nu} = 0$, forgoing the sourcing of the stress-energy tensor $T_{\mu\nu}$.

The usual procedure for the construction of stealth solutions is the following: a) For a given spacetime ansatz, one solves the set of field equations [\(1.2a\)](#page-3-2) generated from a specific action S_q , obtaining the corresponding solution $g_{\mu\nu}$. Then, step b) requires solving the stealth conditions [\(1.2b\)](#page-3-3), with $g_{\mu\nu}$ as a background, to obtain a nontrivial scalar field. This last step, of course, depends on $S_{\rm st}$ and how the assumptions of the corresponding no-hair conjecture are circumvented. We dub this procedure as the "a) $\rightarrow b$ " method or standard procedure.

This paradigm has been applied extensively, where the standard non-minimal coupling of the form $R\phi^2$ has proven to be highly successful in constructing stealth solutions. It was useful not only for dressing Minkowski spacetime with stealth solutions but also for constructing a stealth scalar field compatible with the non-rotating Bañados-Teitelboim-Zanelli (BTZ) black hole in $2+1$ dimensions [\[21](#page-26-1)]. The black hole

 $\overline{^{1}$ By nontrivial, we refer to a non-constant scalar field ϕ .

was supported by a time-dependent scalar field and a self-interacting potential. These works represent one of the simplest ways to obtain stealth solutions. For this reason, it was exploited systematically in other theories that include this term, finding stealth scalar fields in the particular case with conformal coupling [\[39](#page-27-7)[–41](#page-27-8)], with the generic non-minimal coupling in Anti-de Sitter (AdS) spacetimes [\[42](#page-27-9)[–44](#page-28-0)], and even in Lifshitz spacetimes [\[45](#page-28-1)]. In higher dimensional scalar-tensor theories, stealth solutions were also found in Einstein-Gauss-Bonnet gravity non-minimally coupled to a scalar field [\[46](#page-28-2)], its Lovelock extension [\[47](#page-28-3)], and recently in more general contexts [\[48](#page-28-4), [49\]](#page-28-5). Besides, stealth configurations over the Riegert black hole were found for conformal gravity with a conformally coupled scalar field in [\[50](#page-28-6), [51](#page-28-7)].

Since the standard non-minimal coupling admits different specific geometries, it seems natural to ask whether those spacetimes fit into a more general family of solutions. We have found that, to answer this question, it is necessary to reverse the aforementioned "a) $\rightarrow b$ " method. In this way, our starting point is b), which means to satisfy the stealth conditions [\(1.2b\)](#page-3-3), without fixing the background $g_{\mu\nu}$ to any particular solution, or equivalently, without specifying any purely geometric sector S_g . Consequently, only the stealth conditions [\(1.2b\)](#page-3-3) on the scalar field constrain the spacetime geometry. The final step is to find purely gravitational theories compatible with the geometry induced by the stealth conditions. We dub this procedure as the " b) $\rightarrow a$ " method, or reverse procedure.

The reverse procedure in constructing stealth solutions brings new implications that are easy to overlook when the standard procedure is applied. This is because different purely geometric actions can admit the same particular solution for their field equations [\(1.2a\)](#page-3-2) and are compatible with the same non-trivial scalar field obtained in [\(1.2b\)](#page-3-3). In this sense, the mapping that associates elements from the set of solutions obtained in step a) to the elements of the set of solutions obtained in step b) is not bijective. This means that solely the stealth conditions determine the family of spacetimes compatible with a non-minimally coupled scalar field. As a consequence, any purely geometric action whose metric solution fits in this family of metrics can be endowed with a stealth scalar field, broadening the landscape of theories that generate non-minimally coupled stealth solutions. Thus, it is possible not only to recover and extend existing solutions in the literature but also to find new solutions by easily identifying the theories that admit stealth configurations.

The generality of the solution obtained in this work puts us in a position to explore its physical consequences. While it is intuitive to think that no physical imprints are left by "gravitationally undetectable" objects, such as stealth configurations, unexpectedly, we found that the mass of stealth black holes is shifted compared to its trivial counterpart. The latter phenomenon can be seen explicitly in a concrete example by selecting, as the purely geometric theory, the five-dimensional Einstein-Gauss-Bonnet gravity.

This article is organized as follows. To address point b , in Section [2,](#page-6-0) we present the reverse approach in $D \geq 4$ dimensions and construct the stealth solution. We analyze its different branches and discuss their implications. Given its particular nature, Section [3](#page-10-1) is devoted to the three-dimensional case. A scheme of the reverse procedure can be seen in Fig. [1.](#page-5-0) Then, we develop step a) in Section [4.](#page-13-0) We apply the reverse procedure to different gravitational theories, as seen in Table [1,](#page-13-2) recovering and generalizing known results from the literature. We also report novel stealth solutions. In Section [5,](#page-17-0) the Euclidean approach is applied to study thermodynamic aspects of stealth black holes in the five-dimensional Einstein-Gauss-Bonnet theory, demonstrating that the stealth scalar field shifts the black hole mass. In Section [6,](#page-21-0) we summarize, discuss, and explore potential avenues for further research.

Figure 1: Guide route

2 Constructing stealth configurations: The reverse procedure

The reverse approach proposes a generic procedure to construct stealth solutions. However, we must ground this approach in a specific scalar-tensor theory. Accordingly, hereafter, we will focus on the non-minimally coupled scalar field in D dimensions,

$$
S_{\rm st}[\phi, g] = -\int d^D x \sqrt{-g} \left(\frac{1}{2} \nabla_\mu \phi \nabla^\mu \phi + \frac{1}{2} \xi R \phi^2 + U(\phi) \right) , \qquad (2.1)
$$

where $\xi \neq 0$ is the coupling constant and $U(\phi)$ is a self-interacting potential. Inspired by works [\[21,](#page-26-1) [22](#page-26-2), [24,](#page-26-4) [33,](#page-27-2) [42,](#page-27-9) [46,](#page-28-2) [47\]](#page-28-3), we will only consider static and homogeneous spacetimes,

$$
ds^{2} = -f(r)dt^{2} + \frac{dr^{2}}{f(r)} + r^{2}d\Sigma_{D-2,\gamma}^{2},
$$
\n(2.2)

where $d\Sigma_{D-2,\gamma}^2$ denotes the line element of a Euclidean, codimension-two base manifold, locally isometric to the sphere \mathbb{S}^{D-2} , flat space \mathbb{R}^{D-2} , or to the hyperbolic manifold \mathbb{H}^{D-2} , with normalized constant curvature $\gamma = \{+1, 0, -1\}$, respectively. Namely, it can be written as,

$$
d\Sigma_{D-2,\gamma}^{2} = \Omega\left(\vec{x}\right)^{2} dx_{i} dx^{i}, \quad \Omega\left(\vec{x}\right) = \frac{1}{1 + \frac{\gamma}{4}x_{i}x^{i}},\tag{2.3}
$$

where x^i are the Euclidean coordinates of the base manifold with $i = \{1, \ldots, D-2\}$. Any contraction between upper and lower Latin indices is understood to be performed using the Kronecker delta δ^{ij} . If both indexes are at the same level, they are not contracted but fixed.

Here, we construct stealth solutions in $D \geq 4$ dimensions without restricting, in principle, the coordinate dependence of the scalar field. In this context, the Euler-Lagrange equations allow us to define the following stress-energy tensor[2](#page-6-1)

$$
T_{\mu\nu} = \nabla_{\mu}\phi\nabla_{\nu}\phi - g_{\mu\nu}\left(\frac{1}{2}\nabla_{\sigma}\phi\nabla^{\sigma}\phi + U(\phi)\right) + \xi\left(g_{\mu\nu}\Box - \nabla_{\mu}\nabla_{\nu} + G_{\mu\nu}\right)\phi^2.
$$
 (2.4)

We aim to solve $T_{\mu\nu} = 0$ in the ansatz [\(2.2\)](#page-6-2). The potential $U(\phi)$ required to sustain the stealth solution is sensitive to the curvature γ , so it is kept undetermined at the moment.

We begin analyzing the off-diagonal components of the stress-energy tensor. For fixed, but distinct indices μ and ν , it reads

$$
T^{\nu}_{\mu} = \nabla_{\mu}\phi\nabla^{\nu}\phi - \xi\nabla_{\mu}\nabla^{\nu}\phi^{2}, \quad \mu \neq \nu.
$$
 (2.5)

The above can be rewritten in appealing form, according to the following change of variables

$$
\phi = \psi^{\frac{2\xi}{4\xi - 1}},\tag{2.6}
$$

with $\xi \neq 1/4$, obtaining

$$
T^{\nu}_{\mu} = -\frac{4\xi^2}{(4\xi - 1)} \psi^{\frac{1}{4\xi - 1}} \nabla_{\mu} \nabla^{\nu} \psi = 0 \,, \quad \mu \neq \nu \,. \tag{2.7}
$$

²The scalar field equation is satisfied by virtue of the Bianchi identity.

Regarding case $\xi = 1/4$, it is treated in Appendix [A.](#page-22-0) Therefore, the off-diagonal equations [\(2.7\)](#page-6-3) are akin to the sum-separability condition. Because of this, we can obtain a simple generic profile for the scalar field. In terms of the auxiliary function ψ , it reads

$$
\psi(t, r, x_1, \dots, x_{D-2}) = T(t)\sqrt{f(r)} + R(r) + r\Omega(\vec{x})\sum_{i=1}^{D-2} X_i(x_i),
$$
\n(2.8)

where $T(t)$, $R(r)$ and $X_i(x_i)$ are functions that depends only on t, r and the base manifold coordinates x_i , respectively.

Using this expression, the next step is to determine the form of the unknown functions $T(t)$, $R(r)$, and $X_i(x_i)$. Taking the difference $T_i^i - T_j^j = 0$ for fixed but distinct spatial coordinates $i, j = \{1, ..., D-2\}$ (the components of the difference shall not be confused with the trace), we arrive to

$$
T_i^i - T_j^j \propto \frac{\mathrm{d}^2 X_i}{\mathrm{d} x_i^2} - \frac{\mathrm{d}^2 X_j}{\mathrm{d} x_j^2} = 0 \,, \quad i, j = \{1, \dots, D - 2\} \,. \tag{2.9}
$$

The solution is given by

$$
X_i(x_i) = ax_i^2 + b_i x_i + c_i, \quad i = \{1, 2, ..., D-2\},
$$
\n(2.10)

where a, b_i and c_i are integration constants. Thus far, we have $T_1^1 = T_2^2 = \cdots = T_{D-2}^{D-2}$.

The difference $T_t^t - T_r^r$ yields

$$
T_t^t - T_r^r = -\frac{\psi^{\frac{1}{4\xi - 1}}\xi^2}{\sqrt{f}(4\xi - 1)} \left(-4f^{3/2}R'' - 4\ddot{T} - 2Tff'' + Tf'^2 \right) = 0. \tag{2.11}
$$

In the above expression, primes and dots denote radial and time derivatives, respectively.

It is necessary to distinguish in (2.11) that the function $T(t)$ can be constant, generating an additional branch of solutions. These can be seen explicitly by differentiating the expression inside the brackets with respect to the time and the radius, leading to the compatibility condition

$$
2ff'''\dot{T} = 0.\tag{2.12}
$$

The latter admits two nontrivial branches with a time (in)dependent scalar field. After removing redundancies, we get,

$$
R(r) = R_1 r + R_0, \quad \begin{cases} T(t) = 0 & \to \text{Time-independent scalar field} \\ T(t) \neq \text{cst} & \to \text{Time-dependent scalar field} \end{cases}
$$
 (2.13)

At this stage of the procedure, the stress-energy tensor is diagonal, with $T_t^t = T_r^r$, and $T_1^1 = T_2^2 = \cdots =$ T_{D-2}^{D-2} . The next subsection [2.1](#page-7-0) is devoted to developing the time-independent branch of solutions. We must mention that the time-dependent branch in Subsection [2.2](#page-10-0) has some overlap with an unpublished work[3](#page-7-2) .

2.1 Time-independent configuration

The construction of time-independent stealth solutions will require distinguishing between those with a scalar field depending on the spatial coordinates and those with purely radial dependence. To see this, we

 $3A$ yón-Beato, Martínez, Troncoso and Zanelli, private communication.

focus our attention on the difference $T_t^t - T_j^j$ (do not confuse with the trace), which can be schematically written in the form

$$
T_t^t - T_j^j = \frac{\Omega(\vec{x})}{8r^2\sqrt{f}(4\xi - 1)\psi^{\frac{8\xi - 1}{4\xi - 1}}} \left(\Xi_2 x_i x^i + \Xi_1 b_i x^i + \Xi_0 \right) = 0, \qquad (2.14)
$$

where $\Xi_{2,1,0}$ are differential equations detailed in Eq. [\(B.4\)](#page-24-0) of Appendix [B.](#page-23-0) At this point, we see that the time-independent branch provides two sub-branches of solutions:

 $b_i \neq 0$, $\phi = \phi(r, \vec{x})$, \rightarrow Spatial coordinates-dependent scalar field, (2.15)

$$
b_i = 0, \qquad \phi = \phi(r), \qquad \rightarrow \text{Radial-dependent scalar field.} \tag{2.16}
$$

It is worth emphasizing that since the reverse approach does not impose any prior restrictions on the coordinate dependence of the scalar field, the spatial coordinates-dependent scalar field has been overlooked in the literature. This aspect gives rise to novel classes of stealth solutions.

2.1.1 Spatial-coordinates dependent scalar field

The condition $b_i \neq 0$ leads to solve the set of differential equations $\Xi_0 = 0$, $\Xi_1 = 0$ and $\Xi_2 = 0$. We start with $\Xi_1 = 0$, which is a differential equation for the metric function $f(r)$, whose solution is

$$
f(r) = Ar^2 + Br^{\frac{(D-3)-4(D-2)\xi}{4\xi-1}} + \gamma,
$$
\n(2.17)

where A and B are integration constants. Replacing in $\Xi_0 = 0$ and $\Xi_2 = 0$, the following conditions are obtained,

$$
BR_0 = 0, \t\t(2.18)
$$

$$
\gamma \left(2R_1 + \sum_{i=1}^{D-2} c_i \right) + 4a = 0, \qquad (2.19)
$$

where a and c_i are the integration constants of [\(2.10\)](#page-7-3). Note that equation [\(2.18\)](#page-8-1) prevents having nonvanishing constants B and R_0 simultaneously. On the other hand, equation [\(2.19\)](#page-8-2) allows us to write a in terms of γ , R_1 , and c_i . Then, the auxiliary scalar field ψ obtained in [\(2.8\)](#page-7-4) lead to the following profile for the scalar field,

$$
\phi(r,\vec{x}) = \left\{ R_0 + r \left[\left(2\Omega\left(\vec{x}\right) - 1 \right) \left(R_1 + \sum_{i=1}^{D-2} c_i \right) + \Omega\left(\vec{x}\right) b_i x^i \right] \right\}^{\frac{2\xi}{4\xi - 1}}.
$$
\n(2.20)

At this stage, all the diagonal components of the stress-energy tensor are equal. Finally, the stealth conditions $T_{\mu\nu} = 0$ are satisfied by the following self-interacting potential,

$$
U(\phi) = \lambda_1 \phi^2 + \lambda_2 \phi^{-\frac{2\xi - 1}{\xi}} + \lambda_3 \phi^{\frac{1}{2\xi}}, \qquad (2.21)
$$

with

$$
\lambda_1 = \frac{\xi AD(D-1)(\xi - \xi_D)(\xi - \xi_{D+1})}{2(\xi - 1/4)^2},
$$
\n(2.22a)

$$
\lambda_2 = \frac{\xi^2}{8(\xi - 1/4)^2} \left[AR_0^2 + \gamma \left(R_1 + \sum_{i=1}^{D-2} c_i \right)^2 + \sum_{i=1}^{D-2} b_i^2 \right],
$$
\n(2.22b)

$$
\lambda_3 = -\frac{AR_0 \xi^2 (D-1)(\xi - \xi_D)}{(\xi - 1/4)^2} \,. \tag{2.22c}
$$

In general, since the supporting potential determines the theory given in the action principle, its constants must be considered as parameters of the theory. Therefore, they are fixed without variation, implying a relation between the variations of the integration constants. In particular, we see that [\(2.22\)](#page-8-3) implies $\delta\lambda_1 = \delta\lambda_2 = \delta\lambda_3 = 0$. This feature is relevant for the thermodynamic aspects discussed in Section [5.](#page-17-0) As it is expected, the potential in [\(2.21\)](#page-8-4), restore conformal invariance when $\xi = \xi_D = \frac{D-2}{4(D-1)}$.

So far, a spatial-coordinate dependent scalar field dictates a family of admissible geometries given by the metric function [\(2.17\)](#page-8-5). The additional restriction $BR_0 = 0$ [\(2.18\)](#page-8-1) preserves this structure if $R_0 = 0$ and $B \neq 0$. Otherwise, if $R_0 \neq 0$ and $B = 0$, the geometry is isometric to $(A)dS_D$, meaning that any purely gravitational theory whose background solution is maximally symmetric admits a non-minimally coupled stealth scalar field.

2.1.2 Radial-dependent scalar field

In contrast to the previous subsection, the condition $b_i = 0$ implies solving the set of differential equations given by $\Xi_0 = 0$ and $\Xi_2 = 0$. A suitable combination of such equations gives,

$$
\gamma \sum_{i=1}^{D-2} c_i - 4a = 0.
$$
\n(2.23)

This implies, after absorbing redundant constants, a radial-dependent scalar field, whose form is given by

$$
\phi(r) = (R_1r + R_0)^{\frac{2\xi}{4\xi - 1}}.
$$
\n(2.24)

The aforementioned set of differential equations is solved by lengthy expressions for the metric function that depends on Gauss' hypergeometric and Gamma functions. However, familiar solutions are found in the conformal case where $\xi = \xi_D = \frac{D-2}{4(D-1)}$ $\xi = \xi_D = \frac{D-2}{4(D-1)}$ $\xi = \xi_D = \frac{D-2}{4(D-1)}$. Namely ⁴,

$$
f(r) = Ar^2 + 2\gamma \frac{R_1}{R_0} r + \gamma + \frac{B(R_1 r + R_0)^{D-1}}{r^{D-3}}, \quad R_0 \neq 0,
$$
\n(2.25)

from where we find the conformally-invariant potential

$$
U(\phi) = \lambda \phi^{\frac{2D}{D-2}}, \quad \text{with} \quad \lambda = AR_0^2 - \gamma R_1^2. \tag{2.26}
$$

In the conformal case with $R_0 = 0$, the scalar field gets rigid, i.e., its integration constant R_1 is fixed without variation as the potential requires it to be a parameter of the theory. Nevertheless, the stressenergy tensor vanishes for a geometry ruled by $f(r) = Ar^2 + Br - 2\gamma$, and a conformally invariant potential.

With the reverse approach, we can confirm that in dimension $D \geq 4$, the time-independent stealths compatible with General Relativity, with or without cosmological constant, are the stealth solutions in Minkowski [\[22\]](#page-26-2) and $(A)dS_D$ spacetimes¹. Therefore, in this set-up, finding black hole geometries for a given dress necessarily requires going beyond General Relativity (see Section [4](#page-13-0) for examples).

⁴Conformal couplings of different dimensions generate logarithmic contributions in the metric function. We opt not to elaborate further on those cases.

2.2 Time-dependent configuration

Now, we will turn to some comments about the time-dependent branch. From the compatibility condition (2.12) , one can deduce that the metric function is at most a quadratic polynomial in r,

$$
f(r) = Ar^2 + Br + C.
$$
 (2.27)

Plugging the metric function in equations $T_t^t - T_r^r = 0$ and $T_t^t - T_i^i = 0$, determines $C = \gamma$, and the following scalar profile,

$$
\phi(t,r,\vec{x}) = \left\{ R_0 + T(t)\sqrt{f(r)} + R_1r + r\Omega(\vec{x})\sum_{i=1}^{D-2} X_i(x_i) \right\}^{\frac{2\xi}{4\xi - 1}}, \qquad (2.28)
$$

where the nature of this time-dependent configuration is provided by $T(t)$ which is given by,

$$
T(t) = T_1 \cos\left(\frac{\sqrt{-\Delta}}{2}t\right) + \frac{T_0}{\sqrt{-\Delta}}\sin\left(\frac{\sqrt{-\Delta}}{2}t\right), \quad \Delta = B^2 - 4A\gamma. \tag{2.29}
$$

A real scalar field requires a positive leading term in the asymptotic behavior of the metric function $f(r)$. This implies, in particular, that asymptotically AdS metrics can be supported by real stealth configurations, unlike their dS counterparts. The solution for Minkowski stealths from Ref. [\[22](#page-26-2)] is recovered when $f(r)$ is a constant $(A = B = 0$ and $\gamma = 1)$.

The functional form of the self-interacting potential is the same as in the time-dependent configuration $(2.21),$ $(2.21),$

$$
U(\phi) = \lambda_1 \phi^2 + \lambda_2 \phi^{-\frac{2\xi - 1}{\xi}} + \lambda_3 \phi^{\frac{1}{2\xi}}\,,\tag{2.30}
$$

except that the parameters $\lambda_k = \lambda_k (A, R_0, R_1, b_j, c_j, T_0, T_1)$ with $j = \{1, \ldots, D-2\}, k = \{1, 2, 3\}$ obeys new relations with the integration constants. However, these new relations do not give new insights about the solution. It is worth noticing that the case $B = 0$ forces $R_1 = 0$. Otherwise, ξ must be fixed to the conformal value ξ_D , and $R_1 = BR_0/2$.

3 Three-dimensional case

The reverse approach used in the previous section is slightly different in $D = 3$, with $\gamma = 0$, mainly because we lack angular components to perform non-trivial subtractions in the spatial sector of T^{μ}_{ν} . This situation will be reflected in the metric function, whose additive constant might differ from the spatial curvature γ . Apart from this fact, three cases arise depending on the coordinate dependencies for the scalar field, as in the previous section. Again, we focus on homogeneous configurations,

$$
ds2 = -f(r)dt2 + \frac{dr2}{f(r)} + r2 d\theta2
$$

Radial-dependent scalar field: In the purely-radial case, we readily check that

$$
\phi(r) = (R_1r + R_0)^{\frac{2\xi}{4\xi - 1}},\tag{3.1}
$$

.

while the metric function $f(r)$ is generically written in terms of a hypergeometric function. However, as in the $D \geq 4$ case, some particular choices relax the latter behavior. For example, choosing $R_0 = 0$ is consistent with taking $D = 3$ and $\gamma = 0$ in Eq.[\(2.17\)](#page-8-5). The interesting choice is the conformal case with $\xi = 1/8$, where the metric function is a quadratic polynomial,

$$
f(r) = Ar^2 + Br + C,
$$
\n
$$
(3.2)
$$

and the constants B and C are tied with the constants R_0 and R_1 coming from the scalar field via

$$
BR_0 = 2CR_1. \tag{3.3}
$$

For this particular choice, the potential also preserves the conformal invariance,

$$
U(\phi) = \frac{1}{8}\lambda\phi^6, \text{ with } \lambda = AR_0^2 - CR_1^2. \tag{3.4}
$$

Spatial-coordinates dependent scalar field: Now, we deal with the case $\phi = \phi(r, \theta)$. Here, for $\xi \neq 1/8$ we find

$$
\phi(r,\theta) = \left\{ R_0 + r \left[\chi_0 \cos\left(\sqrt{C}\theta\right) + \frac{\chi_1}{\sqrt{C}} \sin\left(\sqrt{C}\theta\right) \right] \right\}^{\frac{2\xi}{4\xi - 1}},\tag{3.5}
$$

while $f(r)$ reads

$$
f(r) = Ar^2 + Br^{-\frac{4\xi}{4\xi - 1}} + C.
$$
\n(3.6)

Similarly to the $D \geq 4$ case, the functional form of the metric function is preserved if $R_0 = 0$. Otherwise, we are restricted to demand $B = 0$. In any case, the potential reads

$$
U(\phi) = \frac{A\xi \left(8\xi - 1\right) \left(6\xi - 1\right)}{\left(4\xi - 1\right)^2} \phi^2 + \frac{2\xi^2 \left[AR_0^2 + C\chi_0^2 + \chi_1^2\right]}{\left(4\xi - 1\right)^2} \phi^{-\frac{2\xi - 1}{\xi}} - \frac{4A\xi^2 R_0 \left(8\xi - 1\right)}{\left(4\xi - 1\right)^2} \phi^{\frac{1}{2\xi}}\,. \tag{3.7}
$$

On the other hand, the conformal case with $\xi = 1/8$ restores the quadratic polynomial [\(3.2\)](#page-11-0) and the relation [\(3.3\)](#page-11-1) between the constants B, C, R_1 and R_0 , so that the scalar field reads

$$
\phi(r,\theta) = \left\{ R_0 + r \left[R_1 + \chi_0 \cos\left(\sqrt{C}\theta\right) + \frac{\chi_1}{\sqrt{C}} \sin\left(\sqrt{C}\theta\right) \right] \right\}^{-\frac{1}{2}},\tag{3.8}
$$

and

$$
U(\phi) = \frac{1}{8} \left[AR_0^2 - CR_1^2 + C\chi_0^2 + \chi_1^2 \right] \phi^6.
$$
 (3.9)

Time-dependent configuration: Finally, some comments regarding the time-dependent branch are in order. Similarly to the $D \geq 4$ case, one easily checks that a time-dependent scalar field is compatible with

$$
f(r) = Ar^2 + Br + C \t\t(3.10)
$$

with a scalar field given by,

$$
\phi(t,r,\theta) = \left(T(t)\sqrt{f(r)} + R_1r + R_0 + rX_1(\theta)\right)^{\frac{2\xi}{4\xi - 1}},
$$
\n(3.11)

where,

$$
T(t) = T_1 \cos\left(\frac{\sqrt{-\Delta}}{2}t\right) + \frac{T_0}{\sqrt{-\Delta}}\sin\left(\frac{\sqrt{-\Delta}}{2}t\right), \quad \Delta = B^2 - 4AC. \tag{3.12}
$$

The function $X_1(\theta)$ fulfills the following differential equation

$$
\frac{\mathrm{d}^2 X_1}{\mathrm{d}\theta^2} + CX_1 + CR_1 - \frac{1}{2} BR_0 = 0. \tag{3.13}
$$

Interestingly, in contrast with the $D \ge 4$ case, here the equation $T_t^t - T_{\theta}^{\theta} = 0$ forces us to choose $B = 0$ unless $\xi = 1/8$, without any restriction on R_0 . For later purposes, we focus on the $C \neq 0$ case, where the scalar field reads

$$
\phi(t,r,\theta) = \left\{ R_0 + \left[T_1 \cos\left(\frac{\sqrt{-\Delta}}{2}t\right) + \frac{T_0}{\sqrt{-\Delta}} \sin\left(\frac{\sqrt{-\Delta}}{2}t\right) \right] \sqrt{Ar^2 + Br + C} + r \left[\frac{BR_0}{2C} + \chi_1 \cos\left(\sqrt{C}\theta\right) + \frac{\chi_0}{\sqrt{C}} \sin\left(\sqrt{C}\theta\right) \right] \right\}^{\frac{2\xi}{4\xi - 1}}.
$$
\n(3.14)

Note that a real scalar field requires $A > 0$ as it is the coefficient of the leading term in the asymptotic region. The self-interacting potential $U(\phi)$ is given by

$$
U(\phi) = \lambda_1 \phi^2 + \lambda_2 \phi^{-\frac{2\xi - 1}{\xi}} + \lambda_3 \phi^{\frac{1}{2\xi}},
$$
\n(3.15)

where

$$
\lambda_1 = \frac{A\xi \left(8\xi - 1\right)\left(6\xi - 1\right)}{\left(4\xi - 1\right)^2},\tag{3.16a}
$$

$$
\lambda_2 = \frac{2\xi^2}{(4\xi - 1)^2} \left[-\frac{1}{4}T_0^2 + \frac{\Delta}{4}T_1^2 + \chi_0^2 + C\chi_1^2 - \frac{\Delta R_0^2}{4C} \right],
$$
\n(3.16b)

$$
\lambda_3 = -\frac{4A\xi^2 R_0 (8\xi - 1)}{(4\xi - 1)^2}.
$$
\n(3.16c)

Summarizing the previous sections, using this reverse approach we have solved the stealth condition $T_{\mu\nu} = 0$ for $D \geq 3$, without any reference to the gravitational sector. In all the cases presented above, the stealth condition fixes the scalar field and the geometry, so the matter content codified in the stress-energy tensor dictates the form of the admissible geometries. In this sense, the scalar field does have an impact on the geometry, even when $T_{\mu\nu} = 0$ is commonly interpreted as "no backreaction". Indeed, recognizing that the scalar field does not need to share the symmetries of the geometry significantly broadens the spectrum of non-trivial scalar profiles. In this respect, three different geometries are determined by a radial-dependent, spatial-coordinates-dependent, and time-dependent scalar field. In the latter case, a quadratic polynomial in the metric function rules the admissible geometries. Taking pertinent limits, this family includes familiar solutions such as the BTZ black hole or the $(A)dS_D$ spacetime, as we will explain in detail in the next section. On the other hand, a time-independent, axisymmetry-breaking scalar field can incorporate an extra term proportional to $r^{\frac{(D-3)-4(D-2)\xi}{4\xi-1}}$, which is different from the usual $1/r^{(D-3)}$ decay in the metric function from GR. In this regard, it is worth emphasizing that within the non-minimal coupling $R\phi^2$, the Schwarzschild and Schwarzschild-AdS black holes are only recovered with a constant scalar field, precluding the existence of a stealth scalar field for such backgrounds.

We stress that, as mentioned in the introduction, we only have addressed point b). In the next section, we move to point a) and complete the construction of a stealth solution, revisiting known results and presenting novel solutions.

4 Applying the reverse procedure: Concrete examples

In this section, we show concrete examples where the reverse approach is used. Following the path from b) to a), we now delve into the search for purely gravitational theories that are compatible with the nongravitating scalar field. This problem is simplified to find gravity theories whose vacuum solutions fit in any of the cases presented in the previous sections, matching the constants from the gravity part with the ones obtained in the scalar-tensor part through this novel procedure. Through this study, we recover and generalize known cases, as well as finding novel stealth configurations. Even though we apply the reverse procedure to specific metric theories, we emphasize that the geometries determined by the stealth conditions presented in the previous section are, in principle, applicable to any other purely metric theory whose vacuum solution matches such geometries. A concrete example of these ideas is the stealth solution in AdS background (c.f. [\(4.5\)](#page-14-0)). Any theory that contains AdS as a vacuum solution is susceptible to being non-minimally coupled to a stealth scalar field.

We begin by conducting a literature review of all cases where the scalar-tensor sector $S_{st}[\phi, g]$ is given by [\(2.1\)](#page-6-4), in the background [\(2.2\)](#page-6-2). These works are presented in Table [1](#page-13-2) for $S_g[g] = \int dx^D \sqrt{-g} \mathcal{L}_g$, given by the Lagrangian density \mathcal{L}_q , in General Relativity with and without cosmological constant [\[21,](#page-26-1) [22\]](#page-26-2), New Massive Gravity (with graviton mass m) [\[33\]](#page-27-2), Einstein–Gauss–Bonnet [\[46\]](#page-28-2), Lovelock [\[47\]](#page-28-3) and Conformal Gravity [\[50](#page-28-6), [51](#page-28-7)].

| \mathcal{L}_g | Set-up | Coupling value of ξ | Reference |
|---|-------------------------------|--|--------------------|
| $R-2\Lambda$ | $D=3$ | Arbitrary | $\left[21\right]$ |
| R | $D \geq 3$, and $\gamma = 1$ | Arbitrary | [22] |
| $R-2\Lambda-\frac{1}{m^2}\left(R_{\mu\nu}R^{\mu\nu}-\frac{3}{8}R^2\right)$ | $D=3$ | 1/8 | $\left[33\right]$ |
| $\frac{1}{2}(R-2\Lambda+\alpha \mathcal{G})$ | $D=5$, and $\gamma=0$ | 1/6 | [46] |
| $\sum_{i} \alpha_{p} \epsilon_{a_{1} \cdots a_{D}} R^{a_{1} a_{2}} \cdots R^{a_{2p-1} a_{2p}} e^{a_{2p+1}} \cdots e^{a_{D}}$ $p=0$ | $D \geq 5$, and $\gamma = 0$ | $(n-1)(D-1)$ $\overline{4(nD}-D+1)$ | $[47]$ |
| $W^{\alpha\beta}_{\mu\nu}W^{\mu\nu}_{\alpha\beta}$ | $D=4$, for any γ | 1/6 | [50, 51] |

Table 1: List of works recovered by the reverse approach. Here, m is the graviton mass, $n = [D/2]$, and α and α_p are the coupling constant of the Gauss-Bonnet and Lovelock densities, respectively

4.1 Stealth solutions in General Relativity

Stealth over BTZ: In three dimensions, General Relativity with negative cosmological constant admits the BTZ black hole [\[52\]](#page-28-8). The stealth overflying this black hole was first introduced in [\[21\]](#page-26-1), and here we revisit the solution from our approach, as a particular case of the time-dependent branch analyzed in Section [3.](#page-10-1) Concretely, the authors considered time and radial scalar configurations over the following metric

$$
ds^{2} = -F(r) dt^{2} + \frac{dr^{2}}{F(r)} + r^{2} \left(d\phi - \frac{J}{2r^{2}} dt \right)^{2},
$$
\n(4.1)

with $F(r) = \frac{r^2}{r^2}$ $\frac{r^2}{\ell^2} - M + \frac{J^2}{4r^2}$ $\frac{a}{4r^2}$. In their analysis, which corresponds to the standard $a) \rightarrow b$) approach under our notation, the equations to solve the stealth condition $T_{\mu\nu} = 0$ force $J = 0$, rendering the metric to be static, with a scalar field obeying the following profile

$$
\phi(t,r) = \left[T(t)\sqrt{F(r)} + h(t) \right]^{\frac{2\xi}{4\xi - 1}}.
$$
\n(4.2)

In the static case, the BTZ metric simply reads $F(r) = \frac{r^2}{r^2}$ $\frac{V}{\ell^2} - M$. From our procedure, including timedependent scalar fields restricts the metric function to be a quadratic polynomial, namely,

$$
\phi = \phi(t, r), \quad f(r) = Ar^2 + Br + C. \tag{4.3}
$$

Hence, we can link the standard and reverse procedures just by fixing,

$$
A = \frac{1}{\ell^2}, \quad B = 0, \quad C = -M.
$$
\n(4.4)

At this point, we explicitly see that our constructive approach is not completed until the gravity theory is included, and gluing both parts is substantial to realize the physical relevance of the parameters or integration constants we have found. In this case, the outcome was that A is a parameter related to the cosmological constant, while −C represents the mass of the black hole. We stress that the scalar field itself cannot provide interpretations regarding the parameters, and there is no guarantee that the ones obtained hold arbitrarily.

We can recover the stealth scalar field from [\[21\]](#page-26-1) as a particular case from the most general structure of the scalar fields presented in Eq. (3.14) , as well as the self-interacting potential $(3.15)-(3.16a)$ $(3.15)-(3.16a)$, just by applying the gluing conditions [\(4.4\)](#page-14-1) and additionally $T_0 = \chi_0 = \chi_1 = 0$.

Stealth overflying Anti-de Sitter: In GR, maximally symmetric spacetimes are uniquely characterized by the cosmological constant Λ . For negative values of Λ , the solution of the Einstein equations with the maximal number of symmetries is Anti-de Sitter (AdS_D) , and it is described by

$$
ds^{2} = -\left(\frac{r^{2}}{\ell^{2}} + 1\right)dt^{2} + \left(\frac{r^{2}}{\ell^{2}} + 1\right)^{-1}dr^{2} + r^{2}d\Sigma_{D-2,1}^{2}, \qquad (4.5)
$$

with $\Lambda = -\frac{(D-1)(D-2)}{2l^2}$ $\frac{2\ell^2}{2\ell^2}$. The reverse approach reduces the difficulty of finding stealth solutions by linking the geometries. This is why it is easy to obtain time-dependent stealth scalar fields over AdS_D for any dimension $D \geq 3$. As remarked in the previous section, time-dependent scalar fields constrain the metric function to be a quadratic polynomial. For $D=3$, the stealth solution is of the form,

$$
\phi = \phi(t, r, \theta), \quad f(r) = Ar^2 + Br + C. \tag{4.6}
$$

Clearly, we can link our procedure to the $AdS₃$ spacetime by fixing

$$
A = \frac{1}{\ell^2}, \quad B = 0, \quad C = 1.
$$
 (4.7)

Applying these conditions to [\(3.14\)](#page-12-0), a novel solution is obtained. This is,

$$
\phi(t,r,\theta) = \left[R_0 + T_1 \cos\left(\frac{t}{\ell}\right) \sqrt{\frac{r^2}{\ell^2} + 1} + \chi_1 \cos(\theta) \right]^{\frac{2\xi}{4\xi - 1}}, \tag{4.8}
$$

after absorbing integration constants T_0 and χ_0 due to the isometries of the metric, while the potential is obtained from [\(3.15\)](#page-12-1)-[\(3.16a\)](#page-12-2). The stealth solution over $(A)dS_D$ can be obtained in $D \geq 4$ just by considering [\(4.7\)](#page-14-2) in the solutions provided in Section [2.](#page-6-0) Finally, our procedure also recovers the Minkowski stealth from Ref. [\[22\]](#page-26-2), by considering $A = B = 0$ and $C = 1$ in the time-dependent branch [\(2.13\)](#page-7-6).

It is noteworthy that the reverse approach allows coupling a stealth scalar field in a non-minimal manner to any metric theory that admits a maximally symmetric spacetime as a solution.

4.2 Stealth solutions beyond General Relativity

New Massive Gravity: A stealth solution from Ref. [\[33\]](#page-27-2) was found in New Massive Gravity, which is also a sensitive metric theory that can be coupled to a stealth scalar field. This means that we consider the following action principle,

$$
S[g,\phi] = \int d^3x \sqrt{-g} \left[R - 2\Lambda - \frac{1}{m^2} \left(R_{\mu\nu} R^{\mu\nu} - \frac{3}{8} R^2 \right) - \left(\frac{1}{2} \nabla_{\mu} \phi \nabla^{\mu} \phi + \frac{1}{2} \xi R \phi^2 + U(\phi) \right) \right] \ . \tag{4.9}
$$

Then, the corresponding stealth solution belongs to the radial dependent configuration in $D = 3$ presented in Section [3.](#page-10-1) In the conformal case, $\xi = 1/8$, we have

$$
\phi = \phi(r), \quad f(r) = Ar^2 + Br + C. \tag{4.10}
$$

Hence, the stealth solution from the above reference can be successfully retrieved by fixing

$$
A = \frac{1}{\ell^2}, \quad B = b, \quad C = -4GM,
$$
\n(4.11)

and the conformal potential

$$
U(\phi) = \lambda \phi^6. \tag{4.12}
$$

From the reverse approach, when the scalar field depends on time, the metric function also behaves as a quadratic polynomial. This simple observation and the linear decay in the vacuum solution of New Massive Gravity make it possible to present an extension of the stealth from Ref. [\[33](#page-27-2)]. Concretely, using

$$
\phi = \phi(t, r), \quad f(r) = Ar^2 + Br + C,
$$
\n(4.13)

and the same conditions [\(4.11\)](#page-15-1), the novel time-dependent scalar field reads

$$
\phi(t,r) = \left[T_1 \cosh\left(\frac{\Delta}{2}t\right) \sqrt{\frac{r^2}{\ell^2} + br - 4GM} + R_0 - \frac{bR_0}{8GM}r \right]^{-\frac{1}{2}},\tag{4.14}
$$

where $\Delta = b^2 + \frac{16GM}{\ell^2}$, and we have already absorbed T_0 by time translations invariance. This timedependent configuration demands the relation for the parameter of the potential

$$
\lambda = \frac{\Delta}{4}T_1^2 - \frac{\Delta R_0^2}{4C},\tag{4.15}
$$

which is fixed without variation. When $T_1 \rightarrow 0$, we recover the purely-radial stealth from Ref. [\[33](#page-27-2)] previously mentioned.

Gauss-Bonnet and Lovelock: Lovelock gravity [\[53](#page-28-9)] is the most general D−dimensional gravity model with second order equations. It coincides with GR in three and four dimensions. The n−th member of the Lovelock family is a topological invariant in $D = 2n$, so they modify GR in $D \geq 2n + 1$ dimensions. To apply the reverse procedure to this theory, we coupled the scalar field in the following form,

$$
S[\phi, g] = \int d^D x \sqrt{-g} \sum_{p=0}^n \alpha_p \delta^{\beta_1 \cdots \beta_{2p}} \gamma_1 \cdots \gamma_{2p} R^{\gamma_1 \gamma_2} \beta_1 \beta_2 \cdots R^{\gamma_{2p-1} \gamma_{2p}} \beta_{2p} - \beta_{2p}
$$

-
$$
\int d^D x \sqrt{-g} \left(\frac{1}{2} \nabla_\mu \phi \nabla^\mu \phi + \frac{1}{2} \xi R \phi^2 + U(\phi) \right) , \quad 1 \le n \le \left[\frac{D-1}{2} \right] .
$$
 (4.16)

In Lovelock gravities, maximally symmetric solutions demand some relations between the coupling constants α_p of the curvature terms. Under the hypothesis that the theory admits a unique AdS-vacua, black hole solutions have been found in Ref. [\[54](#page-28-10)], and the metric function is given by

$$
f(r) = \frac{r^2}{\ell^2} + Br^{-\frac{D-2n-1}{n}} + \gamma.
$$
\n(4.17)

The shape of the metric function allows us to enhance these black hole solutions with a scalar field depending on the transverse coordinates, as shown in Section [2.1.1.](#page-8-0) This is,

$$
\phi = \phi(r, \vec{x}), \quad f(r) = Ar^2 + Br^{\frac{(D-3)-4(D-2)\xi}{4\xi-1}} + \gamma,
$$
\n(4.18)

recovers the metric function [\(4.17\)](#page-16-0) after the following conditions are applied

$$
A = \frac{1}{\ell^2}, \quad \xi = \frac{(n-1)(D-1)}{4(nD - D + 1)}.
$$
\n(4.19)

The integration constant B is related to the mass of the black hole. Stealths in Gauss-Bonnet and Lovelock gravities were found a few years ago [\[46](#page-28-2), [47\]](#page-28-3), where the authors restrict the analysis to flat manifolds $(\gamma = 0)$ only. Our approach not only confirms their result but also generalizes it to base manifolds with non-vanishing curvature and to spatial-coordinates dependent scalar fields.

Conformal gravity: Like the Schwarzschild black hole in General Relativity, Birkhoff's theorem yields the Riegert black hole in conformal gravity, whose stealth was originally found in [\[50\]](#page-28-6), and reconsidered recently as part of a scalar-tensor renormalization program in [\[51](#page-28-7)]. Following the notation of the latter, for $\xi = 1/6$ the solution is given by,

$$
\phi = \phi(r), \quad f(r) = k + \frac{6mG}{r_0} - \frac{2}{r_0} \left(k + \frac{3mG}{r_0} \right) r - \frac{2mG}{r} - \frac{\lambda r^2}{3}.
$$
\n(4.20)

It is clear that from our reverse procedure, this solution is recovered by the four-dimensional timeindependent branch with a metric function [\(2.25\)](#page-9-2), a potential with the form presented in [\(2.26\)](#page-9-3), and a purely radial scalar field given in [\(2.24\)](#page-9-4). This is achieved by imposing the conditions,

$$
A = -\frac{\lambda}{3} - \frac{2mG}{r_0^3}, \qquad B = \frac{2mG\sigma^3}{r_0^3}, \qquad R_1 = \frac{1}{\sigma}, \qquad R_0 = -\frac{r_0}{\sigma}, \qquad \sigma = \sqrt{-\frac{k + \frac{2mG}{r_0} + \frac{\lambda r_0^2}{3}}{2\nu}}, \tag{4.21}
$$

where m, λ and r_0 are integration constants, whereas ν is a parameter that modules the behavior of the associated potential.

Quasitopological gravities: Although they do not possess a topological origin as Lovelock's model, Quasitopological Gravities [\[55](#page-28-11)[–58\]](#page-28-12) are constructed in $D \geq 5$ upon n–th order curvature interactions. Unlike Lovelock, the value of n in Quasitopological Gravities is not restricted by the spacetime dimension and can grow arbitrarily^{[5](#page-17-1)}. These theories emerge as gravitational models with second-order Euler-Lagrange equations when evaluated precisely in constant curvature internal spaces, as in our ansatz [\(2.2\)](#page-6-2), and they satisfy a n−th order algebraic equation for the metric function $f(r)$ [\[60](#page-29-0), [61](#page-29-1)]. It was shown that the n−th Quasitopological gravity fulfills

$$
\frac{D-2}{2}r^{D-2n-1}\left[\gamma - f(r) \right]^n = M ,
$$

where M is an integration constant associated with the ADM mass of the solution. For that reason, the reverse procedure allows straightforwardly accommodating a novel stealth solution considering the spatial-coordinates dependent scalar field [\(2.20\)](#page-8-6),

$$
\phi = \phi(r, \vec{x}), \quad f(r) = Ar^2 + Br^{\frac{(D-3)-4(D-2)\xi}{4\xi-1}} + \gamma,
$$
\n(4.22)

by fixing ξ as

$$
\xi = \frac{(n-1)(D-1)}{4(nD - D + 1)},
$$
\n(4.23)

and the self-interacting potential as shown in [\(2.21\)](#page-8-4).

To end this section, we remark on some general features. As shown through the previous examples and as emphasized in the introduction, the approach a) to b) is not bijective. Furthermore, the reverse construction does not impose restrictions on the purely gravitational sector (i.e., the left-hand side of [\(1.2b\)](#page-3-3)). For example, if $\xi = \xi_{D-1} = \frac{D-3}{4(D-2)}$, the metric function acquires the form

$$
f(r) = Ar^2 - M,
$$
\n(4.24)

which we recognize as a non-rotating BTZ-like black hole [\[52\]](#page-28-8). Thus, any gravitational theory whose vacuum solution admits such a profile is compatible with stealth, with the aforementioned coupling constant.

In sum, we retrieved and generalized known stealth solutions from the literature. Judging by the form of the metric function [\(2.17\)](#page-8-5) in the time-independent branch, the values of ξ and B become significant, with the former controlling the decay of the metric function and the latter being an integration constant that cannot be interpreted solely by solving $T_{\mu\nu} = 0$. Therefore, the physical interpretation of such constants is understood once the gravitational theory is specified.

5 Thermodynamic imprints of stealth black holes

It is well known that gravitational perturbations on the stealth background can backreact, and in consequence, at the perturbative level, the presence of this "gravitationally undetectable" object is exposed. Consequently, perturbations departing from the stealth configuration manifest clear differences with those departing from its trivial counterpart [\[62\]](#page-29-2). However, what is less known and has not yet been explored is whether these differences persist at the thermodynamic level, allowing one to distinguish the stealth black

⁵Recently, it was shown that in $D \geq 5$, the infinite tower of Quasitopological Gravities regularizes the curvature singularity of the Schwarzschild black hole [\[59](#page-29-3)].

hole from the trivial one. The spectrum of stealth solutions obtained by the reverse procedure leaves us in a good position for these purposes. This section aims to show that, indeed, there are differences induced by the presence of a stealth scalar field exposed in the thermodynamic quantities.

Let us choose, as an exemplary case, the five-dimensional Einstein-Gauss-Bonnet action as the purely metric theory $S_g[g]$, endowed with a non-minimally coupled scalar field in $S_{st}[\phi, g]$. This is,

$$
S[\phi, g] = \int d^5 x \sqrt{-g} \left[\frac{1}{2\kappa} \left(R - 2\Lambda + \alpha \mathcal{G} \right) - \frac{1}{2} \nabla_\mu \phi \nabla^\mu \phi - \frac{\xi}{2} R \phi^2 - U(\phi) \right],\tag{5.1}
$$

where the term G stand for the Gauss-Bonnet invariant given by $G = R^2 - 4R_{\alpha\beta}R^{\alpha\beta} + R_{\alpha\beta\mu\nu}R^{\alpha\beta\mu\nu}$, the self-interacting potential is given by,

$$
U(\phi) = \frac{1}{6\ell^2} \phi^2 \;, \tag{5.2}
$$

whose stealth solution was found in [\[46\]](#page-28-2), and it reads,

$$
\phi(r) = \frac{\phi_0}{r}, \quad f(r) = \frac{r^2}{\ell^2} - M. \tag{5.3}
$$

The reverse procedure recovers this solution by setting $D = 5$ and $n = 2$ in [\(4.19\)](#page-16-1).

The Euclidean approach in the calculation of the thermodynamical quantities establishes that the partition function for a thermodynamical ensemble is identified with the Euclidean path integral in the saddlepoint approximation around the classical Euclidean solution [\[63\]](#page-29-4). Once the symmetries of the spacetimes are known, Palais' "principle of symmetric criticality" states that the extremum of the action can be found imposing such symmetries on a minisuperspace metric and then varying the action. These symmetries extremely simplify the Euclidean procedure, as the general Hamiltonian formalism is not always available. However, this principle is not universal, as the symmetries imposed on the minisuperpsace must not over-reduce the degrees of freedom in the action, otherwise the set of field equations emerging from the reduced action will not correspond to the covariant field equations. Accordingly, the Euclidean continuation of this solution reads,

$$
ds_E^2 = N(r)f(r)d\tau^2 + \frac{dr^2}{f(r)} + r^2(dx_1^2 + dx_2^2 + dx_3^2) ,
$$
\n(5.4)

with $\phi = \phi(r)$. The flat coordinates span $x_i \in (0, \beta_i)$ with $i = \{1, 2, 3\}$. The period β of the Euclidean time τ is related to the temperature T of the black hole by $\beta = 1/T$. In consequence, the reduced Euclidean action S_E can be written in Hamiltonian form as follows,

$$
S_E = \beta \sigma \int_{r_+}^{\infty} dr N \mathcal{H} + B_E , \qquad (5.5)
$$

where B_E is a boundary term introduced in action [\(5.1\)](#page-18-0) and $\sigma = \beta_1 \beta_2 \beta_3$ is the volume of the base manifold. The variation of B_E cancels out all the contributions coming from variations of the bulk action, defining a well-posed variational principle to have an extremum on the classical solution. The reduced constraint is given by,

$$
\mathcal{H} = -\frac{3r}{2\kappa}(2f + rf') - \frac{1}{2}r\delta^{ij}\partial_i\phi\partial_j\phi - r^3\left[\frac{1}{2}f(\partial_r\phi)^2 + U(\phi) + \frac{\Lambda}{\kappa}\right] + 6\alpha ff' \n+ \xi\left\{\frac{3}{2}r\phi^2(rf' + 2f) + 2r\delta^{ij}(\partial_i\phi\partial_j\phi + \phi\partial_i\partial_j\phi) + \sqrt{f}\partial_r\left[\sqrt{f}r^3(\phi^2)'\right]\right\}.
$$
\n(5.6)

It is reassuring to check that, despite H being an involved expression, the following off-shell relations are satisfied,

$$
\frac{\delta S_E}{\delta N} = -2\sqrt{g} \frac{\mathcal{E}_{tt}}{N^3 f}, \n\frac{\delta S_E}{\delta f} = \sqrt{g} \left(\mathcal{E}_{rr} - \frac{\mathcal{E}_{tt}}{N^2 f^2} \right), \n\frac{\delta S_E}{\delta \phi} = \sqrt{g} \mathcal{E}_{\phi},
$$
\n(5.7)

confirming the validity of the boundary term. Its variation is given by,

$$
\delta B_E = \beta \sigma \left\{ -\frac{r^2}{2\kappa} \left(3\delta f + 2\kappa r \phi' f \delta \phi \right) + \frac{1}{2} r^2 \xi \left[2r \phi' (2f \delta \phi + \phi \delta f) + \phi \left(4r f \delta \phi' + 3\phi \delta f \right) \right] \Big|_{r_+}^{\infty} \tag{5.8}
$$

The variation of the fields at the horizon r_+ acquires the following form,

$$
\delta f|_{r_{+}} = - f'|_{r_{+}} \delta r_{+} , \quad \delta \phi|_{r_{+}} = \delta \phi (r_{+}) - \phi'|_{r_{+}} \delta r_{+} . \tag{5.9}
$$

Then, the variation of the boundary term at the horizon is

$$
\delta B_E(r_+) = \delta \left\{ \frac{2\pi}{\kappa} \sigma r_+^3 \left[1 - \kappa \xi \phi(r_+)^2 \right] \right\},\tag{5.10}
$$

which can be readily integrated as

$$
B_E(r_+) = \frac{A_+}{4\tilde{G}_+} \quad \text{with} \,, \quad \tilde{G}_+ = \frac{G}{\left(1 - \kappa \xi \phi \left(r_+\right)^2\right)},\tag{5.11}
$$

where $A_+ = \sigma r_+^3$. On the other hand, the variations of the fields at infinity are,

$$
\delta f|_{\infty} = -\delta M, \quad \delta \phi|_{\infty} = \frac{\delta \phi_0}{r}.
$$
\n(5.12)

With these quantities, the variation of the boundary term at infinity is,

$$
\delta B_E(\infty) = \frac{3r^2}{2\ell^2 \kappa} \beta \sigma(\ell^2 - 4\alpha \kappa) \delta M + \beta \sigma \left(6\alpha M - \frac{1}{12} \phi_0^2 \right) \delta M - \frac{1}{3} \beta \sigma M \phi_0 \delta \phi_0 \,. \tag{5.13}
$$

There is a potential quadratic divergence in the first term of this expression. However, solution [\(5.3\)](#page-18-1) requires that $\ell^2 = 4\kappa\alpha$ (the Chern–Simons point), which ensures the finiteness of the variation of the boundary term [\(5.13\)](#page-19-0). This is,

$$
\delta B_E(\infty) = \beta \sigma \left(\frac{3\ell^2}{2\kappa}M - \frac{1}{12}\phi_0^2\right)\delta M - \frac{1}{3}\beta \sigma M \phi_0 \delta \phi_0.
$$
\n(5.14)

Observe that $\delta B_E(\infty)$ is not necessarily a closed 1-form. Condition $\delta^2 B_E(\infty) = 0$ leads,

$$
\phi_0 \delta \phi_0 \wedge \delta M = 0. \tag{5.15}
$$

This equation raises two possibilities: $\phi_0 = 0$ or $\delta \phi_0 \wedge \delta M = 0$. The former switches off the scalar field, while the latter allows the possibility of computing the mass for the stealth solution. This implies to consider ϕ_0 as a function of M, and the variation $\delta B_E(\infty)$ is now exact by Poincaré's lemma. Since we are working in the grand canonical ensemble, the temperature is fixed, and the boundary term at infinity can be integrated as,

$$
B_E(\infty) = \beta \sigma \int dM \left[\frac{3\ell^2}{2\kappa} M - \frac{1}{12} \phi_0^2(M) - \frac{1}{3} M \phi_0(M) \frac{\partial \phi_0(M)}{\partial M} \right].
$$
 (5.16)

It is remarkable to see that the value of the Euclidean action is finite for any differentiable function $\phi_0(M)$ (see below for concrete examples), in which case,

$$
S_E = B(\infty) - B(r_+) = \beta \sigma \int dM \left[\frac{3\ell^2}{2\kappa} M - \frac{1}{12} \phi_0^2(M) - \frac{1}{3} M \phi_0(M) \frac{\partial \phi_0(M)}{\partial M} \right] - \frac{A_+}{4\widetilde{G}_+} \,, \tag{5.17}
$$

up to an arbitrary constant without variation. Since the Gibbs free energy F is related to the Euclidean action by $S_E = \beta F = \beta \mathcal{M} - \mathcal{S}$, the mass $\mathcal M$ and the entropy S are given by the standard thermodynamical relations

$$
\mathcal{M} = \frac{\partial S_E}{\partial \beta} = \sigma \frac{3\ell^2}{4\kappa} M^2 + \Delta \mathcal{M},\qquad(5.18)
$$

$$
S = \left(\beta \frac{\partial}{\partial \beta} - 1\right) S_E = \frac{A_+}{4\widetilde{G}_+}.
$$
\n(5.19)

It is clear from [\(5.18\)](#page-20-0) that the shift in the mass with respect to the trivial configuration is given by,

$$
\Delta \mathcal{M} = \sigma \int dM \left[-\frac{1}{12} \phi_0^2(M) - \frac{1}{3} M \phi_0(M) \frac{\partial \phi_0(M)}{\partial M} \right] , \qquad (5.20)
$$

and the value of the mass is altered by different choices of ϕ_0 as a function of M. This is precisely the case for a hairy black hole with secondary hair. Stealth solutions with primary hair are excluded from a thermodynamic perspective as their mass is not integrable.

In general, the shift in the mass generated by the stealth field must fulfill two conditions: It has to be non-trivial, meaning that $\delta(\Delta \mathcal{M}) \neq 0$; and non-divergent, which in this case is already guaranteed by construction for any differential function $\phi_0(M)$, according to formula [\(5.16\)](#page-20-1). In this regard, consider the following examples.

In the case that the scalar field is rigid, $\partial \phi_0(M)/\partial M = 0$, the mass reads

$$
\mathcal{M} = \sigma \left(\frac{3\ell^2}{4\kappa} M^2 - \frac{1}{12} \phi_0^2 M \right),\tag{5.21}
$$

fulfilling the above criteria. A healthy and non-trivial example can be achieved by $\phi_0(M) = M\ell$, obtaining

$$
\mathcal{M} = \sigma \left(\frac{3\ell^2}{4\kappa} M^2 - \frac{5\ell^2}{36} M^3 \right). \tag{5.22}
$$

It is interesting to analyze some extreme cases at this point. It could be the case that the shift in the mass is tuned in such a way that it cancels out the black hole's mass. To achieve this, and according to [\(5.20\)](#page-20-2), it is straightforward to see that

$$
\mathcal{M} = 0 \tag{5.23}
$$

is achieved by choosing $\phi_0(M) = \sqrt{\frac{6M\ell^2}{n}}$ $\frac{1}{\kappa}$. This kind of black hole with vanishing mass is not new. Black hole configurations with zero mass have already been reported in [\[64](#page-29-5)[–66\]](#page-29-6). However, the novelty of our result relies on the fact that the black hole mass is screened by a non-gravitating source.

6 Summary and outlook

In this article, we presented a novel method for constructing stealth solutions using a reverse approach. The latter proposes a generic procedure to construct stealth solutions by restricting the spacetime geometry only by imposing the stealth conditions $T_{\mu\nu} = 0$. As opposed to the standard procedure of obtaining stealth solutions — from step a) to step b) — starting with step b) by solving $T_{\mu\nu} = 0$ leads to a wide range of geometries compatible with stealth configurations. Then, if such geometries match the vacuum solutions of some gravity theory, such theory can be coupled to a non-gravitating scalar field, completing step a). Another consequence of this approach is that the role of the integration constants in the conserved charges may not be immediately clear. This is expected, as their nature relies on the specific theory to which the stealth solution is coupled. To ground this approach in a specific scalar-tensor theory, we focused on the non-minimally coupled scalar field in D dimensions.

As we show in Section [2](#page-6-0) and [3,](#page-10-1) the reverse approach makes it evident that the coordinate dependence of the scalar profile controls the form of the metric function in the homogeneous ansatz. For a timedependent scalar field, the metric function is determined by $f(r) = Ar^2 + Br + C$. In contrast, in the time-independent case, there are two possible scenarios for the metric function: A spatial-coordinates dependent $\phi(r, \vec{x})$ and a purely radial $\phi(r)$ scalar field. The spatial-coordinates dependent scalar field is compatible with a metric function $f(r) = Ar^2 + Br \frac{(D-3)-4(D-2)\xi}{4\xi-1} + C$ exhibiting a term whose decay depends on the coupling parameter ξ . This term is inherently different from the term found in pure GR, which is proportional to $1/r^{D-3}$ and incompatible with a stealth solution as it is obtained only when $\xi = 0$, yielding a constant scalar field. For the purely radial stealth scalar field, the metric function admits an involved expression of Gauss' hypergeometric and Gamma functions, which, in the conformal case $\xi = \xi_D$, reduces to $f(r) = Ar^2 + 2\gamma R_1 r/R_0 + \gamma + B(R_1 r + R_0)^{D-1}/r^{D-3}$.

In this work, the inclusion of a self-interacting potential in the scalar-tensor sector $S_{st}[\phi, g]$ of the action that supports stealth configurations takes the form

$$
U(\phi) = \lambda_1 \phi^2 + \lambda_2 \phi^{-\frac{2\xi - 1}{\xi}} + \lambda_3 \phi^{\frac{1}{2\xi}}.
$$
\n(6.1)

This potential consists of three terms: one massive $\propto \phi^2$ and two terms involving powers of ξ . The massive term of the potential provides the vacuum of the theory, given by (A)dS or Minkowski spacetimes. The second term $\propto \phi^{-\frac{2\xi-1}{\xi}}$ emerges as an extension of the conformal sector, in the sense that if $\xi = \xi_D$, the associated power reduces to the conformal case. Lastly, the term $\propto \phi^{\frac{1}{2\xi}}$ induces pure deviations from the conformally coupled scalar field. Its effect is controlled by a parameter that modulates the affine profile of the scalar field.

The reverse approach was applied to some concrete examples analyzed in Section [4,](#page-13-0) along with its thermodynamic implications as seen in Section [5.](#page-17-0) Naturally, all the known stealth solutions with nonminimally coupled scalar fields were recovered, and additionally, in some cases, they were generalized. This is because, in this reverse procedure, the problem finally reduces to match the parameters and integration constants in the stealth solution with those generated from the purely metric side. For instance, if one fixes the coupling parameter to its conformal value in one dimension lower, $\xi = \xi_{D-1} = \frac{D-3}{4(D-2)}$, the metric function [\(2.17\)](#page-8-5) acquires a non-rotating BTZ-like form

$$
f(r) = Ar^2 - M,
$$
\n
$$
(6.2)
$$

irrespective of the gravity model under consideration. This means that not only

$$
S[g, \phi] = S_g[g] - \int d^D x \sqrt{-g} \left[\frac{1}{2} \nabla_\mu \phi \nabla^\mu \phi + \frac{1}{2} \xi_{D-1} R \phi^2 + U(\phi) \right],
$$
 (6.3)

with $S_g[g] = \frac{R - 2\Lambda}{2\kappa}$ admits a stealth solution as shown in [\[21\]](#page-26-1), but also any D-dimensional gravity theory $S_q[g]$ whose family of solutions contains a BTZ-like metric function [\(6.2\)](#page-21-1) and in particular AdS_D spacetimes. Furthermore, we have shown that the reverse approach paves a direct path to finding new nongravitating configurations. Concretely, novel stealth solutions have been found in New Massive Gravity, Lovelock, and Quasitopological Gravities (see Section [4.2\)](#page-15-0).

It is important to note that even when the stress-energy tensor vanishes, the stealth solution can still exhibit tangible physical effects. With the solutions provided by the reverse procedure, we focused on studying the thermodynamic imprints left by the non-gravitating configurations. To develop this point, the five-dimensional Einstein-Gauss-Bonnet action with a non-minimally coupled scalar field is considered in Section [5.](#page-17-0) Surprisingly, the black hole mass is screened by the non-gravitating source. Consequently, a stealth scalar field is not only detectable at the perturbative level but also at the thermodynamic level. This result is physically relevant as it may have phenomenological implications: If we measure the mass of a celestial body with our current tools, we would not be able to distinguish whether the measurable mass includes a contribution from a stealth scalar field.

As shown in this article, the reverse approach only requires a $S_{\rm st}$ action and a particular ansatz for the geometry. Here, we explored the non-minimal coupling $R\phi^2$ over a static and spherically symmetric ansatz. In this regard, it is possible to go beyond the aforementioned non-minimal coupling towards, for example, $G^{\mu\nu}\nabla_{\mu}\phi\nabla_{\nu}\phi$, where it is known that a stealth configuration $\phi = \phi(t,r)$ is compatible with the Schwarzschild solution [\[24](#page-26-4)]. Motivated by the existence of this solution, it is feasible to explore the reverse approach within this framework. On the other hand, it is natural to consider an anisotropic ansatz, where stealth solutions are already known on a Lifshitz-type background [\[67\]](#page-29-7). For such an ansatz, the reverse approach could unveil black hole solutions with (bulk or asymptotic) anisotropic scaling.

In this work, we have seen that the stealth scalar field has a physical effect on the mass. Another avenue to pursue is to consider a rotating ansatz and explore the consequences of a stealth configuration on the angular momentum. More generally, one can naturally ask if some electric or color charge is changed under the presence of a stealth configuration. The latter may be achieved through a charged non-minimal coupling.

Acknowledgments

The authors thank E. Ayon-Beato, F. Cánfora, C. Corral, D. Flores-Alfonso, O. Fuentealba, M. Hassaïne, R. Troncoso, and J. Zanelli for useful discussions. L.G. and K.L. especially thank C. Henriquez-Baez for her contribution during the first stage of this project. C.E. is funded by Agencia Nacional de Investigacion y Desarrollo (ANID) through Proyecto FONDECYT Iniciación folio 11221063, Etapa 2024. The work of K.L. is supported by FONDECYT grant 3230618.

A Case $\xi = 1/4$

This appendix deals with the case $\xi = 1/4$ that we omitted in the main article. Starting from the metric ansatz [\(2.2\)](#page-6-2), the off-diagonal components of the T^{ν}_{μ} tensor are simultaneously solved only in the case when the scalar field is separable, and it does not depend on the transverse sector, namely

$$
\phi(t, r, x_i) = T(t)R(r). \tag{A.1}
$$

Furthermore, the T_t^r equation imposes an additional constraint, proportional to

$$
\dot{T}f'=0,\tag{A.2}
$$

which forces purely-radial scalar fields or metrics with $f(r) = cst$. In any case, we automatically have $T_1^1 = T_2^2 = \cdots = T_{D-2}^{D-2}$ for the spatial sector, meaning that we have three effective equations to solve: $T_t^t = 0, T_r^r = 0, \text{ and } T_i^i = 0.$

In the first case, when $\phi = \phi(r)$ is a purely-radial scalar field, the combination $T_t^t - T_r^r = 0$ turns out to be a differential equation for $\phi(r)$, which is solved by

$$
\phi(r) = R_0 e^{R_1 r} \,,\tag{A.3}
$$

but in this case, $f(r)$ is generically given by an elliptic integral.

Regarding the second case, when $\phi = \phi(t, r)$ forces the metric function to be a constant. In this case, we can only dress the Minkowski spacetime, obtained when $f(r) = \gamma = 1$. The scalar field reads

$$
\phi(t,r) = R_0 e^{R_1(r^2 - t^2)},\tag{A.4}
$$

and the potential $U(\phi)$ yields

$$
U(\phi) = \left[\frac{(D-1)R_1}{R_0} + 2\ln\left(\frac{\phi}{R_0}\right)\right]\phi^2,
$$
\n(A.5)

in agreement with the results of Ref. [\[22\]](#page-26-2).

B Explicit form of $T_t^t - T_i^i$ i

In this appendix, we show the subtraction $T_t^t - T_i^i$ for a general ansatz of the form

$$
ds^{2} = -f(r)dt^{2} + \frac{dr^{2}}{f(r)} + h^{2}(r)d\Sigma_{D-2,\gamma}^{2},
$$
\n(B.1)

with $h(r)$ an arbitrary function, where

$$
d\Sigma_{D-2,\gamma}^{2} = \Omega\left(\vec{x}\right)^{2} dx_{i} dx^{i}, \quad \Omega\left(\vec{x}\right) = \frac{1}{1 + \frac{\gamma}{4}x_{i}x^{i}}.
$$
\n(B.2)

The construction of Section [2](#page-6-0) is recovered for the particular case $h(r) = r$.

The difference $T_t^t - T_i^i$ is

$$
T_t^t - T_i^i = \frac{\psi^{\frac{1}{4\xi - 1}} \Omega(\vec{x})}{8\sqrt{f}h^2(4\xi - 1)} \left(\Xi_2 \sum_{i=1}^{D-2} x_i^2 + \Xi_1 \sum_{i=1}^{D-2} b_i x_i + \Xi_0 \right) = 0,
$$
 (B.3)

where $\Xi_{2,1,0}$ are, in general, differential equations for T, h and f. Concretely, Ξ_2 is

$$
\Xi_{2} \equiv 8k\xi^{2}h^{2}\ddot{T} + \left\{\gamma\xi(8\xi - 2)f^{2}hh'' + 2(D - 3)\gamma\xi(4\xi - 1)f^{2}h'^{2}\right\} \n+ \gamma\xi[4(5 - D)\xi + D - 4]ff'hh' + \left[\gamma\xi(1 - 4\xi)ff'' - 2\gamma\xi^{2}f'^{2}\right]h^{2} \n+ 2(D - 3)\gamma^{2}\xi(1 - 4\xi)f\right\}T + 2\xi(4\xi - 1)(4ah + \gamma R)f^{3/2}hh'' \n+ 8(D - 3)\xi\left[\frac{4(D - 2)\xi - D + 3}{D - 3}h + \frac{\gamma(4\xi - 1)}{4}R\right]f^{3/2}h'^{2} \n+ \left\{4a\xi[D - 4 - 4(D - 3)\xi]\sqrt{f}f'h^{2} + \left[\gamma(D - 4)(1 - 4\xi)\sqrt{f}f'R + 8k\xi^{2}f^{3/2}R'\right]h\right\}h' \n+ 4a\xi(1 - 4\xi)\sqrt{f}f''h^{3} + (\gamma\xi(1 - 4\xi)\sqrt{f}f''R - 4k\xi^{2}\sqrt{f}f'R')h^{2} + 4k\xi\left\{2(D - 3)a \n+ \xi\left[\gamma\sum_{i=1}^{D-2}c_{i} - (8D - 20)a\right]\right\}\sqrt{f}h + 2(D - 3)\gamma^{2}\xi(1 - 4\xi)\sqrt{f}R\right\},
$$
\n(B.4a)

whereas Ξ_1 ,

$$
\Xi_1 \equiv 8\xi(4\xi - 1)f^{3/2}h^2h'' + 8\xi[4(D - 2)\xi - D + 3]f^{3/2}hh'^2 - 4\xi[4(D - 3)\xi - D + 4]\sqrt{f}f'h^2h' + 4\xi(1 - 4\xi)\sqrt{f}f''h^3 - 8\gamma\xi(4(D - 2)\xi - D + 3)\sqrt{f}h,
$$
\n(B.4b)

and finally,

$$
\Xi_{0} = 32\xi^{2}h^{2}\ddot{T} + \left\{8\xi(4\xi - 1)f^{2}hh'' + 8(D - 3)\xi(4\xi - 1)f^{2}h'^{2} + 4\xi[4(5 - D)\xi + D - 4]ff'hh' \right\}\n+ \left[-4\xi(4\xi - 1)f f'' - 8\xi^{2}f'^{2}\right]h^{2} - 8(D - 3)\gamma\xi(4\xi - 1)f\right\}T + 8\xi(4\xi - 1)f^{3/2}hh'' \times\n\times\left(R + h\sum_{i=1}^{D-2}c_{i}\right) + 32(D - 2)\xi\left[\left(\xi - \frac{D - 3}{4(D - 2)}\right)h\sum_{i=1}^{D-2}c_{i} + \frac{D - 3}{4(D - 2)}(4\xi - 1)R\right]f^{3/2}h'^{2}\n+ \left\{-16\xi(D - 3)\sum_{i=1}^{D-2}c_{i}\left[\xi - \frac{D - 4}{4(D - 3)}\right]\sqrt{f}f'h^{2} + \left[-4\xi(D - 4)(4\xi - 1)\sqrt{f}f'R\right] (B.4c)\n+32\xi^{2}f^{3/2}R'\right]h\right\}h' - 4\xi(4\xi - 1)\sum_{i=1}^{D-2}c_{i}\sqrt{f}f''h^{3} + \left(-4\xi(4\xi - 1)\sqrt{f}f''R\right)\n-16\xi^{2}\sqrt{f}f'R'\right)h^{2} + \xi\left[-16(2D - 5)\left(\xi - \frac{2(D - 3)}{8D - 20}\right)\sum_{i=2}^{D-2}c_{i}\gamma + 64\xi a\right]\sqrt{f}h\n- 8(D - 3)\gamma\xi(4\xi - 1)\sqrt{f}R.
$$

Particularly, for the time-independent scalar fields of Section [2.1,](#page-7-0) the discussion of Eq. [\(2.14\)](#page-8-7) follows by imposing $T(t) = 0$, $R(r) = R_1r + R_0$, and $g(r) = r$.

References

- [1] Remo Ruffini and John A. Wheeler. "Introducing the black hole". In: Phys. Today 24.1 (1971), p. 30. DOI: [10.1063/1.3022513](https://doi.org/10.1063/1.3022513) (cited on page [3\)](#page-2-1).
- [2] Roy P. Kerr. "Gravitational Field of a Spinning Mass as an Example of Algebraically Special Metrics". In: Phys. Rev. Lett. 11 (5 Sept. 1963), pp. 237–238. DOI: [10.1103/PhysRevLett.11.237](https://doi.org/10.1103/PhysRevLett.11.237) (cited on page [3\)](#page-2-1).
- [3] Werner Israel. "Event Horizons in Static Vacuum Space-Times". In: Phys. Rev. 164 (5 Dec. 1967), pp. 1776–1779. DOI: [10.1103/PhysRev.164.1776](https://doi.org/10.1103/PhysRev.164.1776) (cited on page [3\)](#page-2-1).
- [4] Piotr T. Chrusciel, João Lopes Costa, and Markus Heusler. "Stationary Black Holes: Uniqueness and Beyond". In: Living Reviews in Relativity 15.1 (May 2012). issn: 1433-8351. DOI: [10.12942/lrr-2012-7](https://doi.org/10.12942/lrr-2012-7) (cited on page [3\)](#page-2-1).
- [5] Allen I. Janis, Ezra T. Newman, and Jeffrey Winicour. "Reality of the Schwarzschild Singularity". In: Phys. Rev. Lett. 20 (16 Apr. 1968), pp. 878–880. DOI: [10.1103/PhysRevLett.20.878](https://doi.org/10.1103/PhysRevLett.20.878) (cited on page [3\)](#page-2-1).
- [6] N. Bocharova, K. Bronnikov, and V. Melnikov. "An exact solution of the system of Einstein equations and mass-free scalar field". In: Vestn. Mosk. Univ. Fiz. Astron. 7 (1970), p. 706 (cited on page [3\)](#page-2-1).
- [7] C. Deffayet, G. Esposito-Farese, and A. Vikman. "Covariant Galileon". In: Phys. Rev. D 79 (8 Apr. 2009), p. 084003. DOI: [10.1103/PhysRevD.79.084003](https://doi.org/10.1103/PhysRevD.79.084003) (cited on page [3\)](#page-2-1).
- [8] C. Deffayet, G. Esposito-Farese, and A. Vikman. "Generalized Galileons: All scalar models whose curved background extensions maintain second-order field equations and stress tensors". In: Phys. Rev. D 80 (6 Sept. 2009), p. 064015. DOI: [10.1103/PhysRevD.80.064015](https://doi.org/10.1103/PhysRevD.80.064015) (cited on page [3\)](#page-2-1).
- [9] Tsutomu Kobayashi, Masahide Yamaguchi, and Jun'ichi Yokoyama. "Generalized G-inflation: Inflation with the most general second-order field equations". In: Prog. Theor. Phys. 126 (2011), pp. 511– 529. DOI: [10.1143/PTP.126.511](https://doi.org/10.1143/PTP.126.511). arXiv: [1105.5723 \[hep-th\]](https://arxiv.org/abs/1105.5723) (cited on page [3\)](#page-2-1).
- [10] Gregory Walter Horndeski. "Second-order scalar-tensor field equations in a four-dimensional space". In: Int. J. Theor. Phys. 10 (1974), pp. 363–384. DOI: [10.1007/BF01807638](https://doi.org/10.1007/BF01807638) (cited on page [3\)](#page-2-1).
- [11] Massimiliano Rinaldi. "Black holes with non-minimal derivative coupling". In: Phys. Rev. D 86 (2012), p. 084048.

DOI: [10.1103/PhysRevD.86.084048](https://doi.org/10.1103/PhysRevD.86.084048). arXiv: [1208.0103 \[gr-qc\]](https://arxiv.org/abs/1208.0103) (cited on page [3\)](#page-2-1).

- [12] Thomas P. Sotiriou. "Black Holes and Scalar Fields". In: Class. Quant. Grav. 32.21 (2015), p. 214002. DOI: [10.1088/0264-9381/32/21/214002](https://doi.org/10.1088/0264-9381/32/21/214002). arXiv: [1505.00248 \[gr-qc\]](https://arxiv.org/abs/1505.00248) (cited on page [3\)](#page-2-1).
- [13] Lam Hui and Alberto Nicolis. "No-Hair Theorem for the Galileon". In: Phys. Rev. Lett. 110 (2013), p. 241104. DOI: [10.1103/PhysRevLett.110.241104](https://doi.org/10.1103/PhysRevLett.110.241104). arXiv: [1202.1296 \[hep-th\]](https://arxiv.org/abs/1202.1296) (cited on page [3\)](#page-2-1).
- [14] Andrea Maselli, Hector O. Silva, Masato Minamitsuji, and Emanuele Berti. "Slowly rotating black hole solutions in Horndeski gravity". In: Phys. Rev. D 92 (10 Nov. 2015), p. 104049. DOI: [10.1103/PhysRevD.92.104049](https://doi.org/10.1103/PhysRevD.92.104049) (cited on page [3\)](#page-2-1).
- [15] Marco Crisostomi, Matthew Hull, Kazuya Koyama, and Gianmassimo Tasinato. "Horndeski: beyond, or not beyond?" In: JCAP 03 (2016), p. 038. DOI: [10.1088/1475-7516/2016/03/038](https://doi.org/10.1088/1475-7516/2016/03/038). arXiv: [1601.04658 \[hep-th\]](https://arxiv.org/abs/1601.04658) (cited on page [3\)](#page-2-1).
- [16] Jérôme Gleyzes, David Langlois, Federico Piazza, and Filippo Vernizzi. "Healthy theories beyond Horndeski". In: Phys. Rev. Lett. 114.21 (2015), p. 211101. DOI: [10.1103/PhysRevLett.114.211101](https://doi.org/10.1103/PhysRevLett.114.211101). arXiv: [1404.6495 \[hep-th\]](https://arxiv.org/abs/1404.6495) (cited on page [3\)](#page-2-1).
- [17] Tsutomu Kobayashi. "Horndeski theory and beyond: a review". In: Rept. Prog. Phys. 82.8 (2019), p. 086901. DOI: [10.1088/1361-6633/ab2429](https://doi.org/10.1088/1361-6633/ab2429). arXiv: [1901.07183 \[gr-qc\]](https://arxiv.org/abs/1901.07183) (cited on page [3\)](#page-2-1).
- [18] David Langlois and Karim Noui. "Degenerate higher derivative theories beyond Horndeski: evading the Ostrogradski instability". In: JCAP 02 (2016), p. 034. DOI: [10.1088/1475-7516/2016/02/034](https://doi.org/10.1088/1475-7516/2016/02/034). arXiv: [1510.06930 \[gr-qc\]](https://arxiv.org/abs/1510.06930) (cited on page [3\)](#page-2-1).
- [19] David Langlois. "Degenerate Higher-Order Scalar-Tensor (DHOST) theories". In: 52nd Rencontres de Moriond on Gra 2017, pp. 221–228. arXiv: [1707.03625 \[gr-qc\]](https://arxiv.org/abs/1707.03625) (cited on page [3\)](#page-2-1).
- [20] David Langlois. "Dark energy and modified gravity in degenerate higher-order scalar–tensor (DHOST) theories: A review". In: Int. J. Mod. Phys. D 28.05 (2019), p. 1942006. DOI: [10.1142/S0218271819420069](https://doi.org/10.1142/S0218271819420069). arXiv: [1811.06271 \[gr-qc\]](https://arxiv.org/abs/1811.06271) (cited on page [3\)](#page-2-1).
- [21] Eloy Ayon-Beato, Cristian Martinez, and Jorge Zanelli. "Stealth scalar field overflying a (2+1) black hole". In: Gen. Rel. Grav. 38 (2006), pp. 145–152. DOI: [10.1007/s10714-005-0213-x](https://doi.org/10.1007/s10714-005-0213-x). arXiv: [hep-th/0403228](https://arxiv.org/abs/hep-th/0403228) (cited on pages [3,](#page-2-1) [4,](#page-3-4) [7,](#page-6-5) [14,](#page-13-3) [15,](#page-14-3) [23\)](#page-22-1).
- [22] Eloy Ayon-Beato, Cristian Martinez, Ricardo Troncoso, and Jorge Zanelli. "Gravitational Cheshire effect: Nonminimally coupled scalar fields may not curve spacetime". In: Phys. Rev. D 71 (2005), p. 104037. DOI: [10.1103/PhysRevD.71.104037](https://doi.org/10.1103/PhysRevD.71.104037). arXiv: [hep-th/0505086](https://arxiv.org/abs/hep-th/0505086) (cited on pages [3,](#page-2-1) [7,](#page-6-5) [10,](#page-9-5) [11,](#page-10-2) [14,](#page-13-3) [16,](#page-15-2) [24\)](#page-23-1).
- [23] Christos Charmousis, Theodoros Kolyvaris, Eleftherios Papantonopoulos, and Minas Tsoukalas. "Black Holes in Bi-scalar Extensions of Horndeski Theories". In: JHEP 07 (2014), p. 085. DOI: [10.1007/JHEP07\(2014\)085](https://doi.org/10.1007/JHEP07(2014)085). arXiv: [1404.1024 \[gr-qc\]](https://arxiv.org/abs/1404.1024) (cited on page [3\)](#page-2-1).
- [24] Eugeny Babichev and Christos Charmousis. "Dressing a black hole with a time-dependent Galileon". In: JHEP 08 (2014), p. 106. DOI: [10.1007/JHEP08\(2014\)106](https://doi.org/10.1007/JHEP08(2014)106). arXiv: [1312.3204 \[gr-qc\]](https://arxiv.org/abs/1312.3204) (cited on pages [3,](#page-2-1) [7,](#page-6-5) [23\)](#page-22-1).
- [25] Eugeny Babichev, Christos Charmousis, and Antoine Leh´ebel. "Asymptotically flat black holes in Horndeski theory and beyond". In: JCAP 04 (2017), p. 027. DOI: [10.1088/1475-7516/2017/04/027](https://doi.org/10.1088/1475-7516/2017/04/027). arXiv: [1702.01938 \[gr-qc\]](https://arxiv.org/abs/1702.01938) (cited on page [4\)](#page-3-4).
- [26] Hayato Motohashi and Masato Minamitsuji. "General Relativity solutions in modified gravity". In: Phys. Lett. B 781 (2018), pp. 728–734. DOI: [10.1016/j.physletb.2018.04.041](https://doi.org/10.1016/j.physletb.2018.04.041). arXiv: [1804.01731 \[gr-qc\]](https://arxiv.org/abs/1804.01731) (cited on page [4\)](#page-3-4).
- [27] Masato Minamitsuji and Hayato Motohashi. "Stealth Schwarzschild solution in shift symmetry breaking theories". In: Phys. Rev. D 98.8 (2018), p. 084027. DOI: [10.1103/PhysRevD.98.084027](https://doi.org/10.1103/PhysRevD.98.084027). arXiv: [1809.06611 \[gr-qc\]](https://arxiv.org/abs/1809.06611) (cited on page [4\)](#page-3-4).
- [28] Jibril Ben Achour and Hongguang Liu. "Hairy Schwarzschild-(A)dS black hole solutions in degenerate higher order scalar-tensor theories beyond shift symmetry". In: Phys. Rev. D 99.6 (2019), p. 064042. DOI: [10.1103/PhysRevD.99.064042](https://doi.org/10.1103/PhysRevD.99.064042). arXiv: [1811.05369 \[gr-qc\]](https://arxiv.org/abs/1811.05369) (cited on page [4\)](#page-3-4).
- [29] Christos Charmousis, Marco Crisostomi, Ruth Gregory, and Nikolaos Stergioulas. "Rotating Black Holes in Higher Order Gravity". In: Phys. Rev. D 100.8 (2019), p. 084020. DOI: [10.1103/PhysRevD.100.084020](https://doi.org/10.1103/PhysRevD.100.084020). arXiv: [1903.05519 \[hep-th\]](https://arxiv.org/abs/1903.05519) (cited on page [4\)](#page-3-4).
- [30] Eloy Ayon-Beato, P. Isaac Ramírez-Baca, and César A. Terrero-Escalante. "Cosmological stealths with nonconformal couplings". In: Phys. Rev. D 97.4 (2018), p. 043505. DOI: [10.1103/PhysRevD.97.043505](https://doi.org/10.1103/PhysRevD.97.043505). arXiv: [1512.09375 \[gr-qc\]](https://arxiv.org/abs/1512.09375) (cited on page [4\)](#page-3-4).
- [31] Eloy Ayon-Beato, Alberto A. García, P. Isaac Ramírez-Baca, and César A. Terrero-Escalante. "Conformal stealth for any standard cosmology". In: Phys. Rev. D 88.6 (2013), p. 063523. DOI: [10.1103/PhysRevD.88.063523](https://doi.org/10.1103/PhysRevD.88.063523). arXiv: [1307.6534 \[gr-qc\]](https://arxiv.org/abs/1307.6534) (cited on page [4\)](#page-3-4).
- [32] Hideki Maeda and Kei-ichi Maeda. "Creation of the universe with a stealth scalar field". In: Phys. Rev. D 86 (2012), p. 124045. DOI: [10.1103/PhysRevD.86.124045](https://doi.org/10.1103/PhysRevD.86.124045). arXiv: [1208.5777 \[gr-qc\]](https://arxiv.org/abs/1208.5777) (cited on page [4\)](#page-3-4).
- [33] Mokhtar Hassaine. "Rotating AdS black hole stealth solution in D=3 dimensions". In: Phys. Rev. D 89.4 (2014), p. 044009. DOI: [10.1103/PhysRevD.89.044009](https://doi.org/10.1103/PhysRevD.89.044009). arXiv: [1311.4623 \[hep-th\]](https://arxiv.org/abs/1311.4623) (cited on pages [4,](#page-3-4) [7,](#page-6-5) [14,](#page-13-3) [16\)](#page-15-2).
- [34] Marc Henneaux, Alfredo Pérez, David Tempo, and Ricardo Troncoso. "Extended anti-de Sitter Hypergravity in 2+1 Dimensions and Hypersymmetry Bounds". In: International Workshop on Higher Spin Gauge Theories 2017, pp. 139–157. DOI: [10.1142/9789813144101-0009](https://doi.org/10.1142/9789813144101-0009). arXiv: [1512.08603 \[hep-th\]](https://arxiv.org/abs/1512.08603) (cited on page [4\)](#page-3-4).
- [35] Adolfo Cisterna, Mokhtar Hassaine, Julio Oliva, and Massimiliano Rinaldi. "Static and rotating solutions for Vector-Galileon theories". In: Phys. Rev. D 94.10 (2016), p. 104039. DOI: [10.1103/PhysRevD.94.104039](https://doi.org/10.1103/PhysRevD.94.104039). arXiv: [1609.03430 \[gr-qc\]](https://arxiv.org/abs/1609.03430) (cited on page [4\)](#page-3-4).
- [36] Ivica Smolić. "Spacetimes dressed with stealth electromagnetic fields". In: Phys. Rev. D 97.8 (2018), p. 084041. DOI: [10.1103/PhysRevD.97.084041](https://doi.org/10.1103/PhysRevD.97.084041). arXiv: [1711.07490 \[gr-qc\]](https://arxiv.org/abs/1711.07490) (cited on page [4\)](#page-3-4).
- [37] Cristian Quinzacara, Paola Meza, Almeira Sampson, and Mauricio Valenzuela. "Electromagnetically and gravitationally stealth fields". In: JCAP 03 (2023), p. 032. DOI: [10.1088/1475-7516/2023/03/032](https://doi.org/10.1088/1475-7516/2023/03/032). arXiv: [1904.00494 \[hep-th\]](https://arxiv.org/abs/1904.00494) (cited on page [4\)](#page-3-4).
- [38] J. A. Wheeler and K. Ford. Geons, black holes, and quantum foam: A life in physics. 1998 (cited on page [4\)](#page-3-4).
- [39] Andres Anabalon and Hideki Maeda. "New Charged Black Holes with Conformal Scalar Hair". In: Phys. Rev. D 81 (2010), p. 041501. DOI: [10.1103/PhysRevD.81.041501](https://doi.org/10.1103/PhysRevD.81.041501). arXiv: [0907.0219 \[hep-th\]](https://arxiv.org/abs/0907.0219) (cited on page [5\)](#page-4-0).
- [40] Marcela Cardenas, Oscar Fuentealba, and Cristián Martinez. "Three-dimensional black holes with conformally coupled scalar and gauge fields". In: Phys. Rev. D 90.12 (2014), p. 124072. DOI: [10.1103/PhysRevD.90.124072](https://doi.org/10.1103/PhysRevD.90.124072). arXiv: [1408.1401 \[hep-th\]](https://arxiv.org/abs/1408.1401) (cited on page [5\)](#page-4-0).
- [41] Luis Avilés, Hideki Maeda, and Cristian Martinez. "Exact black-hole formation with a conformally coupled scalar field in three dimensions". In: Class. Quant. Grav. 35.24 (2018), p. 245001. DOI: [10.1088/1361-6382/aaea9f](https://doi.org/10.1088/1361-6382/aaea9f). arXiv: [1808.10040 \[gr-qc\]](https://arxiv.org/abs/1808.10040) (cited on page [5\)](#page-4-0).
- [42] Mokhtar Hassaine. "Analogies between self-duality and stealth matter source". In: J. Phys. A 39 (2006), pp. 8675–8680. DOI: [10.1088/0305-4470/39/27/008](https://doi.org/10.1088/0305-4470/39/27/008). arXiv: [hep-th/0606159](https://arxiv.org/abs/hep-th/0606159) (cited on pages [5,](#page-4-0) [7\)](#page-6-5).
- [43] Marco M. Caldarelli, Christos Charmousis, and Mokhtar Hassaine. "AdS black holes with arbitrary scalar coupling". In: JHEP 10 (2013), p. 015. DOI: [10.1007/JHEP10\(2013\)015](https://doi.org/10.1007/JHEP10(2013)015). arXiv: [1307.5063 \[hep-th\]](https://arxiv.org/abs/1307.5063) (cited on page [5\)](#page-4-0).
- [44] Daniel Flores-Alfonso, Cesar S. Lopez-Monsalvo, and Marco Maceda. "Gravitational Waves from Thurston Geometries". In: (Nov. 2023). arXiv: [2311.12985 \[gr-qc\]](https://arxiv.org/abs/2311.12985) (cited on page [5\)](#page-4-0).
- [45] Moises Bravo-Gaete, Maria Montserrat Juarez-Aubry, and Gerardo Velazquez-Rodriguez. "Lifshitz black holes in four-dimensional critical gravity". In: Phys. Rev. D 105.8 (2022), p. 084009. DOI: [10.1103/PhysRevD.105.084009](https://doi.org/10.1103/PhysRevD.105.084009). arXiv: [2112.01483 \[hep-th\]](https://arxiv.org/abs/2112.01483) (cited on page [5\)](#page-4-0).
- [46] Moises Bravo Gaete and Mokhtar Hassaine. "Topological black holes for Einstein-Gauss-Bonnet gravity with a nonminimal scalar field". In: Phys. Rev. D 88 (2013), p. 104011. DOI: [10.1103/PhysRevD.88.104011](https://doi.org/10.1103/PhysRevD.88.104011). arXiv: [1308.3076 \[hep-th\]](https://arxiv.org/abs/1308.3076) (cited on pages [5,](#page-4-0) [7,](#page-6-5) [14,](#page-13-3) [17,](#page-16-2) [19\)](#page-18-2).
- [47] Moisés Bravo Gaete and Mokhtar Hassaine. "Planar AdS black holes in Lovelock gravity with a nonminimal scalar field". In: JHEP 11 (2013), p. 177. DOI: [10.1007/JHEP11\(2013\)177](https://doi.org/10.1007/JHEP11(2013)177). arXiv: [1309.3338 \[hep-th\]](https://arxiv.org/abs/1309.3338) (cited on pages [5,](#page-4-0) [7,](#page-6-5) [14,](#page-13-3) [17\)](#page-16-2).
- [48] Eugeny Babichev, Christos Charmousis, Mokhtar Hassaine, and Nicolas Lecoeur. "Conformally coupled scalar in Lovelock theory". In: Phys. Rev. D 107.8 (2023), p. 084050. DOI: [10.1103/PhysRevD.107.084050](https://doi.org/10.1103/PhysRevD.107.084050). arXiv: [2302.02920 \[gr-qc\]](https://arxiv.org/abs/2302.02920) (cited on page [5\)](#page-4-0).
- [49] Eloy Ay´on-Beato, Mokhtar Hassaine, and Pedro A. S´anchez. "Non-Noetherian Conformal Cheshire Effect". In: (July 2024). arXiv: [2408.00086 \[hep-th\]](https://arxiv.org/abs/2408.00086) (cited on page [5\)](#page-4-0).
- [50] Y. Brihaye and Y. Verbin. "Spherical structures in conformal gravity and its scalar-tensor extension". In: Phys. Rev. D 80 (12 Dec. 2009), p. 124048. DOI: [10.1103/PhysRevD.80.124048](https://doi.org/10.1103/PhysRevD.80.124048) (cited on pages [5,](#page-4-0) [14,](#page-13-3) [17\)](#page-16-2).
- [51] Giorgos Anastasiou, Ignacio J. Araya, Mairym Busnego-Barrientos, Cristobal Corral, and Nelson Merino. "Conformal renormalization of scalar-tensor theories". In: Phys. Rev. D 107.10 (2023), p. 104049. DOI: [10.1103/PhysRevD.107.104049](https://doi.org/10.1103/PhysRevD.107.104049). arXiv: [2212.04364 \[hep-th\]](https://arxiv.org/abs/2212.04364) (cited on pages [5,](#page-4-0) [14,](#page-13-3) [17\)](#page-16-2).
- [52] Maximo Banados, Claudio Teitelboim, and Jorge Zanelli. "The Black hole in three-dimensional space-time". In: Phys. Rev. Lett. 69 (1992), pp. 1849–1851. DOI: [10.1103/PhysRevLett.69.1849](https://doi.org/10.1103/PhysRevLett.69.1849). arXiv: [hep-th/9204099](https://arxiv.org/abs/hep-th/9204099) (cited on pages [14,](#page-13-3) [18\)](#page-17-2).
- [53] D. Lovelock. "The Einstein tensor and its generalizations". In: J. Math. Phys. 12 (1971), pp. 498– 501. DOI: [10.1063/1.1665613](https://doi.org/10.1063/1.1665613) (cited on page [17\)](#page-16-2).
- [54] Juan Crisostomo, Ricardo Troncoso, and Jorge Zanelli. "Black hole scan". In: Phys. Rev. D 62 (2000), p. 084013.

DOI: [10.1103/PhysRevD.62.084013](https://doi.org/10.1103/PhysRevD.62.084013). arXiv: [hep-th/0003271](https://arxiv.org/abs/hep-th/0003271) (cited on page [17\)](#page-16-2).

- [55] Julio Oliva and Sourya Ray. "A new cubic theory of gravity in five dimensions: Black hole, Birkhoff's theorem and C-function". In: Class. Quant. Grav. 27 (2010), p. 225002. DOI: [10.1088/0264-9381/27/22/225002](https://doi.org/10.1088/0264-9381/27/22/225002). arXiv: [1003.4773 \[gr-qc\]](https://arxiv.org/abs/1003.4773) (cited on page [18\)](#page-17-2).
- [56] Robert C. Myers and Brandon Robinson. "Black Holes in Quasi-topological Gravity". In: JHEP 08 (2010), p. 067. DOI: [10.1007/JHEP08\(2010\)067](https://doi.org/10.1007/JHEP08(2010)067). arXiv: [1003.5357 \[gr-qc\]](https://arxiv.org/abs/1003.5357) (cited on page [18\)](#page-17-2).
- [57] M. H. Dehghani et al. "Black Holes in Quartic Quasitopological Gravity". In: Phys. Rev. D 85 (2012), p. 104009.

DOI: [10.1103/PhysRevD.85.104009](https://doi.org/10.1103/PhysRevD.85.104009). arXiv: [1109.4708 \[hep-th\]](https://arxiv.org/abs/1109.4708) (cited on page [18\)](#page-17-2).

[58] Adolfo Cisterna, Luis Guajardo, Mokhtar Hassaine, and Julio Oliva. "Quintic quasi-topological gravity". In: JHEP 04 (2017), p. 066.

DOI: [10.1007/JHEP04\(2017\)066](https://doi.org/10.1007/JHEP04(2017)066). arXiv: [1702.04676 \[hep-th\]](https://arxiv.org/abs/1702.04676) (cited on page [18\)](#page-17-2).

- [59] Pablo Bueno, Pablo A. Cano, and Robie A. Hennigar. "Regular Black Holes From Pure Gravity". In: (Mar. 2024). arXiv: [2403.04827 \[gr-qc\]](https://arxiv.org/abs/2403.04827) (cited on page [18\)](#page-17-2).
- [60] Pablo Bueno, Pablo A. Cano, and Robie A. Hennigar. "(Generalized) quasi-topological gravities at all orders". In: Class. Quant. Grav. 37.1 (2020), p. 015002. DOI: [10.1088/1361-6382/ab5410](https://doi.org/10.1088/1361-6382/ab5410). arXiv: [1909.07983 \[hep-th\]](https://arxiv.org/abs/1909.07983) (cited on page [18\)](#page-17-2).
- [61] Pablo Bueno, Pablo A. Cano, Robie A. Hennigar, Mengqi Lu, and Javier Moreno. "Generalized quasi-topological gravities: the whole shebang". In: Class. Quant. Grav. 40.1 (2023), p. 015004. DOI: [10.1088/1361-6382/aca236](https://doi.org/10.1088/1361-6382/aca236). arXiv: [2203.05589 \[hep-th\]](https://arxiv.org/abs/2203.05589) (cited on page [18\)](#page-17-2).
- [62] Valerio Faraoni and Andres F. Zambrano Moreno. "Are stealth scalar fields stable?" In: Phys. Rev. D 81 (2010), p. 124050. DOI: [10.1103/PhysRevD.81.124050](https://doi.org/10.1103/PhysRevD.81.124050). arXiv: [1006.1936 \[gr-qc\]](https://arxiv.org/abs/1006.1936) (cited on page [18\)](#page-17-2).
- [63] G. W. Gibbons and S. W. Hawking. "Action integrals and partition functions in quantum gravity". In: Phys. Rev. D 15 (10 May 1977), pp. 2752–2756. DOI: [10.1103/PhysRevD.15.2752](https://doi.org/10.1103/PhysRevD.15.2752) (cited on page [19\)](#page-18-2).
- [64] Rong-Gen Cai, Yan Liu, and Ya-Wen Sun. "A Lifshitz Black Hole in Four Dimensional $R^{**}2$ Gravity". In: JHEP 10 (2009), p. 080. DOI: [10.1088/1126-6708/2009/10/080](https://doi.org/10.1088/1126-6708/2009/10/080). arXiv: [0909.2807 \[hep-th\]](https://arxiv.org/abs/0909.2807) (cited on page [21\)](#page-20-3).
- [65] Andres Anabalon, Fabrizio Canfora, Alex Giacomini, and Julio Oliva. "Black holes with gravitational hair in higher dimensions". In: Phys. Rev. D 84 (2011), p. 084015. DOI: [10.1103/PhysRevD.84.084015](https://doi.org/10.1103/PhysRevD.84.084015). arXiv: [1108.1476 \[hep-th\]](https://arxiv.org/abs/1108.1476) (cited on page [21\)](#page-20-3).
- [66] Francisco Correa and Mokhtar Hassaine. "Thermodynamics of Lovelock black holes with a nonminimal scalar field". In: JHEP 02 (2014), p. 014. DOI: [10.1007/JHEP02\(2014\)014](https://doi.org/10.1007/JHEP02(2014)014). arXiv: [1312.4516 \[hep-th\]](https://arxiv.org/abs/1312.4516) (cited on page [21\)](#page-20-3).
- [67] Eloy Ayón-Beato, Mokhtar Hassaine, and María Montserrat Juárez-Aubry. "Stealths on Anisotropic Holographic Backgrounds". In: (June 2015). arXiv: [1506.03545 \[gr-qc\]](https://arxiv.org/abs/1506.03545) (cited on page [23\)](#page-22-1).