



SCIPP 01/08  
February 2001

## Update on the Measurement of $\alpha_S$ with a 500 GeV Linear Collider

BRUCE A. SCHUMM and ANDREW S. TRUITT

The University of California  
and the  
Santa Cruz Institute for Particle Physics  
Santa Cruz, CA 95064

An update on the prospects for the precise measurement of the strong coupling constant  $\alpha_S$  at a high energy Linear Collider via the three-jet rate is presented. In particular, the issue of the distribution of center-of-mass energies of the identified  $q\bar{q}$  event sample, which can affect the determination of  $\alpha_S$  at the scale  $Q^2 = (500\text{GeV})^2$ , is addressed.

To be published in the proceedings of the LCWS 2000 Linear Collider Workshop,  
Fermilab, Batavia, IL, October 24–28, 2000

This work was supported in part by the Department of Energy, Grant  
#DE-FG03-92ER40689

The idea that a 500 GeV Linear Collider could provide an accurate measurement of the strong coupling constant  $\alpha_S$  at a scale  $Q^2 = (500\text{GeV})^2$  has been under consideration for some time. In Reference [1], the issue of identifying a  $q\bar{q}$  sample of high enough quality (both in terms of its purity as well as its potential bias against  $q\bar{q}$  events with hard gluon radiation) was addressed in the context of the determination of the three-jet rate  $R_3$ . In that study, it was found that selection criteria based on charged particle multiplicity, thrust direction, visible energy, and momentum balance, when combined with a b-quark antitag (to select against  $t\bar{t}$  events) and the use of electron beam polarization (to select against  $W^+W^-$  production) provided a means of identifying a sample of  $e^+e^- \rightarrow q\bar{q}$  ( $q \neq t$ ) at  $E_{cm} = 500$  GeV adequate for the measurement of  $\alpha_S$  to  $\sim \pm 1\%$ .

While the study of Reference [1] did include the effects of initial state radiation and beamstrahlung on the identified event sample, it did not address the issue of how well the differential luminosity spectrum would have to be known in order to achieve an accuracy of  $< \pm 1\%$  on  $\alpha_S$  at  $Q^2 = (500\text{GeV})^2$ . That issue is addressed in this study. A sample of simulated  $e^+e^- \rightarrow q\bar{q}$  events was generated using the PYBMS [2] implementation of PYTHIA [3]. Jets were reconstructed according to the E0 [4] jetfinding algorithm, assuming calorimetric coverage out to  $|\cos\theta| \leq 0.99$ . The limiting effect of calorimeter segmentation was not considered, although it is not expected to substantially change the conclusions of the study.

The differential luminosity spectrum of effective center-of-mass energy of hard  $e^+e^-$  collisions at the Linear Collider can effect the  $\alpha_S$  measurement in two ways. Firstly, due to the accompanying boost, jets can coalesce when analyzed with a fixed jet-mass resolution parameter, leading to an underestimate of the three-jet rate  $R_3$ . Second, the running of  $\alpha_S$  itself leads to an increase of  $R_3$  for events with substantial radiation in the initial state. Both of these effects need to be understood well enough so that their correction introduces as systematic error substantially less than 1%.

Figure 1 shows the ratio of the reconstructed and true three-jet fractions  $R_3$  as a function of the effective center-of-mass energy of the event sample, assuming a nominal beam energy of 250 GeV, and a jet-mass resolution parameter of  $Y_{cut} = 0.003$ . Also shown is the differential luminosity spectrum of  $e^+e^- \rightarrow q\bar{q}$  events after the application of the event selection criteria of Reference [1]. A convolution of the two curves yields a mean value of

$$\left\langle \frac{R_{3recon}}{R_{3true}} \right\rangle = 0.982,$$

or about a 2% difference.

Figure 2 shows the evolved value of  $\alpha_S$  as a function of effective center-of-mass energy, again superimposed over the spectrum of selected events. A convolution of

these two curves yields


$$\left\langle \frac{\alpha_S^{meas}}{\alpha_S^{500}} \right\rangle = 1.012,$$

or about a 1% difference.


In either case, knowledge of the differential luminosity spectrum to about  $\pm 10\%$  or so, a far looser requirement than that imposed by other physics requirements (such as the top threshold study) would lead to a systematic uncertainty substantially below 1%. Thus, it is concluded that neither of these effects presents any problem for the accurate measurement of  $\alpha_S$  at a high energy Linear Collider.

## REFERENCES

1. B. A. Schumm, hep-ex/9612013, November 1996.
2. PYBMS is a package, due to Tim Barklow of SLAC, which implements beamstrahlung and initial state radiation within the PYTHIA framework. It is based on work presented in Reference [3], as well as the following: P. Chen, Phys. Rev. **D46**, (1992), 1186.
3. T. Sjostrand, Computer Physics Commun., **82** (1994), 74.
4. S. Bethke *et al.*, Nucl. Phys. **B370**, (1992), 310.



**FIGURE 1.** Measured vs. true three-jet rate  $R3$  as a function of effective center-of-mass energy  $S^*$  after ISR and beamstrahlung. Also shown is the  $S^*$  distribution of events, again after ISR and beamstrahlung, passing the event selection requirements of Reference [1].



**FIGURE 2.** Evolved  $\alpha_S$  as a function of effective center-of-mass energy  $S^*$  after ISR and beamstrahlung. Also shown is the  $S^*$  distribution of events, again after ISR and beamstrahlung, passing the event selection requirements of Reference [1].