

Shape Functions from $\bar{B} \rightarrow X_c \ell \bar{\nu}_\ell$

H. BOOS, T. FELDMANN, T. MANNEL, B.D. PECJAK
Theoretische Physik 1, Fachbereich Physik, Universität Siegen
D-57068 Siegen, Germany

Abstract

We study inclusive semi-leptonic $\bar{B} \rightarrow X_c \ell \bar{\nu}_\ell$ decay using the power counting $m_c \sim \sqrt{\Lambda_{\text{QCD}} m_b}$. Assuming this scaling for the charm-quark mass, the decay kinematics can be chosen to access the shape-function region even in $b \rightarrow c$ transitions. To apply effective field theory methods in this region we extend SCET to describe massive collinear quarks. We calculate the tree-level decay rate, including $\mathcal{O}(\Lambda_{\text{QCD}}/m_b)$ power corrections, and show that it factorizes into a convolution of jet and shape functions. We identify a certain kinematical variable whose decay spectrum is proportional to the universal leading-order shape function familiar from $b \rightarrow u$ decay, and speculate as to whether information about this shape function can be extracted from data on $b \rightarrow c$ decay.

1 Introduction

The standard tool in heavy quark physics, the expansion in inverse powers of the heavy quark mass, has been extended recently to describe the kinematical situation in which the hadronic jet produced by the decaying heavy quark carries a large energy of order m_b , but has a small invariant mass of order $\sqrt{m_b\Lambda_{\text{QCD}}}$. Although processes occurring in this so-called shape-function region of phase space do not permit a local operator product expansion, they can be described by an expansion in terms of non-local operators evaluated on the light-cone [1, 2]. This expansion is most conveniently discussed within the framework of the soft-collinear effective theory (SCET) [3, 4, 5, 6], as has been done in studies of soft-collinear factorization in $b \rightarrow u$ transitions up to one-loop order [7, 8, 9] and including $1/m_b$ power corrections [10, 11, 12, 13].

Our goal in this work is to examine the effects of final-state quark masses on inclusive semi-leptonic B decay occurring in the shape-function region. For the up, down, and strange quarks, these mass effects are presumably very small. This is clearly not the case for the charm quark, which is in fact most often treated as a heavy quark with mass $m_c \sim m_b$. In this work we suggest a different power counting for the charm-quark mass, namely $m_c \sim \sqrt{\Lambda_{\text{QCD}}m_b}$. This power counting is satisfied numerically and allows us to access the shape-function region even in $b \rightarrow c$ transitions [14, 15]. For decay kinematics restricted to the shape-function region the charm quark can be treated as a collinear quark with finite mass. It is a simple task to include such a collinear quark mass in the SCET Lagrangian [16, 17], but this has not yet been done beyond leading order or applied to a phenomenological example.

In this paper we formulate a version of SCET appropriate for describing processes involving massive collinear quark fields interacting with soft and collinear gluons, and use it as a tool to calculate the hadronic tensor for inclusive semi-leptonic $b \rightarrow c$ decay in the shape-function region. We calculate the hadronic tensor at tree-level, including $\mathcal{O}(\Lambda_{\text{QCD}}/m_b)$ power corrections, and show that it factorizes into a convolution of jet and shape functions. The structure of this factorization formula mirrors closely that of the $b \rightarrow u$ case, the differences being that a slightly different set of shape functions is needed at sub-leading order and that the shape functions depend on a different combination of kinematical variables as compared with the massless case. We identify a certain kinematical variable whose differential decay rate is directly proportional to the universal leading order shape function, and speculate as to whether information on this function can be extracted from a study of $b \rightarrow c$ decay. This variable is a generalization of the P_+ variable suggested in [18], which has been reconsidered in the framework of SCET in [9, 19].

The organization of the paper is as follows. In the next section we discuss our power counting and outline the calculation. We construct the SCET Lagrangian for a massive collinear quark in Section 3 and derive the corresponding transition currents in Section 4. In Section 5 we use this version of SCET to calculate the hadronic tensor for semi-leptonic $b \rightarrow c$ transitions in terms of a factorization formula. Differential decay distributions are discussed in Section 6 and compared with the well-known local

expansion of inclusive $b \rightarrow c$ spectra in Section 7. In Section 8 we discuss possible phenomenological applications of our results and we conclude in Section 9.

2 Power Counting and Factorization in $\bar{B} \rightarrow X_c \ell \bar{\nu}_\ell$

We are interested in calculating decay distributions for inclusive $\bar{B} \rightarrow X_c \ell \bar{\nu}_\ell$ decay in the shape-function region using effective field theory methods. In this section we will define our power counting and explain how a factorization formula arises at tree level. The central quantity of interest is the hadronic tensor, which contains all the QCD effects in the semi-leptonic decay and can be used to derive the differential decay spectrum. We define the hadronic tensor as

$$W^{\mu\nu} = \frac{1}{\pi} \text{Im} \langle \bar{B}(v) | T^{\mu\nu} | \bar{B}(v) \rangle, \quad (1)$$

where we use the state normalization $\langle \bar{B}(v) | \bar{B}(v) \rangle = 1$. The correlator $T^{\mu\nu}$ is given by

$$T^{\mu\nu} = i \int d^4x e^{-iq \cdot x} \text{T} \{ J^\dagger{}^\mu(x) J^\nu(0) \}. \quad (2)$$

Here J^μ is a flavor-changing weak transition current, which is

$$J^\mu = \bar{c} \gamma^\mu (1 - \gamma_5) b \quad (3)$$

for $\bar{B} \rightarrow X_c \ell \bar{\nu}_\ell$ decay. We shall use the notation of [20] and write the hadronic tensor in terms of five scalar structure functions

$$\begin{aligned} W_{\mu\nu} = & W_1 (p_\mu v_\nu + v_\mu p_\nu - g_{\mu\nu} v p - i \epsilon_{\mu\nu\alpha\beta} p^\alpha v^\beta) \\ & - W_2 g_{\mu\nu} + W_3 v_\mu v_\nu + W_4 (p_\mu v_\nu + v_\mu p_\nu) + W_5 p_\mu p_\nu, \end{aligned} \quad (4)$$

where the independent vectors are chosen to be v , the velocity of the \bar{B} meson, and $p \equiv m_b v - q$ with m_b the b quark pole mass and q the momentum of the outgoing lepton pair. We use the convention $\epsilon^{0123} = -1$.

The hadronic tensor is expressed in the effective theory as a double series in the perturbative coupling constant and a small parameter λ , which we define through the relations

$$\frac{\Lambda_{\text{QCD}}}{m_b} \sim \lambda^2 \quad \text{and} \quad \frac{m_c^2}{m_b^2} \sim \lambda^2. \quad (5)$$

This choice correlates the two small scales $m_c/m_b \sim \sqrt{\Lambda_{\text{QCD}}/m_b}$. Quantum fluctuations are described by three widely separated scales, $m_b^2 \gg m_b^2 \lambda^2 \gg m_b^2 \lambda^4$, provided that the jet momentum p satisfies $p^2 - m_c^2 \sim m_b^2 \lambda^2$. We will refer to these scales as the hard, jet, and soft scale respectively. We expect that the fluctuations occurring at these scales can be factorized into a convolution of hard, jet and soft (shape) functions, similarly to the massless case [7, 21]. We will outline the steps used to prove such a factorization

formula to all orders in perturbation theory, under the assumption that there are no additional complications compared to the massless case, but do not give a complete proof for decay into charm quarks. An essential part of a factorization proof would be to show that the IR divergences associated with the renormalization of the leading-power jet function are the same for massive and massless SCET, and that the IR divergences of the new operators at sub-leading power can be treated in dimensional regularization. Our aim here is to explain how our explicit tree-level results can be interpreted as the leading-order term of such a presumed all-orders factorization formula.

We calculate the hadronic tensor by means of a two-step matching procedure, QCD \rightarrow SCET \rightarrow HQET. In the first step, we remove fluctuations at the hard scale by matching the QCD Lagrangian and transition current onto their corresponding expressions in SCET. We derive the SCET Lagrangian in Section 3 and discuss the transition currents in Section 4, to relative order λ^2 in the power counting. In addition to these SCET expressions, we also need the sub-leading HQET Lagrangian

$$\mathcal{L}_{\text{HQET}}^{(2)} = \frac{1}{2m_b} \left[\bar{h}_v (iD_s)^2 h_v + g \frac{C_{\text{mag}}}{2} \bar{h}_v \sigma_{\mu\nu} F_s^{\mu\nu} h_v \right]. \quad (6)$$

In general, $C_{\text{mag}} \neq 1$ due to fluctuations at the hard scale, and the SCET transition currents are multiplied by Wilson coefficients depending on quantities at the hard scale. These Wilson coefficients define the hard functions in the factorization formula. However, the SCET transition currents and HQET Lagrangian have trivial hard coefficients at tree level, so we will not need to discuss these hard functions any further.

After this first step of matching, the correlator (2) is calculated using the effective-theory expressions for the Lagrangian and currents. This will be the subject of Section 5. The effective theory contains a collinear charm-quark field and an HQET field for the b quark, as well as soft and collinear light quarks and gluons. In the next section we will define more precisely what we mean by collinear and soft, for now we simply note that the collinear fields fluctuate at the scale $m_b^2 \lambda^2$, whereas the soft and HQET fields fluctuate at the scale $m_b^2 \lambda^4$. Since the B meson contains only soft degrees of freedom, the matrix element of each term in the correlator factorizes into the form

$$\begin{aligned} \langle \bar{B} | T_i^{\text{eff}} | \bar{B} \rangle &= \langle \bar{B} | \bar{h}_v [\text{soft fields}] h_v | \bar{B} \rangle \otimes \langle 0 | [\text{collinear fields}] | 0 \rangle \\ &\equiv S_i \otimes J_i, \end{aligned} \quad (7)$$

where the \otimes stands for a convolution. The vacuum matrix element of the collinear fields defines a set of perturbatively calculable jet functions J_i containing fluctuations at the jet scale. The tree-level jet functions are given in terms of propagators of the collinear fields. Calculating these jet functions removes the collinear degrees of freedom and defines the second step of matching SCET \rightarrow HQET. The matrix element of the soft fields between the \bar{B} meson states is calculated in HQET and defines a set of non-perturbative shape functions S_i . The differential decay distributions are then written in terms of the W_i , each of which factorizes into a convolution of jet and shape functions.

3 Matching onto the SCET Lagrangian

In this section we derive the SCET Lagrangian for soft and collinear gluons interacting with collinear charm-quark fields carrying a mass $m_c \sim \lambda m_b$. The leading-order result has been given previously in [17]. Here we extend the derivation to sub-leading order, using the position space formalism of [6]. We do not discuss the Yang-Mills Lagrangian, because its matching onto the effective theory is the same as in [22].

The starting point is the QCD Lagrangian for massive quark fields,

$$\mathcal{L}_{\text{QCD}} = \bar{\psi}(i\not{D} - m_c)\psi, \quad (8)$$

where $iD^\mu = i\partial^\mu + gA^\mu$. The derivation of the effective theory proceeds in the usual way. We introduce effective theory fields according to the momentum scaling of their Fourier components. In terms of two light-like vectors n_+, n_- satisfying $n_+n_- = 2$, the components of a collinear momentum p_c^μ scale as $(n_+p_c, p_{c\perp}, n_-p_c) \sim m_b(1, \lambda, \lambda^2)$ and those of a soft momentum as p_s^μ as $\sim m_b(\lambda^2, \lambda^2, \lambda^2)$.

Following the standard procedure, we decompose the QCD gluon field into a collinear field A_c^μ and a soft contribution A_s^μ . With our power counting for the jet momentum, we only have to consider charm-quark momenta which satisfy $p_c^2 - m_c^2 \sim m_b^2\lambda^2$. These are described by massive collinear quark fields. Because of the finite charm-quark mass, loop diagrams do not contain soft divergences related to charm propagators. Therefore charm quarks appear neither in the soft Lagrangian nor in the B -meson states. This is in contrast to the $b \rightarrow u$ decay, where the effective Lagrangian contains interactions between soft and collinear light-quark fields.

Notice that in the region of phase space where $p_c^2 - m_c^2 \sim m_c\Lambda_{\text{QCD}} \sim m_b^2\lambda^3$, we would have to consider another effective theory which describes exclusive hadronic charm resonances. Since $\Lambda/m_c \sim \lambda$ is a small quantity, one can switch to HQET and split the charm-quark momentum into $p_c = m_cv' + k'$, where k' is a residual momentum. Because the energy transfer to the charm quark is still large, it is boosted by a factor of $\mathcal{O}(1/\lambda)$ with respect to the B -meson rest frame. Therefore, the charm velocity scales like $v' \sim (1/\lambda, 1, \lambda)$, and the residual momentum scales as $k' \sim m_b(\lambda, \lambda^2, \lambda^3)$. One can then introduce an effective theory field $h_{v'}$ carrying a dynamical momentum k' , which is described in terms of a “boosted HQET”. Restricting ourselves to collinear quark fields, we have to make sure that the considered phenomenological observables are not sensitive to the exclusive region, see the discussion in Section 8.

For the collinear quark fields we define the projections

$$\xi(x) = \frac{\not{n}_-\not{n}_+}{4}\psi_c(x), \quad \eta(x) = \frac{\not{n}_+\not{n}_-}{4}\psi_c(x). \quad (9)$$

Inserting these into (8) and solving the field equation for η , one finds

$$\eta = -\frac{\not{n}_+}{2} \frac{1}{in_+D} (i\not{D}_\perp - m_c)\xi, \quad (10)$$

so that the Lagrangian becomes

$$\mathcal{L}_{\text{QCD}} = \bar{\xi} \left[in_- D + (i\mathcal{D}_\perp - m_c) \frac{1}{in_+ D} (i\mathcal{D}_\perp + m_c) \right] \frac{\not{n}_+}{2} \xi. \quad (11)$$

The manipulations performed so far are just a rewriting of the QCD Lagrangian. In the next step we expand this Lagrangian as a series in the parameter λ . The expansion is related to the soft gluon fields contained in the covariant derivative $iD^\mu = i\partial^\mu + gA_c^\mu + gA_s^\mu$, as well as the multi-pole expansion of these soft fields. The result for the leading-order Lagrangian is

$$\mathcal{L}_\xi^{(0)} = \bar{\xi} \left(in_- D + (i\mathcal{D}_{\perp c} - m_c) \frac{1}{in_+ D_c} (i\mathcal{D}_{\perp c} + m_c) \right) \frac{\not{n}_+}{2} \xi, \quad (12)$$

while the sub-set of power-suppressed terms proportional to the mass is

$$\begin{aligned} \mathcal{L}_{\xi m}^{(1)} &= m_c \bar{\xi} \left[gA_{\perp s}, \frac{1}{in_+ D_c} \right] \frac{\not{n}_+}{2} \xi, \\ \mathcal{L}_{\xi m1}^{(2)} &= m_c^2 \bar{\xi} \frac{1}{in_+ D_c} g n_+ A_s \frac{1}{in_+ D_c} \frac{\not{n}_+}{2} \xi, \\ \mathcal{L}_{\xi m2}^{(2)} &= m_c \bar{\xi} \left[\frac{1}{in_+ D_c} g n_+ A_s \frac{1}{in_+ D_c}, i\mathcal{D}_{\perp c} \right] \frac{\not{n}_+}{2} \xi, \\ \mathcal{L}_{\xi m3}^{(2)} &= m_c \bar{\xi} \left[(x_\perp \partial_\perp gA_{\perp s}), \frac{1}{in_+ D_c} \right] \frac{\not{n}_+}{2} \xi, \end{aligned} \quad (13)$$

where $\mathcal{L}^{(1)}$ and $\mathcal{L}^{(2)}$ contribute to relative order λ and λ^2 to the effective action. In the terms above the collinear fields are evaluated at x , but the soft fields are multi-pole expanded and depend only on $x_-^\mu = (n_+ x/2) n_-^\mu \equiv x_+ n_-^\mu$. We will not make this explicit in the following, where it is always understood that

$$h_\nu(x) \equiv h_\nu(x_+ n_-^\mu), \quad A_s(x) \equiv A_s(x_+ n_-^\mu). \quad (14)$$

The remaining power-suppressed terms are the same as in the massless case. One can still make the standard field redefinitions ($\xi = Y\xi^{(0)}$, etc.) to decouple the soft gluon field from the leading-order collinear Lagrangian (12), a fact which is crucial to factorization.

The Lagrangian can be written in a manifestly gauge-invariant form by redefining the collinear fields using the methods developed in [22]. Applying these techniques, we find that the result for the leading-order Lagrangian (12) is unchanged, but to relative order λ^2 the power suppressed terms are the same as in the massless case and read [22]

$$\begin{aligned} \mathcal{L}_\xi^{(1)} &= \bar{\xi} x_\perp^\mu n_-^\nu W_c g F_{\mu\nu}^s W_c^\dagger \frac{\not{n}_+}{2} \xi \\ \mathcal{L}_{1\xi}^{(2)} &= \frac{1}{2} \bar{\xi} (n_- x) n_+^\mu n_-^\nu W_c g F_{\mu\nu}^s W_c^\dagger \frac{\not{n}_+}{2} \xi, \\ \mathcal{L}_{2\xi}^{(2)} &= \frac{1}{2} \bar{\xi} x_\perp^\mu x_{\perp\rho} n_-^\nu W_c [D_{\perp s}^\rho, gF_{\mu\nu}^s] W_c^\dagger \frac{\not{n}_+}{2} \xi, \end{aligned}$$

$$\begin{aligned}
\mathcal{L}_{3\xi}^{(2)} &= \frac{1}{2} \bar{\xi} i \not{D}_{\perp c} \frac{1}{in_+ D_c} x_{\perp}^{\mu} \gamma_{\perp}^{\nu} W_c g F_{\mu\nu}^s W_c^{\dagger} \frac{\not{n}_+}{2} \xi \\
&+ \frac{1}{2} \bar{\xi} x_{\perp}^{\mu} \gamma_{\perp}^{\nu} W_c g F_{\mu\nu}^s W_c^{\dagger} \frac{1}{in_+ D_c} i \not{D}_{\perp c} \frac{\not{n}_+}{2} \xi.
\end{aligned} \tag{15}$$

The power-suppressed mass terms first appear at relative order λ^3 . To see this, we note that after making the field redefinitions and working in the collinear light-cone gauge $n_+ A_c = 0$, one effectively replaces

$$\frac{1}{in_+ D} \rightarrow \frac{1}{in_+ \partial} + \mathcal{O}(\lambda^3) \tag{16}$$

in the expansion of (11) (see eq. (23) of [22]). It is then easy to show that all mass terms not contained in the leading-order Lagrangian are suppressed by at least a factor λ^3 .

While no explicit power-suppressed mass effects appear to $\mathcal{O}(\lambda^2)$ accuracy in the effective Lagrangian after making the field redefinitions, we will see later on that they reappear in the time-ordered products of SCET Lagrangians and currents. The calculation with either form of the Lagrangian and currents yields the same result, namely the appearance of one new sub-leading shape function as compared to the massless case.

4 Effective theory currents

In this section we consider the matching of the flavor-changing weak transition current defined in QCD onto its corresponding expression in SCET. For $b \rightarrow c$ transitions, this matching takes the form

$$(\bar{\psi}_c \Gamma b)_{\text{QCD}} \rightarrow e^{-im_b v x} [J^{(0)} + J^{(1)} + J^{(2)} + \dots], \tag{17}$$

where $\bar{\psi}_c$ is a collinear charm-quark field. In general, the J_i are convolutions of a short distance Wilson coefficient with operators built out of SCET and HQET fields. In what follows, however, we will work only at tree level, where the convolutions reduce to simple multiplication. The power-suppressed currents we need here can be deduced from the results in [22] by noting that at tree level all powers of the mass are related to the η field. Using (10) in conjunction with [22], we find for the mass terms

$$J_m^{(1)} = m_c \bar{\xi} \frac{\not{n}_+}{2} \frac{1}{-in_+ \overleftarrow{D}_c} W_c \Gamma h_v, \tag{18}$$

$$J_{m1}^{(2)} = m_c \bar{\xi} \frac{\not{n}_+}{2} \frac{1}{-in_+ \overleftarrow{D}_c} W_c \Gamma x_{\perp} D_{\perp s} h_v, \tag{19}$$

$$J_{m2}^{(2)} = m_c \bar{\xi} \frac{\not{n}_+}{2} \frac{1}{in_+ \overleftarrow{D}_c} \Gamma \frac{\not{n}_-}{2m_b} [i \not{D}_{\perp c} W_c] h_v. \tag{20}$$

The convention is such that derivatives do not act outside of the square brackets, and the soft gluon field and the heavy quark field are multi-pole expanded. We have again

used the field redefinitions of [22] to write the result in a manifestly gauge invariant form. The remaining terms are the same as in the massless case [22, 23, 24]. The sub-set of these that will be needed later on when calculating the tree-level time-ordered products is

$$\begin{aligned}
J^{(0)} &= \bar{\xi} W_c \Gamma h_v, \\
J_1^{(1)} &= \bar{\xi} W_c \Gamma x_{\perp\mu} D_{\perp s}^{\mu} h_v, \quad J_2^{(1)} = -\bar{\xi} i \overleftarrow{D}_{\perp c} \frac{1}{in_+ \overleftarrow{D}_c} W_c \frac{\not{n}_+}{2} \Gamma h_v, \\
J_1^{(2)} &= \bar{\xi} W_c \Gamma \frac{n_- x}{2} n_+ D_s h_v, \quad J_2^{(2)} = \bar{\xi} W_c \Gamma \frac{x_{\mu\perp} x_{\nu\perp}}{2} D_{\perp s}^{\mu} D_{\perp s}^{\nu} h_v, \\
J_3^{(2)} &= -\bar{\xi} i \overleftarrow{D}_{\perp c} \frac{1}{in_+ \overleftarrow{D}_c} W_c \frac{\not{n}_+}{2} \Gamma x_{\perp\mu} D_{\perp s}^{\mu} h_v, \quad J_4^{(2)} = \bar{\xi} W_c \Gamma \frac{i \not{D}_s}{2m_b} h_v.
\end{aligned} \tag{21}$$

5 The hadronic tensor at tree level

In this section we obtain an expression for the hadronic tensor in terms of a factorization formula. We work at tree level and include $\mathcal{O}(\lambda^2)$ power corrections. We start by calculating the time-ordered products appearing in the correlator (2) using the effective theory Lagrangian and currents derived in the previous sections, setting $A_c = 0$ as appropriate at tree level.

Many time-ordered products are possible at $\mathcal{O}(\lambda^2)$. It is convenient to divide these into two groups: those that vanish when $m_c \rightarrow 0$, and those that do not. Since the sub-leading Lagrangian (15) in its gauge-invariant form is actually the same as in the massless case, we can identify the time-ordered products which do not vanish when $m_c \rightarrow 0$ from eq. (54) of [12]. For these terms the difference between the two cases lies only in the leading-order Lagrangian, and we can account for this difference by replacing the propagator for massless collinear quark fields by that for massive fields, which is

$$\begin{aligned}
\langle 0 | T \{ \xi(x)_{a\alpha} \bar{\xi}(y)_{b\beta} \} | 0 \rangle &\equiv i \Delta(x-y) \delta_{ab} \left(\frac{\not{n}_-}{2} \right)_{\alpha\beta} \\
&= \int \frac{d^4 k}{(2\pi)^4} e^{-ik(x-y)} \frac{in_+ k}{k^2 - m_c^2 + i\epsilon} \left(\frac{\not{n}_-}{2} \right)_{\alpha\beta} \delta_{ab}.
\end{aligned} \tag{22}$$

The Latin (Greek) indices refer to color (Dirac) indices. The function $\Delta(z)$ satisfies

$$in_- \partial \Delta(z) = \delta^{(4)}(z) + \frac{1}{in_+ \partial} (m_c^2 - (i\partial_{\perp})^2) \Delta(z). \tag{23}$$

We will always work in the frame where $p_{\perp} = 0$, such that in tree-level expressions ∂_{\perp} can be dropped.

At tree level the factorization formula contains up to a double convolution between the jet and shape functions. The tree-level jet functions are related to (products of)




Figure 1: Representative Feynman diagrams involving soft gluon emission. These give rise to (a) bi-local and (b) tri-local shape functions.

propagators, and some of the convolution integrals can be performed explicitly, after which the results can be written as a convolution over a single variable only. To show how this works and to shorten intermediate expressions, we introduce the integral operators ($m_c^2 \equiv m_c^2 - i\epsilon, n_+p > 0$)

$$\begin{aligned}
I_2 * f &\equiv - \int d^4x e^{ipx} \int \frac{d^4k}{(2\pi)^4} e^{-ikx} \frac{n_+k}{k^2 - m_c^2} f(x_+) \\
&= - \int dx_+ e^{in_-px_+} \int \frac{dn_-k}{2\pi} e^{-in_-kx_+} \frac{1}{n_-k - m_c^2/n_+p} f(x_+) \\
&= i \int_0^\infty dx_+ e^{i(n_-p - m_c^2/n_+p)x_+} f(x_+), \tag{24}
\end{aligned}$$

$$\begin{aligned}
I_3 * g &\equiv - \int d^4x d^4z e^{ipx} \int \frac{d^4k}{(2\pi)^4} \frac{d^4k'}{(2\pi)^4} e^{-ik(x-z)} e^{-ik'z} \frac{n_+k}{(k^2 - m_c^2)(k'^2 - m_c^2)} g(x_+, z_+) \\
&= - \int dx_+ dz_+ e^{in_-px_+} \int \frac{dn_-k}{2\pi} \frac{dn_-k'}{2\pi} e^{-in_-k(x_+-z_+)} e^{-in_-k'z_+} \\
&\quad \frac{1}{n_+p} \left(\frac{1}{n_-k - m_c^2/n_+p} \right) \left(\frac{1}{n_-k' - m_c^2/n_+p} \right) g(x_+, z_+) \\
&= I_2 * \frac{-i}{n_+p} \int_0^{x_+} dz_+ g(x_+, z_+). \tag{25}
\end{aligned}$$

In terms of the factorization formula, the functions f and g represent generic shape functions depending on one or two variables respectively, and the momentum-dependent propagators correspond to tree-level jet functions. These are represented by Feynman diagrams such as Figure 1(a) for $f(x_+)$, and Figure 1(b) for $g(x_+, z_+)$. Performing the integral to arrive at the final lines of (24, 25) is equivalent to performing a convolution integral in the factorization formula. In (24) this is trivial, but in (25) the object

$\int_0^{x_+} dz_+ g(x_+, z_+)$ introduces an “effective shape function” of a single variable only. To keep the structure of factorization as presented in (7) clear, we will be careful to distinguish these effective shape functions from the shape functions defined directly by the multi-local matrix elements of soft operators between \bar{B} meson states.

The integral operators I_2, I_3 are the same as in [12] after replacing $n_-p \rightarrow n_-p - m_c^2/n_{+p}$ in the exponential of I_2^* defined there. We can thus obtain the part of the correlator that does not vanish when $m_c \rightarrow 0$ from eq. (54) of that work by making this replacement, and leaving out the term involving the soft quark field, which does not appear in the massive case. We also need to identify a small number of terms which vanish when $m_c \rightarrow 0$ and cannot be derived from previous results. First, we need insertions of the transition currents J_m . Using the short-hand notation

$$J_A^\dagger J_B i\mathcal{L}_C \equiv i \int d^4x e^{ipx} T \left\{ J_A^\dagger(x) J_B(0) i \int d^4z \mathcal{L}_C(z) \right\}, \quad (26)$$

and working in soft light-cone gauge $n_-A_s = 0$, the relevant time-ordered products are

$$J^{(0)\dagger} J_m^{(1)} + J_m^{(1)\dagger} J^{(0)} = \frac{m_c}{n_{+p}} I_2 * \bar{h}_v(x) \bar{\Gamma} \Gamma h_v(0), \quad (27)$$

$$J_m^{(1)\dagger} J_m^{(1)} = \frac{m_c^2}{(n_{+p})^2} I_2 * \bar{h}_v(x) \bar{\Gamma} \frac{\not{n}_+}{2} \Gamma h_v(0), \quad (28)$$

$$J_{m1}^{(2)\dagger} J^{(0)} = 0. \quad (29)$$

(Recall that the soft fields are multi-pole expanded, see (14).) The only additional mass terms are related to insertions of

$$\mathcal{L}_{1\xi}^{(2)} = \frac{1}{2} \bar{\xi}(n_-x) n_+^\mu n_-^\nu W_c g F_{\mu\nu}^s W_c^\dagger \frac{\not{n}_+}{2} \xi, \quad J_1^{(2)} = \bar{\xi} \Gamma \frac{n_-x}{2} n_+ D_s h_v. \quad (30)$$

They contribute to the correlator through the time-ordered products

$$\begin{aligned} J_1^{(2)\dagger} J^{(0)} + J^{(0)\dagger} J^{(0)} i\mathcal{L}_{1\xi}^{(2)} &= -\frac{m_c^2}{(n_{+p})^2} I_2 * (ix_+) \bar{h}_v(x) in_+ \overleftarrow{\partial} \bar{\Gamma} \frac{\not{n}_-}{2} \Gamma h_v(0) \\ &\quad -\frac{m_c^2}{n_{+p}} I_3 * \bar{h}_v(x) gn_+ A_s(z) \bar{\Gamma} \frac{\not{n}_-}{2} \Gamma h_v(0) \\ &= \frac{im_c^2}{(n_{+p})^2} I_2 * \int_0^{x_+} dz_+ \bar{h}_v(x) (-in_+ \overleftarrow{D}_s)(z) \bar{\Gamma} \frac{\not{n}_-}{2} \Gamma h_v(0). \end{aligned} \quad (31)$$

We used an integration by parts to write the field strength tensor in terms of the soft gluon field and then applied (23) to simplify (31). The simple form of the tree-level propagators has allowed us to replace n_-x with $n_+x m_c^2/(n_{+p})^2$ in this equation, as is easily verified by writing the factors of x in momentum space as derivatives acting on tree-level propagators. These expressions can be made gauge invariant by inserting the

appropriate Wilson lines. After including these Wilson lines, the tree-level result for the correlator can be written entirely in terms of multi-local covariant derivatives of the type in (31). In an arbitrary gauge these are defined by

$$(Y^\dagger iD_s^\mu Y)(z)_{ab} = i\partial^\mu \delta_{ab} + (Y^\dagger [iD_s^\mu Y])(z)_{ab}. \quad (32)$$

The indices refer to color and the derivative does not act outside the square brackets. At tree level the result for the correlator with arbitrary Dirac structure is

$$\begin{aligned} T^{\mu\nu}(u) = & - \int \frac{dx_+ dn_- k}{2\pi} e^{i(u-n_-k)x_+} \frac{1}{n_-k + i\epsilon} \times \left\{ \right. \\ & (\bar{h}_v Y)(x) \bar{\Gamma} \left[\frac{\not{n}_-}{2} + \frac{m_c}{n_+p} + \frac{m_c^2}{(n_+p^2)} \frac{\not{n}_+}{2} \right] \Gamma(Y^\dagger h_v)(0) \\ & + T \left\{ (\bar{h}_v Y)(x) \bar{\Gamma} \frac{\not{n}_-}{2} \Gamma(Y^\dagger h_v)(0) i \int d^4z \mathcal{L}_{\text{HQET}}^{(2)}(z) \right\} \\ & + \frac{1}{2m_b} \left[(\bar{h}_v(-i\overleftarrow{D}_s)Y)(x) \bar{\Gamma} \frac{\not{n}_-}{2} \Gamma(Y^\dagger h_v)(0) + (\bar{h}_v Y)(x) \bar{\Gamma} \frac{\not{n}_-}{2} \Gamma(Y^\dagger i\overleftarrow{D}_s h_v)(0) \right] \\ & - \frac{1}{n_+p} \left[(\bar{h}_v(-i\overleftarrow{D}_s^{\mu\perp})Y)(x) \bar{\Gamma} \frac{\not{n}_+}{2} \gamma_{\mu\perp} \frac{\not{n}_-}{2} \Gamma(Y^\dagger h_v)(0) \right. \\ & \quad \left. + (\bar{h}_v Y)(x) \bar{\Gamma} \frac{\not{n}_-}{2} \gamma_{\mu\perp} \frac{\not{n}_+}{2} \Gamma(Y^\dagger(-i\overleftarrow{D}_s^{\mu\perp})h_v)(0) \right] \\ & + \frac{i}{n_+p} \int_0^{x_+} dz_+ (\bar{h}_v Y)(x) \bar{\Gamma} \frac{\not{n}_-}{2} (Y^\dagger(-i\overleftarrow{D}_{s\perp})(-i\overleftarrow{D}_{s\perp})Y)(z) \Gamma(Y^\dagger h_v)(0) \\ & \left. + \frac{im_c^2}{(n_+p)^2} \int_0^{x_+} dz_+ (\bar{h}_v Y)(x) (Y^\dagger(-in_+\overleftarrow{D}_s)Y)(z) \bar{\Gamma} \frac{\not{n}_-}{2} \Gamma(Y^\dagger h_v)(0) \right\}, \quad (33) \end{aligned}$$

where we have defined $u \equiv n_-p - m_c^2/n_+p$.

In order to calculate the hadronic tensor, we need to take the imaginary part of the correlator and decompose the matrix elements between \bar{B} meson states in terms of scalar functions. The matrix elements of the operators on the final two lines of (33) could be used to directly define a set of effective shape functions depending on the variable u . To keep factorization explicit we do not take this approach, and introduce effective shape functions only in terms of convolutions involving multi-local shape functions defined through \bar{B} meson matrix elements. After contracting with the color indices of the tree-level jet functions, we need the following matrix elements (the Greek subscripts indicate Dirac indices and $\epsilon_{\mu\nu}^\perp \equiv i\epsilon_{\mu\nu\rho\sigma}n_-^\rho v^\sigma$)

$$\langle \bar{B} | (\bar{h}_v Y)(x)_\alpha (Y^\dagger h_v)(0)_\beta | \bar{B} \rangle = \frac{1}{2} \left(\frac{1+\psi}{2} \right)_{\beta\alpha} \tilde{S}(x_+), \quad (34)$$

$$\langle \bar{B} | (\bar{h}_v Y)(x)_\alpha (Y^\dagger h_v)(0)_\beta i \int d^4z \mathcal{L}_{\text{HQET}}^{(2)}(z) | \bar{B} \rangle = \frac{1}{2m_b} \frac{1}{2} \left(\frac{1+\psi}{2} \right)_{\beta\alpha} \tilde{s}(x_+),$$

$$\langle \bar{B} | (\bar{h}_v Y)(x)_\alpha (Y^\dagger i D_s^\mu Y)(z) (Y^\dagger h_v)(0)_\beta | \bar{B} \rangle = \quad (35)$$

$$\begin{aligned} & \frac{1}{2} \left(\frac{1 + \psi'}{2} \right)_{\beta\alpha} \left[-i \tilde{S}'(x_+) v^\mu + \left(i \tilde{S}'(x_+) - \tilde{T}_1(x_+, 0) + \tilde{T}_1(x_+, z_+) \right) n_-^\mu \right] \\ & + \frac{\epsilon_{\perp}^{\mu\rho}}{4} \left(\frac{1 + \psi'}{2} \gamma_{\rho\perp} \gamma_5 \frac{1 + \psi'}{2} \right)_{\beta\alpha} \left[\tilde{t}(x_+) - \tilde{T}_2(x_+, 0) + \tilde{T}_2(x_+, z_+) \right], \end{aligned}$$

$$\langle \bar{B} | (\bar{h}_v Y)(x)_\alpha (Y^\dagger i D_{\mu\perp}^s i D_{\nu\perp}^s Y)(z) (Y^\dagger h_v)(0)_\beta | \bar{B} \rangle = \quad (36)$$

$$\frac{g_{\mu\nu}^\perp}{4} \left(\frac{1 + \psi'}{2} \right)_{\beta\alpha} \left[\tilde{u}_1(x_+) + \tilde{U}_1(x_+, z_+) \right] - \frac{\epsilon_{\mu\nu}^\perp}{4} \left(\frac{1 + \psi'}{2} \not{n}_- \gamma_5 \frac{1 + \psi'}{2} \right)_{\beta\alpha} \tilde{U}_3(x_+, z_+).$$

The results are written in momentum space using the convention

$$\tilde{S}_i(x_{1+}, \dots, x_{n+}) = \int d\omega_1 \dots d\omega_n e^{-i(\omega_1 x_{1+} + \dots + \omega_n x_{n+})} S_i(\omega_1, \dots, \omega_n) \quad (37)$$

for the Fourier transform of a generic shape function S_i . We now introduce effective shape functions defined by the convolutions

$$\begin{aligned} u_s(u) &= \int d\omega_1 d\omega_2 J_2(u; \omega_1, \omega_2) [u_1(\omega_1) \delta(\omega_2) + U_1(\omega_1, \omega_2)], \\ u_a(u) &= \int d\omega_1 d\omega_2 J_2(u; \omega_1, \omega_2) U_3(\omega_1, \omega_2), \\ t_1(u) &= \int d\omega_1 J_2(u; \omega_1, 0) \left[\omega_1 S(\omega_1) - 2 \int d\omega' T_1(\omega_1, \omega') \right] \\ &+ 2 \int d\omega_1 d\omega_2 J_2(u; \omega_1, \omega_2) T_1(\omega_1, \omega_2); \end{aligned} \quad (38)$$

$$\begin{aligned} J_2(u; \omega_1, \omega_2) &= -\frac{1}{\pi} \text{Im} \frac{(n_+ p)^2}{(p_{\omega_1}^2 - m_c^2)(p_{\omega_{12}}^2 - m_c^2)} \\ &= \frac{1}{\omega_2} (\delta(u - \omega_1 - \omega_2) - \delta(u - \omega_1)), \end{aligned} \quad (39)$$

where $p_{\omega_{12\dots j}} = p - \omega_{12\dots j} n_+/2$ and $\omega_{12\dots j} = \omega_1 + \omega_2 + \dots + \omega_j$. J_2 is the tree-level jet function related to the product of propagators in (25).

The contributions to the hadronic tensor which do not vanish when $m_c \rightarrow 0$ can be obtained from eq. (59) of [12] by replacing $n_- p \rightarrow u$. The terms which vanish when $m_c \rightarrow 0$ are

$$\begin{aligned} W_m^{\mu\nu} &= \left(\frac{m_c}{n_+ p} \frac{1}{2} \text{tr} \left[\frac{1 + \psi'}{2} \bar{\Gamma} \Gamma \right] + \frac{m_c^2}{(n_+ p)^2} \frac{1}{2} \text{tr} \left[\frac{1 + \psi'}{2} \bar{\Gamma} \not{n}_+ \Gamma \right] \right) S(u) \\ &- \frac{m_c^2}{(n_+ p)^2} \frac{1}{2} \text{tr} \left[\frac{1 + \psi'}{2} \bar{\Gamma} \not{n}_- \Gamma \right] t_1(u). \end{aligned} \quad (40)$$

To proceed further, we use that the Dirac structures for the semi-leptonic decay in the Standard Model are

$$\bar{\Gamma} = \gamma^\mu(1 - \gamma_5), \quad \Gamma = \gamma^\nu(1 - \gamma_5). \quad (41)$$

The correction linear in m_c vanishes for this V-A structure. Performing the traces, and including now all terms, we find that the components of the hadronic tensor are given by

$$\begin{aligned} W_1 &= \frac{2}{n_+p} \left[\left(1 + \frac{u}{n_+p}\right) S(u) + \frac{s(u)}{2m_b} \right. \\ &\quad \left. - \frac{uS(u) - t(u)}{m_b} - \frac{u_s(u) + u_a(u)}{n_+p} - \frac{m_c^2}{(n_+p)^2} t_1(u) \right], \\ W_2 &= \frac{1}{2} W_3 = -\frac{2u}{n_+p} S(u), \\ W_4 &= -\frac{4}{(n_+p)^2} t(u), \\ W_5 &= \frac{8}{(n_+p)^2} \left[\frac{uS(u) - t(u)}{m_b} + \frac{t(u) + u_a(u)}{n_+p} \right]. \end{aligned} \quad (42)$$

The appearance of the effective shape function $t_1(u)$ within the tree approximation is specific to decay into charm quarks, as is seen by setting $m_c \rightarrow 0$. This shows that the effects of the final-state charm-quark mass are not merely kinematical, but introduce new non-perturbative structure into the factorization formula. This new structure is relevant only in the shape-function region. In Section 7 we will extrapolate our results to the OPE region by performing a moment expansion, and find that the moments of $t_1(u)$ are determined entirely by those of the leading order shape function $S(u)$, so that in this region the effects of the quark mass are purely kinematical.

6 Differential Decay Spectra

From the scalar components W_i of the hadronic tensor one can express the triply differential decay rate as [20]

$$\begin{aligned} &\frac{1}{12\Gamma_0} \frac{d^3\Gamma}{d(n_-\hat{p})d(n_+\hat{p})d\bar{x}} = \\ &(n_+\hat{p} - n_-\hat{p}) \left\{ (1 + \bar{x} - n_-\hat{p} - n_+\hat{p})(n_+\hat{p} + n_-\hat{p} - \bar{x} - n_+\hat{p}n_-\hat{p}) \frac{m_b^2}{2} W_1 \right. \\ &\quad \left. + (1 - n_+\hat{p} + n_+\hat{p}n_-\hat{p}) \frac{m_b}{2} W_2 + [\bar{x}(n_+\hat{p} - \bar{x}) - n_+\hat{p}n_-\hat{p}] \frac{m_b}{4} (W_3 + 2m_b W_4 + m_b^2 W_5) \right\}, \end{aligned} \quad (43)$$

where $\Gamma_0 = G_F^2 m_b^5 |V_{cb}|^2 / 192\pi^3$ and $\bar{x} = 1 - 2E_\ell/m_b$. Here and below hatted quantities are normalized to m_b , i.e. $\hat{p} = p/m_b$, $\hat{m}_c = m_c/m_b$. We have used that only W_1 contains

a leading order term in λ to expand the differential decay rate to $\mathcal{O}(\lambda^2)$. The integration range is given by

$$\frac{\hat{m}_c^2}{n_+\hat{p}} \leq n_-\hat{p} \leq \bar{x} \leq n_+\hat{p} \leq 1. \quad (44)$$

Any decay distribution can be obtained from (43) along with the result for the hadronic tensor in (42). Of special interest for semi-leptonic $b \rightarrow u$ transitions is the singly differential spectrum in the variable $P_+ = n_-p + (M_B - m_b)$, because of the possibilities to extract $|V_{ub}|$ [19]. An examination of (42) shows that $u = n_-p - m_c^2/n_+p$ is the natural partonic analog of P_+ for $b \rightarrow c$ decay, since this is what appears in the argument of the shape functions. (We leave the discussion of an appropriate hadronic variable to Section 8.) Changing variables and performing the integrals over $n_+\hat{p}$ and \bar{x} , we find

$$\begin{aligned} \frac{1}{\Gamma_0} \frac{d\Gamma}{du} = & \left(1 - \frac{14}{3} \frac{u}{m_b} - 8 \frac{m_c^2}{m_b^2} \right) S(u) + \frac{s(u)}{2m_b} - 4 \frac{m_c^2}{m_b^2} t_1(u) \\ & + \frac{1}{3m_b} [t(u) + u_a(u) - 5u_s(u)], \end{aligned} \quad (45)$$

where the allowed phase space is $0 \leq u \leq m_b - m_c^2/m_b$. This result agrees with [11] in the limit $m_c \rightarrow 0$.

A comment is in order concerning the limits of integration used in calculating (45). After changing variables, the integration region is given by

$$\hat{u} + \frac{\hat{m}_c^2}{n_+\hat{p}} \leq \bar{x} \leq n_+\hat{p}, \quad \frac{\hat{u}}{2} + \frac{1}{2} \sqrt{\hat{u}^2 + 4\hat{m}_c^2} \leq n_+\hat{p} \leq 1. \quad (46)$$

The collinear expansion is valid in the region $n_+\hat{p} \sim \mathcal{O}(1)$, and breaks down for the lower limit above, which corresponds to $n_+\hat{p} \sim \hat{m}_c \sim \mathcal{O}(\lambda)$. However, the doubly differential rate behaves like λ^2 in this region of phase space. Hence the contribution to the u spectrum from this region is in total of order λ^3 and does not affect our analysis.

We can also derive the lepton energy spectrum in the endpoint region, where an experimental cut limits the lepton-energy variable \bar{x} to values satisfying $\bar{x} \sim \mathcal{O}(\lambda^2)$. Such a cut restricts the other two kinematic variables to the shape-function region, since $\hat{m}_c^2/n_+\hat{p} \leq n_-\hat{p} \leq \bar{x}$ and $\hat{m}_c^2/(\bar{x} - \hat{u}) \leq n_+\hat{p} \leq 1$ ensure that $n_-\hat{p} \sim \mathcal{O}(\lambda^2)$ and $n_+\hat{p} \sim \mathcal{O}(1)$. Integrating over $n_+\hat{p}$, we find that the leading-order result for the doubly differential spectrum is

$$\frac{1}{\Gamma_0} \frac{d^2\Gamma}{d\bar{x}du} \approx 2S(u) \left[1 - \frac{3\hat{m}_c^4}{(\bar{x} - \hat{u})^2} + \frac{2\hat{m}_c^6}{(\bar{x} - \hat{u})^3} \right] + \mathcal{O}(\lambda^2). \quad (47)$$

We have also calculated the $\mathcal{O}(\lambda^2)$ power-suppressed terms but do not quote them here.

7 Comparison with the local expansion

A more familiar treatment of inclusive $b \rightarrow c$ decay uses the operator product expansion (OPE) to calculate the total rate [25]. Within this approach, the singly differential spec-

trum in the variable u at leading order in α_s and including power-suppressed corrections up to dimension 5 is given by

$$\begin{aligned} \frac{1}{\Gamma_0} \frac{d\Gamma}{du} = & \left(1 - 8 \frac{m_c^2}{m_b^2}\right) \delta(u) \\ & - \left(\frac{17}{18} \frac{\lambda_1}{m_b} + \frac{3}{2} \frac{\lambda_2}{m_b}\right) \delta'(u) + \lambda_1 \left(\frac{8m_c^2}{3m_b^2} - \frac{1}{6}\right) \delta''(u) + \mathcal{O}\left(\frac{1}{m_b^3}\right). \end{aligned} \quad (48)$$

We have not written terms proportional to $\lambda_{1,2} \delta(u)$, since these are Λ^2/m_b^2 corrections which cannot be recovered from our calculation in the shape-function region, which was carried out only to $\mathcal{O}(\Lambda/m_b, m_c^2/m_b^2)$. The OPE result is valid in the kinematical region where all momentum components are of order m_b . We can compare the results calculated in the OPE region with those derived in the shape-function region by considering $u \sim m_b$, in which case moments of the effective shape functions can be expressed in terms of local HQET matrix elements [1] (see also [10, 11, 26]). This expansion is given by

$$\begin{aligned} S(u) &= \delta(u) - \frac{\lambda_1}{6} \delta''(u), & s(u) &= -(\lambda_1 + 3\lambda_2) \delta'(u), \\ u_s(u) &= -\frac{2\lambda_1}{3} \delta'(u), & u_a(u) &= \lambda_2 \delta'(u), \\ t(u) &= -\lambda_2 \delta'(u), & t_1(u) &= -\frac{\lambda_1}{3} \delta''(u). \end{aligned} \quad (49)$$

The higher moments correspond to operators of higher than dimension 5 in the OPE and are neglected.

We will derive the moment expansion for $t_1(u)$ as an explicit example, since it is unique to decay into charm quarks. Examining the definition (38), we see that it is sufficient to derive the moments of the function $T_1(\omega_1, \omega_2)$, since those of $S(\omega)$ are known. At dimension 5, we only need to consider up to the first moment with respect to ω_1 or ω_2 . Moments of the form $\int d\omega_1 d\omega_2 T_1(\omega_1, \omega_2) \omega_1^n$ give a vanishing contribution to $t_1(u)$, as is seen by inserting $T_1(\omega_1, \omega_2) \sim \delta^{(n)}(\omega_1) \delta(\omega_2)$ into (38). This means that we only have to calculate the first moment with respect to ω_2 . To do this, we start with its definition in terms of a tri-local matrix element (see (32) and (35))

$$\frac{n_+^\mu}{2} \langle \bar{B} | (\bar{h}_v Y)(x) [Y^\dagger i D_\mu^s Y](z) (Y^\dagger h_v)(0) | \bar{B} \rangle = \int d\omega_1 d\omega_2 e^{-i\omega_1 x_+ - i\omega_2 z_+} T_1(\omega_1, \omega_2), \quad (50)$$

where the derivative does not act outside the square bracket. The first moment with respect to ω_2 is obtained by acting on both sides of (50) with $-in_- \partial_z$ and then setting x and z to zero to obtain a local HQET matrix element, which is calculated using

$$\langle \bar{B} | \bar{h}_v \Gamma_{\mu\nu} i D^\mu i D^\nu h_v | \bar{B} \rangle = \frac{1}{2} \text{tr} \left(\Gamma_{\mu\nu} \frac{1 + \not{v}}{2} \left[(g^{\mu\nu} - v^\mu v^\nu) \frac{\lambda_1}{3} + i\sigma^{\mu\nu} \frac{\lambda_2}{2} \right] \frac{1 + \not{v}}{2} \right). \quad (51)$$

We then find $\int d\omega_1 d\omega_2 T_1(\omega_1, \omega_2) \omega_2 = 0$, as can be seen by using the identity

$$-in_- \partial_z [Y^\dagger i D_\mu Y](z) = n_-^\nu [Y^\dagger i g F_{\mu\nu} Y](z), \quad (52)$$

and noting that the local HQET matrix element vanishes. We conclude that $T_1(\omega_1, \omega_2)$ is irrelevant for the moment expansion of $t_1(u)$. Inserting the moment expansion for $S(\omega)$ into (38), we find $t_1(u) = -\frac{\lambda_1}{3}\delta''(u) + \dots$, as quoted above.

Replacing the shape functions in (45) by the moments in (49), we find that the local expansion of the shape-function region result reproduces the OPE result (48). We performed the same comparison for the lepton energy spectrum (47), including power-suppressed terms, and confirmed that the moment expansion reproduces the OPE result [25], expanded appropriately for $\hat{m}_c^2 \sim \bar{x} \sim \lambda^2$.

8 Leading-Order Shape-Function Effects

Neglecting sub-leading shape functions and perturbative corrections, the $d\Gamma/du$ spectrum is directly proportional to the leading-order shape function $S(u)$. This raises the prospect of extracting information on the shape function from data on inclusive charm decays, a question which we will now address. Before performing any phenomenological studies we will slightly modify our treatment of m_c^2 corrections. In (45), we kept only terms up to m_c^2 , as is consistent with the SCET power counting. However, higher-order kinematic corrections involving the charm-quark mass arise from phase-space integrals and are easily included. Keeping these additional kinematic corrections, but neglecting all other sub-leading effects, the spectrum becomes

$$\frac{1}{\Gamma_0} \frac{d\Gamma}{du} = f(\rho)S(u), \quad (53)$$

where $f(\rho)$ is the usual phase-space function

$$f(\rho) = (1 - 8\rho + 8\rho^3 - \rho^4 - 12\rho^2 \ln \rho), \quad (54)$$

and we have defined $\rho = m_c^2/m_b^2$. The numerical effect of the phase-space function is sizable. In particular, using $f(\rho) \sim (1 - 8\rho)$ would differ at $\rho \sim 0.1$ significantly from the value obtained from (54). However, once the logarithmic contribution is included, the numerical result is almost exact.

In an experimental analysis one has to rewrite (53) in terms of hadronic variables. To this end, we propose to study the spectrum in the variable

$$U = n_- P - \frac{M_D^2}{n_+ P}, \quad (55)$$

where $n_\pm P = n_\pm p + (M_B - m_b) = n_\pm p + \bar{\Lambda}$ are the hadronic light-cone variables. Since U depends on $n_+ P$ one must make this replacement before integrating over $n_+ P$, in which case the integration region is

$$\frac{U}{2} + \frac{1}{2}\sqrt{U^2 + 4M_D^2} \leq n_+ P \leq M_B, \quad 0 \leq U \leq M_B - \frac{M_D^2}{M_B} \quad (56)$$

One has to be careful when writing the spectrum in terms of U because

$$\begin{aligned} u &= U - \bar{\Lambda} + \frac{M_D^2}{n_+ P} - \frac{m_c^2}{n_+ P - \bar{\Lambda}} \\ &\approx U - \bar{\Lambda} + \frac{M_D^2 - m_c^2}{n_+ P} - \frac{m_c^2}{n_+ P} \frac{\bar{\Lambda}}{n_+ P} + \dots \end{aligned} \quad (57)$$

The last two terms are of order λ^3 and λ^4 respectively as long as $n_+ P \sim M_B$, but become order λ^2 at the lower limit of the $n_+ P$ integration, when $n_+ P \sim M_D$. However, one can show that the contribution of this region to the U spectrum is power-suppressed, so we can ignore this subtlety and write $u = U - \bar{\Lambda}$. The shape function is then a function of $S(U - \bar{\Lambda})$, and the decay distribution is simply

$$\frac{1}{\Gamma_0} \frac{d\Gamma}{dU} = f(\rho) S(U - \bar{\Lambda}). \quad (58)$$

At leading order $\rho = m_c^2/m_b^2 \approx M_D^2/M_B^2$. Eq. (58) opens the possibility to at least cross-check the results for the shape functions obtained from heavy-to-light decays. However, a few comments are in order.

Unlike in the case of heavy-to-light semi-leptonic decays, almost 80% of the total inclusive rate is already exhausted by the two exclusive decays $\bar{B} \rightarrow D\ell\bar{\nu}_\ell$ and $\bar{B} \rightarrow D^*\ell\bar{\nu}_\ell$. This may indicate that the real world is not very close to the limit $\lambda \rightarrow 0$ using the power counting we are suggesting. For the exclusive decay $\bar{B} \rightarrow D\ell\bar{\nu}_\ell$ the U spectrum is concentrated at $U = 0$

$$\frac{d\Gamma(\bar{B} \rightarrow D\ell\bar{\nu}_\ell)}{dU} = \Gamma(\bar{B} \rightarrow D\ell\bar{\nu}_\ell) \delta(U) \quad (59)$$

while for the exclusive decay $\bar{B} \rightarrow D^*\ell\bar{\nu}_\ell$ the decay rate is a function of

$$U = \frac{M_{D^*}^2 - M_D^2}{n_+ P} \sim \mathcal{O}\left(\frac{\lambda^4 M_B^2}{n_+ P}\right).$$

Even when $n_+ P \sim M_D \sim \lambda M_B$ we have $U \sim \lambda^3 M_B$ and this is beyond the sensitivity of the leading-order approach, since the smearing due to the shape function is of the order λ^2 . Thus the contribution of both ground states may be written as

$$\frac{d\Gamma(\bar{B} \rightarrow D\ell\bar{\nu}_\ell)}{dU} + \frac{d\Gamma(\bar{B} \rightarrow D^*\ell\bar{\nu}_\ell)}{dU} = [\Gamma(\bar{B} \rightarrow D\ell\bar{\nu}_\ell) + \Gamma(\bar{B} \rightarrow D^*\ell\bar{\nu}_\ell)] \delta(U). \quad (60)$$

In terms of a moment expansion of the decay rate, we see that these two exclusive decay modes contribute only to the zeroth moment, and conclude that all higher moments of the U spectrum can originate only from excited or non-resonant states. For this reason, it would be interesting to see if and how well the relations for the moments, such as

$$\int dU U \frac{1}{\Gamma_0} \frac{d\Gamma}{dU} = \bar{\Lambda} f(\rho), \quad \int dU U^2 \frac{1}{\Gamma_0} \frac{d\Gamma}{dU} = \left(-\frac{1}{3}\lambda_1 + \bar{\Lambda}^2\right) f(\rho) \quad (61)$$

are satisfied. In particular, we find that to leading order in the SCET expansion the ratio of moments should be the same as the corresponding ratio of P_+ moments in $\bar{B} \rightarrow X_u \ell \bar{\nu}_\ell$ or of photon energy moments in $\bar{B} \rightarrow X_s \gamma$.

9 Conclusions

In this work we examined inclusive semi-leptonic $b \rightarrow c$ decay using the power counting $m_c \sim \sqrt{\Lambda_{\text{QCD}} m_b}$ for the charm-quark mass. With this power counting the decay kinematics can be chosen to access the shape-function region even in the decay $\bar{B} \rightarrow X_c \ell \bar{\nu}_\ell$. We applied effective field theory methods by modifying SCET to include a massive collinear charm quark. This defines a consistent power-counting scheme in terms of the small parameter $\lambda \sim m_c/m_b \sim \sqrt{\Lambda_{\text{QCD}}/m_b}$. We matched the Lagrangian as well as the weak transition current at tree level and including sub-leading terms up to $\mathcal{O}(\lambda^2)$, and used these to derive the hadronic tensor in the shape-function region.

The results are similar to those for $b \rightarrow u$ transitions. This led us to identify a certain variable which is analogous to the variable P_+ in $b \rightarrow u$ decay. As in the heavy-to-light case, at leading power and at tree level the singly differential spectrum in this variable is directly proportional to the leading-order shape function. We showed how the moment expansion of this differential spectrum matches the local OPE result in an appropriate limit.

It remains to be checked whether the power counting we have applied for the charm mass yields a consistent picture or whether the conventional counting $m_c \sim m_b$ is more appropriate. A crosscheck can be performed by measuring the moments of the distribution in this variable and by comparing them to those obtained from the P_+ spectrum in $\bar{B} \rightarrow X_u \ell \bar{\nu}_\ell$ and from the E_γ spectrum in $\bar{B} \rightarrow X_s \gamma$. Our leading-order prediction is that these moments should be equal up to the phase-space factor which appears in the total partonic rates.

Acknowledgements

This work was supported by the DFG Sonderforschungsbereich SFB/TR09 “Computational Theoretical Particle Physics”.

References

- [1] M. Neubert, Phys. Rev. D **49**, 3392 (1994) [hep-ph/9311325].
- [2] I. I. Y. Bigi, M. A. Shifman, N. G. Uraltsev and A. I. Vainshtein, Int. J. Mod. Phys. A **9**, 2467 (1994) [hep-ph/9312359].
- [3] C. W. Bauer, S. Fleming and M. E. Luke, Phys. Rev. D **63**, 014006 (2001) [hep-ph/0005275].
- [4] C. W. Bauer, S. Fleming, D. Pirjol and I. W. Stewart, Phys. Rev. D **63** (2001) 114020 [hep-ph/0011336].
- [5] C. W. Bauer and I. W. Stewart, Phys. Lett. B **516**, 134 (2001) [hep-ph/0107001].

- [6] M. Beneke, A. P. Chapovsky, M. Diehl and T. Feldmann, Nucl. Phys. B **643**, 431 (2002) [hep-ph/0206152].
- [7] C. W. Bauer, D. Pirjol and I. W. Stewart, Phys. Rev. D **65**, 054022 (2002) [hep-ph/0109045].
- [8] C. W. Bauer and A. V. Manohar, Phys. Rev. D **70**, 034024 (2004) [hep-ph/0312109].
- [9] S. W. Bosch, B. O. Lange, M. Neubert and G. Paz, Nucl. Phys. B **699**, 335 (2004) [hep-ph/0402094];
- [10] K. S. M. Lee and I. W. Stewart, [hep-ph/0409045].
- [11] S. W. Bosch, M. Neubert and G. Paz, JHEP **0411**, 073 (2004) [hep-ph/0409115].
- [12] M. Beneke, F. Campanario, T. Mannel and B. D. Pecjak, [hep-ph/0411395].
- [13] F. J. Tackmann, [hep-ph/0503095].
- [14] T. Mannel and M. Neubert, Phys. Rev. D **50**, 2037 (1994) [hep-ph/9402288].
- [15] T. Mannel and F. J. Tackmann, Phys. Rev. D **71**, 034017 (2005) [hep-ph/0408273].
- [16] I. Z. Rothstein, Phys. Rev. D **70** (2004) 054024 [hep-ph/0301240].
- [17] A. K. Leibovich, Z. Ligeti and M. B. Wise, Phys. Lett. B **564**, 231 (2003) [hep-ph/0303099].
- [18] T. Mannel and S. Recksiegel, Phys. Rev. D **60**, 114040 (1999) [hep-ph/9904475].
- [19] S. W. Bosch, B. O. Lange, M. Neubert and G. Paz, Phys. Rev. Lett. **93**, 221801 (2004) [hep-ph/0403223].
- [20] F. De Fazio and M. Neubert, JHEP **9906**, 017 (1999) [hep-ph/9905351].
- [21] G. P. Korchemsky and G. Sterman, Phys. Lett. B **340** (1994) 96 [hep-ph/9407344].
- [22] M. Beneke and T. Feldmann, Phys. Lett. B **553**, 267 (2003) [hep-ph/0211358].
- [23] J. Chay and C. Kim, Phys. Rev. D **65** (2002) 114016 [hep-ph/0201197].
- [24] D. Pirjol and I. W. Stewart, Phys. Rev. D **67**, 094005 (2003) [Erratum-ibid. D **69**, 019903 (2004)] [hep-ph/0211251].
- [25] A. V. Manohar and M. B. Wise, Phys. Rev. D **49**, 1310 (1994) [hep-ph/9308246].
- [26] C. W. Bauer, M. E. Luke and T. Mannel, Phys. Rev. D **68** (2003) 094001 [hep-ph/0102089].