

DAMTP/R-94/7

LA-UR-93-4473

March 1994

THE INSTABILITY OF CHARGED BLACK STRINGS AND p-BRANES

Ruth Gregory

*Department of Applied Mathematics and Theoretical Physics
University of Cambridge, Silver St, Cambridge, CB3 9EW, U.K.*

Raymond Laflamme

*Theoretical Astrophysics, T-6, MSB288, Los Alamos National Laboratory
Los Alamos, NM 87545, USA*

ABSTRACT

We investigate the evolution of small perturbations around charged black strings and branes which are solutions of low energy string theory. We give the details of the analysis for the uncharged case which was summarized in a previous paper. We extend the analysis to the small charge case and give also an analysis for the generic case, following the behavior of unstable modes as the charge is modified. We study specifically a magnetically charged black 6-brane, but show how the instability is generic, and that charge does not in general stabilise black strings and p-branes.

1) Introduction

Black holes have always been a source of fascination and debate, from the nature of the event horizon to the nature of the singularity and whether a full theory of quantum gravity can avert the apparent loss of unitarity in black hole evaporation. Classically, four dimensional black holes are stable, once formed they settle down to a state described solely by their mass, charge and angular momentum; therefore the singularities remain hidden from distant observers. Quantum mechanically, black holes are quite different objects, analogous to a thermal system. They have an entropy (proportional to surface area) and a temperature, since like black bodies they radiate thermally¹. However, Hawking conjectured² that a black hole formed from a pure quantum state would radiate away leaving a mixed state of radiation; this would violate quantum mechanical unitarity. It is certainly true that Hawking's semi-classical description will break down near the final stage of black hole evaporation as planckian curvatures are approached, but if quantum gravity preserves unitarity and information is to be returned, it must do so well before this stage otherwise there is simply not enough energy left in a planck mass black hole to emit all the information stored in a macroscopic black hole. Therefore, in order to resolve this tension between quantum mechanics and thermodynamics, it seems that a semi-classical analysis of sub-Planck size black holes is needed.

Most recently, attention has focussed on low energy string gravity and its implications for black holes. Some of these developments have been quite interesting. In Einstein gravity, charged black holes (the Reissner-Nordström solutions), in addition to having an outer event horizon, have an inner Cauchy horizon which is unstable to matter perturbations in the exterior spacetime³. However, there is no static charged black hole solution in Einstein gravity with only one horizon and a spacelike singularity. On the other hand, a key feature of low energy string gravity is the presence of a dilaton, which greatly changes the causal structure of charged black holes making them like Schwarzschild with one event horizon and a spacelike singularity⁴. This structure is generic, even if the dilaton has a mass⁵, as it must to keep in line with the principle of equivalence.

Of course, most of the analysis of stringy black holes has been performed in low dimensions, namely two or four, whereas string theory tells us there should be ten dimensions,

ideally therefore, one should be examining black holes in ten dimensions. There has been work on black holes in higher dimensions⁶ for a range of horizon topologies⁷. In four dimensions, an event horizon must be topologically spherical⁸, but in higher dimensions this is not necessarily the case, we could have $S^2 \times \mathbb{R}^6$, or $S^3 \times \mathbb{R}^5$ topologies for the horizon. In a previous letter we pointed out that a large class of these black holes are unstable, namely the uncharged ones. In this paper, we present the details of this original argument, as well as providing an extension to cover charged black p-branes.

Why should black holes be stable, yet black strings, say, unstable? Before answering this question in detail, it is worth examining a couple of naive arguments. Without loss of generality, consider a five dimensional black string in Einstein gravity, $\text{Sch}_4 \times \mathbb{R}$. Then the equation governing the metric perturbation $\delta g_{ab} = h_{ab}$, the Lichnerowicz equation, is essentially a wave equation

$$\Delta_L h_{ab} = (\delta_a^c \delta_b^d \square + 2R_a^c{}^d) h_{cd} = 0. \quad (1.1)$$

Because of the symmetries of the background $\text{Sch}_4 \times \mathbb{R}$ metric, this reduces to a four dimensional Lichnerowicz operator plus a ∂_z^2 piece. Performing a fourier decomposition of h_{ab} in the fifth dimension yields

$$\Delta_L h_{ab} = (\Delta_4 - m^2) h_{ab} = 0. \quad (1.2)$$

Since the four dimensional Schwarzschild Lichnerowicz operator has no unstable modes, adding a mass should only increase stability, hence it has been conjectured black strings are stable.

On the other hand, glibly speaking, horizons are like soap bubbles, they have a surface tension, κ - the surface gravity, and soap bubbles do not like being cylindrical! More formally, a portion of horizon of length L contains mass $\mathcal{M} = ML$ and has an entropy proportional to \mathcal{M}^2/L . A five dimensional black hole on the other hand has entropy proportional to $\mathcal{M}^{3/2}$. Thus for large lengths of horizon, the mass contained within the horizon contributes a much lower entropy than if it were in a hyperspherical black hole. This indicates that for large wavelength perturbations in the fifth dimension, we might expect an instability.

As it turns out, the thermodynamic argument is correct. The flaw in the other argument is to assume that the number of physical degrees of freedom of the effective four dimensional tensor gauge field, h_{ab} , remains the same as in Einstein theory. Clearly this is not so, for the fifth dimension adds an effective mass to h_{ab} . A massive tensor field in four dimensions has three degrees of freedom as compared with only two for a massless one. We will show that this extra degree of freedom has a spherically symmetric mode which is responsible for the instability.

The stability of the five dimensional black string was investigated analytically⁹, with the conclusion that there is no non-singular single unstable mode on a Schwarzschild time $t = 0$ surface, however, this argument did not *prove* stability. As emphasized by Vishveshwara¹⁰ in his original Schwarzschild stability argument, the non-existence of a single unstable mode does not preclude the existence of a composite unstable mode, with the combination cancelling the singular behaviour of an inadmissible single singular mode. An example of this sort of situation occurs in the coloured black hole instability, recently confirmed by Bizon and Wald¹¹. That this was probably the case for five dimensional black strings was first pointed out by Whitt¹², who analyzed the stability of Schwarzschild in four dimensional fourth order gravity - a different physical situation, but mathematically identical equations to those studied in ref [9]. The key difference in Whitt's analysis was the choice of an initial data surface ending on the future horizon (see Figure 1.). By avoiding the neck of the Schwarzschild wormhole, one avoids the fixed point of the isometries used to generate the mode decomposition, which avoids in this case issues of superposition. In any case, such a choice of initial data surface is mandated in any real physical scenario, since a black hole or brane must form from collapse, and hence will not possess a Schwarzschild wormhole.

In a previous letter¹³, we showed that using an appropriate initial data surface, uncharged black p -branes were unstable, and conjectured that the same would hold true for charged black p -branes. In this paper, we extend the analysis of [13] to include charge, and show that the instability does indeed persist. It is worth stressing that this instability is not of the Reissner-Nordström form - hidden behind the event horizon, but it is a real *physical* instability of the exterior spacetime which could potentially fragment the horizon.

It is important to emphasize that this can occur classically, for although under regular conditions horizons do not bifurcate¹⁴, if one has a naked singularity, then bifurcation is possible. Since an instability calculation is by its nature linear, it cannot predict the endpoint of an unstable evolution. However, the entropy argument does lend support to the fragmentation scenario and violation of the Cosmic Censorship Hypothesis.

The layout of the paper is as follows. In section two we summarise the black p-branes we are investigating, namely those of Horowitz and Strominger⁷, and derive the general perturbation equations. We discuss gauge constraints and the issue of boundary conditions in detail, then we set up the mode decomposition for our analysis. In section three we specialise to zero charge and show how this greatly simplifies the problem, and present an instability. We then use this ‘zeroth order’ solution to generate an instability for small charge in section four. In section five we present the general case, for arbitrary charge, and present our conclusions in section six.

2) The perturbation problem.

We start this section on the derivation of the perturbation equations by reviewing the particular black p-branes we will be investigating. We focus on solutions of low energy string gravity with symmetry $\mathbb{R}^{10-D} \times \mathbb{R}_t \times S^{D-2}$ and a possible ‘magnetic’ charge. The action for such a theory is

$$\int d^{10}x \sqrt{-g} e^{-2\phi} [R + 4(\nabla\phi)^2 - \frac{2}{(D-2)!} F^2] \quad (2.1)$$

where F is a $(D-2)$ -form field strength. Varying this action yields the equations of motion

$$\square\phi - 2(\nabla\phi)^2 + F^2 \frac{(D-3)}{(D-2)!} = 0 \quad (2.2a)$$

$$\nabla_{a_1} [e^{-2\phi} F^{a_1 \dots a_{D-2}}] = 0 \quad (2.2b)$$

$$R_{ab} + 2\nabla_a \nabla_b \phi - \frac{2}{(D-3)!} F_{aa_2 \dots a_{D-2}} F_b^{a_2 \dots a_{D-2}} = 0 \quad (2.2c)$$

Searching for a solution with the required symmetry indicates a metric of the form

$$ds^2 = -e^A dt^2 + e^F dr^2 + e^B dx_i dx^i + C^2 d\Omega_{D-2}^2 \quad (2.3)$$

where all metric functions are functions of the radial variable only, and the index i runs from $D+1$ to 10, and represents the coordinates in the p-plane of symmetry of the p-brane. Looking for a ‘magnetically’ charged brane,

$$F = Q\epsilon_{D-2} \quad (2.4)$$

where ϵ_{D-2} is the area form on a unit $(D-2)$ -sphere, it is straightforward to show that a solution exists and takes the form⁷

$$e^A = \frac{1 - (\frac{r_+}{r})^{D-3}}{1 - (\frac{r_-}{r})^{D-3}} \quad , \quad e^{-F} = \left(1 - (\frac{r_+}{r})^{D-3}\right) \left(1 - (\frac{r_-}{r})^{D-3}\right) \quad , \quad B = 0 \quad (2.5)$$

$$C(r) = r \quad , \quad e^{-2\phi} = 1 - (\frac{r_-}{r})^{D-3} \quad , \quad Q^2 = \frac{D-3}{2}(r_+r_-)^{D-3}$$

If $r_- = 0$, then $Q^2 = \phi = 0$, and the solution is the uncharged $\text{Sch}_D \times \mathbb{R}^{10-D}$, which we have argued to be unstable. However, in the presence of charge there is also a non-trivial dilaton field, and it is unclear how this will affect the physical instability. We first vary the equations of motion to obtain the bare perturbation equations, then discuss boundary conditions on our perturbations in terms of a non-singular (or generalised Kruskal) coordinate system on the event horizon, then return to the perturbation equations, discussing gauge choices to simplify these, finally discussing an additional simplification to the electromagnetic perturbation.

In order to write the perturbation equations, we use the usual notation

$$\delta g_{ab} = h_{ab} \quad (2.6)$$

and

$$\bar{h}_{ab} = h_{ab} - \frac{1}{2}h g_{ab} \quad (2.7)$$

varying the equations of motion can be seen to give

$$\begin{aligned} \nabla_{a_1}[e^{-2\phi}(\delta F)^{a_1 \dots a_{D-2}}] - 2(\nabla_{a_1} \delta \phi)e^{-2\phi}F^{a_1 \dots a_{D-2}} - (D-3)e^{-2\phi}F^{a_1 b [\dots a_{D-2} \nabla_{a_1} h_b^{a_2}]} \\ - h^{a_1 b} \nabla_{a_1}[e^{-2\phi}F_b^{a_2 \dots a_{D-2}}] - \nabla_{a_1}(\bar{h}_b^{a_1})e^{-2\phi}F^{b \dots a_{D-2}} = 0 \end{aligned} \quad (2.8a)$$

$$\square \delta \phi - h^{ab} \nabla_a \nabla_b \phi + 2h^{ab} \nabla_a \phi \nabla_b \phi - \nabla_c \phi \nabla_b \bar{h}^{bc} - 4 \nabla_a \delta \phi \nabla^a \phi$$

$$-\frac{h^{cd}}{(D-4)!}F_{ca_2\dots a_{D-2}}F_d^{a_2\dots a_{D-2}}+2\frac{(D-3)}{(D-2)!}\delta F_{a_1\dots a_{D-2}}F^{a_1\dots a_{D-2}}=0 \quad (2.8b)$$

$$\begin{aligned} &\square h_{ab}+2R_{cadb}h^{cd}-2R_{e(a}h_{b)}^e-2\nabla_{(a}\nabla_{|e|}\bar{h}_{b)}^e-4\nabla_a\nabla_b\delta\phi-2\nabla_c\phi\nabla^c h_{ab}+4\nabla_c\phi\nabla_{(b}h_{a)}^c \\ &+\frac{4}{(D-3)!}\left[2F_{(a}^{a_2\dots a_{D-2}}\delta F_{b)a_2\dots a_{D-2}}-(D-3)h^{cd}F_{aca_3\dots a_{D-2}}F_{bd}^{a_3\dots a_{D-2}}\right]=0 \quad (2.8c) \end{aligned}$$

These are the bare perturbation equations, we must now examine boundary conditions, and whether there is any choice of gauge which simplifies (2.8).

Boundary conditions.

Now we turn to the question of boundary conditions, which are a key to this problem. Obviously, we want to place initial data on a Cauchy surface for the exterior spacetime, but such a surface necessarily touches the horizon, which is singular in Schwarzschild coordinates. There are therefore two issues here: One is how to define ‘small’ for the perturbation at the horizon, and secondly, which initial data surface to impose these constraints upon.

The first issue is straightforwardly dealt with. Although the horizon is singular in Schwarzschild coordinates, it is not a physical singularity, merely a coordinate singularity. In four dimensions, non-singular coordinates have been known for some time - Kruskal coordinates. In order to generalise these to higher dimensions, the starting point is to identify the generalised tortoise coordinate

$$\begin{aligned} r_D^* &= \int e^{(F-A)/2} dr = \int \frac{r^{D-3}}{(r^{D-3}-r_+^{D-3})} dr = \sum_{n=1}^{D-3} \frac{1}{D-3} \int \frac{dr}{(1-e^{\frac{2\pi i n}{D-3}} r_+/r)} \\ &= r + \sum_{n=1}^{D-3} \frac{e^{\frac{2\pi i n}{D-3}}}{(D-3)} r_+ \log(r - e^{\frac{2\pi i n}{D-3}} r_+) \quad (2.9) \end{aligned}$$

So, for example,

$$\begin{aligned} r_4^* &= r + r_+ \log(r - r_+) \\ r_5^* &= r + \frac{1}{2} r_+ \log\left(\frac{r-r_+}{r+r_+}\right) \\ r_6^* &= r + \frac{1}{3} r_+ \log(r - r_+) - \frac{r_+}{6} \log(r^2 + rr_+ + r_+^2) + \frac{r_+}{\sqrt{3}} \tan^{-1}\left(\frac{\sqrt{3}r_+}{2r+r_+}\right) \end{aligned} \quad (2.10)$$

etc. In particular, as $r \rightarrow r_+$

$$r_D^* \sim \frac{r_+}{D-3} \log(r - r_+) \quad (2.11)$$

Having the tortoise coordinate, we can now define the Kruskal null coordinates

$$\begin{aligned} P_+ &= \exp\{(D-3)(t+r^*)/2r_+\} \\ P_- &= -\exp\{-(D-3)(t-r^*)/2r_+\} \end{aligned} \quad (2.12)$$

In terms of which the radial-time part of the metric looks like

$$ds_2^2 = \frac{4r_+^2(D-3)^2e^A}{P_+P_-}dP_+dP_- \quad (2.13)$$

which is finite as $r \rightarrow r_+$.

Finally, we set

$$\begin{aligned} R &= P_+ - P_- \\ T &= P_+ + P_- \end{aligned} \quad (2.14)$$

in terms of which

$$ds_2^2 = -\frac{r_+^2(D-3)^2e^A}{P_+P_-}(dR^2 - dT^2) = f^2(r)(dR^2 - dT^2) \quad (2.15)$$

in order to readily identify our initial data surface.

This leaves us with the problem of an initial data surface. The domain of dependence must obviously include \mathcal{I}^+ , thus a surface touching the future horizon, or the neck of the Schwarzschild wormhole is acceptable, but a surface touching the past horizon is not, unless it passes through and extends to the opposite horizon on the Penrose diagram. Following Whitt, we impose regularity in the Kruskal system as $r \rightarrow r_+$ at constant non-zero T , i.e., $R \rightarrow T$ from above as depicted on Figure 1. This avoids the issue of mode superposition discussed earlier, and secondly, we believe it to be a better physically motivated choice of surface. This is because in practice a black hole (or brane) would form in a collapse situation, and hence would not have a Schwarzschild wormhole; analyzing stability would necessarily require a surface ending on a future event horizon.

In terms of the Schwarzschild components

$$h^{TT} = -\frac{(D-3)^4}{f^2(r)(T^2 - R^2)} \left[R^2 h^{\hat{t}\hat{t}} + 2RT h^{\hat{t}\hat{r}} + T^2 h^{\hat{r}\hat{r}} \right] \quad (2.16a)$$

$$h^{RR} = -\frac{(D-3)^4}{f^2(r)(T^2-R^2)} \left[T^2 h^{\hat{t}\hat{t}} + 2RT h^{\hat{t}\hat{r}} + R^2 h^{\hat{r}\hat{r}} \right] \quad (2.16b)$$

$$h^{TR} = -\frac{(D-3)^4}{f^2(r)(T^2-R^2)} \left[RT(h^{\hat{t}\hat{t}} + h^{\hat{r}\hat{r}}) + (T^2 + R^2)h^{\hat{t}\hat{r}} \right] \quad (2.16c)$$

$$h^{T*} = \frac{(D-3)^2}{f(r)\sqrt{(R^2-T^2)}} \left[Rh^{\hat{t}*} + Th^{\hat{r}*} \right] \quad (2.16d)$$

$$h^{R*} = \frac{(D-3)^2}{f(r)\sqrt{(R^2-T^2)}} \left[Th^{\hat{t}*} + Rh^{\hat{r}*} \right] \quad (2.16e)$$

all of which should be finite as $r \rightarrow r_+$ on our initial data surface. Since $R = T + O(r - r_+)$ as $R \rightarrow T$ from above, this implies that the various combinations of the normalised $h^{\hat{a}\hat{b}}$ within the square brackets above must be $O(r - r_+)$ as we approach the event horizon. Now we turn to simplifying the perturbation equations by a judicious choice of gauge.

Gauge considerations.

In gravity, physics must be invariant under a re-labelling of coordinates. Such general coordinate transformations (gct's), are generated by vector fields ξ^a , the effect of an infinitesimal gct being to push the coordinates an infinitesimal amount along the integral curves of ξ^a , such that $x^a \rightarrow x^a + \xi^a$. Under such a gauge transformation, physical quantities transform as

$$\mathcal{P} \rightarrow \mathcal{L}_\xi \mathcal{P} \quad (2.17)$$

where \mathcal{L}_ξ denotes the Lie derivative with respect to ξ . Therefore a pure gauge perturbation is of the form

$$\begin{aligned} h_{\xi ab} &= 2\xi_{(a;b)} \\ \delta\phi &= \xi^a \nabla_a \phi \\ \delta F_{a_1 \dots a_{D-2}} &= F_{a_1 \dots a_{D-2};d} \xi^d + (D-2) F_{d[a_2 \dots a_{D-2}} \xi_{a_1]}^d \end{aligned} \quad (2.18)$$

It is straightforward to verify that such a pure gauge perturbation satisfies our generalised Lichnerowicz perturbation equations.

Using such a pure gauge perturbation, we see that

$$\nabla_a \bar{h}_\xi^{ab} = R_a^b \xi^a + \square \xi^b \quad (2.19)$$

However, the right hand side of this equation is the curved space wave operator, which given a source and initial conditions can be integrated to give a solution for ξ^a throughout the manifold¹⁵. Thus, if the divergence of \bar{h}_{ab} is not initially zero, we can change gauge in order to make it so. This simplifies (2.8) to the following:

$$\begin{aligned} \nabla_{a_1}[e^{-2\phi}(\delta F)^{a_1\dots a_{D-2}}] - 2(\nabla_{a_1}\delta\phi)e^{-2\phi}F^{a_1\dots a_{D-2}} - (D-3)e^{-2\phi}F^{a_1b[\dots a_{D-2}}\nabla_{a_1}h_b^{a_2}] \\ - h^{a_1b}\nabla_{a_1}[e^{-2\phi}F_b^{a_2\dots a_{D-2}}] = 0 \end{aligned} \quad (2.20a)$$

$$\begin{aligned} \square\delta\phi - h^{ab}\nabla_a\nabla_b\phi + 2h^{ab}\nabla_a\phi\nabla_b\phi - 4\nabla_a\delta\phi\nabla^a\phi \\ - \frac{h^{cd}}{(D-4)!}F^{ca_2\dots a_{D-2}}F_d^{a_2\dots a_{D-2}} + 2\frac{(D-3)}{(D-2)!}\delta F_{a_1\dots a_{D-2}}F^{a_1\dots a_{D-2}} = 0 \end{aligned} \quad (2.20b)$$

$$\begin{aligned} \square h_{ab} + 2R_{caab}h^{cd} - 2R_{e(a}h_{b)}^e - 4\nabla_a\nabla_b\delta\phi - 2\nabla_c\phi\nabla^c h_{ab} + 4\nabla_c\phi\nabla_{(b}h_{a)}^c \\ + \frac{4}{(D-3)!}\left[2F_{(a}^{a_2\dots a_{D-2}}\delta F_{b)a_2\dots a_{D-2}} - (D-3)h^{cd}F_{aca_3\dots a_{D-2}}F_{bd}^{a_3\dots a_{D-2}}\right] = 0 \end{aligned} \quad (2.20c)$$

The residual gauge freedom are those gct's which satisfy

$$R_a^b\xi^a + \square\xi^b = 0 \quad (2.21)$$

Notice that in a vacuum spacetime, i.e. no dilaton, no ‘electromagnetism’, the trace of the metric perturbation equation becomes

$$\square h = 0 \quad (2.22)$$

and we can choose an harmonic ξ^a to set $h = 0$. Therefore *in vacuo* we may make the additional gauge choice of trace-free. However, more generally it should be noted that transversality is all that can be assumed.

Since a pure gauge perturbation automatically satisfies the perturbation equations, deciding whether a putative instability is physical or not reduces to investigating whether it can be expressed in this form. Alternatively, one can identify physical perturbations by using gauge invariant variables. It is straightforward to adopt Bardeen’s results¹⁶ to the case at hand. The idea is to choose a particular linear combination of variables and their derivatives such that the result becomes independent of ξ^a . The details and a particular set of gauge independent variables are given in Appendix A.

Finally, before beginning the analysis of the equations in detail we will make a further simplification by showing that the ‘electromagnetic’ perturbation, δF_{ab} , can be taken to be zero. In order to perform a stability analysis we will Fourier decompose all the perturbations in terms of the symmetries of the background spacetime. Since Kruskal coordinates are not well suited to such a decomposition, we perform our analysis in Schwarzschild coordinates, transforming to Kruskal at the horizon to check boundary conditions according to (2.16). The Fourier modes in the time and p -brane directions are easy to identify, for an instability they are of the form $e^{\Omega t + i\mu_i x^i}$. The spherical harmonic modes will depend on the number of dimensions, D , that the black hole sits in, as well as on the tensorial nature (scalar, vector, and so on) of the perturbation being analysed. However, since in general higher angular momentum modes are more stable, and since our advertised instability is an s-mode, zero angular momentum modes are all we shall be considering. This clearly means we can make no stability claims if we fail to find an instability, however, since our primary concern is to chart regions of instability, an s-wave analysis will suffice.

Therefore, the spherically symmetric perturbations take the form:

$$\delta\phi = e^{\Omega t + i\mu_i x^i} f(r) \tag{2.23a}$$

$$\delta F = e^{\Omega t + i\mu_i x^i} \begin{cases} q(r)\epsilon_{D-2} & \text{angular components} \\ 0 & \text{mixed angular / t-r-i components} \\ \delta F_T & \text{t-r-i components} \end{cases} \tag{2.23b}$$

$$h^{ab} = e^{\Omega t + i\mu_i x^i} \begin{bmatrix} H^{ij}(r) & H^{it} & H^{ir} & 0 & 0 & \dots \\ H^{tj} & H^{tt} & H^{tr} & 0 & 0 & \dots \\ H^{rj} & H^{tr} & H^{rr} & 0 & 0 & \dots \\ 0 & 0 & 0 & K(r) & 0 & \dots \\ 0 & 0 & 0 & 0 & K/\sin^2\theta & \dots \\ \dots & \dots & \dots & \dots & \dots & \dots \end{bmatrix} \tag{2.23c}$$

Where δF_T can be non-zero only if $D - 2 \leq 5$.

We may now see that the F -perturbation equation, (2.20b), becomes simplified, for noting that the background field $F = Q\epsilon_{D-2}$ has only angular components, and that $h_\beta^\alpha = K(r)\delta_\beta^\alpha/r^2$ for angular components,

$$F^{a_1 b [\dots a_{D-2} \nabla_{a_1} h_b^{a_2}] = F^{a_1 b [\dots a_{D-2} h_{b, a_1}^{a_2}] + F^{a_1 b [\dots a_{D-2} \Gamma_{ca_1}^{a_2}] h_b^c = 0$$

and

$$\nabla_{r/t}[e^{-2\phi}F_{r/t}^{a_2\dots a_{D-2}}] = 0 \Rightarrow h^{a_1b}\nabla_{a_1}[e^{-2\phi}F_b^{a_2\dots a_{D-2}}] \propto g^{ab}\nabla_{a_1}[e^{-2\phi}F_b^{a_2\dots a_{D-2}}] = 0$$

Hence the F -perturbation equation reduces to

$$\nabla_{a_1}[e^{-2\phi}(\delta F)_b^{a_1\dots a_{D-2}}] = 0 \quad (2.24)$$

We can make the further observation that since F and δF are derived from a potential, $d(\delta F) = 0$, hence

$$\delta F_{\theta\phi,\dots,t} = \Omega q(r)\epsilon_{\theta\phi,\dots} = 0 \quad (2.25)$$

provided $\Omega \neq 0$. Thus only δF_T can be non-zero, and this is independent of the angular variables.

We may now argue that δF_T also vanishes. Without loss of generality, we will show this for $D = 4$, the generalisation to higher dimensions being an iterative process.

Consider $d(\delta F) = 0$.

$$F_{ab,c} + F_{bc,a} + F_{ca,b} = 0 \quad (2.26)$$

For $a = t, b = r, c = i$, we have

$$\delta F'_{Tit} + \Omega\delta F_{Trri} + i\mu_i\delta F_{Ttr} = 0 \quad (2.27)$$

Setting $f_t = \Sigma\mu_i\delta F_{Tit}$ and $f_r = \Sigma\mu_i\delta F_{Tir}$ and summing (2.27) appropriately gives

$$f'_t - \Omega f_r + i\mu^2\delta F_{Ttr} = 0 \quad (2.28)$$

where $\mu^2 = \Sigma\mu_i^2$. Now, we return to the equation of motion (2.24) which implies

$$\Omega\delta F_T^{ta} + \sum i\mu_i\delta F_T^{ia} + \delta F_T^{ra'} + \frac{2}{r}\delta F_T^{ra} = 0 \quad (2.29)$$

For $a=r$, we have

$$-\frac{(r-r_-)}{(r-r_+)}\Omega\delta F_{Ttr} + if_r = 0 \quad (2.30)$$

Substituting from (2.28) implies

$$f'_t = \left[\frac{\mu^2(r-r_+)}{\Omega(r-r_-)} + \Omega \right] f_r \quad (2.31)$$

Whereas taking (2.29) for $a = i$, multiplying by μ_i and summing gives

$$-\Omega g^{tt} f_t - (g^{rr} f_r)' - \frac{2}{r} g^{rr} f_r = 0 \quad (2.32)$$

Substituting for f_r from (2.31), and rearranging gives finally

$$\begin{aligned} \frac{(r-r_-)(r-r_+)}{r^2} f_t'' + \left(\frac{(2r-r_+-r_-)}{r^2} - \frac{\mu^2(r_+-r_-)(r-r_+)}{r[\mu^2(r-r_+) + \Omega^2(r-r_-)]} \right) f_t' \\ - \left(\mu^2 + \Omega^2 \frac{(r-r_-)}{(r-r_+)} \right) f_t = 0 \end{aligned} \quad (2.33)$$

By inspection of this equation we can see that there are no regular solutions and thus we must take $f_t = 0$. Hence $f_r = \delta F_{Tr} = 0$.

Now, returning to (2.27) implies

$$\delta F'_{Tit} + \Omega \delta F_{Tri} = 0 \quad (2.34)$$

and (2.26) for $a = i$, $b = t$, $c = j$ implies

$$i\mu_j \delta F_{Tit} + i\mu_i \delta F_{Ttj} + \Omega \delta F_{Tji} = 0 \quad (2.35)$$

Multiplying by μ_i and summing gives

$$i\mu^2 \delta F_{Ttj} - \Omega f_j = 0 \quad (2.36)$$

where $f_j = \Sigma \mu_i \delta F_{Tij}$. Taking the j -component of the equation of motion

$$\Omega \delta F_{Ttj} g^{tt} + i f_j + (\delta F_{Trj} g^{rr})' + \frac{2}{r} \delta F_{Trj} g^{rr} = 0, \quad (2.37)$$

substituting in for f_j , δF_{Trj} , and rearranging gives

$$\frac{(r-r_-)(r-r_+)}{r^2} \delta F''_{Ttj} + \frac{(2r-r_+-r_-)}{r^3} \delta F'_{Ttj} - \left(\mu^2 + \Omega^2 \frac{(r-r_-)}{(r-r_+)} \right) \delta F_{Ttj} = 0 \quad (2.38)$$

As before, this has no regular solutions, hence $\delta F_{Ttj} = 0$. (2.34) and (2.35) then imply that all other components of δF_T vanish, hence the perturbation δF_{ab} may be taken to be zero in all future analysis of the spherically symmetric perturbation equations.

Finally, to summarize, the equations of motion for the spherically symmetric perturbations we are interested in analysing reduce to the following

$$\begin{aligned} & \square \delta \phi - h^{ab} \nabla_a \nabla_b \phi + 2h^{ab} \nabla_a \phi \nabla_b \phi - 4\nabla_a \delta \phi \nabla^a \phi \\ & - \frac{h^{cd}}{(D-4)!} F^{ca_2 \dots a_{D-2}} F_d^{a_2 \dots a_{D-2}} = 0 \end{aligned} \quad (2.39a)$$

$$\begin{aligned} & \square h_{ab} + 2R_{cabd} h^{cd} - 2R_{e(a} h_{b)}^e - 4\nabla_a \nabla_b \delta \phi - 2\nabla_c \phi \nabla^c h_{ab} + 4\nabla_c \phi \nabla_b h_a^c \\ & - \frac{4}{(D-4)!} h^{cd} F_{aca_3 \dots a_{D-2}} F_{bd}^{a_3 \dots a_{D-2}} = 0 \end{aligned} \quad (2.39b)$$

the ‘electromagnetic’ perturbation vanishing in this case. Replacing the background values of the dilaton and Riemann tensor lead to the final versions given in appendix B.

3) The zero charge case.

We will first give a detailed analysis for the case when the black string or brane is uncharged. This simplifies the calculation for two reasons. The first is that the form of the metric is much simpler, leading to more tractable coefficients in the perturbation equations of Appendix B. The second is that as can be seen from appendix B, some equations decouple from each other; in particular, the perturbation of the dilaton can be neglected and we can deal solely with the metric perturbations. In addition, as we have mentioned before, it is consistent to take a vanishing trace of the metric perturbation.

We can first investigate the transverse terms of the metric perturbations in the $10 - D$ dimensions. These perturbations transform like scalars under the D -dimensional coordinate transformations as is well known in Kaluza-Klein decompositions. They are completely decoupled from all other perturbations and obey equations of the form

$$Ah^{ii''} + Bh^{ii'} + Ch^{ii} = 0 \quad (3.1)$$

where A, B and C are function of r . Their explicit form can be found in Appendix B. We

can investigate the behavior of h^{ii} as $r \rightarrow \infty$ and as $r \rightarrow r_+$. We obtain

$$\begin{aligned} h^{ii} &\sim \exp \pm r \sqrt{\Omega^2 + \mu^2} \quad r \rightarrow \infty \\ &\sim (r - r_+)^{\pm \Omega r_+} \quad r \rightarrow r_+ \end{aligned} \tag{3.2}$$

The regular solutions correspond to the minus root at infinity and the positive root at the horizon. It is therefore easy to see that a necessary condition to obtain a regular solution is to have a change in sign in the ratio of coefficient A/C in equation (3.1). From appendix B it is seen that this ratio does not change sign for the transverse part of the metric perturbations. We therefore set these perturbations to zero in our search for an instability.

We can now study the vector perturbations $h^{\mu i}$. An analysis similar to the scalar case shows that they do not lead to an instability, they are therefore set to zero. We are thus left with the tensor perturbations.

The tensor perturbations are more complex as the equations are coupled with each other. It is possible to rewrite them in term of a single variable, h^{tr} say, using the gauge conditions described in appendix B:

$$\begin{aligned} 0 = & \left\{ -\Omega^2 - \mu^2 V + \frac{(D-3)^2 \left(\frac{r_{\pm}}{r}\right)^{2(D-3)}}{4r^2} \right\} H^{tr''} \\ & - \left\{ \frac{\mu^2 [(D-2) - 2\left(\frac{r_{\pm}}{r}\right)^{D-3} + (4-D)\left(\frac{r_{\pm}}{r}\right)^{2(D-3)}]}{rV} \right. \\ & + \left. \frac{\Omega^2 [(D-2) + (2D-7)\left(\frac{r_{\pm}}{r}\right)^{D-3}]}{rV} - \frac{3(D-3)^2 \left(\frac{r_{\pm}}{r}\right)^{2(D-3)} [(D-2) - \left(\frac{r_{\pm}}{r}\right)^{D-3}]}{4r^3 V} \right\} H^{tr'} \\ & + \left\{ \left(\mu^2 + \Omega^2/V \right)^2 + \frac{\Omega^2 [4(D-2) - 8(D-2)\left(\frac{r_{\pm}}{r}\right)^{D-3} - (53 - 34D + 5D^2)\left(\frac{r_{\pm}}{r}\right)^{2(D-3)}]}{4r^2 V^2} \right. \\ & + \left. \frac{\mu^2 [4(D-2) - 4(3D-7)\left(\frac{r_{\pm}}{r}\right)^{D-3} + (D^2 + 2D - 11)\left(\frac{r_{\pm}}{r}\right)^{2(D-3)}]}{4r^2 V} \right. \\ & + \left. \frac{(D-3)^2 \left(\frac{r_{\pm}}{r}\right)^{2(D-3)} [(D-2)(2D-5) - (D-1)(D-2)\left(\frac{r_{\pm}}{r}\right)^{D-3} + \left(\frac{r_{\pm}}{r}\right)^{2(D-3)}]}{4r^4 V^2} \right\} H^{tr} \end{aligned} \tag{3.3}$$

Before dwelling on the existence of an instability for the tensor perturbations we would like to comment on the flat space case, i.e $r_+ = 0$. In this case the ratio of the coefficients

of the first and last term does not change sign and we get the expected result that there is no instability.

When $\Omega > (D - 3)/r_+$ it is possible to use the same analytic argument as for the transverse perturbations to show that no instability exists. However when $\Omega < (D - 3)/r_+$ the ratio of the coefficients of the first and last term does indeed change sign and we must resort to numerical techniques to establish the existence or otherwise of an instability.

It is possible to reduce the set of second order equations to first order ones by using the two gauge conditions (B.10a-b). By differentiating these and using the second order equation we reduce them to first order equations but as we have only three variables and four first order equations we can reduce one of them to a constraint between the fields themselves (without derivatives). One of the variables can thus be reexpressed as a function of the others.

First let us define

$$H_{\pm} = V_+ H^{tt} \pm \frac{H^{rr}}{V_+} \quad (3.4a)$$

$$H = -H^{tr} \quad (3.4b)$$

where $V_+ = (1 - (r_+/r)^{D-3})$. The Lichnerowicz and the gauge equations reduces to the constraint:

$$\begin{aligned} H_+ \frac{V_+}{2} \left[\mu^2 + \frac{(D-3)(D-2)}{2r^2} (1 - V_+) \right] = \\ H_- \left[\Omega^2 + \frac{\mu^2 V_+}{2} + \frac{(D-3)}{4r^2} (-(D-3) + (D-4)V_+ + V_+^2) \right] \\ + H \frac{V_+}{r} \left[-\Omega(D-2) - \frac{\mu^2}{2\Omega} ((D-3) - (D-1)V_+) \right] \end{aligned} \quad (3.5)$$

and the following two first order differential equations

$$\frac{\partial H}{\partial r} = \frac{\Omega}{2V_+} (H_+ + H_-) - \frac{[(D-3) + V_+]}{rV_+} H \quad (3.6)$$

and

$$\frac{\partial H_-}{\partial r} = \frac{\mu^2}{\Omega} H + \frac{(D-2)}{2r} H_+ + \frac{[(D-3) + (-2D+3)V_+]}{2rV_+} H_- \quad (3.7)$$

From this we can deduce that as $r \rightarrow \infty$

$$H = \sqrt{\Omega^2 + \mu^2} F_+ e^{r\sqrt{\Omega^2 + \mu^2}} - \sqrt{\Omega^2 + \mu^2} F_- e^{-r\sqrt{\Omega^2 + \mu^2}} \quad (3.8a)$$

$$H_- = \frac{\mu^2}{\Omega} F_+ e^{r\sqrt{\Omega^2 + \mu^2}} - \frac{\mu^2}{\Omega} F_- e^{-r\sqrt{\Omega^2 + \mu^2}} \quad (3.8b)$$

and as $r \rightarrow r_+$

$$\begin{aligned} H &= (D-3)r_+^{D-4} \left(-\frac{1}{2} + \frac{\Omega r_+}{D-3}\right) G_+ (r^{D-3} - r_+^{D-3})^{-1 + \frac{\Omega r_+}{D-3}} \\ &\quad - (D-3)r_+^{D-4} \left(-\frac{1}{2} - \frac{\Omega r_+}{D-3}\right) G_- (r^{D-3} - r_+^{D-3})^{-1 - \frac{\Omega r_+}{D-3}} \end{aligned} \quad (3.9a)$$

$$\begin{aligned} H_- &= \left(\frac{\mu^2}{\Omega} + \frac{D-2}{r_+}\right) G_+ (r^{D-3} - r_+^{D-3})^{\frac{\Omega r_+}{D-3}} \\ &\quad - \left(\frac{\mu^2}{\Omega} - \frac{D-2}{r_+}\right) G_- (r^{D-3} - r_+^{D-3})^{-\frac{\Omega r_+}{D-3}}. \end{aligned} \quad (3.9b)$$

The behavior of H_+ can be obtained from the constraint (3.5).

The regular solution corresponds to $F_+ = 0$ and $G_- = 0$. We have used a Runge-Kutta algorithm with variable stepsize to integrate these equation from $r_1 = 200$ to $r_2 = 2.000002$ with a tolerance $\epsilon = 10^{-6}$. We do this integration for fixed μ and various values of Ω . Near $r = r_+$ we calculate the ratio $R = G_-/G_+$ and look for a change in sign (as Ω varies). When a change in sign occurs we home in towards the value of Ω for which this change occurs and ensure that the ratio decreases towards such a value (an increase would imply that G_- goes through a zero). Figure 2 shows the behavior of the function H^{tt} , H^{tr} and H^{rr} . In Figure 3. we have plotted Ω as a function of μ for which an unstable mode has been found. This plot takes $r_+ = 2$, but behavior for other values of r_+ can be obtained from the scale transformation

$$\begin{aligned} r_+ &\rightarrow \alpha r_+ \\ \Omega &\rightarrow \frac{\Omega}{\alpha} \\ \mu &\rightarrow \frac{\mu}{\alpha} \end{aligned} \quad (3.10)$$

There are at least two interesting points in Fig.3. The first one is to note that Ω does not seem to go to zero as $\mu \rightarrow 0$. This might be a numerical artifact as our boundary

conditions are different in this limit. However if it is not a numerical artifact it would be surprising at first, as in this particular case the equations reduce exactly to the stability of D-dimensional black holes. We know that black holes in four dimensions are stable, moreover the mode under investigation here is the spherically symmetric mode and we know by Birkhoff's theorem (and its generalization to higher dimensions) that the solution must be Schwarzschild. We must therefore conclude that this mode is pure gauge. We can in fact work out what this gauge transformation looks like in the asymptotic regions.

As the system remains spherically symmetric and has no dependence on the extra dimension, we must have

$$\xi^a = (\xi^t, \xi^r, 0, \dots, 0) \quad (3.11)$$

It is easy to verify that the particular gauge (in the large r limit)

$$\xi^t = \xi^r = e^{\Omega(t-r)} \quad (3.12)$$

corresponding to metric perturbations of the form:

$$h^{tt} = \xi^{t;t} \approx -\Omega e^{\Omega(t-r)} \quad (3.13a)$$

$$h^{rr} = \xi^{r;r} \approx -\Omega e^{\Omega(t-r)} \quad (3.13b)$$

$$h^{tr} = \xi^{(t;r)} \approx -\Omega e^{\Omega(t-r)}. \quad (3.13c)$$

This gives rise to a transverse and traceless metric as can be easily verified. In the $r \rightarrow r_+$ limit we have

$$\xi^t = -r_+(r - r_+)^{-1+\Omega r_+} \quad ; \quad \xi^r = (r - r_+)^{\Omega r_+} \quad (3.14)$$

corresponding to the metric perturbation

$$h^{tt} \approx \left(-\frac{1}{2} + \Omega r_+\right) r_+ (r - r_+)^{-2+\Omega r_+} e^{\Omega t} \quad (3.15a)$$

$$h^{rr} \approx \frac{(-1 + 2\Omega r_+)}{2r_+} (r - r_+)^{\Omega r_+} e^{\Omega t} \quad (3.15b)$$

$$h^{tr} \approx -\left(-\frac{1}{2} + \Omega r_+\right) (r - r_+)^{-1+\Omega r_+} e^{\Omega t} \quad (3.15c)$$

Of course we still have to show that there is a regular solution which links these two asymptotic behavior. This can be done numerically.

The result for $\mu = 0$ might leave the reader worried that the instability found might be pure gauge. However the case where $\mu \neq 0$ is different. In this case it is easy to convince oneself that they cannot be pure gauge. The simplest way is to assume we can write these perturbations in term of ξ^a and find a contradiction. Because perturbations in the $10 - D$ dimensions are zero, we would have

$$\xi^{j;j} = i\mu_j \xi^j = 0 \quad j = D + 1, \dots, 10$$

thus this implies that $\xi^j = 0$ for $\mu_j \neq 0$ as $h^{jj} = 0$. If we now look at $h^{\mu j} = \xi^{\mu;j}$ this equation implies that $\xi^\mu = 0$ and thus $h^{\mu\nu} = 0$, a contradiction. This ensures that the perturbations found in the uncharged case cannot be pure gauge and therefore must be physical. In the charged case the perturbations in the extra dimensions are non-zero and thus such a proof is more difficult. In this latter case we will make use of gauge invariant variables described in appendix A to show that the perturbations are not gauge artifacts.

It is interesting to understand where this extra degree of freedom for the gravitational field has come from. It is a result of the degree of freedom of the gravitons due to the extra dimensions. One easy way to make this clear is to use a Kaluza-Klein analogy. The gauge transformations in D dimensions give the graviton degrees of freedom which are transverse to the direction of propagation. This is related to the masslessness of the graviton. Once the extra dimensions come into play, through the dependence $\exp i\mu_j z^j$, this effectively gives a mass to the graviton and thus a longitudinal component corresponding to a spherically symmetric mode in D dimensions. It is this component which is responsible for the instability. In the limit as $\mu \rightarrow 0$, this mode becomes pure gauge as the mass disappear.

The other interesting point to notice in Fig.3 is the fact that there is a maximum value μ_j^{max} for the instability to exist. The analysis carried above can be repeated in the case where the extra dimensions are periodically identified. The only difference in this case is that the values of μ_j will be quantized as $2\pi n/L_j$, for $n = 0, 1, \dots$ and L being the length of extra dimensions. From this we can see that if the L_j are small enough, the first value of μ_j will be larger than μ_j^{max} and thus the instability will not occur. The instability occurs only when the black holes (in D dimensions) have a Schwarzschild radius of the

order of the size of the extra dimensions as suggested by the entropy argument described in the introduction. Thus if we believe the assumptions of string theory that the $10 - D$ extra dimensions are wrapped in small circles, the instability found here will not have any effect on astrophysical black holes. It is interesting that this instability appears to break the duality of the solutions obtained by replacing the radius L of the extra dimension by its inverse $1/L$. If we choose L appropriately, there is an instability for this spacetime but there will not be one when $L \rightarrow 1/L$, however, as we have considered only momentum modes in this calculation, and not winding modes, it is possible that there will be a winding mode instability for the dual spacetime. However, we have neither verified nor refuted this claim.

We have discussed the existence of the instability but what does it correspond to? All the perturbations have the form $\exp i\mu_j z^j$ and thus implies an oscillatory behavior of the distance scale as a function of the extra dimensions. To make this clearer we can work out the location of the apparent horizon. To do so we introduce ingoing Eddington-Finkelstein coordinates, and for ease of description restrict our attention to $D = 4$:

$$ds^2 = -\frac{(r - r_+)}{r} du^2 + 2du dr + r^2 d\Omega^2 + dz^2 \quad (3.16)$$

where $u = t + r + r_+ \ln(r - r_+)$ in terms of the usual Schwarzschild coordinates. The apparent horizon is located where the outgoing light-cones have zero divergence. The spherically symmetric light rays for this metric are given by

$$\begin{cases} u = \text{const.} & \text{ingoing} \\ \frac{dr}{du} = \frac{(r-r_+)}{2r} & \text{outgoing} \end{cases} \quad (3.17)$$

In the case of the unperturbed black strings the apparent horizon is at $r = r_+$. If we perturb the metric slightly we get for the new outgoing null geodesics

$$h_{uu} - \frac{(r - r_+)}{r} + 2\frac{dr}{du}(1 + h_{ur}) + h_{rr} \left(\frac{dr}{du}\right)^2 = 0$$

which to first order gives

$$\frac{dr}{du} \approx \frac{(r - r_+)}{2r} - \frac{(r - r_+)}{2r} h_{ur} - \frac{1}{2} h_{uu} - \frac{(r - r_+)^2}{8r^2} h_{rr} \quad (3.18)$$

which has a zero for

$$r \approx r_+(1 + h_{uu}) + \lim_{r \rightarrow r_+} \left[(r - r_+)h_{ur} + \frac{(r - r_+)^2}{4r_+} h_{rr} \right]. \quad (3.19)$$

Using the behavior of the solution obtained above we finally get

$$r \approx r_+ + \text{const.} \cos(\mu z) \quad (3.20)$$

This shows that the apparent horizon of the perturbed spacetime oscillates as a function of the extra dimensions. The schematic behavior of the apparent horizon is depicted in Figure 4. Thus it would appear that this perturbation destabilises the event horizon causing it to ripple in the transverse dimensions.

4) The small charge case.

We have seen in the previous section that uncharged black strings and p-branes are unstable. We now investigate the question whether a charge on these black objects might stabilize them. Physically, it seems reasonable that they will also be unstable at least for small charges, this is because for small charge the metric is essentially the same as the uncharged case. As long as we remain outside the event horizon ($r > r_+$), as indeed we do to investigate the instability, the effect of the charge should be negligible. However a non-trivial effect of the charge is that now the $10 - D$ dimensional perturbations that previously vanished become coupled to the D -dimensional non-zero perturbations. This now complicates the issue as we can no longer set the former perturbations to zero; all the perturbations must be solved simultaneously. An additional complication is that it is not possible to take the trace $h = g_{ab}h^{ab}$ to be zero, for as we have already shown, this simplification is only possible in vacuo. The perturbation of the scalar field ϕ couples to h and prevents us from setting that part of the metric to zero.

In order to guide the reader through the maze of these equations we will first focus on the small charge case. Indeed in this case the problem, although more complicated than the uncharged case, is drastically simplified from the general charged case. The simplification come through the observation that the coupling of D -dimensional perturbations to the $10 - D$ ones are always through a factor proportional to the charge $Q^2 = r_- r_+ / 2$. As the

10 – D dimensional perturbations are vanishing for $Q = 0$, we deduce that they are in fact proportional to Q at small charge. This lead us to an expansion of the perturbations in powers of r_- . Using such an expansion $h^{\mu\nu}$ will be zeroth order, and the other variables, $h^{j\mu}$, h^{jk} and $\delta\phi$ will be first order. Notice that h , the trace of h^{ab} , will also be first order.

Before starting the analysis, note that from the form of the perturbation equations in Appendix B, it is easy to see that writing

$$h^{ij} = \frac{i\mu_i\mu_j}{\mu^2} h^{zz} \quad (4.1a)$$

$$h^{\mu i} = \frac{i\mu_i}{\mu} h^{\mu z} \quad (4.1b)$$

removes the $(10 - D)$ i -degrees of freedom to just one transverse degree of freedom, which we call ‘ z ’. Additionally, for calculational simplicity, we will consider the specific case of the black 6-brane ($D = 4$), which will at least remove the variable D from our equations! The analysis can be extended to other black strings and p-branes, however, since we have already shown black p -branes to be unstable for all values of D in the previous section, an analysis for general D at this stage would be neither economical nor illuminating.

We first analyze the equation for the perturbations of scalar field f , h^{zz} and the trace h :

$$\begin{aligned} \frac{(r-r_+)}{r} f'' + \frac{2r-r_+}{r^2} f' - \left(\mu^2 + \frac{\Omega^2 r}{(r-r_+)} \right) f + \frac{r_-(r-2r_+)}{r^2} K \\ + \frac{r_-(4r-5r_+)}{4r^3(-r+r_+)} h^{rr} - \frac{r-r_+(r-r_+)}{4r^5} h^{tt} = 0 \end{aligned} \quad (4.2)$$

$$\frac{(r-r_+)}{r} h^{zz''} + \frac{(2r-r_+)}{r^2} h^{zz'} - \left(\mu^2 + \frac{\Omega^2 r}{(r-r_+)} \right) h^{zz} + 2\mu^2 f(r) = 0 \quad (4.3)$$

$$\frac{(r-r_+)}{r} \left(\frac{h''}{2} - 2f'' \right) + \frac{(2r-r_+)}{r^2} \left(\frac{h'}{2} - 2f' \right) - \left(\mu^2 + \frac{\Omega^2 r}{(r-r_+)} \right) \left(\frac{h}{2} - 2f \right) - \frac{2r-r_+}{r^2} K = 0 \quad (4.4)$$

There is also an equation for h^{rz} and h^{tz} and the gauge condition

$$\nabla_a \bar{h}^{az} \approx h^{rz'} + i\mu h^{zz} + \Omega h^{tz} + \frac{2}{r} h^{rz} - i\frac{\mu}{2} h \approx 0 \quad (4.5)$$

Taking a derivative of this gauge condition and using the h^{tz} equation we get

$$\frac{(r-r_+)}{2r}(i\mu h^{zz'} - i\frac{\mu}{2}h' + \Omega h^{tz'} + 4i\mu f') + \left(\frac{\mu^2}{2} + \frac{\Omega^2 r}{2(r-r_+)}\right)h^{rz} + \frac{i\mu r_-}{2r^2}h^{rr} + \frac{\Omega r_+}{2r^2}h^{tz} \approx 0 \quad (4.6)$$

This gives us two first order equations for h^{rz} and h^{tz}

With these equations we can investigate the behavior of the fields at infinity and near $r = r_+$. In the large r limit all these fields have a behavior of the form $e^{\pm r\sqrt{\Omega^2 + \mu^2}}$, the regular solution having the negative sign. The coefficients of f, h, h^{zz}, h^{tz} are independent of each other but once these are fixed, the coefficient in front of the exponential of h_i^{rz} (denoted by a subscript i) is given by

$$h_i^{rz} \approx \frac{1}{\sqrt{\Omega^2 + \mu^2}} \left(\frac{\mu}{2}h_i + i\Omega h_i^{tz} - \mu h_i^{zz} \right). \quad (4.7)$$

in terms of the coefficients of the exponential of the other perturbations. Near $r = r_+$, the perturbations h, f, h^{zz}, h^{rz} behave as $(r - r_+)^{\pm\Omega r_+}$ and h^{tz} as $(r - r_+)^{-1 \pm \Omega r_+}$.

With these boundary conditions we can integrate eq(4.2-6) to find the unstable mode assuming the zeroth order solutions found previously. We have used a similar technique as in the uncharged case to find the regular solution near $r = r_+$. By adjusting the values at large r we have been able to find regular solutions for all the perturbations. Fig.5 gives the behavior of the modes which lead to the instability.

5) The charged case.

We have now shown that black strings and branes with small charges are unstable by assuming a perturbative analysis in term of the charge. As we have seen in that case, we have a zeroth order equation in a small charge perturbation for the four dimensional metric perturbation. The dilaton and the extra dimensions are first order and could be solved assuming the zeroth order solutions for the four dimensional metric. In this section we give the argument for the general charge. The idea of how to handle the equations is very much the same as in the previous cases but the calculations are much more tedious as the equations will now involve the charge parameter (or rather $r_- = 2Q^2/r_+$).

The equations of motion for the different fields are given in appendix B. As before, we were unable to show the existence or absence of an unstable mode analytically. We have thus resorted to numerical evaluation of the unstable mode. We can rewrite the set of equations in Appendix B as a set of first order equations suitable for numerical integration. First let us define a new variable $q = h/2 - 2f$ where h is the trace of the metric perturbation and rewrite the equations for f, q, h^{zz} as

$$\begin{aligned}
f' &= \pi_f \\
\pi_f' &= -\frac{r^2}{(r-r_-)(r-r_+)} \left(-(\mu^2 + \Omega^2 \frac{(r-r_-)}{(r-r_+)})f + \frac{r_-(r-r_+)(r_- - r_+)}{4r^3(r-r_-)^2} h^{tt} \right. \\
&\quad \left. - \frac{r_-(4r^2 - 5rr_- - 5r_+ + 6r_-r_+)}{4r^2(r-r_-)^2(r-r_+)} h^{rr} \right. \\
&\quad \left. + \frac{r_-(r-2r_+)}{r^2} K + \frac{(2r^2 - rr_+ - r_-r_+)}{r^3} \pi_f \right) \\
q' &= \pi_q \\
\pi_q' &= \frac{r^2}{(r-r_-)(r-r_+)} \left(\frac{2r_-r_+}{r^2} K + (\mu^2 + \Omega^2 \frac{(r-r_-)}{(r-r_+)})q \right. \\
&\quad \left. - \frac{(r-r_-)(2r-r_+)}{r^3} \pi_q \right) \\
h^{zz'} &= \pi_{h^{zz}} \\
\pi_{h^{zz}}' &= \frac{r^2}{(r-r_-)(r-r_+)} \left(-4\mu^2 f + \frac{i2\mu r_-}{r(r-r_-)} h^{rz} \right. \\
&\quad \left. + (\mu^2 + \Omega^2 \frac{(r-r_-)}{(r-r_+)}) h^{zz} + \frac{(-2r+r_-+r_+)}{r^2} \pi_{h^{zz}} \right)
\end{aligned} \tag{5.1}$$

The next step is to use the gauge condition (B.10d), the gauge condition for the index ‘ z ’. The equation for $h^{tz'}$ is obtained by taking the derivative of this and using the equation of motion of h^{rz} to eliminate the second derivative. This gives an equation in terms of the

variables and their first derivatives. Explicitly we have

$$\begin{aligned}
ih^{tz'} &= -\frac{1}{\Omega} \left(-\frac{\Omega(r_- - r_+)}{(r - r_-)(r - r_+)} ih^{tz} - \frac{\mu rr_-}{(r - r_-)^2(r - r_+)} h^{rr} \right. \\
&\quad \left. + \left(\frac{\Omega^2 r^2}{(r - r_+)^2} + \frac{\mu^2 r^2}{(r - r_-)(r - r_+)} \right) ih^{rz} - 2r\mu\pi_f + \mu\pi_q - \mu\pi_{h^{zz}} \right) \quad (5.2) \\
ih^{rz'} &= -\mu q - 2\mu f - \Omega ih^{tz} - \frac{(2r - 3r_-)}{r(r - r_-)} h^{rz} + \mu h^{zz}
\end{aligned}$$

We can use the gauge condition for the indices t and r to get first order equations for h^{tr} and h^{rr}

$$\begin{aligned}
h^{tr'} &= -\frac{\Omega(r - r_-)(q + 2f)}{2(r - r_+)} - \Omega h^{tt} - \frac{(2r^2 - 4rr_- - rr_+ + 3r - r_+)}{r(r - r_-)(r - r_+)} h^{tr} - i\mu h^{tz} \\
h^{rr'} &= \frac{(r - r_-)(r - r_+)}{r^3} (2q + 4f) - \frac{(r - r_+)(-2r^2 + rr_- + 3rr_+ - 2r - r_+)}{2r^3(r - r_-)} h^{tt} \\
&\quad - \Omega h^{tr} - \frac{(6r^2 - 9rr_- - 7rr_+ + 10r - r_+)}{2r(r - r_-)(r - r_+)} h^{rr} - i\mu h^{rz} \\
&\quad - \frac{(r - r_-)(r - r_+)}{r^3} h^{zz} + \frac{(r - r_-)(r - r_+)}{r^2} (\pi_q + 2\pi_f) \quad (5.3)
\end{aligned}$$

Taking a derivative of the gauge condition with index t and getting rid of the second order derivative using the second order equation for h^{tr} we get

$$\begin{aligned}
h^{tt'} &= -\frac{1}{\Omega} \left(-\frac{\Omega(r_- - r_+)}{(r - r_+)^2} q - \frac{\Omega(r_- - r_+)}{(r - r_-)(r - r_+)} h^{tt} + \right. \\
&\quad \left(\frac{\Omega^2 r^2}{(r - r_+)^2} + \frac{\mu^2 r^2}{(r - r_-)(r - r_+)} \right) h^{tr} + \frac{i\mu(r_- - r_+)}{(r - r_-)(r - r_+)} h^{tz} + \\
&\quad \left(\frac{\mu^2 rr_-}{\Omega(r - r_-)^2(r - r_+)} + \frac{\Omega r(-2rr_- + rr_+ + r - r_+)}{(r - r_-)(r - r_+)^3} \right) h^{rr} \quad (5.4) \\
&\quad - \frac{i\mu}{\Omega} \left(\frac{\Omega^2 r^2}{(r - r_+)^2} + \frac{\mu^2 r^2}{(r - r_-)(r - r_+)} \right) h^{rz} - \left(\frac{\mu^2}{\Omega} - \frac{\Omega(r - r_-)}{(r - r_+)} \right) (\pi_q - 2\pi_f) \\
&\quad \left. + \frac{\mu^2}{\Omega} \pi_{h^{zz}} \right)
\end{aligned}$$

We have now the field equations written in terms of first derivatives but we still have an equation left which comes from the time derivative of the gauge condition with index r . This leaves us with a constraint between the variables in equations (5.1-4) given by

$$\begin{aligned}
& \left(\frac{2\Omega^2(r-r_-)^2}{r^2} + \frac{2\mu^2(r-r_-)(r-r_+)}{r^2} + \frac{(r-r_+)(-2r^2+rr_-+3rr_+-2r-r_+)}{2r^5} \right) f \\
& + \left(\frac{(4r^3r_- - 3r^2r_-^2 - 10r^2r_-r_+ + 8rr_-^2r_+ + r^2r_+^2 + 4rr_-r_+^2 - 4r_-^2r_+^2)}{8r^6} \right. \\
& \quad \left. - \frac{\mu^2(r-r_-)(r-r_+)}{2r^2} - \frac{\Omega^2(r-r_-)^2}{2r^2} \right) h \\
& + \left(\frac{(r-r_+)(-r^2r_-^2 + 4r^3r_+ - 8r^2r_-r_+ + 6rr_-^2r_+ - 3r^2r_+^2 + 6rr_-r_+^2 - 4r_-^2r_+^2)}{8r^6(r-r_-)} \right. \\
& \quad \left. - \frac{\Omega^2(r-r_-)(r-r_+)}{2r^2} \right) h^{tt} \\
& + \left(\frac{\mu^2(r-r_+)(2r^2 - rr_- - 3rr_+ + 2r-r_+)}{4\Omega r^3(r-r_-)} + \frac{\Omega(2r-r_-)(r-r_+)}{2r^3} \right) h^{tr} \\
& + \left(-\frac{i\Omega\mu(r-r_-)(r-r_+)}{r^2} + \frac{i\mu(r-r_+)(r-r_+)(2r^2 - rr_- - 3rr_+ + 2r-r_+)}{4\Omega r^5(r-r_-)} \right) h^{tz} \\
& + \left(\frac{(-16r^3r_- + 11r^2r_-^2 + 34r^2r_-r_+ - 24rr_-^2r_+ - r^2r_+^2 - 16rr_-r_+^2 + 12r_-^2r_+^2)}{8r^4(r-r_-)(r-r_+)} \right. \\
& \quad \left. - \frac{\mu^2r_-(r-r_+)(-2r^2+rr_-+3rr_+-2r-r_+)}{4\Omega^2r^4(r-r_-)^2} + \frac{\mu^2}{2} + \frac{\Omega^2(r-r_-)}{2(r-r_+)} \right) h^{rr} \\
& + \left(\frac{i\mu(-4r+3r_-)(r-r_+)}{2r^3} + \frac{i\mu^3(r-r_+)(2r^2 - rr_- - 3rr_+ + 2r-r_+)}{4\Omega^2r^3(r-r_-)} \right) h^{rz} \\
& + \left(\frac{(r-r_-)(r-r_+)(-rr_- - rr_+ + 2r-r_+)}{4r^6} + \frac{\mu^2(r-r_-)(r-r_+)}{2r^2} \right) h^{zz} \\
& + \left(\frac{2(r-r_-)(-3r+2r_-)(r-r_+)^2}{r^5} + \frac{\mu^2(r-r_+)^2(2r^2 - rr_- - 3rr_+ + 2r-r_+)}{\Omega^2r^5} \right) \pi_f \\
& + \frac{\mu^2(r-r_+)^2(-2r^2+rr_-+3rr_+-2r-r_+)}{8\Omega^2r^5} \pi_h \\
& + \left(\frac{(r-r_-)^2(r-r_+)^2}{2r^5} + \frac{\mu^2(r-r_+)^2(2r^2 - rr_- - 3rr_+ + 2r-r_+)}{4\Omega^2r^5} \right) \pi_{h^{zz}} = 0
\end{aligned} \tag{5.5}$$

The initial data (as $r \rightarrow \infty$) must obey this constraint, and then is automatically preserved along the integration. We have used it to verify that the integration errors were indeed small.

In the previous sections we deduced the existence of an instability by finding zeros of a ratio, R , of the irregular to regular solutions near the horizon. As it happened, the functions were essentially independent near the horizon, so we could investigate their ratios independently. Changing the dilaton perturbation, say, and thus R_{dil} , did not affect the ratios for the other perturbations, therefore we could vary our parameters freely to find the instability. However, when the charge becomes large, this is no longer the case, changing the initial value for the dilaton perturbation will now change the ratio of field h_{zz} , say. The task of finding where a 5-dimensional vector (corresponding to the ratio of each of the perturbation fields) is much harder. We have thus resorted to the Newton-Raphson method described in [17]. In a nutshell the idea is to assume that we are not too far from the zero

$$0 = R_i(x + \delta x) \approx R_i(x) + \frac{\partial R_i}{\partial x_j} \delta x_j \quad (5.6)$$

We can then invert this equation to get

$$\delta x_j \approx \frac{\partial R_i}{\partial x_j}^{-1} R_i(x) \quad (5.7)$$

that is we calculate the gradient of R at the value of x we are and follow it in towards the zero of R .

Figure 5 shows Ω for an unstable mode as a function of the charge for different values of μ , the frequency in the orthogonal direction. The important point to note is that the curves cross zero, at which point the unstable mode disappears for the appropriate value of μ .

We have checked using the gauge invariant variables of appendix A that the solutions found correspond to real physical solution and not gauge artifacts. This is obvious for the small charge case as the dominant quantities entering in equations (A.4) and (A.5) are the chargeless terms with the small charge terms being sub-dominant. For large charges we have calculated the quantity (A.4) directly. It is non-zero for the perturbations studied.

Near the extreme caase, the function calculating the ratio between the regular and irregular modes has very large derivatives and thus is not easily amenable to numerical study. As this happens to be also at very small values of Ω we conjecture that the extreme black hole is stable. This important particular case will be investigated elsewhere [18].

6) Conclusion.

We have investigated the instability of ‘magnetically’ charged dilatonic black p -branes in string theory. In particular we have shown that the chargeless instability found in [13] still persists in the presence of a charge on the black string or brane. This demonstrates that this instability is a generic phenomena. We remind the reader that this instability can be pictured as the string horizon collapsing in some regions of the extra dimensions and expanding in others. The unstable mode remains present for large values of charge but in this paper we have not proved its existence or otherwise for the extremal case; from the trend of the time frequency Ω as function of the charge it seems rather plausible that this case might be stable. However, without a more detailed analysis tailored to the extremal case, we cannot be definite. We can however indicate a different way of stabilizing the black p -brane - compactification. Compactification implies that the values of μ_i are quantized. If the compactification is on a scale smaller than the inverse mass of the black hole, the first allowable value of μ would be greater than the maximum one allowed for the instability, so such black doughnuts would be stable. Since there must be compactification of any extra dimensions on an extremely small scale, all but the tiniest black doughnuts would be safe, and those that would not would presumably have evaporated producing their own naked singularities long ago. Thus this instability will have no effect for contemporary astrophysical black holes.

One of the main questions of interest about this instability is the nature of its endpoint - what does the black string become? Of course, since our calculation is linear, we cannot strictly say anything about the final state, however, since in the chargeless case there are no other scales in the problem except the mass per unit length of the black string, it seems very reasonable to conjecture that the endpoint of the instability will be the fragmentation of the black strings into a bunch of small spherical black holes. This suggestion is supported by

the entropy argument given in the first section. Periodic black hole solutions are known¹⁹, so unlike the Reissner-Nordstrom instability, there is a final state solution here. It is quite likely that such a solution will itself exhibit a Jeans-like instability to long wavelength clustering of black holes which itself will be unstable and so on. In any case, a process of fragmentation will produce a naked singularity and hence violate cosmic censorship, however, we should again stress that this is speculative; ideally one should follow the instability numerically into the non-linear régime to see if fragmentation develops, this work is currently underway.

For the charged black p -brane, there are other possibilities. The fact that $\delta F = 0$ suggests that where the horizon shrinks the solution becomes locally closer to extremality. It is possible that when a charge is present on the black string or brane fragmentation will not occur. The electromagnetic repulsion might stabilize the string as the 2-spheres where the charge resides shrink to smaller volume. Unlike the chargeless case, the charged case might not lead to naked singularities and fragmentation. This conclusion assumes both the stability of extremal black p -branes, which is likely, as well as the non-participation of charge in the instability, which is an unknown without following the instability into the non-linear régime. Alternatively, the ‘magnetic’ nature of the charge considered also makes it plausible that as the horizon shrinks in some regions, higher energy physics might come into play, and that in such regions, the relevant monopole solution could ‘pop out’ from behind the horizon. For example, consider $D = 4$. Here an instability is known for the four dimensional monopole black hole, in which for small enough horizons, the t’Hooft Polyakov monopole is the preferred exterior solution²⁰. Thus, in the case of the 6-brane should the horizon volume become too small, there will be sections of the rippled brane that will have t’Hooft Polyakov type monopole fields surrounding them. The charge, having appeared in an exterior smooth form, would then presumably pose no obstruction to the interior collapse of the event horizon, and thence a naked singularity. Again, this assumes some knowledge of the non-linear régime.

We can also comment on black p -branes with topological charge. For example, consider the axionic black holes of Bowick et. al.²¹ These carry ‘quantum’ charge, detectable only by an Aharonov-Bohm scattering process, but as far as classical physics is concerned (hence

our instability) they behave as if uncharged, since the quantum charge they carry does not affect the exterior spacetime which is consequently Schwarzschild. The field strength of the axion field is zero throughout the spacetime, but the gauge field (rather like the gauge field of a local cosmic string) is non-trivial due to the topology of the spacetime, $B = \frac{Q}{\sin\theta} d\theta \wedge d\phi$. This solution can clearly be extended to a five dimensional black string (or ten-dimensional black 6-brane). This solution too will be unstable, however, a five (or ten) dimensional black hole cannot carry the same type of axion charge. Drawing analogy to the gauge field of a local cosmic string, it seems likely that during the fragmentation process, higher energy physics could come into play, producing an axion vortex, which could appear from behind the event horizon by an analogous process discussed above for the charged black p -brane. The endpoint in this case might be a line of black holes threaded by a cosmic axion string (not to be confused with the four-dimensional global string).

To summarize, whether charged or uncharged, black p -branes exhibit a long wavelength instability which causes the apparent horizon to ripple. A plausible endpoint of this instability is fragmentation, and hence violation of cosmic censorship, however, without more evidence in the non-linear régime, this remains speculative. Perhaps a more realistic conclusion is that due to this instability, black strings and p -branes will not form from collapse in the first place. From our work, it appears there are two ways of avoiding this instability. One is if the black brane is extremal, although the methods described here were unable to probe this end of parameter space, all indications were that these solutions were stable. This is also anticipated since extremal branes are supersymmetric. The other way of avoiding the instability is by compactification, indeed, to be totally speculative, since the endpoint of the instability does appear to be some periodic solution, it is tempting to suggest that the instability of some primordial black string could have triggered an effective compactification of our universe!

Acknowledgments.

We are grateful to J.B.Hartle, J.A.Harvey, S.W.Hawking, G.Horowitz, D. Wiltshire, and W.H.Zurek for useful conversations. We appreciate the careful perusal of the manuscript and accompanying remarks by J.Bowcock and P.Laflamme. R.G. is supported by the

S.E.R.C. R.L. thanks Los Alamos National Laboratory for support. R.G. also acknowledges the support of the McCormick Fellowship at the Enrico Fermi Institute where this work was started.

References.

- [1] S.W.Hawking, *Commun. Math. Phys.* **43** 199 (1975).
- [2] S.W.Hawking, *Commun. Math. Phys.* **87** 395 (1982).
- [3] E.Poisson and W. Israel, *Phys. Rev. D***41** 1796 (1990).
- [4] G.Gibbons and K.Maeda, *Nucl. Phys.* **B298** 741 (1988).
D.Garfinkle, G.Horowitz and A.Strominger, *Phys. Rev. D***43** 3140 (1991).
- [5] R.Gregory and J.Harvey, *Phys. Rev. D***47** 2411 (1993).
J.Horne and G.Horowitz, *Nucl. Phys.* **B399** 169 (1993).
- [6] R.Myers and M.J.Perry, *Ann. Phys. (N. Y.)* **172** 304 (1986).
- [7] G.T.Horowitz and A.Strominger, *Nucl. Phys.* **B360** 197 (1991).
- [8] W.Israel, *Commun. Math. Phys.* **8** 245 (1968).
S.W.Hawking, *Commun. Math. Phys.* **25** 152 (1972).
- [9] R.Gregory and R.Laflamme, *Phys. Rev. D***37** 305 (1988).
- [10] C.V. Vishveshwara, *Phys. Rev. D***1** 2870 (1970).
- [11] P.Bizon and R.Wald, *Phys. Lett.* **267B** 173 (1991).
- [12] B. Whitt, Ph.D Thesis, Cambridge, 1988.
- [13] R.Gregory and R.Laflamme, *Phys. Rev. Lett.* **70** 2837 (1993).
- [14] S.W.Hawking and G.F.R.Ellis, *The Large Scale Structure of Spacetime* (Cambridge University Press 1973).
- [15] R.M.Wald *General Relativity* (Chicago University Press).
- [16] J.Bardeen, *Phys. Rev. D***22** 1882 (1980).
- [17] W.H.Press, B.P.Flemming, S.A.Teukolsky and W.T. Vetterling, *Numerical Recipes* (Cambridge University Press 1988).
- [18] R.Gregory and R.Laflamme, The stability analysis of the extremally charged black string, in preparation.
- [19] A.Bogojevic and L.Perivolaropoulos, *Mod. Phys. Lett. A***6** 369 (1991).

[20] K.Lee and E.Weinberg, *Phys. Rev. Lett.* **68** 1100 (1992).

[21] M.Bowick, S.Giddings, J.Harvey, G.Horowitz and A.Strominger, *Phys. Rev. Lett.* **61** 2823 (1988).

Appendix A. Gauge invariant variables.

One of the worries of results in linear perturbations of a system with gauge invariance is to insure that the result are not artifact of the particular gauge chosen. One easy way to deal with this problem is to use gauge invariant variables. In this appendix we explain how to construct these variables.

Under a coordinate transformation

$$x^a \rightarrow x^a + \xi^a \quad (A.1)$$

we have the following transformation of the fields

$$\begin{aligned} h^{ab} &\rightarrow h^{ab} + \xi^{(a;b)} \\ \phi &\rightarrow \phi + \xi^a D_a \phi \end{aligned} \quad (A.2)$$

$$\delta F_{a_1 \dots a_{D-2}} = F_{a_1 \dots a_{D-2}; d} \xi^d + (D-2) F_{d[a_2 \dots a_{D-2} \xi_{a_1}^d]$$

In the metric given in eq. (2.3-2.5), they have the explicit form (for $V_+ = (r - r_+)$ and $V_- = (r - r_-)$)

$$h^{tt} \rightarrow h^{tt} - \frac{V_-}{V_+} \Omega \xi^t - \frac{(r_+ - r_-)}{2V_+^2} \xi^r \quad (A.3a)$$

$$h^{rr} \rightarrow h^{rr} + \frac{V_- V_+}{2} \xi^{r'} + \frac{(-rr_- - rr_+ + 2r_+ r_-)}{2r^3} \xi^r \quad (A.3b)$$

$$h^{rt} \rightarrow h^{rt} + \frac{V_- V_+}{2r^2} \xi^{r'} - \frac{V_-}{2V_+} \Omega \xi^r \quad (A.3c)$$

$$h^{\theta\theta} \rightarrow h^{\theta\theta} + \frac{\xi^r}{r^3} \quad (A.3d)$$

$$h^{zt} \rightarrow h^{zt} - \frac{V_-}{2V_+} \Omega \xi^z + \frac{i\mu}{2} \xi^t \quad (A.3e)$$

$$h^{zr} \rightarrow h^{zr} + \frac{V_- V_+}{2r^2} \xi^{z'} + \frac{i\mu}{2} \xi^r \quad (A.3f)$$

$$h^{zz} \rightarrow h^{zz} + \frac{i\mu}{2} \xi^z \quad (A.3g)$$

It is easy to show that the linear combination

$$Y = h^{zr} + i \frac{V_- V_+}{2\mu r^2} h^{zz'} - \frac{i\mu r^3}{2} h^{\theta\theta} \quad (A.4)$$

is indeed invariant under gauge transformation. However this gauge invariant variable is not well defined when $\mu = 0$, a case under consideration in section 3. A more complicated variable

$$\begin{aligned} X = & h^{tr} + \frac{(r_+ - R_-)V_+}{2r^2 V_- \Omega} h^{tt} + \frac{V_+^2}{2r^2 \Omega} h^{tt'} \\ & - \frac{(r_+ - r_-)(-r^2 + 2rr_+ + 2rr_- - 3r_+ r_-)}{V_+ V_- \Omega} K + \frac{(r_+ - R_-)r}{4\Omega} K' - \frac{V_- \Omega r^3}{2V_+} K \end{aligned} \quad (A.5)$$

can be used to investigate the $\mu \rightarrow 0$ limit in the chargeless case. In this limit we get $X \propto \mu$ as can be shown from the asymptotic region ($r \rightarrow \infty$ and $r \rightarrow r_+$) discussed in section 3. This analysis shows that the perturbations described in sections 3-5 describe real physical instabilities except in the case $\mu = 0$.

Appendix B) Background Quantities.

The background metric is given explicitly by the following:

$$\begin{aligned} g_{tt} &= - \frac{r^{D-3} - r_+^{D-3}}{r^{D-3} - r_-^{D-3}} \\ g_{rr} &= \frac{r^{2(D-3)}}{(r^{D-3} - r_-^{D-3})(r^{D-3} - r_+^{D-3})} \\ g_{\theta_1 \theta_1} &= r^2 \\ g_{\theta_\alpha \theta_\beta} &= r^2 \delta_{\alpha\beta} \prod_{n=1}^{\alpha} \sin^2 \theta_n \quad \text{for } \alpha > 1 \\ g_{ij} &= \delta_{ij} \end{aligned} \quad (B.1)$$

where α, β run from 1 to $D - 2$, and represent the angular variables. The ‘matter’ content has solution:

$$\Phi = -\frac{1}{2} \ln \left\{ 1 - \left(\frac{r_-}{r} \right)^{D-3} \right\} \quad (B.2)$$

$$F = Q\epsilon_{D-2}, \quad Q^2 = \frac{(D-3)}{2}(r_+r_-)^{D-3} \quad (B.3)$$

The form of the perturbation is

$$g^{ac}g^{bd}\delta g_{cd} = h^{ab} = e^{\Omega t + i\mu_i x^i} \begin{bmatrix} H^{ij}(r) & H^{it} & H^{ir} & 0 & 0 & \dots \\ H^{tj} & H^{tt} & H^{tr} & 0 & 0 & \dots \\ H^{rj} & H^{tr} & H^{rr} & 0 & 0 & \dots \\ 0 & 0 & 0 & K(r) & 0 & \dots \\ 0 & 0 & 0 & 0 & K/\sin^2\theta & \dots \\ \dots & \dots & \dots & \dots & \dots & \dots \end{bmatrix} \quad (B.4)$$

$$\delta\Phi = e^{\Omega t + i\mu_i x^i} f(r) \quad (B.5)$$

and finally

$$\delta F = 0 \quad (B.6)$$

The non-zero elements of the Riemann tensor are given by:

$$R^t{}_{rtr} = \frac{(D-3)}{2r^2} \frac{\left[\left(\frac{r_+}{r}\right)^{D-3} - \left(\frac{r_-}{r}\right)^{D-3} \right] \left[D-2 - \left(\frac{r_-}{r}\right)^{D-3} \right]}{\left[1 - \left(\frac{r_+}{r}\right)^{D-3} \right] \left[1 - \left(\frac{r_-}{r}\right)^{D-3} \right]^2} \quad (B.7a)$$

$$R^t{}_{\theta_a l p h a t \theta_\beta} = \frac{-(D-3)}{2r^2} \left[\left(\frac{r_+}{r}\right)^{D-3} - \left(\frac{r_-}{r}\right)^{D-3} \right] g_{\alpha\beta} \quad (B.7b)$$

$$R^{\theta_\alpha}{}_{r\theta_\beta r} = \frac{-(D-3)}{2r^2} \frac{\left[\left(\frac{r_+}{r}\right)^{D-3} + \left(\frac{r_-}{r}\right)^{D-3} - 2\left(\frac{r_+r_-}{r^2}\right)^{D-3} \right]}{\left[1 - \left(\frac{r_+}{r}\right)^{D-3} \right] \left[1 - \left(\frac{r_-}{r}\right)^{D-3} \right]} \delta_\beta^\alpha \quad (B.7c)$$

$$R^{\theta_\alpha}{}_{\theta_\beta \theta_\gamma \theta_\delta} = \frac{1}{r^2} \left[\left(\frac{r_+}{r}\right)^{D-3} + \left(\frac{r_-}{r}\right)^{D-3} - \left(\frac{r_+r_-}{r^2}\right)^{D-3} \right] (\delta_\gamma^\alpha g_{\beta\delta} - \delta_\delta^\alpha g_{\beta\gamma}) \quad (B.7d)$$

Ricci tensor

$$R_{tt} = -\frac{(D-3)^2}{2r^2} \left(\frac{r_-}{r}\right)^{D-3} \frac{\left[\left(\frac{r_+}{r}\right)^{D-3} - \left(\frac{r_-}{r}\right)^{D-3} \right] \left[1 - \left(\frac{r_+}{r}\right)^{D-3} \right]}{\left[1 - \left(\frac{r_-}{r}\right)^{D-3} \right]^2} \quad (B.8a)$$

$$R_{rr} = \frac{(D-3)\left(\frac{r_-}{r}\right)^{D-3} \left[(3D-7)\left(\frac{r_+}{r}\right)^{D-3} + (D-1)\left(\frac{r_-}{r}\right)^{D-3} - 2(D-2)\left(1 + \left(\frac{r_+r_-}{r^2}\right)^{D-3}\right) \right]}{2r^2 \left[1 - \left(\frac{r_-}{r}\right)^{D-3} \right]^2 \left[1 - \left(\frac{r_+}{r}\right)^{D-3} \right]}$$

(B.8b)

$$R_{\theta_\alpha\theta_\beta} = \frac{(D-3)}{r^2} \left(\frac{r_-}{r}\right)^{D-3} g_{\alpha\beta} \quad (B.8c)$$

Ricci scalar

$$R = \frac{(D-3)\left(\frac{r_-}{r}\right)^{D-3}}{r^2\left(1 - \left(\frac{r_-}{r}\right)^{D-3}\right)} \left[-(D-4) - (D-2)\left(\frac{r_+}{r}\right)^{D-3}\left(\frac{r_-}{r}\right)^{D-3} + (2D-5)\left(\frac{r_+}{r}\right)^{D-3} - \left(\frac{r_-}{r}\right)^{D-3} \right] \quad (B.9)$$

The gauge conditions component by component are given by the transverse conditions $\nabla_a \bar{h}^{ab} = 0$. For simplicity in what follows, we will write $R_\pm = (r_\pm/r)^{D-3}$

$$\begin{aligned} \nabla_a \bar{h}^{at} &= \frac{\Omega h^{tt}}{2} + \frac{[(D-2) - (3D-8)R_- - R_+ + (2D-5)R_-R_+]}{r(1-R_+)(1-R_-)} h^{tr} + i\mu_i h^{ti}(r) + \frac{\Omega}{2(1-R_+)^2} h^{rr} \\ &+ \frac{(D-2)\Omega r^2(1-R_-)}{2(1-R_+)} K + \frac{\Omega(1-R_-)}{2(1-R_+)} h^i + h^{tr'}(r) = 0 \end{aligned} \quad (B.10a)$$

$$\begin{aligned} \nabla_a \bar{h}^{ar} &= \frac{(D-3)(1-R_+)(R_+ - R_-)}{r(1-R_-)} h^{tt} - \frac{[(D-2) - (2D-5)R_-]}{r(1-R_-)} h^{rr} \\ &- 2(D-2)r(1-R_-)(1-R_+)K + \Omega h^{tr} + i\mu_i h^{ri} + \frac{(1-R_+)^2}{2} h^{tt'} \\ &+ \frac{h^{rr'}}{2} - \frac{(D-2)}{2}(1-R_-)(1-R_+)r^2 K' - \frac{(1-R_-)(1-R_+)}{2} h_i^{i'} = 0 \end{aligned} \quad (B.10b)$$

$$\nabla_a \bar{h}^{a\theta} = \nabla_a \bar{h}^{a\phi} = 0 \quad (B.10c)$$

$$\begin{aligned} \nabla_a \bar{h}^{ai} &= \frac{i\mu_i(1-R_+)}{2(1-R_-)} h^{tt} + \Omega h^{ti} - \frac{i\mu_i}{2(1-R_+)(1-R_-)} h^{rr} + \frac{[(D-2) - (2D-5)R_-]}{r(1-R_-)} h^{ri} \\ &- \frac{(D-2)}{2} i\mu_i r^2 K + i\mu_j h^{ij} - \frac{i\mu_i}{2} h_j^j + h^{ri'}(r) = 0 \end{aligned} \quad (B.10d)$$

Finally we give equations for the perturbations h^{ab} , the trace $h = g_{ab}h^{ab}$ and $f = \delta\Phi$.

• h^{tt}

$$0 = \frac{-(D-3)h^{rr}}{2r^2} \frac{(R_+ - R_-)[2(D-2) - (D-1)R_+ - (3D-7)R_- + 2(D-2)R_+R_-]}{(1-R_+)^3(1-R_-)}$$

$$\begin{aligned}
& +h^{tt} \left(-\mu^2 - \frac{\Omega^2(1-R_-)}{(1-R_+)} + \frac{(D-3)^2}{2} \frac{(R_+ - R_-)^2}{(1-R_+)(1-R_-)} \right) + (D-2)(D-3)K \frac{(R_+ - R_-)(1-R_-)}{(1-R_+)} \\
& - \frac{2(D-3)\Omega}{r} \frac{(R_+ + R_+R_- - 2R_-)}{(1-R_+)^2} h^{tr} - 4\Omega^2 \frac{(1-R_-)^2}{(1-R_+)^2} f(r) + \frac{2(D-3)}{r} \frac{(1-R_-)(R_+ - R_-)}{(1-R_+)} f' \\
& + [(D-2) - (2D-5)R_- + (2D-7)R_+ - (D-4)R_+R_-] \frac{h^{tt'}}{r} + (1-R_+)(1-R_-)h^{tt''} \quad (B.11a)
\end{aligned}$$

• h^{rr}

$$\begin{aligned}
0 &= \frac{-(D-3)h^{tt}}{2r^2(1-R_-)} (1-R_+)(R_+ - R_-) [2(D-2) - (D-1)(R_+ + R_-) + 2R_+R_-] \\
& + h^{rr} \left(-\mu^2 - \frac{\Omega^2(1-R_-)}{(1-R_+)} - \frac{4(D-2)(1-2R_+ - (D-1)R_-)}{2r^2(1-R_+)(1-R_-)} \right. \\
& + \frac{(3D^2 - 26D + 47)R_-^2 + (D^2 - 10D + 17)R_+^2 + 2(2D^2 - 14D + 23)R_+^2R_-^2}{2r^2(1-R_+)(1-R_-)} \\
& \left. - \frac{2(D^2 - 5)R_+R_- + (D^2 - 12D + 23)R_+^2R_- + (5D - 13)(D-5)R_-^2R_+}{r^2(1-R_+)(1-R_-)} \right) \\
& + (D-2)K(1-R_+)(1-R_-) [2 + 2(D-2)R_+R_- - (D-1)(R_+ + R_-)] \\
& - \frac{2(D-3)\Omega(R_+ - R_-)}{r} h^{tr} + [(D-2) - (2D-5)R_+ - (4D-11)R_- + (5D-14)R_+R_-] \frac{h^{rr'}}{r} \\
& - \frac{2(D-3)}{r} (1-R_+)(1-R_-) [R_+ + R_- - 2R_+R_-] f' - 4(1-R_+)^2(1-R_-)^2 f''(r) + (1-R_+)(1-R_-) h^{rr''} \\
& \hspace{15em} (B.11b)
\end{aligned}$$

• h^{rt}

$$\begin{aligned}
0 &= \frac{\Omega(D-3)}{r} \left(h^{rr} \frac{[2R_- - R_+ - R_+R_-]}{(1-R_+)^2} - h^{tt}(R_+ - R_-) \right) - 2\Omega f \frac{(D-3)(R_+ - R_-)(1-R_-)}{r(1-R_+)} \\
& + h^{rt} \left(-\mu^2 - \frac{\Omega^2(1-R_-)}{(1-R_+)} - \frac{(D-2)[1 - 2(D-2)R_- + (D-5)R_+]}{r^2(1-R_+)(1-R_-)} \right. \\
& \left. + \frac{(D^2 - 10D + 20)R_-^2 - R_+^2 - (2D-5)R_+^2R_-^2}{r^2(1-R_+)(1-R_-)} \right)
\end{aligned}$$

$$\begin{aligned}
& + \frac{(3D^2 - 17D + 26)R_+^2 R_- - (3D^2 - 13D + 16)R_+ R_- - (2D^2 - 18D + 34)R_-^2 R_+}{r^2(1 - R_+)(1 - R_-)} \Big) \\
& + 4\Omega(1 - R_-)^2 f' + [(D - 2) - R_+ - (3D - 8)R_- + (2D - 5)R_+ R_-] \frac{h^{rt'}}{r} + (1 - R_+)(1 - R_-) h^{rt''} \\
& \hspace{15em} (B.11c)
\end{aligned}$$

• $h^{\theta\theta}$

$$\begin{aligned}
0 &= \frac{h^{rr}}{r^4} \frac{[2 - (D - 1)R_+ - (3D - 7)R_- + 2(2D - 5)R_+ R_-]}{(1 - R_+)(1 - R_-)} + \frac{(D - 3)h^{tt}}{r^4} \frac{(R_+ - R_-)(1 - R_+)}{(1 - R_-)} \\
& + K \left(-\mu^2 - \frac{\Omega^2(1 - R_-)}{(1 - R_+)} + \frac{2}{r^2} [(D - 2) + (D - 4)(R_+ + R_-) - (D^2 - 3D - 1)R_+ R_-] \right) \\
& - 4f'(r) \frac{(1 - R_+)(1 - R_-)}{r^3} + [(D + 2) - 5(R_+ + R_-) - (D - 8)R_+ R_-] \frac{K'}{r} + (1 - R_+)(1 - R_-) K'' \\
& \hspace{15em} (B.11d)
\end{aligned}$$

• h^{ti}

$$\begin{aligned}
0 &= -\frac{(D - 3)\Omega(R_+ - 2R_- + R_+ R_-)}{r(1 - R_+)^2} h^{ri} - \frac{i\mu_i(D - 3)R_-}{r(1 - R_-)} h^{tr} + 4i\mu_i\Omega f(r) \frac{(1 - R_-)}{(1 - R_+)} \\
& - h^{ti} \left(\mu^2 + \frac{\Omega^2(1 - R_-)}{(1 - R_+)} \right) + \frac{h^{ti'}}{r} (1 - R_-) [(D - 2) + (D - 4)R_+] + (1 - R_+)(1 - R_-) h^{ti''} \quad (B.11e)
\end{aligned}$$

• h^{ri}

$$\begin{aligned}
0 &= -\frac{(D - 3)\Omega(R_+ - R_-)}{r} h^{ti} - \frac{i\mu_i(D - 3)R_-}{r(1 - R_-)} h^{rr} - 4i\mu_i f'(r) (1 - R_+)(1 - R_-) \\
& + h^{ri} \left(-\mu^2 - \frac{\Omega^2(1 - R_-)}{(1 - R_+)} + \frac{(1 - R_+) [-(D - 2)(1 - (D - 1)R_-) - (2D - 5)R_-^2]}{r^2(1 - R_-)} \right) \\
& + \frac{h^{ri'}}{r} (1 - R_+)(D - 2 - (2D - 5)R_-) + (1 - R_+)(1 - R_-) h^{ri''} \quad (B.11f)
\end{aligned}$$

• h^{ij}

$$0 = -\frac{(D - 3)R_-}{r(1 - R_-)} [i\mu_i h^{rj} + i\mu_j h^{ri}] + 4\mu_i \mu_j f(r) - h^{ij} \left(\mu^2 + \frac{\Omega^2(1 - R_-)}{(1 - R_+)} \right)$$

$$+\frac{h^{ij'}}{r}[(D-2)-R_- - R_+ - (D-4)R_+R_-] + (1-R_+)(1-R_-)h^{ij''} \quad (B.11g)$$

For the trace, h,

$$0 = -2(D-2)(D-3)^2R_+R_-K - \left(\mu^2 + \frac{\Omega^2(1-R_-)}{(1-R_+)}\right)(h-4f) \\ + \frac{(1-R_-)(D-2-R_+)}{r}(h'-4f') + (1-R_-)(1-R_+)(h''-4f'') \quad (B.11h)$$

and finally for the dilaton perturbation:

$$0 = -\frac{(D-3)^2R_-(1-R_+)(R_+-R_-)}{4r^2(1-R_+)^2}h^{tt} + \frac{(D-3)(D-2)}{2}KR_-[1-(D-2)R_+] \\ - \frac{(D-3)R_-[2(D-2)-(3D-7)(R_++R_-)+2(2D-5)R_+R_-]}{4r^2(1-R_-)^2(1-R_+)}h^{rr} - f\left(\mu^2 + \frac{\Omega^2(1-R_-)}{(1-R_+)}\right) \\ + \frac{f'}{r}[(D-2)-R_+ + (D-4)R_- - (2D-7)R_+R_-] + (1-R_+)(1-R_-)f'' \quad (B.11i)$$

Captions.

Figure 1. Penrose diagram for the metric of eq.(1). All but two dimensions have been suppressed; each point of this diagram corresponds to a D-2 sphere times 10-D flat space. The line at $v = v_0$ is the initial hypersurface on which the perturbations are set. It is a Cauchy surface for the spacetime exterior of the black hole.

Figure 2. Plot of the modes H^{tt} , H^{tr} and H^{rr} as a function of $(r - r_+)$ for a regular mode. As $(r - r_+)$ becomes large all the functions decay exponentially, they reach a maximum at small $(r - r_+)$ and go to zero as $(r - r_+) \rightarrow 0$.

Figure 3. Plot of Ω as a function of μ for black strings and branes with $D = 4, \dots, 9$ and $r_+ = 2$ for which an instability has been found. The bold points correspond to value calculated numerically and the lines have been traced to guide the eye.

Figure 4. Schematic diagram of the instability. All the dimensions have been suppressed except the r and z one. The geometry is initially invariant under a z translation. The instability increases the size of the apparent horizon at some values of z and decreases it at other values. In the chargeless case it seems reasonable that the apparent horizon will brake into different parts.

Figure 5. Plot of the modes f, q, H^{zz} and H^{tz} as a function of $(r - r_+)$ for a regular mode. As $(r - r_+)$ becomes large all the functions decay exponentially, they reach a maximum at small $(r - r_+)$ and go to zero as $(r - r_+) \rightarrow 0$.

Figure 6. Plot of Ω as a function of μ for a charged 5d black string with charges corresponding to $r_- = 0, 1.0, 1.5$ and $r_+ = 2$ for which an instability has been found. The bold points correspond to value calculated numerically and the lines have been traced to guide the eye.

This figure "fig1-1.png" is available in "png" format from:

<http://arXiv.org/ps/hep-th/9404071v1>

This figure "fig1-2.png" is available in "png" format from:

<http://arXiv.org/ps/hep-th/9404071v1>

This figure "fig1-3.png" is available in "png" format from:

<http://arXiv.org/ps/hep-th/9404071v1>

Figure 1




Figure 2




Figure 3




Figure 4




Figure 5




Figure 6

